1   Partners in space: Discordant population structure between legume hosts and rhizobium

2   symbionts in their native range

4   Alex B. Riley[1], Michael A. Grillo[2], Brendan Epstein[3], Peter Tiffin[3], Katy D. Heath[1,4,+]

5
Author affiliations:

[1]University of Illinois, Department of Plant Biology, 505 S. Goodwin Ave. Urbana IL 61801

[2]Loyola University Chicago, Department of Biology, 1032 W. Sheridan Rd, Chicago, IL, 60660,

USA

[3]University of Minnesota, Department of Plant and Microbial Biology, 140 Gortner Lab, 1479

Gortner Ave, St. Paul, MN 55108

[4]University of Illinois, Carl R. Woese Institute for Genomic Biology, 1206 W. Gregory Dr.,

Urbana, IL 61801 USA

[+]Corresponding author

15
Running head: Discordant population structure in symbiosis

Keywords: coevolution, mutualism, horizontal gene transfer, divided genome

Acknowledgments: We thank Laurène Gay and Joelle Ronfort for advice and contributions to

field collection, and Amy Marshall-Colón, Julian Catchen, and Rachel Whitaker for feedback on

an early manuscript draft. We thank various sources of funding including NSF IOS-1401864 to

MAG, NSF IOS-1645875 to KDH, NSF PGRP-1856744 to PT and KDH, JGI CSP-503446 to

PT and KDH, and the Department of Plant Biology at the University of Illinois Urbana-

Champaign.

Abstract
27
Coevolution is predicted to depend on how the genetic diversity of interacting species is

geographically structured. Plant-microbe symbioses such as the legume-rhizobium mutualism are

ecologically and economically important, but distinct life history and dispersal mechanisms for

these macrobial and microbial partners, plus dynamic genome composition in bacteria, present

challenges for understanding spatial genetic processes in these systems. Here we study the model

rhizobium *Ensifer meliloti* using a hierarchically-structured sample of 191 strains from 21 sites

in the native range and compare its population structure to that of its host plant *Medicago*

truncatula. We find high local genomic variation and minimal isolation by distance across the

rhizobium genome, particularly at the two symbiosis elements pSymA and pSymB, which have

evolutionary histories and population structures similar to each other and distinct from both the

chromosome and the host. While the chromosome displays weak isolation by distance, it is

uncorrelated with hosts. Patterns of discordant population structure among elements with the

bacterial genome has implications for bacterial adaptation to life in the soil versus symbiosis,

while discordant population genetic structure of hosts and microbes might restrict local

adaptation of species to each other and give rise to phenotypic mismatches in coevolutionary

traits.

Introduction

Coevolution is fundamentally a geographically variable process. This understanding has become widely accepted over the past several decades, most prominently with the inception of Thompson's geographic mosaic theory of coevolution (Thompson 1994, 2005). Coevolving partners can vary in their distributions, abundance, and genetic structure across the landscape and over time (Laine 2005; Thompson 2005; Tack et al. 2014; Carlsson-Granér and Thrall 2015; Fernandes et al. 2019). The GMTC posits that such geographic structure among interacting partners, coupled with isolation by distance, genetic drift, and spatially divergent natural selection will influence the nature of coevolutionary interactions. While there is abundant evidence for the core predictions of the GMTC (*i.e.,* selection mosaics, coevolutionary hot/cold spots, variable trait remixing) it is surprising that comparative analysis of population genetic structure among interacting partners has received relatively little attention (but see Caldera and Currie 2012; Baums et al. 2014; Strobel et al. 2016; Harrison et al. 2017).

Indeed, underlying population genetic structure can, in part, determine the outcomes of coevolution. For example, alignment in genetic structure between interacting species can allow for reciprocal coevolutionary selection (*i.e.,* coevolutionary hotspots) and result in phenotype matching of coevolutionary traits; conversely, discordant population structure, due to different levels of gene flow or other processes, can limit the ability of each species to locally adapt to the other and give rise to coevolutionary trait mismatches (Gandon et al. 1996a; Thompson 2005; Fernandes et al. 2019). Thus, understanding the comparative population genetic structure of interacting species provides key perspective on whether traits are expected to coevolve and how coevolutionary trait variation is maintained through time and space (Heath and Stinchcombe 2014; Hollowell et al. 2016a; Stoy et al. 2020).

Coevolutionary interactions typically occur between species with distinct life histories, range sizes, mating systems, and modes of dispersal – all of which will impact their underlying population genetic structure (Thrall et al. 2007; Revillini et al. 2016; Pita et al. 2018). In general, the majority of coevolutionary research has focused on systems involving eukaryotes and on antagonistic coevolution, with mutualisms between macro and micro-organisms receiving less attention. The investigation of host-microbe interactions poses various logistical and methodological challenges. One challenge for understanding the (co)evolution of bacterial symbionts in nature is the difficulty of isolating populations of a single species from the environment - particularly for soil bacteria in hyper-diverse communities (Heath and Grillo 2016). Plant-microbe mutualisms play critical roles in community and ecosystem processes, yet remain poorly understood in terms of their coevolutionary dynamics, origins, and maintenance (Reynolds et al. 2003; Berg 2009; Revillini et al. 2016).

To add to this complexity, many bacterial genomes are multipartite (Harrison et al. 2010; diCenzo and Finan 2017), including non-chromosomal replicons containing variable amounts of genetic information – ranging from small facultative plasmids that are often lost or transferred to large obligate segments containing core genes (often as megaplasmids or chromids; Harrison et al. 2010). Such divided genomes might represent an adaptation allowing functional division between replicons (reviewed by diCenzo and Finan 2017). Multipartite genome organization presents a challenge in the study of bacterial (co)evolution because different elements may move between individuals independent of each other via horizontal gene transfer, resulting in distinct evolutionary histories and population genetic processes across different elements. These differences are compounded by the fact that these elements can also have widely differing rates of evolution and patterns of recombination due to their gene content ranging from facultative to

obligate, leading to higher rates of evolutionary change in plasmids not required for the survival of the individual in its environment (Cooper et al. 2010; Epstein et al. 2012, 2014; Epstein and Tiffin 2021). We still lack a general understanding of whether and how landscape-scale processes vary among the elements in divided bacterial genomes.

The legume-rhizobium mutualism presents a tractable model system for studying microbial symbiosis due to the economic and ecological importance of biological nitrogen (N) fixation, the relative ease of isolating populations, and genetic resources for several legume-rhizobium partnerships (Kaneko et al. 2002; Yates et al. 2015). In this symbiosis, rhizobia form nodules on plant roots, wherein they fix atmospheric N in exchange for carbon derived from plant photosynthesis; however, rhizobia inhabit the soil as saprophytes when not in symbiosis with a host, thereby experiencing profoundly different environmental conditions and likely selective pressures (Burghardt 2020). In rhizobia with divided genomes (e.g., *Rhizobium* and *Ensifer* spp.), genes for symbiosis are carried on symbiosis elements (one or more megaplasmids or chromids), while the chromosome is thought to be responsible for core metabolic functions. These elements can have distinct evolutionary histories, diversity levels, and recombination rates (e.g., Bailly et al. 2011; Epstein et al. 2012, 2014; Cavassim et al. 2020; Epstein and Tiffin 2021); however, addressing hypotheses about whether these elements play distinct roles in adaptation to hosts versus soil life requires an understanding of how patterns of variation at the symbiosis elements correspond to the chromosome, and if they align with the host within and among natural populations.

Here we examine population structure in *Ensifer meliloti* (Becker et al. 2009; Tang et al. 2014), the rhizobium symbiont of the model legume *Medicago truncatula* (hereafter "*Medicago*"). *Medicago* is a self-fertilizing annual native to the Mediterranean region of Europe

(Bonnin et al. 1996; Siol et al. 2008). Research involving *Medicago* population genetics has identified a pattern of isolation by distance, and structure that is differentiated at the population and regional levels (Bonnin et al. 1996; Ronfort et al. 2006; Siol et al. 2008; Bonhomme et al. 2015; Grillo et al. 2016). Here we use whole genome sequences for 191 isolates of *E. meliloti* sampled from 21 sites in the native range to characterize the population structure of this model mutualist and compare it to the host by reanalyzing *Medicago* sequence data from Grillo et al. (2016) to directly compare population structure in the two species. We ask: 1) how is genetic variation structured in *E. meliloti,* and is there evidence for a pattern of isolation by distance; 2) do the three genomic elements of the rhizobium *E. meliloti* share common patterns of evolutionary history and population structure; and 3) do *Medicago* hosts and their rhizobium symbionts have correlated population genetic structure?

Methods:

Study system:

Ensifer (formerly *Sinorhizobium*) *meliloti* is a rhizobium species in the Alphaproteobacteria (Young and Haukka 1996) that forms N-fixing nodules on the roots of multiple species in the genus *Medicago*, and is one of two *Ensifer* species that forms root nodules on *M. truncatula* (Zribi et al. 2004). The genome of *E. meliloti* is ~6.79Mb, divided between three genomic elements: the chromosome (3.69Mb), megaplasmid pSymA (1.41Mb), and chromid pSymB (1.69Mb), as well as smaller plasmids in some strains (Galibert et al. 2001, Nelson et al. 2018). Metabolic modeling, along with genetic manipulation of the *E. meliloti* genome, have shown that gene content differs functionally between the three genomic elements (Galibert et al. 2001; diCenzo et al. 2014). The chromosome is predicted to carry primarily genes related to core

metabolic function of individuals in bulk soil, while pSymA carries the majority of genes

required for symbiotic N fixation, and pSymB carries primarily genes thought to be important for

life in rhizosphere environments (diCenzo et al. 2014).

Sample collection: We isolated *E. meliloti* strains from 21 sites in the native range of *Medicago*

using a hierarchical sampling design with populations ranging from 1 to approximately 1350 km

apart (Fig. S1; Table S1). Of the 21 sites, eight corresponded to the *Medicago* sites studied in

Grillo *et al*. (2016). At each site we collected soil surrounding the top six inches of the root

system from unearthed host plants. To avoid cross-contamination within each site, the sampling

shovel was wiped clean of excess soil between samples and was pierced into the ground adjacent

to a plant numerous times before sampling soil. Between sampling locations, the shovel was

sterilized with dilute bleach. Soil samples were stored in double wrapped plastic bags and kept at

4°C prior to isolating cultures.

Ensifer strains were isolated or "trapped" in the laboratory from the field samples

following standard protocols (Vincent 1970; Heath 2010). In brief, *Medicago* seeds were nicked

with a razor blade, surface sterilized with 30% bleach, rinsed with sterile water, and imbibed in

sterile water for approximately 30 minutes. Seeds were then directly sown into a given soil

sample housed in a sterilized, fully self-contained Magenta box (see Brown et al. 2020 for

details). Magenta boxes were randomly placed in a temperature-controlled grow room (23° C)

under artificial light set to 12-h days. After four weeks, plants were harvested, and the soil was

washed away from the roots. Individual nodules were removed with forceps and were surface

sterilized by soaking in 30% bleach for 10 minutes and then rinsed with sterilized water. Surface

sterilized nodules were crushed with sterilized forceps and streaked on tryptone-yeast (TY)

media plates. Plates were incubated at 30°C for 48 hours, at which point sterilized glass stir rods were used to streak samples were streaked on to TY plates which were again incubated at 30°C. Individual colonies were then picked and grown in liquid TY media, and these pure cultures were stored in cryotubes in 50% TY 50% glycerol and stored at -80°C. Given variation among host genotypes in rhizobium infection rates due to G x G interactions (Heath and Tiffin 2009; Batstone et al. 2017), we used 24 different *Medicago* host genotypes to trap rhizobia. Because both *E. meliloti* and *E. medicae* infect the roots of *Medicago* in the native range, we used a post-PCR restriction enzyme (RsaI) digestion of the 16S to assign strains to species (following Biondi et al. 2003). Ultimately, we isolated 191 strains of *E. meliloti* presented in the current study. We hereafter refer to *E. meliloti* as simply "*Ensifer*" or "symbiont".

Sequencing: We extracted DNA from cultures of *Ensifer* grown in liquid TY media using the Qiagen DNeasy kit (Hilden, Germany) and sent samples to the DOE Joint Genome Institute (JGI) for sequencing (Berkeley, CA, USA). JGI prepared a paired end sequencing library for each strain, and sequenced samples on an Illumina HiSeq-2500 1TB platform (101nt read length; Illumina, Inc., San Diego, CA, USA). Of the 199 strains submitted to JGI, we received high quality whole genome sequences for 166. We re-grew the remaining 33 strains from frozen cultures (as above), extracted DNA using the Zymo Quick-DNA kit for Fungi or Bacteria (Irvine, CA, USA). These samples were sequenced (2 X 150 or paired end 150 nt read length) on the Novaseq 6000 platform (Illumina, Inc, San Diego, CA, USA) by the Roy J. Carver biotechnology center at the University of Illinois at Urbana-Champaign (USA). We successfully recovered quality genome sequences from 25 of these 33 isolates, for a total of 191 *E. meliloti* strains sequenced.

Genome assembly, annotation, and SNP calling: To ensure high quality SNP calling and genome assembly, we trimmed PCR adaptors and removed PCR duplicates and PhiX contamination using HT-Stream (github.com/s4hts/HTStream), followed by further adaptor removal, removal of bases with quality scores < 30 from the ends of reads, and removal of reads < 80 bp long with TrimGalore! (github.com/FelixKrueger/TrimGalore). For calling SNPs in core genes (genes present in > 80% of strains), we aligned reads to the *E. meliloti* reference genome USDA1106 using BWA with default settings (Li and Durbin 2009). We then used Freebayes (Garrison and Marth 2012) to identify haplotype variants, which we split into SNPs using VCFtools (Danecek et al. 2011). We found 491,277 variable SNPs with variant qualities above 20. We then filtered SNPs to retain only those with depth values between 20 and 230, minor allele frequencies ≥ 0.009 (*i.e.,* present in at least two strains), and that were present in at least 80% of individuals, 72,311 SNPs remained after applying these filter (chromosome: 34,689 SNPs, pSymA: 15,162 SNPs, pSymB: 22,460 SNPs; see Table S2). For variable gene content, we assembled genomes *de novo* using SPADES (Bankevich et al. 2012) with default parameters, followed by annotation using PROKKA (Seemann 2014) and scanned these annotations for presence-absence variants using default settings in ROARY (Page et al. 2015).

Phylogenetic and Statistical Analyses: To explore the population structure of the three genomic elements of *Ensifer,* we first used principal components analysis (PCA). We used the glPca function in the adegenet (Jombart and Ahmed 2011) library in R on a random subsample of 15k SNPs (to equilibrate dataset size) for each element of the *Ensifer* genome to naively cluster individuals by genome-wide similarity. We then plotted the positions of individuals along the

first three axes of variation (using ggplot2; Fig. S2) to visually compare how genetic variation is

distributed across elements (Wickham 2016). To quantify genetic differences we calculated

individual and population-level Dxy (Nei 1972) using the stamppNeisD function from the

package StAMPP (Pembleton et al. 2013) on all variants for each element. To test whether these

patterns of genetic distance among *Ensifer* strains were congruent across the three genome

elements, we used these matrices of individual-based distance metrics in pairwise Mantel tests

comparing the three *Ensifer* genome elements (chromosome vs. pSymA, chromosome vs.

pSymB. pSymA vs. pSymB). Mantel tests were implemented in the R package ade4 (Dray and

Dufour 2007). To explore geographic structure in the variable genome (*i.e.,* genes present in only

some strains), we performed an additional genetic PCA, as above, using the matrix of *Ensifer*

gene presence-absence variants (91,840 genes) and plotted the first three principal components

for visual inspection.

For each of the three elements, we built phylogenies based on the same random

subsample of 15k core genome SNPs used for PCA. We used the neighbor joining (nj) function

in the R package ape (Paradis and Schliep 2019) with 1000 bootstrap replicates and visualized

trees in FigTree (v.1.4.4) (Rambaut 2018). Based on these preliminary results, we treated each

element independently for remaining analyses.

We used AMOVA (poppr.amova) with clone correction and 10,000 random permutations

(Kamvar et al. 2014) to partition the genetic variance among individuals, sites, and regions, as

well as to assess the statistical significance of these levels of spatial genetic structure. To test the

hypothesis of isolation by distance, we calculated Pearson correlations between individual-level

D_{XY} and geographic distances between sampling sites.

To compare the spatial genetic structure of rhizobia and host plants, we reanalyzed RAD-seq data from the 192 *Medicago* genotypes studied in Grillo *et al* (2016). We first called SNPs using Stacks with default parameters (Catchen et al. 2013), then filtered the resulting variants using VCFtools (Danecek et al. 2011) to ensure that all variants were present in at least 80% of lines, had minor allele frequencies > 0.05, and were > 5kb apart. We performed AMOVA, calculated individual and population-based D_{XY}, and tested for isolation by distance in *Medicago* as detailed above for *Ensifer*. For the subset of eight sites for which we had both hosts and symbionts, we used Mantel tests to ask whether the three matrices of pairwise population D_{XY} values from symbionts (chromosome, pSymA, and pSymB) were correlated with that of the host, to test for congruent population genetic structure between hosts and symbionts.

Results

To address the extent of phylogenetic congruence between the chromosome, pSymA, and pSymB of *Ensifer*, we first used pairwise Mantel tests of whether the matrices of genetic distances were correlated. At the individual level, we found non-significant correlations between the chromosome and pSymA and between the chromosome and pSymB, while the two symbiosis elements (pSymA and pSymB) were more strongly correlated (left column, Fig. 1), indicating distinct evolutionary histories of the chromosome versus pSymA and pSymB. Neighbor joining trees also reveal distinct evolutionary histories among the three genome elements. The topology of the chromosomal tree showed five tightly clustered groups of individuals, with almost all strains from Corsica closely related and forming a distinct group (orange cluster; Fig. 2a) and strains from Spain somewhat interspersed with those from France mostly in the blue and yellow clusters. The diversity of strains found in mainland France (the best sampled region) included

representatives from each major chromosomal lineage; indeed, pink and purple clusters were found only in mainland France (except strain 710A from Corsica), mostly on the western Mediterranean coast (Fig. S1; Table S1).

Using these same colors (corresponding to the chromosomal clusters) to annotate the pSym trees (Fig. 2b; Fig. S3) facilitates visualization of differences in tree topology among the three elements, indicative of plasmid transfer (whole or partial) among chromosomal lineages. The internal branches of both symbiotic element trees reveal greater diversity, with chromosomal lineages spread across diverse pSym lineages in both trees (colored outer rings in Fig. 2b; Fig. S3). Some strains were closely related at all three elements of their genomes; for example, the group of closely related chromosomal lineages in pink from the western coast of France appear together in all three trees, indicating little pSym element recombination among these and other lineages (strain 78 in the pSymB tree is a notable exception). On the other hand, the other chromosomal clusters (purple, orange, and yellow clusters in Fig. 2a), appear throughout the pSymA and pSymB tree (see same colors interspersed; Fig. 2b; Fig. S3), indicating that these chromosomal lineages are found with diverse pSym genotypes. For example, the tightly clustered group of closely related chromosomal lineages in orange were found with lineages from across both the pSymA and pSymB trees.

Next, we used genetic PCA to explore population structure and AMOVA to partition the genome-wide genetic variation at the within-population, among-population, and among-region scales. Genetic PCA did not reveal strong differentiation in *Ensifer*, even at the regional scale, in either the core genome SNPs or variable gene content (Fig. S2; Fig. S4). Nevertheless we found significant spatial genetic structure in the core genome at all scales investigated using AMOVA (Table 1), but that the patterns varied among the three *Ensifer* elements. For all three, we found

considerable variation (66-76%) maintained within populations, compared to the among-population and among-region scales (Table 1). All three among-element correlations were stronger at the among-population level (right column, Fig. 1), revealing substantial congruence between pSymA and pSymB, and weakly correlated population structure between the chromosome and both symbiosis elements (right column, Fig. 1). The discrepancy between individual and population level correlations is likely due to high within population diversity (Table 1) and several populations with near-zero genetic distances (*e.g.,* within Spain and France, and between them Table S3).

The *Ensifer* chromosome showed the strongest differentiation at the regional scale (*i.e.,* among Spain, France, and Corsica; Table 1). This finding is likely driven by Corsica, as short branch lengths were found among most Corsica strains on the chromosomal tree (within the orange cluster; Fig 2a) and genetic distances were high between Corsica populations and the rest of the range (Table S3). The chromosome was also the only *Ensifer* element to exhibit significant isolation by distance (IBD) at this spatial scale (Fig 3). Despite overall IBD, there was extremely large variation in the genetic distances between *Ensifer* isolated from even the most distant locations (*i.e.,* D_{XY} distances ranged from 0-0.5 even when strains were found approximately 1350 km apart), consistent with the finding of high proportions of genetic variation maintained at the site level in the face of among-region population structure (Table 1). The two symbiotic elements in *Ensifer* (pSymA and pSymB) were less structured at the among-region and among-population scales (Table 1), and neither exhibited significant IBD at our sampling scale (Fig. 3). Thus although populations differed in genetic composition (significant structure in Table 1), *Ensifer* strains from distant populations up to 1350 km apart were often as closely related to each other as strains from the same sampling site.

Hosts and symbionts had mismatched population genetic structure. We found no significant correlations between the pattern of among-population differentiation (D_{XY}) between *Medicago* hosts and any element in the *Ensifer* genome (all $p > 0.5$; Fig. S5). While both the host and *Ensifer* chromosome exhibit patterns of isolation by distance (see above), their spatial patterns of genetic variation were distinct; while host plants primarily clustered along axes that separated Spain from France and Corsica (Fig. S2; Grillo et al. 2016 and references therein), the *Ensifer* chromosome most clearly clustered into two groups (mainland Europe versus Corsica along PC1, Fig. S2). The lack of IBD and little regional structure in pSymA and pSymB indicate that host plants at a given site have the potential to interact with the diversity of symbiosis plasmids from across this portion of the species range.

Discussion

Evolution, and thus coevolution, is shaped by the interaction of selection with demographic processes such as gene flow occurring across a species range; therefore, understanding the maintenance of genetic variation, the evolution of mutualistic traits, and host-microbe interactions requires that we consider individuals and species in their spatial context. Here we used a hierarchically-structured sample of legume hosts and rhizobial symbionts to show: 1) the elements of the bacterial genome (chromosome versus pSymA and pSymB) have distinct evolutionary histories and population genetic structure at the local and regional scales, 2) host plants have more spatial genetic structure than rhizobia, particularly when compared to the two symbiotic elements (pSymA and pSymB), and 3) patterns of population structure between hosts and symbionts are not correlated across this portion of their native range. We discuss the implications of these main results below.

Spatial genetic processes at the bacterial chromosome and symbiotic elements:

Unlike macrobial systems, the population structure and dispersal abilities of microbial systems remain poorly understood (Martiny et al. 2006; Green et al. 2008; Hanson et al. 2012; Chase et al. 2019; VanInsberghe et al. 2020). Recent advances in sequencing technology have revealed considerable variation in dispersal ability among microbial taxa (Locey and Lennon 2016). Based on metagenomics datasets, rhizobia have many characteristics associated with large geographic ranges and, by extension, large dispersal ability (*e.g.,* being Proteobacteria, plant-associated or soil-borne taxa, and large genome sizes; Choudoir et al. 2018). However, metagenomic studies of microbial communities are limited in their resolution; therefore, population genomic data (Hoetzinger et al. 2021), coupled with spatially-structured samples of both within- and among-population variation (Whitaker and Banfield 2006), are required for inferring population-level processes such as gene flow. It is this population scale that is most relevant to local adaptation and coevolution (Thompson 2005; Hoeksema and Forde 2008; Kraemer and Boynton 2017). Like other diverse microbial systems (Vos and Velicer 2008; Chase et al. 2019; Hoetzinger et al. 2021), *Ensifer* displays both isolation by distance (chromosome) and population genetic structure (all elements); nevertheless, we found extremely closely-related strains up to 1350 km apart – indicating at least some long-distance dispersal. Large dispersal ability, together with the observation of abundant local phenotypic (Heath and Tiffin 2009; Heath 2010) and genomic (Bailly et al. 2006, 2011) variation for symbiosis, suggests that most of the variation relevant to both bacterial adaptation and coevolution can be found at relatively small spatial scales.

At the regional scale sampled here, we identify distinct evolutionary histories and population genetic structures between the chromosome and the symbiosis elements, and overall alignment in population structure between the symbiosis elements. Previous work has shown divergent evolutionary histories of the three elements in this species (Bailly et al. 2006; Galardini et al. 2013; Epstein et al. 2014; Nelson et al. 2018), as well as in other rhizobium taxa with separate symbiosis plasmids (Young et al. 2006; Koppell and Parker 2012; Kumar et al. 2015; Klinger et al. 2016; Pérez Carrascal et al. 2016; Cavassim et al. 2020) or symbiosis gene regions (Hollowell et al. 2016b; Porter et al. 2019). Our analyses support these findings while also revealing how the three genomic elements differ at the individual, among-population, and regional scale using a hierarchically-structured spatial sample of within and among-site variation. Using this approach allowed us to resolve distinct fine-scale spatial genetic processes at individual elements, namely geographically limited chromosomal lineages (e.g., Corsica), but little evidence of this at the symbiosis elements.

One hypothesis for the existence of distinct elements in bacteria is to break up genetic correlations among traits that are important during different phases of the life cycle and/or establish linkage between adaptive gene complexes (diCenzo and Finan 2017). Based on functional genetics and metabolic models (Galibert et al. 2003; diCenzo et al. 2014; diCenzo and Finan 2017), the three major genome elements are thought to play somewhat distinct roles in the life history of *Ensifer:* chromosome (basic metabolism), pSymA (nodulation and N fixation), and pSymB (rhizosphere). The traits, and potentially genes, that underlie fitness in the soil, rhizosphere, and nodules are likely distinct (Friesen and Mathias 2010; Sachs et al. 2011; Heath and Stinchcombe 2014; Burghardt 2019); having these processes segregated into distinct genome elements might allow the buildup of coadapted gene complexes within elements while

facilitating independent adaptation to host and abiotic conditions on separate elements. One

particularly interesting geographic disjunction that distinguishes the chromosome from the other

elements is a cluster of extremely closely-related lineages that was nearly ubiquitous in Corsica

but found elsewhere in low frequency. Although it remains unclear how this pattern arose, it is

possible that this chromosomal lineage has swept to high frequency as the result of recent

selection. Epstein et al. (2012) found low diversity along half of the *E. meliloti* chromosome in a

range-wide sample of 24 strains, likely indicating a recent selective sweep with little

recombination. Given little recombination on the chromosome (Epstein et al. 2012; Nelson et al.

2018), targets of selection on this element can be difficult to distinguish. Nevertheless it would

be interesting to test whether the three genome elements of *Ensifer* are locally adapted in ways

consistent with their functional gene composition, possibly by using host genetic variation

alongside abiotic climatic factors in a landscape genomics model (e.g., Yoder et al. 2014;

Rudman et al. 2018).

Our work suggests that pSymA and pSymB might be inherited together, leading to strong

correlations between them at both the individual and population scales. Recombination in *Ensifer*

is known to occur through conjugation of pSymA and pSymB (Ding and Hynes 2009). Recent

work indicates that pSymB is mobilized by pSymA via a *rctA*-mediated control system, though

tracking these events in real time has proven elusive in the laboratory (Pretorius-Guth et al. 1990;

Galibert et al. 2001; Pérez-Mendoza et al. 2005; Blanca-Ordóñez et al. 2010; diCenzo and Finan

2017). If pSymA and pSymB are coupled in nature, this could give rise to genetic constraint that

could affect *Ensifer*'s ability to adapt to distinct selective forces arising from various

environmental conditions encountered in the rhizosphere versus in association with host plants.

Functional predictions (*e.g.,* gene knockout experiments, functional annotation based on

homology) do not always predict quantitative trait variation; therefore, hypotheses regarding genetic constraint among the elements should be addressed with quantitative genetic correlations between rhizosphere and symbiosis traits (e.g., Ossler and Heath 2018; Wood et al. 2018).

Distinct spatial genetic processes between the chromosome and symbiosis elements, and the coupling of the symbiosis elements, begs the question of how horizontal gene transfer (HGT) and recombination generate population structure in these bacteria. While we summarize the geographical patterns at the whole-element scale in order to compare these to the host (see below), finer-scaled work on gene tree heterogeneity (Degnan and Rosenberg 2009) within each of the three elements is required for a full understanding of whether individual loci, large genomic regions, or even entire plasmids are introgressed through conjugation, and at what spatial scales such events occur. For example, Porter et al. (2019) recently used fine-scale patterns of gene co-occurrence across *Mesorhizobium* strains from California to discover the local transfer and duplication of symbiosis island and associate this variation with symbiotic partner quality.

Mismatched host-symbiont population structure: Our results indicate discordance in population structure between hosts and symbionts due to differences in neutral genetic processes such as gene flow or mutation, and/or spatially-variable selection. *Medicago* host plants are most structured along a longitudinal gradient, indicative of distinct glacial refugia, dividing populations in Spain from France and Corsica (Ronfort et al. 2006; Branca et al. 2011; Yoder et al. 2014; Grillo et al. 2016). By contrast, we found that most genome-wide variation in the *Ensifer* symbionts can be found within individual sites, and our phylogenetic and spatial analyses indicate that regional differentiation and IBD (which was significant for the chromosome) are

different from host plants, since chromosomes from Corsica were distinct from most mainland populations (see discussion above). In contrast to host plants and the *Ensifer* chromosome, pSymA and pSymB showed substantially less regional differentiation and no pattern of IBD, due to increased introgression following long-range dispersal events, higher mutation rates on these symbiotic elements, or both.

A striking feature of mutualisms is the existence of abundant genetic variation for mutualistic traits within and among populations, given models that predict the erosion of trait variation in mutualism (Heath and Stinchcombe 2014; Stoy et al. 2020). Indeed considerable coevolutionary variation (*i.e.,* genetic variation in partner choice, sanctioning, signaling, and partner quality) has been identified in the *Medicago-Ensifer* mutualism (Heath and Tiffin 2009; Heath 2010; Batstone et al. 2017, 2020; Burghardt et al. 2018, 2019). A mismatch in the population genetic structure of interacting species, as we identified here, could restrict populations of host and symbionts from locally adapting to each other and thereby contribute to the maintenance of coevolutionary trait variation. Thus a strong local selective force would be necessary to overcome the homogenizing effects of gene flow (*i.e.,* migration-selection balance; Savolainen et al. 2013) in order for hosts and symbionts to locally adapt to each other. Empirical work in other systems comparing the population structure of interacting species finds a range of outcomes, from correlated spatial genetic patterns between partners (Anderson et al. 2004; Thompson et al. 2005; Smith et al. 2008; Caldera and Currie 2012), to largely discordant patterns, where one species exhibits substantially less population structure across the sampled geographic range, compared to its partner (Dybdahl and Lively 1996; Baums et al. 2014; Strobel et al. 2016).

Spatially-explicit theory on mutualism coevolution to date indicates that the relative rates of gene flow between hosts and symbionts alters the likelihood of trait matching/local adaptation in, and even that gene flow can help explain the existence of maladaptation and abundant genetic variation for mutualism (Parker 1999; Nuismer et al. 2003; Yoder and Nuismer 2010). An important finding of this work, however, is that the outcomes of coevolution depend on how gene flow interacts with forms of natural selection, yet mutualism research often lacks explicit information on how selection in the wild acts on mutualism traits like partner quality and partner choice (Heath and Stinchcombe 2014; Stoy et al. 2020).

The degree of fitness alignment in mutualism, and thus the degree to which these interactions should coevolve like better-studied antagonisms, is an unresolved question (Friesen 2012; Jones et al. 2015; Frederickson 2017; Sachs et al. 2018; Batstone et al. 2020; Gano-Cohen et al. 2020). For antagonistic interactions, which are governed by negative frequency-dependent selection due to interspecific conflict, models suggest that the partner with a higher migration rate will benefit from the influx of new, potentially adaptive alleles that provide a competitive edge (Gandon et al. 1996b; Carlsson-Granér and Thrall 2015). If conflict prevails in mutualisms, and increases in rhizobium fitness occur at the expense of the host (Porter and Simms 2014; Gano-Cohen et al. 2020), incongruent population structures between hosts and symbionts should make it harder for hosts to evolve mechanisms to exclude lower-quality symbionts (*e.g.,* partner choice; Akçay 2017; Younginger and Friesen 2019), though this depends on the strength of selection. On the other hand, alignment between host and symbiont fitness might be common in mutualisms, as suggested by recent lab and experimental evolution studies (Friesen 2012; Batstone et al. 2020). This scenario should ultimately result in positive frequency-dependent selection wherein novelty is disfavored (unless populations are far from their adaptive peak),

leading to stable allele frequencies under purifying selection. Mismatched population structure, as identified here, would restrict the efficacy of positive frequency-dependence and any resulting local adaptation between hosts and symbionts. Our finding of discordant population genetic structure is in agreement with other host-microbe symbioses where microbes are transmitted horizontally and independently of their hosts (Dybdahl and Lively 1996; Baums et al. 2014; Strobel et al. 2016; Harrison et al. 2017). If this is common, then coevolutionary natural selection would have to be quite strong to overcome the demographic force of gene flow and generate patterns of local coadaptation in such microbial symbioses.

Data accessibility:

Raw sequence reads and assemblies are archived at NCBI (accessions ###).

References:

Akçay, E. 2017. Population structure reduces benefits from partner choice in mutualistic symbiosis. Proc. R. Soc. B Biol. Sci. 284: 20162317.

Anderson, B., I. Olivieri, M. Lourmas, and B. A. Stewart. 2004. Comparative population genetic structures and local adaptation of two mutualists. Evolution 58:1730–1747.

Bailly, X., E. Giuntini, M. C. Sexton, R. P. J. Lower, P. W. Harrison, N. Kumar, and J. P. W. Young. 2011. Population genomics of Sinorhizobium medicae based on low-coverage sequencing of sympatric isolates. ISME J. 5:1722–1734.

Bailly, X., I. Olivieri, S. De Mita, J. C. Cleyet-Marel, and G. Béna. 2006. Recombination and selection shape the molecular diversity pattern of nitrogen-fixing Sinorhizobium sp. associated to Medicago. Mol. Ecol. 15:2719–2734.

Bankevich, A., S. Nurk, D. Antipov, A. A. Gurevich, M. Dvorkin, A. S. Kulikov, V. M. Lesin, S. I. Nikolenko, S. Pham, A. D. Prjibelski, A. V. Pyshkin, A. V. Sirotkin, N. Vyahhi, G. Tesler, M. A. Alekseyev, and P. A. Pevzner. 2012. SPAdes: A new genome assembly algorithm and its applications to single-cell sequencing. J. Comput. Biol. 19:455–477.

Batstone, R. T., E. M. Dutton, D. Wang, M. Yang, and M. E. Frederickson. 2017. The evolution of symbiont preference traits in the model legume Medicago truncatula. New Phytol. 213:1850–1861.

Batstone, R. T., A. M. O'Brien, T. L. Harrison, and M. E. Frederickson. 2020. Experimental evolution makes microbes more cooperative with their local host genotype. Science 370:23–26.

Baums, I. B., M. K. Devlin-Durante, and T. C. Lajeunesse. 2014. New insights into the dynamics between reef corals and their associated dinoflagellate endosymbionts from population

genetic studies. Mol. Ecol. 23:4203–4215.

Becker, A., M. J. Barnett, D. Capela, M. Dondrup, P. B. Kamp, E. Krol, B. Linke, S. Rüberg, K. Runte, B. K. Schroeder, S. Weidner, S. N. Yurgel, J. Batut, S. R. Long, A. Pühler, and A. Goesmann. 2009. A portal for rhizobial genomes: RhizoGATE integrates a Sinorhizobium meliloti genome annotation update with postgenome data. J. Biotechnol. 140:45–50.

Berg, G. 2009. Plant-microbe interactions promoting plant growth and health: Perspectives for controlled use of microorganisms in agriculture. Appl. Microbiol. Biotechnol. 84:11–18.

Biondi, E. G., E. Pilli, E. Giuntini, M. L. Roumiantseva, E. E. Andronov, O. P. Onichtchouk, O. N. Kurchak, B. V. Simarov, N. I. Dzyubenko, A. Mengoni, and M. Bazzicalupo. 2003. Genetic relationship of Sinorhizobium meliloti and Sinorhizobium medicae strains isolated from Caucasian region. FEMS Microbiol. Lett. 220:207–213.

Blanca-Ordóñez, H., J. J. Oliva-García, D. Pérez-Mendoza, M. J. Soto, J. Olivares, J. Sanjuán, and J. Nogales. 2010. pSymA-dependent mobilization of the Sinorhizobium meliloti pSymB megaplasmid. J. Bacteriol. 192:6309–6312.

Bonhomme, M., S. Boitard, H. S. Clemente, B. Dumas, N. Young, and C. Jacquet. 2015. Genomic signature of selective sweeps illuminates adaptation of medicago truncatula to root-associated microorganisms. Mol. Biol. Evol. 32:2097–2110.

Bonnin, I., T. Huguet, M. Gherardi, J.-M. Prosperi, and I. Olivieri. 1996. High level of polymorphism and spatial structure in a selfing plant species, Medicago truncatula (Leguminosae), shown using RAPD markers . Am. J. Bot. 83:843–855.

Branca, A., T. D. Paape, P. Zhou, R. Briskine, A. D. Farmer, J. Mudge, A. K. Bharti, J. E. Woodward, G. D. May, L. Gentzbittel, C. Ben, R. Denny, M. J. Sadowsky, J. Ronfort, T. Bataillon, N. D. Young, and P. Tiffin. 2011. Whole-genome nucleotide diversity,

recombination, and linkage disequilibrium in the model legume Medicago truncatula. Proc. Natl. Acad. Sci. U. S. A. 108: E864-870.

Brown, S. P., M. A. Grillo, J. C. Podowski, and K. D. Heath. 2020. Soil origin and plant genotype structure distinct microbiome compartments in the model legume Medicago truncatula. Microbiome 8:1–17.

Burghardt, L. T. 2020. Evolving together, evolving apart: measuring the fitness of rhizobial bacteria in and out of symbiosis with leguminous plants. New Phytol. 228:28–34.

Burghardt, L. T., B. Epstein, J. Guhlin, M. S. Nelson, M. R. Taylor, N. D. Young, M. J. Sadowsky, and P. Tiffin. 2018. Select and resequence reveals relative fitness of bacteria in symbiotic and free-living environments. Proc. Natl. Acad. Sci. U. S. A. 115:2425–2430.

Burghardt, L. T., B. Epstein, and P. Tiffin. 2019. Legacy of prior host and soil selection on rhizobial fitness in planta. Evolution (N. Y). 73:2013–2023.

Caldera, E. J., and C. R. Currie. 2012. The population structure of antibiotic-producing bacterial symbionts of apterostigma dentigerum ants: Impacts of coevolution and multipartite symbiosis. Am. Nat. 180:604–617.

Carlsson-Granér, U., and P. H. Thrall. 2015. Host resistance and pathogen infectivity in host populations with varying connectivity. Evolution 69:926–938.

Catchen, J., P. A. Hohenlohe, S. Bassham, A. Amores, and W. A. Cresko. 2013. StackCatchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). Stacks: an analysis tool set for population genomics. Molecular Ecology 22:3124–40.

Cavassim, M. I. A., S. Moeskjær, C. Moslemi, B. Fields, A. Bachmann, B. J. Vilhjálmsson, M. H. Schierup, J. P. W. Young, and S. U. Andersen. 2020. Symbiosis genes show a unique pattern of introgression and selection within a rhizobium leguminosarum species complex.

Microb. Genomics 6:e000351.

Chase, A., P. Arevalo, E. Brodie, M. Polz, U. Karaoz, and J. B. H. Martiny. 2019. Maintenance of sympatric and allopatric populations in free-living terrestrial bacteria. MBio 10:202361-19.

Choudoir, M. J., A. Barberán, H. L. Menninger, R. R. Dunn, and N. Fierer. 2018. Variation in range size and dispersal capabilities of microbial taxa. Ecology 99:322–334.

Cooper, V. S., S. H. Vohr, S. C. Wrocklage, and P. J. Hatcher. 2010. Why genes evolve faster on secondary chromosomes in bacteria. PLoS Comput. Biol. 6:e1000732.

Danecek, P., A. Auton, G. Abecasis, C. A. Albers, E. Banks, M. A. DePristo, R. E. Handsaker, G. Lunter, G. T. Marth, S. T. Sherry, G. McVean, and R. Durbin. 2011. The variant call format and VCFtools. Bioinformatics 27:2156–2158.

Degnan, J. H., and N. A. Rosenberg. 2009. Gene tree discordance, phylogenetic inference and the multispecies coalescent. Trends Ecol. Evol. 24:332–340.

diCenzo, G. C., and T. M. Finan. 2017. The Divided Bacterial Genome. Microbiol. Mol. Biol. Rev. 81:1–37.

diCenzo, G. C., A. M. MacLean, B. Milunovic, G. B. Golding, and T. M. Finan. 2014. Examination of Prokaryotic Multipartite Genome Evolution through Experimental Genome Reduction. PLoS Genet. 10:e1004742.

Ding, H., and M. F. Hynes. 2009. Plasmid transfer systems in the rhizobia. Can. J. Microbiol. 55:917–927.

Dray, S., and A. B. Dufour. 2007. The ade4 package: Implementing the duality diagram for ecologists. J. Stat. Softw. 22:1–20.

Dybdahl, M. F., and C. M. Lively. 1996. The geography of coevolution: Comparative population

structures for a snail and its trematode parasite. Evolution 50:2264–2275.

Epstein, B., A. Branca, J. Mudge, A. K. Bharti, R. Briskine, A. D. Farmer, M. Sugawara, N. D. Young, M. J. Sadowsky, and P. Tiffin. 2012. Population Genomics of the Facultatively Mutualistic Bacteria Sinorhizobium meliloti and S. medicae. PLoS Genet. 8:1–10.

Epstein, B., M. J. Sadowsky, and P. Tiffin. 2014. Selection on Horizontally Transferred and Duplicated Genes in Sinorhizobium (Ensifer), the Root-Nodule Symbionts of Medicago. Genome Biol. Evol. 6:1199–1209.

Epstein, B., and P. Tiffin. 2021. Comparative genomics reveals high rates of horizontal transfer and strong purifying selection on rhizobial symbiosis genes. Proc. R. Soc. B Biol. Sci. 288: 20201804.

Fernandes, L. D., P. Lemos-Costa, P. R. Guimarães, J. N. Thompson, and M. A. M. de Aguiar. 2019. Coevolution Creates Complex Mosaics across Large Landscapes. Am. Nat. 194:000–000.

Frederickson, M. E. 2017. Mutualisms Are Not on the Verge of Breakdown. Trends Ecol. Evol. 32:727–734.

Friesen, M. L. 2012. Widespread fitness alignment in the legume-rhizobium symbiosis. New Phytol. 194:1096–1111.

Friesen, M. L., and A. Mathias. 2010. Mixed infections may promote diversification of mutualistic symbionts: Why are there ineffective rhizobia? J. Evol. Biol. 23:323–334.

Galardini, M., F. Pini, M. Bazzicalupo, E. G. Biondi, and A. Mengoni. 2013. Replicon-dependent bacterial genome evolution: The case of Sinorhizobium meliloti. Genome Biol. Evol. 5:542–558.

Galibert, F., T. M. Finan, S. R. Long, A. Pühler, P. Abola, F. Ampe, F. Barloy-Hubler, M. J.

Barnet, A. Becker, P. Boistard, G. Bothe, M. Boutry, L. Bowser, J. Buhrmester, E. Cadieu, D. Capela, P. Chain, A. Cowie, R. W. Davis, S. Dréano, N. A. Federspiel, R. F. Fisher, S. Gloux, T. Godrie, A. Goffeau, B. Golding, J. Gouzy, M. Gurjal, I. Hernandez-Lucas, A. Hong, L. Huizar, R. W. Hyman, T. Jones, D. Kahn, M. L. Kahn, S. Kalman, D. H. Keating, E. Kiss, C. Komp, V. Lelaure, D. Masuy, C. Palm, M. C. Peck, T. M. Pohl, D. Portetelle, B. Purnelle, U. Ramsperger, R. Surzycki, P. Thébault, M. Vandenbol, F. J. Vorhölter, S. Weidner, D. H. Wells, K. Wong, K. C. Yeh, and J. Batut. 2001. The composite genome of the legume symbiont Sinorhizobium meliloti. Science 293:668–672.

Gandon, S., Y. Capowiez, Y. Dubois, Y. Michalakis, and I. Olivieri. 1996a. Local adaptation and gene-for-gene coevolution in a metapopulation model. Proc. R. Soc. B Biol. Sci. 263:1003–1009.

Gandon, S., Y. Capowiez, Y. Dubois, Y. Michalakis, and I. Olivieri. 1996b. Local adaptation and gene-for-gene coevolution in a metapopulation model. Proc. R. Soc. B Biol. Sci. 263:1003–1009.

Gano-Cohen, K. A., C. E. Wendlandt, K. Al Moussawi, P. J. Stokes, K. W. Quides, A. J. Weisberg, J. H. Chang, and J. L. Sachs. 2020. Recurrent mutualism breakdown events in a legume rhizobia metapopulation. Proc. R. Soc. B Biol. Sci. 287:20192549.

Garrison, E., and G. Marth. 2012. Haplotype-based variant detection from short-read sequencing. *a*rXiv preprint arXiv:1207.3907

Green, J. L., B. J. M. Bohannan, and R. J. Whitaker. 2008. Microbial biogeography: From taxonomy to traits. Science 320:1039–1043.

Grillo, M. A., S. De Mita, P. V. Burke, K. L. S. Solórzano-Lowell, and K. D. Heath. 2016. Intrapopulation genomics in a model mutualist: Population structure and candidate

symbiosis genes under selection in *Medicago truncatula*. Evolution 70:2704–2717.

Hanson, C. A., J. A. Fuhrman, M. C. Horner-Devine, and J. B. H. Martiny. 2012. Beyond biogeographic patterns: Processes shaping the microbial landscape. Nat. Rev. Microbiol. 10:497–506.

Harrison, P. W., R. P. J. Lower, N. K. D. Kim, and J. P. W. Young. 2010. Introducing the bacterial 'chromid': not a chromosome, not a plasmid. Trends Microbiol. 18:141–148.

Harrison, T. L., C. W. Wood, K. D. Heath, and J. R. Stinchcombe. 2017. Geographically structured genetic variation in the *Medicago lupulina - Ensifer* mutualism. Evolution 71:1787–1801.

Heath, K. D. 2010. Intergenomic epistasis and coevolutionary constraint in plants and rhizobia. Evolution 64:1446–1458.

Heath, K. D., and M. A. Grillo. 2016. Rhizobia: tractable models for bacterial evolutionary ecology. Environ. Microbiol. 18:4307–4311.

Heath, K. D., and J. R. Stinchcombe. 2014a. Explaining mutualism variation: A new evolutionary paradox? Evolution (N. Y). 68:309–317.

Heath, K. D., and J. R. Stinchcombe. 2014b. Explaining mutualism variation: A new evolutionary paradox? Evolution 68:309–317.

Heath, K. D., and P. Tiffin. 2009. Stabilizing mechanisms in a legume-rhizobium mutualism. Evolution 63:652–662.

Hoeksema, J. D., and S. E. Forde. 2008. A meta-analysis of factors affecting local adaptation between interacting species. Am. Nat. 171:275–290.

Hoetzinger, M., A. Pitt, A. Huemer, and M. W. Hahn. 2021. Continental-Scale Gene Flow Prevents Allopatric Divergence of Pelagic Freshwater Bacteria. Genome Biol. Evol. 13:1–

18.

Hollowell, A. C., J. U. Regus, K. A. Gano, R. Bantay, D. Centeno, J. Pham, J. Y. Lyu, D. Moore, A. Bernardo, G. Lopez, A. Patil, S. Patel, Y. Lii, and J. L. Sachs. 2016a. Epidemic Spread of Symbiotic and Non-Symbiotic Bradyrhizobium Genotypes Across California. Microb. Ecol. 71:700–710.

Hollowell, A. C., J. U. Regus, D. Turissini, K. A. Gano-Cohen, R. Bantay, A. Bernardo, D. Moore, J. Pham, and J. L. Sachs. 2016b. Metapopulation dominance and genomicisland acquisition of Bradyrhizobium with superior catabolic capabilities. Proc. R. Soc. B Biol. Sci. 283.

Jombart, T., and I. Ahmed. 2011. adegenet 1.3-1: New tools for the analysis of genome-wide SNP data. Bioinformatics 27:3070–3071.

Jones, E. I., M. E. Afkhami, E. Akçay, J. L. Bronstein, R. Bshary, M. E. Frederickson, K. D. Heath, J. D. Hoeksema, J. H. Ness, M. S. Pankey, S. S. Porter, J. L. Sachs, K. Scharnagl, and M. L. Friesen. 2015. Cheaters must prosper: Reconciling theoretical and empirical perspectives on cheating in mutualism. Ecol. Lett. 18:1270–1284.

Kamvar, Z. N., J. F. Tabima, and N. J. Grünwald. 2014. Poppr: An R package for genetic analysis of populations with clonal, partially clonal, and/or sexual reproduction. PeerJ 2014:1–14.

Kaneko, T., Y. Nakamura, S. Sato, K. Minamisawa, T. Uchiumi, S. Sasamoto, A. Watanabe, K. Idesawa, M. Iriguchi, K. Kawashima, M. Kohara, M. Matsumoto, S. Shimpo, H. Tsuruoka, T. Wada, M. Yamada, and S. Tabata. 2002. Complete genomic sequence of nitrogen-fixing symbiotic bacterium Bradyrhizobium japonicum USDA110. DNA Res. 9:189–197.

Klinger, C. R., J. A. Lau, and K. D. Heath. 2016. Ecological genomics of mutualism decline in

nitrogen-fixing bacteria. Proc. R. Soc. B Biol. Sci. 283:20152563.

Koppell, J. H., and M. A. Parker. 2012. Phylogenetic clustering of Bradyrhizobium symbionts on legumes indigenous to North America. Microbiol. (United Kingdom) 158:2050–2059.

Kraemer, S. A., and P. J. Boynton. 2017. Evidence for microbial local adaptation in nature. Mol. Ecol. 26:1860–1876.

Kumar, N., G. Lad, E. Giuntini, M. E. Kaye, P. Udomwong, N. J. Shamsani, J. P. W. Young, and X. Bailly. 2015. Bacterial genospecies that are not ecologically coherent: population genomics of Rhizobium leguminosarum. Open Biol. 5:140133–140133.

Laine, A. L. 2005. Spatial scale of local adaptation in a plant-pathogen metapopulation. J. Evol. Biol. 18:930–938.

Li, H., and R. Durbin. 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics 25:1754–1760.

Locey, K. J., and J. T. Lennon. 2016. Scaling laws predict global microbial diversity. Proc. Natl. Acad. Sci. U. S. A. 113:5970–5975.

Martiny, J. B. H., B. J. M. Bohannan, J. H. Brown, R. K. Colwell, J. A. Fuhrman, J. L. Green, M. C. Horner-Devine, M. Kane, J. A. Krumins, C. R. Kuske, P. J. Morin, S. Naeem, L. Øvreås, A. L. Reysenbach, V. H. Smith, and J. T. Staley. 2006. Microbial biogeography: Putting microorganisms on the map. Nat. Rev. Microbiol. 4:102–112.

Masatoshi, N. 1972. Genetic Distance between Populations. Am. Nat. 106:283–292.

Nelson, M., J. Guhlin, B. Epstein, P. Tiffin, and M. J. Sadowsky. 2018a. The complete replicons of 16 Ensifer meliloti strains offer insights into intra- and inter-replicon gene transfer, transposon-associated loci, and repeat elements. Microb. Genomics 4:e000174.

Nelson, M., J. Guhlin, B. Epstein, P. Tiffin, and M. J. Sadowsky. 2018b. The complete replicons

of 16 Ensifer meliloti strains offer insights into intra- and inter-replicon gene transfer, transposon-associated loci, and repeat elements. Microb. Genomics 4:1–11.

Nuismer, S. L., J. N. Thompson, and R. Gomulkiewicz. 2003. Coevolution between hosts and parasites with partially overlapping geographic ranges. J. Evol. Biol. 16:1337–1345.

Ossler, J. N., and K. D. Heath. 2018. Shared genes but not shared genetic variation: Legume colonization by two belowground symbionts. Am. Nat. 191:395–406.

Page, A. J., C. A. Cummins, M. Hunt, V. K. Wong, S. Reuter, M. T. G. Holden, M. Fookes, D. Falush, J. A. Keane, and J. Parkhill. 2015. Roary: Rapid large-scale prokaryote pan genome analysis. Bioinformatics 31:3691–3693.

Paradis, E., and K. Schliep. 2019. Ape 5.0: An environment for modern phylogenetics and evolutionary analyses in R. Bioinformatics 35:526–528.

Parker, M. A. 1999. Mutualism in metapopulations of legumes and rhizobia. Am. Nat. 153:48–60.

Pembleton, L. W., N. O. I. Cogan, and J. W. Forster. 2013. StAMPP: An R package for calculation of genetic differentiation and structure of mixed-ploidy level populations. Mol. Ecol. Resour. 13:946–952.

Pérez-Mendoza, D., E. Sepúlveda, V. Pando, S. Muñoz, J. Nogales, J. Olivares, M. J. Soto, J. A. Herrera-Cervera, D. Romero, S. Brom, and J. Sanjuán. 2005. Identification of the rctA gene, which is required for repression of conjugative transfer of rhizobial symbiotic megaplasmids. J. Bacteriol. 187:7341–7350.

Pérez Carrascal, O. M., D. VanInsberghe, S. Juárez, M. F. Polz, P. Vinuesa, and V. González. 2016. Population genomics of the symbiotic plasmids of sympatric nitrogen-fixing *Rhizobium* species associated with *Phaseolus vulgaris*. Environ. Microbiol. 18:2660–2676.

Pita, L., L. Rix, B. M. Slaby, A. Franke, and U. Hentschel. 2018. The sponge holobiont in a changing ocean: from microbes to ecosystems. Microbiome 6:46.

Porter, S. S., J. Faber-Hammond, A. P. Montoya, M. L. Friesen, and C. Sackos. 2019. Dynamic genomic architecture of mutualistic cooperation in a wild population of Mesorhizobium. ISME J. 13:301–315.

Porter, S. S., and E. L. Simms. 2014. Selection for cheating across disparate environments in the legume-rhizobium mutualism. Ecol. Lett. 17:1121–1129.

Pretorius-Guth, I. M., A. Puhler, and R. Simon. 1990. Conjugal transfer of megaplasmid 2 between Rhizobium meliloti strains in alfalfa nodules. Appl. Environ. Microbiol. 56:2354–2359.

Rambaut, A. 2018. FigTree. http://tree.bio.ed.ac.uk/software/figtree/

Revillini, D., C. A. Gehring, and N. C. Johnson. 2016. The role of locally adapted mycorrhizas and rhizobacteria in plant–soil feedback systems. Funct. Ecol. 30:1086–1098.

Reynolds, H. L., A. Packer, J. D. Bever, and K. Clay. 2003. Grassroots ecology: Plant-microbe-soil interactions as drivers of plant community structure and dynamics. Ecology 84:2281–2291.

Ronfort, J., T. Bataillon, S. Santoni, M. Delalande, J. L. David, and J. M. Prosperi. 2006. Microsatellite diversity and broad scale geographic structure in a model legume: Building a set of nested core collection for studying naturally occurring variation in Medicago truncatula. BMC Plant Biol. 6:1–13.

Rudman, S. M., M. A. Barbour, K. Csilléry, P. Gienapp, F. Guillaume, N. G. Hairston, A. P. Hendry, J. R. Lasky, M. Rafajlović, K. Räsänen, P. S. Schmidt, O. Seehausen, N. O. Therkildsen, M. M. Turcotte, and J. M. Levine. 2018. What genomic data can reveal about

eco-evolutionary dynamics. Nat. Ecol. Evol. 2:9–15.

Sachs, J. L., K. W. Quides, and C. E. Wendlandt. 2018. Legumes versus rhizobia : a model for ongoing conflict in symbiosis. New Phytologist 219:1199–1206.

Savolainen, O., M. Lascoux, and J. Merilä. 2013. Ecological genomics of local adaptation. Nat. Rev. Genet. 14:807–820.

Seemann, T. 2014. Prokka: Rapid prokaryotic genome annotation. Bioinformatics 30:2068–2069.

Siol, M., J. M. Prosperi, I. Bonnin, and J. Ronfort. 2008. How multilocus genotypic pattern helps to understand the history of selfing populations: A case study in Medicago truncatula. Heredity 100:517–525.

Smith, C. I., W. K. W. Godsoe, S. Tank, J. B. Yoder, and O. Pellmyr. 2008. Distinguishing coevolution from covicariance in an obligate pollination mutualism: Asynchronous divergence in Joshua tree and its pollinators. Evolution 62:2676–2687.

Stoy, K. S., A. K. Gibson, N. M. Gerardo, and L. T. Morran. 2020. A need to consider the evolutionary genetics of host–symbiont mutualisms. J. Evol. Biol. 33:1656–1668.

Strobel, H. M., F. Alda, C. G. Sprehn, M. J. Blum, and D. C. Heins. 2016. Geographic and host-mediated population genetic structure in a cestode parasite of the three-spined stickleback. Biol. J. Linn. Soc. 119:381–396.

Tack, A. J. M., F. Horns, and A. L. Laine. 2014. The impact of spatial scale and habitat configuration on patterns of trait variation and local adaptation in a wild plant parasite. Evolution 68:176–189.

Tang, H., V. Krishnakumar, S. Bidwell, B. Rosen, A. Chan, S. Zhou, L. Gentzbittel, K. L. Childs, M. Yandell, H. Gundlach, K. F. X. Mayer, D. C. Schwartz, and C. D. Town. 2014.

An improved genome release (version Mt4.0) for the model legume Medicago truncatula. BMC Genomics 15:1–14.

Thompson, A. R., C. E. Thacker, and E. Y. Shaw. 2005. Phylogeography of marine mutualists: Parallel patterns of genetic structure between obligate goby and shrimp partners. Mol. Ecol. 14:3557–3572.

Thompson, J. N. 1994. The Coevolutionary Process. 1st ed. The University Of Chicago Press Books, Chicago, IL, USA.

Thompson, J. N. 2005. The Geographic Mosaic of Coevolution. 1st ed. Chicago Press.

Thrall, P. H., M. E. Hochberg, J. J. Burdon, and J. D. Bever. 2007. Coevolution of symbiotic mutualists and parasites in a community context. Trends Ecol. Evol. 22:120–126.

VanInsberghe, D., P. Arevalo, D. Chien, and M. F. Polz. 2020. How can microbial population genomics inform community ecology? Philos. Trans. R. Soc. B Biol. Sci. 375.

Vincent, J. 1970. A Manual for the Practical Study of Root-Nodule Bacteria. Blackwell Scientific, Oxford and Edinburgh.

Vos, M., and G. J. Velicer. 2008. Isolation by Distance in the Spore-Forming Soil Bacterium Myxococcus xanthus. Curr. Biol. 18:386–391.

Whitaker, R. J., and J. F. Banfield. 2006. Population genomics in natural microbial communities. Trends Ecol. Evol. 21:508–516.

Wickham, H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag, New York.

Wood, C. W., B. L. Pilkington, P. Vaidya, C. Biel, and J. R. Stinchcombe. 2018. Genetic conflict with a parasitic nematode disrupts the legume-rhizobia mutualism. Evol. Lett. 2:233–245.

Yates, R., J. Howieson, S. E. De Meyer, R. Tian, R. Seshadri, A. Pati, T. Woyke, V. Markowitz, N. Ivanova, N. Kyrpides, A. Loi, B. Nutt, G. Garau, L. Sulas, and W. Reeve. 2015. High-

quality permanent draft genome sequence of Rhizobium sullae strain WSM1592; a

Hedysarum coronarium microsymbiont from Sassari, Italy. Stand. Genomic Sci. 10:1–6.

Yoder, J. B., and S. L. Nuismer. 2010. When does coevolution promote diversification? Am.

Nat. 176:802–817.

Yoder, J. B., J. Stanton-Geddes, P. Zhou, R. Briskine, N. D. Young, and P. Tiffin. 2014.

Genomic signature of adaptation to climate in Medicago truncatula. Genetics 196:1263–

1275.

Young, J. P. W., L. C. Crossman, A. W. B. Johnston, N. R. Thomson, Z. F. Ghazoui, K. H. Hull,

M. Wexler, A. R. J. Curson, J. D. Todd, P. S. Poole, T. H. Mauchline, A. K. East, M. A.

Quail, C. Churcher, C. Arrowsmith, I. Cherevach, T. Chillingworth, K. Clarke, A. Cronin,

P. Davis, A. Fraser, Z. Hance, H. Hauser, K. Jagels, S. Moule, K. Mungall, H. Norbertczak,

E. Rabbinowitsch, M. Sanders, M. Simmonds, S. Whitehead, and J. Parkhill. 2006. The

genome of Rhizobium leguminosarum has recognizable core and accessory components.

Genome Biol. 7.

Young, J. P. W., and K. E. Haukka. 1996. Diversity and phylogeny of rhizobia. New Phytol.

133:87–94.

Younginger, B. S., and M. L. Friesen. 2019. Connecting signals and benefits through partner

choice in plant-microbe interactions. FEMS Microbiol. Lett. 366:1–11.

Zribi, K., R. Mhamdi, T. Huguet, and M. E. Aouani. 2004. Distribution and genetic diversity of

rhizobia nodulating natural populations of Medicago truncatula in tunisian soils. Soil Biol.

Biochem. 36:903–908.

Table 1. AMOVA partitioning the genome-wide genetic variation for each of the three elements of the *E. meliloti* genome for 191 strains sampled from 21 populations from three regions (Spain, France, or Corsica) in the native range of the symbiosis, compared to host plant *M. truncatula*. For each level of spatial division, the percent variance explained, phi statistic, and significance are given.

	Chromosome		pSymA		pSymB		*M. truncatula*	
	% variance	*Phi*	*% variance*	*Phi*	*% variance*	*Phi*	*% variance*	*Phi*
Among Region	24.5	0.34**	7.7	0.27**	10.1	0.24***	19.6	0.20***
Among Population within Region	9.7	0.13***	19.2	0.21***	13.8	0.15***	36.5	0.45***
Within Population	65.8	0.25***	73.1	0.08***	76.1	0.10***	44.0	0.56***

*p<0.05; **p<0.01; ***p<0.001; ****p<0.0001

Figure Legends

Figure 1. Tests of correlated population structure between the genomic elements in *E. meliloti*. Shown are correlations between the matrices of pairwise D_{XY} values. Left column are individual D_{XY} values and right column are population-level D_{XY} values.

Figure 2. Neighbor joining trees of rhizobium individuals based on (a) chromosomal and (b) pSymA variant data. For both trees individual tip labels (strain ID followed by soil population) are colored based on region of origin (orange from Corsica, green from mainland France, blue from Spain). Branch support indicated in teal (<70) or pink (<50). The outer ring coloration in both trees represents the five major clusters of the chromosome, allowing for comparison with pSymA. Only pSymA is shown here due to sharing a similar topology to pSymB (see Fig. S3).

Figure 3. Correlations between geographic distance and genetic distance (D_{XY}) for *Medicago* host plants (n=192) (green), and the *E. meliloti* (n=191) chromosome (red), pSymA (blue), and pSymB (purple). Shown are trend lines and Pearson r^2 values from Mantel tests for each comparison ($^{***}p<0.001$).

Figure 1.

Figure. 2a

0.05

Figure. 2b

0.02

Figure. 3

Table S1. Metadata on strains of *Ensifer meliloti*. For each strain the region, sampling site (soil), and specific soil samples are given, along with the genotype of the *Medicago truncatula* plant used to isolate the strain, and the latitude and longitude at which the soil sample was taken.

STRAIN	REGION	SOIL	SOIL_SAMPLE	PLANT GENOTYPE	LAT	LONG
MAG722A	Corsica	C1	C1-3	F83026	42.16748	9.542717
MAG722B	Corsica	C1	C1-3	F83026	42.16748	9.542717
MAG530	Corsica	C2	C2-5	ESP161	42.9687	9.41295
MAG420	Corsica	C3	C3-8	F20089	42.97463	9.365783
MAG421	Corsica	C3	C3-8	F20089	42.97463	9.365783
MAG422	Corsica	C3	C3-8	F20089	42.97463	9.365783
MAG498	Corsica	C3	C3-8	ESP161	42.97463	9.365783
MAG508	Corsica	C3	C3-11	F20089	42.97263	9.366483
MAG701A	Corsica	C3	C3-8	F83026	42.97463	9.365783
MAG701B	Corsica	C3	C3-8	F83026	42.97463	9.365783
MAG714A	Corsica	C3	C3-11	F20093	42.97263	9.366483
MAG714B	Corsica	C3	C3-11	F20093	42.97263	9.366483
MAG513	Corsica	C4	C4-12	ESP161	42.59243	9.04045
MAG514	Corsica	C4	C4-12	ESP161	42.59243	9.04045
MAG533	Corsica	C4	C4-12	F20093	42.59243	9.04045
MAG710A	Corsica	C4	C4-12	F20093	42.59243	9.04045
MAG710B	Corsica	C4	C4-12	F20093	42.59243	9.04045
MAG724A	Corsica	C4	C4-4	F20043	42.59172	9.04035
MAG724B	Corsica	C4	C4-4	F20043	42.59172	9.04035
MAG726A	Corsica	C4	C4-7	F20043	42.59098	9.040233
MAG726B	Corsica	C4	C4-7	F20043	42.59098	9.040233
MAG523	France	1	1-8	ESP009	43.1474	3.000417
MAG524	France	1	1-8	ESP009	43.1474	3.000417
MAG525	France	1	1-8	ESP009	43.1474	3.000417
MAG526	France	1	1-8	ESP009	43.1474	3.000417
MAG141	France	2	2-1	F11005	43.14543	2.999367
MAG183	France	2	2-1	ESP174	43.14543	2.999367
MAG184	France	2	2-1	ESP174	43.14543	2.999367
MAG199	France	2	2-1	F11005	43.14543	2.999367
MAG220	France	2	2-9	ESP174	43.1457	2.9989
MAG32	France	2	2-1	F11005 n	43.14543	2.999367
MAG539	France	2	2-7	F11005	43.14542	2.998683
MAG540	France	2	2-7	F11005	43.14542	2.998683
MAG541	France	2	2-7	F11005	43.14542	2.998683
MAG80	France	2	2-10	ESP174	43.14572	2.998917
MAG82	France	2	2-1	ESP174	43.14543	2.999367
MAG92	France	2	2-9	F11005	43.1457	2.9989

MAG	Country			Site	Latitude	Longitude
MAG93	France	2	2-9	F11005	43.1457	2.9989
MAG123	France	3	3-3	F11005	43.12227	3.077067
MAG177	France	3	3-4	F11005	43.12235	3.075667
MAG194-mel	France	3	3-2	ESP174	43.12233	3.076717
MAG225	France	3	3-2	ESP174	43.12233	3.076717
MAG245	France	3	3-1	ESP174	43.12222	3.07635
MAG246	France	3	3-1	ESP174	43.12222	3.07635
MAG247	France	3	3-1	ESP174	43.12222	3.07635
MAG248	France	3	3-1	ESP174	43.12222	3.07635
MAG34	France	3	3-2	ESP174	43.12233	3.076717
MAG40	France	3	3-3	ESP174	43.12227	3.077067

no

MAG	Country			Site	Latitude	Longitude
MAG69	France	3	3-2	F11005	43.12233	3.076717
MAG90A	France	3	3-7	F11005	43.12232	3.075283
MAG90B	France	3	3-7	F11005	43.12232	3.075283
MAG14	France	4	4-2	ESP174	43.12198	3.095533
MAG16	France	4	4-3	F20089	43.12227	3.096033
MAG204	France	4	4-1	ESP174	43.122	3.095283
MAG205	France	4	4-1	ESP174	43.122	3.095283
MAG206	France	4	4-1	ESP174	43.122	3.095283
MAG24	France	4	4-3	F20089	43.12227	3.096033
MAG27	France	4	4-2	ESP174	43.12198	3.095533

n

MAG	Country			Site	Latitude	Longitude
MAG39	France	4	4-1	F20089	43.122	3.095283
MAG43	France	4	4-1	F20089	43.122	3.095283
MAG48	France	4	4-3	ESP174	43.12227	3.096033
MAG5	France	4	4-3	F20089	43.12227	3.096033
MAG57	France	4	4-3	ESP174	43.12227	3.096033
MAG59	France	4	4-1	ESP174	43.122	3.095283
MAG62	France	4	4-1	F20089	43.122	3.095283
MAG65	France	4	4-3	ESP174	43.12227	3.096033
MAG67	France	4	4-2	F20089	43.12198	3.095533
MAG6A	France	4	4-3	F20089	43.12227	3.096033
MAG6B	France	4	4-3	F20089	43.12227	3.096033
MAG73	France	4	4-3	ESP174	43.12227	3.096033
MAG7A	France	4	4-2	ESP174	43.12198	3.095533
MAG7B	France	4	4-2	ESP174	43.12198	3.095533
MAG17	France	5	5-3	F20089	42.82302	2.9269
MAG209	France	5	5-1	F66009	42.82278	2.927717
MAG36	France	5	5-3	F66009	42.82302	2.9269
MAG46	France	5	5-3	F66009	42.82302	2.9269
MAG71	France	5	5-2	F20089	42.82302	2.927317
MAG201	France	6	6-2	F66009	42.74198	2.81485
MAG219	France	6	6-2	F20089	42.74198	2.81485

MAG38	France	6	6-2	F20089 n	42.74198	2.81485
MAG41	France	6	6-1	F20089	42.74205	2.814917
MAG42	France	6	6-1	F20089	42.74205	2.814917
MAG53	France	6	6-2	F20089	42.74198	2.81485
MAG72	France	6	6-2	F66009	42.74198	2.81485
MAG8	France	6	6-1	F66009	42.74205	2.814917
MAG85	France	6	6-1	F20089	42.74205	2.814917
MAG87	France	6	6-1	F20089	42.74205	2.814917
MAG88	France	6	6-1	F20089	42.74205	2.814917
MAG15	France	7	7-1	F66009 n	42.58827	2.78495
MAG18	France	7	7-1	F66009 n	42.58827	2.78495
MAG189	France	7	7-3	F20089	42.58895	2.785383
MAG191	France	7	7-2	F66009	42.5883	2.785
MAG215	France	7	7-3	F66009	42.58895	2.785383
MAG216	France	7	7-3	F66009	42.58895	2.785383
MAG26	France	7	7-2	F20089	42.5883	2.785
MAG54	France	7	7-2	F20089	42.5883	2.785
MAG78	France	7	7-4	F20089	42.58948	2.784833
MAG182	France	14	14-2	F11005	43.1612	3.05905
MAG200	France	14	14-A	ESP174	43.1612	3.059983
MAG84	France	14	14-A	F11005	43.1612	3.059983
MAG86	France	14	14-3	F11005	43.16108	3.059383
MAG89	France	14	14-1	ESP174	43.16122	3.059017
MAG117B	France	15	15-1	F83026	43.35138	5.169617
MAG145	France	15	15-1	F83026	43.35138	5.169617
MAG101	France	16	16-A	ESP174	43.37568	5.188133
MAG114	France	16	16-B	F83026	43.37568	5.1871
MAG142	France	16	16-3	ESP174	43.37693	5.188133
MAG144A	France	16	16-A	F83026	43.37568	5.188133
MAG144B	France	16	16-A	F83026	43.37568	5.188133
MAG158A	France	16	16-3	ESP174	43.37693	5.188133
MAG158B	France	16	16-3	ESP174	43.37693	5.188133
MAG169	France	16	16-3	ESP174	43.37693	5.188133
MAG97	France	16	16-B	ESP174	43.37568	5.1871
MAG303	France	17	17-5	ESP161	43.12692	6.1252
MAG372	France	17	17-12	F83026	43.12538	6.12535
MAG373	France	17	17-12	F83026	43.12538	6.12535
MAG733A	France	17	17-2	F20043	43.12693	6.125217
MAG762A	France	17	17-12	ESP174	43.12538	6.12535
MAG276	France	18	18-2	ESP161	43.16532	6.4774
MAG277	France	18	18-2	ESP161	43.16532	6.4774
MAG278	France	18	18-2	ESP161	43.16532	6.4774
MAG281	France	18	18-3	ESP161	43.1653	6.4774

MAG282	France	18	18-3	ESP161	43.1653	6.4774
MAG283	France	18	18-3	ESP161	43.1653	6.4774
MAG313	France	18	18-1	F83026	43.16528	6.477433
MAG317	France	18	18-5	F83026	43.1649	6.477233
MAG318	France	18	18-5	F83026	43.1649	6.477233
MAG319	France	18	18-5	F83026	43.1649	6.477233
MAG320	France	18	18-5	F83026	43.1649	6.477233
MAG335	France	18	18-3	F83026	43.1653	6.4774
MAG336	France	18	18-3	F83026	43.1653	6.4774
MAG382	France	18	18-2	F83026	43.16532	6.4774
MAG406	France	18	18-5	ESP161	43.1649	6.477233
MAG286	France	19	19-2	ESP161	43.16932	6.474433
MAG342	France	19	19-3	ESP161	43.16935	6.47445
MAG384	France	19	19-2	F83026	43.16932	6.474433
MAG386	France	19	19-4	F83026	43.16932	6.474483
MAG409	France	19	19-1	F83026	43.16935	6.474417
MAG258	France	20	20-2	F83026	43.53285	6.567617
MAG259	France	20	20-2	F83026	43.53285	6.567617
MAG261	France	20	20-2	F83026	43.53285	6.567617
MAG322	France	20	20-3	ESP161	43.53283	6.567467
MAG325	France	20	20-3	ESP161	43.53283	6.567467
MAG358	France	20	20-1	F83026	43.53403	6.572617
MAG417	France	20	20-3	F83026	43.53283	6.567467
MAG739A	France	20	20-2	F20093	43.53285	6.567617
MAG740A	France	20	20-1	F20093	43.53403	6.572617
MAG749A	France	20	20-2	F83026	43.53285	6.567617
MAG749B	France	20	20-2	F83026	43.53285	6.567617
MAG749C	France	20	20-2	F83026	43.53285	6.567617
MAG753A	France	20	20-3	F20043	43.53283	6.567467
MAG753B	France	20	20-3	F20043	43.53283	6.567467
MAG761A	France	20	20-4	F20043	43.53287	6.566783
MAG10	Spain	8	8-1	ESP174	38.41355	-1.01468
MAG21	Spain	8	8-1	ESP174	38.41355	-1.01468
MAG221	Spain	8	8-4	ESP174	38.4116	-1.01507
MAG230	Spain	8	8-3	ESP174	38.41155	-1.0165
MAG231	Spain	8	8-3	ESP174	38.41155	-1.0165
MAG238	Spain	8	8-2	ESP174	38.41253	-1.01568
MAG242	Spain	8	8-2	ESP174	38.41253	-1.01568
MAG243A	Spain	8	8-2	ESP174	38.41253	-1.01568
MAG243B	Spain	8	8-2	ESP174	38.41253	-1.01568
MAG33	Spain	8	8-4	ESP174	38.4116	-1.01507
MAG35	Spain	8	8-4	ESP174	38.4116	-1.01507
MAG66	Spain	8	8-3	F20050	38.41155	-1.0165
MAG79	Spain	8	8-3	F20050	38.41155	-1.0165
MAG83	Spain	8	8-3	ESP174	38.41155	-1.0165

MAG9	Spain	8	8-1	ESP174 nod	38.41355	-1.01468
MAG192C	Spain	9	8-3	F20050	36.91188	-3.47665
MAG192D	Spain	9	9-3	F20050	36.91188	-3.47665
MAG25A	Spain	9	9-3	ESP174	36.91188	-3.47665
MAG25B	Spain	9	9-3	ESP174	36.91188	-3.47665
MAG31	Spain	9	9-1	ESP174 no	36.91083	-3.47712
MAG746A	Spain	9	9-3	F20093	36.91188	-3.47665
MAG746B	Spain	9	9-3	F20093	36.91188	-3.47665
MAG121	Spain	11	11-3	ESP174	36.93672	-4.3563
MAG146	Spain	11	11-3	F83026	36.93672	-4.3563
MAG148A	Spain	11	11-3	F83026	36.93672	-4.3563
MAG148B	Spain	11	11-3	F83026	36.93672	-4.3563
MAG154	Spain	11	11-1	F83026	36.93678	-4.35608
MAG157	Spain	11	11-1	F83026	36.93678	-4.35608
MAG174	Spain	11	11-1	ESP174	36.93678	-4.35608
MAG176A	Spain	11	11-1	ESP174	36.93678	-4.35608
MAG176B	Spain	11	11-1	ESP174	36.93678	-4.35608
MAG748A	Spain	11	11-1	ESP174	36.93678	-4.35608
MAG748B	Spain	11	11-1	ESP174	36.93678	-4.35608
MAG755A	Spain	11	11-1	F83026	36.93678	-4.35608
MAG755B	Spain	11	11-1	F83026	36.93678	-4.35608
MAG757A	Spain	11	11-3	F20093	36.93672	-4.3563
MAG757C	Spain	11	11-3	F20093	36.93672	-4.3563
MAG758A-mel	Spain	11	11-2	F83026	36.93678	-4.35608

Table S2. SNP counts for each element of the symbiont genome before any variant filtering, and after filtering for quality and depth using VCFtools.

	Chromosome	pSymA	pSymB
Pre-filtering	414,004	179,505	222,935
Quality filters	34,689	15,162	22,460

Table S3. Matrices of population pairwise D_{XY} values for each *E. meliloti* genomic element. The populations are arranged from west to east (Spain-mainland France-Corsica). Cells background coloration indicates the level of D_{XY} (red = low values, green = high values).

***See supplementary Excel file*

Supplemental figure legends

Figure S1. Map of 25 sampling locations in Spain, mainland France, and Corsica. Grey points with black outlines represent sites where both host and symbiont were sampled (n = 8), grey points represent sites where only symbionts were sampled (n = 13), and black circles represent sites where only hosts were sampled (n = 4).

Figure S2. Principal component axis plots showing genome wide similarity on PCs 1-3 for the *Medicago* host plants (n=192) (**A**) and for symbiont (n=191) chromosome (**B**), pSymA (**C**), and pSymB (**D**). The darkest ellipse and points on each plot represent individuals from Spain, the intermediate color dashed line represent individuals from mainland France, and the lightest color represents individuals from Corsica.

Figure S3. Neighbor joining tree of rhizobium individuals based on pSymB variant data. Individual tip labels (strain ID followed by soil population) are colored based on region of origin (orange from Corsica, green from mainland France, blue from Spain). Branch support indicated in teal (<70) or pink (<50). Outer ring represents the chromosomal cluster (from Fig. 2a) for comparison.

Figure S4. Principal component axis plot for all individuals of the symbiont (n=191) based on variable gene content. From top to bottom principal component 1 and 2, 1 and 3, and 2 and 3. The red ellipse and points on each plot represent individuals from Spain, the green dotted line and points represent individuals from mainland France, and the blue points and line represent individuals from Corsica.

Figure S5. Tests of correlated population structure between the genomic elements in *E. meliloti* and the host plant *M. truncatula*

Figure S1.

Figure S2.

Figure S3.

0.03

Figure S4.

Figure S5.