1   Partners in space: Discordant population structure between legume hosts and rhizobium

2   symbionts in their native range

4   Alex B. Riley[1], Michael A. Grillo[2], Brendan Epstein[3], Peter Tiffin[3], Katy D. Heath[1,4,+]

5
Author affiliations:

[1]University of Illinois, Department of Plant Biology, 505 S. Goodwin Ave. Urbana IL 61801

[2]Loyola University Chicago, Department of Biology, 1032 W. Sheridan Rd, Chicago, IL,

60660, USA

[3]University of Minnesota, Department of Plant and Microbial Biology, 140 Gortner Lab, 1479

Gortner Ave, St. Paul, MN 55108

[4]University of Illinois, Carl R. Woese Institute for Genomic Biology, 1206 W. Gregory Dr.,

Urbana, IL 61801 USA

[+]Corresponding author; kheath@illinois.edu

15
16   Running head: Discordant population structure in symbiosis

17   Keywords: coevolution, mutualism, horizontal gene transfer, divided genome

19 **Abstract**
20

Coevolution is predicted to depend on how the genetic diversity of interacting species is

geographically structured. Plant-microbe symbioses such as the legume-rhizobium mutualism are

ecologically and economically important, but distinct life history and dispersal mechanisms for

these host and microbial partners, plus dynamic genome composition in bacteria, present

challenges for understanding spatial genetic processes in these systems. Here we study the model

rhizobium *Ensifer meliloti* using a hierarchically-structured sample of 191 strains from 21 sites

in the native range and compare its population structure to that of its host plant *Medicago*

truncatula. We find high local genomic variation and minimal isolation by distance across the

rhizobium genome, particularly at the two symbiosis elements pSymA and pSymB, which have

evolutionary histories and population structures that are similar to each other but distinct from

both the chromosome and the host. While the chromosome displays weak isolation by distance, it

is uncorrelated with hosts. Patterns of discordant population structure among elements with the

bacterial genome has implications for bacterial adaptation to life in the soil versus symbiosis,

while discordant population genetic structure of hosts and microbes might restrict local

adaptation of species to each other and give rise to phenotypic mismatches in coevolutionary

traits.

Introduction

Coevolution is fundamentally a geographically variable process. This understanding has

become widely accepted over the past several decades, most prominently with the inception of

Thompson's geographic mosaic theory of coevolution (Thompson, 1994, 2005). Coevolving

partners can vary in their distributions, abundance, and genetic structure across the landscape and

over time (Carlsson-Granér & Thrall, 2015; Fernandes, Lemos-Costa, Guimarães, Thompson, &

de Aguiar, 2019; Laine, 2005; Tack, Horns, & Laine, 2014; Thompson, 2005). The evolution of

interacting species will depend on spatial variation in selection, genetic variation, and gene flow.

While there is abundant evidence for the core predictions of the GMTC (*i.e.,* selection mosaics,

coevolutionary hot/cold spots, variable trait remixing) it is surprising that comparative analysis

of population genetic structure among interacting partners has received relatively little attention

(but see Caldera and Currie 2012; Baums et al. 2014; Strobel et al. 2016; Harrison et al. 2017).

Indeed, genetic structure can, in part, determine the outcomes of coevolution. For

example, alignment in the geographic scale and location of population genetic structure between

interacting species can allow for reciprocal selection and adaptation (*i.e.,* coevolutionary

hotspots); conversely, discordant population structure, due to different levels of gene flow or

other processes, can limit the ability of each species to locally adapt to the other and give rise to

trait mismatches (Fernandes et al., 2019; Gandon, Capowiez, Dubois, Michalakis, & Olivieri,

1996a; Thompson, 2005). Thus, understanding the comparative population genetic structure of

interacting species provides a key perspective on whether traits are expected to coevolve and

how coevolutionary trait variation is maintained through time and space (Heath and Stinchcombe

2014; Hollowell et al. 2016a; Stoy et al. 2020).

Coevolutionary interactions typically occur between species with distinct life histories,

range sizes, mating systems, and modes of dispersal ☐ all of which will impact population

genetic structure (Pita, Rix, Slaby, Franke, & Hentschel, 2018; Revillini, Gehring, & Johnson,

2016; Thrall, Hochberg, Burdon, & Bever, 2007). In general, the majority of coevolutionary

research has focused on systems involving antagonistic coevolution of eukaryotes. Mutualisms

between macro and micro-organisms have received less attention, even though these mutualisms

play critical roles in community and ecosystem processes (Berg, 2009; Revillini et al., 2016;

Reynolds, Packer, Bever, & Clay, 2003).

The investigation of host-microbe interactions poses various logistical and

methodological challenges. One challenge is the difficulty of isolating populations of a single

species from the environment - particularly for soil bacteria in hyper-diverse communities (Heath

& Grillo, 2016). In addition, many bacterial genomes are multipartite (Harrison et al. 2010;

diCenzo and Finan 2017), including non-chromosomal replicons containing variable amounts of

genetic information – ranging from small facultative plasmids that are often lost or transferred to

large obligate segments containing core genes (often as megaplasmids or chromids; Harrison et

al. 2010). Such divided genomes might represent an adaptation allowing functional division

between replicons (reviewed by diCenzo and Finan 2017). Multipartite genome organization

presents a challenge in the study of bacterial (co)evolution because different elements may move

between individuals independent of each other, resulting in distinct evolutionary histories and

population structure across different elements. These differences are compounded by the fact that

these elements can also have widely differing rates of evolution and patterns of recombination

leading to higher rates of evolution in plasmids not required for survival (Cooper, Vohr,

Wrocklage, & Hatcher, 2010; Epstein et al., 2012; Epstein, Sadowsky, & Tiffin, 2014; Epstein &

Tiffin, 2021).

The legume-rhizobium mutualism presents a tractable model system for studying

microbial symbiosis due to the relative ease of isolating populations and genetic resources for

several legume-rhizobium partnerships (Kaneko et al., 2002; Yates et al., 2015). In this

symbiosis, rhizobia form nodules on plant roots, wherein they fix atmospheric N in exchange for

carbon derived from plant photosynthesis; however, rhizobia inhabit the soil as saprophytes

when not in symbiosis with a host, thereby experiencing profoundly different environmental

conditions and likely selective pressures (Burghardt, 2020). In rhizobia with divided genomes

(e.g., *Rhizobium* and *Ensifer* spp.), genes for symbiosis are carried on symbiosis elements (one or

more megaplasmids or chromids), while the chromosome is thought to be responsible for core

metabolic functions. These elements can have distinct evolutionary histories, diversity levels,

and recombination rates (e.g., Bailly et al. 2011; Epstein et al. 2012, 2014; Cavassim et al. 2020;

Epstein and Tiffin 2021).

Here we examine population structure in *Ensifer meliloti* (Becker et al., 2009; Tang et al.,

2014), the rhizobium symbiont of the model legume *Medicago truncatula* (hereafter

"*Medicago*"). *Medicago* is a self-fertilizing annual native to the Mediterranean region of Europe

(Bonnin, Huguet, Gherardi, Prosperi, & Olivieri, 1996; Siol, Prosperi, Bonnin, & Ronfort, 2008).

Research involving *Medicago* population genetics has identified a pattern of isolation by

distance, and genetic structure at both the population and regional levels (Bonhomme et al.,

2015; Bonnin et al., 1996; Grillo, De Mita, Burke, Solórzano-Lowell, & Heath, 2016; Ronfort et

al., 2006; Siol et al., 2008). Here we use whole genome sequences for 191 isolates of *E. meliloti*

sampled from 21 sites in the native range to characterize the population structure of this model

mutualist and compare it to the host by reanalyzing *Medicago* sequence data from Grillo et al.

(2016) to directly compare population structure in the two species. We ask: 1) how is genetic

variation structured in *E. meliloti*; 2) do the three genomic elements of the rhizobium *E. meliloti*

share common patterns of evolutionary history and population structure; and 3) do *Medicago*

hosts and their rhizobium symbionts have correlated population genetic structure?

Methods:

Study system:

Ensifer (formerly *Sinorhizobium*) *meliloti* is a rhizobium species in the Alphaproteobacteria

(Young & Haukka, 1996) that forms N-fixing nodules on the roots of multiple species in the

genus *Medicago*, and is one of two *Ensifer* species that forms root nodules on *M. truncatula*

(Zribi, Mhamdi, Huguet, & Aouani, 2004). The genome of *E. meliloti* is ~6.79Mb, divided

between three genomic elements: the chromosome (3.69Mb), megaplasmid pSymA (1.41Mb),

and chromid pSymB (1.69Mb), as well as smaller plasmids in some strains (Galibert et al. 2001,

Nelson et al. 2018). Metabolic modeling, along with genetic manipulation of the *E. meliloti*

genome, have shown that gene content differs functionally between the three genomic elements

(diCenzo, MacLean, Milunovic, Golding, & Finan, 2014; Galibert et al., 2001). The chromosome

is predicted to carry primarily genes related to core metabolic function in soil, pSymA carries the

majority of genes required for symbiotic N fixation, and pSymB carries primarily genes thought

to be important for life in rhizosphere environments (diCenzo et al., 2014).

Sample collection: We isolated *E. meliloti* strains from 21 sites in the native range of *Medicago*

using a hierarchical sampling design with populations ranging from 1 to approximately 1350 km

apart (Fig. S1; Table S1). Of the 21 sites, eight corresponded to the *Medicago* sites studied in

Grillo *et al*. (2016). At each site we collected soil surrounding the top six inches of the root

system from unearthed host plants. In order to generate a hierarchical sampling design, with

individual strains ranging from 0-1350km apart (sites ranging from 1-1350km apart), we

sampled multiple plants from some sites (see table S1). To avoid cross-contamination within

each site, the sampling shovel was wiped clean of excess soil between samples and was pierced

into the ground adjacent to a plant numerous times before sampling soil. Between sampling

locations, the shovel was sterilized with dilute bleach. Soil samples were kept at 4□ prior to

isolating cultures.

Ensifer strains were isolated or "trapped" in the laboratory from the field samples

following standard protocols (Vincent 1970; Heath 2010). In brief, *Medicago* seeds were nicked

with a razor blade, surface sterilized with 30% bleach, rinsed with sterile water, and imbibed in

sterile water for approximately 30 minutes. Seeds were then directly sown into a given soil

sample housed in a sterilized, fully self-contained Magenta box (see Brown et al. 2020 for

details). Magenta boxes were randomly placed in a temperature-controlled grow room (23° C)

under artificial light set to 12-h□days. After four weeks, plants were harvested, and the soil was

washed from the roots. Individual nodules were removed with forceps, surface sterilized by

soaking in 30% bleach for 10 minutes, and then rinsed with sterilized water. Surface sterilized

nodules were crushed with sterilized forceps and streaked on tryptone-yeast (TY) media plates.

Plates were incubated at 30□ for 48 hours, sterilized glass stir rods were then used to streak

samples were streaked on to TY plates which were again incubated at 30□. Individual colonies

were then picked and grown in liquid TY media, and these pure cultures were stored in cryotubes

in 50% TY 50% glycerol and stored at -80□. Given variation among host genotypes in

rhizobium infection rates due to G x G interactions (Batstone, Dutton, Wang, Yang, &

Frederickson, 2017; Heath & Tiffin, 2009), we used 24 *Medicago* host genotypes to trap

rhizobia. Because both *E. meliloti* and *E. medicae* infect the roots of *Medicago* in the native

range, we used a post-PCR restriction enzyme (RsaI) digestion of the 16S to assign strains to

species (following Biondi et al. 2003). Ultimately, we isolated 191 strains of *E. meliloti*

presented in the current study. We hereafter refer to *E. meliloti* as simply *"Ensifer"* or

"symbiont".

Sequencing: We extracted DNA from cultures of *Ensifer* grown in liquid TY media using the

Qiagen DNeasy kit (Hilden, Germany) and sent samples to the DOE Joint Genome Institute

(JGI) for sequencing (Berkeley, CA, USA). JGI prepared a paired end sequencing library for

each strain, and sequenced samples on an Illumina HiSeq-2500 1TB platform (101nt read length;

Illumina, Inc., San Diego, CA, USA). Of the 199 strains submitted to JGI, we received high

quality whole genome sequences for 166. We re-grew the remaining 33 strains from frozen

cultures (as above), extracted DNA using the Zymo Quick-DNA kit for Fungi or Bacteria

(Irvine, CA, USA). These samples were sequenced (2 X 150 or paired end 150 nt read length) on

the Novaseq 6000 platform (Illumina, Inc, San Diego, CA, USA) by the Roy J. Carver

biotechnology center at the University of Illinois at Urbana-Champaign (USA). We successfully

recovered quality genome sequences from 25 of these 33 isolates, for a total of 191 *E. meliloti*

strains sequenced.

Genome assembly, annotation, and SNP calling: To ensure high quality SNP calling and genome

assembly, we trimmed PCR adaptors and removed PCR duplicates and PhiX contamination

using HT-Stream (github.com/s4hts/HTStream), followed by further adaptor removal, removal

of bases with quality scores < 30 from the ends of reads, and removal of reads < 80 bp long with

TrimGalore! (github.com/FelixKrueger/TrimGalore). For calling SNPs in core genes (genes

present in > 80% of strains), we aligned reads to the *E. meliloti* reference genome USDA1106

using BWA with default settings (Li & Durbin, 2009). We then used Freebayes (Garrison &

Marth, 2012) to identify haplotype variants, which we split into SNPs using VCFtools (Danecek

et al., 2011). We found 491,277 variable SNPs with variant qualities above 20. We then filtered

SNPs to retain only those with depth values between 20 and 230, minor allele frequencies ≥

0.009 (*i.e.,* present in at least two strains), and that were present in at least 80% of individuals,

72,311 SNPs remained after applying these filter (chromosome: 34,689 SNPs, pSymA: 15,162

SNPs, pSymB: 22,460 SNPs; see Table S2). For variable gene content, we assembled genomes

de novo using SPADES (Bankevich et al., 2012) with default parameters, followed by annotation

using PROKKA (Seemann, 2014) and scanned these annotations for presence-absence variants

using default settings in ROARY (Page et al., 2015).

Phylogenetic and Statistical Analyses: To explore the population structure of the three genomic

elements of *Ensifer,* we first used principal components analysis (PCA). We used the glPca

function in the adegenet (Jombart & Ahmed, 2011) library in R on a random subsample of 15k

SNPs (to equilibrate dataset size) for each element of the *Ensifer* genome to naively cluster

individuals by genome-wide similarity. We then plotted the positions of individuals along the

first three axes of variation (using ggplot2; Fig. S2) to visually compare how genetic variation is

distributed across elements (Wickham, 2016). To quantify genetic differences we calculated

individual and population-level Dxy (Nei 1972) using the stamppNeisD function from the

package StAMPP (Pembleton, Cogan, & Forster, 2013) on all variants for each element. To test

whether these patterns of genetic distance among *Ensifer* strains were congruent across the three

genome elements, we used these matrices of individual-based distance metrics in pairwise

Mantel tests comparing the three *Ensifer* genome elements (chromosome vs. pSymA,

chromosome vs. pSymB. pSymA vs. pSymB). Mantel tests were implemented in the R package

ade4 (Dray & Dufour, 2007). To explore geographic structure in the variable genome (*i.e.,* genes

present in only some strains), we performed an additional genetic PCA, as above, using the

matrix of *Ensifer* gene presence-absence variants (91,840 genes) and plotted the first three

principal components for visual inspection.

For each of the three elements, we built phylogenies based on the same random

subsample of 15k core genome SNPs used for PCA. We used the neighbor joining (nj) function

in the R package ape (Paradis & Schliep, 2019) with 1000 bootstrap replicates and visualized

trees in FigTree (v.1.4.4) (Rambaut, 2018). Based on these preliminary results, we treated each

element independently for remaining analyses.

We used AMOVA (poppr.amova) with clone correction and 10,000 random permutations

(Kamvar, Tabima, & Gr□unwald, 2014) to partition the genetic variance among individuals,

sites, and regions, as well as to assess the statistical significance of these levels of spatial genetic

structure. To test the hypothesis of isolation by distance, we calculated Pearson correlations

between individual-level D_{XY} and geographic distances between sampling sites.

To compare the spatial genetic structure of rhizobia and host plants, we reanalyzed RAD-

seq data from the 192 *Medicago* genotypes studied in Grillo *et al* (2016). We first called SNPs

using Stacks with default parameters (Catchen, Hohenlohe, Bassham, Amores, & Cresko, 2013),

then filtered the resulting variants using VCFtools (Danecek et al., 2011) to ensure that all

variants were present in at least 80% of lines, had minor allele frequencies > 0.05, and were $>$

5kb apart, 10,814 SNPs remained after these filters. We performed AMOVA, calculated

individual and population-based D_{XY}, and tested for isolation by distance in *Medicago* as

detailed above for *Ensifer*. For the subset of eight sites for which we had both hosts and

symbionts, we used Mantel tests to ask whether the three matrices of pairwise population D_{XY}

values from symbionts (chromosome, pSymA, and pSymB) were correlated with that of the host,

to test for congruent population genetic structure between hosts and symbionts.

Results

To address the extent of phylogenetic congruence between the chromosome, pSymA, and

pSymB of *Ensifer*, we first used pairwise Mantel tests of whether the matrices of genetic

distances were correlated. At the individual level, we found non-significant correlations between

the chromosome and pSymA and between the chromosome and pSymB, while the two symbiosis

elements (pSymA and pSymB) were more strongly correlated (left column, Fig. 1), indicating

distinct evolutionary histories of the chromosome versus pSymA and pSymB. Neighbor joining

trees also reveal distinct evolutionary histories among the three genome elements. The topology

of the chromosomal tree showed five tightly clustered groups of individuals, with almost all

strains from Corsica closely related and forming a distinct group (orange cluster; Fig. 2a) and

strains from Spain somewhat interspersed with those from France mostly in the blue and yellow

clusters. The diversity of strains found in mainland France (the best sampled region) included

representatives from each major chromosomal lineage; indeed, pink and purple clusters were

found only in mainland France (except strain 710A from Corsica), mostly on the western

Mediterranean coast (Fig. S1; Table S1).

Using these same colors (corresponding to the chromosomal clusters) to annotate the

pSym trees (Fig. 2b; Fig. S3) facilitates visualization of differences in tree topology among the

three elements, indicative of plasmid transfer (whole or partial) among chromosomal lineages.

The internal branches of both symbiotic element trees reveal greater diversity, with chromosomal

lineages spread across diverse pSym lineages in both trees (colored outer rings in Fig. 2b; Fig.

S3). Some strains were closely related at all three elements of their genomes; for example, the

group of closely related chromosomal lineages in pink from the western coast of France appear

together in all three trees, indicating little pSym element recombination among these and other

lineages (strain 78 in the pSymB tree is a notable exception). On the other hand, the other

chromosomal clusters (purple, orange, and yellow clusters in Fig. 2a), appear throughout the

pSymA and pSymB tree (see same colors interspersed; Fig. 2b; Fig. S3), indicating that these

chromosomal lineages are found with diverse pSym genotypes. For example, the tightly

clustered group of closely related chromosomal lineages in orange were found with lineages

from across both the pSymA and pSymB trees.

Next, we used genetic PCA to explore population structure and AMOVA to partition the

genome-wide genetic variation at the within-population, among-population, and among-region

scales. Genetic PCA did not reveal strong differentiation in *Ensifer*, even at the regional scale, in

either the core genome SNPs or variable gene content (Fig. S2; Fig. S4). Nevertheless, we found

significant spatial genetic structure in the core genome at all scales investigated using AMOVA

(Table 1), but that the patterns varied among the three *Ensifer* elements. For all three, we found

considerable variation (66-76%) maintained within populations, compared to the among-

population and among-region scales (Table 1). All three among-element correlations were

stronger at the among-population level (right column, Fig. 1), revealing substantial congruence

between pSymA and pSymB, and weakly correlated population structure between the

chromosome and both symbiosis elements (right column, Fig. 1). The discrepancy between

individual and population level correlations is likely due to high within population diversity

(Table 1) and several populations with near-zero genetic distances (*e.g.,* within Spain and

France, and between them Table S3).

The *Ensifer* chromosome showed the strongest differentiation at the regional scale (*i.e.,*

among Spain, France, and Corsica; Table 1). This finding is likely driven by Corsica, as short

branch lengths were found among most Corsica strains on the chromosomal tree (within the

orange cluster; Fig 2a) and genetic distances were high between Corsica populations and the rest

of the range (Table S3). The chromosome was also the only *Ensifer* element to exhibit significant

isolation by distance (IBD) at this spatial scale (Fig 3). Despite overall IBD, there was extremely

large variation in the genetic distances between *Ensifer* isolated from even the most distant

locations (*i.e.,* D_{XY} distances ranged from 0-0.5 even when strains were found approximately

1350 km apart; Table 1). The two symbiotic elements in *Ensifer* (pSymA and pSymB) were less

structured at the among-region and among-population scales (Table 1), and neither exhibited

significant IBD at our sampling scale (Fig. 3). Thus although populations differed in genetic

composition (significant structure in Table 1), *Ensifer* strains from distant populations up to 1350

km apart were often as closely related to each other as strains from the same sampling site.

Hosts and symbionts had mismatched population genetic structure. We found no

significant correlations between the pattern of among-population differentiation (D_{XY}) between

Medicago hosts and any element in the *Ensifer* genome (Fig. S5). While both the host and

Ensifer chromosome exhibit patterns of isolation by distance (see above), their spatial patterns of

genetic variation were distinct; while host plants primarily clustered along axes that separated

Spain from France and Corsica (Fig. S2; Grillo et al. 2016 and references therein), the *Ensifer*

chromosome most clearly clustered into two groups (mainland Europe versus Corsica along PC1,

Fig. S2). The lack of IBD and little regional structure in pSymA and pSymB indicate that host

plants at a given site have the potential to interact with the diversity of symbiosis plasmids from

across this portion of the species range.

Discussion

Evolution, and thus coevolution, is shaped by the interaction of selection with demographic

processes such as gene flow occurring across a species range; therefore, understanding the

maintenance of genetic variation, the evolution of mutualistic traits, and host-microbe

interactions requires that we consider individuals and species in their spatial context. Here we

used a hierarchically-structured sample of legume hosts and rhizobial symbionts to show: 1) the

elements of the bacterial genome (chromosome versus pSymA and pSymB) have distinct

population genetic structure at the local and regional scales, 2) host plants have more spatial

genetic structure than rhizobia, particularly when compared to the two symbiotic elements

(pSymA and pSymB), and 3) patterns of population structure between hosts and symbionts are

not correlated across this portion of their native range.

Spatial genetic processes at the bacterial chromosome and symbiotic elements:

The population structure and dispersal abilities of microbes remain poorly understood (Chase et

al., 2019; Green, Bohannan, & Whitaker, 2008; Hanson, Fuhrman, Horner-Devine, & Martiny,

2012; Martiny et al., 2006; VanInsberghe, Arevalo, Chien, & Polz, 2020). Sequence data have

revealed considerable variation in dispersal ability among microbial taxa (Locey & Lennon,

2016). Based on metagenomics datasets, rhizobia have many characteristics associated with large

geographic ranges and, by extension, large dispersal ability (*e.g.,* Proteobacteria, plant-associated

or soil-borne taxa, and large genome sizes; Choudoir et al. 2018). However, metagenomic

studies of microbial communities are limited in their resolution; therefore, population genomic

data (Hoetzinger, Pitt, Huemer, & Hahn, 2021), coupled with spatially-structured samples of

both within- and among-population variation (Whitaker & Banfield, 2006), are required for

inferring population structure and gene flow, which are key to local adaptation and coevolution

(Thompson 2005; Hoeksema and Forde 2008; Kraemer and Boynton 2017). Like other diverse

microbial systems (Chase et al., 2019; Hoetzinger et al., 2021; Vos & Velicer, 2008), *Ensifer*

displays both isolation by distance (chromosome) and population genetic structure (all elements);

nevertheless, we found extremely closely-related strains up to 1350 km apart – indicating at least

some long-distance dispersal. Large dispersal ability, together with the observation of abundant

local phenotypic (Heath, 2010; Heath & Tiffin, 2009) and genomic (Bailly et al., 2011; Bailly,

Olivieri, De Mita, Cleyet-Marel, & Béna, 2006) variation for symbiosis*,* suggests that most of

the variation relevant to both bacterial adaptation and coevolution can be found at relatively

small spatial scales.

At the regional scale sampled here, we identify distinct evolutionary histories and

population genetic structures between the chromosome and the symbiosis elements, but similar

population structure between the two symbiosis elements. Previous work has shown divergent

evolutionary histories of the three elements in this species (Bailly et al. 2006; Galardini et al.

2013; Epstein et al. 2014; Nelson et al. 2018), as well as in other rhizobium taxa with separate

symbiosis plasmids (Cavassim et al., 2020; Klinger, Lau, & Heath, 2016; Koppell & Parker,

2012; Kumar et al., 2015; Pérez Carrascal et al., 2016; Young et al., 2006) or symbiosis gene

regions (Amanda C. Hollowell et al., 2016; Porter, Faber-Hammond, Montoya, Friesen, &

Sackos, 2019). Our analyses support these findings while also revealing how the three elements

differ at the within-population, among-population, and regional scale using a hierarchically-

structured spatial sample of within and among-site variation. Using this approach allowed us to

resolve distinct fine-scale spatial genetic processes at individual elements, namely

geographically limited chromosomal lineages (e.g., Corsica), but little population structure of

symbiosis elements.

One hypothesis for the existence of distinct elements in bacteria is to break up genetic

correlations among traits that are important during different phases of the life cycle and/or

establish linkage between adaptive gene complexes (diCenzo & Finan, 2017). Based on

functional genetics and metabolic models (Galibert et al. 2003; diCenzo et al. 2014; diCenzo and

Finan 2017), the three major genome elements are thought to play somewhat distinct roles in the

life history of *Ensifer:* chromosome (basic metabolism), pSymA (nodulation and N fixation), and

pSymB (rhizosphere). The traits, and potentially genes, that underlie fitness in the soil,

rhizosphere, and nodules are likely distinct (Friesen and Mathias 2010; Sachs et al. 2011; Heath

and Stinchcombe 2014; Burghardt 2019); having these processes segregated into distinct genome

elements might allow the buildup of coadapted gene complexes within elements while

facilitating independent adaptation to host and abiotic conditions on separate elements. One

particularly interesting geographic disjunction that distinguishes the chromosome from the other

elements is a cluster of extremely closely-related lineages that was nearly ubiquitous in Corsica

but found elsewhere in low frequency. Although it remains unclear how this pattern arose, it is

possible that this chromosomal lineage has swept to high frequency as the result of recent

selection. Epstein et al. (2012) found low diversity along half of the *E. meliloti* chromosome in a

range-wide sample of 24 strains, likely indicating a recent selective sweep with little

recombination. Given little recombination on the chromosome (Epstein et al. 2012; Nelson et al.

2018), targets of selection on this element can be difficult to distinguish. Nevertheless it would

be interesting to test whether the three genome elements of *Ensifer* are locally adapted in ways

consistent with their functional gene composition, possibly by using host genetic variation

alongside abiotic climatic factors in a landscape genomics model (e.g., Yoder et al. 2014;

Rudman et al. 2018).

Our work suggests that pSymA and pSymB might be inherited together, leading to strong

correlations between them at both the individual and population scales. Recombination in *Ensifer*

is known to occur through conjugation of pSymA and pSymB (Ding & Hynes, 2009). Recent

work indicates that pSymB is mobilized by pSymA via a *rctA*-mediated control system, though

tracking these events in real time has proven elusive in the laboratory (Blanca-Ordóñez et al.,

2010; diCenzo & Finan, 2017; Galibert et al., 2001; Pérez-Mendoza et al., 2005; Pretorius-Guth,

Puhler, & Simon, 1990). If pSymA and pSymB are coupled in nature, this could give rise to

genetic constraint that could affect *Ensifer*'s ability to adapt to distinct selective forces arising

from various environmental conditions encountered in the rhizosphere versus in association with

host plants. Functional predictions (*e.g.,* gene knockout experiments, functional annotation based

on homology) do not always predict quantitative trait variation; therefore, hypotheses regarding

genetic constraint among the elements should be addressed with quantitative genetic correlations

between rhizosphere and symbiosis traits (e.g., Ossler and Heath 2018; Wood et al. 2018).

Distinct spatial genetic processes between the chromosome and symbiosis elements, and

the coupling of the symbiosis elements, begs the question of how horizontal transmission of the

symbiosis elements and recombination generate population structure in these bacteria. While we

summarize the geographical patterns at the whole-element scale in order to compare these to the

host (see below), finer-scaled work on gene tree heterogeneity (Degnan & Rosenberg, 2009)

within each of the three elements is required for a full understanding of whether individual loci,

large genomic regions, or even entire plasmids are introgressed through conjugation, and at what

spatial scales such events occur. For example, Porter et al. (2019) recently used fine-scale

patterns of gene co-occurrence across *Mesorhizobium* strains from California to discover the

local transfer and duplication of symbiosis island and associate this variation with symbiotic

partner quality.

Mismatched host-symbiont population structure: Our results indicate discordance in population

structure between hosts and symbionts due to differences in drift, gene flow, mutation, and/or

spatially-variable selection. *Medicago* host plants are most structured along a longitudinal

gradient, indicative of distinct glacial refugia, dividing populations in Spain from France and

Corsica (Branca et al., 2011; Grillo et al., 2016; Ronfort et al., 2006; Yoder et al., 2014). By

contrast, we found that most genome-wide variation in *Ensifer* is found within individual sites,

and our phylogenetic and spatial analyses indicate that regional differentiation and IBD (which

was significant for the chromosome) are different from host plants, since chromosomes from

Corsica were distinct from most mainland populations (see discussion above). In contrast to host

plants and the *Ensifer* chromosome, pSymA and pSymB showed substantially less regional

differentiation and no pattern of IBD, due to increased introgression following long-range

dispersal events, higher effective population size, or both.

A striking feature of mutualisms is the existence of abundant genetic variation for

mutualistic traits within and among populations, given models that predict the erosion of trait

variation in mutualism (Heath and Stinchcombe 2014; Stoy et al. 2020). Indeed considerable

genetic variation in partner choice, sanctioning, signaling, and partner quality has been identified

in the *Medicago-Ensifer* mutualism (Batstone et al., 2017; Batstone, O'Brien, Harrison, &

Frederickson, 2020; Burghardt et al., 2018; Burghardt, Epstein, & Tiffin, 2019; Heath, 2010;

Heath & Tiffin, 2009). A mismatch in the population genetic structure of interacting species, as

we identified here, could restrict populations of host and symbionts from locally adapting to each

other and thereby contribute to the maintenance of coevolutionary trait variation. Thus strong

local selection would be necessary to overcome the homogenizing effects of gene flow (*i.e.,*

migration-selection balance; Savolainen et al. 2013) in order for hosts and symbionts to locally

adapt to each other. Empirical work in other systems comparing the population structure of

interacting species finds a range of outcomes, from correlated spatial genetic patterns between

partners (Anderson, Olivieri, Lourmas, & Stewart, 2004; Caldera & Currie, 2012; Smith,

Godsoe, Tank, Yoder, & Pellmyr, 2008; Thompson, Thacker, & Shaw, 2005), to largely

discordant patterns, where one species exhibits substantially less population structure than its

partner (Baums et al., 2014; Dybdahl & Lively, 1996; Strobel et al., 2016).

Spatially-explicit theory on mutualism coevolution indicates that relative rates of gene

flow between hosts and symbionts alters the likelihood of trait matching/local coadaptation, and

even that gene flow can help explain the existence of maladaptation and abundant genetic

variation for mutualism (Nuismer, Thompson, & Gomulkiewicz, 2003; Parker, 1999; Yoder &

Nuismer, 2010). An important finding of these models, however, is that the outcomes of

coevolution depend on how gene flow interacts with forms of natural selection, yet mutualism

research often lacks explicit information on how selection in the wild acts on mutualism traits

like partner quality and partner choice (Heath and Stinchcombe 2014; Stoy et al. 2020).

The degree of fitness alignment in mutualism, and thus the degree to which these

interactions coevolve like better-studied antagonisms, is an unresolved question (Batstone et al.,

2020; Frederickson, 2017; Maren L. Friesen, 2012; Gano-Cohen et al., 2020; Jones et al., 2015;

Sachs, Quides, & Wendlandt, 2018). For antagonistic interactions, which are governed by

negative frequency-dependent selection due to interspecific conflict, models suggest that the

partner with a higher migration rate will benefit from the influx of new, potentially adaptive

alleles that provide a competitive edge (Carlsson-Granér & Thrall, 2015; Gandon, Capowiez,

Dubois, Michalakis, & Olivieri, 1996b). If conflict prevails in mutualisms, and increases in

rhizobium fitness occur at the expense of the host (Gano-Cohen et al., 2020; Porter & Simms,

2014), incongruent population structures between hosts and symbionts should make it harder for

hosts to evolve mechanisms to exclude lower-quality symbionts (*e.g.,* partner choice; Akçay

2017; Younginger and Friesen 2019), though this depends on the strength of selection. On the

other hand, alignment between host and symbiont fitness might be common in mutualisms, as

suggested by recent lab and experimental evolution studies (Batstone et al., 2020; Friesen, 2012).

This scenario should ultimately result in positive frequency-dependent selection wherein novelty

is disfavored (unless populations are far from their adaptive peak), leading to stable allele

frequencies under purifying selection. Mismatched population structure, as identified here,

would restrict the efficacy of positive frequency-dependence and any resulting local adaptation

between hosts and symbionts. Our finding of discordant population genetic structure is in

agreement with other host-microbe symbioses where microbes are transmitted horizontally and

independently of their hosts (Dybdahl and Lively 1996; Baums et al. 2014; Strobel et al. 2016;

Harrison et al. 2017). If this is common, then coevolutionary selection would have to be quite

strong to overcome the gene flow to generate patterns of local coadaptation.

Acknowledgments:

We thank Laurène Gay and Joelle Ronfort for advice and contributions to field collection, and

Amy Marshall-Colón, Julian Catchen, and Rachel Whitaker for feedback on an early manuscript

draft. We thank various sources of funding including NSF IOS-1401864 to MAG, NSF IOS-

1645875 to KDH, NSF PGRP-1856744 to PT and KDH, JGI CSP-503446 to PT and KDH, and

the Department of Plant Biology at the University of Illinois Urbana-Champaign.

References:

Akçay, E. (2017). Population structure reduces benefits from partner choice in mutualistic symbiosis. *Proceedings of the Royal Society B: Biological Sciences*, *284*(1850). doi: 10.1098/rspb.2016.2317

Anderson, B., Olivieri, I., Lourmas, M., & Stewart, B. A. (2004). Comparative population genetic structures and local adaptation of two mutualists. *Evolution*, *58*(8), 1730–1747. doi: 10.1111/j.0014-3820.2004.tb00457.x

Bailly, X., Giuntini, E., Sexton, M. C., Lower, R. P. J., Harrison, P. W., Kumar, N., & Young, J. P. W. (2011). Population genomics of Sinorhizobium medicae based on low-coverage sequencing of sympatric isolates. *ISME Journal*, *5*(11), 1722–1734. doi: 10.1038/ismej.2011.55

Bailly, X., Olivieri, I., De Mita, S., Cleyet-Marel, J. C., & Béna, G. (2006). Recombination and selection shape the molecular diversity pattern of nitrogen-fixing Sinorhizobium sp. associated to Medicago. *Molecular Ecology*, *15*(10), 2719–2734. doi: 10.1111/j.1365-294X.2006.02969.x

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., … Pevzner, P. A. (2012). SPAdes: A new genome assembly algorithm and its applications to single-cell sequencing. *Journal of Computational Biology*, *19*(5), 455–477. doi: 10.1089/cmb.2012.0021

Batstone, R. T., Dutton, E. M., Wang, D., Yang, M., & Frederickson, M. E. (2017). The evolution of symbiont preference traits in the model legume Medicago truncatula. *New Phytologist*, *213*(4), 1850–1861. doi: 10.1111/nph.14308

Batstone, R. T., O'Brien, A. M., Harrison, T. L., & Frederickson, M. E. (2020). Experimental evolution makes microbes more cooperative with their local host genotype. *Science*, *370*(6515), 23–26. doi: 10.1126/science.abb7222

Baums, I. B., Devlin-Durante, M. K., & Lajeunesse, T. C. (2014). New insights into the dynamics between reef corals and their associated dinoflagellate endosymbionts from population genetic studies. *Molecular Ecology*, *23*(17), 4203–4215. doi: 10.1111/mec.12788

Becker, A., Barnett, M. J., Capela, D., Dondrup, M., Kamp, P. B., Krol, E., … Goesmann, A. (2009). A portal for rhizobial genomes: RhizoGATE integrates a Sinorhizobium meliloti genome annotation update with postgenome data. *Journal of Biotechnology*, *140*(1–2), 45–50. doi: 10.1016/j.jbiotec.2008.11.006

Berg, G. (2009). Plant-microbe interactions promoting plant growth and health: Perspectives for controlled use of microorganisms in agriculture. *Applied Microbiology and Biotechnology*, *84*(1), 11–18. doi: 10.1007/s00253-009-2092-7

Biondi, E. G., Pilli, E., Giuntini, E., Roumiantseva, M. L., Andronov, E. E., Onichtchouk, O. P., … Bazzicalupo, M. (2003). Genetic relationship of Sinorhizobium meliloti and Sinorhizobium medicae strains isolated from Caucasian region. *FEMS Microbiology Letters*, *220*(2), 207–213. doi: 10.1016/S0378-1097(03)00098-3

Blanca-Ordóñez, H., Oliva-García, J. J., Pérez-Mendoza, D., Soto, M. J., Olivares, J., Sanjuán, J., & Nogales, J. (2010). pSymA-dependent mobilization of the Sinorhizobium meliloti pSymB megaplasmid. *Journal of Bacteriology*, *192*(23), 6309–6312. doi: 10.1128/JB.00549-10

Bonhomme, M., Boitard, S., Clemente, H. S., Dumas, B., Young, N., & Jacquet, C. (2015). Genomic signature of selective sweeps illuminates adaptation of medicago truncatula to

root-associated microorganisms. *Molecular Biology and Evolution*, *32*(8), 2097–2110. doi: 10.1093/molbev/msv092

Bonnin, I., Huguet, T., Gherardi, M., Prosperi, J.-M., & Olivieri, I. (1996). High level of polymorphism and spatial structure in a selfing plant species, Medicago truncatula (Leguminosae), shown using RAPD markers . *American Journal of Botany*, *83*(7), 843–855. doi: 10.1002/j.1537-2197.1996.tb12776.x

Branca, A., Paape, T. D., Zhou, P., Briskine, R., Farmer, A. D., Mudge, J., … Tiffin, P. (2011). Whole-genome nucleotide diversity, recombination, and linkage disequilibrium in the model legume Medicago truncatula. *Proceedings of the National Academy of Sciences of the United States of America*, *108*(42). doi: 10.1073/pnas.1104032108

Brown, S. P., Grillo, M. A., Podowski, J. C., & Heath, K. D. (2020). Soil origin and plant genotype structure distinct microbiome compartments in the model legume Medicago truncatula. *Microbiome*, *8*(1), 1–17. doi: 10.1186/s40168-020-00915-9

Burghardt, L. T. (2020). Evolving together, evolving apart: measuring the fitness of rhizobial bacteria in and out of symbiosis with leguminous plants. *New Phytologist*, *228*(1), 28–34. doi: 10.1111/nph.16045

Burghardt, L. T., Epstein, B., Guhlin, J., Nelson, M. S., Taylor, M. R., Young, N. D., … Tiffin, P. (2018). Select and resequence reveals relative fitness of bacteria in symbiotic and free-living environments. *Proceedings of the National Academy of Sciences of the United States of America*, *115*(10), 2425–2430. doi: 10.1073/pnas.1714246115

Burghardt, L. T., Epstein, B., & Tiffin, P. (2019). Legacy of prior host and soil selection on rhizobial fitness in planta. *Evolution*, *73*(9), 2013–2023. doi: 10.1111/evo.13807

Caldera, E. J., & Currie, C. R. (2012). The population structure of antibiotic-producing bacterial symbionts of apterostigma dentigerum ants: Impacts of coevolution and multipartite symbiosis. *American Naturalist*, *180*(5), 604–617. doi: 10.1086/667886

Carlsson-Granér, U., & Thrall, P. H. (2015). Host resistance and pathogen infectivity in host populations with varying connectivity. *Evolution*, *69*(4), 926–938. doi: 10.1111/evo.12631

Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). StackCatchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). Stacks: an analysis tool set for population genomics. *Molecular Ecology*, *22*(11), 3124–3140. doi: 10.1111/mec.12354.Stacks

Cavassim, M. I. A., Moeskjær, S., Moslemi, C., Fields, B., Bachmann, A., Vilhjálmsson, B. J., … Andersen, S. U. (2020). Symbiosis genes show a unique pattern of introgression and selection within a rhizobium leguminosarum species complex. *Microbial Genomics*, *6*(4). doi: 10.1099/mgen.0.000351

Chase, A., Arevalo, P., Brodie, E., Polz, M., Karaoz, U., & Martiny, J. B. H. (2019). Sympatric and allopatric differentiation delineates population structure in free-living terrestrial bacteria. *MBio*. doi: 10.1101/644468

Choudoir, M. J., Barberán, A., Menninger, H. L., Dunn, R. R., & Fierer, N. (2018). Variation in range size and dispersal capabilities of microbial taxa. *Ecology*, *99*(2), 322–334. doi: 10.1002/ecy.2094

Cooper, V. S., Vohr, S. H., Wrocklage, S. C., & Hatcher, P. J. (2010). Why genes evolve faster on secondary chromosomes in bacteria. *PLoS Computational Biology*, *6*(4). doi: 10.1371/journal.pcbi.1000732

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., … Durbin, R. (2011). The variant call format and VCFtools. *Bioinformatics*, *27*(15), 2156–2158. doi:

10.1093/bioinformatics/btr330

Degnan, J. H., & Rosenberg, N. A. (2009). Gene tree discordance, phylogenetic inference and
the multispecies coalescent. *Trends in Ecology and Evolution*, *24*(6), 332–340. doi:
10.1016/j.tree.2009.01.009

diCenzo, G. C., & Finan, T. M. (2017). The Divided Bacterial Genome. *Microbiology and*
Molecular Biology Reviews, *81*(3), 1–37.

diCenzo, G. C., MacLean, A. M., Milunovic, B., Golding, G. B., & Finan, T. M. (2014).
Examination of Prokaryotic Multipartite Genome Evolution through Experimental Genome
Reduction. *PLoS Genetics*, *10*(10). doi: 10.1371/journal.pgen.1004742

Ding, H., & Hynes, M. F. (2009). Plasmid transfer systems in the rhizobia. *Canadian Journal of*
Microbiology, *55*(8), 917–927. doi: 10.1139/W09-056

Dray, S., & Dufour, A. B. (2007). The ade4 package: Implementing the duality diagram for
ecologists. *Journal of Statistical Software*, *22*(4), 1–20. doi: 10.18637/jss.v022.i04

Dybdahl, M. F., & Lively, C. M. (1996). The geography of coevolution: Comparative population
structures for a snail and its trematode parasite. *Evolution*, *50*(6), 2264–2275. doi:
10.1111/j.1558-5646.1996.tb03615.x

Epstein, B., Branca, A., Mudge, J., Bharti, A. K., Briskine, R., Farmer, A. D., … Tiffin, P.
(2012). Population Genomics of the Facultatively Mutualistic Bacteria Sinorhizobium
meliloti and S. medicae. *PLoS Genetics*, *8*(8), 1–10. doi: 10.1371/journal.pgen.1002868

Epstein, B., Sadowsky, M. J., & Tiffin, P. (2014). Selection on Horizontally Transferred and
Duplicated Genes in Sinorhizobium (Ensifer), the Root-Nodule Symbionts of Medicago.
Genome Biology and Evolution, *6*(5), 1199–1209. doi: 10.1093/gbe/evu090

Epstein, B., & Tiffin, P. (2021). Comparative genomics reveals high rates of horizontal transfer
and strong purifying selection on rhizobial symbiosis genes. *Proceedings of the Royal*
Society B: Biological Sciences, *288*(1942). doi: 10.1098/rspb.2020.1804

Fernandes, L. D., Lemos-Costa, P., Guimarães, P. R., Thompson, J. N., & de Aguiar, M. A. M.
(2019). Coevolution Creates Complex Mosaics across Large Landscapes. *The American*
Naturalist, *194*(2), 000–000. doi: 10.1086/704157

Frederickson, M. E. (2017). Mutualisms Are Not on the Verge of Breakdown. *Trends in Ecology*
and Evolution, *32*(10), 727–734. doi: 10.1016/j.tree.2017.07.001

Friesen, M. L., & Mathias, A. (2010). Mixed infections may promote diversification of
mutualistic symbionts: Why are there ineffective rhizobia? *Journal of Evolutionary*
Biology, *23*(2), 323–334. doi: 10.1111/j.1420-9101.2009.01902.x

Friesen, M.L. (2012). Widespread fitness alignment in the legume-rhizobium symbiosis. *New*
Phytologist, *194*(4), 1096–1111. doi: 10.1111/j.1469-8137.2012.04099.x

Galardini, M., Pini, F., Bazzicalupo, M., Biondi, E. G., & Mengoni, A. (2013). Replicon-
dependent bacterial genome evolution: The case of Sinorhizobium meliloti. *Genome*
Biology and Evolution, *5*(3), 542–558. doi: 10.1093/gbe/evt027

Galibert, F., Finan, T. M., Long, S. R., Pühler, A., Abola, P., Ampe, F., … Batut, J. (2001). The
composite genome of the legume symbiont Sinorhizobium meliloti. *Science*, *293*(5530),
668–672. doi: 10.1126/science.1060966

Gandon, S., Capowiez, Y., Dubois, Y., Michalakis, Y., & Olivieri, I. (1996a). Local adaptation
and gene-for-gene coevolution in a metapopulation model. *Proceedings of the Royal*
Society B: Biological Sciences, *263*(1373), 1003–1009. Retrieved from
https://royalsocietypublishing.org/doi/10.1098/rspb.1996.0148

Gandon, S., Capowiez, Y., Dubois, Y., Michalakis, Y., & Olivieri, I. (1996b). Local adaptation

595 and gene-for-gene coevolution in a metapopulation model. *Proceedings of the Royal*
596 *Society B: Biological Sciences*, *263*(1373), 1003–1009. doi:
597 https://doi.org/10.1098/rspb.1996.0148

598 Gano-Cohen, K. A., Wendlandt, C. E., Moussawi, K. Al, Stokes, P. J., Quides, K. W., Weisberg,
599 A. J., … Sachs, J. L. (2020). Recurrent mutualism breakdown events in a legume rhizobia
600 metapopulation. *Proceedings of the Royal Society B: Biological Sciences*, *287*(1919). doi:
601 10.1098/rspb.2019.2549

602 Garrison, E., & Marth, G. (2012). *Haplotype-based variant detection from short-read*
603 *sequencing*. *arXiv* preprint, http://arxiv.org/abs/1207.3907

604 Green, J. L., Bohannan, B. J. M., & Whitaker, R. J. (2008). Microbial biogeography: From
605 taxonomy to traits. *Science*, *320*(5879), 1039–1043. doi: 10.1126/science.1153475

606 Grillo, M. A., De Mita, S., Burke, P. V., Solórzano-Lowell, K. L. S., & Heath, K. D. (2016).
607 Intrapopulation genomics in a model mutualist: Population structure and candidate
608 symbiosis genes under selection in *Medicago truncatula*. *Evolution*, *70*(12), 2704–2717.
609 doi: 10.1111/evo.13095

610 Hanson, C. A., Fuhrman, J. A., Horner-Devine, M. C., & Martiny, J. B. H. (2012). Beyond
611 biogeographic patterns: Processes shaping the microbial landscape. *Nature Reviews*
612 *Microbiology*, *10*(7), 497–506. doi: 10.1038/nrmicro2795

613 Harrison, P. W., Lower, R. P. J., Kim, N. K. D., & Young, J. P. W. (2010). Introducing the
614 bacterial 'chromid': not a chromosome, not a plasmid. *Trends in Microbiology*, *18*(4), 141–
615 148. doi: 10.1016/J.TIM.2009.12.010

616 Harrison, T. L., Wood, C. W., Heath, K. D., & Stinchcombe, J. R. (2017a). Geographically
617 structured genetic variation in the Medicago lupulina–Ensifer mutualism. *Evolution*, *71*(7),
618 1787–1801. doi: 10.1111/evo.13268

619 Harrison, T. L., Wood, C. W., Heath, K. D., & Stinchcombe, J. R. (2017b). Geographically
620 structured genetic variation in the *Medicago lupulina - Ensifer* mutualism. *Evolution*,
621 *71*(7), 1787–1801. doi: 10.1111/evo.13268

622 Heath, K. D. (2010). Intergenomic epistasis and coevolutionary constraint in plants and rhizobia.
623 *Evolution*, *64*(5), 1446–1458. doi: 10.1111/j.1558-5646.2009.00913.x

624 Heath, K. D., & Grillo, M. A. (2016). Rhizobia: tractable models for bacterial evolutionary
625 ecology. *Environmental Microbiology*, *18*(12), 4307–4311. doi: 10.1111/1462-2920.13492

626 Heath, K. D., & Stinchcombe, J. R. (2014a). Explaining mutualism variation: A new
627 evolutionary paradox? *Evolution*, *68*(2), 309–317. doi: 10.1111/evo.12292

628 Heath, K. D., & Tiffin, P. (2009). Stabilizing mechanisms in a legume-rhizobium mutualism.
629 *Evolution*, *63*(3), 652–662. doi: 10.1111/j.1558-5646.2008.00582.x

630 Hoetzinger, M., Pitt, A., Huemer, A., & Hahn, M. W. (2021). Continental-Scale Gene Flow
631 Prevents Allopatric Divergence of Pelagic Freshwater Bacteria. *Genome Biology and*
632 *Evolution*, *13*(3), 1–18. doi: 10.1093/gbe/evab019

633 Hollowell, A. C., Regus, J. U., Gano, K. A., Bantay, R., Centeno, D., Pham, J., … Sachs, J. L.
634 (2016). Epidemic Spread of Symbiotic and Non-Symbiotic Bradyrhizobium Genotypes
635 Across California. *Microbial Ecology*, *71*(3), 700–710. doi: 10.1007/s00248-015-0685-5

636 Hollowell, Amanda C., Regus, J. U., Turissini, D., Gano-Cohen, K. A., Bantay, R., Bernardo, A.,
637 … Sachs, J. L. (2016). Metapopulation dominance and genomicisland acquisition of
638 Bradyrhizobium with superior catabolic capabilities. *Proceedings of the Royal Society B:*
639 *Biological Sciences*, *283*(1829). doi: 10.1098/rspb.2016.0496

640 Jombart, T., & Ahmed, I. (2011). adegenet 1.3-1: New tools for the analysis of genome-wide

SNP data. *Bioinformatics*, *27*(21), 3070–3071. doi: 10.1093/bioinformatics/btr521

Jones, E. I., Afkhami, M. E., Akçay, E., Bronstein, J. L., Bshary, R., Frederickson, M. E., …
Friesen, M. L. (2015). Cheaters must prosper: Reconciling theoretical and empirical
perspectives on cheating in mutualism. *Ecology Letters*, *18*(11), 1270–1284. doi:
10.1111/ele.12507

Kamvar, Z. N., Tabima, J. F., & Grünwald, N. J. (2014). Poppr: An R package for genetic
analysis of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ*,
2014(1), 1–14. doi: 10.7717/peerj.281

Kaneko, T., Nakamura, Y., Sato, S., Minamisawa, K., Uchiumi, T., Sasamoto, S., … Tabata, S.
(2002). Complete genomic sequence of nitrogen-fixing symbiotic bacterium
Bradyrhizobium japonicum USDA110. *DNA Research*, *9*(6), 189–197. doi:
10.1093/dnares/9.6.189

Klinger, C. R., Lau, J. A., & Heath, K. D. (2016). Ecological genomics of mutualism decline in
nitrogen-fixing bacteria. *Proceedings of the Royal Society B: Biological Sciences*,
283(1826), 20152563. doi: 10.1098/rspb.2015.2563

Koppell, J. H., & Parker, M. A. (2012). Phylogenetic clustering of Bradyrhizobium symbionts on
legumes indigenous to North America. *Microbiology (United Kingdom)*, *158*(8), 2050–
2059. doi: 10.1099/mic.0.059238-0

Kumar, N., Lad, G., Giuntini, E., Kaye, M. E., Udomwong, P., Shamsani, N. J., … Bailly, X.
(2015). Bacterial genospecies that are not ecologically coherent: population genomics of
Rhizobium leguminosarum. *Open Biology*, *5*(1), 140133–140133. doi:
10.1098/rsob.140133

Laine, A. L. (2005). Spatial scale of local adaptation in a plant-pathogen metapopulation.
Journal of Evolutionary Biology, *18*(4), 930–938. doi: 10.1111/j.1420-9101.2005.00933.x

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler
transform. *Bioinformatics*, *25*(14), 1754–1760. doi: 10.1093/bioinformatics/btp324

Locey, K. J., & Lennon, J. T. (2016). Scaling laws predict global microbial diversity.
Proceedings of the National Academy of Sciences of the United States of America, *113*(21),
5970–5975. doi: 10.1073/pnas.1521291113

Martiny, J. B. H., Bohannan, B. J. M., Brown, J. H., Colwell, R. K., Fuhrman, J. A., Green, J. L.,
… Staley, J. T. (2006). Microbial biogeography: Putting microorganisms on the map.
Nature Reviews Microbiology, *4*(2), 102–112. doi: 10.1038/nrmicro1341

Masatoshi, N. (1972). Genetic Distance between Populations. *American Naturalist*, *106*(949),
283–292.

Nelson, M., Guhlin, J., Epstein, B., Tiffin, P., & Sadowsky, M. J. (2018a). The complete
replicons of 16 Ensifer meliloti strains offer insights into intra- and inter-replicon gene
transfer, transposon-associated loci, and repeat elements. *Microbial Genomics*, *4*(5). doi:
10.1099/mgen.0.000174

Nelson, M., Guhlin, J., Epstein, B., Tiffin, P., & Sadowsky, M. J. (2018b). The complete
replicons of 16 Ensifer meliloti strains offer insights into intra- and inter-replicon gene
transfer, transposon-associated loci, and repeat elements. *Microbial Genomics*, *4*(5), 1–11.
doi: 10.1099/mgen.0.000174

Nuismer, S. L., Thompson, J. N., & Gomulkiewicz, R. (2003). Coevolution between hosts and
parasites with partially overlapping geographic ranges. *Journal of Evolutionary Biology*,
16(6), 1337–1345. doi: 10.1046/j.1420-9101.2003.00609.x

Ossler, J. N., & Heath, K. D. (2018). Shared genes but not shared genetic variation: Legume

colonization by two belowground symbionts. *American Naturalist*, *191*(3), 395–406. doi:
10.1086/695829

Page, A. J., Cummins, C. A., Hunt, M., Wong, V. K., Reuter, S., Holden, M. T. G., … Parkhill,
J. (2015). Roary: Rapid large-scale prokaryote pan genome analysis. *Bioinformatics*,
31(22), 3691–3693. doi: 10.1093/bioinformatics/btv421

Paradis, E., & Schliep, K. (2019). Ape 5.0: An environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics*, *35*(3), 526–528. doi:
10.1093/bioinformatics/bty633

Parker, M. A. (1999). Mutualism in metapopulations of legumes and rhizobia. *American
Naturalist*, *153*(SUPPL.), 48–60. doi: 10.1086/303211

Pembleton, L. W., Cogan, N. O. I., & Forster, J. W. (2013). StAMPP: An R package for
calculation of genetic differentiation and structure of mixed-ploidy level populations.
Molecular Ecology Resources, *13*(5), 946–952. doi: 10.1111/1755-0998.12129

Pérez-Mendoza, D., Sepúlveda, E., Pando, V., Muñoz, S., Nogales, J., Olivares, J., … Sanjuán, J.
(2005). Identification of the rctA gene, which is required for repression of conjugative
transfer of rhizobial symbiotic megaplasmids. *Journal of Bacteriology*, *187*(21), 7341–
7350. doi: 10.1128/JB.187.21.7341-7350.2005

Pérez Carrascal, O. M., VanInsberghe, D., Juárez, S., Polz, M. F., Vinuesa, P., & González, V.
(2016). Population genomics of the symbiotic plasmids of sympatric nitrogen-fixing
Rhizobium species associated with *Phaseolus vulgaris*. *Environmental Microbiology*, *18*(8),
2660–2676. doi: 10.1111/1462-2920.13415

Pita, L., Rix, L., Slaby, B. M., Franke, A., & Hentschel, U. (2018). The sponge holobiont in a
changing ocean: from microbes to ecosystems. *Microbiome*, *6*(1), 46. doi: 10.1186/s40168-
018-0428-1

Porter, S. S., Faber-Hammond, J., Montoya, A. P., Friesen, M. L., & Sackos, C. (2019). Dynamic
genomic architecture of mutualistic cooperation in a wild population of Mesorhizobium.
ISME Journal, *13*(2), 301–315. doi: 10.1038/s41396-018-0266-y

Porter, S. S., & Simms, E. L. (2014). Selection for cheating across disparate environments in the
legume-rhizobium mutualism. *Ecology Letters*, *17*(9), 1121–1129. doi: 10.1111/ele.12318

Pretorius-Guth, I. M., Puhler, A., & Simon, R. (1990). Conjugal transfer of megaplasmid 2
between Rhizobium meliloti strains in alfalfa nodules. *Applied and Environmental
Microbiology*, *56*(8), 2354–2359. doi: 10.1128/aem.56.8.2354-2359.1990

Rambaut, A. (2018). *FigTree*. Edinburgh. Retrieved from
http://tree.bio.ed.ac.uk/software/figtree/

Revillini, D., Gehring, C. A., & Johnson, N. C. (2016). The role of locally adapted mycorrhizas
and rhizobacteria in plant–soil feedback systems. *Functional Ecology*, *30*(7), 1086–1098.
doi: 10.1111/1365-2435.12668

Reynolds, H. L., Packer, A., Bever, J. D., & Clay, K. (2003). Grassroots ecology: Plant-microbe-
soil interactions as drivers of plant community structure and dynamics. *Ecology*, *84*(9),
2281–2291. doi: 10.1890/02-0298

[dataset] Riley, A.B., Grillo, M.A., Epstein, B., Tiffin, P., & Heath, K.D. (2021). Partners in
space: Discordant population structure between legume hosts and rhizobium symbionts in
their native range. NCBI ###-### and DRYAD ###.

Ronfort, J., Bataillon, T., Santoni, S., Delalande, M., David, J. L., & Prosperi, J. M. (2006).
Microsatellite diversity and broad scale geographic structure in a model legume: Building a
set of nested core collection for studying naturally occurring variation in Medicago

truncatula. *BMC Plant Biology*, *6*, 1–13. doi: 10.1186/1471-2229-6-28

Rudman, S. M., Barbour, M. A., Csilléry, K., Gienapp, P., Guillaume, F., Hairston, N. G., …

Levine, J. M. (2018). What genomic data can reveal about eco-evolutionary dynamics.

Nature Ecology and Evolution, *2*(1), 9–15. doi: 10.1038/s41559-017-0385-2

Sachs, J. L., Quides, K. W., & Wendlandt, C. E. (2018). Legumes versus rhizobia : a model for

ongoing conflict in symbiosis. *New Phytologist, 219,* 1199–1206. doi: 10.1111/nph.15222

Savolainen, O., Lascoux, M., & Merilä, J. (2013). Ecological genomics of local adaptation.

Nature Reviews Genetics, *14*(11), 807–820. doi: 10.1038/nrg3522

Seemann, T. (2014). Prokka: Rapid prokaryotic genome annotation. *Bioinformatics*, *30*(14),

2068–2069. doi: 10.1093/bioinformatics/btu153

Siol, M., Prosperi, J. M., Bonnin, I., & Ronfort, J. (2008). How multilocus genotypic pattern

helps to understand the history of selfing populations: A case study in Medicago truncatula.

Heredity, *100*(5), 517–525. doi: 10.1038/hdy.2008.5

Smith, C. I., Godsoe, W. K. W., Tank, S., Yoder, J. B., & Pellmyr, O. (2008). Distinguishing

coevolution from covicariance in an obligate pollination mutualism: Asynchronous

divergence in Joshua tree and its pollinators. *Evolution*, *62*(10), 2676–2687. doi:

10.1111/j.1558-5646.2008.00500.x

Stoy, K. S., Gibson, A. K., Gerardo, N. M., & Morran, L. T. (2020). A need to consider the

evolutionary genetics of host–symbiont mutualisms. *Journal of Evolutionary Biology*,

33(12), 1656–1668. doi: 10.1111/jeb.13715

Strobel, H. M., Alda, F., Sprehn, C. G., Blum, M. J., & Heins, D. C. (2016). Geographic and

host-mediated population genetic structure in a cestode parasite of the three-spined

stickleback. *Biological Journal of the Linnean Society*, *119*(2), 381–396. doi:

10.1111/bij.12826

Tack, A. J. M., Horns, F., & Laine, A. L. (2014). The impact of spatial scale and habitat

configuration on patterns of trait variation and local adaptation in a wild plant parasite.

Evolution, *68*(1), 176–189. doi: 10.1111/evo.12239

Tang, H., Krishnakumar, V., Bidwell, S., Rosen, B., Chan, A., Zhou, S., … Town, C. D. (2014).

An improved genome release (version Mt4.0) for the model legume Medicago truncatula.

BMC Genomics, *15*(1), 1–14. doi: 10.1186/1471-2164-15-312

Thompson, A. R., Thacker, C. E., & Shaw, E. Y. (2005). Phylogeography of marine mutualists:

Parallel patterns of genetic structure between obligate goby and shrimp partners. *Molecular*

Ecology, *14*(11), 3557–3572. doi: 10.1111/j.1365-294X.2005.02686.x

Thompson, J. N. (1994). *The Coevolutionary Process* (1st ed.). Chicago, IL, USA: The

University Of Chicago Press Books.

Thompson, J. N. (2005). *The Geographic Mosaic of Coevolution* (1st ed.). Chicago Press.

Thrall, P. H., Hochberg, M. E., Burdon, J. J., & Bever, J. D. (2007). Coevolution of symbiotic

mutualists and parasites in a community context. *Trends in Ecology and Evolution*, *22*(3),

120–126. doi: 10.1016/j.tree.2006.11.007

VanInsberghe, D., Arevalo, P., Chien, D., & Polz, M. F. (2020). How can microbial population

genomics inform community ecology? *Philosophical Transactions of the Royal Society B:*

Biological Sciences, *375*(1798). doi: 10.1098/rstb.2019.0253

Vincent, J. (1970). *A Manual for the Practical Study of Root-Nodule Bacteria*. Oxford and

Edinburgh: Blackwell Scientific.

Vos, M., & Velicer, G. J. (2008). Isolation by Distance in the Spore-Forming Soil Bacterium

Myxococcus xanthus. *Current Biology*, *18*(5), 386–391. doi: 10.1016/J.CUB.2008.02.050

Whitaker, R. J., & Banfield, J. F. (2006). Population genomics in natural microbial communities.
Trends in Ecology and Evolution, *21*(9), 508–516. doi: 10.1016/j.tree.2006.07.001
Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. New York: Springer-Verlag.
Retrieved from https://ggplot2.tidyverse.org
Wood, C. W., Pilkington, B. L., Vaidya, P., Biel, C., & Stinchcombe, J. R. (2018). Genetic
conflict with a parasitic nematode disrupts the legume-rhizobia mutualism. *Evolution*
Letters, *2*(3), 233–245. doi: 10.1002/evl3.51
Yates, R., Howieson, J., De Meyer, S. E., Tian, R., Seshadri, R., Pati, A., … Reeve, W. (2015).
High-quality permanent draft genome sequence of Rhizobium sullae strain WSM1592; a
Hedysarum coronarium microsymbiont from Sassari, Italy. *Standards in Genomic*
Sciences, *10*(JULY2015), 1–6. doi: 10.1186/s40793-015-0020-2
Yoder, J. B., & Nuismer, S. L. (2010). When does coevolution promote diversification?
American Naturalist, *176*(6), 802–817. doi: 10.1086/657048
Yoder, J. B., Stanton-Geddes, J., Zhou, P., Briskine, R., Young, N. D., & Tiffin, P. (2014).
Genomic signature of adaptation to climate in Medicago truncatula. *Genetics*, *196*(4),
1263–1275. doi: 10.1534/genetics.113.159319
Young, J. P. W., Crossman, L. C., Johnston, A. W. B., Thomson, N. R., Ghazoui, Z. F., Hull, K.
H., … Parkhill, J. (2006). The genome of Rhizobium leguminosarum has recognizable core
and accessory components. *Genome Biology*, *7*(4). doi: 10.1186/gb-2006-7-4-r34
Young, J. P. W., & Haukka, K. E. (1996). Diversity and phylogeny of rhizobia. *New Phytologist*,
133(1), 87–94. doi: 10.1111/j.1469-8137.1996.tb04344.x
Younginger, B. S., & Friesen, M. L. (2019). Connecting signals and benefits through partner
choice in plant-microbe interactions. *FEMS Microbiology Letters*, *366*(18), 1–11. doi:
10.1093/femsle/fnz217
Zribi, K., Mhamdi, R., Huguet, T., & Aouani, M. E. (2004). Distribution and genetic diversity of
rhizobia nodulating natural populations of Medicago truncatula in tunisian soils. *Soil*
Biology and Biochemistry, *36*(6), 903–908. doi: 10.1016/j.soilbio.2004.02.003
806

Data accessibility:

Raw sequence reads and assemblies are archived at NCBI (accessions ###).

Author contributions:

KDH and MAG conceived of and designed the work. KDH, MAG, and PT acquired funding.

MAG collected soil, performed trapping experiments, and isolated strains, and MAG and ABR

extracted and submitted DNA for sequencing. ABR and BE performed bioinformatic and

evolutionary analyses. ABR, KDH, and MAG drafted the article, and all authors participated in

critical revisions and approved the final version for submission.

Table 1. AMOVA partitioning the genome-wide genetic variation for each of the three elements
of the *E. meliloti* genome for 191 strains sampled from 21 populations from three regions (Spain,
France, or Corsica) in the native range of the symbiosis, compared to host plant *M. truncatula*.
For each level of spatial division, the percent variance explained, phi statistic, and significance
are given.
823

	Chromosome		pSymA		pSymB		*M. truncatula*	
	% variance	*Phi*	*% variance*	*Phi*	*% variance*	*Phi*	*% variance*	*Phi*
Among Region	24.5	0.34[**]	7.7	0.27[**]	10.1	0.24[***]	19.6	0.20[***]
Among Population within Region	9.7	0.13[***]	19.2	0.21[***]	13.8	0.15[***]	36.5	0.45[***]
Within Population	65.8	0.25[***]	73.1	0.08[***]	76.1	0.10[***]	44.0	0.56[***]

824 [*]p<0.05; [**]p<0.01; [***]p<0.001; [****]p<0.0001

828 **Figure 1.** Tests of correlated population structure between the genomic elements in *E. meliloti*.
829 Shown are correlations between the matrices of pairwise D_{XY} values. Left column are individual
830 D_{XY} values and right column are population-level D_{XY} values.
831

Figure 2. Neighbor joining trees of rhizobium individuals based on (a) chromosomal and (b) pSymA variant data. For both trees individual tip labels (strain ID followed by soil population) are colored based on region of origin (orange from Corsica, green from mainland France, blue from Spain). The outer ring coloration in both trees represents the five major clusters of the chromosome, allowing for comparison with pSymA. Only pSymA is shown here due to sharing a similar topology to pSymB (see Fig. S3).

(2a)

848   **(2b)**

0.02

Figure 3. Correlations between geographic distance and individual genetic distance (D_{XY}) for *Medicago* host plants (n=192) (green), and the *E. meliloti* (n=191) chromosome (red), pSymA (blue), and pSymB (purple). Shown are trend lines and Pearson r^2 values from Mantel tests for each comparison ([***] p<0.001).

865   **Table S1.** Metadata on strains of *Ensifer meliloti*. For each strain the region, sampling site (soil),

866   and specific soil samples are given, along with the genotype of the *Medicago truncatula* plant

867   used to isolate the strain, and the latitude and longitude at which the soil sample was taken.

STRAIN	REGION	SOIL	SOIL_SAMPLE	PLANT GENOTYPE	LAT	LONG
MAG722A	Corsica	C1	C1-3	F83026	42.16748	9.542717
MAG722B	Corsica	C1	C1-3	F83026	42.16748	9.542717
MAG530	Corsica	C2	C2-5	ESP161	42.9687	9.41295
MAG420	Corsica	C3	C3-8	F20089	42.97463	9.365783
MAG421	Corsica	C3	C3-8	F20089	42.97463	9.365783
MAG422	Corsica	C3	C3-8	F20089	42.97463	9.365783
MAG498	Corsica	C3	C3-8	ESP161	42.97463	9.365783
MAG508	Corsica	C3	C3-11	F20089	42.97263	9.366483
MAG701A	Corsica	C3	C3-8	F83026	42.97463	9.365783
MAG701B	Corsica	C3	C3-8	F83026	42.97463	9.365783
MAG714A	Corsica	C3	C3-11	F20093	42.97263	9.366483
MAG714B	Corsica	C3	C3-11	F20093	42.97263	9.366483
MAG513	Corsica	C4	C4-12	ESP161	42.59243	9.04045
MAG514	Corsica	C4	C4-12	ESP161	42.59243	9.04045
MAG533	Corsica	C4	C4-12	F20093	42.59243	9.04045
MAG710A	Corsica	C4	C4-12	F20093	42.59243	9.04045
MAG710B	Corsica	C4	C4-12	F20093	42.59243	9.04045
MAG724A	Corsica	C4	C4-4	F20043	42.59172	9.04035
MAG724B	Corsica	C4	C4-4	F20043	42.59172	9.04035
MAG726A	Corsica	C4	C4-7	F20043	42.59098	9.040233
MAG726B	Corsica	C4	C4-7	F20043	42.59098	9.040233
MAG523	France	1	1-8	ESP009	43.1474	3.000417
MAG524	France	1	1-8	ESP009	43.1474	3.000417
MAG525	France	1	1-8	ESP009	43.1474	3.000417
MAG526	France	1	1-8	ESP009	43.1474	3.000417
MAG141	France	2	2-1	F11005	43.14543	2.999367
MAG183	France	2	2-1	ESP174	43.14543	2.999367
MAG184	France	2	2-1	ESP174	43.14543	2.999367
MAG199	France	2	2-1	F11005	43.14543	2.999367
MAG220	France	2	2-9	ESP174	43.1457	2.9989
MAG32	France	2	2-1	F11005 n	43.14543	2.999367
MAG539	France	2	2-7	F11005	43.14542	2.998683
MAG540	France	2	2-7	F11005	43.14542	2.998683
MAG541	France	2	2-7	F11005	43.14542	2.998683
MAG80	France	2	2-10	ESP174	43.14572	2.998917
MAG82	France	2	2-1	ESP174	43.14543	2.999367
MAG92	France	2	2-9	F11005	43.1457	2.9989

MAG93	France	2	2-9	F11005	43.1457	2.9989
MAG123	France	3	3-3	F11005	43.12227	3.077067
MAG177	France	3	3-4	F11005	43.12235	3.075667
MAG194-mel	France	3	3-2	ESP174	43.12233	3.076717
MAG225	France	3	3-2	ESP174	43.12233	3.076717
MAG245	France	3	3-1	ESP174	43.12222	3.07635
MAG246	France	3	3-1	ESP174	43.12222	3.07635
MAG247	France	3	3-1	ESP174	43.12222	3.07635
MAG248	France	3	3-1	ESP174	43.12222	3.07635
MAG34	France	3	3-2	ESP174	43.12233	3.076717
MAG40	France	3	3-3	ESP174 no	43.12227	3.077067
MAG69	France	3	3-2	F11005	43.12233	3.076717
MAG90A	France	3	3-7	F11005	43.12232	3.075283
MAG90B	France	3	3-7	F11005	43.12232	3.075283
MAG14	France	4	4-2	ESP174	43.12198	3.095533
MAG16	France	4	4-3	F20089	43.12227	3.096033
MAG204	France	4	4-1	ESP174	43.122	3.095283
MAG205	France	4	4-1	ESP174	43.122	3.095283
MAG206	France	4	4-1	ESP174	43.122	3.095283
MAG24	France	4	4-3	F20089	43.12227	3.096033
MAG27	France	4	4-2	ESP174 n	43.12198	3.095533
MAG39	France	4	4-1	F20089	43.122	3.095283
MAG43	France	4	4-1	F20089	43.122	3.095283
MAG48	France	4	4-3	ESP174	43.12227	3.096033
MAG5	France	4	4-3	F20089	43.12227	3.096033
MAG57	France	4	4-3	ESP174	43.12227	3.096033
MAG59	France	4	4-1	ESP174	43.122	3.095283
MAG62	France	4	4-1	F20089	43.122	3.095283
MAG65	France	4	4-3	ESP174	43.12227	3.096033
MAG67	France	4	4-2	F20089	43.12198	3.095533
MAG6A	France	4	4-3	F20089	43.12227	3.096033
MAG6B	France	4	4-3	F20089	43.12227	3.096033
MAG73	France	4	4-3	ESP174	43.12227	3.096033
MAG7A	France	4	4-2	ESP174	43.12198	3.095533
MAG7B	France	4	4-2	ESP174	43.12198	3.095533
MAG17	France	5	5-3	F20089	42.82302	2.9269
MAG209	France	5	5-1	F66009	42.82278	2.927717
MAG36	France	5	5-3	F66009	42.82302	2.9269
MAG46	France	5	5-3	F66009	42.82302	2.9269
MAG71	France	5	5-2	F20089	42.82302	2.927317
MAG201	France	6	6-2	F66009	42.74198	2.81485
MAG219	France	6	6-2	F20089	42.74198	2.81485
MAG38	France	6	6-2	F20089	42.74198	2.81485

				n		
MAG41	France	6	6-1	F20089	42.74205	2.814917
MAG42	France	6	6-1	F20089	42.74205	2.814917
MAG53	France	6	6-2	F20089	42.74198	2.81485
MAG72	France	6	6-2	F66009	42.74198	2.81485
MAG8	France	6	6-1	F66009	42.74205	2.814917
MAG85	France	6	6-1	F20089	42.74205	2.814917
MAG87	France	6	6-1	F20089	42.74205	2.814917
MAG88	France	6	6-1	F20089	42.74205	2.814917
MAG15	France	7	7-1	F66009	42.58827	2.78495
				n		
MAG18	France	7	7-1	F66009	42.58827	2.78495
				n		
MAG189	France	7	7-3	F20089	42.58895	2.785383
MAG191	France	7	7-2	F66009	42.5883	2.785
MAG215	France	7	7-3	F66009	42.58895	2.785383
MAG216	France	7	7-3	F66009	42.58895	2.785383
MAG26	France	7	7-2	F20089	42.5883	2.785
MAG54	France	7	7-2	F20089	42.5883	2.785
MAG78	France	7	7-4	F20089	42.58948	2.784833
MAG182	France	14	14-2	F11005	43.1612	3.05905
MAG200	France	14	14-A	ESP174	43.1612	3.059983
MAG84	France	14	14-A	F11005	43.1612	3.059983
MAG86	France	14	14-3	F11005	43.16108	3.059383
MAG89	France	14	14-1	ESP174	43.16122	3.059017
MAG117B	France	15	15-1	F83026	43.35138	5.169617
MAG145	France	15	15-1	F83026	43.35138	5.169617
MAG101	France	16	16-A	ESP174	43.37568	5.188133
MAG114	France	16	16-B	F83026	43.37568	5.1871
MAG142	France	16	16-3	ESP174	43.37693	5.188133
MAG144A	France	16	16-A	F83026	43.37568	5.188133
MAG144B	France	16	16-A	F83026	43.37568	5.188133
MAG158A	France	16	16-3	ESP174	43.37693	5.188133
MAG158B	France	16	16-3	ESP174	43.37693	5.188133
MAG169	France	16	16-3	ESP174	43.37693	5.188133
MAG97	France	16	16-B	ESP174	43.37568	5.1871
MAG303	France	17	17-5	ESP161	43.12692	6.1252
MAG372	France	17	17-12	F83026	43.12538	6.12535
MAG373	France	17	17-12	F83026	43.12538	6.12535
MAG733A	France	17	17-2	F20043	43.12693	6.125217
MAG762A	France	17	17-12	ESP174	43.12538	6.12535
MAG276	France	18	18-2	ESP161	43.16532	6.4774
MAG277	France	18	18-2	ESP161	43.16532	6.4774
MAG278	France	18	18-2	ESP161	43.16532	6.4774
MAG281	France	18	18-3	ESP161	43.1653	6.4774
MAG282	France	18	18-3	ESP161	43.1653	6.4774

MAG283	France	18	18-3	ESP161	43.1653	6.4774
MAG313	France	18	18-1	F83026	43.16528	6.477433
MAG317	France	18	18-5	F83026	43.1649	6.477233
MAG318	France	18	18-5	F83026	43.1649	6.477233
MAG319	France	18	18-5	F83026	43.1649	6.477233
MAG320	France	18	18-5	F83026	43.1649	6.477233
MAG335	France	18	18-3	F83026	43.1653	6.4774
MAG336	France	18	18-3	F83026	43.1653	6.4774
MAG382	France	18	18-2	F83026	43.16532	6.4774
MAG406	France	18	18-5	ESP161	43.1649	6.477233
MAG286	France	19	19-2	ESP161	43.16932	6.474433
MAG342	France	19	19-3	ESP161	43.16935	6.47445
MAG384	France	19	19-2	F83026	43.16932	6.474433
MAG386	France	19	19-4	F83026	43.16932	6.474483
MAG409	France	19	19-1	F83026	43.16935	6.474417
MAG258	France	20	20-2	F83026	43.53285	6.567617
MAG259	France	20	20-2	F83026	43.53285	6.567617
MAG261	France	20	20-2	F83026	43.53285	6.567617
MAG322	France	20	20-3	ESP161	43.53283	6.567467
MAG325	France	20	20-3	ESP161	43.53283	6.567467
MAG358	France	20	20-1	F83026	43.53403	6.572617
MAG417	France	20	20-3	F83026	43.53283	6.567467
MAG739A	France	20	20-2	F20093	43.53285	6.567617
MAG740A	France	20	20-1	F20093	43.53403	6.572617
MAG749A	France	20	20-2	F83026	43.53285	6.567617
MAG749B	France	20	20-2	F83026	43.53285	6.567617
MAG749C	France	20	20-2	F83026	43.53285	6.567617
MAG753A	France	20	20-3	F20043	43.53283	6.567467
MAG753B	France	20	20-3	F20043	43.53283	6.567467
MAG761A	France	20	20-4	F20043	43.53287	6.566783
MAG10	Spain	8	8-1	ESP174	38.41355	-1.01468
MAG21	Spain	8	8-1	ESP174	38.41355	-1.01468
MAG221	Spain	8	8-4	ESP174	38.4116	-1.01507
MAG230	Spain	8	8-3	ESP174	38.41155	-1.0165
MAG231	Spain	8	8-3	ESP174	38.41155	-1.0165
MAG238	Spain	8	8-2	ESP174	38.41253	-1.01568
MAG242	Spain	8	8-2	ESP174	38.41253	-1.01568
MAG243A	Spain	8	8-2	ESP174	38.41253	-1.01568
MAG243B	Spain	8	8-2	ESP174	38.41253	-1.01568
MAG33	Spain	8	8-4	ESP174	38.4116	-1.01507
MAG35	Spain	8	8-4	ESP174	38.4116	-1.01507
MAG66	Spain	8	8-3	F20050	38.41155	-1.0165
MAG79	Spain	8	8-3	F20050	38.41155	-1.0165
MAG83	Spain	8	8-3	ESP174	38.41155	-1.0165
MAG9	Spain	8	8-1	ESP174	38.41355	-1.01468

nod

MAG192C	Spain	9	8-3	F20050	36.91188	-3.47665
MAG192D	Spain	9	9-3	F20050	36.91188	-3.47665
MAG25A	Spain	9	9-3	ESP174	36.91188	-3.47665
MAG25B	Spain	9	9-3	ESP174	36.91188	-3.47665
MAG31	Spain	9	9-1	ESP174	36.91083	-3.47712
				no		
MAG746A	Spain	9	9-3	F20093	36.91188	-3.47665
MAG746B	Spain	9	9-3	F20093	36.91188	-3.47665
MAG121	Spain	11	11-3	ESP174	36.93672	-4.3563
MAG146	Spain	11	11-3	F83026	36.93672	-4.3563
MAG148A	Spain	11	11-3	F83026	36.93672	-4.3563
MAG148B	Spain	11	11-3	F83026	36.93672	-4.3563
MAG154	Spain	11	11-1	F83026	36.93678	-4.35608
MAG157	Spain	11	11-1	F83026	36.93678	-4.35608
MAG174	Spain	11	11-1	ESP174	36.93678	-4.35608
MAG176A	Spain	11	11-1	ESP174	36.93678	-4.35608
MAG176B	Spain	11	11-1	ESP174	36.93678	-4.35608
MAG748A	Spain	11	11-1	ESP174	36.93678	-4.35608
MAG748B	Spain	11	11-1	ESP174	36.93678	-4.35608
MAG755A	Spain	11	11-1	F83026	36.93678	-4.35608
MAG755B	Spain	11	11-1	F83026	36.93678	-4.35608
MAG757A	Spain	11	11-3	F20093	36.93672	-4.3563
MAG757C	Spain	11	11-3	F20093	36.93672	-4.3563
MAG758A-mel	Spain	11	11-2	F83026	36.93678	-4.35608

871 **Table S2.** SNP counts for each element of the symbiont genome before any variant filtering, and

872 after filtering for quality and depth using VCFtools.

	Chromosome	pSymA	pSymB
Pre-filtering	414,004	179,505	222,935
Quality filters	34,689	15,162	22,460

877   **Table S3.** Matrices of population pairwise D_{XY} values for each *E. meliloti* genomic element. The

878   populations are arranged from west to east (Spain-mainland France-Corsica). Cells background

879   coloration indicates the level of D_{XY} (red = low values, green = high values).

880   ***See supplementary Excel file*

Supplemental figures

Figure S1. Map of 25 sampling locations in Spain, mainland France, and Corsica. Grey points

with black outlines represent sites where both host and symbiont were sampled (n = 8), grey

points represent sites where only symbionts were sampled (n = 13), and black circles represent

sites where only hosts were sampled (n = 4).

918 **Figure S2.** Principal component axis plots showing genome wide similarity on PCs 1-3 for the

919 *Medicago* host plants (n=192) (**A**) and for symbiont (n=191) chromosome (**B**), pSymA (**C**), and

920 pSymB (**D**). The darkest ellipse and points on each plot represent individuals from Spain, the

921 intermediate color dashed line represent individuals from mainland France, and the lightest color

922 represents individuals from Corsica.

Figure S3. Neighbor joining tree of rhizobium individuals based on pSymB variant data.

Individual tip labels (strain ID followed by soil population) are colored based on region of origin

(orange from Corsica, green from mainland France, blue from Spain). Branch support indicated

in teal (<70) or pink (<50). Outer ring represents the chromosomal cluster (from Fig. 2a) for

comparison.

Figure S4. Principal component axis plot for all individuals of the symbiont (n=191) based on

variable gene content. From top to bottom principal component 1 and 2, 1 and 3, and 2 and 3.

The red ellipse and points on each plot represent individuals from Spain, the green dotted line

and points represent individuals from mainland France, and the blue points and line represent

individuals from Corsica.

948
949

Figure S5. Tests of correlated population structure between the genomic elements in *E. meliloti* and the host plant *M. truncatula*