Evolution of a minimal cell

Moger-Reischer RZ[1], Glass JI[2], Wise KS[2], Sun L[2,3], Bittencourt D[2,4,], Lynch M[5], Lennon JT[1,*]

[1] Department of Biology, Indiana University, Bloomington, IN 47405, USA

[2] J. Craig Venter Institute, La Jolla, CA 92037, USA

[3] Sorrento Therapeutics, Inc., San Diego, CA 92121 USA

[4] Embrapa Genetic Resources and Biotechnology, Brasília, 70770-917, Brazil

[5] Arizona State University, Tempe, AZ 85287, USA

* Corresponding author: lennonj@indiana.edu

ABSTRACT

Possessing only essential genes, a minimal cell can reveal mechanisms and processes that are critical for the persistence and stability of life. Here, we report on how a synthetically constructed minimal cell contends with the forces of evolution compared to a non-minimized cell from which it was derived. Genome streamlining was costly, but 80% of fitness was regained in 2000 generations. Although selection acted upon divergent sets of mutations, the rates of adaptation in the minimal and non-minimal cell were equivalent. The only apparent constraint of minimization involved epistatic interactions that inhibited the evolution of cell size. Together, our findings demonstrate the power of natural selection to rapidly optimize fitness in the simplest autonomous organism, with implications for the evolution of cellular complexity.

INTRODUCTION

The complexity of an organism is reflected in the number of genes it possesses, a quantity that varies by orders of magnitude across the tree of life. While some obligately endosymbiotic bacteria have fewer than 200 protein-coding genes, many plant and animal genomes contain more than 20,000 genes (Lynch 2007). In principle, the simplest organism would have no functional redundancies and possess only the minimum number of genes essential for life. Any mutation in such an organism could lethally disrupt one or more cellular functions, placing constraints on evolution, as reflected in the fact that essential proteins change more slowly than those encoded by dispensable genes (Graur and Li 2000; Hahn and Kern 2005). Furthermore, organisms with streamlined genomes have fewer targets upon which positive selection can act, thus limiting opportunities for adaptation. Understanding how organisms with simplified genomes overcome these evolutionary challenges has important implications for long-standing problems in biology, including the treatment of clinical pathogens, the persistence of host-associated endosymbionts, the refinement of engineered microorganisms, and the origin of life itself (Leprince et al. 2012; Moran and Bennett 2014; Glass et al. 2017).

The cell is the simplest independent functional unit of life. Yet even unicellular model organisms that are touted for their tractability are complex, possessing thousands of genes and proteins, many of which remain uncharacterized even after decades of in-depth interrogation. The quest for the simplest organism has been aided by advances in synthetic biology, which involves the redesign or novel construction of biological parts and modules (Benner and Sismour 2005; Glass et al. 2017). Promoted as a way to address practical challenges for society (Cameron et al. 2014; DeLisi 2019), synthetic biology also provides a platform for developing powerful simplest-

46   case models through streamlining, whereby nonessential chromosomal sequences are removed

from an organism's genome (Leprince et al. 2012; Hutchison et al. 2016; Glass et al. 2017; Rich-

48   ardson et al. 2017; Lachance et al. 2019). Guided by such strategies, a "minimal cell" was con-

structed with a genome containing only the smallest set of genes required for autonomous cellular

50   life (Hutchison et al. 2016; Breuer et al. 2019). While these efforts succeeded in experimentally

identifying the genetic requirements for basic cellular processes such as metabolism and cell divi-

52   sion, it remains unclear how a minimal cell should respond to the forces of evolution, particularly

given the limited raw materials upon which natural selection can operate as well as the uncharac-

54   terized input of new mutations.

56   To gain insight into the dynamics and outcomes of evolution in a minimal cell, we con-

ducted experiments with synthetic strains of *Mycoplasma mycoides* (Hutchison et al. 2016; Breuer

58   et al. 2019). The minimal cell, *M. mycoides* JCVI-syn3B, was derived from a non-minimal strain,

M. mycoides JCVI-syn1.0, by reducing the chromosome from 901 to 493 genes, resulting in the

60   smallest genome of any known free-living organism (Hutchison et al. 2016; Breuer et al. 2019).

With these two strains, we first investigated whether genome minimization, which included the

62   removal of two DNA replication genes and eight DNA repair genes, altered the rate and spectrum

of new mutations in the minimal cell relative the non-minimal organism. Second, with knowledge

64   of the mutational input, we evaluated whether genome minimization altered the rate and mecha-

nisms of evolution in response to natural selection, as measured using whole-genome sequencing,

66   estimates of population fitness, and phenotypic changes in cell size.

 RESULTS AND DISCUSSION

 Highest recorded mutation rate is robust to genome minimization

Through serial bottlenecking under relaxed selection, we conducted mutation-accumulation ex-

74 periments with populations of *Mycoplasma mycoides* (see Methods). The number of mutations per

nucleotide per generation for the non-minimal cell ($3.13 \pm 0.12 \times 10^{-8}$, mean $\pm$ SEM) was indistin-

76 guishable from that of the minimal cell ($3.25 \pm 0.16 \times 10^{-8}$) ($t_{140} = 0.43$, $P = 0.667$, Fig. 1a). These

mutation rates, which are the highest recorded for any cellular organism, are consistent with other

78 reports where organisms with smaller genomes have higher mutation rates (Kuo and Ochman

2009; Sung et al. 2012; Lynch et al. 2016; Long et al. 2018). For example, *Mesoplasma florum*

80 also has an exceptionally high mutation rate ($1.16 \pm 0.07 \times 10^{-8}$) (Sung et al. 2012), which may

reflect the fact that these closely related bacteria both lack mismatch repair, a system for recogniz-

82 ing and fixing errors that are made during DNA replication (Carvalho et al. 2005). Our data are

also consistent with predictions from the drift-barrier hypothesis. This theory posits that mutation

84 rates evolve downward until the selective advantage of another incremental decrease in the muta-

tion rate is small enough to be effectively neutral and outweighed by genetic drift (Sung et al.

86 2012; Lynch et al. 2016). In other words, populations with lower effective population size (N_e)

experience stronger drift, and hence evolve higher mutation rates (Lynch et al. 2016). Notably, the

88 natural history (obligate pathogen) and genomic features (small genome size and low GC content)

of *M. mycoides* and *Me. florum* suggest these mollicute bacteria most likely have low N_e (Moran

90 2003; Kuo and Ochman 2009; Hershberg 2015; Long et al. 2018).

 Small effect of genome minimization on mutational spectrum

Although the mutation rate was unaffected by minimization, the types of mutations that arise in a

population can still influence evolution. Therefore, we compared the spectra of mutations in the minimal and non-minimal *M. mycoides.* Overall, the composition of mutation types (insertions,

deletions, and single-nucleotide mutations) was unaffected by genome minimization ($\chi^2_2 = 4.16$, $P = 0.125$) (Fig. 1b). Both strains exhibited bias toward deletions over insertions (non-minimal:

$\chi^2_1 = 21.3$, $P = 4.0 \times 10^{-6}$; minimal: $\chi^2_1 = 68.4$, $P < 2.2 \times 10^{-16}$), consistent with findings reported for other bacteria (Kuo and Ochman 2009; Hershberg 2015; Sung et al. 2016). However, the com-

position of single-nucleotide mutations (SNMs), which constituted the largest category of muta­tions (88%), differed between the minimal and non-minimal cell (Monte Carlo $\chi^2 = 69.9$, $P = 1.0$

$\times 10^{-4}$). Mutations from a G or C nucleotide to an A or T nucleotide occurred at a rate ~30-fold higher in the non-minimal cell ($\chi^2_1 = 3736$, $P < 2.2 \times 10^{-16}$) and ~100-fold higher in the minimal

cell ($\chi^2_1 = 1444$, $P < 2.2 \times 10^{-16}$) compared to mutations from A or T to G or C (Fig. 1c). As a consequence, *M. mycoides* exhibits more A:T bias than any known organism (Hershberg 2015;

Long et al. 2018).

There are a few explanations for the extreme A:T mutational bias in the minimal cell. First, *M. mycoides* naturally lacks Dut, a dUTPase protein that draws down the intracellular concentra-

tion of dUTP, thereby avoiding its misincorporation into DNA (Tye and Lehman 1977). Because adenine would pair with a misincorporated uracil, the lack of Dut should lead to an increased

frequency of A:T-biased mutations in the spectrum of cells without Dut (Sedwick et al. 1986). However, since both cells lack Dut, this mechanism is insufficient to explain the discrepancy in

A:T bias between the non-minimal and minimal cells, respectively ($\chi^2_1 = 21.8$, $P = 3.08 \times 10^{-6}$). Instead, the difference in A:T bias is more likely due to the removal of *ung* during minimization

(Hutchison et al. 2016). This nonessential DNA repair gene encodes for the uracil-DNA glycosyl-

ase enzyme Ung, which excises ectopic uracils from DNA via hydrolysis. The process results in a

118   free uracil and an abasic site in the DNA strand (Lindahl et al. 1977), which allows the correct

nucleotide to be inserted using base complementarity. Without Ung, the uracil would remain in

120   place and be paired with A during DNA replication, even if the original nucleotide was not T. One

common cause for the presence of uracil in DNA is the deamination of cytosine (Lindahl 1974).

122   Thus, a cell lacking *ung* would be enriched in C-to-T mutations when C is replaced by U via

deamination and U is replaced by T in subsequent DNA replication via base-pairing with A. In the

124   minimal cell, we observed an excess of these C:G-to-T:A mutations relative to the non-minimal

bacterium ($\chi^2_1 = 89.8$, $P = 1.60 \times 10^{-20}$), consistent with this aspect of the mutation spectrum being

126   related to the minimal cell lacking *ung* (Fig. 1c).

Rapid recovery of fitness in the minimal cell

We next considered how populations of the minimal and non-minimal cell would respond to nat-

130   ural selection. With measured mutation rates of $\sim 3 \times 10^{-8}$ per nucleotide per generation and popu-

lation sizes in excess of 10^7 individuals, we estimated that a new mutation would hit every nucle-

132   otide in the genome >250 times during 2000 generations of experimental evolution. Thus, given

their equal mutation rates per site, populations of either cell type should not be limited by the

134   availability of genetic variation to fuel adaptation, suggesting that any differences in the ways the

two strains adapt would be driven by alterations in genome architecture created by minimization.

To study natural selection, we passaged replicate populations of *M. mycoides* for ~2000

138   generations (see Methods), a period during which rapid adaptation is often observed (Vasi et al.

1994; Gifford et al. 2011). At the end of the experiment, we measured the fitness (competitive growth rates) of all ancestors and evolved populations (see Methods). For the ancestral strains, we determined that genome minimization led to a 50% reduction in fitness (Fig. 2). Despite this major initial cost, the minimal cell rapidly regained fitness during the course of evolution. Approximately 80% of the fitness lost to minimization was recovered during 300 days of serial passaging (Fig. 2). This suggests that a streamlined *M. mycoides* genome is not inherently crippled and can perform almost as well as the non-minimized cell following readaptation.

Contrary to our initial predictions, adaptation was not constrained by genome minimization (Fig. 2). The rate of fitness gain was the same for replicate populations of the non-minimal and minimal cells ($t_6 = 0.40$, $P = 0.704$, Fig. 2). This interpretation was bolstered by results from population-genomic sequencing (see Methods). The relative ratio of nonsynonymous to synonymous fixed SNMs (d_N/d_S) did not differ between the two cell types ($t_6 = 1.18$, $P = 0.859$, Fig. S1), consistent with the interpretation that the rates of molecular evolution were comparable even though all of the genes in the minimal cell are critical for fitness (Graur and Li 2000; Hahn and Kern 2005).

Divergent mechanisms of adaptation

Using a combination of statistical simulation and reverse genetics, we identified mutations that likely contributed to the observed patterns of adaptation. First, we analyzed the gene-by-population matrix for non-synonymous mutations that arose in essential genes during the natural selection experiment (see Methods). The two cell types acquired mutations in different sets of essential genes (PERMANOVA, $F_7 = 4.12$, $P = 0.029$, Fig. 3), suggesting that, despite adapting at similar

162 rates, the populations evolved via divergent routes. To explore this hypothesis, we looked for genes

that acquired a higher number of non-synonymous mutations than expected under assumptions of

164 neutrality (see Methods). We identified 16 genes in the non-minimal genome and 14 in the minimal

genome that were potential targets of positive selection (Table 1). Second, we used reverse genet-

166 ics to experimentally verify that one of the common types of mutation observed in replicate pop-

ulations of both strains was in fact beneficial (Table S1). Using CRISPR editing, we recreated *ftsZ*

168 C-terminal nonsense mutations by inserting an *ftsZ* E315* mutation into the ancestral genomes of

the minimized and non-minimized strains (see Methods). Competition assays with the constructs

170 revealed that this putatively adaptive mutation conferred a 25% fitness increase in the non-minimal

cell and a 14% fitness increase in the minimal cell (Fig. S2).

Comparative analysis of genes under positive selection provided insight into the functional

174 consequences of adaptation in the minimal cell. We hypothesized that mutations in genes related

to membrane transport would be critical for adaptation because the minimal cell relies on the im-

176 port and export of metabolites and other biomolecules for metabolism (Antczak et al. 2019; Breuer

et al. 2019). However, mutations in membrane transport functions were enriched to a similar de-

178 gree in both cell types over 2000 generations of evolution (Fisher's Exact Test $P = 0.934$). Instead,

we detected a signal of enrichment for mutations in biosynthetic genes for the minimal cell

180 (Fisher's Exact Test $P = 0.090$), including those involved in lipid metabolism. Specifically, *fakA*

and *clsA* (Table 1) are considered essential for synthesizing cardiolipin and other lipids from free

182 fatty acids (Breuer et al. 2019), which are important for the construction of cell membranes and

the regulation of cell division. The gene *lgt* is also critical for membrane construction, encoding

184 the protein that transfers diacylglyceryl moieties to anchor surface lipoproteins in the lipid bilayer

(Breuer et al. 2019). Thus, metabolic innovations involving lipid synthesis and distribution may be more important for the minimal cell than enhanced acquisition of metabolites that are already present in the growth medium.

Genome minimization constrains evolution of cell size

The size of single-celled organisms is variable and often linked to fitness in complex ways (Gounand et al. 2016; Mei et al. 2009; Chien et al. 2012). In resource-rich environments, cell size tends to be positively correlated with growth rate, one of the most important components of fitness (Vasi et al. 1994; Mongold and Lenski 1996; Mei et al. 2009; Chien et al. 2012; Hill et al. 2013). For example, in the first 2000 generations of a classic long-term evolution experiment, cell volume and fitness concomitantly increased by 50% and 30%, respectively (Vasi et al. 1994). While an increase in size can accommodate more macromolecules needed for growth and division, it also decreases a cell's surface-to-volume ratio which increases molecular diffusion times. Given these opposing pressures, we evaluated how cell size changed in response to genome streamlining over the course of evolution. Prior to initiating experiments, we observed that the spherically shaped non-minimal cell was 575 ± 28 nm (mean $\pm$ SEM) and the minimal cell was 422 ± 23 nm (see Methods), comparable to measures of cell diameter previously reported (Hutchison et al. 2016; Breuer et al. 2019). Following 2000 generation of evolution, the non-minimal cell exhibited a 73% increase in diameter (993 ± 53 nm, $t_3 = 7.84$, $P = 0.004$) and a corresponding five-fold increase in volume compared to its ancestor (Fig. 4; Table S2). In contrast, the size of the minimal cell did not change during evolution ($t_3 = 0.85$, $P = 0.459$).

While cell size is a complex multigenic trait, previous studies have implicated *ftsZ* with

208   changes in morphology of the minimal cell (Pelletier et al. 2021). It encodes for FtsZ, a protein

that localizes to the midcell and determines the site of membrane constriction during cell division.

210   Prevalent among diverse lineages of bacteria and archaea (McQuillen and Xiao 2020; Liao et al.

2021), *ftsZ* is nevertheless nonessential in *M. mycoides*. However, cells lacking *ftsZ* exhibit aber-

212   rant cell division and morphology (Hutchison et al. 2016; Breuer et al. 2019; Pelletier et al. 2021).

Thus, along with 18 other nonessential genes, *ftsZ* was retained in JCVI-syn3B to aid in culture

214   maintenance and stable growth (Breuer et al. 2019; Pelletier et al. 2021). In our study, *ftsZ* was

consistently mutated over 2000 generations of evolution and was identified as a target of positive

216   selection in both the minimal and non-minimal cell (Table 1, Fig. S2). Introduction of this termi-

nation codon could eliminate the C-terminal region of FtsZ, known to interact with membrane-

218   associated products that recruit FtsZ (Cohan et al. 2020), The early stop codon could also create a

transcriptional polar effect (Graffeuil et al. 2020) that reduces expression of two adjacent down-

220   stream genes within a likely polycistronic operon: MMSYN1_0521, an ortholog of cell division

protein *sepF* and MMSYN1_0520, encoding a hydrolase of unknown functions. Irrespective of

222   mechanism, we demonstrated that mutations in *ftsZ* had a non-additive effect that contributed to

the evolutionary divergence of cell size. Using scanning electron microscopy, we documented that

224   the *ftsZ* E315* nonsense mutation in the ancestral non-minimal cell led to a 25% increase in cell

diameter (t_{118} = 5.56, P = 8.71 × 10^{-8}) and a corresponding two-fold increase in cell volume. In

226   contrast, the same *ftsZ* nonsense mutation in the ancestral minimal cell led to a 19% decrease in

the cell diameter (t_{122} = 3.29, P = 0.001, Fig. S4), which reduced cell volume by half. Thus, the

228   *ftsZ* E315* mutation recapitulated nearly 60% of the evolved divergence in cell size, indicating

that FtsZ plays a central role in cell size in *M. mycoides*.

Although changes in *ftsZ* had opposing effects on the size of the minimal and non-minimal

232   cell, mutations in this gene were adaptive for both strains (Fig. S2, Table 1). Smaller cell size may

be advantageous for the minimal cell. Owing to its slower growth rate, JCVI-syn3B would have

234   spent more time in an exponential growth phase in our evolution experiment. Some studies have

shown that smaller cell size is selected for when bacteria evolve under such conditions (Gounand

236   et al. 2016) even though other studies report a positive association between metabolic rate and

body mass across species (DeLong et al. 2010). Meanwhile, the faster-growing non-minimal cell

238   should experience bouts of feast-or-famine over thousands of generations in a serial batch envi-

ronment. Under such conditions, repeated transitions between exponential and stationary growth

240   phases has been shown to select for an increase in cell size (Vasi et al. 1994; Gounand et al. 2016).

The observed patterns may also reflect constraints imposed by genome streamlining on the ability

242   of the minimal cell to evolve an adaptive increase in cell size (Mei et al. 2009; DeLong et al. 2010;

Chien et al. 2012; Westfall and Levin 2017). With more than 50% of its membrane transport pro-

244   teins removed, the minimal cell may be unable to sequester the resources needed for constructing

and maintaining a larger cell (Chien et al. 2012; Hutchison et al. 2016; Westfall and Levin 2017).

246   However, one might expect that these biosynthetic constraints would be reflected in a reduced rate

of adaptation, which was not the case (Fig. 2). In addition, cell size could evolve as a fitness-

248   neutral byproduct of selection on other traits, such as DNA replication rate (Amir 2017). For ex-

ample, the two strains could have evolved different size trajectories despite similar selection pres-

250   sures, due to epistatic effects of genome minimization such as those demonstrated using the *ftsZ*

E315* mutants (Table S3, Fig. S4). In any case, our findings highlight that cell size, a fundamental

252   feature of biological complexity in multicellular and single-celled organisms alike, evolves in a

way that is highly dependent on genomic context.

Outlook

256   We documented genes, proteins, and traits that are critical for evolutionary performance in the

synthetically constructed *M. mycoides*, a bacterium with the smallest genome of any self-replicat-

258   ing form of life. In its ancestral state, this working approximation of a minimal cell had low fitness,

fewer than 500 protein-coding genes, and the highest mutation rate ever recorded in a cellular

260   organism. A single gene disruption could have proven lethal. Instead, the force of natural selection

outweighed any deleterious consequences of mutation and drift that could have led populations to

262   extinction. Yet, even in one of the simplest organisms, evolution was still complex. Rapid adapta-

tion involved epistatic interactions and selection on divergent targets, 25% of which encoded for

264   proteins of unknown function. Evolution studies like the ones described here can improve gene

characterization and the mapping of regulatory networks, which will inform subsequent rounds of

266   rational engineering and streamlining. In addition, minimal cells can be leveraged to address long-

standing problems in biology. Derived from suites of diverse ancestors, they may serve as new

268   models for investigating innovations of early life (Kacar et al. 2017), while providing bottom-up

insight into the development of multicellularity (Márquez-Zacarías et al. 2021), the mechanisms

270   of coevolution (Hillesland and Stahl 2010), and the assembly of multi-species consortia and their

emergent functions (Goldford et al. 2018).

METHODS

274   **Strains and growth conditions**

We maintained synthetic *Mycoplasma mycoides* JCVI-syn1.0 and synthetic *M. mycoides* JCVI-

276   syn3B in SP4 medium with KnockOut™ Serum Replacement (Gibco) substituted for fetal bovine

serum (Table S4). Cultures were grown in a dark, static growth chamber at 37 °C. The non-minimal JCVI-syn1.0 strain has been described in detail elsewhere (2010). The minimal JVCI-syn3B is identical to the strain synthesized in previous studies (2016) with the following exceptions: JVCI-syn3B possesses a second rRNA operon copy, lacks a gene (MMSYN1_0531) coding for an efflux protein, and has 19 genes that were added back into the minimal genome to render the cell easier to use (Breuer et al. 2019; Pelletier et al. 2021) (Table S5). The strain also contains a landing pad system (*Cre* recombinase and *LoxP*) facilitating genetic manipulation. For competition experiments used to quantify relative fitness, we used a JCVI-syn1.0 strain that expresses mCherry, which allowed us to distinguish it in mixed culture from other strains via flow cytometry (see below).

Mutation accumulation experiment

Overview. Mutation accumulation (MA) experiments are designed to reduce the influence of natural selection through repeated bottlenecks of evolving populations (Lynch et al. 2016). When used with microbial populations, this is typically achieved by transferring single colonies, which have undergone single-cell bottlenecks. Prior to initiating MA experiments, we acclimated JCVI-syn1.0 and JCVI-syn3B to laboratory conditions by maintaining populations in SP4 liquid medium. We took a clone of each acclimated strain to begin the MA experiment. We propagated replicate MA lineages (JCVI-syn1.0: 87 replicates; JCVI-syn3B: 57 replicates) of each strain for 20 to 36 weekly transfers.

Number of generations. In order to compare rates of mutation across replicates, we normalized all rates as per-generation values. To calculate the number of generations per transfer in the MA, we

grew cells on SP4 agar for one week and diluted a sample of 7th day colonies into 1 mL of phosphate-buffered saline (pH = 7.4). Cells were fixed with 20 μL of 25% glutaraldehyde and stained

with 2X SYBR Green, and then counted with a NovoCyte flow cytometer (ACEA Biosciences). We used the dilutions to calculate the number of cells in the original colony, from which we in-

ferred the number of generations ($\log_2(N)$, where N is the number of cells in the undiluted colony) that must have occurred to reach a colony of that size (Dillon et al. 2017), assuming each colony

is formed by a single progenitor cell. Because growth rate and other fitness components can decrease during an MA experiment (Behringer and Hall 2016), we also measured the number of cells

per colony during and at the end of the MA, averaging across time points to estimate the total number of generations. We then used the number of generations per transfer to estimate the effec-

tive population size (N_e) via the harmonic mean method (Behringer and Hall 2016). Specifically, N_e was approximated as the harmonic mean of the series (2^0, 2^1, 2^2, …, 2^f), where f is equal to the

number of generations per transfer inferred from the previous step.

Whole-genome sequencing and sequence analysis. We performed DNA extractions from evolved MA lines using a DNeasy UltraClean Microbial Kit (Qiagen) following the manufacturer's in-

structions, with the additional step of adding 50 μL of 50 mg/mL lysozyme to improve cell lysis. Genomic DNA was sequenced using Illumina MiSeq sequencing to a depth of ≥35X coverage.

Library preparation and DNA sequencing were conducted by the Indiana University Bloomington Center for Genomics and Bioinformatics. Whole-genome sequencing reads were quality con-

trolled using cutadapt (Martin 2011) to trim low-quality base pairs and remove residual adapter sequences. We used breseq with default parameters (Barrick et al. 2014; Deatherage and Barrick

2014) to call mutations using the trimmed reads. We only considered fixed mutations for the MA

lines. We checked for mutations that had arisen in experimental ancestor strains prior to evolution.

324 Ancestral mutations were removed from analysis of all evolved lines derived from that strain using

gdtools (Barrick et al. 2014; Deatherage and Barrick 2014). We used the sequence data to check

326 for contamination or cross-contamination in evolved lines.

328 *Statistical analyses.* To compare mutation rate and spectrum between strains, we used two-sample

t-tests for numerical response variables and two-sample χ^2 tests with continuity correction for com-

330 paring proportions. For comparing proportions to theoretical expectations within a strain, we used

one-sample χ^2 tests with continuity correction.

Adaptive evolution

334 *Overview*. We conducted experiments that allowed bacteria to achieve large population sizes to

increase the efficacy of natural selection. This involved serial passaging of cells in liquid cultures

336 with limited bottlenecking at each transfer. For example, in our experiment, the minimum popula-

tion size was 2-4 $\times 10^7$ for both JCVI-syn1.0 and JCVI-syn3B. We passaged replicate 3 mL liquid

338 cultures of each strain (*n* = 4 per strain) in 13 mm glass culture tubes *via* 1% v/v serial transfer

each day for 300 days in a dark, static incubator held at 37 °C. We calculated the number of gen-

340 erations per day as the $\log_2$ of the dilution factor, i.e., $\log_2(101)$, the number of binary fissions

needed to regenerate the original population size after the 1% v/v transfer (Lenski and Travisano

342 1994). Based on this, we estimated that the *M. mycoides* strains were maintained for 1,997 gener-

ations, which based on other experiments, is long enough for the majority of adaptation to occur

344 (Lenski and Travisano 1994; Lang et al. 2011).

346   *Measurement of fitness.* We measured relative fitness by competing ancestral and evolved strains

against a third-party *M. mycoides* JCVI-syn1.0 strain labeled with mCherry (Wiser and Lenski

348   2015). Each strain was grown in liquid medium to log phase, and then the labeled and unlabeled

strains were simultaneously diluted into a mixed culture in fresh medium. The abundances of the

350   labeled and unlabeled strains in the mixed culture were calculated by immediately (t_0) sampling

the culture, fixing the cells with 20 μL of cold 25% glutaraldehyde incubated at 4 °C for 20 min,

352   and staining with 2X SYBR Green. After 24 hr of growth (t_f), the mixed culture was sampled again

in the same manner. We quantified the abundance of each strain using either our Novocyte flow

354   cytometer (ACEA Biosciences) or an LSR II flow cytometer (BD Biosciences) at Indiana Univer-

sity's Flow Cytometry Core Facility. The abundance of the strain of interest was calculated from

356   the observed number of SYBR Green-only positive cells while the abundance of the third-party

strain was calculated from the number of cells positive for both SYBR Green and mCherry. From

358   these cell counts, we calculated relative fitness as the change in the relative abundance of the strain

of interest during the 24 hr period of competitive growth versus the third-party strain. Specifically,

360   the relative fitness versus the mCherry third-party strain W_C is

$$W_C = \frac{\ln(\frac{N_f}{N_0})}{\ln(\frac{N_{Cf}}{N_{C0}})}$$

362   where N_0 represents the initial abundance of the focal strain, N_f the abundance of the focal strain

after 24 hr, and N_{Cf} and N_{C0} are final and initial abundances of the mCherry third-party strain,

364   respectively (Wiser and Lenski 2015). We standardized fitness values to be relative to the original

Mm JCVI-syn1.0 ancestor strain. In other words, we represent the fitness as *W*: $W = \frac{W_C}{W_{JCVI-syn1.0}}$,

366   where $W_{JCVI-syn1.0}$ is the value of W_C for *M. mycoides* JCVI-syn1.0. Axenic control cultures were

always run contemporaneously in order account for a proportion of mCherry-positive cells that

368   failed to detectably express mCherry.

370   *Whole-genome sequencing and sequence analysis*. DNA extraction, sequencing, and bioinformat-
       ics followed the same methods as for the MA with a few exceptions. Specifically, each replicate
372   population was sequenced to a depth of ≥ 100X coverage, and polymorphic mutations were in-
       cluded in our analyses. In addition, we calculated d_N/d_S for evolved lines using the Jukes-Cantor-
374   corrected equation (Yang 2006) as a possible indicator of molecular constraint among the lines.
       We counted the number of synonymous and nonsynonymous sites using the gdtools module of
376   breseq (Deatherage and Barrick 2014). The observed numbers of synonymous and nonsynony-
       mous substitutions were obtained directly from breseq outputs. Synonymous and nonsynonymous
378   polymorphisms were included in the observed count with probability proportionate to their fre-
       quency in mapped reads. We added a pseudocount of 1 synonymous substitution for all calcula-
380   tions (Shpak et al. 2015) because two of the lines had 0 synonymous substitutions.

382   To identify mutations possibly contributing to adaptation, we looked for genes which had
       mutations in two or more lines. Mutations in the same gene, arising and increasing in frequency in
384   independent lineages, suggests that mutation's rise could be driven by positive selection (Johnson
       et al. 2021). To test this hypothesis, we statistically assessed whether multiply mutated genes had
386   acquired more mutations than would be expected by chance under the assumption that the muta-
       tions were neutral (Johnson et al. 2021). To do this, we recorded all of the polymorphic and fixed
388   mutations that were called within genes. Synonymous mutations were excluded. We then simu-
       lated placing these mutations at random across all genes. The probability of any given gene receiv-
390   ing any given mutation was proportional to the gene's length and relativized to the gene's GC

content using the known mutation rates of G:C nucleotides and A:T nucleotides from the mutation accumulation experiment. We repeated this random placement of mutations 100,000 times. In each simulation, we counted the number of mutations received by each gene, with each fixed mutation increasing the count by 1 and each polymorphism counted at a value equal to its frequency. For each multiply-mutated gene from the real adaptation experiment, we measured the proportion of the 100,000 simulations in which the gene received at least as many mutations as were truly observed and called this proportion the P value. To correct for the multiple tests (Non-minimal cell: 32; Minimal: 22), we report a Benjamini-Hochberg corrected value P_{adj} (Benjamini and Hochberg 1995; Benjamini et al. 2009) for which the false discovery rate $\alpha = 0.05$ for the reported significant genes (Table 1).

Generation of ftsZ *E315* mutant cells.* This process required mutating the bacterial genomes while they were yeast centromeric plasmids (YCPs) followed by genome transplantation of the mutated genomes. The YCPs were mutated using rounds of CRISPR/Cas9 and yeast homologous recombination that is a modification of a method we used previously to mutate *M. mycoides* strains (Kannan et al. 2016).

In the first CRISPR/Cas9 step, the molecule to be mutated was cleaved and the donor DNA comprising sequences from the two flanking genes was recombined with the cut JCVI-syn1.0 or JCVI-syn3B YCP, removing parts of genes of the flanking genes and all of the target gene. The donor DNA had 40 basepair overlaps to both genes flanking the target gene and had a 22 bp *Mycoplasma gallisepticum* 161 CRISPR/Cas9 target sequence with a protospacer adjacent motif (PAM) (5'-GTATAAATACATCCAGGAGtgg-3') that had no homology elsewhere in JCVI-

414   syn1.0 or JCVI-syn3B. The *M. gallisepticum* sequence put a new PAM in the genome, employed

in the second round of CRISPR/Cas9.

The second round of CRISPR/CAS9 cut the JCVI-syn1.0 or JCVI-syn3B YCP at the new

418   *M. gallisepticum* PAM. The cut YCP was then re-circularized using a donor DNA containing the

desired point mutation. The mutagenized regions of the YCPs were PCR amplified and the muta-

420   tion was confirmed by Sanger sequencing. Correctly mutagenized JCVI-syn1.0 or JCVI-syn3B

YCPs were then transplanted into *Mycoplasma capricolum* recipient cells as reported previously

422   (Lartigue et al. 2007, 2009; Hutchison et al. 2016). The mutagenized regions of the transplants

were PCR amplified and sequenced to confirm the presence of the desired mutations.

Microscopy and image analysis. To compare changes in cell size of evolved populations, we cen-

426   trifuged stationary-phase cultures and resuspended the pellet in 1 mL of phosphate-buffered saline

(pH = 7.4). The resuspended cells were fixed by adding 20 μL of cold 25% glutaraldehyde and

428   incubating at 4 °C for 20 min. For microscopic observation, fixed cells were concentrated 50X via

centrifugation and resuspension. The centrifugation steps were performed at room temperature for

430   4 min at 2000 *g*.

432   For phase contrast microscopy, we pipetted 5 μL of concentrated culture onto an agarose

pad (1% agarose in phosphate-buffered saline) and observed cells in the Plan Apo 100X Ph 3

434   objective lens of a Nikon Eclipse 80i phase contrast microscope. Image data were analyzed using

MicrobeJ (Ducret et al. 2016). Cell outlines were drawn using default parameters. Area (number

436   of pixels) within the outline was use as a metric of the area of one face of the spherical cells. Pixel

values were converted to μm^2 using the conversion factor 1 pixel length = 0.0645 μm. Cell diameter

438   was inferred from this area using the assumption of spherical geometry.

440   Scanning electron microscopy (SEM) was performed at the Indiana University Blooming-

ton Electron Microscopy Center. Fixed cells in PBS were pelleted and resuspended in 100 mM

442   sodium cacodylate buffer (pH = 7.2) with 2 mM calcium chloride and 2% sucrose. We coated 12

mm diameter glass coverslips with 0.1% poly-L lysine for 5 min, after which coverslips were

444   washed with a few drops of double distilled water. Resuspended cells were added to the coverslip

surface and allowed to adhere. After 5 min the coverslips were washed twice with 100 mM sodium

446   cacodylate buffer (pH = 7.2) with 2 mM calcium chloride and 2% sucrose. Next, 300 μL of 2%

osmium tetroxide in 100 mM sodium cacodylate buffer (pH = 7.2) with 2% sucrose was added to

448   the surface of the coverslips while on ice. After 30 minutes the coverslips were washed with double

distilled water. Coverslips were placed in a CPD coverslip holder (Electron Microscopy Sciences,

450   catalog number 70193-01). Samples were dehydrated in a graded ethanol series (30%, 50%, 70%,

90%, 95%) while on ice. At room temperature, coverslips were rinsed three times with 100% eth-

452   anol. Each dehydration step lasted 2 min. Critical point drying was done using a Tousimis Samdri

790 critical-point dryer. The dried coverslips were placed on aluminum SEM stubs and sputter

454   coated, using a Safematic CCU-010 with SP-010 Sputter Head, with 45 nm of gold/palladium

(80%/20%). We viewed the samples with a FEI Teneo scanning electron microscope at 2.0 kV, 25

456   pA probe current, and 3.0 mm working distance. The T2 detector was used. We analyzed the SEM

image data using ImageJ (Schneider et al. 2012). We used the "straight" and "measure" features

458   combined with image scale metadata to measure the vertical diameters of imaged cells that met

the following criteria: cells must be round; cells must not have apparent holes or punctures; cells

460   must be completely within the field of view; cells must have an unambiguous perimeter; there

must be no suggestion that a cell is currently or has recently undergone binary fission; cells must

462   ≥ 0.1 µm across. Each image was processed counterclockwise starting from east. The samples were

processed in a randomized order.

Statistical analyses. We compared relative fitness, d_N/d_S, and cell size between evolved popula-

466   tions using two-sample *t*-tests, and to ancestral values using one-sample *t*-tests. We also used two-

sample *t*-tests to compare the relative fitness and cell size of *ftsZ* E315* mutants to the respective

468   ancestral values.

470   We compared the composition of genes acquiring mutations among the evolved replicate

lines by first constructing a gene-by-population matrix. Herein, each row represented an evolved

472   line and each column represented a gene that had acquired at least one mutation among all of the

lines. Each cell of the matrix was filled with the sum value of mutations occurring in that gene in

474   that line, where fixed mutations were valued at 1 and polymorphisms were valued equal to the

allele frequency. Only essential genes, shared between JCVI-syn1.0 and JCVI-syn3B, were con-

476   sidered. We used a permutational multivariate analysis of variance (PERMANOVA) on the Bray-

Curtis distances generated from the gene-by-population matrix to test for the significance of cell

478   type (minimal vs. non-minimal) on the composition of mutations using the adonis function in the

R package vegan (Oksanen et al. 2020). For visualization, the Bray-Curtis distances were decom-

480   posed into two dimensions using principal coordinate analysis via the cmdscale function.

ACKNOWLEDGEMENTS

482   We acknowledge financial support from the US National Science Foundation (DEB-1442246 and 1934554 JTL; MCB-1818344 and 1840320 JIG, LS, and KSW), US Army Research Office Grant

484   (W911NF-14-1-0411 JTL and ML), the National Aeronautics and Space Administration (80NSSC20K0618 JTL), the National Institutes of Health (R35-GM122566-01 and 2017-202

486   ML), and the Brazilian Agricultural Research Corporation (DMCB). We also acknowledge B. Stein for assistance with scanning electron microscopy (NIH 1S10OD023501-01); J. French, E.

488   Snider, K. McKenzie, B. Lehmkuhl for assistance in the laboratory; and M. Behringer for discussion regarding sequence analyses. Data and code are available at https://github.com/LennonLab/MinimalCell.

LITERATURE CITED

492   Amir, A. 2017. Is cell size a spandrel? eLife 6:e22186.

Antczak, M., M. Michaelis, and M. N. Wass. 2019. Environmental conditions shape the nature of a minimal

494   bacterial genome. Nat. Commun. 10:3100.

Barrick, J. E., G. Colburn, D. E. Deatherage, C. C. Traverse, M. D. Strand, J. J. Borges, D. B. Knoester, A.

496   Reba, and A. G. Meyer. 2014. Identifying structural variation in haploid microbial genomes from

short-read resequencing data using breseq. BMC Genomics 15:1039.

498   Behringer, M. G., and D. W. Hall. 2016. Genome-wide estimates of mutation rates and spectrum in *Schiz-*

osaccharomyces pombe indicate CpG sites are highly mutagenic despite the absence of DNA meth-

500   ylation. G3 Bethesda Md 6:149–160.

Benjamini, Y., R. Heller, and D. Yekutieli. 2009. Selective inference in complex research. Philos. Trans.

502   R. Soc. Math. Phys. Eng. Sci. 367:4255–4271.

Benjamini, Y., and Y. Hochberg. 1995. Controlling the false discovery rate: a practical and powerful ap-

504   proach to multiple testing. J. R. Stat. Soc. Ser. B Methodol. 57:289–300.

Benner, S. A., and A. M. Sismour. 2005. Synthetic biology. Nat. Rev. Genet. 6:533–543.

506   Breuer, M., T. M. Earnest, C. Merryman, K. S. Wise, L. Sun, M. R. Lynott, C. A. Hutchison, H. O. Smith,

J. D. Lapek, D. J. Gonzalez, V. de Crécy-Lagard, D. Haas, A. D. Hanson, P. Labhsetwar, J. I. Glass,

508   and Z. Luthey-Schulten. 2019. Essential metabolism for a minimal cell. eLife 8:e36842.

Cameron, D. E., C. J. Bashor, and J. J. Collins. 2014. A brief history of synthetic biology. Nat. Rev. Micro-

510   biol. 12:381–390.

Carvalho, F. M., M. M. Fonseca, S. Batistuzzo De Medeiros, K. C. Scortecci, C. A. G. Blaha, and L. F.

512   Agnez-Lima. 2005. DNA repair in reduced genome: the Mycoplasma model. Gene 360:111–119.

Chien, A.-C., N. S. Hill, and P. A. Levin. 2012. Cell size control in bacteria. Curr. Biol. 22:R340–R349.

514   Cohan, M. C., A. M. P. Eddelbuettel, P. A. Levin, and R. V. Pappu. 2020. Dissecting the functional contri-

butions of the intrinsically disordered C-terminal tail of *Bacillus subtilis* FtsZ. J. Mol. Biol.

516   432:3205–3221.

Deatherage, D. E., and J. E. Barrick. 2014. Identification of mutations in laboratory-evolved microbes from next-generation sequencing data using breseq. Methods Mol. Biol. Clifton NJ 1151:165–188.

DeLisi, C. 2019. The role of synthetic biology in climate change mitigation. Biol. Direct 14:14.

DeLong, J. P., J. G. Okie, M. E. Moses, R. M. Sibly, and J. H. Brown. 2010. Shifts in metabolic scaling, production, and efficiency across major evolutionary transitions of life. Proc. Natl. Acad. Sci. 107:12941–12945.

Dillon, M. M., W. Sung, R. Sebra, M. Lynch, and V. S. Cooper. 2017. Genome-wide biases in the rate and molecular spectrum of spontaneous mutations in *Vibrio cholerae* and *Vibrio fischeri*. Mol. Biol. Evol. 34:93–109.

Ducret, A., E. M. Quardokus, and Y. V. Brun. 2016. MicrobeJ, a tool for high throughput bacterial cell detection and quantitative analysis. Nat. Microbiol. 1:16077.

Gibson, D. G., J. I. Glass, C. Lartigue, V. N. Noskov, R.-Y. Chuang, M. A. Algire, G. A. Benders, M. G. Montague, L. Ma, M. M. Moodie, C. Merryman, S. Vashee, R. Krishnakumar, N. Assad-Garcia, C. Andrews-Pfannkoch, E. A. Denisova, L. Young, Z.-Q. Qi, T. H. Segall-Shapiro, C. H. Calvey, P. P. Parmar, C. A. Hutchison, H. O. Smith, and J. C. Venter. 2010. Creation of a bacterial cell controlled by a chemically synthesized genome. Science 329:52–56.

Gifford, D. R., S. E. Schoustra, and R. Kassen. 2011. The length of adaptive walks is insensitive to starting fitness in *Aspergillus nidulans*. Evol. Int. J. Org. Evol. 65:3070–3078.

Glass, J. I., C. Merryman, K. S. Wise, C. A. Hutchison, and H. O. Smith. 2017. Minimal cells—real and imagined. Cold Spring Harb. Perspect. Biol. 9:a023861.

Goldford, J. E., N. Lu, D. Bajić, S. Estrela, M. Tikhonov, A. Sanchez-Gorostiaga, D. Segrè, P. Mehta, and A. Sanchez. 2018. Emergent simplicity in microbial community assembly. Science 361:469–474.

Gounand, I., T. Daufresne, D. Gravel, C. Bouvier, T. Bouvier, M. Combe, C. Gougat-Barbera, F. Poly, C. Torres-Barceló, and N. Mouquet. 2016. Size evolution in microorganisms masks trade-offs predicted by the growth rate hypothesis. Proc. R. Soc. B Biol. Sci. 283:20162272.

Graffeuil, A., B. E. Uhlin, and D. A. Cisneros. 2020. "Polar mutagenesis of bacterial transcriptional units using Cas12a." Microbiology.

Graur, D., and W.-H. Li. 2000. Fundamentals of molecular evolution. 2nd ed. Sinauer Associates, Sunderland, Mass.

Hahn, M. W., and A. D. Kern. 2005. Comparative genomics of centrality and essentiality in three eukaryotic protein-interaction networks. Mol. Biol. Evol. 22:803–806.

Hershberg, R. 2015. Mutation—The engine of evolution: studying mutation and its role in the evolution of bacteria. Cold Spring Harb. Perspect. Biol. 7:a018077.

Hill, N. S., P. J. Buske, Y. Shi, and P. A. Levin. 2013. A moonlighting enzyme links *Escherichia coli* cell size with central metabolism. PLoS Genet. 9:e1003663.

Hillesland, K. L., and D. A. Stahl. 2010. Rapid evolution of stability and productivity at the origin of a microbial mutualism. Proc. Natl. Acad. Sci. 107:2124–2129.

Hutchison, C. A., R.-Y. Chuang, V. N. Noskov, N. Assad-Garcia, T. J. Deerinck, M. H. Ellisman, J. Gill, K. Kannan, B. J. Karas, L. Ma, J. F. Pelletier, Z.-Q. Qi, R. A. Richter, E. A. Strychalski, L. Sun, Y.

Suzuki, B. Tsvetanova, K. S. Wise, H. O. Smith, J. I. Glass, C. Merryman, D. G. Gibson, and J. C. Venter. 2016. Design and synthesis of a minimal bacterial genome. Science 351:aad6253.

Johnson, M. S., S. Gopalakrishnan, J. Goyal, M. E. Dillingham, C. W. Bakerlee, P. T. Humphrey, T. Jagdish, E. R. Jerison, K. Kosheleva, K. R. Lawrence, J. Min, A. Moulana, A. M. Phillips, J. C. Piper,

R. Purkanti, A. Rego-Costa, M. J. McDonald, A. N. Nguyen Ba, and M. M. Desai. 2021. Phenotypic and molecular evolution across 10,000 generations in laboratory budding yeast populations.

eLife 10:e63910.

Kacar, B., X. Ge, S. Sanyal, and E. A. Gaucher. 2017. Experimental evolution of *Escherichia coli* harboring

564   an ancient translation protein. J. Mol. Evol. 84:69–84.

Kannan, K., B. Tsvetanova, R.-Y. Chuang, V. N. Noskov, N. Assad-Garcia, L. Ma, C. A. Hutchison III, H.

566   O. Smith, J. I. Glass, C. Merryman, J. C. Venter, and D. G. Gibson. 2016. One step engineering of the small-subunit ribosomal RNA using CRISPR/Cas9. Sci. Rep. 6.

568   Kuo, C.-H., and H. Ochman. 2009. Deletional bias across the three domains of life. Genome Biol. Evol. 1:145–152.

570   Lachance, J.-C., S. Rodrigue, and B. O. Palsson. 2019. Minimal cells, maximal knowledge. eLife 8:e45379.

Lang, Gregory. I., D. Botstein, and M. M. Desai. 2011. Genetic variation and the fate of beneficial mutations

572   in asexual populations. Genetics 188:647–661.

Lartigue, C., J. I. Glass, N. Alperovich, R. Pieper, P. P. Parmar, C. A. Hutchison, H. O. Smith, and J. C.

Venter. 2007. Genome transplantation in bacteria: changing one species to another. Science 317:632–638.

Lartigue, C., S. Vashee, M. A. Algire, R.-Y. Chuang, G. A. Benders, L. Ma, V. N. Noskov, E. A. Denisova, D. G. Gibson, N. Assad-Garcia, N. Alperovich, D. W. Thomas, C. Merryman, C. A. Hutchison, H.

O. Smith, J. C. Venter, and J. I. Glass. 2009. Creating bacterial strains from genomes that have been cloned and engineered in yeast. Science 325:1693–1696.

Lenski, R. E., and M. Travisano. 1994. Dynamics of adaptation and diversification: a 10,000-generation experiment with bacterial populations. Proc. Natl. Acad. Sci. U. S. A. 91:6808–6814.

Leprince, A., M. W. van Passel, and V. A. M. dos Santos. 2012. Streamlining genomes: toward the generation of simplified and stabilized microbial systems. Curr. Opin. Biotechnol. 23:651–658.

Liao, Y., S. Ithurbide, C. Evenhuis, J. Löwe, and I. G. Duggin. 2021. Cell division in the archaeon *Haloferax volcanii* relies on two FtsZ proteins with distinct functions in division ring assembly and con-

striction. Nat. Microbiol. 6:594–605.

Lindahl, T. 1974. An N-glycosidase from *Escherichia coli* that releases free uracil from DNA containing

588   deaminated cytosine residues. Proc. Natl. Acad. Sci. U. S. A. 71:3649–3653.

Lindahl, T., S. Ljungquist, W. Siegert, B. Nyberg, and B. Sperens. 1977. DNA N-glycosidases: properties

590   of uracil-DNA glycosidase from *Escherichia coli*. J. Biol. Chem. 252:3286–3294.

Long, H., W. Sung, S. Kucukyildirim, E. Williams, S. F. Miller, W. Guo, C. Patterson, C. Gregory, C.

592   Strauss, C. Stone, C. Berne, D. Kysela, W. R. Shoemaker, M. E. Muscarella, H. Luo, J. T. Lennon,

Y. V. Brun, and M. Lynch. 2018. Evolutionary determinants of genome-wide nucleotide composi-

594  tion. Nat. Ecol. Evol. 2:237–240.

Lynch, M. 2007. The origins of genome architecture. Sinauer Associates, Sunderland, Mass.

596 Lynch, M., M. S. Ackerman, J.-F. Gout, H. Long, W. Sung, W. K. Thomas, and P. L. Foster. 2016. Genetic

drift, selection and the evolution of the mutation rate. Nat. Rev. Genet. 17:704–714.

598 Márquez-Zacarías, P., P. L. Conlin, K. Tong, J. T. Pentz, and W. C. Ratcliff. 2021. Why have aggregative

multicellular organisms stayed simple? Curr. Genet., doi: 10.1007/s00294-021-01193-0.

600 Martin, M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMB-

net.journal 17:10.

602 McQuillen, R., and J. Xiao. 2020. Insights into the structure, function, and dynamics of the bacterial cyto-

kinetic FtsZ-ring. Annu. Rev. Biophys. 49:309–341.

604 Mei, Z.-P., Z. V. Finkel, and A. J. Irwin. 2009. Light and nutrient availability affect the size-scaling of

growth in phytoplankton. J. Theor. Biol. 259:582–588.

606 Mongold, J. A., and R. E. Lenski. 1996. Experimental rejection of a nonadaptive explanation for increased

cell size in *Escherichia coli*. J. Bacteriol. 178:5333–5334.

608 Moran, N. A. 2003. Tracing the evolution of gene loss in obligate bacterial symbionts. Curr. Opin. Micro-

biol. 6:512–518.

610 Moran, N. A., and G. M. Bennett. 2014. The tiniest tiny genomes. Annu. Rev. Microbiol. 68:195–215.

Oksanen, J., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R. Minchin, R. B. O'Hara,

612  G. L. Simpson, P. Solymos, M. H. H. Stevens, E. Szoecs, and H. Wagner. 2020. R package 'vegan':

community ecology package.

614 Pelletier, J. F., L. Sun, K. S. Wise, N. Assad-Garcia, B. J. Karas, T. J. Deerinck, M. H. Ellisman, A. Mershin,

N. Gershenfeld, R.-Y. Chuang, J. I. Glass, and E. A. Strychalski. 2021. Genetic requirements for

616  cell division in a genomically minimal cell. Cell S0092867421002932.

Richardson, S. M., L. A. Mitchell, G. Stracquadanio, K. Yang, J. S. Dymond, J. E. DiCarlo, D. Lee, C. L. V. Huang, S. Chandrasegaran, Y. Cai, J. D. Boeke, and J. S. Bader. 2017. Design of a synthetic yeast genome. Science 355:1040–1044.

Schneider, C. A., W. S. Rasband, and K. W. Eliceiri. 2012. NIH Image to ImageJ: 25 years of image analysis. Nat. Methods 9:671–675.

Sedwick, W. D., O. E. Brown, and B. W. Glickman. 1986. Deoxyuridine misincorporation causes site-specific mutational lesions in the lacI gene of *Escherichia coli*. Mutat. Res. 162:7–20.

Shpak, M., M. M. Goldberg, and M. C. Cowperthwaite. 2015. Rapid and convergent evolution in the Glioblastoma multiforme genome. Genomics 105:159–167.

Sung, W., M. S. Ackerman, M. M. Dillon, T. G. Platt, C. Fuqua, V. S. Cooper, and M. Lynch. 2016. Evolution of the insertion-deletion mutation rate across the tree of life. G3 Genes Genomes Genet. 6:2583–2591.

Sung, W., M. S. Ackerman, S. F. Miller, T. G. Doak, and M. Lynch. 2012. Drift-barrier hypothesis and mutation-rate evolution. Proc. Natl. Acad. Sci. 109:18488–18492.

Tye, B.-K., and I. R. Lehman. 1977. Excision repair of uracil incorporated in DNA as a result of a defect in dUTPase. J. Mol. Biol. 117:293–306.

Vasi, F., M. Travisano, and R. E. Lenski. 1994. Long-term experimental evolution in *Escherichia coli*. II. Changes in life-history traits during adaptation to a seasonal environment. Am. Nat. 144:432–456.

Westfall, C. S., and P. A. Levin. 2017. Bacterial cell size: multifactorial and multifaceted. Annu. Rev. Microbiol. 71:499–517.

Wiser, M. J., and R. E. Lenski. 2015. A comparison of methods to measure fitness in *Escherichia coli*. PLoS ONE 10:e0126210.

Yang, Z. 2006. Computational molecular evolution. Oxford University Press, Oxford.

Table 1. Mutations in genes that were identified as being under positive selection during 2000 generations of experimental evolution. Statistical simulations were performed to find genes acquiring more mutations than expected to occur by chance, indicative of positive selection increasing the allelic proportion of such mutations. P_{adj} corresponds to significance following Benjamin-Hochberg correction to account for multiple comparisons. Genes are assigned to categories based on the secondary functional classifications (Breuer et al. 2019). Note that "Central metabolism" corresponds to "Central carbon metabolism" in the original source (Breuer et al. 2019). While most nonessential genes are absent from the minimal cell (*), a few nonessential genes were retained in *M. mycoides* JCVI-syn3B to facilitate cultivation and robust growth (†). Uncharacterized genes did not fall into defined category (N/A). For mutations that were found in both strains, "non" refers to non-minimal and "min" refers to minimal.

Non-minimal cell only

Gene	Annotation	Category	P_{adj}
lpdA	Dihydrolipoyl dehydrogenase	Central metabolism	<0.0001
pyk	Pyruvate kinase	Central metabolism	0.016
dnaA_1	Chromosomal replication initiator protein	DNA maintenance	0.0005
ecfT	ECF transporter T component	Membrane transport	<0.0001
Locus 0339*	Na$^+$ ABC transporter, ATP-binding component	Membrane transport	0.016
Locus 0030	Uncharacterized ABC transporter ATP-binding	Membrane transport	0.016
rpoA	DNA-directed RNA polymerase, alpha subunit	Transcription	0.0090
Locus 0187*	Transcriptional regulator, GntR family	Transcription	0.0098
tnpA_1*	IS1296 transposase protein A	DNA maintenance	0.0015
tnpB_1*	IS1296 transposase protein B	DNA maintenance	0.0023
Locus 0892*	Putative membrane protein	N/A	0.016

Locus 0898*	Conserved hypothetical protein	N/A	0.029
Locus 0471*	Hypothetical protein	N/A	0.0003
Locus 0751*	Putative viral A-type inclusion protein	N/A	0.016

Minimal cell only

Gene	Annotation	Category	P_{adj}
atpD	F0F1 ATP synthase subunit beta	Central metabolism	<0.0001
pyrG	CTP synthase	Central metabolism	0.0002
dnaN	DNA polymerase III subunit beta	DNA maintenance	<0.0001
lgt	Diacylglyceryl transferase	Biosynthesis	0.0038
clsA	Cardiolipin synthase	Biosynthesis	0.015
fakA	Fatty acid kinase subunit A	Biosynthesis	0.0025
amiF	Oligopeptide ABC transporter ATP-binding	Membrane transport	0.0036
rpoC	DNA-directed RNA polymerase, beta subunit	Transcription	0.0024
rpsC	30S ribosomal protein S3	Translation	0.021
Locus 0373	Uncharacterized protein	N/A	0.015
Locus 0691	Uncharacterized protein	N/A	0.027
Locus 0430	Uncharacterized DNA-binding protein	N/A	0.0012

Minimal and non-minimal cell

Gene	Annotation	Category	P_{adj}
ftsZ †	Cell division protein ftsZ	Cell growth	non <0.0001 min <0.0001
tetM	Tetracycline resistance ribosomal protection	Drug resistance	non = 0.029 min = 0.0079

FIGURE CAPTIONS

656   **Fig. 1.** Mutation rate and spectrum of the minimal and non-minimal cell estimated from mutation

accumulation experiments. (**A**) Although *M. mycoides* has the highest recorded mutation rate

658   (base substitutions and indels), it was not affected by genome minimization. (**B**) The proportions

of insertions, deletions, and single nucleotide mutations (SNMs) were also the same for the mini-

660   mal and non-minimal cell. (**C**) Among SNMs, which accounted for 88% of all mutations, the

minimal cell exhibited a stronger AT bias in its mutation spectrum than the non-minimal cell,

662   particularly in the C:G-to-A:T category. In this panel, ns = not significant and *** = $P < 0.001$.

664   **Fig. 2.** Genome minimization resulted in a 50% reduction in relative fitness, but 80% of this cost

was regained over 2000 generations of evolution. Despite removal of nearly half of its genome,

666   the minimal cell adapted at the same rate as the non-minimal cell. Data represent mean ± SEM.

Because the experiment was initiated with a single clone, error bars for the ancestral time point

668   were calculated from experimental replicates, while error bars for evolved lines were calculated

from replicate populations.

    Fig. 3. The non-minimal cell and minimal cell acquired adaptive mutations in different sets of

672   shared essential genes ($P = 0.029$) over 2000 generations of evolution (Table 1) as depicted in

ordination from a principal coordinates analysis (PCoA) created with a gene-by-population ma-

674   trix using Bray-Curtis distance metric. Dashed lines represent 95% confidence ellipses around

replicate populations (solid symbols).

Fig. 4. Effect of genome minimization on the evolution of cell size. (**A**) The size of non-minimal cells increased by 73% over 2000 generations of evolution, while the size of minimal cell remained the same. Data represent means ± SEM based on images obtained via phase contrast microscopy. Scanning electron micrographs obtained from evolved replicate populations of the (**B**) non-minimal and (**C**) minimal cell.

Fig. 1

Fig. 1. Mutation rate and spectrum of the minimal and non-minimal cell estimated from mutation accumulation experiments. (**A**) Although *M. mycoides* has the highest recorded mutation rate (base substitutions and indels), it was not affected by genome minimization. (**B**) The proportions of insertions, deletions, and single nucleotide mutations (SNMs) were also the same for the minimal and non-minimal cell. (**C**) Among SMNs, which accounted for 88% of all mutations, the minimal cell exhibited a stronger AT bias in its mutation spectrum than the non-minimal cell, particularly in the C:G-to-A:T category. In this panel, ns = not significant and *** = P < 0.001.

Fig. 2. Genome minimization resulted in a 50% reduction in relative fitness, but 80% of this cost was regained over
2000 generations of evolution. Despite removal of nearly half of its genome, the minimal cell adapted at the same
rate as the non-minimal cell. Data represent mean ± SEM. Because the experiment was initiated with a single clone,
error bars for the ancestral time point were calculated from experimental replicates, while error bars for evolved lines
were calculated from replicate populations.

694
696
698

706 **Fig. 3**

710 **Fig. 3.** The non-minimal cell and minimal cell acquired adaptive mutations in different sets of shared essential genes (P = 0.029) over 2000 generations of evolution (Table 1) as depicted in ordination from a principal coordinates analy-
712 sis (PCoA) created with a gene-by-population matrix using Bray-Curtis distance metric. Dashed lines represent 95% confidence ellipses around replicate populations (solid symbols).

Fig. 4

Fig. 4. Effect of genome minimization on the evolution of cell size. (**A**) The size of non-minimal cells increased by
728   73% over 2000 generations of evolution, while the size of minimal cell remained the same. Data represent means ±
SEM based on images obtained via phase contrast microscopy. Scanning electron micrographs obtained from
730   evolved replicate populations of the (**B**) non-minimal and (**C**) minimal cell.