Intrinsic Network Connectivity Reflects the Cyclic Trajectory of Migraine Attacks

Anne Stankewitz [1,2]*, Enrico Schulz [1,3]*

[1]Department of Neurology, Ludwig-Maximilians-Universität München, Munich, Germany

[2]Department of Neurology, Klinikum rechts der Isar, Technische Universität München, Munich, Germany

[3]Department of Medical Psychology, Ludwig-Maximilians-Universität München, Munich, Germany

Corresponding author:
Dr Enrico Schulz
Klinikum der Universität München, Department of Neurology
Ludwig-Maximilians-Universität München
A: Fraunhoferstr. 20, 82152 Martinsried, Germany
O: +49 89 4400 74826
E: es@pain.sc

Abstract

Background. Episodic migraine is considered to be cyclic in nature, triggered by the hypothalamus. To assess the natural trajectory of intrinsic networks over an entire migraine cycle, we designed a longitudinal intra-individual study using functional magnetic resonance imaging (fMRI).

Methods. Intrinsic network connectivity was assessed for 12 migraineurs in 82 sessions including spontaneous, untriggered headache attacks and follow-up recordings towards the next attack.

Results. We found cyclic changes in the visual, auditory, and somatosensory networks, in limbic networks (e.g. thalamo-insular, parahippocampal), and in the salience network (anterior insula and dorsal anterior cingulate cortex). Connectivity changes also extended to further cortical networks, such as the central executive network, the default mode network, as well as subcortical networks.

Almost all of these network connectivity changes followed the trajectory of a linear increase over the pain-free interval that peaked immediately prior to the headache, and "dropped" to the baseline level during the headache. These network alterations are associated with a number of cortical functions that may explain the variety of ictal and pre-ictal physiological and psychological migraine symptoms.

Conclusion. Our results suggest that migraine disease is associated with widespread cyclic alterations of intrinsic networks that develop before the headache is initiated, i.e. during the interictal and premonitory phase. The increasing magnitude of connectivity within these networks towards the next attack may reflect an increasing effort to maintain network integrity.

Introduction

Migraine is a cyclic disease that affects approximately 10% of the population. Recurring headache episodes are the most disabling symptom, but many patients also suffer from a variety of sensory and autonomous symptoms [1].

Migraine pathophysiology is complex and, thus far, not fully understood. Resting-state functional magnetic resonance imaging (rs-fMRI) has highlighted the presence of dysfunctional connectivities in migraineurs, particularly in somatosensory [2], dorsal attention [3,4], salience [2], executive control [4], and default mode networks [4,5].

Such a broad reorganisation of brain networks is thought to influence multisensory integration processes. In particular, alterations of limbic and sensory networks may lead to an increased sensory susceptibility that might make the migraineurs' brain vulnerable to intrinsic and external factors that trigger migraine attacks. Despite the cyclic nature of the migraine disease, longitudinal studies are sparse [6,7]. The majority of neuroimaging studies have used cross-sectional designs and found deviant cortical activity [8,9], seed-based connectivity [10], and resting-state network connectivity in migraineurs.

Here, we conducted an intra-individual rs-fMRI study to follow the trajectory of intrinsic cortical networks over the entire migraine cycle. Due to the clinical features of the disease and previous findings using neurophysiological and neuroimaging techniques [11,12], we hypothesised substantial alterations of sensory, limbic, and the salience networks during the migraine cycle.

Materials and Methods

Subjects

Twenty-two episodic migraine patients were included in the study, twelve of these data sets were complete and suitable for the statistical analysis (11 females and one male). Drop-outs from the initially-recruited subjects resulted from incidental MRI findings (n=2), technical problems with the scanner hardware (n=3) or software (n=1), illness of patients (n=2), or taking analgesic medication due to an acute painful event (n=1) during the scanning period. One patient decided to withdraw prematurely from the study. The exclusion of subjects was required due to the within-subject design, as the inclusion of incomplete data could have led to an inaccurate estimation of the subjects' random effects in the statistical model. Moreover, an incomplete time series would miss mandatory data for the period immediately before the migraine attack.

Migraine diagnosis was based upon the classification of the International Headache Society [13]. The patients did not report any other neurological or psychiatric disorders, were not taking preventative medication for migraine for at least six months, but were allowed to take their regular acute migraine medication after the recording of the headache attack (non-steroidal anti-inflammatory drugs or triptans). All patients gave their written, informed consent. The study was conducted according to the Declaration of Helsinki and approved by the Ethics Committee of the Technische Universität München, Germany. All patients were remunerated for participation.

Study design

Using a longitudinal, intra-individual study design, migraine patients were tested repeatedly over an entire migraine cycle (Figure 1). The imaging time series for each patient started with the recording of a spontaneous, untriggered and untreated headache attack within the first 6 hours after the beginning of the headache. We only recorded headache attacks which were reported with an intensity of middle to strong, predefined as a minimum of "4" on a numerical rating scale with the endpoints zero (no pain) and 10 (extremely intense pain). Brain data were then recorded every 1-4 days at the same time of day until patients informed us by phone about the following headache attack (which was not scanned). The time series was completed with the last attack-free recording.

Image acquisition

MRI data were collected on a 3 Tesla scanner (Ingenia, Philips, The Netherlands) using a 32-channel head coil. Patients were instructed to remain awake and relaxed with their eyes closed. For the 300 volumes of resting state data, we used the following parameters: TR = 2000 ms; time to echo (TE) = 30 ms; FOV = 192 × 192 mm^2; flip angle = 90°; number of slices = 37; voxel size = 3 × 3 x 3 mm^3 (0.29 mm gap). For image registration, a high resolution T1-weighted anatomical image was collected with: TR = 9000 ms, TE = 4 ms, flip angle = 8°, FOV = 240 x 240 x 170 mm^3; number of slices = 170; voxel size = 1.0 x 1.0 × 1.0 mm^3). Field maps were acquired in each session to control for B0-effects; 64 slices, TR = 960 ms, FOV = 192 × 192 mm^2; voxel size = 2.0 x 2.0 × 2.0 mm^3, 0.2 mm gap between slices. TE = 6 ms / 10.55 ms, flip angle 60°.

Image preprocessing

The data were preprocessed with FSL. The Melodic toolbox was used to execute brain extraction, high-pass filtering with a frequency cutoff of 1/100 Hz, spatial registration to the MNI template, corrections for head motion during scanning, and - due to the potential contribution of smaller subcortical structures - a spatial smoothing (5 mm FWHM). The relatively small smoothing kernel was chosen to enable the detection of nucleic connectivity in subcortical regions. A distortion correction of the images was used based on field maps. The data were further semi-automatically cleaned of artefacts with ICA through Melodic. The number of components had been automatically estimated by Melodic and artefact-related components were removed from the data. Head movement during scanning did not exceed 2 mm or 2° in any direction. The movement during the migraine recording was within the range of the movement of the pre-ictal recordings (relative: 0.07±0.02 vs. 0.08±0.02; absolute: 0.2±0.7 vs 0.19±0.07).

Statistical analyses

Following preprocessing, we ran a group ICA with temporally concatenated MNI-registered data of all 81 recordings using Melodic. Sixty-one networks (components) were determined by the ICA using automatic dimensionality estimation. Nineteen of these components were identified as artefacts by visual inspection (see Supplementary Material). Thus, we selected 42 resting-state networks. A dual regression analysis computed the individual maps for all 42 networks and each of the 81 sessions. We were running a voxel-wise statistics within the boundaries of the group network maps (as defined by Melodic) for the time series of 81 recordings. We aimed to investigate whether the trajectory of the time series is following two predefined time courses. In order to do so, we created two time vectors for each patient's migraine cycle and encoded the day of the recording by assigning numbers between one and two. The numbers one or two were assigned to the measurements during the headache attack, depending on the following two possible trajectories of cortical processing (Fig. 2):

Trajectory one ("reset mode"): In this predefined model, we aimed to detect the resting-state connectivity that has its peak (or trough) just *before* the headache attack starts (during the premonitory phase) and then drops (or jumps) back to normal during the headache [14–17]. From here, we model a linear increase (or decrease) over the migraine cycle to the next attack (Fig. 2, upper part). In this hypothetical time course, the magnitude of the cortical map on the first day after the attack would be similar to the brain connectivity during the attack. This trajectory can be interpreted as a cortical "reset" mechanism and is in line with neurophysiological and imaging studies suggesting altered cortical processes in migraineurs during the interictal interval that normalise just before or during headache attacks [14–17]. For example, for five measurements over 10 days, the following vector is used to reflect trajectory one: 1 (=attack), 1.2, 1.4, 1.6, 1.8.

Trajectory two ("pain mode"): This model detects the cortical processes that have their peak (or trough) *during* the headache attack then drops (or jumps) back to normal the next day. From there, we assume a linear increase (or decrease) over the migraine cycle towards the next attack (Fig. 2, lower part). We hypothesised increased magnitude of brain connectivity in regions that contribute to the processing of migraine symptoms, e.g. pain, increased sensitivity to light, sound, and odours, and vegetative complaints. In this hypothetical time course, the brain connectivity on the day prior to the attack would be similar to the brain connectivity during the headache attack. Similar to the above-mentioned example with five measurements over 10 days, the following vector is used for trajectory two: 2 (=attack), 1.2, 1.4, 1.6, 1.8.

We explored the cyclic change of cortical rsfMRI connectivity over the migraine interval separately for each network. We computed voxel-wise linear mixed effects models (LME; [18] in Matlab (Version R2018a, Mathworks, USA) within the z-masks of the network map as generated by Melodic. We related the time points within the migraine cycle to the session-specific cortical map of interest:

(1) *ICAcmp_timecourse ~ time + (1| subject)*

T-values of the fixed effect were computed voxel-wise as quotients between the beta estimates and the standard errors of the equation. The statistical results were corrected for multiple testing using the FDR procedure [19].

Data availability

Inquiries for additional data are available upon reasonable request.

Results

Clinical characteristics

Twelve migraine patients were included in the study (mean age: 28; range: 21–40 years; 11 females). Migraine patients reported a mean disease duration of 13.6 years (range, 6-29 years). The attack frequency was recorded as the number of episodes per month and ranged between one and 10 per month. Nine patients reported migraine without aura. Laterality of headache pain was reported as either unilateral (five right-sided, three left-sided) or bilateral (four). Attack severity of the scanned headache was recorded on a numerical rating scale ranging from zero (no pain) to 10 (highest imaginable pain) and ranged between four and nine. The time points of the attacks were equally distributed across the menstrual cycles of the female patients.

Imaging results

We observed connectivity changes over the migraine cycle in 28 networks; 27 network changes apply to trajectory one and one network change applies to trajectory two. An overview of network changes depending on the trajectories is shown in Table 1; detailed information is given in Table 2. Maps of all networks are included in the Supplementary Material.

Table 1. Intrinsic networks that showed activity changes over the migraine cycle.

	Increasing cycle	Decreasing cycle
Trajectory 1	visual (basal visual areas), auditory, somatosensory, central executive, salience, cerebellar, basal ganglia, pDMN, thalamic, frontal, temporal, sensory-motor, motor, limbic, insular, and cortically distributed (occipito-parietal and fusiform-parietal) networks	visual (higher order areas), cerebellar networks
Trajectory 2	-	limbic

Trajectory one: Changing connectivity predominantly applies to trajectory one, where 25 functional networks showed a linear increase of the magnitude of network connectivity over the interictal interval and a "drop" during the headache (Table 2a):

(I) sensory: two visual networks (intra-calcarine cortex and occipital pole (=basal/ primary visual areas), lingual gyrus as well as precuneus), two somatosensory networks (post- and precentral gyrus, supplementary motor area, and cerebellum) and one auditory network (planum temporale and central operculum) networks;

(II) two sensory-motor networks (precentral gyrus and supplementary motor area); III) central executive network (lateral occipital cortex and cerebellum);

(IV) salience network (dorsal anterior cingulate cortex, anterior insula, frontal pole, and central operculum);

(V) cortically distributed networks: two frontal networks (orbitofrontal cortex, frontal pole, and precentral gyrus), one temporo-supramarginal network (superior temporal cortex and orbitofrontal cortex), one occipito-lingual network (lateral occipital cortex and middle temporal gyrus) and one fusiform-parietal-frontal network (e.g. supramarginal gyrus and inferior frontal cortex);

(VI) two limbic networks (hippocampus and the dorsolateral prefrontal cortex (DLPFC);

(VII) insular network (planum polare);

(VIII) thalamo-insular network;

(IX) basal ganglia network (putamen and thalamus);

(X) posterior cingulate-precuneus network (precuneus);

(XI) lingual network (posterior cingulate cortex, precuneus, supracalcarine cortex); XII) three cerebellar networks: intra-cerebellar (cerebellum), cerebello-insular (hippocampus) and cerebello-ponto-thalamic (temporal and lateral occipital cortices, cerebellum and lingual gyrus);

(XIII) posterior default mode network (DMN) (posterior cingulate cortex).

In two of these 25 networks, we also found brain regions that showed the inverse behaviour, namely a linear decrease of the magnitude of network connectivity over the pain-free interval with peak magnitude during the headache:

(I) visual network (lateral occipital cortex (= higher order visual area) and the posterior cingulate cortex); and

(II) cerebello-ponto-thalamic network (occipital fusiform cortex and cerebellum) (Table 2b).

Figure 3 shows exemplary results in the visual (Fig. 3a), thalamo-insular (Fig. 3b), and the salience networks (Fig. 3c). All statistical tests were thresholded at $p < 0.05$ (FDR corrected).

Trajectory two: One limbic network followed trajectory two: the connectivity of the parahippocampal gyrus linearly decreased towards the next attack and reached its lowest magnitude during the headache (Table 2c). We did not observe any significant effect of the opposite contrast (linear increases over the pain-free interval with peak magnitude during the headache). All statistical tests were thresholded at $p < 0.05$ (FDR corrected).

Table 2a-c. Cyclic changes of intrinsic networks

a. Trajectory 1 - linear *decreasing* activity over the pain-free interval with a *peak* during the headache

C	cluster sizes	t-values	MNI coordinates			Brain regions	Functional networks
			x	**y**	**z**		
5	16	4.03	26	-90	18	Lateral Occipital Cortex	Visual
43	100	6.1	22	-90	-18	Occipital Fusiform Gyrus	Cerebellar
	21	4.6	0	-52	-22	Cerebellum	

b. Trajectory 1 - linear *increasing* activity over the pain-free interval with a *"drop"* during the headache

C	cluster sizes	t-values	MNI coordinates			Brain regions	Functional networks
			x	**y**	**z**		
4	21	4.36	-28	-80	-48	Cerebellum	Attention
	18	4.6	40	-60	42	Lateral Occipital Cortex	
5	105	4.64	14	-64	28	Precuneus Cortex	Visual
	29	4.32	-26	-62	4	Intracalcarine Cortex	
9	72	4.32	28	-90	30	Occipital Pole	Visual
13	17	4.15	46	-70	-36	Cerebellum	Cerebellar
14	96	4.41	-16	-70	28	Precuneus Cortex	Posterior DMN
	60	4.17	22	-64	32	Precuneus Cortex	
19	89	4.95	-52	-10	18	Central Opercular Cortex	Temporal
	18	4.05	64	-12	6	Planum Temporale	
20	35	5.24	-60	-36	42	Supramarginal Gyrus	Cortically distributed
21	19	3.88	4	-8	58	Juxtapositional Lobule Cortex	Motor
22	22	4.68	-36	10	-20	Temporal Pole	Auditory
25	20	4.22	-2	-30	26	Posterior Cingulate Gyrus	Posterior DMN
27	437	6.04	-2	-20	2	Thalamus	Thalamic
	34	4.5	18	-8	-6	Pallidum	
29	21	4.23	-44	-78	26	Lateral Occipital Cortex	Cortically distributed
	18	4.48	-62	-60	-6	Middle Temporal Gyrus	
30	23	4.91	-22	60	-2	Frontal Pole	Frontal
32	31	4.24	24	0	-8	Putamen	Basal ganglia
	28	4.47	-12	0	8	Caudatus	
34	136	5.04	0	34	30	Paracingulate Gyrus	Salience
	23	5.02	28	36	28	Frontal Pole	

C	cluster size	t-values	x	y	z	Brain regions	Functional networks
	17	4.16	-44	-22	18	Central Opercular Cortex	
	16	4.71	40	12	-6	Insular Cortex	
35	29	4.53	56	-34	4	Superior Temporal Gyrus	Temporal
	20	4.7	30	20	-20	Frontal Orbital Cortex	
36	23	4.13	-44	-18	38	Postcentral Gyrus	Sensory-motor
37	39	4.46	48	26	0	Frontal Orbital Cortex	Frontal
	18	4.01	52	-8	46	Precentral Gyrus	
	18	5.15	18	20	68	Superior Frontal Gyrus	
39	26	4.85	44	-2	-6	Insular Cortex	Cerebellar
43	126	5.44	36	-44	-28	Temporal Occipital Fusiform Cortex	Cerebellar
	95	4.69	-52	-80	2	Lateral Occipital Cortex	
	50	4.33	-16	-60	-12	Lingual Gyrus	
	20	4.37	-50	-68	20	Lateral Occipital Cortex	
44	49	4.73	-30	-24	56	Precentral Gyrus	Sensory-motor
	48	4.21	-36	-28	40	Postcentral Gyrus	
	22	4.58	12	-52	-16	Cerebellum	
51	37	4.6	36	34	18	Dorsolateral Prefrontal Cortex	Limbic
58	152	5.52	32	-6	46	Precentral Gyrus	Sensory
	39	5.13	0	-2	52	Juxtapositional Lobule Cortex	

c. Trajectory 2 - linear *decreasing* activity over the pain-free interval with a *"drop"* during the headache

C	cluster size	t-values	MNI coordinates			Brain regions	Functional networks
			x	y	z		
54	40	4.92	-18	-4	-26	Parahippocampal Gyrus	Limbic

Discussion

The present longitudinal fMRI study aimed to study the cyclic connectivity of intrinsic functional networks in episodic migraine. We analysed two possible network trajectories over the migraine interval.

Sensory networks. Alterations of sensory processing in migraine have been shown in various studies [11,20]. Sensory thresholds have been suggested to depend on the phase within the migraine cycle, where the lowest thresholds were detected during the headache phase [11,12]. Here, we mainly observed linearly increasing sensory network

connectivity over the pain-free interval with a peak prior to the beginning of the headache that "dropped" to the initial level during the headache in brain regions belonging to visual (e.g. intracalcarine cortex and occipital pole), auditory (e.g. planum temporale), and somatosensory networks (e.g. postcentral gyrus). For a visual network we also observed the inverse effect, where the connectivity of the lateral occipital cortex decreased over the pain-free interval and reached its peak during the headache. An increasing magnitude of network connectivity in sensory networks over the migraine cycle towards the attack, as observed in our study, is likely to reflect an increased sensitivity to sensory input. Additionally, it may reflect an increasing vulnerability to internal and external stressful events during the pain-free interval. External triggers for migraine attacks in some patients, such as bright or flickering light [21] and odours [22], highlight dysfunctional sensory processing. The origin of such hypersensitivity in migraineurs is not yet understood [23]. Genetic predispositions, such as ion channelopathies, can affect excitability thresholds making the cortex more vulnerable to diencephalic (thalamus, hypothalamus) and pontine inflow [24]. Consequently, repetitive stimulation may result in an overload of cortical sensory neurons. The "drop" of network connectivity during the headache suggests a "rebooting" of cortical processes.

Salience networks. In line with our hypotheses, we found cyclic changes in the salience network, e.g. in the anterior cingulate cortex and the anterior insula. Similar to the sensory networks, connectivity increased over the interictal interval, reached the maximum prior to the headache, and "dropped" to the baseline level during the attack. These findings are corroborated by previous work on ictal and pre-ictal alterations of the salience network in migraine [2].
It could be speculated that the deviances in the salience network are associated with an interictally-impaired habituation in migraine, which could be explained either by reduced intra-cortical inhibition or by increased cortical excitability. Indeed, some studies report an abnormal level of both inhibitory and excitatory neurotransmitters, such as in GABA (gamma-aminobutyric acid) and glutamate [25]. Specifically, an increased amount of glutamate has been observed in migraine patients in the anterior paracingulate cortex, a region that is an essential part of the salience network [26]. Numerous neurophysiological studies confirm the impaired inhibition by showing increased responsiveness to sensory stimulation in the visual, auditory, and somatosensory domain (for a comprehensive review see [27]).

Limbic networks. Altered limbic connectivity is likely to be responsible for several migraine symptoms during the prodromal, ictal and postictal phases of the migraine cycle (e.g. fatigue, irritability, yawning, polyuria, and food cravings). Furthermore, many trigger factors for migraine attacks are associated with limbic circuits, e.g. psycho-physical distress and emotions, homeostatic changes or circadian rhythms [28,29].

Here, we revealed connectivity changes of the right DLPFC. This region is integrated in a limbic network, which also includes the hippocampus, thalamus, hypothalamus, cingulate cortex, and the insula. The DLPFC connectivity increased over the pain-free interval, reached its peak immediately prior to the headache, and "dropped" to the

lowest level during the headache. The DLPFC is a key node of cognitive circuits and contributes to executive function, attention, decision-making, and emotional regulation [30]. The DLPFC further contributes to the top-down control of nociceptive input [31]. Increasing connectivity of the DLPFC over the pain-free interval may reflect the increasing prefrontal effort to control cognitive, emotional, and sensory processes, whereas a "drop" in DLPFC connectivity during the ictal phase is likely to reflect the altered pain and cognitive processes resulting in headache and cognitive deficits.

In a further limbic network, the connectivity of the left hippocampus (as part of a bilateral hippocampus-amygdala network) increased interictally and "dropped" during the headache. There is growing evidence showing a cortical effect of the hippocampus in migraineurs in comparison to healthy controls, e.g. a reduced volume [32] and a dependency of its connectivity on the attack frequency [33]. The amygdala-hippocampus network has been associated with the processing of mood [34], memory encoding [35], psychiatric diseases [36], and further emotion-related cognitive functions [37]. Our findings are in line with these recent studies and could reflect an increase in anxiety, decrease in mood, and pain-related rumination towards the next attack. Indeed, a previous study reported growing connectivity between the amygdala and the hippocampus after stress [38].

In addition, a limbic network, composed of the bilateral parahippocampal gyri, hippocampi and the amygdalae, showed a linearly decreasing magnitude of connectivity of the left parahippocampal gyrus over the pain-free interval and reached its lowest level during the headache. The network that comprises these three regions has been associated predominantly with the processing of memory [39] and object novelty [40]. Further, lesions in the parahippocampal gyrus were found to cause severe memory deficits [41] and impaired emotional responses [42].
The migraineurs' imperative focus on the headache is suggested to explain the low connectivity of the parahippocampus during the attack. This focus may impair cognitive functions *per se*, but can also prevent the processing of any past- or future-related memory during attacks [43].

Contrary to our hypothesis, we did not find any cyclic alterations of hypothalamic networks. One explanation might be that the hypothalamus is not a permanent part of the intrinsic networks, but may instead orchestrate different networks at different time points during the migraine interval as a rhythm generator.

Thalamo-insular network. There is increasing neurophysiological and neuroimaging evidence for the role of the thalamus within the migraine cycle [44]. As the thalamus is connected to the sensory, salience, and limbic networks in an upstream direction, almost every stimulus that reaches the cortex is first processed and gated by the thalamus [45]. We observed bilateral connectivity changes of the thalamus within a thalamo-insular network. Thalamic connectivity changes followed the trajectory of a linear increase over the interictal interval with a peak immediately prior to the headache and a "drop" to the baseline level during the headache. Structural [46], functional [44], and neurochemical thalamic alterations [47] in migraine have been highlighted in recent

imaging studies. During spontaneous attacks, altered functional connectivities between the thalamus and cortical regions contributing to pain processing and modulation (e.g. the insula and the orbitofrontal cortex) have been observed [48]. Furthermore, using diffusion tensor MRI it has been shown that migraine patients exhibited a higher fractional anisotropy in the thalamus during the interictal interval, which normalised during the attack [17].

Through the connections between the thalamus and the principal structures of the salience network (anterior insula, anterior cingulate cortex), the current results may suggest a maladaptive gating mechanism in the migraineurs' brain that usually prevents an overload of irrelevant and unnecessarily salient information to higher cortical areas. In migraine, internal and external input to the cortex is likely to be insufficiently filtered or inhibited by GABAerge thalamo-cortical and cortico-thalamic circuits [12].

Other networks. We also found cyclic connectivity changes in various other cortical (temporal and frontal networks, as well as in cortically distributed networks) and subcortical networks (cerebellar and basal ganglia). Most of these networks followed the trajectory of a linear increase over the interictal interval with a peak prior to the headache and a "drop" to the baseline level during the headache. This applies to the posterior DMN, which has previously been shown to be altered in migraineurs [5] and other chronic pain states [49]. DMN dysfunctions have been related to maladaptive stress responses, which seems to characterise chronic pain patients in general [49]. Similarly, we found cyclic changes of the central executive network. Although cognitive symptoms are not included as core symptoms to diagnose migraine disease, they are frequently reported by patients during the premonitory, ictal, and postdrome phases [50].

Summary. The present results suggest that migraine disease is associated with cyclic alterations of various functional networks, underlining the complexity of the clinical picture. Importantly, network changes evolve early during the migraine cycle, long before the headache is initiated.

Our findings provide further evidence for the use of psychological approaches during the pain-free migraine interval. Cognitive-behavioural interventions, relaxation techniques, or biofeedback, which are aimed at normalising network functions, help patients to regulate major trigger factors of migraine attacks (e.g. psycho-physical stress, sleep-awake rhythms). By modulating sympathetic activity or opioidergic activity, these interventions are likely to prevent a recurring loss of network synchronicity over the migraine cycle.

Acknowledgements

We thank Dr Stephanie Irving for providing comments and copy-editing the manuscript. We thank Prof Dr Andreas Straube and Dr Daniel Keeser for their comments.

Funding

The study has been funded by the Else-Kröner-Fresenius Stiftung (2014-A85).

Competing Interests

The authors report no competing interests.

Key Findings

- Cortical processes of migraineurs undergo cyclic changes, already detectable in the pain-free phase.
- The increasing magnitude of connectivity towards the next attack in several networks reflects the increasing effort to keep the network integrity intact.
- Network decoupling in the ictal phase can explain the variety of ictal and pre-ictal migraine symptoms.
- These findings provide further evidence for the need of early therapeutic approaches during the pain-free interval.

References

1. Blau JN. Migraine: theories of pathogenesis. *Lancet* 1992; 339: 1202–1207.
2. Amin FM, Hougaard A, Magon S, et al. Change in brain network connectivity during PACAP38-induced migraine attacks: A resting-state functional MRI study. *Neurology* 2016; 86: 180–187.
3. Niddam DM, Lai K-L, Fuh J-L, et al. Reduced functional connectivity between salience and visual networks in migraine with aura. *Cephalalgia* 2016; 36: 53–66.
4. Coppola G, Di Renzo A, Petolicchio B, et al. Aberrant interactions of cortical networks in chronic migraine: A resting-state fMRI study. *Neurology* 2019; 92: e2550–e2558.
5. Coppola G, Di Renzo A, Tinelli E, et al. Resting state connectivity between default mode network and insula encodes acute migraine headache. *Cephalalgia* 2018; 38: 846–854.
6. Schulte LH, Menz MM, Haaker J, et al. The migraineur's brain networks: Continuous resting state fMRI over 30 days. *Cephalalgia* 2020; 333102420951465.
7. Schulte LH, May A. Of generators, networks and migraine attacks. *Curr Opin Neurol* 2017; 30: 241–245.
8. Stankewitz A, Aderjan D, Eippert F, et al. Trigeminal nociceptive transmission in migraineurs predicts migraine attacks. *J Neurosci* 2011; 31: 1937–1943.
9. Stankewitz A, May A. Increased limbic and brainstem activity during migraine attacks following olfactory stimulation. *Neurology* 2011; 77: 476–482.
10. Schwedt TJ, Chong CD. Functional imaging and migraine: new connections? *Curr Opin Neurol* 2015; 28: 265–270.
11. Harriott AM, Schwedt TJ. Migraine is associated with altered processing of sensory stimuli. *Curr Pain Headache Rep* 2014; 18: 458.
12. Peng K-P, May A. Migraine understood as a sensory threshold disease. *Pain* 2019; 160: 1494–1501.
13. Headache Classification Committee of the International Headache Society (IHS). The

International Classification of Headache Disorders, 3rd edition. *Cephalalgia* 2018; 38: 1–211.

14. Judit A, Sándor PS, Schoenen J. Habituation of visual and intensity dependence of auditory evoked cortical potentials tends to normalize just before and during the migraine attack. *Cephalalgia* 2000; 20: 714–719.

15. Chen W-T, Wang S-J, Fuh J-L, et al. Peri-ictal normalization of visual cortex excitability in migraine: an MEG study. *Cephalalgia* 2009; 29: 1202–1211.

16. Kropp P, Gerber WD. Prediction of migraine attacks using a slow cortical potential, the contingent negative variation. *Neurosci Lett* 1998; 257: 73–76.

17. Coppola G, Tinelli E, Lepre C, et al. Dynamic changes in thalamic microstructure of migraine without aura patients: a diffusion tensor magnetic resonance imaging study. *Eur J Neurol* 2014; 21: 287–e13.

18. Witkovský V. Estimation, testing, and prediction regions of the fixed and random effects by solving the Henderson's mixed model equations. *Measurement science review* 2012; 12: 234–248.

19. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J R Stat Soc Series B Stat Methodol* 1995; 57: 289–300.

20. de Tommaso M, Ambrosini A, Brighina F, et al. Altered processing of sensory stimuli in patients with migraine. *Nat Rev Neurol* 2014; 10: 144–155.

21. Hougaard A, Amin FM, Hauge AW, et al. Provocation of migraine with aura using natural trigger factors. *Neurology* 2013; 80: 428–431.

22. Kelman L. The triggers or precipitants of the acute migraine attack. *Cephalalgia* 2007; 27: 394–402.

23. Demarquay G, Mauguière F. Central Nervous System Underpinnings of Sensory Hypersensitivity in Migraine: Insights from Neuroimaging and Electrophysiological Studies. *Headache* 2016; 56: 1418–1438.

24. Kullmann DM. Neurological channelopathies. *Annu Rev Neurosci* 2010; 33: 151–172.

25. Bigal ME, Hetherington H, Pan J, et al. Occipital levels of GABA are related to severe headaches in migraine. *Neurology* 2008; 70: 2078–2080.

26. González de la Aleja J, Ramos A, Mato-Abad V, et al. Higher glutamate to glutamine ratios in occipital regions in women with migraine during the interictal state. *Headache* 2013; 53: 365–375.

27. Coppola G, Pierelli F, Schoenen J. Is the cerebral cortex hyperexcitable or hyperresponsive in migraine? *Cephalalgia* 2007; 27: 1427–1439.

28. Karsan N, Bose P, Newman J, et al. Are some patient-perceived migraine triggers simply early manifestations of the attack? *J Neurol*. Epub ahead of print 5 January 2021. DOI: 10.1007/s00415-020-10344-1.

29. van Casteren DS, Verhagen IE, Onderwater GLJ, et al. Sex differences in prevalence of migraine trigger factors: A cross-sectional study. *Cephalalgia* 2020; 333102420974362.

30. Carlén M. What constitutes the prefrontal cortex? *Science* 2017; 358: 478–482.

31. Eippert F, Bingel U, Schoell ED, et al. Activation of the opioidergic descending pain control system underlies placebo analgesia. *Neuron* 2009; 63: 533–543.

32. Chong CD, Dumkrieger GM, Schwedt TJ. Structural Co-Variance Patterns in Migraine: A Cross-Sectional Study Exploring the Role of the Hippocampus. *Headache: The Journal of Head and Face Pain* 2017; 57: 1522–1531.

33. Maleki N, Becerra L, Brawn J, et al. Common hippocampal structural and functional changes in migraine. *Brain Struct Funct* 2013; 218: 903–912.

34. Kirkby LA, Luongo FJ, Lee MB, et al. An Amygdala-Hippocampus Subnetwork that Encodes Variation in Human Mood. *Cell* 2018; 175: 1688–1700.e14.

35. Erin S. Predictors of successful memory encoding in the human hippocampus and amygdala. *Frontiers in Systems Neuroscience*; 3. Epub ahead of print 2009. DOI: 10.3389/conf.neuro.06.2009.03.092.

36. Sylvester CM, Yu Q, Srivastava AB, et al. Individual-specific functional connectivity of the amygdala: A substrate for precision psychiatry. *Proc Natl Acad Sci U S A* 2020; 117: 3808–3818.

37. Zheng J, Anderson KL, Leal SL, et al. Amygdala-hippocampal dynamics during salient information processing. *Nature Communications*; 8. Epub ahead of print 2017. DOI: 10.1038/ncomms14413.

38. Ghosh S, Laxmi TR, Chattarji S. Functional connectivity from the amygdala to the hippocampus grows stronger after stress. *J Neurosci* 2013; 33: 7234–7244.

39. Kilpatrick L, Cahill L. Amygdala modulation of parahippocampal and frontal regions during emotionally influenced memory storage. *Neuroimage* 2003; 20: 2091–2099.

40. Kaplan R, Horner AJ, Bandettini PA, et al. Human hippocampal processing of environmental novelty during spatial navigation. *Hippocampus* 2014; 24: 740–750.

41. Zola-Morgan S, Squire LR, Amaral DG, et al. Lesions of perirhinal and parahippocampal cortex that spare the amygdala and hippocampal formation produce severe memory impairment. *J Neurosci* 1989; 9: 4355–4370.

42. Gosselin N, Samson S, Adolphs R, et al. Emotional responses to unpleasant music correlates with damage to the parahippocampal cortex. *Brain* 2006; 129: 2585–2592.

43. Gil-Gouveia R, Oliveira AG, Martins IP. Cognitive dysfunction during migraine attacks: a study on migraine without aura. *Cephalalgia* 2015; 35: 662–674.

44. Younis S, Hougaard A, Noseda R, et al. Current understanding of thalamic structure and function in migraine. *Cephalalgia* 2019; 39: 1675–1682.

45. LaBerge D. Attention, awareness, and the triangular circuit. *Conscious Cogn* 1997; 6: 149–181.

46. Shin KJ, Lee H-J, Park KM. Alterations of individual thalamic nuclei volumes in patients with migraine. *J Headache Pain* 2019; 20: 112.

47. Bathel A, Schweizer L, Stude P, et al. Increased thalamic glutamate/glutamine levels in migraineurs. *J Headache Pain* 2018; 19: 55.

48. Amin FM, Hougaard A, Magon S, et al. Altered thalamic connectivity during spontaneous attacks of migraine without aura: A resting-state fMRI study. *Cephalalgia* 2018; 38: 1237–1244.

49. Baliki MN, Mansour AR, Baria AT, et al. Functional reorganization of the default mode network across chronic pain conditions. *PLoS One* 2014; 9: e106133.

50. Vuralli D, Ayata C, Bolay H. Cognitive dysfunction and migraine. *J Headache Pain* 2018; 19: 109.

Figure Legends

Figure 1 | Time course of individual recordings. Solid vertical lines indicate recording days. The first and the last line represent the days of migraine attacks, the colourful lines in between represent the recordings within the migraine cycle relative to the attack days. Each patient has its own colour. The number of days between the first recorded migraine attack and the subsequent migraine attack indicates the different length of each patient's migraine cycle (7 days for patient 11 and 21 days for patient 4).

Figure 2 | Time series of migraine-related resting-state maps. Two hypothetical time series of migraine-related cortical processes were modelled in the statistical analysis. In the first time course (upper part), the cortical processes drop during the headache attacks; the brain processes would be "reset" during attacks, then would resemble the processes on the day after the attacks. In the second time course (lower part), the cortical processes would reach their climax during the attacks and are similar to the days before attacks. These processes could be used as a biomarker for an impending migraine attack. **The figure is intended to illustrate the cyclic nature of migraine attacks and time-varying magnitude of two potential cortical processes; we recorded only one migraine cycle (grey area).**

Figure 3 | Statistical analysis across the migraine cycle. The Figure shows 3 exemplary intrinsic networks (visual-sensory network, thalamo-insular network, and the salience network). The dark red and blue colours represent the entire extension of the networks. The results of the change of network connectivity throughout the migraine cycle are superimposed in lighter colours. For network 5 (A), we found a main effect in the primary visual cortex in the calcarine sulcus; for network 27 (B), we found an effect predominantly in the thalamus; and for network 34 (C), we observed an effect in the anterior insular cortex and the dorsal ACC. R = right hemisphere, L = left hemisphere.

migraine attack "resets" brain activity

cortical activity peaks at migraine attack

time (days)

16d **9d** **8d** **21d** **15d** **15d** **13d** **9d** **8d** **17d** **7d** **10d**