Isolation and characterization of novel phages targeting pathogenic *Klebsiella*

pneumoniae

Na Li[1#], Yigang Zeng[2#], Rong Bao[1], Tongyu Zhu[1], Demeng Tan[2*], Bijie Hu[1*],

[1]Zhongshan Hospital, Fudan University, Shanghai, China

[2]Shanghai Public Health Clinical Center, Fudan University, Shanghai, China

[#]These authors contributed equally: Na Li, Yigang Zeng

*Correspondence: Demeng Tan (demengtan@gmail.com) and Bijie Hu (hu.bijie@zs-
hospital.sh.cn)

Abstract

Klebsiella pneumoniae is a dominant cause of community-acquired and nosocomial

infections, specifically among immunocompromised individuals. The increasing

occurrence of multidrug-resistant (MDR) isolates has significantly impacted the

effectiveness of antimicrobial agents. As antibiotic resistance is becoming prevalent, the

use of bacteriophages to treat pathogenic bacterial infections is becoming revitalized.

Elucidating the details of phage-bacteria interactions will provide insights into phage

biology and the better development of phage therapies. In this study, a total of 22 *K.*

pneumoniae isolates were assessed for their genetic and phenotypic relatedness by

multi-locus sequence typing (MLST), endonuclease S1 nuclease pulsed-field gel

electrophoresis (S1-PFGE), and *in vitro* antibiotic susceptibility testing. In addition, the

beta-lactamase gene (*bla*$_{KPC}$) was characterized to determine the spread and outbreak

of *K. pneumoniae* carbapenemase (KPC)-producing enterobacterial pathogens. Using

these *K. pneumoniae* isolates, three phages were isolated and characterized to evaluate

the application of lytic phages against these 22 ST11 *K. pneumoniae* isolates. *In vitro*

inhibition assays with three phages and *K. pneumoniae* strain ZS15 demonstrated the

strong lytic potential of the phages, however, followed by the rapid growth of phage-

resistant mutants. Together, this data adds more comprehensive knowledge to known

phage biology and further emphasizes their complexity and future challenges to

overcome prior to using phages for controlling this important MDR bacterium.

Keywords: Bacteriophage; Phage therapy; *Klebsiella pneumoniae*; Phage-host

interactions; Multidrug-resistance

Introduction

The ongoing explosion of the antimicrobial resistance crisis has been accelerated by the

overuse and misuse of antibiotics, causing the emergence and increased prevalence of

MDR bacteria (Ventola, 2015). *Klebsiella pneumoniae*, a Gram-negative

Enterobacterales, can be found in the environment and has been associated with

community-acquired infections (Pendleton et al., 2013). The increasingly common

mechanism of carbapenem resistance in *K. pneumoniae* is mediated by extended-

spectrum β-lactamases (ESBLs), which are able to break the β-lactam structure in the

most commonly used antibiotics (Pitout et al., 1997). Thus, infections of KPC-producing

bacteria are consistently associated with poor clinical outcomes and represent a

significant and continued threat to the healthcare system. As this problem continues to

grow, there is a strong need to find an alternative means to conventional antimicrobials

in order to combat this clinical pathogen.

Bacteriophages (phages for short) are the most common and diverse biological entities

in the biosphere, playing an important role in promoting bacterial diversity, structuring

community composition, and being a key driver in nutrient turnover (Suttle, 2005). Lytic

phages infect and lyse bacterial hosts and are an attractive alternative towards

combating the fast development of antibiotic-resistant *K. pneumoniae* (Sulakvelidze et

al., 2001;Hung et al., 2011). The use of phages has gained interest over the past years

and phages that are specific for MDR bacterial pathogens, including *Pseudomonas*

aeruginosa, *Enterococcus* spp., *Mycobacterium abscessus*, *Staphylococcus aureus*, *A.*

baumannii, and *Escherichia coli* have been identified and used to treat bacterial

infections (Chibani-Chennoufi et al., 2004b;Capparelli et al., 2007;McVay et al.,

2007;Letkiewicz et al., 2009;Schooley et al., 2017;Abd El-Aziz et al., 2019;Dedrick et al.,

2019;Cano et al., 2020). These studies demonstrated that lytic phages can be harnessed

to prevent and eradicate pathogenic bacteria, and in some cases, alleviate current

medical challenges posed by antibacterial resistance and thus further reduce

mortalities. While previous applications of phage therapy have shown promising results,

the technology is still in a stage of development due to rapid evolution dynamics

between phages and their hosts (Chibani-Chennoufi et al., 2004a). It is, therefore,

crucial to characterize both phage and host communities prior to any intended phage

applications.

In this study, we analyzed the molecular epidemiological feature of a collection of 22

clinical *K. pneumoniae* isolates, specifically on the characteristics of MLST, plasmid, and

antibiotic susceptibility patterns. Using these *K. pneumoniae* isolates, we isolated and

characterize three phages native to the hospital sewage. Details of phage host range,

genome, morphological characters, and phage-host interactions were established to

assess the application of phage therapies against the carbapenem-resistant isolates.

Together, our study provides important information regarding the treatment and

prevention of this important human pathogen.

Materials and methods

Bacterial strains and growth conditions

Twenty-two nonduplicate bacterial isolates of *K. pneumoniae* were collected from

Zhongshan Hospital, Shanghai, China and identified with MALDI-TOF Mass Spectrometry

(Bruker, USA) from June 2019 - May 2020. Isolates were routinely grown in either Luria-

Bertani broth (LB) containing per liter: tryptophan 10 g, NaCl 10 g, and Yeast extract 5 g

with aeration, or, on LB agar (15 g agar per liter) plates at 37°C. The beta-lactamase

gene (bla_{KPC}) was detected by PCR Master Mix (Thermo, CA, USA) with forward primer

5'-TGTAAGTTACCGCGCTGAGG-3' and reverse primer 5'-CCAGACGACGGCATAGTCAT-3'.

All strains were stored at -80°C in LB broth with 25% glycerol (vol/vol) and sub-

cultivated before each use.

Multi-locus sequence typing (MLST) and antibiotic susceptibility of bacterial isolates

In order to study population genetics and micro-evolution of this epidemic bacteria,

MLST was applied to characterize these 22 isolates using sequences of internal

fragments of seven house-keeping genes, according to the protocol available on the

MLST Pasteur website (https://bigsdb.pasteur.fr/klebsiella/klebsiella.html) (Diancourt et

al., 2005). The purified MLST amplicons were sent for Sanger sequencing, followed by

further comparison with those available on the MLST database to determine the

sequence type (ST). The VITEK2 compact automated system (bioMérieux, Lyon, France)

was used to determine antibiotic susceptibility profiles according to the manufacturer's

instructions. The susceptibility breakpoints were interpreted with reference to the latest

documents from the Clinical and Laboratory Standards Institute (CLSI).

S1-Pulsed Field Gel Electrophoresis (S1-PFGE)

Klebsiella with plasmid-mediated carbapenem resistance poses significant challenges in

clinical practice. To characterize the plasmid content in each strain, S1 nuclease

digestion analyzed with PFGE was performed using a CHEF Mapper XA apparatus (Bio-

Rad Laboratories, Hercules, CA). Briefly, the adjusted bacterial suspension mixed in a 1%

agarose gel was digested with a proteinase K solution for 2 h at 54°C, then further

treated with S1 nuclease. The electrophoresis conditions used in this study were as

follows: initial switch time, 3 s; a final switch time, 36 s; gradient, 6 V/cm; angle, 120;

temperature, 14°C; and time, 18.5 h. After the electrophoresis run was complete, the

gel was stained with 1 mg/mL ethidium bromide and visualized with the ChemiDoc XRS

system (Bio-Rad Laboratories, CA, USA). Genetic similarities were measured using the

Dice coefficient and the clustering was carried out by the unweighted-pair group

method with arithmetic averages (UPGMA) in BioNumerics (Applied Maths).

Isolation and proliferation of phage

Phages were isolated using an enrichment protocol as described previously (Tan et al.,

2014). Briefly, infiltrated hospital sewage was mixed with the same volume of 2 × LB

broth and incubated at 37°C for 6 h to stimulate proliferation of total abundances of

specific phages targeting the *K. pneumoniae* isolates above. Aliquots of enrichment

samples were centrifuged (12,000×*g*, 10 min, 4°C) and sterilized with 0.1% chloroform.

The presence of *K. pneumoniae*-phages in the hospital sewage samples was detected by

double-layer agar assays with different fold dilutions of the cell-free spent supernatant

from the enrichment assay. In order to obtain individual phages, single plaques were

picked and purified three times with their corresponding hosts. Phage stocks were

enriched and titrated using double-layer agar assays as described previously and stored

at 4°C before use (Kropinski et al., 2009).

Phage-host range assay

The host range of obtained phages was determined against 22 clinical isolates by

standard spot assays. Aliquots of 2 μL diluted phage lysate (~10^5 PFU ml^{-1}) was spotted

on lawns containing 4 mL top agar mixed with 300 μL mid-log phase bacteria. After

incubating at 37°C for 12 h, the spots were assessed by the turbidity of plaques and

characterized as strong infection (clear), weak infection (turbid), or no inhibition (no

plaque). The experiment was undertaken as independent duplicates.

Susceptibility of strain ZS15 to phage infection and isolation of phage-resistant

mutants

Overnight bacterial cultures were 1,000-fold diluted into LB broth and grown to an

exponential phase with an OD600 at 0.3 and infected with its corresponding phages at

an multiplicity Of infection (MOI) of 0.01. The suspension was incubated at 37°C with

aeration and samples were harvested at 1-h intervals. The phage lysis efficiency was

determined by optical density measurements. The experiment was done in triplicate

along with parallel control cultures without phages. At time point 8-h, phage lysates

were streaked onto LB agar plates and individual colonies were subsequentially purified

to determine phage susceptibility by spot test assays as described above.

Phage DNA extraction and genomic analysis

For the extraction of phage DNA, 2 mL aliquots of phage lysate with a titer of

approximately 10^9 PFU mL^{-1} was digested with DNase I (Thermo, CA, USA) and RNase A

(Promega, USA) to remove any residual bacterial genomic DNA and RNA according to

the manufacturer's protocol. After digesting for 1.5 h at 37°C, EDTA (Thermo, CA, USA)

was added to a final concentration of 20 mM to inactivate DNase I and RNase A. Total

phage DNA was extracted using the Qiagen DNeasy Blood and Tissue kit (Qiagen, CA,

USA) following the manufacturer's instructions. Genomic DNA was sent to Sangon

Biotech (Shanghai, China) and sequenced using an Illumina HiSeq sequencer.

The raw sequence containing the phage genome was quality controlled and trimmed

with FastQC 0.11.2 (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and

Trimmomatic 0.36 (http://www.usadellab.org/cms/?page=trimmomatic). The trimmed

reads were assembled into a single raw contig via SPAdes 3.5.0.

(http://cab.spbu.ru/software/spades/). Genes were predicted and annotated using

Rapid Annotation using Subsystem Technology (RAST, http://rast.nmpdr.org/). Phage

virulence factor was examined using Virulence Factor Database (VFDB,

http://www.mgc.ac.cn/VFs/). Phylogenetic relationship between the genome of phage

NL_ZS_1, NL_ZS_2, and NL_ZS_3 and the genomes of related phages were analyzed

using CLC Main Workbench (CLC, Qiagen). To construct the phylogenetic tree of phage

terminase large subunits, the reference amino acid sequences of were collected from

the NCBI database. The terminase large subunits were aligned, and neighbor-joining

tree were created using Jukes-Cantor model with a bootstrap analysis of 1000

replicates.

Bacteriophage morphology by TEM

Based on the genomic differences, phages were selected for examination by

transmission electron microscopy (TEM). Phage stocks were back-diluted in a SM buffer

(50 mM Tris-Cl, pH 7.5, 99 mM NaCl, 8 mM MgSO$_4$, 0.01% gelatin) to a titer of 10^8

PFU/mL. Diluted phage lysates were then placed on the surface of Formvar carbon-

coated grids (Sigma-Aldrich, St. Louis, MO, USA) for 5 min to allow adsorption. Grids

were stained with 2% sodium phosphotungstate for 2 min prior to being washed in a

drop of distilled water and air-dried before being examined using a JEM-2100

microscope operated at 80 kV.

Results

Isolation and characterization of *K. pneumoniae* strains

The emergence of carbapenem-resistant *K. pneumoniae* (CRKP) has become one major

burden of healthcare-associated infection due to the dispersion of bacteria carrying the

bla$_{KPC}$ gene (Bush and Bradford, 2016). KPC was first found in carbapenem-resistant *K.*

pneumoniae strains in 2001 in North America, and now KPC-producing strains can be

frequently isolated from nosocomial samples. It is, therefore, essential to determine and

characterize the nosocomial spread of KPC clones and their corresponding phenotype

and genotype. Together, 22 *K. pneumoniae* strains were recovered from various clinical

specimens. According to the *K. pneumoniae* MLST database, all of the 22 clinical isolates

belonged to ST11 (allelic profile 3311114), which is the predominant clone of KPC-

producing *K. pneumoniae* in China (Table 1). Despite more than 100 different sequence

types (STs) being reported worldwide, the mass dissemination of carbapenemase-

producing *K. pneumoniae* has been restricted largely to ST258, ST11, ST340, and ST512

(Pitout et al., 2015).

In addition, the bla_{KPC} gene was detected in 19 out of 22 isolates, indicating the

prevalence of β-lactamase nosocomial pathogens (Table 1). The bla_{KPC} gene can be

disseminated between species by horizontal (lateral) gene transfer (HGT) via various

means, such as plasmids, transposons, prophages, and integrative conjugative elements

(ICEs), which play an important effect in generating genotypic heterogeneity and

phenotypic variation among bacterial populations (Hardiman et al., 2016;Botelho and

Schulenburg, 2020). For instance, transfer of interspecies KPC can occur through the

dissemination of mobile genetic elements between strains of *S. aureus* (Goren et al.,

2010).

Since pathogenicity and antimicrobial resistance genes are mainly coded by plasmids, in

order to identify the subtypes of *K. pneumoniae* ST11, pulsed-field gel electrophoresis

with S1 nuclease (S1-PFGE) were utilized to resolve the prevalence of plasmid

fingerprints of all the 22 isolates, particularly large ones, which clustered into 6 primary

clades (A, B, C, D, E, and F) (Figure 1). Clade A and B were the most common types,

while the other clades (C, D, E, and F) accounted for 31.8%. All groups harbored one to

two plasmids ranging from 20 Kbp and 250 Kbp. Thus, it appears that S1-PFGE is a highly

discriminative molecular typing method capable of effectively discriminating ST11

clones into several different subtypes, indicating the prevalence of ST11 was not clonally

homogenous.

Imipenem and meropenem resistance involve the production of β-lactamase

Individual antimicrobial data can be found in Table 1. For ticarcillin/clavulanic acid,

piperacillin/tazobactam, cefepime, aztreonam, ciprofloxacin, and levofloxacin, drug

resistance was detected in all strains. In addition, low percentages of resistance to

doxycycline, minocycline, tigecycline, colistin, and trimethoprim/sulfamethoxazole were

also observed for most strains. Of these, 17 out of the 22 isolates become resistant to

both imipenem and meropenem with a MIC of ≥16 μg/mL, while the other 5 were both

sensitive to imipenem and meropenem, except for strain ZS18 which was resistant to

imipenem but intermediate to meropenem. For these 5 sensitive isolates, 2 of them

were confirmed by PCR and sequencing to have the bla_{KPC} gene. Interestingly, strain ZS4

harboring bla_{KPC} was susceptible to imipenem and meropenem, perhaps indicating the

resistance is associated with changes in gene expression at the mRNA levels (Senchyna

et al., 2019).

Isolation and characterization of phages

Three phages (NL_ZS_1, NL_ZS_2, and NL_ZS_3) were isolated from local hospital

sewage using the enrichment protocol to evaluate the potential of phage therapy as an

effective alternative against *K. pneumoniae* strains. The host range of the three phages

and their lytic efficiency against all of the 22 bacterial isolates were assessed using the

spot test assay. According to the host range profile, the phages were able to infect all of

the 22 bacterial isolates, however, the phages displayed a unique profile in terms of

host range patterns and lytic efficiency (Figure 2). Based on their host susceptibilities,

phages were divided into two major groups. Specifically, phage NL_ZS_3 displayed the

broadest host range, while phage NL_ZS_1 and phage NL_ZS_2 showed a relatively

narrow host range, infecting only ~50% of the tested *K. pneumoniae* strains (Figure 2).

The host range differences between the phages observed in this study are very likely

due to the different phage characterization, i.e., efficiency of adsorption and lytic

efficiency in its potential hosts.

Morphology and genome sequence of phages

All phage lysates formed transparent plaques on strain ZS15, which are considered as an

indicator of strong lysis. Plaque morphologies of all phages presented a zone of clear

lysis on the host strain ZS15, while a halo zone was detected on the plaques of both

phages NL_ZS_1 and NL_ZS_2, indicating the production of polysaccharide

depolymerase (Figure 3). Depolymerase activity are usually encoded by their tail fibers

or tail spikes protein on the baseplate, which has the potential to provide an alternative

treatment to conventional antibiotics.

Based on transmission electron microscopy (TEM) analysis, all the phages belong to the

family *Podoviridae* of *Caudovirales* order based on the International Committee on

Taxonomy of Viruses (ICTV) definitions, having short, non-contractile tails (~11 nm in

length) and icosahedral heads (~61 nm diameter) (Figure 3 and Table 2). In addition,

total genomic DNA was extracted from each of the three phages, and their genomes

were sequenced by Illumina MiSeq. The assembly sizes of these three phages ranged

from 38 kb to 40 kb, having a G+C content ranging between 50% to 53%, with 45 to 49

predicted protein-coding genes (Table 2). All phage genomes (GenBank accession: phage

NL_ZS_1, MT813140; phage NL_ZS_2, MT813141; and phage NL_ZS_3, MT813142) were

submitted to RAST (Rapid Annotation using Subsystem Technology) and further

annotated by the available pipeline. No identifiable virulent genes, antibiotic resistance

genes, or genes involved in lysogenic cycles were detected. Therefore, it appears that

those phages isolated in this study can satisfy the safety assessment based on their

complete genome sequences. Progressive multiple genome alignments were analyzed

using Easyfig software (https://mjsull.github.io/Easyfig/) (Figure 4) to determine the

relatedness of three *K. pneumoniae* phages. A considerable relation was shown

between phage NL_ZS_1 and phage NL_ZS_2, while phage NL_ZS_2 displays a relatively

low homologous region with phage NL_ZS_3. Likely, phylogenetic analysis of the large

subunit of the terminase of phage NL_ZS_1, NL_ZS_2, and NL_ZS_3 reveals their close

relation to large terminase subunits of other phages, specifically, phage NL_ZS_1 and

phage NL_ZS_2 display high similarities to *Klebsiella* phage K11, BIS33, and KpV289,

while phage NL_ZS_3 was closely related to *Salmonella* phage phiSG-JL2 (Figure 5).

Together, these results showed high relation to the phage-host range pattern,

demonstrating phage susceptibility correlates with phage core genome, such as genes

involved in the phage tail fibers. However, further studies are needed to probe the

genetic determinants that contribute to the host-range difference among these three

phage isolates.

Phage growth characteristics in LB broth

To examine the efficiency of phage infection, *K. pneumoniae* ZS15 was infection in liquid

cultures with individual phages (NL_ZS_1, NL_ZS_2, and NL_ZS_3) at an MOI of ~0.01,

respectively, and their inhibition was assessed by bacterial growth following the

measurement of the optical density of the bacterial cultures with and without phages. A

similar inhibitory effect of phage NL_ZS_1 and phage NL_ZS_2 was observed in the *in*

vitro inhibition assay with a drastic decline in optical density on the initial control of the

host population for 1.5 h followed by regrowth of the resistant population (Figure 6). A

similar trend in the phage-amended cultures of ZS15+ NL_ZS_3 was also observed in the

first 5 h, however, it was followed by less pronounced decrease and finally stabilized

after 6 h (Figure 6), After 8 h, the OD600 nm of the phage-amended cultures of group 1

(ZS15+phage NL_ZS_1 and ZS15+phage NL_ZS_2) and group 2 (ZS15+phage NL_ZS_3)

stabilized at 1.96 and 0.09, corresponding to 52% to 96% of the final OD in the control

group, respectively (Figure 6). Together, these results demonstrate that phage NL_ZS_3

was slightly more effective in reducing bacterial densities than the rest of the phages.

Despite *K. pneumoniae* ZS15 displaying a strong susceptibility to infection by all three

phages, phage NL_ZS_3 was better at suppressing resistance to host strain ZS15 over

phage NL_ZS_1 and NL_ZS_2. Bacteria are known to rapidly develop phage resistance

when subjected to predation by lytic phages in a laboratory medium (Labrie et al.,

2010). By screening phage resistance mutants, we attempted to better understand the

inhibition differences among each of the phage-amended cultures. Phage insensitive

mutants were isolated from phage lysate cultures during the incubation for the analysis

of changes in their susceptibility upon phage selection. Based on the results of spot test

assays, we observed that phage-amended cultures (ZS15+phage NL_ZS_1, ZS15+phage

NL_ZS_2, and ZS15+phage NL_ZS_3) were dominated by resistant non-mucoid clones

after 8 h of incubation, indicating conversion of mucoid *K. pneumoniae* strain ZS15 to

non-mucoid variants by consequence of selective pressure operated by phages (Table

3). However, in the ZS15+phage NL_ZS_3 cultures, heterogeneity in phage susceptibility

was observed (Table 3). Out of 30 *K. pneumoniae* ZS15 isolates, 2 (~6.7%) were fully

sensitive to phage NL_ZS_3 and displayed the hypermucoviscosity (HV) phenotype as

the parental strain ZS15, and partially sensitive (i.e., reduced phage susceptibly when

compared to the parental strain) subpopulations (~50%) were isolated as well and did

not revert back to the wild-type phenotype despite three rounds of purification.

Interestingly, 43.3% were fully resistant to phage NL_ZS_3, however, a small fraction

(~30%) of phage NL_ZS_3 resistant mutants reverted back to the parental phenotype

after three rounds of purification, suggesting different anti-phage mechanisms were

allowing coexistence of phage sensitive mutants and that their lytic phages could exist

between bacterial host ZS15 and phage NL_ZS_3. Since phage susceptibility was only

measured at 8 h, we do not know the dynamics of phenotype heterogeneity over time

or how long such persistence can last before heritage mutations dominate the entire

population.

Anti-phage defense-mediated altered sensitivity to antibiotics

Previous studies have shown that evolutionary phage-host interactions affect phage

resistance and antibiotic susceptibility both positively and negatively, and are highly

genotype-dependent in bacterial populations (Comeau et al., 2007;Chan et al.,

2016;Burmeister et al., 2020). We started to question whether phage-resistant mutants

changed their antibiotic susceptibilities via antagonistic coevolution between hosts and

parasites in LB broth. To investigate this further, we performed Vitek 2 assays for

antimicrobial susceptibility, testing phage-resistant mutants (ZS15-3R) and their

parental phage susceptible strains (ZS15). Interestingly, for tigecycline susceptibility

testing of carbapenem-resistant *K. pneumoniae* ZS15 strain, we noticed that strain ZS15

evolved resistance to phage NL_ZS_3, while simultaneously becoming more sensitive to

tigecycline with the minimum inhibitory concentration shifting from 8 µg/mL to 2

µg/mL. Unfortunately, we were unable to identify potential genes related to phage

selection and antibiotic sensitivity during the interactions between strain ZS15 and

phage NL_ZS_3. Further, comparative genomic sequence analyses are needed to

elucidate the underlying mechanisms.

Discussion

Klebsiella pneumoniae is a common nosocomial pathogen that causes difficult-to-treat

infections worldwide. The prevalence and distribution of bla_{KPC} gene in *K. pneumoniae* is

increasing, making carbapenem resistance isolates difficult to treat due to the lack of

effective and safe alternative options. Successful application of phage therapy in

treating *K. pneumoniae* caused infections requires a detailed understanding of the

phenotypic and genotypic diversity printers of *K. pneumoniae* pathogens as well as well-

characterized phages that covers host diversities. In this study, we characterized 3

phages and 22 *K. pneumoniae* strains, covering considerable temporal variability.

Further, *K. pneumoniae* strain ZS15 was selected for more detailed studies of phage lytic

potential and phage-host interactions.

Diversity of *K. pneumoniae* ST11 isolates

The clonal relationship between *K. pneumoniae* strains examined in the present study

belonged to ST11 irrespective of time or place of isolation. ST11 is among the most

important clinical dominant in Asia and South America and has been reported in

different regions worldwide since its emergence (Qi et al., 2011). In 2018, a fatal

outbreak of ST11 carbapenem-resistant hypervirulent *K. pneumoniae* in a Chinese

hospital has caused high mortality and morbidity (Gu et al., 2018;Yao et al., 2018).

Likely, ST258, single-locus variants of ST11 (the *tonB* allele distinguishes the two

sequence types), has emerged as an important cause of hospital death in the United

States and other countries, and is largely responsible for the global spread of KPC (Toth

et al., 2010). However, the molecular linage between ST11 and ST258 largely remains

unknown, recent comparative genomic analyses showed ST258 is a hybrid with ~80% of

the chromosome homology to ST11 strains, while the remaining 20% display high

similarity to the ST442 isolates (Chen et al., 2014). Despite MLST does not provide

sufficient discrimination for the 22 pathogenic isolates or resolve differences across

spatial and temporal sac ales, understanding the molecular evolutionary history of

bacterial isolates is an important step towards the development of diagnostic and

potential therapeutic strategies to combat infections caused by multidrug-resistant *K.*

pneumoniae.

Carbapenem resistance was seen among all *Klebsiella* isolates. Since carbapenem-

resistant *K. pneumoniae* (CRKp) are endemic to local hospitals, where colistin has been

largely used as the last treatment option to fight against these infections. Out of 22

strains, 2 isolates (strain ZS2 and ZS16) display resistance to colistin. Together, there 22

isolates could be regarded as a real superbug that could pose a major public health

challenge. There is currently little understanding of molecular mechanisms underlying

antibiotic resistance and its impact on virulence remain unclear, despite ~86% of strains

were confirmed to be carbapenemase gene positive. A recent sequence of complete *K.*

pneumoniae genomes showed that genes coding for resistance to antibiotics or

virulence factors is located in the plasmid. Moreover, the spread of these resistant

plasmids into the normal gut flora poses an even great public health threat. In order to

recapitulate the contents and diversity of the plasmids of genetically related *K.*

pneumoniae strains, it is important to track the spread of its plasmids between each

isolate. In our study, S1-PFGE showed four clusters of the CRKp producing strains that

corresponded to ST11, which demonstrates a high degree of genetic diversity. Future

work will seek to understand the dissemination of these plasmids and their potentials of

converting any *K. pneumoniae* strains with no existing antimicrobial resistance or

virulence determinants into CRKp strain immediately. Together, these results provide

important insights into relatively narrow phylogenetic and phenotypic groups of

bacteria, emphasizing the complexity for the development of alternatives to treat this

pathogen.

Phage isolation and diversity

Generally, phages are widespread and genetically diverse in almost all environments,

especially in sewage, known as a fertility source for many pathogens. The high

occurrence of *K. pneumoniae* phages in the collected samples (~72%) suggested the *K.*

pneumoniae phages are widespread in the local hospital sewage supply. The phages

isolated from the local hospital sewage displayed two distinguished host ranges when

tested against the collection of 22 *K. pneumoniae* isolates. Only 1 morphological family,

Podoviridae, was represented, and the phages ranged in genome size from 38 kb to 40

kb. Due to the limited sample numbers, we do not know to what extent these isolated

phages represent the true diversity and structure of local hospital sewage phage

communities. Further studies of viral metagenomics that are not dependent on cell

culture approaches are needed to expand our understanding of phage genomic diversity

from the evolutionary perspective and perhaps genetic exchanges in response to the

selection pressure from the bacterial host.

Phage NL_ZS_3 displayed a relatively broad host range, while phage NL_ZS_1 and phage

NL_ZS_2 shared identical relatively narrow host ranges, suggesting that phage NL_ZS_1

and phage NL_ZS_2 might use similar phage binding receptors to recognize bacterial

phage receptors in their susceptible hosts. This is interesting as bioinformatic analysis

showed that phage-binding receptors of phage NL_ZS_1 and phage NL_ZS_2 were

identical based on the amino acid sequence. The possible explanation for host range

difference could be phage NL_ZS_3 used a conservative apparatus as its phage

receptors, such as outer membrane proteins, lipopolysaccharides, flagella, or pili.

Moreover, several copies of genes encoding phage tails or tail fibers were present in the

phage NL_ZS_3, which opens the possibility that phage NL_ZS_3 may use a primary

receptor to initiate attachment but require a secondary cell surface receptor to

irreversibly bind to the bacterial host for injection, permitting more flexibility to develop

on a wider range of hosts. For example, the T4 phage uses its long tail fibers to

recognize lipopolysaccharide and the outer membrane protein C, present on the

bacterial surface (Tétart et al., 1996;Baslé et al., 2006). This reversible attachment

allows the phage to move along the surface and find a suitable site to inject its DNA.

Phage-host interactions in strain ZS15

Isolation and characterization of bacterial isolates during the LB broth experiments

confirmed the rapid development of phage resistance and subsequently dominated the

bacterial populations in the phage NL_ZS_1 and phage NL_ZS_2, which is regarded as a

significant barrier to the successful application of phage therapies (Koskella and

Meaden, 2013). Interestingly, in the current experiment with strain ZS15, phage NL_ZS_

3 exposures led to the rapid development of phenotypic diversity (i.e., resistant,

partially sensitive, and resistant) within a population that enables survival of fluctuating,

suggesting different anti-phage mechanisms were allowing coexistence of phage

sensitive mutants and that their lytic phages could exist. Since phage susceptibility was

only measured at 8 h, we do not know the dynamics of phenotype heterogeneity over

time or how long such persistence can last before heritage mutations dominate the

entire population. It is, therefore, crucial to characterize the phage-host interaction

regarding the phenotypic and genetic diversification in bacteria and phages prior to any

intended phage applications. The exact resistant mechanism by which strain ZS15 avoids

phages infection, however, remains to be discovered.

A common form of developing phage resistance for bacterial hosts is associated with

loss or altered structure of the receptors to block the potential phage adsorption in

order to avoid infection (Labrie et al., 2010). Such changes are directly linked to the key

step in recognition between phage receptor-binding proteins (RBP) located on tail

fibers, and phage receptors encoded on the susceptible bacterial host cell surface

(Bertozzi Silva et al., 2016). In general, the fitness costs resulting from receptor mutation

(i.e., motility, nutrient transport, or antimicrobial susceptibility) make the mutants

unable to compete with phage-susceptible strains in both environmental samples and

infection models, even in the presence of phage predation (Seed et al., 2012;Sumrall et

al., 2019). Understanding the evolutionary implication of phage-receptor interactions

will be beneficial for predicting how phages interact with their host, as phage receptor

mutation-mediated phenotypic variabilities may result in exopolysaccharide associated

fitness cost. Such characterization of phage resistance evolution will be used as a guide

for selecting proper phage targeting of specific receptors with a significant clinical

application. Our data reveal when phage resistance evolves, the resistant mutants

showed increased tigecycline sensitivity, despite the mechanism underlying potential is

not fully understood in this study.

Tigecycline, known as a tetracycline antibiotic, has shown the greatest antimicrobial

activity against CRKP. Genes e.g. *tet*(B), *tet*(A), *tet*(K), *tet*(M) and *tet*(S) coding for

multidrug efflux pumps are membrane-related proteins and can efflux tetracyclines

from the bacteria intracellularly, thus decreasing the MIC and subsequently allowing

protein synthesis without affecting cell viability (Linkevicius et al., 2016). Using

membrane-related proteins as a surface structure is one of the most common

components functioning as a phage receptor. Therefore, it makes sense that phage-

resistant defense strategies would confer a fitness cost for bacteria, i.e., becoming more

sensitive to certain antibiotics. For instance, in *P. aeruginosa*, phage OMKO1 evolved to

become less resistant to antibiotics. Likewise, in *E. coli*, the TolC protein evolved in a

variety of diverse cellular functions, including antibiotic efflux and phage TLS receptor

(Chan et al., 2016). However, unlike previous studies, the anti-phage TLS mediated TolC

mutation did not influence its role in antibiotic efflux. Overall, these results reveal that a

phage-host arm race can result in different outcomes in terms of antibiotic

susceptibilities, highlighting the diversity and complexity of host interactions within

bacterial populations.

Potential applications of phage therapy

Successful application of phage therapy requires collaboration between microbiology

and preclinical development; however, little is known about their alignment to guide

future development. Since the discovery of phages, many phages have been isolated

and identified as potential alternatives to antibiotics with different efficacy profiles.

Phage specificity is both a benefit and an obstacle for clinical trials, as phage receptor

diversity affects phage-host interaction. When the bacterial population is under phage

attack, only a small fraction of cells can survive these selective pressures, and those that

survive determine the fate of the population. Therefore, elucidating details of phage-

host interactions and how these would influence bacterial physiological heterogeneity is

fundamental to the development of phage therapies. In this study, the three highlighted

phages showed promising *in vitro* effects that perhaps these phages can be tamed for

our benefits without negatively affecting their natural balance. The main limitation of

this study is that only a small number of bacterial isolates and phages have been

investigated. A large-scale study on the interactions between carbapenem-resistant K

pneumoniae strains and phages covering a considerable spatial and temporal variability

are essential for the successful application of phage-based pathogen control to fill our

knowledge gap between *in vitro* and *in vivo* models.

Funding

This project was supported by the National Natural Science Foundation of China (NSFC)

(No. 42006136), Shanghai Municipal Commission of Health (No.20204Y0336), and

Shanghai Pujiang Talents Plan Project (No. 2020PJ054) to DT.

Acknowledgments

We thank Xiao Wang and Ruoyu Li for technical assistance. We thank all members from

Zhu's lab for thoughtful discussions and grateful to Brett C. Gonzalez for proofreading

the manuscript.

Declaration of competing interest

The authors declare no competing financial interests.

References

Abd El-Aziz, A.M., Elgaml, A., and Ali, Y.M. (2019). Bacteriophage Therapy Increases
Complement-Mediated Lysis of Bacteria and Enhances Bacterial Clearance After Acute
Lung Infection With Multidrug-Resistant Pseudomonas aeruginosa. *J Infect Dis* 219**,**
1439-1447.
Baslé, A., Rummel, G., Storici, P., Rosenbusch, J.P., and Schirmer, T.J.J.O.M.B. (2006). Crystal
structure of osmoporin OmpC from E. coli at 2.0 Å. 362**,** 933-942.
Bertozzi Silva, J., Storms, Z., and Sauvageau, D. (2016). Host receptors for bacteriophage
adsorption. *FEMS Microbiol Lett* 363.
Botelho, J., and Schulenburg, H. (2020). The Role of Integrative and Conjugative Elements in
Antibiotic Resistance Evolution. *Trends Microbiol*.
Burmeister, A.R., Fortier, A., Roush, C., Lessing, A.J., Bender, R.G., Barahman, R., Grant, R., Chan,
B.K., and Turner, P.E.J.P.O.T.N.a.O.S. (2020). Pleiotropy complicates a trade-off between
phage resistance and antibiotic resistance. 117**,** 11207-11216.
Bush, K., and Bradford, P.a.J.C.S.H.P.I.M. (2016). β-Lactams and β-lactamase inhibitors: an
overview. 6**,** a025247.
Cano, E.J., Caflisch, K.M., Bollyky, P.L., Van Belleghem, J.D., Patel, R., Fackler, J., Brownstein,
M.J., Horne, B.A., Biswas, B., and Henry, M.J.C.I.D. (2020). Phage Therapy for Limb-
threatening Prosthetic Knee Klebsiella pneumoniae Infection: Case Report and In Vitro
Characterization of Anti-biofilm Activity.
Capparelli, R., Parlato, M., Borriello, G., Salvatore, P., and Iannelli, D. (2007). Experimental phage
therapy against Staphylococcus aureus in mice. *Antimicrob Agents Chemother* 51**,** 2765-
2773.
Chan, B.K., Sistrom, M., Wertz, J.E., Kortright, K.E., Narayan, D., and Turner, P.E. (2016). Phage
selection restores antibiotic sensitivity in MDR Pseudomonas aeruginosa. *Sci Rep* 6**,**
26717.
Chen, L., Mathema, B., Pitout, J.D., Deleo, F.R., and Kreiswirth, B.N.J.M. (2014). Epidemic
Klebsiella pneumoniae ST258 is a hybrid strain. 5**,** e01355-01314.
Chibani-Chennoufi, S., Bruttin, A., Dillmann, M.L., and Brussow, H. (2004a). Phage-host
interaction: an ecological perspective. *J Bacteriol* 186**,** 3677-3686.
Chibani-Chennoufi, S., Sidoti, J., Bruttin, A., Kutter, E., Sarker, S., and Brussow, H. (2004b). In
vitro and in vivo bacteriolytic activities of Escherichia coli phages: implications for phage
therapy. *Antimicrob Agents Chemother* 48**,** 2558-2569.
Comeau, A.M., Tétart, F., Trojet, S.N., Prere, M.-F., and Krisch, H.J.P.O. (2007). Phage-antibiotic
synergy (PAS): β-lactam and quinolone antibiotics stimulate virulent phage growth. 2**,**
e799.
Dedrick, R.M., Guerrero-Bustamante, C.A., Garlena, R.A., Russell, D.A., Ford, K., Harris, K.,
Gilmour, K.C., Soothill, J., Jacobs-Sera, D., and Schooley, R.T. (2019). Engineered
bacteriophages for treatment of a patient with a disseminated drug-resistant
Mycobacterium abscessus. *Nature medicine* 25**,** 730.
Diancourt, L., Passet, V., Verhoef, J., Grimont, P.A., and Brisse, S. (2005). Multilocus sequence
typing of Klebsiella pneumoniae nosocomial isolates. *J Clin Microbiol* 43**,** 4178-4182.
Goren, M.G., Carmeli, Y., Schwaber, M.J., Chmelnitsky, I., Schechner, V., and Navon-Venezia, S.
(2010). Transfer of carbapenem-resistant plasmid from Klebsiella pneumoniae ST258 to
Escherichia coli in patient. *Emerg Infect Dis* 16**,** 1014-1017.
Gu, D., Dong, N., Zheng, Z., Lin, D., Huang, M., Wang, L., Chan, E.W.-C., Shu, L., Yu, J., and Zhang,
R.J.T.L.I.D. (2018). A fatal outbreak of ST11 carbapenem-resistant hypervirulent
Klebsiella pneumoniae in a Chinese hospital: a molecular epidemiological study. 18**,** 37-
46.

Hardiman, C.A., Weingarten, R.A., Conlan, S., Khil, P., Dekker, J.P., Mathers, A.J., Sheppard, A.E.,
Segre, J.A., and Frank, K.M. (2016). Horizontal Transfer of Carbapenemase-Encoding
Plasmids and Comparison with Hospital Epidemiology Data. *Antimicrob Agents*
Chemother 60**,** 4910-4919.
Hung, C.H., Kuo, C.F., Wang, C.H., Wu, C.M., and Tsao, N. (2011). Experimental phage therapy in
treating Klebsiella pneumoniae-mediated liver abscesses and bacteremia in mice.
Antimicrob Agents Chemother 55**,** 1358-1365.
Koskella, B., and Meaden, S. (2013). Understanding bacteriophage specificity in natural microbial
communities. *Viruses* 5**,** 806-823.
Kropinski, A.M., Mazzocco, A., Waddell, T.E., Lingohr, E., and Johnson, R.P. (2009). "Enumeration
of Bacteriophages by Double Agar Overlay Plaque Assay," in *Bacteriophages: Methods*
and Protocols, Volume 1: Isolation, Characterization, and Interactions, eds. M.R.J. Clokie
& A.M. Kropinski. (Totowa, NJ: Humana Press), 69-76.
Labrie, S.J., Samson, J.E., and Moineau, S. (2010). Bacteriophage resistance mechanisms. *Nat*
Rev Microbiol 8**,** 317-327.
Letkiewicz, S., Międzybrodzki, R., Fortuna, W., Weber-Dąbrowska, B., and Górski, A.J.F.M.
(2009). Eradication of Enterococcus faecalis by phage therapy in chronic bacterial
prostatitis—case report. 54**,** 457.
Linkevicius, M., Sandegren, L., and Andersson, D.I. (2016). Potential of Tetracycline Resistance
Proteins To Evolve Tigecycline Resistance. *Antimicrob Agents Chemother* 60**,** 789-796.
Mcvay, C.S., Velasquez, M., and Fralick, J.A. (2007). Phage therapy of Pseudomonas aeruginosa
infection in a mouse burn wound model. *Antimicrob Agents Chemother* 51**,** 1934-1938.
Pendleton, J.N., Gorman, S.P., and Gilmore, B.F. (2013). Clinical relevance of the ESKAPE
pathogens. *Expert Rev Anti Infect Ther* 11**,** 297-308.
Pitout, J.D., Nordmann, P., and Poirel, L. (2015). Carbapenemase-Producing Klebsiella
pneumoniae, a Key Pathogen Set for Global Nosocomial Dominance. *Antimicrob Agents*
Chemother 59**,** 5873-5884.
Pitout, J.D., Sanders, C.C., and Sanders Jr, W.E.J.T.a.J.O.M. (1997). Antimicrobial resistance with
focus on β-lactam resistance in gram-negative bacilli. 103**,** 51-59.
Qi, Y., Wei, Z., Ji, S., Du, X., Shen, P., and Yu, Y.J.J.O.a.C. (2011). ST11, the dominant clone of KPC-
producing Klebsiella pneumoniae in China. 66**,** 307-312.
Schooley, R.T., Biswas, B., Gill, J.J., Hernandez-Morales, A., Lancaster, J., Lessor, L., Barr, J.J.,
Reed, S.L., Rohwer, F., Benler, S.J.a.A., and Chemotherapy (2017). Development and use
of personalized bacteriophage-based therapeutic cocktails to treat a patient with a
disseminated resistant Acinetobacter baumannii infection. 61.
Seed, K.D., Faruque, S.M., Mekalanos, J.J., Calderwood, S.B., Qadri, F., and Camilli, A. (2012).
Phase variable O antigen biosynthetic genes control expression of the major protective
antigen and bacteriophage receptor in Vibrio cholerae O1. *PLoS Pathog* 8**,** e1002917.
Senchyna, F., Gaur, R.L., Sandlund, J., Truong, C., Tremintin, G., Kültz, D., Gomez, C.A.,
Tamburini, F.B., Andermann, T., Bhatt, A.J.D.M., and Disease, I. (2019). Diversity of
resistance mechanisms in carbapenem-resistant Enterobacteriaceae at a health care
system in Northern California, from 2013 to 2016. 93**,** 250-257.
Sulakvelidze, A., Alavidze, Z., and Morris, J.G., Jr. (2001). Bacteriophage therapy. *Antimicrob*
Agents Chemother 45**,** 649-659.
Sumrall, E.T., Shen, Y., Keller, A.P., Rismondo, J., Pavlou, M., Eugster, M.R., Boulos, S., Disson, O.,
Thouvenot, P., and Kilcher, S.J.P.P. (2019). Phage resistance at the cost of virulence:
Listeria monocytogenes serovar 4b requires galactosylated teichoic acids for InlB-
mediated invasion. 15**,** e1008032.

Suttle, C.A. (2005). Viruses in the sea. *Nature* 437**,** 356-361.

Tan, D., Gram, L., Middelboe, M.J.A., and Microbiology, E. (2014). Vibriophages and their

589   interactions with the fish pathogen Vibrio anguillarum. 80**,** 3128-3140.

590   Tétart, F., Repoila, F., Monod, C., and Krisch, H. (1996). "Bacteriophage T4 host range is

591   expanded by duplications of a small domain of the tail fiber adhesin". Elsevier).

592   Toth, A., Damjanova, I., Puskas, E., Jánvári, L., Farkas, M., Dobák, A., Böröcz, K., Paszti,

593   J.J.E.J.O.C.M., and Diseases, I. (2010). Emergence of a colistin-resistant KPC-2-producing

594   Klebsiella pneumoniae ST258 clone in Hungary. 29**,** 765-769.

595   Ventola, C.L. (2015). The antibiotic resistance crisis: part 1: causes and threats. *P T* 40**,** 277-283.

596   Yao, H., Qin, S., Chen, S., Shen, J., and Du, X.-D.J.T.L.I.D. (2018). Emergence of carbapenem-

597   resistant hypervirulent Klebsiella pneumoniae. 18**,** 25.

Fig. 1. Dendrogram generated by BioNumerics software (Version7.1, Applied Maths, St-Martens-Latem, Belgium) showing cluster results of molecular epidemiology S1-PFGE analysis of 21 *K. pneumoniae* isolates. Band-based similarity coefficients were calculated using Dice coefficient and unweighted pair group average method algorithm (UPGMA).

Fig. 2. Phage-host range assay of the 3 phages (columns) against the 22 *K. pneumoniae* isolates (rows). *Klebsiella pneumoniae* phages displayed broad and narrow host ranges. Black, grey and white indicate infection, weak infection and no plaque formation, respectively.

Fig. 3. Plaque images (up) and their corresponding TEM micrographs (below). Phages were stained with 2% sodium phosphotungstate and were found to have short small podophage morphology. Scale bar, 200 nm.

Fig. 4. The comparison of the whole genome sequences among three *K. pneumoniae* phages (NL_ZS_1, NL_ZS_2, and NL_ZS_3) with Easyfig using the BLASTN algorithm. Grey lines connect regions indicates nucleotide identity >66%, and are gradient-colored according to their similarity; darker shade of grey represents higher identify. The designations of the open reading frames of NL_ZS_1 are shown on the top.

Fig. 5. Phylogenetic tree was constructed based on the amino acid sequence of the larger terminase subunit of phages NL_ZS_1, NL_ZS_2, and NL_ZS_3 and phages sharing homology sequence identity retrieved from GenBank (https://www.ncbi.nlm.nih.gov/). The evolutionary history of 20 larger terminase subunit amino acid sequences were aligned and inferred using Neighbor Joining method and 1000 bootstrap replicates (CLC Genomics Workbench 8).

Fig. 6. Inhibition assay of phage NL_ZS_1, NL_ZS_2, and NL_ZS_3 in the treatment at an MOI of 0.1 and control without phage in cultures of *K. pneumoniae* ZS15. Optical density (OD600) was

measured at 1-h intervals over an 8-h incubation period. Error bars represent the standard

deviations from all experiments, carried out in triplicate.

Dice (Opt:1.50%) (Tol 1.5%-1.5%) (H>0.0% S>0.0%) [0.0%-100.0%]

PFGE-Xbal-6~36S

PFGE-Xbal-6~36S

- A
  - ZS5
  - ZS7
  - ZS13
  - ZS15
  - ZS16
  - ZS17
  - ZS21
- B
  - ZS1
  - ZS2
  - ZS4
  - ZS8
  - ZS9
  - ZS10
  - ZS11
  - ZS18
- C
  - ZS3
  - ZS14
- D
  - ZS19
  - ZS22
- E
  - ZS6
  - ZS20
- F
  - ZS12

Strain	Phage		
	NL_ZS_1	NL_ZS_2	NL_ZS_3
ZS1	(white)	(white)	(light gray)
ZS2	(white)	(white)	(light gray)
ZS3	(light gray)	(dark gray)	(dark gray)
ZS4	(dark gray)	(dark gray)	(dark gray)
ZS5	(light gray)	(light gray)	(dark gray)
ZS6	(white)	(white)	(light gray)
ZS7	(dark gray)	(dark gray)	(dark gray)
ZS8	(light gray)	(dark gray)	(dark gray)
ZS9	(white)	(white)	(dark gray)
ZS10	(dark gray)	(dark gray)	(dark gray)
ZS11	(white)	(white)	(light gray)
ZS12	(white)	(white)	(light gray)
ZS13	(light gray)	(dark gray)	(dark gray)
ZS14	(light gray)	(dark gray)	(dark gray)
ZS15	(dark gray)	(dark gray)	(dark gray)
ZS16	(dark gray)	(dark gray)	(dark gray)
ZS17	(white)	(white)	(light gray)
ZS18	(white)	(white)	(dark gray)
ZS19	(white)	(white)	(light gray)
ZS20	(white)	(white)	(light gray)
ZS21	(white)	(white)	(light gray)
ZS22	(light gray)	(dark gray)	(dark gray)

NL_ZS_1
NL_ZS_2
NL_ZS_3

1 Kbp

100%

66%

Phage-associated ATP-dependent DNA ligase
Phage protein Gp1.6
Phage nucleotide kinase Gp1.7
Phage protein Gp1.8
Phage protein Gp2
Phage single-stranded DNA-binding proteinGp2.5
Phage endonuclease I
Phage endolysin
Phage protein
Phage protein
Phage primase/helicase protein Gp4A
Phage protein Gp4.3
Phage protein Gp4.5
Phage protein Gp4.7
Phage DNA-directed DNA polymerase

hypothetical protein
Phage protein Gp5.7
hypothetical protein
Phage exonuclease
Phage protein Gp6.3
Phage protein Gp6.5
Phage virion protein Gp6.7
Phage minor assembly protein Gp7.3
Phage collar
Phage capsid assembly scaffolding protein Gp5
Phage major capsid protein Gp10A
Phage protein
Phage non-contractile tail tubular proteinGp11
Phage non-contractile tail tubular proteinGp12
Phage scaffold protein Gp13

Phage DNA ejectosome component
Phage DNA ejectosome component
Phage DNA ejectosome component Gp16
tail fibers protein
Phage holin, class II Gp17.5
Phage terminase small subunit Gp18, DNApackaging
Phage Rz-like lysis protein Gp18.5
Phage terminase large subunit Gp19, DNApackaging
Phage protein Gp7.8/Gp7.7 contains HNHendonuclease motif
Phage S-adenosyl-L-methionine hydrolase
Phage protein
Phage serine/threonine kinase
Phage DNA-directed RNA polymerase
Phage protein
Phage protein

YP_004678767.1 Escherichia phage K30
NL_ZS_2
YP_009787501.1 Klebsiella phage vB_KpnP_BIS33
YP_009215503.1 Klebsiella phage vB_KpnP_KpV289
YP_002003836.1 Klebsiella phage K11
NL_ZS_1
QEQ94702.1 Erwinia phage pEp_SNUABM_09
YP_009801107.1 Pectobacterium phage Jarilo
YP_006990246.1 Stenotrophomonas phage IME15
QMP81570.1 Aeromonas phage PZL-Ah1
YP_009799428.1 Escherichia phage Ebrios
NP_523347.1 Enterobacteria phage T3
WP_154159299.1 Xylella fastidiosa
YP_009786120.1 Escherichia phage ECA2
NP_052122.1 Yersinia phage phiYeO3-12
QNN97382.1 Serratia phage SALSA
YP_009786915.1 Serratia phage SM9-3Y
YP_001949795.1 Salmonella phage phiSG-JL2
NL_ZS_3
YP_009949166.1 Enterobacter phage KNP3

0.080

Table 1. MLST types and antibiotic susceptibility profiles of the 22 *Klebsiella pneumoniae* isolates and their corresponding phage-resistant mutants. Antibiotic abbreviations: ticarcillin/clavulanic acid (TIM), piperacillin/tazobactam (TZP), ceftazidime (CAZ), cefoperazone (CPZ), cefepime (FEP), aztreonam (ATM), imipenem (IPM), meropenem (MEM), amikacin (AMK), tobramycin (TOB), ciprofloxacin (CIP), levofloxacin (LVX), doxycycline (DOX), minocycline (MIN), tigecycline (TGC), colistin (CST), trimethoprim-sulfamethoxazole (SXT).

Table 2. Morphological and genomic characterization of phages isolated from the local hospital sewage.

Table 3. Phage susceptibility properties of *K. pneumoniae* ZS15 isolates obtained from cultures enriched with the phages. "Resistant" indicates that isolates were not susceptible to phage infection in the spot test. "Partially sensitive" indicates that isolates had a reduced susceptibility to the phage. "Sensitive" indicates that isolates were fully sensitive to the phage.

										Minimum inhibitory concentration (μg/mL)									
Strain	MLST	bla_{KPC}	TIM	TZP	CAZ	CPZ	FEP	ATM	IMP	MEM	AMK	TOB	CIP	LVX	DOX	MIN	TGC	CST	SXT
ZS1	ST11	Yes	≥128	≥128	≥64	≥64	≥32	≥64	≥16	≥16	≥64	≥16	≥4	≥8	2	4	≤0.5	≤0.5	≤20
ZS2	ST11	Yes	≥128	≥128	≥64	32	≥32	≥64	≥16	≥16	4	8	≥4	≥8	2	8	1	≥16	≤20
ZS3	ST11	No	≥128	≥128	≥64	≥64	≥32	≥64	1	1	≥64	≥16	≥4	≥8	4	≥16	2	≤0.5	≤20
ZS4	ST11	Yes	≥128	≥128	32	≥64	≥32	≥64	0.5	≤0.25	≥64	≥16	≥4	≥8	2	8	1	≤0.5	≤20
ZS5	ST11	Yes	≥128	≥128	≥64	≥64	≥32	≥64	≥16	≥16	4	≤1	≥4	≥8	8	≥16	4	≤0.5	≤20
ZS6	ST11	Yes	≥128	≥128	≥64	≥64	≥32	≥64	≥16	≥16	≥64	≥16	≥4	≥8	1	2	≤0.5	≤0.5	≤20
ZS7	ST11	Yes	≥128	≥128	≥64	≥64	≥32	≥64	≥16	≥16	4	≤1	≥4	≥8	≥16	≥16	≥8	≤0.5	≤20
ZS8	ST11	Yes	≥128	≥128	≥64	≥64	≥32	≥64	≥16	≥16	≥64	≥16	≥4	≥8	4	8	2	≤0.5	≤20
ZS9	ST11	Yes	≥128	≥128	≥64	≥64	≥32	≥64	≥16	≥16	≥64	≥16	≥4	≥8	4	8	2	≤0.5	≤20
ZS10	ST11	Yes	≥128	≥128	≥64	≥64	≥32	≥64	≥16	≥16	≥64	≥16	≥4	≥8	≥16	≥16	4	4	≥320
ZS11	ST11	Yes	≥128	≥128	≥64	≥64	≥32	≥64	≥16	≥16	≥64	≥16	≥4	≥8	2	8	2	≤0.5	≤20
ZS12	ST11	Yes	≥128	≥128	≥64	≥64	≥32	≥64	≥16	≥16	≥64	≥16	≥4	≥8	2	8	1	≤0.5	≤20
ZS13	ST11	Yes	≥128	≥128	≥64	≥64	≥32	≥64	≥16	≥16	≥64	≥16	≥4	≥8	2	8	≤0.5	≤0.5	≤20
ZS14	ST11	No	≥128	≥128	32	≥64	≥32	≥64	1	0.5	≥64	≥16	≥4	≥8	2	8	≤0.5	≤0.5	≤20
ZS15	ST11	Yes	≥128	≥128	≥64	≥64	≥32	≥64	≥16	≥16	≥64	≥16	≥4	≥8	2	8	8	≤0.5	≤20
ZS16	ST11	Yes	≥128	≥128	≥64	≥64	≥32	≥64	≥16	≥16	≥64	≥16	≥4	≥8	4	≥16	2	≥16	≤20
ZS17	ST11	Yes	≥128	≥128	≥64	≥64	≥32	≥64	≥16	≥16	≥64	≥16	≥4	≥8	4	8	1	≤0.5	≤20
ZS18	ST11	Yes	≥128	≥128	≥64	32	≥32	≥64	≥16	8	8	≤1	≥4	≥8	≤0.5	2	≤0.5	≤0.5	≤20
ZS19	ST11	No	≥128	≥128	8	≥64	≥32	≥64	≤0.25	≤0.25	≥64	≥16	≥4	≥8	2	4	≤0.5	≤0.5	≤20
ZS20	ST11	Yes	≥128	≥128	≥64	≥64	≥32	≥64	≥16	≥16	≥64	≥16	≥4	≥8	1	4	≤0.5	≤0.5	≤20
ZS21	ST11	Yes	≥128	≥128	≥64	≥64	≥32	≥64	≥16	≥16	≥64	≥16	≥4	≥8	1	4	≤0.5	≤0.5	≤20
ZS22	ST11	Yes	≥128	≥128	≥64	≥64	≥32	≥64	≥16	≥16	≥64	≥16	≥4	≥8	1	2	≤0.5	≤0.5	≤20

Phage	Family	Host	Head diam (nm)	Head length (nm)	Tail diam (nm)	Tail length (nm)	Genome size (bp)	GC%	Genome type	Predicted genes	Source
NL_ZS_1	Podoviridae	ZS15	63	58	13	14	40,428	53	dsDNA	45	Sewage
NL_ZS_2	Podoviridae	ZS15	58	63	13	9	40,222	52.8	dsDNA	49	Sewage
NL_ZS_3	Podoviridae	ZS15	63	63	13	11	38,741	50.7	dsDNA	46	Sewage

Percentage of phage susceptibility properties of *K. pneumoniae* ZS15 isolates (%)

Phage	Resistant	Partially sensitive	Sensitive
NL_ZS_1	100%	0	0
NL_ZS_2	100%	0	0
NL_ZS_3	43.3%	50%	6.7%