Parallel genetic changes underlie integrated craniofacial traits in an adaptive radiation of

trophic specialist pupfishes

Michelle E. St. John[1,2], Julia C. Dunker[1], Emilie J. Richards[1,2], Stephanie Romero[3], Christopher

H. Martin[1,2]

[1]Department of Integrative Biology, University of California, Berkeley, CA 94720

[2]Museum of Vertebrate Zoology, University of California, Berkeley, CA 94720

[3]Department of Evolution and Ecology, University of California, Davis, CA 95616

Keywords: quantitative trait loci, ecological speciation, functional morphology, ecomorph,

lepidophagy, durophagy

Correspondence: chmartin@berkeley.edu

Abstract

Many factors such as divergence time, shared standing genetic variation, frequency of

introgression, and mutation rates can influence the likelihood of whether populations adapt to

similar environments via parallel or non-parallel genetic changes. However, the frequency of

parallel *vs* non-parallel genetic changes resulting in parallel phenotypic evolution is still

unknown. In this study, we used a QTL mapping approach to investigate the genetic basis of

highly divergent craniofacial traits between scale- and snail-eating trophic specialist species

across similar hypersaline lake environments in an adaptive radiation of pupfishes endemic to

San Salvador Island, Bahamas. We raised F2 intercrosses of scale- and snail-eaters from two

different lake populations of sympatric specialists, estimated linkage maps, scanned for

significant QTL for 30 skeletal and craniofacial traits, and compared the location of QTL

between lakes to quantify parallel and non-parallel genetic changes. We found strong support for

parallel genetic changes in both lakes for five traits in which we detected a significant QTL in at

least one lake. However, many of these shared QTL affected different, but highly correlated

craniofacial traits in each lake, suggesting that pleiotropy and trait integration should not be

neglected when estimating rates of parallel evolution. We further observed a 23-52% increase in

adaptive introgression within shared QTL, suggesting that introgression may be important for

parallel evolution. Overall, our results suggest that the same genomic regions contribute to

parallel integrated craniofacial phenotypes across lakes. We also highlight the need for more

expansive searches for shared QTL when testing for parallel evolution.

Introduction

Convergent evolution describes the independent evolution of similar phenotypes in response to similar selective pressures and provides strong support for ecological adaptation (Losos 2009; Schluter 2000). This includes both non-parallel genetic changes, such as the evolution of antifreeze glycoproteins in icefishes or the 'thunniform' body shape of lamnid sharks and tunas (Chen et al. 1997; Donley et al. 2004), and parallel genetic changes such as tetrodotoxin resistance in snakes and pufferfishes or the evolution of voltage-gated sodium channels in mormyrid and gymnotiform electric fishes (Hopkins 1995; Katz 2006; Jost et al. 2008; Feldman et al. 2009; Zakon et al. 2006). Instances of convergence across independent lineages (i.e., across groups that lack a recent common ancestor and shared genetic backgrounds) provide the strongest evidence for adaptation; however, repeated evolution of similar phenotypes in response to similar selective pressures among lineages derived from the same ancestral population can also provide insight into the process of adaptation. Understanding this process, traditionally known as parallel evolution (Futuyman 1986), is important because it can help to tease apart the contributions of natural selection and shared genetic constraints to similar phenotypes (Schluter et al. 2004; Stuart et al. 2017). Parallel phenotypic evolution can also occur via parallel or non-parallel genetic changes (e.g., Cresko et al. 2004), but even non-parallel genetic changes occurring in the same ancestral genetic background (e.g. Chan et al. 2010; Xie et al. 2019) provide weaker evidence for adaptation than convergence across independent lineages due this shared history. Despite substantial attention, the frequency and likelihood of parallel phenotypic evolution via parallel or non-parallel genetic changes is still relatively unknown (Stern and Orgogozo 2008; Stern 2013; Rosenblum et al. 2014).

Many factors influence whether parallel phenotypic evolution in similar environments is

produced by parallel or non-parallel genetic mechanisms. First, recently diverged species exhibit

increased probabilities of genetic parallelism when adapting to similar environments. Recently

diverged taxa may inhabit similar environments more frequently or they may have similar

genetic architectures, similar genetic variance-covariance matrices, or similar genetic

backgrounds that produce similar epistatic interactions (Conte et al. 2012; Rosenblum et al.

2014). Second, any mechanism that allows the use of the same adaptive genetic mechanism

should increase the likelihood of convergence via parallelism, including the availability of shared

standing genetic variation and introgression (Rosenblum et al. 2014). For example, threespine

sticklebacks colonized freshwater thousands of times and converged on similar phenotypes

largely due to selection on an ancient shared pool of marine standing genetic variation (Jones et

al. 2012; Feulner et al. 2013; Nelson and Cresko 2018; Haenel et al. 2019; but see: Chan et al.

2010; Stuart et al. 2017). Similarly, increased adaptive introgression should also make genetic

parallelism more likely (Grant et al. 2004; Morjan and Rieseberg 2004; Hedrick 2013; Taylor et

al. 2020). Third, adaptive genetic variation with larger effect sizes and fewer pleiotropic effects

should be reused more frequently across populations, particularly when a population is far from a

new adaptive optimum (Linnen et al., 2013; Orr, 2005; Stern, 2013). Finally, de novo mutations,

large mutational target sizes, and polygenic adaptive phenotypes are more likely to result in

parallel phenotypic evolution via non-parallel genetic pathways (Wittkopp et al. 2003; Kowalko

et al. 2013; Bolnick et al. 2018, but see: Colosimo et al. 2004; Chan et al. 2010; Xie et al. 2019).

Quantitative trait locus (QTL) mapping is often used to infer whether parallel or non-

parallel genetic changes underlie parallel phenotypes. However, many QTL studies only

investigate a limited number of traits that are controlled by large effect loci, which may bias the

literature towards supporting genetic parallelism (Conte et al. 2012). This bias may be

exacerbated by the fact that in many QTL studies the genomic regions associated with a parallel

phenotype are large, contain many genes, and their effects on phenotypic variance are

overestimated in under-powered studies (Beavis 1998). These methodological and experimental

limitations reduce confidence in the specific genomic regions associated with a parallel

phenotype and, by extension, reduce confidence in whether parallel evolution was due to parallel

or non-parallel genetic changes. One possible solution is to compare the genomic regions

associated with many different phenotypes across populations (Erickson et al. 2016). In this

scenario, shared genomic regions across populations provide strong support for genetic

parallelism, except in the likely rare instances of independent de novo mutations within the same

region (O'Brown et al. 2015; Xie et al. 2019; Chan et al. 2010).

The San Salvador Island (SSI), Bahamas pupfish radiation is an excellent system for

investigating the genetic underpinnings of parallel ecomorph phenotypes because novel trophic

specialists occur in sympatry across multiple hypersaline lake populations on the island. The

radiation includes three pupfish species: a generalist pupfish (*Cyprinodon variegatus*), a scale-

eating (lepidophagous) pupfish (*C. desquamator*), and a snail-eating (durophagous) pupfish (*C.*

brontotheroides; Martin and Wainwright 2013). The snail- and scale-eating pupfishes are

endemic to SSI and occur in sympatry with one another and the generalist pupfish.

Among lakes, specialists have converged on multivariate phenotypes that are adaptive for

their given ecological niche. For example, scale-eaters across all lakes exhibit increased oral jaw

size (Martin & Wainwright, 2013; Hernandez et al. 2018) and reduced lower jaw angles during

scale-eating strikes which may play a critical role in scale-biting performance during high-speed

strikes on their prey (St. John et al. 2020b). Similarly, the snail-eating pupfish exhibits a novel

nasal protrusion which may improve oral snail-shelling performance or result from sexual

selection (Martin and Wainwright 2013; St. John et al. 2020a). Furthermore, the nasal protrusion

of the snail-eating species varies substantially among lake populations (Martin and Feinstein

2014; Hernandez et al. 2018). Despite the importance of these species characteristics, we still do

not understand how their genetic architecture varies across populations.

There is some evidence to suggest that parallel genetic changes underlie specialist

phenotypes on SSI. First, the SSI radiation is very young, diverging only about 10 kya (Hagey

and Mylroie 1995). Second, previous genomic analyses show that many of the alleles associated

with trophic specialization arrived on SSI from Caribbean-wide standing genetic variation within

generalist pupfish populations, but there are also some de novo adaptive mutations associated

with scale-eating (Richards et al. 2021). Scale-eaters form a monophyletic group, suggesting a

shared genetic component to the scale-eating phenotype across lakes (Richards and Martin

2017). In contrast, snail-eaters and generalists often genetically cluster together by lake instead

of by species—suggesting that non-parallel genetic changes could underlie parallel snail-eater

phenotypes across lakes (Martin and Feinstein 2014; Richards and Martin 2017). Furthermore,

previous studies have documented strong genetic divergence between scale-eaters from Crescent

Pond and all other populations of scale-eater (Richards & Martin, 2017; Richards et al., 2021).

In this study we mapped the genetic basis of 30 skeletal craniofacial and body traits

associated with snail- and scale-eating using lab-reared F2 intercrosses from Crescent Pond and

Little Lake. We called variants, estimated linkage maps, and performed QTL analyses

independently for each F2 population. We found that only one trait— cranial height—mapped to

the same genomic region in both Crescent Pond and Little Lake, but 4 of the 5 remaining

significant QTL detected in one lake mapped to the same genomic region as a highly correlated

craniofacial trait in the second lake. Ultimately, we conclude that parallel evolution through

reuse of introgressed adaptive alleles is acting to produce similar snail- and scale-eating

phenotypes across lake populations on SSI.

Methods

Genetic cross

Currently, pupfish species can been found in 12 hypersaline lakes across the island: generalist

pupfish are found in allopatry in five lakes, the generalist and snail-eating pupfish are found in

sympatry without the scale-eater in two lakes, the generalist and scale-eater are found in

sympatry in a single lake, and all three species are found in sympatry in four lakes (Martin and

Feinstein 2014). We collected wild-caught scale-eating and snail-eating pupfishes from two

different sympatric populations (containing all three species) on SSI – Crescent Pond and Little

Lake—during the years of 2011 and 2013-2015 using seine nets. We brought individuals back to

the University of California, Davis or the University of California, Berkeley and a single wild-

caught scale-eating female from each lake was allowed to breed freely with a single wild-caught

snail-eating male from the same lake resulting in two separate genetic crosses (one cross from

Crescent Pond and one cross from Little Lake). At least four F1 offspring from each hybrid

population were crossed to produce F2 intercrosses, resulting in 354 individuals from Crescent

Pond and 287 individuals from Little Lake included in this study. All fish were maintained in 40-

L tanks at 5-10ppt salinity at the University of California, Davis or the University of California,

Berkeley. We fed fry a diet of newly hatched *Artemia* nauplii for approximately 40 days post

fertilization, after which they were switched to the adult diet of frozen and pellet foods. We

euthanized fish in an overdose of MS-222 (Finquel, Inc.) according to the approved University of

California, Davis Institutional Animal Care and Use Protocol #17455 or University of California,

Berkeley IACUC Protocol AUP-2015-01-7053, and stored them in 95% ethanol.

Phenotyping

Sex and Mate Preferences

For individuals from Crescent Pond, we recorded their sex using their sexually dimorphic body

and fin coloration. Male pupfish develop a blue iridescent coloration along their anteriodorsal

surface and a black marginal band along their caudal fin (Echelle and Echelle 2020).

Once F2 hybrids reached sexual maturity, we performed mating assays using a subset of

the hybrid females from Crescent Pond to estimate mate preferences for snail- or scale-eating

mates (N=74). Prior to the mating assays, female fish were isolated for at least twelve hours and

conditioned on frozen bloodworms with a 12:12 light:dark cycle. Mating assays occurred in three

1.1 m diameter kiddie pools (5-10ppts salinity). Pools were covered with gravel substrate and

divided in half. In each half, we placed three clear plastic 7.5-L Kritter Keepers in a row

containing three conspecific males housed individually to avoid aggression. Size-matched scale-

eater males were placed on one side of each arena and snail-eating males on the other. Once the

males were placed in individually in clear boxes, a female F2 hybrid from Crescent Pond was

placed into the center of one of the three kiddie pools, chosen at random. We considered females

acclimated to the pool once they had visited both rows of males, after which we started the

seven-minute trial period. During each trial we recorded the amount of time a female spent

within one body-length of each species. Each female was tested consecutively in all three pools,

and we used the mean of her association time (scale-eater association time / total association

time during each 7-minute trial) across all three pools for QTL analysis. Size-matched males

were periodically rotated into and among kiddie pools during the 12-month testing period.

Morphological Traits

To measure skeletal phenotypes in our F2 intercrosses, we cleared and double-stained each

specimen with alizarin red and alcian blue. Before clearing and staining, each fish was skinned

and fixed in 95% ethanol. We then fixed specimens in 10% buffered formalin for at least one

week and stained batches of individually labeled specimens following Dingerkus and Uhler's

(1977) protocol. We suspended cleared and stained specimens in glycerin, and photographed

their left lateral side using a Canon EOS 60D digital SLR camera with a 60 mm macro lens. For

each individual, we took two types of photographs: first, we took a whole-body photograph to

calculate fin and body measurements and second, a lateral skull image to calculate craniofacial

measurements (Figure 1). We used DLTdv8 software (Hedrick 2008) to digitize 11 landmarks

on each whole body image and 19 landmarks on each lateral skull image following the

morphometric methods described in Martin et al. (2017). For individuals from Crescent Pond, we

also weighed the adductor mandibulae muscle mass. Each image included a standardized grid

background which we used to calibrate and transform our measurements from pixels into

millimeters. In total, we measured 354 individuals from Crescent Pond and 287 individuals from

Little Lake. We used R to convert the 30 landmarks into linear distances. To reduce

measurement error due to the lateral positioning of the specimens, we took the mean distances

from the two clearest skull and whole-body photographs for each individual when possible. If an

individual did not have two clear photographs for each orientation, we measured the single

clearest photograph. Finally, we size-corrected each trait by using the residuals from a linear

model including the log-transformed measurement of each trait as the response variable and log-

transformed standard length as the predictor variable. We investigated whether size-corrected

traits varied between the two populations using a PCA and a MANOVA test, but found no

appreciable difference between them (Fig. S1, num df = 28, approximate F-value= 0.34, $P = 1$)

Genotyping

We genotyped individuals using three different methods: First, we used whole genome

resequencing for the wild-caught F0 parental generation of our Crescent Pond and Little Lake

intercrosses. We used DNeasy Blood and Tissue Kits (Qiagen, Inc.) to extract DNA from the

muscle tissue of each fish and quantified it on a Qubit 3.0 fluoromether (Thermofisher Scientific,

Inc.). Genomic libraries were then prepared at the Vincent J. Coates Genomic Sequencing Center

(QB3) using the automated Apollo 324 system (WaterGen Biosystems, Inc.). Samples were

fragmented using Covaris sonication and barcoded with Illumina indices. A quality check was

also performed on all samples using a Fragment Analyzer (Advanced Analytical Technologies,

Inc.). We used 150 paired-end sequencing on an Illumina Hiseq4000 for these four parental

samples along with an additional 38 samples that were included in a previous study (Richards

and Martin 2017).

Second, in addition to the 190 previously sequenced individuals from Crescent Pond used

for a QTL mapping study (Martin et al. 2017), we included an additional 164 F2 individuals

from Crescent Pond sequenced using double-digest restriction site associated sequencing

(ddRADseq) following similar library prep and sequencing methods described in Martin et al.

(2015, 2016, 2017). Briefly, we prepared four indexed libraries each containing 96 barcoded

individuals. We sequenced these using 100 single-end high-output mode on two lanes of

Illumina Hiseq4000 at the Vincent J. Coates Genomic Sequencing Center (QB3).

Finally, we sequenced all F2 individuals from Little Lake and a subset of previously

sequenced, but low-coverage Crescent Pond F2's (N=84), using Nextera-tagmented reductively-

amplified DNA (NextRad) sequencing (Russello et al. 2015). We followed the above methods

for DNA extraction and sent samples to SNPsaurus (SNPsaurus, LLC) for quality checking,

NextRad library preparation, and 150 single-end sequencing on two lanes of Illumina Hiseq4000

at the University of Oregon sequencing core.

Calling Variants

We used the following methods to call variants separately for: 1) the Crescent Pond intercross (2

parents and 354 F2 hybrids), and 2) the Little Lake intercross (2 parents and 285 F2 hybrids):

First, we inspected raw read quality using FastQC (Babraham Bioinformatics Institute, v0.11.7)

and trimmed reads to their appropriate length (100bp for samples sequenced with ddRAD, and

150bp for samples sequenced with NextRAD) using TrimGalore! (v0.6.4). For samples that were

sequenced using both ddRAD and NextRad methods, we concatenated trimmed raw reads into a

single file. We next used bwa-mem to map reads from all individuals in an intercross, both

parents and offspring, to the *Cyprinodon brontotheroides* reference genome (v 1.0; total

sequence length = 1,162,855,435 bp; number of scaffolds = 15,698, scaffold N50 = 32 Mbp;

(Richards et al. 2021)). We identified duplicate reads using MarkDuplicates and created BAM

indices using BuildBamIndex in the Picard package (http://picard.sourceforge.net(v.2.0.1)).

Following the best practices guide from the Genome Analysis Toolkit (v 3.5; (Depristo et al.

2011)), we called and refined our single nucleotide polymorphism (SNP) variant data set using

the program HaplotypeCaller. Pupfish lack high-quality known alleles because they are a non-

model organism; we therefore used the recommended hard filter criteria (QD < 2.0; FS <; 60;

MQRankSum < -12.5; ReadPosRankSum < -8; (Depristo et al. 2011; Marsden et al. 2014)) to

filter our SNP variant dataset. Ultimately, we detected 13.7 million variants in our Crescent Pond

dataset and 14.4 million variants in our Little Lake dataset.

We used the program STACKS to further filter our dataset and convert our vcf files into

phenotype and genotype comma-separated values files that could be imported into the Rqtl

program. Specifically, we used the populations program to filter out variants that were not

present in both the parental and F2 populations, and to filter out variants found in 10% or less of

the population. From this filtering step we retained 36,318 variants with 46.5 mean mappable

progeny per site in Crescent Pond and 87,579 variants with 85.984 mean mappable progeny per

site in Little Lake.

We continued to filter our datasets using the Rqtl (v1.46-2), and ASMap (v1.0-4)

packages (Broman et al. 2003; Taylor and Butler 2017). We started filtering by removing

individuals that did not contain any filtered variants and any duplicate individuals. This reduced

our Crescent Pond data set to 227 individuals, and our Little Lake data set to 281 individuals.

Next, we filtered markers that had >0.98 or <0.1 heterozygosity (Crescent Pond: markers

=15,247, Little Lake: markers=14,661). This step also filtered out 13 individuals from Crescent

Pond which only contained markers with >0.98 or <0.1 heterozygosity. Before constructing our

genetic maps, we set aside markers that appeared to suffer from segregation distortion. We used

the pullCross() function from the ASmap package to set aside markers in both data sets that were

missing in >75% of individuals, departed from Mendelian ratios (1:2:1), or any co-located

markers for the initial construction of the linkage maps. This filtering retained more than twice

the number of markers for Crescent Pond than Little Lake. We therefore used a stricter filtering

threshold for missing data (i.e., removing markers with >72% missing data) for our Crescent

Pond dataset to construct linkage maps of comparable sizes for downstream comparative

analyses. At the end of this filtering process the Crescent Pond dataset contained 214 individuals

and 657 markers and the Little Lake dataset contained 281 individuals with 490 markers.

Linkage Map Construction

We used the mstmap.cross() function to form initial linkage groups and order markers, using the

kosambi method for calculating genetic distances and a clustering threshold of $P = 1 \times 10^{-14}$ for

Little Lake and $P = 1 \times 10^{-20}$ for Crescent Pond. After forming these initial linkage groups, we

used the pushCross() function from the ASmap package to integrate previously set aside markers

back into our map. We pushed markers back based on a segregation ratio of 3:4:3 and we pushed

back any markers that had previously been designated as co-located. This increased our map

sizes to 817 markers for Crescent Pond and 580 markers for Little Lake. With these additional

markers, we re-estimated our linkage map using the est.rf() and formLinkageGroups() functions

from the Rqtl package. We used a max recombination fraction of 0.35 and a minimum LOD

threshold of 5 to estimate linkage groups for both data sets. We used the

droponemarker() command from Rqtl with an error probability of 0.01 to identify and drop

problematic markers from the genetic maps, including dropping linkage groups with 3 or fewer

markers. Finally, we visually inspected our linkage groups using plotRF() from the Rqtl package,

and merged linkage groups which had been incorrectly split up using the mergeCross() function

from the ASmap package. Ultimately our final genetic maps included: 1) Crescent Pond: 214

individuals, 743 markers, 24 linkage groups and 2) Little Lake: 281 individuals, 540 markers,

and 24 linkage groups (**Figure *2***).

QTL Analyses

We mapped QTL for 29 skeletal traits for both populations, and additional morphological

(adductor mandibulae muscle mass) and behavioral traits (mate preference) for Crescent Pond.

We used the Rqtl2 package (v0.22-11) to calculate genotype probabilities with a multipoint

hidden Markov model using an error probability of 0.0001 and a Kosambi map function. We

calculated kinship matrices to account for the relationship among individuals in two ways: 1)

overall kinship, which represents the proportion of shared alleles between individuals, and 2)

kinship calculated using the leave-one-chromosome-out method (LOCO). We used the scan1()

function to perform three separate genome scans using a single-qtl model by: 1) Haley-Knott

regression, 2) a linear mixed model using the overall kinship matrix, and 3) a linear mixed model

using the LOCO kinship matrix. For our Crescent Pond data set we also included sex as an

additive covariate. We assessed the significance of all three models using two significance

thresholds $P < 0.1$ and 0.05 based on 1000 permutations each, using the scan1perm() function.

As noted above the scan1() function can use several different methods to determine if a region is

significantly associated with a given phenotype (Broman et al., 2019; Haley & Knott, 1992;

Yang, Zaitlen, Goddard, Visscher, & Price, 2014; Yu et al., 2006), however, it is clear from

previous theoretical work that many of these methods may suffer from type II error depending on

the size of an organism's genome, the density of markers in a linkage map, or the complexity of

the phenotypic traits being measured (Lander and Botstein 1989; Risch 1990). We therefore

relaxed the *P*-value cut off from 0.05 to 0.1 to capture potentially important genomic regions.

This relaxation is further supported by the LOD scores associated with regions significant at the

$P<0.1$ level because they all exceed the traditional threshold of 3 (Nyholt 2000), the more

conservative threshold of ~3.3 (Lander and Kruglyak 1995; Nyholt 2000), the suggestive

threshold of 1.86 (Lander and Kruglyak 1995), and are in line with estimates of significant LOD

thresholds in previous studies (Erickson et al. 2016). All three of these methods detected similar

QTLs and moving forward we only used the Haley-Knott regression method.

For each trait, we calculated the location of the maximum LOD score, and used the fit1()

function to re-fit a single-QTL model at this location. We used the newly calculated LOD score

to estimate the proportion of variance explained by the QTL and to calculate a P-value

associated with each significant QTL (x^2-test). We also used the location of the maximum LOD

score to calculate 95% Bayes credible intervals using the bayes_int() function from the Rqtl2

package. We note that the maximum LOD score associated with every trait across both ponds

exceeded the suggestive threshold of 1.86 (Lander and Kruglyak 1995). We used the

find.markerpos() function from Rqtl to determine where markers in each linkage map fell within

the reference genome. With this information we were able to determine the scaffolds/positions

from the reference genome that fell within the 95% credible intervals for each putative QTL.

Finally, we used the maxmarg() function from the Rqtl2 package to find the genotype with the

maximum marginal probability at the location of the maximum LOD. We used these genotypes

to visualize the relationship between genotype and phenotypes.

Identifying adaptive alleles within QTL regions

For each scaffold that fell within a QTL's credible interval we calculated the minimum and

maximum position for that scaffold (that was identified in the putative QTL region) and searched

the *C. brontotheroides* reference genome for annotated genes within the region. We then

compared this list to a previously published list of genes that 1) contained or were adjacent to

(within 20 kbp) fixed or nearly fixed (*Fst* > 0.95) SNPs between specialist species on SSI, and 2)

showed significant evidence of a hard selective sweep in both the site frequency spectrum-based

method SweeD (Pavlidis et al. 2013) and the linkage-disequilibrium-based method OmegaPlus

(Alachiotis et al. 2012). We hereafter refer to these loci as adaptive alleles. We also noted

whether adaptive alleles within QTL regions were classified as de novo, introgressed, or as

standing genetic variation on SSI (Richards et al. 2021). We used a bootstrap resampling method

to determine whether the observed proportions of adaptive alleles originating from de novo,

introgression, or standing genetic variation found within QTL 95% credible intervals were

different than the proportions expected when drawn from the genome at random. We used the

boot (v. 1.3-25) package (Buckland et al. 1998; Canty and Ripley 2021) to resample our entire

adaptive allele dataset (with replacement) 10,000 times. We then used the boot.ci() command

from the boot package to calculated the 95% credible intervals around expected proportions of de

novo, introgressed, and standing adaptive alleles. We performed these calculations separately for

scale-eater and snail-eater adaptive alleles.

Results

Linkage Map Construction

We identified 24 linkage groups from 743 markers for Crescent Pond, and 24 linkage groups

from 540 markers for Little Lake (**Figure *2***). Previous karyotypes of *Cyprinodon* species

estimated 24 diploid chromosomes, matching the linkage groups in this study (Liu & Echelle,

2013; Stevenson, 1981). The total map length for Crescent Pond was 7335 cM and the total map

length for Little Lake was 5330; the largest linkage groups for each map were 740 cM and 380

cM, respectively, and inter-marker map distance did not exceed 20cM in either map. To compare

our maps and to determine if the same genomic regions were being reused across lakes, we

identified where each marker was located in our reference genome. Overall, we found 324

markers in both maps that were within 10 Kbp of one another, indicating that 60% of the Little

Lake map was also present in the Crescent Pond map and 44% of the Crescent Pond map was

present in the Little Lake map (**Figure 3**).

Craniofacial QTL

We detected three significant QTL in Crescent Pond and five QTL in Little Lake (**Table _1_**,

Table _2_). In Crescent Pond, we identified QTL associated with the depth of the dentigerous arm

of the premaxilla, cranial height, and adductor mandibulae muscle mass. Cranial height in

Crescent Pond mapped to linkage group (LG) 10. Dentigerous arm depth and adductor

mandibulae muscle mass both mapped to LG 13, which also contained the max LOD scores for

two additional jaw traits (jaw opening in-lever and maxillary head height; **Table _2_**). The 95%

credible intervals for all these traits overlapped, suggesting that LG 13 may contain a single

pleiotropic locus or many loci that affect all four traits.

In Little Lake, we detected significant QTL associated with jaw closing in-lever (i.e.

height of the coronoid process on the articular: LG9), width and depth of the dentigerous arm of

the premaxilla (LG3 and LG6), maxillary head protrusion (LG10), and cranial height (LG1;

Table _1_,**Table _2_**). The 95% credible interval for dentigerous arm width on LG3 also contained

the max LOD score for lower jaw length, suggesting that either a single pleiotropic locus or a

cluster of loci in this region may be controlling both traits.

Candidate genes and adaptive alleles within QTL regions

Cranial height

Cranial height was the only trait with statistically significant or marginally significant QTL in both lakes (Figure 4, $P < 0.1$). While the QTL occurred on different linkage groups between maps, we found a high degree of synteny between these linkage groups indicating that the QTL is located in the same genomic region in both lakes (**Table *2***, **Figure 3**). We also found the same overdominant genetic pattern in both lake crosses: heterozygotes showed increased cranial height relative to homozygous individuals (**Figure *5***).

We found 44 genes within scaffold 33 that fell partially or fully within the 95% credible intervals of the QTL in both lakes (Table 1, Table S1). Only three of these genes contained adaptive alleles within 20 kb: *wdr31*, *bri3bp*, and *gnaq* (**Table *3***). Interestingly, *gnaq* is well known to be associated with craniofacial development (Hall et al. 2007; Shirley et al. 2013) and is differentially expressed between our specialist species in developing larvae (McGirr and Martin 2020).

Dentigerous Arm Width

We found that regions on scaffolds 58 and 24 were associated with a significant QTL for dentigerous arm width in Little Lake and contained the max LOD scores for maxillary head protrusion and female mate preference in Crescent Pond (**Table *1***, **Table *2***). We found 161 genes which fell partially or completely within these shared regions, but only 2 genes, *dysf* and *cyp26b1*, which contained adaptive alleles within 20 kbp (**Table *3***). The *dysf* gene provides instructions for making a protein called dysferlin, which is found in the sarcolemma membrane

that surrounds muscle fibers (Liu et al. 1998). This could indicate a role for muscle development

in affecting skeletal development of the maxilla and premaxilla.

Dentigerous Arm Depth

The QTL for dentigerous arm depth in Little Lake was associated with LG 6, which corresponds

to LG 7 in Crescent Pond, however, no traits from Crescent Pond mapped to this linkage group

(**Table 2**, **Figure 3**). Instead, dentigerous arm depth in Crescent Pond was associated with LG 13

and did not share any similar genomic regions with those associated with dentigerous arm depth

in Little Lake. We found 80 genes completely or partially within the 95% credible region for this

QTL in Little Lak, but none contained adaptive alleles based on our criteria (Figure S1). In fact,

only a single adaptive allele was found in this QTL region, but it was in an unannotated region of

the genome (**Table 3**).

Maxillary Head Protrusion

Maxillary head protrusion in Little Lake mapped to a QTL region on LG10 which corresponds to

the max LOD scores for both lower jaw length and caudal peduncle height in Crescent Pond

(**Table 2**, **Figure 3**). Across lakes, all three traits were associated with scaffolds 53, 2336, and

6275. We found 528 genes partially or fully within these shared regions, but only 21 of these

genes contained adaptive alleles within 20 kbp (**Table 3**). One of these genes, *twist1*, contains a

non-synonymous substitution fixed in scale-eating pupfish on San Salvador Island, Bahamas

(Richards et al. 2021). *Twist1* is a transcription factor and oncogene associated with palate

development and oral jaw size in model organisms (Parsons et al. 2014; Teng et al. 2018).

Jaw Closing In-Lever

The QTL for jaw closing in-lever was associated with LG 9 in Little Lake, which corresponds to

the max LOD scores for orbit diameter and anterior body depth in Crescent Pond (**Table *2***,

Figure 3). Scaffolds 8 and 8020 were associated with all three of these traits. We found 13 genes

which partially or completely fell within these shared regions, and only two genes, *map2k6* and

galr2, which contained adaptive alleles within 20 kbp (**Table *3***). *Galr2* was also previously

detected within a significant QTL for lower jaw length in pupfish (Martin et al. 2017).

Dentigerous Arm Depth and Adductor Mandibulae Muscle Mass

Finally, in Crescent Pond the QTL for dentigerous arm depth and adductor mandibulae muscle

mass mapped to the exact same location on LG 13 (95% CI dentigerous arm depth (0, 250),

adductor mandibulae muscle mass (0,70). This linkage group corresponds to LG14 in Little

Lake, which contains the max LOD scores for both palatine height and suspensorium length

(**Table *3***). We found 52 genes that overlapped between these regions, 18 of which contained

adaptive alleles. Furthermore, three of the genes— *ube2w, ncoa2, and prlh*—contained adaptive

alleles that introgressed from Laguna Bavaro in the Dominican Republic to snail-eating pupfish

(*ube2w*), from Lake Cunningham, New Providence Island to scale-eating pupfish (*ncoa2*), or

from North Carolina, USA to scale-eating pupfish (*prlh)*. We also found four genes that

contained adaptive alleles within 20 kbp that arose from de novo mutations: *cd226, cmbl, slc51a,*

and *zfhx;* however*,* only one adaptive allele in *slc51a* is found within a coding region.

Origins of adaptive alleles

Adaptive alleles originating from standing genetic variation across the Caribbean were most

common within shared QTL regions between lakes (86.03% within scale-eater populations, and

53.32% within snail-eating populations; **Table *3***). However, observed proportions within shared

QTL were significantly less than expected by chance (scale-eater expected 95% CI: (88.33%-

90.37%), snail-eater expected 95% CI: (62%-67%;10,000 bootstrapped iterations). Instead, we

found more introgressed scale-eater and snail-eater adaptive variants in shared QTL regions than

expected by chance (Scale-eater observed: 12.13% introgressed, scale-eater expected 95% CI:

(7.96%-9.88%); Snail-eater observed: 46.67% introgressed, snail-eater expected 95% CI:

(32.22%-37.06%)). Finally, we found that about 1.83% of adaptive alleles within overlapping

regions between lakes originated from de novo mutations in scale-eaters, however, this fell

within the predicted null range (95% CI: (1.29%-2.17%)).

Discussion

Parallel genetic changes underlie 5 out of 6 of craniofacial QTL

We found evidence supporting both parallel and non-parallel genetic changes in an adaptive

radiation of trophic specialist pupfishes. A single significant QTL was associated with cranial

height in both lakes and mapped to the same genomic region, suggesting that parallel genetic

changes are responsible for variation in this trait in both lakes. On the other hand, significant

QTLs were identified for premaxilla dentigerous arm depth in each lake, but they mapped to

different locations, indicating that this trait is associated with non-parallel genetic changes. We

found an additional three traits with significant QTLs (dentigerous arm width, jaw closing in-

lever, maxillary head protrusion) in the Little Lake population that were not detected in Crescent

Pond. However, all genomic regions associated with these traits in Little Lake also mapped to

the max LOD score for this same integrated suite of craniofacial traits in Crescent Pond.

Therefore, rather than assume independent QTL for each trait, we conservatively conclude that

the same genomic regions are being reused in each lake and affect a highly integrated suite of

craniofacial traits. Overall, we found that 5 out of the 6 significant QTLs were reused in some

way across lakes suggesting that parallel genetic changes underly adaptive phenotypes in the San

Salvador Island pupfish radiation.

High level of QTL reuse across ponds

Overall, we found that about 16% (1 out of 6) of the identified QTL regions corresponded to

non-parallel changes and 84% (5 out of 6) corresponded to parallel genetic changes—either

affecting the same phenotypic trait or a tightly correlated craniofacial trait— across populations.

The presence of both non-parallel and parallel genetic changes leading to convergent phenotypes

across lakes has been documented previously. For example, Colosimo et al. (2004) investigated

the genetic basis of armor plate morphology in two independent threespine stickleback

populations and found a single large effect locus on LG 4 in the two populations. However, they

also noted a potential difference in the dominance relationships of alleles across ponds at this

location, and found additional differences in modifier QTLs between populations, suggesting

that both parallel and non-parallel genetic changes could lead to the loss of armor plating.

Similarly, Erickson et al. (2016) found evidence for both parallel (43% of QTL regions

overlapped between at least two populations) and non-parallel (57% of QTL regions were found

in only a single population) evolution in a QTL study investigating the genetic basis of 36

skeletal phenotypes in three independent threespine stickleback populations. However, our

findings suggest that pupfish exhibit a much higher proportion of parallel evolution than

previously documented in stickleback. In fact, Conte et al. (2012) estimated that the probability

of convergence via gene reuse is only 32-55% —which is 1.5 to 2.5 times lower than our current

finding— although this may be underestimated (Stern 2013).

Pupfish may have a higher rate of parallel evolution than other model fish speciation

systems for a few reasons. First, the pupfish radiation is recent, although comparable in age to

glacial stickleback populations, with specialist species diverging less than 10kya (Hagey and

Mylroie 1995; Martin and Wainwright 2013), and parallel evolution is predicted to be more

likely when populations or species have recently diverged (Rosenblum et al. 2014). This may be

because recently diverged species are more likely to experience similar environments, have

access to similar pools of genetic variation (either due to standing genetic variation or

introgression), or similar genetic constraints. Second, the genomic basis of pupfish skeletal traits

may be primarily controlled by cis-regulatory elements, which evolve more quickly and have

less negative pleiotropy which may make them more likely to undergo parallel evolution (Stern

and Orgogozo 2008). However, a previous study of allele-specific expression in the pupfish

system found strong evidence that two cis-regulatory alleles were associated with skeletal

development, but trans-acting elements predominated overall (McGirr and Martin 2021).

In part, the increased proportion of parallel evolution estimated in this study results from

our relaxed thresholds for detecting and categorizing shared QTL regions. Previous QTL studies

have typically searched for evidence of parallel evolution by only looking for one-to-one

mapping in which the same genomic regions are associated with the same trait across

populations at a genome-wide level of significance in each (Colosimo et al. 2004; Conte et al.

2012). While this method provides the most clear-cut examples of parallel evolution, we argue

that it vastly underestimates its frequency in nature. For example, this method would not

consider reuse of the same genomic regions for integrated morphological traits as parallel

evolution, a pattern seen in this study and in Erickson et al. (2016). Furthermore, the strict one-

to-one significance method for detecting parallel evolution does not include consideration of the

hierarchy and diversity of convergence and parallel evolution, which can span morphological

traits, ecotypes, performance, or even fitness (James et al. 2020; Rosenblum et al., 2014; Stern,

2013; Martin and Wainwright 2013). Ultimately, we argue that our method of quantifying

parallel evolution provides a more wholistic view of the process and better captures the

frequency of reuse of adaptive genetic variation in nature.

Few QTL may affect many highly integrated craniofacial traits

There are several processes that may cause the same genomic regions to be associated with

different traits between lakes. First, these genomic regions may be highly pleiotropic and affect

several traits simultaneously. For example, Albert et al. (2007) found that that on average a

single QTL affected 3.5 phenotypic traits in an analysis of 54 body traits in three-spine

stickleback. Wagner et al. (2008) found a similar pattern in QTL analyses of 70 skeletal traits in

mice, where a single QTL affected on average 7.8 phenotypic traits (the maximum being 30).

A second possibility is that a single QTL region may contain several tightly linked

causative variants that are responsible for variation in many traits. Correlated phenotypic traits

are generally assumed to have a shared genetic basis, but it is extremely difficult to determine if

this is due to pleiotropy or tight linkage between genomic regions (Lynch and Walsh 1998;

Gardner and Latta 2007; Paaby and Rockman 2013; Wright et al. 2010).

Finally, it may be that differences in methodology or sample sizes between lakes enable

us to detect significant QTL for some traits in one lake and not the other. For example, our

analyses of Little Lake allowed us to detect significant QTL for effect sizes greater than 6.54

PVE at 80% power, but we could only detect significant QTL for effect sizes greater than 8.41

PVE at 80% power in Crescent Pond due to our lower sample size for this cross (Sen et al.

2007). However, this level of power is typical in many non-model QTL studies (Ashton et al.

2017). The ability to detect a significant QTL in one lake but not the other may be further

explained by our use of different sequencing methods between populations. However, a critical

component of our analyses involved searching for regions within 10 kbp of one another across

maps to provide confidence that if we detected a significant QTL in one lake and not the other

that it was not simply because that genomic region was not captured by the sequencing. For

example, in Little Lake we detected a significant QTL associated with dentigerous arm depth on

LG 6 but did not find any traits associated with this region of the genome in Crescent Pond.

QTL are associated with different craniofacial traits across different lakes

In this study we found an intriguing pattern of different traits mapping to the same region of the

genome across lake populations. One potential explanation for this is that there are different

relationships between traits in each lake, and we find some evidence of this in our phenotypic

data. Figure S2 depicts correlation matrices between traits in 1) Little Lake and 2) Crescent

Pond, and X^2 comparisons of these two matrices reveals that the relationship between traits

varies significantly between lakes (X^2=3135.99, df = 756, $P<$ 3.6e-29). For example, the

relationship between maxillary head protrusion and lower jaw length is more than two times

stronger in Little Lake compared to Crescent Pond (Pearson's r_{LL}=0.27, Pearson's r_{CP} = 0.12),

the relationship between dentigerous arm depth and suspensorium length is 1.8 times stronger in

Little Lake than in Crescent Pond (Pearson's r_{LL}=0.45, Pearson's r_{CP} = 0.24), and the relationship

between jaw closing in-lever and anterior body depth is more than two times stronger in Crescent

Pond than in Little Lake (Pearson's r_{CP} = 0.23, Pearson's r_{LL}=0.11).

565   This pattern may be explained by different epistatic interactions in each lake. For

example, Juenger et al. (2005) detected significant QTL-QTL interactions in one mapping

population of *Arabidopsis* but found no evidence of the same interactions in the other population.

When we investigated the relationship between phenotype and genotype for cranial height, we

found the same overdominance pattern in both lakes (Figure 5). However, the presence of

epistatic interactions may also be an obstacle for QTL detection. In a mapping study of body

weight in chicken, Carlborg et al. (2006) were only able to detect a single weak QTL despite the

extremely divergent phenotypes between parental lines. However, when accounting for epistatic

interactions, Carlborg et al. identified several additional significant QTL regions that explained a

large amount of variation in body weights.

575   Finally, our method for searching for putative QTL regions may have led to this pattern.

Similar studies have searched for influential genomic regions by first identifying a putative QTL

in a single population, and then searching the already identified linkage group in the second

population for any signal of a QTL associated with the same phenotype, often using relaxed

LOD thresholds closer to the suggestive cut-off (LOD> 1.8, e.g., Erickson et al. 2016). Our

approach, however, independently identified the positions of maximum LOD for all traits across

the entire linkage map before searching for similar implicated regions between populations. We

argue that our approach minimizes bias, because there are no prior expectations about which

traits should be associated with a given genomic region within a suite of integrated traits, and

reduces false positives because we only examine the maximum LOD position for each trait.

Identifying causative regions within QTL

Multiple mapping populations across lakes may also be particularly useful for identifying

candidate causal alleles. We found that one out of our six unique QTL regions mapped to the

same genomic location across lakes and was associated with the same phenotypic trait—cranial

height (Figure 4). In Crescent Pond, we found that a region of 110 cM was associated with this

trait (LG10, position: 204, 95% CI (130,340)), which contained 426 genes. However, when we

compared this region to the region independently identified in our Little Lake analysis, we found

that the overlapping region was reduced to 20cM (LG1, position: 259, 95% CI (250-270)) and

contained only 44 genes—a reduction of more than 80%. We found a similar pattern in the

additional four QTL regions that mapped to the same genomic location across maps but were

associated with different phenotypic traits and observed an average 56% reduction in region size.

As noted above, Erickson et al. (2016) used a similar method of identifying candidate QTL

regions across three hybrid populations of stickleback, and found that 43% of identified QTL

regions were shared across two or more populations; however, they did not investigate whether

these QTL regions completely or partially overlapped.

We also searched for adaptive alleles within QTL region that were identified in a

previous study as 1) nearly fixed between species (F_{st} >0.95) and 2) showed significant evidence

of a hard selective sweep (Richards et al. 2021). Overall, we found 789 shared genes within

shared QTL regions across lakes, and that 45 of those genes contained adaptive variants (5.7%).

This is a six-fold increase from the genome-wide expectation of 0.91% (176 genes associated

with at least one adaptive variant / 19304 annotated genic regions), suggesting that these specific

regions are important for adaptation to scale- and snail-feeding in wild pupfish. For example, a

variant in *twist1* was found within the region associated with maxillary head protrusion in Little

Lake (which also overlapped with lower jaw length and caudal peduncle height in Crescent

Pond). In model organisms, *twist1* is associated with palate and jaw development (Parsons et al.

2014; Teng et al. 2018), and previous genome-wide association scans in pupfish showed that a

region containing *twist1* was significantly associated with oral jaw size in the system (Richards

et al. 2021). Similarly, we found that variants associated with *galr2* fell within the QTL region

associated with jaw closing in-lever in Little Lake (which also overlapped with regions

associated with orbit diameter and anterior body depth in Crescent Pond; scaffolds 8 and 8020),

and previous QTL mapping studies, gene expression studies, and genome-wide association

analyses have all implicated regions containing *galr2* with oral jaw development in pupfish

(McGirr and Martin 2016; Martin et al. 2017; Richards et al. 2021).

Increased use of introgressed adaptive variants in QTL regions

We found that most genetic variation within shared QTL regions was also segregating across

outgroup Caribbean generalist populations characterized by Richards et al. (2021; 86.04% within

scale-eater populations, and 53.32% within snail-eating populations). Furthermore, we found

more introgressed adaptive alleles from both scale-eater (observed: 12.13% introgressed,

expected 95% CI: (7.96%-9.88%)) and snail-eater populations in shared QTL regions than

expected by chance (observed: 46.67%, expected 95% CI: (32.22%-37.06%)). This supports the

prediction that standing genetic variation and introgressed variation should underlie parallel

genetic changes (Stern 2013; Thompson et al. 2019). Finally, we found that only 1.83% of

adaptive alleles within shared QTL regions across both lakes originated from de novo mutations

on San Salvador Island. While this percentage did not differ significantly from the expected

estimates (expected 95% CI: 1.3%-2.17%) it does not eliminate the possibility that de novo

mutations play an important adaptive role in pupfish evolution.

Conclusion

In conclusion, we found that a single QTL region was responsible for variation in cranial height

in both populations, and an additional four QTL regions were responsible for variation in

different craniofacial traits across lakes, suggesting that parallel genetic changes underlie

integrated suites of adaptive craniofacial phenotypes on San Salvador Island. Adaptive alleles

were more commonly found within these detected QTL regions, and more of these adaptive

alleles arrived on SSI via introgression than expected by chance. Finally, we argue that

investigating QTL regions across populations in concert with estimation of hard selective sweeps

in wild populations is a powerful tool for identifying potential causative regions of the genome

affecting adaptive divergence.

Acknowledgements

We thank the University of California, Davis, the University of California, Berkeley, the

University of North Carolina at Chapel Hill, NSF CAREER 1749764, NIH 5R01DE027052-02,

and BSF 2016136 for funding to CHM. The Bahamas Environmental Science and Technology

Commission and the Ministry of Agriculture provided permission to export fish and conduct this

research. Rochelle Hanna, Velda Knowles, Troy Day, and the Gerace Research Centre provided

logistical assistance in the field. All animal care protocols were approved by the University of

California, Davis and the University of California, Berkeley Animal Care

Author Contributions

MESJ and CHM designed research; MESJ, CHM, JCD, and SR performed data collection; MESJ

and EJR performed data analysis; MESJ and CHM wrote the paper. CHM provided funding.

Data Accessibility

Data will be deposited to Dryad and NCBI. Genomes are archived at the National Center for

Biotechnology Information BioProject Database (Accessions:

PRJNA690558; PRJNA394148, PRJNA391309; and PRJNA305422).

References

Alachiotis, N., A. Stamatakis, and P. Pavlidis. 2012. OmegaPlus: A scalable tool for rapid detection of selective sweeps in whole-genome datasets. Bioinformatics 28:2274–2275. Oxford Academic.

Albert, A. Y. K., S. Sawaya, T. H. Vines, A. K. Knecht, C. T. Miller, B. R. Summers, S. Balabhadra, D. M. Kingsley, and D. Schluter. 2007. The genetics of adaptive shape shift in stickleback: pleiotropy and effect size. Evolution (N. Y). 0:071115145922005-??? John Wiley & Sons, Ltd.

Ashton, D. T., P. A. Ritchie, and M. Wellenreuther. 2017. Fifteen years of quantitative trait loci studies in fish: challenges and future directions. Mol. Ecol. 26:1465–1476.

Beavis, W. D. 1998. QTL analyses: power, precision, and accuracy. Pp. 145–162 *in* Molecular dissection of complex traits.

Besnard, G., A. M. Muasya, F. Russier, E. H. Roalson, N. Salamin, and P. A. Christin. 2009. Phylogenomics of C4 photosynthesis in sedges (Cyperaceae): Multiple appearances and genetic convergence. Mol. Biol. Evol. 26:1909–1919. Oxford Academic.

Bolnick, D. I., R. D. H. Barrett, K. B. Oke, D. J. Rennison, and Y. E. Stuart. 2018. (Non)Parallel evolution. Annual Reviews Inc.

Bricelj, V. M., L. Connell, K. Konoki, S. P. MacQuarrie, T. Scheuer, W. A. Catterall, and V. L. Trainer. 2005. Sodium channel mutation leading to saxitoxin resistance in clams increases risk of PSP. Nature 434:763–767. Nature Publishing Group.

Broman, K. W., D. M. Gatti, P. Simecek, N. A. Furlotte, P. Prins, Ś. Sen, B. S. Yandell, and G. A. Churchill. 2019. R/qtl2: Software for mapping quantitative trait loci with high-dimensional data and multiparent populations. Genetics 211:495–502. Genetics Society of America.

Broman, K. W., H. Wu, S. Sen, and G. A. Churchill. 2003. R/qtl: QTL mapping in experimental crosses. Bioinformatics 19:889–890. Oxford Academic.

Buckland, S. T., A. C. Davison, and D. V. Hinkley. 1998. Bootstrap Methods and Their Application. Biometrics 54:795.

Canty, A., and B. D. Ripley. 2021. boot: Bootstrap R (S-Plus) Functions. R Packag. version 1.3-28.

Carlborg, Ö., L. Jacobsson, P. Åhgren, P. Siegel, and L. Andersson. 2006. Epistasis and the release of genetic variation during long-term selection. Nat. Genet. 38:418–420. Nature Publishing Group.

Chan, Y. F., M. E. Marks, F. C. Jones, G. Villarreal, M. D. Shapiro, S. D. Brady, A. M. Southwick, D. M. Absher, J. Grimwood, J. Schmutz, R. M. Myers, D. Petrov, B. Jónsson, D. Schluter, M. A. Bell, and D. M. Kingsley. 2010. Adaptive evolution of pelvic reduction in sticklebacks by recurrent deletion of a pitxl enhancer. Science (80-.). 327:302–305. American Association for the Advancement of Science.

Chen, L., A. L. Devries, and C. H. C. Cheng. 1997. Convergent evolution of antifreeze glycoproteins in Antarctic notothenioid fish and Arctic cod. Proc. Natl. Acad. Sci. U. S. A. 94:3817–3822.

Christin, P. A., N. Salamin, V. Savolainen, M. R. Duvall, and G. Besnard. 2007. C4 Photosynthesis Evolved in Grasses via Parallel Adaptive Genetic Changes. Curr. Biol. 17:1241–1247. Cell Press.

Colosimo, P. F., C. L. Peichel, K. Nereng, B. K. Blackman, M. D. Shapiro, D. Schluter, and D. M. Kingsley. 2004. The genetic architecture of parallel armor plate reduction in threespine sticklebacks. PLoS Biol. 2:e109. Public Library of Science.

Conte, G. L., M. E. Arnegard, C. L. Peichel, and D. Schluter. 2012. The probability of genetic parallelism and convergence in natural populations. Proc. R. Soc. B Biol. Sci. 279:5039–5047. Royal Society.

Cresko, W. A., A. Amores, C. Wilson, J. Murphy, M. Currey, P. Phillips, M. A. Bell, C. B. Kimmel, and J. H. Postlethwait. 2004. Parallel genetic basis for repeated evolution of armor loss in Alaskan threespine stickleback populations. Proc. Natl. Acad. Sci. U. S. A. 101:6050–6055.

Depristo, M. A., E. Banks, R. Poplin, K. V Garimella, J. R. Maguire, C. Hartl, A. A. Philippakis, G. Del Angel, M. A. Rivas, M. Hanna, A. McKenna, T. J. Fennell, A. M. Kernytsky, A. Y. Sivachenko, K. Cibulskis, S. B. Gabriel, D. Altshuler, and M. J. Daly. 2011. A framework for variation discovery and genotyping using next-generation DNA sequencing data. Nat. Genet. 43:491–501.

Dingerkus, G., and L. D. Uhler. 1977. Stain Technology Enzyme Clearing of Alcian Blue Stained Whole Small Vertebrates for Demonstration of Cartilage. , doi: 10.3109/10520297709116780.

Donley, J. M., C. A. Sepulveda, P. Konstantinidis, S. Gemballa, and R. E. Shadwick. 2004. Convergent evolution in mechanical design of lamnid sharks and tunas. Nature 429:61–65.

Echelle, A. A., and A. E. Echelle. 2020. Cyprinodontidae: Pupfishes. Pp. 609–673 *in* Freshwater Fishes of North America. The Johns Hopkins University Press.

Erickson, P. A., A. M. Glazer, E. E. Killingbeck, R. M. Agoglia, J. Baek, S. M. Carsanaro, A. M. Lee, P. A. Cleves, D. Schluter, and C. T. Miller. 2016. Partially repeatable genetic basis of benthic adaptation in threespine sticklebacks. Evolution (N. Y). 70:887–902.

Feldman, C. R., E. D. Brodie, E. D. Brodie, and M. E. Pfrender. 2012. Constraint shapes

convergence in tetrodotoxinresistant sodium channels of snakes. Proc. Natl. Acad. Sci. U. S. A. 109:4556–4561.

Feldman, C. R., E. D. Brodie, E. D. Brodie, and M. E. Pfrender. 2009. The evolutionary origins of beneficial alleles during the repeated adaptation of garter snakes to deadly prey. Proc. Natl. Acad. Sci. U. S. A. 106:13415–13420.

Feulner, P. G. D., F. J. J. Chain, M. Panchal, C. Eizaguirre, M. Kalbe, T. L. Lenz, M. Mundry, I. E. Samonte, M. Stoll, M. Milinski, T. B. H. Reusch, and E. Bornberg-Bauer. 2013. Genome-wide patterns of standing genetic variation in a marine population of three-spined sticklebacks. Mol. Ecol. 22:635–649.

Futuyman, D. J. 1986. Evolutionary Biology. 2d ed. Sinauer, Sunderland, Mass.

Gardner, K. M., and R. G. Latta. 2007. Shared quantitative trait loci underlying the genetic correlation between continuous traits. Mol. Ecol. 16:4195–4209. John Wiley & Sons, Ltd.

Grant, P. R., B. R. Grant, J. A. Markert, L. F. Keller, and K. Petren. 2004. Convergent evolution of Darwin's finches caused by introgressive hybridization and selection. Evolution (N. Y). 58:1588–1599. Society for the Study of Evolution.

Haenel, Q., M. Roesti, D. Moser, A. D. C. MacColl, and D. Berner. 2019. Predictable genome-wide sorting of standing genetic variation during parallel adaptation to basic versus acidic environments in stickleback fish. Evol. Lett. 3:28–42. Wiley.

Hagey, F. M., and J. E. Mylroie. 1995. Pleistocene lake and lagoon deposits, San Salvador island, Bahamas. Spec. Pap. Soc. Am. 77–90.

Haley, C. S., and S. A. Knott. 1992. A simple regression method for mapping quantitative trait loci in line crosses using flanking markers. Heredity (Edinb). 69:315–324.

Hall, B. D., R. G. Cadle, S. M. Morrill-Cornelius, and C. A. Bay. 2007. Phakomatosis pigmentovascularis: Implications for severity with special reference to Mongolian spots associated with Sturge-Weber and Klippel-Trenaunay syndromes. Pp. 3047–3053 *in* American Journal of Medical Genetics, Part A. John Wiley & Sons, Ltd.

Hedrick, P. W. 2013. Adaptive introgression in animals: Examples and comparison to new mutation and standing variation as sources of adaptive variation. John Wiley & Sons, Ltd.

Hedrick, T. L. 2008. Software techniques for two- and three-dimensional kinematic measurements of biological and biomimetic systems. Bioinspiration and Biomimetics 3.

Hernandez, L. P., D. Adriaens, C. H. Martin, P. C. Wainwright, B. Masschaele, and M. Dierick. 2018. Building trophic specializations that result in substantial niche partitioning within a young adaptive radiation. J. Anat. 232:173–185.

Hopkins, C. D. 1995. Convergent designs for electrogenesis and electroreception. Curr. Opin. Neurobiol. 5:769–777.

James, M. E., M. J. Wilkinson, H. L. North, J. Engelstädter, and D. Ortiz-Barrientos. 2020. A framework to quantify phenotypic and genotypic parallel evolution. bioRxiv, doi: 10.1101/2020.02.05.936450.

Jones, F. C., M. G. Grabherr, Y. F. Chan, P. Russell, E. Mauceli, J. Johnson, R. Swofford, M. Pirun, M. C. Zody, S. White, E. Birney, S. Searle, J. Schmutz, J. Grimwood, M. C. Dickson, R. M. Myers, C. T. Miller, B. R. Summers, A. K. Knecht, S. D. Brady, H. Zhang, A. A. Pollen, T. Howes, C. Amemiya, J. Baldwin, T. Bloom, D. B. Jaffe, R. Nicol, J. Wilkinson, E. S. Lander, F. Di Palma, K. Lindblad-Toh, and D. M. Kingsley. 2012. The genomic basis of adaptive evolution in threespine sticklebacks. Nature 484:55–61.

Jost, M. C., D. M. Hillis, Y. Lu, J. W. Kyle, H. A. Fozzard, and H. H. Zakon. 2008. Toxin-resistant sodium channels: Parallel adaptive evolution across a complete gene family. Mol.

Biol. Evol. 25:1016–1024. Oxford Academic.

Juenger, T. E., S. Sen, K. A. Stowe, and E. L. Simms. 2005. Epistasis and genotype-environment interaction for quantitative trait loci affecting flowering time in Arabidopsis thaliana. Pp. 87–105 *in* Genetica.

Katz, P. S. 2006. Comparative Neurophysiology: An Electric Convergence in Fish.

Kowalko, J. E., N. Rohner, T. A. Linden, S. B. Rompani, W. C. Warren, R. Borowsky, C. J. Tabin, W. R. Jeffery, and M. Yoshizawa. 2013. Convergence in feeding posture occurs through different genetic loci in independently evolved cave populations of Astyanax mexicanus. Proc. Natl. Acad. Sci. U. S. A. 110:16933–16938.

Lander, E., and L. Kruglyak. 1995. Genetic dissection of complex traits: Guidelines for interpreting and reporting linkage results. Nat. Genet. 11:241–247.

Lander, E. S., and S. Botstein. 1989. Mapping mendelian factors underlying quantitative traits using RFLP linkage maps. Genetics 121:185.

Linnen, C. R., Y. P. Poh, B. K. Peterson, R. D. H. Barrett, J. G. Larson, J. D. Jensen, and H. E. Hoekstra. 2013. Adaptive evolution of multiple traits through multiple mutations at a single gene. Science (80-.). 339:1312–1316. American Association for the Advancement of Science.

Liu, J., M. Aoki, I. Illa, C. Wu, M. Fardeau, C. Angelini, C. Serrano, J. Andoni Urtizberea, F. Hentati, M. Ben Hamida, S. Bohlega, E. J. Culper, A. A. Amato, K. Bossie, J. Oeltjen, K. Bejaoui, D. McKenna-Yasek, B. A. Hosler, E. Schurr, K. Arahata, P. J. De Jong, and R. H. Brown. 1998. Dysferlin, a novel skeletal muscle gene, is mutated in Miyoshi myopathy and limb girdle muscular dystrophy. Nat. Genet. 20:31–36.

Liu, R. K., and A. A. Echelle. 2013. Behavior of the Catarina pupfish (cyprinodontidae: Megupsilon aporus), a severely imperiled species. Southwest. Nat. 58:1–7. Southwestern Association of Naturalists.

Losos, J. B. 2009. Lizards in an Evolutionary Tree: Ecology and Adaptive Radiation of Anoles. University of California Press, Berkeley.

Lynch, M., and B. Walsh. 1998. Genetics and Analysis of Quantitative Traits. Vol. 1. Sinauer Associates, Sunderland, MA, Sunderland, MA.

Marsden, C. D., Y. Lee, K. Kreppel, A. Weakley, A. Cornel, H. M. Ferguson, E. Eskin, and G. C. Lanzaro. 2014. Diversity, differentiation, and linkage disequilibrium: Prospects for association mapping in the malaria vector anopheles arabiensis. G3 Genes, Genomes, Genet. 4:121–131. Genetics Society of America.

Martin, C. H., J. E. Crawford, B. J. Turner, and L. H. Simons. 2016. Diabolical survival in Death Valley: recent pupfish colonization, gene flow and genetic assimilation in the smallest species range on earth. Proc. R. Soc. B Biol. Sci. 283:20152334.

Martin, C. H., J. S. Cutler, J. P. Friel, C. Dening Touokong, G. Coop, and P. C. Wainwright. 2015. Complex histories of repeated gene flow in Cameroon crater lake cichlids cast doubt on one of the clearest examples of sympatric speciation. Evolution (N. Y). 69:1406–1422.

Martin, C. H., P. A. Erickson, and C. T. Miller. 2017. The genetic architecture of novel trophic specialists: larger effect sizes are associated with exceptional oral jaw diversification in a pupfish adaptive radiation. Mol. Ecol. 26:624–638.

Martin, C. H., and L. C. Feinstein. 2014. Novel trophic niches drive variable progress towards ecological speciation within an adaptive radiation of pupfishes. Mol. Ecol. 23:1846–1862.

Martin, C. H., and P. C. Wainwright. 2013. A Remarkable Species Flock of Cyprinodon Pupfishes Endemic to San Salvador Island, Bahamas. Bull. Peabody Museum Nat. Hist.

54:231–241.

McGirr, J. A., and C. H. Martin. 2020. Ecological divergence in sympatry causes gene misexpression in hybrids. Mol. Ecol. 29:2707–2721.

McGirr, J. A., and C. H. Martin. 2021. Few Fixed Variants between Trophic Specialist Pupfish Species Reveal Candidate Cis-Regulatory Alleles Underlying Rapid Craniofacial Divergence. Mol. Biol. Evol. 38:405–423.

McGirr, J. A., and C. H. Martin. 2016. Novel candidate genes underlying extreme trophic specialization in Caribbean pupfishes. Mol. Biol. Evol. 34:msw286. Narnia.

Morjan, C. L., and L. H. Rieseberg. 2004. How species evolve collectively: implications of gene flow and selection for the spread of advantageous alleles. Mol. Ecol. 13:1341–1356. John Wiley & Sons, Ltd.

Nelson, T. C., and W. A. Cresko. 2018. Ancient genomic variation underlies repeated ecological adaptation in young stickleback populations. Evol. Lett. 2:9–21.

Nyholt, D. R. 2000. All LODs are not created equal. Am. J. Hum. Genet. 67:282–288. University of Chicago Press.

O'brown, N. M., B. R. Summers, F. C. Jones, S. D. Brady, and D. M. Kingsley. 2015. A recurrent regulatory change underlying altered expression and Wnt response of the stickleback armor plates gene EDA. Elife 2015.

Orr, H. A. 2006. The distribution of fitness effects among beneficial mutations in Fisher's geometric model of adaptation. J. Theor. Biol. 238:279–285.

Paaby, A. B., and M. V. Rockman. 2013. The many faces of pleiotropy. Elsevier Current Trends.

Parsons, T. E., S. M. Weinberg, K. Khaksarfard, R. N. Howie, M. Elsalanty, J. C. Yu, and J. J. Cray. 2014. Craniofacial shape variation in Twist1+/- mutant mice. Anat. Rec. 297:826–833. Blackwell Publishing Inc.

Pavlidis, P., D. Živković, A. Stamatakis, and N. Alachiotis. 2013. SweeD: Likelihood-based detection of selective sweeps in thousands of genomes. Mol. Biol. Evol. 30:2224–2234. Mol Biol Evol.

Richards, E. J., and C. H. Martin. 2017. Adaptive introgression from distant Caribbean islands contributed to the diversification of a microendemic adaptive radiation of trophic specialist pupfishes. PLoS Genet. 13:e1006919.

Richards, E. J., J. A. McGirr, J. R. Wang, M. E. St. John, J. W. Poelstra, M. J. Solano, D. C. O'Connell, B. J. Turner, and C. H. Martin. 2021. A vertebrate adaptive radiation is assembled from an ancient and disjunct spatiotemporal landscape. Proc. Natl. Acad. Sci. 118:e2011811118. Proceedings of the National Academy of Sciences.

Risch, N. 1990. Linkage strategies for genetically complex traits. I. Multilocus models. Am. J. Hum. Genet. 46:222–228.

Rosenblum, E. B., C. E. Parent, and E. E. Brandt. 2014. The molecular basis of phenotypic convergence. Annu. Rev. Ecol. Evol. Syst. 45:203–226.

Russello, M. A., M. D. Waterhouse, P. D. Etter, and E. A. Johnson. 2015. From promise to practice: Pairing non-invasive sampling with genomics in conservation. PeerJ 2015:e1106. PeerJ Inc.

Schluter, D., E. A. Clifford, M. Nemethy, and J. S. McKinnon. 2004. Parallel evolution and inheritance of quantitative traits. Am. Nat. 163:809–822. The University of Chicago Press.

Sen, Ś., J. M. Satagopan, K. W. Broman, and G. A. Churchill. 2007. R/qtlDesign: Inbred line cross experimental design. Mamm. Genome 18:87–93.

Shirley, M. D., H. Tang, C. J. Gallione, J. D. Baugher, L. P. Frelin, B. Cohen, P. E. North, D. A.

Marchuk, A. M. Comi, and J. Pevsner. 2013. Sturge–Weber Syndrome and Port-Wine Stains Caused by Somatic Mutation in GNAQ . N. Engl. J. Med. 368:1971–1979. New England Journal of Medicine (NEJM/MMS).

St. John, M. E., K. Dixon, and C. H. Martin. 2020a. Oral shelling within an adaptive radiation of pupfishes: testing the adaptive function of novel nasal protrusion and behavioral preference. J. Fish Biol. 1–9.

St. John, M. E., R. Holzman, and C. H. Martin. 2020b. Rapid adaptive evolution of scale-eating kinematics to a novel ecological niche. J. Exp. Biol. jeb.217570.

Stern, D. L. 2013. The genetic causes of convergent evolution. Nat. Rev. Genet. 14:751–764. Nature Publishing Group.

Stern, D. L., and V. Orgogozo. 2008. The loci of evolution: How predictable is genetic evolution?

Stevenson, M. M. 1981. Karyomorphology of Several Species of Cyprinodon. Copeia 1981:494. JSTOR.

Stuart, Y. E., T. Veen, J. N. Weber, D. Hanson, M. Ravinet, B. K. Lohman, C. J. Thompson, T. Tasneem, A. Doggett, R. Izen, N. Ahmed, R. D. H. Barrett, A. P. Hendry, C. L. Peichel, and D. I. Bolnick. 2017. Contrasting effects of environment and genetics generate a continuum of parallel evolution. Nat. Ecol. Evol. 1:158. Nature Publishing Group.

Taylor, J., and D. Butler. 2017. R package ASMap: Efficient genetic linkage map construction and diagnosis. J. Stat. Softw., doi: 10.18637/jss.v079.i06.

Taylor, R. S., M. Manseau, R. L. Horn, S. Keobouasone, G. B. Golding, and P. J. Wilson. 2020. The role of introgression and ecotypic parallelism in delineating intraspecific conservation units. Mol. Ecol. 29:2793–2809. Blackwell Publishing Ltd.

Teng, C. S., M. C. Ting, D. T. Farmer, M. Brockop, R. E. Maxson, and J. G. Crump. 2018. Altered bone growth dynamics prefigure craniosynostosis in a zebrafish model of Saethre-Chotzen syndrome. Elife 7. eLife Sciences Publications Ltd.

Thompson, K. A., M. M. Osmond, and D. Schluter. 2019. Parallel genetic evolution and speciation from standing variation. Evol. Lett. 3:129–141. Wiley.

Wagner, G. P., J. P. Kenney-Hunt, M. Pavlicev, J. R. Peck, D. Waxman, and J. M. Cheverud. 2008. Pleiotropic scaling of gene effects and the "cost of complexity." Nature 452:470–472. Nature Publishing Group.

Wittkopp, P. J., B. L. Williams, J. E. Selegue, and S. B. Carroll. 2003. Drosophila pigmentation evolution: Divergent genotypes underlying convergent phenotypes. Proc. Natl. Acad. Sci. U. S. A. 100:1808–1813. National Academy of Sciences.

Xie, K. T., G. Wang, A. C. Thompson, J. I. Wucherpfennig, T. E. Reimchen, A. D. C. MacColl, D. Schluter, M. A. Bell, K. M. Vasquez, and D. M. Kingsley. 2019. DNA fragility in the parallel evolution of pelvic reduction in stickleback fish. Science (80-.). 363:81–84.

Yang, J., N. A. Zaitlen, M. E. Goddard, P. M. Visscher, and A. L. Price. 2014. Advantages and pitfalls in the application of mixed-model association methods.

Yu, J., G. Pressoir, W. H. Briggs, I. V. Bi, M. Yamasaki, J. F. Doebley, M. D. McMullen, B. S. Gaut, D. M. Nielsen, J. B. Holland, S. Kresovich, and E. S. Buckler. 2006. A unified mixed-model method for association mapping that accounts for multiple levels of relatedness. Nat. Genet. 38:203–208. Nature Publishing Group.

Tables and Figures

Table 1. Maximum LOD scores for all 29 traits measured in Little Lake and Crescent Pond mapping crosses. A genome scan with a single-QTL model by Haley-Knott regression was used to identify the position with the highest LOD score, 95% Bayesian credible intervals, and the genome-wide significance level for each trait ($P < 0.1$: $\cdot$; $P < 0.05$: *). We also report the scaffold numbers of genomic regions that fell within the 95% credible intervals associated with the maximum LOD position for each trait, the number of individuals phenotyped, percent variance explained (PVE) by the max LOD region, and the uncorrected P-value associated with each max LOD region.

Trait	Population	Scaffold	Max LOD	Genome-wide Significance	n	PVE	x^2 P-Value
Lower Jaw Length	Crescent Pond	53, 7087, 2336, 6275, 26, 7335	2.89		205	6.29	0.0013
	Little Lake	24, 4028, 58, 16	3.30		228	6.45	0.0005
Jaw closing In-Lever	Crescent Pond	31, 4, 451, 19	3.60		204	7.81	0.0002
	Little Lake	8, 9588, 8020	4.11	$\cdot$	227	7.99	0.0001
Jaw Opening In-Lever	Crescent Pond	6086, 11	2.43		205	5.32	0.0037
	Little Lake	43	2.98		227	5.87	0.0010
Palatine Height	Crescent Pond	34, 22, 6304	2.90		205	6.31	0.0013
	Little Lake	11	2.73		228	5.36	0.0019
Suspensorium Length	Crescent Pond	46, 37, 31, 26, 60, 7556, 10198, 22	3.54		204	7.68	0.0003
	Little Lake	11	3.51		227	6.88	0.0003
Dentigerous Arm Width	Crescent Pond	52, 13137, 40	2.19		202	4.87	0.0065
	Little Lake	24, 4028, 58, 16	4.05	$\cdot$ *	228	7.85	0.0001
Maxilla Length	Crescent Pond	27, 593, 4, 31, 451, 19	2.67		204	5.85	0.0021
	Little Lake	56	3.03		228	5.94	0.0009
Dentigerous Arm Base	Crescent Pond	27, 593, 4, 31, 451, 19	2.98		205	6.47	0.0011
	Little Lake	26	3.70		228	7.21	0.0002

Measurement	Population						
Dentigerous Arm Depth	Crescent Pond	6086, 11, 46	4.20	.*	205	9.00	0.0001
	Little Lake	5	3.70	.	217	7.55	0.0002
Ascending Process Length	Crescent Pond	27, 593, 4, 31, 451, 19	2.70		201	6.00	0.0020
	Little Lake	46, 37	3.70		210	7.79	0.0002
Maxillary Head Height	Crescent Pond	6086, 11, 46	2.33		205	5.11	0.0046
	Little Lake	7, 30	2.19		228	4.33	0.0064
Ectopterygoid	Crescent Pond	14, 9, 16, 5405, 11419	2.81		205	6.11	0.0016
	Little Lake	9	3.36		228	6.56	0.0004
Maxillary Head Protrusion	Crescent Pond	58, 24, 41, 47	2.70		205	5.88	0.0020
	Little Lake	7431, 53, 6275, 2336, 25	4.03	.*	228	7.82	0.0001
Nasal Tissue Protrusion	Crescent Pond	46, 37, 31, 26, 60, 7556, 10198, 22	2.25		205	4.93	0.0056
	Little Lake	9	3.69		228	7.18	0.0002
Orbit Diameter	Crescent Pond	9588, 8, 8020	2.34		205	5.13	0.0045
	Little Lake	52, 40, 41	2.58		227	5.10	0.0026
Cranial Height	Crescent Pond	33, 39	3.59	.	205	7.74	0.0003
	Little Lake	33	3.94	.	224	7.78	0.0001
Head Depth	Crescent Pond	16, 40	2.98		204	6.51	0.0010
	Little Lake	52, 40, 41	2.71		223	5.45	0.0019
Pelvic Girdle Length	Crescent Pond	31, 18, 15, 11057, 55, 52	2.68		203	5.90	0.0021
	Little Lake	27, 37, 7	2.87		226	5.68	0.0014
Premaxilla Pelvic Girdle	Crescent Pond	37, 46, 7556, 10198	3.15		202	6.92	0.0007
	Little Lake	35, 38, 20, 8508, 10278, 33	2.63		231	5.10	0.0024
Standard	Crescent	14, 9, 16, 5405,	2.90		204	6.34	0.0013

Length (mm)	Pond	11419					
	Little Lake	31, 46, 37	3.48		231	6.69	0.0003
Cranium Dorsal Fin	Crescent Pond	6704, 52, 13137, 40	2.84		205	6.18	0.0014
	Little Lake	37, 22, 7556	3.45		231	6.65	0.0004
Dorsal Fin Width	Crescent Pond	43, 26, 14743	2.18		205	4.78	0.0066
	Little Lake	842, 44, 1074, 6, 30	3.00		230	5.83	0.0010
Dorsal Fin Height	Crescent Pond	18, 31, 15, 11057, 55	2.84		203	6.23	0.0015
	Little Lake	43	3.50		222	7.00	0.0003
Anterior Body Depth	Crescent Pond	8, 8020	2.94		204	6.43	0.0011
	Little Lake	6094, 5, 4	3.33		230	6.45	0.0005
Posterior Body Depth	Crescent Pond	20, 471, 39, 8508, 33	2.86		203	6.27	0.0014
	Little Lake	18, 15	3.02		228	5.92	0.0009
Caudal Peduncle Length	Crescent Pond	31, 18, 15, 11057, 55, 52	2.87		203	6.30	0.0014
	Little Lake	24, 4028, 58, 16	2.16		230	4.23	0.0070
Anal Fin Width	Crescent Pond	18, 15, 11057, 55	2.89		201	6.41	0.0013
	Little Lake	6, 842, 44, 1074, 30	2.40		229	4.71	0.0040
Anal Fin Height	Crescent Pond	43, 26, 14743	2.93		201	6.48	0.0012
	Little Lake	8, 9588, 8020	3.15		229	6.14	0.0007
Caudal Peduncle Height	Crescent Pond	53, 7087, 2336, 6275, 26, 7335	1.97		205	4.32	0.0108
	Little Lake	47, 1962	3.32		230	6.44	0.0005
Adductor	Crescent Pond	6086, 11	3.56	•	170	9.18	0.0003
	Little Lake	-	-	-	-	-	-
Proportion Time Spent Near Scale-Eater Mates	Crescent Pond	58, 24, 41, 47	2.05		74	12.00	0.0089
	Little Lake	-	-	-	-	-	-

Table 2. Position of maximum LOD score and 95% credible intervals for each trait in the Little Lake linkage map and the Crescent Pond linkage map. Colors represent corresponding linkage groups across lakes. Asterisks represent traits that were marginally significant at the *P* < 0.1 level in the genome scan.

Little Lake						Crescent Pond				
				Position genomewide						Position genomewide
	Trait	Sig.	LG	Max LOD score	95% CI		Trait	Sig.	LG	Max LOD score	95% CI
	Cranial Height	*	1	259	(250,270)		Suspensorium Length		1	566	(20,730)
	Premaxilla to Pelvic Girdle		1	146	(0,350)		Nasal Tissue Protrusion		1	570	(0,740)
	Cranium to Dorsal Fin		2	303	(160,380)		Premaxilla to Pelvic Girdle		1	568	(310,600)
	Lower Jaw Length		3	9	(0,340)		Ectopterygoid		3	272	(0,560)
	Dentigerous Arm Width	*	3	9	(0,340)		Standard Length (mm)		3	50	(40,500)
	Caudal Peduncle Length		3	168	(0,340)		Dentigerous Arm Width		4	317	(40,510)
	Dorsal Find width		4	187	(10,310)		Cranium to Dorsal Fin		4	89	(30,510)
	Anal Fin Width		4	14	(0,280)		Lower Jaw Length		5	136	(0,470)
	Dentigerous Arm Depth	*	6	79	(20,90)		Caudal Peduncle height		5	381	(0,470)
	Anterior Body Depth		6	289	(0,300)		Jaw Closing In-Lever		6	380	(150,410)
	Orbit Diameter		8	266	(0,290)		Maxilla Length		6	468	(0,480)
	Head Depth		8	206	(0,290)		Dentigerous Arm Base		6	107	(0,480)
	Jaw Closing In-Lever	*	9	54	(40,90)		Ascending Proces Length		6	106	(0,470)
	Anal Fin Height		9	100	(70,240)		Pelvic Girdle Length		8	370	(20,435)
	Maxillary Head Protrusion	*	10	35	(0,260)		Dorsal Fin Height		8	91	(0,380)
	Ascending Process Length		12	119	(90,150)		Caudal Peduncle Length		8	258	(30,425)
	Standard Length (mm)		12	200	(50,210)		Anal Fin Width		8	190	(110,400)
	Ectopterygoid		13	170	(150,180)		Maxillary Head Protrusion		9	300	(0,350)
	Nasal Tissue Protrusion		13	193	(20,200)		Proportion Time Spent Near Scale-Eater Males		9	166	(50,340)
	Palatine Height		14	147	(110,210)		Cranial Height	*	10	204	(130,340)
	Suspensorium Length		14	153	(70,180)		Posterior Body Depth		10	270	(0,330)
	Jaw Opening In-Lever		16	58	(40,140)		Palatine Height		11	70	(0,310)
	Dorsal Fin Height		16	52	(40,60)		Head Depth		12	111	(100,280)
	Pelvic Girdle Length		17	50	(10,160)		Opening In-Lever		13	10	(0,90)
	Maxillary Head Height		18	122	(30,160)		Dentigerous Arm Depth	*	13	2	(0,250)
	Caudal Peduncle Height		19	44	(20,90)		Maxillary Head Height		13	170	(0,280)
	Dentigerous Arm Base		21	74	(0,100)		Adductor Mandibulae Mass	*	13	2	(0,70)
	Maxilla Length		22	40	(20,50)		Dorsal Fin Width		14	305	(30,330)
	Posterior Body Depth		24	30	(10,30)		Anal Fin Width		14	330	(280,330)
							Orbit Diameter		16	107	(0,190)
							Anterior Body Depth		16	170	(10,220)

Table 3. Number of adaptive alleles and any genes within 20 kbp found in trait QTL with maximum LOD scores for both lakes. Adaptive alleles were categorized as either standing genetic variation (SGV), introgression (Intro.), or de novo mutations, and were estimated independently for snail-eaters and scale-eaters in a previous study (Richards et al. 2021). Asterisks represent traits that were significant at the P < 0.1 level in the genome-wide scan, while crosses show traits that corresponded to the same locations in the alternate lake.

Traits	Gene	Snail-Eater		Scale-Eater		
		SGV	Intro.	SGV	Intro.	de novo
Cranial Height*	*bri3bp*	-	26	28	-	-
	gnaq	9	-	9	-	-
	wdr31	18	2	20	-	-
	Unannotated Regions	1	-	11	-	-
Dentigerous Arm Width* **Female mate preference†** **Maxillary Head Protrusion†**	*cyp26b1*	-	8	8	-	-
	dysf	-	-	1	-	-
	Unannotated Regions	-	67	216	-	1
Dentigerous Arm Depth*	Unannotated Regions	-	-	1	-	-
Maxillary Head Protrusion* **Lower Jaw Length†** **Caudal Peduncle Height†**	*cox6b1*	8	-	8	-	-
	cyp21a2	-	-	2	-	-
	eva1b	-	-	2	-	-
	fhod3	-	-	2	-	-
	galnt1	-	-	-	17	-
	glipr2	-	-	3	-	-
	hdac9b	-	-	-	1	-
	mag	-	-	2	-	-
	map7d1	25	-	25	-	-
	mindy3	-	-	8	-	-
	nacad	-	-	2	-	-
	pxn1	-	-	1	-	-
	rasip1	13	-	13	-	-
	slc2a3	15	-	15	-	-
	steap4	-	-	-	26	-
	tbrg4	-	-	2	-	-
	them4	-	-	5	-	-
	tnc	-	-	1	-	-
	twist1	-	-	-	-	1

Trait	Gene					
	zhx2	5	-	6	-	-
	znf628	5	-	6	-	-
	Unannotated Regions	29	68	93	64	-
Jaw closing In-Lever* **Orbit Diameter†** **Anterior Body Depth†**	*galr2*	-	-	-	2	-
	map2k6	-	-	-	3	-
Dentigerous Arm Depth* **Adductor Mandibulae Masst*** **Palatine Height†** **Suspensorium Length†**	*atp8a2*	92	-	92	-	-
	cd226	6	-	6	-	1
	cdk8	-	-	1	-	-
	cmbl	-	-	4	-	7
	crispld1	-	-	7	-	-
	dok6	-	-	50	-	-
	fbxl7	-	-	6	-	-
	hnf4g	-	-	1	-	-
	med1	-	-	26	-	-
	mtrr	-	-	2	-	-
	ncoa2	7	-	-	4	-
	prlh	-	-	12	6	-
	rnf6	-	-	4	-	-
	shisa2	18	-	38	-	-
	slc51a	-	-	22	-	7
	spice1	4	-	2	-	-
	ube2w	-	48	-	-	-
	zfhx4	-	-	-	-	1
	Unannotated Regions	34	34	131	3	1

Figure 1 A) Representative photographs of F2 intercross cleared and double-stained specimen used for skeletal morphometrics. Points represent landmarks used to measure linear distances between skeletal traits. B) Table containing the two landmarks that correspond to each trait.

A)

B)

Head			Body		
Point 1	Point 2	Trait	Point 1	Point 2	Trait
1	2	Lower Jaw Length	20	21	Premaxilla to Pelvic Girdle
2	3	Jaw closing In-Lever	20	31	Standard Length
2	4	Jaw Opening In-Lever	22	23	Cranium to Dorsal Fin
2	11	Palatine Height	23	24	Dorsal Fin Width
2	18	Suspensorium Length	23	25	Dorsal Fin Height
5	8	Dentigerous Arm Width	23	26	Anterior Body Depth
6	11	Maxilla Length	24	27	Posterior Body Depth
7	5	Dentigerous Arm Base	24	30	Caudal Peduncle Length
8	9	Dentigerous Arm Depth	26	27	Anal Fin Width
9	10	Ascending Process Length	27	28	Anal Fin Height
11	13	Maxillary Head Height	29	30	Caudal Peduncle Height
11	14	Ectopterygoid			
12	13	Maxillary Head Protrusion			
12	19	Nasal Tissue Protrusion			
14	15	Orbit Diameter			
15	16	Cranial Height			
16	18	Head Depth			
17	18	Pelvic Girdle Length			

For the Crescent Pond individuals, we recorded sex and the mass of the adductor mandibulae muscle before clearing and staining each specimen.

Figure 2 Linkage maps for A) Crescent Pond and B) Little Lake crosses. The Crescent Pond linkage map was estimated from 743 markers and the Little Lake linkage map was estimated from 540 markers. Both maps were generated from crosses between a scale-eater (*C. desquamator*) and snail-eater (*C. brontotheroides*) from the respective lakes.

Figure 3 Circos plot depicting the relationship between the Crescent Pond (red) and Little La
linkage maps (blue), which share 324 markers within 10 kbp of one another. Numbers
surrounding each semi-circle represent linkage group numbers in each lake. Markers that are
shared across lakes are connected via the colored lines.

Figure 4 LOD profile for cranial height in Crescent Pond (red) and Little Lake (blue) F2 hybrids. LOD profiles were estimated by a Haley-Knott regression and are plotted relative to the position along the implicated linkage group (LG 10 for Crescent Pond, LG 1 for Little Lake) which are represented along the X-axis. Genome wide significance levels of P = 0.1, 0.05, and 0.01 are shown by the grey horizontal lines. Linkage groups along the X-axis also show the position of maximum LOD along with 95% Bayesian credible intervals. The orange fill color within the linkage groups corresponds to overlapping regions of scaffold 33 between crosses.

Figure 5 Cranial height phenotypes (size-corrected residuals) for each genotype in Crescent Pond (red) and Little Lake (blue). Both lakes show that heterozygotes (AB) exhibit greater cranial heights than homozygous parental genotypes.