Hybridization alters the shape of the genotypic fitness landscape, increasing access to novel fitness peaks during adaptive radiation

Austin H. Patton[1,2], Emilie J. Richards[1,2], Katelyn J. Gould[3], Logan K. Buie[3], Christopher H. Martin[1,2]

[1]*Department of Integrative Biology, University of California, Berkeley, CA*

[2]*Museum of Vertebrate Zoology, University of California, Berkeley, CA*

[3]*Department of Biology, University of North Carolina, Chapel Hill, NC*

Keywords: fitness landscape; genotypic fitness network; adaptive radiation; introgression; standing genetic variation; *de novo* mutation

Correspondence: austinhpatton@berkeley.edu, chmartin@berkeley.edu

Abstract

Estimating the complex relationship between fitness and genotype or phenotype (i.e. the adaptive

landscape) is one of the central goals of evolutionary biology. However, adaptive walks connecting

genotypes to organismal fitness, speciation, and novel ecological niches are still poorly understood

and processes for surmounting fitness valleys remain controversial. One outstanding system for

addressing these connections is a recent adaptive radiation of ecologically and morphologically

novel pupfishes (a generalist, molluscivore, and scale-eater) endemic to San Salvador Island,

Bahamas. We leveraged whole-genome sequencing of 139 hybrids from two independent field

fitness experiments to identify the genomic basis of fitness, estimate genotypic fitness networks,

and measure the accessibility of adaptive walks on the fitness landscape. We identified 132 SNPs

that were significantly associated with fitness in field enclosures. Six out of the 13 regions most

strongly associated with fitness contained differentially expressed genes and fixed SNPs between

trophic specialists; one gene (*mettl21e*) was also misexpressed in lab-reared hybrids, suggesting a

potential intrinsic genetic incompatibility. We then constructed genotypic fitness networks from

adaptive alleles and show that scale-eating specialists are the most isolated of the three species on

these networks. Intriguingly, introgressed and *de novo* variants reduced fitness landscape

ruggedness as compared to standing variation, increasing the accessibility of genotypic fitness

paths from generalist to specialists. Our results suggest that adaptive introgression and *de novo*

mutations alter the shape of the fitness landscape, providing key connections in adaptive walks

circumventing fitness valleys and triggering the evolution of novelty during adaptive radiation.

Introduction

First conceptualized by Sewell Wright in 1932, the adaptive landscape describes the complex

relationship between genotype or phenotype and fitness (1). The landscape is a concept, a

metaphor, and an empirical measurement that exerts substantial influence over all evolutionary

dynamics (2–6). Fitness landscapes were originally depicted as high-dimensional networks

spanning genotypic space in which each genotype is associated with fitness (1). Simpson (7)

later described phenotypic evolution of populations through time on a rugged landscape, in

which isolated clusters of fitness peaks represent 'adaptive zones' relative to adjacent regions of

low fitness (8). Lande and Arnold formalized the analysis of selection and estimation of

phenotypic fitness landscapes (9–11), leading to empirical studies of fitness landscapes in

numerous systems (12–18). Fitness surfaces are also central components of speciation models

and theory (19–21).

A central focus of fitness landscape theory is the characterization of the shape of the

fitness landscape. Theoretical and empirical studies frequently attempt to describe its

topography, such as quantifying the number of fitness peaks, one component of landscape

ruggedness that affects the predictability of evolution (5, 22–24). Importantly, the existence of

multiple peaks and valleys on the fitness landscape implies epistasis for fitness, or non-additive

effects on fitness resulting from genotypic interactions (23, 25–28). Fitness epistasis reduces the

predictability of evolution because the resultant increase in the number of peaks increases the

number of viable evolutionary outcomes (8, 29). Increasing fitness epistasis also increases

landscape ruggedness, thus reducing the probability of converging on any one fitness peak and

ultimately diversifying potential evolutionary outcomes (8, 30).

This leads to a fundamental concept in fitness landscape theory: Not all genotypic

pathways are evolutionarily accessible (5, 26, 31–36). In large populations, paths through

genotype space that monotonically increase in fitness at each mutational step are favored over

alternatives with neutral or deleterious steps (37). These accessible genotypic paths can be

considered adaptive walks under Fisher's geometric model, by which adaptation proceeds

towards a phenotypic optimum via additive mutations of small phenotypic effect (37, 38). On

rugged landscapes as originally envisioned by Wright (23), greater numbers of peaks (i.e. the

ruggedness) increase the mean length of potential adaptive walks to any one fitness optimum,

while decreasing the length of accessible paths to the nearest peak. Ultimately, this leads to a

decrease in the probability that any one fitness optimum is reached. Simultaneously, increasing

landscape ruggedness decreases the length of adaptive walks to the nearest local optimum, owing

to the corresponding increase in peak density.

There are a growing number of experimental studies of adaptive walks in nature,

including the evolution of toxin resistance in monarch butterflies (39), alcohol tolerance in

Drosophila (40, 41), and host-shift in aphids (42). Likewise, the accessibility of genotypic fitness

networks has now been explored in numerous microbial systems, including the evolution of

antibiotic resistance (31), metabolism (43), citrate exploitation (44), and glucose limitation in *E.*

coli (45), and adaptation to salinity in yeast via evolution of heat shock protein *Hsp90* (22).

However, these studies are still limited to the investigation of specific coding substitutions and

their effects on fitness in laboratory environments. Nosil et al. (46) estimated genotypic fitness

networks for *Timema* stick insects based on a field experiment. Similarly, this study focused on a

single large-effect locus underlying dimorphic coloration between ecotypes. These studies

represent significant advances, but extension of fitness landscape theory to empirical systems

including multiple species remains an underexplored area of future research at the intersection of

micro- and macroevolution. Such studies can provide insight into the topography of fitness

landscapes in natural systems, the accessibility of interspecific adaptive walks, and ultimately the

predictability of evolution.

One promising system for estimating fitness landscapes is a recent adaptive radiation of

Cyprinodon pupfishes endemic to San Salvador Island, Bahamas (17, 18, 47, 48). This radiation

is comprised of two trophic specialists, a molluscivore (durophage: *Cyprinodon brontotheroides*)

and a scale-eater (lepidophage: *C. desquamator*), derived from a Caribbean-wide generalist (*C.*

variegatus) which also coexists in the same habitats. These three species all occur in sympatry in

the hypersaline lakes of San Salvador Island, Bahamas (Fig 1a). Found in the benthic littoral

zone of each lake, all three species forage within the same benthic microhabitat; indeed, no

habitat segregation has been observed in 14 years of field studies. Originating less than 10,000

years ago (based on geological age estimates for the lakes: (49)), the functional and trophic

novelty harbored within this radiation is the product of exceptional rates of craniofacial

morphological evolution (50–53). Furthermore, species boundaries persist across multiple lake

populations, despite persistent admixture among species (54, 55). We previously estimated

fitness landscapes in these hypersaline lakes from two independent field experiments measuring

the growth and survival of hybrids placed in field enclosures (Figure 1b). Selection analyses

revealed a multi-peaked phenotypic fitness landscape that is stable across lake populations, year

of study, and manipulation of the frequency of rare hybrid phenotypes (17, 18, 48)). One of the

strongest and most persistent trends across studies and treatments was that hybrid phenotypes

resembling the scale-eater were isolated in the lowest fitness region for both growth and survival

relative to the other two species (17, 18). In contrast, hybrids resembling the generalist occupied

a fitness peak and were separated by a smaller fitness valley from hybrids resembling the

molluscivore, which occurred on a second peak of higher fitness.

107   Evolutionary trajectories through regions of low fitness should be inaccessible to natural

selection. How then did an ancestral generalist population cross these phenotypic fitness valleys

to reach new fitness peaks and adapt to novel ecological niches? A growing theoretical and

empirical literature on fitness landscapes has demonstrated the limited conditions for crossing

fitness valleys (56–59). Fitness peaks and valleys in morphospace may result only from the

reduction of the adaptive landscape to two phenotypic dimensions (60). Additional phenotypic

and genotypic dimensions may reveal fitness ridges that entirely circumvent fitness valleys (48,

61, 62). Indeed, owing to nonlinearity in the association between phenotype and fitness (63, 64),

even a single-peaked phenotypic fitness landscape may be underlaid by a multi-peaked genotypic

fitness landscape (65, 66). In this respect, investigating the high-dimensional genotypic fitness

landscape is key to understanding the origins of novelty in this system, particularly given the rare

evolution of lepidophagy (scale-eating), a niche occupied by less than 0.3% of all fishes (67).

119   Furthermore, the relative contributions of standing genetic variation, *de novo* mutations,

and adaptive introgression to the tempo and mode of evolution are now of central interest to the

field of speciation genomics (68–72). The three-dimensional adaptive landscape metaphor is

often invoked to explain how the genetic, phenotypic, and ecological diversity introduced to

populations by hybridization facilitates the colonization of neighboring fitness peaks that are

unoccupied by either hybridizing species (73–75). However, extension of these ideas to more

high-dimensional genotypic fitness landscapes remains underexplored. For instance, we have yet

to learn how the appearance of novel adaptive genetic variation through introgressive

hybridization or *de novo* mutation alters the realized epistatic interactions among loci, thus

potentially altering the shape of the fitness landscape and the accessibility of interspecific

adaptive walks.

130
Figure 1. San Salvador Island pupfishes and their hybrids. a. From top to bottom: the generalist,
Cyprinodon variegatus, the molluscivore *C. brontotheroides*, and the scale-eater *C. desquamator*. **b.**
Representative images of experimental field enclosures. **c.** Principal component analysis of 1,129,771 LD-
pruned SNPs genotyped in hybrids and the three parental species. **d.** Unsupervised ADMIXTURE analyses
for Crescent Pond (top) and Little Lake (bottom). G, M and S indicate individual samples of Generalists
(G), Molluscivores (M), and Scale-eaters (S), respectively, followed by all resequenced hybrid individuals
from field experiments. Colors indicate ancestry proportions in each population (K = 3).

The adaptive radiation of San Salvador Island pupfishes, like many others (76–80),

appears to have originated from a complex interplay of abundant standing genetic variation,

adaptive introgression from neighboring islands, and several *de novo* single-nucleotide mutations

and deletions found only in the scale-eater (55, 81). Notably, both specialists harbor numerous

introgressed SNPs showing evidence of hard selective sweeps in the regulatory regions of known

craniofacial genes (55, 81). In contrast, hard selective sweeps of *de novo* mutations only appear

in the scale-eating species, *C. desquamator*. Here, we leverage whole genome sequencing of 139

hybrids measured in field experiments to identify the genomic basis of fitness differences, infer

genotypic fitness networks, summarize their topography, and quantify the accessibility of novel

fitness peaks and the influence of each source of genetic variation on interspecific adaptive

walks.

Results

Sample collection and genomic resequencing

We resequenced 139 hybrids (86 survivors, 56 deaths (Appendix 1—table 1)) from two

independent field experiments across a total of six field enclosures and two lake populations

(2011: two high-density 3 m diameter enclosures exposed for three months: Crescent Pond $n =$

796; Little Lake $n = 875$ F2 hybrids (17); 2014/2015: four high-density 4 m diameter enclosures

exposed for three months in Crescent Pond, $n = 923$ F4/F5 hybrids and eleven months in Little

Lake, $n = 842$ F4/F5 hybrids (18)). We then characterized patterns of genetic variation among

parental species in each lake and their lab-reared hybrids used in field experiments. We

genotyped 1,129,771 SNPs with an average coverage of 9.79x per individual.

Population structure and ancestry associations with fitness

Principal components analysis (PCA) of genetic variation strongly differentiated pupfishes

sampled from Little Lake/Osprey Lake and Crescent Pond (PC1: 22.7% variance explained) and

among species within each lake (PC2: 15.9% variance explained: Figure 1d; Figure 1—figure

supplement 1-2). These results were supported by ADMIXTURE analyses (82, 83) (Figure 1e).

However, some hybrids were genotypically transgressive, falling outside the genotypic

distributions of the three parental species (Figure 1—figure supplement 2), leading

ADMIXTURE to assign the third cluster to these hybrids, rather than generalists which often

contain segregating variation found in trophic specialists (67). This pattern persisted in a

supervised ADMIXTURE analysis, in which we assigned individuals from the three parental

species *a priori* to their own population and estimated admixture proportions for the remaining

hybrids (Figure 1—figure supplement 3). Pairwise genetic distances did not predict pairwise

morphological distances (Figure 1—figure supplement 4).

We analyzed three measures of fitness (growth, survival, and their composite: see

Methods and Supplement for details), but focus herein on composite fitness, which is equal to

growth for survivors and zero for non-survivors. Growth could not be measured for tagged

hybrids that died in field enclosures and thus were not recovered. Because reproductive success

was not possible to quantify in field experiments (due to continuous egg-laying and very small,

newly hatched fry), composite fitness included only measurements of growth and survival.

Interestingly, in no case were genome-wide patterns of parental ancestry in hybrids

(estimated from unsupervised ADMIXTURE analyses) associated with hybrid composite fitness

(generalist $P = 0.385$; scale-eater $P = 0.439$; molluscivore $P = 0.195$), growth (generalist $P =$

0.119; scale-eater $P = 0.283$; molluscivore $P = 0.328$), or survival probability (generalist $P =$

0.440; scale-eater $P = 0.804$; molluscivore $P = 0.313$) while controlling for effects of lake and

experiment (Figure 1—figure supplement 5; Appendix 1—table 2). Similar results were obtained

when repeating these analyses using admixture proportions estimated from a supervised

ADMIXTURE analysis (Appendix 1—table 3), using only samples from the second field

experiment (Appendix 1—table 4), or using principal component axes estimated from genome-

wide SNPs (Appendix 1—table 5: See Supplementary Results). Therefore, in contrast to

previous studies (84–86), in this system genome-wide ancestry is not consistently associated

with fitness, highlighting the complex nonlinear relationship between genotype, phenotype, and

fitness within this nascent adaptive radiation. We must look to local ancestry to understand

fitness relationships (e.g. (87)).

194
Figure 2. The genetic basis of fitness variation and improved inference of adaptive landscapes. a) Per-
SNP log10 *p*-values from a genome-wide association test with GEMMA for composite fitness (survival x
growth). Lake and experiment were included as covariates in the linear mixed model. SNPs that were
significant at FDR < 0.05 are indicated in blue; red SNPs above dashed red line cross the threshold for
Bonferroni significance at α = 0.05. The first twenty-four scaffolds are sorted from largest to smallest and
remaining scaffolds were pooled. The six genes associated with composite fitness which were both strongly
differentiated ($F_{ST} > 0.95$) and differentially expressed between specialists (88) are annotated. **b-c)** Best-fit
adaptive landscape for composite fitness using either morphology alone (**b:** flat surface with only
directional selection) or morphology in combination with fitness-associated SNPs (**c:** highly nonlinear
surface). Best-fit model in **c** was a generalized additive model (GAM) including a thin-plate spline for both
LD axes, fixed effects of experiment and lake, and fixed effects of the seven (see supplementary methods)
SNPs most strongly associated with fitness shown in red in panel **a**. **d)** Three-dimensional view of **c** with
relative positions of the three parental phenotypes indicated.

Genome-wide association mapping of fitness

From our LD-pruned dataset we used a linear mixed model in GEMMA to identify 132 SNPs in

regions that were strongly associated with composite fitness, including 13 which remained

significant at the conservative Bonferroni-corrected threshold (Figure 2a, Appendix 1—table 6-

7; see supplement for results for survival and growth alone [Appendix 1—table 8-9; Figure 2—

figure supplement 1]). Gene ontologies for these 132 fitness-associated regions were

significantly enriched for synaptic signaling and chemical synaptic transmission [False discovery

rate (FDR) rate < 0.01; Figure 2—figure supplement 2; Appendix 1—table 7]. Ontologies

enriched at an FDR rate < 0.05 were related to signaling and regulation of cell communication

(for growth, see Figure 2—figure supplement 3). We did not identify any enrichment for

ontologies related to craniofacial development which have previously been identified to play a

significant role in the adaptive divergence of these fishes (55, 81, 88). This suggests that fitness-

associated regions in our field experiments captured additional components of fitness beyond the

external morphological phenotypes measured in previous studies.

We characterized whether genes in or near fitness-associated regions were implicated in

adaptive divergence of the specialists. Surprisingly, no fitness-associated regions overlapped

with regions showing significant evidence of a hard selective sweep (55). However, six fitness-

associated genes were previously shown to contain either fixed divergent SNPs (*csad, glcci1,*

ino80c, mag, pim2, mettl21e), or a fixed deletion between specialists (*med25*) (88). *Med25*

(Mediator Complex Subunit 25) is a craniofacial transcription factor associated with cleft palate

in humans and zebrafish (89, 90); a precursor of *mag* (Myelin Associated Glycoprotein) was also

associated with the parallel evolution of the thick-lipped phenotype in Midas cichlids based on

differential expression among morphs (91). Three of the six remaining fitness-associated genes

containing divergent SNPs (88) were associated with growth and/or body size measurements in

other fishes. First, *csad* plays an important role in synthesizing taurine which is a rate-limiting

enzyme affecting growth rate in parrotfishes (92), rainbow trout (93), and Japanese flounder

(94). Second, *glcci1* is associated with the body depth/length ratio in yellow croaker (95). Third,

ino80c is associated with measures of body size in Nile tilapia (96). Finally, *mettl21e* was

differentially expressed among specialists and also misexpressed in F1 hybrids between scale-

eaters and molluscivores at eight days post-fertilization and thus is a putative genetic

incompatibility in this system that may impact their fitness in field enclosures (88, 97). Although

it has not been associated with growth or body size in fishes, *mettl21e* is associated with

intramuscular fat deposition in cattle (98). Taken together, these findings support the

interpretation that fitness-associated regions are associated with unmeasured traits, particularly

physiological growth rate, or craniofacial shape in the case of the deletion in *med25*, that affect

fitness in our hybrid field experiments. However, the fitness associated loci we identified appear

not to have the subject of selective sweeps in either specialist.

Fitness-associated SNPs improve inference of the adaptive landscape

Fitness landscapes in past studies were estimated using slightly different sets of morphological

traits; thus, to enable inclusion of all hybrids on a single fitness landscape, a single observer

(AHP) remeasured all sequenced hybrids for 31 morphological trats (Figure 2—figure

supplement 4; Appendix 1—table 10). We used linear discriminant axes and generalized additive

modelling (GAM) to estimate phenotypic fitness landscapes for the sequenced hybrids on a two-

dimensional morphospace indicating similarity to each of the three parental populations

following previous studies (17, 18)(Figure 2—figure supplements 5; Appendix 1—table 11-13).

We then tested whether the inclusion of the 13 genomic regions most strongly associated with

fitness (red: Figure 2a) in GAM models improved our inference of the underlying adaptive

landscape. Models including fitness-associated SNPs were invariably favored over models with

external morphology alone (ΔAICc > 8.6: Appendix 1—table 14-15). Morphology-only models

predicted a flat fitness surface (Figure 2b, Figure 2—figure supplement 6; predictions restricted

to observed hybrid morphospace). In contrast, models including fitness-associated SNPs

predicted a complex and nonlinear fitness landscape, despite our limited dataset of 139

sequenced hybrids relative to samples in previous morphology-only studies of around 800

hybrids per enclosure.

To reduce complexity of the full model estimated from 31 morphological traits including

all 13 fitness-associated SNPs, we fit an additional model including only the seven most

significant fitness-associated SNPs in the full model. This reduced model was the best-fit; the

inferred adaptive landscape was complex and characterized by a fitness peak near hybrids

resembling the generalist phenotype separated by a small fitness valley from a second region of

high fitness for hybrids resembling the molluscivore phenotype. Hybrids resembling the scale-

eater phenotype again occurred in a large fitness valley (Figure 2b-2d: For results pertaining to

growth or survival see Appendix 1: Figure 2—figure supplement 6, Appendix 1—table 11-15).

Each of these fitness peaks and valleys were frequently recovered across 10,000 bootstrap

replicates; landscapes inferred from bootstrap replicates were often more complex with increased

curvature relative to inferences from our observed dataset (Figure 2—figure supplement 7).

Thus, the fitness landscape estimated from our observed dataset appears robust to sampling

uncertainty.

276
Figure 3. Scale-eaters are isolated on the fitness landscape. a. Most nearly fixed or fixed variants (F_{ST}
≥ 0.95) experiencing hard selective sweeps (hereafter 'adaptive alleles') originated as standing genetic
variation (SGV: molluscivores = 96%, scale-eaters = 92%), followed by introgression (molluscivores =
4%, scale-eaters = 6%), and *de novo* mutation (scale-eaters = 2%)(55). Pie charts show adaptive alleles
retained in our study for each species; networks are constructed from either set of adaptive alleles. **b.**
Genotypic network constructed from a random sample of ten SNPs, sampled from all SNPs shown in **a**.
Each edge between nodes is up five mutational steps away; edge width is proportional to mutational
distance: wider edges connect closer haplotypes; hybrid node size is proportional to fitness (larger nodes
are of greater fitness value). **c.** Median number of mutational steps within or between species (e.g. Figure
4a). All pairwise comparisons using Tukey's HSD test (after FDR correction) were significant.

Compared to previous studies, the highest fitness optimum was shifted from the

molluscivore to the generalist phenotype. This suggests that fitness-associated SNPs increased

the fitness of hybrids resembling generalists beyond expectations based on their morphology

alone, consistent with the hypothesis that fitness-associated SNPs are associated with

unmeasured non-morphological traits affecting fitness. Indeed, visualization of observed

haplotypes in hybrids across the fitness landscape supported this interpretation; one of the most

common haplotypes was most frequent in hybrids resembling generalists near the peak of high

fitness and rare in hybrids resembling either trophic specialist (Figure 2—figure supplement 8).

Regardless, this two-dimensional phenotypic fitness landscape did not reveal fitness ridges

connecting generalists to specialists, further emphasizing the need to investigate the genotypic

fitness landscape.

Trophic novelty is associated with isolation on the genotypic fitness network

The adaptive radiation of pupfishes on San Salvador island originated within the last 10,000

years through a combination of selection on standing genetic variation, adaptive introgression,

and *de novo* mutations (55). However, it is unclear how each source of genetic variation aided in

the traversal of fitness paths or contributed to the colonization of novel fitness peaks. To address

this knowledge gap, we first sought to visualize genotypic fitness networks and gain insight into

how isolated the three species are in genotypic space. Understanding the relative isolation of

each specialist from the generalist can reveal the relative accessibility of their respective adaptive

walks on the genotypic fitness landscape.

To accomplish this, we reconstructed genotypic fitness networks from 1,498 candidate

adaptive alleles previously identified in this system (e.g. Figure 3a (55)). These regions displayed

significant evidence of a hard selective sweep using both site frequency spectrum and LD-based

methods, SweeD (99) and OmegaPlus (100), and contained fixed or nearly fixed SNPs ($F_{ST} >$

0.95) differentiating trophic specialists across lakes (55). Adaptive alleles were classified as

standing variation, introgressed, or *de novo* mutations based on extensive sampling of focal and

related *Cyprinodon* pupfish species across San Salvador Island and neighboring Caribbean

islands, as well as North and South American outgroups (55). We note, however, that adaptive

alleles designated as *de novo* on San Salvador Island may be segregating at low frequencies in

other sampled populations or present in unsampled populations.

These fitness networks depict both hybrids and parental species in genotypic space, with

nodes representing SNP haplotypes and edges connecting mutational neighbors (Figure 3b).

Genotypic space is immense; using SNPs coded as homozygous reference, heterozygote, or

homozygous alternate, the number of potential haplotypes is equal to $3^{\# \, SNPs \, in \, network}$. For

instance, to construct a reduced network of 100 SNPs, there are a total of $3^{100} = 5.17 \, X \, 10^{57}$

possible nodes. Thus, unlike experimental studies of individual proteins in haploid *E. coli* (31,

45) or yeast (22), it is not possible for us to investigate the full breadth of genotypic space.

Instead, to understand the distribution of parental species and their hybrids in genotypic

space, we began by using a random sample of ten SNPs drawn from our set of candidate adaptive

alleles in this system. Here, we plotted edges between nodes up to five mutational steps away

(e.g. Figure 3b) and found that generalists and molluscivores are closer on the genotypic fitness

network than either is to scale-eaters (Figure 3c), as expected based on their genetic distance.

Most scale-eaters appear quite isolated in genotypic space, separated from the generalist cluster

of nodes by 12.6 ± 0.091 (mean $\pm$ SE: $P < 0.001$) mutational steps and from molluscivores by

16.3 ± 0.060 steps ($P < 0.001$). In contrast, molluscivores were separated from generalists by

5.37 ± 0.103 steps ($P < 0.001$). Generalists show the greatest intrapopulation distances, separated

from each other by 6.08 ± 0.088 steps ($P < 0.001$). In contrast, molluscivores exhibited the

smallest intrapopulation distances, separated by 1.75 ± 0.021 steps ($P < 0.001$). Scale-eater

intrapopulation distances were intermediate (4.71 ± 0.088 steps: $P < 0.001$).

337
Figure 4. Molluscivore genotypes were more accessible to generalists on the genotypic fitness
landscape than scale-eater genotypes. a. Diagram illustrating genotypic fitness networks and adaptive
walks between species for a hypothetical two-SNP genotypic fitness landscape. Species A & B are
separated by four mutational steps. Dashed lines indicate inaccessible paths that decrease in fitness
leaving a single possible accessible evolutionary trajectory between species A and B (indicated by bold
arrows). Each node in our study is associated with an empirical measure of hybrid fitness from field
experiments (17, 18). Edges are always drawn as directed from low to high fitness. **b.** The same network
as in (**a.**), with fitness plotted on the Y-axis and number of mutational steps from species A to B on the X-
axis. The only accessible path between species A and B is indicated by solid arrows. **c.** Number of
accessible paths between generalists and either specialist, scaled by network size. **d.** Length (# of nodes)
of the shortest accessible paths. Means (large points) ± 2 standard errors are plotted. **e.** Ruggedness, as
measured by the number of peaks (genotypes with no fitter neighbors within a single mutational step
(32)). **f.** Number of accessible paths to peaks, scaled by network size. **g.** Length of the shortest accessible
path to the nearest peak. **h.** Odds ratios (OR: ML estimate and 95% CI) for each measure of accessibility
(x-axis corresponds to panel letters); molluscivore networks have significantly greater summary statistics
when OR > 1. Molluscivore genotypes are more accessible to generalists than scale-eater genotypes due
to a significantly greater number of accessible paths separating them (**c.**) that are significantly shorter (**d.**).
Molluscivore genotypic networks were also less rugged, i.e. they contained significantly fewer peaks (**e.**),
each of which were in turn more accessible from the generalist genotypes (**f., g.**).

Molluscivore genotypes are more accessible to generalists than scale-eater genotypes on the

genotypic fitness landscape

The most accessible paths through genotypic fitness networks are characterized by

monotonically increasing fitness at each mutational step and the smallest possible number of

steps between two states (5, 31, 33) (Figure 4a-b). Furthermore, as described earlier, the

accessibility of individual fitness peaks is predicted to be reduced on increasingly rugged fitness

landscapes that are characterized by a greater number of fitness peaks (8, 30, 33, 101). This

provides three useful metrics of evolutionary accessibility for genotypic trajectories: 1) the total

number of accessible paths relative to network size (Figure 4—figure supplement 1; Appendix

1—table 16), 2) the length of the shortest accessible paths, and 3) the number of fitness peaks

(ruggedness). Here, we define peaks as genotypes with no fitter neighbors and within a single

mutational step (32). With these three metrics, we can quantify the accessibility of interspecific

genotypic pathways.

We used these measures of accessibility to ask: 1) whether molluscivore or scale-eater

genotypes were more accessible to generalists on the fitness landscape (Figure 4c-d), and 2)

whether molluscivore and scale-eater genotypic fitness networks differed in their ruggedness,

characterized by peak number (Figure 4e-g). These measures provide insight into the

predictability of evolution and the role that epistasis plays in their evolution (8, 23, 29, 102).

We constructed 5,000 genotypic fitness networks from a random sample of five species-

specific candidate adaptive SNPs (Figure 3a) for either molluscivores or scale-eaters, requiring

that at least one SNP of each source of genetic variation be present in the sample. We used odds

ratios to compare the relative accessibility and ruggedness of molluscivore fitness networks

compared to scale-eater networks (Figure 4h). Thus, odds ratios (OR) greater than 1 imply

summary statistics are greater for molluscivores than for scale-eaters.

We found that molluscivore genotypes were significantly more accessible to generalists

on the fitness landscape than scale-eaters (Appendix 1—table 17); molluscivore networks had

significantly more accessible paths [OR: (95% CI)= 2.095: (1.934, 2.274)] that were

significantly shorter [OR and 95% CI = 0.253: (0.231, 0.277)]. Not only were molluscivore

genotypes more accessible to generalists, but molluscivore fitness networks were significantly

less rugged than scale-eater networks, comprised of fewer peaks [OR and 95% CI = 0.604:

(0.575, 0.634)], and connected by significantly more accessible paths [OR and 95% CI = 1.514:

(1.404, 1.635)], that contained fewer mutational steps [OR and 95% CI = 0.539: (0.500, 0.579)].

Adaptive introgression and de novo *mutations increase accessibility of novel fitness peaks*

We further used our two metrics of accessibility and landscape ruggedness to ask how different

sources of adaptive genetic variation may influence the topography of the fitness landscape, the

traversal of fitness paths separating generalists from specialists, and ultimately colonization of

novel fitness peaks. We constructed genotypic fitness networks limited to only one of the three

main sources of adaptive genetic variation: standing genetic variation, introgression from one of

four focal Caribbean generalist populations, or *de novo* mutations unique to San Salvador Island.

We also examined all combinations of these three sources to better reflect the actual process of

adaptive divergence originating from only standing genetic variation, then adaptive introgression

plus standing genetic variation, and finally the refinement stage of de novo mutations (55).

Figure 5. Adaptive introgression and *de novo* mutations increase access to specialist fitness peaks.
Odds ratios (maximum likelihood estimate and 95% CI) indicate the effect of each source of variation on
accessibility compared to networks estimated from standing variation alone. Asterisks denote significance
($p < 0.0001$ = ****, < 0.001 = ***). **a.** The number of accessible (i.e. monotonically increasing in fitness)
paths per network, scaled by the size of the network (# of nodes in network). Significance was assessed
using a likelihood ratio test, corrected for the false discovery rate (reported in Appendix 1—table 18).
Dashed lines correspond to the median estimate for standing genetic variation to aid comparison to other
sources of adaptive variation. **b.** Number of mutational steps in the shortest accessible path. Means are
plotted as large circles, with two standard errors shown; dashed horizontal lines correspond to the mean
for standing genetic variation. **c.** Ruggedness of molluscivore and scale-eater genotypic fitness networks
constructed from each source of genetic variation measured by the number of peaks (genotypes with no
fitter neighbors).

We compared sets of 5,000 random 5-SNP genotypic networks drawn from different

sources of adaptive variation (Figure 4a) and compared the effect of each source of variation on

measures of accessibility and landscape ruggedness relative to standing genetic variation. We

treated standing variation as our basis for comparison because this is the source of genetic

variation first available to natural selection (103).

We discovered that genotypic trajectories between generalists and either trophic specialist

in genotypic fitness networks constructed from introgressed or *de novo* adaptive mutations were

significantly more accessible than networks constructed from standing genetic variation (Figure

5). Specifically, random networks that included alternate sources of adaptive variation contained

significantly more accessible fitness paths from generalist to specialists than networks

constructed from standing genetic variation alone, while controlling for differences in overall

network size (Figure 5a; Appendix 1—table 18). Furthermore, accessible paths between

generalists and specialists in networks constructed from introgressed or *de novo* adaptive loci

were significantly shorter in length (Figure 5b). We recovered the same pattern whether

constructing fitness networks from these sources of variation alone or in combination. These

results held across all measures of fitness and for analyses repeated using only hybrids sampled

from the second field experiment (Figure 5—figure supplement 1-2, Appendix 1—table 18-19).

Our finding of increased accessibility of interspecific genotypic trajectories suggests that

fitness landscapes constructed from adaptive standing genetic variation alone are more rugged

than networks including adaptive loci originating from either introgression or *de novo* mutation.

Quantification of landscape ruggedness supported this hypothesis in all cases (Figure 5c;

Appendix 1—table 18-19). Additionally, increasing landscape ruggedness significantly

decreased the length of accessible paths to the nearest local peak [glm(Min. Path Length ~ # of

Peaks, family = "poisson"): $P < 0.0001$, β = -0.088, 95% CI = -0.095 – 0.081].

Scale-eater fitness genotypic fitness landscapes constructed from a combination of

adaptive loci sourced from standing variation, introgression, and *de novo* mutations had

significantly more accessible paths (scaled by network size) separating generalists from scale-

eaters [OR and 95% CI = 1.879: (1.743, 2.041); LRT $P < 0.0001$; Figure 5a] and these paths

were significantly shorter in length compared to networks constructed from standing variation

alone [OR and 95% CI = 0.876: (0.823, 0.932); LRT $P < 0.0001$; Figure 5b]. The only exception

to this pattern across all three fitness measures was for growth rate in genotypic fitness networks

constructed for molluscivore adaptive loci; no significant difference was observed in the length

of the shortest accessible path between networks constructed using standing variation alone or

those constructed using introgressed alleles [OR and 95% CI = 0.994: (0.915, 1.079); LRT $P =$

0.8826; Appendix 1—table 18]. Interestingly, however, for networks constructed from standing

variation and introgressed alleles, we again observed a significant reduction in length of the

shortest accessible paths [OR and 95% CI = 0.897: (0.835, 0.962); LRT $P = 0.0050$; Appendix

1—table 18].

Discussion

We developed a new approach for estimating genotypic fitness landscapes for diploid organisms

and applied it to a system in which phenotypic fitness landscapes have been extensively

investigated. We were able to address long-standing questions posed by fitness landscape theory

in an empirical system and assess the extent to which the shape of the fitness landscape and

accessibility of adaptive walks are contingent upon the source of adaptive genetic variation. We

show that not only are scale-eaters more isolated than molluscivores from generalists on the

fitness landscape, but that the scale-eater fitness landscape is more rugged than molluscivores.

This indicates that epistasis is more pervasive on the scale-eater fitness landscape, leading to less

predictable evolutionary outcomes and fewer accessible trajectories from generalist to scale-eater

genotypes. Overall, we found that most genotypic trajectories were inaccessible and included one

or more mutational steps that decreased in fitness from generalist to specialist. This finding is

consistent with the patterns observed by Weinreich et al. (31), who constructed combinatorially

complete fitness networks for five mutations contributing to antibiotic resistance in *E. coli* and

found that only 18 of 120 possible genotypic trajectories were evolutionarily accessible. In

contrast, Khan et al. (45) estimated that over half of all trajectories were accessible on a complete

fitness landscape constructed using the first five adaptive mutations to fix in an experimental

population of *E. coli*.

We also show that fitness landscapes are most rugged, and therefore epistasis is most

pervasive, when constructed from standing genetic variation alone, ultimately leading to a

reduction in the accessibility of fitness peaks on these landscapes (Figure 5). This finding has

significant implications for the predictability of evolution in the earliest stages of the speciation

process. Adaptation from standing genetic variation is thought to initially be more rapid due to

its initial availability and potentially reduced genetic load within a population (103–105). In

contrast, we consistently found that networks constructed from a combination of adaptive

standing variation, introgression, and *de novo* mutations reduced the ruggedness of fitness

landscapes and thus increased accessibility of interspecific evolutionary trajectories (Figure 5).

This would suggest that adaptive introgression or *de novo* mutations reduce the impacts of

epistasis, resulting in a smoother fitness landscape with a greater number of accessible adaptive

481 walks, facilitating the colonization of new adaptive zones. Future studies testing the generality of

482 these findings will be invaluable for our understanding of the speciation process.

483  Furthermore, our results shed light on the classic problem of crossing fitness valleys on

484 three-dimensional phenotypic fitness landscapes. We show that phenotypic fitness valleys may

485 be circumvented by rare accessible paths on the genotypic fitness landscape. These results are

486 consistent with increasing recognition that three-dimensional depictions of the fitness landscape

487 may lead to incorrect intuitions about how populations evolve (3, 5, 106).

488  Our study represents a significant contribution to the growing body of work applying

489 fitness landscape theory to empirical systems (39, 46, 107–109). Unlike previous studies that

490 experimentally generated combinatorically complete fitness landscapes (22, 31, 45), we

491 subsampled loci across the genome, enabling us to quantify aspects of the genotypic fitness

492 landscape, despite the limitations imposed by large genome sizes and non-model vertebrates.

493 One limitation of this approach is that subsampled fitness networks may not directly correspond

494 to the full landscape (5, 110). For instance, a given subsampled fitness landscape may be present

495 on multiple global, fully sampled fitness landscapes (110). Secondly, nodes (here, SNP

496 haplotypes) can appear disconnected in a subsampled fitness landscape, but may be connected in

497 the full fitness landscape (5). Nevertheless, given that there are more possible genotypes for a

498 gene of 1,000 base-pairs than particles in the known universe (1, 111), nearly all empirical

499 fitness landscapes must necessarily be subsampled at some scale.

500  Although inferences from subsampled fitness networks have their limitations, so too do

501 those obtained from combinatorically complete fitness landscapes, which may themselves be

502 misleading (102). By including mutations that are not segregating in natural populations, the

503 shape of the "complete" fitness landscape and thus accessibility of fitness peaks may be quite

different from what occurs in nature. The shape of fitness landscapes in nature is dictated by the

"realized" epistasis that occurs among naturally segregating loci (102). Changes to "realized"

epistasis induced by introgression or *de novo* mutations appears to be the mechanism altering the

shape of the fitness landscape and thus accessibility of fitness peaks. Our findings that adaptive

introgression and *de novo* mutations make fitness peaks more accessible points towards a

pervasive role of epistasis in determining the predictability of evolution and the speciation

process (5, 8, 22–24, 30).

In the present study we have taken snapshots of the fitness landscape from loci that have

already undergone hard selective sweeps. Consequently, we cannot directly assess the influence

of each adaptive allele on the fitness landscape through time as it increases in frequency.

However, so far we have failed to detect evidence of frequency-dependent selection in this

system after experimental manipulations, at least for morphological traits (18). Future

experimental or simulation studies may track how novel adaptive alleles affect fitness landscape

topography as they increase in frequency.

Conclusion

Our findings are consistent with a growing body of evidence that *de novo* and introgressed

adaptive variation may contribute to rapid speciation and evolution towards novel fitness peaks

(44, 70, 78, 112–116). We demonstrate that adaptive introgression smooths the fitness landscape

and increases the accessibility of fitness peaks. This provides an alternative mechanism to

explain why hybridization appears to play such a pervasive role in adaptive radiation and

speciation. There are many examples of hybridization promoting or inducing rapid speciation

and adaptive radiation. Whether in Galapagos finches (117), African cichlids (77, 78, 113, 118,

119), or *Heliconius* butterflies (75, 120), hybridization has been shown to play a generative role

in adaptive radiation and the evolution of novelty. One mechanism is the increased genotypic,

phenotypic, and ecological diversity generated by hybridization in the form of transgressive

phenotypes (74, 121–124). This diversity in turn facilitates the colonization of novel fitness

peaks and ecological niches, particularly after colonization of a new environment rich in

ecological opportunity (73, 74, 125). However, this model often assumes that the fitness

landscape remains static after adaptive introgression. Here we show that adaptive introgression

directly alters the shape of the fitness landscape, making novel fitness peaks more accessible to

natural selection. Thus, hybridization not only generates genetic diversity, but this diversity can

alter the shape of the fitness landscape, changing which genotypic combinations are favored by

natural selection along with the adaptive walks that lead to them.

Acknowledgements:

We thank Michelle St. John, David Tian, Jacqueline Galvez, and Maria Fernanda Palominos for

insightful discussion of the results; the Gerace Research Centre and Troy Day for logistical

support; the BEST Commission and the government of the Bahamas for permission to collect,

import, and export samples; the Vincent J. Coates Genomics Sequencing Center and Functional

Genomics Laboratory at UC Berkeley for performing whole genome library preparation and

sequencing (supported by NIH S10 OD018174 Instrumentation Grant), and the University of

California, Berkeley and University of North Carolina at Chapel Hill for computational

resources. This work was funded by the National Science Foundation DEB CAREER grant

#1749764, National Institutes of Health grant 5R01DE027052-02, the University of North

Carolina at Chapel Hill, and the University of California, Berkeley to CHM, graduate research

funding from the Museum of Vertebrate Zoology to EJR, and an NSF Postdoctoral Research

Fellowship in Biology under Grant No. 2109838 to AHP.

Data availability:

554   Genomic data are archived at the National Center for Biotechnology Information BioProject

555   Database (Accessions: PRJNA690558; PRJNA394148, PRJNA391309). Sample metadata

556   including morphological measurements and admixture proportions have been uploaded to Dryad:

557   https://doi.org/10.5061/dryad.0vt4b8h0m.

Methods:

Sampling

Our final genomic dataset was comprised of 139 hybrid samples used in two separate field

experiments (17, 18) on San Salvador Island. Experiments were conducted in two lakes: Little

Lake (N = 71) and Crescent Pond (N = 68). Hybrids used in the first field experiment (17) were

comprised of F2 and backcrossed outbred juveniles resulting from crosses between all three

species. Juveniles were raised for 2 months in the lab, individually tagged by injecting a stainless

steel sequential coded wire tag (Northwest Marine Technologies, Inc.) into their left dorsal

musculature, and photographed pre-release for morphometric analyses. Experimental field

enclosures consisted of high- and low-density treatments; density was varied by the number of

tagged juveniles released into each enclosure. Hybrids in the second field experiment (18) were

comprised of F4-F5 outbred juveniles resulting from crosses between all three species.

Individuals were spawned, raised, tagged, and photographed in the same way prior to release.

The second field experiment consisted of high- and low-frequency treatments of approximately

equal densities. The frequency of rare transgressive hybrid phenotypes was manipulated between

treatments in each lake, such that the high- and low-frequency treatments harbored an artificially

increased and decreased frequency of transgressive phenotypes, respectively (18).

All hybrids were measured for 32 external morphological traits (see below). Additionally,

we sequenced parental species of the generalist (N = 17), molluscivores (N = 27), and scale-

eaters (N = 25) sampled from these two lakes and previously included in Richards et al. (55).

Note that we treated samples from Little Lake and Osprey Lake as the same population because

these two lakes are connected through a sand bar and fish from these populations are genetically

undifferentiated (54, 55). For morphological analyses, we additionally measured samples of 60

generalists, 38 molluscivores, and 60 scale-eaters raised in the same laboratory common garden

environment as the hybrids used in field experiments. A full list of samples is included in the

supplement (Appendix 1—table 1).

Sequencing, genotyping, and filtering

Raw reads from a combined set of 396 samples (see supplement) were first mapped to the *C.*

brontotheroides reference genome (genome size = 1.16 Gb; scaffold N50 = 32 Mb) (55) using

bwa-mem (v. 0.7.2). Duplicate reads were identified using MarkDuplicates and BAM indices

were subsequently generated using the Picard software package (126). Samples were genotyped

following Richards et al. (55) according to GATK best practices (127). Specifically, Single

Nucleotide Polymorphisms (SNPs) were called and filtered using hard-filtering criteria in

HaplotypeCaller. We used the following criteria in our filtering approach: QD < 2.0; QUAL <

20; FS < 60; MQRankSum < -12.5; ReadPosRankSum < -8 (127–129).

Following initial genotyping with GATK, we subsequently filtered our data further using

VCFtools (130). Specifically, we filtered using the following flags: --maf 0.05; --min-alleles 2; --

max-alleles 2; --min-meanDP 7; --max-meanDP 100; --max-missing 0.85. Indels were removed.

To reduce non-independence among sites in our final dataset, we conservatively removed sites in

strong linkage disequilibrium using plink v1.9 (--indep-pairwise 10['kb'] 50 0.5: (131)). This

resulted in the retention of 1,129,771 SNPs across 139 hybrid samples and the 69 wild-caught

samples from Richards et al (55). Unless otherwise specified, these SNPs were used for all

downstream analyses.

Hybrid fitness measures

We used three proxies for fitness: survival, growth, or a composite measure of the two. Survival

was a binary trait indicating whether a fish survived (i.e. a tagged fish was recovered) or not

during its exposure period in field enclosures. Growth was a continuous measure, defined as the

proportional increase in Standard Length $\left(\frac{Final\ SL-Starting\ SL}{Starting\ SL}\right)$. Lastly, we defined composite

fitness as survival * growth, similar to the metric used in (132), and analogous to composite

fitness in Hereford (133) who used fecundity as their second fitness measure, rather than growth.

Composite fitness is equal to growth for survivors and equals zero for non-survivors because

growth could not be assessed for non-surviving individuals. Because composite fitness represents

the most information-rich metric of fitness, we report composite fitness results in the main text;

results for growth and survival are included in the supplement.

Population genetic variation

To visualize genetic variation present in hybrids and across lakes (Crescent Pond and Little

Lake), we first used a Principal Components Analysis (PCA) of genetic variation using plink

v1.90 ((131), Figure 1), plotting the first two principal component axes using R (version 3.6.3 (R

Core team 2020). We then estimated admixture proportions in hybrids using ADMIXTURE

v1.3.0 (82). Populations of each species were substantially differentiated between Crescent Pond

and Little Lake (54, 55); thus, independent ADMIXTURE analyses were conducted for each

lake. Because we were primarily interested in admixture proportions of hybrids, we set K = 3 in

these analyses, corresponding to the three parental species used in hybrid crosses. Using

admixture proportions of hybrid individuals, we tested the hypothesis that ancestry predicts

hybrid composite fitness in experimental field enclosures by fitting a generalized additive model

including either 1) scale-eater ancestry or 2) molluscivore ancestry with fixed effects for

experiment and lake. This was repeated for survival and growth separately. Composite fitness

was analyzed using a tobit (zero-censored) model to account for zero-inflation using the censReg

R package (134), survival was analyzed using a binomial model, and growth was analyzed using

a gaussian model. We conducted additional ADMIXTURE analyses that either 1) were

supervised, with generalist, molluscivore, and scale-eater parentals *a priori* assigned to one of

three populations, with only hybrid ancestry proportions being estimated by admixture, or 2)

using only samples from the second field experiment. The same linear models described above

were subsequently repeated using these alternative admixture proportions.

Genome-wide association tests

To identify SNPs that were most strongly associated with fitness (survival, growth, or

composite), we implemented the linear mixed model (LMM) approach in GEMMA (v. 0.98.1:

(135)). This analysis was repeated using each fitness measure as the response variable. To

account for relatedness among samples we estimated the kinship matrix among all 139 hybrid

samples, which in turn were used in downstream LMMs. To account for the potentially

confounding effect of year/experiment and lake on estimated fitness measures, we included each

as covariates in the LMMs. To ensure rare variants were not included in these analyses, we only

included sites that had a minor allele frequency greater than 5% across all hybrids. A total of

933,520 SNPs were analyzed; 196,251 SNPS were excluded due to allele frequency change

following removal of parental species. SNPs strongly associated with fitness were identified with

1) a False Discovery Rate (FDR: Benjamini and Hochberg 1995) less than 0.05, or a 2) *P*-value

< 0.05 following Bonferroni correction. We focused primarily on the sites identified by the

conservative Bonferroni correction, however.

Gene ontology enrichment

We annotated sites that were significantly associated with fitness using snpEff (136) and the

annotated *C. brontotheroides* reference genome (55). We constructed a custom database within

snpEff using the functional annotations originally produced by Richards et al. (55), and

subsequently extracted information on the annotations and putative functional consequences of

each variant.

Using genes identified for each SNP that was significantly associated with one of the

fitness measures, we performed Gene Ontology (GO) enrichment analyses using ShinyGO v0.61

(137). For genes identified as being intergenic, we included both flanking genes. As in Richards

et al. (2021), the gene symbol (abbreviation) database that had the greatest overlap with ours was

that of the human database; thus, we tested for enrichment of biological process ontologies

curated for human gene functions, based on annotations from Ensembl. Results are reported for

biological processes that were significantly enriched with FDR < 0.05. We then compared this

list of candidate loci to those identified in past studies of San Salvador Island pupfishes (55, 88,

138).

Morphometrics

We measured 31 external morphological traits for all 139 hybrids and 69 parental individuals

from Crescent Pond (30 generalists, 19 molluscivores, and 30 scale-eaters) and 85 from Little

Lake (30 generalists, 25 molluscivores, and 30 scale-eaters). We digitally landmarked dorsal and

lateral photographs (both sides) of each lab-reared hybrid (pre-release) or parent using DLTdv8

(139). Measurements included 27 linear distances and three angles. For nearly all individuals,

lateral measurements were collected from both lateral photographs and averaged. Morphological

variables were size-corrected using the residuals of a log10(trait) ~ log10(standard length)

regression standardized for selection analyses as outlined in the supplement. We used these 31

morphological traits to estimate two linear discriminant (LD) axes that best distinguished the

generalist, molluscivore, and scale-eater using the LDA function in the mass package in R. We

then used the resultant LD model to predict LD-scores for the 139 sequenced hybrids for later

use in generalized additive models.

Estimation of adaptive landscapes

We fit generalized additive models (GAMs) using the mgcv package v. 1.8.28 (140) in R to

estimate fitness landscapes for the two discriminant axes (LD1-2) and fitness. All models

included a thin-plate spline fit to the two linear discriminant axes and we included both lake and

experiment in all models as fixed effects. Lake by experiment interaction terms were also

included in some models. Models were ranked using the corrected Akaike Information Criterion

for small sample sizes (AICc) and were considered to be a substantially worse fit to the data if

ΔAICc > 4 from the best-fit model (141). The best-fit model from the above approach was in

turn used to visualize fitness landscapes, plotting predicted values of fitness measures on the two

discriminant axes in R (Figure 2).

Using these results, we tested whether inclusion of SNPs that were strongly associated

with fitness (i.e., those that surpassed the 0.05 Bonferroni threshold) improved estimation of

fitness landscapes. We first extracted genotypes for the highly significant SNPs identified by

GEMMA (13 for composite fitness, four for only growth: see section *Fitness-genotype*

association test), and coded these as either reference, single, or double mutants using VCFtools

(130). We then used the best-fit models identified above and fit a range of models that included

one or all SNPs. Individual fitness-associated SNPs were treated as ordered factors (i.e. transition

from homozygous reference to heterozygote to homozygous alternate) and modeled using a

factor-smooth in the generalized additive models. Note that factor "smooths" are effectively

modeled as step-functions.

To quantify whether the local features of the complete fitness landscape constructed

using all morphological variables and the most strongly fitness-associated SNPs were robust to

sampling uncertainty, we conducted a bootstrapping procedure for this model. Specifically, we

resampled hybrids with replacement 10,000 times and refit the full model. We then calculated

the mean predicted composite fitness for each linear discriminant (LD) axis in slices across the

fitness landscape, both for our observed dataset, and for each bootstrap replicate. Slices divided

the fitness landscape into thirds for each LD axis. We then quantified the mean and standard

deviation of the predicted composite fitness for each position along the other LD axis.

We quantified uncertainty (mean ± SD) around local features of the bootstrapped fitness

landscapes as compared to the observed values of predicted fitness for the same 'slice' of the

fitness landscape. We predicted values at the same 30 points along each LD axis. We then

plotted the locations of parents along the x-axis (LD1 or LD2) to enable relation of features on

the fitness landscape to parental phenotypic distributions.

Estimation of genotypic fitness networks

We first estimated genotypic networks using sites previously shown to be highly divergent (F_{ST}

> 0.95) and showing significant evidence of a hard selective sweep in one of the trophic

specialists (based on evidence from both SweeD and OmegaPlus: (55, 99, 100)). We identified

the SNPs in our unpruned full dataset overlapping with sites inferred to have undergone selective

sweeps (55), resulting in 380 SNPs for molluscivores and 1,118 SNPs for scale-eaters. We

subsequently constructed genotypic fitness networks in igraph v. 1.2.4.1 (142) following the

procedure outlined in the supplement.

To visualize the high-dimensional genotypic fitness network, we randomly sampled ten

adaptive loci 100 times and plotted haplotypes connected by edges if they were within five

mutational steps of one another (Figure 3C). Then we calculated the mean distance between all

species pairs (in number of mutational steps). We used pairwise Tukey's HSD tests to test whether

inter-species distances differed.

Estimation of evolutionary accessibility

We tested whether the evolutionary accessibility of genotypic fitness trajectories through

observed hybrid genotypes from generalist to each specialist species differed based on the source

of genetic variation. We restricted our investigation to networks composed of adaptive loci as

previously described (Figure 3A: Richards et al. 2021). This included a total of 380 SNPs in the

molluscivores, and 1,118 in the scale-eaters. The reduced number of adaptive SNPs sites in our

dataset as compared to that of (55) is due primarily to the increased stringency of our filtering.

We further partitioned these SNPs by their respective sources: standing genetic variation

(molluscivore N = 364; scale-eater N = 1,029), *de novo* mutation (scale-eater N = 24), or

introgression (molluscivore N = 16; scale-eater N = 65), again using the assignments from

Richards et al. (55). For analyses of trajectories between generalists and molluscivores, we

included only SNPs found to be sweeping in molluscivores; likewise, we included only SNPs

sweeping in scale-eaters for analysis of trajectories between generalists and scale-eaters.

The full procedure for constructing genotypic fitness networks, identifying accessible

paths, and quantifying accessibility is outlined in the supplement. Briefly, we randomly

generated 5,000 datasets of five SNPs comprised of either 1) standing genetic variation, 2)

adaptive introgression, 3) *de novo* mutation (scale-eaters only), 4) standing genetic variation +

adaptive introgression, 5) standing genetic variation + *de novo* mutation, or 6) standing genetic

variation + adaptive introgression + *de novo* mutation (scale-eaters only). We additionally

repeated this procedure using both classes of SNPs for molluscivores to determine whether

genotypic trajectories separating generalists to molluscivores are more accessible than those

between generalists and scale-eaters. Because different sets of sites are sweeping in each

specialist, we conducted these analyses separately for each species. We then constructed

genotypic networks, in which nodes are haplotypes of SNPs encoded in 012 format (0 =

homozygous reference, 1 = heterozygote, 2 = homozygous alternate), and edges link mutational

neighbors. When determining whether a path was accessible or not, we only included paths for

which each mutational step (i.e. each intervening haplotype) between generalist to specialist was

observed in at least one hybrid sample.

With these networks, we sought to ask 1) whether molluscivores or scale-eaters are more

accessible to generalists on their respective genotypic fitness landscapes, 2) whether the

ruggedness of the genotypic fitness landscape varied among specialists, and 3) whether

accessibility is contingent upon the source of genetic variation available to natural selection. For

each random network sampled and for each measure of fitness we calculated 1) the minimum

length of accessible paths between a random generalist and specialist sampled from our

sequenced individuals, 2) the number of accessible paths between the same generalist and

specialist pair, 3) the number of nodes, 4) the number of edges in the network, 5) the number of

peaks on the landscape (genotypes with no fitter neighbors (32)), 6) the distance of parental

nodes to these peaks, and 7) the number of accessible paths separating them. Larger networks

often have a greater number of potential paths, including both accessible and inaccessible paths

(Figure 4—figure supplement 1), and we were interested in the relative availability of accessible

adaptive pathways. Consequently, we divided the number of accessible paths in each random

network sampled by the number of nodes. Using our six summary statistics, we tested whether

accessibility and landscape ruggedness differed between networks constructed from

SGV/Introgression/*De novo* mutations (for scale-eaters) or SGV/Introgression (for

molluscivores). To do so, we calculated the mean and standard error of each summary statistic

across all 5,000 replicates. We then modeled the association between each summary statistic and

species using a logistic regression, whereby species was modeled as a binary response variable

(i.e. scale-eater networks = 0, molluscivore networks = 1), with each measure of accessibility as

the predictor. We arbitrarily treated scale-eater networks as the control, and using the estimated

coefficients obtained an odds ratio (OR) that corresponds to the extent to which molluscivore

networks either have increased (OR > 1) or decreased (OR < 1) accessibility measures relative to

scale-eater networks. Significance was similarly assessed using a likelihood ratio test. Additional

details on this procedure may be found in the supplement. Using the fitted logistic model, we

conducted a likelihood ratio test to quantify significance. To explicitly test the hypothesis that

increasing landscape ruggedness reduced the length of accessible paths to the nearest fitness

peak, we fit a Poisson regression model in R in which the number of fitness peaks predicts the

length of the shortest accessible path between any generalist or specialist node and any fitness

peak on that landscape: glm(Min. Distance to Peak ~ Number of Peaks, family = "poisson").

A similar procedure was used to assess whether measures of accessibility (scaled number

of accessible paths, length of the shortest accessible path) and landscape ruggedness (number of

peaks) differed within species among networks constructed from different sources of genetic

variation. Here, networks constructed from SGV were treated as the control, to which all other

networks were compared. For example, to test whether accessibility of the generalist-to-scale-

eater paths are greater in networks constructed from *de novo* mutations than those from SGV, a

logistic model was fitted wherein the response variable for SGV networks was assigned to be 0,

and 1 for *de novo* networks. As before, significance was similarly assessed using a likelihood

ratio test, but here *P*-values were corrected for multiple testing using the false discovery rate

(143). We assessed whether differences in these measures among the two alternate generalist to

specialist trajectories in networks constructed from all three sources of variation were significant

using an ANOVA in R (144). Due to the highly skewed nature of these distributions, post-hoc

pairwise significance was assessed using a nonparametric Kruskal-Wallis one-way analysis of

variance in the agricolae package (145) in R.

Literature Cited

1. S. Wright, The roles of mutation, inbreeding, crossbreeding and selection in evolution. *Sixth Int. Congr. Genet.*, 356–366 (1932).

2. S. Gavrilets, Evolution and speciation on holey adaptive landscapes. *Trends Ecol. Evol.* **12**, 307–312 (1997).

3. S. Gavrilets, "High-dimensional fitness landscapes and speciation" in *Evolution—the Extended Synthesis*, M. Pigliucci, G. B. Müller, Eds. (MIT Press, 2010), pp. 51–64.

4. E. Svensson, R. Calsbeek, *The adaptive landscape in evolutionary biology*, E. Svensson, R. Calsbeek, Eds. (Oxford University Press, 2012).

5. I. Fragata, A. Blanckaert, M. A. Dias Louro, D. A. Liberles, C. Bank, Evolution in the light of fitness landscape theory. *Trends Ecol. Evol.* **34**, 69–82 (2019).

6. K. K. Fear, T. Price, The adaptive surface in ecology. *Oikos* **82**, 440–448 (1998).

7. G. G. Simpson, *Tempo and mode in evolution* (Columbia University Press, 1944).

8. S. Kauffman, S. Levin, Towards a general theory of adaptive walks on rugged landscapes. *J. Theor. Biol.* **128**, 11–45 (1987).

9. R. Lande, S. J. Arnold, The measurement of selection on correlated characters. *Evolution* **37**, 1226 (1983).

10. S. J. Arnold, M. E. Pfrender, A. G. Jones, "The adaptive landscape as a conceptual bridge between micro- and macroevolution" in *Microevolution Rate, Pattern, Process*, (Springer, Dordrecht, 2001), pp. 9–32.

11. S. J. Arnold, Performance surfaces and adaptive landscapes. *Integr. Comp. Biol.* **43**, 367–375 (2003).

12. D. Schluter, P. R. Grant, Determinants of morphological patterns in communities of Darwin's finches. *Am. Nat.* **123**, 175–196 (1984).

13. D. Schluter, Estimating the form of natural selection on a quantitative trait. *Evolution* **42**, 849–861 (1988).

14. A. P. Hendry, S. K. Huber, L. F. De León, A. Herrel, J. Podos, Disruptive selection in a bimodal population of Darwin's finches. *Proc. R. Soc. B Biol. Sci.* **276**, 753–759 (2009).

15. M. O. Beausoleil, *et al.*, Temporally varying disruptive selection in the medium ground finch (Geospiza fortis). *Proc. R. Soc. B Biol. Sci.* **286** (2019).

16. C. W. Benkman, Divergent selection drives the adaptive radiation of crossbills. *Evolution* **57**, 1176–1181 (2003).

17. C. H. Martin, P. C. Wainwright, Multiple fitness peaks on the adaptive landscape drive adaptive radiation in the wild. *Science* **339**, 208–211 (2013).

18. C. H. Martin, K. J. Gould, Surprising spatiotemporal stability of a multi-peak fitness landscape revealed by independent field experiments measuring hybrid fitness. *Evol. Lett.* **4**, 530–544 (2020).

19. S. Gavrilets, *Fitness landscapes and the origin of species* (Princeton University Press, 2004).

20. M. Turelli, N. H. Barton, J. A. Coyne, Theory and speciation. *Trends Ecol. Evol.* **16**, 330–343 (2001).

21. M. R. Servedio, J. W. Boughman, The role of sexual selection in local adaptation and speciation. *Annu. Rev. Ecol. Evol. Syst.* **48**, 85–109 (2017).

22. C. Bank, S. Matuszewski, R. T. Hietpas, J. D. Jensen, On the (un)predictability of a large intragenic fitness landscape. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 14085–14090 (2016).

23. S. Wright, Evolution in Mendelian Populations. *Genetics* **16**, 97–159 (1931).

24. T. Aita, M. Iwakura, Y. Husimi, A cross-section of the fitness landscape of dihydrofolate
reductase. *Protein Eng. Des. Sel.* **14**, 633–638 (2001).
25. M. C. Whitlock, P. C. Phillips, F. B. G. Moore, S. J. Tonsor, Multiple fitness peaks and
epistasis. *https://doi.org/10.1146/annurev.es.26.110195.003125* **26**, 601–629 (1995).
26. F. J. Poelwijk, D. J. Kiviet, D. M. Weinreich, S. J. Tans, Empirical fitness landscapes
reveal accessible evolutionary paths. *Nature* **445**, 383–386 (2007).
27. F. J. Poelwijk, S. Tănase-Nicola, D. J. Kiviet, S. J. Tans, Reciprocal sign epistasis is a
necessary condition for multi-peaked fitness landscapes. *J. Theor. Biol.* **272**, 141–144
(2011).
28. F. J. Poelwijk, S. Tănase-Nicola, D. J. Kiviet, S. J. Tans, Reciprocal sign epistasis is a
necessary condition for multi-peaked fitness landscapes. *J. Theor. Biol.* **272**, 141–144
(2011).
29. J. Neidhart, I. G. Szendro, J. Krug, Adaptation in tunably rugged fitness landscapes: The
Rough Mount Fuji model. *Genetics* **198**, 699–721 (2014).
30. J. Neidhart, I. G. Szendro, J. Krug, Adaptation in tunably rugged fitness landscapes: The
Rough Mount Fuji model. *Genetics* **198**, 699–721 (2014).
31. D. M. Weinreich, N. F. Delaney, M. A. DePristo, D. L. Hartl, Darwinian evolution can
follow only very few mutational paths to fitter proteins. *Science* **312**, 111–114 (2006).
32. L. Ferretti, D. Weinreich, F. Tajima, G. Achaz, Evolutionary constraints in fitness
landscapes. *Heredity (Edinb).* **121**, 466–481 (2018).
33. J. Franke, A. Klözer, J. A. G. M. de Visser, J. Krug, Evolutionary accessibility of
mutational pathways. *PLoS Comput. Biol.* **7**, 1002134 (2011).
34. J. A. G. M. De Visser, J. Krug, Empirical fitness landscapes and the predictability of
evolution. *Nat. Rev. Genet. 2014 157* **15**, 480–490 (2014).
35. J. Maynard Smith, Natural selection and the concept of a protein space. *Nat. 1970
2255232* **225**, 563–564 (1970).
36. C. Brandon Ogbunugafor, A reflection on 50 years of John Maynard Smith's "Protein
Space." *Genetics* **214**, 749–754 (2020).
37. R. A. Fisher, *The genetical theory of natural selection* (Oxford University Press, 1930).
38. H. A. Orr, The genetic theory of adaptation: A brief history. *Nat. Rev. Genet.* **6**, 119–127
(2005).
39. M. Karageorgi, *et al.*, Genome editing retraces the evolution of toxin resistance in the
monarch butterfly. *Nature* **574**, 409–412 (2019).
40. M. A. Siddiq, D. W. Loehlin, K. L. Montooth, J. W. Thornton, Experimental test and
refutation of a classic case of molecular adaptation in Drosophila melanogaster. *Nat. Ecol.
Evol.* **1**, 25 (2017).
41. M. A. Siddiq, J. W. Thornton, Fitness effects but no temperature-mediated balancing
selection at the polymorphic Adh gene of Drosophila melanogaster. *Proc. Natl. Acad. Sci.
U. S. A.* **116**, 21634–21640 (2019).
42. K. S. Singh, *et al.*, The genetic architecture of a host shift: An adaptive walk protected an
aphid and its endosymbiont from plant chemical defenses. *Sci. Adv.* **6**, eaba1070 (2020).
43. F. Peng, *et al.*, Effects of beneficial mutations in pykF gene vary over time and across
replicate populations in a long-term experiment with bacteria. *Mol. Biol. Evol.* **35**, 202–
210 (2018).
44. Z. D. Blount, C. Z. Borland, R. E. Lenski, Historical contingency and the evolution of a
key innovation in an experimental population of Escherichia coli. *Proc. Natl. Acad. Sci.*

U. S. A. **105**, 7899–7906 (2008).

45.   A. I. Khan, D. M. Dinh, D. Schneider, R. E. Lenski, T. F. Cooper, Negative epistasis
between beneficial mutations in an evolving bacterial population. *Science* **332**, 1193–1196
(2011).

46.   P. Nosil, *et al.*, Ecology shapes epistasis in a genotype–phenotype–fitness map for stick
insect colour. *Nat. Ecol. Evol.* **4**, 1673–1684 (2020).

47.   C. H. Martin, P. C. Wainwright, A remarkable species flock of Cyprinodon pupfishes
endemic to San Salvador Island, Bahamas. *Bull. Peabody Museum Nat. Hist.* **54**, 231–241
(2013).

48.   C. H. Martin, Context dependence in complex adaptive landscapes: frequency and trait-
dependent selection surfaces within an adaptive radiation of Caribbean pupfishes.
Evolution **70**, 1265–1282 (2016).

49.   B. J. Turner, D. D. Duvernell, T. M. Bunt, M. G. Barton, Reproductive isolation among
endemic pupfishes (Cyprinodon) on San Salvador Island, Bahamas: Microsatellite
evidence. *Biol. J. Linn. Soc.* **95**, 566–582 (2008).

50.   C. H. Martin, P. C. Wainwright, Trophic novelty is linked to exceptional rates of
morphological diversification in two adaptive radiations of cyprinodon pupfish. *Evolution*
65, 2197–2212 (2011).

51.   M. E. St. John, K. E. Dixon, C. H. Martin, Oral shelling within an adaptive radiation of
pupfishes: Testing the adaptive function of a novel nasal protrusion and behavioural
preference. *J. Fish Biol.* **97**, 163–171 (2020).

52.   C. H. Martin, J. A. McGirr, E. J. Richards, M. E. St. John, How to investigate the origins
of novelty: Insights gained from genetic, behavioral, and fitness perspectives. *Integr. Org.*
Biol. **1** (2019).

53.   M. E. St. John, R. Holzman, C. H. Martin, Rapid adaptive evolution of scale-eating
kinematics to a novel ecological niche. *J. Exp. Biol.* **223** (2020).

54.   C. H. Martin, L. C. Feinstein, Novel trophic niches drive variable progress towards
ecological speciation within an adaptive radiation of pupfishes. *Mol. Ecol.* **23**, 1846–1862
(2014).

55.   E. J. Richards, *et al.*, A vertebrate adaptive radiation is assembled from an ancient and
disjunct spatiotemporal landscape. *Proc. Natl. Acad. Sci.* **118**, e2011811118 (2021).

56.   D. B. Weissman, M. W. Feldman, D. S. Fisher, The rate of fitness-valley crossing in
sexual populations. *Genetics* **186**, 1389–1410 (2010).

57.   D. B. Weissman, M. M. Desai, D. S. Fisher, M. W. Feldman, The rate at which asexual
populations cross fitness valleys. *Theor. Popul. Biol.* **75**, 286–300 (2009).

58.   Y. Iwasa, F. Michor, M. A. Nowak, Stochastic tunnels in evolutionary dynamics. *Genetics*
166, 1571–1579 (2004).

59.   A. F. Bitbol, D. J. Schwab, Quantifying the role of population subdivision in evolution on
rugged fitness landscapes. *PLoS Comput. Biol.* **10**, e1003778 (2014).

60.   A. Wagner, "High-dimensional adaptive landscapes facilitate evolutionary innovation" in
The Adaptive Landscape in Evolutionary Biology, E. I. Svensson, R. Calsbeek, Eds.
(Oxford University Press, 2012), pp. 271–282.

61.   M. Conrad, The geometry of evolution. *Biosystems* **24**, 61–81 (1990).

62.   A. C. Whibley, *et al.*, Evolutionary paths underlying flower color variation in
Antirrhinum. *Science* **313**, 963–966 (2006).

63.   G. Martin, S. F. Elena, T. Lenormand, Distributions of epistasis in microbes fit predictions

from a fitness landscape model. *Nat. Genet. 2007 394* **39**, 555–560 (2007).

64.   P. A. Gros, H. Le Nagard, O. Tenaillon, The evolution of epistasis and its links with
genetic robustness, complexity and drift in a phenotypic model of adaptation. *Genetics*
182, 277–293 (2009).

65.   S. Hwang, S. C. Park, J. Krug, Genotypic complexity of Fisher's Geometric Model.
Genetics **206**, 1049–1079 (2017).

66.   S. C. Park, S. Hwang, J. Krug, Distribution of the number of fitness maxima in Fisher's
geometric model. *J. Phys. A Math. Theor.* **53**, 385601 (2020).

67.   R. Froese, D. Pauldy, FishBase. *FishBase* (2021).

68.   O. Seehausen, others, Genomics and the origin of species. *Nat. Rev. Genet.* **15**, 176–192
(2014).

69.   S. H. Martin, C. D. Jiggins, Interpreting the genomic landscape of introgression. *Curr.*
Opin. Genet. Dev. **47**, 69–74 (2017).

70.   D. A. Marques, J. I. Meier, O. Seehausen, A Combinatorial View on Speciation and
Adaptive Radiation. *Trends Ecol. Evol.* **34**, 531–544 (2019).

71.   T. C. Nelson, W. A. Cresko, Ancient genomic variation underlies repeated ecological
adaptation in young stickleback populations. *Evol. Lett.* **2**, 9–21 (2018).

72.   K. A. Thompson, M. M. Osmond, D. Schluter, Parallel genetic evolution and speciation
from standing variation. *Evol. Lett.* **3**, 129–141 (2019).

73.   J. Mallet, Hybrid speciation. *Nature* **446**, 279–83 (2007).

74.   O. Seehausen, Hybridization and adaptive radiation. *Trends Ecol. Evol.* **19**, 198–207
(2004).

75.   C. Pardo-Diaz, *et al.*, Adaptive introgression across species boundaries in Heliconius
butterflies. *PLoS Genet.* **8**, e1002752 (2012).

76.   J. B. Pease, D. C. Haak, M. W. Hahn, L. C. Moyle, Phylogenomics reveals three sources
of adaptive variation during a rapid radiation. *PLoS Biol.* **14**, e1002379 (2016).

77.   E. J. Richards, J. W. Poelstra, C. H. Martin, Don't throw out the sympatric speciation with
the crater lake water: fine-scale investigation of introgression provides equivocal support
for causal role of secondary gene flow in one of the clearest examples of sympatric
speciation. *Evol. Lett.* **2**, 524–540 (2018).

78.   J. I. Meier, *et al.*, Ancient hybridization fuels rapid cichlid fish adaptive radiations. *Nat.*
Commun. **8**, 14363 (2017).

79.   M. D. McGee, *et al.*, The ecological and genomic basis of explosive adaptive radiation.
Nature **586**, 75–79 (2020).

80.   I. Irisarri, *et al.*, Phylogenomics uncovers early hybridization and adaptive loci shaping the
radiation of Lake Tanganyika cichlid fishes. *Nat. Commun.* **9**, 1–12 (2018).

81.   E. J. Richards, C. H. Martin, Adaptive introgression from distant Caribbean islands
contributed to the diversification of a microendemic adaptive radiation of trophic
specialist pupfishes. *PLoS Genet.* **13**, e1006919 (2017).

82.   D. H. Alexander, J. Novembre, K. Lange, Fast model-based estimation of ancestry in
unrelated individuals. *Genome Res.* **19**, 1655–1664 (2009).

83.   D. H. Alexander, K. Lange, Enhancements to the ADMIXTURE algorithm for individual
ancestry estimation. *BMC Bioinformatics* **12**, 1–6 (2011).

84.   M. E. Arnegard, *et al.*, Genetics of ecological divergence during speciation. *Nature* **511**,
307–311 (2014).

85.   S. Wang, J. J. Hard, F. Utter, Genetic variation and fitness in salmonids. *Conserv. Genet.*

3, 321–333 (2002).

86.   R. Leimu, P. Mutikainen, J. Koricheva, M. Fischer, How general are positive relationships
between plant population size, fitness and genetic variation? *J. Ecol.* **94**, 942–952 (2006).

87.   D. Schluter, *et al.*, Fitness maps to a large-effect locus in introduced stickleback
populations. *Proc. Natl. Acad. Sci. U. S. A.* **118**, 1914889118 (2021).

88.   J. A. McGirr, C. H. Martin, Ecological divergence in sympatry causes gene misexpression
in hybrids. *Mol. Ecol.* **29**, 2707–2721 (2020).

89.   Y. Nakamura, *et al.*, Wwp2 is essential for palatogenesis mediated by the interaction
between Sox9 and mediator subunit 25. *Nat. Commun.* **2**, 1–10 (2011).

90.   L. Mork, G. Crump, "Zebrafish craniofacial development: A Window into early
patterning" in *Current Topics in Developmental Biology*, Y. Chai, Ed. (Academic Press
Inc., 2015), pp. 235–269.

91.   T. Manousaki, *et al.*, Parsing parallel evolution: ecological divergence and differential
gene expression in the adaptive radiations of thick-lipped Midas cichlid fishes from
Nicaragua. *Mol. Ecol.* **22**, 650–669 (2013).

92.   S. J. Lim, *et al.*, Taurine is an essential nutrient for juvenile parrot fish Oplegnathus
fasciatus. *Aquaculture* **414**–**415**, 274–279 (2013).

93.   T. G. Gaylord, A. M. Teague, F. T. Barrows, Taurine supplementation of all-plant protein
diets for rainbow trout (Oncorhynchus mykiss). *J. World Aquac. Soc.* **37**, 509–517 (2006).

94.   M. Yokoyama, T. Takeuchi, G. S. Park, J. Nakazoe, Hepatic cysteinesulphinate
decarboxylase activity in fish. *Aquac. Res.* **32**, 216–220 (2001).

95.   Z. Zhou, *et al.*, Genome-wide association study of growth and body-shape related traits in
large yellow croaker (Larimichthys crocea) using ddRAD sequencing. *Mar. Biotechnol.*
21, 655–670 (2019).

96.   G. M. Yoshida, J. M. Yáñez, Multi-trait GWAS using imputed high-density genotypes
from whole-genome sequencing identifies genes associated with body traits in Nile tilapia.
BMC Genomics **22**, 1–13 (2021).

97.   J. Kulmuni, A. M. Westram, Intrinsic incompatibilities evolving as a by-product of
divergent ecological selection: Considering them in empirical studies on divergence with
gene flow. *Mol. Ecol.* **26**, 3093–3103 (2017).

98.   L. F. S. Fonseca, *et al.*, Gene expression profiling and identification of hub genes in
Nellore cattle with different marbling score levels. *Genomics* **112**, 873–879 (2020).

99.   P. Pavlidis, D. Živković, A. Stamatakis, N. Alachiotis, SweeD: Likelihood-based
detection of selective sweeps in thousands of genomes. *Mol. Biol. Evol.* **30**, 2224–2234
(2013).

100.   N. Alachiotis, A. Stamatakis, P. Pavlidis, OmegaPlus: A scalable tool for rapid detection
of selective sweeps in whole-genome datasets. *Bioinformatics* **28**, 2274–2275 (2012).

101.   J. A. G. M. De Visser, J. Krug, Empirical fitness landscapes and the predictability of
evolution. *Nat. Rev. Genet. 2014 157* **15**, 480–490 (2014).

102.   M. C. Whitlock, P. C. Phillips, F. B. G. Moore, S. J. Tonsor, Multiple fitness peaks and
epistasis. *https://doi.org/10.1146/annurev.es.26.110195.003125* **26**, 601–629 (1995).

103.   R. D. H. Barrett, D. Schluter, Adaptation from standing genetic variation. *Trends Ecol.*
Evol. **23**, 38–44 (2008).

104.   J. Hermisson, P. S. Pennings, Soft sweeps and beyond: understanding the patterns and
probabilities of selection footprints under rapid adaptation. *Methods Ecol. Evol.* **8**, 700–
716 (2017).

105. P. W. Hedrick, Adaptive introgression in animals: examples and comparison to new
mutation and standing variation as sources of adaptive variation. *Mol. Ecol.* **22**, 4606–
4618 (2013).
106. J. Kaplan, The end of the adaptive landscape metaphor? *Biol. Philos. 2008 235* **23**, 625–
638 (2008).
107. V. O. Pokusaeva, *et al.*, An experimental assay of the interactions of amino acids from
orthologous sequences shaping a complex fitness landscape. *PLOS Genet.* **15**, e1008079
(2019).
108. L. I. Gong, M. A. Suchard, J. D. Bloom, Stability-mediated epistasis constrains the
evolution of an influenza protein. *Elife* **2013** (2013).
109. L. I. Gong, J. D. Bloom, Epistatically interacting substitutions are enriched during
adaptive protein evolution. *PLOS Genet.* **10**, e1004328 (2014).
110. F. Blanquart, T. Bataillon, Epistasis and the structure of fitness landscapes: Are
experimental fitness landscapes compatible with fisher's geometric model? *Genetics* **203**,
847–862 (2016).
111. I. G. Szendro, M. F. Schenk, J. Franke, J. Krug, J. A. G. M. De Visser, Quantitative
analyses of empirical fitness landscapes. *J. Stat. Mech. Theory Exp.*, P01005 (2013).
112. T. C. Nelson, *et al.*, Ancient and recent introgression shape the evolutionary history of
pollinator adaptation and speciation in a model monkeyflower radiation (Mimulus section
Erythranthe). *PLoS Genet.* **17**, e1009095 (2021).
113. H. Svardal, *et al.*, Ancestral hybridization facilitated species diversification in the lake
malawi cichlid fish adaptive radiation. *Mol. Biol. Evol.* **37**, 1100–1113 (2020).
114. S. Lamichhaney, *et al.*, Rapid hybrid speciation in Darwin's Finches. *Science* **359**, 224–
228 (2018).
115. P. R. Grant, B. R. Grant, Hybridization increases population variation during adaptive
radiation. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 23216–23224 (2019).
116. N. B. Edelman, J. Mallet, Prevalence and adaptive impact of introgression. *Annu. Rev.*
Genet. **55**, 265–283 (2021).
117. P. R. Grant, B. R. Grant, Hybridization increases population variation during adaptive
radiation. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 23216–23224 (2019).
118. J. I. Meier, *et al.*, The coincidence of ecological opportunity with hybridization explains
rapid adaptive radiation in Lake Mweru cichlid fishes. *Nat. Commun.* **10** (2019).
119. J. W. Poelstra, E. J. Richards, C. H. Martin, Speciation in sympatry with ongoing
secondary gene flow and a potential olfactory trigger in a radiation of Cameroon cichlids.
Mol. Ecol. **27**, 4270–4288 (2018).
120. M. Moest, *et al.*, Selective sweeps on novel and introgressed variation shape mimicry loci
in a butterfly adaptive radiation. *PLoS Biol.* **18**, e3000597 (2020).
121. R. C. Lewontin, L. C. Birch, Hybridization as a Source of Variation for Adaptation to
New Environments. *Evolution* **20**, 315–336 (1966).
122. L. H. Rieseberg, M. A. Archer, R. K. Wayne, Transgressive segregation, adaptation and
speciation. *Heredity (Edinb).* **83**, 363 (1999).
123. K. Kagawa, G. Takimoto, Hybridization can promote adaptive radiation by means of
transgressive segregation. *Ecol. Lett.* **21**, 264–274 (2018).
124. R. Abbott, *et al.*, Hybridization and speciation. *J. Evol. Biol.* **26**, 229–46 (2013).
125. N. B. Edelman, J. Mallet, Prevalence and adaptive impact of introgression. *Annu. Rev.*
Genet. **55**, 265–283 (2021).

126.   Broad Institute, Picard Toolkit (2018).

127.   M. A. Depristo, *et al.*, A framework for variation discovery and genotyping using next-
1078   generation DNA sequencing data. *Nat. Genet.* **43**, 491–501 (2011).

1079   128.   R. Poplin, *et al.*, Scaling accurate genetic variant discovery to tens of thousands of
1080   samples. *bioRxiv*, 201178 (2017).

1081   129.   C. D. Marsden, *et al.*, Diversity, differentiation, and linkage disequilibrium: Prospects for
1082   association mapping in the malaria vector anopheles arabiensis. *G3 Genes, Genomes,*
1083   *Genet.* **4**, 121–131 (2014).

1084   130.   P. Danecek, *et al.*, The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158
1085   (2011).

1086   131.   S. Purcell, *et al.*, PLINK: A tool set for whole-genome association and population-based
1087   linkage analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).

1088   132.   C. T. DiVittorio, *et al.*, Natural selection maintains species despite frequent hybridization
1089   in the desert shrub Encelia. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 33373–33383 (2021).

1090   133.   J. Hereford, A quantitative survey of local adaptation and fitness trade-offs. *Am. Nat.* **173**,
1091   579–588 (2009).

1092   134.   A. Henningsen, censReg: Censored regression (Tobit) models. (2020).

1093   135.   X. Zhou, M. Stephens, Genome-wide efficient mixed-model analysis for association
1094   studies. *Nat. Genet.* **44**, 821–824 (2012).

1095   136.   P. Cingolani, *et al.*, Using Drosophila melanogaster as a model for genotoxic chemical
1096   mutational studies with a new program, SnpSift. *Front. Genet.* **3** (2012).

1097   137.   S. X. Ge, D. Jung, D. Jung, R. Yao, ShinyGO: A graphical gene-set enrichment tool for
1098   animals and plants. *Bioinformatics* **36**, 2628–2629 (2020).

1099   138.   J. A. McGirr, C. H. Martin, Few fixed variants between trophic specialist pupfish species
1100   reveal candidate cis-regulatory alleles underlying rapid craniofacial divergence. *Mol. Biol.*
1101   *Evol.* **38**, 405–423 (2021).

1102   139.   T. L. Hedrick, Software techniques for two- and three-dimensional kinematic
1103   measurements of biological and biomimetic systems. *Bioinspiration and Biomimetics* **3**,
1104   034001 (2008).

1105   140.   S. N. Wood, Fast stable restricted maximum likelihood and marginal likelihood estimation
1106   of semiparametric generalized linear models. *J. R. Stat. Soc. Ser. B (Statistical Methodol.*
1107   **73**, 3–36 (2011).

1108   141.   K. P. Burnham, D. R. Anderson, *Model Selection and Multimodel Inference* (Springer
1109   New York, 2002).

1110   142.   G. Csardi, T. Nepusz, The igraph software package for complex network research.
1111   *InterJournal Complex Syst.* **1695**, 1–9 (2006).

1112   143.   Y. Benjamini, Y. Hochberg, Controlling the false discovery rate: a practical and powerful
1113   approach to multiple testing. *J. R. Stat. Soc. Ser. B*, 289–300 (1995).

1114   144.   R. C. Team, R: A language and environment for statistical computing. (2019).

1115   145.   F. de Mendiburu, agricolae: Statistical procedures for agricultural research (2020).

1116   146.   T. Guillerme, dispRity: A modular R package for measuring disparity. *Methods Ecol.*
1117   *Evol.* **9**, 1755–1763 (2018).

1118   147.   B. J. Knaus, N. J. Grünwald, VCFR: a package to manipulate and visualize variant call
1119   format data in R. *Mol. Ecol. Resour.* **17**, 44–53 (2017).

1120   148.   S. Der Yang, T. S. Lin, F. G. Liu, C. H. Liou, Influence of dietary phosphorus levels on
1121   growth, metabolic response and body composition of juvenile silver perch (Bidyanus

bidyanus). *Aquaculture* **253**, 592–601 (2006).

149. J. M. Costa, *et al.*, Inadequate dietary phosphorus levels cause skeletal anomalies and alter
osteocalcin gene expression in Zebrafish. *Int. J. Mol. Sci.* **19**, 364 (2018).

150. H. Yu, *et al.*, Effects of stocking density and dietary phosphorus levels on the growth
performance, antioxidant capacity, and nitrogen and phosphorus emissions of juvenile
blunt snout bream (Megalobrama amblycephala). *Aquac. Nutr.* **27**, 581–591 (2021).

1128

Appendix 1:
Supplementary Methods
Sampling of hybrid individuals
Samples of hybrid *Cyprinodon* pupfish included herein were first collected following two
separate fitness experiments, conducted on San Salvador Island in 2011 (17) and 2016 (18)
respectively. Experiments were carried out in two lakes: Little Lake (LL), and Crescent Pond
(CP). Following their initial collection at the conclusion of their respective experiments (see (17)
and (18) for protocols), samples were stored in ethanol. In late 2018, 149 hybrid samples were
selected for use in this experiment. Of these, 27 are from the experiment conducted in 2011 (14
from LL, 13 from CP), and the remaining 122 are from the 2016 experiment (58 from LL, 64
from CP). Due to reduced sample size for some species within Little Lake, we include fish
obtained from Osprey Lake for downstream analyses comparing hybrids to Little Lake, as the
two comprise a single, interconnected body of water.
1142
Genomic Library Prep
DNA was extracted from the muscle tissue of hybrids using DNeasy Blood and Tissue kits
(Qiagen, Inc.); these extractions were then quantified using a Qubit 3.0 fluorometer (Thermo
Scientific, Inc). Genomic libraries were prepared by the Vincent J. Coates Genomic Sequencing
Center (QB3) on the automated Apollo 324 system (WaterGen Biosystems, Inc.). Samples were
fragmented using a Covaris sonicator, and barcoded with Illumina indices. Samples were quality
checked using a Fragment Analyzer (Advanced Analyical Technologies, Inc.). All samples were
sequenced to approximately 10x raw coverage on an Illumina NovaSeq.
1151
Genotyping and filtering
Because we are interested in genomic variants found not only within our hybrid samples but
across San Salvador island and the Caribbean, we conducted genotyping including all 247
samples from Richards et al. (55). These samples include members of the three species (*C.*
variegatus, *C. brontotheroides*, *C. desquamator*) found on San Salvador Island, as well as
individuals of *C. variegatus* found throughout the Caribbean, and numerous outgroups (*C.*
laciniatus, *C. higuey*, *C. dearborni*, *Megupsilon aporus*, and *Cualac tesselatus*). We then
excluded *M. aporus* and *C. tesselatus* along with 18 additional samples for which necessary data
for downstream analyses were missing (e.g. quality photographs for the collection of
morphological data). This approach led to the retention of a total of 4,206,786 total SNPs and
139 hybrid individuals.
1163
Morphometrics
Because the morphological measurements used in the 2013 and 2016 experiments differ slightly,
we remeasured all sequenced hybrid individuals and up to 30 individuals of each parent species
for the 30 morphological characters described in Martin & Gould (18), as well as for standard
length (SL).
For each photograph, unit-scale was obtained by additionally landmarking points on a
regular grid included in each photograph using DLTdv8a (139). Landmark data (x-y coordinates
in units of pixels) were subsequently uploaded into R and converted to millimeters (in the case of
linear measurements) or degrees (for angular measurements) using a custom script.
We then assessed, for each trait, the need for size-correction. That is, we sought to avoid
an outsized role of body size in downstream interpretation. Thus, if a trait was colinear with SL,

we regressed the two (treating SL as the predictor) and took the residuals. In each case, both SL
and the response were Log-10 transformed. Subsequently, residuals were scaled such that their
distribution had a mean of 0, and a standard deviation of 1. Traits that did not need size
correction were also Log-transformed and unit-scaled.
We used these morphological measurements to estimate two linear discriminant (LD)
axes that distinguish the generalist, molluscivore, and scale-eater using the LDA function in R.
That is, we used morphological data from the 165 parental fish to estimate LD scores for each
individual. Doing so, we were able to correctly assign individual fish to their corresponding
species with 99.4% accuracy (Figure 2—figure supplement 4-5). Attempting to predict species
assignment by lake did not improve this prediction accuracy (instead reducing prediction
accuracy to 98.7%); consequently, we proceeded with the LD axes estimated without accounting
for lake.
We additionally asked whether 1) specialists were more morphologically constrained
than generalists and 2) if hybrids were less constrained than the three parental species. To do so,
we calculated morphological disparity per group using the dispRity.per.group function
implemented in the R package dispRity v1.3.3 (146). Specifically, this function calculates the
median distance between each row (sample) and the centroid of the matrix per group. By
bootstrapping the data 100 times, we tested the hypothesis that each of the three parental species
differed significantly in their morphological disparity, first using an ANOVA, followed by a
post-hoc pairwise t-test to assess pairwise significance in R. We corrected for multiple tests
using an FDR correction.
To test whether genetic distance predicts morphological distance, we calculated two
distance matrices. First, we calculated pairwise Euclidean distances between all hybrids using all
morphological variables. Then, we calculated pairwise genetic distances between all hybrids
using the final set of SNPs described above with the genet.dist function implemented in vcfR
(147). We then fit a simple linear model, regressing genetic distance on morphological distance.
1201
Estimation of Adaptive Landscapes
We sought to characterize the extent to which the three measures of fitness are predicted by
morphology alone and to, in turn, visualize fitness landscapes for our sequenced hybrids. To do
so, we fitted six generalized additive models (GAMs) using the mgcv package v. 1.8.28 (140) in
R. All models included a thin-plate spline fitted for the two linear discriminant axes. Because we
have strong *a priori* knowledge that fitness outcomes will be contingent to some extent on
experiment year and on the lake in which hybrids were placed, we include experiment and lake
in all fitted models, modeled either as fixed effects, or as an interaction term between the two.
Additionally, we fitted models that included individual splines for each linear discriminant axis,
either with or without experiment or lake as a factor smooth. The full list of models and their
respective fits are included in the supplement (Appendix 1—table 11-13).
For composite fitness, we excluded three SNPs that were within close proximity (i.e. <
1000bp) to a SNP that was more significantly associated. Because of the reduced number of
significantly associated SNPs identified for growth (four) as compared to composite fitness (ten),
we were able to fit and compare all combinations of significantly associated SNPs for the former.
The best-fit model for composite fitness was also the most complex, including all fitness
associated SNPs. Thus, to reduce model complexity, we fit one additional model, excluding any
of the three SNP (fixed effect) that was not significant in the full model. The full range of models

and their associated fits are reported in Appendix 1—table 14-15. As before, predicted fitness
values across LD space were extracted from the best-fit model and plotted using R.
1222
Genotypic Fitness Networks
Recent work has shown that the adaptive radiation of the pupfish of San Salvador island involved
selection on standing genetic variation, adaptive introgression, and *de novo* mutation (55).
Furthermore, the specialists on San Salvador Island received approximately twice as much
adaptive introgression as did generalists on neighboring islands. These findings imply that each
source of genetic variation may exert a unique influence on the fitness landscape, in turn
facilitating the radiation. We sought to explore this possibility and so estimated genotypic fitness
networks using sites previously shown to have undergone hard selective sweeps in specialists. To
do so, we identified the SNPs in our un-thinned dataset overlapping with sites inferred to have
undergone selective sweeps (55) to produce two datasets (one for each specialist). We then
constructed networks using the following procedure:

1234

1235   1) Target SNPs were extracted from the vcf file of all sequenced hybrids and parental species
1236   from Crescent Pond and Little Lake in 0/1/2 format using VCFtools. That is, individuals
1237   genotyped as homozygote reference were coded as 0, heterozygotes as 1, and homozygote
1238   alternatives as 2. Note again that the reference genome used was that of the molluscivore,
1239   *C. brontotheroides*.
1240   2) These SNPs were subsequently loaded into R, and concatenated into haplotypes, such that
1241   each sequenced individual has a haplotype. These per-sample haplotypes were
1242   subsequently associated with metadata, namely the observed binary survival, growth,
1243   assignment as hybrid or one of the three parental species, and lake of origin.
1244   3) Each haplotype was subsequently collapsed and summarized, such that mean survival,
1245   growth, and composite fitness are retained for each unique haplotype. Additionally, the
1246   number of hybrids, generalists, molluscivores and scale-eaters that have the haplotype were
1247   recorded.
1248   4) The distance, in number of mutational steps was then calculated for each pairwise
1249   combination of haplotypes. For example, the distance between haplotype 000 and 001 is a
1250   single mutational step, whereas haplotypes 000 and 002 are two steps away.
1251   5) Lastly, we constructed networks using the R package igraph v. 1.2.4.1 (142). Specifically,
1252   nodes represent haplotypes, and edges are drawn between haplotypes that are mutational
1253   neighbors (i.e. are a single mutational step away). Nodes present in hybrids were colored
1254   and sized proportional to their respective mean fitness. Nodes present only in parental
1255   species were colored according to the species in which that haplotype is unique to.

1256

Estimation of evolutionary accessibility
The large number of SNPs in our dataset above raises numerous challenges in the visualization
and summarization of fitness networks. Perhaps most significant, is that as the number of sites
assessed increases, the sequence space increases vastly; the number of potential haplotypes is
defined by 3 to the power of the number of SNPs, and the number of potential edges is defined
by the number of haplotypes choose 2. Consequently, networks constructed from a larger
number of focal SNPs are comprised of haplotypes that are separated on average by more
mutational steps than those constructed from fewer SNPs. Because we can only interpret the
fitness consequences of evolutionary trajectories for which we have data along each mutational

step, we restricted analysis to haplotypes that are mutational neighbors (i.e., separated by a single
mutational step).
To do so, we developed a permutation approach to construct fitness networks from SNPs
that were sourced from the three sources of genetic variation defined above as well as all
possible combinations including standing variation (i.e. standing genetic variation + adaptive
introgression and/or de novo mutation). Specifically, from each set we sampled five SNPs up to
5000 times. For networks constructed from combinations of sources (e.g. SGV + introgression),
we ensured that at least one of each source was present. To do so, we generated 1000 random
sets of SNPs for all possible combinations (e.g. 1 SGV - 4 introgression, 2 SGV – 3
introgression, etc). Then, we sampled up to 5000 of these combinations; these samples comprise
our permutations. Then, from each permutation, we constructed fitness networks using the five
steps defined above; these networks served as the subsequent assessment of evolutionary
accessibility of genotypic trajectories separating the generalists from either specialist. Edges
were only drawn such that the network is directed; that is, edges were drawn from low to higher
or equal fitness nodes.
We constructed networks using both parental species and hybrids. For each trajectory
under consideration (generalist → molluscivore & generalist → scale-eater), we first identified
all generalist to specialist trajectories that are connected by accessible (monotonically increasing)
paths. From these connected generalist-specialist node pairs, we randomly sampled a single pair.
We then identified the number and length of accessible paths separating the generalist node from
the specialist node and recorded these values using the "all_simple_paths" function in igraph and
excluded any paths that traversed through haplotypes not found in any hybrid (i.e., exclusive to
parental species) and thus had no information on fitness.
Specifically, two node paths (a single mutational step from generalist to specialist) were
allowed only if both parental nodes (SNP haplotypes) were also observed in hybrids, with fitness
data. Three-node paths were allowed if hybrid fitness data was present for at least one of the
parental nodes and the intervening node between the two parental nodes. Paths that were four
nodes or longer were allowed only if all intervening nodes between parental nodes had
associated hybrid fitness data.
We additionally calculated the number of peaks on the genotypic fitness landscape, as
well as the number of accessible paths between parental nodes and these peaks, and the
minimum distance from parental nodes to a peak on the landscape. We define 'peaks' on the
genotypic fitness landscape as genotypes (SNP haplotypes) with no-fitter neighboring genotype,
follow the definition of Ferretti et al. (32). This definition is inclusive of nodes/genotypes that
are equal in fitness to their neighbors, which may be fitter than all other neighboring nodes. We
conservatively excluded nodes that shared only a single neighbor.
1302
Supplementary Results
Population ancestry associations with fitness
When repeating our test for an association between fitness measures and ancestry proportions as
estimated from a supervised ADMIXTURE analyses, we recovered similar results with one
exception; generalist ancestry was significantly associated with growth rate (generalist: $P =$
0.021). Admixture proportions estimated from an unsupervised analysis did not significantly
predict any measure of fitness when only using hybrids from the second field experiment
(Appendix 1—table 4) (18). Genome-wide PC1 was associated with composite fitness ($P =$
0.004) and survival ($P < 0.001$), whereas PC2 was not (Appendix 1—table 5). However, PC1

largely explains differences among lakes (Figure 1c); thus, the positive correlation between PC1
and fitness is likely explained by the previously described overall differences in survival
observed between the two lakes in past experiments (17, 18).
1315
Genomic associations recovered for composite fitness and growth, not survival
Whereas we identified 132 SNPs that were associated with composite fitness, only 58 were
associated with growth and none were associated with survival. Of the SNPs associated with
growth, only four remained significant using the conservative Bonferroni threshold. Across all
significant sites (either via FDR or Bonferroni correction) a total of 11 were shared across
analyses. The only gene proximate to a growth associated SNP was *csad.* Lastly, we found a
single gene shared between our study and the 125 ecological DMIs (putative genetic
incompatibilities that are differentially expressed among specialists and misexpressed in F1
hybrids) presented in McGirr & Martin (88). This gene, associated with growth (but not
composite fitness) in our study, is *mettl21e*.
When considering SNPs found to be associated with growth, we did not identify any gene
ontologies that were significantly enriched at a False Discovery Rate < 0.01. However, looking
at those enriched at an FDR < 0.05, we do observe a number of ontologies related to biosynthetic
processes, and regulation of metabolic processes (Figure 2—figure supplement 3). Specifically,
the greatest (and most significant) enrichment was for that of phosphorus and phosphate
containing compound metabolic processes and their regulation. Phosphorous deficiencies have
previously been associated with poor growth in silver perch (*Bidyanus bidyanus*: (148)) and
skeletal deformaties (including vertebral compression and craniofacial deformaties) in zebrafish
(*Danio rerio*: (149)). Similarly, blunt snout bream (*Megalobrama amblycephala*) exhibited
greater growth rates with increasing phosphorous levels in their diets (150). In short, enrichment
of growth-associated SNPs for ontologies pertaining to phosphorous metabolism is consistent
with the substantial literature documenting that phosphorous availability and metabolism is a
determinant of growth in fishes.
1339
Morphological variation within sampled hybrids
As in the previous two experiments, there is a relative paucity of hybrids exhibiting the
morphologies that characterize either specialist. Rather, most hybrids fall near the generalists,
with a number exhibiting transgressive morphologies (Figure 2—figure supplement 5a). As
expected, both specialists exhibit reduced morphological disparity as compared to generalists,
and hybrids show the greatest (Figure 2—figure supplement 5b. That is, the specialists appear
more morphologically constrained than generalists, falling on average closer to the group
centroid. Interestingly molluscivores exhibit the least disparity, even less so than scale-eaters.
1348
Fitness-associated SNPs influence shape of the adaptive landscape
Using morphology alone, the best fit generalized additive model for survival, growth, and
composite fitness were simpler than the model for composite fitness using both morphology and
fitness-associated SNPs (Appendix 1—table 11-13). For survival and composite fitness (Figure
2—figure supplement 6a-6b), this model included a thin-plate spline for LD1 & LD2, with
experiment and lake included as fixed effects. The resultant landscape was also similar for these
two analyses, supporting an interpretation of directional selection, favoring molluscivores. For
growth, the best-fit model had the thin-plate spline for LD1 & LD2, but included an interaction
term between experiment and lake (Figure 2—figure supplement 6c; Appendix 1—table 12). In

contrast to the previous two models, the landscape predicted using growth as our proxy of fitness
supported an interpretation of directional selection in favor or hybrids most similar to generalists,
and to a lesser extent, scale-eaters.
Notably, model selection using AICc invariably supported the inclusion of fitness
associated SNPs for growth and composite fitness (Appendix 1—table 14-15). For growth, the
best-fit model including genotypes was an improvement of 22.99 AICc over the model including
morphology alone, whereas for composite fitness, the improvement was 94.527 AICc.
Interestingly, the best-fit models and growth including associated SNPs was similar to that of the
landscape without fitness-associated SNPs, but largely supported an interpretation of directional
selection in favor of scale-eaters, and to a lesser extent generalists.

Figure 1—figure supplement 1. Proportion (%) genetic variance explained by the first 20 Principal Components obtained using all SNPs and individuals from Crescent Pond, Little Lake, and Osprey Lake, as well as experimental hybrids. The first two principal component axes are plotted in Figure 1.

Figure 1—figure supplement 2. Principal components two, three, and four.

Figure 1—figure supplement 3. Supervised ADMIXTURE analyses for Crescent Pond (top) and Little Lake (bottom). Sampled individuals of each species (leftmost) individuals were assigned to one of three populations, whereas ancestry proportions were estimated for all resequenced hybrid individuals from field experiments. Colors correspond to probability of assignment to one of three assumed populations/species (K = 3) in this analysis.

Figure 1—figure supplement 4. Genetic distance predicts morphological distance among sampled hybrids.

Figure 1—figure supplement 5. The proportion of generalist or specialist ancestry in hybrids did not predict in experimental hybrids using either A) composite fitness (tobit/zero-censored), B) survival (binomial) or C) growth (gaussian). Results of a generalized in which survival (modeled as a binomial) is predicted by ancestry proportion, including lake and experiment as fixed effects. Points represent individual hybrids, with each individual represented by two points, one indicating their respective scale-eater (salmon, and molluscivore (blue) ancestry proportions. *P*-values correspond to the effect of each type of ancestry (scale-eater – red, molluscivore – blue) on survival probability. Lines are predicted values and are colored according to species.

Figure 2—figure supplement 1. Manhattan plots illustrating the strength of association between individual SNPs and either Survival (A) or Growth (B) as inferred by GEMMA. Significant associations are highlighted in blue (FDR < 0.05) or red (Bonferroni correction P < 0.05). The dashed red line indicates the threshold for significance following Bonferroni p-value adjustment. METTL21E and CSAD are both highly differentiated ($F_{ST} > 0.95$) among specialists, differentially expressed. METTL21E is also misexpressed in F1 hybrids, meaning it exhibits gene expression that is higher or lower than observed in both parental species (88), indicating it is involved in intrinsic incompatibilities.

Figure 2—figure supplement 2. Gene ontology enrichment for SNPs found to be associated with composite fitness. Darker colors indicate ontologies that are more significantly enriched following FDR correction. Length of bar is proportional to the number of genes assigned to an ontology.

Figure 2—figure supplement 3. Gene ontology enrichment for SNPs found to be associated with growth. Darker colors indicate ontologies that are more significantly enriched following FDR correction. Length of bar is proportional to the number of genes assigned to an ontology.

Figure 2—figure supplement 4. The 29 landmarks used to digitally measure thirty traits plus standard length using DLTDV8a (139)**.** The corresponding traits are shown in Appendix 1—table 7.

Figure 2—figure supplement 5. Morphological variation in the three San Salvador Island pupfish species and their experimentally produced hybrids. A. Linear discriminant (LD) morphospace. Parent species and hybrids are plotted using the two LD axes that together serve to distinguish species with 99.4% accuracy. **B.** Within-group disparity calculated as the median distance between each individual and the groups centroid. Small, semitransparent points are the result of 100 bootstrap replicates, and are summarized by box plots, which in turn show the median and interquartile ranges of these bootstrap replicates. Large, opaque points are the observed disparities using the full dataset per group. All pairwise comparisons using t-tests following correction for the false discovery rate were significant ($P < 0.001$).

Figure 2—figure supplement 6. Best-fit fitness landscapes for composite fitness (a) survival (b), and growth without associated SNPs (c) and growth including associated SNPs (d). Colored points indicate locations of parent species in LD morphospace, and ellipses indicate their 50% (solid) and 95% (dotted) confidence intervals. Grey points indicate location of hybrid individuals, with size proportional to their fitness measure. Cooler colors on the adaptive landscape indicate lower predicted fitness.

Figure 2—figure supplement 7. Comparison of 10,000 bootstrapped estimates of predicted mean composite fitness to estimations from observed data across slices of the fitness landscape. The focal fitness landscape (Figure 3c-d) was estimated using composite fitness, linear discriminant axes obtained from all morphological traits, and the most-strongly fitness associated SNPs. The mean value of predicted fitness across all bootstrap replicates is plotted as a solid black line; the dashed black line indicates ± 1 standard deviation. The observed predicted fitness is plotted as the solid red line. Observed parental morphological LD scores are plotted as colored vertical hashes: see legend at bottom. Subplots **a.**, **b.**, and **c.**, are estimates along LD1, as calculated from the bottom, middle and top third of LD2 values respectively. Subplots **d.**, **e.**, and **f.**, are estimates along LD2, as calculated from the bottom, middle and top third of LD1 values respectively.

Figure 2—figure supplement 8. The topography of the composite fitness adaptive landscape is influenced by the distribution of a common SNP haplotype. Shown is the same landscape as in Figure 2, but the plotted points are unique SNP haplotypes for the thirteen most strongly fitness-associated SNPs. One haplotype is particularly frequent among hybrids; individuals with this haplotype are closer in morphospace to generalists and drive the emergence of a local fitness optimum for generalists. All other haplotypes (points) are plotted with a distinct color per haplotype.

Figure 4—figure supplement 1. The raw number of accessible paths increases with network size. The number of edges (and thus number of potential paths) is strongly positively correlated with the observed number of nodes (unique SNP haplotypes) in a network (**a.**). Correspondingly, both the number of nodes (b.) and number of edges (c.) positively correlate with the number of accessible paths between generalists and specialists in a given network. Only results for composite fitness are plotted; results are consistent across fitness measures. Models correspond to those in Appendix 1—table 13. Poisson regression was chosen as each response variable correspond to count-data. Because Poisson regression models are log-linear, we report the exponentiated coefficient which corresponds to the expected multiplicative increase in the mean of Y per unit-value of X.

Figure 5—figure supplement 1. Adaptive loci sourced from introgression and *de novo* mutation reduce fitness landscape ruggedness and increase accessibility as compared to standing genetic variation (SGV) using survival as our proxy for fitness. Odds ratios (maximum likelihood estimate and 95% CI) indicate the effect of each source of variation on accessibility as compared to networks estimated from standing variation alone. **a.** The number of accessible (i.e. monotonically increasing in fitness) paths per network, scaled by the size of the network (# of nodes in network). Significance was assessed using a likelihood ratio test, corrected for the false discovery rate (reported in Appendix 1—table 13). Dashed lines correspond to the median estimate for standing genetic variation to aid comparison to other sources of adaptive variation. **b.** Number of mutational steps in the shortest accessible path. Means are plotted as large circles, with two standard errors shown; dashed horizontal lines correspond to the mean for standing genetic variation. **c.** Ruggedness of molluscivore and scale-eater fitness landscapes constructed from each source of genetic variation as measured by the number of peaks (genotypes with no fitter neighbors).

Figure 5—figure supplement 2. Adaptive loci sourced from introgression and *de novo* mutation reduce fitness landscape ruggedness and increase accessibility as compared to standing genetic variation (SGV) using growth as our proxy for fitness. Odds ratios (maximum likelihood estimate and 95% CI) indicate the effect of each source of variation on accessibility as compared to networks estimated from standing variation alone. **a.** The number of accessible (i.e. monotonically increasing in fitness) paths per network, scaled by the size of the network (# of nodes in network). Significance was assessed using a likelihood ratio test, corrected for the false discovery rate (reported in Appendix 1—table 13). Dashed lines correspond to the median estimate for standing genetic variation to aid comparison to other sources of adaptive variation. **b.** Number of mutational steps in the shortest accessible path. Means are plotted as large circles, with two standard errors shown; dashed horizontal lines correspond to the mean for standing genetic variation. **c.** Ruggedness of molluscivore and scale-eater fitness landscapes constructed from each source of genetic variation as measured by the number of peaks (genotypes with no fitter neighbors).

Appendix 1—Table 1. Samples of hybrids and parental studies used either in genomic or morphological analyses, along with associated metadata.

ID	Sequenced	Morphology	Survivorship	Experiment	Lake	Species
CP04E02	This study	Yes	Non-Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP07F05	This study	Yes	Non-Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP08H06	This study	Yes	Non-Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP09C02	This study	Yes	Non-Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP09D01	This study	Yes	Non-Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP09F10	This study	Yes	Non-Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP10B05	This study	Yes	Non-Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP11C10	This study	Yes	Non-Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP13E03	This study	Yes	Non-Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP13F01	This study	Yes	Non-Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
LL01F03	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL01F05	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL01G05	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL02A08	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL02D09	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL02D11	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL02E04	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL02E07	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL02E09	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL04E03	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL04E09	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL04F02	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL04F07	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL05A01	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL05C06	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL05C09	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL05C10	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL05D10	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL05E06	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL05E08	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL05H10	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL06A10	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL06B03	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL06C10	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL06D12	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL06E10	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL06F08	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL08B09	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL08B11	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL08G07	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL08G10	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL08H01	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL09A09	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL09A10	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL09B05	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL09C03	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL09D01	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL09D10	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL09E01	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL09F03	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL09H07	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL09H10	This study	Yes	Non-Survivor	Martin & Gould 2020	Little Lake	Hybrid

CP06G01	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP06H09	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP07E08	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP07H01	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP07H11	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP08F07	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP08G08	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP09D02	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP10A03	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP10C04	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP10C07	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP10C12	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP10F12	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP10H01	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP11B03	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP11C08	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP11D04	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP11G06	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP11G12	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP12E05	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP12E07	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP12H04	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP13B02	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP13B08	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP14C04	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP15A10	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP15B08	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP15E01	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP15E03	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP17A01	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP18B11	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP19B02	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP19C02	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP19C07	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP19F01	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CP19G03	This study	Yes	Survivor	Martin & Gould 2020	Crescent Pond	Hybrid
CPH01	This study	Yes	Survivor	Martin & Wainwright 2013	Crescent Pond	Hybrid
CPH02	This study	Yes	Survivor	Martin & Wainwright 2013	Crescent Pond	Hybrid
CPH03	This study	Yes	Survivor	Martin & Wainwright 2013	Crescent Pond	Hybrid
CPH04	This study	Yes	Survivor	Martin & Wainwright 2013	Crescent Pond	Hybrid
CPH05	This study	Yes	Survivor	Martin & Wainwright 2013	Crescent Pond	Hybrid
CPH07	This study	Yes	Survivor	Martin & Wainwright 2013	Crescent Pond	Hybrid
CPH08	This study	Yes	Survivor	Martin & Wainwright 2013	Crescent Pond	Hybrid
CPH09	This study	Yes	Survivor	Martin & Wainwright 2013	Crescent Pond	Hybrid
CPH10	This study	Yes	Survivor	Martin & Wainwright 2013	Crescent Pond	Hybrid
CPH100	This study	Yes	Survivor	Martin & Wainwright 2013	Crescent Pond	Hybrid
CPH11	This study	Yes	Survivor	Martin & Wainwright 2013	Crescent Pond	Hybrid
CPH123	This study	Yes	Survivor	Martin & Wainwright 2013	Crescent Pond	Hybrid
LL01H04	This study	Yes	Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL02G09	This study	Yes	Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL06B06	This study	Yes	Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL06E04	This study	Yes	Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL07E12	This study	Yes	Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL07G04	This study	Yes	Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL08A09	This study	Yes	Survivor	Martin & Gould 2020	Little Lake	Hybrid
LL08B04	This study	Yes	Survivor	Martin & Gould 2020	Little Lake	Hybrid

LL175	This study	Yes	Survivor	Martin & Wainwright 2013	Little Lake	Hybrid
LL23	This study	Yes	Survivor	Martin & Wainwright 2013	Little Lake	Hybrid
LL247	This study	Yes	Survivor	Martin & Wainwright 2013	Little Lake	Hybrid
LL251	This study	Yes	Survivor	Martin & Wainwright 2013	Little Lake	Hybrid
LL271	This study	Yes	Survivor	Martin & Wainwright 2013	Little Lake	Hybrid
LLH12	This study	Yes	Survivor	Martin & Wainwright 2013	Little Lake	Hybrid
LLH34	This study	Yes	Survivor	Martin & Wainwright 2013	Little Lake	Hybrid
LLH41	This study	Yes	Survivor	Martin & Wainwright 2013	Little Lake	Hybrid
LLH51	This study	Yes	Survivor	Martin & Wainwright 2013	Little Lake	Hybrid
LLH94	This study	Yes	Survivor	Martin & Wainwright 2013	Little Lake	Hybrid
CRPA1	Richards et al. 2021	No	NA	NA	Crescent Pond	Generalist
CRPA1000	Richards et al. 2021	No	NA	NA	Crescent Pond	Generalist
CRPA1001	Richards et al. 2021	No	NA	NA	Crescent Pond	Generalist
CRPA1003	Richards et al. 2021	No	NA	NA	Crescent Pond	Generalist
CRPA3	Richards et al. 2021	No	NA	NA	Crescent Pond	Generalist
LILA1	Richards et al. 2021	No	NA	NA	Little Lake	Generalist
OSPA1	Richards et al. 2021	No	NA	NA	Osprey Pond (Little Lake)	Generalist
OSPA1000	Richards et al. 2021	No	NA	NA	Osprey Pond (Little Lake)	Generalist
OSPA1001	Richards et al. 2021	No	NA	NA	Osprey Pond (Little Lake)	Generalist
OSPA11	Richards et al. 2021	No	NA	NA	Osprey Pond (Little Lake)	Generalist
OSPA12	Richards et al. 2021	No	NA	NA	Osprey Pond (Little Lake)	Generalist
OSPA13	Richards et al. 2021	No	NA	NA	Osprey Pond (Little Lake)	Generalist
OSPA4	Richards et al. 2021	No	NA	NA	Osprey Pond (Little Lake)	Generalist
OSPA5	Richards et al. 2021	No	NA	NA	Osprey Pond (Little Lake)	Generalist
OSPA6	Richards et al. 2021	No	NA	NA	Osprey Pond (Little Lake)	Generalist
OSPA8	Richards et al. 2021	No	NA	NA	Osprey Pond (Little Lake)	Generalist
OSPA9	Richards et al. 2021	No	NA	NA	Osprey Pond (Little Lake)	Generalist
CRPM1	Richards et al. 2021	No	NA	NA	Crescent Pond	Molluscivore
CRPM10	Richards et al. 2021	No	NA	NA	Crescent Pond	Molluscivore
CRPM1000	Richards et al. 2021	No	NA	NA	Crescent Pond	Molluscivore
CRPM1001	Richards et al. 2021	No	NA	NA	Crescent Pond	Molluscivore
CRPM11	Richards et al. 2021	No	NA	NA	Crescent Pond	Molluscivore
CRPM2	Richards et al. 2021	No	NA	NA	Crescent Pond	Molluscivore
CRPM3	Richards et al. 2021	No	NA	NA	Crescent Pond	Molluscivore
CRPM5	Richards et al. 2021	No	NA	NA	Crescent Pond	Molluscivore
CRPM6	Richards et al. 2021	No	NA	NA	Crescent Pond	Molluscivore
CRPM7	Richards et al. 2021	No	NA	NA	Crescent Pond	Molluscivore
CRPM8	Richards et al. 2021	No	NA	NA	Crescent Pond	Molluscivore
CRPM9	Richards et al. 2021	No	NA	NA	Crescent Pond	Molluscivore
LILM-QTL	Richards et al. 2021	No	NA	NA	Little Lake	Molluscivore
LILM3	Richards et al. 2021	No	NA	NA	Little Lake	Molluscivore
LILM4	Richards et al. 2021	No	NA	NA	Little Lake	Molluscivore
LILM5	Richards et al. 2021	No	NA	NA	Little Lake	Molluscivore
OSPM1	Richards et al. 2021	No	NA	NA	Osprey Pond (Little Lake)	Molluscivore
OSPM1000	Richards et al. 2021	No	NA	NA	Osprey Pond (Little Lake)	Molluscivore
OSPM1001	Richards et al. 2021	No	NA	NA	Osprey Pond (Little Lake)	Molluscivore
OSPM11	Richards et al. 2021	No	NA	NA	Osprey Pond (Little Lake)	Molluscivore
OSPM2	Richards et al. 2021	No	NA	NA	Osprey Pond (Little Lake)	Molluscivore
OSPM3	Richards et al. 2021	No	NA	NA	Osprey Pond (Little Lake)	Molluscivore
OSPM4	Richards et al. 2021	No	NA	NA	Osprey Pond (Little Lake)	Molluscivore
OSPM5	Richards et al. 2021	No	NA	NA	Osprey Pond (Little Lake)	Molluscivore
OSPM7	Richards et al. 2021	No	NA	NA	Osprey Pond (Little Lake)	Molluscivore
OSPM8	Richards et al. 2021	No	NA	NA	Osprey Pond (Little Lake)	Molluscivore
OSPM9	Richards et al. 2021	No	NA	NA	Osprey Pond (Little Lake)	Molluscivore
CRPP-QTL	Richards et al. 2021	No	NA	NA	Crescent Pond	Scale-eater
CRPP1000	Richards et al. 2021	No	NA	NA	Crescent Pond	Scale-eater

LILP4	Richards et al. 2021	No	NA	NA		Little Lake	Scale-eater
LILP5	Richards et al. 2021	No	NA	NA		Little Lake	Scale-eater
OSPP1	Richards et al. 2021	No	NA	NA		Osprey Pond (Little Lake)	Scale-eater
OSPP10	Richards et al. 2021	No	NA	NA		Osprey Pond (Little Lake)	Scale-eater
OSPP1000	Richards et al. 2021	No	NA	NA		Osprey Pond (Little Lake)	Scale-eater
OSPP1001	Richards et al. 2021	No	NA	NA		Osprey Pond (Little Lake)	Scale-eater
OSPP11	Richards et al. 2021	No	NA	NA		Osprey Pond (Little Lake)	Scale-eater
OSPP2	Richards et al. 2021	No	NA	NA		Osprey Pond (Little Lake)	Scale-eater
OSPP3	Richards et al. 2021	No	NA	NA		Osprey Pond (Little Lake)	Scale-eater
OSPP4	Richards et al. 2021	No	NA	NA		Osprey Pond (Little Lake)	Scale-eater
OSPP5	Richards et al. 2021	No	NA	NA		Osprey Pond (Little Lake)	Scale-eater
OSPP7	Richards et al. 2021	No	NA	NA		Osprey Pond (Little Lake)	Scale-eater
OSPP9	Richards et al. 2021	No	NA	NA		Osprey Pond (Little Lake)	Scale-eater
CPA01	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA03	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA05	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA07	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA09	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA11	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA13	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA15	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA17	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA19	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA21	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA23	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA25	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA27	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA29	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA31	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA33	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA35	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA37	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA39	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA41	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA43	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA45	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA47	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA49	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA51	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA53	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA55	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA57	NA	Yes	Parental	NA		Crescent Pond	Generalist
CPA59	NA	Yes	Parental	NA		Crescent Pond	Generalist
LLA20	NA	Yes	Parental	NA		Little Lake	Generalist
LLA21	NA	Yes	Parental	NA		Little Lake	Generalist
LLA22	NA	Yes	Parental	NA		Little Lake	Generalist
LLA23	NA	Yes	Parental	NA		Little Lake	Generalist
LLA24	NA	Yes	Parental	NA		Little Lake	Generalist
LLA25	NA	Yes	Parental	NA		Little Lake	Generalist
LLA26	NA	Yes	Parental	NA		Little Lake	Generalist
LLA27	NA	Yes	Parental	NA		Little Lake	Generalist
LLA28	NA	Yes	Parental	NA		Little Lake	Generalist
LLA29	NA	Yes	Parental	NA		Little Lake	Generalist
LLA30	NA	Yes	Parental	NA		Little Lake	Generalist
LLA31	NA	Yes	Parental	NA		Little Lake	Generalist
LLA32	NA	Yes	Parental	NA		Little Lake	Generalist

LLA43	NA	Yes	Parental	NA	Little Lake	Generalist
LLA44	NA	Yes	Parental	NA	Little Lake	Generalist
LLA45	NA	Yes	Parental	NA	Little Lake	Generalist
LLA46	NA	Yes	Parental	NA	Little Lake	Generalist
LLA47	NA	Yes	Parental	NA	Little Lake	Generalist
LLA48	NA	Yes	Parental	NA	Little Lake	Generalist
LLA49	NA	Yes	Parental	NA	Little Lake	Generalist
CPM01	NA	Yes	Parental	NA	Crescent Pond	Molluscivore
CPM02	NA	Yes	Parental	NA	Crescent Pond	Molluscivore
CPM03	NA	Yes	Parental	NA	Crescent Pond	Molluscivore
CPM04	NA	Yes	Parental	NA	Crescent Pond	Molluscivore
CPM05	NA	Yes	Parental	NA	Crescent Pond	Molluscivore
CPM06	NA	Yes	Parental	NA	Crescent Pond	Molluscivore
CPM07	NA	Yes	Parental	NA	Crescent Pond	Molluscivore
CPM08	NA	Yes	Parental	NA	Crescent Pond	Molluscivore
CPM09	NA	Yes	Parental	NA	Crescent Pond	Molluscivore
CPM10	NA	Yes	Parental	NA	Crescent Pond	Molluscivore
CPM11	NA	Yes	Parental	NA	Crescent Pond	Molluscivore
CPM12	NA	Yes	Parental	NA	Crescent Pond	Molluscivore
CPM13	NA	Yes	Parental	NA	Crescent Pond	Molluscivore
CPM14	NA	Yes	Parental	NA	Crescent Pond	Molluscivore
CPM15	NA	Yes	Parental	NA	Crescent Pond	Molluscivore
CPM16	NA	Yes	Parental	NA	Crescent Pond	Molluscivore
CPM17	NA	Yes	Parental	NA	Crescent Pond	Molluscivore
CPM18	NA	Yes	Parental	NA	Crescent Pond	Molluscivore
CPM19	NA	Yes	Parental	NA	Crescent Pond	Molluscivore
CPM20	NA	Yes	Parental	NA	Crescent Pond	Molluscivore
LLM01	NA	Yes	Parental	NA	Little Lake	Molluscivore
LLM02	NA	Yes	Parental	NA	Little Lake	Molluscivore
LLM03	NA	Yes	Parental	NA	Little Lake	Molluscivore
LLM04	NA	Yes	Parental	NA	Little Lake	Molluscivore
LLM05	NA	Yes	Parental	NA	Little Lake	Molluscivore
LLM06	NA	Yes	Parental	NA	Little Lake	Molluscivore
LLM07	NA	Yes	Parental	NA	Little Lake	Molluscivore
LLM08	NA	Yes	Parental	NA	Little Lake	Molluscivore
LLM09	NA	Yes	Parental	NA	Little Lake	Molluscivore
LLM10	NA	Yes	Parental	NA	Little Lake	Molluscivore
LLM11	NA	Yes	Parental	NA	Little Lake	Molluscivore
LLM12	NA	Yes	Parental	NA	Little Lake	Molluscivore
LLM13	NA	Yes	Parental	NA	Little Lake	Molluscivore
LLM14	NA	Yes	Parental	NA	Little Lake	Molluscivore
LLM15	NA	Yes	Parental	NA	Little Lake	Molluscivore
LLM16	NA	Yes	Parental	NA	Little Lake	Molluscivore
LLM17	NA	Yes	Parental	NA	Little Lake	Molluscivore
LLM18	NA	Yes	Parental	NA	Little Lake	Molluscivore
CPP01	NA	Yes	Parental	NA	Crescent Pond	Scale-eater
CPP02	NA	Yes	Parental	NA	Crescent Pond	Scale-eater
CPP03	NA	Yes	Parental	NA	Crescent Pond	Scale-eater
CPP04	NA	Yes	Parental	NA	Crescent Pond	Scale-eater
CPP05	NA	Yes	Parental	NA	Crescent Pond	Scale-eater
CPP06	NA	Yes	Parental	NA	Crescent Pond	Scale-eater
CPP07	NA	Yes	Parental	NA	Crescent Pond	Scale-eater
CPP08	NA	Yes	Parental	NA	Crescent Pond	Scale-eater
CPP09	NA	Yes	Parental	NA	Crescent Pond	Scale-eater
CPP10	NA	Yes	Parental	NA	Crescent Pond	Scale-eater
CPP11	NA	Yes	Parental	NA	Crescent Pond	Scale-eater

CPP22	NA	Yes	Parental	NA	Crescent Pond	Scale-eater
CPP23	NA	Yes	Parental	NA	Crescent Pond	Scale-eater
CPP24	NA	Yes	Parental	NA	Crescent Pond	Scale-eater
CPP25	NA	Yes	Parental	NA	Crescent Pond	Scale-eater
CPP26	NA	Yes	Parental	NA	Crescent Pond	Scale-eater
CPP27	NA	Yes	Parental	NA	Crescent Pond	Scale-eater
CPP28	NA	Yes	Parental	NA	Crescent Pond	Scale-eater
CPP29	NA	Yes	Parental	NA	Crescent Pond	Scale-eater
CPP30	NA	Yes	Parental	NA	Crescent Pond	Scale-eater
LLP01	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP02	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP03	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP04	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP05	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP06	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP07	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP08	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP09	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP10	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP11	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP12	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP13	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP14	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP15	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP16	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP17	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP18	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP19	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP20	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP21	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP22	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP23	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP24	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP25	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP26	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP27	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP28	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP29	NA	Yes	Parental	NA	Little Lake	Scale-eater
LLP30	NA	Yes	Parental	NA	Little Lake	Scale-eater

Appendix 1—Table 2. Models tested to assess the extent to which specialist ancestry predicts measures of fitness and their respective fits using all samples and an unsupervised admixture analysis. Best-fit models are bolded.

Fitness Measure	Model Family	Species	Model	AIC	Ancestry Coefficient	Ancestry *P*-value
Composite	**Tobit**	**Generalist**	Composite ~ AncestryProp	202.40	0.02	0.8525
			Composite ~ AncestryProp + Lake	**196.02**	**-0.11**	**0.3847**
			Composite ~ AncestryProp + Experiment	203.38	0.08	0.5223
			Composite ~ AncestryProp + Lake + Experiment	197.56	-0.06	0.6610
			Composite ~ AncestryProp + Lake * Experiment	197.77	-0.10	0.5071
		Molluscivore	Composite ~ AncestryProp	202.26	0.06	0.6745
			Composite ~ AncestryProp + Lake	**195.08**	**0.20**	**0.1945**
			Composite ~ AncestryProp + Experiment	203.74	0.03	0.8131
			Composite ~ AncestryProp + Lake + Experiment	196.54	0.17	0.2700
			Composite ~ AncestryProp + Lake * Experiment	196.66	0.20	0.2129
		Scale-eater	Composite ~ AncestryProp	201.64	-0.17	0.3725
			Composite ~ AncestryProp + Lake	196.69	-0.06	0.7619
			Composite ~ AncestryProp + Experiment	201.62	-0.33	0.1463
			Composite ~ AncestryProp + Lake + Experiment	196.90	-0.22	0.3595
			Composite ~ AncestryProp + Lake * Experiment	**197.61**	**-0.19**	**0.4393**
Growth	**Gaussian**	**Generalist**	Growth ~ AncestryProp	42.47	0.08	0.4084
			Growth ~ AncestryProp + Lake	18.85	0.32	0.0010
			Growth ~ AncestryProp + Experiment	13.43	-0.32	0.0032
			Growth ~ AncestryProp + Lake + Experiment	-23.42	-0.07	0.4413
			Growth ~ AncestryProp + Lake * Experiment	**-89.90**	**0.10**	**0.1187**
		Molluscivore	Growth ~ AncestryProp	42.52	0.11	0.4231
			Growth ~ AncestryProp + Lake	29.24	-0.13	0.3383
			Growth ~ AncestryProp + Experiment	13.91	0.35	0.0041
			Growth ~ AncestryProp + Lake + Experiment	-23.57	0.09	0.3932
			Growth ~ AncestryProp + Lake * Experiment	**-88.32**	**-0.07**	**0.3280**
		Scale-eater	Growth ~ AncestryProp	36.71	-0.41	0.0122
			Growth ~ AncestryProp + Lake	12.25	-0.64	0.0000
			Growth ~ AncestryProp + Experiment	21.98	0.14	0.4870
			Growth ~ AncestryProp + Lake + Experiment	-22.80	0.00	0.9950
			Growth ~ AncestryProp + Lake * Experiment	**-88.53**	**-0.11**	**0.2825**
Survival	**Binomial**	**Generalist**	Survival ~ AncestryProp	188.70	-0.13	0.7769
			Survival ~ AncestryProp + Lake	151.13	-1.59	0.0144
			Survival ~ AncestryProp + Experiment	159.06	0.92	0.0793
			Survival ~ AncestryProp + Lake + Experiment	**123.49**	**-0.54**	**0.4400**
			Survival ~ AncestryProp + Lake * Experiment	125.49	-0.54	0.4400
		Molluscivore	Survival ~ AncestryProp	188.78	0.03	0.9643
			Survival ~ AncestryProp + Lake	155.11	1.13	0.0936
			Survival ~ AncestryProp + Experiment	161.41	-0.53	0.3709
			Survival ~ AncestryProp + Lake + Experiment	**123.06**	**0.76**	**0.3127**
			Survival ~ AncestryProp + Lake * Experiment	125.06	0.76	0.3127
		Scale-eater	Survival ~ AncestryProp	188.61	0.34	0.6756
			Survival ~ AncestryProp + Lake	154.93	1.60	0.0852
			Survival ~ AncestryProp + Experiment	158.99	-1.69	0.0828
			Survival ~ AncestryProp + Lake + Experiment	**124.04**	**-0.28**	**0.8041**
			Survival ~ AncestryProp + Lake * Experiment	126.04	-0.28	0.8041

Appendix 1—Table 3. Models tested to assess the extent to which specialist ancestry predicts measures of fitness and their respective fits using all samples and a supervised admixture analysis. Best-fit models are bolded.

Fitness Measure	Model Family	Species	Model	AIC	Ancestry Coefficient	Ancestry *P*-value
Composite	**Tobit**	**Generalist**	Composite ~ AncestryProp	201.70	0.09	0.3883
			Composite ~ AncestryProp + Lake	**196.79**	**0.00**	**0.9774**
			Composite ~ AncestryProp + Experiment	201.56	0.19	0.1375
			Composite ~ AncestryProp + Lake + Experiment	197.39	0.08	0.5475
			Composite ~ AncestryProp + Lake * Experiment	198.06	0.06	0.6961
		Molluscivore	Composite ~ AncestryProp	202.39	-0.03	0.8382
			Composite ~ AncestryProp + Lake	**196.49**	**0.08**	**0.5870**
			Composite ~ AncestryProp + Experiment	203.59	-0.06	0.6576
			Composite ~ AncestryProp + Lake + Experiment	197.68	0.04	0.7787
			Composite ~ AncestryProp + Lake * Experiment	198.03	0.07	0.6638
		Scale-eater	Composite ~ AncestryProp	201.06	-0.22	0.2395
			Composite ~ AncestryProp + Lake	196.33	-0.13	0.4993
			Composite ~ AncestryProp + Experiment	200.40	-0.41	0.0717
			Composite ~ AncestryProp + Lake + Experiment	**195.82**	**-0.32**	**0.1705**
			Composite ~ AncestryProp + Lake * Experiment	196.60	-0.29	0.2091
Growth	**Gaussian**	**Generalist**	Growth ~ AncestryProp	39.30	0.17	0.0525
			Growth ~ AncestryProp + Lake	11.24	0.37	0.0000
			Growth ~ AncestryProp + Experiment	19.52	-0.18	0.0915
			Growth ~ AncestryProp + Lake + Experiment	-22.80	0.01	0.9525
			Growth ~ AncestryProp + Lake * Experiment	**-93.03**	**0.14**	**0.0206**
		Molluscivore	Growth ~ AncestryProp	43.11	-0.03	0.8005
			Growth ~ AncestryProp + Lake	26.45	-0.23	0.0581
			Growth ~ AncestryProp + Experiment	18.62	0.23	0.0541
			Growth ~ AncestryProp + Lake + Experiment	-22.89	0.03	0.7653
			Growth ~ AncestryProp + Lake * Experiment	**-89.91**	**-0.10**	**0.1177**
		Scale-eater	Growth ~ AncestryProp	32.80	-0.50	0.0015
			Growth ~ AncestryProp + Lake	6.53	-0.69	0.0000
			Growth ~ AncestryProp + Experiment	22.48	-0.02	0.9330
			Growth ~ AncestryProp + Lake + Experiment	-23.14	-0.09	0.5710
			Growth ~ AncestryProp + Lake * Experiment	**-89.59**	**-0.15**	**0.1427**
Survival	**Binomial**	**Generalist**	Survival ~ AncestryProp	188.75	-0.09	0.8445
			Survival ~ AncestryProp + Lake	152.86	-1.30	0.0292
			Survival ~ AncestryProp + Experiment	157.53	1.13	0.0334
			Survival ~ AncestryProp + Lake + Experiment	**124.09**	**-0.08**	**0.9038**
			Survival ~ AncestryProp + Lake * Experiment	126.09	-0.08	0.9038
		Molluscivore	Survival ~ AncestryProp	188.78	-0.05	0.9246
			Survival ~ AncestryProp + Lake	156.09	0.89	0.1683
			Survival ~ AncestryProp + Experiment	160.74	-0.70	0.2264
			Survival ~ AncestryProp + Lake + Experiment	**123.85**	**0.36**	**0.6142**
			Survival ~ AncestryProp + Lake * Experiment	125.85	0.36	0.6142
		Scale-eater	Survival ~ AncestryProp	188.58	0.35	0.6522
			Survival ~ AncestryProp + Lake	155.46	1.36	0.1160
			Survival ~ AncestryProp + Experiment	158.78	-1.70	0.0770
			Survival ~ AncestryProp + Lake + Experiment	**123.77**	**-0.64**	**0.5753**
			Survival ~ AncestryProp + Lake * Experiment	125.77	-0.64	0.5753

Appendix 1—Table 4. Models tested to assess the extent to which specialist ancestry predicts measures of fitness and their respective fits using only samples from the second field experiment (Martin and Gould 2020) and an unsupervised admixture analysis. Best-fit models are bolded.

Fitness Measure	Model Family	Species	Model	AIC	Ancestry Coefficient	Ancestry *P*-value
Composite	**Tobit**	**Generalist**	Composite ~ AncestryProp	187.62	0.11	0.5151
			Composite ~ AncestryProp + Lake	**179.83**	**-0.13**	**0.5004**
			Composite ~ AncestryProp * Lake	181.65	-0.04	0.8924
		Molluscivore	Composite ~ AncestryProp	188.04	0.02	0.9170
			Composite ~ AncestryProp + Lake	**178.97**	**0.24**	**0.2536**
			Composite ~ AncestryProp * Lake	179.95	-0.04	0.9036
		Scale-eater	Composite ~ AncestryProp	186.12	-0.43	0.1743
			Composite ~ AncestryProp + Lake	**179.90**	**-0.20**	**0.5361**
			Composite ~ AncestryProp * Lake	180.85	0.28	0.6288
Growth	**Gaussian**	**Generalist**	Growth ~ AncestryProp	27.50	-0.34	0.0070
			Growth ~ AncestryProp + Lake	-51.14	0.08	0.2452
			Growth ~ AncestryProp * Lake	**-53.47**	**0.01**	**0.9339**
		Molluscivore	Growth ~ AncestryProp	28.21	0.38	0.0103
			Growth ~ AncestryProp + Lake	**-50.78**	**-0.08**	**0.3149**
			Growth ~ AncestryProp * Lake	-49.16	-0.05	0.6565
		Scale-eater	Growth ~ AncestryProp	34.00	0.29	0.3072
			Growth ~ AncestryProp + Lake	-50.03	-0.08	0.5823
			Growth ~ AncestryProp * Lake	**-52.24**	**0.09**	**0.5772**
Survival	**Binomial**	**Generalist**	Survival ~ AncestryProp	157.06	0.92	0.0793
			Survival ~ AncestryProp + Lake	**121.49**	**-0.54**	**0.4400**
			Survival ~ AncestryProp * Lake	123.46	-0.36	0.7660
		Molluscivore	Survival ~ AncestryProp	159.41	-0.53	0.3709
			Survival ~ AncestryProp + Lake	**121.06**	**0.76**	**0.3127**
			Survival ~ AncestryProp * Lake	122.70	-0.04	0.9788
		Scale-eater	Survival ~ AncestryProp	156.99	-1.69	0.0828
			Survival ~ AncestryProp + Lake	**122.04**	**-0.28**	**0.8041**
			Survival ~ AncestryProp * Lake	123.24	2.00	1.0000

Appendix 1—Table 5. Models tested to assess the extent to which genome-wide variation (PC1/PC2) predicts measures of fitness and their respective fits using all samples and an unsupervised admixture analysis. Best-fit models are bolded.

Fitness Measure	Model Family	Principle Component	Model	AIC	Ancestry Coefficient	Ancestry *P*-value
Composite	Tobit	PC1	**Composite ~ PC1**	**193.80**	**1.67**	**0.0043**
			Composite ~ PC1 + Lake	195.52	3.07	0.2586
			Composite ~ PC1 + Experiment	194.27	1.80	0.0030
			Composite ~ PC1 + Lake + Experiment	195.36	4.52	0.1233
			Composite ~ PC1 + Lake * Experiment	197.07	3.62	0.2831
		PC2	Composite ~ PC2	202.44	-0.01	0.9907
			Composite ~ PC2 + Lake	195.45	1.20	0.2456
			Composite ~ PC2 + Experiment	203.75	0.21	0.8294
			Composite ~ PC2 + Lake + Experiment	**195.21**	**1.79**	**0.1130**
			Composite ~ PC2 + Lake * Experiment	196.35	1.57	0.1738
Growth	Gaussian	PC1	Growth ~ PC1	30.97	-1.62	0.0006
			Growth ~ PC1 + Lake	29.37	1.57	0.3723
			Growth ~ PC1 + Experiment	-32.37	-2.92	0.0000
			Growth ~ PC1 + Lake + Experiment	-30.92	-3.87	0.0055
			Growth ~ PC1 + Lake * Experiment	**-87.40**	**0.33**	**0.7638**
		PC2	Growth ~ PC2	23.61	2.88	0.0002
			Growth ~ PC2 + Lake	23.61	2.00	0.0119
			Growth ~ PC2 + Experiment	14.04	2.04	0.0044
			Growth ~ PC2 + Lake + Experiment	-23.65	-0.61	0.3677
			Growth ~ PC2 + Lake * Experiment	**-87.49**	**0.20**	**0.6724**
Survival	Binomial	PC1	Survival ~ PC1	156.24	15.09	0.0000
			Survival ~ PC1 + Lake	157.87	6.04	0.6943
			Survival ~ PC1 + Experiment	**121.52**	**17.78**	**0.0000**
			Survival ~ PC1 + Lake + Experiment	123.51	15.92	0.4386
			Survival ~ PC1 + Lake * Experiment	125.51	15.92	0.4386
		PC2	Survival ~ PC2	157.96	-9.44	0.0268
			Survival ~ PC2 + Lake	157.96	-1.21	0.7988
			Survival ~ PC2 + Experiment	160.72	-5.63	0.2248
			Survival ~ PC2 + Lake + Experiment	**123.39**	**4.89**	**0.4047**
			Survival ~ PC2 + Lake * Experiment	125.39	4.89	0.4047

Appendix 1—Table 6. SNPs found to be strongly associated with composite fitness using SnpEff (136)**.** SNPs that were identified as being strongly associated with both growth and composite fitness are italicized, and those that remain significant after a Bonferroni correction are bolded.

Scaffold	Position	Significance	REF	ALT	Variant Type	Gene Identifier	Gene Card
HiC_scaffold_1	32071263	FDR	A	C	intergenic	CBRO_00000660-CBRO_00000661	Znf250-Ptgdr2
HiC_scaffold_1	**43866598**	**Bonferroni**	**G**	**A**	**intergenic**	**CBRO_00000910-CBRO_00000911**	**PPM1K-OVCH2**
HiC_scaffold_1	43867614	FDR	C	T	intergenic	CBRO_00000910-CBRO_00000911	PPM1K-OVCH2
HiC_scaffold_3	2658774	FDR	T	A	intergenic	CBRO_00017856-CBRO_00017857	KMT2E-Magi2
HiC_scaffold_3	2658775	FDR	T	C	intergenic	CBRO_00017856-CBRO_00017857	KMT2E-Magi2
HiC_scaffold_3	2658793	FDR	C	A	intergenic	CBRO_00017856-CBRO_00017857	KMT2E-Magi2
HiC_scaffold_4	18899496	FDR	T	C	intergenic	CBRO_00012427-CBRO_00012428	UNKNOWN-edc4
HiC_scaffold_5	18306419	FDR	C	G	upstream; intergenic	CBRO_00001232; CBRO_00001231-CBRO_00001232	xlrs1; PPEF2-xlrs1
HiC_scaffold_5	18306428	FDR	A	T	upstream; intergenic	CBRO_00001232; CBRO_00001231-CBRO_00001232	xlrs1; PPEF2-xlrs1
HiC_scaffold_5	18307019	FDR	G	T	intronic	CBRO_00001232	xlrs1
HiC_scaffold_5	18307030	FDR	G	A	intronic	CBRO_00001232	xlrs1
HiC_scaffold_5	18311696	FDR	C	T	intronic	CBRO_00001232	xlrs1
HiC_scaffold_5	40475116	FDR	T	A	intergenic	CBRO_00001627-CBRO_00001628	SLC25A44-UBE2Q2
HiC_scaffold_7	10290141	FDR	T	C	downstream; intergenic	CBRO_00009717; CBRO_00009716-CBRO_00009717	Nfkbie; SLC35B2-Nfkbie
HiC_scaffold_7	10290142	FDR	G	C	downstream; intergenic	CBRO_00009717; CBRO_00009716-CBRO_00009717	Nfkbie; SLC35B2-Nfkbie
HiC_scaffold_7	10290165	FDR	A	G	downstream; intergenic	CBRO_00009717; CBRO_00009716-CBRO_00009717	Nfkbie; SLC35B2-Nfkbie
HiC_scaffold_7	10290166	FDR	T	C	downstream; intergenic	CBRO_00009717; CBRO_00009716-CBRO_00009717	Nfkbie; SLC35B2-Nfkbie
HiC_scaffold_7	10290168	FDR	T	C	downstream; intergenic	CBRO_00009717; CBRO_00009716-CBRO_00009717	Nfkbie; SLC35B2-Nfkbie
HiC_scaffold_7	13815058	FDR	A	G	synonymous; downstream	CBRO_00009805; CBRO_00009804	Aloxe3; UNKNOWN
HiC_scaffold_7	13815326	FDR	C	T	intronic	CBRO_00009805	Aloxe3
HiC_scaffold_7	15349830	FDR	C	A	intergenic	CBRO_00009834-CBRO_00009835	Adcy8-efr3b

HiC_scaffold_7	18378061	FDR	T	A	intergenic	CBRO_00009902-CBRO_00009903	Fam84a-DDX1
HiC_scaffold_8	**20263964**	**Bonferroni**	**G**	**A**	**upstream; intergenic**	**CBRO_00010647; CBRO_00010647-CBRO_00010648**	**Srcin1; Srcin1-Srcin1**
HiC_scaffold_8	31539262	FDR	G	A	upstream; intergenic	CBRO_00010835; CBRO_00010834-CBRO_00010835	Gjd3; Gjd3-Gjd3
HiC_scaffold_11	3106766	FDR	T	A	intronic	CBRO_00013361	prkdc
HiC_scaffold_11	3138733	FDR	G	A	intergenic	CBRO_00013361-CBRO_00013362	prkdc-arhgap29
HiC_scaffold_11	5658921	FDR	A	G	intronic	CBRO_00013404	KAZN
HiC_scaffold_14	17556007	FDR	C	T	intergenic	CBRO_00014134-CBRO_00014135	UNKNOWN-Abr
HiC_scaffold_14	17556026	FDR	C	A	intergenic	CBRO_00014134-CBRO_00014135	UNKNOWN-Abr
HiC_scaffold_16	32837191	FDR	T	C	intronic	CBRO_00003226	KIF1B
HiC_scaffold_16	35727503	FDR	G	C	intergenic	CBRO_00003289-CBRO_00003290	Pip5k1c-Polr2e
HiC_scaffold_16	40215889	FDR	G	A	intergenic	CBRO_00003383-CBRO_00003384	chst10-UNKNOWN
HiC_scaffold_16	40300592	FDR	C	A	intergenic	CBRO_00003384-CBRO_00003385	UNKNOWN-Carmil3
HiC_scaffold_18	26969972	FDR	T	G	intronic	CBRO_00013239	CSAD
HiC_scaffold_18	26970123	FDR	C	T	synonymous	CBRO_00013239	CSAD
HiC_scaffold_18	26970601	FDR	T	A	intronic	CBRO_00013239	CSAD
HiC_scaffold_18	26978410	FDR	G	A	intronic	CBRO_00013240	Znf740
HiC_scaffold_20	332642	FDR	G	A	missense	CBRO_00016084	GTF3C4
HiC_scaffold_20	332689	FDR	A	G	missense	CBRO_00016084	GTF3C4
HiC_scaffold_20	14820437	FDR	A	T	intronic	CBRO_00016304	MALT1
HiC_scaffold_20	14820448	FDR	T	C	intronic	CBRO_00016304	MALT1
HiC_scaffold_24	1469530	FDR	G	T	intergenic	CBRO_00014375-CBRO_00014376	UNKNOWN-UNKNOWN
HiC_scaffold_24	3223583	FDR	T	A	intergenic	CBRO_00014421-CBRO_00014422	Gal3st3-RIN2
HiC_scaffold_24	11618442	FDR	T	G	intronic	CBRO_00014601	ABCA4
HiC_scaffold_24	**15964553**	**Bonferroni**	**C**	**T**	**intergenic**	**CBRO_00014635-CBRO_00014636**	**Lrfn2-SNX15**
HiC_scaffold_27	1898180	FDR	G	A	downstream; intergenic	CBRO_00005887; CBRO_00005886-CBRO_00005887	UNKNOWN; hoxb13a-UNKNOWN
HiC_scaffold_27	8137335	FDR	G	T	intronic	CBRO_00005998	SMARCA4
HiC_scaffold_27	9065056	FDR	C	A	intergenic	CBRO_00006026-CBRO_00006027	ANKFN1-ccdc134
HiC_scaffold_27	12370585	FDR	C	T	intergenic	CBRO_00006131-CBRO_00006132	SHISA9-Desi1

HiC_scaffold_27	32078665	FDR	C	G	intergenic	CBRO_00006685-CBRO_00006686	med25-Lrrc4b
HiC_scaffold_27	34904388	FDR	A	C	intergenic	CBRO_00006756-CBRO_00006757	Grin2c-nog3
HiC_scaffold_27	35431570	FDR	G	C	intergenic	CBRO_00006756-CBRO_00006757	Grin2c-nog3
HiC_scaffold_27	35431578	FDR	C	T	intergenic	CBRO_00006756-CBRO_00006757	Grin2c-nog3
HiC_scaffold_27	35431585	FDR	A	G	intergenic	CBRO_00006756-CBRO_00006757	Grin2c-nog3
HiC_scaffold_34	16654675	FDR	G	C	intergenic	CBRO_00001997-CBRO_00001998	MDFIC2-foxp1b
HiC_scaffold_34	19393244	FDR	A	T	upstream; intergenic	CBRO_00002027; CBRO_00002027-CBRO_00002028	SUOX; SUOX-SUOX
HiC_scaffold_34	22010499	FDR	C	T	intronic	CBRO_00002117	GNAI2
HiC_scaffold_34	31220916	FDR	T	A	upstream; intergenic	CBRO_00002389; CBRO_00002388-CBRO_00002389	CTTNBP2NL; Kcnd3-CTTNBP2NL
HiC_scaffold_34	37769304	FDR	A	C	intergenic	CBRO_00002519-CBRO_00002520	ASIC2-asic1
HiC_scaffold_37	5963405	FDR	T	C	intergenic	CBRO_00011020-CBRO_00011021	C14orf93-pim2
HiC_scaffold_37	11017168	FDR	G	A	intergenic	CBRO_00011155-CBRO_00011156	GALNT12-elp2
HiC_scaffold_37	13822135	FDR	A	G	upstream; intergenic	CBRO_00011221; CBRO_00011220-CBRO_00011221	SATB1; KCNH8-SATB1
HiC_scaffold_37	13823678	FDR	G	A	upstream; intergenic	CBRO_00011221; CBRO_00011220-CBRO_00011221	SATB1; KCNH8-SATB1
HiC_scaffold_37	13832007	FDR	T	A	missense	CBRO_00011221	SATB1
HiC_scaffold_37	16920863	FDR	T	C	intergenic	CBRO_00011259-CBRO_00011260	CSMD1-UNKNOWN
HiC_scaffold_37	**18591438**	**Bonferroni**	**G**	**A**	**intergenic**	**CBRO_00011301-CBRO_00011302**	**cck-trim71**
HiC_scaffold_37	18591463	FDR	C	T	intergenic	CBRO_00011301-CBRO_00011302	cck-trim71
HiC_scaffold_37	18596716	FDR	A	T	intergenic	CBRO_00011301-CBRO_00011302	cck-trim71
HiC_scaffold_40	5885291	FDR	G	T	downstream; intergenic	CBRO_00016614; CBRO_00016613-CBRO_00016614	C14orf93 homolog; HTR2A-C14orf93 homolog
HiC_scaffold_43	2568535	FDR	A	G	intergenic	CBRO_00008246-CBRO_00008247	UNKNOWN-Gpr68
HiC_scaffold_44	25886134	FDR	A	C	intergenic	CBRO_00007276-CBRO_00007277	RAPGEF2-QDPR
HiC_scaffold_45	885834	FDR	A	C	upstream; intergenic	CBRO_00018766; CBRO_00018766-CBRO_00018767	Mog; Mog-EPHB4
HiC_scaffold_45	2213441	FDR	G	T	intronic	CBRO_00018814	Nlrp12
HiC_scaffold_46	856248	FDR	A	G	intergenic	CBRO_00007439-CBRO_00007440	UNKNOWN-NLRP12
HiC_scaffold_46	1232350	FDR	G	A	intronic	CBRO_00007448	NEB

HiC_scaffold_46	30183758	FDR	C	G	intergenic	CBRO_00008059-CBRO_00008060	TFRC-Rgs11
HiC_scaffold_46	32048848	FDR	A	C	intronic	CBRO_00008105	Hsd17b7
HiC_scaffold_46	32050109	FDR	G	T	synonymous	CBRO_00008105	Hsd17b7
HiC_scaffold_46	**35151009**	**Bonferroni**	**T**	**C**	**downstream; intergenic**	**CBRO_00008165; CBRO_00008165-CBRO_00008166**	**Klf9; Klf9-Tsen15**
HiC_scaffold_46	35163681	FDR	T	C	downstream; intergenic	CBRO_00008166; CBRO_00008167; CBRO_00008166-CBRO_00008167	Tsen15; UNKNOWN; Tsen15-UNKNOWN
HiC_scaffold_46	35164267	FDR	A	G	downstream; intergenic	CBRO_00008166; CBRO_00008167; CBRO_00008166-CBRO_00008167	Tsen15; UNKNOWN; Tsen15-UNKNOWN
HiC_scaffold_47	787141	FDR	T	G	upstream; intergenic	CBRO_00008838; CBRO_00008837-CBRO_00008838	Nlrc3; NLRC3-Nlrc3
HiC_scaffold_47	4300295	FDR	G	A	downstream; intergenic	CBRO_00008933; CBRO_00008933-CBRO_00008934	NEK6; NEK6-Psmb7
HiC_scaffold_47	6767766	FDR	T	C	intergenic	CBRO_00008991-CBRO_00008992	TACR1-Grk5
HiC_scaffold_52	20791197	FDR	T	A	upstream; intronic	CBRO_00011892; CBRO_00011891	rabl3; GTF2E1
HiC_scaffold_52	22551701	FDR	G	C	intergenic	CBRO_00011922-CBRO_00011923	ALS2-Serp2
HiC_scaffold_52	31021517	FDR	C	T	intergenic	CBRO_00012051-CBRO_00012052	Tmeff2-slc39a10
HiC_scaffold_53	**11317840**	**Bonferroni**	**A**	**G**	**intergenic**	**CBRO_00005178-CBRO_00005179**	**Fucolectin-1-Fucolectin-5**
HiC_scaffold_53	11326035	FDR	C	T	intergenic	CBRO_00005178-CBRO_00005179	Fucolectin-1-Fucolectin-5
HiC_scaffold_53	11326410	FDR	G	A	intergenic	CBRO_00005178-CBRO_00005179	Fucolectin-1-Fucolectin-5
HiC_scaffold_53	11331079	FDR	A	G	intergenic	CBRO_00005178-CBRO_00005179	Fucolectin-1-Fucolectin-5
HiC_scaffold_53	15966447	FDR	G	A	upstream; downstream; intergenic	CBRO_00005235; CBRO_00005234; CBRO_00005234-CBRO_00005235	Tmem222; WDTC1; WDTC1-Tmem222
HiC_scaffold_53	17413090	FDR	A	G	missense	CBRO_00005274	Mag
HiC_scaffold_53	20715576	FDR	C	A	intergenic	CBRO_00005380-CBRO_00005381	Scrt2-Ino80c
HiC_scaffold_53	20715623	FDR	G	A	intergenic	CBRO_00005380-CBRO_00005381	Scrt2-Ino80c
HiC_scaffold_53	20715851	FDR	A	G	intergenic	CBRO_00005380-CBRO_00005381	Scrt2-Ino80c
HiC_scaffold_53	27386094	FDR	G	T	intronic	CBRO_00005625	Arhgef1
HiC_scaffold_53	27396961	FDR	T	A	intronic	CBRO_00005625	Arhgef1
HiC_scaffold_53	27398605	FDR	T	G	downstream; intergenic	CBRO_00005625; CBRO_00005625-CBRO_00005626	Arhgef1; Arhgef1-CD79A
HiC_scaffold_53	33282501	FDR	A	G	intergenic	CBRO_00005732-CBRO_00005733	mios-GLCCI1

HiC_scaffold_53	39228193	FDR	T	C	intergenic	CBRO_00005830-CBRO_00005831	UNKNOWN-GnRHR2
HiC_scaffold_53	**39228264**	**Bonferroni**	**C**	**T**	**intergenic**	**CBRO_00005830-CBRO_00005831**	**UNKNOWN-GnRHR2**
HiC_scaffold_53	39228942	FDR	A	T	intergenic	CBRO_00005830-CBRO_00005831	UNKNOWN-GnRHR2
HiC_scaffold_53	39279263	FDR	G	T	intronic	CBRO_00005832	IGDCC3
HiC_scaffold_53	39770914	FDR	A	T	intronic	CBRO_00005849	Cpne4
HiC_scaffold_611	5621	FDR	A	T	intergenic	CHR_START-CBRO_00020243	CHR_START-fzdz-a
HiC_scaffold_611	5625	FDR	G	T	intergenic	CHR_START-CBRO_00020243	CHR_START-fzdz-a
HiC_scaffold_611	5634	FDR	T	A	intergenic	CHR_START-CBRO_00020243	CHR_START-fzdz-a
HiC_scaffold_1053	3494	FDR	A	G	downstream; intergenic	CBRO_00020503; CHR_START-CBRO_00020503	UBE2G2; CHR_START-UBE2G2
HiC_scaffold_1133	9946	FDR	A	G	intergenic	.	.
HiC_scaffold_1371	7318	FDR	C	T	upstream; intergenic	CBRO_00021026; CHR_START-CBRO_00021026	UNKNOWN; CHR_START-UNKNOWN
HiC_scaffold_1848	**40119**	**Bonferroni**	**C**	**A**	**intergenic**	**.**	**.**
HiC_scaffold_1848	**40465**	**Bonferroni**	**T**	**A**	**intergenic**	**.**	**.**
HiC_scaffold_1848	**40590**	**Bonferroni**	**T**	**C**	**intergenic**	**.**	**.**
HiC_scaffold_1848	40877	FDR	T	C	intergenic	.	.
HiC_scaffold_1848	41351	FDR	T	C	intergenic	.	.
HiC_scaffold_2220	10128	FDR	G	T	intergenic	.	.
HiC_scaffold_4461	**12939**	**Bonferroni**	**T**	**A**	**intergenic**	**.**	**.**
HiC_scaffold_4665	13941	FDR	C	T	intergenic	.	.
HiC_scaffold_6275	6000	FDR	C	A	intergenic	.	.
HiC_scaffold_6337	5745	FDR	T	G	intronic	CBRO_00021217	PKP3
HiC_scaffold_6769	2796	FDR	G	A	intergenic	.	.
HiC_scaffold_6963	5970	FDR	C	A	intergenic	.	.
HiC_scaffold_6963	6101	FDR	A	G	intergenic	.	.
HiC_scaffold_9280	3448	FDR	G	A	intergenic	.	.
HiC_scaffold_9949	52	FDR	C	T	intergenic	.	.
HiC_scaffold_10928	3575	FDR	G	A	downstream; intergenic	CBRO_00021896; CHR_START-CBRO_00021896	CYP2A10; CHR_START-CYP2A10
HiC_scaffold_11921	5560	FDR	A	T	intergenic	.	.
HiC_scaffold_12068	2929	FDR	G	T	intergenic	.	.

HiC_scaffold_1277 8	**1456**	**Bonferroni**	**T**	**A**	**missense**	**CBRO_00022026**	**COL8A1**
HiC_scaffold_17578	180	FDR	T	G	intergenic	.	.
HiC_scaffold_18999	**1084**	**Bonferroni**	**A**	**G**	**intergenic**	**.**	**.**

Appendix 1—Table 7. Gene ontology term enrichment for genes associated with composite fitness.

Functional Category	Enrichment FDR	Genes in list	Total genes	Genes
Anterograde trans-synaptic signaling	0.0060	13	746	GRIN2C, SHISA9, ALS2, KIF1B, ASIC1, GNAI2, TACR1, ADCY8, LRFN2, ABR, RAPGEF2, ASIC2, PIP5K1C
Synaptic signaling	0.0060	13	762	GRIN2C, SHISA9, ALS2, KIF1B, ASIC1, GNAI2, TACR1, ADCY8, LRFN2, ABR, RAPGEF2, ASIC2, PIP5K1C
Trans-synaptic signaling	0.0060	13	756	GRIN2C, SHISA9, ALS2, KIF1B, ASIC1, GNAI2, TACR1, ADCY8, LRFN2, ABR, RAPGEF2, ASIC2, PIP5K1C
Chemical synaptic transmission	0.0060	13	746	GRIN2C, SHISA9, ALS2, KIF1B, ASIC1, GNAI2, TACR1, ADCY8, LRFN2, ABR, RAPGEF2, ASIC2, PIP5K1C
Regulation of I-kappaB kinase/NF-kappaB signaling	0.0179	7	241	NEK6, SLC35B2, MALT1, NLRP12, NLRC3, DDX1, PIM2
Regulation of trans-synaptic signaling	0.0185	9	468	GRIN2C, SHISA9, ASIC1, GNAI2, TACR1, ADCY8, LRFN2, ABR, RAPGEF2
Regulation of cell communication	0.0185	32	3903	NEK6, SLC35B2, MALT1, NLRP12, PPEF2, GRIN2C, NLRC3, SLC39A10, SHISA9, ALS2, RAPGEF2, SMARCA4, MAGI2, RGS11, ARHGEF1, DDX1, PIM2, ASIC1, GNAI2, TACR1, ADCY8, LRFN2, ABR, PTGDR2, CCK, GPR68, MIOS, SCRT2, PSMB7, ARHGAP29, PIP5K1C, GRK5
Regulation of signaling	0.0185	32	3952	NEK6, SLC35B2, MALT1, NLRP12, PPEF2, GRIN2C, NLRC3, SLC39A10, SHISA9, ALS2, RAPGEF2, SMARCA4, MAGI2, RGS11, ARHGEF1, DDX1, PIM2, ASIC1, GNAI2, TACR1, ADCY8, LRFN2, ABR, PTGDR2, CCK, GPR68, MIOS, SCRT2, PSMB7, ARHGAP29, PIP5K1C, GRK5
Modulation of chemical synaptic transmission	0.0185	9	467	GRIN2C, SHISA9, ASIC1, GNAI2, TACR1, ADCY8, LRFN2, ABR, RAPGEF2
Sensory perception of sour taste	0.0185	2	5	ASIC2, ASIC1
Regulation of biological quality	0.0185	34	4319	GRIN2C, KCNH8, SLC39A10, SHISA9, PRKDC, TACR1, SMARCA4, KCND3, ABCA4, TRIM71, KMT2E, TFRC, PIM2, MAG, ASIC2, ASIC1, GNAI2, LRRC4B, HSD17B7, WDTC1, ADCY8, LRFN2, ABR, ALOXE3, CSMD1, CCK, SRCIN1, ALS2, GPR68, RAPGEF2, NEB, PSMB7, NFKBIE, PIP5K1C
I-kappaB kinase/NF-kappaB signaling	0.0185	7	280	NEK6, SLC35B2, MALT1, NLRP12, NLRC3, DDX1, PIM2
Positive regulation of catalytic activity	0.0185	17	1514	PPM1K, ALS2, RAPGEF2, MAGI2, SRCIN1, RGS11, ARHGEF1, GNAI2, GTF3C4, RIN2, ARHGAP29, ABR, CCK, SLC39A10, MALT1, ADCY8, NLRP12
Regulation of protein tyrosine phosphatase activity	0.0216	2	6	GNAI2, SLC39A10
Multicellular organismal response to stress	0.0216	4	74	ALS2, ASIC1, TACR1, CCK
Positive regulation of molecular function	0.0270	19	1901	PPM1K, MALT1, ALS2, RAPGEF2, SMARCA4, MAGI2, SRCIN1, RGS11, ARHGEF1, GNAI2, GTF3C4, RIN2, ARHGAP29, ADCY8, ABR, CCK, SLC39A10, MED25, NLRP12
Cell-cell signaling	0.0311	18	1774	GRIN2C, SHISA9, ALS2, KIF1B, ASIC1, GNAI2, TACR1, ADCY8, LRFN2, ABR, GRK5, GPR68, SMARCA4, RAPGEF2, MAGI2, ASIC2, PSMB7, PIP5K1C
Tachykinin receptor signaling pathway	0.0314	2	8	TACR1, GRK5
Cell proliferation	0.0355	20	2165	TFRC, PIM2, RAPGEF2, CD79A, GNAI2, TACR1, COL8A1, MALT1, SATB1, PTGDR2, CCK, SLC39A10, TRIM71, PRKDC, MED25, KLF9, GRK5, NLRC3, MAGI2, ARHGEF1
Behavioral fear response	0.0355	3	39	ALS2, ASIC1, CCK
Response to pH	0.0355	3	41	ASIC1, ASIC2, GPR68
Behavioral defense response	0.0355	3	40	ALS2, ASIC1, CCK
Fear response	0.0355	3	41	ALS2, ASIC1, CCK
Positive regulation of hydrolase activity	0.0401	11	833	ALS2, RAPGEF2, MAGI2, RGS11, ARHGEF1, RIN2, ARHGAP29, ABR, CCK, SLC39A10, NLRP12
NcRNA transcription	0.0457	4	110	GTF3C4, SMARCA4, POLR2E, GTF2E1
Hexose mediated signaling	0.0457	2	13	SMARCA4, ADCY8
Sugar mediated signaling pathway	0.0457	2	13	SMARCA4, ADCY8
Glucose mediated signaling pathway	0.0457	2	13	SMARCA4, ADCY8
Positive regulation of cell communication	0.0457	18	1937	NEK6, SLC35B2, MALT1, GRIN2C, SLC39A10, ALS2, RAPGEF2, SMARCA4, NLRP12, DDX1, PIM2, GNAI2, TACR1, ADCY8, PTGDR2, GPR68, MIOS, PSMB7
Positive regulation of signaling	0.0457	18	1945	NEK6, SLC35B2, MALT1, GRIN2C, SLC39A10, ALS2, RAPGEF2, SMARCA4, NLRP12, DDX1, PIM2, GNAI2, TACR1, ADCY8, PTGDR2, GPR68, MIOS, PSMB7

Appendix 1—Table 8. SNPs found to be strongly associated with growth SnpEff (136)**.** SNPs that were identified as being strongly associated with both growth and composite fitness are italicized, and those that remain significant after a Bonferroni correction are bolded.

Scaffold	Position	Significance	REF	ALT	Variant Type	Gene Identifier	Gene Card
HiC_scaffold_4	8897671	FDR	A	T	intergenic	CBRO_00012254-CBRO_00012255	Megf10-UNKNOWN
HiC_scaffold_4	16057698	FDR	C	A	intergenic	CBRO_00012386-CBRO_00012387	CCND2-Mlycd
HiC_scaffold_4	29273446	FDR	T	C	synonymous	CBRO_00012620	CKAP5
HiC_scaffold_4	29273458	FDR	C	T	synonymous	CBRO_00012620	CKAP5
HiC_scaffold_5	18259775	FDR	C	A	upstream; intergenic	CBRO_00001229; CBRO_00001229-CBRO_00001230	AP1S2; AP1S2-phka2
HiC_scaffold_5	*18306419*	*FDR*	*C*	*G*	*upstream; intergenic*	*CBRO_00001232; CBRO_00001231-CBRO_00001232*	*xlrs1; PPEF2-xlrs1*
HiC_scaffold_5	*18306428*	*FDR*	*A*	*T*	*upstream; intergenic*	*CBRO_00001232; CBRO_00001231-CBRO_00001232*	*xlrs1; PPEF2-xlrs1*
HiC_scaffold_5	*18307019*	*FDR*	*G*	*T*	*intronic*	*CBRO_00001232*	*xlrs1*
HiC_scaffold_5	*18307030*	*FDR*	*G*	*A*	*intronic*	*CBRO_00001232*	*xlrs1*
HiC_scaffold_5	36619253	FDR	T	C	intergenic	CBRO_00001583-CBRO_00001584	Chst12-ZDHHC13
HiC_scaffold_7	*13815058*	*FDR*	*A*	*G*	*synonymous; downstream*	*CBRO_00009805; CBRO_00009804*	*Aloxe3; UNKNOWN*
HiC_scaffold_7	13823565	FDR	T	C	intronic	CBRO_00009806	Fbxo30
HiC_scaffold_7	13824467	FDR	A	G	missense	CBRO_00009806	Fbxo30
HiC_scaffold_7	19371997	FDR	C	T	upstream; intergenic	CBRO_00009922; CBRO_00009921-CBRO_00009922	ELOVL4; TENT5A-ELOVL4
HiC_scaffold_7	19372002	FDR	T	G	upstream; intergenic	CBRO_00009922; CBRO_00009921-CBRO_00009922	ELOVL4; TENT5A-ELOVL4
HiC_scaffold_8	20265076	FDR	T	C	upstream; intergenic	CBRO_00010647; CBRO_00010647-CBRO_00010648	Srcin1; Srcin1-Srcin1
HiC_scaffold_8	20265098	FDR	G	C	upstream; intergenic	CBRO_00010647; CBRO_00010647-CBRO_00010648	Srcin1; Srcin1-Srcin1
HiC_scaffold_8	20278571	FDR	G	A	intergenic	CBRO_00010647-CBRO_00010648	Srcin1-Srcin1
HiC_scaffold_9	15585466	FDR	C	G	missense	CBRO_00004552	Tmem260
HiC_scaffold_9	18453213	FDR	A	G	intergenic	CBRO_00004639-CBRO_00004640	Bub1b-PAK6
HiC_scaffold_9	28127377	FDR	C	T	intergenic	CBRO_00004857-CBRO_00004858	METTL21E-RASA3
HiC_scaffold_10	193812	FDR	C	G	intergenic	CBRO_00018931-CBRO_00018932	Spsb4-UNKNOWN

HiC_scaffold_11	25632195	FDR	C	T	intergenic	CBRO_00013717-CBRO_00013718	CDH10-Cdh6
HiC_scaffold_11	25632258	FDR	C	T	intergenic	CBRO_00013717-CBRO_00013718	CDH10-Cdh6
HiC_scaffold_11	25632641	FDR	A	G	intergenic	CBRO_00013717-CBRO_00013718	CDH10-Cdh6
HiC_scaffold_14	14624430	FDR	T	C	intergenic	CBRO_00014084-CBRO_00014085	KDM6B-FGF11
HiC_scaffold_18	26970449	FDR	C	T	intronic	CBRO_00013239	CSAD
HiC_scaffold_27	*8137335*	*FDR*	*G*	*T*	*intronic*	*CBRO_00005998*	*SMARCA4*
HiC_scaffold_27	21919164	FDR	G	T	synonymous	CBRO_00006396	SSTR2
HiC_scaffold_29	2136546	FDR	A	T	upstream; intergenic	CBRO_00016347; CBRO_00016347-CBRO_00016348	RAPGEF6; RAPGEF6-ACSL6
HiC_scaffold_29	2147361	FDR	C	T	intergenic	CBRO_00016347-CBRO_00016348	RAPGEF6-ACSL6
HiC_scaffold_31	6226140	FDR	A	C	intronic	CBRO_00017224	ranbp9
HiC_scaffold_34	7439419	FDR	G	T	intergenic	CBRO_00001828-CBRO_00001829	rnf152-CAMTA1
HiC_scaffold_34	30571660	FDR	C	A	intronic	CBRO_00002374	SHMT2
HiC_scaffold_40	4693917	FDR	G	A	intronic	CBRO_00016589	SLC37A1
HiC_scaffold_40	4694015	FDR	T	C	synonymous	CBRO_00016589	SLC37A1
HiC_scaffold_40	4723358	FDR	A	G	intronic	CBRO_00016591	UBXN4
HiC_scaffold_40	4732538	FDR	T	C	intronic	CBRO_00016591	UBXN4
HiC_scaffold_40	4734825	FDR	G	T	intronic	CBRO_00016591	UBXN4
HiC_scaffold_40	4780518	FDR	G	C	intronic	CBRO_00016592	ITGB2
HiC_scaffold_40	5016538	FDR	C	A	upstream; intergenic	CBRO_00016599; CBRO_00016599-CBRO_00016600	RRP1B; RRP1B-ITGB2
HiC_scaffold_43	26869231	FDR	C	G	synonymous	CBRO_00008677	lrpprc
HiC_scaffold_43	27800569	FDR	C	T	downstream; intergenic	CBRO_00008685; CBRO_00008685-CBRO_00008686	timp3; timp3-ETV6
HiC_scaffold_44	19344526	FDR	C	T	intergenic	CBRO_00007178-CBRO_00007179	ATP11C-sox3
HiC_scaffold_46	16495639	FDR	G	A	intronic	CBRO_00007743	PHLPP1
HiC_scaffold_46	16510323	FDR	G	A	intronic	CBRO_00007743	PHLPP1
HiC_scaffold_46	16512668	FDR	T	A	intronic	CBRO_00007743	PHLPP1
HiC_scaffold_46	**16512886**	**Bonferroni**	**T**	**A**	**intronic**	**CBRO_00007743**	**PHLPP1**
HiC_scaffold_46	16513809	FDR	T	C	synonymous	CBRO_00007743	PHLPP1
HiC_scaffold_52	19031083	FDR	T	A	intergenic	CBRO_00011872-CBRO_00011873	NRP2-MREG

HiC_scaffold_52	19031216	FDR	A	T	intergenic	CBRO_00011872-CBRO_00011873	NRP2-MREG
HiC_scaffold_1848	***40119***	***Bonferroni***	***C***	***A***	***intergenic***	***.***	***.***
HiC_scaffold_1848	***40465***	***Bonferroni***	***T***	***A***	***intergenic***	***.***	***.***
HiC_scaffold_1848	*40590*	*FDR*	*T*	*C*	*intergenic*	*.*	*.*
HiC_scaffold_1848	*40877*	*FDR*	*T*	*C*	*intergenic*	*.*	*.*
HiC_scaffold_1848	*41351*	*FDR*	*T*	*C*	*intergenic*	*.*	*.*
HiC_scaffold_7644	**5971**	**Bonferroni**	**T**	**G**	**intergenic**	**.**	**.**
HiC_scaffold_12681	4019	FDR	T	C	upstream; intergenic	CBRO_00022068; CBRO_00022068-CHR_END	UNKNOWN; UNKNOWN-CHR_END

Appendix 1—Table 9. Gene ontology term enrichment for genes associated with growth.

Functional Category	Enrichment FDR	Genes in list	Total genes	Genes
Negative regulation of intracellular signal transduction	0.0259	6	537	PPEF2, RNF152, RASA3, RANBP9, PHLPP1, TIMP3
Phosphorus metabolic process	0.0259	17	3597	PHLPP1, CCND2, PAK6, PPEF2, SHMT2, MLYCD, RRP1B, ITGB2, SRCIN1, PHKA2, ELOVL4, BUB1B, RANBP9, TIMP3, CAMTA1, ACSL6, RASA3
Phosphate-containing compound metabolic process	0.0259	17	3570	PHLPP1, CCND2, PAK6, PPEF2, SHMT2, MLYCD, RRP1B, ITGB2, SRCIN1, PHKA2, ELOVL4, BUB1B, RANBP9, TIMP3, CAMTA1, ACSL6, RASA3
Serine family amino acid catabolic process	0.0259	2	17	SHMT2, CSAD
Organic acid biosynthetic process	0.0259	6	466	ELOVL4, SHMT2, MLYCD, CHST12, ALOXE3, CSAD
Regulation of phosphate metabolic process	0.0259	11	1870	PHLPP1, CCND2, PAK6, PPEF2, SHMT2, RRP1B, ITGB2, SRCIN1, RANBP9, TIMP3, CAMTA1
Nucleoside bisphosphate biosynthetic process	0.0259	3	82	MLYCD, ELOVL4, ACSL6
Ribonucleoside bisphosphate biosynthetic process	0.0259	3	82	MLYCD, ELOVL4, ACSL6
Purine nucleoside bisphosphate biosynthetic process	0.0259	3	82	MLYCD, ELOVL4, ACSL6
Long-chain fatty-acyl-CoA biosynthetic process	0.0259	2	19	ELOVL4, ACSL6
Thioester biosynthetic process	0.0259	3	66	MLYCD, ELOVL4, ACSL6
Positive regulation by host of viral transcription	0.0259	2	18	SMARCA4, RRP1B
Sulfur compound biosynthetic process	0.0259	5	207	MLYCD, CHST12, ELOVL4, CSAD, ACSL6
Regulation of phosphorus metabolic process	0.0259	11	1872	PHLPP1, CCND2, PAK6, PPEF2, SHMT2, RRP1B, ITGB2, SRCIN1, RANBP9, TIMP3, CAMTA1
Acyl-CoA biosynthetic process	0.0259	3	66	MLYCD, ELOVL4, ACSL6
Hypothalamus development	0.0291	2	22	NRP2, SOX3
Sulfur compound metabolic process	0.0312	5	393	MLYCD, CHST12, ELOVL4, CSAD, ACSL6
Long-chain fatty-acyl-CoA metabolic process	0.0343	2	26	ELOVL4, ACSL6
Regulation of mitochondrial translation	0.0343	2	26	LRPPRC, SHMT2
Limbic system development	0.0405	3	115	NRP2, KDM6B, SOX3
Acyl-CoA metabolic process	0.0406	3	119	MLYCD, ELOVL4, ACSL6
Thioester metabolic process	0.0406	3	119	MLYCD, ELOVL4, ACSL6
Protein phosphorylation	0.0412	11	2093	PHLPP1, CCND2, PAK6, PPEF2, ITGB2, SRCIN1, PHKA2, BUB1B, RANBP9, TIMP3, RASA3
Modulation by host of viral transcription	0.0412	2	33	SMARCA4, RRP1B
Carboxylic acid biosynthetic process	0.0412	5	465	ELOVL4, SHMT2, MLYCD, CHST12, ALOXE3
Modulation of transcription in other organism involved in symbiotic interaction	0.0412	2	34	SMARCA4, RRP1B
Modulation by host of symbiont transcription	0.0412	2	33	SMARCA4, RRP1B
Cellular protein modification process	0.0495	17	4434	PHLPP1, CCND2, PAK6, PPEF2, RNF152, ITGB2, SPSB4, SRCIN1, PHKA2, FBXO30, KDM6B, BUB1B, RANBP9, TIMP3, CAMTA1, SHMT2, RASA3
Protein modification process	0.0495	17	4434	PHLPP1, CCND2, PAK6, PPEF2, RNF152, ITGB2, SPSB4, SRCIN1, PHKA2, FBXO30, KDM6B, BUB1B, RANBP9, TIMP3, CAMTA1, SHMT2, RASA3

Appendix 1—Table 10. List of the 31 morphological traits measured for this study, and standard length; corresponding landmark ID's match those shown in Figure 2—figure supplement 3.

Trait Index	Trait Description	Trait Shorthand	Points
1	Nasal protrusion	nose	3-4
2	Nasal length	foresnout	2-5
3	Orbit to anal fin insertion	bellylen	6-15
4	Lateral facial length	snoutlen	2-6
5	Upper jaw to pectoral girdle	jaw2pect	2-14
6	Lateral skull length	pmx2add	2-11
7	Premaxilla length	pmxlen	2-9
8	Lower mandible length	jawlen	1-9
9	Jaw joint to orbit	foreeyewidth	6-9
10	Horizontal orbit diameter	eyewidth	6-8
11	Vertical orbit diameter	eyeht	7-10
12	Head height	headht	7-9
13	Suspensorium length	suspensorium	9-11
14	Adductor height	adductorht	11-12
15	Subopercle to pectoral girdle	ad2pect	11-14
16	Pectoral fin insertion width	pectinsertion	13-14
17	Anal to caudal distance	analtocaudal	15-16
18	Caudal peduncle height	caudalpedht	16-18
19	Dorsal to caudal distance	dorsaltocaudal	18-19
20	Body depth	bodydepth	15-19
21	Nasal protrusion angle	nasalangle	7-5-3
22	Premaxilla to orbit angle	topeyeangle	7-2-10
23	Premaxilla to adductor angle	lowereyeangle	7-2-11
24	Dorsal facial length	dorsalsnoutlen	23-24, 25-26
25	Adductor to premaxilla	eyetosnout	21-24, 25-28
26	Neurocranium to premaxilla	headlen	24-20, 25-29
27	Orbit to premaxilla	innereyetosnout	22-24, 25-27
28	Interorbital width	cranialwidth	22-27
29	Orbital neurocranium width	hindeyewidth	21-28
30	Max. neurocranium width	headwidth	20-29
31	Standard length (SL)	SL	2-17

Appendix 1—Table 11. Generalized additive models fitted to composite fitness. Model fit was assessed using AICc, and Akaike Weights represent proportional model support. A thin plate spline for the two linear discriminant axes s(LD1, LD2) is always included, as is a fixed effect of either experiment (i.e. Martin & Wainwright 2013, Martin and Gould 2020) or Lake (Crescent Pond/Little Lake) or an interaction between the two. In the last two models, Experiment and Lake are included as splines, modeled using a factor smooth (bs = "fs"). The best fit model had 5 estimated degrees of freedom.

Model	AICc	ΔAICc	Akaike Weights
Composite ~ s(LD1, LD2) + Experiment + Lake	*99.114*	*0.000*	*0.825*
Composite ~ s(LD1, LD2) + Experiment * Lake	102.210	3.096	0.175
Composite ~ s(LD1, LD2) + s(LD1) + s(LD2) + Experiment + Lake	131.456	32.342	< 0.001
Composite ~ s(LD1, LD2) + s(LD1) + s(LD2) + Experiment * Lake	135.894	36.781	< 0.001
Composite ~ s(LD1, LD2) + s(LD1, Experiment, bs = "fs") + s(LD2, Experiment, bs = "fs") + Lake	230.428	131.314	< 0.001
Composite ~ s(LD1, LD2) + s(LD1, Lake, bs = "fs") + s(LD2, Lake, bs = "fs") + Experiment	230.868	131.754	< 0.001

Appendix 1—Table 12.
Generalized additive models fitted to growth. Model fit was assessed using AICc, and Akaike Weights represent proportional model support. A thin plate spline for the two linear discriminant axes s(LD1, LD2) is always included, as is a fixed effect of either experiment (i.e. Martin & Wainwright 2013, Martin and Gould 2020) or Lake (Crescent Pond/Little Lake) or an interaction between the two. In the last two models, Experiment and Lake are included as splines, modeled using a factor smooth (bs = "fs"). The best fit model had 8.93 estimated degrees of freedom.

Model	AICc	ΔAICc	Akaike Weights
*Growth ~ s(LD1, LD2) + Experiment * Lake*	*-44.658*	*0.000*	*1*
Growth ~ s(LD1, LD2) + Experiment + Lake	3.904	48.562	< 0.001
Growth ~ s(LD1, LD2) + s(LD1) + s(LD2) + Experiment * Lake	46.249	90.907	< 0.001
Growth ~ s(LD1, LD2) + s(LD1) + s(LD2) + Experiment + Lake	89.121	133.779	< 0.001
Growth ~ s(LD1, LD2) + s(LD1, Lake, bs = "fs") + s(LD2, Lake, bs = "fs") + Experiment	690.379	735.038	< 0.001
Growth ~ s(LD1, LD2) + s(LD1, Experiment, bs = "fs") + s(LD2, Experiment, bs = "fs") + Lake	693.748	738.406	< 0.001

Appendix 1—Table 13. Generalized additive models fitted to survival. Model fit was assessed using AICc, and Akaike Weights represent proportional model support. A thin plate spline for the two linear discriminant axes s(LD1, LD2) is always included, as is a fixed effect of either experiment (i.e. Martin & Wainwright 2013, Martin and Gould 2020) or Lake (Crescent Pond/Little Lake) or an interaction between the two. In the last two models, Experiment and Lake are included as splines, modeled using a factor smooth (bs = "fs"). The best fit model had 5 estimated degrees of freedom.

Model	AICc	ΔAICc	Akaike Weights
Survival ~ s(LD1, LD2) + Experiment + Lake	*141.057*	*0.000*	*0.849*
Survival ~ s(LD1, LD2) + Experiment * Lake	144.504	3.447	0.151
Survival ~ s(LD1, LD2) + s(LD1) + s(LD2) + Experiment + Lake	173.399	32.342	< 0.001
Survival ~ s(LD1, LD2) + s(LD1) + s(LD2) + Experiment * Lake	178.189	37.132	< 0.001
Survival ~ s(LD1, LD2) + s(LD1, Experiment, bs = "fs") + s(LD2, Experiment, bs = "fs") + Lake	273.547	132.490	< 0.001
Survival ~ s(LD1, LD2) + s(LD1, Lake, bs = "fs") + s(LD2, Lake, bs = "fs") + Experiment	273.694	132.637	< 0.001

Appendix 1—Table 14. Generalized additive models fitted to growth including SNPs most strongly associated with composite fitness. Model fit was assessed using AICc, and Akaike Weights represent proportional model support. The best fit model for composite fitness using morphology alone (see Table 8) was used as the base model. The SNPs that were most strongly associated with composite fitness (following a Bonferroni correction) were included as fixed effects, modeled as splines using a factor smooth, treating genotype as an ordered factor. Note that three SNPs were excluded due to their close proximity to other SNPs that were more strongly associated. All SNPs were considered individually, as well as all SNPs together. We were unable to assess all possible combinations of SNPs due to the vast number of potential models given the number of SNPs under consideration; rather, we fit one final model that only included SNPs found to be significant in the full model. In turn this model led to a substantial improvement in AICc. The best fit model had 20.29 estimated degrees of freedom.

Model	AICc	ΔAICc	Akaike Weights
Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site1) + s(Site2) + s(Site6) + s(Site7) + s(Site8) + s(Site9) + s(Site10)	*4.586*	*0.000*	*0.999*
Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site1) + s(Site2) + s(Site3) + s(Site4) + s(Site5) + s(Site6) + s(Site7) + s(Site8) + s(Site9) + s(Site10)	40.876	36.290	< 0.001
Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site3)	55.588	51.001	< 0.001
Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site7)	58.386	53.800	< 0.001
Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site4)	65.453	60.867	< 0.001
Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site2)	71.245	66.658	< 0.001
Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site5)	72.329	67.743	< 0.001
Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site1)	73.671	69.085	< 0.001
Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site8)	74.413	69.827	< 0.001
Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site9)	74.680	70.094	< 0.001
Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site10)	88.977	84.391	< 0.001
Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site6)	90.427	85.841	< 0.001
Composite Fitness ~ s(LD1, LD2) + Experiment + Lake	99.114	94.527	< 0.001

Note: Site1 = HiC_Scaffold_1:43866598, Site2 = HiC_Scaffold_53:11317840, Site3 = HiC_Scaffold_46:35151009, Site4 = HiC_Scaffold_8:20263964, Site5 = 37:18591438, Site5 = HiC_Scaffold_37:18591438, Site6 = HiC_Scaffold_24:15964553, Site7 = HiC_Scaffold_1848:40590, Site8 = HiC_Scaffold_4461:12939, Site9 = HiC_Scaffold_12778:1456, Site10 = HiC_Scaffold_18999:1084.

Appendix 1—Table 15. Generalized additive models fitted to growth including SNPs most strongly associated with growth. Model fit was assessed using AICc, and Akaike Weights represent proportional model support. The best fit model for growth using morphology alone (see Table 9) was used as the base model. Each of the four SNPs that were most strongly associated with growth (following a Bonferroni correction) were included as fixed effects, modeled as splines using a factor smooth, treating genotype as an ordered factor. All SNPs were considered individually, as well as all possible combinations. This was only feasible due to the small number of SNPs assessed (four). The best fit model had 7.97 estimated degrees of freedom.

Model	AICc	ΔAICc	Akaike Weights
Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site3) + s(Site4)	-67.649	0.000	0.490
Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site3)	-65.634	2.015	0.179
Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site1)	-64.161	3.488	0.086
Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site1) + s(Site3)	-63.926	3.723	0.076
Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site1) + s(Site2)	-63.861	3.788	0.074
Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site2) + s(Site4)	-63.503	4.146	0.062
Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site2)	-61.849	5.800	0.027
Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site1) + s(Site4)	-58.044	9.604	0.004
Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site4)	-56.068	11.581	0.001
Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site1) + s(Site3) + s(Site4)	-54.878	12.770	< 0.001
Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site1) + s(Site2) + s(Site4)	-54.509	13.140	< 0.001
Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site2) + s(Site3)	-47.602	20.047	< 0.001
Growth ~ s(LD1, LD2) + Experiment * Lake	-44.658	22.990	< 0.001
Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site1) + s(Site2) + s(Site3)	-41.689	25.960	< 0.001
Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site1) + s(Site2) + s(Site3) + s(Site4)	-29.801	37.847	< 0.001

Note: Site1 = HiC_Scaffold_46:16512886, Site2 = HiC_Scaffold_1848:40119, Site3 = HiC_Scaffold_1848:40465, Site4 = HiC_Scaffold_7644:5971

Appendix 1—Table 16. General linear models fitted to examine the relationship between aspects of network size (i.e. number of nodes, number of edges linking neighboring nodes) and the number of accessible paths between generalists and specialists. Models were fitted using each of the three different fitness measures; bolded lines correspond to the best-fit model for each response variable, within each measure of fitness. Poisson regression was chosen as each response variable correspond to count-data. Because Poisson regression models are log-linear, we report both the estimated coefficient, as well as it's exponentiated value which corresponds to the expected multiplicative increase in the mean of Y per unit-value of X.

Fitness Measure	Model	Family	AIC	Coefficient	exp(Coefficient)	P-value
	# Edges in Network ~ # Nodes in Network	Poisson	34260.63	0.0342	1.0348	< 0.0001
	# Edges in Network ~ # Nodes in Network + Trajectory	Poisson	34260.93	0.0344	1.0350	< 0.0001
	# Edges in Network ~ # Nodes in Network * Trajectory	**Poisson**	**33599.17**	**0.0386**	**1.0394**	**< 0.0001**
	# Accessible Paths ~ # Nodes In Network	Poisson	12207.29	0.0082	1.0083	< 0.0001
Composite	**# Accessible Paths ~ # Nodes In Network + Trajectory**	**Poisson**	**12203.83**	**0.0061**	**1.0062**	**< 0.0001**
	# Accessible Paths ~ # Nodes In Network * Trajectory	Poisson	12204.89	0.0071	1.0071	< 0.0001
	# Accessible Paths ~ # Edges In Network	Poisson	12203.07	0.0044	1.0044	< 0.0001
	# Accessible Paths ~ # Edges In Network + Trajectory	**Poisson**	**12200.31**	**0.0034**	**1.0034**	**< 0.0001**
	# Accessible Paths ~ # Edges In Network * Trajectory	Poisson	12201.69	0.0039	1.0039	< 0.0001
	# Edges in Network ~ # Nodes in Network	Poisson	26739.96	0.0478	1.0489	< 0.0001
	# Edges in Network ~ # Nodes in Network + Trajectory	Poisson	26711.02	0.0489	1.0501	< 0.0001
	# Edges in Network ~ # Nodes in Network * Trajectory	**Poisson**	**26507.72**	**0.0532**	**1.0546**	**< 0.0001**
	# Accessible Paths ~ # Nodes In Network	Poisson	10162.51	0.0106	1.0107	< 0.0001
Growth	**# Accessible Paths ~ # Nodes In Network + Trajectory**	**Poisson**	**10160.39**	**0.0083**	**1.0083**	**0.0002**
	# Accessible Paths ~ # Nodes In Network * Trajectory	Poisson	10162.28	0.0088	1.0089	0.0011
	# Accessible Paths ~ # Edges In Network	Poisson	10159.39	0.0062	1.0062	< 0.0001
	# Accessible Paths ~ # Edges In Network + Trajectory	**Poisson**	**10157.09**	**0.0050**	**1.0050**	**< 0.0001**
	# Accessible Paths ~ # Edges In Network * Trajectory	Poisson	10159.09	0.0050	1.0050	0.0010
	# Edges in Network ~ # Nodes in Network	Poisson	32986.67	0.0346	1.0352	< 0.0001
	# Edges in Network ~ # Nodes in Network + Trajectory	Poisson	32978.31	0.0350	1.0356	< 0.0001
	# Edges in Network ~ # Nodes in Network * Trajectory	**Poisson**	**32437.55**	**0.0384**	**1.0392**	**< 0.0001**
	# Accessible Paths ~ # Nodes In Network	**Poisson**	**11725.19**	**0.0058**	**1.0058**	**< 0.0001**
Survival	# Accessible Paths ~ # Nodes In Network + Trajectory	Poisson	11727.13	0.0056	1.0056	0.0001
	# Accessible Paths ~ # Nodes In Network * Trajectory	Poisson	11727.11	0.0068	1.0068	< 0.0001
	# Accessible Paths ~ # Edges In Network	**Poisson**	**11722.24**	**0.0032**	**1.0032**	**< 0.0001**
	# Accessible Paths ~ # Edges In Network + Trajectory	Poisson	11724.23	0.0031	1.0031	< 0.0001
	# Accessible Paths ~ # Edges In Network * Trajectory	Poisson	11725.27	0.0036	1.0036	< 0.0001

Appendix 1—Table 17. Accessibility of specialists to generalists and the ruggedness of their respective fitness landscapes. Odds ratios were obtained by modelling the association between each summary statistic and the species from which adaptive loci were used to construct the fitness network. Scale-eaters were treated as the baseline of comparison in the comparison of odds ratios; thus, positive odds ratios imply that summary statistics for molluscivore fitness networks are greater than those constructed from scale-eater adaptive loci and vice versa. For generalist to specialist comparisons, accessible paths were identified between one randomly sampled generalist node and one randomly sampled specialist node. For comparison of the peaks in networks, these summary statistics were calculated from either molluscivore or scale-eater fitness networks, identifying the number of peaks (nodes with no fitter neighbors – see Methods), and the scaled (total divided by number of nodes in the network) number of accessible paths separating all focal specialist nodes and all peaks in the network.

Comparison	Summary Statistic	Mean / SE *Molluscivore Network*	Mean / SE *Scale-Eater Network*	Odds Ratio: (95% CI) *Molluscivore / Scale Eater*	LRT *P*-value
Generalist to Specialist	*Number of nodes in network*	22.994 / 0.106	31.000 / 0.177	0.818: (0.807, 0.829)	< 0.0001
	Number of accessible paths	1.105 / 0.007	1.268 / 0.018	0.515: (0.449, 0.588)	< 0.0001
	Scaled number of accessible paths	0.051 / 0.001	0.042 / 0.001	2.095: (1.934, 2.274)	< 0.0001
	Length of shortest accessible path	2.410 / 0.012	3.444 / 0.029	0.253: (0.231, 0.277)	< 0.0001
Peaks in Network	*Number of peaks*	3.274 / 0.035	4.637 / 0.046	0.604: (0.575, 0.634)	< 0.0001
	Scaled number of accessible paths to peaks	0.095 / 0.001	0.087 / 0.001	1.514: (1.404, 1.635)	< 0.0001
	Length of shortest accessible path to nearest peak	0.823 / 0.022	1.482 / 0.029	0.539: (0.500, 0.579)	< 0.0001

Appendix 1—Table 18. Influence of different sources of adaptive genetic variation on accessibility of fitness paths separating either generalists from molluscivores, or generalists and scale-eaters using all samples. Results for networks using all three measures of fitness (composite fitness, survival, and growth) are reported. Networks were constructed from random draws of five SNPs from either Standing genetic variation (SGV), introgression, or *de novo* mutations, as well as their combinations. Odds ratios were obtained by modelling the association between each accessibility measure and the source of genetic variation used to construct the fitness network, relative to networks constructed from standing variation. Thus, positive odds ratios imply that networks from standing variation have measures of accessibility that are smaller as compared to the alternative (e.g. introgression, *de novo* mutations, etc).

Trajectory	Source	Accessibility	Composite Fitness			Growth			Survival		
			Mean / SE	LRT P-value	Odds Ratio: (95% CI)	Mean / SE	LRT P-value	Odds Ratio: (95% CI)	Mean / SE	LRT P-value	Odds Ratio: (95% CI)
Generalist to Molluscivore	*Introgression*	# Nodes in network	22.095 / 0.118	< 0.0001	0.627: (0.606, 0.646)	17.630 / 0.098	< 0.0001	0.556: (0.535, 0.577)	21.845 / 0.108	< 0.0001	0.628: (0.608, 0.646)
		# Accessible paths	1.071 / 0.005	< 0.0001	0.442: (0.378, 0.515)	1.099 / 0.007	< 0.0001	0.721: (0.613, 0.842)	1.039 / 0.004	< 0.0001	0.214: (0.174, 0.259)
		# Accessible paths / # nodes in network	0.052 / 0.0004	< 0.0001	17.131: (15.076, 19.556)	0.066 / 0.0006	< 0.0001	10.272: (9.105, 11.636)	0.051 / 0.0003	< 0.0001	11.256: (10.121, 12.560)
		Length of shortest accessible path	2.480 / 0.013	< 0.0001	0.768: (0.717, 0.823)	2.457 / 0.016	0.8826	0.994: (0.915, 1.079)	2.475 / 0.012	< 0.0001	0.662: (0.619, 0.707)
		Ruggedness (# of peaks)	1.444 / 0.025	< 0.0001	0.250: (0.234, 0.268)	0.706 / 0.018	< 0.0001	0.252: (0.233, 0.271)	1.863 / 0.033	< 0.0001	0.397: (0.378, 0.416)
	SGV + Introgression	# Nodes in network	32.167 / 0.146	< 0.0001	0.820: (0.813, 0.827)	22.994 / 0.106	< 0.0001	0.777: (0.768, 0.786)	32.386 / 0.144	< 0.0001	0.826: (0.819, 0.833)
		# Accessible paths	1.117 / 0.007	< 0.0001	0.694: (0.624, 0.771)	1.105 / 0.006	< 0.0001	0.773: (0.684, 0.873)	1.140 / 0.007	< 0.0001	0.666: (0.608, 0.728)
		# Accessible paths / # nodes in network	0.037 / 0.0002	< 0.0001	3.974: (3.707, 4.266)	0.051 / 0.0003	< 0.0001	3.538: (3.297, 3.802)	0.038 / 0.0003	< 0.0001	3.108: (2.922, 3.309)
		Length of shortest accessible path	2.524 / 0.012	< 0.0001	0.843: (0.793, 0.894)	2.410 / 0.011	0.0050	0.897: (0.835, 0.962)	2.573 / 0.012	< 0.0001	0.798: (0.755, 0.843)
		Ruggedness (# of peaks)	3.906 / 0.034	< 0.0001	0.523: (0.509, 0.537)	2.218 / 0.028	< 0.0001	0.556: (0.539, 0.574)	5.651 / 0.056	< 0.0001	0.690: (0.680, 0.700)
Generalist to Scale-eater	*Introgression*	# Nodes in network	45.783 / 0.261	< 0.0001	0.866: (0.858, 0.874)	33.569 / 0.232	< 0.0001	0.838: (0.827, 0.849)	45.423 / 0.270	< 0.0001	0.870: (0.862, 0.879)
		# Accessible paths	1.190 / 0.012	< 0.0001	0.632: (0.564, 0.706)	1.178 / 0.013	< 0.0001	0.637: (0.556, 0.728)	1.191 / 0.012	< 0.0001	0.668: (0.593, 0.751)
		# Accessible paths / # nodes in network	0.027 / 0.0003	< 0.0001	1.790: (1.658, 1.936)	0.037 / 0.0004	< 0.0001	2.011: (1.837, 2.206)	0.028 / 0.0003	< 0.0001	1.931: (1.778, 2.101)
		Length of shortest accessible path	3.033 / 0.021	< 0.0001	0.552: (0.513, 0.593)	3.040 / 0.023	< 0.0001	0.570: (0.525, 0.618)	3.103 / 0.021	< 0.0001	0.679: (0.629, 0.733)
		Ruggedness (# of peaks)	7.879 / 0.036	< 0.0001	0.752: (0.739, 0.766)	5.516 / 0.033	< 0.0001	0.724: (0.709, 0.740)	15.020 / 0.088	< 0.0001	0.838: (0.831, 0.845)
	De novo	# Nodes in network	46.292 / 0.177	< 0.0001	0.835: (0.826, 0.844)	35.650 / 0.129	< 0.0001	0.796: (0.783, 0.808)	46.251 / 0.199	< 0.0001	0.844: (0.835, 0.854)
		# Accessible paths	1.234 / 0.012	< 0.0001	0.736: (0.673, 0.803)	1.196 / 0.011	< 0.0001	0.686: (0.612, 0.766)	1.172 / 0.011	< 0.0001	0.628: (0.559, 0.703)
		# Accessible paths / # nodes in network	0.027 / 0.0002	< 0.0001	1.908: (1.770, 2.061)	0.034 / 0.0003	< 0.0001	1.764: (1.622, 1.922)	0.026 / 0.0002	< 0.0001	1.818: (1.678, 1.972)
		Length of shortest accessible path	3.088 / 0.018	< 0.0001	0.589: (0.553, 0.628)	3.029 / 0.018	< 0.0001	0.555: (0.515, 0.597)	2.920 / 0.017	< 0.0001	0.518: (0.477, 0.560)
		Ruggedness (# of peaks)	8.381 / 0.034	< 0.0001	0.806: (0.792, 0.819)	6.145 / 0.031	< 0.0001	0.809: (0.792, 0.825)	15.615 / 0.079	< 0.0001	0.836: (0.829, 0.843)
	SGV + Introgression	# Nodes in network	56.559 / 0.272	< 0.0001	0.952: (0.945, 0.959)	40.858 / 0.238	< 0.0001	0.925: (0.915, 0.935)	55.717 / 0.302	< 0.0001	0.950: (0.943, 0.957)
		# Accessible paths	1.331 / 0.019	0.0199	0.904: (0.831, 0.981)	1.304 / 0.021	0.1646	0.928: (0.835, 1.031)	1.312 / 0.022	0.1469	0.932: (0.850, 1.018)
		# Accessible paths / # nodes in network	0.024 / 0.0003	< 0.0001	1.167: (1.089, 1.252)	0.033 / 0.0005	< 0.0001	1.270: (1.172, 1.379)	0.024 / 0.0004	< 0.0001	1.209: (1.121, 1.304)
		Length of shortest accessible path	3.415 / 0.026	< 0.0001	0.828: (0.777, 0.882)	3.419 / 0.031	< 0.0001	0.851: (0.793, 0.913)	3.263 / 0.026	< 0.0001	0.823: (0.763, 0.887)
		Ruggedness (# of peaks)	9.319 / 0.036	< 0.0001	0.943: (0.928, 0.957)	6.731 / 0.030	< 0.0001	0.912: (0.895, 0.930)	19.924 / 0.089	< 0.0001	0.954: (0.948, 0.960)
	SGV + De novo	# Nodes in network	56.723 / 0.214	< 0.0001	0.947: (0.941, 0.954)	42.261 / 0.177	< 0.0001	0.936: (0.926, 0.946)	56.488 / 0.230	< 0.0001	0.951: (0.943, 0.958)
		# Accessible paths	1.344 / 0.019	0.0522	0.931: (0.865, 1.001)	1.287 / 0.018	0.0473	0.901: (0.816, 0.994)	1.315 / 0.023	0.1632	0.949: (0.879, 1.021)
		# Accessible paths / # nodes in network	0.024 / 0.0003	< 0.0001	1.155: (1.081, 1.235)	0.031 / 0.0004	0.0017	1.128: (1.046, 1.218)	0.024 / 0.0004	0.0004	1.134: (1.057, 1.218)
		Length of shortest accessible path	3.431 / 0.024	< 0.0001	0.842: (0.794, 0.892)	3.391 / 0.027	< 0.0001	0.834: (0.780, 0.891)	3.299 / 0.023	< 0.0001	0.859: (0.802, 0.920)
		Ruggedness (# of peaks)	9.525 / 0.035	0.0006	0.973: (0.958, 0.988)	6.975 / 0.029	0.0006	0.967: (0.949, 0.986)	19.849 / 0.085	< 0.0001	0.950: (0.944, 0.956)
	SGV + De novo + Introgression	# Nodes in network	46.882 / 0.200	< 0.0001	0.826: (0.816, 0.837)	31.000 / 0.177	< 0.0001	0.722: (0.703, 0.739)	46.875 / 0.222	< 0.0001	0.835: (0.823, 0.845)
		# Accessible paths	1.322 / 0.019	0.0125	0.896: (0.824, 0.971)	1.268 / 0.018	0.0102	0.855: (0.762, 0.956)	1.247 / 0.018	0.0001	0.811: (0.726, 0.900)
		# Accessible paths / # nodes in network	0.029 / 0.0004	< 0.0001	1.879: (1.734, 2.041)	0.042 / 0.0006	< 0.0001	2.798: (2.511, 3.128)	0.027 / 0.0004	< 0.0001	1.785: (1.640, 1.948)
		Length of shortest accessible path	3.484 / 0.026	< 0.0001	0.876: (0.823, 0.932)	3.444 / 0.029	< 0.0001	0.861: (0.800, 0.925)	3.221 / 0.024	< 0.0001	0.775: (0.716, 0.838)
		Ruggedness (# of peaks)	7.440 / 0.029	< 0.0001	0.650: (0.637, 0.664)	4.555 / 0.025	< 0.0001	0.485: (0.471, 0.500)	12.933 / 0.062	< 0.0001	0.725: (0.716, 0.734)

Appendix 1—Table 19. Influence of different sources of adaptive genetic variation on accessibility of fitness paths separating either generalists from molluscivores, or generalists and scale-eaters using only samples from the second field experiment (Martin & Gould 2021). Results for networks using all three measures of fitness (composite fitness, survival, and growth) are reported. Networks were constructed from random draws of five SNPs from either Standing genetic variation (SGV), introgression, or *de novo* mutations, as well as their combinations. Odds ratios were obtained by modelling the association between each accessibility measure and the source of genetic variation used to construct the fitness network, relative to networks constructed from standing variation. Thus, positive odds ratios imply that networks from standing variation have measures of accessibility that are smaller as compared to the alternative (e.g. introgression, *de novo* mutations, etc).

Trajectory	Source	Accessibility	Composite Fitness			Growth			Survival		
			Mean / SE	LRT P-value	Odds Ratio: (95% CI)	Mean / SE	LRT P-value	Odds Ratio: (95% CI)	Mean / SE	LRT P-value	Odds Ratio: (95% CI)
Generalist to Molluscivore	*Introgression*	# Nodes in network	22.062 / 0.118	< 0.0001	0.628: (0.608, 0.648)	17.632 / 0.099	< 0.0001	0.556: (0.535, 0.577)	21.836 / 0.107	< 0.0001	0.617: (0.596, 0.637)
		# Accessible paths	1.067 / 0.005	< 0.0001	0.402: (0.341, 0.471)	1.107 / 0.008	< 0.0001	0.721: (0.618, 0.836)	1.039 / 0.004	< 0.0001	0.205: (0.167, 0.248)
		# Accessible paths / # nodes in network	0.052 / 0.0004	< 0.0001	16.263: (14.350, 18.515)	0.066 / 0.0006	< 0.0001	9.411: (8.375, 10.617)	0.051 / 0.0003	< 0.0001	10.698: (9.642, 11.907)
		Length of shortest accessible path	2.483 / 0.013	< 0.0001	0.751: (0.700, 0.804)	2.465 / 0.016	0.7601	0.987: (0.908, 1.073)	2.473 / 0.013	< 0.0001	0.652: (0.610, 0.695)
		Ruggedness (# of peaks)	1.437 / 0.025	< 0.0001	0.247: (0.230, 0.264)	0.702 / 0.018	< 0.0001	0.257: (0.238, 0.276)	1.853 / 0.033	< 0.0001	0.405: (0.386, 0.425)
	SGV + Introgression	# Nodes in network	32.301 / 0.146	< 0.0001	0.824: (0.818, 0.831)	23.042 / 0.103	< 0.0001	0.769: (0.760, 0.779)	32.488 / 0.144	< 0.0001	0.826: (0.819, 0.832)
		# Accessible paths	1.134 / 0.007	< 0.0001	0.753: (0.681, 0.832)	1.115 / 0.007	< 0.0001	0.778: (0.693, 0.871)	1.146 / 0.008	< 0.0001	0.664: (0.609, 0.723)
		# Accessible paths / # nodes in network	0.037 / 0.0002	< 0.0001	3.929: (3.671, 4.210)	0.051 / 0.0003	< 0.0001	3.505: (3.267, 3.765)	0.037 / 0.0002	< 0.0001	3.053: (2.874, 3.245)
		Length of shortest accessible path	2.538 / 0.012	< 0.0001	0.842: (0.793, 0.893)	2.426 / 0.011	0.0136	0.907: (0.846, 0.974)	2.597 / 0.013	< 0.0001	0.819: (0.777, 0.863)
		Ruggedness (# of peaks)	3.954 / 0.034	< 0.0001	0.518: (0.504, 0.532)	2.243 / 0.028	< 0.0001	0.560: (0.542, 0.577)	5.742 / 0.056	< 0.0001	0.690: (0.680, 0.700)
Generalist to Scale-eater	*Introgression*	# Nodes in network	45.775 / 0.254	< 0.0001	0.864: (0.855, 0.872)	33.658 / 0.219	< 0.0001	0.837: (0.826, 0.849)	45.306 / 0.264	< 0.0001	0.862: (0.853, 0.870)
		# Accessible paths	1.202 / 0.013	< 0.0001	0.723: (0.647, 0.803)	1.180 / 0.013	< 0.0001	0.733: (0.639, 0.835)	1.208 / 0.013	< 0.0001	0.743: (0.661, 0.830)
		# Accessible paths / # nodes in network	0.027 / 0.0003	< 0.0001	1.959: (1.809, 2.125)	0.037 / 0.0004	< 0.0001	2.243: (2.038, 2.476)	0.028 / 0.0003	< 0.0001	2.087: (1.916, 2.279)
		Length of shortest accessible path	3.108 / 0.022	< 0.0001	0.605: (0.564, 0.648)	3.063 / 0.022	< 0.0001	0.568: (0.523, 0.615)	3.132 / 0.022	< 0.0001	0.713: (0.662, 0.768)
		Ruggedness (# of peaks)	7.928 / 0.036	< 0.0001	0.760: (0.747, 0.773)	5.572 / 0.033	< 0.0001	0.727: (0.712, 0.743)	15.072 / 0.087	< 0.0001	0.838: (0.831, 0.845)
	De novo	# Nodes in network	46.510 / 0.177	< 0.0001	0.843: (0.834, 0.851)	35.882 / 0.129	< 0.0001	0.808: (0.796, 0.820)	46.454 / 0.196	< 0.0001	0.843: (0.833, 0.852)
		# Accessible paths	1.223 / 0.011	< 0.0001	0.775: (0.708, 0.846)	1.202 / 0.011	0.0002	0.806: (0.723, 0.896)	1.168 / 0.010	< 0.0001	0.639: (0.566, 0.717)
		# Accessible paths / # nodes in network	0.027 / 0.0002	< 0.0001	2.010: (1.861, 2.176)	0.034 / 0.0003	< 0.0001	1.934: (1.768, 2.121)	0.026 / 0.0002	< 0.0001	1.874: (1.730, 2.034)
		Length of shortest accessible path	3.100 / 0.018	< 0.0001	0.602: (0.566, 0.640)	3.049 / 0.018	< 0.0001	0.564: (0.524, 0.605)	2.935 / 0.017	< 0.0001	0.553: (0.512, 0.596)
		Ruggedness (# of peaks)	8.336 / 0.034	< 0.0001	0.798: (0.784, 0.812)	6.147 / 0.030	< 0.0001	0.803: (0.787, 0.820)	15.630 / 0.079	< 0.0001	0.837: (0.830, 0.844)
	SGV + Introgression	# Nodes in network	56.208 / 0.268	< 0.0001	0.952: (0.946, 0.959)	41.204 / 0.243	< 0.0001	0.937: (0.928, 0.947)	55.697 / 0.277	< 0.0001	0.948: (0.940, 0.955)
		# Accessible paths	1.339 / 0.021	0.6195	0.980: (0.907, 1.060)	1.285 / 0.020	0.922	0.985: (0.885, 1.097)	1.307 / 0.020	0.256	0.946: (0.866, 1.031)
		# Accessible paths / # nodes in network	0.024 / 0.0004	< 0.0001	1.234: (1.151, 1.325)	0.032 / 0.0005	< 0.0001	1.299: (1.196, 1.413)	0.024 / 0.0004	< 0.0001	1.231: (1.144, 1.327)
		Length of shortest accessible path	3.465 / 0.026	< 0.0001	0.860: (0.809, 0.914)	3.460 / 0.030	< 0.0001	0.860: (0.802, 0.922)	3.306 / 0.025	0.0001	0.867: (0.807, 0.932)
		Ruggedness (# of peaks)	9.231 / 0.036	< 0.0001	0.930: (0.915, 0.944)	6.666 / 0.030	< 0.0001	0.902: (0.885, 0.920)	19.815 / 0.090	< 0.0001	0.951: (0.945, 0.957)
	SGV + De novo	# Nodes in network	56.259 / 0.212	< 0.0001	0.947: (0.940, 0.953)	41.991 / 0.173	< 0.0001	0.936: (0.925, 0.946)	56.339 / 0.224	< 0.0001	0.949: (0.942, 0.956)
		# Accessible paths	1.329 / 0.017	0.4575	0.966: (0.895, 1.042)	1.290 / 0.020	0.922	0.995: (0.905, 1.097)	1.329 / 0.018	0.5596	0.976: (0.900, 1.059)
		# Accessible paths / # nodes in network	0.024 / 0.0003	< 0.0001	1.233: (1.153, 1.319)	0.031 / 0.0004	< 0.0001	1.225: (1.133, 1.326)	0.024 / 0.0003	< 0.0001	1.243: (1.158, 1.335)
		Length of shortest accessible path	3.452 / 0.023	< 0.0001	0.851: (0.803, 0.902)	3.379 / 0.025	< 0.0001	0.800: (0.748, 0.856)	3.288 / 0.022	< 0.0001	0.852: (0.797, 0.912)
		Ruggedness (# of peaks)	9.406 / 0.035	< 0.0001	0.953: (0.938, 0.968)	6.898 / 0.029	< 0.0001	0.948: (0.930, 0.966)	19.782 / 0.084	< 0.0001	0.947: (0.941, 0.953)
	SGV + De novo + Introgression	# Nodes in network	47.220 / 0.207	< 0.0001	0.841: (0.831, 0.851)	31.051 / 0.183	< 0.0001	0.740: (0.723, 0.757)	47.117 / 0.224	< 0.0001	0.840: (0.829, 0.850)
		# Accessible paths	1.298 / 0.018	0.074	0.918: (0.842, 0.998)	1.289 / 0.022	0.922	0.994: (0.894, 1.104)	1.221 / 0.016	< 0.0001	0.790: (0.705, 0.879)
		# Accessible paths / # nodes in network	0.028 / 0.0004	< 0.0001	1.945: (1.792, 2.116)	0.043 / 0.0007	< 0.0001	3.307: (2.936, 3.741)	0.027 / 0.0004	< 0.0001	1.784: (1.639, 1.946)
		Length of shortest accessible path	3.497 / 0.025	< 0.0001	0.879: (0.827, 0.935)	3.479 / 0.030	0.0001	0.869: (0.809, 0.933)	3.269 / 0.024	< 0.0001	0.826: (0.766, 0.891)
		Ruggedness (# of peaks)	7.361 / 0.031	< 0.0001	0.650: (0.637, 0.664)	4.530 / 0.026	< 0.0001	0.490: (0.475, 0.504)	12.869 / 0.062	< 0.0001	0.725: (0.716, 0.734)