1   **The first genotypic fitness network in a vertebrate reveals that hybridization**

2   **increases access to novel fitness peaks**

4   Austin H. Patton[1,2], Emilie J. Richards[1,2], Katelyn J. Gould[3], Logan K. Buie[3], Christopher H.

5   Martin[1,2]

6
[1]*Department of Integrative Biology, University of California, Berkeley, CA*

[2]*Museum of Vertebrate Zoology, University of California, Berkeley, CA*

[3]*Department of Biology, University of North Carolina, Chapel Hill, NC*

**Keywords:** fitness landscape; genotypic fitness network adaptive radiation; introgression;

standing genetic variation; *de novo* mutation

**Correspondence:** chmartin@berkeley.edu

**Abstract**

Estimating the complex relationship between fitness and genotype or phenotype (i.e. the adaptive

landscape) is one of the central goals of evolutionary biology. Empirical fitness landscapes have

now been estimated for numerous systems, from phage to proteins to finches. However, adaptive

walks connecting genotypes to organismal fitness, speciation, and novel ecological niches are still

poorly understood and the process by which fitness valleys are crossed remains controversial. One

outstanding system for addressing these connections is a recent adaptive radiation of ecologically

and morphologically novel pupfishes (a generalist, molluscivore, and scale-eater) endemic to San

Salvador Island, Bahamas. Here, we leveraged whole-genome sequencing of 139 hybrids from

two independent field fitness experiments to identify the genomic basis of fitness, visualize the

first genotypic fitness networks in a vertebrate system, and measure the accessibility of adaptive

walks on the fitness landscape. We identified 132 SNPs that were significantly associated with

fitness in field enclosures. Six fitness-associated regions contained differentially expressed genes

and fixed SNPs between trophic specialists; one gene (*mettle21e*) was also misexpressed in

hybrids, suggesting a potential intrinsic genetic incompatibility. We then constructed genotypic

fitness networks from adaptive alleles and showed that scale-eating specialists are most isolated

on these networks. Furthermore, only introgressed and *de novo* variants, not standing genetic

variation, increased the accessibility of genotypic fitness paths from generalist to specialists. Our

results suggest that adaptive introgression and *de novo* mutations provided key connections in

adaptive walks necessary for crossing fitness valleys and triggering the evolution of novelty during

adaptive radiation.

**Significance**

*The fitness landscape describes the complex relationship between genotype or phenotype and*

*fitness and is the conceptual bridge between micro- and macroevolution. There are many studies*

*of phenotypic fitness landscapes across a range of animal and plant taxa, but studies of the*

*genotypic fitness landscape remain limited to microbial systems and the function of individual*

*proteins. Here, we conducted the first study investigating the genotypic fitness landscape for*

*organismal fitness in an exceptionally young vertebrate adaptive radiation including a generalist*

*algivore, molluscivore, and scale-eating pupfish. We show that hybridization and de novo*

*mutations increase the availability of accessible pathways to new fitness peaks on the fitness*

*landscape, enabling the evolution of novelty.*

**Introduction**

First conceptualized by Sewell Wright in 1932, the adaptive landscape describes the complex

relationship between genotype or phenotype and fitness (1). The landscape is a concept,

metaphor, and an empirical measurement that exerts substantial influence over all evolutionary

dynamics (2–6). Fitness landscapes were originally depicted as high-dimensional networks

spanning genotypic space in which each genotype is associated with fitness (1). Simpson (7)

later described phenotypic evolution of populations through time on a rugged landscape, in

which isolated clusters of fitness peaks represent 'adaptive zones' relative to adjacent regions of

low fitness (8). Lande and Arnold formalized the analysis of selection and estimation of

phenotypic fitness landscapes (9–11), leading to empirical studies of fitness landscapes in

numerous empirical systems (12–18). Fitness surfaces are also central components of speciation

models and theory (19–21). Speciation can result from divergent selection (22), either on static

landscapes or through negative frequency-dependent disruptive selection (23–28).

A fundamental concept in fitness landscape theory is that not all genotypic trajectories

are accessible (5, 29, 30). Rather, only paths that monotonically increase in fitness at each

mutational step are evolutionarily accessible, an expectation stemming from Fisher's

fundamental theorem (31). These accessible genotypic trajectories can thus be considered

potential adaptive walks under Fisher's geometric model (31, 32). There are a growing number

of experimental studies of adaptive walks in nature, including the evolution of toxin resistance in

monarch butterflies (33), alcohol tolerance in Drosophila (34, 35), and host-shift in aphids (36).

Likewise, the accessibility of genotypic fitness networks has now been explored in numerous

microbial systems, including the evolution of antibiotic resistance (29) and metabolism (37),

citrate exploitation (38), and glucose limitation in *E. coli* (39), as well as adaptation to salinity in

yeast via evolution of heat shock protein *Hsp90* (40). However, these studies are still limited to

the investigation of specific coding substitutions and their effects on fitness in laboratory

environments. Nosil et al. (41) estimated genotypic fitness networks for *Timema* stick insects

based on a field experiment. However, this study still focused on a single large-effect locus

underlying dimorphic coloration between ecotypes. To our knowledge, no study has estimated

genotypic fitness landscapes in a vertebrate system among multiple species as originally

envisioned by Wright (1) despite the clear utility of this framework in bridging micro- and

macroevolutionary dynamics.

One promising system for estimating fitness landscapes is a recent adaptive radiation of

*Cyprinodon* pupfishes (17, 18, 42, 43). This radiation is comprised of two endemic specialists, a

molluscivore (durophage: *Cyprinodon brontotheroides*) and a scale-eater (lepidophage: *C.*

*desquamator*), derived from a Caribbean-wide generalist ancestor (*C. variegatus*). These three

species all occur in sympatry in the hypersaline lakes of San Salvador Island, Bahamas (Fig 1A).

Originating less than 10,000 years ago (based on geological age estimates for the lakes: (44)), the

functional and trophic novelty harbored within this radiation is the product of exceptional rates

of craniofacial morphological evolution (45–49). Furthermore, species boundaries appear to

persist across multiple lake populations, despite persistent admixture among species (50, 51). We

previously estimated fitness landscapes in these hypersaline lakes from two independent field

experiments measuring the growth and survival of hybrids placed in field enclosures. Selection

analyses revealed a stable, multi-peaked fitness landscape across lake populations, year of study,

and manipulation of the frequency of rare hybrid phenotypes (Fig. 1B: (17, 18, 43)). One of the

strongest and most persistent trends across studies and treatments was the apparent isolation of

hybrid phenotypes resembling the scale-eater by a large fitness valley from the other two species

(17, 18). Hybrids resembling the generalist occupied a fitness peak and were isolated by a

smaller fitness valley from hybrids resembling the molluscivore in a second region of high

fitness. Understanding how an ancestral generalist population crossed these fitness valleys is key

to understanding the origins of novelty in this system, particularly given the rare evolution of

lepidophagy (scale-eating), a niche occupied by less than 0.3% of all fishes (52, 53).

111
**Figure 1. San Salvador Island pupfishes and their hybrids. a.** From top to bottom: the generalist,
*Cyprinodon variegatus*, the molluscivore *C. brontotheroides*, and the scale-eater *C. desquamator*. **b.**
Representative images of experimental field enclosures. **c.** Principal component analysis of 1,129,771 SNPs
genotyped in hybrids and the three parental species. **d.** ADMIXTURE analyses for Crescent Pond (top) and
Little Lake (bottom). G, M and S indicate individual samples of Generalists (G), Molluscivores (M), and
Scale-eaters (S), respectively, followed by all resequenced hybrid individuals from field experiments.
Colors correspond to probability of assignment to one of three assumed populations/species (K = 3 in this
analysis).
120

The relative contributions of standing genetic variation, *de novo* mutation, and

introgression to the tempo and mode of adaptive radiation are now of central interest to the field

of speciation genomics (54–59). Yet to date, no study has investigated the relative influence of

each on the accessibility of the fitness landscape, a critical determinant of the dynamics of

speciation. The adaptive radiation of San Salvador Island pupfishes, like many others (60–64),

appears to have resulted from a complex interplay of abundant standing genetic variation,

adaptive introgression from neighboring islands, and several *de novo* substitutions and deletions

found only in the scale-eater (51, 65). We have now identified many adaptive alleles showing

evidence of hard selective sweeps in each of the trophic specialists (51). Notably, both specialists

harbor numerous introgressed adaptive alleles in the regulatory regions of known craniofacial

genes (51, 65). In contrast, hard selective sweeps of *de novo* mutations only appear in the scale-

eating species, *C. desquamator*; many of these novel mutations also fall in or near craniofacial

genes, such as a single non-synonymous fixed substitution in the transcription factor and

oncogene *twist1* (51). Here, we leverage whole genome sequencing of 139 hybrids used in these

field experiments to identify the genomic basis of fitness differences, infer genotypic fitness

networks, and quantify the accessibility of novel fitness peaks and the influence of each source

of genetic variation on these adaptive walks.

**Results and Discussion**

*Population structure and ancestry association with fitness*

We resequenced 139 hybrids (86 survivors, 56 deaths (Table S1)) from two independent field

experiments across a total of six field enclosures and two lake populations (2011: two high-

density 3 m diameter enclosures for three months: Crescent Pond n = 796; Little Lake = 875 F2

hybrids (17); 2014/2015: four high-density 4 m diameter enclosures: three months in Crescent

Pond n = 923 F4/F5 hybrids; eleven months in Little Lake = 842 F4/F5 hybrids (66)). We then

characterized patterns of genetic variation among parental species in each lake and their lab-

reared hybrids used in field experiments. We genotyped 1,129,771 SNPs with an average

coverage of 9.79x per individual. Principal components analysis (PCA) of genetic variation

strongly differentiated pupfishes sampled from Little Lake/Osprey Lake and Crescent Pond

(PC1: 22.7% variance explained) and among species within each lake (PC2: 15.9% variance

explained: Fig. 1D; Fig. S1-S2). These results were supported by ADMIXTURE analyses (67,

68) (Fig. 1E). However, some hybrids were genotypically transgressive, falling outside the

genotypic distributions of the three parental species (Fig. S2), leading ADMIXTURE to attribute

the third population to these hybrids, rather than generalists which often contain segregating

variation found in the trophic specialists (67).

Neither scale-eater nor molluscivore ancestry estimated by ADMIXTURE analysis was

significantly associated with hybrid composite fitness (Fig. S3A, Table S2: Scale-eater $P$ =

0.164; molluscivore $P$ = 0.595), survival probability (Fig. S3B: Scale-eater $p$ = 0.804;

molluscivore $P$ = 0.313), or growth (Fig. S3C: Scale-eater $P$ = 0.283; molluscivore $P$ = 0.328),

while controlling for effects of lake and experiment. Similarly, neither PC1 nor PC2 was

associated with composite fitness (Table S2). Furthermore, pairwise genetic distance did not

predict pairwise morphological distance (Fig. S4). Therefore, in contrast to previous studies (70–

72), in this system genome-wide ancestry is not associated with fitness, highlighting the complex

nonlinear relationship between genotype, phenotype, and fitness within this nascent adaptive

radiation. We must look to local ancestry to understand fitness relationships (e.g. (73)).

**Figure 2. The genetic basis of fitness variation and improved inference of adaptive landscapes. a)** Per-SNP -log10 *p*-values from a genome-wide association test with GEMMA for composite fitness (survival x growth). Lake and experiment were included as covariates in the linear mixed model. SNPs that were significant at FDR < 0.05 are indicated in blue; red SNPs above dashed red line cross the threshold for Bonferroni significance at α = 0.05. First twenty-four scaffolds are sorted from largest to smallest and remaining scaffolds were pooled. The six genes associated with composite fitness which were both strongly differentiated ($F_{ST} > 0.95$) and differentially expressed between specialists (69) are annotated. **b-c)** Best-fit adaptive landscape for composite fitness using either morphology alone (**b:** flat surface with only directional selection) or morphology in combination with fitness associated SNPs (**c:** highly nonlinear surface). Best-fit model in C was a GAM including a thin plate spline for both LD axes, fixed effects of experiment and lake, and fixed effects of seven (see supplementary methods) SNPs most strongly associated with fitness shown in red. **c)** Three-dimensional view of **C** with relative positions of the three parental species indicated.

*Genome-wide association mapping of fitness*

From our LD-pruned dataset we used a linear mixed model in GEMMA to identify 132 SNPs in

regions that were strongly associated with composite fitness, including 13 which remained

significant at the conservative Bonferroni threshold (Fig. 2A, Table S3: See supplement for

results for survival and growth alone [Table S4-S5; Figure S5-6]). Gene ontologies for these 132

fitness-associated regions were significantly enriched for synaptic signaling and chemical

synaptic transmission (FDR rate < 0.01; Fig. S7; Table S5). Ontologies enriched at an FDR rate

< 0.05 were related to signaling and regulation of cell communication. We did not identify any

enrichment for ontologies related to craniofacial development which have previously been

identified to play a significant role in the adaptive divergence of these fishes (51, 65, 69, 74, 75).

This suggests that fitness-associated regions in our field experiments capture other components

of fitness in addition to the external phenotypes measured in previous studies. Surprisingly, our

fitness associated regions did not overlap with regions showing significant evidence of a hard

selective sweep (51). However, we did find six fitness-associated SNPs near genes differentially

expressed between specialists in addition to one that was also misexpressed in F1 hybrids (69).

This is particularly intriguing because gene misexpression in hybrids, i.e. genes differentially

expressed relative to both parents, could indicate the breakdown of genetic regulatory networks

in hybrids and intrinsic genetic incompatibilities that may impact their fitness in field enclosures

(69, 76).

We characterized whether genes in or near fitness-associated regions are implicated in

adaptive divergence of the specialists. Five fitness-associated genes have previously been shown

to include fixed divergent SNPs between specialists (*csad, glcci1, ino80c, mag, pim2*), and one

contained a fixed deletion between specialists (*med25*) (69). *med25* (Mediator Complex Subunit

25) is a craniofacial transcription factor associated with cleft palate in humans and zebrafish (77,

78). Four of the remaining genes shared across our study and McGirr et al. (69) are associated

with growth and/or body size measurements in other fishes. For example, *csad* plays an

important role in synthesizing taurine which is a rate-limiting enzyme affecting growth rate in

parrotfishes (79), rainbow trout (80), and Japanese flounder (81). Second, *glcci1* is associated

with the body depth/length ratio in the large yellow croaker (82). Third, *ino80c* is associated with

measures of body size in the Nile tilapia (83). The only gene associated with growth but not

composite fitness, *mettl21e*, was also misexpressed in F1 hybrids between scale-eaters and

molluscivores at eight days post-fertilization, and thus is a putative genetic incompatibility in this

system (69). Although it has not been associated with growth-related or body-size traits in fishes,

it is associated with intramuscular fat deposition in cattle (84). Taken together, these findings

support the interpretation that our fitness-associated regions are associated with unmeasured

traits, particularly physiological growth rates, or craniofacial shape in the case of the deletion in

*med25*, that affect fitness in our hybrid field experiments.

*Fitness-associated SNPs improve inference of the adaptive landscape*

We used linear discriminant axes and generalized additive modelling (GAM) to estimate

phenotypic fitness landscapes for the sequenced hybrids on a two-dimensional morphospace

indicating similarity to each of the three parental populations following previous studies (17, 18).

We then tested whether the inclusion of the 13 genomic regions most strongly associated with

fitness (red: Fig. 2A) in GAM models improved our inference of the underlying adaptive

landscape. Models with fitness-associated SNPs were invariably favored over models with

external morphology alone ($\Delta$AICc > 8.6: Tables S8-S9). Morphology-only models predicted a

flat linear landscape (Fig. 2B, Fig. S10); whereas including fitness-associated SNPs resulted in

inferences of increased curvature and complexity of the fitness surface, despite our limited

dataset of 139 hybrids relative to sample sizes in previous studies around 800 hybrids per

enclosure.

To reduce complexity of the full model estimated from 31 morphological traits including

all 13 fitness-associated SNPs, we fit an additional model including only the seven significant

fitness-associated SNPs in the full model. This reduced model was the best-fit; the inferred

adaptive landscape was complex and characterized by a fitness peak near hybrids resembling

generalist phenotypes separated by a small fitness valley from a second region of high fitness for

molluscivore hybrid phenotypes. Scale-eater hybrid phenotypes again corresponded to a large

fitness valley (Fig. 2B-C: For results pertaining to growth or survival see Supplemental Materials

– Fig. S10, Table S10-12). Compared to previous studies, the highest fitness optimum was

shifted from molluscivore phenotypes to generalists. This suggests that fitness-associated SNPs

increased the fitness of hybrids resembling generalists beyond expectations based on their

morphology alone, consistent with the hypothesis that fitness-associated SNPs are associated

with unmeasured traits affecting fitness. Indeed, visualization of haplotype frequency on the

fitness landscape supported this interpretation: one of the most common haplotypes associated

with high fitness was most frequent in hybrids resembling generalists and rare in hybrids

resembling specialists (Fig. S11).

*Trophic novelty is associated with isolation on the genotypic fitness landscape*

We reconstructed genotypic fitness networks from 1,498 candidate adaptive alleles previously

identified in this system (e.g. Fig. 3). These regions displayed significant evidence of a hard

selective sweep using both site frequency spectrum and LD-based methods, SweeD (85) and

OmegaPlus (86), and contained fixed or nearly fixed SNPs (Fst > 0.95) differentiating trophic

specialists across lakes (51). These fitness networks depict both hybrids and parental species in

genotypic space, with nodes representing SNP haplotypes and edges connecting mutational

neighbors (Fig. 3B). Genotype space is immense; using SNPs coded as homozygous reference,

heterozygote, or homozygous alternate, the number of potential haplotypes is equal to

$3^{\# \ SNPs \ in \ network}$. For instance, to construct a reduced network of even 100 SNPs, there would

be a total of $3^{100} = 5.17\ X\ 10^{57}$ possible nodes. Thus, unlike experimental studies of individual

proteins in haploid *E. coli* (29, 39) or yeast (40), it is not possible for us to investigate the full

breadth of genotypic space.

262
**Figure 3. Genotypic fitness networks connecting San Salvador Island pupfishes. a)** The majority of
nearly fixed or fixed variants ($F_{ST} \geq 0.95$) experiencing hard selective sweeps (hereafter 'adaptive alleles')
originated as standing genetic variation (SGV: molluscivores = 96%, scale-eaters = 92%), followed by
introgression (molluscivores = 4%, scale-eaters = 6%), and lastly *de novo* mutation (scale-eaters =
2%)(51). Pie charts shows adaptive alleles retained in our study for each species. **b)** Diagram illustrating
genotypic fitness networks and adaptive walks in a pair of species for two SNPs concatenated into
haplotypes. Species A and B are separated by four mutational steps. Dashed lines are inaccessible as they
result in a decrease in fitness; this leaves a single possible accessible evolutionary trajectory between
species A and B (indicated by bold arrows). Each node is associated with an empirical measure of hybrid
fitness. Edges are always drawn as directed, from low to high fitness. **c)** The same network as in B, with
fitness plotted on the Y-axis and number of mutational steps from species A to B on the X-axis. Dashed
lines are inaccessible (decreasing in fitness), solid lines are accessible (increasing in fitness), and the only
accessible path between species A and B is indicated by solid arrows. **d)** Genotypic network constructed
using a random sample of ten SNPs, sampled from all SNPs shown in **a**. Each edge between nodes is up
to five mutational steps away; edge width is proportional to mutational distance: wider edges connect
closer haplotypes. Node size (for hybrids) is scaled proportional to mean composite fitness; larger circles
correspond to fitter haplotypes. **e)** Median number of mutational steps within or between species (e.g. Fig.
3C). All pairwise comparisons using Tukey's HSD test (after FDR correction) were significant.
281

Instead, to understand the distribution of parental species and their hybrids in genotypic

space, we began by using a random sample of ten SNPs drawn from our set of candidate adaptive

alleles in this system. We chose ten SNPs for this first analysis for a larger sample of SNPs to

characterize the overall structure of the genotypic fitness network, while not sampling too vast a

space; already, the number of possible genotypes exceeds 59,000. Here, we plotted edges

between nodes up to five mutational steps away and found that generalists and molluscivores are

closer in the genotypic fitness network than either is to scale-eaters (Fig 4A), as expected based

on their genetic distance. Most scale-eaters appear quite isolated in genotypic space, separated

from the generalist cluster of nodes by $12.6 \pm 0.091$ (mean $\pm$ SE: $P < 0.001$) mutational steps and

from molluscivores by $16.3 \pm 0.060$ steps ($P < 0.001$). In contrast, molluscivores were separated

from generalists by $5.37 \pm 0.103$ steps ($P < 0.001$). Generalists show the greatest intrapopulation

distances, separated from each other by $6.08 \pm 0.088$ steps ($P < 0.001$). In contrast,

molluscivores exhibited the smallest intrapopulation distances, separated by $1.75 \pm 0.021$ steps

($P < 0.001$). Scale-eater intrapopulation distances were intermediate ($4.71 \pm 0.088$ steps: $P <$

0.001).

*Adaptive introgression and* de novo *mutations increase accessibility of novel fitness peaks*

Evolutionary accessibility reflects the probability that one genotypic trajectory will be favored by

natural selection over another, and thus can be related to the predictability of evolution (5, 29,

30, 87, 88). Genotypic trajectories with the minimum number of steps between two states that are

characterized by monotonically increasing fitness at each step are considered to be the most

accessible (5, 29, 30). That is, each mutational step is favored by natural selection and longer

paths are less probable (30). This provides two useful metrics of fitness network accessibility: 1)

The number of accessible paths and 2) the shortest length of accessible paths.

The adaptive radiation of pupfishes on San Salvador island originated within the last

10,000 years through a combination of selection on standing genetic variation, adaptive

introgression, and *de novo* mutations (51). However, it is unclear how each source of genetic

variation aided in the traversal of fitness paths or contributed to the colonization of novel fitness

peaks. To address this question, we constructed genotypic fitness networks composed of each of

the three main sources of adaptive genetic variation: standing genetic variation, introgression

from one of four focal Caribbean generalist populations, or *de novo* mutations unique to San

Salvador Island. We also examined combinations of these categories to better reflect the stages

of adaptive divergence. We compared sets of 1,000 random 5-SNP genotypic networks drawn

from different sources of adaptive variation and identified and scored the lengths of accessible

paths (i.e. those monotonically increasing in fitness) from generalist to each specialist. We asked

whether source of genetic variation affected 1) the number of accessible paths or 2) the shortest

length of accessible paths.

We discovered that both scale-eater and molluscivore specialist fitness peaks in

genotypic networks constructed from introgressed or *de novo* adaptive mutations were

significantly more accessible than networks constructed from standing genetic variation (Fig. 4C,

Table S13, S16). Random networks drawn from these sources of adaptive variation contained

more accessible fitness paths from generalist to specialists than networks constructed from

standing genetic variation, while controlling for differences in overall network size (Figs. S12-

S13, Table S13). Specifically, networks constructed from adaptive introgression had 126% more

accessible paths than those constructed from standing variation for molluscivores (median: $P <$

0.0001: Fig. 4C), and 39% more for scale-eaters ($P < 0.0001$: Fig. 4C). Likewise, networks

constructed from *de novo* mutations had 42% more accessible paths ($P < 0.0001$: Fig. 4C).

Consistent with past experimental studies in *E. coli* (29, 39) and yeast (40), accessible

paths were quite rare; the vast majority of genotypic trajectories were inaccessible, i.e. they

included one or more mutational steps that decreased in fitness from generalist to specialist.

Compared to standing variation, accessible paths were also significantly shorter in introgressed

(mean: 9% shorter in molluscivores: $P < 0.0001$, 23% shorter in scale-eaters: $P < 0.0001$) and *de*

*novo* mutation networks (14% shorter in scale-eaters $P < 0.0001$: Fig. 4D. Fig S12, Tables S13,

S17). Thus, introgressed and *de novo* adaptive mutations present more possible trajectories and

reduce the number of steps needed to traverse between species and access novel trophic

specialist fitness peaks (See Supplemental Materials Fig. S14-S15). These results are consistent

with a growing body of evidence that *de novo* and introgressed variation may contribute to rapid

speciation and evolution towards novel fitness peaks (38, 56, 62, 89–92).

**Figure 4. Adaptive introgression and *de novo* mutations increase access to specialist fitness peaks. a)** Example genotypic fitness networks constructed from five SNPs randomly drawn from introgressed adaptive loci for molluscivores (left) or scale-eaters (right). Edges connect nodes (haplotypes) that are separated by a single mutational step, arrows indicate the direction of mutational steps (directed towards higher fitness or parental haplotypes if not observed in hybrids), and red arrows indicate an example accessible trajectory. **b)** Accessibility of genotypic trajectories separating generalists from specialists as measured using networks constructed from all adaptive loci. Left: Proportion of accessible (i.e. monotonically increasing in fitness) paths per network, scaled by the total size of the network. Right: Number of mutational steps in the shortest accessible path. Because of the discrete nature of these data, means are plotted as large circles, with two standard errors shown. Due to the highly skewed nature of these distributions, pairwise significance among all network sets was determined using a nonparametric Kruskal-Wallis one-way analysis of variance following correction for multiple tests using an estimated False Discovery Rate. **c-d)** The same plots as in **b**, but for fitness landscapes constructed from adaptive loci originating from standing genetic variation (SGV), introgression, *de novo* mutation, or a combination.

*Conclusion*

Here, we have analyzed the first multi-species genotypic fitness landscape based on hybrid

fitness across multiple field experiments. We have shown that adaptive introgression and *de novo*

mutation increase the accessibility of novel fitness peaks, thus promoting the evolution of

novelty. Adaptation from standing genetic variation may initially be more rapid due to its higher

frequency, initial availability, and potentially reduced genetic load within a population (93–95).

Novel fitness peaks are not inaccessible from standing genetic variation alone – rather, they are

simply *less* accessible and thus we predict that standing genetic variation alone is less likely to

have triggered adaptive radiation in this system (93).

Once genetic variation arrives from introgression or arises *de novo* in a population,

colonization of new adaptive zones may be expedited due to the reduced number of mutational

steps in accessible paths. Furthermore, *de novo* mutations that undergo selective sweeps are also

predicted to have larger effect sizes (96). These findings are consistent with an interpretation of

adaptive radiation unfolding in stages (51, 97), first via standing genetic variation, next

introgression from across the Caribbean, and finally *de novo* mutations.

Our findings also provide additional support for the isolation of scale-eaters on the fitness

landscape and the rare evolution of scale-eating specialists despite comparable environments

across the Caribbean with nearly identical levels of adaptive standing genetic variation. That is,

adaptive *de novo* mutations may enable the crossing of a large fitness valley by increasing the

number of accessible fitness pathways through genotype space. This in turn may then trap scale-

eater populations on this peak by reducing the number of accessible fitness pathways back to the

generalist peak. In short, by increasing accessibility of adaptive walks to novel trophic specialist

niches, *de novo* and introgressed adaptive alleles simultaneously reduce the accessibility of

reverting to the generalist niche after the population reaches the new adaptive optimum.

**Acknowledgements:**

We thank Michelle St. John, David Tian, and Jacqueline Galvez for insightful discussion of the

results; the Gerace Research Centre and Troy Day for logistical support; the government of the

Bahamas for permission to collect, import, and export samples; the Vincent J. Coates Genomics

Sequencing Center and Functional Genomics Laboratory at UC Berkeley for performing whole

genome library preparation and sequencing (supported by NIH S10 OD018174 Instrumentation

Grant), and the University of California, Berkeley and University of North Carolina at Chapel

Hill for computational resources. This work was funded by the National Science Foundation

DEB CAREER grant #1749764, National Institutes of Health grant 5R01DE027052-02, the

University of North Carolina at Chapel Hill, and the University of California, Berkeley to CHM,

graduate research funding from the Museum of Vertebrate Zoology to EJR, and an NSF

Postdoctoral Research Fellowship in Biology under Grant No. 2109838 to AHP. Genomic data

are archived at the National Center for Biotechnology Information BioProject Database

(Accessions: PRJNA690558; PRJNA394148, PRJNA391309).

**Methods:**

*Sampling*

Our final dataset is comprised of 139 hybrid samples used in two separate field experiments (17,

18) on San Salvador island. Experiments were conducted in two lakes: Little Lake (N = 71) and

Crescent Pond (N = 68). Hybrids used in the first field experiment (17) were spawned in the lab

between December 2010 and January 2011; juveniles were raised for 2 months, individually

tagged by injecting a stainless steel sequential coded wire tag (Northwest Marine Technologies,

Inc.) into their left dorsal musculature, and photographed pre-release for morphometric analyses.

Experimental field enclosures consisted of high- and low-density treatments; density was varied

by the number of tagged juveniles released into each enclosure. Hybrids in the second field

experiment (18) were comprised of F4-F5 outbred juveniles resulting from crosses between all

three species. Individuals were spawned, raised, and tagged in an equivalent manner and

photographed prior to release. Experimental enclosures included high- and low-frequency

treatments in which the frequency of rare transgressive hybrid phenotypes was altered between

treatments in each lake.

All hybrids were measured for 32 external morphological traits (see below). Additionally,

we included sequenced parental species of the generalist (N = 17), molluscivores (N = 27) and

scale-eaters (N = 25) sampled from these two lakes previously included in Richards et al. (51).

Note that we treat samples from Little Lake and Osprey Lake as the same population because

these two lakes are connected within one hypersaline lake system and fish from these

populations are genetically undifferentiated (50, 51). For morphological analyses, we

additionally measured samples of 60 generalists, 38 molluscivores, and 60 scale-eaters raised in

the same laboratory common garden environment. A full list of samples is included in the

supplement (Table S1).

***Sequencing, genotyping and filtering***

Raw reads from a combined set of 396 samples (see supplement) were first mapped to the *C.*

*brontotheroides* reference genome (genome size = 1.16 Gb; scaffold N50 = 32 Mb) (51) using

bwa-mem (v. 0.7.2). Duplicate reads were identified and reads using MarkDuplicates, and BAM

424 indices were subsequently generated using the Picard software package (98). Samples were

425 genotyped following Richards et al., (51) according to GATK best practices (99). Specifically,

426 Single Nucleotide Polymorphisms (SNPs) were called and filtered using hard-filtering criteria in

427 HaplotypeCaller. We used the following criteria in our filtering approach: QD < 2.0; QUAL <

428 20; FS < 60; MQRankSum < -12.5; ReadPosRankSum < -8 (99–101).

429   Following initial genotyping with GATK, we subsequently filtered our data further using

430 VCFtools (102). Specifically, we filtered using the following flags: --maf 0.05; --min-alleles 2; --

431 max-alleles 2; --min-meanDP 7; --max-meanDP 100; --max-missing 0.85. Indels were removed.

432 To reduce non-independence among sites in our final dataset, we conservatively removed sites in

433 strong linkage disequilibrium using plink v1.9 (--indep-pairwise 10['kb'] 50 0.5: (103)). Doing so

434 resulted in the retention of 1,129,771 SNPs across 139 hybrid samples and the 69 samples from

435 Richards et al (51). Unless otherwise specified, these SNPs were used for all downstream

436 analyses.

438 ***Hybrid fitness measures***

439 We used three measures of fitness: survival, growth, or a composite measure of the two. Survival

440 was a binary trait indicating whether a fish survived (i.e. a tagged fish was recovered) or not

441 during its exposure period in field enclosures. Growth was a continuous measure, defined as the

442 proportional increase in Standard Length $\left(\frac{Final\ SL-Starting\ SL}{Starting\ SL}\right)$. Lastly, we defined composite

443 fitness as survival * growth, similar to the metric used in (104), and analogous to composite

444 fitness in Hereford (105) who used fecundity as their second fitness measure, rather than growth.

445 Because composite fitness represents the most information rich metric of fitness, we report

composite fitness results in the main text; results for growth and survival are included in the

supplement.

***Population genetic variation***

To visualize genetic variation present in hybrids and across lakes (Crescent Pond and Little

Lake), we first used a Principal Components Analysis (PCA) of genetic variation using plink

v1.90 ((103), Fig. 1), plotting the first two principal component axes using R (version 3.6.3 (R

Core team 2020). We then estimated admixture proportions in hybrids using ADMIXTURE

v1.3.0 (67). Parental species found on San Salvador Island were substantially differentiated

among Crescent Pond and Little Lake (50, 51); thus, independent ADMIXTURE analyses were

conducted for each lake. Because we were primarily interested in admixture proportions of

hybrids, we set K = 3 in these analyses, corresponding to the three parental species used in

hybrid crosses. Using admixture proportions of hybrid individuals, we tested the hypothesis that

ancestry predicts hybrid composite fitness in experimental field enclosures by fitting a

generalized additive model including either 1) scale-eater ancestry or 2) molluscivore ancestry

with fixed effects for experiment and lake. This was repeated for survival and growth separately.

***Genome-wide association tests***

To identify SNPs that were most strongly associated with fitness (survival, growth, or

composite), we implemented the linear mixed model (LMM) approach in GEMMA (v. 0.98.1:

(106)). This analysis was repeated using each fitness measure as the response variable. To

account for relatedness among samples we estimated the kinship matrix among all 139 hybrid

samples, which in turn were used in downstream LMMs. To account for the potentially

confounding effect of year/experiment and lake on estimated fitness measures, we included each

as covariates in the LMMs. To ensure rare variants were not included in these analyses, we only

included sites that had a minor allele frequency greater than 5% across all hybrids. A total of

933,520 SNPs were analyzed; 196,251 SNPS were excluded due to allele frequency change

following removal of parental species. SNPs strongly associated with fitness were identified as

1) a False Discovery Rate (FDR: Benjamini and Hochberg 1995) less than 0.05, or a 2) *P*-value

< 0.05 following a Bonferroni correction. We focus primarily on the sites identified by the

conservative Bonferroni correction, however.

***Gene ontology enrichment***

We annotated sites that were significantly associated with fitness using snpEff (107) and the

annotated *C. brontotheroides* reference genome (51). That is, we constructed a custom database

within snpEff using the functional annotations originally produced by Richards et al. (51), and

subsequently extracted information on the annotations and putative functional consequences of

each variant.

Using genes identified for each SNP that is significantly associated with one of the fitness

measures, we performed Gene Ontology (GO) enrichment analyses using ShinyGO v0.61 (108).

For genes identified as being intergenic, we included both flanking genes. As in Richards et al.

(2021), the gene symbol (abbreviation) database that had the greatest overlap with ours was that

of the human database; thus, we tested for enrichment of biological process ontologies curated

for human gene functions, based on annotations from Ensembl. Results are reported for

biological processes that were significantly enriched with FDR < 0.05. We then compared this

list of candidate loci to those identified in past studies of San Salvador Island pupfishes (51, 69,

109).

**Morphometrics**

We measured 31 external morphological traits for all 139 hybrids and 69 parental individuals

from Crescent Pond (30 generalists, 19 molluscivores, and 30 scale-eaters) and 85 from Little

Lake (30 generalists, 25 molluscivores, and 30 scale-eaters). We digitally landmarked dorsal and

lateral photographs (both sides) of each lab-reared hybrid (pre-release) or parent using DLTdv8

(110). Measurements include 27 linear characters and three angles. For nearly all individuals,

lateral measurements were collected from both lateral photographs and averaged. Morphological

variables were size-corrected using the residuals of a log10(trait) – log10(standard length)

regression standardized for selection analyses as outlined in the supplement. We used these 31

morphological traits to estimate two linear discriminant (LD) axes that best distinguished the

generalist, molluscivore, and scale-eater using the LDA function in the mass package in R. We

then used the resultant LD model to predict LD-scores for the 139 sequenced hybrids for later

use in generalized additive models.

**Estimation of adaptive landscapes**

We fit generalized additive models (GAMs) using the mgcv package v. 1.8.28 (111) in R to

estimate fitness landscapes relating the two discriminant axes (LD1-2) to fitness. All models

included a thin-plate spline fitted for the two linear discriminant axes, and we included both lake

and experiment in all models as fixed effects. Lake x experiment interaction terms were also

included in some models. Models were ranked using the corrected Akaike Information Criterion

(AICc) and were considered to be a significantly worse fit if ΔAICc > 4 from the best-fit model

(112). The best-fit model from the above approach was in turn used to visualize fitness

landscapes, plotting predicted values of fitness measures across LD space in R.

Using these results, we tested whether inclusion of SNPs that were strongly associated

with fitness (i.e., those that surpassed the 0.05 Bonferroni threshold) improved estimation of

fitness landscapes. We first extracted genotypes for the highly significant SNPs identified by

GEMMA (13 for composite fitness, four for growth: see section *Fitness-genotype association*

*test*), and coded these as either reference, single or double mutants using VCFtools (102). We

then used the best-fit models identified above and fit a range of models that included one or all

SNPs. Individual fitness-associated SNPs were treated as ordered factors (i.e. transition from

homozygous reference to heterozygote to homozygous alternate) and modeled using a factor-

smooth in the generalized additive models. Note that factor "smooths" are effectively modeled as

step-functions.

***Estimation of genotypic fitness networks***

We first estimated genotypic networks using sites previously shown to be highly divergent ($F_{ST}$

> 0.95) and having undergone hard selective sweeps in specialists. To do so, we identified the

SNPs in our unpruned full dataset overlapping with sites inferred to have undergone selective

sweeps (51), resulting in 380 SNPs for molluscivores and 1,118 SNPs for scale-eaters. We

subsequently constructed genotypic fitness networks in igraph v. 1.2.4.1 (113) following the

procedure outlined in the supplement.

To visualize the high-dimensional genotypic fitness network, we randomly sampled ten

adaptive loci 100 times and plotted haplotypes connected by edges if they were within five

mutational steps of one another (Figure 3C). Then we calculated the mean distance between all

species pairs (in number of mutational steps). We used pairwise Tukey's HSD tests to test whether

inter-species distances differed.

***Estimation of evolutionary accessibility***

We tested whether the evolutionary accessibility of genotypic fitness trajectories through

observed hybrid genotypes from generalist to each specialist species differed based on the source

of genetic variation. We restricted our investigation to networks composed of adaptive loci as

previously described (Figure 3A: Richards et al. 2021). This included a total of 380 SNPs in the

molluscivores, and 1,118 in the scale-eaters. The reduced number of adaptive SNPs sites in our

dataset as compared to that of (51) is due in large part to the increased stringency of our filtering.

We further partitioned these SNPs by their respective sources: standing genetic variation

(molluscivore N = 364; scale-eater N = 1,029), *de novo* mutation (scale-eater N = 24), or

introgression (molluscivore N = 16; scale-eater N = 65), again using the assignments from

Richards et al. (51). For analyses of trajectories between generalists and molluscivores, we

included only SNPs found to be sweeping in molluscivores; likewise, we included only SNPs

sweeping in scale-eaters for analysis of trajectories between generalists and scale-eaters.

The full procedure for constructing genotypic fitness networks, identifying accessible

paths, and quantifying accessibility is outlined in the supplement. Briefly, we randomly

generated 1,000 datasets of five SNPs comprised of either 1) standing genetic variation, 2)

adaptive introgression, 3) *de novo* mutation, 4) standing genetic variation + adaptive

introgression, or 5) standing genetic variation + *de novo* mutation. We additionally repeated this

procedure using all three classes of SNPs for scale-eaters to determine whether genotypic

trajectories separating generalists to molluscivores are more accessible than those between

generalists and scale-eaters. Because different sets of sites are sweeping in each specialist, we

conducted these analyses separately for each species. We then constructed genotypic networks,

in which nodes are haplotypes of SNPs encoded in 012 format (0 = homozygous reference, 1 =

heterozygote, 2 = homozygous alternate), and edges link mutational neighbors. When

determining whether a path was accessible or not, we only included paths for which each

mutational step (i.e. each intervening haplotype) between generalist to specialist was observed in

at least one hybrid sample.

For each random network sample and for each measure of fitness we calculated 1) the

minimum length of accessible paths between a random generalist and specialist sampled from

our sequenced individuals, 2) the number of accessible paths between the same generalist and

specialist pair, 3) the number of nodes and 4) the number of edges in the network. Larger

networks will necessarily have a greater number of potential paths, including both accessible and

inaccessible paths (Fig. S16), and we were interested in the relative proportion of accessible

adaptive pathways, not just the raw number. For this reason and because network size differs

among permutations, we divided the number of accessible paths in each random network

sampled by the number of nodes, thus normalizing the counts and making them comparable

across replicates and sources of variation. We assessed whether differences in these measures

among the three sources of variation or between the two alternate generalist to specialist

trajectories were significant using an ANOVA in R; due to the highly skewed nature of these

distributions, post-hoc pairwise significance was assessed using a nonparametric Kruskal-Wallis

one way analysis of variance in the agricolae package (114) in R.

**Literature Cited**

1. S. Wright, The roles of mutation, inbreeding, crossbreeding and selection in evolution. *Sixth Int. Congr. Genet.*, 356–366 (1932).

2. S. Gavrilets, Evolution and speciation on holey adaptive landscapes. *Trends Ecol. Evol.* **12**, 307–312 (1997).

3. S. Gavrilets, "High-dimensional fitness landscapes and speciation" in *Evolution—the Extended Synthesis*, M. Pigliucci, G. B. Müller, Eds. (MIT Press, 2010), pp. 51–64.

4. E. Svensson, R. Calsbeek, *The adaptive landscape in evolutionary biology*, E. Svensson, R. Calsbeek, Eds. (Oxford University Press, 2012).

5. I. Fragata, A. Blanckaert, M. A. Dias Louro, D. A. Liberles, C. Bank, Evolution in the light of fitness landscape theory. *Trends Ecol. Evol.* **34**, 69–82 (2019).

6. K. K. Fear, T. Price, The adaptive surface in ecology. *Oikos* **82**, 440–448 (1998).

7. G. G. Simpson, *Tempo and mode in evolution* (Columbia University Press, 1944).

8. S. Kauffman, S. Levin, Towards a general theory of adaptive walks on rugged landscapes. *J. Theor. Biol.* **128**, 11–45 (1987).

9. R. Lande, S. J. Arnold, The measurement of selection on correlated characters. *Evolution* **37**, 1226 (1983).

10. S. J. Arnold, M. E. Pfrender, A. G. Jones, "The adaptive landscape as a conceptual bridge between micro- and macroevolution" in *Microevolution Rate, Pattern, Process*, (Springer, Dordrecht, 2001), pp. 9–32.

11. S. J. Arnold, Performance surfaces and adaptive landscapes. *Integr. Comp. Biol.* **43**, 367–375 (2003).

12. D. Schluter, P. R. Grant, Determinants of morphological patterns in communities of Darwin's finches. *Am. Nat.* **123**, 175–196 (1984).

13. D. Schluter, Estimating the form of natural selection on a quantitative trait. *Evolution* **42**, 849–861 (1988).

14. A. P. Hendry, S. K. Huber, L. F. De León, A. Herrel, J. Podos, Disruptive selection in a bimodal population of Darwin's finches. *Proc. R. Soc. B Biol. Sci.* **276**, 753–759 (2009).

15. M. O. Beausoleil, *et al.*, Temporally varying disruptive selection in the medium ground finch (Geospiza fortis). *Proc. R. Soc. B Biol. Sci.* **286** (2019).

16. C. W. Benkman, Divergent selection drives the adaptive radiation of crossbills. *Evolution* **57**, 1176–1181 (2003).

17. C. H. Martin, P. C. Wainwright, Multiple fitness peaks on the adaptive landscape drive adaptive radiation in the wild. *Science* **339**, 208–211 (2013).

18. C. H. Martin, K. J. Gould, Surprising spatiotemporal stability of a multi-peak fitness landscape revealed by independent field experiments measuring hybrid fitness. *Evol. Lett.*, evl3.195 (2020).

19. S. Gavrilets, *Fitness landscapes and the origin of species* (Princeton University Press, 2004).

20. M. Turelli, N. H. Barton, J. A. Coyne, Theory and speciation. *Trends Ecol. Evol.* **16**, 330–343 (2001).

21. M. R. Servedio, J. W. Boughman, The role of sexual selection in local adaptation and speciation. *Annu. Rev. Ecol. Evol. Syst.* **48**, 85–109 (2017).

22. N. H. Barton, "Founder effect speciation" in *Speciation and Its Consequences*, D. Otte, J. A. Endler, Eds. (Sinauer, 1989), pp. 229–256.

23. U. Dieckmann, M. Doebeli, On the origin of species by sympatric speciation. *Nature* **400**,

354–357 (1999).

24. D. I. Bolnick, Can intraspecific competition drive disruptive selection? An experimental test in natural populations of sticklebacks. *Evolution* **58**, 608–618 (2004).

25. F. J. Weissing, P. Edelaar, G. S. van Doorn, Adaptive speciation theory: A conceptual review. *Behav. Ecol. Sociobiol.* **65**, 461–480 (2011).

26. J. E. Cresswell, C. Galen, Frequency-dependent selection and adaptive surfaces for floral character combinations: the pollination of Polemonium viscosum. *Am. Nat.* **138**, 1342–1353 (1991).

27. S. P. Otto, M. R. Servedio, S. L. Nuismer, Frequency-dependent selection and the evolution of assortative mating. *Genetics* **179**, 2091–2112 (2008).

28. B. C. Haller, A. P. Hendry, Solving the paradox of stasis: Squashed stabilizing selection and the limits of detection. *Evolution* **68**, 483–500 (2014).

29. D. M. Weinreich, N. F. Delaney, M. A. DePristo, D. L. Hartl, Darwinian evolution can follow only very few mutational paths to fitter proteins. *Science* **312**, 111–114 (2006).

30. J. Franke, A. Klözer, J. A. G. M. de Visser, J. Krug, Evolutionary accessibility of mutational pathways. *PLoS Comput. Biol.* **7**, 1002134 (2011).

31. R. A. Fisher, *The genetical theory of natural selection* (Oxford University Press, 1930).

32. H. A. Orr, The genetic theory of adaptation: A brief history. *Nat. Rev. Genet.* **6**, 119–127 (2005).

33. M. Karageorgi, *et al.*, Genome editing retraces the evolution of toxin resistance in the monarch butterfly. *Nature* **574**, 409–412 (2019).

34. M. A. Siddiq, D. W. Loehlin, K. L. Montooth, J. W. Thornton, Experimental test and refutation of a classic case of molecular adaptation in Drosophila melanogaster. *Nat. Ecol. Evol.* **1**, 25 (2017).

35. M. A. Siddiq, J. W. Thornton, Fitness effects but no temperature-mediated balancing selection at the polymorphic Adh gene of Drosophila melanogaster. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 21634–21640 (2019).

36. K. S. Singh, *et al.*, The genetic architecture of a host shift: An adaptive walk protected an aphid and its endosymbiont from plant chemical defenses. *Sci. Adv.* **6**, eaba1070 (2020).

37. F. Peng, *et al.*, Effects of beneficial mutations in pykF gene vary over time and across replicate populations in a long-term experiment with bacteria. *Mol. Biol. Evol.* **35**, 202–210 (2018).

38. Z. D. Blount, C. Z. Borland, R. E. Lenski, Historical contingency and the evolution of a key innovation in an experimental population of Escherichia coli. *Proc. Natl. Acad. Sci. U. S. A.* **105**, 7899–7906 (2008).

39. A. I. Khan, D. M. Dinh, D. Schneider, R. E. Lenski, T. F. Cooper, Negative epistasis between beneficial mutations in an evolving bacterial population. *Science* **332**, 1193–1196 (2011).

40. C. Bank, S. Matuszewski, R. T. Hietpas, J. D. Jensen, On the (un)predictability of a large intragenic fitness landscape. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 14085–14090 (2016).

41. P. Nosil, *et al.*, Ecology shapes epistasis in a genotype–phenotype–fitness map for stick insect colour. *Nat. Ecol. Evol.* **4**, 1673–1684 (2020).

42. C. H. Martin, P. C. Wainwright, A remarkable species flock of Cyprinodon pupfishes endemic to San Salvador Island, Bahamas. *Bull. Peabody Museum Nat. Hist.* **54**, 231–241 (2013).

43. C. H. Martin, Context dependence in complex adaptive landscapes: frequency and trait-

674   dependent selection surfaces within an adaptive radiation of Caribbean pupfishes.
675   *Evolution* **70**, 1265–1282 (2016).

676 44. B. J. Turner, D. D. Duvernell, T. M. Bunt, M. G. Barton, Reproductive isolation among
677   endemic pupfishes (Cyprinodon) on San Salvador Island, Bahamas: Microsatellite
678   evidence. *Biol. J. Linn. Soc.* **95**, 566–582 (2008).

679 45. C. H. Martin, P. C. Wainwright, Trophic novelty is linked to exceptional rates of
680   morphological diversification in two adaptive radiations of cyprinodon pupfish. *Evolution*
681   **65**, 2197–2212 (2011).

682 46. C. H. Martin, The cryptic origins of evolutionary novelty: 1000-fold faster trophic
683   diversification rates without increased ecological opportunity or hybrid swarm. *Evolution*
684   **70**, 2504–2519 (2016).

685 47. M. E. St. John, K. E. Dixon, C. H. Martin, Oral shelling within an adaptive radiation of
686   pupfishes: Testing the adaptive function of a novel nasal protrusion and behavioural
687   preference. *J. Fish Biol.* **97**, 163–171 (2020).

688 48. C. H. Martin, J. A. McGirr, E. J. Richards, M. E. St. John, How to investigate the origins
689   of novelty: Insights gained from genetic, behavioral, and fitness perspectives. *Integr. Org.*
690   *Biol.* **1** (2019).

691 49. M. E. St. John, R. Holzman, C. H. Martin, Rapid adaptive evolution of scale-eating
692   kinematics to a novel ecological niche. *J. Exp. Biol.* **223** (2020).

693 50. C. H. Martin, L. C. Feinstein, Novel trophic niches drive variable progress towards
694   ecological speciation within an adaptive radiation of pupfishes. *Mol. Ecol.* **23**, 1846–1862
695   (2014).

696 51. E. J. Richards, *et al.*, A vertebrate adaptive radiation is assembled from an ancient and
697   disjunct spatiotemporal landscape. *Proc. Natl. Acad. Sci.* **118**, e2011811118 (2021).

698 52. C. H. Martin, P. C. Wainwright, On the measurement of ecological novelty: Scale-eating
699   pupfish are separated by 168 my from other scale-eating fishes. *PLoS One* **8**, e71164
700   (2013).

701 53. R. Froese, D. Pauldy, FishBase. *FishBase* (2021).

702 54. O. Seehausen, others, Genomics and the origin of species. *Nat. Rev. Genet.* **15**, 176–192
703   (2014).

704 55. S. H. Martin, C. D. Jiggins, Interpreting the genomic landscape of introgression. *Curr.*
705   *Opin. Genet. Dev.* **47**, 69–74 (2017).

706 56. D. A. Marques, J. I. Meier, O. Seehausen, A Combinatorial View on Speciation and
707   Adaptive Radiation. *Trends Ecol. Evol.* **34**, 531–544 (2019).

708 57. J. W. Poelstra, E. J. Richards, C. H. Martin, Speciation in sympatry with ongoing
709   secondary gene flow and a potential olfactory trigger in a radiation of Cameroon cichlids.
710   *Mol. Ecol.* **27**, 4270–4288 (2018).

711 58. T. C. Nelson, W. A. Cresko, Ancient genomic variation underlies repeated ecological
712   adaptation in young stickleback populations. *Evol. Lett.* **2**, 9–21 (2018).

713 59. K. A. Thompson, M. M. Osmond, D. Schluter, Parallel genetic evolution and speciation
714   from standing variation. *Evol. Lett.* **3**, 129–141 (2019).

715 60. J. B. Pease, D. C. Haak, M. W. Hahn, L. C. Moyle, Phylogenomics reveals three sources
716   of adaptive variation during a rapid radiation. *PLoS Biol.* **14**, e1002379 (2016).

717 61. E. J. Richards, J. W. Poelstra, C. H. Martin, Don't throw out the sympatric speciation with
718   the crater lake water: fine-scale investigation of introgression provides equivocal support
719   for causal role of secondary gene flow in one of the clearest examples of sympatric

speciation. *Evol. Lett.* **2**, 524–540 (2018).

62. J. I. Meier, *et al.*, Ancient hybridization fuels rapid cichlid fish adaptive radiations. *Nat. Commun.* **8**, 14363 (2017).

63. M. D. McGee, *et al.*, The ecological and genomic basis of explosive adaptive radiation. *Nature* **586**, 75–79 (2020).

64. I. Irisarri, *et al.*, Phylogenomics uncovers early hybridization and adaptive loci shaping the radiation of Lake Tanganyika cichlid fishes. *Nat. Commun.* **9**, 1–12 (2018).

65. E. J. Richards, C. H. Martin, Adaptive introgression from distant Caribbean islands contributed to the diversification of a microendemic adaptive radiation of trophic specialist pupfishes. *PLoS Genet.* **13**, e1006919 (2017).

66. C. H. Martin, K. J. Gould, Surprising spatiotemporal stability of a multi-peak fitness landscape revealed by independent field experiments measuring hybrid fitness. *Evol. Lett.* **4**, 530–544 (2020).

67. D. H. Alexander, J. Novembre, K. Lange, Fast model-based estimation of ancestry in unrelated individuals. *Genome Res.* **19**, 1655–1664 (2009).

68. D. H. Alexander, K. Lange, Enhancements to the ADMIXTURE algorithm for individual ancestry estimation. *BMC Bioinformatics* **12**, 1–6 (2011).

69. J. A. McGirr, C. H. Martin, Ecological divergence in sympatry causes gene misexpression in hybrids. *Mol. Ecol.* **29**, 2707–2721 (2020).

70. M. E. Arnegard, *et al.*, Genetics of ecological divergence during speciation. *Nature* **511**, 307–311 (2014).

71. S. Wang, J. J. Hard, F. Utter, Genetic variation and fitness in salmonids. *Conserv. Genet.* **3**, 321–333 (2002).

72. R. Leimu, P. Mutikainen, J. Koricheva, M. Fischer, How general are positive relationships between plant population size, fitness and genetic variation? *J. Ecol.* **94**, 942–952 (2006).

73. D. Schluter, *et al.*, Fitness maps to a large-effect locus in introduced stickleback populations. *Proc. Natl. Acad. Sci. U. S. A.* **118**, 1914889118 (2021).

74. C. H. Martin, P. A. Erickson, C. T. Miller, The genetic architecture of novel trophic specialists: larger effect sizes are associated with exceptional oral jaw diversification in a pupfish adaptive radiation. *Mol. Ecol.* **26**, 624–638 (2017).

75. J. A. McGirr, C. H. Martin, Parallel evolution of gene expression between trophic specialists despite divergent genotypes and morphologies. *Evol. Lett.* **2**, 62–75 (2018).

76. J. Kulmuni, A. M. Westram, Intrinsic incompatibilities evolving as a by-product of divergent ecological selection: Considering them in empirical studies on divergence with gene flow. *Mol. Ecol.* **26**, 3093–3103 (2017).

77. Y. Nakamura, *et al.*, Wwp2 is essential for palatogenesis mediated by the interaction between Sox9 and mediator subunit 25. *Nat. Commun.* **2**, 1–10 (2011).

78. L. Mork, G. Crump, "Zebrafish craniofacial development: A Window into early patterning" in *Current Topics in Developmental Biology*, Y. Chai, Ed. (Academic Press Inc., 2015), pp. 235–269.

79. S. J. Lim, *et al.*, Taurine is an essential nutrient for juvenile parrot fish Oplegnathus fasciatus. *Aquaculture* **414–415**, 274–279 (2013).

80. T. G. Gaylord, A. M. Teague, F. T. Barrows, Taurine supplementation of all-plant protein diets for rainbow trout (Oncorhynchus mykiss). *J. World Aquac. Soc.* **37**, 509–517 (2006).

81. M. Yokoyama, T. Takeuchi, G. S. Park, J. Nakazoe, Hepatic cysteinesulphinate decarboxylase activity in fish. *Aquac. Res.* **32**, 216–220 (2001).

82. Z. Zhou, *et al.*, Genome-wide association study of growth and body-shape related traits in
large yellow croaker (Larimichthys crocea) using ddRAD sequencing. *Mar. Biotechnol.*
**21**, 655–670 (2019).
83. G. M. Yoshida, J. M. Yáñez, Multi-trait GWAS using imputed high-density genotypes
from whole-genome sequencing identifies genes associated with body traits in Nile tilapia.
*BMC Genomics* **22**, 1–13 (2021).
84. L. F. S. Fonseca, *et al.*, Gene expression profiling and identification of hub genes in
Nellore cattle with different marbling score levels. *Genomics* **112**, 873–879 (2020).
85. P. Pavlidis, D. Živković, A. Stamatakis, N. Alachiotis, SweeD: Likelihood-based
detection of selective sweeps in thousands of genomes. *Mol. Biol. Evol.* **30**, 2224–2234
(2013).
86. N. Alachiotis, A. Stamatakis, P. Pavlidis, OmegaPlus: A scalable tool for rapid detection
of selective sweeps in whole-genome datasets. *Bioinformatics* **28**, 2274–2275 (2012).
87. F. J. Poelwijk, D. J. Kiviet, D. M. Weinreich, S. J. Tans, Empirical fitness landscapes
reveal accessible evolutionary paths. *Nature* **445**, 383–386 (2007).
88. L. Ferretti, D. Weinreich, F. Tajima, G. Achaz, Evolutionary constraints in fitness
landscapes. *Heredity (Edinb)*. **121**, 466–481 (2018).
89. T. C. Nelson, *et al.*, Ancient and recent introgression shape the evolutionary history of
pollinator adaptation and speciation in a model monkeyflower radiation (Mimulus section
Erythranthe). *PLoS Genet.* **17**, e1009095 (2021).
90. H. Svardal, *et al.*, Ancestral hybridization facilitated species diversification in the lake
malawi cichlid fish adaptive radiation. *Mol. Biol. Evol.* **37**, 1100–1113 (2020).
91. S. Lamichhaney, *et al.*, Rapid hybrid speciation in Darwin's Finches. *Science* **359**, 224–
228 (2018).
92. P. R. Grant, B. R. Grant, Hybridization increases population variation during adaptive
radiation. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 23216–23224 (2019).
93. R. D. H. Barrett, D. Schluter, Adaptation from standing genetic variation. *Trends Ecol.*
*Evol.* **23**, 38–44 (2008).
94. J. Hermisson, P. S. Pennings, Soft sweeps and beyond: understanding the patterns and
probabilities of selection footprints under rapid adaptation. *Methods Ecol. Evol.* **8**, 700–
716 (2017).
95. P. W. Hedrick, Adaptive introgression in animals: examples and comparison to new
mutation and standing variation as sources of adaptive variation. *Mol. Ecol.* **22**, 4606–
4618 (2013).
96. S. Matuszewski, J. Hermisson, M. Kopp, Catch me if you can: Adaptation from standing
genetic variation to a moving phenotypic optimum. *Genetics* **200**, 1255–1274 (2015).
97. J. T. Streelman, P. D. Danley, The stages of vertebrate evolutionary radiation. *Trends*
*Ecol. Evol.* **18**, 126–131 (2003).
98. Broad Institute, Picard Toolkit (2018).
99. M. A. Depristo, *et al.*, A framework for variation discovery and genotyping using next-
generation DNA sequencing data. *Nat. Genet.* **43**, 491–501 (2011).
100. R. Poplin, *et al.*, Scaling accurate genetic variant discovery to tens of thousands of
samples. *bioRxiv*, 201178 (2017).
101. C. D. Marsden, *et al.*, Diversity, differentiation, and linkage disequilibrium: Prospects for
association mapping in the malaria vector anopheles arabiensis. *G3 Genes, Genomes,*
*Genet.* **4**, 121–131 (2014).

102. P. Danecek, *et al.*, The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158
(2011).
103. S. Purcell, *et al.*, PLINK: A tool set for whole-genome association and population-based
linkage analyses. *Am. J. Hum. Genet.* **81**, 559–575 (2007).
104. C. T. DiVittorio, *et al.*, Natural selection maintains species despite frequent hybridization
in the desert shrub Encelia. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 33373–33383 (2021).
105. J. Hereford, A quantitative survey of local adaptation and fitness trade-offs. *Am. Nat.* **173**,
579–588 (2009).
106. X. Zhou, M. Stephens, Genome-wide efficient mixed-model analysis for association
studies. *Nat. Genet.* **44**, 821–824 (2012).
107. P. Cingolani, *et al.*, Using Drosophila melanogaster as a model for genotoxic chemical
mutational studies with a new program, SnpSift. *Front. Genet.* **3** (2012).
108. S. X. Ge, D. Jung, D. Jung, R. Yao, ShinyGO: A graphical gene-set enrichment tool for
animals and plants. *Bioinformatics* **36**, 2628–2629 (2020).
109. J. A. McGirr, C. H. Martin, Few fixed variants between trophic specialist pupfish species
reveal candidate cis-regulatory alleles underlying rapid craniofacial divergence. *Mol. Biol.*
*Evol.* **38**, 405–423 (2021).
110. T. L. Hedrick, Software techniques for two- and three-dimensional kinematic
measurements of biological and biomimetic systems. *Bioinspiration and Biomimetics* **3**,
034001 (2008).
111. S. N. Wood, Fast stable restricted maximum likelihood and marginal likelihood estimation
of semiparametric generalized linear models. *J. R. Stat. Soc. Ser. B (Statistical Methodol.*
**73**, 3–36 (2011).
112. K. P. Burnham, D. R. Anderson, *Model Selection and Multimodel Inference* (Springer
New York, 2002).
113. G. Csardi, T. Nepusz, The igraph software package for complex network research.
*InterJournal Complex Syst.* **1695**, 1–9 (2006).
114. F. de Mendiburu, agricolae: Statistical procedures for agricultural research (2020).
115. T. Guillerme, dispRity: A modular R package for measuring disparity. *Methods Ecol.*
*Evol.* **9**, 1755–1763 (2018).
116. B. J. Knaus, N. J. Grünwald, VCFR: a package to manipulate and visualize variant call
format data in R. *Mol. Ecol. Resour.* **17**, 44–53 (2017).
117. S. Der Yang, T. S. Lin, F. G. Liu, C. H. Liou, Influence of dietary phosphorus levels on
growth, metabolic response and body composition of juvenile silver perch (Bidyanus
bidyanus). *Aquaculture* **253**, 592–601 (2006).
118. J. M. Costa, *et al.*, Inadequate dietary phosphorus levels cause skeletal anomalies and alter
osteocalcin gene expression in Zebrafish. *Int. J. Mol. Sci.* **19**, 364 (2018).
119. H. Yu, *et al.*, Effects of stocking density and dietary phosphorus levels on the growth
performance, antioxidant capacity, and nitrogen and phosphorus emissions of juvenile
blunt snout bream (Megalobrama amblycephala). *Aquac. Nutr.* **27**, 581–591 (2021).
852

**Supplementary Methods**
*Sampling of hybrid individuals*
Samples of hybrid *Cyprinodon* pupfish included herein were first collected following two
separate fitness experiments, conducted on San Salvador Island in 2011 (17) and 2016 (18)
respectively. Experiments were carried out in two lakes: Little Lake (LL), and Crescent Pond
(CP). Following their initial collection at the conclusion of their respective experiments (see (17)
and (18) for protocols), samples were stored in ethanol. In late 2018, 149 hybrid samples were
selected for use in this experiment. Of these, 27 are from the experiment conducted in 2011 (14
from LL, 13 from CP), and the remaining 122 are from the 2016 experiment (58 from LL, 64
from CP). Due to reduced sample size for some species within Little Lake, we include fish
obtained from Osprey Lake for downstream analyses comparing hybrids to Little Lake, as the
two comprise a single, interconnected body of water.
865
*Genomic Library Prep*
DNA was extracted from the muscle tissue of hybrids using DNeasy Blood and Tissue kits
(Qiagen, Inc.); these extractions were then quantified using a Qubit 3.0 fluorometer (Thermo
Scientific, Inc). Genomic libraries were prepared by the Vincent J. Coates Genomic Sequencing
Center (QB3) on the automated Apollo 324 system (WaterGen Biosystems, Inc.). Samples were
fragmented using a Covaris sonicator, and barcoded with Illumina indices. Samples were quality
checked using a Fragment Analyzer (Advanced Analyical Technologies, Inc.). All samples were
sequenced to approximately 10x raw coverage on an Illumina NovaSeq.
874
*Genotyping and filtering*
Because we are interested in genomic variants found not only within our hybrid samples but
across San Salvador island and the Caribbean, we conducted genotyping including all 247
samples from Richards et al. (51). These samples include members of the three species (*C.*
*variegatus*, *C. brontotheroides*, *C. desquamator*) found on San Salvador Island, as well as
individuals of *C. variegatus* found throughout the Caribbean, and numerous outgroups (*C.*
*laciniatus*, *C. higuey*, *C. dearborni*, *Megupsilon aporus*, and *Cualac tesselatus*). We then
excluded *M. aporus* and *C. tesselatus* along with 18 additional samples for which necessary data
for downstream analyses were missing (e.g. quality photographs for the collection of
morphological data). This approach led to the retention of a total of 4,206,786 total SNPs and
139 hybrid individuals.
886
*Morphometrics*
Because the morphological measurements used in the 2013 and 2016 experiments differ slightly,
we remeasured all sequenced hybrid individuals and up to 30 individuals of each parent species
for the 30 morphological characters described in Martin & Gould (18), as well as for standard
length (SL).
For each photograph, unit-scale was obtained by additionally landmarking points on a
regular grid included in each photograph using DLTdv8a (110). Landmark data (x-y coordinates
in units of pixels) were subsequently uploaded into R and converted to millimeters (in the case of
linear measurements) or degrees (for angular measurements) using a custom script.
We then assessed, for each trait, the need for size-correction. That is, we sought to avoid
an outsized role of body size in downstream interpretation. Thus, if a trait was colinear with SL,
we regressed the two (treating SL as the predictor) and took the residuals. In each case, both SL

and the response were Log-10 transformed. Subsequently, residuals were scaled such that their
distribution had a mean of 0, and a standard deviation of 1. Traits that did not need size
correction were also Log-transformed and unit-scaled.
We used these morphological measurements to estimate two linear discriminant (LD)
axes that distinguish the generalist, molluscivore, and scale-eater using the LDA function in R.
That is, we used morphological data from the 165 parental fish to estimate LD scores for each
individual. Doing so, we were able to correctly assign individual fish to their corresponding
species with 99.4% accuracy (Fig. S8-S9: Table S7). Attempting to predict species assignment
by lake did not improve this prediction accuracy (instead reducing prediction accuracy to
98.7%); consequently, we proceeded with the LD axes estimated without accounting for lake.
We additionally asked whether 1) specialists were more morphologically constrained
than generalists and 2) if hybrids were less constrained than the three parental species. To do so,
we calculated morphological disparity per group using the dispRity.per.group function
implemented in the R package dispRity v1.3.3 (115). Specifically, this function calculates the
median distance between each row (sample) and the centroid of the matrix per group. By
bootstrapping the data 100 times, we tested the hypothesis that each of the three parental species
differed significantly in their morphological disparity, first using an ANOVA, followed by a
post-hoc pairwise t-test to assess pairwise significance in R. We corrected for multiple tests
using an FDR correction.
To test whether genetic distance predicts morphological distance, we calculated two
distance matrices. First, we calculated pairwise Euclidean distances between all hybrids using all
morphological variables. Then, we calculated pairwise genetic distances between all hybrids
using the final set of SNPs described above with the genet.dist function implemented in vcfR
(116). We then fit a simple linear model, regressing genetic distance on morphological distance.
923
***Estimation of Adaptive Landscapes***
We sought to characterize the extent to which the three measures of fitness are predicted by
morphology alone and to, in turn, visualize fitness landscapes for our sequenced hybrids. To do
so, we fitted six generalized additive models (GAMs) using the mgcv package v. 1.8.28 (111) in
R. All models included a thin-plate spline fitted for the two linear discriminant axes. Because we
have strong *a priori* knowledge that fitness outcomes will be contingent to some extent on
experiment year and on the lake in which hybrids were placed, we include experiment and lake
in all fitted models, modeled either as fixed effects, or as an interaction term between the two.
Additionally, we fitted models that included individual splines for each linear discriminant axis,
either with or without experiment or lake as a factor smooth. The full list of models and their
respective fits are included in the supplement (Supplementary Tables S8-S10).
For composite fitness, we excluded three SNPs that were within close proximity (i.e. <
1000bp) to a SNP that was more significantly associated. Because of the reduced number of
significantly associated SNPs identified for growth (four) as compared to composite fitness (ten),
we were able to fit and compare all combinations of significantly associated SNPs for the former.
The best-fit model for composite fitness was also the most complex, including all fitness
associated SNPs. Thus, to reduce model complexity, we fit one additional model, excluding any
of the three SNP (fixed effect) that was not significant in the full model. The full range of models
and their associated fits are reported in Tables S11-S12. As before, predicted fitness values
across LD space were extracted from the best-fit model and plotted using R.
944

***Genotypic Fitness Networks***
Recent work has shown that the adaptive radiation of the pupfish of San Salvador island involved
selection on standing genetic variation, adaptive introgression, and *de novo* mutation (51).
Furthermore, the specialists on San Salvador Island received approximately twice as much
adaptive introgression as did generalists on neighboring islands. These findings imply that each
source of genetic variation may exert a unique influence on the fitness landscape, in turn
facilitating the radiation. We sought to explore this possibility and so estimated genotypic fitness
networks using sites previously shown to have undergone hard selective sweeps in specialists. To
do so, we identified the SNPs in our un-thinned dataset overlapping with sites inferred to have
undergone selective sweeps (51) to produce two datasets (one for each specialist). We then
constructed networks using the following procedure:
956
1) Target SNPs were extracted from the vcf file of all sequenced hybrids and parental species
from Crescent Pond and Little Lake in 0/1/2 format using VCFtools. That is, individuals
genotyped as homozygote reference were coded as 0, heterozygotes as 1, and homozygote
alternatives as 2. Note again that the reference genome used was that of the molluscivore,
*C. brontotheroides*.
2) These SNPs were subsequently loaded into R, and concatenated into haplotypes, such that
each sequenced individual has a haplotype. These per-sample haplotypes were
subsequently associated with metadata, namely the observed binary survival, growth,
assignment as hybrid or one of the three parental species, and lake of origin.
3) Each haplotype was subsequently collapsed and summarized, such that mean survival,
growth, and composite fitness are retained for each unique haplotype. Additionally, the
number of hybrids, generalists, molluscivores and scale-eaters that have the haplotype were
recorded.
4) The distance, in number of mutational steps was then calculated for each pairwise
combination of haplotypes. For example, the distance between haplotype 000 and 001 is a
single mutational step, whereas haplotypes 000 and 002 are two steps away.
5) Lastly, we constructed networks using the R package igraph v. 1.2.4.1 (113). Specifically,
nodes represent haplotypes, and edges are drawn between haplotypes that are mutational
neighbors (i.e. are a single mutational step away). Nodes present in hybrids were colored
and sized proportional to their respective mean fitness. Nodes present only in parental
species were colored according to the species in which that haplotype is unique to.
978
***Estimation of evolutionary accessibility***
The large number of SNPs in our dataset above raises numerous challenges in the visualization
and summarization of fitness networks. Perhaps most significant, is that as the number of sites
assessed increases, the sequence space increases vastly; the number of potential haplotypes is
defined by 3 to the power of the number of SNPs, and the number of potential edges is defined
by the number of haplotypes choose 2. Consequently, networks constructed from a larger
number of focal SNPs are comprised of haplotypes that are separated on average by more
mutational steps than those constructed from fewer SNPs. Because we can only interpret the
fitness consequences of evolutionary trajectories for which we have data along each mutational
step, we restricted analysis to haplotypes that are mutational neighbors (i.e., separated by a single
mutational step).

To do so, we developed a permutation approach to construct fitness networks from SNPs
that were sourced from the three sources of genetic variation defined above as well as their
combinations (i.e. standing genetic variation + adaptive introgression or de novo mutation).
Specifically, from each set we sampled five SNPs up to 1000 times. Then, from each
permutation, we constructed fitness networks using the five steps defined above; these networks
served as the subsequent assessment of evolutionary accessibility of genotypic trajectories
separating the generalists from either specialist. Edges were only drawn such that the network is
directed; that is, edges were drawn from low to high fitness nodes.
We constructed networks using both parental species and hybrids. For each trajectory
under consideration (generalist → molluscivore & generalist → scale-eater), we first identified
all generalist to specialist trajectories that are connected by accessible (monotonically increasing)
paths. From these connected generalist-specialist node pairs, we randomly sampled a single pair.
We then identified the number and length of accessible paths separating the generalist node from
the specialist node and recorded these values using the "all_simple_paths" function in igraph and
excluded any paths that traversed through haplotypes not found in any hybrid (i.e., exclusive to
parental species) and thus had no information on fitness.
1006
**1007   Supplementary Results**
**1008   *Genomic associations recovered for composite fitness and growth, not survival***
Whereas we identified 132 SNPs that were associated with composite fitness, only 58 were
associated with growth and none were associated with survival. Of the SNPs associated with
growth, only four remained significant using the conservative Bonferroni threshold. Across all
significant sites (either via FDR or Bonferroni correction) a total of 11 were shared across
analyses. The only gene proximate to a growth associated SNP was *csad.* Lastly, we found a
single gene shared between our study and the 125 ecological DMIs (putatitve genetic
incompatibilities that are differentially expressed among specialists and misexpressed in F1
hybrids) presented in McGirr & Martin (69). This gene, associated with growth (but not
composite fitness) in our study, is *mettl21e.*
When considering SNPs found to be associated with growth, we did not identify any gene
ontologies that were significantly enriched at a False Discovery Rate < 0.01. However, looking
at those enriched at an FDR < 0.05, we do observe a number of ontologies related to biosynthetic
processes, and regulation of metabolic processes (Fig. S6). Specifically, the greatest (and most
significant) enrichment was for that of phosphorus and phosphate containing compound
metabolic processes and their regulation. Phosphorous deficiencies have previously been
associated with poor growth in silver perch (*Bidyanus bidyanus*: (117)) and skeletal deformaties
(including vertebral compression and craniofacial deformaties) in zebrafish (*Danio rerio*: (118)).
Similarly, blunt snout bream (*Megalobrama amblycephala*) exhibited greater growth rates with
increasing phosphorous levels in their diets (119). In short, enrichment of growth-associated
SNPs for ontologies pertaining to phosphorous metabolism is consistent with the substantial
literature documenting that phosphorous availability and metabolism is a determinant of growth
in fishes.
1031
**1032   *Morphological variation within sampled hybrids***
As in the previous two experiments, there is a relative paucity of hybrids exhibiting the
morphologies that characterize either specialist. Rather, most hybrids fall near the generalists,
with a number exhibiting transgressive morphologies (Fig. S9A). As expected, both specialists

exhibit reduced morphological disparity as compared to generalists, and hybrids show the
greatest (Fig. S9B). That is, the specialists appear more morphologically constrained than
generalists, falling on average closer to the group centroid. Interestingly molluscivores exhibit
the least disparity, even less so than scale-eaters.
1040
***Fitness-associated SNPs improve estimation of the adaptive landscape***
Using morphology alone, the best fit generalized additive model for survival, growth, and
composite fitness were simpler than the model for composite fitness using both morphology and
fitness-associated SNPs (Tables S8-10). For survival and composite fitness (Figure S10A &
S10B), this model included a thin-plate spline for LD1 & LD2, with experiment and lake
included as fixed effects. The resultant landscape was also similar for these two analyses,
supporting an interpretation of directional selection, favoring molluscivores. For growth, the
best-fit model had the thin-plate spline for LD1 & LD2, but included an interaction term between
experiment and lake (Table S9). In contrast to the previous two models, the landscape predicted
using growth as our proxy of fitness supported an interpretation of directional selection in favor
or hybrids most similar to generalists, and to a lesser extent, scale-eaters.
Notably, model selection using AICc invariably supported the inclusion of fitness
associated SNPs for growth and composite fitness (Tables S11-S12). For growth, the best-fit
model including genotypes was an improvement of 22.99 AICc over the model including
morphology alone, whereas for composite fitness, the improvement was 94.527 AICc.
Interestingly, the best-fit models and growth including associated SNPs was similar to that of the
landscape without fitness-associated SNPs, but largely supported an interpretation of directional
selection in favor of scale-eaters, and to a lesser extent generalists.
1059
***Evolutionary accessibility is influenced by the source of genetic variation***
As with composite fitness, we find that fitness landscapes including either adaptive introgression
or *de novo* mutation are always smaller (i.e. fewer nodes and edges in the network) than
networks constructed from standing genetic variation (SGV) alone (Fig. S13; Table S13-S15). In
fact, for networks constructed using growth as our fitness proxy, the influence appears even
stronger. These results hold even when rarefying our standing genetic variation SNPs to 65, the
number of introgressed variants in scale-eaters and the second largest set of variants (data not
shown).
Similar to composite fitness, we find that fitness landscapes estimated using both survival
(but see generalist to scale-eater) and growth including adaptive introgression have both
proportionally more accessible paths between generalists and specialists that are shorter (Fig.
S14-S15, Table S13, S16-S17). However, there are several exceptions both for fitness landscapes
constructed using survival as our proxy of fitness and looking at genotypic trajectories separating
generalists from scale-eaters. First, we found that fitness landscapes constructed using adaptive
introgression alone actually had relatively fewer accessible paths than those constructed using
standing variation (Fig. S14A, Tables S16), however as with other analyses, these accessible
paths were significantly shorter (Fig. S14B, Table S16). Second, we did not observe significant
differences between any other group in the relative number of accessible paths (Fig. S14A, Table
S16). Lastly, we did not find that accessible paths in networks constructed from a combination of
standing genetic variation and adaptive introgression were significantly shorter than those from
standing genetic variation along (Fig. S14, Table S17). One potential explanation for these
contrasting findings for accessible paths between generalists and scale-eaters is that because the

measure of fitness used to obtain these results, survival, is a binary, the fitness landscape
essentially converges on that of a holey adaptive landscape (2, 3, 19). Thus, adaptation between
generalists and scale eaters, which are particularly isolated on these fitness landscapes, may
occur through predominantly neutral evolution, erasing some of the signal detected for other
measures of fitness.

1087
1088 **Figure S1. Proportion (%) genetic variance explained by the first 20 Principal Components**
1089 **obtained using all SNPs and individuals from Crescent Pond, Little Lake, and Osprey Lake, as well**
1090 **as experimental hybrids**. The first two principal component axes are plotted in Figure 1.
1091

1092
1093    **Figure S2. Principal components two, three, and four.**

1094
**Figure S3. The proportion of specialist ancestry in hybrids did not predict in experimental hybrids**
**using either A) composite fitness, B) survival or C) growth.** Results of a generalized in which survival
(modeled as a binomial) is predicted by ancestry proportion, including lake and experiment as fixed
effects. Points represent individual hybrids, with each individual represented by two points, one
indicating their respective scale-eater (salmon, and molluscivore (blue) ancestry proportions. *P*-values
correspond to the effect of each type of ancestry (scale-eater – red, molluscivore – blue) on survival
probability. Lines are predicted values and are colored according to species.

1102
1103      **Figure S4. Genetic distance predicts morphological distance among sampled hybrids.**

1104
1105 **Figure S5. Manhattan plots illustrating the strength of association between individual SNPs and either Survival (A) or Growth (B) as**
1106 **inferred by GEMMA.** Significant associations are highlighted in blue (FDR < 0.05) or red (Bonferroni correction P < 0.05). The dashed red line
1107 indicates the threshold for significance following Bonferroni p-value adjustment. METTL21E and CSAD are both highly differentiated ($F_{ST}$ >
1108 0.95) among specialists, differentially expressed. METTL21E is also misexpressed in F1 hybrids, meaning it exhibits gene expression that is
1109 higher or lower than observed in both parental species (69), indicating it is involved in intrinsic incompatibilies.

1110

**Figure S6. Gene ontology enrichment for SNPs found to be associated with growth.** Darker colors indicate ontologies that are more significantly enriched following FDR correction. Length of bar is proportional to the number of genes assigned to an ontology.

1113

**Figure S7. Gene ontology enrichment for SNPs found to be associated with composite fitness.** Darker colors indicate ontologies that are more significantly enriched following FDR correction. Length of bar is proportional to the number of genes assigned to an ontology,

1116

**Figure S8. The 29 landmarks used to digitally measure thirty traits plus standard length using DLTDV8a** (110)**. The corresponding traits are shown in Table S7.

1119

**Figure S9. Morphological variation in the three San Salvador Island pupfish species and their experimentally produced hybrids. A.** Linear discriminant (LD) morphospace. Parent species and hybrids are plotted using the two LD axes that together serve to distinguish species with 99.4% accuracy. **B.** Within-group disparity calculated as the median distance between each individual and the groups centroid. Small, semitransparent points are the result of 100 bootstrap replicates, and are summarized by box plots, which in turn show the median and interquartile ranges of these bootstrap replicates. Large, opaque points are the observed disparities using the full dataset per group. All pairwise comparisons using t-tests following correction for the false discovery rate were significant ($P < 0.001$).

**Figure S10. Best-fit fitness landscapes for composite fitness without fitness associated SNPs (A) survival (B), growth without associated SNPs (C) and growth including associated SNPs (D).** Colored points indicate locations of parent species in LD morphospace, and ellipses indicate their 50% (solid) and 95% (dotted) confidence intervals. Grey points indicate location of hybrid individuals, with size proportional to their fitness measure. Cooler colors on the adaptive landscape indicate lower predicted fitness.

1136
1137
**Figure S11. The topography of the composite fitness adaptive landscape is influenced by the**
**distribution of a common SNP haplotype.** Shown is the same landscape as in Figure 2, but the
plotted points are unique SNP haplotypes for the thirteen most strongly fitness-associated SNPs.
One haplotype is particularly frequent among hybrids; individuals with this haplotype are closer
in morphospace to generalists and drive the emergence of a local fitness optimum for generalists.
All other haplotypes (points) are plotted with a distinct color per haplotype.

1144
**Figure S12. Genotypic fitness networks (using composite fitness or survival) are smaller both with respect to number of nodes (A) and**
**number of edges (B) when SNPs sourced from introgression or *de novo* mutation are included than networks constructed from standing**
**genetic variation (SGV) alone.** Compact letter display reports pairwise significance according to a nonparametric Kruskal-Wallis one-way analysis
of variance following correction for multiple tests using an estimated False Discovery Rate. Letters are ordered (i.e. **a** to **c**) in decreasing values of
the median. These results are constant whether survival or composite fitness are used because they comprise the same set of samples (both survivors
and non-survivors)

**Figure S13. Genotypic fitness networks (using growth) are smaller both with respect to number of nodes (A) and number of edges (B) when SNPs sourced from introgression or *de novo* mutation are included than networks constructed from standing genetic variation (SGV) alone.** Compact letter display reports pairwise significance according to a nonparametric Kruskal-Wallis one-way analysis of variance follwing correction for multiple tests using an estimated False Discovery Rate. Letters are ordered (i.e. **a** to **c**) in decreasing values of the median. These results include only survivors and thus differ from those shown in Figure S12.

1157

1158
1159 **Figure S14. Accessibility of between-species genotypic trajectories in survival fitness landscapes constructed from SNPs exhibiting**
1160 **signatures of divergent selection (Richards et al. 2021) that are sourced from either standing genetic variation (SGV), introgression,** *de novo*
1161 **mutation, or a combination. A)** The number of accessible (i.e. monotonically increasing in fitness) paths per network, scaled by the size of the
1162 network. Compact letter display reports pairwise significance according to a nonparametric Kruskal-Wallis one-way analysis of variance following
1163 correction for multiple tests using an estimated False Discovery Rate. **B)** Number of mutational steps in the shortest accessible path.

1164
1165 **Figure S15. Accessibility of between-species genotypic trajectories in growth fitness landscapes constructed from SNPs exhibiting signatures**
1166 **of divergent selection (Richards et al. 2021) that are sourced from either standing genetic variation (SGV), introgression, *de novo* mutation,**
1167 **or a combination. A)** The number of accessible (i.e. monotonically increasing in fitness) paths per network, scaled by the size of the network.
1168 Compact letter display reports pairwise significance according to a nonparametric Kruskal-Wallis one-way analysis of variance following correction
1169 for multiple tests using an estimated False Discovery Rate. **B)** Number of mutational steps in the shortest accessible path.

1170
1171 **Figure S16. The number of edges and thus potential number of paths (A) and accessible paths (B) increases as network size (number of**
1172 **nodes) increases.**

1173 **Table S1. Samples of hybrids and parental studies used either in genomic or morphological**
1174 **analyses, along with associated metadata.**
1175

| ID | Sequenced | Morphology | Survivorship | Experiment | Lake | Species |
|----|-----------|------------|--------------|------------|------|---------|
| CP04E02 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP07F05 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP08H06 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP09C02 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP09D01 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP09F10 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP10B05 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP11C10 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP13E03 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP13F01 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| LL01F03 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL01F05 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL01G05 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL02A08 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL02D09 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL02D11 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL02E04 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL02E07 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL02E09 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL04E03 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL04E09 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL04F02 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL04F07 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL05A01 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL05C06 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL05C09 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL05C10 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL05D10 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL05E06 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL05E08 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL05H10 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL06A10 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL06B03 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL06C10 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL06D12 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL06E10 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL06F08 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL08B09 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL08B11 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL08G07 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL08G10 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL08H01 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL09A09 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL09A10 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL09B05 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL09C03 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL09D01 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL09D10 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL09E01 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL09F03 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL09H07 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL09H10 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL10A08 | This study | Yes | Non-Survivor | Martin & Gould 2020 | Little Lake | Hybrid |

| CP02D01 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
|---------|-----------|-----|----------|---------------------|---------------|--------|
| CP02E09 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP03D02 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP03G03 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP03G07 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP05B08 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP05E02 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP05G12 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP05H10 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP06E09 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP06G01 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP06H09 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP07E08 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP07H01 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP07H11 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP08F07 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP08G08 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP09D02 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP10A03 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP10C04 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP10C07 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP10C12 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP10F12 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP10H01 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP11B03 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP11C08 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP11D04 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP11G06 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP11G12 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP12E05 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP12E07 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP12H04 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP13B02 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP13B08 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP14C04 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP15A10 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP15B08 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP15E01 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP15E03 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP17A01 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP18B11 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP19B02 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP19C02 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP19C07 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP19F01 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CP19G03 | This study | Yes | Survivor | Martin & Gould 2020 | Crescent Pond | Hybrid |
| CPH01 | This study | Yes | Survivor | Martin & Wainwright 2013 | Crescent Pond | Hybrid |
| CPH02 | This study | Yes | Survivor | Martin & Wainwright 2013 | Crescent Pond | Hybrid |
| CPH03 | This study | Yes | Survivor | Martin & Wainwright 2013 | Crescent Pond | Hybrid |
| CPH04 | This study | Yes | Survivor | Martin & Wainwright 2013 | Crescent Pond | Hybrid |
| CPH05 | This study | Yes | Survivor | Martin & Wainwright 2013 | Crescent Pond | Hybrid |
| CPH07 | This study | Yes | Survivor | Martin & Wainwright 2013 | Crescent Pond | Hybrid |
| CPH08 | This study | Yes | Survivor | Martin & Wainwright 2013 | Crescent Pond | Hybrid |
| CPH09 | This study | Yes | Survivor | Martin & Wainwright 2013 | Crescent Pond | Hybrid |
| CPH10 | This study | Yes | Survivor | Martin & Wainwright 2013 | Crescent Pond | Hybrid |
| CPH100 | This study | Yes | Survivor | Martin & Wainwright 2013 | Crescent Pond | Hybrid |
| CPH11 | This study | Yes | Survivor | Martin & Wainwright 2013 | Crescent Pond | Hybrid |

| CPH123 | This study | Yes | Survivor | Martin & Wainwright 2013 | Crescent Pond | Hybrid |
|---|---|---|---|---|---|---|
| LL01H04 | This study | Yes | Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL02G09 | This study | Yes | Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL06B06 | This study | Yes | Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL06E04 | This study | Yes | Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL07E12 | This study | Yes | Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL07G04 | This study | Yes | Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL08A09 | This study | Yes | Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL08B04 | This study | Yes | Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL08D05 | This study | Yes | Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL10F08 | This study | Yes | Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL12B04 | This study | Yes | Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL13B06 | This study | Yes | Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL13D04 | This study | Yes | Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL17H03 | This study | Yes | Survivor | Martin & Gould 2020 | Little Lake | Hybrid |
| LL106 | This study | Yes | Survivor | Martin & Wainwright 2013 | Little Lake | Hybrid |
| LL114 | This study | Yes | Survivor | Martin & Wainwright 2013 | Little Lake | Hybrid |
| LL124 | This study | Yes | Survivor | Martin & Wainwright 2013 | Little Lake | Hybrid |
| LL129 | This study | Yes | Survivor | Martin & Wainwright 2013 | Little Lake | Hybrid |
| LL175 | This study | Yes | Survivor | Martin & Wainwright 2013 | Little Lake | Hybrid |
| LL23 | This study | Yes | Survivor | Martin & Wainwright 2013 | Little Lake | Hybrid |
| LL247 | This study | Yes | Survivor | Martin & Wainwright 2013 | Little Lake | Hybrid |
| LL251 | This study | Yes | Survivor | Martin & Wainwright 2013 | Little Lake | Hybrid |
| LL271 | This study | Yes | Survivor | Martin & Wainwright 2013 | Little Lake | Hybrid |
| LLH12 | This study | Yes | Survivor | Martin & Wainwright 2013 | Little Lake | Hybrid |
| LLH34 | This study | Yes | Survivor | Martin & Wainwright 2013 | Little Lake | Hybrid |
| LLH41 | This study | Yes | Survivor | Martin & Wainwright 2013 | Little Lake | Hybrid |
| LLH51 | This study | Yes | Survivor | Martin & Wainwright 2013 | Little Lake | Hybrid |
| LLH94 | This study | Yes | Survivor | Martin & Wainwright 2013 | Little Lake | Hybrid |
| CRPA1 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Generalist |
| CRPA1000 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Generalist |
| CRPA1001 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Generalist |
| CRPA1003 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Generalist |
| CRPA3 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Generalist |
| LILA1 | Richards et al. 2021 | No | NA | NA | Little Lake | Generalist |
| OSPA1 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Generalist |
| OSPA1000 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Generalist |
| OSPA1001 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Generalist |
| OSPA11 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Generalist |
| OSPA12 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Generalist |
| OSPA13 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Generalist |
| OSPA4 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Generalist |
| OSPA5 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Generalist |
| OSPA6 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Generalist |
| OSPA8 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Generalist |
| OSPA9 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Generalist |
| CRPM1 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Molluscivore |
| CRPM10 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Molluscivore |
| CRPM1000 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Molluscivore |
| CRPM1001 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Molluscivore |
| CRPM11 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Molluscivore |
| CRPM2 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Molluscivore |
| CRPM3 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Molluscivore |
| CRPM5 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Molluscivore |
| CRPM6 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Molluscivore |
| CRPM7 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Molluscivore |
| CRPM8 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Molluscivore |

| | | | | | | |
|---|---|---|---|---|---|---|
| CRPM9 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Molluscivore |
| LILM-QTL | Richards et al. 2021 | No | NA | NA | Little Lake | Molluscivore |
| LILM3 | Richards et al. 2021 | No | NA | NA | Little Lake | Molluscivore |
| LILM4 | Richards et al. 2021 | No | NA | NA | Little Lake | Molluscivore |
| LILM5 | Richards et al. 2021 | No | NA | NA | Little Lake | Molluscivore |
| OSPM1 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Molluscivore |
| OSPM1000 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Molluscivore |
| OSPM1001 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Molluscivore |
| OSPM11 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Molluscivore |
| OSPM2 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Molluscivore |
| OSPM3 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Molluscivore |
| OSPM4 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Molluscivore |
| OSPM5 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Molluscivore |
| OSPM7 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Molluscivore |
| OSPM8 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Molluscivore |
| OSPM9 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Molluscivore |
| CRPP-QTL | Richards et al. 2021 | No | NA | NA | Crescent Pond | Scale-eater |
| CRPP1000 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Scale-eater |
| CRPP1001 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Scale-eater |
| CRPP2 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Scale-eater |
| CRPP3 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Scale-eater |
| CRPP4 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Scale-eater |
| CRPP5 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Scale-eater |
| CRPP7 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Scale-eater |
| CRPP8 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Scale-eater |
| CRPP9 | Richards et al. 2021 | No | NA | NA | Crescent Pond | Scale-eater |
| LILP-QTL | Richards et al. 2021 | No | NA | NA | Little Lake | Scale-eater |
| LILP3 | Richards et al. 2021 | No | NA | NA | Little Lake | Scale-eater |
| LILP4 | Richards et al. 2021 | No | NA | NA | Little Lake | Scale-eater |
| LILP5 | Richards et al. 2021 | No | NA | NA | Little Lake | Scale-eater |
| OSPP1 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Scale-eater |
| OSPP10 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Scale-eater |
| OSPP1000 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Scale-eater |
| OSPP1001 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Scale-eater |
| OSPP11 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Scale-eater |
| OSPP2 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Scale-eater |
| OSPP3 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Scale-eater |
| OSPP4 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Scale-eater |
| OSPP5 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Scale-eater |
| OSPP7 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Scale-eater |
| OSPP9 | Richards et al. 2021 | No | NA | NA | Osprey Pond (Little Lake) | Scale-eater |
| CPA01 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA03 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA05 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA07 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA09 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA11 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA13 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA15 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA17 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA19 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA21 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA23 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA25 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA27 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA29 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA31 | NA | Yes | Parental | NA | Crescent Pond | Generalist |

| | | | | | | |
|---|---|---|---|---|---|---|
| CPA33 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA35 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA37 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA39 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA41 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA43 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA45 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA47 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA49 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA51 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA53 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA55 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA57 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| CPA59 | NA | Yes | Parental | NA | Crescent Pond | Generalist |
| LLA20 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA21 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA22 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA23 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA24 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA25 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA26 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA27 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA28 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA29 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA30 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA31 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA32 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA33 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA34 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA35 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA36 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA37 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA38 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA39 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA40 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA41 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA42 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA43 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA44 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA45 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA46 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA47 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA48 | NA | Yes | Parental | NA | Little Lake | Generalist |
| LLA49 | NA | Yes | Parental | NA | Little Lake | Generalist |
| CPM01 | NA | Yes | Parental | NA | Crescent Pond | Molluscivore |
| CPM02 | NA | Yes | Parental | NA | Crescent Pond | Molluscivore |
| CPM03 | NA | Yes | Parental | NA | Crescent Pond | Molluscivore |
| CPM04 | NA | Yes | Parental | NA | Crescent Pond | Molluscivore |
| CPM05 | NA | Yes | Parental | NA | Crescent Pond | Molluscivore |
| CPM06 | NA | Yes | Parental | NA | Crescent Pond | Molluscivore |
| CPM07 | NA | Yes | Parental | NA | Crescent Pond | Molluscivore |
| CPM08 | NA | Yes | Parental | NA | Crescent Pond | Molluscivore |
| CPM09 | NA | Yes | Parental | NA | Crescent Pond | Molluscivore |
| CPM10 | NA | Yes | Parental | NA | Crescent Pond | Molluscivore |
| CPM11 | NA | Yes | Parental | NA | Crescent Pond | Molluscivore |
| CPM12 | NA | Yes | Parental | NA | Crescent Pond | Molluscivore |
| CPM13 | NA | Yes | Parental | NA | Crescent Pond | Molluscivore |

| CPM14 | NA | Yes | Parental | NA | Crescent Pond | Molluscivore |
|---|---|---|---|---|---|---|
| CPM15 | NA | Yes | Parental | NA | Crescent Pond | Molluscivore |
| CPM16 | NA | Yes | Parental | NA | Crescent Pond | Molluscivore |
| CPM17 | NA | Yes | Parental | NA | Crescent Pond | Molluscivore |
| CPM18 | NA | Yes | Parental | NA | Crescent Pond | Molluscivore |
| CPM19 | NA | Yes | Parental | NA | Crescent Pond | Molluscivore |
| CPM20 | NA | Yes | Parental | NA | Crescent Pond | Molluscivore |
| LLM01 | NA | Yes | Parental | NA | Little Lake | Molluscivore |
| LLM02 | NA | Yes | Parental | NA | Little Lake | Molluscivore |
| LLM03 | NA | Yes | Parental | NA | Little Lake | Molluscivore |
| LLM04 | NA | Yes | Parental | NA | Little Lake | Molluscivore |
| LLM05 | NA | Yes | Parental | NA | Little Lake | Molluscivore |
| LLM06 | NA | Yes | Parental | NA | Little Lake | Molluscivore |
| LLM07 | NA | Yes | Parental | NA | Little Lake | Molluscivore |
| LLM08 | NA | Yes | Parental | NA | Little Lake | Molluscivore |
| LLM09 | NA | Yes | Parental | NA | Little Lake | Molluscivore |
| LLM10 | NA | Yes | Parental | NA | Little Lake | Molluscivore |
| LLM11 | NA | Yes | Parental | NA | Little Lake | Molluscivore |
| LLM12 | NA | Yes | Parental | NA | Little Lake | Molluscivore |
| LLM13 | NA | Yes | Parental | NA | Little Lake | Molluscivore |
| LLM14 | NA | Yes | Parental | NA | Little Lake | Molluscivore |
| LLM15 | NA | Yes | Parental | NA | Little Lake | Molluscivore |
| LLM16 | NA | Yes | Parental | NA | Little Lake | Molluscivore |
| LLM17 | NA | Yes | Parental | NA | Little Lake | Molluscivore |
| LLM18 | NA | Yes | Parental | NA | Little Lake | Molluscivore |
| CPP01 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP02 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP03 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP04 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP05 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP06 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP07 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP08 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP09 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP10 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP11 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP12 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP13 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP14 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP15 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP16 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP17 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP18 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP19 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP20 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP21 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP22 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP23 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP24 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP25 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP26 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP27 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP28 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP29 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| CPP30 | NA | Yes | Parental | NA | Crescent Pond | Scale-eater |
| LLP01 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP02 | NA | Yes | Parental | NA | Little Lake | Scale-eater |

| LLP03 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
|---|---|---|---|---|---|---|
| LLP04 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP05 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP06 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP07 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP08 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP09 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP10 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP11 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP12 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP13 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP14 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP15 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP16 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP17 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP18 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP19 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP20 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP21 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP22 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP23 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP24 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP25 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP26 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP27 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP28 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP29 | NA | Yes | Parental | NA | Little Lake | Scale-eater |
| LLP30 | NA | Yes | Parental | NA | Little Lake | Scale-eater |

1176

1177 **Table S2. Models tested to assess the extent to which specialist ancestry are genome-wide**
1178 **variation (PC1/PC2) predicts measures of fitness and their respective fits.** Best-fit models are
1179 bolded

| Fitness Measure | Species | Model | AIC | Ancestry/PC *P*-value |
|---|---|---|---|---|
| **Composite Fitness** | Molluscivore | **Composite ~ AncestryProp** | **77.69** | **0.5951** |
| | | Composite ~ AncestryProp + Lake | 78.19 | 0.4048 |
| | | Composite ~ AncestryProp + Experiment | 78.28 | 0.4435 |
| | | Composite ~ AncestryProp + Lake + Experiment | 78.67 | 0.2845 |
| | | Composite ~ AncestryProp + Lake * Experiment | 80.62 | 0.2797 |
| | Scale-eater | **Composite ~ AncestryProp** | **76.01** | **0.1641** |
| | | Composite ~ AncestryProp + Lake | 77.41 | 0.2283 |
| | | Composite ~ AncestryProp + Experiment | 77.78 | 0.3000 |
| | | Composite ~ AncestryProp + Lake + Experiment | 79.12 | 0.4013 |
| | | Composite ~ AncestryProp + Lake * Experiment | 81.12 | 0.4079 |
| | PC1 | **Composite ~ PC1** | **76.32** | **0.2017** |
| | | Composite ~ PC1 + Lake | 77.04 | 0.1778 |
| | | Composite ~ PC1 + Experiment | 77.43 | 0.2341 |
| | | Composite ~ PC1 + Lake + Experiment | 78.64 | 0.2786 |
| | | Composite ~ PC1 + Lake * Experiment | 80.39 | 0.2389 |
| | PC2 | **Composite ~ PC2** | **76.32** | **0.2740** |
| | | Composite ~ PC2 + Lake | 76.32 | 0.1124 |
| | | Composite ~ PC2 + Experiment | 78.15 | 0.3985 |
| | | Composite ~ PC2 + Lake + Experiment | 77.99 | 0.1793 |
| | | Composite ~ PC2 + Lake * Experiment | 79.96 | 0.1797 |
| **Growth** | Molluscivore | Growth ~ AncestryProp | 42.52 | 0.4231 |
| | | Growth ~ AncestryProp + Lake | 29.24 | 0.3383 |
| | | Growth ~ AncestryProp + Experiment | 13.91 | 0.0041 |
| | | Growth ~ AncestryProp + Lake + Experiment | -23.57 | 0.3932 |
| | | **Growth ~ AncestryProp + Lake * Experiment** | **-88.32** | **0.3280** |
| | Scale-eater | Growth ~ AncestryProp | 36.71 | 0.0122 |
| | | Growth ~ AncestryProp + Lake | 12.25 | 0.0000 |
| | | Growth ~ AncestryProp + Experiment | 21.98 | 0.4870 |
| | | Growth ~ AncestryProp + Lake + Experiment | -22.80 | 0.9950 |
| | | **Growth ~ AncestryProp + Lake * Experiment** | **-88.53** | **0.2825** |
| | PC1 | Growth ~ PC1 | 30.97 | 0.0006 |
| | | Growth ~ PC1 + Lake | 29.37 | 0.3723 |
| | | Growth ~ PC1 + Experiment | -32.37 | 0.0000 |
| | | Growth ~ PC1 + Lake + Experiment | -30.92 | 0.0055 |
| | | **Growth ~ PC1 + Lake * Experiment** | **-87.40** | **0.7638** |
| | PC2 | Growth ~ PC2 | 23.61 | 0.0002 |
| | | Growth ~ PC2 + Lake | 23.61 | 0.0119 |
| | | Growth ~ PC2 + Experiment | 14.04 | 0.0044 |
| | | Growth ~ PC2 + Lake + Experiment | -23.65 | 0.3677 |
| | | **Growth ~ PC2 + Lake * Experiment** | **-87.49** | **0.6724** |
| **Survival** | Molluscivore | Survival ~ AncestryProp | 188.78 | 0.9643 |
| | | Survival ~ AncestryProp + Lake | 155.11 | 0.0936 |
| | | Survival ~ AncestryProp + Experiment | 161.41 | 0.3709 |
| | | **Survival ~ AncestryProp + Lake + Experiment** | **123.06** | **0.3127** |
| | | Survival ~ AncestryProp + Lake * Experiment | 125.06 | 0.3127 |
| | Scale-eater | Survival ~ AncestryProp | 188.61 | 0.6756 |
| | | Survival ~ AncestryProp + Lake | 154.93 | 0.0852 |
| | | Survival ~ AncestryProp + Experiment | 158.99 | 0.0828 |
| | | **Survival ~ AncestryProp + Lake + Experiment** | **124.04** | **0.8041** |
| | | Survival ~ AncestryProp + Lake * Experiment | 126.04 | 0.8041 |
| | PC1 | **Survival ~ PC1** | **156.24** | **0.0000** |
| | | Survival ~ PC1 + Lake | 157.87 | 0.6943 |
| | | Survival ~ PC1 + Experiment | 121.52 | 0.0000 |
| | | Survival ~ PC1 + Lake + Experiment | 123.51 | 0.4386 |
| | | Survival ~ PC1 + Lake * Experiment | 125.51 | 0.4386 |
| | PC2 | **Survival ~ PC2** | **157.96** | **0.0268** |
| | | Survival ~ PC2 + Lake | 157.96 | 0.7988 |
| | | Survival ~ PC2 + Experiment | 160.72 | 0.2248 |
| | | Survival ~ PC2 + Lake + Experiment | 123.39 | 0.4047 |
| | | Survival ~ PC2 + Lake * Experiment | 125.39 | 0.4047 |

1180

1181  **Table S3. SNPs found to be strongly associated with composite fitness using SnpEff** (107)**.** SNPs that were identified as being strongly associated with both growth and composite fitness are
1182  italicized, and those that remain significant after a Bonferroni correction are bolded.

| Scaffold | Position | Significance | REF | ALT | Variant Type | Gene Identifier | Gene Card |
|---|---|---|---|---|---|---|---|
| HiC_scaffold_1 | 32071263 | FDR | A | C | intergenic | CBRO_00000660-CBRO_00000661 | Znf250-Ptgdr2 |
| **HiC_scaffold_1** | **43866598** | **Bonferroni** | **G** | **A** | **intergenic** | **CBRO_00000910-CBRO_00000911** | **PPM1K-OVCH2** |
| HiC_scaffold_1 | 43867614 | FDR | C | T | intergenic | CBRO_00000910-CBRO_00000911 | PPM1K-OVCH2 |
| HiC_scaffold_3 | 2658774 | FDR | T | A | intergenic | CBRO_00017856-CBRO_00017857 | KMT2E-Magi2 |
| HiC_scaffold_3 | 2658775 | FDR | T | C | intergenic | CBRO_00017856-CBRO_00017857 | KMT2E-Magi2 |
| HiC_scaffold_3 | 2658793 | FDR | C | A | intergenic | CBRO_00017856-CBRO_00017857 | KMT2E-Magi2 |
| HiC_scaffold_4 | 18899496 | FDR | T | C | intergenic | CBRO_00012427-CBRO_00012428 | UNKNOWN-edc4 |
| HiC_scaffold_5 | 18306419 | FDR | C | G | upstream; intergenic | CBRO_00001232; CBRO_00001231-CBRO_00001232 | xlrs1; PPEF2-xlrs1 |
| HiC_scaffold_5 | 18306428 | FDR | A | T | upstream; intergenic | CBRO_00001232; CBRO_00001231-CBRO_00001232 | xlrs1; PPEF2-xlrs1 |
| HiC_scaffold_5 | 18307019 | FDR | G | T | intronic | CBRO_00001232 | xlrs1 |
| HiC_scaffold_5 | 18307030 | FDR | G | A | intronic | CBRO_00001232 | xlrs1 |
| HiC_scaffold_5 | 18311696 | FDR | C | T | intronic | CBRO_00001232 | xlrs1 |
| HiC_scaffold_5 | 40475116 | FDR | T | A | intergenic | CBRO_00001627-CBRO_00001628 | SLC25A44-UBE2Q2 |
| HiC_scaffold_7 | 10290141 | FDR | T | C | downstream; intergenic | CBRO_00009717; CBRO_00009716-CBRO_00009717 | Nfkbie; SLC35B2-Nfkbie |
| HiC_scaffold_7 | 10290142 | FDR | G | C | downstream; intergenic | CBRO_00009717; CBRO_00009716-CBRO_00009717 | Nfkbie; SLC35B2-Nfkbie |
| HiC_scaffold_7 | 10290165 | FDR | A | G | downstream; intergenic | CBRO_00009717; CBRO_00009716-CBRO_00009717 | Nfkbie; SLC35B2-Nfkbie |
| HiC_scaffold_7 | 10290166 | FDR | T | C | downstream; intergenic | CBRO_00009717; CBRO_00009716-CBRO_00009717 | Nfkbie; SLC35B2-Nfkbie |
| HiC_scaffold_7 | 10290168 | FDR | T | C | downstream; intergenic | CBRO_00009717; CBRO_00009716-CBRO_00009717 | Nfkbie; SLC35B2-Nfkbie |
| HiC_scaffold_7 | 13815058 | FDR | A | G | synonymous; downstream | CBRO_00009805; CBRO_00009804 | Aloxe3; UNKNOWN |
| HiC_scaffold_7 | 13815326 | FDR | C | T | intronic | CBRO_00009805 | Aloxe3 |
| HiC_scaffold_7 | 15349830 | FDR | C | A | intergenic | CBRO_00009834-CBRO_00009835 | Adcy8-efr3b |
| HiC_scaffold_7 | 18378061 | FDR | T | A | intergenic | CBRO_00009902-CBRO_00009903 | Fam84a-DDX1 |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| **HiC_scaffold_8** | **20263964** | **Bonferroni** | **G** | **A** | **upstream; intergenic** | **CBRO_00010647; CBRO_00010647-CBRO_00010648** | **Srcin1; Srcin1-Srcin1** |
| HiC_scaffold_8 | 31539262 | FDR | G | A | upstream; intergenic | CBRO_00010835; CBRO_00010834-CBRO_00010835 | Gjd3; Gjd3-Gjd3 |
| HiC_scaffold_11 | 3106766 | FDR | T | A | intronic | CBRO_00013361 | prkdc |
| HiC_scaffold_11 | 3138733 | FDR | G | A | intergenic | CBRO_00013361-CBRO_00013362 | prkdc-arhgap29 |
| HiC_scaffold_11 | 5658921 | FDR | A | G | intronic | CBRO_00013404 | KAZN |
| HiC_scaffold_14 | 17556007 | FDR | C | T | intergenic | CBRO_00014134-CBRO_00014135 | UNKNOWN-Abr |
| HiC_scaffold_14 | 17556026 | FDR | C | A | intergenic | CBRO_00014134-CBRO_00014135 | UNKNOWN-Abr |
| HiC_scaffold_16 | 32837191 | FDR | T | C | intronic | CBRO_00003226 | KIF1B |
| HiC_scaffold_16 | 35727503 | FDR | G | C | intergenic | CBRO_00003289-CBRO_00003290 | Pip5k1c-Polr2e |
| HiC_scaffold_16 | 40215889 | FDR | G | A | intergenic | CBRO_00003383-CBRO_00003384 | chst10-UNKNOWN |
| HiC_scaffold_16 | 40300592 | FDR | C | A | intergenic | CBRO_00003384-CBRO_00003385 | UNKNOWN-Carmil3 |
| HiC_scaffold_18 | 26969972 | FDR | T | G | intronic | CBRO_00013239 | CSAD |
| HiC_scaffold_18 | 26970123 | FDR | C | T | synonymous | CBRO_00013239 | CSAD |
| HiC_scaffold_18 | 26970601 | FDR | T | A | intronic | CBRO_00013239 | CSAD |
| HiC_scaffold_18 | 26978410 | FDR | G | A | intronic | CBRO_00013240 | Znf740 |
| HiC_scaffold_20 | 332642 | FDR | G | A | missense | CBRO_00016084 | GTF3C4 |
| HiC_scaffold_20 | 332689 | FDR | A | G | missense | CBRO_00016084 | GTF3C4 |
| HiC_scaffold_20 | 14820437 | FDR | A | T | intronic | CBRO_00016304 | MALT1 |
| HiC_scaffold_20 | 14820448 | FDR | T | C | intronic | CBRO_00016304 | MALT1 |
| HiC_scaffold_24 | 1469530 | FDR | G | T | intergenic | CBRO_00014375-CBRO_00014376 | UNKNOWN-UNKNOWN |
| HiC_scaffold_24 | 3223583 | FDR | T | A | intergenic | CBRO_00014421-CBRO_00014422 | Gal3st3-RIN2 |
| HiC_scaffold_24 | 11618442 | FDR | T | G | intronic | CBRO_00014601 | ABCA4 |
| **HiC_scaffold_24** | **15964553** | **Bonferroni** | **C** | **T** | **intergenic** | **CBRO_00014635-CBRO_00014636** | **Lrfn2-SNX15** |
| HiC_scaffold_27 | 1898180 | FDR | G | A | downstream; intergenic | CBRO_00005887; CBRO_00005886-CBRO_00005887 | UNKNOWN; hoxb13a-UNKNOWN |
| HiC_scaffold_27 | 8137335 | FDR | G | T | intronic | CBRO_00005998 | SMARCA4 |
| HiC_scaffold_27 | 9065056 | FDR | C | A | intergenic | CBRO_00006026-CBRO_00006027 | ANKFN1-ccdc134 |
| HiC_scaffold_27 | 12370585 | FDR | C | T | intergenic | CBRO_00006131-CBRO_00006132 | SHISA9-Desi1 |
| HiC_scaffold_27 | 32078665 | FDR | C | G | intergenic | CBRO_00006685-CBRO_00006686 | med25-Lrrc4b |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| HiC_scaffold_27 | 34904388 | FDR | A | C | intergenic | CBRO_00006756-CBRO_00006757 | Grin2c-nog3 |
| HiC_scaffold_27 | 35431570 | FDR | G | C | intergenic | CBRO_00006756-CBRO_00006757 | Grin2c-nog3 |
| HiC_scaffold_27 | 35431578 | FDR | C | T | intergenic | CBRO_00006756-CBRO_00006757 | Grin2c-nog3 |
| HiC_scaffold_27 | 35431585 | FDR | A | G | intergenic | CBRO_00006756-CBRO_00006757 | Grin2c-nog3 |
| HiC_scaffold_34 | 16654675 | FDR | G | C | intergenic | CBRO_00001997-CBRO_00001998 | MDFIC2-foxp1b |
| HiC_scaffold_34 | 19393244 | FDR | A | T | upstream; intergenic | CBRO_00002027; CBRO_00002027-CBRO_00002028 | SUOX; SUOX-SUOX |
| HiC_scaffold_34 | 22010499 | FDR | C | T | intronic | CBRO_00002117 | GNAI2 |
| HiC_scaffold_34 | 31220916 | FDR | T | A | upstream; intergenic | CBRO_00002389; CBRO_00002388-CBRO_00002389 | CTTNBP2NL; Kcnd3-CTTNBP2NL |
| HiC_scaffold_34 | 37769304 | FDR | A | C | intergenic | CBRO_00002519-CBRO_00002520 | ASIC2-asic1 |
| HiC_scaffold_37 | 5963405 | FDR | T | C | intergenic | CBRO_00011020-CBRO_00011021 | C14orf93-pim2 |
| HiC_scaffold_37 | 11017168 | FDR | G | A | intergenic | CBRO_00011155-CBRO_00011156 | GALNT12-elp2 |
| HiC_scaffold_37 | 13822135 | FDR | A | G | upstream; intergenic | CBRO_00011221; CBRO_00011220-CBRO_00011221 | SATB1; KCNH8-SATB1 |
| HiC_scaffold_37 | 13823678 | FDR | G | A | upstream; intergenic | CBRO_00011221; CBRO_00011220-CBRO_00011221 | SATB1; KCNH8-SATB1 |
| HiC_scaffold_37 | 13832007 | FDR | T | A | missense | CBRO_00011221 | SATB1 |
| HiC_scaffold_37 | 16920863 | FDR | T | C | intergenic | CBRO_00011259-CBRO_00011260 | CSMD1-UNKNOWN |
| **HiC_scaffold_37** | **18591438** | **Bonferroni** | **G** | **A** | **intergenic** | **CBRO_00011301-CBRO_00011302** | **cck-trim71** |
| HiC_scaffold_37 | 18591463 | FDR | C | T | intergenic | CBRO_00011301-CBRO_00011302 | cck-trim71 |
| HiC_scaffold_37 | 18596716 | FDR | A | T | intergenic | CBRO_00011301-CBRO_00011302 | cck-trim71 |
| HiC_scaffold_40 | 5885291 | FDR | G | T | downstream; intergenic | CBRO_00016614; CBRO_00016613-CBRO_00016614 | C14orf93 homolog; HTR2A-C14orf93 homolog |
| HiC_scaffold_43 | 2568535 | FDR | A | G | intergenic | CBRO_00008246-CBRO_00008247 | UNKNOWN-Gpr68 |
| HiC_scaffold_44 | 25886134 | FDR | A | C | intergenic | CBRO_00007276-CBRO_00007277 | RAPGEF2-QDPR |
| HiC_scaffold_45 | 885834 | FDR | A | C | upstream; intergenic | CBRO_00018766; CBRO_00018766-CBRO_00018767 | Mog; Mog-EPHB4 |
| HiC_scaffold_45 | 2213441 | FDR | G | T | intronic | CBRO_00018814 | Nlrp12 |
| HiC_scaffold_46 | 856248 | FDR | A | G | intergenic | CBRO_00007439-CBRO_00007440 | UNKNOWN-NLRP12 |
| HiC_scaffold_46 | 1232350 | FDR | G | A | intronic | CBRO_00007448 | NEB |
| HiC_scaffold_46 | 30183758 | FDR | C | G | intergenic | CBRO_00008059-CBRO_00008060 | TFRC-Rgs11 |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| HiC_scaffold_46 | 32048848 | FDR | A | C | intronic | CBRO_00008105 | Hsd17b7 |
| HiC_scaffold_46 | 32050109 | FDR | G | T | synonymous | CBRO_00008105 | Hsd17b7 |
| **HiC_scaffold_46** | **35151009** | **Bonferroni** | **T** | **C** | **downstream; intergenic** | **CBRO_00008165; CBRO_00008165-CBRO_00008166** | **Klf9; Klf9-Tsen15** |
| HiC_scaffold_46 | 35163681 | FDR | T | C | downstream; downstream; intergenic | CBRO_00008166; CBRO_00008167; CBRO_00008166-CBRO_00008167 | Tsen15; UNKNOWN; Tsen15-UNKNOWN |
| HiC_scaffold_46 | 35164267 | FDR | A | G | downstream; downstream; intergenic | CBRO_00008166; CBRO_00008167; CBRO_00008166-CBRO_00008167 | Tsen15; UNKNOWN; Tsen15-UNKNOWN |
| HiC_scaffold_47 | 787141 | FDR | T | G | upstream; intergenic | CBRO_00008838; CBRO_00008837-CBRO_00008838 | Nlrc3; NLRC3-Nlrc3 |
| HiC_scaffold_47 | 4300295 | FDR | G | A | downstream; intergenic | CBRO_00008933; CBRO_00008933-CBRO_00008934 | NEK6; NEK6-Psmb7 |
| HiC_scaffold_47 | 6767766 | FDR | T | C | intergenic | CBRO_00008991-CBRO_00008992 | TACR1-Grk5 |
| HiC_scaffold_52 | 20791197 | FDR | T | A | upstream; intronic | CBRO_00011892; CBRO_00011891 | rabl3; GTF2E1 |
| HiC_scaffold_52 | 22551701 | FDR | G | C | intergenic | CBRO_00011922-CBRO_00011923 | ALS2-Serp2 |
| HiC_scaffold_52 | 31021517 | FDR | C | T | intergenic | CBRO_00012051-CBRO_00012052 | Tmeff2-slc39a10 |
| **HiC_scaffold_53** | **11317840** | **Bonferroni** | **A** | **G** | **intergenic** | **CBRO_00005178-CBRO_00005179** | **Fucolectin-1-Fucolectin-5** |
| HiC_scaffold_53 | 11326035 | FDR | C | T | intergenic | CBRO_00005178-CBRO_00005179 | Fucolectin-1-Fucolectin-5 |
| HiC_scaffold_53 | 11326410 | FDR | G | A | intergenic | CBRO_00005178-CBRO_00005179 | Fucolectin-1-Fucolectin-5 |
| HiC_scaffold_53 | 11331079 | FDR | A | G | intergenic | CBRO_00005178-CBRO_00005179 | Fucolectin-1-Fucolectin-5 |
| HiC_scaffold_53 | 15966447 | FDR | G | A | upstream; downstream; intergenic | CBRO_00005235; CBRO_00005234; CBRO_00005234-CBRO_00005235 | Tmem222; WDTC1; WDTC1-Tmem222 |
| HiC_scaffold_53 | 17413090 | FDR | A | G | missense | CBRO_00005274 | Mag |
| HiC_scaffold_53 | 20715576 | FDR | C | A | intergenic | CBRO_00005380-CBRO_00005381 | Scrt2-Ino80c |
| HiC_scaffold_53 | 20715623 | FDR | G | A | intergenic | CBRO_00005380-CBRO_00005381 | Scrt2-Ino80c |
| HiC_scaffold_53 | 20715851 | FDR | A | G | intergenic | CBRO_00005380-CBRO_00005381 | Scrt2-Ino80c |
| HiC_scaffold_53 | 27386094 | FDR | G | T | intronic | CBRO_00005625 | Arhgef1 |
| HiC_scaffold_53 | 27396961 | FDR | T | A | intronic | CBRO_00005625 | Arhgef1 |
| HiC_scaffold_53 | 27398605 | FDR | T | G | downstream; intergenic | CBRO_00005625; CBRO_00005625-CBRO_00005626 | Arhgef1; Arhgef1-CD79A |
| HiC_scaffold_53 | 33282501 | FDR | A | G | intergenic | CBRO_00005732-CBRO_00005733 | mios-GLCCI1 |
| HiC_scaffold_53 | 39228193 | FDR | T | C | intergenic | CBRO_00005830-CBRO_00005831 | UNKNOWN-GnRHR2 |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| **HiC_scaffold_53** | **39228264** | **Bonferroni** | **C** | **T** | **intergenic** | **CBRO_00005830-CBRO_00005831** | **UNKNOWN-GnRHR2** |
| HiC_scaffold_53 | 39228942 | FDR | A | T | intergenic | CBRO_00005830-CBRO_00005831 | UNKNOWN-GnRHR2 |
| HiC_scaffold_53 | 39279263 | FDR | G | T | intronic | CBRO_00005832 | IGDCC3 |
| HiC_scaffold_53 | 39770914 | FDR | A | T | intronic | CBRO_00005849 | Cpne4 |
| HiC_scaffold_611 | 5621 | FDR | A | T | intergenic | CHR_START-CBRO_00020243 | CHR_START-fzdz-a |
| HiC_scaffold_611 | 5625 | FDR | G | T | intergenic | CHR_START-CBRO_00020243 | CHR_START-fzdz-a |
| HiC_scaffold_611 | 5634 | FDR | T | A | intergenic | CHR_START-CBRO_00020243 | CHR_START-fzdz-a |
| HiC_scaffold_1053 | 3494 | FDR | A | G | downstream; intergenic | CBRO_00020503; CHR_START-CBRO_00020503 | UBE2G2; CHR_START-UBE2G2 |
| HiC_scaffold_1133 | 9946 | FDR | A | G | intergenic | . | . |
| HiC_scaffold_1371 | 7318 | FDR | C | T | upstream; intergenic | CBRO_00021026; CHR_START-CBRO_00021026 | UNKNOWN; CHR_START-UNKNOWN |
| **HiC_scaffold_1848** | **40119** | **Bonferroni** | **C** | **A** | **intergenic** | **.** | **.** |
| **HiC_scaffold_1848** | **40465** | **Bonferroni** | **T** | **A** | **intergenic** | **.** | **.** |
| **HiC_scaffold_1848** | **40590** | **Bonferroni** | **T** | **C** | **intergenic** | **.** | **.** |
| HiC_scaffold_1848 | 40877 | FDR | T | C | intergenic | . | . |
| HiC_scaffold_1848 | 41351 | FDR | T | C | intergenic | . | . |
| HiC_scaffold_2220 | 10128 | FDR | G | T | intergenic | . | . |
| **HiC_scaffold_4461** | **12939** | **Bonferroni** | **T** | **A** | **intergenic** | **.** | **.** |
| HiC_scaffold_4665 | 13941 | FDR | C | T | intergenic | . | . |
| HiC_scaffold_6275 | 6000 | FDR | C | A | intergenic | . | . |
| HiC_scaffold_6337 | 5745 | FDR | T | G | intronic | CBRO_00021217 | PKP3 |
| HiC_scaffold_6769 | 2796 | FDR | G | A | intergenic | . | . |
| HiC_scaffold_6963 | 5970 | FDR | C | A | intergenic | . | . |
| HiC_scaffold_6963 | 6101 | FDR | A | G | intergenic | . | . |
| HiC_scaffold_9280 | 3448 | FDR | G | A | intergenic | . | . |
| HiC_scaffold_9949 | 52 | FDR | C | T | intergenic | . | . |
| HiC_scaffold_10928 | 3575 | FDR | G | A | downstream; intergenic | CBRO_00021896; CHR_START-CBRO_00021896 | CYP2A10; CHR_START-CYP2A10 |
| HiC_scaffold_11921 | 5560 | FDR | A | T | intergenic | . | . |
| HiC_scaffold_12068 | 2929 | FDR | G | T | intergenic | . | . |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| **HiC_scaffold_1277 8** | **1456** | **Bonferroni** | **T** | **A** | **missense** | **CBRO_00022026** | **COL8A1** |
| HiC_scaffold_17578 | 180 | FDR | T | G | intergenic | . | . |
| **HiC_scaffold_18999** | **1084** | **Bonferroni** | **A** | **G** | **intergenic** | **.** | **.** |

1183

1184 **Table S4. SNPs found to be strongly associated with growth SnpEff (107).** SNPs that were identified as being strongly associated with both growth and composite fitness are italicized, and those
1185 that remain significant after a Bonferroni correction are bolded.

| Scaffold | Position | Significance | REF | ALT | Variant Type | Gene Identifier | Gene Card |
|---|---|---|---|---|---|---|---|
| HiC_scaffold_4 | 8897671 | FDR | A | T | intergenic | CBRO_00012254-CBRO_00012255 | Megf10-UNKNOWN |
| HiC_scaffold_4 | 16057698 | FDR | C | A | intergenic | CBRO_00012386-CBRO_00012387 | CCND2-Mlycd |
| HiC_scaffold_4 | 29273446 | FDR | T | C | synonymous | CBRO_00012620 | CKAP5 |
| HiC_scaffold_4 | 29273458 | FDR | C | T | synonymous | CBRO_00012620 | CKAP5 |
| HiC_scaffold_5 | 18259775 | FDR | C | A | upstream; intergenic | CBRO_00001229; CBRO_00001229-CBRO_00001230 | AP1S2; AP1S2-phka2 |
| *HiC_scaffold_5* | *18306419* | *FDR* | *C* | *G* | *upstream; intergenic* | *CBRO_00001232; CBRO_00001231-CBRO_00001232* | *xlrs1; PPEF2-xlrs1* |
| *HiC_scaffold_5* | *18306428* | *FDR* | *A* | *T* | *upstream; intergenic* | *CBRO_00001232; CBRO_00001231-CBRO_00001232* | *xlrs1; PPEF2-xlrs1* |
| *HiC_scaffold_5* | *18307019* | *FDR* | *G* | *T* | *intronic* | *CBRO_00001232* | *xlrs1* |
| *HiC_scaffold_5* | *18307030* | *FDR* | *G* | *A* | *intronic* | *CBRO_00001232* | *xlrs1* |
| HiC_scaffold_5 | 36619253 | FDR | T | C | intergenic | CBRO_00001583-CBRO_00001584 | Chst12-ZDHHC13 |
| *HiC_scaffold_7* | *13815058* | *FDR* | *A* | *G* | *synonymous; downstream* | *CBRO_00009805; CBRO_00009804* | *Aloxe3; UNKNOWN* |
| HiC_scaffold_7 | 13823565 | FDR | T | C | intronic | CBRO_00009806 | Fbxo30 |
| HiC_scaffold_7 | 13824467 | FDR | A | G | missense | CBRO_00009806 | Fbxo30 |
| HiC_scaffold_7 | 19371997 | FDR | C | T | upstream; intergenic | CBRO_00009922; CBRO_00009921-CBRO_00009922 | ELOVL4; TENT5A-ELOVL4 |
| HiC_scaffold_7 | 19372002 | FDR | T | G | upstream; intergenic | CBRO_00009922; CBRO_00009921-CBRO_00009922 | ELOVL4; TENT5A-ELOVL4 |
| HiC_scaffold_8 | 20265076 | FDR | T | C | upstream; intergenic | CBRO_00010647; CBRO_00010647-CBRO_00010648 | Srcin1; Srcin1-Srcin1 |
| HiC_scaffold_8 | 20265098 | FDR | G | C | upstream; intergenic | CBRO_00010647; CBRO_00010647-CBRO_00010648 | Srcin1; Srcin1-Srcin1 |
| HiC_scaffold_8 | 20278571 | FDR | G | A | intergenic | CBRO_00010647-CBRO_00010648 | Srcin1-Srcin1 |
| HiC_scaffold_9 | 15585466 | FDR | C | G | missense | CBRO_00004552 | Tmem260 |
| HiC_scaffold_9 | 18453213 | FDR | A | G | intergenic | CBRO_00004639-CBRO_00004640 | Bub1b-PAK6 |
| HiC_scaffold_9 | 28127377 | FDR | C | T | intergenic | CBRO_00004857-CBRO_00004858 | METTL21E-RASA3 |
| HiC_scaffold_10 | 193812 | FDR | C | G | intergenic | CBRO_00018931-CBRO_00018932 | Spsb4-UNKNOWN |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| HiC_scaffold_11 | 25632195 | FDR | C | T | intergenic | CBRO_00013717-CBRO_00013718 | CDH10-Cdh6 |
| HiC_scaffold_11 | 25632258 | FDR | C | T | intergenic | CBRO_00013717-CBRO_00013718 | CDH10-Cdh6 |
| HiC_scaffold_11 | 25632641 | FDR | A | G | intergenic | CBRO_00013717-CBRO_00013718 | CDH10-Cdh6 |
| HiC_scaffold_14 | 14624430 | FDR | T | C | intergenic | CBRO_00014084-CBRO_00014085 | KDM6B-FGF11 |
| HiC_scaffold_18 | 26970449 | FDR | C | T | intronic | CBRO_00013239 | CSAD |
| *HiC_scaffold_27* | *8137335* | *FDR* | *G* | *T* | *intronic* | *CBRO_00005998* | *SMARCA4* |
| HiC_scaffold_27 | 21919164 | FDR | G | T | synonymous | CBRO_00006396 | SSTR2 |
| HiC_scaffold_29 | 2136546 | FDR | A | T | upstream; intergenic | CBRO_00016347; CBRO_00016347-CBRO_00016348 | RAPGEF6; RAPGEF6-ACSL6 |
| HiC_scaffold_29 | 2147361 | FDR | C | T | intergenic | CBRO_00016347-CBRO_00016348 | RAPGEF6-ACSL6 |
| HiC_scaffold_31 | 6226140 | FDR | A | C | intronic | CBRO_00017224 | ranbp9 |
| HiC_scaffold_34 | 7439419 | FDR | G | T | intergenic | CBRO_00001828-CBRO_00001829 | rnf152-CAMTA1 |
| HiC_scaffold_34 | 30571660 | FDR | C | A | intronic | CBRO_00002374 | SHMT2 |
| HiC_scaffold_40 | 4693917 | FDR | G | A | intronic | CBRO_00016589 | SLC37A1 |
| HiC_scaffold_40 | 4694015 | FDR | T | C | synonymous | CBRO_00016589 | SLC37A1 |
| HiC_scaffold_40 | 4723358 | FDR | A | G | intronic | CBRO_00016591 | UBXN4 |
| HiC_scaffold_40 | 4732538 | FDR | T | C | intronic | CBRO_00016591 | UBXN4 |
| HiC_scaffold_40 | 4734825 | FDR | G | T | intronic | CBRO_00016591 | UBXN4 |
| HiC_scaffold_40 | 4780518 | FDR | G | C | intronic | CBRO_00016592 | ITGB2 |
| HiC_scaffold_40 | 5016538 | FDR | C | A | upstream; intergenic | CBRO_00016599; CBRO_00016599-CBRO_00016600 | RRP1B; RRP1B-ITGB2 |
| HiC_scaffold_43 | 26869231 | FDR | C | G | synonymous | CBRO_00008677 | lrpprc |
| HiC_scaffold_43 | 27800569 | FDR | C | T | downstream; intergenic | CBRO_00008685; CBRO_00008685-CBRO_00008686 | timp3; timp3-ETV6 |
| HiC_scaffold_44 | 19344526 | FDR | C | T | intergenic | CBRO_00007178-CBRO_00007179 | ATP11C-sox3 |
| HiC_scaffold_46 | 16495639 | FDR | G | A | intronic | CBRO_00007743 | PHLPP1 |
| HiC_scaffold_46 | 16510323 | FDR | G | A | intronic | CBRO_00007743 | PHLPP1 |
| HiC_scaffold_46 | 16512668 | FDR | T | A | intronic | CBRO_00007743 | PHLPP1 |
| **HiC_scaffold_46** | **16512886** | **Bonferroni** | **T** | **A** | **intronic** | **CBRO_00007743** | **PHLPP1** |
| HiC_scaffold_46 | 16513809 | FDR | T | C | synonymous | CBRO_00007743 | PHLPP1 |
| HiC_scaffold_52 | 19031083 | FDR | T | A | intergenic | CBRO_00011872-CBRO_00011873 | NRP2-MREG |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| HiC_scaffold_52 | 19031216 | FDR | A | T | intergenic | CBRO_00011872-CBRO_00011873 | NRP2-MREG |
| *HiC_scaffold_1848* | *40119* | *Bonferroni* | *C* | *A* | *intergenic* | *.* | *.* |
| *HiC_scaffold_1848* | *40465* | *Bonferroni* | *T* | *A* | *intergenic* | *.* | *.* |
| *HiC_scaffold_1848* | *40590* | *FDR* | *T* | *C* | *intergenic* | *.* | *.* |
| *HiC_scaffold_1848* | *40877* | *FDR* | *T* | *C* | *intergenic* | *.* | *.* |
| *HiC_scaffold_1848* | *41351* | *FDR* | *T* | *C* | *intergenic* | *.* | *.* |
| **HiC_scaffold_7644** | **5971** | **Bonferroni** | **T** | **G** | **intergenic** | **.** | **.** |
| HiC_scaffold_12681 | 4019 | FDR | T | C | upstream; intergenic | CBRO_00022068; CBRO_00022068-CHR_END | UNKNOWN; UNKNOWN-CHR_END |

1186

1187 **Table S5. Gene ontology term enrichment for genes associated with growth.**

| Functional Category | Enrichment FDR | Genes in list | Total genes | Genes |
|---|---|---|---|---|
| Negative regulation of intracellular signal transduction | 0.0259 | 6 | 537 | PPEF2, RNF152, RASA3, RANBP9, PHLPP1, TIMP3 |
| Phosphorus metabolic process | 0.0259 | 17 | 3597 | PHLPP1, CCND2, PAK6, PPEF2, SHMT2, MLYCD, RRP1B, ITGB2, SRCIN1, PHKA2, ELOVL4, BUB1B, RANBP9, TIMP3, CAMTA1, ACSL6, RASA3 |
| Phosphate-containing compound metabolic process | 0.0259 | 17 | 3570 | PHLPP1, CCND2, PAK6, PPEF2, SHMT2, MLYCD, RRP1B, ITGB2, SRCIN1, PHKA2, ELOVL4, BUB1B, RANBP9, TIMP3, CAMTA1, ACSL6, RASA3 |
| Serine family amino acid catabolic process | 0.0259 | 2 | 17 | SHMT2, CSAD |
| Organic acid biosynthetic process | 0.0259 | 6 | 466 | ELOVL4, SHMT2, MLYCD, CHST12, ALOXE3, CSAD |
| Regulation of phosphate metabolic process | 0.0259 | 11 | 1870 | PHLPP1, CCND2, PAK6, PPEF2, SHMT2, RRP1B, ITGB2, SRCIN1, RANBP9, TIMP3, CAMTA1 |
| Nucleoside bisphosphate biosynthetic process | 0.0259 | 3 | 82 | MLYCD, ELOVL4, ACSL6 |
| Ribonucleoside bisphosphate biosynthetic process | 0.0259 | 3 | 82 | MLYCD, ELOVL4, ACSL6 |
| Purine nucleoside bisphosphate biosynthetic process | 0.0259 | 3 | 82 | MLYCD, ELOVL4, ACSL6 |
| Long-chain fatty-acyl-CoA biosynthetic process | 0.0259 | 2 | 19 | ELOVL4, ACSL6 |
| Thioester biosynthetic process | 0.0259 | 3 | 66 | MLYCD, ELOVL4, ACSL6 |
| Positive regulation by host of viral transcription | 0.0259 | 2 | 18 | SMARCA4, RRP1B |
| Sulfur compound biosynthetic process | 0.0259 | 5 | 207 | MLYCD, CHST12, ELOVL4, CSAD, ACSL6 |
| Regulation of phosphorus metabolic process | 0.0259 | 11 | 1872 | PHLPP1, CCND2, PAK6, PPEF2, SHMT2, RRP1B, ITGB2, SRCIN1, RANBP9, TIMP3, CAMTA1 |
| Acyl-CoA biosynthetic process | 0.0259 | 3 | 66 | MLYCD, ELOVL4, ACSL6 |
| Hypothalamus development | 0.0291 | 2 | 22 | NRP2, SOX3 |
| Sulfur compound metabolic process | 0.0312 | 5 | 393 | MLYCD, CHST12, ELOVL4, CSAD, ACSL6 |
| Long-chain fatty-acyl-CoA metabolic process | 0.0343 | 2 | 26 | ELOVL4, ACSL6 |
| Regulation of mitochondrial translation | 0.0343 | 2 | 26 | LRPPRC, SHMT2 |
| Limbic system development | 0.0405 | 3 | 115 | NRP2, KDM6B, SOX3 |
| Acyl-CoA metabolic process | 0.0406 | 3 | 119 | MLYCD, ELOVL4, ACSL6 |
| Thioester metabolic process | 0.0406 | 3 | 119 | MLYCD, ELOVL4, ACSL6 |
| Protein phosphorylation | 0.0412 | 11 | 2093 | PHLPP1, CCND2, PAK6, PPEF2, ITGB2, SRCIN1, PHKA2, BUB1B, RANBP9, TIMP3, RASA3 |
| Modulation by host of viral transcription | 0.0412 | 2 | 33 | SMARCA4, RRP1B |
| Carboxylic acid biosynthetic process | 0.0412 | 5 | 465 | ELOVL4, SHMT2, MLYCD, CHST12, ALOXE3 |
| Modulation of transcription in other organism involved in symbiotic interaction | 0.0412 | 2 | 34 | SMARCA4, RRP1B |
| Modulation by host of symbiont transcription | 0.0412 | 2 | 33 | SMARCA4, RRP1B |
| Cellular protein modification process | 0.0495 | 17 | 4434 | PHLPP1, CCND2, PAK6, PPEF2, RNF152, ITGB2, SPSB4, SRCIN1, PHKA2, FBXO30, KDM6B, BUB1B, RANBP9, TIMP3, CAMTA1, SHMT2, RASA3 |
| Protein modification process | 0.0495 | 17 | 4434 | PHLPP1, CCND2, PAK6, PPEF2, RNF152, ITGB2, SPSB4, SRCIN1, PHKA2, FBXO30, KDM6B, BUB1B, RANBP9, TIMP3, CAMTA1, SHMT2, RASA3 |

1188

1189 **Table S6. Gene ontology term enrichment for genes associated with composite fitness.**

| Functional Category | Enrichment FDR | Genes in list | Total genes | Genes |
|---|---|---|---|---|
| Anterograde trans-synaptic signaling | 0.0060 | 13 | 746 | GRIN2C, SHISA9, ALS2, KIF1B, ASIC1, GNAI2, TACR1, ADCY8, LRFN2, ABR, RAPGEF2, ASIC2, PIP5K1C |
| Synaptic signaling | 0.0060 | 13 | 762 | GRIN2C, SHISA9, ALS2, KIF1B, ASIC1, GNAI2, TACR1, ADCY8, LRFN2, ABR, RAPGEF2, ASIC2, PIP5K1C |
| Trans-synaptic signaling | 0.0060 | 13 | 756 | GRIN2C, SHISA9, ALS2, KIF1B, ASIC1, GNAI2, TACR1, ADCY8, LRFN2, ABR, RAPGEF2, ASIC2, PIP5K1C |
| Chemical synaptic transmission | 0.0060 | 13 | 746 | GRIN2C, SHISA9, ALS2, KIF1B, ASIC1, GNAI2, TACR1, ADCY8, LRFN2, ABR, RAPGEF2, ASIC2, PIP5K1C |
| Regulation of I-kappaB kinase/NF-kappaB signaling | 0.0179 | 7 | 241 | NEK6, SLC35B2, MALT1, NLRP12, NLRC3, DDX1, PIM2 |
| Regulation of trans-synaptic signaling | 0.0185 | 9 | 468 | GRIN2C, SHISA9, ASIC1, GNAI2, TACR1, ADCY8, LRFN2, ABR, RAPGEF2 |
| Regulation of cell communication | 0.0185 | 32 | 3903 | NEK6, SLC35B2, MALT1, NLRP12, PPEF2, GRIN2C, NLRC3, SLC39A10, SHISA9, ALS2, RAPGEF2, SMARCA4, MAGI2, RGS11, ARHGEF1, DDX1, PIM2, ASIC1, GNAI2, TACR1, ADCY8, LRFN2, ABR, PTGDR2, CCK, GPR68, MIOS, SCRT2, PSMB7, ARHGAP29, PIP5K1C, GRK5 |
| Regulation of signaling | 0.0185 | 32 | 3952 | NEK6, SLC35B2, MALT1, NLRP12, PPEF2, GRIN2C, NLRC3, SLC39A10, SHISA9, ALS2, RAPGEF2, SMARCA4, MAGI2, RGS11, ARHGEF1, DDX1, PIM2, ASIC1, GNAI2, TACR1, ADCY8, LRFN2, ABR, PTGDR2, CCK, GPR68, MIOS, SCRT2, PSMB7, ARHGAP29, PIP5K1C, GRK5 |
| Modulation of chemical synaptic transmission | 0.0185 | 9 | 467 | GRIN2C, SHISA9, ASIC1, GNAI2, TACR1, ADCY8, LRFN2, ABR, RAPGEF2 |
| Sensory perception of sour taste | 0.0185 | 2 | 5 | ASIC2, ASIC1 |
| Regulation of biological quality | 0.0185 | 34 | 4319 | GRIN2C, KCNH8, SLC39A10, SHISA9, PRKDC, TACR1, SMARCA4, KCND3, ABCA4, TRIM71, KMT2E, TFRC, PIM2, MAG, ASIC2, ASIC1, GNAI2, LRRC4B, HSD17B7, WDTC1, ADCY8, LRFN2, ABR, ALOXE3, CSMD1, CCK, SRCIN1, ALS2, GPR68, RAPGEF2, NEB, PSMB7, NFKBIE, PIP5K1C |
| I-kappaB kinase/NF-kappaB signaling | 0.0185 | 7 | 280 | NEK6, SLC35B2, MALT1, NLRP12, NLRC3, DDX1, PIM2 |
| Positive regulation of catalytic activity | 0.0185 | 17 | 1514 | PPM1K, ALS2, RAPGEF2, MAGI2, SRCIN1, RGS11, ARHGEF1, GNAI2, GTF3C4, RIN2, ARHGAP29, ABR, CCK, SLC39A10, MALT1, ADCY8, NLRP12 |
| Regulation of protein tyrosine phosphatase activity | 0.0216 | 2 | 6 | GNAI2, SLC39A10 |
| Multicellular organismal response to stress | 0.0216 | 4 | 74 | ALS2, ASIC1, TACR1, CCK |
| Positive regulation of molecular function | 0.0270 | 19 | 1901 | PPM1K, MALT1, ALS2, RAPGEF2, SMARCA4, MAGI2, SRCIN1, RGS11, ARHGEF1, GNAI2, GTF3C4, RIN2, ARHGAP29, ADCY8, ABR, CCK, SLC39A10, MED25, NLRP12 |
| Cell-cell signaling | 0.0311 | 18 | 1774 | GRIN2C, SHISA9, ALS2, KIF1B, ASIC1, GNAI2, TACR1, ADCY8, LRFN2, ABR, GRK5, GPR68, SMARCA4, RAPGEF2, MAGI2, ASIC2, PSMB7, PIP5K1C |
| Tachykinin receptor signaling pathway | 0.0314 | 2 | 8 | TACR1, GRK5 |
| Cell proliferation | 0.0355 | 20 | 2165 | TFRC, PIM2, RAPGEF2, CD79A, GNAI2, TACR1, COL8A1, MALT1, SATB1, PTGDR2, CCK, SLC39A10, TRIM71, PRKDC, MED25, KLF9, GRK5, NLRC3, MAGI2, ARHGEF1 |
| Behavioral fear response | 0.0355 | 3 | 39 | ALS2, ASIC1, CCK |
| Response to pH | 0.0355 | 3 | 41 | ASIC1, ASIC2, GPR68 |
| Behavioral defense response | 0.0355 | 3 | 40 | ALS2, ASIC1, CCK |
| Fear response | 0.0355 | 3 | 41 | ALS2, ASIC1, CCK |
| Positive regulation of hydrolase activity | 0.0401 | 11 | 833 | ALS2, RAPGEF2, MAGI2, RGS11, ARHGEF1, RIN2, ARHGAP29, ABR, CCK, SLC39A10, NLRP12 |
| NcRNA transcription | 0.0457 | 4 | 110 | GTF3C4, SMARCA4, POLR2E, GTF2E1 |
| Hexose mediated signaling | 0.0457 | 2 | 13 | SMARCA4, ADCY8 |
| Sugar mediated signaling pathway | 0.0457 | 2 | 13 | SMARCA4, ADCY8 |
| Glucose mediated signaling pathway | 0.0457 | 2 | 13 | SMARCA4, ADCY8 |
| Positive regulation of cell communication | 0.0457 | 18 | 1937 | NEK6, SLC35B2, MALT1, GRIN2C, SLC39A10, ALS2, RAPGEF2, SMARCA4, NLRP12, DDX1, PIM2, GNAI2, TACR1, ADCY8, PTGDR2, GPR68, MIOS, PSMB7 |
| Positive regulation of signaling | 0.0457 | 18 | 1945 | NEK6, SLC35B2, MALT1, GRIN2C, SLC39A10, ALS2, RAPGEF2, SMARCA4, NLRP12, DDX1, PIM2, GNAI2, TACR1, ADCY8, PTGDR2, GPR68, MIOS, PSMB7 |

1190

**Table S7. List of the 31 morphological traits measured for this study, and standard length; corresponding landmark ID's match those shown in Figure S8.**

| Trait Index | Trait Description | Trait Shorthand | Points |
|---|---|---|---|
| 1 | Nasal protrusion | nose | 3-4 |
| 2 | Nasal length | foresnout | 2-5 |
| 3 | Orbit to anal fin insertion | bellylen | 6-15 |
| 4 | Lateral facial length | snoutlen | 2-6 |
| 5 | Upper jaw to pectoral girdle | jaw2pect | 2-14 |
| 6 | Lateral skull length | pmx2add | 2-11 |
| 7 | Premaxilla length | pmxlen | 2-9 |
| 8 | Lower mandible length | jawlen | 1-9 |
| 9 | Jaw joint to orbit | foreeyewidth | 6-9 |
| 10 | Horizontal orbit diameter | eyewidth | 6-8 |
| 11 | Vertical orbit diameter | eyeht | 7-10 |
| 12 | Head height | headht | 7-9 |
| 13 | Suspensorium length | suspensorium | 9-11 |
| 14 | Adductor height | adductorht | 11-12 |
| 15 | Subopercle to pectoral girdle | ad2pect | 11-14 |
| 16 | Pectoral fin insertion width | pectinsertion | 13-14 |
| 17 | Anal to caudal distance | analtocaudal | 15-16 |
| 18 | Caudal peduncle height | caudalpedht | 16-18 |
| 19 | Dorsal to caudal distance | dorsaltocaudal | 18-19 |
| 20 | Body depth | bodydepth | 15-19 |
| 21 | Nasal protrusion angle | nasalangle | 7-5-3 |
| 22 | Premaxilla to orbit angle | topeyeangle | 7-2-10 |
| 23 | Premaxilla to adductor angle | lowereyeangle | 7-2-11 |
| 24 | Dorsal facial length | dorsalsnoutlen | 23-24, 25-26 |
| 25 | Adductor to premaxilla | eyetosnout | 21-24, 25-28 |
| 26 | Neurocranium to premaxilla | headlen | 24-20, 25-29 |
| 27 | Orbit to premaxilla | innereyetosnout | 22-24, 25-27 |
| 28 | Interorbital width | cranialwidth | 22-27 |
| 29 | Orbital neurocranium width | hindeyewidth | 21-28 |
| 30 | Max. neurocranium width | headwidth | 20-29 |
| 31 | Standard length (SL) | SL | 2-17 |

1232 **Table S8. Generalized additive models fitted to composite fitness.** Model fit was assessed using AICc, and Akaike Weights represent proportional
1233 model support. A thin plate spline for the two linear discriminant axes s(LD1, LD2) is always included, as is a fixed effect of either experiment (i.e.
1234 Martin & Wainwright 2013, Martin and Gould 2020) or Lake (Crescent Pond/Little Lake) or an interaction between the two. In the last two models,
1235 Experiment and Lake are included as splines, modeled using a factor smooth (bs = "fs"). The best fit model had 5 estimated degrees of freedom.

| Model | AICc | ΔAICc | Akaike Weights |
|---|---|---|---|
| *Composite ~ s(LD1, LD2) + Experiment + Lake* | *99.114* | *0.000* | *0.825* |
| Composite ~ s(LD1, LD2) + Experiment * Lake | 102.210 | 3.096 | 0.175 |
| Composite ~ s(LD1, LD2) + s(LD1) + s(LD2) + Experiment + Lake | 131.456 | 32.342 | < 0.001 |
| Composite ~ s(LD1, LD2) + s(LD1) + s(LD2) + Experiment * Lake | 135.894 | 36.781 | < 0.001 |
| Composite ~ s(LD1, LD2) + s(LD1, Experiment, bs = "fs") + s(LD2, Experiment, bs = "fs") + Lake | 230.428 | 131.314 | < 0.001 |
| Composite ~ s(LD1, LD2) + s(LD1, Lake, bs = "fs") + s(LD2, Lake, bs = "fs") + Experiment | 230.868 | 131.754 | < 0.001 |

1236

1237 **Table S9. Generalized additive models fitted to growth.** Model fit was assessed using AICc, and Akaike Weights represent proportional model
1238 support. A thin plate spline for the two linear discriminant axes s(LD1, LD2) is always included, as is a fixed effect of either experiment (i.e. Martin
1239 & Wainwright 2013, Martin and Gould 2020) or Lake (Crescent Pond/Little Lake) or an interaction between the two. In the last two models,
1240 Experiment and Lake are included as splines, modeled using a factor smooth (bs = "fs"). The best fit model had 8.93 estimated degrees of freedom.

| Model | AICc | ΔAICc | Akaike Weights |
|---|---|---|---|
| *Growth ~ s(LD1, LD2) + Experiment * Lake* | *-44.658* | *0.000* | *1* |
| Growth ~ s(LD1, LD2) + Experiment + Lake | 3.904 | 48.562 | < 0.001 |
| Growth ~ s(LD1, LD2) + s(LD1) + s(LD2) + Experiment * Lake | 46.249 | 90.907 | < 0.001 |
| Growth ~ s(LD1, LD2) + s(LD1) + s(LD2) + Experiment + Lake | 89.121 | 133.779 | < 0.001 |
| Growth ~ s(LD1, LD2) + s(LD1, Lake, bs = "fs") + s(LD2, Lake, bs = "fs") + Experiment | 690.379 | 735.038 | < 0.001 |
| Growth ~ s(LD1, LD2) + s(LD1, Experiment, bs = "fs") + s(LD2, Experiment, bs = "fs") + Lake | 693.748 | 738.406 | < 0.001 |

1241
1242
1243
1244

1245
1246
1247

1248 **Table S10. Generalized additive models fitted to survival.** Model fit was assessed using AICc, and Akaike Weights represent proportional model
1249 support. A thin plate spline for the two linear discriminant axes s(LD1, LD2) is always included, as is a fixed effect of either experiment (i.e. Martin
1250 & Wainwright 2013, Martin and Gould 2020) or Lake (Crescent Pond/Little Lake) or an interaction between the two. In the last two models,
1251 Experiment and Lake are included as splines, modeled using a factor smooth (bs = "fs"). The best fit model had 5 estimated degrees of freedom.

| Model | AICc | ΔAICc | Akaike Weights |
|---|---|---|---|
| *Survival ~ s(LD1, LD2) + Experiment + Lake* | *141.057* | *0.000* | *0.849* |
| Survival ~ s(LD1, LD2) + Experiment * Lake | 144.504 | 3.447 | 0.151 |
| Survival ~ s(LD1, LD2) + s(LD1) + s(LD2) + Experiment + Lake | 173.399 | 32.342 | < 0.001 |
| Survival ~ s(LD1, LD2) + s(LD1) + s(LD2) + Experiment * Lake | 178.189 | 37.132 | < 0.001 |
| Survival ~ s(LD1, LD2) + s(LD1, Experiment, bs = "fs") + s(LD2, Experiment, bs = "fs") + Lake | 273.547 | 132.490 | < 0.001 |
| Survival ~ s(LD1, LD2) + s(LD1, Lake, bs = "fs") + s(LD2, Lake, bs = "fs") + Experiment | 273.694 | 132.637 | < 0.001 |

1274
**Table S11. Generalized additive models fitted to growth including SNPs most strongly associated with composite fitness.** Model fit was
assessed using AICc, and Akaike Weights represent proportional model support. The best fit model for composite fitness using morphology alone
(see Table S8) was used as the base model. The SNPs that were most strongly associated with composite fitness (following a Bonferroni correction)
were included as fixed effects, modeled as splines using a factor smooth, treating genotype as an ordered factor. Note that three SNPs were excluded
due to their close proximity to other SNPs that were more strongly associated. All SNPs were considered individually, as well as all SNPs together.
We were unable to assess all possible combinations of SNPs due to the vast number of potential models given the number of SNPs under
consideration; rather, we fit one final model that only included SNPs found to be significant in the full model. In turn this model led to a substantial
improvement in AICc. The best fit model had 20.29 estimated degrees of freedom.

| Model | AICc | ΔAICc | Akaike Weights |
|---|---|---|---|
| *Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site1) + s(Site2) + s(Site6) + s(Site7) + s(Site8) + s(Site9) + s(Site10)* | *4.586* | *0.000* | *0.999* |
| Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site1) + s(Site2) + s(Site3) + s(Site4) + s(Site5) + s(Site6) + s(Site7) + s(Site8) + s(Site9) + s(Site10) | 40.876 | 36.290 | < 0.001 |
| Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site3) | 55.588 | 51.001 | < 0.001 |
| Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site7) | 58.386 | 53.800 | < 0.001 |
| Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site4) | 65.453 | 60.867 | < 0.001 |
| Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site2) | 71.245 | 66.658 | < 0.001 |
| Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site5) | 72.329 | 67.743 | < 0.001 |
| Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site1) | 73.671 | 69.085 | < 0.001 |
| Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site8) | 74.413 | 69.827 | < 0.001 |
| Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site9) | 74.680 | 70.094 | < 0.001 |
| Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site10) | 88.977 | 84.391 | < 0.001 |
| Composite Fitness ~ s(LD1, LD2) + Experiment + Lake + s(Site6) | 90.427 | 85.841 | < 0.001 |
| Composite Fitness ~ s(LD1, LD2) + Experiment + Lake | 99.114 | 94.527 | < 0.001 |

1283
1284 Note: Site1 = HiC_Scaffold_1:43866598, Site2 = HiC_Scaffold_53:11317840, Site3 = HiC_Scaffold_46:35151009, Site4 = HiC_Scaffold_8:20263964, Site5 =
1285 37:18591438, Site5 = HiC_Scaffold_37:18591438, Site6 = HiC_Scaffold_24:15964553, Site7 = HiC_Scaffold_1848:40590, Site8 = HiC_Scaffold_4461:12939,
1286 Site9 = HiC_Scaffold_12778:1456, Site10 = HiC_Scaffold_18999:1084
1287

1288
1289
**Table S12. Generalized additive models fitted to growth including SNPs most strongly associated with growth.** Model fit was assessed using
AICc, and Akaike Weights represent proportional model support. The best fit model for growth using morphology alone (see Table S9) was used as
the base model. Each of the four SNPs that were most strongly associated with growth (following a Bonferroni correction) were included as fixed
effects, modeled as splines using a factor smooth, treating genotype as an ordered factor. All SNPs were considered individually, as well as all
possible combinations. This was only feasible due to the small number of SNPs assessed (four). The best fit model had 7.97 estimated degrees of
freedom.

| Model | AICc | ΔAICc | Akaike Weights |
|---|---|---|---|
| Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site3) + s(Site4) | -67.649 | 0.000 | 0.490 |
| Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site3) | -65.634 | 2.015 | 0.179 |
| Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site1) | -64.161 | 3.488 | 0.086 |
| Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site1) + s(Site3) | -63.926 | 3.723 | 0.076 |
| Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site1) + s(Site2) | -63.861 | 3.788 | 0.074 |
| Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site2) + s(Site4) | -63.503 | 4.146 | 0.062 |
| Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site2) | -61.849 | 5.800 | 0.027 |
| Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site1) + s(Site4) | -58.044 | 9.604 | 0.004 |
| Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site4) | -56.068 | 11.581 | 0.001 |
| Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site1) + s(Site3) + s(Site4) | -54.878 | 12.770 | < 0.001 |
| Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site1) + s(Site2) + s(Site4) | -54.509 | 13.140 | < 0.001 |
| Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site2) + s(Site3) | -47.602 | 20.047 | < 0.001 |
| Growth ~ s(LD1, LD2) + Experiment * Lake | -44.658 | 22.990 | < 0.001 |
| Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site1) + s(Site2) + s(Site3) | -41.689 | 25.960 | < 0.001 |
| Growth ~ s(LD1, LD2) + Experiment * Lake + s(Site1) + s(Site2) + s(Site3) + s(Site4) | -29.801 | 37.847 | < 0.001 |

1296
1297 Note: Site1 = HiC_Scaffold_46:16512886, Site2 = HiC_Scaffold_1848:40119, Site3 = HiC_Scaffold_1848:40465, Site4 = HiC_Scaffold_7644:5971

1298 **Table S13. Summary statistics for fitness networks constructed using adaptive loci.** Results for networks using all three measures of fitness
1299 (composite fitness, survival, and growth) are reported. Networks were constructed from random draws of five SNPs from either Standing genetic
1300 variation (SGV), introgression, or *de novo* mutations. The scaled number of accessible paths was calculated by dividing the number of accessible
1301 paths between generalists and specialists by the number of nodes present in a given fitness network.

| Fitness Metric | Trajectory | Source | Number of Nodes in Network | | | Number of Edges in Network | | | Number of Accessible Paths | | | Scaled Number of Accessible Paths | | | Distance of Shortest Accesible Path | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | | Median | Mean | SE | Median | Mean | SE | Median | Mean | SE | Median | Mean | SE | Median | Mean | SE |
| Composite | AtoM | SGV | 59.000 | 59.233 | 10.955 | 110.000 | 110.254 | 28.461 | 1.000 | 2.585 | 8.488 | 0.019 | 0.043 | 0.128 | 3.000 | 3.185 | 1.048 |
| | | Introgression | 28.000 | 28.026 | 7.456 | 43.000 | 44.459 | 16.704 | 1.000 | 1.890 | 6.686 | 0.043 | 0.069 | 0.220 | 3.000 | 2.914 | 0.892 |
| | | SGV + Introgression | 51.000 | 50.180 | 12.763 | 92.000 | 92.442 | 30.504 | 1.000 | 5.404 | 75.895 | 0.026 | 0.087 | 0.930 | 3.000 | 3.107 | 0.945 |
| | AtoP | SGV | 61.000 | 61.324 | 9.101 | 116.000 | 117.164 | 24.262 | 1.000 | 4.015 | 21.691 | 0.019 | 0.063 | 0.293 | 4.000 | 3.978 | 1.316 |
| | | Introgression | 46.000 | 46.785 | 11.442 | 77.000 | 80.066 | 28.504 | 1.000 | 1.664 | 2.741 | 0.026 | 0.036 | 0.047 | 3.000 | 3.248 | 1.025 |
| | | *De novo* | 48.000 | 47.373 | 9.243 | 83.000 | 83.082 | 24.048 | 1.000 | 2.039 | 5.624 | 0.026 | 0.042 | 0.098 | 3.000 | 3.495 | 1.052 |
| | | SGV + Introgression | 57.000 | 56.858 | 12.003 | 107.000 | 105.897 | 31.286 | 1.000 | 3.589 | 19.955 | 0.020 | 0.059 | 0.295 | 4.000 | 3.701 | 1.088 |
| | | SGV + *De novo* | 58.000 | 57.200 | 9.633 | 108.500 | 107.732 | 25.493 | 1.000 | 4.396 | 40.886 | 0.020 | 0.076 | 0.728 | 4.000 | 3.768 | 1.159 |
| Survival | AtoM | SGV | 62.000 | 61.752 | 9.036 | 136.000 | 136.282 | 29.143 | 1.000 | 309.508 | 4922.357 | 0.022 | 4.189 | 65.732 | 3.000 | 3.502 | 1.469 |
| | | Introgression | 27.500 | 27.920 | 7.145 | 46.000 | 48.345 | 17.452 | 1.000 | 2.491 | 7.478 | 0.043 | 0.089 | 0.240 | 3.000 | 2.922 | 0.952 |
| | | SGV + Introgression | 51.000 | 50.655 | 12.569 | 103.000 | 104.784 | 35.887 | 1.000 | 31.986 | 284.777 | 0.029 | 0.477 | 3.852 | 3.000 | 3.264 | 1.204 |
| | AtoP | SGV | 61.000 | 61.313 | 9.191 | 131.500 | 134.025 | 28.533 | 3.000 | 358.756 | 3156.552 | 0.054 | 5.721 | 51.543 | 5.000 | 5.163 | 2.077 |
| | | Introgression | 47.000 | 47.954 | 11.559 | 92.000 | 94.270 | 33.744 | 1.000 | 15.053 | 113.473 | 0.031 | 0.244 | 1.630 | 4.000 | 4.081 | 1.626 |
| | | *De novo* | 48.000 | 47.875 | 9.159 | 96.000 | 95.836 | 28.763 | 1.000 | 99.206 | 1933.358 | 0.031 | 1.550 | 29.317 | 4.000 | 4.058 | 1.690 |
| | | SGV + Introgression | 58.000 | 57.379 | 11.275 | 124.000 | 123.244 | 35.495 | 2.000 | 153.381 | 1199.043 | 0.038 | 2.255 | 17.255 | 5.000 | 4.861 | 1.948 |
| | | SGV + *De novo* | 58.000 | 57.572 | 9.406 | 127.000 | 124.970 | 29.966 | 2.000 | 143.683 | 1432.083 | 0.045 | 2.121 | 20.358 | 4.000 | 4.696 | 1.848 |
| Growth | AtoM | SGV | 38.000 | 38.166 | 6.678 | 48.000 | 47.959 | 12.495 | 1.000 | 1.142 | 0.443 | 0.027 | 0.031 | 0.012 | 2.000 | 2.347 | 0.655 |
| | | Introgression | 17.000 | 17.670 | 4.154 | 16.000 | 16.227 | 5.546 | 1.000 | 1.011 | 0.120 | 0.059 | 0.061 | 0.021 | 2.000 | 2.151 | 0.440 |
| | | SGV + Introgression | 31.000 | 31.104 | 7.998 | 35.000 | 36.299 | 13.914 | 1.000 | 1.111 | 0.376 | 0.033 | 0.038 | 0.016 | 2.000 | 2.265 | 0.553 |
| | AtoP | SGV | 38.000 | 37.975 | 6.154 | 52.000 | 51.120 | 12.469 | 1.000 | 1.316 | 0.749 | 0.028 | 0.035 | 0.020 | 2.000 | 2.861 | 1.143 |
| | | Introgression | 27.500 | 28.513 | 7.232 | 31.000 | 33.080 | 12.378 | 1.000 | 1.115 | 0.359 | 0.037 | 0.042 | 0.016 | 2.000 | 2.248 | 0.453 |
| | | *De novo* | 32.000 | 31.402 | 5.488 | 38.000 | 38.112 | 9.874 | 1.000 | 1.104 | 0.339 | 0.032 | 0.036 | 0.011 | 2.000 | 2.329 | 0.624 |
| | | SGV + Introgression | 33.000 | 34.295 | 7.648 | 44.500 | 43.807 | 14.444 | 1.000 | 1.205 | 0.590 | 0.031 | 0.037 | 0.018 | 2.000 | 2.500 | 0.816 |
| | | SGV + *De novo* | 36.000 | 35.389 | 5.713 | 45.000 | 45.326 | 11.109 | 1.000 | 1.154 | 0.519 | 0.029 | 0.033 | 0.015 | 2.000 | 2.571 | 0.812 |

1302

1303 **Table S14. Results of pairwise comparisons using a non-parametric Kruskal-Wallis test for measures of network size (number of nodes and**
1304 **number of edges), using the samples for composite fitness and survival.** *P*-values corrected for multiple tests using the false discovery rate.
1305 Medians, means, and SE for each group reported in Table S13.

| | Pairwise Comparison | Number of Nodes in Network | | Number of Edges in Network | |
|---|---|---|---|---|---|
| | | Generalist To Molluscivore | Generalist to Scale-eater | Generalist To Molluscivore | Generalist to Scale-eater |
| | SGV - Introgression | < 0.0001 | < 0.0001 | < 0.0001 | < 0.0001 |
| | SGV - SGV + Introgression | < 0.0001 | < 0.0001 | < 0.0001 | < 0.0001 |
| | Introgression - SGV + Introgression | < 0.0001 | < 0.0001 | < 0.0001 | < 0.0001 |
| Composite | SGV - *De novo* | - | < 0.0001 | - | < 0.0001 |
| Fitness & | SGV - SGV + *De novo* | - | < 0.0001 | - | < 0.0001 |
| Survival | *De novo* - SGV + *De novo* | - | < 0.0001 | - | < 0.0001 |
| | *De novo* - SGV + Introgression | - | < 0.0001 | - | < 0.0001 |
| | *De novo* - Introgression | - | 0.6921 | - | 0.1128 |
| | Introgression - SGV + *De novo* | - | < 0.0001 | - | < 0.0001 |
1306 | | SGV + *De novo* - SGV + Introgression | - | 0.3068 | - | 0.0831 |

1307

1308 **Table S15. Results of pairwise comparisons using a non-parametric Kruskal-Wallis test for measures of network size (number of nodes and**
1309 **number of edges), using the samples for growth.** *P*-values corrected for multiple tests using the false discovery rate. Medians, means, and SE for
1310 each group reported in Table S13.

| | Pairwise Comparison | Number of Nodes in Network | | Number of Edges in Network | |
|---|---|---|---|---|---|
| | | Generalist To Molluscivore | Generalist to Scale-eater | Generalist To Molluscivore | Generalist to Scale-eater |
| | SGV - Introgression | < 0.0001 | < 0.0001 | < 0.0001 | < 0.0001 |
| | SGV - SGV + Introgression | < 0.0001 | < 0.0001 | < 0.0001 | < 0.0001 |
| | Introgression - SGV + Introgression | < 0.0001 | < 0.0001 | < 0.0001 | < 0.0001 |
| | SGV - *De novo* | - | < 0.0001 | - | < 0.0001 |
| Growth | SGV - SGV + *De novo* | - | 0.0014 | - | 0.0002 |
| | *De novo* - SGV + *De novo* | - | < 0.0001 | - | < 0.0001 |
| | *De novo* - SGV + Introgression | - | 0.0002 | - | 0.0001 |
| | *De novo* - Introgression | - | < 0.0001 | - | < 0.0001 |
| | Introgression - SGV + *De novo* | - | < 0.0001 | - | < 0.0001 |
1311 | | *SGV + De novo* - SGV + Introgression | - | 0.0590 | - | 0.1498 |

1312
1313

**Table S16. Results of pairwise comparisons using a non-parametric Kruskal-Wallis test for differences in the relative number of accessible paths between generalists and specialists in networks constructed using different sources of genetic variation using either composite fitness, growth, or survival as our measure of fitness.** *P*-values corrected for multiple tests using the false discovery rate. Medians, means, and SE for each group reported in Table S13.

| | Pairwise Comparison | Generalist To Molluscivore | Generalist to Scale-eater |
|---|---|---|---|
| Composite Fitness | SGV - Introgression | < 0.0001 | < 0.0001 |
| | SGV - SGV + Introgression | < 0.0001 | 0.7060 |
| | Introgression - SGV + Introgression | < 0.0001 | < 0.0001 |
| | SGV - *De novo* | - | < 0.0001 |
| | SGV - SGV + *De novo* | - | 0.0679 |
| | *De novo* - SGV + *De novo* | - | < 0.0001 |
| | *De novo* - SGV + Introgression | - | < 0.0001 |
| | *De novo* - Introgression | - | 0.0911 |
| | Introgression - SGV + *De novo* | - | < 0.0001 |
| | *SGV + De novo* - SGV + Introgression | - | 0.1232 |
| | Pairwise Comparison | Generalist To Molluscivore | Generalist to Scale-eater |
| Growth | SGV - Introgression | < 0.0001 | < 0.0001 |
| | SGV - SGV + Introgression | < 0.0001 | 0.0145 |
| | Introgression - SGV + Introgression | < 0.0001 | < 0.0001 |
| | SGV - *De novo* | - | < 0.0001 |
| | SGV - SGV + *De novo* | - | 0.5854 |
| | *De novo* - SGV + *De novo* | - | < 0.0001 |
| | *De novo* - SGV + Introgression | - | 0.0996 |
| | *De novo* - Introgression | - | 0.0001 |
| | Introgression - SGV + *De novo* | - | < 0.0001 |
| | *SGV + De novo* - SGV + Introgression | - | 0.0385 |
| | Pairwise Comparison | Generalist To Molluscivore | Generalist to Scale-eater |
| Survival | SGV - Introgression | < 0.0001 | 0.0425 |
| | SGV - SGV + Introgression | 0.0004 | 0.3872 |
| | Introgression - SGV + Introgression | < 0.0001 | 0.0468 |
| | SGV - *De novo* | - | 0.0757 |
| | SGV - SGV + *De novo* | - | 0.7967 |
| | *De novo* - SGV + *De novo* | - | 0.0148 |
| | *De novo* - SGV + Introgression | - | 0.1450 |
| | *De novo* - Introgression | - | 0.5277 |
| | Introgression - SGV + *De novo* | - | 0.0036 |
| | *SGV + De novo* - SGV + Introgression | - | 0.3122 |

**Table S17. Results of pairwise comparisons using a non-parametric Kruskal-Wallis test for differences in the length of the shortest accessible paths between generalists and specialists in networks constructed using different sources of genetic variation using either composite fitness, growth, or survival as our measure of fitness.** *P*-values corrected for multiple tests using the false discovery rate. Medians, means, and SE for each group reported in Table S13.

|  | Pairwise Comparison | Generalist To Molluscivore | Generalist to Scale-eater |
|---|---|---|---|
| Composite Fitness | SGV - Introgression | < 0.0001 | < 0.0001 |
|  | SGV - SGV + Introgression | 0.0002 | 0.0020 |
|  | Introgression - SGV + Introgression | 0.3294 | < 0.0001 |
|  | SGV - *De novo* | - | < 0.0001 |
|  | SGV - SGV + *De novo* | - | 0.0232 |
|  | *De novo* - SGV + *De novo* | - | < 0.0001 |
|  | *De novo* - SGV + Introgression | - | 0.0024 |
|  | *De novo* - Introgression | - | 0.0002 |
|  | Introgression - SGV + *De novo* | - | < 0.0001 |
|  | SGV + *De novo* - SGV + Introgression | - | 0.3535 |
|  | Pairwise Comparison | Generalist To Molluscivore | Generalist to Scale-eater |
| Growth | SGV - Introgression | < 0.0001 | < 0.0001 |
|  | SGV - SGV + Introgression | 0.0182 | 0.0073 |
|  | Introgression - SGV + Introgression | < 0.0001 | 0.0215 |
|  | SGV - *De novo* | - | < 0.0001 |
|  | SGV - SGV + *De novo* | - | 0.0309 |
|  | *De novo* - SGV + *De novo* | - | 0.0003 |
|  | *De novo* - SGV + Introgression | - | 0.0775 |
|  | *De novo* - Introgression | - | 0.3384 |
|  | Introgression - SGV + *De novo* | - | 0.0001 |
|  | SGV + *De novo* - SGV + Introgression | - | 0.3562 |
|  | Pairwise Comparison | Generalist To Molluscivore | Generalist to Scale-eater |
| Survival | SGV - Introgression | < 0.0001 | < 0.0001 |
|  | SGV - SGV + Introgression | 0.0028 | 0.1279 |
|  | Introgression - SGV + Introgression | < 0.0001 | < 0.0001 |
|  | SGV - *De novo* | - | < 0.0001 |
|  | SGV - SGV + *De novo* | - | 0.0152 |
|  | *De novo* - SGV + *De novo* | - | < 0.0001 |
|  | *De novo* - SGV + Introgression | - | < 0.0001 |
|  | *De novo* - Introgression | - | 0.4422 |
|  | Introgression - SGV + *De novo* | - | < 0.0001 |
|  | SGV + *De novo* - SGV + Introgression | - | 0.1454 |