Short title: Hydropower impacts on Amazonian wildlife

Understanding hydropower impacts on Amazonian wildlife is limited by a lack of robust

evidence: results from a systematic review

Eduardo Rodrigues dos Santos[1,2], Fernanda Michalski[1,2,3], Darren Norris[*1,2,4]

[1] Postgraduate Programme in Tropical Biodiversity, Federal University of Amapá, Macapá,
Amapá, Brazil
[2] Ecology and Conservation of Amazonian Vertebrates Research Group, Federal University of
Amapá, Macapá, Amapá, Brazil
[3] Instituto Pró-Carnívoros, Atibaia, São Paulo, Brazil
[4] School of Environmental Sciences, Federal University of Amapá, Macapá, Amapá, Brazil
12
*Corresponding Author:
Darren Norris,
Ecology and Conservation of Amazonian Vertebrates Research Group
Federal University of Amapá
Rod. Juscelino Kubitschek Km 02, 68903-419 Macapá, Brazil.
Email: dnorris75@gmail.com
19

21   Keywords: Amazon, dam, evidence based conservation, hydropower, impact evaluation, study

22   design, vertebrates

Abstract

Background and Research Aims: Although hydropower provides energy to fuel economic development across Amazonia, strategies to minimize or mitigate impacts in highly biodiverse Amazonian environments remain unclear. The growing number of operational and planned hydroelectrics requires robust scientific evidence to evaluate impacts of these projects on Amazonian vertebrates. Here we investigated the existing scientific knowledge base documenting impacts of hydropower developments on vertebrates across Brazilian Amazonia.

Methods: We reviewed the scientific literature from 1945 to 2020 published in English, Spanish and Portuguese to assess the temporal and spatial patterns in publications and the types of study design adopted as well as scientific evidence presented.

Results: A total of 24 published articles documented impacts on fish (n = 20), mammals (n = 3) and freshwater turtles (n = 1). Most study designs (87.5%) lacked appropriate controls and only three studies adopted more robust Before-After-Control-Impact designs. The published evidence did not generally support causal inference with only two studies (8.3%) including appropriate controls and/or confounding variables.

Conclusion: Decades of published assessments (54.2% of which were funded by hydropower developers or their subsidiaries) do not appear to have established robust evidence of impacts of hydropower developments on Amazonian vertebrates. This lack of robust evidence could limit the development of effective minimization and mitigation actions for the diverse vertebrate groups impacted by hydroelectrics across Brazilian Amazonia.

Implications for Conservation: To avoid misleading inferences there is a need to integrate more robust study designs into impact assessments of hydropower developments in the Brazilian Amazon.

Introduction

The development and operation of hydroelectric power plants generates multiple environmental and social impacts across tropical regions, ranging from habitat destruction to changes in river flow, habitat fragmentation, and overhunting (Aurelio-Silva et al., 2016; Benchimol & Peres, 2015; Bueno & Peres, 2019; Cosson et al., 1999; Palmeirim et al., 2017). The increasing number of hydroelectrics in tropical rivers means there is an urgent need to understand impacts to establish minimization and mitigation actions necessary to ensure sustainability of these developments.

In South America, hydropower projects with reservoirs and run-of-river dams are common (Finer & Jenkins, 2012). For example, in 2021 Brazilian Amazonia has 29 operational hydroelectric power plants (including only those with installed power > 30 MW) and an additional 93 in process of regularization and construction (SIGEL, 2021). Projects with reservoir storage (e.g. Balbina dam in Brazil), make it possible to adjust the level of water to produce energy during periods of water scarcity, which can make substantial changes to both the landscape and water flow (Egré & Milewski, 2002; Fearnside, 1989). Projects using run-of-river dams use the natural river flow to generate energy and reduce environmental impacts (Egré & Milewski, 2002). Yet due to highly seasonal rainfall and river flow rates the vast majority of Amazonian run-of-river dams include reservoirs e.g. Belo Monte (Fearnside, 2006; Hall & Branford, 2012) and as such generate drastic impacts on flowrates (Mendes et al., 2021).

The Amazon rainforest is reknowned for its globally important biodiversity and availability of hydric resources (Dirzo & Raven, 2003; Malhi et al., 2008). The Amazon basin has a large vertebrate biodiversity (Silva et al., 2005). For example, the total number of freshwater fish species present in the Amazon basin represents ~15% of all freshwater fishes

described worldwide (Jézéquel et al., 2020). Similarly, for three groups of terrestrial vertebrates (birds, mammals and amphibians), the Brazilian Amazon has a higher overall species richness compared with other Brazilian biomes (Jenkins et al., 2015). Vertebrates have great importance in the management of tropical forest ecosystems (Janzen, 1970). This includes seed dispersal, predation, regulation of water quality, and nutrient and carbon cycles in both terrestrial and aquatic ecosystems (Böhm et al., 2013; Fletcher et al., 2006; Raxworthy et al., 2008).

Amazon biodiversity is increasingly threatened by several factors, including habitat loss and fragmentation and climate change (Dudgeon et al., 2006; Laurance et al., 2011; Li et al., 2013; Malhi et al., 2008; Michalski & Peres, 2007; Schneider et al., 2021). One of the major threats to Amazonian biodiversity identified by the International Union for Conservation of Nature is the construction of hydroelectric power plants (IUCN, 2020). These constructions make a direct impact on the local environment and an indirect impact on a large scale, extending through the entire hydrology basin that is inserted (Carvalho et al., 2018). Expansion of hydropower developments in the Brazilian Amazon started in the 1980s (Fearnside, 2001; Junk et al., 1981), but only since 1986 does Brazilian legislation requires that developers need to produce a mandatory Environmental Impact Assessment (EIA), that evaluates the impact of the project and provides necessary minimization and mitigation actions. Although millions of dollars were invested, these EIAs are widely criticized as overly simplistic and generalist (Fearnside, 2014; Gerlak et al., 2020; Simões et al., 2014).

Systematic reviews summarize and evaluate studies, making evidence available for decision-makers (Gopalakrishnan & Ganeshkumar, 2013). A number of reviews document impacts of dams across the Amazon (Athayde, Mathews, et al., 2019; Ferreira et al., 2014; Lees et al., 2016). Recently several studies evaluated the impacts of hydroelectrics on water flow,

sediments, and on aquatic Amazonian species, mostly fishes (Athayde, Mathews, et al., 2019;

Castello et al., 2013; Latrubesse et al., 2017; Turgeon et al., 2021). But these and other reviews

did not evaluate the quality of evidence presented in the primary studies. Indeed, to date there

have been no systematic reviews on the impacts of hydroelectrics on Amazonian vertebrates.

In this review, we evaluated the scientific literature reporting hydroelectric impacts on

vertebrates in Brazilian Amazonia. Specifically we addressed the following questions: (1) what

are the temporal and spatial patterns of articles, (2) study designs adopted and (3) evidence types

generated.

Methods

Study identification and selection

We focused on vertebrates as this group includes fish which is perhaps the most intensively

studied wilfdlife group in terms of hydropower impacts globally (Algera et al., 2020; Arantes et

al., 2019; Turgeon et al., 2021). As such vertebrates should present a best case scenario for the

scientific evidence documenting hydropower impacts on Amazonian wildlife. Searches were

conducted for articles published from 1945 to 2020 using four different databases: ISI Web of

Science, SCOPUS, PubMed and Scielo. The databases were searched using the following

combination of terms: (Amazon*) and (hydroelectric or hydropower or dam) and (mammal or

fish or bird or reptile or amphibian or vertebrate) and (impact* or effect*). The same terms were

translated and searches repeated in Portuguese and Spanish. Searches were conducted twice,

once on 28 March 2020 and again on 29 March 2021 to update publications from 2020.

Studies were selected following guidelines established by the Preferred Reporting Items for a

Systematic Review and Meta-analysis [PRISMA (Moher et al., 2015; Shamseer et al., 2015),

Figure 1]. First, we screened all titles, keywords and abstracts and excluded duplicates and any

studies that were not related to hydroelectric developments and vertebrates within the legal

Brazilian Amazon. The full-text of all articles that passed initial screening was then read to

establish eligibility.

As our focus was on evaluating impacts, the studies needed to include results from comparisons

with at least one of the following: control areas (including space-for-time) and/or the impacted

area after the hydroelectric was operational. Selected articles needed to present basic

data/primary studies (Salafsky et al., 2019) from operational hydroelectrics, as such laboratory

experiments, simulations, reviews and meta-analysis were not included. Studies that used novel

reservoir environments to test theories (e.g. species-area relationships on reservoir islands) were

not included. In addition, studies with lists of species compared with other areas in only a

qualitative narrative form or where comparisons were only discussed (not included as part of the

sampling methodology or analysis) were also excluded at this stage.

Figure 1. PRISMA flow chart. Showing process used to assess and select studies.

Study data extraction

Each study was evaluated by one reviewer, who compiled: publication year, vertebrate groups,

period of data collection, study design, geographic coordinates for the studied dams [obtained by

joining dam name with coordinates provided by SIGEL (2021)], evidence type and whether the

study received funding/data from the developer/operator (Supplemental Material Appendix 1).

Study design typology followed definitions in Christie et al. (2019) and evidence types were

classified following Burivalova et al. (2019) (Table 1). Finally the PRISMA process and data

extraction stages were independently reviewed by two researchers (DN and FM) and corrections

made to ensure reproducibility and consistency.

Table 1. Study Designs and Evidence Types. Typology used to classify selected studies. Descriptions summarized from Christie et al. (2019) and Burivalova et al. (2019).

Study Design	Description
After	Sampling data post-impact without a control or data before.
Before-After	Sampling data before and post impact without a control.
Control-Impact	Sampling data from a control area and compare with post-impact data.
Before-After Control-Impact	Sampling data before and post impact with a control.

Evidence type	Description
Case Report	Descriptive data from the intervention and its effects, made by interviews, perception or sense of fairness.
Case-Control I	Studies that compare a metric before and after an intervention.
Case-Control II	Studies that compare a metric before and after an intervention taking cofounding variables into account.
Quasi-Experimental	Studies that compare a metric before and after with a control unit similar as possible to treatment units.

Hydroelectric data

To contextualize the literature review we compiled data on the operational hydroelectric plants in

the legal Brazilian Amazon. For each hydroelectric plant we obtained geographic coordinates,

operational start date and power output from the Brazilian Electric Sector Geographic

Information (SIGEL – "Sistema de Informações Georreferenciadas do Setor Elétrico"), provided

and maintained by the Brazilian National Agency of Electricity (ANEEL – "Agência Nacional

151 de Energia Elétrica", downloaded from: https://sigel.aneel.gov.br/Down/, accessed on 30 March

152 2021). We retained only hydroelectric power plants (HPPs) with an installed power greater than

153 30 MW (Supplemental Material Appendix 2). We used ArcGIS 10.3 (ESRI, 2015) in order to

154 produce the final distribution map of the hydroelectric plants and study locations.

Data Analysis

All analyses were performed in R (R Development Core Team, 2020). Patterns in the geographic

and temporal distribution of publications were evaluated using maps and descriptive analysis. As

Brazilian states are an important administrative and legislative unit for the management of

environmental resources, we compared the distribution of hydroelectrics and publications

between the nine states of the 5 Mkm^2 Legal Brazilian Amazon [Acre, Amapá, Amazonas, Mato

Grosso, Maranhão, Pará, Rondônia, Roraima and Tocantins, (IBGE, 2020)]. The distribution of

study designs and evidence types was compared between studies that i) received funding and/or

data from the hydroelectric developer/operator and ii) independent research studies without any

declared association with the hydroelectric developer/operator.

Results

Temporal and spatial distribution of studies

A total of 24 peer-reviewed studies were included in our review most of which (n = 16) were

published between 2015 and 2020 (Figure 2). The first article found in our review was published

in 1981 (Junk et al., 1981). This was four years after the hydroelectric plant under study ("Curuá-

Una") became operational in 1977 and six years after the first hydroelectric plant became

operational in the legal Brazilian Amazon in 1975 (Figure 2). Although the number of

operational hydroelectrics increased stedily in the subsequent decades, the number of published

articles started to increase only recently (Figure 2). After the first published study there was a 12

year gap until the next publication and few studies (n = 4) were published by 2012, despite there

being 15 operational hydroelectrics in 2010.

179
Figure 2. Temporal distribution of published studies and operational hydroelectrics. Annual
frequency of A) published articles documenting impacts on vertebrates (n = 24) and B) newly
operational hydroelectrics (n = 29) across the legal Brazilian Amazon. Dashed lines show
cumulative totals.

Based on our inclusion criteria we were able to identify studies assessing impacts on only three

groups of vertebrates (Figure 2): fish (n = 20), mammals (n = 3) and turtles (n = 1). The major

research interest was related to fish (83.3% of studies) with the four articles published during the

first three decades (1981 – 2013) focusing exclusively on this group (Figure 2). The three

mammal studies (Calaça & de Melo, 2017; Calaça et al., 2015; Palmeirim et al., 2014) were

published between 2014 and 2017 and all focused on the semi-aquatic Giant Otter (*Pteronura*

brasiliensis). The study assessing impacts on turtles (Norris et al., 2018) focused on the Yellow-

spotted River Turtle (*Podocnemis unifilis*).

Figure 3. Spatial distribution of published studies and operational hydroelectrics. Geographic
location of A) operational hydroelectrics (circles, n = 29) and B) studies documenting impacts on
vertebrates (triangles, n = 24) across the legal Brazilian Amazon. The size of the circles showing
hydroelectric locations is proportional to the power output of each hydroelectric, and light grey
lines represent major rivers. Plots show distribution of power output (MW) by C) State of all 29
operational hydroelectric and D) The 12 hydroelectrics included in 24 studies. The sequence of
States is ordered by total power output of operational hydroelectrics in each state (high to low
from left to right).

The studies assessed impacts caused by 12 of the 29 operational hydroelectric plants. The distribution of studies tended to follow the power output of the dams in each state (Figure 3) and we found a positive but insignificant correlation between power output and number of studies per hydroelectric power plant (Spearman Correlation rho = 0.41, p = 0.181). Nearly half of studies (n = 11) investigated impacts of three power plants, namely Jirau and Santo Antônio (n = 7, with 6 studies including both) in the state of Rondônia and Peixe Angical (n = 4) in Tocantins. With the two most intensely studied hydroelectrics (Jirau and Santo Antonio, power output 3750 and 3568 MW respectively) accounting for 7 of the 13 studies published since 2017. The remaining 9 hydroelectric plants had one or two studies each. We also found a weak positive correlation between the number of hydroelectrics and number of published studies per state (Spearman Correlation rho = 0.21, p = 0.686). Mato Grosso was the state with most hydroelectric power plants (n = 13), but was severely under-represented with only two published studies (Figure 3), both of which focused around the recently operational Teles Pires dam [1,819 MW, operational in November 2015, (Calaça & de Melo, 2017; Calaça et al., 2015)].

Study Design and Evidence Type

Most studies (87.5%) adopted either "After" (n = 6) or "Before-After" (n = 15) study designs (Figure 4). Only three studies used a Before-After Control-Impact design, two with fish (Araújo et al., 2013; Lima et al., 2018) and one with turtles (Norris et al., 2018).

Figure 4. Temporal distribution of study designs and evidence types. The A) study design used and B) type of evidence produced by 24 published articles documenting impacts hydroelectric developments on vertebrates across the legal Brazilian Amazon. Classification follows previously published definitions of study designs (Christie et al., 2019) and evidence types (Burivalova et al., 2019). Studies are grouped into those conducted without financial support from the developer/operator ("independent") and those that received financial support or data from the developer/operator ("operator").

Most publications (91.7%, n = 22) did not support causal inference, with evidence coming from

either Case-report (n = 6) or Case-Control I (n = 16) studies (Figure 4). Only one Quasi-

Experimental study was found, which included data collected pre and post reservoir formation

with both impacted and control areas and analysis to explicitly test the Before-After Control-

Impact interaction (Norris et al., 2018). The proportion of independent (n = 11) and operator

funded (n = 13) studies was similar (Chi-squared = 0.17, df = 1, p = 0.683) and there was no

significant difference in the frequencies of study designs or evidence types between

independently or operator funded studies (Figure 4, Fisher's Exact Test p = 0.725 and 0.288 for

study designs and evidence types respectively).

Discussion

Our systematic review showed that (1) studies focused on understanding the impacts of hydroelectrics on Amazonian vertebrates are increasing, but weak sampling designs resulted in a lack of robust evidence, (2) the majority of studies focused on fish, and (3) there was a tendency for studies to be concentrated on high potency "mega" hydropower plants. We first turn to discuss the lack of evidence due to weak sampling designs and then explore the focus on selected vertebrate groups, discrepancy on studies focused on large dams and lack of integrated studies.

The lack of robust evidence was surprising considering hydropower development impacts are so strong and well known at a global scale (Grill et al., 2019; Liermann et al., 2012; Maavara et al., 2020). We found that studies across Brazilian Amazonia were biased by a focus on mega-dams. A major part of the increasing number of studies since 2012 can be attributed to studies of only two dams (Jirau and Santo Antonio). Although the sustainability of both projects was questioned (Fearnside, 2014, 2015), both received certification by Hydropower Sustainability Assessment Protocol (https://www.hydrosustainability.org/published-assessments/santo-antonio and https://www.hydrosustainability.org/published-assessments/jirau , accessed 23 June 2021). Our results show that scientific evidence documenting the impacts of both was generally weak (i.e. below expected best practice). A finding that supports recent analysis showing a link between superficial impact assessments and a lack of social and environmental sustainability of Amazonian hydropower developments (Fearnside, 2018; Gerlak et al., 2020).

We found that studies generally adopted weak sampling designs (e.g. lacking controls) and lacked evidence necessary to generate reliable inference (Christie et al., 2021; Christie et al., 2019; Salafsky et al., 2019). Most of the studies found in our review focused on fishes and are therefore likely to represent best-case scenario in terms of scientific knowledge and evidence

base. In fact, this finding follows global patterns where fishes were one of the most frequently

studied groups used to evaluate effects of hydroelectric dams in both temperate (Algera et al.,

2020) and tropical regions (Arantes et al., 2019). But, impacts of run-of-river dams are poorly

studied even for fish the most intensively studied group (Turgeon et al., 2021). Moreover, there

is a lack of studies on mutiple vertebrate groups, which is essential to understand hydroelectric

effects on complex hydrological systems such as the Amazon (Park & Latrubesse, 2017).

Our review showed a lack of studies assessing multiple hydroeletrics and/or multiple vertebrate

groups along the same river. In Brazil, several hydroelectric plants belonging to different

operators are commonly arranged in the same river, creating "cascades" (Athayde, Duarte, et al.,

2019; Mendes et al., 2017). Although many studies focus on mega-dams,the combined effect of

multiple hydroelectrics, which can cuase cumulative impacts (Athayde, Duarte, et al., 2019)

remains poorly documented. For example, Coaracy Nunes was the first dam installed in the legal

Brazilian Amazon in 1975, since then two additional dams have become operational along the

same river, providing a toal of three dams with a combined oputput of 549 MW (78, 252 and 219

MW) within a 18 km stretch of river. The impact of these multiple dams is thought to have

drasticly altered both upstream and downstream flow rates and following the installation of the

second dam (Ferreira Gomes) in 2014 the rivers downstream course became divided, draining

predominantly to the Amazon river not the Atlantic Ocean (Silva dos Santos, 2017). Whilst

individual studies focus on fish (Sá-Oliveira et al., 2015; Sá-Oliveira et al., 2016) and turtles

(Norris et al., 2018; Norris et al., 2020) along the impacted river, these studies focused on

different dams and adopted different sampling designs, which limits the ability to integrate

results for important basin wide analysis necessary to inform mitigation actions.

We failed to find studies including important cofounding impacts such as deforestation (Stickler

et al., 2013). Although deforestation and tree mortality have been widely documented as

important impacts of Amazonian dams (Athayde, Mathews, et al., 2019; Resende et al., 2019;

Stickler et al., 2013) no studies included these important cofounding variables in the assessments

of vertebrates. For example, the lack of studies in Mato Grosso was particularly surprising

considering previous studies on effects of forest fragmentation on vertebrates in this state

(Michalski & Peres, 2007; Norris & Michalski, 2009).

We found few studies considering the overall number and investment in hydropower projects

across the Legal Brazilian Amazonia. Even fewer studies were found when considering only

those with a robust design and able to establish causal inference. It could be suggested that weak

evidence is a reflection of a lack of investment in science and technology, together with a

reduction in investment in the Brazilian Ministry of the Environemnt over the past twenty years

(de Area Leão Pereira et al., 2019). Although there is undoubtedly support for such

considerations, the lack of robust survey designs can also perhaps be attributed more simply to a

failure of researchers to adopt robust designs (Christie et al., 2021; Christie et al., 2019).

However, we need to highlight that our review has some limitations, as we did not include "grey

literature" in our searchers. Thus, it is important to recognize the potential for gaps or missing

studies that were not published in peer-reviewed journals. On the other hand, as we would expect

published studies to have more robust designs and anlaysis compared with the grey literature, our

review, performed in searches across four different databases and in three languages is likely to

be a best-case representation of the scientific evidence base documenting hydroelectric impacts

on vertebrates in the Brazilian Amazonia.

Implications for conservation

There is an urgent need to take advantage of freely available data to organize and plan effective surveys and sampling strategies to evaluate sustainability of current and future hydroelectric across the Brazilian Amazon. Below we provide recommendations to help develop a more robust evidence base.

1. **Geographical distribution of studies.**

   Research gaps: Studies were focused within specific regions

   Future directions: Increase the number of studies all around Brazilian Amazon with a focus in Mato Grosso state, which has more than 50% of operational and planned hydroelectrics.

2. **Study groups.**

   Research gaps: The majority of studies focus on understanding the impacts on fish.

   Future directions: Increase studies focusing on other threatened vertebrate groups including amphibians, birds, mammals, and reptiles.

3. **Hydroelectric power plants.**

   Research gaps: Most of our reviewed studies were concentrated in three large hydroelectric power plants.

   Future directions: Increase number of studies to represent the distribution of operational and planned power output. This should include closer integration with university research teams to develop robust evidence as part of the necessary Environmental Impact Assessments.

4. **Study design and evidence.**

   Research gaps: There is currently a lack of robust evidence to evaluate impacts of hydroelectric power plants on Amazonian wildlife.

   Future directions: Studies need to include more robust designs (e.g. Before-After Control-Impact) to establish causal inference.

Acknowledgements

The Federal University of Amapá (UNIFAP) provided logistical support.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The data presented here were collected during ERS's master study, which was funded by a studentship from the Brazilian Federal Agency for Support and Evaluation of Graduate Education, Ministry of Education ("Coordenação de Aperfeiçoamento de Pessoal de Nível Superior" – CAPES —Grant: 675849). FM receives a productivity scholarship from CNPq (Grant: Process 302806/2018-0) and was funded by CNPq (Grant: 403679/2016-8).

Referrences

Algera, D. A., Rytwinski, T., Taylor, J. J., Bennett, J. R., Smokorowski, K. E., Harrison, P. M., .
. . Cooke, S. J. (2020). What are the relative risks of mortality and injury for fish during
downstream passage at hydroelectric dams in temperate regions? A systematic review.
Environmental Evidence, 9(1), 3. doi:https://doi.org/10.1186/s13750-020-0184-0
Arantes, C. C., Fitzgerald, D. B., Hoeinghaus, D. J., & Winemiller, K. O. (2019). Impacts of
hydroelectric dams on fishes and fisheries in tropical rivers through the lens of functional
traits. *Current Opinion in Environmental Sustainability, 37*, 28-40.
doi:https://doi.org/10.1016/j.cosust.2019.04.009
Araújo, E. S., Marques, E. E., Freitas, I. S., Neuberger, A. L., Fernandes, R., & Pelicice, F. M.
(2013). Changes in distance decay relationships after river regulation: similarity among
fish assemblages in a large Amazonian river. *Ecology of Freshwater Fish, 22*(4), 543-
552. doi:https://doi.org/10.1111/eff.12054
Athayde, S., Duarte, C. G., Gallardo, A. L. C. F., Moretto, E. M., Sangoi, L. A., Dibo, A. P. A., .
. . Sánchez, L. E. (2019). Improving policies and instruments to address cumulative
impacts of small hydropower in the Amazon. *Energy Policy, 132*, 265-271.
doi:https://doi.org/10.1016/j.enpol.2019.05.003
Athayde, S., Mathews, M., Bohlman, S., Brasil, W., Doria, C., Dutka-Gianelli, J., . . . Kaplan, D.
(2019). Mapping research on hydropower and sustainability in the Brazilian Amazon:
advances, gaps in knowledge and future directions. *Current Opinion in Environmental*
Sustainability, 37, 50-69. doi:https://doi.org/10.1016/j.cosust.2019.06.004
Aurelio-Silva, M., Anciaes, M., Henriques, L. M. P., Benchimol, M., & Peres, C. A. (2016).
Patterns of local extinction in an Amazonian archipelagic avifauna following 25 years of
insularization. *Biological Conservation, 199*, 101-109.
doi:https://doi.org/10.1016/j.biocon.2016.03.016
Benchimol, M., & Peres, C. A. (2015). Widespread Forest Vertebrate Extinctions Induced by a
Mega Hydroelectric Dam in Lowland Amazonia. *Plos One, 10*(7).
doi:https://doi.org/10.1371/journal.pone.0129818
Böhm, M., Collen, B., Baillie, J., Bowles, P., Chanson, J., Cox, N., . . . Zug, G. (2013). The
Conservation Status of the World's Reptiles. *Biological Conservation, 157*, 372-385.
doi:https://doi.org/10.1016/j.biocon.2012.07.015
Bueno, A. S., & Peres, C. A. (2019). Patch-scale biodiversity retention in fragmented landscapes:
Reconciling the habitat amount hypothesis with the island biogeography theory. *Journal*
of Biogeography, 46(3), 621-632. doi:https://doi.org/10.1111/jbi.13499
Burivalova, Z., Miteva, D., Salafsky, N., Butler, R. A., & Wilcove, D. S. (2019). Evidence Types
and Trends in Tropical Forest Conservation Literature. *Trends in Ecology & Evolution,*
34(7), 669-679. doi:https://doi.org/10.1016/j.tree.2019.03.002
Calaça, A. M., & de Melo, F. R. (2017). Reestablishment of giant otters in habitats altered by the
filling of the Teles Pires hydroelectric dam in the Amazonia. *IUCN Otter Specialist*
Group Bulletin, 34(2), 73-78.
Calaça, A. M., Faedo, O. J., & de Melo, F. R. (2015). Hydroelectric Dams: The First Responses
from Giant Otters to a Changing Environment. *IUCN Otter Spec. Group Bull, 32*(1), 48-
58.
Carvalho, D. N., Boniolo, M. R., Santos, R. G., Batista, L. V., Malavazzi, A. A., Reis, F. A. G.
V., & Giordano, L. d. C. (2018). Criteria applied in the definition of influence areas,

impacts and programmes in environmental impact studies of Brazilian hydroelectric power plants. *Geociencias - UNESP, 37*, 15.

Castello, L., McGrath, D. G., Hess, L. L., Coe, M. T., Lefebvre, P. A., Petry, P., . . . Arantes, C. C. (2013). The vulnerability of Amazon freshwater ecosystems. *Conservation Letters, 6*(4), 217-229. doi:https://doi.org/10.1111/conl.12008

Christie, A. P., Amano, T., Martin, P. A., Petrovan, S. O., Shackelford, G. E., Simmons, B. I., . . . Sutherland, W. J. (2021). The challenge of biased evidence in conservation. *Conservation Biology, 35*(1), 249-262. doi:https://doi.org/10.1111/cobi.13577

Christie, A. P., Amano, T., Martin, P. A., Shackelford, G. E., Simmons, B. I., & Sutherland, W. J. (2019). Simple study designs in ecology produce inaccurate estimates of biodiversity responses. *Journal of Applied Ecology, 56*(12), 2742-2754. doi:https://doi.org/10.1111/1365-2664.13499

Cosson, J. F., Ringuet, S., Claessens, O., de Massary, J. C., Dalecky, A., Villiers, J. F., . . . Pons, J. M. (1999). Ecological changes in recent land-bridge islands in French Guiana, with emphasis on vertebrate communities. *Biological Conservation, 91*(2-3), 213-222. doi:https://doi.org/10.1016/s0006-3207(99)00091-9

de Area Leão Pereira, E. J., Silveira Ferreira, P. J., de Santana Ribeiro, L. C., Sabadini Carvalho, T., & de Barros Pereira, H. B. (2019). Policy in Brazil (2016–2019) threaten conservation of the Amazon rainforest. *Environmental Science & Policy, 100*, 8-12. doi:https://doi.org/10.1016/j.envsci.2019.06.001

Dirzo, R., & Raven, P. H. (2003). Global State of Biodiversity and Loss. *Annual Review of Environment and Resources, 28*(1), 137-167. doi:https://doi.org/10.1146/annurev.energy.28.050302.105532

Dudgeon, D., Arthington, A. H., Gessner, M. O., Kawabata, Z. I., Knowler, D. J., Leveque, C., . . . Sullivan, C. A. (2006). Freshwater biodiversity: importance, threats, status and conservation challenges. *Biological Reviews, 81*(2), 163-182. doi:https://doi.org/10.1017/s1464793105006950

Egré, D., & Milewski, J. C. (2002). The diversity of hydropower projects. *Energy Policy, 30*(14), 1225-1230. doi:https://doi.org/10.1016/S0301-4215(02)00083-6

ESRI. (2015). ArcGIS Desktop: Release 10.3. Redlands, CA: Environmental Systems Research Institute.

Fearnside, P. M. (1989). Brazil's Balbina Dam: Environment versus the legacy of the Pharaohs in Amazonia. *Environ Manage, 13*(4), 401-423. doi:https://doi.org/10.1007/BF01867675

Fearnside, P. M. (2001). Environmental impacts of Brazil's Tucurui Dam: unlearned lessons for hydroelectric development in Amazonia. *Environ Manage, 27*(3), 377-396. doi:https://doi.org/10.1007/s002670010156

Fearnside, P. M. (2006). Dams in the Amazon: Belo Monte and Brazil's Hydroelectric Development of the Xingu River Basin. *Environ Manage, 38*(1), 16. doi:https://doi.org/10.1007/s00267-005-0113-6

Fearnside, P. M. (2014). Impacts of Brazil's Madeira River Dams: Unlearned lessons for hydroelectric development in Amazonia. *Environmental Science & Policy, 38*, 164-172. doi:https://doi.org/10.1016/j.envsci.2013.11.004

Fearnside, P. M. (2015). Tropical hydropower in the clean development mechanism: Brazil's Santo Antônio Dam as an example of the need for change. *Climatic Change, 131*(4), 575-589. doi:https://doi.org/10.1007/s10584-015-1393-3

Fearnside, P. M. (2018). Challenges for sustainable development in Brazilian Amazonia.
Sustainable Development, 26(2), 141-149. doi:https://doi.org/10.1002/sd.1725
Ferreira, J., Aragão, L. E. O. C., Barlow, J., Barreto, P., Berenguer, E., Bustamante, M., . . .
Zuanon, J. (2014). Brazil's environmental leadership at risk. *Science, 346*(6210), 706.
doi:https://doi.org/10.1126/science.1260194
Finer, M., & Jenkins, C. N. (2012). Proliferation of hydroelectric dams in the Andean Amazon
and implications for Andes-Amazon connectivity. *Plos One, 7*(4), e35126-e35126.
doi:https://doi.org/10.1371/journal.pone.0035126
Fletcher, D., Hopkins, W., Saldaña, T., Baionno, J., Arribas, C., Standora, M., & Fernandez-
Delgado, C. (2006). Geckos as indicators of mining pollution. *Environmental toxicology*
and chemistry / SETAC, 25, 2432-2445. doi:https://doi.org/10.1897/05-556R.1
Gerlak, A. K., Saguier, M., Mills-Novoa, M., Fearnside, P. M., & Albrecht, T. R. (2020). Dams,
Chinese investments, and EIAs: A race to the bottom in South America? *Ambio, 49*(1),
156-164. doi:https://doi.org/10.1007/s13280-018-01145-y
Gopalakrishnan, S., & Ganeshkumar, P. (2013). Systematic reviews and meta-analysis:
Understanding the best evidence in primary healthcare. *2*(1), 9-14.
doi:https://doi.org/10.4103/2249-4863.109934
Grill, G., Lehner, B., Thieme, M., Geenen, B., Tickner, D., Antonelli, F., . . . Zarfl, C. (2019).
Mapping the world's free-flowing rivers. *Nature, 569*(7755), 215-221.
doi:https://doi.org/10.1038/s41586-019-1111-9
Hall, A., & Branford, S. (2012). Development, Dams and Dilma: the Saga of Belo Monte.
Critical Sociology, 38(6), 851-862. doi:https://doi.org/10.1177/0896920512440712
IBGE. (2020). Legal Amazon Boundaries for 2019 [Press release]. Retrieved from
https://censos.ibge.gov.br/en/2185-news-agency/releases-en/28109-ibge-updates-map-of-
the-legal-amazon.html
IUCN. (2020). The IUCN Red List of Threatened Species. .
Janzen, D. H. (1970). Herbivores and the Number of Tree Species in Tropical Forests. *104*(940),
501-528. doi:https://doi.org/10.1086/282687
Jenkins, C. N., Alves, M. A. S., Uezu, A., & Vale, M. M. (2015). Patterns of Vertebrate
Diversity and Protection in Brazil. *Plos One, 10*(12), e0145064.
doi:https://doi.org/10.1371/journal.pone.0145064
Jézéquel, C., Tedesco, P. A., Bigorne, R., Maldonado-Ocampo, J. A., Ortega, H., Hidalgo, M., . .
. Oberdorff, T. (2020). A database of freshwater fish species of the Amazon Basin.
Scientific Data, 7(1), 96. doi:https://doi.org/10.1038/s41597-020-0436-4
Junk, W. J., Robertson, B. A., Darwich, A. J., & Vieira, I. (1981). Investigações limnológicas e
ictiológicas em Curuá-Una, a primeira represa hidrelétrica na Amazônia Central. *Acta*
Amazonica, 11(4), 689-717. doi:https://doi.org/10.1590/1809-43921981114689
Latrubesse, E. M., Arima, E. Y., Dunne, T., Park, E., Baker, V. R., d'Horta, F. M., . . . Stevaux, J.
C. (2017). Damming the rivers of the Amazon basin. *Nature, 546*(7658), 363-369.
doi:https://doi.org/10.1038/nature22333
Laurance, W. F., Camargo, J. L. C., Luizao, R. C. C., Laurance, S. G., Pimm, S. L., Bruna, E.
M., . . . Lovejoy, T. E. (2011). The fate of Amazonian forest fragments: A 32-year
investigation. *Biological Conservation, 144*(1), 56-67.
doi:https://doi.org/10.1016/j.biocon.2010.09.021

Lees, A. C., Peres, C. A., Fearnside, P. M., Schneider, M., & Zuanon, J. A. S. (2016).
Hydropower and the future of Amazonian biodiversity. *Biodiversity and Conservation,*
25(3), 451-466. doi:https://doi.org/10.1007/s10531-016-1072-3
Li, J. S., Lin, X., Chen, A. P., Peterson, T., Ma, K. P., Bertzky, M., . . . Poulter, B. (2013). Global
Priority Conservation Areas in the Face of 21st Century Climate Change. *Plos One, 8*(1),
9. doi:https://doi.org/10.1371/journal.pone.0054839
Liermann, C. R., Nilsson, C., Robertson, J., & Ng, R. Y. (2012). Implications of Dam
Obstruction for Global Freshwater Fish Diversity. *BioScience, 62*(6), 539-548.
doi:10.1525/bio.2012.62.6.5
Lima, A. C., Sayanda, D., Agostinho, C. S., Machado, A. L., Soares, A. M. V. M., & Monaghan,
K. A. (2018). Using a trait-based approach to measure the impact of dam closure in fish
communities of a Neotropical River. *Ecology of Freshwater Fish, 27*(1), 408-420.
doi:https://doi.org/10.1111/eff.12356
Maavara, T., Chen, Q., Van Meter, K., Brown, L. E., Zhang, J., Ni, J., & Zarfl, C. (2020). River
dam impacts on biogeochemical cycling. *Nature Reviews Earth & Environment, 1*(2),
103-116. doi:https://doi.org/10.1038/s43017-019-0019-0
Malhi, Y., Roberts, J. T., Betts, R. A., Killeen, T. J., Li, W. H., & Nobre, C. A. (2008). Climate
change, deforestation, and the fate of the Amazon. *Science, 319*(5860), 169-172.
doi:https://doi.org/10.1126/science.1146961
Mendes, C. A. B., Beluco, A., & Canales, F. A. (2017). Some important uncertainties related to
climate change in projections for the Brazilian hydropower expansion in the Amazon.
Energy, 141, 123-138. doi:https://doi.org/10.1016/j.energy.2017.09.071
Mendes, Y. A., Oliveira, R. S., Montag, L. F. A., Andrade, M. C., Giarrizzo, T., Rocha, R. M., &
Auxiliadora P. Ferreira, M. (2021). Sedentary fish as indicators of changes in the river
flow rate after impoundment. *Ecological Indicators, 125*, 107466.
doi:https://doi.org/10.1016/j.ecolind.2021.107466
Michalski, F., & Peres, C. A. (2007). Disturbance-Mediated Mammal Persistence and
Abundance-Area Relationships in Amazonian Forest Fragments. *Conservation Biology,*
21(6), 1626-1640. doi:https://doi.org/10.1111/j.1523-1739.2007.00797.x
Moher, D., Shamseer, L., Clarke, M., Ghersi, D., Liberati, A., Petticrew, M., . . . Group, P.-P.
(2015). Preferred reporting items for systematic review and meta-analysis protocols
(PRISMA-P) 2015 statement. *Systematic Reviews, 4*(1), 1.
doi:https://doi.org/10.1186/2046-4053-4-1
Norris, D., & Michalski, F. (2009). Are otters an effective flagship for the conservation of
riparian corridors in an Amazon deforestation frontier. *IUCN Otter Spec. Group Bull,*
26(2), 72-76.
Norris, D., Michalski, F., & Gibbs, J. P. (2018). Beyond harm's reach? Submersion of river turtle
nesting areas and implications for restoration actions after Amazon hydropower
development. *PeerJ, 6*, e4228. doi:https://doi.org/10.7717/peerj.4228
Norris, D., Michalski, F., & Gibbs, J. P. (2020). Community based actions save Yellow-spotted
river turtle (Podocnemis unifilis) eggs and hatchlings flooded by rapid river level rises.
PeerJ, 8, e9921. doi:10.7717/peerj.9921
Palmeirim, A. F., Peres, C. A., & Rosas, F. C. W. (2014). Giant otter population responses to
habitat expansion and degradation induced by a mega hydroelectric dam. *Biological*
Conservation, 174, 30-38. doi:https://doi.org/10.1016/j.biocon.2014.03.015

Palmeirim, A. F., Vieira, M. V., & Peres, C. A. (2017). Non-random lizard extinctions in land-
bridge Amazonian forest islands after 28 years of isolation. *Biological Conservation, 214*,
55-65. doi:https://doi.org/10.1016/j.biocon.2017.08.002
Park, E., & Latrubesse, E. M. (2017). The hydro-geomorphologic complexity of the lower
Amazon River floodplain and hydrological connectivity assessed by remote sensing and
field control. *Remote Sensing of Environment, 198*, 321-332.
doi:https://doi.org/10.1016/j.rse.2017.06.021
R Development Core Team. (2020). R: A language and enviroment for statistical computing.
Vienna, Austria: R Fundation for Statistical Computing.
Raxworthy, C., Pearson, R., Zimkus, B., Reddy, S., Deo, A., Nussbaum, R., & Ingram, C.
(2008). Continental speciation in the tropics: Contrasting biogeographic patterns of
divergence in the Uroplatus leaf-tailed gecko radiation of Madagascar. *Journal of
Zoology, 275*, 423-440. doi:https://doi.org/10.1111/j.1469-7998.2008.00460.x
Resende, A. F. d., Schöngart, J., Streher, A. S., Ferreira-Ferreira, J., Piedade, M. T. F., & Silva,
T. S. F. (2019). Massive tree mortality from flood pulse disturbances in Amazonian
floodplain forests: The collateral effects of hydropower production. *Science of The Total
Environment, 659*, 587-598. doi:https://doi.org/10.1016/j.scitotenv.2018.12.208
Sá-Oliveira, J. C., Hawes, J. E., Isaac-Nahum, V. J., & Peres, C. A. (2015). Upstream and
downstream responses of fish assemblages to an eastern Amazonian hydroelectric dam.
Freshwater Biology, 60(10), 2037-2050. doi:https://doi.org/10.1111/fwb.12628
Sá-Oliveira, J. C., Isaac, V. J., Araújo, A. S., & Ferrari, S. F. (2016). Factors Structuring the Fish
Community in the Area of the Coaracy Nunes Hydroelectric Reservoir in Amapá,
Northern Brazil. *Tropical Conservation Science, 9*(1), 16-33.
doi:https://doi.org/10.1177/194008291600900103
Salafsky, N., Boshoven, J., Burivalova, Z., Dubois, N. S., Gomez, A., Johnson, A., . . . Wordley,
C. F. R. (2019). Defining and using evidence in conservation practice. *Conservation
Science and Practice, 1*(5), e27. doi:https://doi.org/10.1111/csp2.27
Schneider, M., Biedzicki de Marques, A. A., & Peres, C. A. (2021). Brazil's Next Deforestation
Frontiers. *Tropical Conservation Science, 14*, 19400829211020472.
doi:https://doi.org/10.1177/19400829211020472
Shamseer, L., Moher, D., Clarke, M., Ghersi, D., Liberati, A., Petticrew, M., . . . Stewart, L. A.
(2015). Preferred reporting items for systematic review and meta-analysis protocols
(PRISMA-P) 2015: elaboration and explanation. *British Medical Journal, 349*, g7647.
doi:https://doi.org/10.1136/bmj.g7647
SIGEL. (2021). Sistema de Informações Georreferenciadas do Setor Elétrico. Available from
Agência Nacional de Energia Elétrica Sistema de Informações Georreferenciadas do
Setor Elétrico Retrieved 30 March 2021, from ANEEL https://sigel.aneel.gov.br/Down/
Silva dos Santos, E. (2017). *Alterações geomorfológicas no baixo rio Araguari e seus impactos
na hidrodinâmica e na qualidade da água.* (PhD), Universidade Federal do Amapa.
Retrieved from https://www2.unifap.br/ppgbio/files/2018/03/Santos-2017-Tese-de-
Doutorado.pdf
Silva, J. M. C. d., Rylands, A. B., & Gustavo, A. B. d. F. (2005). The Fate of the Amazonian
Areas of Endemism. *Conservation Biology, 19*(3), 689-694.
doi:https://doi.org/10.1111/j.1523-1739.2005.00705.x
Simões, P. I., Stow, A., Hödl, W., Amézquita, A., Farias, I. P., & Lima, A. P. (2014). The Value
of Including Intraspecific Measures of Biodiversity in Environmental impact Surveys is

Highlighted by the Amazonian Brilliant-Thighed Frog (Allobates Femoralis). *Tropical Conservation Science, 7*(4), 811-828. doi:https://doi.org/10.1177/194008291400700416

Stickler, C. M., Coe, M. T., Costa, M. H., Nepstad, D. C., McGrath, D. G., Dias, L. C. P., . . . Soares-Filho, B. S. (2013). Dependence of hydropower energy generation on forests in the Amazon Basin at local and regional scales. *Proceedings of the National Academy of Sciences, 110*(23), 9601. doi:https://doi.org/10.1073/pnas.1215331110

Turgeon, K., Trottier, G., Turpin, C., Bulle, C., & Margni, M. (2021). Empirical characterization factors to be used in LCA and assessing the effects of hydropower on fish richness. *Ecological Indicators, 121*, 107047. doi:https://doi.org/10.1016/j.ecolind.2020.107047

Supplemental Material

Understanding hydropower impacts on Amazonian wildlife is limited by a lack of robust evidence: results from a systematic review

Eduardo Rodrigues dos Santos, Fernanda Michalski, Darren Norris

Appendix 1. List of studies that evaluated hydropower impacts on vertebrates in the Legal Brazilian Amazon.

	Reference	Title	Focal Group	Dam Count	Study Design	Evidence Type
1	Junk et al. (1981)	Investigações limnológicas e ictiológicas em Curuá-Una, a primeira represa hidrelétrica na Amazônia Central	fish	1	After	Case report
2	Santos (1995)	IMPACTOS DA HIDRELÉTRICA SAMUEL SOBRE AS COMUNIDADES DE PEIXES DO RIO JAMARI (RONDÔNIA, BRASIL)	fish	1	Before-After	Case–control I
3	de Merona et al. (2001)	Short term effects of Tucuruí Dam (Amazonia, Brazil) on the trophic organization of fish communities	fish	1	Before-After	Case–control I
4	Pelicice and Agostinho (2012)	Deficient downstream passage through fish ladders: the case of Peixe Angical Dam, Tocantins River, Brazil	fish	1	After	Case report
5	Araújo et al. (2013)	Changes in distance decay relationships after river regulation: Similarity among fish assemblages in a large Amazonian river	fish	1	Before-After Control-Impact	Case–control I
6	Hallwass et al. (2013)	Fishers' knowledge identifies environmental changes and fish abundance trends in impounded tropical rivers	fish	1	Before-After	Case–control I
7	Palmeirim et al. (2014)	Giant otter population responses to habitat expansion and degradation induced by a mega hydroelectric dam	mammal	1	After	Case report
8	Medeiros et al. (2014).	Short-term changes in energy allocation by Hemiodontidae fish after the construction of a large reservoir (Lajeado Dam, Tocantins River)	fish	1	Before-After	Case–control I
9	Sa-Oliveira et al. (2015)	Upstream and downstream responses of fish assemblages to an eastern Amazonian hydroelectric dam	fish	1	After	Case–control I
10	Sá-Oliveira et al. (2015)	Fish community structure as an indicator of the long-term effects of the damming of an Amazonian	fish	1	After	Case–control I

	Reference	Title	Focal Group	Dam Count	Study Design	Evidence Type
		river				
11	Calaça et al. (2015)	Hydroelectric Dams: The First Responses from Giant Otters to a Changing Environment.	mammal	1	Before-After	Case–control I
12	Cella-Ribeiro et al. (2017)	Temporal fish community responses to two cascade run-of-river dams in the Madeira River, Amazon basin.	fish	2	Before-After	Case–control I
13	Calaça and de Melo (2017)	Reestablishment of giant otters in habitats altered by the filling of the Teles Pires hydroelectric dam in the Amazonia.	mammal	1	Before-After	Case–control I
14	Lima et al. (2018)	Using a trait-based approach to measure the impact of dam closure in fish communities of a Neotropical River.	fish	1	Before-After Control-Impact	Case–control II
15	Santos et al. (2018)	The decline of fisheries on the Madeira River, Brazil: The high cost of the hydroelectric dams in the Amazon Basin	fish	2	Before-After	Case report
16	Norris et al. (2018)	Beyond harm's reach? Submersion of river turtle nesting areas and implications for restoration actions after Amazon hydropower development	turtle	1	Before-After Control-Impact	Quasi-experimental
17	Fitzgerald et al. (2018)	Diversity and community structure of rapids-dwelling fishes of the Xingu River: Implications for conservation amid large-scale hydroelectric development	fish	1	After	Case–control I
18	Melo et al. (2019)	Flipped reducetarianism: A vegan fish subordinated to carnivory by suppression of the flooded forest in the Amazon	fish	1	Before-After	Case–control I
19	Van Damme et al. (2019)	Upstream dam impacts on gilded catfish brachyplatystoma rousseauxii (Siluriformes: Pimelodidae) in the Bolivian amazon	fish	2	Before-After	Case–control I
20	Monaghan et al. (2020)	The impact of a hydroelectric dam on Neotropical fish communities: A spatio-temporal analysis of the Trophic Upsurge Hypothesis.	fish	1	Before-After	Case–control I
21	Peronico et al. (2020)	Community reassembly after river regulation: rapid loss of fish diversity and the emergence of a new state	fish	1	Before-After	Case–control I
22	Santos et al. (2020)	Damming Amazon Rivers: Environmental impacts of hydroelectric dams on Brazil's	fish	2	Before-After	Case report

	Reference	Title	Focal Group	Dam Count	Study Design	Evidence Type
		Madeira River according to local fishers' perception				
23	Lima et al. (2020)	Fisheries and trophic structure of a large tropical river under impoundment	fish	2	Before-After	Case-control I
24	Sant'Anna et al. (2020)	Fishing Production of Pinirampus pirinampu and Brachyplatystoma platynemum Catfish has been Affected by Large Dams of the Madeira River (Brazilian Amazon).	fish	2	Before-After	Case report

Appendix 2. Operational hydroelectrics in the Legal Brazilian Amazon. Data for hydroelectrics with installed capcity > 30 MW obtained from the online database maintained by the Brazilian National Agency of Electricity [ANEEL – "Agência Nacional de Energia Elétrica", downloaded from: https://sigel.aneel.gov.br/Down/, accessed on 30 March 2021, (SIGEL, 2021)].

	Name	State	Operational	Number of studies	Installed Capacity(MW)
1	Balbina	Amazonas (AM)	1989	1	249.75
2	Coaracy Nunes	Amapá (AP)	1975	2	78.00
3	Santo Antônio do Jari	Amapá (AP)	2014	0	392.95
4	Ferreira Gomes	Amapá (AP)	2014	0	252.00
5	Cachoeira Caldeirão	Amapá (AP)	2016	1	219.00
6	Juba II	Mato Grosso (MT)	1995	0	42.00
7	Juba I	Mato Grosso (MT)	1995	0	42.00
8	Manso	Mato Grosso (MT)	2000	0	210.00
9	Itiquira (Casas de Forças I e II)	Mato Grosso (MT)	2002	0	157.37
10	Guaporé	Mato Grosso (MT)	2003	0	120.00
11	Jauru	Mato Grosso (MT)	2003	0	121.50
12	Ponte de Pedra	Mato Grosso (MT)	2005	0	176.10
13	Dardanelos	Mato Grosso (MT)	2011	0	261.00
14	Teles Pires	Mato Grosso (MT)	2015	2	1819.80
15	Salto Apiacás	Mato Grosso (MT)	2016	0	45.00
16	São Manoel	Mato Grosso (MT)	2017	0	700.00
17	Colíder	Mato Grosso (MT)	2019	0	300.00
18	Sinop	Mato Grosso (MT)	2019	0	401.88
19	Curuá-Una	Pará (PA)	1977	1	42.80
20	Tucuruí	Pará (PA)	1984	2	8535.00
21	Belo Monte	Pará (PA)	2016	1	11233.10
22	Samuel	Rondônia (RO)	1989	1	216.75
23	Rondon II	Rondônia (RO)	2011	0	73.50
24	Santo Antônio*	Rondônia (RO)	2012	7*	3568.00
25	Jirau*	Rondônia (RO)	2013	6*	3750.00
26	Luís Eduardo Magalhães (Lajeado)	Tocantins (TO)	2001	2	902.50
27	Peixe Angical	Tocantins (TO)	2006	4	498.75
28	São Salvador	Tocantins (TO)	2009	0	243.20
29	Estreito	Tocantins (TO)	2011	0	1087.00
	Totals			**24***	**35738.95**

* 6 studies included both Jirau and Santo Antônio dams.

References

Araújo, E. S., Marques, E. E., Freitas, I. S., Neuberger, A. L., Fernandes, R., & Pelicice, F. M. (2013). Changes in distance decay relationships after river regulation: Similarity among fish assemblages in a large Amazonian river. *Ecology of Freshwater Fish, 22*(4), 543-552. doi:10.1111/eff.12054

Calaça, A. M., & de Melo, F. R. (2017). Reestablishment of giant otters in habitats altered by the filling of the Teles Pires hydroelectric dam in the Amazonia. *IUCN Otter Specialist Group Bulletin, 34*(2), 73-78.

Calaça, A. M., Faedo, O. J., & de Melo, F. R. (2015). Hydroelectric Dams: The First Responses from Giant Otters to a Changing Environment. *IUCN Otter Spec. Group Bull, 32*(1), 48-58.

Cella-Ribeiro, A., Doria, C. R. D., Dutka-Gianelli, J., Alves, H., & Torrente-Vilara, G. (2017). Temporal fish community responses to two cascade run-of-river dams in the Madeira River, Amazon basin. *Ecohydrology, 10*(8). doi:10.1002/eco.1889

de Merona, B., Mendes dos Santos, G., & Gonccalves de Almeida, R. (2001). Short term effects of Tucurui Dam (Amazonia, Brazil) on the trophic organization of fish communities. *Environmental Biology of Fishes, 60*(4), 375-392. doi:10.1023/a:1011033025706

Fitzgerald, D. B., Perez, M. H. S., Sousa, L. M., Goncalves, A. P., Py-Daniel, L. R., Lujan, N. K., . . . Lundberg, J. G. (2018). Diversity and community structure of rapids-dwelling fishes of the Xingu River: Implications for conservation amid large-scale hydroelectric development. *Biological Conservation, 222*, 104-112. doi:10.1016/j.biocon.2018.04.002

Hallwass, G., Lopes, P. F., Juras, A. A., & Silvano, R. A. (2013). Fishers' knowledge identifies environmental changes and fish abundance trends in impounded tropical rivers. *Ecol Appl, 23*(2), 392-407. doi:10.1890/12-0429.1

Junk, W. J., Robertson, B. A., Darwich, A. J., & Vieira, I. (1981). Investigações limnológicas e ictiológicas em Curuá-Una, a primeira represa hidrelétrica na Amazônia Central. *Acta Amazonica, 11*(4), 689-717. doi:https://doi.org/10.1590/1809-43921981114689

Lima, A. C., Sayanda, D., Agostinho, C. S., Machado, A. L., Soares, A. M. V. M., & Monaghan, K. A. (2018). Using a trait-based approach to measure the impact of dam closure in fish communities of a Neotropical River. *Ecology of Freshwater Fish, 27*(1), 408-420. doi:https://doi.org/10.1111/eff.12356

Lima, M. A. L., Doria, C. R., Carvalho, A. R., & Angelini, R. (2020). Fisheries and trophic structure of a large tropical river under impoundment. *Ecological Indicators, 113*. doi:10.1016/j.ecolind.2020.106162

Medeiros, E. R., Pelicice, F. M., Agostinho, C. S., & Marques, E. E. (2014). Short-term changes in energy allocation by Hemiodontidae fish after the construction of a large reservoir (Lajeado Dam, Tocantins River). *Neotropical Ichthyology, 12*(3), 649-658. doi:10.1590/1982-0224-20130186

Melo, T., Torrente-Vilara, G., & Ropke, C. P. (2019). Flipped reducetarianism: A vegan fish subordinated to carnivory by suppression of the flooded forest in the Amazon. *Forest Ecology and Management, 435*, 138-143. doi:10.1016/j.foreco.2018.12.050

Monaghan, K. A., Agostinho, C. S., Pelicice, F. M., & Soares, A. M. V. M. (2020). The impact of a hydroelectric dam on Neotropical fish communities: A spatio-temporal analysis of the Trophic Upsurge Hypothesis. *Ecology of Freshwater Fish, 29*(2), 384-397. doi:https://doi.org/10.1111/eff.12522

Norris, D., Michalski, F., & Gibbs, J. P. (2018). Beyond harm's reach? Submersion of river turtle nesting areas and implications for restoration actions after Amazon hydropower development. *PeerJ, 6*, e4228. doi:https://doi.org/10.7717/peerj.4228

Palmeirim, A. F., Peres, C. A., & Rosas, F. C. W. (2014). Giant otter population responses to habitat expansion and degradation induced by a mega hydroelectric dam. *Biological Conservation, 174*, 30-38. doi:10.1016/j.biocon.2014.03.015

Pelicice, F. M., & Agostinho, C. S. (2012). Deficient downstream passage through fish ladders: the case of Peixe Angical Dam, Tocantins River, Brazil. *Neotropical Ichthyology, 10*(4), 705-713. doi:10.1590/s1679-62252012000400003

Peronico, P. B., Agostinho, C. S., Fernandes, R., & Pelicice, F. M. (2020). Community reassembly after river regulation: rapid loss of fish diversity and the emergence of a new state. *Hydrobiologia, 847*(2), 519-533. doi:10.1007/s10750-019-04117-9

Sa-Oliveira, J. C., Hawes, J. E., Isaac-Nahum, V. J., & Peres, C. A. (2015). Upstream and downstream responses of fish assemblages to an eastern Amazonian hydroelectric dam. *Freshwater Biology, 60*(10), 2037-2050. doi:10.1111/fwb.12628

Sá-Oliveira, J. C., Isaac, V. J., & Ferrari, S. F. (2015). Fish community structure as an indicator of the long-term effects of the damming of an Amazonian river. *Environmental Biology of Fishes, 98*(1), 273-286. doi:10.1007/s10641-014-0288-x

Sant'Anna, I. R. A., Freitas, C. E. d. C., Sousa, R. G. C., Beltrão dos Anjos, H. D., & Doria, C. R. d. C. (2020). Fishing Production of Pinirampus pirinampu and Brachyplatystoma platynemum Catfish has been Affected by Large Dams of the Madeira River (Brazilian Amazon). *Boletim do Instituto de Pesca, 46*(2). doi:https://doi.org/10.20950/1678-2305.2020.46.2.581

Santos, G. M. d. (1995). Impactos da hidrelétrica Samuel sobre as comunidades de peixes do rio Jamari (Rondônia, Brasil). *Acta Amazonica, 25*(3-4), 247-280. doi:https://doi.org/10.1590/1809-43921995253280

Santos, R. E., Pinto-Coelho, R. M., Drumond, M. A., Fonseca, R., & Zanchi, F. B. (2020). Damming Amazon Rivers: Environmental impacts of hydroelectric dams on Brazil's Madeira River according to local fishers' perception. *Ambio*. doi:10.1007/s13280-020-01316-w

Santos, R. E., Pinto-Coelho, R. M., Fonseca, R., Simoes, N. R., & Zanchi, F. B. (2018). The decline of fisheries on the Madeira River, Brazil: The high cost of the hydroelectric dams in the Amazon Basin. *Fisheries Management and Ecology, 25*(5), 380-391. doi:10.1111/fme.12305

SIGEL. (2021). Sistema de Informações Georreferenciadas do Setor Elétrico. Available from Agência Nacional de Energia Elétrica Sistema de Informações Georreferenciadas do Setor Elétrico Retrieved 30 March 2021, from ANEEL https://sigel.aneel.gov.br/Down/

Van Damme, P. A., Córdova-Clavijo, L., Baigún, C., Hauser, M., da Costa Doria, C. R., & Duponchelle, F. (2019). Upstream dam impacts on gilded catfish brachyplatystoma rousseauxii (Siluriformes: Pimelodidae) in the Bolivian amazon. *Neotropical Ichthyology, 17*(4). doi:10.1590/1982-0224-20190118