Strong phylogenetic signal despite high phylogenomic complexity in an Andean plant

radiation (*Freziera,* Pentaphylacaceae)

Laura Frost[1,2,3]*, Ana M. Bedoya[1,3], & Laura Lagomarsino[1]*

[1]Shirley C. Tucker Herbarium, Department of Biological Sciences, Louisiana State University,

Baton Rouge, LA 70808, United States

[2]Biology Department, University of South Alabama, Mobile, AL 36688, United States

[3] These co-first authors contributed equally

*Corresponding author:

Laura Frost

Tel: +1 2565205813

Email: lafrost@southalabama.edu

Laura Lagomarsino

Tel: +1 2255788555

Email: llagomarsino1@lsu.edu

ABSTRACT

The Andes mountains of western South America are a globally important biodiversity hotspot,

yet there is a paucity of resolved phylogenies for plant clades from this region. Filling an

important gap to our understanding of the World's richest flora, we present the first phylogeny of

Freziera (Pentaphylacaceae), an Andean-centered, cloud forest radiation. Our dataset was

obtained via hybrid-enriched target sequence capture of Angiosperms353 universal loci for 50 of

the ca. 75 spp., obtained almost entirely from herbarium specimens. We identify high

phylogenomic complexity in *Freziera*, including the presence of data artifacts. Via by-eye

observation of gene trees, detailed examination of warnings from recently improved assembly

pipelines, and gene tree filtering, we identified that artifactual orthologs (i.e., the presence of

only one copy of a multi-copy gene due to differential assembly) were an important source of

gene tree heterogeneity that had a negative impact on phylogenetic inference and support. These

artifactual orthologs may be common in plant phylogenomic datasets, where multiple instances

of genome duplication are common. After accounting for artifactual orthologs as source of gene

tree error, we identified a significant, but non-specific signal of introgression using Patterson's D

and f4 statistics. Despite phylogenomic complexity, we were able to resolve *Freziera* into nine

well-supported subclades whose evolution has been shaped by multiple evolutionary processes,

including incomplete lineage sorting, historical gene flow, and gene duplication. Our results

highlight the complexities of plant phylogenomics, which are heightened in Andean radiations,

and show the impact of filtering data processing artifacts and standard filtering approaches on

phylogenetic inference.

KEYWORDS: Andean radiation; Angiosperms353; data artifacts; gene tree filtering;

introgression; locus filtering; museomics; paralogy

The Andean mountains in South America are one of the most species-rich areas of the

world and serve as a center of diversity for many plant groups (Gentry 1982; Mutke and

Barthlott 2005). The recent uplift of the Andes has resulted in some of the fastest plant

evolutionary radiations reported to date (Madriñán et al. 2013; Hughes 2016), with some greatly

phenotypically diverse clades (Hughes and Eastwood 2006; Lagomarsino et al. 2016). A

significant portion of Andean plant biodiversity is yet to be described (Ulloa Ulloa et al. 2017)

and many, if not most, Andean plant species, have never been included in a phylogeny.

However, these phylogenies are fundamental toward understanding the evolutionary patterns that

contribute to the origin and diversification of biodiversity in the World's richest flora.

Establishing well-supported phylogenies for Andean plant groups is challenging for many

reasons. Short divergence times between speciation events results in incomplete lineage sorting

(ILS), incipient speciation, and introgression all contribute to poor phylogenetic resolution and

high gene tree-species tree discordance (Vargas et al. 2017; Morales-Briones et al. 2018;

Lagomarsino et al. 2022). This is further complicated by repeated whole genome duplication

events throughout the evolutionary history of plants, at both deep and shallow scales (One

Thousand Plant Transcriptomes Initiative 2019). There are additional practical limitations for

phylogenetic inference in Andean systems. It is difficult to achieve full taxon sampling as

species are often narrowly endemic and distributed in remote locations, and members of clades

occur in many countries, each with different policies concerning collection and exportation of

samples. As a result, achieving dense sampling of Andean-centered lineages commonly requires

the use of herbarium specimens as a source of genetic material, which is associated with lower

quality and quantity DNA than in freshly-collected or silica-dried leaf tissue (Bakker et al. 2015).

Improvements in methodology in the past decade bring us closer to achieving resolved

phylogenies in previously intractable groups. Advancements in genomic sequencing, including

hybrid-enriched target sequence capture, allow for the collection of hundreds to thousands of

loci, even from degraded DNA from natural history collections (Bakker et al. 2015; McKain et

al. 2018). Further, the development of universal probe sets facilitates sequencing of hundreds of

loci for any system, regardless of genomic resources available (Johnson et al. 2019). Analytical

methods are also increasingly able to accommodate multiple biological sources of gene tree

discordance (Ogilvie et al. 2017; Solís-Lemus et al. 2017; Zhang et al. 2018).

Still, a major barrier to phylogenomic inference in many plant clades is the difficulty in

efficient identification of orthologous and paralogous sequences from multi-copy loci (Yang and

Smith 2014; Morales-Briones et al. 2022). The presence of multiple sequences for a single locus

within an individual may result from allelic variation or gene/genome duplication. The latter

process has repeatedly taken place during the evolutionary history of plant lineages, and results

in the presence of non-orthologous gene copies within the same genome (i.e., paralogs) (Li and

Barker 2020). Automated detection of multi-copy genes in phylogenomic datasets could be

hindered when sequence data fails to meet contig number, contig length, or read depth thresholds

set during assembly, a scenario that is more likely when is DNA obtained from herbarium

specimens (Bakker et al. 2015).

The presence of only one copy of a multi-copy gene in a sequenced dataset (i.e., hidden

paralogy) may be due to biological processes or to data processing artifacts. "Pseudo-orthologs"

are a type of hidden paralogy that results from a biological process: differential loss after gene

duplication leads to the presence of a single but non-orthologous copy across species in nature

(Smith and Hahn 2021a, 2021b). While there is some evidence that coalescent-based

phylogenetic methods are relatively robust to pseudo-orthologs (Smith and Hahn 2021a, 2021b),

this finding is applicable only under certain conditions that do not include whole genome

duplication followed by rediploidization— a phenomenon common in plants (Li et al. 2021).

Hidden paralogy can also be artifactual, including when differential assembly of copies in a

multi-copy gene results in the recovery of non-orthologous sequences for a given locus across

taxa. We refer to these as "artifactual orthologs." While it is well-documented that unrecognized

paralogy can have negative impacts on species tree inference (Brown and Thomson 2017), the

presence of artifactual orthologs in plant phylogenomic datasets and their impact on phylogenetic

inference remains underexplored.

We combat the many challenges of Andean plant phylogenomics to infer the first

phylogeny and introgression history of *Freziera* (Pentaphylacaceae), a group that constitutes a

cloud forest plant radiation that previously lacked any phylogenetic information. *Freziera*

includes 75 spp. of trees and shrubs that are widely distributed throughout montane regions of

the Neotropics, from southern Mexico to Bolivia, with a center of diversity (61 spp.) in Andean

cloud forests (Santamaría-Aguilar and Monro 2019; Fig. 1).There has been at least one whole

genome duplication event in an ancestor of Pentaphylacaceae with the inclusive order Ericales

(Larson et al. 2020), but ploidy levels in *Freziera* are unknown due to scarcity of genomic

resources and chromosome counts. Chromosome counts in Pentaphylacaceae are highly variable

(i.e., *Adinandra* [2 spp., n=42], *Cleyera* [1 spp., n=45], *Eurya* [4 spp., n=21, 29, 42], and

Ternstroemia [2 spp., n=20, 25]; from Chromosome Counts Database [CCDB;(Rice et al. 2015);

ccdb.tau.ac.il]).

Using Angiosperms353 (Johnson et al. 2019) target enrichment data derived almost

entirely from herbarium specimens, we establish a phylogenetic baseline of *Freziera* despite

widespread artifactual orthology and find evidence for historical introgression. We further

explore how various types of data curation help remove artifactual orthologs and their impact on

species tree inference. We finally provide suggestions to explore complex empirical

phylogenomic datasets, especially those with a history of genome duplication and that are

obtained with a high reliance on natural history collections.

125
126   **Figure 1.** Diversity and distribution of *Freziera*. *Freziera* has significant variation in leaf morphology and
127   pubescence, as illustrated by branches of a) *F. candicans* and b) *Freziera sp.*; meanwhile, c) flower and d) fruit
128   morphology are relatively stable. *Freziera* is an Andean radiation; e) most species are distributed in this mountain
129   chain in western South America, with some species in Central America, the Guiana Shield, the Caribbean, and the
130   Atlantic Forest of Brazil. Photos by L. Lagomarsino.
131

MATERIALS AND METHODS

Taxon Sampling

Ninety-four accessions representing 55 *Freziera* species—approximately 73% of the

species diversity —were sampled for the ingroup. All but five came from herbarium specimens,

which had an average age of 31.6 years (range: 7.4– 82.8; Supplementary Table 1). Nine

accessions from other Pentaphylacaceae were sampled for the outgroup, including *Eurya*

japonica (in the genus sister to *Freziera)*, *Cleyera albopunctata* (a member of tribe Frezierieae),

and 7 species of *Ternstroemia* (belonging to the sister tribe Ternstroemieae; (Weitzman et al.

2004; Tsou et al. 2016).

DNA extraction, library prep, target enrichment, and sequencing

Detailed descriptions of laboratory methods are provided in online Appendix 1. Briefly,

DNA extraction followed a modified sorbitol extraction protocol (Štorchová et al. 2000). Library

preparation used the KAPA Hyper Prep and KAPA HiFi HS Library Amplification kits with iTru

i5 and i7 dual-indexing primers. Target enrichment was carried out using the MyBaits

Angiosperms353 universal probe set (Johnson et al. 2019; Hale et al. 2020). DNA libraries were

sequenced by Novogene in one Illumina Hiseq 3000 lane with 150 bp paired-end reads.

Although most samples come from herbarium specimens, DNA concentration met minimum

standards for sequencing.

Raw data processing

Demultiplexed raw sequence reads were trimmed with illumiprocessor v2.0.9 (Faircloth

2013, 2016), a wrapper for Trimmomatic v0.39 (Bolger et al. 2014). Default settings were used

and reads with a minimum length of 40 bp kept. Trimmed reads were assembled into

supercontigs (exons and flanking intronic regions) with HybPiper v1.3.1 (Johnson et al. 2016)

using the Angiosperm353 target file as reference (Johnson et al. 2019). In addition, we used a

taxon-specific target file for Pentaphylacaceae using Easy353 (Zhang et al. 2022), a reference-

guided assembly tool for recovery of Angiosperms353 gene sets. We used the transcriptome of

Ternstroemia gymnanthera (One Thousand Plant Transcriptomes Initiative 2019) for sequence

retrieval with Easy353. Additional details of data processing prior to final gene tree inference are

available in online Appendix 1.

Gene tree inference

Preliminary gene trees from the HybPiper v1.3.1 supercontig alignments were generated

from aligned sequences for the 322 loci lacking paralog flags with RAxML v.8.2.12 (Stamatakis

2014) under the GTRCAT nucleotide substitution model with 200 rapid bootstraps. The resulting

trees were processed with TreeShrink v1.3.3 (Mai and Mirarab 2018) to detect and remove

unusually long branches (i.e., potential cross contaminants). Processing in TreeShrink was

performed on a "per-gene" and "all-gene" basis. The identified branches (most of which

corresponded to samples with few sequenced reads; Supplementary Table S1) were removed

from final alignments. We then removed alignments with fewer than 25 ingroup samples from

further analyses. Accessions present in <10% of processed gene trees were trimmed using the R

package ape (Paradis and Schliep 2019a). Gene trees were inferred with IQ-TREE multicore

v2.1.1 (Nguyen et al. 2015) including model section via ModelFinder (Kalyaanamoorthy et al.

2017), tree search, ultrafast bootstraps (Hoang et al. 2018), and Shimodaira-Hasegawa-like

approximate likelihood-ratio test (SH-aLRT; (Guindon et al. 2010; Anisimova et al. 2011).

Paralog warnings and detection of artifactual orthologs

Despite assembling a single sequence per sample for the majority of loci and raising

paralog warnings for <9% loci with HybPiper v1.3.1 (Supplementary Table S1), visual

observation of gene trees and alignments suggested the presence of undetected paralog sequences

(i.e., artifactual orthologs) in the dataset. Evidence for this included split clades of outgroup

184 species and distinctive motifs in alignments that reflected higher sequence divergence than

185 expected by allelic variation alone (Fig. 2b). Gene trees without paralog warnings were first

186 examined by-eye in FigTree v1.4.3 (Rambaut 2014) and sorted as putative orthologs or

187 artifactual orthologs. Artifactual orthologs are identified when different accessions for the same

188 ingroup species, each represented by only a single sequence in the alignment, cluster into two

189 subclades separated by relatively long internal branches (Fig. 2b). Visual detection of artifactual

190 paralogs from gene trees in phylogenomic datasets thus benefits from sampling multiple

191 individuals per species. We grouped loci with no paralog warnings from HybPiper v1.3.1. and

192 confirmed by-eye as putative orthologs into a gene tree set called *orthologs.by.eye* (Table 1).

193   We compared our by-eye identification of artifactual orthologs with two automated

194 paralogy detection strategies: HybPiper v2.0 (Johnson et al. 2016) and HybPhaser (Nauheimer et

195 al. 2021). Both versions of HybPiper could detect paralogs if multiple assembled contigs each

196 cover ≥75% of a reference sequence; this returns a long paralog warning in HybPiper2.

197 HybPiper2 also raises paralog warnings when multiple contigs are assembled across ≥75% of the

198 reference sequence even when individual contigs are <75% the full sequence length; this returns

199 a paralog-by-depth warning. In addition, HybPiper2 reports stitched contigs (i.e., those derived

200 from multiple SPAdes contigs), which can be used to identify chimeric sequences (i.e., stitched

201 contigs in which the two separate sequences are derived from different gene copies); HybPiper2

202 returns chimera warnings when a sufficient number of read pairs from stitched contigs map to

203 different SPAdes contigs and have sequence mismatch above a given threshold for one read but

204 not the other (see HybPiper2 documentation for additional details;

205 https://github.com/mossmatters/HybPiper/). HybPhaser is an automated strategy to identify

206 hybrids, polyploids, contamination, and paralog sequences by identifying samples with a high

proportion of heterozygous loci and allele divergence, and loci with a high proportion of SNPs

(Nauheimer et al. 2021).

In addition, we applied a method for automated orthology detection (i.e., gene tree

pruning) using homologs with monophyletic outgroup (MO; Table 1; Yang and Smith 2014;

Morales-Briones et al. 2021a). MO identifies clusters in gene trees with monophyletic outgroups

and searches from root to tips for duplications. When duplicated taxa are found on either side of

a bifurcation, the subtree with the fewest ingroup taxa is pruned. MO does not directly identify

artifactual orthologs but has the potential to remove their effect by selecting orthogroups and

allowing the inclusion of more loci in a final dataset.

We generated five sets of gene trees to compare artifactual orthology identification using

the above automated pipelines; a description of filtering criteria and thresholds for each dataset

are in Table 1. Loci that received a paralog warning in HybPiper v1.3.1 were excluded to

generate the *orthologs.unfiltered* dataset. The remaining filtered datasets (i.e.,

orthologs.HybPiper2.long, *orthologs.HybPiper2.no.warnings, HybPhaser,* and *MO*) are subsets

of the *orthologs.unfiltered* dataset. For the *HybPhaser* dataset, we did not filter loci by missing

data (we kept loci recovered in at least 0.01 proportion of taxa) to ensure that only loci with

excess heterozygosity (loci with more than 1.5× the interquartile range above the third quartile),

as expected in putative paralogs, were removed. We kept ortholog groups with ≥25 ingroup taxa

in the *MO* dataset for comparison with the other filtering strategies.

Impact of standard gene tree filtering strategies on the presence of artifactual orthologs

We explored how standard gene tree filtering affects the presence of artifactual orthologs

(as identified by-eye) in our dataset. Gene tree filtering is typically applied to minimize gene tree

estimation error, gene tree discordance (Molloy and Warnow 2018), and for phylogenomic

subsampling (Mongiardino Koch 2021) but applying these metrics may also favor the selection

of artifactual orthologs in a dataset (Fig. 2d,e). Gene trees were filtered using five common

empirical criteria for phylogenomic subsampling (Table 1): one alignment-based metric

(*proportion.PI*), two tree length metrics (*tree.length* and *proportion.internal.branch.lengths*), and

two gene tree support metrics (*average.BS* and *bipartition*). Due to the presence of non-

orthologous copies in alignments including artifactual orthologs, a high degree of genetic

variation is expected relative to true orthologs, increasing the proportion of informative sites and,

because of well-supported clusters of non-orthologous sequences, we expected an elevated

average bootstrap support in gene trees (Fig. 2d). Filtering by bootstrap support is standard

practice, collapsing low supported nodes in gene trees prior to species tree inference to remove

gene tree error (Zhang et al. 2018). The two tree length criteria were selected because, while a

high percentage of internal branch length can signal high phylogenetic signal (Shen et al. 2016),

it can also indicate biological pseudo-orthologs (Smith and Hahn, 2016b) and likely, artifactual

orthologs (Fig. 2). Finally, filtering by bipartition support may remove artifactual orthologs by

favoring loci that are more concordant with the inferred species tree (Smith et al. 2018; Fig. 2e).

Standard gene tree filtering strategies were applied to the *orthologs.unfiltered* dataset.

The *proportion.PI* was obtained using AMAS (Borowiec 2016) with outgroups removed.

Estimates of *proportion.internal.branch.lengths* were extracted from IQ-TREE '*.iqtree' output

files. Bootstrap support values were extracted from IQ-TREE maximum likelihood tree files and

averaged to calculate *average.BS*. For *tree.length*, input gene trees were rooted using pxrr in

phyx (Brown et al. 2017) and tree length was calculated with the script 'get_var_length.py' of

SortaDate (Smith et al. 2018) excluding outgroups. *Bipartition* support was calculated with

SortaDate (Smith et al. 2018)'get_bp_genetrees.py' script(Smith et al. 2018) against a species

tree generated with ASTRAL-III (Zhang et al. 2018) from the *orthologs.unfiltered* dataset.

Cutoff thresholds for each filter were selected near the average value in our dataset to keep a

similar number of loci for each criterion.

Table 1. Descriptions of the filtering criteria used, names applied to each of the datasets, number of loci removed
and selected by each, and the impact of filtering criteria on gene tree discordance, branch support, and species tree
topology. Values for datasets that performed equally or better than the *orthologs.unfiltered* dataset for each metric
are bolded. Major topological conflicts between species trees and a consensus topology of the 11 filtered datasets
(Fig. 3b) are shown, including instances where the species tree is compatible (+) or in disagreement (-) with the
consensus topology. Datasets from which the Elaphoglossifolia group was not resolved as monophyletic are
indicated with "n/a". Asterisks mark disagreement from the alternative placement of only one species. The
"Branching order of the Elaphoglossifolia group" includes conflict in the placement of *F. magnibracteolata.*

Description of filtering criteria (for keeping loci)	Putative orthologs dataset name after filtering	No. of loci removed by filtering criteria	No. of loci that passed filtering criteria	Normalized Quartet Score	Average branch support	Proportion of highly supported branches (PP > 0.97)	Two clades in Candicans group	Mono-phyletic Elapho-glossifolia group	Branching order of Elapho-glossifolia group	Placement of Arbutifolia clade
loci not flagged as paralogs by HybPiper v.1.3.1	*orthologs.unfiltered*	31	313	0.567	0.774	0.390	+	−	n/a	+
loci identified as putative orthologs after by-eye inspection of gene trees and alignments	*orthologs.by.eye*	131 (90)	182	**0.634**	**0.792**	0.377	+	+	+	+
loci with 1 contig at ≥75% reference sequence (no long paralog warnings from HybPiper v.2.0)	*orthologs.HybPiper2.long*	51	262	**0.582**	**0.780**	0.377	+	+	+	+
loci without paralog warnings by depth or length from HybPiper v.2.0	*orthologs.HybPiper2.no.warnings*	126	187	**0.619**	**0.783**	0.351	+	+	+	+
loci that have a lower proportion of SNPs than 1.5x the interquartile range above the 3rd quartile for any given sample	*HybPhaser*	71	242	**0.597**	**0.783**	0.364	+	+	+	+
orthology inference via gene tree pruning using homologs with monophyletic, non-repeating outgroups	*MO*	91	222	**0.662**	0.664	0.221	+*	+*	−	−
alignment contains ≥ 7.5% parsimony informative sites in ingroup (average value across all alignments)	*proportion.PI*	153	160	0.528	0.715	0.247	−	−	n/a	+
gene trees with internal branch lengths comprising an above average proportion of the total tree length across all gene trees	*proportion.internal.branch.lengths*	164	148	0.542	0.685	0.234	−	+	+	+
gene trees with average bootstrap support across all nodes above the average value across all gene trees	*average.BS*	150	163	0.560	0.727	0.234	+	+	−	+
gene tree with total tree length above the average value across all gene trees	*tree.length*	173	139	0.524	0.688	0.260	−	−	n/a	−
gene trees with above average bipartition support relative to the species tree inferred from the orthologs.unfiltered dataset	*bipartition*	147	166	**0.601**	**0.792**	**0.390**	+	+	+	+

268   *Phylogenetic relationships and introgression in Freziera*

269   *Impact of filtering criteria on species tree inference.*—Species trees were inferred in

ASTRAL-III from the 11 datasets listed in Table 1. A majority-rule consensus tree was

generated from the species trees for the 11 filtered datasets (*consensus* function of the R package

ape; Paradis and Schliep 2019b) to identify major clades that were concordant across datasets.

Gene tree discordance was assessed for all species tree topologies using the final normalized

quartet score (NQS; proportion of quartet trees in gene trees that are present in the species tree

(Mirarab et al. 2014b) estimated with ASTRAL-III. Impact of filtering criteria on branch support

was assessed by calculating the average branch support and proportion of well-supported

branches (ppl≥0.97; Rabiee and Mirarab 2020) across the 11 resulting species trees. We inferred

a second consensus tree from the species trees generated from the datasets that showed the

greatest improvements to both normalized quartet score and average branch support (*bipartition,*

by.eye, HybPhaser, and *orthologs.HybPiper2.no.warnings*), which also had substantial

reductions in the number of artifactual orthologs (Fig. 2c). The resulting tree was compared to

the *orthologs.unfiltered* dataset to explore the impact that the removal of paralogs, including

artifactual orthologs, has on species tree inference.

Signals of introgression in Freziera.—To assess genomic evidence of introgression in

species of *Freziera*, we calculated Patterson's D and f4 statistics using the *Dtrios* function in

Dsuite v.0.4 r43 (Malinsky et al. 2021). The input VCF file was generated with dDocent (Puritz

et al. 2014) by mapping all *Freziera* trimmed reads to the sequences of *T. tepezapote* for the loci

in the *bipartition*, *by.eye*, and *hybiper2.no.warnings* datasets. We called SNPs using default

values for all mapping parameters. The resulting VCF files were filtered with vcftools (Danecek

et al. 2011). We retained SNPs with <50% missing data and retained only one SNP per 100 bp

window to decrease the likelihood of including linked SNPs. The statistical significance of the D

and f4 was assessed using block jackknife on windows of 75–78 SNPs followed by Benjamini-

Hochberg correction as implemented in R (Benjamini and Hochberg 1995) to assess family-wise

error rate (following (Malinsky et al. 2018). The D and f4 statistics were estimated for all

possible trios across three datasets (13,244, 15,180, 14,190 trios for the *bipartition*, *by.eye*, and

hybiper2.no.warnings respectively). The ASTRAL-III species trees inferred from each of the

three datasets were specified in Dsuite so that the D and f4 estimated values were arranged

according to the tree.

The relatively small number of loci in our dataset limited the power to detect introgression

along a phylogeny. The f-branch statistic, which accounts for the correlation of D and f4 due to

shared ancestry among multiple potential introgression donor species (Malinsky et al. 2018),

allows for a better interpretation of introgression patterns across a tree (Malinsky et al. 2021).

However, simulation analyses have shown that for a relatively small number of unlinked SNPs

(<10,000), the proportion of cases where the strongest inferred f-branch signal corresponds to the

correct simulated gene flow recipient and donor branches is <20% (Malinsky et al. 2021). Due to

the limited number of unlinked SNPs in our targeted sequence capture dataset, we did not apply

this metric.

RESULTS

Sequence capture efficiency

Voucher information for accessions and per sample data for read trimming and contig

assembly are available in Supplementary Table S1. Of the 102 accessions for which target

enrichment libraries were prepared and sequenced, 22 (21 *Freziera* and 1 outgroup) were

excluded either because no sequences were assembled for any of the 353 target loci, or because

they were identified to have exceedingly long branches with TreeShrink (most of the latter

corresponded to samples with few sequenced reads; Supplementary Table S1). Target

enrichment efficiency across all sequenced samples is shown in Supplementary Figure S1.

Seventy-two accessions representing 50 of the 75 species of *Freziera* and 8 members of the

outgroup remained (see methods; Supplementary Table S1), which had assembled sequences for

52–348 genes (average: ~296). Standard assembly statistics from both versions of HybPiper are

in Supplementary Tables S1-S2. No loci were flagged by HybPiper2 as putatively chimeric.

However, all but 13 loci included at least one sequence with stitched contigs (Supplementary

Table S3). All sequences without stitched contigs correspond to regions spanning a single exon.

Off-target data collection was insufficient to assemble plastomes.

Paralog warnings and detection of artifactual orthologs

The 11 paralogy detection and gene tree filtering criteria that we applied generated

datasets with 140–313 loci (Table 1). Results of artifactual ortholog detection by-eye and from

automated strategies are shown in Fig. 2a and Supplementary Table S3. HybPiper v.1.3.1 raised

paralog warnings for 31 loci (i.e., >1 contig was assembled for at least one sample for a given

locus). Of the 322 loci that were not flagged as paralogs, nine were removed because they

contained fewer than 25 ingroup samples, leaving 313 with a single assembled contig per

species. By-eye inspection of these 313 loci resulted in the identification of 90 artifactual

orthologs. Loci flagged as potentially paralogous in HybPiper v.1.3.1 also had long paralog

warnings and paralog warnings by depth with HybPiper2. Of the 313 loci without paralog

warnings from HybPiper v.1.31, 51 had long paralog warnings, of which 46 also received

paralog warnings by contig depth with HybPiper2. An additional 75 loci were not flagged as

long paralogs but did receive paralog warning by contig depth for at least one sample, totaling

121 loci with warnings by contig depth and 126 loci with any kind of paralog warning issued by

HybPiper2. HybPhaser identified 73 loci as putative paralogs (Supplementary Table S3; online

Appendix 2), most of which also had paralog warnings by depth. There was substantial overlap

of loci identified by these methods (Fig. 2c). Three of the methods we compared— HybPiper2

paralog-by-depth warnings, HybPiper2 paralog-by-length warnings, and HybPhaser— flagged

70 (78%) of the loci that we identified as a artifactual orthologs in our by-eye assessment (Fig.

2c). Of these, HybPiper2 paralog-by-depth warnings had the greatest overlap with our by-eye

assessment (Fig. 2c; Supplementary Table S3). Assembly with a taxon-specific target file

resulted in similarly low levels of paralog detection as the universal target file: <8% of

assembled loci recovered >1 long contig for at least one sample (Supplementary Table S4).

There was a correlation between the number of reads mapped and the number of paralog

warnings from HybPiper in a sample (Supplementary Fig. S3).

Impact of standard gene tree filtering strategies on the presence of artifactual orthologs

Summary statistics for alignments and gene trees are available in Supplementary Table

S3. The impact of gene tree filtering strategies on removal of artifactual and putative orthologs

(as identified by-eye) is shown in Fig. 2e. Putative orthologs are removed with all filtering

strategies, and only filtering gene trees by bipartition support resulted in the removal of most

artifactual orthologs. As commonly applied, the remaining four gene tree filtering criteria all

resulted in datasets in which there were either similar proportions of putative orthologs and

artifactual orthologs in the final datasets (*average.BS, proportion.internal.branch.length*), or that

favored the inclusion of artifactual orthologs (*tree.length*, *proportion.PI*).

Figure 2. Exemplary gene trees and alignment subsets for a) a putative ortholog, and b) an artifactual ortholog. Putative ortholog gene trees exhibit monophyletic genera for outgroups and appropriate relationships to the ingroup, a relatively shallow backbone in the ingroup, and relationships similar to the preliminary species tree. A relatively low level of variation is present in alignments for putative orthologs, which is fitting of a locus in a universal probe set being applied within a genus. Artifactual ortholog gene trees often have deep divergences between subsets of the ingroup, including between samples from the same species. Within those subclades, patterns of relationships from the preliminary species tree are repeated. In some cases, as in b), outgroups are also non-monophyletic and subsets of outgroup samples are recovered as sister to the ingroup subclades. Alignments for artifactual orthologs exhibit relatively high variation with observable motifs between the suspected copies, as indicated by the black and white dots. c) Overlap between artifactual orthologs identified by-eye, and paralog warnings raised by automated paralogy detection strategies. d) Expectations for informative, single-copy orthologs change for many gene tree filtering metrics when data sets include artifactual orthologs; arrows indicate whether expected values for metrics are high or low. e) Impact of gene tree filtering strategies on removal of artifactual and putative orthologs as identified by-eye.

The dashed lines indicate the various thresholds above which gene trees were kept in filtered datasets, as indicated in
Table 1.

Impact of data curation on phylogenetic performance

We found that removing artifactual orthologs from our datasets improved phylogenetic

performance metrics, and that, with the exception of filtering by bipartition support, gene tree

filtering did not. Datasets in which curation reduced the proportion of artifactual orthologs

(*orthologs.by.eye, orthologs.HybPiper2, HybPhaser*, and *bipartition*) had the highest normalized

quartet scores and average branch support (Table 1). Except for *bipartition*, these values were

consistently lower in datasets resulting from standard gene tree filtering methods (Table 1),

likely due to an increased proportion of artifactual orthologs relative to the unfiltered dataset

(Fig. 2b). Relative to the unfiltered dataset, the number of highly supported branches was similar

in datasets in which artifactual orthologs were filtered out, and lower in datasets that underwent

conventional gene tree filtering, again apart from *bipartition* (Table 1). While there was some

impact on removing artifactual orthologs on branch support, it is possible that these results were

relatively small due to the high degree of gene tree heterogeneity in our dataset (Table 1).

We also found consistent impacts of removing artifactual orthologs on the species tree

topology. Datasets where artifactual orthologs were removed (*by.eye, HybPiper2.long,*

HybPiper2.depth, HybPhaser, bipartition) consistently recovered the Elaphoglossifolia group as

monophyletic. Contrastingly, species tree topologies generated by gene tree filtering differed

from any of those generated from datasets that were curated to remove artifactual orthologs, and

sometimes introduced relationships that were found in no other analyses, including of the

unfiltered dataset (e.g., non-monophyletic Callophylla and Karsteniana clades and placement of

the Arbutifolia clade; Table 1; Fig. 3b).

Phylogenetic relationships and introgression in Freziera

The majority-rule consensus tree of all 11 species trees resulting from data filtering and

the consensus tree of the four datasets with the lowest proportions of artifactual orthologs were

congruent in the major clades recovered and the relationships between those clades. Seven clades

that were frequently inferred across analyses were identified—the Humiriifolia, Canescens,

Incana, Arbutifolia, Lanata, Karsteniana, and Calophylla clades—along with *F. grisebachii* and

F. magnibracteolata, which formed monotypic clades (Fig. 3a). The Elaphoglossifolia group,

comprising the Canescens and Incana clades and F. magnibracteolata, was additionally recovered

as monophyletic in both consensus trees. Five of the 11 datasets produced species trees

consistent with monophyly of the nine clades and their branching order in the consensus tree

(Table 1; Supplementary Fig. S2). Among species trees consistent with the consensus topology,

some branches at deep nodes were inferred with high support (ppl≥0.97) across all analyses: the

common ancestor of all *Freziera*, the common ancestor of core *Freziera* (all *Freziera* excluding

the Humiriifolia clade), and the successive node within core *Freziera*, the Humiriifolia,

Canescens, and Incana clades, and the Candicans group, which comprises the Karsteniana, and

Calophylla clades (Fig. 3). Species-level relationships within most subclades were consistent

across analyses except for the species-rich Candicans group (Supplementary Fig. S4).

Regions of lowest support in the consensus tree tended to be in conflict between species

trees and the consensus topology. These regions often involved two notable sets of taxa: the

Elaphoglossifolia group, comprising the Incana and Canescens clades and *F. magnibracteolata*,

and the Candicans group, which comprises the Calophylla and Karsteniana clades (Table 1, Fig.

3; Supplementary Fig. S4). While the Elaphoglossifolia group was monophyletic in most

datasets, including all in which the proportion of artifactual orthologs was reduced (Table 1), the

unstable placement of *F. magnibracteolata* rendered it non-monophyletic three datasets:

orthologs.unfiltered, *proportion.PI*, and *tree.length* (Table 1; Fig. 3b; Supplementary Fig.

S2a,g,j). Contrasting with the Elaphoglossifolia group, the monophyly of the Candicans group

was identified in all analyses, generally with high support despite a high degree of gene tree

heterogeneity; however, the resolution of its constituent Calophylla and Karsteniana subclades

varied across analyses (Table 1). While all datasets in which the proportion of artifactual

orthologs was reduced resolved the Calophylla and Karsteniana subclades as monophyletic,

species-level relationships within these groups varied (Supplementary Fig. S4).

For the *bipartition*, *by.eye*, and *orthologs.HybPiper2.no.warnings* datasets, p-values for

the estimated D statistics could not be calculated for 75.9%, 85.5%, and 93.6%, respectively, of

all evaluated trios due to lack of ABBA-BABA variants (Supplementary Fig. S5). Estimates of

Patterson's D >0 were significant (corrected p-value ≤ 0.05, Z-score >3) for 104 (3.27%), 162

(7.38%), and 46 (5.08%) trios out of the remaining evaluated trios. The number of statistically

significant D statistics was probably underestimated in our study given the large proportion of

trios for which p-values could not be calculated due to a lack of ABBA-BABA variants (~76%–

94%). Resulting f4-ratio statistics across the three datasets also show evidence for multiple

instances of introgression in *Freziera* (Supplementary Fig. S5). These results were further

supported by the results from HybPhaser, which identified several samples with >80% locus

heterozygosity (LH) and >1% allele divergence (AD) across all major groups recovered (online

Appendix 2; Supplementary Fig. S6). This is indicative of prevalence of introgression and

polyploidy (Nauheimer et al. 2021) in *Freziera* species included in this study.

444
Figure 3. Names of clades consistently recovered and species tree topologies for a) The consensus tree of the four
datasets in which the proportion of artifactual orthologs was reduced (*bipartition*, *by.eye*, *HybPhaser*,
orthologs.HybPiper.no.warnings; left) versus the orthologs.unfiltered species tree (right). The placement of *F.*
magnibracteolata– the major topological conflict between the two trees– is highlighted with the blue line connecting
the tip labels in each. Branches with full support (LPP from ASTRAL = 1) are indicated by an asterisk; elsewhere,
the LPP×100 is provided. Support for each of the four datasets in the consensus tree are given clockwise from the
top left: *bipartition*, *by.eye*, *HybPhaser*, *orthologs.HybPiper.no.warnings*. b) Cartoon trees displaying the most

frequent differences in relationships between clades recovered by different datasets and the consensus tree of all
datasets. Cartoons depict one topological scenario but not the only topological outcome.
454

DISCUSSION

Relying on DNA extracted primarily from herbarium specimens, we inferred the first

phylogeny of *Freziera* (Pentaphylacaceae), an understudied tropical plant lineage. Our museomic

dataset, representing hybrid-enriched target sequence capture of Angiosperms353 loci, highlights

the many challenges of working with understudied clades even in the genomic era: no a priori

phylogenetic hypothesis, a universal bait set, poor-quality DNA, and high proportion of paralogs,

many of which we identified as artifactual orthologs. Despite phylogenomic complexity

including a high degree of gene tree heterogeneity, we resolve *Freziera* into nine clades whose

histories have been shaped by myriad evolutionary processes, including incomplete lineage

sorting, introgression, and gene or genome duplication. In the face of these complexities, we

identify that the biggest improvements to phylogenetic inference in *Freziera* did not come from

filtering gene trees to maximize phylogenetic informativeness. Instead, they came from reducing

the noise from artifactual orthologs, which was accomplished in a variety of ways: observing

data by-eye, implementing automated pipelines, and identifying gene tree filtering mechanisms

that are consistent with reducing this artifact. We offer recommendations on strategies for

removal of similar data processing artifacts for phylogenetic inference of groups where multi-

copy loci are expected to be prevalent.

Identification of artifactual orthologs using automated pipelines

We observed a widespread pattern of artifactual orthology in our dataset, in which multi-

copy genes were recovered as single copy due to errors in the assembly process. These loci had

multiple identifiable motifs in their alignments, and their resulting gene trees typically exhibited

477 polyphyly of species and deep divergences between clades, including outgroup taxa and

478 members of those spuriously polyphyletic species (Fig. 2b). As troubling as the prospect of

479 unfiltered paralogs may be for empiricists, artifactual orthologs may be easier to detect than

480 biological pseudo-orthologs (Smith and Hahn 2021b) because of these striking, easily

481 identifiable patterns.

482   Using our visual inspection as a baseline, we were able to assess the ability of automated

483 paralogy detection pipelines to identify artifactual orthologs. There was substantial overlap in the

484 loci that we identified by-eye and those removed through various automated paralog detection

485 mechanisms (Fig. 2a). While artifactual orthologs result from the assembly of one contig for a

486 truly multi-copy locus, they were often identifiable if more than one short contig covered >75%

487 of the target length (the strategy used in *orthologs.HybPiper2.no.warnings*). Similarly, but to a

488 lesser extent, artifactual orthologs were removed by filtering out loci with high proportion of

489 SNPs (*HybPhaser*) and with the assembly of multiple contigs at at least 75% of sequence length

490 (*orthologs.HybPiper2.long*). We recommend by-eye identification as an initial strategy to

491 explore whether artifactual orthologs are present in an individual empirical dataset, and

492 emphasize the utility of sampling multiple individuals per species.

493   Our dataset relied heavily on degraded DNA extracted from herbarium tissue, which we

494 believe contributes to the assembly of artifactual orthologs. Though high-molecular weight DNA

495 is fragmented during library preparation, the desired fragment size for Illumina libraries is 200-

496 500 base pairs (bp; Bronner and Quail 2019). As is common when working with specimens from

497 the wet tropics (Bakker et al. 2015), many of our samples had a high proportion of fragments

498 <200 bp (unpublished). These short fragments restrict data collection in non-coding regions, as

499 fragments are less likely to span the regions flanking targeted exons. Using HybPiper v2.0.1, we

identified a high proportion of stitched contigs—separate contigs concatenated into a single

sequence. The assembly of multiple contigs belonging to different paralog copies into a single

sequence (i.e., chimeric sequences) is an artifact that is possible in the presence of stitched

contigs. The preferential selection of the longest contig with the highest coverage for a given

gene region during assembly further contributes to decreased paralog detection and increased

chimeric assembly. Despite the high proportion of stitched contigs, no genes were flagged by

HybPiper v2.0.1 with a chimera warning for any sample (Supplementary Table S2). This is

likely due to current limitations and stringency in chimeric sequence detection. Specifically,

paired ends must completely map to separate contigs and reads must map entirely within an exon

on the separate contigs to flag a chimera warning. Short fragments from degraded DNA reduce

the likelihood of finding read pairs that will meet these criteria, and, therefore, the likelihood of

detecting chimeric sequences, though they may be present in the dataset.

Effectively identifying chimeric sequences from single or stitched contigs (as even a

single assembled contig for a given locus may be a chimera of different alleles) remains a

challenge in the assembly of target capture data from short read sequences, especially as

assembly pipelines cannot distinguish between reads derived from different paralogs or alleles.

Detection of chimeric sequences and its impact on phylogenetic inference remains a fundamental

problem in phylogenomics, particularly in groups where polyploidization and hybridization are

suspected to be prevalent (Morales-Briones et al. 2018). While not possible with degraded DNA

from herbarium specimens, a potential solution includes generating long-read sequence data at a

sufficient depth for accurate phasing of copies.

We believe the lack of data spanning introns, resulting in a high proportion of stitched

contigs, and the potential for greater disparity between capture of both copies from herbarium

DNA, is a primary source of artifactual orthologs. Low sequencing depth resulting in poor

coverage of non-coding regions is another factor that can increase the proportion of stitched

contigs and/or differential success capturing copies. However, sequencing depth is unlikely to be

the main source of artifactual orthologs in our dataset as the number of reads mapped to target

loci (Supplementary Table S1) in our dataset is higher per sample than studies that have

successfully examined paralogs in Angiosperms353 datasets (Johnson et al. 2019; Gardner et al.

2020). Taken together, this highlights caveats that remain with museomic data: long-read

sequence data is not an option in many cases and deeper sequencing cannot recover missing data.

Herbarium specimens are an invaluable resource for improving sampling and filling gaps in the

tree of life, and we do not advocate for the exclusion of herbarium specimens in phylogenomic

datasets. Rather, we recommend careful assessment of datasets considering these artifactual

complications in assembly.

Impact of standard gene tree filtering strategies in the presence of artifactual orthologs

Gene tree filtering is a commonly applied strategy to minimize gene tree estimation error

(GTEE) and its impact on species tree estimation (Molloy and Warnow 2018). Without an

explicit method to estimate GTEE in empirical data, multiple criteria including alignment length,

the number/proportion of variable or parsimony informative sites, total tree length, the

proportion of internal branch lengths, and average bootstrap support of gene trees have all been

used as implicit proxies to account for GTEE (Leaché et al. 2014; Liu et al. 2015; Shen et al.

2016; Blom et al. 2017). It is argued that these metrics select for "higher quality" gene trees,

however, this assumption is violated in the presence of artifactual orthologs, which are

associated with higher values of many standard metrics (Fig. 2c).

Only one filtering criterion successfully reduced the proportion of artifactual orthologs in

the *Freziera* dataset: bipartition support. The high efficiency of bipartition support (Fig. 2e) is

likely due to the major topological differences between the consensus species tree and gene trees

of artifactual orthologs (Fig. 2). However, a significant proportion of putative orthologs were

also removed through this filtering mechanism. This curation strategy should be used with

caution and may require additional assessment of loci falling below the threshold if investigating

biological processes such as ILS or introgression, as some highly discordant, single-copy loci

will also be filtered. Given that the most successful methods of automated paralog detection

require assembly with HybPiper, bipartition support could provide a useful metric by which

users can remove artifactual orthologs in datasets assembled using other pipelines.

Both artifactual orthologs and putative orthologs were at removed similar levels from

datasets by the remaining four gene tree filtering criteria, in particular *tree.length* and

proportion.PI. This is consistent with our predictions about characteristics of artifactual

orthologs (Fig. 2d,e) and demonstrates that filtering by alignment or gene tree characteristics

using standard techniques may actually result in a relative increase of data artifacts in complex

phylogenomic datasets.

While gene tree filtering metrics have the potential to significantly improve phylogenetic

support (Doyle et al. 2015) and clarify relationships, it may be an inappropriate strategy for some

studies (Molloy and Warnow 2018). This is true in this case for *Freziera*, in which gene tree

filtering resulted in lower branch support and spurious topological inference for all gene tree

filtering criteria except bipartition support (Table 1, Supplementary Fig. S2). This is likely the

result of removal of a large proportion of single-copy loci relative to artifactual orthologs (Fig.

2b). The impact of the presence of artifactual orthologs and gene tree filtering strategies on

species tree inference will likely vary across datasets. We recommend careful observation of

phylogenomic datasets before applying these criteria of data curation.

Impact of data curation on phylogenetic performance in the face of artifactual orthologs

Species tree inference of *Freziera* with a two-step, coalescent-aware species tree

inference algorithm (i.e., ASTRAL-III) was relatively robust to artifactual orthology. This is

consistent with recent studies demonstrating that species tree methods are robust to the presence

of paralogs (Yan et al. 2022), though undetected paralogs have also been documented to mislead

species tree inference (Brown and Thomson 2017; Siu-Ting et al. 2019). Across analyses of our

eleven datasets, backbone relationships were largely consistent and topological differences were

primarily concentrated in a few portions of the phylogeny, at least one of which (the Candicans

group) corresponds to a rapid radiation where levels of ILS and introgression are likely high. We

found that the unfiltered dataset (i.e., *orthologs.unfiltered*) had a slightly higher proportion of

well-supported nodes, though this number was close in absolute value to those from curated

datasets with a reduced proportion of artifactual orthologs (Table 1). This may be simply a result

of a larger dataset, as coalescent-based phylogenetic inference algorithms require a large number

of independent loci to resolve challenging relationships (Leaché and Rannala 2011; Mirarab et

al. 2014a).

Reducing artifactual orthologs had an overall positive impact on phylogenetic inference

and support (Table 1; Fig. 3). Curated datasets where artifactual orthologs were removed

(*orthologs.by.eye, orthologs.HybPiper2, HybPhaser, bipartition*; Table 1) had higher normalized

quartet scores and average bootstrap support relative to the unfiltered dataset, and consistently

recovered the monophyly of the Elaphoglossifolia group via the placement of *F*.

magnibracteolata, a relationship that was not present in the unfiltered dataset (Table 1).

Improvements in these datasets are likely the result of the reduction in gene tree heterogeneity

with the removal of the noise introduced by artifactual orthologs, though gene tree heterogeneity

persists in curated datasets and likely reflects biological processes that have shaped *Freziera*'s

evolutionary history (Table 1). Ensuring that paralogs, including artifactual orthologs, are

appropriately handled in phylogenetic analyses is not only essential for accurate phylogenetic

estimation, it is also crucial to downstream analyses that require accurate branch lengths,

including divergence date estimation (Siu-Ting et al. 2019).

Orthology inference using monophyletic outgroup (*MO*) was not associated with

improved phylogenetic performance, despite being successfully applied in datasets where

paralogs are represented as multiple copy loci in assembled datasets (Morales-Briones et al.

2022). The *MO* species tree had reduced gene tree discordance (i.e., higher normalized quartet

sampling score) relative to the unfiltered dataset, likely from the removal of non-orthologous

copies. However, it failed to recover some of the major clades identified from curated data where

artifactual orthologs were removed. This may be due to information loss: the high number of

multicopy loci in our dataset resulted in many trees being extensively pruned by MO, resulting in

smaller subtrees with fewer taxa. MO is a valuable tool to identify orthologous clusters in

complex phylogenomic datasets with multi-copy paralogs and can increase phylogenetic

resolution in groups with a history of polyploidy (Morales-Briones et al. 2022); however, it

cannot be used to identify artifactual orthologs and did not improve phylogenetic performance in

Freziera, where paralogs were predominantly single-copy.

Phylogenetic relationships and introgression in Freziera

One of the barriers to the study of Neotropical diversification is the difficulty resolving

phylogenies of recent, rapid Andean radiations. Despite *Freziera*'s low species richness

compared to many cloud forest plant clades, we find similar hallmarks of explosive radiation in

our phylogenetic results. In the face of phylogenomic complexity—including the presence of

multiple copy paralogs, artifactual orthologs, rapid radiations, and DNA extracted from

herbarium specimens—relationships among species and subclades of *Freziera* were consistently

inferred across curated datasets. Most topological differences across species trees in datasets

with a reduced proportion of artifactual orthologs were within the Candicans group, a rapid

radiation with short internal branch lengths. We resolve *Freziera* into nine subclades, most of

which are moderately to well supported (Fig. 3). Despite their monophyly, these clades generally

lack morphological synapomorphies and are not geographically structured. The Humiriifolia

clade, which is sister to core *Freziera*, best represents the wide morphological diversity in each

subclade: its species have among the largest (*F. humiriifolia*) and smallest (*F. yanachagensis*)

leaves in the genus, despite occurring in close proximity in the Cordillera del Cóndor region of

southern Ecuador. While our phylogenetic backbone is generally well-supported, there are three

non-mutually exclusive biological sources that explain the very high levels of gene tree

discordance in our dataset: introgression, ILS, and gene (or genome) duplication. Despite a

genome-wide phylogenomic dataset, we are unable to pinpoint exactly when and where along the

phylogeny each of these processes has occurred. However, our resulting phylogenetic framework

provides a robust starting point from which to understand the non-bifurcating nature of

diversification of this Andean shrub clade.

Incomplete lineage sorting is common in rapid Andean radiations (Morales-Briones et al.

2018, Murillo-A. et al. 2022). Within *Freziera*, the Candicans group, especially its substituent

Calophylla clade, carries the hallmarks of ILS due to rapid radiation, including conflicting

species relationships with short branch lengths between close relatives (Fig 3a; Supplementary

Fig. S4). Not only is gene tree-species tree discordance higher in this clade compared to other

regions of *Freziera*'s phylogeny, but species relationships also differ across analyses

(Supplementary Fig. S4). Further indication of ILS in this clade is the fact that the Calophylla

clade is the most widespread of *Freziera*'s subclades and includes species with some of the

broadest distributions in the genus. These distributional patterns allow a greater possibility that

ancestral populations were large and widespread, contributing to ILS, amplifying the effect of

short times between speciation events in this group.

Even after applying data curation that may inflate support for bifurcating relationships, a

strong signal of introgression was identifiable in *Freziera*. This was evidenced by multiple

significant f4 and D statistics (Supplementary Fig. S5). Due to the limited number of unlinked

SNPs in our target capture data and the correlation of D and f4 introgression statistics when trios

share internal branches, we were not able to isolate the exact branches along which introgression

has occurred— a challenge even among the most complete datasets (Tricou et al. 2022).

However, our resulting phylogenetic framework provides a robust starting point from which to

understand the non-bifurcating nature of *Freziera*'s diversification. Areas of conflict across

species trees in our curated datasets are strong candidates for lineages that have been directly

shaped by past introgression (Fig. 3, Supplementary Fig. S4). A species that is particularly

promising for future investigation is *F. magnibracteolata*, whose unstable placement along the

backbone of *Freziera* (Table 1. Fig. 3; Supplementary Figs. S2,S4) suggests a potential history of

introgression (MacGuigan and Near 2019; Cai et al. 2021). Notably, the placement of this

species was the only major relationship to be impacted by removing artifactual orthologs relative

to the unfiltered dataset (Table 1, *Monophyletic Elaphoglossifolia group*). To further assess the

extent of gene flow and identify the branches involved in introgression events in this rapid

radiation, future research will target a much larger portion of the genome (Malinsky et al. 2021)

and include deeper taxon with multiple individuals per species.

Finally, either gene or genome duplication has resulted in paralogs in *Freziera*. We

identify both paralogs for which multiple copies are identifiable during assembly and artifactual

orthologs that are represented by only a single copy in our dataset (Fig. 2b, Supplementary Table

S3). While the processes that gave rise to these paralogs are yet to be examined in detail, it is

likely that they are product of allopolyploidy, especially considering the extensive history of

genome duplication via polyploidy in Ericales, the order to which *Freziera* belongs (Larson et al.

2020), as well as chromosome count variation within Pentaphylacaceae (Rice et al. 2015).

CONCLUSION

A major current challenge in phylogenomics is the difficulty in teasing apart specific

sources of gene tree discordance in empirical datasets, and to account for these in phylogenetic

inference (Morales-Briones et al. 2022; Tricou et al. 2022). It is a significant challenge to

accurately identify paralogs, pinpoint specific instances of introgression, disentangle incomplete

lineage sorting from historical gene flow, and reduce the impact of gene tree estimation error in a

single empirical phylogenomic dataset in which all of these sources of discordance are present.

In addition to these challenges, phylogenomic data have the potential to be very complex,

particularly for clades that are well-understood to be recalcitrant like Andean plant radiations

(Pease et al. 2016; Vargas et al. 2017). Here, we showed that careful data curation allowed us to

detect a high proportion of artifactual orthologs, which we were able to reduce with multiple,

non-mutually exclusive methods: heeding paralog warnings, removing gene trees with a high

proportion of heterozygous sites, and filtering gene trees using bipartition support. These data

curation strategies were subsequently associated with higher support, lower gene tree conflict,

and a more stable species tree— the first for an understudied tropical plant clade that previously

lacked any phylogenetic information. We advocate for the observation of empirical

phylogenomic data, including gene tree alignments and topologies, and that data curation be

tailored to unique properties of individual datasets to better address the above-mentioned

complexities in phylogenetic inference.

While commonly used filtering techniques, assembly parameters, and other automated

aspects of phylogenomics are powerful tools to improving phylogenetic inference, we have

shown that they can also increase the proportion of data artifacts (i.e., artifactual paralogs) and

have negative impacts on phylogenetic support and inference. Automated filtering techniques are

not a replacement for a deep understanding of a dataset. Selecting filtering strategies for

individual datasets should be informed by the latter since the decision is likely to be a balance

between minimizing the presence of data artifacts while maximizing the number of loci useful

for phylogenetic inference. While targeted sequence capture of universal loci offers potential,

especially for phylogenetic studies relying heavily on DNA from natural history collections,

these datasets are not without limitations related to the nature of the data themselves and to the

algorithms we use to process and analyze them. Combining exploration of datasets with deep

knowledge of the organismal biology of targeted clades is crucial toward overcoming these

limitations and inferring robust phylogenetic hypotheses.

DATA AVAILABILITY

All data generated in this study are publicly accessible. Raw sequence reads are deposited on

NCBI's Sequence Read Archive: (*web address added upon acceptance*). Supplementary

materials cited in this paper and processed data are available from the Dryad Digital Repository

(*https://datadryad.org/stash/share/X6HXB23fQmPqODdRRdXNJnGePrw29vUPbm1JgIijjFo*).

ACKNOWLEDGEMENTS

This research was funded by a Louisiana Board of Regents Research Competitiveness

Subprogram grant, the LSU College of Science and Office of Research and Economic

Development, and NSF Award DEB-2055525. We would like to thank the Missouri Botanical

Garden (MO) for their access to their collections. We thank Brant Faircloth, Matthew Johnson,

Carl Oliveros, and Jessie Salter for their guidance in library preparation, and Brant Faircloth for

access to laboratory equipment. Computational analyses were performed on LSU High

Performance Computing's SuperMike cluster. We thank three anonymous reviewers for their

comments and criticism, which greatly improved previous versions of this work. This manuscript

benefited from feedback from Laymon Ball, Janet Mansaray, and Diego Paredes-Burneo. The

taxonomic expertise of Daniel Santamaría-Aguilar was critical throughout the design and

implementation of this research.

REFERENCES

Anisimova M., Gil M., Dufayard J.-F., Dessimoz C., Gascuel O. 2011. Survey of branch support

methods demonstrates accuracy, power, and robustness of fast likelihood-based

approximation schemes. Syst. Biol. 60:685–699.

Bakker F.T., Lei D., Yu J., Mohammadin S., Wei Z., van de Kerke S., Gravendeel B.,

Nieuwenhuis M., Staats M., Alquezar-Planas D.E., Holmer R. 2015. Herbarium genomics:

plastome sequence assembly from a range of herbarium specimens using an Iterative

Organelle Genome Assembly pipeline. Biol. J. Linn. Soc. Lond. 117:33–43.

Benjamini Y., Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. J. R. Stat. Soc. Series B Stat. Methodol. 57:289–

300.

Blom M.P.K., Bragg J.G., Potter S., Moritz C. 2017. Accounting for uncertainty in gene tree

estimation: summary-coalescent species tree inference in a challenging radiation of

Australian lizards. Syst. Biol. 66:352–366.

Bolger A.M., Lohse M., Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence

data. Bioinformatics. 30:2114–2120.

Borowiec M.L. 2016. AMAS: a fast tool for alignment manipulation and computing of summary

statistics. PeerJ. 4:e1660.

Bronner I.F., Quail, M.A. 2019. Best practices for Illumina library preparation. *Curr. Prot. Hum.*

Genet. s 102:e86.

Brown J.M., Thomson R.C. 2017. Bayes Factors Unmask Highly Variable Information Content,

Bias, and Extreme Influence in Phylogenomic Analyses. Syst. Biol. 66:517–530.

Brown J.W., Walker J.F., Smith S.A. 2017. Phyx: phylogenetic tools for unix. Bioinformatics.

33:1886–1888.

Cai L., Xi Z., Lemmon E.M., Lemmon A.R., Mast A., Buddenhagen C.E., Liu L., Davis C.C.

2021. The Perfect Storm: Gene Tree Estimation Error, Incomplete Lineage Sorting, and

Ancient Gene Flow Explain the Most Recalcitrant Ancient Angiosperm Clade,

Malpighiales. Syst. Biol. 70:491–507.

Danecek P., Auton A., Abecasis G., Albers C.A., Banks E., DePristo M.A., Handsaker R.E.,

Lunter G., Marth G.T., Sherry S.T., McVean G., Durbin R., 1000 Genomes Project Analysis

Group. 2011. The variant call format and VCFtools. Bioinformatics. 27:2156–2158.

Doyle V.P., Young R.E., Naylor G.J.P., Brown J.M. 2015. Can We Identify Genes with

Increased Phylogenetic Reliability? Syst. Biol. 64:824–837.

Faircloth B.C. 2013. Illumiprocessor: a trimmomatic wrapper for parallel adapter and quality

trimming. http://dx.doi.org/10.6079/J9ILL.

Faircloth B.C. 2016. PHYLUCE is a software package for the analysis of conserved genomic

loci. Bioinformatics. 32:786–788.

Gardner E.M., Johnson M.G., Pereira J.T., Puad A.S.A., Arifiani D., Wickett N.J., Zerega N.J.C.

2020. Paralogs and off-target sequences improve phylogenetic resolution in a densely-

sampled study of the breadfruit genus (Artocarpus, Moraceae). Syst. Biol. 70:558–575.

Gentry A.H. 1982. Neotropical floristic diversity: phytogeographical connections between

Central and South America, Pleistocene climatic fluctuations, or an accident of the Andean

orogeny? Ann. Mo. Bot. Gard. 69:557–593.

Guindon S., Dufayard J.-F., Lefort V., Anisimova M., Hordijk W., Gascuel O. 2010. New

algorithms and methods to estimate maximum-likelihood phylogenies: assessing the

performance of PhyML 3.0. Syst. Biol. 59:307–321.

Hale H., Gardner E.M., Viruel J., Pokorny L., Johnson M.G. 2020. Strategies for reducing per-

sample costs in target capture sequencing for phylogenomics and population genomics in

plants. Appl. Plant Sci. 8:e11337.

Hughes C.E. 2016. The tropical Andean plant diversity powerhouse. New Phytol. 210:1152–

1154.

Hughes C.E., Eastwood R. 2006. Island radiation on a continental scale: exceptional rates of

plant diversification after uplift of the Andes. Proc. Natl. Acad. Sci. U. S. A. 103:10334–

10339.

Johnson M.G., Gardner E.M., Liu Y., Medina R., Goffinet B., Shaw A.J., Zerega N.J.C., Wickett

N.J. 2016. HybPiper: Extracting coding sequence and introns for phylogenetics from high-

throughput sequencing reads using target enrichment. Appl. Plant Sci. 4:1600016.

Johnson M.G., Pokorny L., Dodsworth S., Botigué L.R., Cowan R.S., Devault A., Eiserhardt

W.L., Epitawalage N., Forest F., Kim J.T., Leebens-Mack J.H., Leitch I.J., Maurin O., Soltis

D.E., Soltis P.S., Wong G.K.-S., Baker W.J., Wickett N.J. 2019. A universal probe set for

targeted sequencing of 353 nuclear genes from any flowering plant designed using k-

Medoids clustering. Syst. Biol. 68:594–606.

Kalyaanamoorthy S., Minh B.Q., Wong T.K.F., von Haeseler A., Jermiin L.S. 2017.

ModelFinder: fast model selection for accurate phylogenetic estimates. Nat. Methods.

14:587–589.

Lagomarsino L.P., Condamine F.L., Antonelli A., Mulch A., Davis C.C. 2016. The abiotic and

biotic drivers of rapid diversification in Andean bellflowers (Campanulaceae). New Phytol.

Lagomarsino L.P., Frankel L., Uribe-Convers S., Antonelli A., Muchhala N. 2022. Increased

resolution in the face of conflict: phylogenomics of the Neotropical bellflowers

(Campanulaceae: Lobelioideae), a rapid plant radiation. Ann. Bot. 129:723–736.

Larson D.A., Walker J.F., Vargas O.M., Smith S.A. 2020. A consensus phylogenomic approach

highlights paleopolyploid and rapid radiation in the history of Ericales. Am. J. Bot.

107:773–789.

Leaché A.D., Rannala B. 2011. The accuracy of species tree estimation under simulation: a

comparison of methods. Syst. Biol. 60:126–137.

Leaché A.D., Wagner P., Linkem C.W., Böhme W., Papenfuss T.J., Chong R.A., Lavin B.R.,

Bauer A.M., Nielsen S.V., Greenbaum E., Rödel M.-O., Schmitz A., LeBreton M., Ineich I.,

Chirio L., Ofori-Boateng C., Eniang E.A., Baha El Din S., Lemmon A.R., Burbrink F.T.

2014. A hybrid phylogenetic–phylogenomic approach for species tree estimation in African

Agama lizards with applications to biogeography, character evolution, and diversification.

Mol. Phylogenet. Evol. 79:215–230.

Liu L., Xi Z., Wu S., Davis C.C., Edwards S.V. 2015. Estimating phylogenetic trees from

genome-scale data. Ann. N. Y. Acad. Sci. 1360:36–53.

Li Z., Barker M.S. 2020. Inferring putative ancient whole-genome duplications in the 1000

Plants (1KP) initiative: access to gene family phylogenies and age distributions.

Gigascience. 9.

Li Z., McKibben M.T.W., Finch G.S., Blischak P.D., Sutherland B.L., Barker M.S. 2021.

Patterns and Processes of Diploidization in Land Plants. Annu. Rev. Plant Biol. 72:387–410.

MacGuigan D.J., Near T.J. 2019. Phylogenomic Signatures of Ancient Introgression in a Rogue

Lineage of Darters (Teleostei: Percidae). Syst. Biol. 68:329–346.

Madriñán S., Cortés A.J., Richardson J.E. 2013. Páramo is the world's fastest evolving and

coolest biodiversity hotspot. Front. Genet. 4:192.

Mai U., Mirarab S. 2018. TreeShrink: fast and accurate detection of outlier long branches in

collections of phylogenetic trees. BMC Genomics. 19:272.

Malinsky M., Matschiner M., Svardal H. 2021. Dsuite - Fast D-statistics and related admixture

evidence from VCF files. Mol. Ecol. Resour. 21:584–595.

Malinsky M., Svardal H., Tyers A.M., Miska E.A., Genner M.J., Turner G.F., Durbin R. 2018.

Whole-genome sequences of Malawi cichlids reveal multiple radiations interconnected by

gene flow. Nat Ecol Evol. 2:1940–1955.

McKain M.R., Johnson M.G., Uribe-Convers S., Eaton D., Yang Y. 2018. Practical

considerations for plant phylogenomics. Appl. Plant Sci. 6:e1038.

Mirarab S., Bayzid M.S., Warnow T. 2014a. Evaluating Summary Methods for Multilocus

Species Tree Estimation in the Presence of Incomplete Lineage Sorting. Syst. Biol. 65:366–

380.

Mirarab S., Reaz R., Bayzid M.S., Zimmermann T., Swenson M.S., Warnow T. 2014b.

ASTRAL: genome-scale coalescent-based species tree estimation. Bioinformatics. 30:i541–

8.

Molloy E.K., Warnow T. 2018. To Include or Not to Include: The Impact of Gene Filtering on

Species Tree Estimation Methods. Systematic Biology. 67:285–303.

Mongiardino Koch N. 2021. Phylogenomic Subsampling and the Search for Phylogenetically

Reliable Loci. Mol. Biol. Evol. 38:4025–4038.

Morales-Briones D.F., Gehrke B., Huang C.-H., Liston A., Ma H., Marx H.E., Tank D.C., Yang

Y. 2021a. Analysis of paralogs in target enrichment data pinpoints multiple ancient

polyploidy events in Alchemilla s.L. (Rosaceae). Syst. Biol. 71:190–207.

Morales-Briones D.F., Kadereit G., Tefarikis D.T., Moore M.J., Smith S.A., Brockington S.F.,

Timoneda A., Yim W.C., Cushman J.C., Yang Y. 2021b. Disentangling Sources of Gene

Tree Discordance in Phylogenomic Data Sets: Testing Ancient Hybridizations in

Amaranthaceae s.l. Syst. Biol. 70:219–235.

Morales-Briones D.F., Liston A., Tank D.C. 2018. Phylogenomic analyses reveal a deep history

of hybridization and polyploidy in the Neotropical genus *Lachemilla* (Rosaceae). New

Phytol. 218:1668–1684.

Murillo-A J., Valencia-D J., Orozco C. I., Parra-O C., Neubig, K. M. 2022. Incomplete lineage

sorting and reticulate evolution mask species relationships in Brunelliaceae, an Andean

family with rapid, recent diversification. *Am. J. Bot.* 109:1139-1156.

Mutke J., Barthlott W. 2005. Patterns of vascular plant diversity at continental to global scales.

Biol. Skr. 55:521–531.

Nauheimer L., Weigner N., Joyce E., Crayn D., Clarke C., Nargar K. 2021. HybPhaser: A

workflow for the detection and phasing of hybrids in target capture data sets. Appl. Plant

Sci. 9.

Nguyen L.-T., Schmidt H.A., von Haeseler A., Minh B.Q. 2015. IQ-TREE: a fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. Mol. Biol. Evol.

32:268–274.

Ogilvie H.A., Bouckaert R.R., Drummond A.J. 2017. StarBEAST2 brings faster species tree

inference and accurate estimates of substitution rates. Mol. Biol. Evol. 34:2101–2114.

One Thousand Plant Transcriptomes Initiative. 2019. One thousand plant transcriptomes and the

phylogenomics of green plants. Nature. 574:679–685.

Paradis E., Schliep K. 2019a. ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. Bioinformatics. 35:526–528.

Paradis E., Schliep K. 2019b. ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. Bioinformatics. 35:526–528.

Pease J.B., Haak D.C., Hahn M.W., Moyle L.C. 2016. Phylogenomics Reveals Three Sources of

Adaptive Variation during a Rapid Radiation. PLoS Biol. 14:e1002379.

Puritz J.B., Hollenbeck C.M., Gold J.R. 2014. dDocent: a RADseq, variant-calling pipeline

designed for population genomics of non-model organisms. PeerJ. 2:e431.

Rabiee M., Mirarab S. 2020. Forcing external constraints on tree inference using ASTRAL.

BMC Genomics. 21:218.

Rambaut. 2014. FigTree v1. 4.2, a graphical viewer of phylogenetic trees. //tree. bio. ed. ac.

uk/software/figtree/right angle bracket.

Rice A., Glick L., Abadi S., Einhorn M., Kopelman N.M., Salman-Minkov A., Mayzel J., Chay

O., Mayrose I. 2015. The Chromosome Counts Database (CCDB) - a community resource

of plant chromosome numbers. New Phytol. 206:19–26.

Santamaría-Aguilar D., Monro A.K. 2019. Compendium of *Freziera* (Pentaphylacaceae) of

South America including eleven new species and the typification of 22 names. Kew Bull.

74:14.

Shen X.-X., Salichos L., Rokas A. 2016. A genome-scale investigation of how sequence,

function, and tree-based gene properties influence phylogenetic inference. Genome Biol.

Evol. 8:2565–2580.

Siu-Ting K., Torres-Sánchez M., San Mauro D., Wilcockson D., Wilkinson M., Pisani D.,

O'Connell M.J., Creevey C.J. 2019. Inadvertent Paralog Inclusion Drives Artifactual

Topologies and Timetree Estimates in Phylogenomics. Mol. Biol. Evol. 36:1344–1356.

Smith M.L., Hahn M.W. 2021a. New approaches for inferring phylogenies in the presence of

paralogs. Trends Genet. 37:174–187.

Smith M.L., Hahn M.W. 2021b. The frequency and topology of pseudoorthologs.

bioRxiv.:2021.02.17.431499.

Smith S.A., Brown J.W., Walker J.F. 2018. So many genes, so little time: A practical approach

to divergence-time estimation in the genomic era. PLoS One. 13:e0197433.

Solís-Lemus C., Bastide P., Ané C. 2017. PhyloNetworks: a package for phylogenetic networks.

Mol. Biol. Evol. 34:3292–3298.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of

large phylogenies. Bioinformatics. 30:1312–1313.

Štorchová H., Hrdličková R., Chrtek J. Jr, Tetera M., Fitze D., Fehrer J. 2000. An improved

method of DNA isolation from plants collected in the field and conserved in saturated

NaCl/CTAB solution. Taxon. 49:79–84.

Tricou T., Tannier E., de Vienne D.M. 2022. Ghost Lineages Highly Influence the Interpretation

of Introgression Tests. Syst. Biol. 71:1147–1158.

Tsou C.-H., Li L., Vijayan K. 2016. The intra-familial relationships of Pentaphylacaceae s.l. as

revealed by DNA sequence analysis. Biochem. Genet. 54:270–282.

Ulloa Ulloa C., Acevedo-Rodríguez P., Beck S., Belgrano M.J., Bernal R., Berry P.E., Brako L.,

Celis M., Davidse G., Forzza R.C., Gradstein S.R., Hokche O., León B., León-Yánez S.,

Magill R.E., Neill D.A., Nee M., Raven P.H., Stimmel H., Strong M.T., Villaseñor J.L.,

Zarucchi J.L., Zuloaga F.O., Jørgensen P.M. 2017. An integrated assessment of the vascular

plant species of the Americas. Science. 358:1614–1617.

Vargas O.M., Ortiz E.M., Simpson B.B. 2017. Conflicting phylogenomic signals reveal a pattern

of reticulate evolution in a recent high-Andean diversification (Asteraceae: Astereae:

Diplostephium). New Phytol. 214:1736–1750.

Weitzman A.L., Dressler S., Stevens P.F. 2004. Ternstroemiaceae. In: Kubitzki K., editor.

Flowering Plants. Dicotyledons: Celastrales, Oxalidales, Rosales, Cornales, Ericales. Berlin,

Heidelberg: Springer Berlin Heidelberg. p. 450–460.

Yang Y., Smith S.A. 2014. Orthology inference in nonmodel organisms using transcriptomes

and low-coverage genomes: improving accuracy and matrix occupancy for phylogenomics.

Mol. Biol. Evol. 31:3081–3092.

Yan Z., Smith M.L., Du P., Hahn M.W., Nakhleh L. 2022. Species Tree Inference Methods

Intended to Deal with Incomplete Lineage Sorting Are Robust to the Presence of Paralogs.

Syst. Biol. 71:367–381.

Zhang C., Rabiee M., Sayyari E., Mirarab S. 2018. ASTRAL-III: polynomial time species tree

reconstruction from partially resolved gene trees. BMC Bioinformatics. 19:153.

Zhang Z., Xie P., Guo Y., Zhou W., Liu E., Yu Y. 2022. Easy353: A Tool to Get

Angiosperms353 Genes for Phylogenomic Research. Mol. Biol. Evol. 39.

FIGURE AND TABLE CAPTIONS

FIGURE 1. Diversity and distribution of *Freziera*. *Freziera* has significant variation in leaf

morphology and pubescence, as illustrated by branches of a) *F. candicans* and b) *Freziera sp.*;

meanwhile, c) flower and d) fruit morphology are relatively stable. *Freziera* is an Andean

radiation; e) most species are distributed in this mountain chain in western South America, with

some species in Central America, the Guiana Shield, the Caribbean, and the Atlantic Forest of

Brazil. Photos by L. Lagomarsino.

932
FIGURE 2. Exemplary gene trees and alignment subsets for a) a putative ortholog, and b) an

artifactual ortholog. Putative ortholog gene trees exhibit monophyletic genera for outgroups and

appropriate relationships to the ingroup, a relatively shallow backbone in the ingroup, and

relationships similar to the preliminary species tree. A relatively low level of variation is present

in alignments for putative orthologs, which is fitting of a locus in a universal probe set being

applied within a genus. Artifactual ortholog gene trees often have deep divergences between

subsets of the ingroup, including between samples from the same species. Within those

subclades, patterns of relationships from the preliminary species tree are repeated. In some cases,

as in b), outgroups are also non-monophyletic and subsets of outgroup samples are recovered as

sister to the ingroup subclades. Alignments for artifactual orthologs exhibit relatively high

variation with observable motifs between the suspected copies, as indicated by the black and

white dots. c) Overlap between artifactual orthologs identified by-eye, and paralog warnings

raised by automated paralogy detection strategies. d) Expectations for informative, single-copy

orthologs change for many gene tree filtering metrics when data sets include artifactual

orthologs; arrows indicate whether expected values for metrics are high or low. e) Impact of gene

tree filtering strategies on removal of artifactual and putative orthologs as identified by-eye. The

dashed lines indicate the various thresholds above which gene trees were kept in filtered datasets,

as indicated in Table 1.

951
FIGURE 3. Names of clades consistently recovered and species tree topologies for a) The

consensus tree of the four datasets in which the proportion of artifactual orthologs was reduced

(*bipartition*, *by.eye*, *HybPhaser*, *orthologs.HybPiper.no.warnings*; left) versus the

orthologs.unfiltered species tree (right). The placement of *F. magnibracteolata*– the major

topological conflict between the two trees– is highlighted with the blue line connecting the tip

labels in each. Branches with full support (LPP from ASTRAL = 1) are indicated by an asterisk;

elsewhere, the LPP * 100 is provided. Support for each of the four datasets in the consensus tree

are given clockwise from the top left: *bipartition*, *by.eye*, *HybPhaser*,

orthologs.HybPiper.no.warnings. b) Cartoon trees displaying the most frequent differences in

relationships between clades recovered by different datasets and the consensus tree of all

datasets. Cartoons depict one topological scenario but not the only topological outcome.

963
TABLE 1. Descriptions of the filtering criteria used, names applied to each of the datasets,

number of loci removed and selected by each, and the impact of filtering criteria on gene tree

discordance, branch support, and species tree topology. Values for datasets that performed

equally or better than the orthologs.unfiltered dataset for each metric are bolded. Major

topological conflicts between species trees and a consensus topology of the 11 filtered datasets

(Fig. 3b) are shown, including instances where the species tree is compatible (+) or in

disagreement (-) with the consensus topology. Datasets from which the Elaphoglossifolia group

was not resolved as monophyletic are indicated with "n/a". Asterisks mark disagreement from

the alternative placement of only one species. The "Branching order of the Elaphoglossifolia

group" includes conflict in the placement of F. magnibracteolata.

974
975   ONLINE APPENDIX 1. Supplementary Materials and Methods

976
977   ONLINE APPENDIX 2. Paralogs identified by HybPhaser for all samples and for each sample.

978
979   SUPPLEMENTARY FIGURE S1. Heat map showing gene recovery efficiency for all submitted

980   samples (rows) across the 353 targeted genes (columns).

981
982   SUPPLEMENTARY FIGURE S2. ASTRAL-III species trees for the 11 datasets: a)

983   *orthologs.unfiltered*, b) *orthologs.by.eye*, c) *orthologs.HybPiper.long*, d)

984   *orthologs.HybPiper.no.warnings*, e) *HybPhaser*, f) *MO*, g) *proportion.PI*, h)

985   *proportion.internal.branch.lengths*, i) *average.BS*, j) *tree.length*, k) *bipartition*

986
987   SUPPLEMENTARY FIGURE S3. Number of reads mapped to target regions in silica (blue) and

988   herbarium (red) samples. There is a correlation between the number of reads mapped and the

989   number of paralog warnings from HybPiper in a sample ($R^2 = 0.55$, $F(df=1) = 123.6$, $p < 0.0001$).

990   Assembly and paralog detection are improved with increased sequencing effort.

991
SUPPLEMENTARY FIGURE S4. Tanglegram plots showing pairwise comparisons between

ASTRAL-III species tree topologies for a) *consensus* vs. *orthologs.unfiltered*, b) *consensus* vs.

orthologs.by.eye, c) *consensus* vs. *orthologs.HybPiper2.long*, d) *consensus* vs.

orthologs.HybPiper2.no.warnings, e) *consensus* vs. *HybPhaser*, f) *consensus* vs. *MO*, g)

consensus vs. *proportion.PI*, h) *consensus* vs. *proportion.internal.branch.lengths*, i) *consensus*

vs. *average.BS*, j) *consensus* vs. *tree.length*, k) *consensus* vs. *bipartition*, l) *orthologs.by.eye* vs.

orthologs.unfiltered, m) *orthologs.by.eye* vs. *orthologs.HybPiper2.long*, n) *orthologs.by.eye* vs.

orthologs.HybPiper2.no.warnings, o) *orthologs.by.eye* vs. *HybPhaser*, p) *orthologs.by.eye* vs.

MO, q) *orthologs.by.eye* vs. *proportion.PI*, r) *orthologs.by.eye* vs.

proportion.internal.branch.lengths, s) *orthologs.by.eye* vs. *average.BS*, t) *orthologs.by.eye* vs.

tree.length, u) *orthologs.by.eye* vs. *bipartition*, v) *orthologs.HybPiper2.no.warnings* vs.

orthologs.unfiltered, w) *orthologs.HybPiper2.no.warnings* vs. *orthologs.HybPiper2.long*, x)

orthologs.HybPiper2.no.warnings vs. *HybPhaser*, y) *orthologs.HybPiper2.no.warnings* vs. MO,

z) *orthologs.HybPiper2.no.warnings* vs. *proportion.PI*, aa) *orthologs.HybPiper2.no.warnings* vs.

proportion.internal.branch.lengths, bb) *orthologs.HybPiper2.no.warnings* vs. *average.BS*, cc)

orthologs.HybPiper2.no.warnings vs. *tree.length*, dd) *orthologs.HybPiper2.no.warnings* vs.

bipartition, ee) *HybPhaser* vs. *orthologs.unfiltered*, ff) *HybPhaser* vs. *orthologs.HybPiper2.long*,

gg) *HybPhaser* vs. MO, hh) *HybPhaser* vs. *proportion.PI*, ii) *HybPhaser* vs.

proportion.internal.branch.lengths, jj) *HybPhaser* vs. *average.BS*, kk) *HybPhaser* vs.

tree.length, ll) *HybPhaser* vs. *bipartition*, mm) *bipartition* vs. *orthologs.unfiltered*, nn)

bipartition vs. *orthologs.HybPiper2.long*, oo) *bipartition* vs. *MO*, pp) *bipartition* vs.

proportion.PI, qq) *bipartition* vs. *proportion.internal.branch.lengths*, rr) *bipartition* vs.

average.BS, ss) *bipartition* vs. *tree.length*, tt) *orthologs.HybPiper2.long* vs. *orthologs.unfiltered*,

uu) *MO* vs. *orthologs.unfiltered*, vv) *proportion.PI* vs. *orthologs.unfiltered*, xx)

proportion.internal.branch.lengths vs. *orthologs.unfiltered*, yy) *average.BS* vs.

orthologs.unfiltered, zz) *tree.length* vs. *orthologs.unfiltered.*

1018

1019   SUPPLEMENTARY FIGURE S5. Results of D and f4 statistics calculated with Dsuite on the

1020   "bipartition support", "by eye", and "HybPiper2 no warnings" datasets. The columns indicate 1)

1021   values of Patterson's D-statistic for evaluated trios, 2) highly significant p-values associated with

1022 the D-statistics after Benjamini-Hochberg correction, and 3) f4 ratio statistics. Introgression is

1023 detected across all datasets even after assessing family-wise error rate.

1024
1025 SUPPLEMENTARY FIGURE S6. Scatterplot showing locus heterozygosity (proportion of loci with

1026 SNPs) versus allele divergence calculated with HybPhaser for all included accessions.

1027
1028 SUPPLEMENTARY TABLE S1. Per sample data--species name, voucher information and herbarium

1029 code for tissue gathered from specimens (codes follow Index Herbariorum:

1030 http://sweetgum.nybg.org/science/ih/), collection date, provenance, and sample ID used in

1031 phylogenetic analyses--and summary statistics generated with HybPiper v.1.3.1.

1032
1033 SUPPLEMENTARY TABLE S2. HybPiper v.2.0.1 assembly statistics for all samples included in this

1034 study. Paralog warnings per individual are shown.

1035
1036 SUPPLEMENTARY TABLE S3. Paralog identification results from the by-eye and automated

1037 strategies per locus. The proportion of stitched contigs, paralog warning by depth, and long

1038 paralog warnings from HybPiper v.2.0.1, paralog warnings from HybPiper v.1.3.1, removed loci

1039 with HybPhaser as well as per locus statistics for alignments and gene trees used to explore how

1040 standard gene tree filtering affects the presence of artifactual orthologs in our dataset are shown.

1041 Details of by-eye identification of paralogs, and the handling of reports from HybPiper v,2.0.1

1042 are in Appendix 1.

1043
1044 SUPPLEMENTARY TABLE S4. HybPiper v.2.0.1 paralog report from assembly using a taxon-

1045 specific target file generated with Easy353. Reference sequences were extracted from

1046 transcriptome data of Ternstroemia gymnanthera (Pentaphylacaceae). Numbers on the table

1047 correspond to the number of contigs assembled per locus.

1048

1049   SUPPLEMENTARY TABLE S5. Per sample summary statistics from HybPhaser, including the

1050   recovered sequence length (bp), the recovered sequence length as the proportion of the target

1051   sequences from HybPiper (bpoftarget), number of paralogs found in all samples, number of

1052   additional paralogs found for each sample, number of loci recovered, allele divergence, locus

1053   heterozygosity, loci with >0.5% SNPs, loci with >1% SNPs, loci with >2% SNPs