We get by with a little help from our friends: shared adaptive variation provides a bridge to novel ecological specialists during adaptive radiation

Emilie J. Richards[1,2] ,Christopher H. Martin[1,2]

[1]Department of Integrative Biology, University of California, Berkeley

[2]Museum of Vertebrate Zoology, University of California, Berkeley

Correspondence: *chmartin@berkeley.edu*

Keywords: ecological speciation, adaptive radiation, novelty, speciation genomics, adaptive introgression, trophic innovation

Abstract

Adaptive radiations involve astounding bursts of phenotypic, ecological, and species diversity. However, the microevolutionary processes that underlie the origins of these bursts are still poorly understood. We report the discovery of an intermediate 'wide-mouth' scale-eating ecomorph in a sympatric radiation of *Cyprinodon* pupfishes, illuminating the transition from a widespread algae-eating generalist to a novel microendemic scale-eating specialist. We first show that this ecomorph occurs in sympatry with generalist *C. variegatus* and scale-eating specialist *C. desquamator* on San Salvador Island, Bahamas, but is genetically differentiated, morphologically distinct, and often consumes scales. We then compared the timing of selective sweeps on shared and unique adaptive variants in trophic specialists to characterize their adaptive walk. Shared adaptive regions swept first in both the specialist *desquamator* and the intermediate 'wide-mouth' ecomorph, followed by unique sweeps of introgressed variation in 'wide-mouth' and de novo variation in *desquamator*. The two scale-eating populations additionally shared 9% of their hard selective sweeps with molluscivores *C. brontotheroides*, despite no single common ancestor among specialists. Our work provides a new microevolutionary framework for investigating how major ecological transitions occur and illustrates how both shared and unique genetic variation can provide a bridge for multiple species to access novel ecological niches.

Introduction

Rapid bursts of diversification and repeated bouts of speciation like those seen in adaptive

radiations contradict current mechanistic speciation models that predict diversification should

slow with time as available niche space becomes increasingly subdivided and disruptive

selection becomes weaker with each recurrent speciation event (e.g. [1–3]). Diversification on

complex adaptive landscapes with multiple empty fitness peaks corresponding to different niches

provides an alternative mechanism to niche subdivision [4–6]. However, these landscapes

present a new problem to our mechanistic understanding of adaptive radiations: How do

populations manage to escape local optima, cross fitness valleys, and access new fitness peaks

[7–10]? Colonizing new fitness peaks on the adaptive landscape presents challenges because it

requires transitions in behaviors, morphological traits, or a combination of the two that allow

organisms to adapt to new ecological niches [11]. Spectacular ecological transitions do often

occur during adaptive radiation, such as blood-drinking [12] or plant carnivory [13,14], yet it is

still poorly understood how such seemingly discontinuous transitions occur.

Recent conceptual frameworks for understanding adaptation to novel fitness peaks

suggest that these major ecological transitions likely occur in stages of potentiation, actualization

and refinement [15,16]. The initial emergence of a novel trait likely requires further refinement

to become successfully incorporated into the functional ecology of an organism. Several

experimental lab studies suggest that novel ecological transitions are highly contingent on

accruing a series of mutations that incrementally refine adaptations to colonize new fitness peaks

[16,17]. This idea that genetic background is important in setting the stage for adaptation also

underlies many hypotheses for adaptive radiation, such as the hybrid swarm and syngameon

hypotheses – in which radiations are driven by acquiring novel combinations of alleles through

the exchange of genetic variation either from distinct lineages outside the radiation or within the

radiation itself [18]. However, we are only just beginning to explore how gene flow and shared

genetic variation gives recipient lineages access to new fitness peaks in the wild and generates

adaptive radiations [6].

An adaptive radiation of trophic specialist pupfishes on San Salvador Island (SSI) in the

Bahamas is an excellent system for understanding how the rapid evolution of major ecological

transitions occurs in nature. This radiation contains a widespread generalist pupfish species

(*Cyprinodon variegatus*) that occurs in sympatry with two previously described trophic

specialists that are endemic to the hypersaline lakes on the island: a molluscivore (*C.*

bronotheroides) with a novel nasal protrusion which is an oral-sheller of gastropods [19] and a

scale-eating specialist (*C. desquamator*) with two-fold larger oral jaws [20]. The evolutionary

novelties in this system originated recently; the lakes on SSI were dry during the last glacial

maximum 6-20 kya years ago [21,22]. Intriguingly, we recently discovered a fourth species of

pupfish living in sympatry with the two specialists and generalist on SSI [23]. This species

exhibits intermediate jaw morphology between *C. desquamator* and *C. variegatus* (figure 1).

Here we refer to this new ecomorph as the 'wide-mouth' because its mouth is wider than any

other species in the radiation. The multi-peak fitness landscape driving this radiation suggests

that *C. desquamator* is isolated by a large fitness valley from *C. variegatus* and *C.*

brontotheroides [9] and this intermediate 'wide-mouth' may provide clues about the

microevolutionary processes underlying how the observed novel fitness peaks are traversed in

the wild.

Here we first investigate the position of the 'wide-mouth' on the ecological spectrum

from generalist to scale-eating specialist using a combination of morphological, behavioral,

dietary, and genomic data. We then estimated the demographic history of the 'wide-mouth' and

explored the spatial origins and timing of selection on shared and unique genetic variation

involved in adaptation to scale-eating to better understand this ecological transition. Our results

suggest that while intermediate in jaw length known to be relevant for the highly specialized

scale-eater *C. desquamator*, *C sp.* 'wide-mouth' demonstrates transgressive morphology and a

distinct genetic background. Our investigation of the timing of selection and genetic origins of

the adaptive alleles shared and unique between the two scale-eating species indicates divergent

adaptive walks that are highly dependent on their genetic background. Despite shared origins,

access to unique genetic variation in each of the two scale-eating sister species likely resulted in

distinct adaptive walks and ultimately contributed to the diversity of ecological specialists

observed in this radiation.

Figure 1. The San Salvador Island radiation of pupfish**.** Pie charts indicate the presence of

sympatric *Cyprinodon* species in each lake and are color-coded with representative pictures of

generalist *C. variegatus* (gold), recently discovered *C.* sp. 'wide-mouth' ecomorph (red-orange)

with intermediate jaws, scale-eater *C. desquamator* (teal) with the largest oral jaws, and

molluscivore *C. brontotheroides* (purple) with characteristic nasal protrusion. Labeled lakes

contain all known *C.* sp. 'wide-mouth' populations sampled for this study. Satellite image from

Google Earth.

Methods

Ecological and morphological characterization of 'wide-mouth' scale-eater

C. variegatus, C. desquamator, and *C.* sp. 'wide-mouth' individuals from 3 lake populations

(Osprey Lake, Great Lake, and Oyster Pond) in which we had sufficient specimens (n=84; *C. brontotheroides* not shown) were measured for 9 external morphological traits using digital

calipers. Traits were selected for specific connections to foraging performance which differed

across the three species in a previous study [9]. We also characterized diet for *C. variegatus, C.*

desquamator, and 'wide-mouth' in Osprey Lake from stomach content analyses (*n*=10 per

species) and stable isotope analyses of muscle tissue from wild-collected samples (*n*=75).

Dietary overlap was characterized by comparison of population mean scale count from gut

contents using ANOVA and ellipse areas and bivariate means on isotope biplots using SIBER

[24]. See supplemental methods for more details on sample sizes and analyses.

Genomic library preparation and variant filtration

To explore the evolutionary history of *C.* sp. 'wide-mouth', we sequenced whole genomes of 24

individuals following protocols used in a previous study [25] that included genomes from *C.*

variegatus, C. desquamator, and *C. brontotheroides*. Our final genetic dataset after filtering

contained 6.4 million variants across 110 individuals from the four species (7.9x median

coverage). See supplemental methods for the full sequencing and genotyping protocol.

Genomic origins of the 'wide-mouth' scale-eater

We first tested whether these *C*. sp. 'wide-mouth' individuals represented recent (e.g. F1/F2)

hybrids of *C. variegatus* and *C. desquamator* in the wild using principal component and

ADMIXTURE analyses to look for the genome-wide pattern expected in PCAs when recent

hybrids between two populations are included. We also used formal tests for introgression and

admixed populations, f_3 and f_4-statistics [26], to assess whether 'wide-mouth' are the byproduct

of recent admixture. Finally, we used *fastsimcoal2* (v2.6.0.3;[27]), a demographic modeling

approach based on the folded minor allele frequency spectrum (mSFS), to discriminate among

alternative evolutionary scenarios for the origin of 'wide-mouth' and estimated divergence times

among all four species based on the best model fit from an AIC test (see supplementary methods

for more detail).

<u>Characterization of unique and shared adaptive alleles among specialists</u>

Across all four populations in Osprey Lake, we looked for regions that showed evidence of a

hard selective sweep using SweeD (v.3.3.4;[28]). The composite likelihood ratio (CLR) for a

hard selective sweep was calculated in 50-kb windows across scaffolds that were at least 100-kb

in length (99 scaffolds; 85.6% of the genome). Significance thresholds were determined using

CLR values from neutral sequences simulated under MSMC inferred demographic scenarios of

historical effective population size changes (Supplemental methods; figure S1; table S3).

Next, we searched for candidate adaptive alleles associated with species divergence by

overlapping selective sweep regions with regions of high genetic divergence based on fixed or

nearly fixed SNPs between species. We chose to also look at regions with nearly fixed SNPs (F_{st}

≥ 0.95) to accommodate ongoing gene flow between these young species. F_{st} between the

populations and species was calculated per variant site using –weir-pop-fist function in vcftools

(v.0.1.15;[29]).

<u>Timing of selection on candidate adaptive alleles</u>

We also determined the relative age of candidate adaptive alleles by generating estimates of

coalescent times using starTMRCA (v0.6.1;[30]). For each candidate adaptive allele that was

unique to the three specialists and the 16 shared alleles between *C. desquamator* and 'wide-

mouth', a 1-Mb window surrounding the variant was extracted into separate vcfs for each

species. These sets of variants were then analyzed in starTMRCA with a mutation rate of 1.56 x

10^{-8} substitutions per base pair (from Caribbean pupfishes [25]) and a recombination rate of

3.11×10^{-8} (from stickleback; [31]). Each analysis was run three times per focal adaptive allele

and all runs were checked for convergence between and within runs. Most runs rapidly

converged within the initial 6000 steps, but 5 runs did not converge after an additional 4000

steps and were discarded from further analysis. See supplementary methods for more details on

timing analyses.

<u>Characterization of adaptive introgression adaptive alleles in *C*. sp 'wide-mouth'</u>

Lastly, we investigated the spatial origins of adaptive alleles shared and unique to the two scale-

eating specialists by searching in our previous study spanning Caribbean-wide outgroup

populations for these alleles [25]. Adaptive alleles were assigned as standing genetic variation if

observed in any population outside SSI or de novo if they were only observed within populations

on SSI. Additionally, we investigated signatures of introgression across the genome of *C*. sp

'wide-mouth' and *C. desquamator* to determine if they showed evidence of adaptive

introgression from outgroup generalist populations as observed previously [25]. See

supplementary methods for more details on introgression analyses.

Results

<u>'Wide-mouth' ecomorph is ecologically intermediate and morphologically distinct</u>

We found that the 'wide-mouth' ecomorph is morphologically distinct from *C. desquamator* and

C. variegatus across a suite of craniofacial traits (figure 2A-B). The lower jaw length of 'wide-

mouth' was intermediate between *C. desquamator* and *C. variegatus* (figure 2C), while the

buccal width and adductor mandibulae height were 8% larger in 'wide-mouth' than *C.*

desquamator (figure 2E-F). These morphological differences were consistent across multiple

lakes (figure S2). Small modifications in craniofacial morphology among these species have

major impacts on scale-eating performance in this system by altering kinematic traits such as

peak gap size which is partially controlled by the length of the lower jaw, jaw protrusion

distance, and the angle of the lower jaw relative to the suspensorium [32].

C. sp 'wide-mouth' also did not show morphological divergence comparable to that

observed in the molluscivore *C. brontotheroides.* The molluscivore specialist presents an

opposing axis of morphological divergence to the scale-eating specialists, with shorter oral jaw

length and larger eye diameter than even the generalist *C. variegatus*, in addition to a novel nasal

protrusion of the maxilla not observed in any other Cyprinodontidae species [33].

Morphological traits were heritable in a common garden laboratory environment after

one generation: lab-reared *C.* sp. 'wide-mouth' displayed significantly larger buccal width than

C. desquamator (t-test; $P = 0.003$) and maintained their characteristic intermediate jaw lengths

(ANOVA; $P = 0.03$, figure S3). There was also some evidence of phenotypic plasticity in both *C.*

desquamator and *C. sp* 'wide-mouth' compared to wild individuals likely caused by the common

lab diet. See supplementary results for more details.

Figure 2. *C.* sp. 'wide-mouth' has distinct morphology within the San Salvador Island adaptive

radiation. A) First two principal components of morphological diversity for 8 size-corrected

traits and 95% confidence ellipses by species (*C. variegatus* : gold; *C.* sp. 'wide-mouth': red-

orange; *C. desquamator*: teal; *C. brontotheriodes* not shown). PC1 is mainly described by lower

jaw length and PC2 by adductor mandibulae insertion height, buccal width, and neurocranium

width. B) Depictions of the three external measurements that best distinguished *C.* sp. 'wide-

mouth' from both *C. desquamator* and *C. variegatus*, measured using digital calipers. C-E) The

relationship between standard length (mm) of individuals and their C) lower jaw length, D)

buccal cavity width, and E) adductor mandibulae insertion height (AM insertion) across

individuals of the three species in Osprey Lake. 95% confidence bands for linear models in gray.

<u>'Wide-mouth' occupies a distinct intermediate scale-eating niche</u>

We found that 'wide-mouth' ingested scales, but at a significantly lower frequency than C.

desquamator (Wilcoxon Rank Sum test, $P = 0.004$; figure 3A). We did not detect any scales in

C. variegatus guts (figure 3A). Detritus made up the rest of the *C. sp.* 'wide-mouth' and *C.*

desquamator diets and was the dominant component of *C. variegatus* gut contents, except for a

single individual with one mollusc shell. A previous study that characterized contents of *C.*

variegatus, *C. brontotheroides*, and *C. desquamator* populations across several ponds also found

detritus to be the dominant component of each species' diet (49-71%) and nearly zero scales in

the gut contents of *C. variegatus* and *C. brontotheroides* [33].

The intermediate scale-eating dietary niche of the wide-mouth ecomorph is

complemented by our stable isotope analyses, which provide long-term snapshots of the carbon

sources and relative trophic levels in these species. Osprey Lake individuals collected on the

same day from the same site differed in δ15N levels across species (ANOVA, $P = 4.55 \times 10^{-6}$;

figure 3B and S4); 'wide-mouth' δ15N was intermediate between *C. variegatus* and *C.*

desquamator (Tukey HSD; $P = 1.34 \times 10^{-5}$ & 1.11×10^{-4} respectively), supporting its intermediate

trophic position. SIBER analyses of trophic niche position indicate distinct positioning of the

wide-mouth ecomorph based on the lack of extensive overlap in niche space measured by

standard ellipse areas and bivariate means with 95% confidence intervals of isotope values

among the species (figure 3B).

Figure 3. *C.* sp. 'wide-mouth' ingests scales. A) Scale counts from gut content analysis of the

hindgut of Osprey pupfish populations (10 individuals per species). B) Relative trophic position

(δ15N stable isotope ratio) and dietary carbon source (δ13C stable isotope ratio) with 95%

confidence ellipses for generalist and scale-eating species. Solid lines represent 95% confidence

intervals around bivariate mean, dotted lines represent standard ellipse areas corrected for sample

size (contain 40% of data; SEAc), shaded circles represent ellipse area that contain 95% of the

data calculated using the R package SIBER. C) Still images of scale-eating strikes in *C.*

desquamator and *C.* sp. 'wide-mouth' filmed at 1100 fps on a Phantom VEO 440S camera.

<u>*C.* sp. 'wide-mouth' did not result from hybridization between *C. variegatus* and *C. desquamator*</u>

Several lines of genomic evidence from PCA, ADMIXTURE, and *f-statistic* analyses support the

'wide-mouth' ecomorph as a genetically distinct species rather than a recent hybrid between *C.*

desquamator and *C. variegatus* on SSI as their intermediate jaw morphology might suggest

(figure 4A-C & S5-6; see Supplementary material for more details). Demographic modeling of

divergence and gene flow on SSI places *C. sp* 'wide-mouth' as sister to *C. desquamator*,

supporting previous phylogenetic inference [34]. In the best supported model of 28 demographic

models tested (table S2), 'wide-mouth' and *C. desquamator* diverged 11,658 years ago (95 CI:

8,257-20,113 years; figure S7B; table S2) with ongoing gene flow. Additionally, the amount of

genetic divergence between populations indicates that *C. desquamator* and *C.* sp. 'wide-mouth'

are more genetically diverged from each other than to the generalist *C. variegatus* (e.g. F_{st} in

figure 4C).

Figure 4. Patterns of selection and genetic divergence in specialist genomes**.** A) Principal

components analysis of the four focal groups on San Salvador Island based on an LD-pruned

subset of genetic variants (78,840 SNPs). B) Ancestry proportions across individuals of the four

focal groups. Proportions were inferred from ADMIXTURE analyses with 2 values of K with the

highest likelihood on the same LD-pruned dataset in A. C) Selective sweep length distributions

across generalist and scale-eating species. Rug plot below each histogram represents the counts

of selective sweeps in different length bins. D) The total number of fixed or nearly-fixed SNPs

($F_{st} \geq 0.95$) between each group in Osprey Pond. E) The number of adaptive alleles (fixed or

nearly-fixed SNPS [$F_{st} \geq 0.95$] relative to *C. variegatus* and under selection in each population of

specialists in Osprey Lake. Venn diagram highlights those adaptive alleles that are unique to

each specialist and shared with other specialists.

<u>*C.* sp 'wide-mouth' is comprised of both shared and unique adaptive alleles</u>

Next we looked at regions of the genome in both *C. desquamator* and *C.* sp. wide-mouth that

showed strong evidence of hard selective sweeps. We found 6 shared hard selective sweeps in

both species containing a total of 15 SNPS that were fixed or nearly-fixed compared to the

sympatric generalist *C. variegatus* (figure 4E, 5): 10 SNPs were in unannotated regions, two

were in the introns of the gene *daam2,* and three were in putative regulatory regions (with 20-kb)

of the genes *usp50, atp8a1,* and *znf214* (one variant each). Shared adaptive alleles in the gene

daam2, a wnt signaling regulator, are intriguing because knockdown of this gene causes

abnormal snout morphology in mice [35] and abnormal cranial and skeletal development in

zebrafish [36].

We also found unique sets of adaptive alleles in *C.* sp 'wide-mouth' and *C. desquamator*

(figure 4E;figure 5). None of the adaptive alleles unique to *C.* sp. 'wide-mouth' were in or near

genes annotated for craniofacial phenotypes in model organisms, despite its distinctive

craniofacial morphologies. In *C. desquamator,* three of 12 unique adaptive alleles were in or near

genes associated with or known to affect craniofacial phenotypes: a *de novo* non-synonymous

coding substitution in the gene *twist1*, several putative regulatory variants near the gene *gnaq,*

and 8 variants in or near the gene *bri3bp*, which is located inside a QTL region for cranial height

in pupfish [37]. *C. brontotheriodes* also contained at least one unique candidate craniofacial

adaptive allele: a non-synonymous coding substitution in the gene *kat6b* (figure 5), which is

associated with abnormal craniofacial morphologies, including shorter mandibles, in mice [38].

This pattern of unique alleles relevant to craniofacial phenotypes in specialists *C.*

brontotheriodes and *C. desquamator*, but not *C.* sp 'wide-mouth', holds even if we lower the

threshold to the top 1 percentile of F_{st} outliers between specialists and generalist (see

supplemental results; figure S10).

The origins of adaptive alleles in C. sp 'wide-mouth' and desquamator

The adaptive alleles shared by *C. desquamator* and *C.* sp. 'wide-mouth' occurred as low

frequency standing genetic variation in the Caribbean, with the exception of a single de novo

allele on SSI located in an unannotated region on scaffold 6 (figure 5). The adaptive alleles

unique to *C. desquamator* and *C.* sp 'wide-mouth' also predominantly came from standing

genetic variation (84% and 81%, respectively). 14% of adaptive alleles unique to *C.*

desquamator were de novo mutations to SSI and 2% occurred in candidate introgression regions

(table S7). We found the opposite in *C. sp* 'wide-mouth': only 4% of their unique adaptive

alleles were de novo mutations whereas 15% occurred in candidate introgression regions (table

S8). This adaptive introgression was detected for generalist populations sampled from North

Carolina and Laguna Bavaro in the Dominican Republic (table S8; figure S11). Using the

Relative Node Depth (RND) statistic, we also discovered that 5 of the 6 shared adaptive alleles

(all except for the unannotated region on scaffold 43;table S6) appear introgressed between *C.*

desquamator and *C.* sp. 'wide-mouth', suggesting a substantial contribution of introgression to

the adaptive alleles observed in *C.* sp. 'wide-mouth'.

<u>Timing of selection on adaptive alleles reveals features of the adaptive walk to scale-eating</u>

Selective sweeps occurred much more recently in both populations than their inferred divergence

times (figure 5). Intriguingly, selection on 4 of the 6 adaptive alleles occurred significantly

earlier in *C. desquamator* than 'wide-mouth'. Only a single adaptive allele had an older median

age estimate in 'wide-mouth' than *C. desquamator*, although the 95% HPD intervals overlapped

between the species (figure 5). Additionally, overall we found a significant difference in timing

of selection between shared and unique adaptive alleles in the two scale-eater populations

(ANOVA *P*-value = 0.00478).   In *C. desquamator*, shared adaptive alleles swept before any

unique adaptive alleles (Tukey HSD *P*-value = 0.003217; figure 5). For the 'wide-mouth', shared

adaptive alleles with *C. desquamator* also generally swept before those unique to the species,

despite these unique alleles being standing and introgressed variation from the Caribbean (figure

5). However, this difference was not significant due to one unique adaptive allele (*slitrk5)* that

swept first (figure 5; ANOVA, Tukey HSD; *P* = 0.8367). This adaptive allele resides in a region

that appears to be introgressed with the Laguna Bavaro generalist population in the Dominican

Republic where this allele also show signs of a hard selective sweep [25]. The older age estimate

of this sweep in *C*. sp 'wide-mouth' might be due to older shared selection for the alleles in other

Caribbean populations before introgression with *C*. sp 'wide-mouth'. All other introgressed

adaptive alleles in *C*. sp. 'wide-mouth' swept more recently than shared sweeps with

desquamator, including the shared de novo allele, and were not under selection in outgroup

generalist populations.

Intriguingly, all but one of the de novo adaptive alleles in *desquamator* swept at the same

time in the recent past (figure 5). Only one of these adaptive alleles in *olfm1* region has 95%

HPD age range that overlaps with the next oldest selective sweep on standing genetic variation

(*gnaq*; figure 5), suggesting a discrete stage of selection on new mutations in *C. desquamator*.

Figure 5. Timing of selection on adaptive alleles in trophic specialists nested within the

demographic history of the radiation. The mean and 95% HPD estimates for the timing of

selection on sets of fixed or nearly fixed SNPs (named by the gene they are in or within 20-kb

of) for the three specialist populations found in sympatry in Osprey Lake (sweeps in *C.*

variegatus not shown). The age of each beneficial allele is color coded by the species and the

inferred demographic history is displayed in the background for comparison. Gene names

highlighted in bold are associated with oral jaw size. Gene names are colored by source of

genetic variation (de novo: red; introgressed with outgroup: blue; standing genetic variation:

black). Gene names with asterisk indicate those inferred as introgressed between *C. desquamator*

and *C.* sp 'wide-mouth'.

<u>Shared signatures of selection across the three specialists in the radiation</u>

Lastly, we compared the genetic divergence and selection patterns observed in the two scale-

eating specialists to the divergent molluscivores specialist *C. brontotheroides* to investigate the

extent of allele sharing among all three trophic specialists in this adaptive radiation. We found

that no fixed or nearly-fixed alleles relative to the generalist *C. variegatus* were shared across all

three specialists (figure S9-S10; supplementary results). However, we did find evidence of 44

shared selective sweeps across all three specialist populations that were not shared with *C.*

variegatus populations (figure S12C). These shared regions were significantly enriched for genes

annotated for metabolic processes (figure S12D), suggesting shared selection for metabolizing

the more protein-rich diet across all three trophic specialists (also see [39]).

Discussion

<u>Discovery of a new cryptic intermediate scale-eater highlights the power of reusing genetic</u>

<u>variation to access novel niches</u>

The hallmark of adaptive radiation is a rapid burst of diversification which is predicted by theory

to slow down over time as niche subdivision increases [6]. An alternative possibility is that

radiations can be self-propagating and that the diversity generated within the first stages of

radiation helps beget further diversity [40]. This could happen through exploitation of new

trophic levels created by new species or physical alterations of the environment by new species

that may create additional opportunities for speciation (reviewed in [6,41]). The diversity begets

diversity hypothesis can also be visualized as the exploration of a complex multi-peaked fitness

landscape; as species in the radiation colonize new peaks, this provides access to additional

neighboring fitness peaks to fuel rapid radiation. Our discovery of a cryptic new scale-eating

species through morphological, dietary, and genomic analyses revealed shared nearly-fixed or

fixed adaptive alleles in both scale-eating species relative to the generalist *C. variegatus*. While

C. sp. 'wide-mouth' is ecologically intermediate in its scale-eating behavior, our estimates of the

relative timing of selective sweeps suggest that these shared alleles were first selected upon in

the more specialized scale-eater *C. desquamator*, although unaccounted for demographic

differences may also be contributing to differences in estimated timing between species.

378   Intriguingly, the shared adaptive alleles between *C. desquamator* and *C.* sp. 'wide-

mouth' have potentially introgressed more recently rather than selected upon in their shared

common ancestor. Five of the six regions surrounding these shared adaptive alleles showed

patterns of high genetic similarity consistent with introgression (table S10). Alternatively, this

genetic similarity may also be caused by strong background selection on shared ancestral

variation. Effective population sizes are not drastically different between the two species and

exon density is not in the upper tail of the genome wide-distribution (figure S1;table S10), two

factors found in other studies where background selection tends to confound adaptive

introgression inferences [42,43]. However, we do not have extensive knowledge of

recombination breakpoints in this non-model system to distinguish between strong background

selection on shared ancestral variation and adaptive introgression scenarios for each candidate

adaptative introgression region.

We also found strong signatures of introgression in *C.* sp. 'wide-mouth' genomes from

outgroup generalist populations that were not shared with *C. desquamator* (figure S11; table

S11). Craniofacial morphology is a major axis of diversification between trophic specialists in

this system [44], yet *C.* sp. 'wide-mouth' appears to have little unique genetic variation relevant

for craniofacial traits compared to the other two specialists (figure S10). Despite this, they do

exhibit transgressive craniofacial phenotypes not seen in the other specialists. An intriguing

implication of these findings is that hybridization may allow different species to share many of

the same adaptive alleles to occupy distinct but similar niches, in line with the syngameon and

'diversity begets diversity' hypotheses of adaptive radiation [18,40].

<u>An adaptive walk underlies the major ecological transition from generalist to scale-eating</u>

<u>specialist</u>

The foundational model of adaptation is that it proceeds in 'adaptive walks' towards fitness

optima that involve the sequential fixation of adaptive alleles that move a population in the

phenotypic direction of the local optimum [45]. The distinct timing of selection across different

adaptive alleles in both *C. desquamator* and *C.* sp 'wide-mouth' suggests that the ecological

transition from generalist to novel scale-eating specialist involved an adaptive walk in which

selection on a beneficial allele was contingent on prior fixation of other adaptive alleles in each

specialists' genetic background (see supplemental materials for further discussion). This is best

highlighted by the pattern observed in *C. desquamator* in which nearly all de novo mutations

swept at the same time in a distinct selective stage from other adaptive variants rather than being

selected upon as they originated (figure 5).

The (un)predictability of adaptive walks to novel ecological niches

A recent study characterizing genotypic fitness landscapes underlying the transition from *C.*

variegatus and *C. desquamator* revealed a rugged landscape with many local fitness peaks, likely

due to epistatic interactions among alleles [46]. The staggered timing of selection on alleles lends

support to this finding. Epistasis can reduce the number of adaptive walks selection will promote

[47], and might explain why the same adaptive alleles were the first to undergo hard selective

sweeps in both 'wide-mouth' and *desquamator*.

We also found evidence for shared selective sweeps across all three specialists in regions

that are enriched for genes annotated for metabolic processes such as short chain fatty acid and

propionate metabolism (figure S12D). The lack of fixed alleles in these regions relevant to

dietary specialization suggests polygenic selection (see supplemental for more discussion).

Subtle shifts of allele frequencies across the genome can lead to divergent genomic backgrounds

that give populations access to different ecological niches (e.g. [48,49]).

While both shared sweeps among all specialists and shared adaptive alleles among the

two scale-eating species suggest constrained adaptive walks along overlapping genotypic

pathways, we still see most selective sweeps are unique to each species in this radiation (figure

4; figure S11). Curiously, some adaptive standing genetic variation rose to high frequency in *C.*

desquamator but did not similarly undergo selection in *C.* sp 'wide-mouth', despite its adaptation

to a similar ecological niche and the presence of these alleles segregating at low frequency in the

'wide-mouth' population. This highlights the dual influence of epistatic interactions on adaptive

walks in rugged landscapes – epistasis reduces number of available paths but increases the

number of local fitness peaks populations can get stuck on [50]. Selection on the same adaptive

alleles may have allowed both scale-eating species access to the same area of the fitness

landscape but epistatic interactions with private mutations and introgressed variation in each

lineage may have resulted in divergent paths to scale-eating, ultimately contributing to diverse

evolutionary outcomes even from a shared starting point.

The use of adaptive alleles from distinct spatial sources, the distinct morphologies and

divergent genomic backgrounds, and potential introgression of adaptive alleles from the more

specialized scale-eater *C. desquamator* into *C.* sp 'wide-mouth' reveals a tangled path for novel

ecological transitions in nature. The complex epistatic interactions at microevolutionary scales

implicated in this study make it all the more fascinating that novel ecological transitions are a

common macroevolutionary feature of adaptive radiation.

Acknowledgements

We thank Priya Moorjani, Michelle St. John, Joseph McGirr, Jacquelyn Galvez, David Tian,

Austin Patton, and Joseph Heras for helpful comments on the manuscript; the Gerace Research

Centre and Troy Day for logistical support; the governments of the Bahamas for permission to

collect and export samples. This research was funded by the National Science Foundation DEB

CAREER grant #1749764, National Institutes of Health grant 5R01DE027052-02, the University

of North Carolina at Chapel Hill, and the University of California, Berkeley to CHM.

Author Contributions

Conceptualization: EJR,CHM; Data Collection: EJR,CHM; Statistical analyses: EJR; Resources:

CHM; Visualization: EJR,CHM; Original draft: EJR; Revising: EJR, CHM

Declaration of Interests

The authors declare no competing interests.

Ethics Statement

Fishes were euthanized in an overdose of buffered MS-222 (Finquel, Inc.) following approved

protocols from the University of North Carolina at Chapel Hill Animal Care and Use Committee

(#18-061.0), and the University of California, Berkeley Animal Care and Use Committee (AUP-

2015-01-7053) and preserved in 95-100% ethanol. All animals were collected and exported with

2017 and 2018 permits from the Bahamas Environmental Science and Technology commission

and Ministry of Agriculture.

Data Accessibility

All genomic data are archived at the National Center for Biotechnology Information BioProject

Database (Accessions:

PRJNA690558; PRJNA394148, PRJNA391309; and PRJNA305422). Ecological and

morphological data are provided in the supplemental materials.

References

1.   Dieckmann U, Doebeli M. 1999 On the origin of species by sympatric speciation. *Nature*
400, 354.
2.   Polechová J, Barton NH, Gavrilefs S. 2005 Speciation thrrough competition: a critical
review. *Evolution (N. Y).* **59**, 1194–1210. (doi:10.1554/04-691)
3.   Bolnick DI. 2006 Multi-species outcomes in a common model of sympatric speciation. *J.*
Theor. Biol. **241**, 734–744. (doi:10.1016/j.jtbi.2006.01.009)
4.   Kondrashov AS, Kondrashov FA. 1999 Interactions among quantitative traits in the
course of sympatric speciation. *Nature* **400**, 351–354.
5.   Gavrilets S. 2014 Models of speciation: where are we now? *J. Hered.* **105**, 743–755.
(doi:10.1093/jhered/esu045)
6.   Martin CH, Richards EJ. 2019 The Paradox behind the Pattern of Rapid Adaptive
Radiation: How Can the Speciation Process Sustain Itself through an Early Burst? *Annu.*

Rev. Ecol. Evol. Syst.

7.   Arnold SJ, Pfrender ME, G. JA. 2001 The Apaptive Landscape as a conceptual bridge
between mico- and macroevolution. *Genetica* **112**–**113**, 9–32.

8.   Svennson E, Calsbeek R. 2012 *The adaptive landscape in evolutionary biology*. Oxford
University Press.

9.   Martin CH, Wainwright PC. 2013 Multiple fitness peaks on the adaptive landscape drive
adaptive radiation in the wild. *Science* **339**, 208–11. (doi:10.1126/science.1227710)

10.   Martin CH, Gould KJ. 2020 Surprising spatiotemporal stability of a multi-peak fitness
landscape revealed by independent field experiments measuring hybrid fitness. *Evol. Lett.*
, 530–544. (doi:10.1002/evl3.195)

11.   Calsbeek R, Irschick DJ. 2007 The quick and the dead: Correlational selection on
morphology, performance, and habitat use in island lizards. *Evolution (N. Y).* **61**, 2493–
2503. (doi:10.1111/j.1558-5646.2007.00206.x)

12.   Grant PR, Grant RB. 2008 *How and Why Species Multiply*. Princeton University Press.

13.   Givnish TJ, Sytsma KJ, Smith J, Hahn W, DH B, Burkhardt E. 1997 Molecular evolution
and adaptive radiation in *Brocchinia* (Bromeliaceae: Pitcairnioideae) atop tepuis of the
Guyana Shield. In *Molecular evolution and adaptive radiation*, pp. 259–311. Cambridge:
Cambridge University Press.

14.   Givnish TJ, Burkhardt EL, Happel RE, Weintraub JD. 1984 Carnivory in the Bromeliad
Brocchinia reducta , with a Cost / Benefit Model for the General Restriction of
Carnivorous Plants to Sunny , Moist , Nutrient-Poor Habitats. *Am. Nat.* **124**, 479–497.

15.   Erwin DH. 2021 A conceptual framework of evolutionary novelty and innovation. *Biol.*
Rev. **96**, 1–15. (doi:10.1111/brv.12643)

16.   Blount ZD, Barrick JE, Davidson CJ, Lenski RE. 2012 Genomic analysis of a key
innovation in an experimental *Escherichia coli* population. *Nature* **489**, 513–518.
(doi:10.1038/nature11514)

17.   Quandt EM, Deatherage DE, Ellington AD, Georgiou G, Barrick JE. 2014 Recursive
genomewide recombination and sequencing reveals a key refinement step in the evolution
of a metabolic innovation in Escherichia coli. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 2217–
2222. (doi:10.1073/pnas.1314561111)

18.   Seehausen O. 2004 Hybridization and adaptive radiation. *Trends Ecol. Evol.* **19**, 198–207.
(doi:10.1016/j.tree.2004.01.003)

19.   John MES, Dixon KE, Martin CH. 2020 Oral shelling within an adaptive radiation of
pupfishes : Testing the adaptive function of a novel nasal protrusion and behavioural
preference. , 1–9. (doi:10.1111/jfb.14344)

20.   Martin CH, Wainwright PC. 2013 A remarkable species flock of *Cyprinodon* pupfishes
endemic to San Salvador Island, Bahamas. *Bull. Peabody Museum Nat. Hist.* **54**, 231–240.

21.   Turner BJ, Duvernell DD, Bunt TM, Barton MG. 2008 Reproductive isolation among
endemic pupfishes (Cyprinodon) on San Salvador Island, Bahamas: Microsatellite
evidence. *Biol. J. Linn. Soc.* **95**, 566–582. (doi:10.1111/j.1095-8312.2008.01079.x)

22.   Clark PU, Dyke AS, Shakun JD, Carlson AE, Clark J, Wohlfarth B, Mitrovica JX,
Hostetler SW, McCabe AM. 2009 The Last Glacial Maximum. *Science (80-.).* **325**, 710–
714. (doi:10.1126/science.1172873)

23.   Richards EJ, Martin CH. 2017 Adaptive introgression from distant Caribbean islands
contributed to the diversification of a microendemic adaptive radiation of trophic
specialist pupfishes. *PLoS Genet.* **13**, 1–35. (doi:10.1371/journal.pgen.1006919)

24.   Jackson AL, Inger R, Parnell AC, Bearhop S. 2011 Comparing isotopic niche widths
among and within communities: SIBER - Stable Isotope Bayesian Ellipses in R. *J. Anim.*
Ecol. **80**, 595–602. (doi:10.1111/j.1365-2656.2011.01806.x)

25.   Richards EJ, McGirr JA, Wang JR, St. John ME, Poelstra JW, Solano MJ, O'Connell DC,
Turner BJ, Martin CH. 2021 A vertebrate adaptive radiation is assembled from an ancient
and disjunct spatiotemporal landscape. *Proc. Natl. Acad. Sci. U. S. A.* **118**.
(doi:10.1073/pnas.2011811118)

26.   Peter BM. 2016 Admixture, population structure, and f-statistics. *Genetics* **202**, 1485–
1501. (doi:10.1534/genetics.115.183913)

27.   Excoffier L, Dupanloup I, Huerta-Sánchez E, Sousa VC, Foll M. 2013 Robust
Demographic Inference from Genomic and SNP Data. *PLOS Genet.* **9**, e1003905.
(doi:10.1371/journal.pgen.1003905)

28.   Pavlidis P, Živković D, Stamatakis A, Alachiotis N. 2013 SweeD: Likelihood-based
detection of selective sweeps in thousands of genomes. *Mol. Biol. Evol.* **30**, 2224–2234.
(doi:10.1093/molbev/mst112)

29.   Danecek P *et al.* 2011 The variant call format and VCFtools. *Bioinformatics* **27**, 2156–
2158. (doi:10.1093/bioinformatics/btr330)

30.   Smith J, Coop G, Stephens M, Novembre J. 2018 Estimating Time to the Common
Ancestor for a Beneficial Allele. *Mol. Biol. Evol.* **35**, 1003–1017. (doi:10.1101/071241)

31.   Roesti M, Moser D, Berner D. 2013 Recombination in the threespine stickleback genome
- Patterns and consequences. *Mol. Ecol.* **22**, 3014–3027. (doi:10.1111/mec.12322)

32.   St. John ME, Holzman R, Martin CH. 2020 Rapid adaptive evolution of scale-eating
kinematics to a novel ecological niche. *J. Exp. Biol.* **223**, jeb217570.
(doi:10.1242/jeb.217570)

33.   Martin CH, Wainwright PC. 2013 On the measurement of ecological novelty: scale-eating
pupfish are separated by 168 my from other scale-eating fishes. *PLoS One* **8**, e71164.
(doi:10.1371/journal.pone.0071164)

34.   Richards EJ, Martin CH. 2017 Adaptive introgression from distant Caribbean islands
contributed to the diversification of a microendemic adaptive radiation of trophic
specialist pupfishes. *PLOS Genet.* **13**, e1006919.

35.   Dickinson ME *et al.* 2016 High-throughput discovery of novel developmental phenotypes.
Nature **537**, 508–514. (doi:10.1038/nature19356)

36.   Kida YS, Sato T, Miyasaka KY, Suto A, Ogura T. 2007 Daam1 regulates the endocytosis
of EphB during the convergent extension of the zebrafish notochord. *Proc. Natl. Acad.*
Sci. U. S. A. **104**, 6708–6713. (doi:10.1073/pnas.0608946104)

37.   St. John, Michelle E., Dunker JC, Richards, Emilie J., Romero S, Martin CH. 2021
Parallel genetic changes underlie integrated craniofacial traits in an adaptive radiation of
trophic specialist pupfishes. , 1–46.

38.   Thomas T, Voss AK, Chowdhury K, Gruss P. 2000 Querkopf, a MYST family histone
acetyltransferase, is required for normal cerebral cortex development. *Development* **127**,
2537–2548.

39.   Mcgirr JA, Martin CH. 2018 Parallel evolution of gene expression between trophic
specialists despite divergent genotypes and morphologies. *Evol. Lett.* **2**, 62–75.
(doi:10.1002/evl3.41)

40.   Whittaker R. 1977 Evolution of species diversity in land communities. *Evol. Biol.*

41.   Stroud JT, Losos JB. 2016 Ecological Opportunity and Adaptive Radiation. *Annu. Rev.*

Ecol. Evol. Syst. **47**, 507–532. (doi:10.1146/annurev-ecolsys-121415-032254)

42. Kim BY, Huber CD, Lohmueller KE. 2018 Deleterious variation shapes the genomic
landscape of introgression. *PLoS Genet.* **14**, 1–30. (doi:10.1371/journal.pgen.1007741)

43. Zhang X, Kim B, Lohmueller KE, Huerta-Sánchez E. 2020 The impact of recessive
deleterious variation on signals of adaptive introgression in human populations. *Genetics*
215, 799–812. (doi:10.1534/genetics.120.303081)

44. Martin CH, Wainwright PC. 2011 Trophic novelty is linked to exceptional rates of
morphological diversification in two adaptive radiations of cyprinodon pupfish. *Evolution*
(N. Y). **65**, 2197–2212. (doi:10.1111/j.1558-5646.2011.01294.x)

45. Orr HA. 2005 The genetic theory of adaptation: A brief history. *Nat. Rev. Genet.* **6**, 119–
127. (doi:10.1038/nrg1523)

46. Patton AH, Richards EJ, Gould KJ, Buie LK, Christopher H. 2021 Adaptive introgression
and de novo mutations increase access to novel fitness peaks on the fitness landscape
during a vertebrate adaptive radiation.

47. Weinreich DM, Delaney NF, DePristo MA, Hartl DL. 2006 Darwinian Evolution Can
Follow Only Very Few Mutational Paths to Fitter Proteins. *Science (80-.).* **312**, 111 LP –
114. (doi:10.1126/science.1123539)

48. Otte KA, Nolte V, Mallard F, Schlötterer C. 2021 The genetic architecture of temperature
adaptation is shaped by population ancestry and not by selection regime. *Genome Biol.* **22**,
1–25. (doi:10.1186/s13059-021-02425-9)

49. Hansen TF. 2013 Why epistasis is important for selection and adaptation. *Evolution (N.*
Y). **67**, 3501–3511. (doi:10.1111/evo.12214)

50. Fragata I, Blanckaert A, Dias Louro MA, Liberles DA, Bank C. 2019 Evolution in the
light of fitness landscape theory. *Trends Ecol. Evol.* **34**, 69–82.
(doi:https://doi.org/10.1016/j.tree.2018.10.009)

Supplemental Text, Figures and Tables:

We get by with a little help from our friends: shared

adaptive variation provides a bridge to novel ecological

specialists during adaptive radiation

Emilie J. Richards[1,2] Christopher H. Martin[1,2]

[1]Department of Integrative Biology, University of California, Berkeley

[2]Museum of Vertebrate Zoology, University of California, Berkeley

Correspondence: *chmartin@berkeley.edu*

Keywords: ecological speciation, adaptive radiation, novelty, speciation genomics, adaptive

introgression, trophic innovation

Methods

 Sampling

C. sp. 'wide-mouth' individuals were collected on San Salvador Island in the Bahamas using

hand and seine nets between 2017 and 2018. 'Wide-mouth' individuals were collected from 5

lakes in which taxa occurs in sympatry with just *C. variegatus* (Great Lake, and Stout's Lake) or

with generalist and *C. desquamator* (Oyster Pond, Osprey Lake, and Mermaid's Pond,; Fig. 1).

In Oyster Pond and Little Lake, *C. desquamator* is extremely rare (only a single Oyster Pond

individual grouped genetically and morphologically with *C. desquamator*), so Osprey Lake

appears to be the only lake where all three species and the recently discovered 'wide-mouth'

coexist at appreciable frequencies in sympatry. 'Wide-mouth' individuals have also been

collected in Little Lake in the past (Bruce Turner, pers. obs. 1990s).

Fishes were euthanized in an overdose of buffered MS-222 (Finquel, Inc.) following

approved protocols from the University of North Carolina at Chapel Hill Animal Care and Use

Committee (#18-061.0), and the University of California, Berkeley Animal Care and Use

Committee (AUP-2015-01-7053) and preserved in 95-100% ethanol.

Ecological and morphological characterization of 'wide-mouth' scale-eater

Multiple individuals from 3 lake populations (Osprey Lake, Great Lake, and Oyster Pond) in

which we had measurable specimens of the species (n=84) were measured for 9 traits using

digital calipers. Traits were selected for specific connections to foraging performance [1,2] and

that differed across the three groups in a previous study when externally measured [3]. For each

specimen we measured (1) standard length from the anterior tip of the premaxilla to the posterior

margin on the midline of the caudal peduncle, (2) lower jaw length from the medial point of the

dentary on the lower jaw to the distal point of rotation on the quadrate-articular joint, (3) the

width of the buccal from the distance between the proximal sides of the buccal cavity in dorsal

view, (4) the height of the adductor mandibulae insertion from the vertex between the vertical

and horizontal arms of the preopercle to the dorsal margin of the hyomandibula, (5) the distance

from the insertion of the first ray of the dorsal fin to the insertion of the first ray of the anal fin,

(6) the height of the caudal peduncle from the dorsal and ventral procurrent rays on the caudal

fin, (7) neurocranium width from the narrowest distance between the eyes from the dorsal view,

(8) body width from just behind the posterior margins of each operculum in dorsal view, and (9)

diameter of the orbit. Measurements were made by a single observer (EJR). Repeatability of

measurements was assessed by remeasuring a random subset of 10 individuals per species twice.

Measurements on the full dataset were not taken until repeatability was above 85% per trait (r^2 in

a linear regression of both sets of measurements). To remove the effects of body size on trait

size, we used the residuals from a linear regression of the log-transformed trait on log

transformed standard length. We also compared these morphological measurements between lab-

reared F2 and wild caught specimens to rule out strong plasticity in traits of interest.

For a subset of individuals collected from each population (*C. variegatus, C.

desquamator,* and 'wide-mouth') in Osprey Lake, diet was characterized from stomach content

analyses (*n*=10 each) and stable isotope analyses (*n*=75; 42 *C. variegatus*, 16 *C. desquamator*, 17

'wide-mouth' individuals). These are two complementary approaches that reflect short-term and

long-term snapshots of the dietary differences among these groups. For stomach content analysis,

a single observer (EJR) dissected out a 1 cm portion of the gut from the posterior hindgut region

and counted the number of scales found. For stable isotope analyses, muscle tissue samples were

taken from the caudal peduncle region of each individual immediately after euthanasia in an

overdose of MS-222. Muscle samples were individually labeled and dried at 60° C for at least 24

hours. Foil-wrapped tissue samples were then sent to the UC Davis Stable isotope facility and

analyzed on a PDZ Europa ANCA-GSL elemental analyzer interfaced to a PDZ Europa 20-20

isotope ratio mass spectrometer to characterize the natural abundances of δ13C and δ15N among

individuals. δ13C and δ15N levels were summarized across each of the three species to look at

overlap among species by 40% and 95% Standard Area Ellipses and 95% confidence ellipses

around bivariate means on isotope biplots (figure 3) of the data using R package SIBER [4].

Additionally we assessed whether there were significant differences in δ15N values between

species through an ANOVA using aov() function from the base statistics package in R (v 4.1.0).

<u>Genomic library preparation</u>

We sequenced 24 'wide-mouth' individuals following protocols used in a previous study [5] in

which we sequenced genomes from *Cyprinodon variegatus*, *C. desquamator*, and *C.*

brontotheroides. Raw reads were mapped from these 24 individuals to a de-novo assembly of

Cyprinodon brontotheroides reference genome (v 1.0; total sequence length = 1,162,855,435 bp;

number of scaffolds = 15,698, scaffold N50 = 32 Mbp) with bwa-mem (v 0.7.12;[6]). Duplicate

reads were identified using MarkDuplicates and BAM indices were created using

BuildBamIndex in the Picard software package (http://picard.sourceforge.net(v.2.0.1)). We

followed the best practices guide recommended in the Genome Analysis Toolkit (v 3.5;[7]) to

call and refine our single nucleotide polymorphism (SNP) variant dataset using the program

HaplotypeCaller. Variants in 'wide-mouth' individuals were called jointly with the 202

individuals sequenced from a previous study [5]. We filtered SNPs based on the recommended

hard filter criteria (i.e. QD < 2.0; FS > 60; MQRankSum < -12.5; ReadPosRankSum < -8;[7,8])

because we lacked high-quality known variants for these non-model species. After selecting for

only individuals from San Salvador Island, variants were additionally filtered to remove SNPs

with a minor allele frequency below 0.05, genotype quality below 20, or containing more than

10% missing data across all individuals at the site using vcftools (v.0.1.15;[9]). Variants in

poorly mapped regions were then removed using a mask file generated from the program

SNPable (http://bit.ly/snpable; k-mer length =50, and 'stringency'=0.5). Our final dataset after

filtering contained 6.4 million variants with 7.9x median coverage per individual.

<u>Testing for signatures of recent hybrids</u>

We first tested whether these 'wide-mouth' individuals represented recent (e.g. F1/F2) hybrids of

C. variegatus and *C. desquamator* in the wild. These two young species do reproduce with each

other in the lab and rare hybrid spawning events have been observed in the wild (unpublished

data ME St. John). As a first visual assessment, we conducted principal component analysis on

C. desquamator, generalist, and 'wide-mouth' individuals to look for the genome-wide pattern

expected in PCAs in which recent hybrids between two populations are included. To perform the

PCA, the genetic dataset was first pruned for SNPs in strong linkage disequilibrium using the LD

pruning function (--indep-pairwise 50 5 0.5) in plink (v1.9;[10]), leaving 2.6 million variants.

We then ran a principal component analysis using the eigenvectors output by plink's pca

function (--pca). The first two principal components were plotted in R (R Core Team 2018

v3.5.0).

As a complementary assessment, we used an admixture model-based approach to

estimate the proportion of shared ancestry among individuals in our dataset using ADMIXTURE

(v.1.3.0)[11]. The number of populations (K) was decided upon using ADMIXTURE's cross-

validation method (--cv) across 1-10 population values of K. K = 4 was then chosen using the

broken-stick method. Ancestry proportions estimated by ADMIXTURE were plotted in R for the

K value with the highest likelihood and the two K values surrounding to explore whether the

strong signatures of population structure in Crescent Pond *C. desquamator* individuals was

masking hybridization signatures in any of the 'wide-mouth' populations.

Lastly, to discriminate among alternative evolutionary scenarios for the origin of 'wide-

mouth' and to estimate divergence time among all four species, we used demographic modeling

based on the folded minor allele frequency spectrum (mSFS). The observed mSFS was computed

using the SFStools script from D. Marques available on github

(https://github.com/marqueda/SFS-scripts). For these demographic model comparisons, we used

only individuals from Osprey Lake (*C. desquamator*: n=10; *C. variegatus* : n=12; *C.* sp. 'wide-

mouth': n=11; *C. brontotheriodes*: n=13) to avoid additional complex population structure across

ponds. We contrasted 28 demographic models of different topologies and gene flow scenarios

across the four groups (Fig S5;Table 1 and S2). For each model, the fit to the observed

multidimensional mSFS was maximized using the composite-likelihood method fastsimcoal

(v2.6.0.3;[12] with 100,000 coalescent simulations, 40 expectation-maximization cycles, and

pooling all entries with less than 20 SNPs). For parameter estimates, we used a wide search

range with log-uniform distributions with the range of some priors informed by previous

estimates of effective population size from MSMC and the age of the last glacial maximum

(~20kya) prior to which the lakes would have been dry on SSI. These ranges were not upper

bounded by these specified priors and so the simulations were free to explore parameter space

that exceeded the priors.

For each demographic model, we ran 100 independent *fastsimcoal2* runs to determine

the parameter estimates with the maximum likelihood. The best fitting demographic model was

identified using Akaike information criteria (AIC). To get confidence intervals for the parameter

estimates from the best fitting model, we simulated 100 mSFS based on the maximum likelihood

parameter estimates from our best fitting model and ran 50 independent runs with this model on

each simulated SFS. The parameter point estimates from the run with the highest likelihood (of

the 50 independent runs) from each simulated SFS were then used to compute 95 percentile

confidence intervals as a measure of uncertainty in the parameter estimates from the observed

SFS.

<u>Testing for shared adaptive alleles</u>

Patterns of directional selection

Various demographic histories can shift the distribution of low- and high-frequency derived

variants to falsely resemble signatures of hard selective sweeps. In a previous study, we used

MSMC analyses to infer histories of population size changes in *C. variegatus*, *C. desquamator*

and *C. brontotheriodes* (Richards et al. 2021). In this study, we repeated the same analysis for

the 'wide-mouth'. In order to account for demography of the 'wide-mouth' population in

downstream analyses, we used the MSMC (v. 1.0.1;[13] to infer historical effective population

size (N_e) changes in the 'wide-mouth'. We ran MSMC on unphased GATK-called genotypes

separately for a single individual 'wide-mouth' from Osprey Lake with 16x mean coverage

across its genome (Fig S4). As recommended in the MSMC documentation, we masked sites

with less than half or more than double the mean coverage for that individual or with a genotype

quality below 20. We also excluded sites with <10 reads as recommended by Nadachowska-

Brzyska et al. [14]. To scale the output of MSMC to real time and effective population sizes, we

used a one-year generation time [15] and the estimated spontaneous mutation rate of 1.56 x10^{-8}

estimated from high coverage sequencing of two independent pedigreed parent-offspring crosses

of San Salvador Island species from a previous study (Richards et al. 2021).

Across all four populations in Osprey Lake, we looked for regions that appeared to be

under strong divergent selection in the form of a hard selective sweep from the site frequency

spectrum using SweeD (v.3.3.4;[16]). In this calculation of the composite likelihood ratio (CLR)

of a sweep, we incorporated our empirical estimate of the decrease in population size for each

focal population estimated from MSMC analyses in 50-kb windows across scaffolds that were at

least 100-kb in length (99 scaffolds; 85.6% of the genome). We also calculated CLRs across

100,000 scaffolds consisting of neutrally evolving sequences simulated with ms-move [17],

controlling for the impact of the inferred population size decreases over time for each population

from MSMC runs mentioned above (Fig. S4; Table S3). The CLR ratios for the simulated

datasets were then used to assess outlier CLR ratios from the empirical dataset. Regions with

CLR ratios above the 95th percentile value of CLR from the neutral simulated dataset were

considered candidate hard selective sweep regions (Table S3). We compared selective sweeps

across *C. variegatus, C. desquamator, C. brontotheriodes*, and 'wide-mouth' to look for shared

and unique selective sweeps among the four groups.

Characterization of candidate adaptive alleles

We searched for candidate adaptive alleles underlying divergent traits among the species by

overlapping selective sweeps regions with regions of high genetic divergence based on fixed or

nearly fixed SNPs between groups. We chose to look at nearly fixed SNPs over only fixed

variation to accommodate the ongoing geneflow occurring between these young species. We

took two approaches to finding fixed or nearly-fixed variants: 1) a fixed threshold of $F_{st} \geq 0.95$

across all comparisons and 2) and threshold of the 99.9th percentile of F_{st}, which varied among

comparisons (range of F_{st}: 0.73-0.83).

We made the following pairwise comparisons for F_{st} calculations including a) between *C.*

variegatus and each of the specialists, b) each specialist against all other groups and c) shared

between two or three specialists against *C. variegatus*. For nearly fixed variants with the $F_{st} \geq$

0.95 threshold, we looked for putative function of the candidate adaptive alleles by looking at

gene annotations of any gene the variant was in or near (within 20-kb of the gene, which is

within the 50-kb LD decay estimate). We used available gene annotations from model organisms

of mice and zebrafish from MGI, ZFIN, and we checked other annotation databases and studies

for verification of putative function, including Phenoscape Knowledgebase

(https://kb.phenoscape.org/#/home), NCBI's PubMed (https://www.ncbi.nlm.nih.gov/pubmed),

and the Gene Ontology database using AMIGO2 [18] .

For the 99.9th percentile F_{st} variants: We performed gene ontology (GO) enrichment analyses

for genes near candidate adaptive variants using ShinyGo (v.0.51;[19]). In the *C. brontotheroides*

reference genome annotations (described in *de novo* genome assembly and annotation section),

gene symbols largely match human gene symbols. Thus, the best matching species when we

searched for enrichment across biological process ontologies curated mostly human gene

functions.

Timing of selection on candidate adaptive alleles

Lastly, we also determined the relative age of candidate adaptive alleles by generating estimates

of coalescent times using the R package starTMRCA (v0.6-1;[20]). For each candidate adaptive

allele that was unique to the three specialists and the 16 shared alleles between *C. desquamator*

and 'wide-mouth', a 1-Mb window surrounding the variant was extracted into separate vcfs for

individuals within each species. The program requires complete genotype data so we first filtered

out any individuals with more than 10% missing data (1 *C. brontotheriodes* and 2 'wide-mouth'

individuals). With Tassel5 [21], we then used the LD KKNI command to infer missing sites

based on LD for remaining individuals with less than 10% missing data. Subsequently we

removed the small number of sites with any missing data across individuals within each

population. Since several adaptive alleles fall within the same 1-Mb window, we chose a single

adaptive allele at random within each 1-Mb window as the focal allele to estimate timing for.

This lead to 6 time estimates for the 16 shared adaptive alleles between *C. desquamator* and *C.*

sp. 'wide-mouth', 5 time estimates for the unique adaptive alleles to *C.* sp. 'wide-mouth', X time

estimates for the unique adaptive alleles to *C. desquamator* and 3 time estimates for the unique

adaptive alleles to *C. brontotheroides*.

These sets of variants were then analyzed in starTMRCA with the mutation rate of

1.56×10^{-8} substitutions per base pair, and a recombination rate of 3.11×10^{-8} (from genome-wide

recombination estimate for stickleback; [22]). Each analysis was run three times per focal

adaptive allele and all runs were checked for convergence between and within runs. Most runs

rapidly converged within the initial 6000 steps, but 5 runs did not converge after an additional

4000 steps and were discarded from further analysis.

840
841   Characterization of introgression in C. sp 'wide-mouth' genome

Introgression with C. desquamator in candidate adaptive allele regions

To assess whether shared adaptive alleles between *C.* sp 'wide-mouth' and *C. desquamator* may

be due to introgession between the two populations, we calculated Relative Node Depth (RND)

statistics [23] in 10-kb sliding windows across the genome. RND statistics measure the relative

node depth of two taxa compared to an outgroup and is calculated as the quotient of D_{xy} between

two species to the average distance from each to an outgroup. This relative amount of divergence

compared to an outgroup allows the statistic to be robust to mutation rate variation across the

genome, in which low neutral mutation rate regions might be mistaken for a locus that has

experienced a recent introgression event. RND was calculated using a custom script that used R

package PopGenome (v2.7.5;[24]) first calculate D_{xy} values calculated for *C. desquamator*, *C.*

sp. 'wide-mouth' and *C. artifrons* in 10-kb windows across the genome and then RND for each

scaffold that contained a candidate adaptive variant that was unique or shared between *C.*

desquamator and *C.* sp wide-mouth. Significance of RND values was evaluated using

simulations with no migration using ms-move [17]. We used estimates of changes in effective

population size and divergence times for each population from our *fastsimcoal2* analyses and

ms-move to simulate neutral genetic divergence between two isolated populations. The threshold

for significant introgression regions was determined by simulating genetic sequences under a

coalescent model with no gene flow, consisting of 150,000 10-kb windows each containing the

mean number of alleles observed in our dataset and running these simulated sequences through

the same custom PopGenome script (RND_PopGenome_Artout_null.R) to calculate RND

values. Empirical windows were considered candidates for introgression if the RND statistic was

in the bottom 5[th] percentile of simulated values (RND < 0.22; Table S10). To assess whether the

sympatric coexistence of four incompletely reproductively isolated species in Osprey Lake has

lead to introgression across specialist and generalist population boundaries as well, we assessed

whether RND test statistics calculated between *C. variegatus* and *C. desquamator* in Osprey

Lake were similarly detected as introgressed in these regions. None of these regions appeared to

be introgressed with the generalist *C. variegatus* (Table S10), suggesting the introgressed

signature between *C. desquamator* and *C*. sp. 'wide-mouth' is not likely due to neutral

introgression likely occurring between all species in the lake due to their incomplete

reproductive isolation.

Characterizing introgression with outgroup Caribbean generalist populations in candidate

adaptive allele regions

We also investigated whether any of these candidate adaptive alleles in *C. desquamator* and *C.*

sp 'wide-mouth' fell in regions that appear to be introgressed from an outgroup generalist

population (similar to analyses done in our previous study [25]). First, we extracted the genomes

of individuals assigned to our 5 focal outgroup generalist populations in which we had 8 or more

individuals sequenced in that previous study. We filtered variants across these 5 populations and

our San Salvador Island samples down to those that has a quality of 20, and no more than 10%

missing in any population for a dataset of 10,876,882 SNPs across 10 populations (including the

four SSI species and the outgroup *C. artifrons*) and the 26 scaffolds that candidate adaptive

alleles were found on. To detect introgression involving outgroup populations we used the

differential test of introgression, *df*-statistic [26], which is designed to look for signatures of

introgression across sliding genomic windows [27]. The *df*-statistic is a modified version of the

D-statistic, looks at allele frequencies fitting two allelic patterns referred to as ABBA and BABA

based on the tree (((P1,P2),P3),O), where O is an outgroup species in which no gene flow is

thought to occur with the other populations [27]. We used 2 individuals of *C. artifrons* from

Cancun, Mexico as our outgroup population for this test, which forms the deepest divergence

event with *C. variegatus* within the *Cyprinodon* clade [28], and focused on introgression

between *C. desquamator* or *C. sp.* 'wide-mouth' and outgroup Caribbean generalist populations

that was not shared with *C. brontotheroides*. Based on the tree (((P1,P2),P3), *C. artifrons*), the

df statistic was calculated for the combinations of populations in which the focal population (P2)

was either the scale-eater or the molluscivore, the other specialist population was the sister group

(P1), and P3 was one of the Caribbean outgroup populations sequenced in a previous study (Fort

Fisher, North Carolina Coast;Isla Margarita, Venezuela; Lagunas Bavoros, Dominican Republic,

and New Providence Island, Bahamas. *Fd*-statistics were calculated from 10-kb sliding windows

using R package PopGenome. Empirical windows were considered candidates for introgression

if the *df*-statistic was in the top 95[th] percentile of simulated values (*df* > 0.55).

Characterization of admixture history on San Salvador Island with outgroup generalist

populations

The results from sliding window tests for introgression surrounding candidate adaptive

alleles for *C.* sp. 'wide-mouth' indicated that this cryptic species may be the product of

admixture between SSI species and outgroup generalist lineage related to our North Carolina

coast population (Table S8 and S10). To further explore this, we created genome-wide admixture

graphs using the program qpgraph [29] in R package admixtools2 (v2.0.0;

https://uqrmaie1.github.io/admixtools/). Qpgraph can accommodate many populations and

admixture events and creates an admixture graph that summarizes all pairwise f_2- and f_3-statistics

across included populations and estimates admixture proportions and drift weights between

populations. Admixture graphs model the number of admixture events specified by the user and

evaluates the fit of the observed *f*-statistic values between populations to those expected from

this model to produce a negative log likelihood score that measures the total amount by which

the modeled f-statistics failed to match the observed values (where larger likelihood values

indicate poorer fit).

Normally in qpgraph analyses, the user manually specifies the topology of the model and

the program then solves for the optimal values of the parameters. The recent R package

admixtools2 also includes the function find_graphs() that generates and evaluates admixture

graphs across user-defined number of iterations to find the a range of feasible and best fitting

graph topologies for a set of *f*-statistics. Using the four Osprey Lake populations on San Salvador

Island (*C. variegatus, C. brontotheroides, C. desquamator* and *C.* sp. 'wide-mouth', and the 5

Caribbean generalist populations and *C. artifrons* outrgroup mentioned in the above two

sections, we ran find_graphs() 10 times to model 1-10 admixture events.

The input data consisted of 20,381 SNPs that remained after filtering the dataset that

included the outgroup generalist populations used in the sliding-window tests above and an

additional filters for heterozygous sites that had an allele balance between 0.3 and 0.7, minimum

3x depth of coverage, and quality filter of 30. The final dataset used as input for qpgraph

included 20,381 SNPs across 10 populations and each find_graphs() run was allowed to explore

topology space for 1000 iterations. The topology with the best fit was extracted from these runs

and assigned to be the admixture graph for that admixture event. The model and admixture graph

with the lowest likelihood of all admixture events tested (LnL: range 2.16-16.23 across all 10

models), included 7 admixture events. One of these admixture events depicted admixture

between *C. desquamator* and a lineage related to the North Carolina coast generalist population

resulting in the C. sp 'wide-mouth' population. Other admixture events in this admixture graph,

such as an admixture event at the base of the SSI radiation and a separate admixture event

involving *C. desquamator* and an outgroup generalist, supported signatures inferred using

independent program TreeMix from a previous study[25].

The inference of *C.* sp. 'wide-mouth' as admixed between *C. desquamator* and a distantly

related generalist species appeared in the 8 admixture graphs modeling 3-10 admixtures events,

suggesting it's a strong signature in the dataset. However, the outgroup Caribbean generalist

population that acted as one of the donors varied across different runs from being sister to North

Carolina coast population to falling somewhere along the branch that divides the Venezuela

population from the rest of the Caribbean populations. This may suggest that we are missing the

true donor population in our sample. To further support this signature of *C.* sp. 'wide-mouth'

experiencing introgression following a potential secondary contact event between San Salvador

Island and an outgroup generalist population, we used f_4-statistics to look for genome-wide

signatures of differential introgression *between C. desquamator* and *C.* sp 'wide-mouth'.

Differential signatures of introgression between the two would likely indicate introgression

that *C.* sp. 'wide-mouth' contains in its genome but is not shared with *C. desquamator* and

therefore likely not a signature of shared introgression in the ancestor of the two scale-eating

populations on SSI. Genome-wide f_4 statistics were calculated using the f4() function in

admixtools2 which calculates f_4-statistics, Z scores and p-values for the specified populations.

Following the tree ((P1,P2),(P3,P4)), *C. desquamator* was used as sister species P1, *C.* sp. 'wide-

mouth' as focal species P2, the 5 generalist populations from Fort Fisher, North Carolina Coast;

Isla Margarita, Venezuela; Lagunas Bavoros, Dominican Republic, Rum Cay, Bahamams and

New Providence Island, Bahamas were rotated in as the potential donor population P3 and *C.*

artifrons used as population P4 (Table S11). The f_4-statistic that included North Carolina coast

population had highest Z-score and indicated a significant amount of gene flow (p-

value=0.004;Table S11). However, tests that included Rum Cay and New Providence Island also

resulted in f_4-statistics with marginally significant p-values (p-value=0.056 and 0.067

respectively; Table S11), perhaps mirroring that admixture occurred with an unsampled

generalist population somewhere North of SSI and most closely related to North Carolina

population.

Results
969
970
<u>'Wide-mouth' ecomorph is ecologically intermediate and morphologically distinct</u>

Morphological traits were heritable in a common garden laboratory environment after one

generation: lab-reared *C.* sp. 'wide-mouth' displayed significantly larger buccal width than *C.*

desquamator (t-test; $P = 0.003$) and maintained their characteristic intermediate jaw lengths

(ANOVA; $P = 0.03$, figure S3). There was also some evidence of phenotypic plasticity in both *C.*

desquamator and *C.* sp 'wide-mouth' compared to wild individuals. Two of the focal three traits

that distinguished wild populations of *C.* sp. 'wide-mouth' from the other pupfish species, lower

jaw length and buccal width, remain significantly distinct in the lab-reared populations (based on

non-overlapping 95% confidence intervals around mean residuals of each species). However,

one trait that distinguishes *C.* sp. 'wide-mouth' in the wild appears to be quite plastic under

laboratory rearing conditions; there were no significant differences in mean adductor mandibulae

height residuals among species in laboratory environments. This is interesting because adductor

mandibulae height is a proxy measurement for the cross-sectional width of the adductor muscle

directly beneath the skin. The plasticity observed in this proxy may be due to muscle plasticity.

In the lab-reared colonies, all species are reared on the same diet, such that the scale-eating

populations do not scale-feed as often as they would in the wild and thus lab-reared populations

of scale-eaters may not develop the notably increased adductor muscle mass. Interestingly, the

distinct differences in adductor mandibulae insertion height among wild populations disappeared

in the lab-reared populations (figure S3), indicating phenotypic plasticity associated with the

common diet of pellet foods and lack of scale-eating attacks in the lab.

We found that 'wide-mouth' ingest scales, but at a significantly lower frequency than *C.*

desquamator (Wilcoxon Rank Sum test, $P = 0.004$; figure 3A). We did not detect any scales in

C. variegatus guts (figure 3A). Detritus made up the only other gut contents present besides

scales in all the specimen of *C.* sp 'wide-mouth' from Osprey Lake that we dissected. This was

similar for the other scale-eater *C. desquamator* individuals from Osprey Lake, in which we also

found only scales and detritus in their gut. For the generalist species *C. variegatus*, their guts

contained only detritus except for a single individual in which we found a snail shell. The low

diversity in gut contents among species may be due to the relatively small sample size of

individuals we had that we could perform gut content analysis on for this study (n=10 for each

species). However, the predominance of detritus in the gut contents of Osprey Lake populations

of all three species is not surprising given the thick (<1m) bottom layer of mud and flocculent in

this lake[30]. Additionally, detritus made the largest percentage (49-71%) of San Salvador Island

pupfish diet in other lakes across the island as well. A previous study that conducted gut content

analyses with 2-4X larger sample sizes also found that the majority of *C. variegatus, C.*

brontotheroides, and *C. desquamator*'s diet is detritus. Except for one population of *C.*

desquamator in Crescent Pond where scales made up 50% of their gut contents and detritus only

made up 30%. Therefore, the ecological intermediacy of *C.* sp 'wide-mouth' is supported by

their intermediate ratio of scales to detritus, which is the predominant axis of dietary divergence

between *C. variegatus* and *C. desquamator* in this lake. The lack of other dietary items in the gut

contents of *C.* sp 'wide-mouth' further supports our arguments that their ecological divergence is

along the same specialist axis of scale-eater *C. desquamator* and that they share little ecological

overlap with the other molluscivore specialist *C. brontotheroides.*

1013
1014   'Wide-mouth' did not result from hybridization between *C. variegatus* and *C. desquamatory*

1015

Several lines of genomic evidence support the 'wide-mouth' ecomorph as a distinct species.

First, 'wide-mouth' individuals do not occupy an intermediate position between *C. desquamator*

and *C. variegatus* along either of the first two principal components that represent the major axes

of genetic variation in the radiation (figure S4A). In the ADMIXTURE analysis, 'wide-mouth'

individuals share more ancestry with each other than with either *C. desquamator* or *C. variegatus*

under the two most likely K values (K=6 and 7; figure S4B&5). Lastly, significant positive f_3-

statistics and the non-significant f_4-statistics support our inference that the 'wide-mouth' is not a

recent admixed population of any pairwise combinations of the *C. brontotheriodes*, *C.*

variegatus, or *C. desquamator* populations (table S1).

1025
C. sp. 'wide-mouth' did not result from hybridization between *C. variegatus* and *C. desquamator*

We compared 28 different demographic models of divergence and gene flow on San Salvador

Island (table S2) that represented all possible topologies among the four species and explored

zero, early, and current gene flow scenarios. In the best supported model, 'wide-mouth' was

sister to *C. desquamator* with current gene flow and a divergence time estimate of 11,658 years

ago (95 CI: 8,257-20,113 years; figure 4; table 1 and S2). This model indicates that the ancestor

to 'wide-mouth' and *C. desquamator* populations first diverged from *C. variegatus* 15,390 years

ago (95% CI: 10,722-23,927 years; figure 4B). This estimate of the origin of the radiation

overlaps well with geological age estimates based on the filling of hypersaline lakes on San

Salvador Island after the end of the last glacial maximum period (~6-19K years ago; [31,32]). *C.*

variegatus and *C. brontotheriodes* populations in Osprey Lake diverged recently about 462 years

ago (95% CI: 411-1,121 years; figure 4B).

1038
<u>Shared and unique adaptive alleles in *C.* sp. 'wide-mouth' and *desquamator*</u>

We discovered that the two scale-eating populations shared a set of the same adaptive alleles not

found in *C. brontotheriodes* or *C. variegatus* (Fig 5D). This consisted of 15 alleles in 6 shared

hard selective sweeps in *C. desquamator* and 'wide-mouth': 10 SNPs were in unannotated

regions, two were in the introns of the gene *daam2,* and three were in regulatory regions of the

genes *usp50, atp8a1,* and *znf214* (one variant each).

Shared adaptive alleles in the gene *daam2,* a wnt signaling regulator, are intriguing

because knockdown of this gene causes abnormal snout morphology, osteoporosis, and changes

in insulin and alkaline phosphate levels in mice [33], and abnormal cranial and skeletal

development in zebrafish [34]. Craniofacial morphology is one of the rapidly diversifying traits

in this system, suggesting that divergence in *daam2* may play a role in the shared craniofacial

divergence of the two scale-eating species. Similarly, *usp50* functions in protein metabolism and

deubiquination [35] and may play a role in shared metabolic adaptations to the higher-protein

content of a scale-eating or snail-eating diet (also note microbiome divergence of *C.*

desquamator with increased prevalence of collagenase-digesting bacteria when reared in a

common garden:[36]). *atp8a1* is an ion transporter [37], *znf214* is a transcription binding factor,

and the ten unannotated variants were not associated with lower jaw size variation in a previous

genome-wide association study of the radiation (Richards et al. 2021).

Despite the divergent craniofacial features of the 'wide-mouth', none of the adaptive

alleles unique to the 'wide-mouth' appear to be in or near genes annotated for craniofacial

phenotypes in model organisms. In *C. desquamator,* three of the 13 sets of unique adaptive

alleles are in or near genes annotated for craniofacial phenotypes: a *de novo* non-synonymous

coding substitution in the gene *twist1*, several putative regulatory variants near the gene *gnaq,*

and 8 variants in and near the gene *bri3bp*, which is located inside a QTL region for cranial

height in pupfish (St. John et al. 2021). In *C. brontotheriodes*, there is also a candidate

craniofacial adaptive allele: a non-synonymous coding substitution in the gene *kat6b*, which is

associated with abnormal craniofacial morphologies, including shorter mandibles in mice [38,39]

and Ohdo Syndrome and bulbous noses in humans [40].

This pattern of unique alleles relevant to craniofacial phenotypes in *C. brontotheriodes*

and *C. desquamator*, but not 'wide-mouth', holds even if we lower the threshold to the top 1

percentile of F_{st} between specialists and generalist (see supplemental for more details). 'Roof of

mouth development', 'bone development' and 'skeletal system development' are in the top 20

most significantly enriched GO terms for *desquamator* candidate adaptive alleles, with 6

significantly enriched terms relevant to cranial and skeletal development (out of 165 enriched

terms total). Similarly, 'embryonic skeletal system development' and 'skeletal system

development' were significantly enriched terms for *brontotheriodes* candidate adaptive SNP

alleles (out of 8 enriched terms total). However, 'wide-mouth' adaptive alleles at this lower F_{st}

threshold were not significantly enriched for any GO terms related to craniofacial or skeletal

morphology (n=52 terms; Fig S7).

1078

<u>Substantial history of admixture in *C.* sp 'wide-mouth'</u>

To further investigate the signatures of adaptive introgression in this species, we

modeled admixture across San Salvador Island and several key generalist populations in the

Caribbean with f_3- and f_4-statistics (table S11;figure S10). Previous signatures of admixture

involving the base of SSI radiation and outgroup generalist populations [25] were detected

again in our admixture graph, alongside a new signature of admixture that appears

to be secondary contact between a pupfish lineage most closely related to North Carolina in our

dataset and *C.* sp 'wide-mouth' (table S11; figure 5 & S10).

1087
<u>Shared signatures of selection across the three specialists in the radiation</u>

We found that a higher percentage of the genome was under positive selection in all specialist

populations compared to the *C. variegatus* population (figure 5A). This pattern might be

expected given the divergent selection pressures the specialists face to adapt to different trophic

niches. Of all the specialists, *C. brontotheriodes* exhibited both the most selective sweeps and the

longest selective sweeps (figure 5A). This suggests that *C. brontotheriodes* have undergone

selection most recently among all the species in Osprey Lake and is supported by the recent

divergence time between the *C. brontotheriodes* and *C. variegatus* populations (figure 4D). The

shared selective sweeps were shorter than any of the selective sweeps unique to each of the

specialists, suggesting that selection for these shared regions was not the most recent or strongest

in any of the specialists (figure S11C).

Additionally, all specialists, including the two scale-eating species, were more genetically

diverged from each other than to *C. variegatus* species based on the number of SNPs fixed or

nearly-fixed between them (figure 5B). This pattern of stronger genetic divergence between

specialists also held across fixed SNPs, the top 1% of F_{st}, and genome-wide average F_{st} (table

S4). Next, we identified a set of candidate adaptive alleles for each specialist by filtering for

SNPs that were fixed or nearly fixed (F_{st}, ≥ 0.95) relative to *C. variegatus* and found within a

hard selective sweep in the focal specialist. In this set of candidate adaptive alleles, *C.*

brontotheriodes had the fewest adaptive alleles while *C. desquamator* had the most (figure 5B).

All three specialists, including the two scale-eating species, were more genetically diverged from

each other than to *C. variegatus* based on the number of SNPs fixed or nearly-fixed between

them across multiple F_{st} thresholds (figure S11B; table S4).

Discussion
1111
The hallmark of adaptive radiation is a rapid burst of diversification which is predicted by theory

to slow down over time as niche subdivision increases [41]. An alternative possibility is that

radiations can be self-propagating and that the diversity generated within the first stages of

radiation helps beget further diversity [42]. This could happen through exploitation of new

trophic levels created by new species or physical alterations of the environment by new species

that may create additional opportunities for speciation (reviewed in [41,43]). The diversity begets

diversity hypothesis can also be visualized as the exploration of a complex multi-peaked fitness

landscape; as species in the radiation colonize new peaks, this provides access to additional

neighboring fitness peaks to fuel rapid radiation. At the microevolutionary level, shared adaptive

variants can also help other populations colonize new or ecologically similar areas of the fitness

landscape. However, the genetic basis and microevolutionary processes underlying major

ecological transitions are still poorly understood in nature.

1124

An adaptive walk underlies the major ecological transition from generalist to scale-eating

specialist

One of the foundational models of adaptation is that it proceeds in 'adaptive walks' towards

fitness optima that involve the sequential fixation of adaptive alleles that move a population in

the phenotypic direction of the local optimum [44–48]. The distinct timing of selection across

different adaptive alleles suggests that the ecological transition from generalist to novel scale-

eating specialist involved such an adaptive walk rather than a sudden burst of concurrent

selection events after some major environmental shift. These distinct, multiple bouts of selection

could be caused by mutation-limited (i.e. waiting for new beneficial alleles; [49–51]) or

mutation-order processes (i.e. epistatic interactions; [52–54].

We found that all of the shared adaptive alleles between the scale-eating populations

occurred at low frequency in generalist populations on other Caribbean islands in a previous

study ([25]; Table S7). For example, three copies of the shared *usp50* adaptive allele were found

outside of San Salvador Island (Table S7). This indicates that initial bouts of selection occurred

on available standing genetic variation and that staggered timing of hard selective sweeps in each

trophic specialist most likely reflects mutation-order processes in which selection on a beneficial

allele was contingent on prior fixation of other adaptive alleles in each specialists' genetic

background. However, several adaptive alleles originated from introgression or de novo

mutations found only on San Salvador Island (Table S7), so part of the adaptive walk may have

also been mutation-limited.

1145

1146

Cryptic scale-eating species reveals features of the adaptive walk towards scale-eating

specialization

Although the two scale-eating species shared a set of adaptive alleles and a recent common

ancestor, 5 of 6 shared adaptive regions swept at significantly different times between the two

species. This difference in timing may result from several different scenarios: 1) independent

adaptive walks to the same scale-eating niche, 2) independent adaptive walks to different scale-

eating niches or 3) the adaptive walk of one population depends on the adaptive walk of the other

population. We explore each of these scenarios in turn.

First, the difference in timing of selection on the same shared adaptive alleles could

indicate independent adaptive walks to the same scale-eating peak that occurred at different

times and/or by slightly different routes. *C. desquamator* and 'wide-mouth' populations have

predominantly abutting ranges, with only a small amount of geographic overlap on San Salvador

Island in four lakes, Osprey, Oyster, Little Lake, and Mermaid's Pond (Fig 1). If this current

distribution is representative of their historical ranges and the two lineages began diverging

about 11,000 years ago, it is possible that the adaptive walks took place in different lakes and

were largely independent of one another. The differences in timing on the shared adaptive alleles

and the presence of unique alleles observed in each population are reminiscent of mutation-order

speciation [53,55]. In mutation-order speciation the same alleles are favored in populations that

are adapting to similar environments, yet by chance and/or epistatic interactions with different

genetic backgrounds, similar adaptive alleles fix in just one population. At least one set of alleles

near the gene *slitrk5* are unique to the 'wide-mouth' scale-eater and fixed well before selection

occurred on the shared adaptive variants between the two scale-eater populations (Fig 5).

Epistatic interactions of these *slitrk5* alleles may have prevented the 'wide-mouth' from fixing

additional segregating alleles that are uniquely fixed in *C. desquamator.*

Second, the difference in timing of selection may have occurred because the two scale-

eating populations are adapting to two different scale-eating niches. The intermediate diet,

distinctly sized morphological traits, and smaller body size in the 'wide-mouth' scale-eater

compared to *C. desquamator* may indicate the 'wide-mouth' scale-eater is adapting to a different

scale-eating niche. While the two scale-eating populations do share some overlap in adaptive

alleles and selective sweeps, the majority of selective sweeps are unique to each species (Fig 5 &

6), including neurogenesis, brain, and nervous system development (*slitrk5, sema4g, and*

smarce1), whereas unique adaptive alleles in *desquamator* include craniofacial development

annotations (*olfm1, gnaq, twist1*) as well. This difference in gene annotations can also be seen at

the broader level of regions of the genome under selection (Fig S7). Therefore, the differences in

timing at shared adaptive alleles might reflect differences in relative strength of selection due to

different selective regimes experienced by the two scale-eating populations. While starTMRCA

is fairly robust in its timing estimates to different strengths of the selection [20], we cannot rule

out that differences in relative timing of selection might partially reflect differences in strength of

selection occurring in the two scale-eating populations.

Third, another scenario that could have led to the older timing of selection across shared

adaptive alleles in *desquamator* than 'wide mouth' is one in which the adaptive walk of one

population depended on adaptive alleles from another. The significantly older timing of

selection in *desquamator* on shared adaptive alleles and the low absolute genetic divergence

between these two scale-eating population in these regions (Fig 6) suggests that these alleles may

have introgressed between the two populations. Given that these shared adaptive alleles appear

as standing genetic variation across Caribbean populations, albeit at very low frequency in our

sampling, it is likely that the shared alleles were present in the ancestor and were segregating in

both populations after they diverged. There are several reasons why adaptive introgression may

have been necessary for adaptive divergence of 'wide-mouth' despite the alleles initially

segregating in the ancestral population. In line with mutation-order processes (e.g. Mani and

Clarke 1990; Schluter 2009; Good et al. 2017), these alleles might not have been adaptive until

the right genetic background was present in the 'wide-mouth'. Introgression from *C.*

desquamator in which the allele was already swept to high frequency could have raised the

frequency of these alleles in 'wide-mouth' and increased the likelihood of fixation for the

beneficial alleles (reviewed in [56]). This is consistent with previously proposed hypotheses for

how diversity begets further diversity in adaptive radiations (Whittaker 1977; Stroud and Losos

2016; Martin and Richards 2019).

1204

Shared aspects of genomic landscape of selection among all specialists additionally supports

specialization promoting further specialization

Intriguingly we also find evidence of selective sweeps shared across all three specialists despite

their divergent adaptations and lack of a shared specialist ancestor. However, there are no

commonly shared adaptive alleles fixed against *C. variegatus* in these shared selective sweep

regions. Hard selective sweeps indicate a selection scenario in which a single beneficial allele of

large effect on a trait is swept to high frequency (reviewed in [57]). The lack of highly divergent

alleles in these regions might indicate polygenic selection events underlie the shared signatures

across specialists. The broad-scale transition from dietary generalist to dietary specialist may

involve polygenic selection with small shifts in the frequency of many alleles. Despite very

different expectations about the genetic basis of adaptation, polygenic selection events may have

been detected as a hard selective sweep in this study as the two selection types can be

challenging to distinguish based solely on patterns of genetic variation [58–61]. Recently

developed frameworks provide additional criteria to help distinguish between the two selective

regimes [59] but are beyond the scope of this current study.

These shared selective sweeps across all specialists are not the longest selective sweeps

in any of the specialist genomes (> 50-kb; Fig 5C), indicating they are not the most recent or

strongest selection events. However, these shared selective sweeps appear to be relevant to

dietary specialization as these regions are enriched for genes annotated for metabolic processes

such as short chain fatty acid and propionate metabolism (Fig 5). Propionates and other short

chain fatty acids are common microbiome metabolites [62]. A recent microbiome study in lab-

reared populations found that *C. desquamator* microbiomes appear enriched for

Burkholderiaceae bacteria that can digest collagen, further supporting an adaptive role for

microbiomes in the radiation [36].

We also see pairwise shared selective sweeps between all combinations of two

specialists. In contrast to the sweeps shared between all specialists, we see stronger genetic

divergence among species in these regions and a proportion of the selective sweeps shared

between two specialists do contain fixed or nearly-fixed alleles. However, the specialists that

share selective sweeps in these regions are fixed for different adaptive alleles, consistent with

previously observed patterns of parallel differential gene expression in specialists despite

divergent genotypes [63]. These alleles were possibly under balancing selection in the ancestor

(most of these alleles are segregating at intermediate frequency in *C. variegatus* currently) and

the alternate alleles were driven to fixation between the specialists during their respective

divergences. While the species have discrete phenotypes across several morphological traits

important to divergence in this system, these traits often lie on a continuous axis (i.e. shorter oral

jaw lengths in *C. brontotheriodes*, intermediate jaw lengths in *C. variegatus*, and longer oral jaw

lengths in 'wide-mouth' and *C. desquamator*). Some of the alleles that are alternatively fixed

between specialists at the same position may have incomplete dominance effects on such traits.

In line with adaptation to divergent trophic niches across the species, we do still see many

unique signatures of selection among the specialists. The *C. variegatus* population had the fewest

selective sweep signatures. All three specialists had more selective sweeps potentially resulting

from strong directional selection for adapting to new fitness peaks. However, while we might

expect more selective sweeps in *C. desquamator* genomes given the deeper valley on the fitness

landscape isolating the scale-eater phenotypes [3,64], C. *brontotheriodes* had the most selective

sweeps detected in their genome. They also have the longest sweeps detected (e.g. 100-kb),

suggesting that selection may have occurred most recently in the *C. brontotheriodes*. This

matches with the most recent divergence event being between generalist and *C. brontotheriodes*

in Osprey Lake. An intriguing implication of this result is the potentially long wait time for

speciation of *C. brontotheriodes* species compared *C. desquamator* despite the shallower fitness

valley isolating *C. brontotheroides* from generalists. However, we cannot yet rule out that the

recent divergence time estimated in Osprey Lake reflects a more recent arrival of *C.*

brontotheriodes to that lake after initially diverging long ago in another lake not incorporated in

our demographic model.

1258

1259

1260

1261
Fig S1. Changes in effective population size over time for Osprey populations. Inferred using
MSMC [13] on high-coverage (18-36X) genomes from single individuals of each of the four
species in Osprey Lake, San Salvador Island Bahamas: *C. variegatus* (gold), *C. brontotheroides*
(purple), *C. desquamator* (teal), and C. sp. 'wide-mouth' (red-orange). Time was scaled using a
mutation rate of 1.56x10-8 mutations/basepair/generation and a generation time of 1 year.
1267

1268

1269

Figure S2. Relationships between three divergent traits and standard length across ponds.
The relationship between standard length (mm) of individuals and their D) lower jaw length, E)
buccal cavity width, and F) adductor mandibulae muscle insertion height across individuals the
three species *C. desquamator* (teal), *C. variegatus* (gold), and *C.* sp. 'wide-mouth' (red-orange)
across different ponds (Osprey, Oyster, and Great Lake). Colored lines represent linear model of
these relationships for each species with their 95% confidence bands in gray.
1276
1277

1278

1279

1280

1281

1282

1283

1284

Figure S3. Comparison of the three focal divergent traits in lab and wild populations. A. 95% CI of the standardized sizes of 8 traits measured (see y-axis label on each panel for particular trait) for *C. variegatus* (gold;n=5), *C.* sp. 'wide-mouth' (red-orange;n=20) and *C. desquamator* (teal;n=20). Each plot is broken into two panels of wild caught and lab reared individuals (wild: circle; lab: triangle) represent the mean values of traits from each population. Confidence intervals that overlap between species are highlight in black. Lab raised individuals include F1, F2 and F3 generations. AM insertion height is a proxy for cross-sectional area of the adductor mandibulae

1296

1297
Figure S4. Trophic level comparison among populations in Osprey Lake. d15N values from
isotope analysis of muscle tissue from 48 individuals across the three species that span the
ecological spectrum from generalist to specialized scale-eater: generalist *C. variegatus*
(var;gold), *C. sp.* 'wide-mouth' (wid;red-orange), and *C. desquamator* (des; teal). ANOVA
results indicate significant differences in d15N levels across populations indicating that each
occupies a distinct trophic level.
1304
1305

1306
1307 **Figure S5. *C.* sp. 'wide-mouth' population did not result from recent hybridization.** A)
1308 Principal components analysis of the four focal groups on San Salvador Island based on an LD-
1309 pruned subset of genetic variants (78,840 SNPs). B) Ancestry proportions across individuals of
1310 the four focal groups. Proportions were inferred from ADMIXTURE analyses with 2 values of K
1311 with the highest likelihood on the same LD-pruned dataset in A.

1312
Figure S6. *C.* **sp. 'wide-mouth' is not an admixed population**. ADMIXTURE analysis of San
Salvador Island populations using an LD-pruned subset of genetic variants from 109 individuals
(78,840 variants). A) Cross validation error estimates from ADMIXTURE that indicate a model
of K=6-7 was the best fit. B) Ancestry proportions across individuals of the three focal groups.
Proportions were inferred from ADMIXTURE analyses with 2 values of K with the highest
likelihood on the same LD-pruned dataset in A.
1319
1320
1321

1322
Fig S7. Demographic history of San Salvador Island radiation. A) Best supported
demographic model in which *C. desquamator* diverged from the *C.* sp. 'wide-mouth' based on
an LD-pruned dataset with no missing information across all Osprey Lake individuals (67,400
SNPs). Divergence time is shown with the maximum likelihood point estimate from the run with
the best fit and the 95% confidence interval for that parameter estimate based on 100 bootstrap
replicates. B) Maximum likelihood point estimate and 95% confidence intervals for divergence
time based on 100 bootstrap replicates from this model. C) Maximum likelihood point estimate
and 95% confidence intervals for migration rate parameters involved in the best fitting model
depicted in units of the number of migrants per generation.

1341
1342 **Fig S8. Top five best fitting demographic models for Osprey Lake.** From a genetic dataset
1343 that was LD-pruned and had no missing information across individuals comprised of 67,400
1344 SNPs. Listed in order of lowest AIC scores (Table 1 and S2) with best supported model in A. All
1345 top models were of recent bidirectional gene flow but varied in topology among the four species.

1346
1347
Fig S9. Genetic divergence among populations in Osprey Lake in regions of shared
selection across two or more specialists. Each panel represents pairwise F_{st} comparison
between two populations for every SNP found in regions under shared selection across two or
more specialist species on San Salvador Island: C. brontotheroides (bro); C. desquamator (des),
C. sp. 'wide-mouth' (wid). Regions shared across all three specialists (bro+des+wid) do not
contain highly divergent SNPs between any of the species in the radiation. Highly divergent
SNPs (Fst > 0.75) are only found in regions shared between one or two specialists.
1355

1356
Fig S10. Top 20 enriched GO categories for divergent alleles in each of the three specialists.
GO enrichment analyses were performed on genes in or near (within 20-kb) of a SNP that was
under a hard selective sweep and strongly diverged from generalist species (top 1% of Fst values
across genome) for a) 'wide-mouth' b) *C. desquamator*, and C) *C. brontotheriodes*. GO
categories that were significantly enriched for relevant terms corresponding to craniofacial
development, the major axes of morphological divergence in this radiation, are highlighted in
red. All terms included were significant at an FDR < 0.05 and full list of terms in Data S1-3.
1364
1365
1366

1367
1368 Figure S11. Caribbean pupfish admixture graph. Best-fiting admixture graph from
1369 *qpGraph* approach that estimated f_3-statistics across Caribbean populations. This admixture
1370 model features 7 admixture events represented by admixture edges and proportions (percentages
1371 on dotted branches) and drift edges (numbers on solid branches) that minimize the difference
1372 between fitted and estimated f_3-statistics.
1373

1374

1375

Figure S12. Patterns of selection and genetic divergence in specialist genomes. A) Selective
sweep length distributions across all four San Salvador Island species. Rug plot below each
histogram represents the counts of selective sweeps in different length bins. B) The total number
of fixed or nearly-fixed SNPs ($F_{st} \geq 0.95$) between each group in Osprey Pond. C) Ridgeline
plots for length distributions of selective sweeps shared between different combinations of
specialists (*C. brontotheriodes*: bro, *C. desquamator*: des, *C.* sp. 'wide-mouth': wid). D) GO
enrichment analysis of the 57 genes contained in the shared selective sweep regions across all
specialists. E) The number of adaptive alleles (fixed or nearly-fixed SNPS [$F_{st} \geq 0.95$] relative to
C. variegatus and under selection in each population of specialists in Osprey Lake. F) Venn
diagram highlights those adaptive alleles that are unique to each specialist and shared with other
specialists.

1387

1388

1389

1390

1391

Table S1. No significant genome-wide signature of admixture among populations on San Salvador Islands. Results from three formal tests for introgression to assess whether 'wide-mouth' populations are hybrids between generalist *C. variegatus* and scale-eater *C. desquamator*: D-, *f₃*- and *f₄*-statistics across all possible combinations of Osprey Lake populations of generalist species. Significant *f₃*-statistic had Z-scores > -2; significant *f₄* and D-statistic Z-scores were smaller than -2 and greater than 2. The only significant signature of admixture (bolded Z-scores) comes from D- and *f₄*-statistics based on relationships that violate the expected tree (((C. desquamator; 'wide-mouth'),Generalist),Outgroup) and therefore should not be interpreted as evidence of 'wide-mouth' being an admixed population. *C. variegatus* population from Rum Cay, the nearest neighbor island in the Bahamas to San Salvador Island was used as an outgroup population for D- and *f₄*-statistics.

f_3-statistic						
A	B	C	--	f_3	stderr	Z
C. variegatus	*C. desquamator*	C. sp. 'wide-mouth'		0.004626	0.00088	5.257
f_4- statistic						
P1	P2	P3	P4	f_4	stderr	Z
C. sp 'wide-mouth'	*C. desquamator*	*C. variegatus*	Rum Cay	-0.000196	0.000164	-1.2
C. sp 'wide-mouth'	*C. variegatus*	*C. brontotheroides*	Rum Cay	0.000186	0.000201	0.927
D-statistic						
P1	P2	P3	Outgroup	D	stderr	Z
C.sp.'wide-mouth'	*C. desquamator*	*C. variegatus*	Rum Cay	-0.0018	0.001498	-1.199
C. sp. 'wide-mouth'	*C. variegatus*	*C. brontotheroides*	Rum Cay	0.0017	0.001792	0.931

Table S2. Support for the 28 demographic models for the evolution of the 'wide-mouths' from the site frequency spectrum. The likelihood and AIC scores all demographic models

Divergence Model	LnL	ΔLnL	AIC	ΔAIC
(1) ((des,wid),(bro,var)) + (des,wid) diverges first + recent gene flow	-11701	1077	54133	0
(2) ((des,bro),var)) ,wid)) + recent gene flow	-11780	1112	54296	35
(3) ((des,bro),wid),var)) + recent gene flow	-11806	1138	54418	61
(4) ((bro,wid),des),var)) + recent gene flow	-11808	1140	54423	63
(5) ((des,wid),(bro,var)) + (bro,var) diverges first + recent gene flow	-11846	1178	54602	101
(6) ((des,wid), bro),var)) + recent gene flow	-11855	1187	54642	110
(7) ((des,bro),var)) ,wid))+ early gene flow	-12016	1348	55364	271
(8) ((des,var),bro), wid)) + recent gene flow	-12024	1356	55426	279
(9) ((wid,bro),des),var)) + early gene flow	-12086	1418	55688	341
(10) ((wid,des),bro),var)) + early gene flow	-11855	1419	55692	342
(11) ((des,wid),(bro,var)) + early gene flow	-12139	1471	55929	394
(12) ((des,wid), bro),var)) + no gene flow	-12315	1648	56738	570
(13) ((wid,bro),des),var)) + no gene flow	-12326	1658	56787	581
(14) ((des,bro),var)) ,wid))+ no gene flow	-12334	1667	56826	589
(15) ((des,wid),(bro,var)) + recent sister gene flow only	-12394	1726	57106	649
(16) ((des,wid),(bro,var))+ no gene flow	-12396	1728	57109	651
(17) ((wid,var)),bro),des)) + recent gene flow	-12399	1731	57148	654
(18) ((bro,var)) ,wid),des)) + recent gene flow	-12401	1733	57157	656
(19) ((wid,bro),var)),des)) + recent gene flow	-12401	1734	57163	657
(20) ((des,wid),var),bro)) + recent gene flow	-12420	1752	57244	675
(21) ((wid,var)),des),bro)) + recent gene flow	-12425	1757	57265	680
(22) ((des,var),wid),bro)) + recent gene flow	-12425	1757	57267	680
(23) ((wid,var)),(bro,des)) + no gene flow	-23245	12577	107070	11500
(24) ((wid,var)),(bro,des)) + early gene flow	-23658	12990	108976	11913
(25) ((wid,var)),(bro,des)) + recent gene flow	-23659	12991	109003	11914
(26) ((des,var),(bro,wid)) + early gene flow	-34983	24315	161129	23238
(27) ((des,var),(bro,wid)) + no gene flow	-34983	24315	161125	23238

| (28) ((des,var),(bro,wid)) + recent gene flow | -34983 | 24315 | 161149 | 23238 |

Table S3. Parameters for selective sweep analyses in SweeD.
The 95[th] percentile of composite likelihood ratio threshold based on neutral simulations under the
demographic scenario of decreasing population size through time inferred with MSMC (Fig S4),
and the population size change and haplotype number information parameters required by
SweeD for each species.

Species	CLR threshold	SweeD Commands
C. variegatus	0.40320	-folded -strictPolymorphic -G 0.407 -eN 5.45 181.8 -s 24
C. brontotheroides	0.43473	-folded -strictPolymorphic -G 0.389 -eN 5.88 196 -s 32
C. desquamator	0.40537	-folded -strictPolymorphic -G 0.218 -eN 8.11 270 -s 16
C. sp. 'wide-mouth'	0.39262	-folded -strictPolymorphic -G 0.203 -eN 8.57 276 -s 22

**Table S4. Genetic divergence among the four populations in Osprey Lake, San Salvador
Island across different thresholds of F_{st}.** Relative measure of genetic divergence was

1472 calculated in pairwise combinations of the different species as the number of fixed SNPs
1473 between them, the number of fixed or nearly fixed SNPs between them, the top 1% of F_{st}
1474 between them from the distribution of F_{st} between SNPs and the genome-wide average F_{st} across
1475 all SNPs in the genome.

F_{st} Comparison	Number of fixed SNPs ($F_{st} =1$)	Number of nearly fixed SNPS ($F_{st} \geq 0.95$)	Top 1%	Genome-wide average
C. sp. 'wide-mouth' vs *C. desquamator*	5,212	5,915	0.84	0.131
C. sp. 'wide-mouth' vs *C. brontotheroides*	5238	14,602	0.85	0.15
C. sp. 'wide-mouth' vs *C. variegatus*	247	869	0.68	0.104
C. desquamator vs *C. brontotheroides*	6941	22,088	0.86	0.17
C. desquamator vs *C. variegatus*	414	1,964	0.71	0.116
C. variegatus vs *C. brontotheroides*	110	567	0.637	0.09

1476
1477 **Table S5. Comparison of the average absolute genetic divergence (D_{xy}) between the four**
1478 **species in Osprey Lake 'wide-mouth', *C. variegatus*, *C. desquamator* and *C. brontotheriodes***
1479 **and the shared adaptive alleles between scale-eater populations.**
1480

Comparison	Mean D_{xy}
C. variegatus vs *C. brontotheroides*	0.167
C. variegatus vs *C. desquamator*	0.169
C. variegatus vs *C.* sp. 'wide-mouth'	0.165
C. brontotheroides vs *C. desquamator*	0.162
C. brontotheroides vs *C.* sp. 'wide-mouth'	0.156
C. desquamator vs *C.* sp. 'wide-mouth'	0.157
Selective sweeps in *C.* sp. 'wide-mouth'	0.123
Shared genetic variants with *C. desquamator*	0.0814
Unique genetic variants in *C.* sp. 'wide-mouth'	0.24

Table S6. The distribution of shared adaptive alleles between the two scale-eating species
across the Caribbean. Numbers represent the copies of the *C. desquamator* and 'wide-mouth'
allele present in outgroup populations collected from
San Salvador Island (SS), Rum Cay (RC), Cat Island (CT), Exumas Islands (EX), Long Island
(LG), Mayaguana (MG), New Providence Island (NPI) in the Bahamas. Lagunas Bavaros
(BAV) and Etang Saumautre (ETA) in Dominican Republic, Sarasota Estuary in Florida (FL),
Curacoa (CUR), Caicos Island (CAI) and Isla Margarita (MAR) in Venezuela.
1493

| Allele | Gene | Distribution | *C. variegatus* | | | | | | | | *C. laciniatus* | *C. higuey* | *C. bondi* | *C. dearborni* | | |
			SS	RC	CT	EX	LG	MG	FL	NC	NPI	BAV	ETA	CUR	CAI	MAR
HiC_scaffold_6:907101	NA	SGV	0	0	1	0	0	0	0	12	0	0	0	0	0	0
HiC_scaffold_6:907425	NA	SGV	1	0	0	0	0	0	0	7	0	0	0	0	0	0
HiC_scaffold_6:907665	NA	SGV	0	0	0	0	0	0	0	7	0	0	0	0	0	0
HiC_scaffold_6:908229	NA	SGV	1	0	1	0	0	0	0	6	3	0	0	0	0	0
HiC_scaffold_6:908889	NA	SGV	1	0	0	0	0	0	0	6	0	0	0	0	0	0
HiC_scaffold_6:923182	NA	SGV	0	5	0	1	0	0	1	0	0	0	0	0	0	0
HiC_scaffold_6:923590	NA	SGV	0	5	0	2	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_6:924203	NA	de novo	0	0	0	0	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_6:929621	NA	SGV	1	2	1	0	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_9:31763392	daam2	SGV	6	3	0	1	0	1	1	1	6	1	0	0	0	0
HiC_scaffold_9:31763400	daam2	SGV	8	5	0	1	1	0	1	8	10	8	0	0	0	0
HiC_scaffold_24:5390450	usp50	SGV	1	1	0	0	0	0	0	0	1	0	0	0	1	0
HiC_scaffold_43:27385265	NA	SGV	3	1	0	2	1	1	0	5	1	0	1	2	0	11
HiC_scaffold_43:27389200	NA	SGV	5	1	0	0	1	1	1	12	12	2	1	3	1	12
HiC_scaffold_44:14963656	atp8a1	SGV	3	1	0	1	0	1	1	12	12	2	1	3	1	12
HiC_scaffold_60:1798737	znf214	SGV	3	0	0	0	0	1	0	0	2	0	0	0	0	0

1519
1520

Allele	Gene	Distribution	*C. variegatus*								*C. laciniatus*	*C. higuey*	*C. bondi*	*C. dearborni*		
			SS	RC	CT	EX	LG	MG	FL	NC	NP	BAV	ETA	CUR	CAI	MAR
HiC_scaffold_1:29498971	NA	SGV	2	1	0	1	1	2	2	0	0	0	2	2	0	22
HiC_scaffold_1:29505055	NA	SGV	6	0	0	0	0	0	0	24	0	0	0	0	0	0
HiC_scaffold_1:29518877	NA	de novo	4	0	0	0	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_1:36701070	NA	SGV	2	0	0	0	0	0	0	24	7	9	0	0	0	0
HiC_scaffold_1:36701336	NA	SGV	2	3	0	0	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_1:36704196	NA	SGV	4	1	0	0	0	0	0	24	5	10	0	0	0	0
HiC_scaffold_1:36705430	NA	SGV	3	1	0	0	0	0	0	24	5	9	0	0	0	0
HiC_scaffold_1:36705978	NA	SGV	1	1	0	0	0	0	0	22	7	10	0	0	0	0
HiC_scaffold_1:36707870	NA	SGV	4	4	2	2	0	2	0	0	0	0	0	0	2	0
HiC_scaffold_1:36713418	NA	SGV	1	0	0	0	0	0	0	24	7	12	2	4	0	0
HiC_scaffold_1:36715711	NA	Intro NC	4	6	3	4	0	2	0	23	9	11	2	4	2	0
HiC_scaffold_1:36715712	NA	Intro NC	4	6	3	4	0	2	2	23	9	11	0	4	2	0
HiC_scaffold_1:36740602	NA	SGV	5	5	2	4	1	1	2	20	4	20	2	4	0	22
HiC_scaffold_1:36741004	NA	SGV	2	4	2	2	1	0	0	22	7	7	0	4	0	24
HiC_scaffold_1:36742540	NA	SGV	2	0	0	0	0	0	0	0	0	1	0	0	0	0
HiC_scaffold_1:36743115	NA	SGV	4	6	2	0	1	0	2	24	6	7	0	4	0	24
HiC_scaffold_1:36744639	NA	SGV	0	1	0	0	0	0	0	0	0	2	0	0	1	0
HiC_scaffold_1:36744717	NA	SGV	1	2	0	2	1	2	2	24	5	5	2	4	2	23
HiC_scaffold_1:36744821	NA	SGV	1	1	0	1	0	0	1	0	0	2	0	0	1	0
HiC_scaffold_1:36745174	NA	SGV	2	6	2	6	1	4	2	22	5	18	2	4	2	24
HiC_scaffold_1:36745293	NA	SGV	3	4	4	6	1	2	2	24	6	20	2	4	2	24
HiC_scaffold_1:36748002	NA	SGV	0	0	0	0	0	0	0	24	0	5	0	0	1	0
HiC_scaffold_5:5681542	NA	de novo	1	0	0	0	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_5:6082906	NA	SGV	3	9	0	2	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_5:6082926	NA	SGV	3	9	0	2	1	0	1	24	0	6	2	0	0	24
HiC_scaffold_5:8296047	NA	de novo	1	0	0	0	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_5:8465173	NA	de novo	2	0	0	0	0	0	0	0	0	0	0	0	0	0

HiC_scaffold_5:21455838	NA	SGV	6	0	0	2	0	0	0	8	0	6	0	2	1	0
HiC_scaffold_8:14526358	NA	de novo	0	0	0	0	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_11:10913509	NA	de novo	3	0	0	0	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_11:10948683	cd226	de novo	5	0	0	0	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_11:12963188	NA	SGV	1	0	0	4	2	1	2	6	1	3	0	0	0	0
HiC_scaffold_21:10041965	cntn5	SGV	7	0	0	0	0	2	0	0	17	0	0	0	1	0
HiC_scaffold_21:31993865	slc35e1	de novo	1	0	0	0	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_24:7400912	NA	SGV	0	4	0	4	0	2	0	0	0	0	0	0	0	0
HiC_scaffold_24:7402072	NA	SGV	1	2	0	2	1	0	1	0	0	0	2	0	0	0
HiC_scaffold_33:12638684	bri3bp	SGV	11	15	1	0	0	2	0	0	4	2	0	4	0	0
HiC_scaffold_33:12644904	bri3bp	SGV	2	9	0	0	0	1	0	0	4	0	0	0	0	0
HiC_scaffold_33:12645055	bri3bp	SGV	3	7	0	2	0	1	1	11	8	15	2	4	2	0
HiC_scaffold_33:12645107	bri3bp	SGV	1	7	0	0	0	0	0	0	4	0	0	0	0	0
HiC_scaffold_33:12645423	bri3bp	SGV	5	8	0	0	0	0	0	0	4	0	0	0	0	0
HiC_scaffold_33:12645669	bri3bp	SGV	1	8	0	0	0	0	0	10	5	0	0	0	0	0
HiC_scaffold_33:12646217	bri3bp	SGV	2	9	0	0	0	0	0	0	6	0	0	0	0	0
HiC_scaffold_33:12651722	wdr31	SGV	4	0	0	0	0	0	0	0	0	1	0	0	0	0
HiC_scaffold_33:12656779	wdr31	SGV	7	12	1	0	0	1	0	3	16	9	2	2	0	0
HiC_scaffold_33:12656781	wdr31	SGV	7	12	1	0	0	1	0	3	16	9	2	2	0	0
HiC_scaffold_33:12656782	wdr31	SGV	7	12	1	0	0	1	0	3	16	9	2	2	0	0
HiC_scaffold_33:12663470	wdr31	SGV	4	9	0	0	1	0	2	8	16	6	2	2	0	21
HiC_scaffold_33:12663497	wdr31	SGV	4	8	0	0	0	0	0	0	13	0	0	0	0	0
HiC_scaffold_33:12663527	wdr31	SGV	4	8	0	0	0	0	0	0	12	0	0	0	0	0
HiC_scaffold_33:12664211	wdr31	SGV	3	11	0	0	0	0	0	0	16	0	0	0	0	0
HiC_scaffold_33:12665844	wdr31	SGV	2	12	0	0	0	0	0	0	18	0	0	0	0	0
HiC_scaffold_33:12666096	wdr31	SGV	7	14	1	4	1	1	1	24	23	2	0	4	0	24
HiC_scaffold_33:12666381	wdr31	SGV	4	11	0	4	0	1	0	0	14	5	0	2	0	0
HiC_scaffold_33:12666394	wdr31	SGV	4	11	0	0	0	0	0	0	17	0	0	0	0	0
HiC_scaffold_33:12670109	wdr31	SGV	5	11	0	0	0	0	0	0	14	0	0	0	0	0
HiC_scaffold_33:12670176	wdr31	SGV	5	10	0	0	2	0	0	6	15	6	2	0	0	18
HiC_scaffold_33:12892230	gnaq	SGV	2	5	0	0	0	0	0	0	19	0	0	0	0	0
HiC_scaffold_33:12893080	gnaq	SGV	1	5	0	0	0	0	0	0	22	1	0	0	0	0
HiC_scaffold_43:26601424	NA	SGV	2	0	0	0	1	0	0	0	0	0	0	0	0	0
HiC_scaffold_43:26602031	NA	SGV	1	1	0	0	1	0	2	21	0	0	0	0	0	1
HiC_scaffold_43:26602094	NA	SGV	0	0	0	0	2	0	0	13	0	0	0	4	0	0
HiC_scaffold_43:26603077	NA	SGV	2	0	0	0	0	2	0	12	0	0	2	0	0	0
HiC_scaffold_43:26603083	NA	SGV	2	0	0	0	0	2	0	12	0	0	2	0	0	0
HiC_scaffold_43:26606120	NA	de novo	1	0	0	0	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_43:26606125	NA	de novo	1	0	0	0	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_44:37672920	NA	SGV	4	0	0	3	0	0	0	6	0	0	0	0	0	0

HiC_scaffold_44:37672992	NA	SGV	6	0	0	3	0	0	2	0	0	0	2	4	0	0
HiC_scaffold_44:37673070	NA	SGV	2	0	0	0	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_44:37673124	NA	SGV	2	0	0	3	0	0	1	0	0	0	0	0	0	0
HiC_scaffold_44:37674761	NA	SGV	3	0	0	3	0	0	2	0	0	0	0	0	0	0
HiC_scaffold_44:37674976	NA	SGV	3	0	0	3	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_44:37675863	NA	SGV	5	0	0	3	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_44:37693328	NA	SGV	4	0	0	0	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_47:14782939	olfm1	SGV	32	17	2	3	1	2	1	12	14	10	1	2	1	12
HiC_scaffold_52:13758756	NA	SGV	0	0	0	0	0	0	0	0	0	0	1	0	0	0
HiC_scaffold_52:16733488	NA	SGV	14	7	0	1	1	0	0	0	0	11	0	0	0	0
HiC_scaffold_52:16737799	NA	SGV	14	7	0	1	1	3	0	15	13	10	0	4	0	20
HiC_scaffold_52:16737807	NA	SGV	14	6	0	0	1	2	0	0	0	8	0	0	0	0
HiC_scaffold_52:16737812	NA	SGV	14	7	0	1	1	3	0	15	13	8	0	4	0	0
HiC_scaffold_52:16737828	NA	SGV	14	6	0	0	1	0	0	0	0	8	0	0	0	0
HiC_scaffold_52:21897888	NA	de novo	1	0	0	0	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_53:18968932	twist1	de novo	1	0	0	0	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_8238:1360	NA	de novo	0	0	0	0	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_9898:5097	NA	SGV	6	1	0	0	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_9898:5770	NA	SGV	5	0	0	4	1	4	0	0	20	0	2	0	0	22
HiC_scaffold_10137:3210	NA	SGV	4	0	0	0	0	0	0	10	1	0	0	0	0	2
HiC_scaffold_10137:5068	NA	SGV	5	0	0	0	0	0	0	9	3	0	2	0	0	0

Table S8. The distribution of unique adaptive alleles to the intermediate scale-eater *C. sp.*
'wide-mouth' across the Caribbean. Numbers represent the copies of the *C. desquamator*
allele present in outgroup populations collected from

San Salvador Island (SS), Rum Cay (RC), Cat Island (CT), Exumas Islands (EX), Long Island
(LG), Mayaguana (MG), New Providence Island (NPI) in the Bahamas. Lagunas Bavaros
(BAV) and Etang Saumautre (ETA) in Dominican Republic, Sarasota Estuary in Florida (FL),
Curacoa (CUR), Caicos Island (CAI) and Isla Margarita (MAR) in Venezuela. The spatial
distribution of each allele is summarized into three categories: standing genetic variation (SGV);
de novo to San Salvador Island populations (de novo) and alleles found in introgressed regions
(Intro). Putative donor populations for introgressed regions are indicated by the same location
codes.

1551
1552

			C. variegatus								*C. laciniatus*	*C. higuey*	*C. bondi*	*C. dearborni*		
Allele	Gene	Distribution	SS	RC	CT	EX	LG	MG	FL	NC	NP	BAV	ETA	CUR	CAI	MAR
HiC_scaffold_1:34344664	smarce1	Intro NC	1	0	0	4	1	2	2	24	2	11	2	4	0	20
HiC_scaffold_1:34346071	smarce1	Intro NC	1	0	0	0	0	1	0	24	0	0	0	0	0	0
HiC_scaffold_1:34346073	smarce1	Intro NC	1	0	0	0	0	1	0	24	0	0	0	0	0	0
HiC_scaffold_5:20660128	NA	SGV	1	0	0	0	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_5:20941055	NA	SGV	1	1	0	3	0	2	0	0	5	6	0	0	1	0
HiC_scaffold_5:20941063	NA	SGV	1	1	0	3	0	2	0	0	5	6	0	0	1	0
HiC_scaffold_5:20945163	NA	SGV	3	1	0	3	0	2	0	0	8	11	0	0	1	2
HiC_scaffold_5:20945417	NA	SGV	1	1	0	4	0	2	0	0	5	7	0	0	1	0
HiC_scaffold_5:20945569	NA	SGV	1	1	0	3	0	3	0	0	4	9	0	4	2	0
HiC_scaffold_5:20964282	NA	SGV	1	1	0	2	1	2	0	15	5	9	0	0	0	0
HiC_scaffold_5:20964283	NA	SGV	1	1	0	2	1	2	0	15	5	9	0	0	0	0
HiC_scaffold_5:20968573	NA	SGV	0	1	0	4	1	3	0	0	7	7	0	4	0	0
HiC_scaffold_5:20969017	NA	SGV	1	1	0	4	1	4	0	0	7	7	0	0	0	0
HiC_scaffold_5:20969195	NA	SGV	2	1	0	2	1	2	0	0	7	6	0	0	0	0
HiC_scaffold_5:20969509	NA	SGV	0	1	0	3	1	2	0	16	6	6	0	4	0	0
HiC_scaffold_5:20969704	NA	SGV	1	2	0	3	1	4	0	0	6	6	0	0	0	0
HiC_scaffold_5:20969759	NA	SGV	1	1	0	3	1	2	0	10	5	5	0	4	0	0
HiC_scaffold_5:20970496	NA	SGV	0	2	0	3	0	2	0	14	6	6	0	4	1	0
HiC_scaffold_5:20971701	NA	SGV	1	1	0	3	1	4	0	0	8	5	0	0	0	1
HiC_scaffold_5:20973584	NA	SGV	1	2	0	4	1	4	0	0	9	6	0	0	0	0
HiC_scaffold_5:20974017	NA	SGV	0	1	0	1	0	4	2	3	8	6	0	0	0	0
HiC_scaffold_5:20977516	NA	SGV	1	2	0	2	1	4	0	0	4	7	0	0	0	0
HiC_scaffold_5:20979898	NA	SGV	0	2	2	3	1	2	1	0	6	5	0	0	1	0
HiC_scaffold_5:20986384	NA	SGV	0	2	0	2	1	2	0	0	8	0	0	0	2	0
HiC_scaffold_5:20988210	NA	SGV	0	3	0	2	1	2	0	0	7	0	0	0	0	0
HiC_scaffold_5:20988292	NA	SGV	0	4	0	3	1	4	0	0	6	1	0	0	2	0
HiC_scaffold_5:20988346	NA	SGV	0	5	0	3	0	2	0	0	8	0	0	0	1	0
HiC_scaffold_11:21559293	NA	SGV	1	22	0	4	0	2	1	0	13	2	0	0	0	0
HiC_scaffold_11:21571679	NA	SGV	1	17	0	1	1	2	1	5	13	2	0	0	1	0

Region	Gene	Category														
HiC_scaffold_11:21571720	NA	SGV	1	22	0	1	1	0	0	5	0	3	0	0	0	0
HiC_scaffold_11:21571742	NA	SGV	1	22	0	1	1	0	0	5	0	3	0	0	1	0
HiC_scaffold_24:15255378	NA	Intro NC	0	0	0	0	1	0	1	23	0	0	0	0	0	24
HiC_scaffold_43:19331844	semag4	SGV	7	0	2	0	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_44:14922408	NA	SGV	4	11	1	0	0	0	0	0	0	1	1	0	0	0
HiC_scaffold_44:14922640	NA	SGV	2	9	0	0	0	0	0	0	0	0	2	0	2	0
HiC_scaffold_44:14922976	NA	SGV	2	11	0	0	0	0	0	0	3	1	0	4	2	0
HiC_scaffold_52:9589311	slitrk5	Intro BAV	1	0	0	2	1	0	0	21	7	20	0	0	0	20
HiC_scaffold_52:9589319	slitrk5	Intro BAV	1	0	0	2	1	0	0	21	7	20	0	0	0	20
HiC_scaffold_52:9592382	slitrk5	Intro BAV	0	1	0	0	2	0	0	2	1	1	2	0	0	26
HiC_scaffold_5743:1358	NA	SGV	0	4	0	2	1	2	0	0	9	0	0	0	2	0
HiC_scaffold_5789:9634	NA	SGV	0	3	0	2	0	2	0	0	7	0	0	0	0	0
HiC_scaffold_5789:10010	NA	de novo	0	0	0	0	0	0	0	0	0	0	0	0	0	0
HiC_scaffold_5789:11490	NA	SGV	0	4	0	2	2	2	0	0	5	0	0	0	0	0
HiC_scaffold_5789:12093	NA	SGV	0	4	0	2	0	2	0	0	7	0	0	0	1	0
HiC_scaffold_13976:3987	NA	SGV	0	0	0	2	1	2	0	0	5	0	0	0	2	0

Table S10. Assessment of introgression signatures between population in Osprey Lake, San
Salvador Island in regions that contain putative adaptive alleles unique to *C. sp*. wide-
mouth, unique to *C. desquamator* and shared between both scale-eating populations.
Introgressed regions are highlighted in bold. Introgressed regions were those that had a relative
node depth (RND) value between two populations. RND statistics were calculated in 10-kb
windows between *C. desquamator* and *C.* sp. 'wide-mouth' (des vs wid) and *C. desquamator* and

C. sp. 'wide-mouth' (des vs var), with the latter test serving as a control for recent and ongoing
gene exchange due to sympatric overlap in the lake that is not relevant for scale-eating
adaptation. Adaptive allele, gene names, and RND values that are bolded represent candidate
introgressed regions with adaptive alleles in them. Candidate introgressed regions were regions
that fell below a significance threshold value (RND < 0.28) based on the lower 5[th] percentile of
RND values calculated from simulations of two populations that experience no gene flow during
divergence. The number of exons present in each 10-kb window is also included in this table.
1588

Adaptive allele	Gene	RND des vs wid	RND des v var	Introgressed Region	Number of exons
Unique *C. sp.* 'wide-mouth' adaptive alleles					
HiC_scaffold_5:20941063	NA	1.48663438	1.11402561	20940001-20950000	0
HiC_scaffold_5:20964282	NA	1.53346729	1.28660246	20960001-20970000	0
HiC_scaffold_5:20970496	NA	1.52647461	1.29854143	20970001-20980000	0
HiC_scaffold_5:20986384	NA	1.34806367	0.90675156	20980001-20990000	0
HiC_scaffold_11:21559293	NA	1.24490382	0.78312989	21550001-21560000	0
HiC_scaffold_11:21571679	NA	1.74507532	0.87626263	21570001-21580000	0
HiC_scaffold_24:15255378	NA	0.60467498	0.54123148	15250001-15260000	0
HiC_scaffold_43:19331844	Sema4g	1.18018089	0.94111453	19330001-19340000	0
HiC_scaffold_44:14922408	atp8a1	1.60947983	0.97866515	14920001-14930000	0
HiC_scaffold_52:9589311	Slitrk5	1.47013193	1.14495019	9580001-9590000	0
C. desquamator and *C. sp.* 'wide-mouth' adaptive alleles					
HiC_scaffold_6:907101	**NA**	**0.01764957**	0.304609	900001-910000	0
HiC_scaffold_6:923182	**NA**	**0.03560806**	0.41432253	920001-930000	0
HiC_scaffold_6:934123	**NA**	**0.01180417**	0.29362216	930001-940000	0
HiC_scaffold_6:941221	**NA**	**0.03304484**	0.52001902	940001-950000	0
HiC_scaffold_6:951028	NA	0.33613727	0.34973512	950001-960000	**0**
HiC_scaffold_9:31763392	**DAAM2**	**0.0241461**	0.07809267	31750001-31760000	4
HiC_scaffold_9:31763392	**DAAM2**	**0.06961605**	0.83262139	31760001-31770000	5
HiC_scaffold_24:5390450	**Usp50**	**0.0833701**	0.90291628	5390001-5400000	0
HiC_scaffold_43:27385265	NA	1.30551148	2.06180406	27380001-27390000	0
HiC_scaffold_44:14963656	**ATP8A1**	**0.13299985**	1.26725109	14960001-14970000	2
HiC_scaffold_60:1798737	**ZNF214**	**0.08551992**	1.60256069	1790001-1800000	1
Unique *C. desquamator* adaptive alleles					
HiC_scaffold_5:6082906	NA	1.61515973	1.39739811	6080001-6090000	0
HiC_scaffold_11:10913509	cd226	0.25420561	1.04030645	10910001-10920000	2
HiC_scaffold_11:10948683	cd226	1.11058873	0.75472269	10940001-10950000	2
HiC_scaffold_21:10041965	cntn5	1.51795767	1.5514369	10040001-10050000	3
HiC_scaffold_21:31993865	SLC35E1	4.35	4.81944444	31990001-32000000	3
HiC_scaffold_24:7401348	NA	0.62028279	0.62001004	7400001-7410000	0
HiC_scaffold_33:12638684	bri3bp	1.87610126	1.80369995	12630001-12640000	1

HiC_scaffold_33:12641481	bri3bp	2.05234267	1.93074547	12640001-12650000	2
HiC_scaffold_33:12656933	wdr31	2.08801498	1.91525971	12650001-12660000	8
HiC_scaffold_33:12664211	wdr31	2.04327131	1.88029401	12660001-12670000	0
HiC_scaffold_33:12670176	wdr31	2	1.87301587	12670001-12680000	0
HiC_scaffold_33:12891903	gnaq	1.19586694	1.21312932	12890001-12900000	0
HiC_scaffold_43:26606125	NA	1.7651081	1.60827345	26600001-26610000	0
HiC_scaffold_44:37673070	NA	1.3785441	1.29438333	37670001-37680000	0
HiC_scaffold_44:37693328	NA	1.2303647	1.09117949	37690001-37700000	0
HiC_scaffold_47:14782939	**olfm1**	**0.19804878**	0.7647762	14780001-14790000	1
HiC_scaffold_52:16737828	NA	1.19630072	1.25855174	16730001-16740000	0
HiC_scaffold_53:18968932	**twist1**	**0.05856256**	0.54868019	18960001-18970000	2

Table S11.Genome-wide signatures of differential introgression between *C.* sp. 'wide-mouth' and *C. desquamator*. f_4-statistics, standard error, z-scores and p-values for genome-wide test with 5 outgroup generalist populations from across the Caribbean. *C. artifrons* used as outgroup species in which we expect minimal gene flow.

–

P1	P2	P3	Outgroup	f4	Standard Error	Z-score	P-value
wid	des	Dominican Republic	*Artifrons*	-0.0003	0.000173	-1.7136	0.082
wid	des	Rum Cay	*Artifrons*	0.00035	0.000188	1.90484	0.05
wid	des	North Carolina	*Artifrons*	0.00053	0.000186	2.85	0.0044
wid	des	Venezuela	*Artifrons*	0.00011	0.000143	0.78891	0.43
wid	des	New Providence Island	*Artifrons*	0.00027	0.000152	1.81370	0.069

References

1.   Wainwright PC, Bellwood DR, Westneat MW, Grubich JR, Hoey AS. 2004 A functional morphospace for the skull of labrid fishes: Patterns of diversity in a complex biomechanical system. *Biol. J. Linn. Soc.* **82**, 1–25. (doi:10.1111/j.1095-

8312.2004.00313.x)

2.   Patricia Hernandez L, Gibb AC, Ferry-Graham L. 2009 Trophic apparatus in
cyprinodontiform fishes: Functional specializations for picking and scraping behaviors. *J.*
Morphol. **270**, 645–661. (doi:10.1002/jmor.10711)

3.   Martin CH, Wainwright PC. 2013 Multiple fitness peaks on the adaptive landscape drive
adaptive radiation in the wild. *Science* **339**, 208–11. (doi:10.1126/science.1227710)

4.   Jackson AL, Inger R, Parnell AC, Bearhop S. 2011 Comparing isotopic niche widths among
and within communities: SIBER - Stable Isotope Bayesian Ellipses in R. *J. Anim. Ecol.* **80**,
595–602. (doi:10.1111/j.1365-2656.2011.01806.x)

5.   Richards E, McGirr J, Wang J, St. John M, Poelstra J, Solano M, O'Connell D, Turner B,
Martin C. 2020 Major stages of vertebrate adaptive radiation are assembled from a
disparate spatiotemporal landscape. (doi:10.1101/2020.03.12.988774)

6.   Li H, Durbin R. 2011 Inference of human population history from individual whole-
genome sequences. *Nature* **475**, 493–496. (doi:10.1038/nature10231)

7.   DePristo MA *et al.* 2011 A framework for variation discovery and genotyping using next-
generation DNA sequencing data. *Nat. Genet.* **43**, 491–8. (doi:10.1038/ng.806)

8.   Marsden CD, Lee Y, Kreppel K, Weakley A, Cornel A, Ferguson HM, Eskin E, Lanzaro GC.
2014 Diversity, differentiation, and linkage disequilibrium: prospects for association
mapping in the malaria vector Anopheles arabiensis. *G3 (Bethesda).* **4**, 121–31.
(doi:10.1534/g3.113.008326)

9.   Danecek P *et al.* 2011 The variant call format and VCFtools. *Bioinformatics* **27**, 2156–
2158. (doi:10.1093/bioinformatics/btr330)

10.   Purcell S *et al.* 2007 PLINK: A Tool Set for Whole-Genome Association and Population-
Based Linkage Analyses. *Am. J. Hum. Genet.* **81**, 559–575. (doi:10.1086/519795)

11.   Alexander DH, Novembre J, Lange K. 2009 Fast model-based estimation of ancestry in
unrelated individuals. *Genome Res.* **19**, 1655–1664.

12.   Excoffier L, Dupanloup I, Huerta-Sánchez E, Sousa VC, Foll M. 2013 Robust Demographic
Inference from Genomic and SNP Data. *PLOS Genet.* **9**, e1003905.
(doi:10.1371/journal.pgen.1003905)

13.   Schiffels S, Durbin R. 2014 Inferring human population size and separation history from
multiple genome sequences. *Nat. Genet.* **46**, 919. (doi:10.1038/ng.3015)

14.   Nadachowska-Brzyska K, Burri R, Smeds L, Ellegren H. 2016 PSMC analysis of effective
population sizes in molecular ecology and its application to black-and-white Ficedula
flycatchers. *Mol. Ecol.* **25**, 1058–1072. (doi:10.1111/mec.13540)

15.   Martin CH, Crawford JE, Turner BJ, Simons LH. 2016 Diabolical survival in Death Valley:
recent pupfish colonization, gene flow and genetic assimilation in the smallest species
range on earth. *Proc. R. Soc. B Biol. Sci.* **283**, 20152334. (doi:10.1098/rspb.2015.2334)

16.   Pavlidis P, Živković D, Stamatakis A, Alachiotis N. 2013 SweeD: Likelihood-based
detection of selective sweeps in thousands of genomes. *Mol. Biol. Evol.* **30**, 2224–2234.
(doi:10.1093/molbev/mst112)

17.   Garrigan D, Geneva A. 2014 msmove: A modified version of Hudson's coalescent
simulator ms allowing for finer control and tracking of migrant genealogies.
(doi:10.6084/m9.figshare.1060474)

18.   Balsa-Canto E, Henriques D, Gabor A, Banga JR. 2016 AMIGO2, a toolbox for dynamic

modeling, optimization and control in systems biology. *Bioinformatics* **32**, 1–2.
(doi:10.1093/bioinformatics/btw411)

19.   Ge SX, Jung D. 2018 ShinyGO: a graphical enrichment tool for animals and plants. *bioRxiv*
, 2. (doi:10.1101/315150)

20.   Smith J, Coop G, Stephens M, Novembre J. 2018 Estimating Time to the Common
Ancestor for a Beneficial Allele. *Mol. Biol. Evol.* **35**, 1003–1017. (doi:10.1101/071241)

21.   Bradbury PJ, Zhang Z, Kroon DE, Casstevens TM, Ramdoss Y, Buckler ES. 2007 TASSEL:
Software for association mapping of complex traits in diverse samples. *Bioinformatics* **23**,
2633–2635. (doi:10.1093/bioinformatics/btm308)

22.   Roesti M, Moser D, Berner D. 2013 Recombination in the threespine stickleback genome
- Patterns and consequences. *Mol. Ecol.* **22**, 3014–3027. (doi:10.1111/mec.12322)

23.   Feder JL, Xie X, Rull J, Velez S, Forbes A, Leung B, Dambroski H, Filchak KE, Aluja M. 2005
Mayr, Dobzhansky, and Bush and the complexities of sympatric speciation in Rhagoletis.
Proc. Natl. Acad. Sci. U. S. A. **102 Suppl**, 6573–80. (doi:10.1073/pnas.0502099102)

24.   Pfeifer B, Wittelsbürger U, Ramos-Onsins SE, Lercher MJ. 2014 PopGenome: An efficient
swiss army knife for population genomic analyses in R. *Mol. Biol. Evol.* **31**, 1929–1936.
(doi:10.1093/molbev/msu136)

25.   Richards EJ, McGirr JA, Wang JR, St. John ME, Poelstra JW, Solano MJ, O'Connell DC,
Turner BJ, Martin CH. 2021 A vertebrate adaptive radiation is assembled from an ancient
and disjunct spatiotemporal landscape. *Proc. Natl. Acad. Sci. U. S. A.* **118**.
(doi:10.1073/pnas.2011811118)

26.   Pfeifer B, Kapan DD. 2019 Estimates of introgression as a function of pairwise distances.
BMC Bioinformatics **20**, 1–11. (doi:10.1186/s12859-019-2747-z)

27.   Martin SH, Davey JW, Jiggins CD. 2015 Evaluating the use of ABBA-BABA statistics to
locate introgressed loci. *Mol. Biol. Evol.* **32**, 244–257. (doi:10.1093/molbev/msu269)

28.   Echelle AA, Carson EW, Echelle AF, Bussche D, Dowling TE. 2005 Historical Biogeography
of the New-World Pupfish Genus Cyprinodon (Teleostei: Cyprinodontidae). *Copeia* **2005**,
320–339.

29.   Patterson N, Moorjani P, Luo Y, Mallick S, Rohland N, Zhan Y, Genschoreck T, Webster T,
Reich D. 2012 Ancient admixture in human history. *Genetics* **192**, 1069–1093.
(doi:10.1534/genetics.112.145037)

30.   Edwards CD. 2001 Effect of Salinity on the Ecology of Molluscs in the Inland Saline
Waters of San Salvador Island: A Experiment in Progress. *Proc. Eight Symp. Nat. Hist.*
Bahamas. , 15–26.

31.   Turner BJ, Duvernell DD, Bunt TM, Barton MG. 2008 Reproductive isolation among
endemic pupfishes (Cyprinodon) on San Salvador Island, Bahamas: Microsatellite
evidence. *Biol. J. Linn. Soc.* **95**, 566–582. (doi:10.1111/j.1095-8312.2008.01079.x)

32.   Clark PU, Dyke AS, Shakun JD, Carlson AE, Clark J, Wohlfarth B, Mitrovica JX, Hostetler
SW, McCabe AM. 2009 The Last Glacial Maximum. *Science (80-.).* **325**, 710–714.
(doi:10.1126/science.1172873)

33.   Dickinson ME *et al.* 2016 High-throughput discovery of novel developmental phenotypes.
Nature **537**, 508–514. (doi:10.1038/nature19356)

34.   Kida YS, Sato T, Miyasaka KY, Suto A, Ogura T. 2007 Daam1 regulates the endocytosis of
EphB during the convergent extension of the zebrafish notochord. *Proc. Natl. Acad. Sci.*

U. S. A. **104**, 6708–6713. (doi:10.1073/pnas.0608946104)

35.   Lee JY, Seo D, You J, Chung S, Park JS, Lee JH, Jung SM, Lee YS, Park SH. 2017 The
deubiquitinating enzyme, ubiquitin-specific peptidase 50, regulates inflammasome
activation by targeting the ASC adaptor protein. *FEBS Lett.* **591**, 479–490.
(doi:10.1002/1873-3468.12558)

36.   Heras J, Martin C. 2021 Nonadaptive radiation of the gut microbiome in an adaptive
radiation of Cyprinodon pupfishes with minor shifts for scale-eating. *bioRxiv*

37.   Soupene E. 2008 ATP8A1 activity and phosphatidylserine transbilayer movement. *J.*
Receptor. Ligand Channel Res. **Volume 1**, 1–10. (doi:10.2147/jrlcr.s3773)

38.   Thomas T, Voss AK, Chowdhury K, Gruss P. 2000 Querkopf, a MYST family histone
acetyltransferase, is required for normal cerebral cortex development. *Development*
127, 2537–2548.

39.   Kraft M *et al.* 2011 Disruption of the histone acetyltransferase MYST4 leads to a noonan
syndrome-like phenotype and hyperactivated MAPK signaling in humans and mice. *J.*
Clin. Invest. **121**, 3479–3491. (doi:10.1172/JCI43428)

40.   Clayton-Smith J *et al.* 2011 Whole-exome-sequencing identifies mutations in histone
acetyltransferase gene KAT6B in individuals with the say-barber-biesecker variant of
Ohdo syndrome. *Am. J. Hum. Genet.* **89**, 675–681. (doi:10.1016/j.ajhg.2011.10.008)

41.   Martin CH, Richards EJ. 2019 The Paradox behind the Pattern of Rapid Adaptive
Radiation: How Can the Speciation Process Sustain Itself through an Early Burst? *Annu.*
Rev. Ecol. Evol. Syst.

42.   Whittaker R. 1977 Evolution of species diversity in land communities. *Evol. Biol.*

43.   Stroud JT, Losos JB. 2016 Ecological Opportunity and Adaptive Radiation. *Annu. Rev. Ecol.*
Evol. Syst. **47**, 507–532. (doi:10.1146/annurev-ecolsys-121415-032254)

44.   Orr HA. 1998 The Population Genetics of Adaptation: The Distribution of Factors Fixed
during Adaptive Evolution. *Evolution (N. Y).* **52**, 935. (doi:10.2307/2411226)

45.   Orr HA. 2005 The genetic theory of adaptation: A brief history. *Nat. Rev. Genet.* **6**, 119–
127. (doi:10.1038/nrg1523)

46.   Gillespie JH. 1984 Molecular Evolution Over the Mutational Landscape. *Evolution (N. Y).*
38, 1116–1129. (doi:10.2307/2408444)

47.   Kauffman S, Levin S. 1987 Towards a general theory of adaptive walks on rugged
landscapes. *J. Theor. Biol.* **128**, 11–45. (doi:https://doi.org/10.1016/S0022-
5193(87)80029-2)

48.   Maynard Smith J. 1970 Natural Selection and the Concept of a Protein Space. *Nature* **225**,
563–564. (doi:10.1038/225563a0)

49.   Bell G. 2013 Evolutionary rescue and the limits of adaptation. *Philos. Trans. R. Soc. B Biol.*
Sci. **368**, 20120080. (doi:10.1098/rstb.2012.0080)

50.   Lindsey HA, Gallie J, Taylor S, Kerr B. 2013 Evolutionary rescue from extinction is
contingent on a lower rate of environmental change. *Nature* **494**, 463–467.
(doi:10.1038/nature11879)

51.   Rousselle M, Simion P, Tilak M-K, Figuet E, Nabholz B, Galtier N. 2020 Is adaptation
limited by mutation? A timescale-dependent effect of genetic diversity on the adaptive
substitution rate in animals. *PLOS Genet.* **16**, e1008668.

52.   Mani GS, Clarke BC. 1990 Mutational order: a major stochastic process in evolution. *Proc.*

1743  *R. Soc. London. B. Biol. Sci.* **240**, 29–37. (doi:10.1098/rspb.1990.0025)

1744 53. Schluter D. 2009 Evidence for Ecological Speciation and Its Alternative. *Science (80-.).*
1745  **323**, 737 LP – 741. (doi:10.1126/science.1160006)

1746 54. Good BH, McDonald MJ, Barrick JE, Lenski RE, Desai MM. 2017 The dynamics of
1747  molecular evolution over 60,000 generations. *Nature* **551**, 45–50.
1748  (doi:10.1038/nature24287)

1749 55. Schluter D, Conte GL. 2009 Genetics and ecological speciation. *Proc. Natl. Acad. Sci.* **106**,
1750  9955–9962. (doi:10.1073/pnas.0901264106)

1751 56. Patwa Z, Wahl LM. 2008 The fixation probability of beneficial mutations. *J. R. Soc.*
1752  *Interface* **5**, 1279–1289. (doi:10.1098/rsif.2008.0248)

1753 57. Pavlidis P, Alachiotis N. 2017 A survey of methods and tools to detect recent and strong
1754  positive selection. *J. Biol. Res.* **24**, 7. (doi:10.1186/s40709-017-0064-0)

1755 58. Höllinger I, Pennings PS, Hermisson J. 2019 Polygenic adaptation: From sweeps to subtle
1756  frequency shifts. *PLOS Genet.* **15**, e1008035.

1757 59. Barghi N, Hermisson J, Schlötterer C. 2020 Polygenic adaptation: a unifying framework to
1758  understand positive selection. *Nat. Rev. Genet.* **21**, 769–781. (doi:10.1038/s41576-020-
1759  0250-z)

1760 60. Chevin LM, Hospital F. 2008 Selective sweep at a quantitative trait locus in the presence
1761  of background genetic variation. *Genetics* **180**, 1645–1660.
1762  (doi:10.1534/genetics.108.093351)

1763 61. Thornton KR. 2019 Polygenic adaptation to an environmental shift: Temporal dynamics
1764  of variation under Gaussian stabilizing selection and additive effects on a single trait.
1765  *Genetics* **213**, 1513–1530. (doi:10.1534/genetics.119.302662)

1766 62. Flint HJ, Scott KP, Louis P, Duncan SH. 2012 The role of the gut microbiota in nutrition
1767  and health. *Nat. Rev. Gastroenterol. Hepatol.* **9**, 577–589.
1768  (doi:10.1038/nrgastro.2012.156)

1769 63. Mcgirr JA, Martin CH. 2018 Parallel evolution of gene expression between trophic
1770  specialists despite divergent genotypes and morphologies. *Evol. Lett.* **2**, 62–75.
1771  (doi:10.1002/evl3.41)

1772 64. Martin CH, Gould KJ. 2020 Surprising spatiotemporal stability of a multi-peak fitness
1773  landscape revealed by independent field experiments measuring hybrid fitness. *Evol.*
1774  *Lett.* , 530–544. (doi:10.1002/evl3.195)

1775
1776