Title: Living with high potassium: how plants balance nutrient acquisition during K-induced

salt stress

Pramod Pantha[1], Dong-Ha Oh[1], David Longstreth[1] and Maheshi Dassanayake[1,*]

[1]Department of Biological Sciences, Louisiana State University, Baton Rouge, LA 70803, USA

*Address correspondence to: maheshid@lsu.edu

Short title: Avoiding N starvation upon high K stress

Abstract

K is more toxic in plants than Na at similar concentrations but the molecular mechanisms

governing excess K-induced salt stress are unknown. We used *Arabidopsis thaliana* and its

extremophyte relative *Schrenkiella parvula*, to explore this largely unexplored question using

comparative physiological, ionomic, transcriptomic, and metabolomic approaches aimed at

understanding how plants can develop resilience to excess K. Our results showed that the stress

responses exhibited by the two plants diverged at a decisive step where the stress-sensitive *A.*

thaliana could not limit excess K influx and suffered severe nutrient depletion. The stress

adapted model, *S. parvula*, was able to independently regulate reduction in K uptake while

sustaining uptake of other major nutrients including N. Maintaining N uptake and its

uninterrupted assimilation into primary metabolites allowed *S. parvula* to sustain growth and

concurrently boost its antioxidant and osmolyte pools, facilitated by a targeted transcriptomic

response. In contrast, *A. thaliana* descended into mismanaged transcriptional cascades

including induction of biotic and abiotic stress responses and autophagy accompanied by

inhibited growth, reduced photosynthesis, and increased ROS. This study provides a basic

framework to select key pathways to target in the development of building plant resilience

towards nutrient imbalance caused by K-induced salt stress.

Keywords

K-induced salt stress, potassium transport, transcriptomics, metabolomics, ionomics, molecular

phenotype, abiotic stress, nitrogen metabolism, osmoprotectants, antioxidants, antagonistic

pleiotropy

Introduction

Can excess potassium (K) in soil be too much of a good thing for plants? Its role as an

essential macronutrient for plants is well established and K constitutes 2-10% of plant dry

weight (Wang and Wu, 2013). The cytosolic concentration of K^+ is around 100 mM and the

vacuolar concentrations generally range between 20-200 mM depending on tissue,

developmental age, or species (Leigh and Wyn Jones, 1984; Ashley et al., 2006). These

concentrations are typically higher than soil concentrations found in most agricultural soils

(Maathuis, 2009) and therefore the vast majority of studies have only investigated uptake into

plants from low K^+ concentrations in the soil. As a result, we hardly know if excess K^+ in soils

would be as toxic as high Na^+ (Pantha and Dassanayake, 2020), and if it is, how it exerts plant

toxicity effects that may be different from Na-induced salt stress seen for high salt

concentrations known to cause plant toxicity (>100 mM NaCl). While toxicity in plants growing

at high concentration of other nutrients, such as boron (Aquea et al., 2012; Wang et al., 2021),

copper (Lequeux et al., 2010), and nitrogen (Yoshitake et al., 2021), have been investigated, the

molecular mechanisms behind high K^+ responses are virtually unknown.

The need to use alternative agricultural lands and recycled wastewater amidst

diminishing freshwater resources is intense (IPCC, 2019). These needs cannot be addressed

without foundational knowledge of how plant nutrient balance is achieved when conventional

land use and irrigation practices no longer provide adequate solutions. Worldwide there are

many regions that are naturally high in soil K^+ content (Duval et al., 2005; Warren, 2016). Many

industrial as well as agricultural processing plants produce wastewater with exceptionally high

K^+ concentrations considered excessive for plant growth (Arienzo et al., 2009). During

wastewater treatments, unlike N, P, or organic matter which are typically processed using

microbial activity, K is concentrated due to evaporation (Arienzo et al., 2009). Therefore,

knowing the tolerance mechanisms against high K, which can cause osmotic stress similar to Na

on top of additional stresses because of its role as a major nutrient in plants, are imperative

toward designing robust crops in our quest to convert marginal lands into productive

agricultural lands and reclaim land left unusable due to irrigation practices such as the use of

recycled wastewater.

How high K^+ in soils affect plant growth is not a new question for plant scientists (Eijk,

1939). Past studies have reported that high concentrations of K^+ severely affected growth and

other developmental processes when tested in a handful of species. For example, growth of

Salicornia herbacea was induced by 166, 250, and 333 mM NaCl but reduced when treated with

the same concentrations of KCl, implying ion specific detrimental effects of K^+ in a halophyte

known to be able to cope with high salinity stress (Eijk, 1939). Additional studies using other

halophytes including, *Suaeda aegyptiaca* (Eshel, 1985), *S. salsa* (Wang et al., 2001), and several

Atriplex spp. (Ashby and Beadle, 1957; Matoh et al., 1986; Ramos et al., 2004) have shown that

KCl is more toxic than NaCl at similar osmotic strengths. Recently, Richter et al., (2019)

compared the effects of 100 mM NaCl, 50 mM Na_2SO_4, and 100 mM KCl and reported that K^+

caused more negative physiological and metabolic responses then Na^+ in *Vicia faba*. These

studies suggest that K^+ may not elicit the same physiological, metabolic, or genetic responses

Na^+ does and plants may require distinct genetic pathways in addition to canonical salt

response pathways (Pantha and Dassanayake, 2020) to survive high K-induced salt stress.

In this study, we aimed to identify K-induced salt stress responses and deduce the

underlying cellular mechanisms plants have evolved to adapt to K^+ toxicity. We compared

Arabidopsis thaliana (Col-0 ecotype), a model plant sensitive to high K^+, to its extremophyte

relative, *Schrenkiella parvula* (Lake Tuz ecotype) that thrives in high K^+ soils (Nilhan et al., 2008;

Oh et al., 2014). We examined multi-omics level features and responses exhibited by the two

model species to high K^+ treatments to identify the relevant genetic pathways influential in

regulating or are affected by K^+ toxicity responses. Our results revealed an extensive ionomic,

metabolic, and transcriptomic reprogramming during high potassium stress in the stress-

sensitive model while providing novel insights into how a stress-adapted model had evolved

cellular mechanisms to tolerate K^+ toxicity via metabolic and transcriptomic adjustments.

Materials and methods

Plant material, growth conditions, and stress treatments

Plants were grown at 125-150 µmol m^{-2}s^{-1} photon flux density at a photoperiod of 12 h light, 12 h dark, 23°C, and 60% relative humidity for all the experiments unless described otherwise. For transcriptome, ionome, and metabolome profiling, *Arabidopsis thaliana* (Col-0) and *Schrenkiella parvula* (Lake Tuz ecotype) plants were grown hydroponically in 1/5th strength Hoagland's solution as described in Wang et al., 2021. 25-day-old seedlings (days after germination) were transferred to fresh hydroponic growth media at control conditions or supplemented with 150 mM KCl. Samples were harvested at 0, 3, 24, and 72 hours after treatment (HAT) as summarized in Fig S1. Shoot and root tissue samples were harvested on a randomized basis from the same growth chamber, at the same time of day for control and salt-treated plants. Roots were briefly dried with a paper towel to soak any excess growth solution. At least five plants per sample were harvested in at least three biological replicates (Fig S1). All treatment and harvest times were set at 4 h after the beginning of the light cycle to avoid variation due to circadian effects. The salt treatment was non-lethal to both *A. thaliana* and *S. parvula* plants based on preliminary tests using a series of salt concentrations.

The primary root length, lateral root number and density, and root hair length were quantified using plate-grown seedlings. For this, surface-sterilized seeds were germinated in Petri plates on 1/4th Murashige and Skoog (MS) medium (Murashige and Skoog, 1962) solidified with 0.8% Phyto agar (PlantMedium) and grown for 5 days. The seedlings were transferred to 13 cm-diameter large Petri plates with the same media supplemented with different concentration of salts and further incubated in a vertical position to allow better root growth. All measurements were completed before the primary root reached the bottom of the plates. Microscopic images of root hairs were obtained using a Zeiss Lumar brightfield microscope with a 1.5 X FWD 30 mm objective. Plants were imaged for lengths of roots, root hair, and leaf area measurements and quantified using ImageJ (Ferreira and Rasband, 2012). Lateral root density was calculated by dividing the total lateral root number by the total length of the primary root.

For the plants grown in soil, seeds were directly germinated in

13.75cmx13.75cmx6.87cm pots with SunGro MetroMix 360 soil (SunGro Horticulture, Agawam,

MA). The plants were watered with 1/5th strength Hoagland's solution once every week and

with tap water every other day. 21 days-old *A. thaliana* and *S. parvula* plants were watered

with tap water or 50-400 mM NaCl or KCl salt solutions for control and salt treated plants,

respectively, every other day for 14 days.

CO_2 assimilation rates

CO_2 assimilation rates were measured on hydroponically grown plants using a LiCor

6400XT portable system attached to a 6400-17 whole plant Arabidopsis chamber mounted with

a 6400-18 RGB light source (LiCor Inc, Lincoln Nebraska). Rates were measured on 25-day old *A.*

thaliana and *S. parvula* plants at 0, 3, 24, and 72 h after exposure to 150 mM KCl. Individual

plants were placed in a 15 ml centrifuge tube containing the appropriate treatment solution

and the centrifuge tube was inserted into a sealed 38 mm Cone-tainer^{TM} mounted in the whole

plant Arabidopsis chamber. The chamber was kept at a slight positive pressure and CO_2

exchange was minimal for centrifuge tubes containing nutrient solution with no plants. CO_2

assimilation rates were measured at 400 μL L^{-1} CO_2, 1,000 μmol photons m^{-2} s^{-1}, a chamber

temperature of 23°C and an air flow rate of 500 mL min^{-1} after CO_2 assimilation rates were

constant (usually within 5 minutes for *A. thaliana* and 10 minutes for *S. parvula*).

Measurements were made on 3 replicate plants.

Ionome and Total C and N profiling

For each time point and condition described in Fig S1, four biological replicates were

harvested and dried for 1 week at 40°C and weighed daily until the dry weights stabilized

before grinding. The sample preparation for ionome measurements was performed following

Ziegler et al. (2013). Briefly, 10-30 mg of dried samples were digested with 2.5 mL concentrated

HNO$_3$ (AR Select Grade, VWR International, LLC) at room temperature overnight and heated at

105°C for 2 h. Each sample was diluted to 10 mL with ultrapure 18MΩ water (UPW) and 0.9 mL

of the diluted extract was further mixed with 4.1 mL UPW to a 5 mL final volume, of which 1.2

mL was used for ICP-MS (inductively coupled plasma mass spectrometry). Concentrations of 21

elements (K, P, Mg, S, Ca, Al, Zn, Co, Ni, Fe, Se, Cu, B, Mn, Mo, As, Rb, Cd, Na, Li, and Sr) were

quantified at USDA-ARS-Plant Genetics Facility, Donald Danforth Plant Science Center, as

described in Baxter et al. (2014), using a Perkin Elmer NexION ICP-MS platform with 20 µg L^{-1} In

as an internal standard (Aristar Plus, BDH Chemicals). Results were visualized as heatmaps using

the 'pheatmap' R package (v. 1.0.12) (Kolde, 2012). Total carbon and nitrogen were quantified

using a Costech 4010 elemental analyzer (Costech, Valencia, CA, US). Each sample ranged from

2-10 mg ground tissue. Acetanilide and 4 mg of wheat flour were used as internal controls.

One-way ANOVA, followed by Tukey's post hoc test ($p<0.05$) was performed using the agricolae

R package for 3-4 biological replicates to determine elements with significant differences

between samples.

Metabolite profiling

To obtain high-throughput metabolite profiles on samples described in Fig S1, we

performed non-targeted gas chromatography-mass spectrometry (GC-MS) at the Fiehn

Laboratory, West-coast Metabolomics Center at UC-Davis. Following harvest, both root and

shoot samples were immediately frozen in liquid nitrogen and processed as described in Fiehn

et al. (2008). A Leco Pegasus IV mass spectrometer with the following parameters were used:

unit mass resolution at 17 spectra per second with a scan mass range of 85-500 Da at 70 V

ionization energy and 1800 V detector voltage with a 230°C transfer line and a 250°C ion

source. Raw data files were processed using ChromaTOF software (Version 2.32, Leco) with

automatic mass spectral deconvolution followed by peak detection at 5:1 signal/noise levels for

all chromatograms. The BinBase database was used for automated annotation of detected

peaks to metabolites (https://bitbucket.org/fiehnlab/binbase/src/master/). To generate

relative abundance values, we used peak heights of specific retention indices which allows

greater separation between metabolites and higher detection capacity for less abundant

metabolites. The raw abundance/intensity for each identified compound was normalized with a

series of fatty acid methyl esters (xylulose, oleamide, mannonic acid, enolpyruvate,

dihydroxymalonic acid, coniferin, butyrolactam, 2-ketoglucose dimethylacetal, and 2,5-

dihydroxypyrazine) used as National Institute of Standards and Technology (NIST) internal

standards. Significance tests were performed using one-way ANOVA followed by Tukey's post

hoc test at $p<0.05$ in the agricolae R package for at least three biological replicates and circular

plots to visualize selected metabolites were generated using the circlize package in R (Gu et al.,

2014).

RNA-seq analyses

Total RNA from three biological replicates of root and shoot tissues harvested at 0, 3,

24, and 72 h after treating with 150 mM KCl (Fig S1) was extracted (~6 µg) using RNeasy Plant

Mini Kit (Qiagen, Hilden, Germany) with column digestion to remove DNA contamination. Six to

eight plants were pooled for each replicate. A total of 48 RNA-seq libraries were prepared using

the TruSeq RNA library preparation kit (Illumina, San Diego, USA) and sequenced on an Illumina

HISeq 4000 platform at the Roy J. Carver Biotechnology Center, University of Illinois at Urbana-

Champaign. A total of 12-23 million 50-bp reads was obtained from each library (available at

NCBI-SRA database under BioProject PRJNA63667). Following quality checks using FastQC

(Andrews, 2010), RNA-seq reads uniquely mapped to *A. thaliana* TAIR10 or *S. parvula*

(Dassanayake et al., 2011) v2.2 gene models (https://phytozome-next.jgi.doe.gov/), were

quantified using Bowtie (Langmead et al., 2009) and a custom script as described in (Pantha et

al., 2021).

Differentially expressed genes (DEGs) between K^+ treated and control samples were

identified using DESeq2 with an adjusted p-value cutoff 0.01 (Love et al., 2014). DEGs shared

between treatments were plotted using UpSetR (Conway et al., 2017). Orthologous gene pairs

between the two species were identified using the CLfinder-OrthNet pipeline with default

settings (Oh and Dassanayake, 2019) and further refined as described in Wang et al. (2021) to

include lineage-specific genes and duplications. For ortholog expression comparisons between

S. parvula and *A. thaliana*, regularized log (rlog) values estimated by DESeq2 were normalized

per each gene. Ortholog pairs compared to the median across all samples were used to

determine temporal co-expression clusters based on fuzzy k-means clustering (Gasch and Eisen,

2002) with a membership cutoff set to 0.5. From initially identified 10 root and 11 shoot

clusters (Table S11), we filtered out clusters that did not show a response to K treatments in

both species and identified 5 root and 3 shoot co-expression super clusters with distinct

response trends. All expressed genes were assigned Gene Ontology (GO) annotations when

available and DEGs with GO annotations were used to identify enrichment processes using

BinGO (Maere et al., 2005). To summarize functional groups enriched among DEGs, redundant

GO terms with > 50% overlap were further clustered using GOMCL (Wang et al., 2020). DEGs

were also annotated based on KEGG pathways (Kanehisa et al., 2016). Functional processes

identified with specific GO terms which included less than 500 genes to represent a function

were used for functional enrichment analysis. Additionally, processes with less than 20 DEGs

were removed from downstream analysis. This avoided inclusion of functions represented by

fewer genes and any process that had large number of genes that resulted in generic

descriptions of the processes. This avoided inclusion of functions represented by a few genes

and any process that had large number of genes that resulted in broad or generic descriptions

from the comparison. Heatmaps for gene expression visualization were generated using the

'pheatmap' R package (v. 1.0.12) (Kolde, 2012).

Histochemical detection of hydrogen peroxide and superoxide

Histochemical detection of H_2O_2 was performed on 25 days-old hydroponically grown *A.*

thaliana and *S. parvula* plants using 3,3'-diaminobenzidine (DAB, Sigma-Aldrich, Poland) which

is oxidized in the presence of H_2O_2 and forms a brown stain (Daudi and O'Brien, 2012). We

further assessed the formation of superoxide during high K^+ stress using nitroblue tetrazolium

(NBT) (Thermo Fisher Scientific) which in the presence of O_2^- forms a purple stain (Jabs et al.,

1996). The DAB and NBT solutions were applied on leaves under a gentle vacuum infiltration for

5 minutes followed by shaking at 100 rpm for 4 h. Five mL of 10 mM Na_2HPO_4 without staining

solutions was used as a negative control. Staining solutions were replaced with a bleaching

solution (ethanol: acetic acid: glycerol = 3:1:1) for destaining chlorophyll by boiling for 15

minutes. Destained leaves were imaged using a Zeiss Lumar microscope with x0.8

magnification. We tested three plants per condition and two leaves per plant.

Results

KCl is more toxic than NaCl at the same osmotic strengths

Independent of plant age (Fig 1A, S2) and tissue type (Fig 1C-E), K^+ exerts more severe

growth disturbances than observed for Na^+ at the same concentrations in both species. For

example, *A. thaliana* treated with 100 mM KCl did not survive compared to 100 mM NaCl

treated plants while *S. parvula* treated with 400 mM KCl did not flower compared to 400 mM

NaCl treated plants (Fig S2). It should be noted that the growth media used in the current study

included 1.2 mM K^+ to support plant growth in all conditions and the ≥50 mM KCl treatments

provide excess K^+ beyond its expected range serving as a nutrient (0.1-6 mM) (Ashley et al.,

2006). The more tolerant species, *S. parvula*, was more resilient to higher concentrations of KCl

than *A. thaliana* before it showed significant defects in both primary and lateral growth (Fig 1B,

S3A, and S3B). *S. parvula* root hair length was also less affected compared to that in *A. thaliana*

(Fig 1C, S3C). Compared to the same concentration of NaCl treatments, KCl seemed to be more

toxic even to the more salt stress-tolerant *S. parvula* (Fig 1A, B, D). These results suggest that

despite similar levels of osmotic stress elicited by K^+ and Na^+ at equal concentrations, K^+ may

exert additional stresses different from Na^+ that may initiate physiological responses specific to

K^+ stress.

Because plant growth is tightly coupled to photosynthesis, we tested how the rate of

CO_2 assimilation was affected by K^+ toxicity. CO_2 assimilation was reduced in *A. thaliana* in

response to 150 mM KCl within 24 hours after treatment (HAT), but *S. parvula* did not show any

short-term effects (Fig 1E). Under longer-term treatments, there was a significant reduction in

total leaf area in both species and the reduction was more pronounced with high K^+ (Fig 1D).

Consequently, in *A. thaliana*, the total carbon content in shoots declined within 72 HAT (Fig 1F).

Overall, 150 mM KCl was sufficient to induce physiological stress phenotypes in roots

and shoots of *A. thaliana* within a 3-day period, while impacting long-term growth in *S. parvula*

(Fig 1 and S2). Notably, 150 mM KCl treatment given for more than a week was lethal to *A.*

thaliana in all tested growth conditions. Therefore, we selected 150 mM KCl for our cross-

species comparative-omics analyses to investigate K^+ toxicity responses, when both species are

expected to show active cellular responses to minimize or tolerate stress during the early stages

of the treatment at 0, 3, 24, and 72 HAT (Fig S1).

Excess K^+ causes nutrient depletions in *A. thaliana* but not in *S. parvula*

Given the role of K as an essential macronutrient, plants have evolved multiple

transporters to facilitate entry of K^+ into roots rather than its exit pathways (Shabala and Cuin,

2008). Therefore, we first hypothesized that high K^+ in soils will lead to high K^+ contents in

plants and shoots will accumulate more of the excess K^+ in the transport sequence from soil-

root-shoot. Second, we predict that the differential accumulation of K^+ in roots and shoots will

alter many cellular processes and thereby affect transport and retention of other nutrients. To

test our hypotheses, we quantified the root and shoot contents of 16 established plant

nutrients (K, N, P, Mg, S, Ca, Al, Zn, Co, Ni, Fe, Se, Cu, B, Mn, and Mo) and six toxic elements

found in most soils at low concentrations (Fig 2) in hydroponically grown *A. thaliana* and *S.*

parvula. Nitrogen was measured separately but from the same plant samples (see methods and

Fig S1). With ionomic profiling, we aimed to test if (a) K^+ would accumulate in plants as an

isolated process decoupled from other element accumulation; (b) high K^+ treatment would

cause nutrient imbalances; or (c) it allowed entry of other toxic elements and could indirectly

cause toxicity symptoms.

We observed three notable trends in how the two species partitioned K between roots

and shoots (Fig 2A). First, both species kept higher K^+ levels in shoots than in roots, but the

difference between shoots and roots was strikingly larger in *S. parvula* than in *A. thaliana,*

under control conditions (*i.e.* 0 HAT before exposure to excess K^+). Second, *A. thaliana* reduced

the root K^+ levels by 24 HAT compared to 0 HAT, but was unable to maintain the lower K^+ level

with prolonged stress, leading to significantly increased K^+ levels in both roots and shoots by 72

HAT. Third, *S. parvula* maintained a significantly lower K^+ content in the roots compared to *A.*

thaliana throughout the stress treatment even when the root K^+ levels increased over time

compared to the control. Importantly, *S. parvula* was able to maintain shoot K^+ levels

comparable to the control (Fig 2A). While both species had a higher total K^+ content in shoots

than in roots, only *A. thaliana* allowed a further increase in its shoot K^+ content under the

current experimental conditions. *S. parvula* appeared to avoid using shoots as a sink for excess

K^+ compared to *A. thaliana*.

Higher accumulation of K^+ in *A. thaliana* was not an isolated process and it was coupled

to a severe nutrient imbalance leading to the depletion of seven nutrients (Fig 2B-D). In line

with uninterrupted root growth observed under high K^+ stress (Fig 1A), *S. parvula* did not show

a nutrient imbalance. As summarized in Figure 2D, *S. parvula* showed a remarkable capacity in

maintaining major macronutrients (C, N, and P) at steady states upon exposure to excess K^+,

while in *A. thaliana* all three major nutrients were depleted as K^+ significantly accumulated over

time in both roots and shoots.

We further observed that excess K^+ did not cause accumulation of any other element

that could cause indirect toxicity effects in either species (Fig 2C). Therefore, we can rule out

the possibility of observing indirect toxicity effects from other non-nutrients (heavy metals) in

downstream analyses (metabolome and transcriptome analyses) to deduce cellular processes

affected by excess K^+. Depletion of Na in roots (Fig 2C) in both species indirectly showed the

effort by both species to limit the entry of K^+ and/or increase export using K/Na transporters or

channels that can act non-selectively on both ions. However, the overall effects on the ionomic

profiles indicated that a greater capacity to extrude K^+ from the soil-root interface or prevent

entry into the roots from soil cannot be the only major mechanism underlying K^+ toxicity

tolerance in *S. parvula*. For example, *S. parvula* showed higher fold-increase of root K upon KCl

treatment although the overall K content was still lower than the K content in *A. thaliana* roots

(Fig 2A).

***S. parvula* is more responsive than *A. thaliana* at the metabolome level to high K^+ stress**

We obtained 472 metabolites of which 145 are known and 327 were unannotated

(Table S1) to expand our search to identify cellular processes influencing responses to high K^+

stress (Fig 3). Interestingly, we found that *S. parvula* roots, which showed minimal changes at

the physiological and ionomic levels compared to *A. thaliana* during high K^+ stress, was more

responsive at the metabolomic level (Fig 3A). Considering the higher fraction of unannotated

and hence unknown metabolites in metabolite profiles (Table S1), we tested if the unknown

metabolite pool had a similar response trend to the known metabolites identified between the

two species (Fig 3B). In all tissue/time points, we observed a high correlation of relative

abundances of metabolites, whether known metabolites (Fig 3B, r) or all metabolites including

unknowns (Fig 3B r') were considered. For downstream analysis, we only considered the known

metabolites with significant abundance changes (abbreviated as MACs for this study) between

control and KCl treated conditions (Fig 3A). We further categorized them into (1) amino acids

and immediate precursors or derivatives, (2) fatty acids and lipids, (3) nucleic acids, (4) organic

acids and their immediate derivatives, and (5) sugars, sugar alcohols, and their associated

derivatives (Table S1, Fig 3C).

Overall, the metabolic profile comparisons between the two species showed three

distinct trends. First, with longer stress durations, the number of MACs increased in all tissues

for both species (Fig 3A, C). Second, shoots had more MACs in *A. thaliana*, while *S. parvula*

contained more MACs in roots in response to excess K^+ (Fig 3A, C). The higher responsiveness of

the shoot metabolome in *A. thaliana* is aligned with our initial hypothesis that we would expect

more metabolic disturbances in photosynthetic tissues when excess K^+ accumulated more in

shoots (Fig 2A). In contrast, the *S. parvula* root metabolome was more responsive compared to

both the metabolomes of *A. thaliana* root and *S. parvula* shoot (Fig 3C), suggesting an

alternative response strategy of adjusting predominantly the root metabolites in a species more

tolerant to K^+ toxicity. Third, increased abundances of multiple primary metabolite pools

especially in *S. parvula* roots (Fig 3D, Root, outer circle) were conspicuous against the overall

depletion of primary metabolites in *A. thaliana* in all tissues/time points (Fig 3D, Root and

Shoot, middle circle). One of the largest groups that changed in metabolite abundance during

high K^+ stress was amino acids and their derivatives in *S. parvula*. Further, sugars and the

associated metabolite pools changed in larger proportion in roots of both species and in *A.*

thaliana shoots. This suggested that the two species primarily used different metabolic

groups/pathways to respond to high K^+ stress, a notion further supported by few cross-species

overlaps among MACs (Fig 3A). Therefore, we further examined how amino acids and sugars

significantly changed from control to high K^+ stress conditions over time.

Species specific MACs can indicate metabolic pathways or metabolite groups that are

responding to minimize cellular toxicity, severely interrupted by stress, or both. Excess K

detrimentally affected *A. thaliana* but hardly interrupted the growth of *S. parvula* (Fig 1).

Therefore, we explored the possibility that metabolic pathways or groups represented by MACs

in *S. parvula* were enriched in pathways acting towards minimizing cellular stress, while MACs

in *A. thaliana* were more representative of pathways or groups disrupted by high K^+ stress.

More sugars and amino acids that can serve as osmolytes (raffinose, myo-inositol, sucrose, and

proline) (Fig 3D) (Gong et al., 2005) during salt stress increased in *S. parvula* compared to *A.*

thaliana. When we tested for enriched metabolic pathways or metabolite groups among MACs,

galactose metabolism was enriched in *A. thaliana* shoots and *S. parvula* roots (Fig S4A). The

metabolites in galactose metabolism (KEGG Pathway ID:ath00052) included several key

metabolites (for e.g. galactinol, myo-inositol, and raffinose) (Fig S4B) which are known to

protect plants against oxidative and osmotic stresses (Taji et al., 2002; Nishizawa et al., 2008).

Interestingly, these increased uniquely in *S. parvula* roots (Fig S4B), suggesting coordinated

metabolic adjustments balancing C and N sources to maintain growth while minimizing damage

from oxidative bursts in *S. parvula* roots. Overall, the more stress tolerant species, *S.parvula*,

seemed to boost the protective metabolite pools while the more stress sensitive species, *A.*

thaliana, was depleting its initial pools for such protective metabolites during stress.

A. thaliana shows wide-ranging misregulation of transcriptional resources during peak stress

response times reflective of its critical failures to maintain cellular processes

We next examined the transcriptional stress response landscapes and sought to deduce

specific cellular processes initiated by *A. thaliana* and *S. parvula* after exposure to high external

K^+. Such cellular processes could further build on time-dependent physiological, ionomic, and

metabolic processes we identified earlier in this study as responses to excess K^+ accumulation.

We obtained 12-23 million reads per RNA-Seq library prepared from three biological replicates

of root and shoot samples of both species, harvested at four time points (0, 3, 24, and 72 HAT)

(Fig S1). A total of ~70 - 90% reads were uniquely mapped to primary protein-coding gene

models in *A. thaliana* (TAIR10) and *S. parvula* (V2.2) reference genomes (Table S2).

The Principal Component Analysis (PCA) on shoot and root transcriptomes clustered

samples based on the treatment time points with *A. thaliana* showing a greater separation

between conditions compared to *S. parvula* (Fig S5A). The individual transcriptomes in pairwise

comparisons showed a higher correlation within *S. parvula* samples than for *A. thaliana*

samples (Fig S5B). Overall, *S. parvula* had fewer transcriptome-wide changes in response to

high K^+ stress than *A. thaliana.* PCA on 23,281 ortholog pairs identified between the two species

further supported that *A. thaliana* transcriptomes had a greater variance along a temporal

trajectory for duration of exposure to excess K^+, while *S. parvula* showed a similar trajectory but

with a much smaller variance (Fig 4A). The total number of differentially expressed genes

(DEGs) that showed a significant response in at least one time point was remarkably higher in *A. thaliana* (9907 in roots and 12574 in shoots) compared to that in *S. parvula* (1377 in roots and 494 in shoots) (Fig S6 and Table S3). This observation led us to question whether the stress sensitive *A. thaliana* is transcriptionally adjusting and concurrently failing to maintain steady levels of critical cellular processes, while the more stress tolerant *S. parvula* is only responding with minimal adjustments to a subset of processes *A. thaliana* may be adjusting suboptimally. Therefore, we first identified the key cellular processes and pathways *A. thaliana* was transcriptionally adjusting, to deduce processes interrupted or misregulated due to excess K^+ as well as processes likely activated for protective roles.

We annotated all DEGs with a representative Gene Ontology (GO) function (Table S4, S5, S6) and examined their time dependent progression in *A. thaliana* (Fig 4B and S6). The temporal transcriptomic response climaxed at 24 HAT as shown by having the highest number of DEGs and the largest number of enriched cellular functions. Additionally, the shoots lagged behind the root responses (Fig 4B). This lag is reflected by the most number of DEGs and enriched functions observed during 3-24 HAT in roots and 24-72 HAT in shoots. Despite the response time lag, the enriched processes are largely shared between roots and shoots in *A. thaliana* (Fig 4B).

One of the most emergent cellular process categories enriched in both roots and shoots in *A. thaliana* was response to stress (Fig 4B). Within this category, the prevalent specific responses across all time points were responses to salt, oxidative, and ionic stresses. Unexpectedly, biotic stress-related GO terms (e.g. response to bacterium, response to chitin, and immune response) were also prominently enriched at all time points (Fig 4B). Induction of biotic stress responses in *A. thaliana* seemed counterintuitive when the *S. parvula* orthologs of these *A. thaliana* high K^+-induced DEGs were mostly suppressed in response to high K^+ stress (Table S3 and S7). Given that *A. thaliana* shows a peak transcriptional response at 24 HAT, this made us question whether we were observing transcriptional chaos rather than targeted transcriptional resource allocation between biotic and abiotic stresses to manage the extant stress experienced by *A. thaliana*.

404 We observe three indicators to suggest that there is overall misregulation in critical

405 signaling pathways that highlight chaotic and inefficient responses to overcome cellular toxicity

406 intensified due to excess K^+ accumulation in *A. thaliana* (Fig 2). First, the cellular response to

407 stress in shoots included enriched functions representing all major plant hormone pathways

408 (Fig 4B). In addition to the ABA response enriched at all time points, all other major plant

409 hormone responses are elicited concurrently during 24-72 HAT in shoots in the absence of

410 canonical developmental or biotic stress cues, suggesting disruptions in hormone signaling.

411 Second, we find autophagy as the most enriched process with the highest proportion of DEGs

412 (24 DEGs or 60% of all annotated, adjusted p-value 9.39E-07), accompanied by additional

413 processes including cell death and leaf senescence, enriched during 24-72 HAT in shoots. Third,

414 we see enriched responses to cold, heat, wounding, drought, and hypoxia which all share

415 common genetic mechanisms with responses to oxidative stress also prevalent during 24-72

416 HAT (Fig 4B). This suggested unmitigated oxidative stresses. We compared the ROS scavenging

417 capacity between *A. thaliana* and *S. parvula* leaves during high K^+ stress, by assessing the

418 accumulation of oxidative stress markers, H_2O_2 and O_2^-. As predicted by the transcriptional

419 response, leaves of *A. thaliana* showed greater levels of oxidative stress compared to *S. parvula*

420 (Fig 4C). While the reduced numbers of DEGs at 72 HAT especially in roots indicated a degree of

421 acclimation in *A. thaliana* (Fig 4B and S6B), the prolonged ABA signaling together with non-

422 specific activation of hormone pathways, autophagy, and oxidative stresses serve as molecular

423 phenotypes that indicate erratic and inefficient responses especially in *A. thaliana* shoot, which

424 could lead to eventual failures in cellular homeostasis after prolonged exposure to high K^+

425 stress as supported by the observed growth phenotypes (Fig 1). These molecular phenotypes

426 become even more compelling when compared to their respective transcriptional profiles in *S.*

427 *parvula* orthologs which remained mostly unchanged (Fig 4D).

428 Primary metabolic processes involving carboxylic acid/carbohydrate and amino acid

429 metabolism form the second largest group among the most affected transcriptional categories

430 following stress responses (Fig 4B). These processes were enriched in all time points during high

431 K^+ stress in both *A. thaliana* roots and shoots. This further supports the previous physiological

432 and metabolic responses that showed primary C and N metabolism pathways were severely

affected (Fig 1E and F, Fig 3C and D). Notably, most transcriptional adjustments to C and N

metabolism were detected at later time points. For example, interruptions to photosynthesis at

24 and 72 HAT (Fig 1E) were aligned with the corresponding transcriptional processes enriched

at the same time points in *A. thaliana* shoots (Fig 4B). On the contrary, growth responses

influenced by nutrient homeostasis during stress, such as root development, was detected as a

transcriptionally enriched process early at 3 HAT (Fig 4B), although the interruptions to root

hair growth was detected later at 72 HAT (Fig 1C).

S. parvula shows a confined yet a targeted transcriptomic response steered towards stress

tolerance while supporting enhanced C-N metabolism

We searched for DEGs among *A. thaliana* whose *S. parvula* orthologs also showed active

responses (Fig 5A). These ortholog pairs likely represent cellular processes that require active

transcriptional adjustments to survive the accumulation of excess K^+ in plant tissues even in a

more stress tolerant species. We predicted that the transcriptional adjustments in diametric

responses (*i.e.* genes that are induced in one species when their orthologs are suppressed in

the other species) between the species will be detrimental to the stress sensitive species, while

shared responses in similar directions will be beneficial yet likely underdeveloped or

unsustainable to survive the stress by the stress sensitive species. There were fewer ortholog

pairs showing diametric responses (256 and 60 ortholog pairs in roots and shoots, respectively)

compared to those with shared responses (305 and 283 in roots and shoots respectively)

between *A. thaliana* and *S. parvula* (Fig 5A, S6).

Among the diametric responses, the largest functional cluster identified as induced in *A.*

thaliana, while suppressed in *S. parvula* roots was response to stress (Fig S6A). Most of the

subprocesses in this cluster are associated with biotic stress (Table S9). Therefore, we further

assessed this transcriptional divergence between the two species as a proportional effort

invested in biotic vs abiotic stress responses out of the total DEGs responded in at least one

time point within each species (Fig 5B). The effort to suppress biotic stress responses in *S.*

parvula roots (10%) was similar to the proportional induction for biotic stress responses in *A.*

thaliana (9%) within total non-redundant DEGs (Fig 5B). Contrarily, orthologs that were

suppressed in *A. thaliana*, but induced in *S. parvula* roots were associated with ion transport

and cell wall organization (Fig S6A). These transcriptional adjustments support the physiological

response in *S. parvula* where new roots grew uninterrupted maintaining ion uptake during

exposure to excess K$^+$, compared to *A. thaliana* roots that failed to keep nutrient uptake and

subsequently stopped growing (Fig 1B and 2C). In shoots, orthologs induced in *A. thaliana* but

suppressed in *S. parvula* include multiple genes associated with defense responses,

corroborating our observation of the erratic transcriptional resource allocation to biotic stress

responses in *A. thaliana* (Fig S6B). The orthologs suppressed in *A. thaliana* but induced in *S.*

parvula shoots were enriched in carboxylic acid and amine metabolism and associated

transport functions (Fig S6B). This reflects the transcriptional undertaking in *S. parvula* that

supports continued growth (Fig 1B, D, E, and S2), upheld nutrient balance (Fig 2B, C, and D), and

induction of metabolite pools associated with C and N metabolism (Fig 3C, D) during high K$^+$

stress, while *A. thaliana* shows opposite physiological, ionomic, and metabolic responses.

The ortholog pairs that showed shared induction (156 and 199 in root and shoot,

respectively) in *A. thaliana* and *S. parvula* were largely represented by abiotic stress responses

mediated by hormones including ABA (Fig S6 and Table S10). The ortholog pairs with shared

suppression (149 and 84 in root and shoot, respectively) were enriched in functional processes

involved in biotic stress in roots and photosynthesis in shoots. Therefore, at least in roots, *A.*

thaliana seems to actively reduce its transcriptional allocation to biotic stress via subsets of

genes regulated congruently with *S. parvula* (Fig S6A)*.* However, genes showing this pattern

were negligible based on a proportional effort (0.004% in *A. thaliana*) to reduce transcriptional

allocation to biotic stress responses compared to *S. parvula* (10%) in roots (Fig 5B).

Over 50% of orthologs differently expressed in response to excess K$^+$ in *S. parvula* roots

and ~30% in shoots show unique expression trends different from *A. thaliana* (Fig 5A). Both *S.*

parvula specific responses and diametric responses between the two species were more

pronounced in roots than in shoots. We predicted that *S. parvula* activates decisive

transcriptional regulatory circuits that are either absent (*i.e. S. parvula*-specific responses) or

wired differently (*i.e.* diametric responses) in roots than in shoots when responding to excess K$^+$

stress.

The overall transcriptomic response of *S. parvula* encapsulates induction of more

targeted salt stress responses compared to *A. thaliana*, including especially oxidative stress

responses, sugar and amino acid metabolism, and associated ion transport, with concordant

induction in growth promoting processes and transcriptional resource recuperation by

suppressing biotic stress responses (Fig 5C). The transcriptional effort to facilitate growth

amidst excess K^+ accumulation in tissues is reflected by induced transcripts involved in cell wall

biogenesis, RNA processing, and development along with concurrent suppression for rapid

growth limiting processes such as cell wall thickening and callose deposition (Table S10).

Differential expression of K^+ transporters lead to differential compartmentalization of excess

K^+ concurrent to extensive stress signaling cascades triggered in *A. thaliana*

Our ionome profile comparisons demonstrated that *A. thaliana* exposed to high external

K^+ over time accumulate K^+ in its tissue concurrently with the depletion of C, N, and P at a

whole plant level (Fig 2). Additionally, transcriptional responses to excess K^+ had ion transport

among the most represented functions within and between species comparisons (Fig 4B, 5C,

and S6A). The ability in *S. parvula* to curb K^+ accumulation and prevent the depletion of major

nutrients led us to hypothesize that K^+ transporters are differentially expressed in the two

species as a first line of defense against a severe nutrient imbalance during excess K^+ stress.

Among over 70 transporters reported to transport K^+ in *A. thaliana* (Shabala and Cuin, 2008),

we predicted that the transcripts coding for K^+ transporters which allow K^+ into roots and

upload K^+ into the xylem or phloem would be primary targets for down-regulation in the effort

to restrain K^+ accumulation. On the other hand, transcripts sequestering K^+ into the vacuoles,

especially when tissue $[K^+]$ is high, will be induced. To assess these predictions, we checked the

transcript profiles of all known K^+ transporters.

We first searched for K^+ transporter transcripts that showed significantly different basal

level abundances (Fig 6A) and responses to high K^+ stress (Fig 6B) between *A. thaliana* and *S.*

parvula. We categorized four transport routes that would limit accumulation of K^+ in plants and

compartmentalize excess K^+ within tissues or cells: a. limit entry into roots, b. promote efflux

from roots, c. constrain long distance transport between root and shoot, and d. enhance

sequestration into vacuoles. We then sought to find out routes that are potentially failing to

keep up with the mounting stress in *A. thaliana* and/or more strategically exploited by *S.*

parvula.

a. Limit entry into roots. The most suppressed gene in *A. thaliana* under excess K^+ stress

is the *high-affinity K^+ transporter 5* (*HAK5*) that showed a 13-fold reduction at 24 and 72 HAT. At

low soil K^+ levels, the major transporter for K^+ uptake in roots is HAK5. This mode of K^+

transport switches to low affinity transport mediated by the K channel AKT1 that requires the

channel subunit KC1 to function at sufficient K^+ levels in soil (Gierth et al., 2005; Xu et al., 2006;

Wang et al., 2016) These two major transporters are activated post-translationally by the

protein kinase complex calcineurin B-like proteins 1 and 9 (CBL1&9)/CBL-interacting protein

kinase 23 (CIPK23) (Ragel et al., 2015). Under excess K^+ stress both transport complexes and

their core post-translational regulatory unit in *A. thaliana* roots are suppressed (Fig 6B, C, and

S7A). The transcripts coding for the transporters are unchanged in *S. parvula* roots but the

interacting partners of the kinase complex are suppressed (Fig 6B and S7A). The transcriptional

effort to limit entry of K^+ into roots during high K^+ stress is further exemplified by the

suppression of *RAP2.11*, a transcriptional activator of *HAK5* (Kim et al., 2012) and the induction

of *ARF2*, a repressor of *HAK5* transcription (Zhao et al., 2016), in *A. thaliana* roots (Fig 6C).

Similarly, *ARF2* is also induced in *S. parvula* roots. Other K^+ transporters that are reported to

function in K^+ uptake into roots are also suppressed in *A. thaliana* (e.g. KUP6 and KUP8) under

excess K^+ stress (Fig 6B). While a similar transcriptional suppression for KUP transporters is

absent in *S. parvula* roots*,* it shows concerted transcriptional suppression of cyclic nucleotide

gated channels, *CNGC3/*10/12/13 (Fig 6B and C), suggesting down-regulation of non-selective

uptake of monovalent cations including K^+ into roots (Gobert et al., 2006; Guo et al., 2008).

b. Promote efflux from roots. We do not know of any K^+ efflux transporter that

functions to extrude excess K^+ from roots back to the soil under excess K^+ conditions. However,

S. parvula, which has evolved in soils naturally high in K^+, induced transcription of a K^+ outward

rectifying channel, *GORK* (Ivashikina et al., 2001) and the Na^+ exporter *SOS1* (Shi et al., 2000) in

roots (Fig 6B, C and S8A). Induction of *GORK* is known to cause K^+ leakage from roots under

biotic and abiotic stresses in plants leading to programmed cell death (Demidchik et al., 2014).

We propose that *S. parvula* has evolved to allow export of excess K^+ via induction of GORK

without the destructive downstream consequences of cell death as expected in *A. thaliana* (Fig

6B, C, 4B, and 4D). We also note that the basal expression level of *GORK* in *S. parvula* roots is

higher than in *A. thaliana* roots (Fig 6A). Next, SOS1, the antiporter with the highest Na^+ efflux

capacity in roots, is known for its Na^+ specificity (Oh et al., 2009). Therefore, induction of *SOS1*

in *S. parvula* under high K^+ stress (Fig S8A) is likely an effort to counterbalance the increasing

osmotic stress due to elevated K^+ by exporting available Na^+ from roots. This explanation fits

with Na^+ being the only ion depleted in *S. parvula* roots during excess K^+ stress (Fig 2C).

Similarly, a well-documented Na^+/K^+ transporter family in plants encoded by *HKT1* (Uozumi et

al., 2000; Ali et al., 2016) known for its key roles in minimizing Na-induced salt stress, is present

as a single copy (AtHKT1) for selective Na^+ transport in *A. thaliana*. In *S. parvula*, among the two

duplicated *HKT1* paralogs, *SpHKT1;2* has been demonstrated to show selective-K^+ transport

while Sp*HKT1;1*, orthologous to *A. thaliana HKT1,* shows selective-Na^+ transport (Ali et al.,

2018). We observe basal-level expression biases for the *S. parvula HKT1* paralogs, however the

K^+-selective paralog, despite its much higher transcript abundance, did not significantly respond

to excess K^+ (Fig S8B).

c. Constrain long distance transport between root and shoot. The long-distance

transport of K^+ via xylem loading is mediated by SKOR, NRT1.5, and KUP7 in *A. thaliana*

(Gaymard et al., 1998; Han et al., 2016; Li et al., 2017). *SKOR* and *NRT1.5* were suppressed in *A.*

thaliana roots as predicted. However, *KUP7* showed induction in *A. thaliana* roots at 72 HAT

concordantly when K^+ accumulation was observed in shoots following exposure to high K^+ (Fig

6B and 2A). Contrastingly, none of these transporters were differently regulated in *S. parvula*

roots. AKT2 is the dominant channel protein regulating long distance transport via loading and

unloading to the phloem (Dreyer et al., 2017). It too is significantly suppressed in *A. thaliana*

shoots, but unchanged in *S. parvula* roots and shoots (Fig 6B and C).

d. Enhance sequestration into vacuoles. Vacuolar K^+ concentration can fluctuate widely,

and this sequestration capacity is tightly controlled with pH of the vacuole primarily regulated

by the $Na^+,K^+/H^+$ antiporters, NHX1 and NHX2. These are spatiotemporally regulated with

overlapping functions. Additionally, NHX4 is a more selective K^+/H^+ antiporter functioning to

store K^+ in vacuoles at a smaller scale (Bassil et al., 2019). In *A. thaliana* during high K^+ stress,

NHX2 is induced in roots, whereas *NHX1* is induced in shoots. However, the transcriptional

signal to promote K$^+$ sequestration in *A. thaliana* either in roots or shoots is unclear, because in

roots *NHX1* is suppressed while in shoots *NHX4* is suppressed (Fig 6B and C). This tissue-specific

mixed regulatory pattern observed in *A. thaliana* is absent in *S. parvula*. Nonetheless, *S. parvula*

does not show any pronounced transcriptional commitment to enhance transport of excess K$^+$

into vacuoles in shoots. The nonselective slow-activating vacuolar (SV) channel encoded by

TPC1 (tandem-pore calcium channel) and K$^+$-selective vacuolar channels (TPK/KCO) play primary

roles in regulating the release of K$^+$ into the cytosol from the vacuole (Voelker et al., 2006;

Gobert et al., 2007). Under high K$^+$ stress, *A. thaliana* induces *KCO6* in roots while keeping *TPC1*

unchanged. However, *S. parvula*, through suppression of *KCO5*, seems to attempt to

preferentially retain excess K$^+$ sequestered in vacuoles in roots (Fig 6B and C). Such an attempt

to compartmentalize excess K$^+$ in vacuoles is further suggested by the suppression of tonoplast

localized nonselective cation channels *CNGC19* and *CNGC20* (Yuen and Christopher, 2013) in *S.*

parvula roots.

Our initial physiological assessments showed that photosynthesis was decreased at 24

HAT in *A. thaliana* (Fig 1E). The transcriptional signal of K$^+$ transporters associated with

maintaining turgor of guard cells corroborate the physiological observation, by showing a bias

towards closed stomata via an induction of *GORK* coordinated with a suppression of guard cell

localized *KAT1/2* in *A. thaliana* shoots (Fig 6B and C) (Ivashikina et al., 2001; Szyroki et al.,

2001). The comparable regulatory unit for closed stomata is not induced in *S. parvula* shoots as

a transcriptomic signal.

Osmotic and oxidative stresses combined in plant tissues entail regulation of water

transport mediated by aquaporins and multiple signaling cascades regulated by calcium

signaling pathways (Srivastava et al., 2014; Takahashi and Shinozaki, 2019). Accumulation of

excess K$^+$ leads to a sweeping array of differentially regulated aquaporins and calcium signaling

pathway genes in both roots and shoots of *A. thaliana* compared to a limited regulation

detected for *S. parvula* (Fig S7). This reinforces our overall depiction of the stress response in *S.*

parvula to reflect a restrained set of transcriptional responses during excess K$^+$ stress.

Excess K⁺ induced nitrogen starvation and failure to activate key pathways in N-metabolism are critical flaws in *A. thaliana* compared to *S. parvula*

The steep drop in total N content in *A. thaliana* compared to *S. parvula* (Fig 2B and D); reduction in amino acids and derivatives in *A. thaliana* while these metabolites increased in *S. parvula* (Fig 3C and D); followed by suppression of transcriptomic signals associated with amine metabolism in *A. thaliana* when these were induced in *S. parvula* (Fig 4B, 5C, and S6B) necessitated further examination on how excess K⁺ may alter N-metabolism in plants. Under low soil K levels, N uptake in the form of nitrate is tightly coupled to K uptake and translocation within the plant. Many of the K and N transporters or their immediate post-transcriptional activators are co-regulated at the transcriptional level (Coskun et al., 2017). We searched for specific transcriptomic cues to determine how N transport was disrupted under high K⁺, leading to a deficiency in physiological processes needed to maintain growth and development or a shortage of protective metabolites against oxidative and osmotic stress initiated by excess K⁺ in tissues.

Does excess K⁺ induce N starvation in *A. thaliana*? The dual affinity nitrate transporter, NRT1.1 (NPF6.3/CHL1) is the main NO_3^- sensor in plant roots in addition to accounting for up to 80% of NO_3^- uptake from roots (Feng et al., 2020). Within 3 HAT and onwards, *NRT1.1* in *A. thaliana* roots is down-regulated (Fig 7A). At low soil NO_3^- levels, *NRT1.1* is activated by *CIPK23* to function as a high affinity NO_3^- transporter (Coskun et al., 2017). We note that in *A. thaliana* (and not in *S. parvula*) roots, *CIPK23* is concurrently suppressed with the main K uptake system formed of *HAK5* and *AKT1-KC1* (Fig 6C). This potentially limits the N content in roots within 24 HAT (Fig 2B) and we see *A. thaliana* roots activating N starvation signals by inducing the expression of high affinity NO_3^- transporters, *NRT2.1* and *NRT2.4* upon high K⁺ treatment (Fig 7A) even when the growth medium in the current experimental condition has sufficient NO_3^- levels. NRT2.1 and NRT2.4 account for over 75% of influx when soil NO_3^- levels are low (O'Brien et al., 2016).

The long-distance transport from root to shoot via xylem loading of NO_3^- in roots is primarily regulated via NRT1.5 (NPF7.3) which is a NO_3^-/K⁺ cotransporter (Li et al., 2017). We observe that in *A. thaliana* (and not in *S. parvula*) roots, *NRT1.5* is suppressed possibly in an

attempt to limit excess K^+ accumulation in shoots, but consequently depriving NO_3^- in shoots

(Fig 2B and 7A). This high K^+ induced N starvation environment generated in *A. thaliana* is

further reflected by its additional transcriptional effort to remobilize NO_3^- internally. For

example, *NRT1.7* and *NRT1.8* are induced to promote translocation of NO_3^- from old to young

leaves and from xylem back into roots respectively, while *NRT1.9, NRT1.11,* and *NRT1.12* are

suppressed to restrict transport via phloem and limit NO_3^- movement in shoots under low NO_3^-

levels (Fig 7A) (O'Brien et al., 2016). The transcriptional regulatory emphasis on *NRT1.8* is

outstanding during excess K^+ stress, given that it is the highest induced (104-fold and 73-fold at

24 and 72 HAT, respectively) in the entire *A. thaliana* transcriptomic response (Fig 7A and Table

S3). Even in *S. parvula* which has overall a more restrained transcriptomic response, an ortholog

copy of *AtNRT1;8* is induced in roots at 24 HAT. Additionally, *NRT1.8* is triplicated in *S. parvula*

(Oh and Dassanayake, 2019), allowing additional regulatory flexibility to the species to

redistribute NO_3^- via the xylem back to the roots.

NH_4^+ provides the second major source of N following NO_3^- and the growth medium

provided in the current experimental conditions includes NH_4^+ at 0.2 mM compared to 1.4 mM

NO_3^-. Therefore, we expected to see an induction in NH_4^+ transporters as an alternative N

nutrition response to overcome the high K^+-induced N-starvation, primarily caused through the

co-suppression of NO_3^- and K^+ uptake, in *A. thaliana* roots. However, there is a widely

documented reciprocal antagonistic activity between NH_4^+ and K^+ transport contrasting to the

interdependent cotransport activity between NO_3^- and K^+. Excess NH_4^+ limits root uptake of K^+

and its translocation to shoots and an external supply of K^+ or induction of K^+ transporters

relieves ammonium toxicity (Coskun et al., 2017). Further, the high affinity NH_4^+ specific

ammonium transporters (AMT) are inhibited by CIPK23 when the same is known to activate

NO_3^- and K^+ transporters (Straub et al., 2017). Nonetheless, the antagonistic transport functions

between NH_4^+ and K^+ are not established under excess K^+ and non-toxic NH_4^+ levels.

Counterintuitive to our expectations, *AMT1;1/2/3*, which encode transporters that account for

>90% of high-affinity ammonium uptake into roots (Yuan et al., 2007), were co-suppressed in *A.*

thaliana concurrent to suppression of K^+ transporters that regulate soil to root influx upon

excess K^+ (Fig 7A). The exception to this coordinated suppression was seen with *AMT1;5* which

was induced upon high K^+ in *A. thaliana* roots (Fig 7A). Notably, *AMT1;5* has been shown to

have an opposite activation mode compared to the other *AMTs* (Neuhauser et al., 2007).

We next checked whether the N assimilation pathway from NO_3^- to glutamine via NH_4^+

was also suppressed in *A. thaliana* experiencing a high K^+-induced N-starvation. Indeed, the

genes coding for nitrate reductase (*NIA1/NR1, NIA2/NR2*) and nitrite reductase (*NIR*) were

coordinately down-regulated in *A. thaliana* root and shoot under high K^+ (Fig 7B). In

angiosperms including *A. thaliana*, the main assimilation point of inorganic N to organic

compounds is the GS-GOGAT (glutamine synthetase-glutamate synthase) pathway which is

tightly coupled to the N and C metabolic state of the plant (O'Brien et al., 2016). The GS enzyme

coded by the *GLN* family uses NH_4^+ derived from primary N uptake; recycled from

photorespiration; or remobilized from protein hydrolysis. The GOGAT enzyme comprises

plastid-targeted *GLU* and cytosolic *GLT* to form a net output of glutamate from glutamine and

2-oxoglutarate, which can be converted to various amino acids and organic acids (O'Brien et al.,

2016; Ji et al., 2019). The cytosolic *GLN1;2* and plastidial *GLN2* (coding GS enzymes) together

with *GLT* and *GLU1* (coding GOGAT enzymes) were suppressed under a low N and C metabolic

environment created by excess K^+ (Fig 2D and 7B). Contrastingly, the induction of *GLN1;1* and

GLN1;3 together with *GLU2* especially in *A. thaliana* shoots likely reflects the transcriptional

effort to recycle N under a high K^+-induced N-starved condition (Fig 7B).

We predicted that the suppression of the overall N-assimilation pathway would be

reflected in the change in primary metabolite pools derived from glutamate in *A. thaliana*

during excess K-induced N starvation. In our comparative transcriptome profiling, *S. parvula*

boosted its transcriptional resource allocation to defend against osmotic and oxidative stresses

while *A. thaliana* showed a setback in similar transcriptional efforts (Fig 5B, C and S6). We

wanted to see if this setback in *A. thaliana* to mount appropriate defenses against osmotic and

oxidative stress coincided with the depletion of metabolites directly derived from glutamate

that are known to function as organic osmolytes and antioxidants. Proline is a primary

metabolite with dual functions serving as an osmoprotectant and an antioxidant (Hayat et al.,

2012). We observed a coordinated effort to accumulate proline and its immediate precursors in

the induction of transcripts coding for the key proline biosynthesis enzymes (Fig 7B, *P5CS1/2*) in

both species. However only *S. parvula* was able to significantly accumulate proline during

exposure to excess K^+ (Fig 7C). We see similar pronounced efforts in increasing antioxidant

capacity via GABA and beta-alanine (Fig 7B and C), concordant to increasing synthesis towards

raffinose and myo-inositol only in *S. parvula* (Fig S4B) to protect against osmotic stress. Overall,

S. parvula is not hindered in its capacity to accumulate carbon and nitrogen-rich antioxidants

and osmoprotectants by maintaining N uptake from roots and N-assimilation pathways

separated from the suppressed K-uptake pathways. Contrastingly, the two processes were

jointly suppressed in *A. thaliana* leading to the depleted N resources (Fig 2B) and, in turn,

failure to accumulate C and N-rich protective metabolites (Fig 7C).

Co-expressed gene clusters between *S. parvula* and *A. thaliana* indicate stress preparedness

in *S. parvula*

Given the more targeted transcriptomic responses (Fig 5) and its niche adaptations to

survive high K^+ levels found in its native habitat (Oh et al., 2014), we predicted that, in *S.*

parvula, a large proportion of stress adapted transcripts would be constitutively expressed, as

previously reported for stress preparedness at the basal level for B toxicity (Wang et al., 2021).

To deduce constitutively expressed transcripts in *S. parvula* with decisive roles in preadaptation

toward high K^+, we generated co-expressed ortholog pair clusters using 14,318 and 14,903

ortholog pairs expressed in root and shoot respectively, in *A. thaliana* and *S. parvula*. Among

them, we identified five root and three shoot co-expressed clusters (Fig 8, RC1-5 and SC1-3,

respectively) that had significant basal expression differences and high K^+ stress transcriptomic

responses significantly regulated only in one of the species, a pattern that may signify stress-

prepared transcriptomes unique to each species (Fig 8 and Table S11).

In three out of five root co-expression clusters, only *A. thaliana* orthologs showed

induction or suppression with a maximum response at 24 HAT (Fig 8A, RC1-3). These clusters

largely represented transcripts associated with stress responses, C and N metabolism,

transport, and root development we discussed earlier (Fig 4). Interestingly, the 4th and 5th

clusters (Fig 8A, RC4-5) of which *S. parvula* showed a response, accounted for 203 ortholog

pairs. Yet, 37% of these orthologs had no functional annotations that could be deduced from

Gene Ontologies and therefore could not be meaningfully summarized into representative

processes. This highlights the extent of functional obscurity or novelty of genes that respond to

specific ionic stresses even in *A. thaliana* (Fig 8A, RC5), and the novel regulatory modes

detected in orthologs of closely related species whose functional assignment may have been

overlooked due to the lack of responses in the model plant *A. thaliana* (Fig 8A, RC4). In all three

shoot co-expressed clusters, *A. thaliana* showed a response that peaked at 24 HAT, while the *S.*

parvula orthologs showed a constitutive expression either lower than the basal level in *A.*

thaliana (Fig 8B, SC1), higher than the induced level in *A. thaliana* (Fig 8B, SC2), or equaled the

control (0 HAT) level in *A. thaliana* (Fig 8B, SC3). The enriched functions in shoot clusters largely

overlapped to include stress responses and C and N metabolic processes. In all, 9836 ortholog

pairs, all clusters except RC4 and RC5, showed constitutive expression in *S. parvula* while only

76 ortholog pairs (RC4) showed a unique response from *S. parvula* when *A. thaliana* showed

constitutive expression (Fig 8). Overall, these co-expressed clusters between *A. thaliana* and *S.*

parvula demonstrate the stability in transcriptional resource allocation (*i.e.* stress

preparedness) in *S. parvula* to facilitate growth and development during excess K^+ stress, while

significant alterations are observed for *A. thaliana* orthologs coinciding with its growth

interruptions during high K^+ stress and erratic transcriptional resource management.

Discussion

Salt stress mechanisms induced by high K^+ are largely unknown compared to the

collective understanding for high Na^+ tolerance in plants. Our results demonstrate that high K^+

is more detrimental to plants than Na^+ given at the same external concentrations (Fig 1).

Previous studies support this observation noting that excess KCl caused more severe salt stress

symptoms than NaCl in multiple crops and even among halophytes (Eijk, 1939; Ashby and

Beadle, 1957; Eshel, 1985; Matoh et al., 1986; Cramer et al., 1990; Wang et al., 2001; Ramos et

al., 2004; Richter et al., 2019). Generally, halophytes have evolved to survive high levels of Na

and the subsequent toxicity symptoms entailing oxidative, osmotic, and ionic stresses, while

being able to take up K (Kazachkova et al., 2018; Pantha and Dassanayake, 2020). However, the

canonical adaptations described for salt tolerance mechanisms associated with NaCl-induced

salt stress (Pantha and Dassanayake, 2020) are insufficient to explain adaptations required for

KCl-induced salt stress.

The molecular mechanisms underlying K^+ toxicity, excess K^+ sensing, and how those

signals are transduced to manage overall stress avoidance or mitigation have been poorly

investigated in plants. The extremophyte model, *S. parvula,* amidst high K^+ at unfavorable

growth levels can sustain its growth and development; compartmentalize excess K more in

roots than in shoots; maintain uninterrupted nutrient uptake; increase its antioxidant and

osmoprotectant pools; decouple transcriptional regulation between K and N uptake; and

coordinately induce abiotic stress response pathways along with growth promoting pathways

(Fig 9). Contrastingly, the more stress-sensitive plant, *A. thaliana*, shows interrupted root and

shoot growth; excessive accumulation of K^+ in both roots and shoots; depletion of essential

nutrients; depletion of N-containing primary metabolites; and sweeping transcriptomic

adjustments suggesting initiation of autophagy, ROS accumulation, response to both abiotic

and biotic stresses, and responses to all major hormone pathways (Fig 9). We propose two

deterministic steps in the overall stress response sequence to survive high K^+ stress.

Surviving high K toxicity by avoiding N starvation

Unlike in *A. thaliana,* even when excess K^+ was accumulating in *S. parvula* roots over

time, the overall root growth was not severely affected and nutrient balance was maintained

(Fig 1 and 2). Halophytes are known for their ability to maintain nutrient balance and prevent

excess salt from accumulating in shoots under salt stresses exerted by NaCl (Kazachkova et al.,

2018; Zhao et al., 2020). However, unlike Na, K is a macronutrient and plants have evolved

many more functionally redundant transporters to take up K^+ into roots and redistribute into

shoots via xylem and phloem (Shabala and Cuin, 2008). When external K^+ levels exceed

physiologically optimal conditions, it is not surprising that the immediate response from both

plants was to suppress expression of selective K^+ transporters that primarily control K^+ influx

from the root soil interface (Fig 6). Additionally, *S. parvula* down-regulates non-selective CNGC

cation channels that may be permeable to K^+ especially in roots. Several CNGC type channels

are reported to allow Na^+ or K^+ transport and have been implicated in their functions during Na-

induced salt stress by controlling Na influx into roots. However, their functional and

spatiotemporal specificity remains largely unresolved (Dietrich et al., 2020) and needs to be

determined before evaluating how selected CNGCs may be involved in also limiting excess K

flow into roots under K-induced salt stress.

Concurrent to suppression of K^+ uptake, *A. thaliana* co-suppresses long distance

transport of K^+. One of the key transporters used in this process is NRT1.5 which is a co-

transporter of NO_3^- and K^+ (Li et al., 2017). NO_3^- is transported as a counterion with K^+ in plant

root to shoot translocation as described by the 'Dijkshoorn–Ben Zioni model' (Dijkshoorn et al.,

1968; Zioni et al., 1971; Coskun et al., 2017). The suppression of *NRT1.5* limits NO_3^-

remobilization in plants (Chen et al., 2012). This obstruction to N transport within the plant is

compounded by the transcriptional co-suppression of NRT and AMT transporters known to

operate at the root soil interface limiting N intake into plants (Fig 7) (Tegeder and Masclaux-

Daubresse, 2018). This creates an N-starved condition for *A. thaliana* not observed for *S.*

parvula. During limiting K^+ conditions, N-uptake is down regulated to prevent excess N-induced

toxicity in plants as a favorable mechanism to adapt to K^+ starvation (Armengaud et al., 2004).

This interdependent between N and K transport and regulation that is favorable at low K^+ levels

appear to be detrimental at high K^+ levels as it functions as an antagonistic pleiotropic effect

(condition dependent traits that can cause positive as well as negative impacts).

We observed a significant transcriptional induction of *NRT1.8/NPF7.2* (the most induced

gene in *A. thaliana* during excess K^+ stress) in *A. thaliana* and thereby avoid N depletion in roots

by N-reimport from the stele (Li et al., 2010). However, this transcriptional effort did not

cascade to the ionomic level (Fig 2B and D). N remobilization via induction of *NRT1.8* while

concurrently suppressing *NRT1.5* (Fig 7A) during N starvation is regulated by ethylene-jasmonic

acid signaling together with low N-sensing by nitrate reductase (Chen et al., 2012; Zhang et al.,

2014). Both ethylene and jasmonic acid signaling are among the enriched differently regulated

transcriptional processes in *A. thaliana* (Fig 4B). Interestingly, *S. parvula* appears to have a more

flexible and effective regulatory capacity to allow N-uptake decoupled from restricted K-uptake

and also it does not suppress internal remobilization of NO_3^- and K^+ via NRT1.5 (Fig 6 and 7).

This prevents *S. parvula* from experiencing a high-K induced N-starvation.

The depletion of N uptake in *A. thaliana* further descends into depletion of primary

metabolic pools rich in N (Fig 3D) with a concomitant transcriptional suppression observed in N

assimilation via the GS-GOGAT pathway (Fig 7) (O'Brien et al., 2016; Ji et al., 2019). This not

only creates a shortage of essential primary metabolic pools required for growth and

development, but also depletes essential antioxidant and osmolyte pools to defend against the

mounting oxidative and osmotic stresses (Fig 3, 7, 9). High K in the growth medium is known to

exert osmotic stress at a comparable level to Na and can also lead to oxidative damage

(Osakabe et al., 2013; Zheng et al., 2013). This creates an overall need to boost osmotic and

antioxidant defense systems to successfully survive high K^+ toxicity stress.

Accumulation of compatible osmolytes during osmotic stress reduces water loss from

tissues and helps to maintain turgor pressure which in turn would allow plants to regulate

water transport and photosynthesis (Apse and Blumwald, 2000). Synergistic transcriptional and

metabolic resource allocation to increase osmoprotectants during high K^+ stress is much more

pronounced in *S. parvula* than in *A. thaliana*. Among several osmolytes highlighted in our

results earlier, *S. parvula* roots and shoots notably accumulated significantly higher levels of

proline. Further, the precursors to proline, glutamine and glutamate, accumulated in *S. parvula*

roots (Fig 3 and 7). Stress tolerance upon increased accumulation of proline resulting from both

increased synthesis and reduced catabolism has been previously shown as a key adaptive

mechanism in *E. salsugineum* during salt stress, an extremophyte relative of *S. parvula* (Kant et

al., 2006). Proline plays an important role by serving as an osmolyte, and an antioxidant, and is

reported to facilitate increased photosynthesis during salt stress in many plants (Kishor et al.,

1995; Gong et al., 2005; Kumar et al., 2010; Ghanti et al., 2011; Hayat et al., 2012).

K and N together and independently regulate phosphorus (P) uptake into plants (Coskun

et al., 2017; Maeda et al., 2018; Cui et al., 2019). Additionally, high K^+ toxicity has been reported

to induce P-starvation (Ródenas et al., 2019). Therefore, K and N status of a plant serve as a key

determinant that controls the overall nutrient uptake processes including P uptake in plants

and availability of defense compounds required during salt stress. During high K-induced N-

starved conditions, *A. thaliana* experienced severe shortages of multiple key nutrients (Fig 2),

which *S. parvula* seemingly avoided by having independent regulatory capacity of K and N

uptake and remobilization (Fig 6, 7, and 9). Therefore, we propose that the ability to regulate

independent K^+ uptake is the first key deterministic step towards building resilience to high

external K^+.

Efficient transcriptional resource allocation vs transcriptional mismanagement

Pleiotropic K signaling beneficial under low K conditions lead to wasteful transcriptional

resources during high K^+ conditions in *A. thaliana* and leads to futile activation of multiple

hormone signaling pathways. K serves as a key signaling molecule for hormonal regulation

required for developmental processes, biotic stress signaling, and abiotic stress signaling (Zhang

et al., 2014; Hauser et al., 2017; Shabala, 2017; Hughes et al., 2020; Hetherington et al., 2021).

Plants constantly sense and adjust growth and development based on nutrient availability, and

in the context of K availability, all canonical mechanisms described for hormone-dependent

growth modifications that involve sensing K^+ are based on sensing external K^+ at low or growth

favorable conditions. When supplied with toxic levels of K^+ for most plants, we found that *A.*

thaliana was unable to adjust the K-dependent signaling to fit the extant specific stress

condition and allowed non- selective hormone signaling pathways to be transcriptionally

regulated (Fig 4). This would have led to transcriptional resource mismanagement especially

when biotic stress response pathways were activated concurrent to depleted N-sources and

when it was at a comparable level of proportional allocation assigned to abiotic stress

management (Fig 2, 3, 4B, 5, and S5). This level of non-selective transcriptional activity is

indicative of transcriptional mismanagement in *A. thaliana* seen at 24 HAT and it is not a

signature of deregulation or lack of regulation expected from dying plants as we observe a clear

decline in such non-selective pathway regulation by 72 HAT of exposure to high K^+ (Fig 4B).

Therefore, we propose that the capacity to avoid non-selective transcriptional mismanagement

is the second major deterministic step in surviving high K^+ stress. If unavoided, it can lead to

systemic damage via activation of ROS and autophagy pathways, as demonstrated by *A.*

thaliana with its increased ROS accumulation perhaps resulting from induction of biotic stress

responses or unmitigated oxidative stresses directly induced by high K^+ detrimental especially

at a nutrient starved environment created by high K^+ (Fig 4). ROS signals are known to induce

autophagy often associated with abiotic and biotic stress responses, and during nutrient

recycling, in addition to several plant developmental processes (Liu et al., 2005; Thompson et

al., 2005; Lv et al., 2014; Pantha and Dassanayake, 2020). However, uncontrolled or

misregulated initiation of autophagy signifies a failed stress response strategy (Floyd et al.,

2015). In *A. thaliana* shoots, autophagy is the most enriched transcriptional pathway active at

24 and 72 HAT. The collective transcriptional signal enriched for pathways associated with lipid

catabolism, protein degradation, DNA repair, cell death, and leaf senescence (Das and

Roychoudhury, 2014) (Fig 4B) is further indicative of the maladaptive stress response strategies

shown by *A. thaliana* contrasted against *S. parvula* which shows a more refined transcriptional

response to survive high K^+ stress. The *S. parvula* transcriptome shows a pre-adapted state to

high K^+ stress when examined together with co-expressed orthologs in *A. thaliana. S. parvula*

mostly activates a limited yet focused response towards achieving stress tolerance while

supporting overall growth and metabolism (Fig 5 and 8). Extremophyte transcriptomes and

metabolomes including previous reports on *S. parvula* have shown similar stress-ready states

constant stresses frequently associated with their native environments (Kant et al., 2006; Lugan

et al., 2010; Oh et al., 2014; Lee et al., 2021; Wang et al., 2021).

In conclusion, upon exposure to high K^+, plants undergo physiological, metabolic, and

transcriptional changes and a subset of those changes lead to stress adaptive traits while the

other responses are indicative of failed cellular responses unable to meet the increasing

systemic stress exerted by excess K^+ retention in tissues. The deterministic step whether a plant

would be able to survive K-induced salt stress or descend into unmitigated stress responses was

primarily set early on in the ability to regulate K uptake independent from other nutrient

uptake pathways. This decoupled regulation of K sensing and stress signaling can be leveraged

to design improved crops that are better able to dynamically adjust to a wide array of soils or

irrigation water sources with different salt compositions.

Acknowledgements

We thank Dr. Guannan Wang, Kieu-Nga Tran, Chathura Wijesinghege, Dr. Aaron Smith, and Dr.

John Larkin for providing feedback on the manuscript and facilitating helpful discussions; Prava

Adhikari for additional assistance with phenotyping, and undergraduate students Saad

Chaudhari, Megan Guilbeau, and August Steinkamp for their assistance to grow plants. This

work was supported by the US National Science Foundation awards MCB-1616827 and IOS-

EDGE-1923589, US Department of Energy BER-DE-SC0020358, and the Next-Generation

BioGreen21 Program of Republic of Korea (PJ01317301) awarded to MD and DHO. PP was

supported by an Economic Development Assistantship award from Louisiana State University.

We acknowledge the LSU High Performance Computing services for providing computational

resources needed for data analyses.

Author Contributions

902 PP conducted experiments and performed data analyses. DL supervised and assisted with

903 measuring CO_2 assimilation rates. DHO provided bioinformatics assistance. MD developed the

904 experimental design and supervised the overall project. PP and MD interpreted results and

905 wrote the article with input from all co-authors who revised and approved the final manuscript.

References

Ali A, Khan IU, Jan M, Khan HA, Hussain S, Nisar M, Chung WS, Yun D-J (2018) The High-
Affinity Potassium Transporter EpHKT1;2 From the Extremophile Eutrema parvula
Mediates Salt Tolerance. Front Plant Sci **9**: 1–11

Ali A, Raddatz N, Aman R, Kim S, Park HC, Jan M, Baek D, Khan IU, Oh D, Lee SY, et al (2016) A
Single Amino-Acid Substitution in the Sodium Transporter HKT1 Associated with Plant Salt
Tolerance. Plant Physiol **171**: 2112–2126

Andrews S (2010) FastQC: A quality control tool for high throughput sequence data. Babraham
Bioinforma http://www.bioinformatics.babraham.ac.uk/projects/

Apse MP, Blumwald E (2000) Engineering salt tolerance in plants. Curr Opin Biotechnol **13**:
146–150

Aquea F, Federici F, Moscoso C, Vega A, Jullian P, Haseloff J, Arce-Johnson P (2012) A
molecular framework for the inhibition of Arabidopsis root growth in response to boron
toxicity. Plant, Cell Environ **35**: 719–734

Arienzo M, Christen EW, Quayle W, Kumar A (2009) A review of the fate of potassium in the
soil-plant system after land application of wastewaters. J Hazard Mater **164**: 415–422

Armengaud P, Breitling R, Amtmann A (2004) The Potassium-Dependent Transcriptome of
Arabidopsis Reveals a Prominent Role of Jasmonic Acid in Nutrient Signaling. Plant Physiol
136: 2556–2576

Ashby WC, Beadle NCW (1957) Studies in Halophytes : III . Salinity Factors in the Growth of
Australian Saltbushes Author (s): W . C . Ashby and N . C . W . Beadle Published by : Wiley
on behalf of the Ecological Society of America Stable URL :

https://www.jstor.org/stable/1931695. Ecology **38**: 344–352

Ashley MK, Grant M, Grabov A (2006) Plant responses to potassium deficiencies: a role for
potassium transport proteins. J Exp Bot **57**: 425–436

Bassil E, Zhang S, Gong H, Tajima H, Blumwald E (2019) Cation specificity of vacuolar NHX-type
cation/H + Antiporters 1. Plant Physiol **179**: 616–629

Baxter IR, Ziegler G, Lahner B, Mickelbart M V, Foley R, Danku J, Armstrong P, Salt DE,
Hoekenga OA (2014) Single-Kernel Ionomic Profiles Are Highly Heritable Indicators of
Genetic and Environmental Influences on Elemental Accumulation in Maize Grain (Zea
mays). PLoS One. doi: 10.1371/journal.pone.0087628

Chen CZ, Lv XF, Li JY, Yi HY, Gong JM (2012) Arabidopsis NRT1.5 is another essential component
in the regulation of nitrate reallocation and stress tolerance. Plant Physiol **159**: 1582–1590

Conway JR, Lex A, Gehlenborg N (2017) UpSetR : an R package for the visualization of
intersecting sets and their properties. **33**: 2938–2940

Coskun D, Britto DT, Kronzucker HJ (2017) The nitrogen–potassium intersection: membranes,
metabolism, and mechanism. Plant Cell Environ **40**: 2029–2041

Cramer GR, Epstein E, Lauchli A (1990) Effects of sodium , potassium and calcium on salt-
stressed barley . I . Growth analysis. 83–88

Cui Y, Li X, Yuan J, Wang F, Wang S (2019) Biochemical and Biophysical Research
Communications Nitrate transporter NPF7 . 3 / NRT1 . 5 plays an essential role in
regulating phosphate de fi ciency responses in Arabidopsis. Biochem Biophys Res Commun
508: 314–319

Das K, Roychoudhury A (2014) Reactive oxygen species (ROS) and response of antioxidants as
ROS-scavengers during environmental stress in plants. Front Environ Sci **2**: 1–13

Dassanayake M, Oh D-H, Haas JS, Hernandez A, Hong H, Ali S, Yun D-J, Bressan R a, Zhu J-K,
Bohnert HJ, et al (2011) The genome of the extremophile crucifer Thellungiella parvula.
Nat Genet **43**: 913–918

Daudi A, O'Brien JA (2012) Detection of Hydrogen Peroxide by DAB Staining in. Bio-protocol **2**:
e263

Demidchik V, Straltsova D, Medvedev SS, Pozhvanov GA, Sokolik A (2014) Stress-induced
electrolyte leakage : the role of K+ -permeable channels and involvement in programmed
cell death and metabolic adjustment. J Exp Bot **65**: 1259–1270

Dietrich P, Moeder W, Yoshioka K (2020) Plant cyclic nucleotide-gated channels: New insights
on their functions and regulation. Plant Physiol **184**: 27–38

Dijkshoorn W, Lathwell DJ, Wit CTDE (1968) Temporal changes in carboxylate content of
ryegrass with stepwise change in nutrition. Plant Soil

Dreyer I, Gomez-Porras JL, Riedelsberger J (2017) The potassium battery: a mobile energy
source for transport processes in plant vascular tissues. New Phytol **216**: 1049–1053

Duval JS, Carson JM, Holman PB, Darnley AG (2005) Terrestrial Radioactivity and Gamma-ray
Exposure in the United States and Canada. US Geol Surv Open-File Rep 1413:

Eijk VM (1939) Analysis of the effect of NaCl on development, succulence and transpiration in
Salicornia herbacea, as well as studies on the influence of salt uptake on root respiration in
Aster Tripolium. Recl des Trav Bot néerlandais **36**: 559–657

Eshel A (1985) Response of Suaeda aegyptiaca to KCl, NaCl and Na2SO4 treatments. Physiol
Plant **64**: 308–315

Feng H, Fan X, Miller AJ, Xu G (2020) Plant nitrogen uptake and assimilation: Regulation of
cellular pH homeostasis. J Exp Bot **71**: 4380–4392
Ferreira T, Rasband W (2012) ImageJ User Guide User Guide ImageJ.
Fiehn O, Wohlgemuth G, Scholz M, Kind T, Lee DY, Lu Y, Moon S, Nikolau B (2008) Quality
control for plant metabolomics: Reporting MSI-compliant studies. Plant J **53**: 691–704
Floyd BE, Pu Y, Soto-Burgos J, Bassham DC (2015) To Live or Die: Autophagy in Plants. In:
Gunawardena A., McCabe P. (eds) Plant Programmed Cell Death. Springer, Cham
Gasch AP, Eisen MB (2002) Exploring the conditional coregulation of yeast gene expression
through fuzzy k-means clustering. Genome Biol. doi: 10.1186/gb-2002-3-11-research0059
**Gaymard F, Pilot G, Lacombe B, Bouchez D, Bruneau D, Boucherez J, Michaux-Ferrière N,
Thibaud JB, Sentenac H** (1998) Identification and disruption of a plant shaker-like outward
channel involved in K+release into the xylem sap. Cell **94**: 647–655
Ghanti SKK, Sujata KG, Kumar BMV, Karba NN (2011) Heterologous expression of P5CS gene in
chickpea enhances salt tolerance without affecting yield. Biol Plant **55**: 634–640
Gierth M, Mäser P, Schroeder JI (2005) The potassium transporter AtHAK5 functions in K(+)
deprivation-induced high-affinity K(+) uptake and AKT1 K(+) channel contribution to K(+)
uptake kinetics in Arabidopsis roots. Plant Physiol **137**: 1105–14
Gobert A, Isayenkov S, Voelker C, Czempinski K, Maathuis FJM (2007) The two-pore channel
TPK1 gene encodes the vacuolar K ⅛ conductance and plays a role in K ⅛ homeostasis.
Proc Natl Acad Sci USA **104**: 10726–10731
Gobert A, Park G, Amtmann A, Sanders D, Maathuis FJM (2006) Arabidopsis thaliana Cyclic
Nucleotide Gated Channel 3 forms a non-selective ion transporter involved in germination
and cation transport. J Exp Bot **57**: 791–800
Gong Q, Li P, Ma S, Indu Rupassara S, Bohnert HJ (2005) Salinity stress adaptation competence
in the extremophile *Thellungiella halophila* in comparison with its relative *Arabidopsis
thaliana*. Plant J **44**: 826–839
Gu Z, Gu L, Eils R, Schlesner M, Brors B (2014) circlize implements and enhances circular
visualization in R. Bioinformatics **30**: 2811–2812
Guo K, Babourina O, Christopher DA, Borsics T, Rengel Z (2008) The cyclic nucleotide-gated
channel, AtCNGC10, influences salt tolerance in Arabidopsis. Physiol Plant **134**: 499–507
Han M, Wu W, Wu WH, Wang Y (2016) Potassium Transporter KUP7 Is Involved in K+
Acquisition and Translocation in Arabidopsis Root under K+-Limited Conditions. Mol Plant
9: 437–446
Hauser F, Li Z, Waadt R, Schroeder JI (2017) SnapShot: Abscisic Acid Signaling. Cell **171**: 1708-
1708.e0
**Hayat S, Hayat Q, Alyemeni MN, Wani AS, Pichtel J, Ahmad A, Hayat S, Hayat Q, Alyemeni
MN, Wani AS, et al** (2012) Role of proline under changing environments A review Role of
proline under changing environments A review. doi: 10.4161/psb.21949
Hetherington FM, Kakkar M, Topping JF, Lindsey K (2021) Gibberellin signaling mediates lateral
root inhibition in response to K+ -deprivation. Plant Physiol **185**: 1198–1215
Hughes AM, Zwack PJ, Cobine PA, Rashotte AM (2020) Cytokinin-regulated targets of Cytokinin
Response Factor 6 are involved in potassium transport. Plant Direct 1–15
IPCC (2019) Climate Change and Land: an IPCC special report on climate change, desertification,
land degradation, sustainable land management, food security, and greenhouse gas fluxes

in terrestrial ecosystems [P.R. Shukla, J. Skea, E. Calvo Buendia, V. Masson-De.

Ivashikina N, Dirk B, Ache P, Meyerho O, Felle HH, Hedrich R (2001) K+ channel profile and
electrical properties of Arabidopsis root hairs. FEBS Lett **508**: 463–469

Jabs T, Dietrich RA, Dangl JL (1996) Initiation of runaway cell death in an Arabidopsis mutant by
extracellular superoxide. Science (80-) **273**: 1853–1856

Ji Y, Li Q, Liu G, Selvaraj G, Zheng Z, Zou J, Wei Y (2019) Roles of Cytosolic Glutamine
Synthetases in Arabidopsis Development and Stress Responses. Plant Cell Physiol **60**: 657–
671

Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M (2016) KEGG as a reference
resource for gene and protein annotation. Nucleic Acids Res **44**: D457–D462

Kant S, Kant P, Raveh E, Barak S (2006) Evidence that differential gene expression between the
halophyte, Thellungiella halophila, and Arabidopsis thaliana is responsible for higher levels
of the compatible osmolyte proline and tight control of Na+ uptake in T. halophila. Plant,
Cell Environ **29**: 1220–1234

Kazachkova Y, Eshel G, Pantha P, Cheeseman JM, Dassanayake M, Barak S (2018)
Halophytism: What Have We Learnt From Arabidopsis Relative Model Systems? Plant
Physiol **178**: 972–988

Kim MJ, Ruzicka D, Shin R, Schachtman DP (2012) The Arabidopsis AP2 / ERF Transcription
Factor RAP2 . 11 Modulates Plant Response to Low-Potassium Conditions. Mol Plant **5**:
1042–1057

Kishor PBK, Hong Z, Miao C-H, Hu C-AA, Verma DPS (1995) Overexpression of A1-Pyrroline-5-
Carboxylate Synthetase Increases Proline Production and Confers Osmotolerance in
Transgenic Plants '. Plant Physiol **108**: 1387–1394

Kolde R (2012) pheatmap v.1.0.8. https://cran.r-project.org/package=pheatmap 1–7

Kumar V, Shriram V, Kishor PBK (2010) Enhanced proline accumulation and salt stress
tolerance of transgenic indica rice by over-expressing P5CSF129A gene. Plant Biotechnol
Rep **4**: 37–48

Langmead B, Trapnell C, Pop M, Salzberg SL (2009) Ultrafast and memory-efficient alignment
of short DNA sequences to the human genome. Genome Biol **10**: R25

Lee G, Ahmadi H, Quintana J, Syllwasschy L, Preite V, Anderson JE, Genetics M, Information C,
Footnotes A (2021) Constitutively enhanced genome integrity maintenance and direct
stress mitigation characterize transcriptome of extreme stress-adapted Arabidopsis halleri.
bioRxiv

Leigh R, Wyn Jones RG (1984) A hypothesis relating critical potassium concentrations for
growth to the distribution and function of this ion in the plant cell. New Phytol. 97. 1–13

Lequeux H, Hermans C, Lutts S, Verbruggen N (2010) Response to copper excess in Arabidopsis
thaliana: Impact on the root system architecture, hormone distribution, lignin
accumulation and mineral profile. Plant Physiol Biochem **48**: 673–682

Li H, Yu M, Du XQ, Wang ZF, Wu WH, Quintero FJ, Jin XH, Li HD, Wang Y (2017) NRT1.5/NPF7.3
functions as a proton-coupled H+/K+ antiporter for K+ loading into the xylem in
arabidopsis. Plant Cell **29**: 2016–2026

Li JY, Fu YL, Pike SM, Bao J, Tian W, Zhang Y, Chen CZ, Zhang Y, Li HM, Huang J, et al (2010)
The Arabidopsis nitrate transporter NRT1.8 functions in nitrate removal from the xylem
sap and mediates cadmium tolerance. Plant Cell **22**: 1633–1646

Liu Y, Schiff M, Czymmek K, Tallóczy Z, Levine B, Dinesh-Kumar SP (2005) Autophagy regulates
programmed cell death during the plant innate immune response. Cell **121**: 567–577

Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol **15**: 550

Lugan R, Niogret MF, Leport L, Guégan JP, Larher FR, Savouré A, Kopka J, Bouchereau A (2010)
Metabolome and water homeostasis analysis of Thellungiella salsuginea suggests that
dehydration tolerance is a key response to osmotic stress in this halophyte. Plant J **64**:
215–229

Lv X, Pu X, Qin G, Zhu T, Lin H (2014) The roles of autophagy in development and stress
responses in Arabidopsis thaliana. Apoptosis **19**: 905–921

Maathuis FJ (2009) Physiological functions of mineral macronutrients. Curr Opin Plant Biol **12**:
250–258

**Maeda Y, Konishi M, Kiba T, Sakuraba Y, Sawaki N, Kurai T, Ueda Y, Sakakibara H, Yanagisawa
S** (2018) A NIGT1-centred transcriptional cascade regulates nitrate signalling and
incorporates phosphorus starvation signals in Arabidopsis. Nat Commun. doi:
10.1038/s41467-018-03832-6

Maere S, Heymans K, Kuiper M (2005) BiNGO: A Cytoscape plugin to assess overrepresentation
of Gene Ontology categories in Biological Networks. Bioinformatics **21**: 3448–3449

Matoh T, Watanabe J, Takahashi E (1986) Effects of Sodium and Potassium Salts on the Growth
of a Halophyte. Soil Sci Plant Nutr **32**: 451–459

Murashige T, Skoog F (1962) A Revised Medium for Rapid Growth and Bio Assays with Tobacco
Tissue Cultures. Physiol Plant **15**: 473–497

Neuhauser B, Dynowski M, Mayer M, Ludewig U (2007) Regulation of NH 4 1 Transport by
Essential Cross Talk between AMT Monomers through the. Plant Physiol **143**: 1651–1659

Nilhan TG, Emre YA, Osman K (2008) Soil Determinants for distribution of Halocnemum
strobilaceum Bieb. (Chenopodiaceae)Around Lake Tuz, Turkey. Pakistan J Biol Sci **11**: 565–
570

Nishizawa A, Yabuta Y, Shigeoka S (2008) Galactinol and Raffinose Constitute a Novel Function
to Protect Plants from Oxidative Damage. Plant Physiol **147**: 1251–1263

O'Brien JAA, Vega A, Bouguyon E, Krouk G, Gojon A, Coruzzi G, Gutiérrez RAA (2016) Nitrate
Transport, Sensing, and Responses in Plants. Mol Plant **9**: 837–856

Oh D-H, Dassanayake M (2019) Landscape of gene transposition-duplication within the
Brassicaceae family. DNA Res **26**: 21–36

Oh D-H, Hong H, Lee SY, Yun D-J, Bohnert HJ, Dassanayake M (2014) Genome structures and
transcriptomes signify niche adaptation for the multi-ion tolerant extremophyte
Schrenkiella parvula. Plant Physiol **164**: 2123–2138

**Oh D-H, Leidi E, Zhang Q, Hwang S-M, Li Y, Quintero FJ, Jiang X, D'Urzo MP, Lee SY, Zhao Y, et
al** (2009) Loss of halophytism by interference with SOS1 expression. Plant Physiol **151**:
210–222

**Osakabe Y, Arinaga N, Umezawa T, Katsura S, Nagamachi K, Tanaka H, Ohiraki H, Yamada K,
Seo S-U, Abo M, et al** (2013) Osmotic Stress Responses and Plant Growth Controlled by
Potassium Transporters in Arabidopsis. Plant Cell **25**: 609–624

Pantha P, Chalivendra S, Oh DH, Elderd BD, Dassanayake M (2021) A tale of two transcriptomic
responses in agricultural pests via host defenses and viral replication. Int J Mol Sci **22**: 1–27

Pantha P, Dassanayake M (2020) Living with Salt. Innov **1**: 100050

Ragel P, Ródenas R, García-martín E, Andrés Z, Villalta I, Nieves-cordones M, Rivero RM,
Martínez V, Pardo JM, Quintero FJ, et al (2015) The CBL-Interacting Protein Kinase CIPK23
Regulates HAK5-Mediated High-Af fi nity K + Uptake in Arabidopsis Roots 1 [OPEN]. Plant
Physiol **169**: 2863–2873

Ramos J, López MJ, Benlloch M (2004) Effect of NaCl and KCl salts on the growth and solute
accumulation of the halophyte Atriplex nummularia. Plant Soil **259**: 163–168

Richter JA, Behr JH, Erban A, Kopka J, Zörb C (2019) Ion-dependent metabolic responses of
Vicia faba L. to salt stress. Plant Cell Environ **42**: 295–309

Ródenas R, Martinez V, Nieves-cordones M, Rubio F (2019) High External K + Concentrations
Impair Pi Nutrition , Induce the Phosphate Starvation Response , and Reduce Arsenic
Toxicity in Arabidopsis Plants. Int J Mol Sci **20**: 1–18

Shabala S (2017) Signalling by potassium: Another second messenger to add to the list? J Exp
Bot **68**: 4003–4007

Shabala S, Cuin TA (2008) Potassium transport and plant salt tolerance. Physiol Plant **133**: 651–
669

Shi H, Ishitani M, Kim C, Zhu J-K (2000) The Arabidopsis thaliana salt tolerance gene SOS1
encodes a putative Na+/H+ antiporter. Proc Natl Acad Sci USA **97**: 6896–6901

Srivastava AK, Penna S, Nguyen D Van, Tran LP (2014) Multifaceted roles of aquaporins as
molecular conduits in plant responses to abiotic stresses. Crit Rev Biotechnol **8551**: 1–10

Straub T, Ludewig U, Neuhäuser B (2017) The Kinase CIPK23 Inhibits Ammonium Transport in
Arabidopsis thaliana. Plant Cell **29**: 409–422

Szyroki A, Ivashikina N, Dietrich P, Roelfsema MRG, Ache P, Reintanz B, Deeken R, Godde M,
Felle H, Steinmeyer R, et al (2001) KAT1 is not essential for stomatal opening. Proc Natl
Acad Sci U S A **98**: 2917–2921

Taji T, Ohsumi C, Iuchi S, Seki M, Kasuga M, Kobayashi M, Yamaguchi―Shinozaki K,
Shinozaki K (2002) Important roles of drought- and cold-inducible genes for galactinol
synthase in stress tolerance in Arabidopsis thaliana. Plant J **29**: 417–426

Takahashi F, Shinozaki K (2019) Long-distance signaling in plant stress response. Curr Opin
Plant Biol **47**: 106–111

Tegeder M, Masclaux-Daubresse C (2018) Source and sink mechanisms of nitrogen transport
and use. New Phytol **217**: 35–53

Thompson AR, Doelling JH, Suttangkakul A, Vierstra RD (2005) Autophagic Nutrient Recycling
in Arabidopsis Directed by the ATG8 and ATG12 Conjugation Pathways 1. Plant Physiol
138: 2097–2110

Uozumi N, Kim EJ, Rubio F, Yamaguchi T, Muto S, Tsuboi a, Bakker EP, Nakamura T,
Schroeder JI (2000) The Arabidopsis HKT1 gene homolog mediates inward Na(+) currents
in xenopus laevis oocytes and Na(+) uptake in Saccharomyces cerevisiae. Plant Physiol **122**:
1249–1259

Voelker C, Schmidt D, Mueller-roeber B, Czempinski K (2006) Members of the Arabidopsis
AtTPK / KCO family form homomeric vacuolar channels in planta. Plant J **48**: 296–306

Wang B, Lüttge U, Ratajczak R (2001) Effects of salt treatment and osmotic stress on V-ATPase
and V-PPase in leaves of the halophyte Suaeda salsa. J Exp Bot **52**: 2355–2365

Wang G, DiTusa SF, Oh DH, Herrmann AD, Mendoza-Cozatl DG, O'Neill MA, Smith AP,

Dassanayake M (2021) Cross species multi-omics reveals cell wall sequestration and
elevated global transcription as mechanisms of boron tolerance in plants. New Phytol. doi:
10.1101/2020.10.01.321760
Wang G, Oh D, Dassanayake M (2020) GOMCL: a toolkit to cluster, evaluate, and extract non-
redundant associations of Gene Ontology-based functions. BMC Bioinformatics **21**: 1–9
Wang X-P, Li-MeiChen, Liu W-X, Shen L-K, Wang F-L, Zhou Y, Zhang Z, Wu W-H, Wang Y (2016)
AtKC1 and CIPK23 Synergistically Modulate AKT1-Mediated Low-Potassium Stress. Plant
Physiol **170**: 2264–2277
Wang Y, Wu W-H (2013) Potassium Transport and Signaling in Higher Plants. Annu Rev Plant
Biol **64**: 451–476
Warren JK (2016) Potash Resources: Occurrences and Controls. Evaporites A Geol Compend.
doi: 10.1007/978-3-319-13512-0_11
Xu J, Li HD, Chen LQ, Wang Y, Liu LL, He L, Wu WH (2006) A Protein Kinase, Interacting with
Two Calcineurin B-like Proteins, Regulates K+ Transporter AKT1 in Arabidopsis. Cell **125**:
1347–1360
Yoshitake Y, Nakamura S, Shinozaki D, Izumi M (2021) RCB-mediated chlorophagy caused by
oversupply of nitrogen suppresses phosphate-starvation stress in plants. Plant Physiol **185**:
318–330
Yuan L, Loqué D, Kojima S, Rauch S, Ishiyama K, Inoue E, Takahashi H, Von Wirén N (2007) The
organization of high-affinity ammonium uptake in Arabidopsis roots depends on the
spatial arrangement and biochemical properties of AMT1-type transporters. Plant Cell **19**:
2636–2652
Yuen CCY, Christopher DA (2013) The group IV-A cyclic nucleotide-gated channels, CNGC19 and
CNGC20, localize to the vacuole membrane in Arabidopsis thaliana. AoB Plants **5**: 1–14
Zhang G Bin, Yi HY, Gong JM (2014) The Arabidopsis Ethylene/Jasmonic acid-NRT signaling
module coordinates nitrate reallocation and the trade-off between growth and
environmental adaptation. Plant Cell **26**: 3984–3998
Zhao C, Zhang H, Song C, Zhu J-K, Shabala S (2020) Mechanisms of Plant Responses and
Adaptation to Soil Salinity. Innov **1**: 100017
Zhao S, Zhang M, Wang Y (2016) Phosphorylation of ARF2 Relieves Its Repression of
Transcription of the K + Transporter Gene HAK5 in Response to Low Potassium Stress.
Plant Cell **28**: 3005–3019
Zheng S, Pan T, Fan L, Qiu Q (2013) A Novel AtKEA Gene Family , Homolog of Bacterial K+/ H+
Antiporters , Plays Potential Roles in K + Homeostasis and Osmotic Adjustment in
Arabidopsis. PLoS One **8**: e81463
Ziegler G, Terauchi A, Becker A, Armstrong P, Hudson K, Baxter I (2013) Ionomic Screening of
Field-Grown Soybean Identifies Mutants with Altered Seed Elemental Composition. Plant
Genome **6**: 0
Zioni A BEN, Vaadia Y, Lips SH (1971) Nitrate Uptake by Roots as Regulated by Nitrate
Reduction Products of the Shoot. Physiol Plant **24**: 288–290

 Figures

Figure 1. KCl is more toxic than NaCl at the same osmotic strength. [A] Seedlings of *Arabidopsis thaliana* (on the left for each plate) and *Schrenkiella parvula* (on the right for each plate) on 1/4 MS media supplemented with 0 to 200 mM NaCl and KCl. **[B]** Primary root length, lateral root number, and lateral root density measured on 17-day-old seedlings grown under conditions used in [A] based on a 12-day treatment of NaCl or KCl. **[C]** Root hair length measured under the same growth conditions used in [A] for variable KCl treatments and monitored for a week. **[D]** Leaf area measured for 5- and 6-week-old hydroponically grown *A. thaliana* and *S. parvula* treated with increasing levels of NaCl and KCl for 1- and 3-weeks respectively. **[E]** Photosynthesis measured as the rate of CO_2 assimilation of the entire shoot/rosette in 25-day-old hydroponically grown plants and monitored up to 72 HAT. **[F]** Total carbon as a % weight based on total dry mass for root and shoot tissue under conditions used in [E]. Minimum 5 plants per condition used in B and C and a minimum of 3 plants per condition used for D and E. Asterisks indicate significant changes between the treated samples to its respective control samples (t-test with $p < 0.05$). Data are presented as mean of at least 3 independent biological replicates ± SD. DAT-Days after treatment, HAT- Hours after treatment.

Figure 2. Excess K accumulation caused severe nutrient imbalance in *A. thaliana* compared to *S. parvula*.
[A] K accumulated differently between *A. thaliana* and *S. parvula*. *A. thaliana* promoted shoot retention while *S. parvula* promoted root retention of K. **[B]** Increased accumulation of K resulted in N depletion in *A. thaliana* compared to *S. parvula*. **[C]** Differential accumulation of K caused imbalances of multiple nutrients in *A. thaliana* roots even in early time points and caused hardly any noticeable differences in *S. parvula*. **[D]** Percent change in CNPK elemental content in root and shoot of *A. thaliana* and *S. parvula* under high [K⁺]. *A. thaliana* fails to maintain major macronutrient levels preventing depletion of those pools contrasted to *S. parvula*. Data are represented as mean of at least 4 (for A and C) and 3 (for B) independent replicates with ± SD given using ≥ 5-8 hydroponically grown plants per replicate. Elemental compositions were quantified using ICP-MS and total N was obtained using an elemental combustion system. The total elemental composition is reported after normalization for dry weight. All quantitative measurements were evaluated by one-way ANOVA followed by Tukey's post-hoc test, *p-value* <0.05 and same letters in the bar graph does not differ statistically significantly and asterisks indicate significant changes between the treated samples to its respective control samples. HAT- Hours after treatment.

1191

1192

1193

1194

1195

1196

1197

1198

Figure 3. *S. parvula* root metabolome is more responsive than *A. thaliana* and induces specific antioxidants and osmoprotectants to adapt to high K$^+$ stress in both roots and shoots. **[A]** Metabolites that significantly changed in abundance at 24 and 72 hours after treatment (HAT). **[B]** *A. thaliana* and *S. parvula* metabolome profiles are highly corelated with each other upon high K$^+$ in both roots and shoots. Pearson correlation coefficient is calculated for 145 known metabolites (r) and all metabolites including the unannotated metabolites (r'). **[C]** Overview of the temporal changes in primary metabolic pools based on known metabolite annotations. Metabolites that significantly changed in abundance at respective time points are shown followed by a "/" and the total number of metabolites counted in that category. **[D]** Individual metabolites in each functional group mapped to represent their abundance starting at basal level (inner circles) to 24 and 72 h of exposure to high K shown in concentric outer rings. Metabolites mostly highlighted in Results and Discussion are marked with arrow heads in the outer circle in both plots. Significance test for metabolite abundance was performed with one-way ANOVA followed by Tukey's post-hoc test, *p-value* <0.05. Data are represented as mean of at least 3 independent replicates ± SD given using ≥ 5 hydroponically grown plants per replicate.

Figure 4. *A. thaliana* shows non-selective transcriptional modifications during excess K⁺ stress than *S. parvula*. [A] Principal component analysis (PCA) of ortholog expression between *A. thaliana* and *S. parvula* root and shoot transcriptomes at 0, 3, 24, and 72 hours after treatment (HAT). **[B]** Enriched functional processes based on GO annotations associated with differently expressed genes (DEGs) in *A. thaliana*. The temporal sequence of enriched functions is given as 3 h specific, 3 and 24 h together with 24 h specific, and 72 h, 72 h specific, and present at all-time points from 3-24-72 h. Only functional processes that were detected at least in two time points are shown and the processes are sorted based on their functional hierarchy when applicable. **[C]** *A. thaliana* accumulated higher levels of hydrogen peroxide (H_2O_2) and superoxide ions (O_2^-) in leaves under excess K⁺ stress compared to *S. parvula*. Hydroponically grown plants were used treated similar to the plants used for the RNA-Seq study and stained with DAB and NBT. **[D]** Transcriptional profiles of selected pathways that serve as indicators of transcriptional mismanagement in *A. thaliana* during excess K⁺ stress compared to *S. parvula*. DEGs were called using DESeq2 with a p-adj value based on Benjamini-Hochberg correction for multiple testing set to <0.01.

1201

1202

1203

1204

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

Figure 5. *S. parvula* **shows a confined transcriptomic response geared toward concurrent induction of abiotic stress responses and enhanced transcriptional allocation to C and N metabolism. [A]** The overall expression specificity and response direction of orthologs in *A. thaliana* and *S. parvula.* Selected orthologs are differentially expressed genes (DEGs) at least in one time point compared to the respective control condition and then counted as a non-redundant set when all 3, 24, and 72 HAT samples were considered for total counts. **[B]** The proportion contributing to abiotic and biotic stress stimuli within non-redundant DEGs. The effort to suppress biotic stress responses in *S. parvula* roots (10%) was similar to the proportional induction for biotic stress responses in *A. thaliana* (9%). **[C]** The functionally enriched processes represented by DEGs in *S. parvula* that responded to high K^+. A non-redundant set from all time points (3, 24, and 72 HAT) are given. A node in each cluster represents a gene ontology (GO) term; size of a node represents the number of genes included in that GO term; the clusters that represent similar functions share the same color and are given a representative cluster name and ID; and the edges between nodes show the DEGs that are shared between functions. All clusters included in the network have adj p-values <0.05 with false discovery rate correction applied. More significant values are represented by darker node colors. The functional enrichment network was created using GOMCL. DEGs were called using DESeq2 with a p-adj value based on Benjamini-Hochberg correction for multiple testing set to <0.01.

1218

Figure 6. Differential expression of K⁺ transporters in *A. thaliana* and *S. parvula*. [A] Basal level expression comparison of orthologs between *A. thaliana* and *S. parvula* in roots and shoots. The dash-gray diagonal line marks identical expression in both species and solid red lines represent a 2-fold change in one species compared to the other species. Transporters/channels with >2-fold change basal differences between the species are labeled. **[B]** The temporal expression of functionally established K⁺ transporters and channels in *A. thaliana* and their *S. parvula* orthologs upon high K⁺ treatment. **[C]** Key K⁺ transporters and channels differently regulated between roots and shoots in *A. thaliana* and *S. parvula*. DEGs at each time point were called using DESeq2 compared to 0 h with a p-adj value based on Benjamini-Hochberg correction for multiple testing set to <0.01.

1219

1220

1221

1222

1223

1224

Figure 7. Excess K⁺ induced nitrogen starvation and suppressed associated N-assimilation pathways in *A. thaliana* compared to *S. parvula*. [A] Nitrogen uptake and distribution was severely affected in *A. thaliana* upon high K⁺. The expression changes associated with major nitrogen transporters in *A. thaliana* that regulate root uptake and long-distance transport of N are summarized. **[B]** Coordinated transcriptional regulation showing high K⁺-induced suppression of nitrogen assimilation and efforts to accumulate glutamate-derived osmoprotectants. The arrows in front of heatmap blocks indicate the direction of the reaction. **[C]** The primary metabolite pools derived from glutamate in roots and shoots. DEGs at each time point were called using DESeq2 compared to 0 h with a p-adj value based on Benjamini-Hochberg correction for multiple testing set to <0.01. Significance test for metabolite abundances was performed with one-way ANOVA followed by Tukey's post-hoc test, *p-value* <0.05. Data are represented as mean of at least 3 independent replicates ± SD (≥ 5 hydroponically grown plants per replicate).

1225

1226

1227

1228

1229

Figure 8. Co-expressed ortholog gene modules highlight stress preparedness towards high K+ stress in *S. parvula* orthologs that are constitutively expressed compared to induction or suppression of orthologs in *A. thaliana*. Normalized gene expression clusters of ortholog pairs (OP) between *A. thaliana* and *S. parvula* in **[A]** roots and **[B]** shoots. Fuzzy K-means clustering was used to find temporally co-regulated clusters with a membership cutoff of ≥0.5. Box and whisker plots mark the median expression at each time point with the thick line within the box; interquartile range between first and third quartile are shown with the box; and interquartile range x 1.5 are marked by whiskers for lower and upper extremes. Basal level for 0 h in *A. thaliana* is marked by a grey line in all plots. Each cluster was used for a functional enrichment analysis represented by GOMCL summaries placed below co-expression plots. A node in each cluster represents a gene ontology (GO) term; size of a node represents the number of genes included in that GO term; the clusters that represent similar functions share the same color and are given a representative cluster name and the edges between nodes show the orthologs that are shared between functions. All clusters included in the network have adj p-values <0.05 with false discovery rate correction applied. More significant values are represented by darker nodes.

1230

Figure 9. Deterministic cellular processes in surviving high K⁺ stress.

1231

1232

1233

1234

1235

1236

1237

1238

1239

1240

1241

1242

1243

1244

1245

1246   **Supplementary figures**

Figure S1. Sampling scheme for the ionome, metabolome, and transcriptome experiments performed in this study. 25-day-old *Arabidopsis thaliana* and *Schrenkiella parvula* plants were grown in 1/5th Hoagland's solution with/without 150 mM KCl for up to 72 hours after treatment (HAT). All samples were 28-day-old at the time of sample harvest. Both ionome and metabolome profiles have 4 biological replicates and transcriptome samples have 3 biological replicates. Each replicate contains tissues from at least 5 different plants.

1247

1248

Figure S2. KCl is more toxic than NaCl at the same osmotic strength. [A] 21-day-old plants treated for 35 days every other day. Yellow arrow heads indicate visibly wilting shoots only in KCl treated plants. **[B]** The same plants imaged after 10 days show 100 mM KCl treated plants did not survive while 100 mM NaCl treated plants appear to be stressed but not dead in *A. thaliana*. In *S. parvula* 400 mM KCl treated plants did not develop any flower while 400 mM NaCl treated plants flowered and developed seeds (indicated with purple arrow heads).

1249

1250

Figure S3. Reduction in primary root length **[A]** and leaf area **[B]** upon Na and K treatment. Primary root length was measured on 17-day-old seedlings grown on 1/4 MS media supplemented with 0 to 200 mM NaCl and KCl based on a 12-day treatment of NaCl or KCl. Leaf area was measured for 5- and 6-week-old hydroponically grown *A. thaliana* and *S. parvula* treated with increasing levels of NaCl and KCl for 1- and 3-weeks respectively. Minimum 5 plants per condition used in A and a minimum of 3 plants per condition used for B. **[C]** Representative images showing the effect of KCl on root hair growth in *A. thaliana* and *S. parvula*. Scale bars are 1 mm for all the panels. Seeds were germinated in 1/4th MS and transferred to the corresponding plates with 100 or 150 mM KCl 5 days after germination. Same roots were photographed on 3, 5, and 7 DAT. DAT-Days after treatment. Asterisks indicate significant changes between the treated samples to its respective control samples (t-test with p <0.05). DAT-Days after treatment.

1251

1252

[A]

Root

A. thaliana Enriched processes	Total # of metabolites	# of MACs*	P-value	P-adj
Galactose Metabolism	38	4	0.004	0.411
Lactose Degradation	9	2	0.011	0.533

S. parvula Enriched processes	Total # of metabolites	# of MACs*	P-value	P-adj
Galactose Metabolism	38	7	0.0002	0.016
Urea Cycle	29	6	0.0003	0.016
Aspartate Metabolism	35	5	0.006	0.199
Glucose-Alanine Cycle	13	3	0.009	0.22
Ammonia Recycling	32	4	0.023	0.449
Arginine and Proline Metabolism	53	5	0.034	0.519
Lactose Degradation	9	2	0.037	0.519
Malate-Aspartate Shuttle	10	2	0.045	0.555

Shoot

A. thaliana Enriched processes	Total # of metabolites	# of MACs*	P-value	P-adj
Galactose Metabolism	38	7	0.0003	0.025
Lactose Degradation	9	3	0.003	0.154
Glycerolipid Metabolism	25	4	0.011	0.26
Glycolysis	25	4	0.011	0.26
Lactose Synthesis	20	3	0.032	0.553
Gluconeogenesis	35	4	0.034	0.553

S. parvula Enriched processes	Total # of metabolites	# of MACs*	P-value	P-adj
Galactose Metabolism	38	5	0.002	0.196
Lactose Degradation	9	2	0.02	0.981
Glucose-Alanine Cycle	13	2	0.041	1

Significantly enriched process

*Metabolites that significantly changed in abundance

[B]

Figure S4. Carbohydrate and nitrogen metabolism related processes were enriched in *S. parvula* roots. **[A]** Enriched metabolic pathways in *A. thaliana* and *S. parvula* in response to excess K stress. Metabolite Enrichment Analysis was performed with MetaboAnalyst with a p-adj cut off set to < 0.05. p-values were adjusted using Benjamini-Hochberg correction for multiple testing. **[B]** Simplified galactose metabolism pathway. The seven metabolites that significantly changed in abundance and quantified using GC-MS are indicated in bold font.

1253

1254

1255

1256

1257

1258

1259

1260

Figure S5. **Overview of transcriptomes in each condition tested in response to high K stress in _A. thaliana_ and _S. parvula_ roots. [A]** Principal component analysis (PCA) for 0, 3, 24, and 72 hours after treatment (HAT) for _A. thaliana_ (left panel) and _S. parvula_ roots and shoots. **[B]** Overall transcript level correlation between conditions. Correlation plots were generated using PerformanceAnalytics library in R 4.0.2. Pearson correlation coefficient is given within plots for each comparison.

1261

1262

Figure S6. **The majority of differentially expressed genes (DEGs) show species specific responses followed by tissue and response time specificity.** Differentially expressed genes (DEGs) between *A. thaliana* and *S. parvula* for **[A]** roots and **[B]** shoots. Enriched functions are highlighted for diametric and shared responses between differently regulated ortholog pairs (OP). DEGs at each time point were called using DESeq2 compared to 0 h with a p-adj value based on Benjamini-Hochberg correction for multiple testing set to <0.01. The shared genes were plotted using UpsetR in R 4.0.2.

1263

Figure S7. Genes coding for calcium signaling and aquaporins are differently regulated during high K⁺ stress. Genes associated with **[A]** calcium signaling pathway and **[B]** aquaporins during high K⁺ in root and shoot of *A. thaliana* and *S. parvula*. The significantly changed genes at least in one condition are presented in the heatmap. DEGs at each time point were called using DESeq2 compared to 0 h with a p-adj value based on Benjamini-Hochberg correction for multiple testing set to <0.01.

1264

Figure S8. Normalized expression of [A] *SOS1* **and [B]** *HKT1* **in roots and shoots of** *A. thaliana* **and** *S. parvula* **under 150 mM KCl treatments.** *HKT1* has undergone a tandem duplication in *S. parvula,* and *SpHKT1;2* is known to function in selective K^+ transport. *AtHKT1* and *SpHKT1;1* are selective Na transporters. * represent DEGs at each time point called using DESeq2 compared to 0 h with a p-adj value based on Benjamini-Hochberg correction for multiple testing set to <0.01.

1265

1266

Supplementary tables

Table S1. Relative abundance of metabolites for *A. thaliana* and *S. parvula* roots and shoots

sample. 25-days-old hydroponically grown seedlings were treated for 24 and 72 hours after 150

mM KCl treatment and control samples were harvested together with the treated samples.

Data are represented as the mean of at least 3 independent replicates. ≥5 plants per replicate

were used.

Table S2. Number of total reads and percentage of uniquely mapped reads to *A. thaliana*

(TAIR10) or *S. parvula* v2.2 gene models for root and shoot transcriptomes under high K^+. At- *A.*

thaliana, Sp- *S. parvula*, C- control, 3- 3 hours after treatment, 24- 24 hours after treatment, 72-

72 hours after treatment, R- root samples, S- shoot samples. 25-days-old hydroponically grown

seedlings were treated for 3, 24, and 72 h after 150 mM KCl treatment. Control samples were

harvested together with the treated samples. Data are represented as the mean of at least 3

independent replicates. ≥5 plants per replicate were used.

Table S3. List of differentially expressed genes (DEGs) in 3, 24, and 72 hours after treatment (HAT) in *A. thaliana* and *S. parvula* root and shoot. DEGs were called using DESeq2 with a *p*-adj value based on Benjamini-Hochberg correction for multiple testing set to <0.01. Data are represented as mean of 3 independent replicates ± SD given using ≥ 5 hydroponically grown plants per replicate.

Table S4. Enriched biological processes for a non-redundant set of induced and suppressed DEGs for *A. thaliana* roots and shoots sample under 150 mM KCl. DEGs were functionally annotated by a Gene Ontology (GO) enrichment test using BinGO in Cytoscape and enriched biological processes were further clustered based on shared genes using GOMCL with adj *p*-value <0.05 after false discovery rate correction with Benjamini-Hochberg correction.

Table S5. Enriched biological processes for time point-specific DEGs (3, 3&24+24, 24&72, 72, and 3&24&72) for *A. thaliana* roots sample under 150 mM KCl. DEGs were functionally annotated by a Gene Ontology (GO) enrichment test using BinGO in Cytoscape and enriched biological processes were further clustered based on shared genes using GOMCL with adj *p*-value <0.05 after false discovery rate correction with Benjamini-Hochberg correction.

Table S6. Enriched biological processes for time point-specific DEGs (3, 3&24+24, 24&72, 72, and 3&24&72) for *A. thaliana* shoots sample under 150 mM KCl. DEGs were functionally annotated by a Gene Ontology (GO) enrichment test using BinGO in Cytoscape and enriched biological processes were further clustered based on shared genes using GOMCL with adj *p*-value <0.05 after false discovery rate correction with Benjamini-Hochberg correction.

Table S7. Enriched biological processes for time point-specific (3, 3&24, 24, 24&72, 72, and 3&24&72) induced and suppressed DEGs for *A. thaliana* roots sample under 150 mM KCl. DEGs were functionally annotated by a Gene Ontology (GO) enrichment test using BinGO in Cytoscape and enriched biological processes were further clustered based on shared genes using GOMCL with adj *p*-value <0.05 after false discovery rate correction with Benjamini-Hochberg correction.

Table S8. Enriched biological processes for time point-specific (3, 3&24, 24, 24&72, 72, and 3&24&72) induced and suppressed DEGs for *A. thaliana* shoots sample under 150 mM KCl.

DEGs were functionally annotated by a Gene Ontology (GO) enrichment test using BinGO in

Cytoscape and enriched biological processes were further clustered based on shared genes

using GOMCL with adj *p*-value <0.05 after false discovery rate correction with Benjamini-

Hochberg correction.

Table S9. Enriched biological processes for diametric responses (*i.e.* genes that are induced in

one species when their orthologs are suppressed in the other species) in *A. thaliana* and *S.*

parvula roots and shoots sample under 150 mM KCl. The pattern is extracted from Fig S6. DEGs

were functionally annotated by a Gene Ontology (GO) enrichment test using BinGO in

Cytoscape and enriched biological processes were further clustered based on shared genes

using GOMCL with adj *p*-value <0.05 after false discovery rate correction with Benjamini-

Hochberg correction.

Table S10. Enriched biological processes for a non-redundant set of induced and suppressed

DEGs for *S. parvula* roots and shoots sample under 150 mM KCl. The *S. parvula* DEG orthologs

with *A. thaliana* were functionally annotated by a Gene Ontology (GO) enrichment test using

BinGO in Cytoscape and enriched biological processes were further clustered based on shared

genes using GOMCL with adj *p*-value <0.05 after false discovery rate correction with Benjamini-

Hochberg correction.

Table S11. Normalized gene expression clusters of ortholog pairs (OP) between *A. thaliana* and

S. parvula in roots and shoots sample. Fuzzy K-means clustering was used to find temporally co-

regulated clusters with a membership cutoff of >0.5. From initially identified 10 root and 11

shoot clusters, we filtered out clusters that did not show a response to K treatments in both

species and identified 5 root (RC1, RC2, RC3, RC4, and RC5) and 3 shoot (SC1, SC2, and SC3) co-

expression superclusters with distinct response trends. The orthologs from each cluster were

functionally annotated by a Gene Ontology (GO) enrichment test using BinGO in Cytoscape and

enriched biological processes were further clustered based on shared genes using GOMCL with

adj *p*-value <0.05 after false discovery rate correction with Benjamini-Hochberg correction.