1

Dissociation of β$_2$m from MHC Class I

Triggers Formation of Noncovalent, Transient Heavy Chain Dimers

6   Cindy Dirscherl[1,x], Sara Löchte[2,x], Zeynep Hein[1], Janine-Denise Kopicki[3], Antonia Regina Harders[1],
7   Noemi Linden[1], Julian Weghuber[4], Maria Garcia-Alai[5,6], Charlotte Uetrecht[3,7], Martin Zacharias[8],
8   Jacob Piehler[2,y], Peter Lanzerstorfer[6,y], and Sebastian Springer[1,y]*
9

[1]Department of Life Sciences and Chemistry, Jacobs University Bremen, Germany;

[2]Department of Biology and Center of Cellular Nanoanalytics, University of Osnabrück, 49076
Osnabrück, Germany;

[3]Heinrich Pette Institute, Leibniz Institute for Experimental Virology, Hamburg, Germany;

[4]University of Applied Sciences Upper Austria, 4600 Wels, Austria;

[5]European Molecular Biology Laboratory, Hamburg Outstation, Hamburg, Germany;

[6]Centre for Structural Systems Biology, Hamburg, Germany;

[7]European XFEL, Schenefeld, Germany;

[8]Physics Department, Technical University of Munich, Garching, Germany.

[x]Equal contribution.

[y]Equal contribution.

*   To whom correspondence should be addressed:

25   Department of Life Sciences and Chemistry

26   Jacobs University Bremen

27   28759 Bremen

28   s.springer@jacobs-university.de

29   +49 421 2003243

Abstract

Previously, we demonstrated that major histocompatibility complex class I molecules (MHC-I) that

lose the antigenic peptide and the light chain beta-2 microglobulin (β_2m) to become free heavy chains

(FHCs) associate to form dimers or higher-order oligomers in the plasma membrane (Dirscherl et al.,

2018). Here, we investigate this homotypic interaction of MHC-I FHCs by combining a micropattern

assay with fluorescence recovery after photobleaching (FRAP) and with single molecule co-tracking

to elucidate their molecular structure, abundance, and dynamics. We find that MHC-I FHC complexes

are dimeric, transient, non-covalent, and mediated by the α_3 domain. FHC interaction correlates with

a decrease in the diffusion coefficient and an increase in the number of immobile molecules at the cell

surface. Molecular docking and dynamics simulations suggest that in the complexes, the α_3 domain of

one FHC binds to another FHC in a manner similar to the β_2m light chain. We propose distinct

functions of the MHC-I FHC dimers.

Significance Statement

MHC class I molecules are cell surface transmembrane proteins. In addition to binding receptors on other cells (*in trans*), such as the T cell receptor or inhibitory receptors of Natural Killer cells, they also bind to proteins on the same cell (*in cis*), including other class I molecules. The functions of such *in cis* associations are not well understood, and investigation is difficult because class I molecules exist on the cell membrane in three different states, depending on whether they are associated with their light chain and with peptide. Using three independent approaches, we report here the binding of class I free heavy chains (but not the other states) to each other *in cis* to form non-covalent dimers at the cell surface. The roles of these dimers can now be investigated; we propose that they function in signaling, endocytic sorting, or both.

1   **Introduction**

Major histocompatibility class I molecules (MHC-I)[*] are central to the immune response against

infections and malignancies by presenting antigenic peptides to T cells (Comber and Philip, 2014;

Kaufman, 2018; Townsend and Bodmer, 1989). MHC-I heterotrimers consist of the polymorphic

transmembrane heavy chain (HC), the non-polymorphic soluble light chain beta-2 microglobulin

(β_2m), and the peptide (Townsend et al., 1990, 1989). In addition to this trimer, two more states of

MHC-I occur in cells, the 'empty' HC/β_2m heterodimer that lacks peptide (Ljunggren et al., 1990;

Sebastián Montealegre et al., 2015), and the monomeric 'free' heavy chain (FHC) (Edidin et al., 1997;

Geng et al., 2018). Since the binding of peptide and β_2m is cooperative (Elliott et al., 1991; Gakamsky

et al., 1996), HC/β_2m heterodimers are conformationally unstable, and loss of peptide leads to the rapid

formation of FHCs (**Figure 1A**) and to the subsequent endocytic removal of FHCs by a sorting

mechanism that is not understood (Sebastián Montealegre et al., 2015). For FHCs present at the cell

surface, important regulatory functions mediated by homo- and heteromeric interactions in *cis* and

trans have been proposed (Arosa et al., 2007), which suggest a defined spatiotemporal organization

and dynamics of MHC-I and in particular FHC in the plasma membrane. Indeed, clustering and

dimerization of MHC-I under different conditions have been described (Allen et al., 1999; Armony et

al., 2021; Baia et al., 2016; Blumenthal et al., 2016; Bodnar et al., 2003; Capps et al., 1993; Chakrabarti

et al., 1992; Fassett et al., 2001; Ferez et al., 2014; Lu et al., 2012; Makhadiyeva et al., 2012; Matko

---

[*]**Abbreviations: AUC**, area under the curve; **β_2m**, beta-2 microglobulin; **BFA**, brefeldin A; **BSA**, buried surface area; **D^b**, the murine MHC-I molecule H-2D^b; **ER**, endoplasmic reticulum; **FHC**, free heavy chain; **FRAP**, fluorescence recovery after photobleaching; **GFP**, green fluorescent protein; **HA**, hemagglutinin (tag); **HC**, heavy chain; **K^b**, the murine MHC-I molecule H-2K^b; **MD**, molecular dynamics; **MHC-I**, major histocompatibility complex class I; **MS**, mass spectrometry; **RMSD**, root mean square deviation; **SL8**, ligand peptide cognate to K^b (single-letter amino acid code SIINFEKL); **SMCT**, single-molecule co-tracking; **SMT**, single-molecule tracking; **TAP**, transporter associated with antigen processing; **TIRF**, total internal reflection fluorescence; **TMD**, transmembrane domain.

et al., 1994; Triantafilou et al., 2000), the basic molecular principles of which, however, have remained

obscure.

Recently, we have achieved direct detection of homomeric FHC interactions in the intact plasma

membrane by means of a live-cell two-hybrid micropattern assay (**Figure 1B, C**): micrometer-sized

patterns of anti-hemagglutinin tag (HA) monoclonal antibody are printed onto glass coverslips

(Dirscherl et al., 2018; Schwarzenbacher et al., 2008; Sevcsik et al., 2015). Onto these micropatterns,

cells are seeded that express two different MHC-I HC constructs. One construct has an N-terminal

(extracellular) HA tag and thus, while diffusing laterally in the plasma membrane, is captured into the

printed antibody pattern. The other has no HA tag but a C-terminal (intracellular) green fluorescent

protein (GFP) fusion domain (**Figure 1B**). Interaction of GFP-tagged HC with micropatterned HC is

detected by an increased GFP fluorescence in the micropattern (**Figure 1C**). With TAP2 (transporter

associated with antigen processing)-deficient fibroblasts (which cannot transport peptide into the

endoplasmic reticulum (ER)), empty HC/β_2m heterodimers are present at the cell surface. By adding

peptide, shifting cells to 25 °C, or incubating at 37 °C, we therefore can accumulate trimers, HC/β_2m

heterodimers, or FHCs, respectively, at the cell surface (**Figure 1B** and described below). These

experiments revealed that formation of homomeric MHC-I only takes place in the absence of β_2m, *i.e.*,

only between FHCs.

We now demonstrate the molecular principles that govern such homomeric MHC-I FHC association

in the plasma membrane. Cell micropatterning in conjunction with fluorescence recovery after

photobleaching (FRAP) was used to probe dynamics, stability, and prominence of FHC complexes.

We furthermore directly demonstrate FHC association in the plasma membrane under physiological

conditions by using real-time single-molecule tracking (SMT) and co-tracking. We find that FHC

transiently associate into non-covalent dimers with lifetimes in the sub-second range and that they

diffuse with reduced mobility at the cell surface. Based on our findings that the HC/HC complexes

contain no β2m and that the α3 domain of the HC is sufficient for dimerization *in vitro* and in cells, we

propose a molecular model structure of MHC-I HC/HC dimers by *in silico* docking and molecular

dynamics (MD) simulation. Our findings clearly differentiate the cell surface dynamics and properties

of empty FHCs from the peptide-loaded trimers of MHC-I, pointing to the formation of structurally

well-defined HC/HC homodimers that may be responsible for distinct endosomal trafficking and other

biological functions previously ascribed to FHCs.

Results

A micropattern assay reveals non-covalent association of MHC class I free heavy chains

In TAP-deficient STF1 cells, which are unable to load MHC-I with peptides, incubation at 25 °C

accumulates murine HC/β2m heterodimers at the plasma membrane, since at that temperature,

dissociation of β2m and subsequent endocytosis are inhibited (Day et al., 1995; Ljunggren et al., 1990;

Sebastián Montealegre et al., 2015). When the temperature is shifted to 37 °C, β2m dissociates to yield

FHCs, and lateral *in cis* interactions between HA-K^b and K^b-GFP (both hybrids of the HC of the murine

MHC-I molecule H-2K^b) become visible in the micropattern two-hybrid assay (**Figure 1C**). When

cognate peptide is added to the cells, HC/β2m/peptide trimers are stable and do not associate (**Figure**

2B) (Dirscherl et al., 2018).

Up to six different MHC-I allotypes are present at the cell surface of human and murine cells. To

date, though, an interaction of different MHC-I allotypes in the same plasma membrane has not been

shown. We therefore tested for such heterotypic interactions between K^b and H-2D^b (D^b). Just as for

K^b/K^b, also K^b and D^b FHCs (at 37 °C) interacted with each other, whereas β2m-bound heterodimers

at 25 °C did not (**Figure 1D**). Hence, our micropattern assay confirmed that formation of heterotypic

FHC interactions is possible.

We next asked which molecular characteristics are required for HC/HC interactions, and we first tested whether the HCs are held together by cytosolic disulfide bonds as previously suggested for certain allotypes under oxidizing conditions (Baía et al., 2016; Capps et al., 1993; Makhadiyeva et al., 2012). We replaced the single cysteine in the cytosolic tail of K^b-GFP (residue 332) with a serine and tested for association between this mutant and wild type HA-K^b (**Figure 1E**). K^b(C332S)-GFP and wild type HA-K^b interacted in over 90% of cells, demonstrating that cysteine 332 is not required for HC/HC interactions.

To test more generally for any disulfide bonding in MHC-I interactions, we immunoprecipitated HA-K^b molecules from STF1 cell lysate with anti-HA antibody. By non-reducing gel electrophoresis, covalent homodimers of K^b were not observed, while the well-described disulfide-linked homodimers of the human MHC-I allotype HLA-B*27:05 (Dangoria et al., 2002) were readily detected (**Figure 1F**). We conclude that the K^b FHCs are non-covalently associated and that they do not undergo intramolecular disulfide bonding under the conditions of our assay.

Since the cysteines in the cytosolic tail of K^b are not required for HC/HC interaction, we hypothesized that FHCs non-covalently associate via their extracellular domains, and that the loss of β_2m was a prerequisite for HC/HC interaction. As anticipated, a disulfide-stabilized variant of K^b–GFP (Y84C/A139C), in which β_2m dissociation is dramatically decreased (Hein et al., 2014), did not interact with HA–K^b (**Figure 1G**). In agreement with our earlier finding that covalent attachment of β_2m to the HC prevents HC/HC association (Dirscherl et al., 2018), this result demonstrates that HC/β_2m heterodimers do not interact with other HCs.

Free heavy chains associate on TAP-proficient cells

So far, we used TAP2-deficient STF1 cells to obtain homogeneous populations of free heavy chains at the cell surface. Since MHC-I HC/β_2m heterodimers and FHCs both exist at the cell surface of wild

type cells (Day et al., 1995; Ljunggren et al., 1990; Ortiz-Navarrete and Hammerling, 1991, p.), we

next tested whether FHCs also associate in cells with wild type TAP function (**Figure 2A**). Following

the 25 °C to 37 °C temperature shift that triggers FHC formation, we quantified HC/HC interaction in

STF1/TAP2 over time by GFP fluorescence contrast analysis between pattern elements and

interspaces. Interactions increased in TAP2-positive and in TAP2-deficient cells with the same

dynamics and reached a maximum after ca. 60 min of incubation at 37 °C (**Figure 2B, C**), with the

kinetics likely governed by the dissociation of β_2m from the HC/β_2m heterodimers (Sebastián

Montealegre et al., 2015). As expected, HC/β_2m/peptide trimers, stabilized by cognate SIINFEKL

peptide, did not show any interaction (**Figure 2B**). This experiment confirms that HC/HC interactions

take place at the surface of TAP2-proficient cells.

FHC association slows down cell surface diffusion of MHC-I

Since dimers and oligomers of FHCs have more transmembrane domains (TMDs) than single

HC/β_2m/peptide complexes, we hypothesized that they should diffuse more slowly in the plasma

membrane (Lippincott-Schwartz et al., 2018). We therefore carried out total internal reflection

fluorescence (TIRF) fluorescence recovery after photobleaching (FRAP) experiments on STF1 cells

with HA-K^b and K^b-GFP on surfaces with or without micropatterns (**Figure 2D**) and quantified the

recovery dynamics (**Figure 2E**). Diffusion constants were significantly higher for HC/β_2m/peptide

complexes (+ peptide) than for FHCs (– peptide; **Figure 2F**), suggesting that FHCs form complexes

with two or more TMDs. Since this effect occurred both in cells seeded on the pattern elements (on

pattern) and in cells seeded off the micropatterns (off pattern), we conclude that the micropatterns are

not required for HC/HC interactions, i.e., that the interactions are not an artefact of the two-hybrid

micropattern assay. This is also supported by the co-immunoprecipitation of MHC-I molecules in the

absence of micropatterns (Dirscherl et al., 2018; Triantafilou et al., 2000). Diffusion constants of

HC/β2m/peptide trimers on pattern and off pattern were not significantly different, which demonstrates

that the antibody micropatterns themselves do not impede the diffusion of plasma membrane proteins.

In the FRAP experiments, a portion of K^b-GFP appeared immobile on a timescale of seconds as

evidenced by the incomplete fluorescence recovery (**Fig. 2E**). This immobile fraction is significantly

greater for FHCs than for HC/β2m/peptide trimers (**Fig. 2G**). This observation is readily explained for

K^b-GFP bound to HA-K^b molecules immobilized within micropatterns, but it is remarkable for the

FHCs on cells outside of the micropattern (gray column in **Fig. 2G**). This observation may be ascribed

either to the formation of large oligomers, or – perhaps more likely – to an association of FHCs with

immobile components of the plasma membrane, e.g., related to endocytosis. From the FRAP curves,

the exchange rate of the freely diffusing pool of K^b-GFP into and out of the bleached ROIs is obtained

through a bi-exponential fit as the slow recovery rate (k_{slow}) (**Figure 2H**, solid bars); the fast recovery

rate k_{fast} represents free diffusion (Sprague and Mcnally, 2005)(see the Materials and Methods). On

the pattern elements, the k_{slow} of FHCs was much smaller than in cells outside the patterns, suggesting

a half-time of dissociation from the pattern-bound immobile associations of about 140 seconds.

Likewise, the fluorescence signal of pattern elements in the immediate vicinity of the bleached region

remained unaltered, indicating that no detectable exchange of K^b-GFP between the enriched HA-K^b

regions occurs on the second timescale (**Figure 2S3**). This very slow exchange of K^b-GFP molecules

associated on pattern elements suggests either multiple association and dissociation events in a small

radius due to densely immobilized binding partners, or a very stable association of the immobile FHCs

(see the discussion). In conclusion, K^b FHCs show reduced diffusion rates on the cell surface, which

is most easily explained by homotypic association.

Transient FHC dimerization directly observed at single molecule level

We therefore turned to directly visualizing FHC diffusion and interaction in the plasma membrane

under physiological conditions by single molecule tracking (SMT) and co-tracking (SMCT) (Moraga

et al., 2015; Sevcsik et al., 2015; Wilmes et al., 2020a). STF-1 cells transiently expressing K^b with its

N-terminus fused to monomeric GFP (GFP-K^b) were imaged by TIRF microscopy. The GFP tag was

labeled with photostable fluorescent dyes by using equal concentrations of anti-GFP nanobodies

conjugated to either ATTO Rho11 (^{Rho11}NB) or ATTO 643 (^{ATTO643}NB), ensuring selective imaging

of K^b in the plasma membrane and tracking with high fidelity (**Figure 3A**). Imaging was performed at

37 °C to induce formation of FHC in the absence of the peptide, while replenishment of GFP-K^b/β_2m

dimers from the ER was inhibited by Brefeldin A (BFA).

After labeling with ^{Rho11}NB and ^{ATTO643}NB, individual K^b subunits were observed at densities of <1

molecule/µm² diffusing randomly in the plasma membrane (**Movie S1, Figure 3S1**). Co-tracking

analysis revealed homomeric interaction of K^b FHCs in the absence of peptide, whereas these events

were very rare for peptide-loaded K^b (**Figure 3B-D**). As a positive control for the formation of

homodimers, a crosslinker based on the tandem anti-GFP nanobody LaG16 (Fridy et al., 2014), which

recognizes a different epitope than the labeled nanobodies, was added to the medium. While

dimerizing of K^b via the GFP tag resulted in a nominal fraction of ~38% (median) co-locomoting

molecules, only ~3% of the FHCs were found associated, indicating weak interaction of FHCs at these

low cell surface expression levels of K^b (~1 molecule/µm² in total, corresponding to <5000

molecules/cell).

From the trajectory analysis, we determined the diffusion coefficients of individual MHC-I

molecules by mean square displacement analysis (**Figure 3E, Figure 3S2**). In the absence of peptide,

diffusion of FHCs was significantly slower than in the presence of the peptide. This decrease in single

molecule diffusion coefficients supports interaction of FHCs in the plasma membrane. A similar

decrease in the diffusion coefficient was observed upon dimerization of peptide-loaded K^b with the tandem nanobody, suggesting that FHCs associate into dimers. Indeed, largely identical diffusion coefficients were found for the fraction of molecules identified as dimers by SMCT (**Figure 3S3, Table 1**). Strikingly, the diffusion constants obtained by SMT for peptide-loaded K^b and FHC from SMT are consistent with those obtained by FRAP experiments under the same conditions (**Figure 2E, Table 1**), highlighting that similar phenomena are being probed by these complementary techniques. Tracking analysis revealed a slightly elevated immobile fraction of ~20% for K^b in the absence of the peptide compared to ~17% for peptide-loaded K^b (**Figure 3F**). Again, a similar effect was observed for artificially dimerized, peptide-loaded K^b, corroborating the dimeric stoichiometry of associated FHCs. Importantly, the "immobile fraction" in SMT refers to much shorter time and length scales as compared to FRAP (cf. methods section), and therefore, the absolute numbers are not comparable. Within the immobile fraction, a higher level of FHC was found associated, as compared to the mobile fraction (**Figure 3G**).

SMCT analysis clearly revealed dissociation of FHC homomers, confirming its transient nature (**Figure 3B, Movie S2**). We estimated the lifetime of FHC complexes from co-trajectory length histograms. Fitting of an exponential decay revealed a significantly shorter lifetime for FHC co-trajectories compared to the average co-tracking lifetime determined for stably NB-crosslinked K^b, which is limited by co-tracking fidelity and photobleaching (**Figure 3H**). Taken together, SMT directly confirmed a transient homotypic HC/HC interaction in the plasma membrane that leads to reduced diffusion velocity and immobilization, in line with the FRAP experiments. Since the degree of association was low, and higher-order oligomerization was not observed, we propose that HC/HC interactions are weak and have dimeric stoichiometry.

The α₃ domain of K^b forms dimers and is sufficient for FHC association

We next explored the molecular mechanism of HC/HC interaction. Since cytosolic cysteines are

not involved (**Figure 1E, F**) and dissociation of β_2m is necessary (**Figure 1G**), we hypothesized that

HC/HC interactions involve the extracellular portion of the HCs. To determine more precisely which

domains are involved, we used STF1 cells in the micropattern assay that expressed constructs of K^b

that lacked the α_1/α_2 domain. Remarkably, α_3-GFP showed excellent copatterning with HA-K^b-RFP

(fused to red fluorescent protein; **Figure 4A**). Likewise, HA-α_3 and α_3-GFP showed a significant

copatterning, demonstrating that the α_3 domain is sufficient (**Figure 4B**). Indeed, the isolated, soluble

α_3 domain of K^b, when refolded *in vitro*, showed significant formation of homodimers in size

exclusion chromatography (SEC; **Figure 4C**), while no prominent higher order complexes were

observed. These α_3 homodimers were not linked by disulfide bonds as shown by nonreducing gel

electrophoresis (**Figure 4D**), just like the interaction of full-length FHC in the plasma membrane

(**Figure 1F**). Given the 1:1 peak ratio observed at 10 µM protein concentration, a binding affinity of

10 to 20 µM can be estimated for this interaction. The non-covalent α_3 homodimers were also

detectable by native mass spectrometry (MS) in a concentration-dependent manner, and they easily

separated into the monomers when the collision cell voltage was increased in an MS/MS experiment,

suggesting a low-affinity interaction (**Figure 4E-G, 4S1**, **Table 2**). These observations suggest that

FHC dimerization in the plasma membrane is at least partly based on the intrinsic affinity between the

α_3 domains.

A molecular model of the K^b HC dimer

To image one possible arrangement of the K^b heavy chains in a dimer, we used molecular docking.

A starting geometry was obtained by placing the α_3 domain of one K^b HC in the same position as the

β_2m in the original K^b HC/β_2m heterodimer and adjusting the α_1/α_2 superdomain to avoid sterical

overlap. The structure was then energy-minimized and further refined by Molecular Dynamics (MD) simulations (**Figure 5A**). During MD simulations of 400 ns, no signs of dissociation were observed, and a stable root-mean-square deviation (RMSD) from the start structure was reached (**Figure 5S1**). In the model, the α_3 domain of one FHC substitutes for β_2m, binding both the α_3 domain and the α_1/α_2 superdomain of another FHC with a buried interface area (BSA) of 2480 Å^2, which is comparable to the BSA of 2740 Å^2 between β_2m and the HC in heterodimers (**Figure 5B**, bottom), and to other stable protein complexes (Bahadur and Zacharias, 2008). Similar results were obtained with other arrangements of the two heavy chains (not shown).

Discussion

Homo- and heterotypic interactions of MHC-I FHC in *cis* and *trans* have been proposed to play an important role in regulating adaptive and innate immune responses (Arosa et al., 2007) (see below), but the molecular principles governing FHC interactions have remained unclear. By combining live cell interaction and diffusion analysis using cell micropatterning and FRAP as well as SMT and SMCT, we have here shown that, after losing β_2m, murine H-2K^b MHC-I molecules at the cell surface interact in a homotypic and heterotypic manner to form dimers, which are transient with a stability on the second timescale. For these interactions, the α_3 domains of the FHCs are sufficient.

While we cannot entirely rule out formation of higher oligomers, our data rather point toward dimers being the prevalent species. We analyzed the diffusion properties of FHCs and HC/β_2m/peptide trimers at different cell surface expression levels by FRAP and SMT. FRAP experiments were carried out at high surface densities, which probably exceeded the typical endogenous cell surface expression levels of 10^5 copies/cell (Spack and Edidin, 1986). By contrast, SMT was performed at densities of ~1 molecule/µm², *i.e.*, <5000 molecules/cell. Despite these differences in expression levels, we found highly consistent diffusion coefficients D by both FRAP and SMT, which revealed significantly

decreased mobility upon dissociation of β_2m (**Table 1**). However, under both conditions, the decrease

in D was ~50%, which perfectly agrees to the change in D upon dimerizing intact MHC-I by the

tandem nanobody (**Figure 3E**). Likewise, we and others have previously observed very similar

decreases of 30-50% upon dimerization of cell surface receptors (Gambin et al., 2006; Ho et al., 2017;

Low-Nam et al., 2011; Moraga et al., 2015; Richter et al., 2017; Váradi et al., 2019; Wilmes et al.,

2020a, 2015a), corroborating that the HC/HC complexes are mostly dimers.

Direct detection of FHC interaction by SMCT suggested weak, non-covalent dimerization while

higher oligomers and clustering could not be detected. Estimated lifetime of the FHC/FHC dimer ($t_{1/2} =$

220 ± 150 ms, **Figure 3G, H**) were well within the measurable range, *i.e.*, substantially below the

apparent half-life obtained for the quasi-irreversibly, nanobody-crosslinked control ($t_{1/2} > 10\ s$), which

defined the limit of co-tracking fidelity. Together with the immunoprecipitation experiments (**Figure**

1F) and micropatterning of FHC lacking free cysteines (**Figure 1E**), these observations clearly

establish that the FHC molecules at the plasma membrane are not covalently linked. The short half-

live and the non-covalent monomer-dimer equilibrium observed by size exclusion chromatography at

higher micromolar concentration for the isolated α_3 domain (**Figure 4CD**), as well as the mass

spectrometry data (**Figure 4E-G**), point to a low-affinity interaction with a dimerization K_d of perhaps

10 to 20 µM. Previous systematic studies on affinity-dimerization correlation of heterodimeric

cytokine receptors (Wilmes et al., 2015) would predict efficient FHC dimerization at physiological

densities above 10 molecules/µm², which is in line with the endogenous MHC-I expression level. The

relatively low level of dimerization we observed by SMCT can be rationalized with the high

background of endogenous MHC-I that are unlabeled and invisible to us. In line with this observation,

the decrease of the diffusion coefficient in SMT was much more prominent than the dimer fraction

identified by SMCT. This interpretation is in line with the observation that more efficient interaction

was observed at the highly elevated cell surface expression levels used in micropatterning experiments.

The weak and transient nature of HC/HC dimerization seen by SMT is similar to the nature of cell surface protein-protein association measured in other systems (Lin et al., 2014). It does not conflict with the clear and stringent patterning of the GFP fusion in the micropattern/FRAP experiments. In the latter, the concentration of HA-tagged bait HCs is considerably higher due to their immobilization by the antibodies, which prevents their endocytosis, and this probably creates an affinity matrix for the GFP fusions that can retain them for many seconds due to rebinding events. This also suggests that in endosomes, whose internal volume is very small, HC/HC interaction might be potentiated due to the increase in the concentration of the monomers compared to the plasma membrane.

In line with the observation of non-covalent dimerization in the plasma membrane, we obtained a robust structural model of a self-contained K^b FHC homodimer. The atomistic model (**Figure 5**) was derived from the experimental findings that dissociation of β_2m is required, that the α_3 domains are sufficient, and that a direct α_3/α_3 interaction exists (**Figures 1E, 4A-C**). We propose that β_2m dissociation exposes a binding site on the FHC for the α_3 domain of another FHC. However, other arrangements of the two FHCs in a dimer are theoretically possible, and only experimental data will give a definitive answer.

Several findings in the literature are consistent with the formation of complexes of MHC class I HCs in the absence of β_2m and peptide (Armony et al., 2021; Bodnar et al., 2003; Chakrabarti et al., 1992; Matko et al., 1994; Triantafilou et al., 2000). Still, it is important to differentiate these HC/HC dimers from class I associations described elsewhere (partially reviewed in (Arosa et al., 2007; Campbell et al., 2012)), namely HC/β_2m/peptide trimers that are covalently dimerized via disulfide bonds in their cytosolic tails (Capps et al., 1993; Makhadiyeva et al., 2012) and that may play a role in binding the LILRB NK cell receptor (Baia et al., 2016); the non-covalent nano- and microscale clusters of HC/β_2m/peptide trimers (not detected in our system) that may stem from the fusion of exocytic vesicles with the plasma membrane and that may play a role in TCR recognition (Blumenthal et al.,

2016; Ferez et al., 2014; Fooksman et al., 2006; Lu et al., 2012); the macroscopic 'clusters' of class I molecules at the signaling interface between cells (Fassett et al., 2001) and the covalent dimers of the HLA- B*27:05 heavy chain that are linked by disulfide bonds through Cys-67 (Chen et al., 2017), though our non-covalent HC/HC dimers may be a precursor to the formation of the latter.

The unexpected discovery that FHCs non-covalently associate into defined dimers allows exciting hypotheses of their distinct functional properties. FHC dimers might be responsible for the immunomodulatory functions of cell surface heavy chains, *i.e.,* the stabilization of MHC-I trimers to assist T cell activation (Geng et al., 2018; Schell, 2002) and direct binding of FHCs to receptors on other cells (for example, FHCs of HLA-F binding to activating receptors on NK cells (Dulberger et al., 2017, 2017; Goodridge et al., 2013)). Furthermore, dimerization of FHCs might enhance endocytosis in order to remove the non-functional FHCs, which themselves cannot activate T cells and are known to be short-lived (Mahmutefendic et al., 2011; S. Montealegre et al., 2015); alternatively or additionally, such associated HCs may bind to other proteins *in cis* and promote their removal from the plasma membrane. Such endocytic removal might be achieved by altered endosomal routing, since the local density of membrane proteins in endosomes is higher than at the plasma membrane, and thus, efficient dimerization of MHC-I FHCs is expected. In such a scenario, even transient oligomerization in endosomes might prevent the return of internalized MHC-I FHCs to the cell surface (S. Montealegre et al., 2015). Taken together, the presence of non-covalent, transient FHC dimers points to exciting new aspects in the regulation of MHC-I functions with much potential for further investigation.

1   Materials and Methods

2   Key resources table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional Information
gene (*Mus (M.) musculus*)	H-2K^b	NCBI GenBank	NM_001001892.2	Used for plasmid design
gene (*M. musculus*)	H-2D^b	NCBI GenBank	NM_010380.3	Used for plasmid design
gene (*M. musculus*)	TAP2	NCBI GenBank	NM_011530.3	Used for plasmid design
strain, strain background (*Escherichia* (*E.*) *coli Rosetta*)	BL21(*DE3*) *pLysS*	Novagen	Cat. # 70956	Used for expression of recombinant α_3 domain of K^b
cell line (*homo sapiens*)	STF1	PMID:10074495	N/A	Used for all experiments
recombinant DNA reagent	pET3a (plasmid)	Novagen	Cat. # 69418-3	Used for transformation of *E. coli*
recombinant DNA reagent	puc2CL6IPwo (lentiviral vector)	PMID: 21248040	N/A	Used for transduction and transfection
recombinant DNA reagent (*M. musculus*)	puc2CL6IPwo/ E3-HA-K^b	DOI: 10.7554/eLife.34150.001	N/A	For stable transduction of STF1 cells
recombinant DNA reagent (*M. musculus*)	puc2CL6IPwo/ K^b-GFP	DOI: 10.7554/eLife.34150.001	N/A	
recombinant DNA reagent (*M. musculus*)	puc2CL6IPwo/ D^b-GFP	DOI: 10.7554/eLife.34150.001	N/A	For stable co-transduction of HA-K^b expressing STF1 cells
recombinant DNA reagent (*M. musculus*)	puc2CL6IPwo/ TAP2	This paper	N/A	
recombinant DNA reagent (*M. musculus*)	puc2CL6IPwo/K^b(Y84C/A139C)-GFP (plasmid)	DOI: 10.1242/jcs.145334	N/A	
recombinant DNA reagent (*M. musculus*)	puc2CL6IPwo/K^b(C332S)-GFP	This paper	N/A	10 µg for transfection of one 10-cm plate of confluent cells
recombinant DNA reagent (*M. musculus*)	puc2CL6IPwo/α_3 domain (of K^b)-GFP	This paper	N/A	
antibody	12CA5 (anti-hemagglutinin (HA))	PMID: 6192445. (Niman et al., 1983)[7]	N/A	Produced and purified in house from hybridoma cells 0.6 mg/mL for printing; 1:100 dilution of hybridoma supernatant for Western blotting.
antibody	polyclonal rabbit anti-HA	Abcam	Cat. # ab9110	Used as primary antibody in Western blot, according to manufacturer's recommendations
antibody	alkaline phosphatase-conjugated goat anti-rabbit serum	Biorad	Cat. #1706518	Used as primary antibody in Western blot, according to manufacturer's recommendations
peptide, recombinant protein	SIINFEKL peptide	GeneCust Ellange, Luxemburg	N/A	2 mM final concentration
commercial assay or kit	HiLoad® 16/600 Superdex® 200 pg (column for SEC)	GE Healthcare	Cat. # GE28-9893-35	Used for purification of recombinant α_3 domain of K^b
chemical compound, drug	Alexa Fluor 647 NHS ester	Thermo Fisher Scientific	Cat. # A37566	Used for labelling anti-HA antibodies for micropattern assay according to the manufacturer's protocol.
chemical compound, drug	Brefeldin A Solution (1,000X)	BioLegend	Cat. # 420601	10 µg/ml

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional Information
software, algorithm	ImageJ	National Institutes of Health	https://imagej.nih.gov/ij/	Used for image processing (cropping, rotation and adjustment of brightness and contrast levels)
software, algorithm	GraphPad Prism version 8.4.0	GraphPad Software	https://www.graphpad.com	

Cells and cell lines

TAP-deficient human STF1 fibroblasts (kindly provided by Henri de la Salle, Etablissement de

Transfusion Sanguine de Strasbourg, Strasbourg, France) were cultivated at 37 °C and 5% CO_2 in

Earle's minimum Essential Medium (MEM) with stable glutamine supplemented with 10% fetal

bovine serum (FBS), non-essential amino acids and HEPES buffer without addition of antibiotics.

Genes, vectors, and gene expression

HA-K^b and K^b-GFP constructs were described previously.(Dirscherl et al., 2018) HA-K^b carries an

influenza hemagglutinin (HA) tag at the N terminus of the full-length murine H-2K^b, whereas K^b-GFP

carries a GFP domain at the C terminus of H-2K^b. The D^b-GFP construct is analogous to the K^b-GFP

construct. The α_3-GFP construct consisted of the H-2K^b signal sequence and residues 204-369 of H-

2K^b, including transmembrane and cytosolic domains. The GFP-K^b construct used in single-molecule

imaging consists of a signal sequence and GFP fused to the N terminus of H-2K^b itself lacking a signal

sequence.

Stable cell lines were generated by lentiviral transduction as described(Hein et al., 2014), and

transient transfection was achieved by electroporation(Garstka et al., 2007) or by calcium-phosphate-

precipitation(Graham and van der Eb, 1973) as described.

Micropattern assay

Photolithography. Silicon master molds were prepared by semiconductor photolithography as

described previously (Dirscherl et al., 2017).

PDMS stamps and Antibody Patterns. PDMS stamps were generated from basic elastomer and

curing agent (Sylgard 184 Silicone Elastomer Kit) as described previously (Dirscherl et al., 2017).

Patterning cell surface proteins. Coverslips with antibody pattern were placed into sterile 6-well

plates. Cells were immediately seeded as indicated at a concentration of ca. 50 000 cells per well.

Usually, cells were incubated for 4-6 hours at 37 °C for adhesion and then shifted to 25 °C to

accumulate MHC-I molecules at the cell surface. Samples were then kept at 25 °C to increase cell

surface heterodimer levels or shifted back to 37 °C for 3-4 hours to induce FHCs by dissociation of

β_2m.

Dyes. Purified antibodies were labeled with Alexa Fluor 647 NHS ester (Thermo Fisher Scientific,

Darmstadt, Germany) according to the manufacturer's protocol.

Peptides. The K^b-specific peptide SL8 (SIINFEKL in the single-letter amino acid code) was

synthesized by GeneCust (Ellange, Luxemburg) and emc microcollections (Tübingen, Germany) and

purified by HPLC (90% purity).) Peptides were added to the cells at a final concentration of 2 µM for

15-30 min at 37 °C to induce peptide binding (Dirscherl et al., 2018).

Washing and fixation. Cells were washed with phosphate buffered saline (PBS), 10 mM phosphate

pH 7.5, 150 mM NaCl), fixed with 3% paraformaldehyde (PFA), and observed by confocal laser

scanning microscopy (cLSM).

Microscopy. We used a confocal laser scanning microscope (LSM 510 Meta, Carl Zeiss Jena

GmbH, Germany) equipped with argon and helium-neon lasers at 488, 543, and 633 nm. Images were

recorded with a 63× Plan Apochromat oil objective (numerical aperture 1.4) at a resolution of 1596 ×

1596 pixels. Data acquisition was performed with the LSM 510 META software, release 3.2 (Carl

Zeiss Jena). During image acquisition, patterns and cells were imaged in the same focal plane at a

pinhole of 1 Airy unit. Image analysis and processing were performed using ImageJ (National

Institutes of Health, Bethesda, USA). Image processing comprised cropping, rotation and adjustment

of brightness and contrast levels. Experiments in Figure 1 were repeated at least three times each.

Recombinant α3 domain of H-2K^b

The α3 domain of H-2K^b (residues 205-295) was cloned into pET3a (preceded by the residues

MAIQR and followed by DRDM) and expressed in *E. coli* BL21(*DE3*) *pLysS*, refolded *in vitro* as

described, and isolated by size exclusion chromatography (SEC) on a Cytiva Hiload Superdex 200

16/600 column (Anjanappa et al., 2020). Molecular weights of the peaks were determined by

comparison to SEC protein standards (Cytiva), namely bovine thyroglobulin (670 kDa), bovine

gamma globulin (158 kDa), chicken ovalbumin (44 kDa), horse myoglobin (17 kDa), and vitamin B12

(1.35 kDa). The D5 fraction, corresponding to the elution peak at approximately 20-30 kDa, was boiled

with or without DTT (0.6 M final concentration) in sample buffer (LSB) (350 mM Tris-Cl pH 6.8,

10.28% sodium dodecyl sulfate (SDS), 36% glycerol 0.012% bromophenol blue). Inclusion body

extract boiled without DTT (non-reducing) served as positive control for the formation of covalent

oligomers. Protein quality control after refolding and SEC was performed by monitoring nDSF runs

acquired with a Nanotemper Prometheus NT.48 fluorimeter (Nanotemper, Munich) controlled by

PR.ThermControl (version 2.1.2).

Precipitation of surface class I

TAP2-deficient STF1 cells expressing either CE3-HA-K^b or CE3-HA-B*27:05 were kept overnight

at 25 °C and then pretreated with tris(2-carboxyethyl)phosphine (TCEP; 1 mM, 10 min), labeled with

nm of Bio-MPAA-K3 (5 min) at room temperature (RT) (Reinhardt et al., 2014). Lysis was

performed in native lysis buffer (50 mM Tris Cl (pH7.4), 150 mM NaCl, 5 mM EDTA, and 1% Triton

X100) for 1 hour at 4 °C. Biotinylated surface proteins were then isolated with neutravidin-coated

agarose beads (Thermo Fisher Scientific, Darmstadt Germany). The isolates were boiled at 95 °C for

7 min in the presence (reducing) or absence (non-reducing) of 10 mM dithiothreitol (DTT) in sample

buffer as described above. Samples were separated by SDS-PAGE and transferred onto polyvinylidene

fluoride (PVDF) membranes. MHC molecules were visualized on the membranes with polyclonal

rabbit anti-HA antibody as primary antibody (ab9110, Abcam, Cambridge, United Kingdom) and

alkaline phosphatase-conjugated anti-rabbit serum from goat as secondary antibody (1706518, Biorad,

Munich, Germany). The signals were visualized by treating the blot with BCIP/NBT substrate (B1911,

Sigma Aldrich, St. Louis, Missouri, United States).

Flow cytometry

For verification of cell surface levels of MHC-I, flow cytometry was performed with anti-HLA

class I antibody W6/32 (Barnstable, 1978) and anti-HA antibody (12CA5, described in key resources

table). Antibody-antigen complexes were labeled with goat secondary antibody against mouse IgG

conjugated with allophycocyanin (APC) (115-135-164, Dianova, Hamburg, Germany). Fluorescent

signal was recorded by a CyFlow1Space flow cytometer (Sysmex- Partec, Norderstedt, Germany) and

analyzed by Flowjo, LLC software.

FRAP

Microcontact printing and antibody patterning for total internal reflection fluorescence

(TIRF) microscopy was performed as described previously.(Lanzerstorfer et al., 2020) In short, a

field of a large-area PFPE elastomeric stamp (1 µM grid size), obtained by the EV-Group (St. Florian

am Inn, Upper Austria, Austria), was cut out, and washed by flushing with ethanol (100%) and distilled

water. After drying with nitrogen, the stamp was incubated in 50 mL bovine serum albumin (BSA)

solution (1 mg/mL) for 30 min. This step was followed by washing the stamp again with PBS and

distilled water. After drying with nitrogen, the stamp was placed with homogeneous pressure onto the

clean epoxy-coated glass bottom of a 96-well plate and incubated overnight at 4 °C. The next day, the

stamp was stripped from the glass with forceps, and the glass bottom was bonded to a 96-well plastic

casting with adhesive tape (3M) and closed with an appropriate lid. For the live cell experiments, a

reaction chamber was incubated with 100 µL streptavidin solution (50 µg/mL) and incubated for

30 min at room temperature. After washing two times with PBS, 100 µL biotinylated antibody solution

(10 µg/mL) was added for 30 min at room temperature. Finally, the incubation wells were washed

twice with PBS, and cells were seeded at defined cell density for the live cell microscopy analysis.

The cells were allowed to attach to the surface for at least 3-4 h prior to imaging to ensure a

homogeneous cell membrane/substrate interface, which is a prerequisite for quantitative TIRF

microscopy. For negative control to test the adhesion of the antibodies, anti-HA antibody (Abcam,

ab26228) was labeled with a Zenon Alexa Fluor 488 IgG Labeling Kit (Thermofisher, Z25102),

printed, bleached, and fluorescence recovery was quantified as described below (**Figure 2S1**). In a

second control experiment, binding and dissociation of a construct with both tags (HA-K^b-GFP) from

the antibody micropattern was tested, which only accounted for less than 20% of the mobility of the

K^b fraction (**Figure 2S2**). Thus, in our experiments, k_{slow} was determined by the binding events

between K^b-GFP and other K^b molecules.

Live-cell TIRF microscopy. The detection system was set up on an epi-fluorescence microscope

(Nikon Eclipse Ti2). A multi-laser engine (Toptica Photonics, Munich, Germany) was used for

selective fluorescence excitation of GFP at 488 nm and RFP at 568 nm. The samples were illuminated

in total internal reflection (TIR) configuration (Nikon Ti-LAPP) using a 60x oil immersion objective

(NA = 1.49, APON 60XO TIRF). After appropriate filtering with standard filter sets, the fluorescence

was imaged onto a sCMOS camera (Zyla 4.2, Andor, Northern Ireland). The samples were mounted

on an x-y-stage (CMR-STG-MHIX2-motorized table, Märzhäuser, Germany), and scanning of the

larger areas was supported by a laser-guided automated Perfect Focus System (Nikon PFS).

TIR-FRAP experiments and calculation of diffusion coefficients. FRAP experiments were

carried out on an epi-fluorescence microscope as described above. Single patterns (or equivalent ROIs

with 1 µM in diameter in unpatterned cells) were photobleached (Andor FRAPPA) with a high-

intensity laser pulse (488 nm) applied for 500 ms. Recovery images were recorded at indicated time

intervals. Normalization of data was done by pre-bleach images, and first data analysis was carried out

using NIS Elements software package (Nikon). Further data processing was done in Graphpad Prism.

Resulting FRAP curves were plotted based on the standard error of the mean and fitted using a bi-

exponential equation. Kinetic FRAP parameters were directly obtained from curve fitting with the

following model assumptions: Since unhindered MHC-I diffusion (e.g., off pattern and/or after peptide

treatment; Fig. 2E) was found to be much faster than the recovery within our pattern elements, we

applied a diffusion-uncoupled recovery scheme (Sprague and Mcnally, 2005). Here, after the

photobleaching step, fluorescent molecules rapidly diffuse throughout the bleached micropattern, and

only bound bleached molecules remain inside the spot. The bleached molecules then gradually

dissociate from their binding sites. Unbleached molecules can then replace the bleached molecules at

the binding sites as they become vacant. The diffusion-uncoupled FRAP recovery curve consists of

two separable components: the early recovery due to diffusion and the slower recovery due to exchange

at binding sites, and can be fitted using a diffusion-uncoupled two-component fit:

$$Y = Y_{t=0} + A_1(1 - e^{-kfast*x}) + A_2(1 - e^{-kslow*x}),$$

where A_1 is the amplitude of the fast-diffusing population, A_2 the amplitude of the slow diffusing

population (binding reaction), and k_{fast} and k_{slow} are the rate constants of A_1 and A_2, respectively.

Diffusion coefficients were obtained using the initial image recordings and the simFRAP plugin for

ImageJ (Blumenthal *et al.*, 2015).

Temperature-induced K^b associations. Temperature-dependent experiments were carried out on

an epi-fluorescence microscope as described above further equipped with a cage incubator (Okolab,

Shanghai, China). Cells were grown at 25 °C overnight on antibody-patterned surfaces and treated

with SIINFEKL peptide as indicated. For induction of K^b FHC association, cells were mounted on pre-

warmed microscopy stage, and imaging of the GFP signal was started when the medium reached

37 °C.

Fluorescence contrast quantitation. Contrast analysis was performed as described previously

(Lanzerstorfer et al., 2014) In short, initial imaging recording was supported by the Nikon NIS

Elements software. Images were exported as TIFF frames and fluorescence contrast analysis was

performed with the Spotty framework (Borgmann et al., 2012). The fluorescence contrast <c> was

calculated as

14
$$\langle c \rangle = \frac{(F^+ - F^-)}{\left(F^+ - F_{bg}\right)}$$

, where F$^+$ is the intensity of the inner pixels of the pattern, F$^-$ the intensity of the surrounding pixels

of the micropattern, and F$_{bg}$ the intensity of the global background.

Single molecule microscopy

Imaging. Cells were transferred 48 h post transfection onto glass coverslips coated with a poly-L-

lysine-graft-(polyethylene glycol) copolymer functionalized with RGD tripeptide to minimize non-

specific binding of fluorescent nanobodies (You et al., 2014). Cells imaged in presence of SL8 peptide

(sequence SIINFEKL) were pre-incubated with 1 µM of SL8 peptide 12 h before imaging. Single-

molecule imaging experiments were conducted by total internal reflection fluorescence (TIRF)

microscopy with an inverted microscope (Olympus IX83) equipped with a motorized 4-Line TIR

illumination condenser (Olympus) and a back-illuminated electron multiplying (EM) CCD camera

(iXon Ultra 897, Andor Technology). A 100× magnification objective with a numerical aperture of

1.45 (UPLAPO 100× HR, NA 1.5, Olympus) together with a 1.6 × magnification changer was used

for TIR illumination of the sample. Imaging was conducted with or without 1 µM of fresh SL8 peptide.

The sample was preincubated with 10 µg/ml of Brefeldin A (BFA) for 15 min in order to inhibit protein

transport of GFP-K^b/β_2m heterodimers to the plasma membrane. All experiments were carried out at

37 °C in medium without phenol red supplemented with an oxygen scavenger and a redox-active

photoprotectant to minimize photobleaching (Vogelsang et al., 2008) and penicillin and streptomycin

(PAA). For cell surface labeling of GFP-K^b, a 1:1 mixture of anti-GFP nanobodies (2 nM each) site-

specifically conjugated with ATTO 643 and ATTO Rho11 (Wilmes et al., 2020b), respectively, were

added to the medium, thus ensuring >90 % binding given the 0.3 nM binding affinity (Kirchhofer et

al., 2010). After incubation for at least 5 min, image acquisition was started with the labeled

nanobodies kept in the bulk solution during the whole experiment in order to ensure high equilibrium

binding. Dimerization of the positive control was induced by applying 0.3 nM of a tandem nanobody

crosslinker (LaG16V2) binding to an orthogonal epitope (Fridy et al., 2014). For single molecule co-

localization and co-tracking experiments, orange (ATTO Rho11) and red (ATTO 643) emitting

fluorophores were simultaneously excited by illumination with a 561 nm laser (MPB

Communications) and a 642 nm laser (MPB Communications). Fluorescence was detected with a

spectral image splitter (QuadView QV2, Photometrics) with a dichroic beam splitter (Chroma)

combined with the bandpass filter 600/37 (BrightLine HC) for detection of ATTO Rho11 and 685/40

(Brightline HC) for detection of ATTO 643 dividing each emission channel into 256 × 256 pixels.

Image stacks of 150 frames were recorded for each cell at a time resolution of 32 ms/frame. Diffusion

constants were determined by mean square displacement analysis within a time window of 320 ms (10

frames).

Single molecule analysis. MHC homodimerization was quantified based on sequential co-localization and co-tracking analysis using self-written Matlab code "SlimFAST " as described in detail previously (Roder et al., 2014; Wilmes et al., 2015b). After aligning ATTO 643 and ATTO Rho11 channels with sub-pixel precision through spatial transformation based on a calibration measurement with multicolor fluorescent beads (TetraSpeck microspheres 0.1 μm, Invitrogen), individual molecules detected in both spectral channels of the same frame within a distance threshold of 100 nm were considered to be co-localized. For SMCT analysis, the multiple-target tracing (MTT) algorithm was applied to this dataset of co-localized molecules to reconstruct co-locomotion trajectories (co-trajectories) from the identified population of co-localizations. For the co-tracking analysis, only co-trajectories with a minimum of 10 consecutive steps (320 ms) were considered. This cut-off was determined based on systematic analysis of a negative control experiment with non-interacting model transmembrane proteins (Wilmes et al., 2020b) in order to minimize background from random co-localization. The relative fraction of co-tracked molecules was determined with respect to the absolute number of trajectories from both channels and corrected for homodimers stochastically double-labeled with the same fluorophore species as follows:

$$AB^* = \frac{AB}{2 \times \left[\left(\frac{A}{A+B}\right) \times \left(\frac{B}{A+B}\right)\right]}$$

$$rel.\,co-locomotion = \frac{2 \times AB^*}{(A+B)}$$

where A, B, AB and AB* are the numbers of trajectories observed for ATTO Rho11, ATTO 643, co-trajectories and corrected co-trajectories, respectively.

For **Figures 3F and 3G**, the relative fraction of co-localized mobile and immobile molecules was determined with respect to the absolute number of mobile or immobile molecules from both channels and corrected for homodimers stochastically double-labeled with the same fluorophore species as follows:

$$1 \qquad AB^* = \frac{AB/A + AB/B}{2}$$

$$2 \qquad rel.\,fraction = \frac{AB^*}{2 \times \left[\left(\frac{A}{A+B}\right) \times \left(\frac{B}{A+B}\right)\right]}$$

where A, B, AB and AB* are the numbers of molecules part of a trajectory observed for

ATTO Rho11, ATTO 643, co-localized molecules and corrected co-localized molecules, respectively.

Immobile molecules were classified by their appearance within a radius described by the localization

precision and molecular observation probability.

Statistics and statistical analysis. For **Figure 3D-H**, each data point represents the analysis from

one cell, with ≥14 cells measured per experiment and many trajectories analyzed per cell (**3D**, 238-

2513; **3E**, 238-2513; **3F**, 314-2732; **3G**, 41-354 tracked immobile particles and 362-3141 mobile

particles; **3H**, 2-301 co-trajectories). Statistical significances were calculated by two-sample

Kolmogorov-Smirnov test, Kruskal-Wallis test with multiple comparisons, unpaired *t* test and two-

way ANOVA with multiple comparisons as indicated in the figure legends, using version Prism 8.4.0

for MacOS (GraphPad, San Diego, USA). Box plots were used for visualization and indicate the data

distribution of second and third quartile (box), median (line), mean (square) and data range (whiskers).

Native mass spectrometry

In advance of native MS measurements, a small amount (<1%) of covalent dimers of the α_3 domain,

which had formed as side product during refolding, were removed by size exclusion chromatography

on Superdex 75 10/300 GL (Cytiva). Amicon Ultra 0.5 mL centrifugal filter units (molecular weight

cut-off 3 kDa; Merck Millipore) were used at 14,000 x g and 4 °C to exchange purified protein samples

to 150 mM ammonium acetate (99.99 %; Sigma-Aldrich), pH 7.2. The final concentration of the

α_3 domain monomer was 5 µM, 10 µM or 20 µM. Native MS analysis was implemented on a Q-Tof

II mass spectrometer in positive electrospray ionization mode. The instrument was modified to enable

high mass experiments (Waters and MS Vision, (van den Heuvel et al., 2006)). Sample ions were

introduced into the vacuum using homemade capillaries via a nano-electrospray ionization source in

positive ion mode (source pressure: 10 mbar). Borosilicate glass tubes (inner diameter: 0.68 mm, outer

diameter: 1.2 mm; World Precision Instruments) were pulled into closed capillaries in a two-step

program using a squared box filament (2.5 mm × 2.5 mm) within a micropipette puller (P-1000, Sutter

Instruments). The capillaries were then gold-coated using a sputter coater (5.0×10^{-2} mbar, 30.0 mA,

100 s, 3 runs to vacuum limit 3.0×10^{-2} mbar argon, distance of plate holder: 5 cm; CCU-010,

safematic). Capillaries were opened directly on the sample cone of the mass spectrometer. In regular

MS mode, spectra were recorded at a capillary voltage of 1.45 kV and a cone voltage of 100 V to

150 V. Protein species with quaternary structure were assigned by MS/MS analysis. These experiments

were carried out using argon as collision gas (1.2×10^{-2} mbar). The acceleration voltage ranged from

10 V to 100 V. Comparability of results was ensured as MS quadrupole profiles and pusher settings

were kept constant in all measurements. A spectrum of cesium iodide (25 g/L) was recorded on the

same day of the particular measurement to calibrate the data.

All spectra were evaluated regarding experimental mass (MassLynx V4.1, Waters) and area under

the curve (AUC; UniDec (Marty et al., 2015)) of the detected mass species. The values of the shown

averaged masses and AUC of the different species as well as the corresponding standard deviation

result from at least three independent measurements. Exact experimental masses are presented in

Figure 4S1.

Molecular model of the K^b heavy chain dimer

The crystal structure of K^b in complex with a chicken ovalbumin epitope (PDB 3p9m) served as

template to generate a model of the heavy chain dimer. Using the Pymol program (DeLano, 2002), it

was possible to superimpose a copy of the 3p9m heavy chain on the $\beta_2 m$ subunit of a second 3p9m

structure (FHC dimer). The superposition involved only the matching of the α_3 domain backbone of

the K^b molecule (residues 181-277) onto the $\beta_2 m$ subunit resulting in a small root-mean-square

deviation (RMSD) of 1.25 Å. Very little steric overlap of the superimposed α_3 domain with the second

heavy chain was detected. This minimal overlap of the α_1/α_2 domain of the superimposed heavy chain

was removed by adjusting backbone dihedral angles of residues 179-181 in the linker between α_1/α_2

domain and the α_3 domain. The initial structural model was energy-minimized to remove any residual

steric overlap and was prepared for molecular dynamics (MD) simulations using the Amber18 package

(Case et al., 2018).

Molecular docking

For comparison of FHC homodimers and K^b HC/$\beta_2 m$ heterodimers, MD simulations were

performed starting from the coordinates of the K^b structure in PDB 3p9m. Proteins were solvated in

octahedral boxes including explicit sodium and chloride ions (0.15 M) and explicit TIP3P water

molecules keeping a minimum distance of 10 Å between protein atoms and box boundaries (Jorgensen

et al., 1983). The parm14SB force field was used for the proteins and peptides (Maier et al., 2015).

The simulation systems were again energy minimized (5000 steps) after solvation followed by heating

up to 300 K in steps of 100 K with position restraints on all heavy atoms of the proteins. Subsequently,

positional restraints were gradually removed from an initial 25 kcal·mol^{-1}·Å^{-2} to 0.5 kcal·mol^{-1}·Å^{-2}

within 0.5 ns followed by a 1 ns unrestrained equilibration at 300 K. All production simulations were

performed at a temperature of 300 K and a pressure of 1 bar. The hydrogen mass repartition option of

Amber was used to allow a time step of 4 fs (Hopkins et al., 2015). Unrestrained production

simulations for up to 400 ns were performed. The interface packing was analyzed by calculation of the

buried surface area using the Shrake method, (Shrake and Rupley, 1973); analysis of root-mean-square

deviations (RMSD) was performed using the cpptraj module of the Amber18 package.

Acknowledgements

10 We thank the donors of reagents as mentioned in the Materials, Venkat Raman Ramnarayan for

11 comments on the manuscript, Christian P. Richter for support with SMT/SMCT evaluation, Ankur

12 Saikia and Christian Guenther for performing protein chromatography, the iBiOs staff for technical

13 support with single molecule microscopy, the SPC facility at EMBL Hamburg for technical support,

14 and Uschi Wellbrock for excellent technical assistance.

Funding

Deutsche Forschungsgemeinschaft (DFG SP583/7-2), Bundesministerium für Bildung und

Forschung (BMBF, 031A153A); Tönjes Vagt Foundation (XXXII), iNEXT-Discovery (11911),

Jacobs University (all to SSp); Deutsche Forschungsgemeinschaft (SFB 944, projects P8 and Z,

Facility iBiOs, PI 405/14-1 to JP). PL and JW acknowledge funding from the province of Upper

Austria as a part of the FH Upper Austria Center of Excellence for Technological Innovation in

Medicine (TIMed Center) and the Christian Doppler Forschungsgesellschaft (Josef Ressel Center for

Phytogenic Drug Research). CU acknowledges funding from the Leibniz Association through SAW-

2014-HPI-4 grant. The Heinrich-Pette-Institute, Leibniz Institute for Experimental Virology is

supported by the Free and Hanseatic City Hamburg and the Federal Ministry of Health

(Bundesministerium für Gesundheit, BMG).

Author Contributions

The experimental work was performed as follows: CD, Fig. 1BCDEG; ZH, Fig. 1F; PL, Fig. 2,

2S1-3, and 4AB; SL, Fig. 3, 3S1-3, movies; ARH, Fig. 4CD; JDK, Fig. 4E-G, 4S1. MGA, CU, JW

and JP supervised and interpreted experiments. MZ performed all MD simulations (Fig. 5, 5S1). SSp

conceived the work, supervised and coordinated experiments, and wrote the manuscript in cooperation

with JP and PL and assistance by NL with input from all authors. Funding was acquired by CU, JP,

JW, MGA, MZ, PL, and SSp.

Competing Interests:

13   The authors declare that no competing interests exist.

References

Allen RL, O'Callaghan CA, McMichael AJ, Bowness P. 1999. Cutting edge: HLA-B27 can form a novel beta 2-microglobulin-free heavy chain homodimer structure. *J Immunol Baltim Md 1950* **162**:5045–5048.

Anjanappa R, Garcia-Alai M, Kopicki J-D, Lockhauserbäumer J, Aboelmagd M, Hinrichs J, Nemtanu IM, Uetrecht C, Zacharias M, Springer S, Meijers R. 2020. Structures of peptide-free and partially loaded MHC class I molecules reveal mechanisms of peptide selection. *Nat Commun* **11**:1314. doi:10.1038/s41467-020-14862-4

Armony G, Heck AJR, Wu W. 2021. Extracellular crosslinking mass spectrometry reveals HLA class I – HLA class II interactions on the cell surface. *Mol Immunol* **136**:16–25. doi:10.1016/j.molimm.2021.05.010

Arosa FA, Santos SG, Powis SJ. 2007. Open conformers: the hidden face of MHC-I molecules. *Trends Immunol* **28**:115–123. doi:10.1016/j.it.2007.01.002

Bahadur RP, Zacharias M. 2008. The interface of protein-protein complexes: Analysis of contacts and prediction of interactions. *Cell Mol Life Sci* **65**:1059–1072. doi:10.1007/s00018-007-7451-x

Baia D, Pou J, Jones D, Mandelboim O, Trowsdale J, Muntasell A, Lopez-Botet M. 2016. Interaction of the LILRB1 inhibitory receptor with HLA class Ia dimers. *Eur J Immunol* **46**:1681–90. doi:10.1002/eji.201546149

Baía D, Pou J, Jones D, Mandelboim O, Trowsdale J, Muntasell A, López-Botet M. 2016. Interaction of the LILRB1 inhibitory receptor with HLA class Ia dimers. *Eur J Immunol* **46**:1681–1690. doi:10.1002/eji.201546149

Barnstable C. 1978. Production of monoclonal antibodies to group A erythrocytes, HLA and other human cell surface antigens-new tools for genetic analysis. *Cell* **14**:9–20. doi:10.1016/0092-8674(78)90296-9

Blumenthal D, Edidin M, Gheber LA. 2016. Trafficking of MHC molecules to the cell surface creates dynamic protein patches. *J Cell Sci* **129**:3342–3350. doi:10.1242/jcs.187112

Bodnar A, Bacso Z, Jenei A, Jovin TM, Edidin M, Damjanovich S, Matko J. 2003. Class I HLA oligomerization at the surface of B cells is controlled by exogenous beta(2)-microglobulin: implications in activation of cytotoxic T lymphocytes. *Int Immunol* **15**:331–9.

Borgmann DM, Weghuber J, Schaller S, Jacak J, Winkler SM. 2012. Identification of patterns in microscopy images of biological samples using evolution strategies.

Campbell EC, Antoniou AN, Powis SJ. 2012. The multi-faceted nature of HLA class I dimer molecules. *Immunology* **136**:380–384. doi:10.1111/j.1365-2567.2012.03593.x

Capps GG, Robinson BE, Lewis KD, Zúñiga MC. 1993. In vivo dimeric association of class I MHC heavy chains. Possible relationship to class I MHC heavy chain-beta 2-microglobulin dissociation. *J Immunol* **151**:159–169.

Case DA, Ben-Shalom IY, Brozell SR, Cerutti DS, Cheatham TE III, Cruzeiro VWD, Darde TA, Duke RE, Ghoreishi D, Gilson MK, Gohlke H, Goetz AW, Greene D, Harris R, Homeyer N, Izadi S, Kovalenko A, Kurtzman T, Lee TS, LeGrand S, Li P, Lin C, Liu J, Luchko T, Luo R, Mermelstein DJ, Merz KM, Miao Y, Monard G, Nguyen C, Nguyen H, Omelyan I, Onufriev A, Pan F, Qi R, Roe DR, Roitberg A, Sagui C, Schott-Verdugo S, Shen J, Simmerling CL, Smith J, Salomon-Ferrer R, Swails J, Walker RC, Wang J, Wei H, Wolf RM, Wu X, Xiao L, York DM, Kollman PA. 2018. Amber18. San Fransisco: University of California.

Chakrabarti A, Matko J, Rahman NA, Barisas BG, Edidin M. 1992. Self-association of class I major histocompatibility complex molecules in liposome and cell surface membranes. *Biochemistry* **31**:7182–9.

Chen B, Li J, He C, Li D, Tong W, Zou Y, Xu W. 2017. Role of HLA-B27 in the pathogenesis of ankylosing spondylitis. *Mol Med Rep* **15**:1943–1951. doi:10.3892/mmr.2017.6248

Comber JD, Philip R. 2014. MHC class I antigen presentation and implications for developing a new generation of therapeutic vaccines. *Ther Adv Vaccines* **2**:77–89. doi:10.1177/2051013614525375

Dangoria NS, DeLay ML, Kingsbury DJ, Mear JP, Uchanska-Ziegler B, Ziegler A, Colbert RA. 2002. HLA-B27 Misfolding Is Associated with Aberrant Intermolecular Disulfide Bond Formation (Dimerization) in the Endoplasmic Reticulum. *J Biol Chem* **277**:23459–23468. doi:10.1074/jbc.M110336200

Day PM, Esquivel F, Lukszo J, Bennink JR, Yewdell JW. 1995. Effect of TAP on the generation and intracellular trafficking of peptide-receptive major histocompatibility complex class I molecules. *Immunity* **2**:137–147. doi:10.1016/S1074-7613(95)80014-X

DeLano WL. 2002. Pymol: An open-source molecular graphics tool. *CCP4 Newsl Protein Crystallogr* 82–92.

Dirscherl C, Hein Z, Ramnarayan VR, Jacob-Dolan C, Springer S. 2018. A two-hybrid antibody micropattern assay reveals specific in cis interactions of MHC I heavy chains at the cell surface. *eLife* **7**:e34150. doi:10.7554/eLife.34150

Dirscherl C, Palankar R, Delcea M, Kolesnikova TA, Springer S. 2017. Specific Capture of Peptide-Receptive Major Histocompatibility Complex Class I Molecules by Antibody Micropatterns Allows for a Novel Peptide-Binding Assay in Live Cells. *Small* **13**:1602974. doi:10.1002/smll.201602974

Dulberger CL, McMurtrey CP, Hölzemer A, Neu KE, Liu V, Steinbach AM, Garcia-Beltran WF, Sulak M, Jabri B, Lynch VJ, Altfeld M, Hildebrand WH, Adams EJ. 2017. Human Leukocyte Antigen F Presents Peptides and Regulates Immunity through Interactions with NK Cell Receptors. *Immunity* **46**:1018-1029.e7. doi:10.1016/j.immuni.2017.06.002

Edidin M, Achilles S, Zeff R, Wei T. 1997. Probing the stability of class I major histocompatibility complex (MHC) molecules on the surface of human cells. *Immunogenetics* **46**:41–45. doi:10.1007/s002510050240

Elliott T, Cerundolo V, Elvin J, Townsend A. 1991. Peptide-induced conformational change of the class I heavy chain. *Nature* **351**:402–406. doi:10.1038/351402a0

Fassett MS, Davis DM, Valter MM, Cohen GB, Strominger JL. 2001. Signaling at the inhibitory natural killer cell immune synapse regulates lipid raft polarization but not class I MHC clustering. *Proc Natl Acad Sci U S A* **98**:14547–14552. doi:10.1073/pnas.211563598

Ferez M, Castro M, Alarcon B, van Santen HM. 2014. Cognate peptide-MHC complexes are expressed as tightly apposed nanoclusters in virus-infected cells to allow TCR crosslinking. *J Immunol* **192**:52–8. doi:10.4049/jimmunol.1301224

Fooksman DR, Grönvall GK, Tang Q, Edidin M. 2006. Clustering Class I MHC Modulates Sensitivity of T Cell Recognition. *J Immunol* **176**:6673–6680. doi:10.4049/jimmunol.176.11.6673

Fridy PC, Li Y, Keegan S, Thompson MK, Nudelman I, Scheid JF, Oeffinger M, Nussenzweig MC, Fenyö D, Chait BT, Rout MP. 2014. A robust pipeline for rapid production of versatile nanobody repertoires. *Nat Methods* **11**:1253–1260. doi:10.1038/nmeth.3170

Gakamsky DM, Bjorkman PJ, Pecht I. 1996. Peptide Interaction with a Class I Major Histocompatibility Complex-Encoded Molecule: Allosteric Control of the Ternary Complex Stability [†]. *Biochemistry* **35**:14841–14848. doi:10.1021/bi961707u

Gambin Y, Lopez-Esparza R, Reffay M, Sierecki E, Gov NS, Genest M, Hodges RS, Urbach W. 2006. Lateral mobility of proteins in liquid membranes revisited. *Proc Natl Acad Sci U S A* **103**:2098–2102. doi:10.1073/pnas.0511026103

Garstka M, Borchert B, Al-Balushi M, Praveen P, Kühl N, Majoul I, Duden R, Springer S. 2007. Peptide-receptive Major Histocompatibility Complex Class I Molecules Cycle between Endoplasmic Reticulum and *cis* -Golgi in Wild-type Lymphocytes. *J Biol Chem* **282**:30680–30690. doi:10.1074/jbc.M701721200

Geng J, Altman JD, Krishnakumar S, Raghavan M. 2018. Empty conformers of HLA-B preferentially bind CD8 and regulate CD8+ T cell function. *eLife* **7**:e36341. doi:10.7554/eLife.36341

Goodridge JP, Lee N, Burian A, Pyo C-W, Tykodi SS, Warren EH, Yee C, Riddell SR, Geraghty DE. 2013. HLA-F and MHC-I Open Conformers Cooperate in a MHC-I Antigen Cross-Presentation Pathway. *J Immunol* **191**:1567–1577. doi:10.4049/jimmunol.1300080

Graham FL, van der Eb AJ. 1973. A new technique for the assay of infectivity of human adenovirus 5 DNA. *Virology* **52**:456–467. doi:10.1016/0042-6822(73)90341-3

Hein Z, Uchtenhagen H, Abualrous ET, Saini SK, Janssen L, Van Hateren A, Wiek C, Hanenberg H, Momburg F, Achour A, Elliott T, Springer S, Boulanger D. 2014. Peptide-independent stabilization of MHC class I molecules breaches cellular quality control. *J Cell Sci* **127**:2885–2897. doi:10.1242/jcs.145334

Ho CCM, Chhabra A, Starkl P, Schnorr P-J, Wilmes S, Moraga I, Kwon H-S, Gaudenzio N, Sibilano R, Wehrman TS, Gakovic M, Sockolosky JT, Tiffany MR, Ring AM, Piehler J, Weissman IL, Galli SJ, Shizuru JA, Garcia KC. 2017. Decoupling the Functional Pleiotropy of Stem Cell Factor by Tuning c-Kit Signaling. *Cell* **168**:1041-1052.e18. doi:10.1016/j.cell.2017.02.011

Hopkins CW, Le Grand S, Walker RC, Roitberg AE. 2015. Long-Time-Step Molecular Dynamics through Hydrogen Mass Repartitioning. *J Chem Theory Comput* **11**:1864–1874. doi:10.1021/ct5010406

Jorgensen WL, Chandrasekhar J, Madura JD, Impey RW, Klein ML. 1983. Comparison of simple potential functions for simulating liquid water. *J Chem Phys* **79**:926–935. doi:10.1063/1.445869

Kaufman J. 2018. Generalists and Specialists: A New View of How MHC Class I Molecules Fight Infectious Pathogens. *Trends Immunol* **39**:367–379. doi:10.1016/j.it.2018.01.001

Kirchhofer A, Helma J, Schmidthals K, Frauer C, Cui S, Karcher A, Pellis M, Muyldermans S, Casas-Delucchi CS, Cardoso MC, Leonhardt H, Hopfner K-P, Rothbauer U. 2010. Modulation of protein properties in living cells using nanobodies. *Nat Struct Mol Biol* **17**:133–138. doi:10.1038/nsmb.1727

Lanzerstorfer P, Borgmann D, Schütz G, Winkler SM, Höglinger O, Weghuber J. 2014. Quantification and kinetic analysis of Grb2-EGFR interaction on micro-patterned surfaces for the characterization of EGFR-modulating substances. *PloS One* **9**:e92151. doi:10.1371/journal.pone.0092151

Lanzerstorfer P, Müller U, Gordiyenko K, Weghuber J, Niemeyer CM. 2020. Highly Modular Protein Micropatterning Sheds Light on the Role of Clathrin-Mediated Endocytosis for the Quantitative Analysis of Protein-Protein Interactions in Live Cells. *Biomolecules* **10**:540. doi:10.3390/biom10040540

Lin W-C, Iversen L, Tu H-L, Rhodes C, Christensen SM, Iwig JS, Hansen SD, Huang WYC, Groves JT. 2014. H-Ras forms dimers on membrane surfaces via a protein-protein interface. *Proc Natl Acad Sci U S A* **111**:2996–3001. doi:10.1073/pnas.1321155111

Lippincott-Schwartz J, Snapp EL, Phair RD. 2018. The Development and Enhancement of FRAP as a Key Tool for Investigating Protein Dynamics. *Biophys J* **115**:1146–1155. doi:10.1016/j.bpj.2018.08.007

Ljunggren H-G, Stam NJ, Öhlén C, Neefjes JJ, Höglund P, Heemels M-T, Bastin J, Schumacher TNM, Townsend A, Kärre K, Ploegh HL. 1990. Empty MHC class I molecules come out in the cold. *Nature* **346**:476–480. doi:10.1038/346476a0

Low-Nam ST, Lidke KA, Cutler PJ, Roovers RC, van Bergen en Henegouwen PMP, Wilson BS, Lidke DS. 2011. ErbB1 dimerization is promoted by domain co-confinement and stabilized by ligand binding. *Nat Struct Mol Biol* **18**:1244–1249. doi:10.1038/nsmb.2135

Lu X, Gibbs JS, Hickman HD, David A, Dolan BP, Jin Y, Kranz DM, Bennink JR, Yewdell JW, Varma R. 2012. Endogenous viral antigen processing generates peptide-specific MHC class I cell-surface clusters. *Proc Natl Acad Sci U S A* **109**:15407–15412. doi:10.1073/pnas.1208696109

Mahmutefendic H, Blagojevic G, Tomas MI, Kucic N, Lucin P. 2011. Segregation of open Major Histocompatibility Class I conformers at the plasma membrane and during endosomal trafficking reveals conformation-based sorting in the endosomal system. *Int J Biochem Cell Biol* **43**:504–15. doi:10.1016/j.biocel.2010.12.002

Maier JA, Martinez C, Kasavajhala K, Wickstrom L, Hauser KE, Simmerling C. 2015. ff14SB: Improving the Accuracy of Protein Side Chain and Backbone Parameters from ff99SB. *J Chem Theory Comput* **11**:3696–3713. doi:10.1021/acs.jctc.5b00255

Makhadiyeva D, Lam L, Moatari M, Vallance J, Zheng Y, Campbell EC, Powis SJ. 2012. MHC class I dimer formation by alteration of the cellular redox environment and induction of apoptosis: Redox control of MHC class I dimer formation. *Immunology* **135**:133–139. doi:10.1111/j.1365-2567.2011.03518.x

Marty MT, Baldwin AJ, Marklund EG, Hochberg GKA, Benesch JLP, Robinson CV. 2015. Bayesian Deconvolution of Mass and Ion Mobility Spectra: From Binary Interactions to Polydisperse Ensembles. *Anal Chem* **87**:4370–4376. doi:10.1021/acs.analchem.5b00140

Matko J, Bushkin Y, Wei T, Edidin M. 1994. Clustering of class I HLA molecules on the surfaces of activated and transformed human cells. *J Immunol* **152**:3353–60.

Montealegre Sebastián, Venugopalan V, Fritzsche S, Kulicke C, Hein Z, Springer S. 2015. Dissociation of β_2-microglobulin determines the surface quality control of major histocompatibility complex class I molecules. *FASEB J* **29**:2780–2788. doi:10.1096/fj.14-268094

Montealegre S., Venugopalan V, Fritzsche S, Kulicke C, Hein Z, Springer S. 2015. Dissociation of beta2-microglobulin determines the surface quality control of major histocompatibility complex class I molecules. *Faseb J* **29**:2780–8. doi:10.1096/fj.14-268094

Moraga I, Wernig G, Wilmes S, Gryshkova V, Richter CP, Hong W-J, Sinha R, Guo F, Fabionar H, Wehrman TS, Krutzik P, Demharter S, Plo I, Weissman IL, Minary P, Majeti R, Constantinescu SN, Piehler J, Garcia KC. 2015. Tuning cytokine receptor signaling by re-orienting dimer geometry with surrogate ligands. *Cell* **160**:1196–1208. doi:10.1016/j.cell.2015.02.011

Niman HL, Houghten RA, Walker LE, Reisfeld RA, Wilson IA, Hogle JM, Lerner RA. 1983. Generation of protein-reactive antibodies by short peptides is an event of high frequency:

implications for the structural basis of immune recognition. *Proc Natl Acad Sci* **80**:4949–4953. doi:10.1073/pnas.80.16.4949

Ortiz-Navarrete V, Hammerling GJ. 1991. Surface appearance and instability of empty H-2 class I molecules under physiological conditions. *Proc Natl Acad Sci* **88**:3594–3597. doi:10.1073/pnas.88.9.3594

Reinhardt U, Lotze J, Zernia S, Mörl K, Beck-Sickinger AG, Seitz O. 2014. Peptide-Templated Acyl Transfer: A Chemical Method for the Labeling of Membrane Proteins on Live Cells. *Angew Chem Int Ed* **53**:10237–10241. doi:10.1002/anie.201403214

Richter D, Moraga I, Winkelmann H, Birkholz O, Wilmes S, Schulte M, Kraich M, Kenneweg H, Beutel O, Selenschik P, Paterok D, Gavutis M, Schmidt T, Garcia KC, Müller TD, Piehler J. 2017. Ligand-induced type II interleukin-4 receptor dimers are sustained by rapid re-association within plasma membrane microcompartments. *Nat Commun* **8**:15976. doi:10.1038/ncomms15976

Roder F, Wilmes S, Richter CP, Piehler J. 2014. Rapid Transfer of Transmembrane Proteins for Single Molecule Dimerization Assays in Polymer-Supported Membranes. *ACS Chem Biol* **9**:2479–2484. doi:10.1021/cb5005806

Schell TD. 2002. The assembly of functional beta2-microglobulin-free MHC class I molecules that interact with peptides and CD8+ T lymphocytes. *Int Immunol* **14**:775–782. doi:10.1093/intimm/dxf041

Schwarzenbacher M, Kaltenbrunner M, Brameshuber M, Hesch C, Paster W, Weghuber J, Heise B, Sonnleitner A, Stockinger H, Schutz GJ. 2008. Micropatterning for quantitative analysis of protein-protein interactions in living cells. *Nat Methods* **5**:1053–60. doi:10.1038/nmeth.1268

Sevcsik E, Brameshuber M, Folser M, Weghuber J, Honigmann A, Schutz GJ. 2015. GPI-anchored proteins do not reside in ordered domains in the live cell plasma membrane. *Nat Commun* **6**:6969. doi:10.1038/ncomms7969

Shrake A, Rupley JA. 1973. Environment and exposure to solvent of protein atoms. Lysozyme and insulin. *J Mol Biol* **79**:351–371. doi:10.1016/0022-2836(73)90011-9

Spack E, Edidin M. 1986. The class I MHC antigens of erythrocytes: a serologic and biochemical study. *J Immunol Baltim Md 1950* **136**:2943–2952.

Sprague B, Mcnally J. 2005. FRAP analysis of binding: proper and fitting. *Trends Cell Biol* **15**:84–91. doi:10.1016/j.tcb.2004.12.001

Townsend A, Bodmer H. 1989. Antigen recognition by class I -restricted cytotoxic T lymphocytes. *Annu Rev Immunol* 601–24.

Townsend A, Elliott T, Cerundolo V, Foster L, Barber B, Tse A. 1990. Assembly of MHC class I molecules analyzed in vitro. *Cell* **62**:285–295. doi:10.1016/0092-8674(90)90366-M

Townsend A, Öhlén C, Bastin J, Ljunggren H-G, Foster L, Kärre K. 1989. Association of class I major histocompatibility heavy and light chains induced by viral peptides. *Nature* **340**:443–448. doi:10.1038/340443a0

Triantafilou K, Triantafilou M, Wilson KM, Fernandez N. 2000. Human major histocompatibility molecules have the intrinsic ability to form homotypic associations. *Hum Immunol* **61**:585–598. doi:10.1016/S0198-8859(00)00112-9

van den Heuvel RHH, van Duijn E, Mazon H, Synowsky SA, Lorenzen K, Versluis C, Brouns SJJ, Langridge D, van der Oost J, Hoyes J, Heck AJR. 2006. Improving the Performance of a Quadrupole Time-of-Flight Instrument for Macromolecular Mass Spectrometry. *Anal Chem* **78**:7473–7483. doi:10.1021/ac061039a

Váradi T, Schneider M, Sevcsik E, Kiesenhofer D, Baumgart F, Batta G, Kovács T, Platzer R, Huppa
JB, Szöllősi J, Schütz GJ, Brameshuber M, Nagy P. 2019. Homo- and Heteroassociations Drive
Activation of ErbB3. *Biophys J* **117**:1935–1947. doi:10.1016/j.bpj.2019.10.001

Vogelsang J, Kasper R, Steinhauer C, Person B, Heilemann M, Sauer M, Tinnefeld P. 2008. A
Reducing and Oxidizing System Minimizes Photobleaching and Blinking of Fluorescent Dyes.
Angew Chem Int Ed **47**:5465–5469. doi:10.1002/anie.200801518

Wilmes S, Beutel O, Li Z, Francois-Newton V, Richter CP, Janning D, Kroll C, Hanhart P, Hötte K,
You C, Uzé G, Pellegrini S, Piehler J. 2015a. Receptor dimerization dynamics as a regulatory
valve for plasticity of type I interferon signaling. *J Cell Biol* **209**:579–593.
doi:10.1083/jcb.201412049

Wilmes S, Beutel O, Li Z, Francois-Newton V, Richter CP, Janning D, Kroll C, Hanhart P, Hötte K,
You C, Uzé G, Pellegrini S, Piehler J. 2015b. Receptor dimerization dynamics as a regulatory
valve for plasticity of type I interferon signaling. *J Cell Biol* **209**:579–593.
doi:10.1083/jcb.201412049

Wilmes S, Hafer M, Vuorio J, Tucker JA, Winkelmann H, Löchte S, Stanly TA, Pulgar Prieto KD,
Poojari C, Sharma V, Richter CP, Kurre R, Hubbard SR, Garcia KC, Moraga I, Vattulainen I,
Hitchcock IS, Piehler J. 2020a. Mechanism of homodimeric cytokine receptor activation and
dysregulation by oncogenic mutations. *Science* **367**:643–652. doi:10.1126/science.aaw3242

Wilmes S, Hafer M, Vuorio J, Tucker JA, Winkelmann H, Löchte S, Stanly TA, Pulgar Prieto KD,
Poojari C, Sharma V, Richter CP, Kurre R, Hubbard SR, Garcia KC, Moraga I, Vattulainen I,
Hitchcock IS, Piehler J. 2020b. Mechanism of homodimeric cytokine receptor activation and
dysregulation by oncogenic mutations. *Science* **367**:643–652. doi:10.1126/science.aaw3242

You C, Richter CP, Löchte S, Wilmes S, Piehler J. 2014. Dynamic Submicroscopic Signaling Zones
Revealed by Pair Correlation Tracking and Localization Microscopy. *Anal Chem* **86**:8593–
8602. doi:10.1021/ac501127r

28
29

1   **Figure legends**

2   With the figures.

1   **Tables**

3   **Table 1:** Comparison of diffusion coefficients (mean $\pm$ SD in $\mu m^2/s$). NA, not applicable.

	FRAP	SMT
HC/β₂m/peptide trimers (+peptide)	0.35 ±0.12	0.35±0.06
FHCs (–peptide)	0.19 ±0.05	0.25±0.04
FHCs (–peptide) [a]	0.19 ±0.05	0.18±0.10
Dimerized HC/β₂m/peptide trimers (+peptide, + CL)	NA	0.20±0.05
Dimerized HC/β₂m/peptide trimers (+peptide, + CL)[a]	NA	0.15±0.04

4   [a]only dimers identified by SMCT

1 **Supplement figure legends**

2 With the supplement figures.

Supplement Movie legends

Movie S1: Dual-color SMCT identifies FHC association in live cell plasma membranes.

GFP-K^b dimer formation was probed by co-locomotion analysis. Individual GFP-K^b molecules

labeled with Rho11 (magenta) and Dy647 (green) in STF1 cells were imaged by TIRF microscopy

(scale bar, 2.5 µm; frame rate, 10 Hz; playback speed, 1/3 real time). Movies were acquired in the

absence of peptide (FHCs, left), in the presence of peptide (HC/β_2m/peptide complexes; center) and

in the presence of peptide and a crosslinker consisting of a dimeric anti-GFP nanobody (right).

Localized molecules in each channel are encircled, co-locomoting molecules are highlighted by white

diamonds.

Movies S2: SMCT demonstrates transient association of FHCs.

Individual GFP-K^b molecules labeled with Rho11 (magenta) and Dy647 (green) imaged by TIRF

microscopy (scale bar, 450 nm; frame rate, 10 Hz; playback, 1/3 real time). Selected image section

showing SMCT of an individual FHC complex (*i.e.*, in the absence of peptide). Localized molecules

in each channel are encircled, co-locomoting molecules are highlighted by white diamonds, with the

dissociation occurring between 4.03 s and 4.06 s.

Movies S3A-B: Molecular dynamics simulations.

A, 400 ns MD simulation of an FHC dimer as in **Figure 4E**. The extracellular domains of the FHCs

are shown. The α_3 domain of one K^b molecule is depicted in yellow, the α_3 domain of the other K^b

molecule is depicted in red. The α_1/α_2 superdomains are shown in yellow and purple for both FHCs.

B, 400 ns MD simulation of the K^b FHC/β_2m heterodimer. The extracellular domains of the FHCs

are shown. The light chain β_2m (red) and the α_1/α_2 superdomain (yellow and purple) show a favorable

interface.

Figure 1: MHC I HC/HC association requires dissociation of β₂m but no disulfide bonds.

A, Schematic of MHC class I states at the plasma membrane (PM). Dissociation of peptide from the HC/β2m/peptide trimer results in an 'empty' HC/β_2m heterodimer. Dissociation of β_2m then produces 'free' HCs, which can form FHC associations (HC/HC dimer shown).

B, Schematic representation of the two-hybrid antibody micropattern assay. Cells expressing a class I GFP fusion (green) and an N-terminally HA-tagged class I (gray) are seeded onto glass slides that are printed with micrometer-sized patterns of fluorescently labelled anti-HA antibodies. Dissociation of β_2m generates FHCs of both constructs, which diffuse freely in the plasma membrane and even-tually associate with each other (center). The HC/HC associations, which contain both HA-tagged and GFP-fused FHCs, localize in the pattern elements and are visible as pattern-shaped GFP fluorescence on the plasma membrane.

C, Representative fluorescence micrograph showing one single STF1 cell expressing both HA-tagged and GFP-fused H-2K^b in phase contrast (left), the purple anti-HA antibody pattern on the glass slide (middle), and green K^b-GFP colocalizing with the antibody pattern (right). The arrows point to the fluorophore observed in the respective panel and emphasize the plane of the image. Scale bar, 20 μm.

D, Interaction occurs between HA-tagged K^b and Db-GFP FHCs (37 °C) but not between HC/β_2m heterodimers (25 °C). Scale bar, 20 μm.

E, HC/HC interaction does not involve intracellular disulfide bond formation. No interaction is visible between FHCs (37 °C) of HA–K^b and K^b(C332S) GFP, which lacks the cytosolic cysteine. Scale bar, 20 μm.

F, B27, but not K^b, forms covalent dimers. HA-K^b (K^b in the label) and HA-B*27:05 (B27) molecules were immunoprecipitated from the lysate of transduced STF1 cells with an anti-HA monoclonal antibody, separated by reducing (+) or nonreducing (–) SDS-PAGE, and monomers (heavy chain) and covalent homodimers as indicated were detected by Western blotting with an anti-HA antiserum. * de-notes a background band.

G, No interaction of K^b–GFP with F pocket-stabilized HA–K^b(Y84C/A139C), a construct with stable β_2m association. Scale bar, 20 μm.

Figure 2: FHCs interact in TAP-proficient cells and in the absence of patterns.
(continued on next page)

Figure 2: FHCs interact in TAP-proficient cells and in the absence of patterns.

A, Schematic of cell surface interaction between FHCs in TAP2-deficient (left) and TAP2-proficient cells.

B, Time course of K^b HC/HC interaction in the antibody micropattern assay upon temperature shift from 25 °C to 37 °C in TAP2-proficient (right) and TAP2-deficient cells (left). Representative TIRF microscopy images are shown. 2 µM of cognate SIINFEKL peptide were added as indicated. Scale bar, 8 µm. Insets show enlarged intensity-adjusted regions within the selected cells.

C, Quantification of fluorescence contrast from B. Error bars are standard error of the mean (SEM) of >10 cells measured in ≥2 independent experiments.

D, FRAP analysis of K^b cell surface dynamics. Cells expressing K^b constructs as in B were monitored in the absence of patterns (top) or on anti-HA antibody patterns (bottom) with or without SIINFEKL peptide as indicated. Regions in the yellow circle were bleached, and recovery of fluorescence was followed over time. Scale bar, 5 µm. Insets show the enlarged intensity-adjusted region of bleached spot.

E, Normalized mean fluorescence recovery curves from FRAP experiments, error bars represent SEM.

F, Diffusion coefficients calculated from E. Significant increase in diffusion coefficients upon peptide addition, both in the absence and presence of antibody patterns. ****, $p \leq 0.0001$ by two-way ANOVA. ns, not significant. Error bars are SEM of >20 cells for each condition measured in ≥2 inde-pendent experiments.

G, Immobile fraction of K^b–GFP molecules, i.e. molecules that remain in the bleached area during the time of the experiment. With peptide vs. without peptide ****, $p \leq 0.0001$ and on pattern vs. in the absence of pattern ****, $p \leq 0.0001$ by two-way ANOVA. Error bars are SEM of >20 cells for each condition measured in ≥2 independent experiments.

H, Exchange rates of K^b molecules calculated from fluorescence recovery curves. k_{fast} (left Y axis) is the rate of diffusion of K^b–GFP molecules. k_{slow} (right Y axis) is the rate of K^b–GFP exchange on (= binding to and dissociation from) HA–K^b. With peptide vs. without peptide ****, $p \leq 0.0001$ on anti-HA pattern by two-way ANOVA. Error bars are SEM of >20 cells for each condition measured in ≥2 independent experiments.

Figure 3: Single-molecule microscopy shows K^b dimer formation.
(continued on next page)

Figure 3: Single-molecule microscopy shows K^b dimer formation.

A, Schematic: analyzing FHC association by single-molecule microscopy. GFP-K^b molecules at the plasma membrane were labeled with two different fluorophores (ATTO Rho11, magenta, and ATTO 643, green) using anti-GFP nanobodies. Diffusion and interaction of individual molecules was quantified by tracking and co-tracking of individual molecules.

B, Formation and dissociation of an individual GFP-K^b dimer detected by co-tracking analysis. Co-locomoting molecules are highlighted by a white square. Scale bar, 400 nm.

C, Single molecule trajectories (green, magenta) and co-trajectories (black) observed for GFP-K^b FHCs (no peptide, top) and HC/β_2m/peptide trimers (With peptide, middle). As a control, the same experiment was carried out after crosslinking peptide-loaded K^b with a tandem anti-GFP nanobody (With crosslinker, bottom). Scale bar, 2 µm.

D-H, statistical analyses by two-sample Kolmogorov-Smirnov test (****, p ≤0.0001; ***, p ≤0.001; **, p ≤0.01). Sample numbers see the methods section.

D, Fraction of detected co-trajectories out of the total number of observed trajectories for K^b in the absence and in the presence of peptide and crosslinker as indicated.

E, Diffusion coefficients of K^b molecules determined by single-molecule tracking in the absence and presence of peptide and crosslinker.

F, The fraction of immobile molecules out of the total GFP-K^b surface molecules determined by single-molecule tracking. SIINFEKL peptide and crosslinker were added as indicated.

G, Comparison of the fractions of mobile and immobile GFP-K^b molecules that are associated. The fraction of (associated immobile K^b)/(total immobile K^b) is shown vs. the fraction of (associated mobile K^b)/(total mobile K^b).

H, Quantification of co-trajectory half-life for GFP-K^b as obtained from single molecule co-trajectories with SIINFEKL peptide and crosslinker as indicated; the dissociation rates calculated from the means are 3.1 (no peptide), 17 (peptide, no crosslinker), and 1.3 s^{-1} (with peptide and crosslinker).

Figure 4: The K^b α$_3$ domains form dimers.

A, HC/HC association between HA-K^b-RFP and α_3-GFP. Distance between pattern elements, 2 μm.

B, HC/HC association between HA-α_3 and α_3-GFP. Distance between pattern elements, 2 μm.

C, Size exclusion chromatography of the K^b α_3 domain expressed in E.coli and folded in vitro.

D, Nonreducing SDS-PAGE of the α_3 domain homodimer fraction from C shows absence of disulfide linkage between the monomers. The positive control for disulfide oligomer formation is α_3 inclusion bodies, where some molecules are linked by disulfide bonds.

E, MS/MS analysis of the α_3 domain at 10 μM. The 9+ peak at 2551 *m/z* corresponding to the α_3 dimer was selected for MS/MS analysis at 10 V, 50 V and 100 V in the collision cell. The dimer (double spheres) easily dissociates in the gas-phase indicating a non-covalent rather low-affinity binding event. At 50 V, no dimer is detectable anymore displaced by monomeric signal (single spheres).

F, Overall area under the curve (AUC) for the detected α_3 domain dimers. Native mass spectra were recorded using 5 μM, 10 μM or 20 μM α_3 domain at 10 V (dark red) and 25 V (light red), respectively. The AUC was determined over the entire spectrum for the dimeric mass species and is shown as mean (*n*=3) ±SD. The dimeric fraction is concentration-dependent.

G, Representative mass spectra (20 μM α_3 domain) show the charge distributions of monomer (single sphere) and dimer (double spheres) at 10 V (dark red) and 25 V (light red) in the collision cell.

Figure 5: One hypothetical model of a Kᵇ FHC dimer.
In this hypothetical structure of the extracellular portion of an FHC dimer **(A)**, one of several structures predicted by molecular docking and molecular dynamics simulation, the α_3 domain of one FHC (orange) binds to another FHC (yellow/purple/lavender) in a manner resembling the binding of β_2m in the crystal structure of peptide-loaded Kᵇ (**B**, PDB 3P9L).

A

B

Figure 2 Supplement 1: Micropattern negative antibody control.
The fluorescent signal of the bleached area in an Alexa Fluor 488-labeled anti-HA antibody micropattern remains low. A pattern element was bleached (**A**), and fluorescence recovery was quantified over a time of 2 min (**B**). Pattern element distance, 2 μm.

Figure 2 Supplement 2: Binding and dissociation of HA-K^b-GFP to and from antibody micropattern is negligible.

STF1 cells expressing HA-K^b-GFP (a construct with dual tags: HA at the N terminus, and GFP at the C terminus) were seeded onto anti-HA antibody patterns, GFP was bleached over one pattern element, and fluorescence recovery was measured over time (red curve). The data sets of HA-K^b + K^b-GFP samples of Figure 2E with or without peptide are also shown as a comparison. Error bars represent SEM (*n* = 9 cells for HA-K^b-GFP cells).

Figure 2 Supplement 3: Dissociation of K^b-GFP from micropattern elements is negligible.
GFP bleaching over one pattern element (yellow frame) does not affect the fluorescence of neighboring pattern elements (blue, red, orange frames) over a time period of 250 s.

Dirscherl *et al.*
Figure 3 Supplement 1

Figure 3 Supplement 1: Density of molecules in SMT and SMCT experiments.
Mean initial density of localized GFP-K^b molecules in each detection channel for the different conditions.
Each data point corresponds to the analysis from one single cell with ≥14 cells measured for each con-dition.
*, p ≤ 0.05 by a two-sample Kolmogorov-Smirnov test.

Figure 3 Supplement 2: Random diffusion of mobile fraction.
Mean squared displacement (MSD) analysis of GFP-K^b molecules under different conditions and linear fit.
Mean of the MSD values obtained from 10 cells, error bars indicate standard deviations (SD).

Figure 3 Supplement 3: Similar diffusion coefficient of FHC and MHC I dimers.
Comparison of the diffusion constants of the entire mobile fractions and the dimers only for of GFP-K^b molecules in the absence of peptide and in the presence of peptide and the tandem nanobody crosslinker (CL). ***, p ≤ 0.001 by a two-sample Kolmogorov-Smirnov test.

Figure 4 Supplementary Table 1: Results from native mass spectrometry.
Experimental masses (m_{exp}) of the different α_3 domain protein species were determined from at least three independent mass spectrometry measurements. They are listed together with the respective values for standard deviation (SD) along with the theoretically calculated molecular weight (M).

mass species	M [Da]	M_{exp} [Da] ± SD
α_3 monomer	11,475	11,472.7 ± 0.4
α_3 monomer$_{short}$	10,623	10,0621 ± 1
α_3 dimer	22,950	22,947 ± 1
α_3 dimer$_{short}$	22,098	22,094 ± 1

A

B

Figure 5 Supplement 1: MD simulation snapshots of an FHC dimer.
A, The α_3 domain of one K^b molecule is depicted in orange, the α_3 domain of the other K^b molecule is depicted in lavender. The α_1/α_2 superdomains are shown in yellow/purple for both FHCs. A favorable interaction of the α_3 domain of one K^b molecule (orange) with the α_1/α_2 superdomain (yellow/purple) of another K^b molecule (with α_3 domain in lavender) remains stable after 200 ns (left) and 400 ns (right). The panels represent the conformations sampled at 200 ns and 400 ns time points indicating the exposure and high mobility of the α_1/α_2 superdomain of one of the HCs.
B, Root-mean-square deviation (RMSD) of the protein backbone with respect to the starting structure vs. simulation time for the simulation of the K^b HC/β_2m heterodimer (black line) and for the simulation of the predicted FHC dimer (HC/α_3 domain, red line) is low. In the FHC dimer, the relative mobility of the α_1/α_2 superdomain of the second HC (green line) after superposition of the trajectory on the first HC is high.