1

Dissociation of β_2m from MHC Class I
Triggers Formation of Noncovalent, Transient Heavy Chain Dimers

Cindy Dirscherl[1,x], Sara Löchte[2,x], Zeynep Hein[1], Janine-Denise Kopicki[3], Antonia Regina Harders[1], Noemi Linden[1], Andreas Karner[9], Johannes Preiner[9], Julian Weghuber[4], Maria Garcia-Alai[5,6], Charlotte Uetrecht[3,7], Martin Zacharias[8], Jacob Piehler[2,*], Peter Lanzerstorfer[4,*], and Sebastian Springer[1,*]

[1]Department of Life Sciences and Chemistry, Jacobs University Bremen, Germany;

[2]Department of Biology and Center for Cellular Nanoanalytics, Osnabrück University, 49076 Osnabrück, Germany;

[3]Heinrich Pette Institute, Leibniz Institute for Experimental Virology, Hamburg, Germany;

[4]University of Applied Sciences Upper Austria, 4600 Wels, Austria;

[5]European Molecular Biology Laboratory, Hamburg Outstation, Hamburg, Germany;

[6]Centre for Structural Systems Biology, Hamburg, Germany;

[7]European XFEL, Schenefeld, Germany;

[8]Physics Department, Technical University of Munich, Garching, Germany;

[9]University of Applied Sciences Upper Austria, 4020 Linz, Austria.

[x]Equal contribution.

* To whom correspondence should be addressed:

Sebastian **Springer**	Peter **Lanzerstorfer**	Jacob **Piehler**
Department of Life Sciences and Chemistry	School of Engineering	Department of Biology/Chemistry
Jacobs University Bremen	University of Applied Sciences Austria	Osnabrück University
Campus Ring 1	Stelzhamerstraße 23	Barbarastr. 11
28759 Bremen, Germany	A-4600 Wels, Austria	49076 Osnabrück, Germany
s.springer@jacobs-university.de	peter.lanzerstorfer@fh-wels.at	piehler@uos.de
+49 421 2003243	+43 5 080444402	+49 541 969 2800

1 **Abstract**

At the plasma membrane of mammalian cells, major histocompatibility complex class I

molecules (MHC-I) present antigenic peptides to cytotoxic T cells. Following the loss of the

peptide and the light chain beta-2 microglobulin (β_2m), the resulting free heavy chains

(FHCs) can associate into homotypic complexes in the plasma membrane. Here, we investi-

gate the stoichiometry and dynamics of MHC-I FHCs assemblies by combining a

micropattern assay with fluorescence recovery after photobleaching (FRAP) and with single

molecule co-tracking. We identify non-covalent MHC-I FHC dimers mediated by the α_3 do-

main as the prevalent species at the plasma membrane, leading a moderate decrease in the

diffusion coefficient. MHC-I FHC dimers show increased tendency to cluster into higher or-

der oligomers as concluded from an increased immobile fraction with higher single molecule

co-localization. In *vitro* studies with isolated proteins in conjunction with molecular docking

and dynamics simulations suggest that in the complexes, the α_3 domain of one FHC binds to

another FHC in a manner similar to the β_2m light chain.

Significance Statement

MHC class I molecules are cell surface transmembrane proteins with key functions in adaptive immunity against viral infections. The spatiotemporal organization of fully assembled MHC I at the cell surface and its function with respect to trans-interactions with T and NK cells has been studied in detail. By contrast, the consequences of peptide and β_2m dissociation yielding to formation of free heavy chains (FHC) have remained unclear. We have discovered that class I free heavy chains form distinct non-covalent dimers at the cell surface rather than non-specific clustering, and we have identified a dimerization interface mediated by the α_3 domain. We propose that these non-covalent dimers are the basis of distinct signaling and endocytic sorting of MHC I FHC. This is to be explored in further work.

1 Introduction

Major histocompatibility class I molecules (MHC-I) fulfil central tasks of the adaptive

immune response against infections and malignancies by presenting antigenic peptides to T

cells (Comber and Philip, 2014; Kaufman, 2018; Townsend and Bodmer, 1989). MHC-I

heterotrimers consist of the polymorphic transmembrane heavy chain (HC), the non-

polymorphic soluble light chain beta-2 microglobulin (β_2m), and the peptide (Townsend et

al., 1990, 1989). In addition to this trimer, two more states of MHC-I occur at the cell sur-

face, the 'empty' HC/β_2m heterodimer that lacks peptide (Ljunggren et al., 1990; Sebastián

Montealegre et al., 2015), and the monomeric 'free' heavy chain (FHC) (Edidin et al., 1997;

Geng et al., 2018). Since the binding of peptide and β_2m is cooperative (Elliott et al., 1991;

Gakamsky et al., 1996), HC/β_2m heterodimers are conformationally unstable, and loss of

peptide leads to the rapid formation of FHCs (schematic in **Figure 1A**) and to the subsequent

endocytic removal of FHCs by a sorting mechanism that is not understood (Sebastián Mon-

tealegre et al., 2015). For FHCs present at the cell surface, important regulatory functions

mediated by homo- and heteromeric interactions in *cis* and *trans* have been proposed (Arosa

et al., 2021, 2007; Campbell et al., 2012), which suggest defined spatiotemporal organization

and dynamics of FHC in the plasma membrane. Indeed, clustering and covalent dimerization

of MHC-I have been identified using a variety of approaches including recombinant proteins

and live cells (Allen et al., 1999; Antoniou et al., 2011; Armony et al., 2021; Baia et al.,

2016; Blumenthal et al., 2016; Bodnar et al., 2003; Capps et al., 1993; Chakrabarti et al.,

1992; Fassett et al., 2001; Ferez et al., 2014; Lu et al., 2012; Makhadiyeva et al., 2012; Matko

et al., 1994; Triantafilou et al., 2000); these represent complexes of different and largely un-

clear composition, size, and type of intermolecular bonding.

Recently, we have achieved direct detection of homomeric FHC interactions in the intact

plasma membrane by means of a live-cell two-hybrid micropattern assay (**Figure 1B, C**)

(Dirscherl et al., 2018): micrometer-sized patterns of anti-hemagglutinin tag (HA) monoclo-

nal antibody are printed onto glass coverslips (Schwarzenbacher et al., 2008; Sevcsik et al.,

2015). Onto these micropatterns, cells are seeded that express two different MHC-I HC con-

structs: one construct has an N-terminal (extracellular) HA tag and thus, while diffusing lat-

erally in the plasma membrane, is captured into the printed antibody pattern. The other has no

HA tag but a C-terminal (intracellular) green fluorescent protein (GFP) fusion domain (**Fig-**

ure 1B). Interaction of GFP-tagged HC with micropatterned HA-HC is detected by an in-

creased GFP fluorescence of the pattern elements (**Figure 1C**). ==The system allows the obser-==

==vation of different defined conformational states of class I:== with TAP2 (transporter associated

with antigen processing)-deficient fibroblasts (which cannot transport peptide into the endo-

plasmic reticulum (ER)), empty HC/β_2m heterodimers are present at the cell surface. By add-

ing peptide, shifting cells to 25 °C, or incubating at 37 °C, we therefore can accumulate

trimers, HC/β_2m heterodimers, or FHCs, respectively, at the cell surface (**Figure 1B** and de-

scribed below). These experiments revealed that formation of homomeric MHC-I only takes

place in the absence of β_2m, *i.e.*, only between FHCs.

We now have uncovered the molecular principles that govern such homomeric MHC-I

FHC association in the plasma membrane. Cell micropatterning in conjunction with fluores-

cence recovery after photobleaching (FRAP) was used to probe dynamics, stability, and

prominence of FHC complexes. We directly demonstrate FHC association in the plasma

membrane under physiological conditions by using real-time single-molecule tracking (SMT)

and co-tracking (SMCT). Surprisingly, we find that FHCs transiently associate into non-

covalent dimers with lifetimes in the sub-second range. Based on our findings that the

HC/HC complexes contain no β_2m and that the α_3 domain of the HC is sufficient for dimeri-

zation *in vitro* and in cells, we propose a molecular model structure of MHC-I HC/HC dimers

supported by *in silico* docking and molecular dynamics (MD) simulation. Our findings clear-

ly differentiate the cell surface dynamics and properties of empty FHCs from the peptide-

loaded trimers of MHC-I, pointing to the formation of structurally well-defined HC/HC

homodimers that may be responsible for distinct endosomal trafficking and other biological

functions previously ascribed to FHCs.

Results

A micropattern assay reveals non-covalent association of MHC class I free heavy chains

In STF1 cells, which are fibroblasts that cannot load MHC-I with peptides due to a defi-
ciency in the TAP peptide transporter, incubation at 25 °C accumulates murine HC/β_2m het-
erodimers at the plasma membrane, since at that temperature, dissociation of β_2m and subse-
quent endocytosis are inhibited (Day et al., 1995; Ljunggren et al., 1990; Sebastián Mon-
tealegre et al., 2015). When the temperature is shifted to 37 °C, β_2m dissociates to yield
FHCs, and lateral *in cis* interactions between HA-K^b and K^b-GFP (both hybrids of the HC of
the murine MHC-I molecule H-2K^b) become visible in the micropattern two-hybrid assay,
where the GFP fluorescence arranges in the shapes of the antibody micropattern (**Figure 1C**).
When cognate peptide is added to the cells, HC/β_2m/peptide trimers are stable and do not
associate with each other (**Figure 2B**) (Dirscherl et al., 2018).

Up to six different MHC-I allotypes are present at the cell surface of human and murine
cells. To date, though, an interaction of different MHC-I allotypes in the same plasma mem-
brane has not been shown. We therefore tested for such heterotypic interactions between K^b

and H-2D^b (D^b). Just as for K^b/K^b, also K^b and D^b FHCs (at 37 °C) interacted with each other, whereas β$_2$m-bound heterodimers at 25 °C did not (**Figure 1D**). Hence, our micropattern assay confirmed that formation of heterotypic FHC interactions is possible.

We next asked which molecular characteristics are required for HC/HC interactions, and we first tested whether the HCs are held together by cytosolic disulfide bonds as previously suggested for certain allotypes under oxidizing conditions (Baía et al., 2016; Capps et al., 1993; Makhadiyeva et al., 2012). We replaced the single cysteine in the cytosolic tail of K^b-GFP (residue 332) with a serine and tested for association between this mutant and wild type HA□K^b (**Figure 1E**). K^b(C332S)-GFP and wild type HA-K^b interacted in over 90% of cells, just as the wild type K^b-GFP, demonstrating that cysteine 332 is not required for HC/HC interactions.

To test more generally for any disulfide bonding in MHC-I interactions, we immunoprecipitated HA-K^b molecules from STF1 cell lysate with an anti-HA antibody. By non-reducing gel electrophoresis, covalent homodimers of K^b were not observed, while the well-described disulfide-linked homodimers of the human MHC-I allotype HLA-B*27:05 (Dangoria et al., 2002) were readily detected (**Figure 1F**). We conclude that the K^b FHCs are non-covalently associated and that they do not undergo intramolecular disulfide bonding under the conditions of our assay.

Since the cysteines in the cytosolic tail of K^b are not required for HC/HC interaction, we hypothesized that FHCs non-covalently associate via their extracellular domains, and that the loss of β$_2$m is a prerequisite for HC/HC interaction. As anticipated, a disulfide-stabilized variant of K^b–GFP (Y84C/A139C), in which β$_2$m dissociation is dramatically decreased (Hein et al., 2014), did not interact with HA–K^b (**Figure 1G**). In agreement with our earlier finding that covalent attachment of β$_2$m to the HC also prevents HC/HC association (Dirscherl et al.,

2018), this result demonstrates that HC/β_2m heterodimers do not interact with other HC/β_2m

heterodimers, nor with FHCs. ==These findings demonstrate that the MHC class I complexes==

==observed in our system are non-covalent in nature and comprise free heavy chains, devoid of==

==β_2m and peptide.==

Free heavy chains associate on TAP-proficient cells

So far, we used TAP2-deficient STF1 cells to obtain homogeneous populations of free

heavy chains at the cell surface. Since MHC-I HC/β_2m heterodimers and FHCs both exist at

the cell surface of wild type cells (Day et al., 1995; Ljunggren et al., 1990; Ortiz-Navarrete

and Hammerling, 1991, p.), we next tested whether FHCs also associate in cells with wild

type TAP function (schematic in **Figure 2A**). Following the 25 °C to 37 °C temperature shift

that triggers FHC formation, we performed the same anti-HA antibody two-hybrid

micropatterning assay with HA-K^b and K^b-GFP as above (in **Figure 1B and C**). Again, the

punctate GFP fluorescence signifies the recruitment of K^b-GFP fusion to the printed patterns

of the anti-HA antibodies, mediated by the HA-K^b fusion. We quantified K^b HC/HC interac-

tion over time in STF1/TAP2 cells, which are able to load class I molecules with peptides, by

GFP fluorescence contrast analysis between pattern elements and interspaces. Interactions

increased in TAP2-positive and in TAP2-deficient cells with the same dynamics and reached

a maximum after ca. 60 min of incubation at 37 °C (**Figure 2B, C**), with the kinetics likely

governed by the dissociation of β_2m from the HC/β_2m heterodimers, which occurs on this

timescale (Sebastián Montealegre et al., 2015). As expected, HC/β_2m/peptide trimers, stabi-

lized by cognate SIINFEKL peptide, did not show any interaction (**Figure 2B**). This experi-

ment confirms that HC/HC interactions ==indeed== take place at the surface of TAP2-proficient

cells. ==In addition, these experiments again confirm our earlier observations== (Dirscherl et al.,

2018) that the HC/HC interaction occurs only between FHCs: when dissociation of β_2m is inhibited by low temperature (**Figure 1 DEG**) or by added peptide (**Figure 2 BD**), the interaction does not occur. The same experiments also strongly suggest that the recruitment of the K^b-GFP fusion to the pattern elements specifically reports on the interaction between MHC class I molecules (since it depends on their conformation) and is not due to an attraction between K^b-GFP and other surface proteins, e.g., adhesion molecules, that might have been attracted to the plasma membrane above the pattern elements.

FHC association slows down cell surface diffusion of MHC-I

Since dimers and oligomers of FHCs have more transmembrane domains (TMDs) than single HC/β_2m/peptide complexes, we hypothesized that they should diffuse more slowly in the plasma membrane (Gambin et al., 2006; Wilmes et al., 2015a). We therefore carried out total internal reflection fluorescence (TIRF) fluorescence recovery after photobleaching (FRAP) experiments on STF1 cells with HA-K^b and K^b-GFP on surfaces with or without micropatterns (**Figure 2D**) and quantified the recovery dynamics (**Figure 2E**). Diffusion constants of FHCs (– peptide) were significantly decreased as compared to HC/β_2m/peptide complexes (+ peptide; **Figure 2F**), suggesting that FHCs form complexes. The moderate decrease of ≈40% is in line with the formation of dimers rather than clustering into larger complexes. Since this effect occurred both in cells seeded on the pattern elements (on pattern) and in cells seeded off the micropatterns (off pattern), we conclude that the micropatterns are not required for HC/HC interactions, i.e., that the interactions are not an artefact of the two-hybrid micropattern assay. This statement is also supported by the co-immunoprecipitation of MHC-I molecules in the absence of micropatterns (Dirscherl et al., 2018; Triantafilou et al., 2000). Diffusion constants of HC/β_2m/peptide trimers on pattern and off pattern were not

significantly different, which demonstrates that the antibody micropatterns themselves do not impede the diffusion of plasma membrane proteins.

In the FRAP experiments, a portion of K^b-GFP appeared immobile on a timescale of seconds as evidenced by the incomplete fluorescence recovery (**Fig. 2E**). This immobile fraction is significantly higher for FHCs than for HC/β_2m/peptide trimers (**Fig. 2G**). This observation is readily explained for K^b-GFP bound to HA-K^b molecules immobilized within micropatterns, but it is remarkable for the FHCs on cells outside of the micropattern (gray column in **Fig. 2G**). There, it may be ascribed to the formation of large oligomers, and/or to the association of FHCs with immobile structures, such as the cytoskeleton and/or due to sequestration in endocytic membrane compartments (Bondar et al., 2020; Ibach et al., 2015; Mylvaganam et al., 2018; Vámosi et al., 2019). From the FRAP curves, the exchange rate of the freely diffusing pool of K^b-GFP into and out of the bleached regions of interest (ROIs) is obtained through a bi-exponential fit as the slow recovery rate (k_{slow}) (**Figure 2H**, solid bars); the fast recovery rate k_{fast} represents free diffusion (Sprague and Mcnally, 2005)(see the Materials and Methods). On the pattern elements, the k_{slow} of FHCs was much smaller than in cells outside the patterns, suggesting a half-time of dissociation from the pattern-bound immobile associations of about 140 seconds. Likewise, the fluorescence signal of pattern elements in the immediate vicinity of the bleached region remained unaltered, indicating that no detectable exchange of K^b-GFP between the enriched HA-K^b regions occurs on the second timescale (**Figure 2S3**). This very slow exchange of K^b-GFP molecules associated on pattern elements suggests either multiple association and dissociation events in a small radius due to densely immobilized binding partners, or a very stable association of the immobile FHCs (see the discussion). Taken together, the data from **Figure 2** show reduced diffusion rates of K^b

FHCs at the cell surface compared to HC/β_2m/peptide complexes, which is most easily ex-
plained by homotypic association of the former.

Transient FHC dimerization directly observed at single molecule level

We therefore turned to directly visualizing FHC diffusion and interaction in the plasma membrane under physiological conditions by single molecule tracking (SMT) and co-tracking (SMCT) (Moraga et al., 2015; Sevcsik et al., 2015; Wilmes et al., 2020a). STF-1 cells transiently expressing K^b with its N-terminus fused to monomeric GFP (GFP-K^b) were imaged by TIRF microscopy. The GFP tag was labeled with photostable fluorescent dyes by using equal concentrations of anti-GFP nanobodies conjugated to either ATTO Rho11 (^{Rho11}NB) or ATTO 643 (^{ATTO643}NB), with one nanobody binding one GFP molecule, ensuring selective imaging of K^b in the plasma membrane and tracking with high fidelity (**Figure 3A**). Imaging was performed at 37 °C to induce formation of FHC in the absence of the peptide, while replenishment of GFP-K^b/β_2m dimers from the ER was inhibited by Brefeldin A (BFA).

After labeling with ^{Rho11}NB and ^{ATTO643}NB, individual K^b subunits were observed at densities of <1 molecule/µm² diffusing randomly in the plasma membrane (**Supplementary Movie S1, Supplementary Figure S3A**). Co-tracking analysis revealed homomeric interaction of K^b FHCs in the absence of peptide, whereas these events were very rare for peptide-loaded K^b (**Figure 3B-D**). As a positive control for the formation of homodimers, a crosslinker based on the tandem anti-GFP nanobody LaG16 (Fridy et al., 2014), which recognizes a different epitope than the labeled nanobodies, was added to the medium. While dimerizing of K^b via the GFP tag resulted in a nominal fraction of ≈38% (median) co-locomoting molecules, only ≈3% of the FHCs were found associated, indicating weak interaction of FHCs at these

low cell surface expression levels of K^b ($\approx$1 molecule/μm² in total, corresponding to <5000 molecules/cell).

From the trajectory analysis, we determined the diffusion coefficients of individual MHC-I molecules by mean square displacement analysis (**Figure 3E, Supplementary Figure S3B**). In the absence of peptide, diffusion of FHCs was significantly slower than in the presence of the peptide. This decrease in single molecule diffusion coefficients supports interaction of FHCs in the plasma membrane. A similar decrease in the diffusion coefficient was observed upon dimerization of peptide-loaded K^b with the tandem nanobody, suggesting that FHCs associate into dimers. Indeed, largely identical diffusion coefficients were found for the fraction of molecules identified as dimers by SMCT (**Supplementary Figure S3C, Table 1**). Strikingly, the diffusion constants obtained by SMT for peptide-loaded K^b and for FHC from SMT are consistent with those obtained by FRAP experiments under the same conditions (**Figure 2E, Table 1**), highlighting that similar phenomena are being probed by these complementary techniques. Tracking analysis revealed a slightly elevated immobile fraction of ~20% for K^b in the absence of the peptide compared to $\approx$17% for peptide-loaded K^b (**Figure 3F**). Again, a similar effect was observed for artificially dimerized, peptide-loaded K^b, corroborating the dimeric stoichiometry of associated FHCs. Importantly, the "immobile fraction" in SMT refers to much shorter time and length scales as compared to FRAP (cf. methods section), and therefore, the absolute numbers are not comparable. Within the immobile fraction, a higher level of FHC was found associated, as compared to the mobile fraction (**Figure 3G**). Overall, the single molecule diffusion analyses suggest that FHC dimers have an increased propensity to be immobilized at the plasma membrane, probably by clustering that may be related to endocytosis. Interestingly, this feature is reproduced by artificial dimerization of intact MHC-I by a crosslinker, suggesting that FHC dimerization is a switch regulating its cell surface dynamics.

SMCT analysis moreover revealed dissociation of FHC homomers, confirming its transi-

ent nature (**Figure 3B, Supplementary Movies S2-5**). We estimated the lifetime of FHC

complexes from co-trajectory length histograms. Fitting of an exponential decay revealed a

significantly shorter lifetime for FHC co-trajectories compared to the average co-tracking

lifetime determined for stably NB-crosslinked K^b, which is limited by co-tracking fidelity and

photobleaching (**Figure 3H**). Taken together, the SMT and SMCT results of **Figure 3** direct-

ly confirm a transient homotypic HC/HC interaction in the plasma membrane that leads to

reduced diffusion velocity and immobilization, in line with the FRAP experiments. Since the

degree of association was low, and higher-order oligomerization was not observed, we pro-

pose that HC/HC interactions are weak and have dimeric stoichiometry.

The α_3 domain of K^b forms dimers and is sufficient for FHC association

We next explored the molecular mechanism of HC/HC interaction. Since cytosolic

cysteines are not involved (**Figure 1E, F**) and dissociation of β_2m is necessary (**Figure 1G**),

we hypothesized that HC/HC interactions involve the extracellular portion of the HCs. To

determine more precisely which domains are involved, we used STF1 cells in the

micropattern assay that expressed constructs of K^b that lacked the α_1/α_2 domain. Remarka-

bly, α_3-GFP showed excellent copatterning with HA-K^b-RFP (fused to red fluorescent pro-

tein; **Figure 4A**) and with HA-α_3 (**Figure 4B**), demonstrating that isolated α_3 domains can

bind to each other on the surface of live cells. Indeed, the isolated, soluble α_3 domain of K^b,

when refolded *in vitro*, showed (after testing of its folded state by fluorescence spectroscopy,

Supplementary Figure S4C) significant formation of homodimers in size exclusion chroma-

tography (SEC; **Figure 4C**), while no prominent higher-order complexes were observed.

These α_3 homodimers were not linked by disulfide bonds as shown by nonreducing gel elec-

trophoresis (**Figure 4D**), just like the interaction of full-length FHC in the plasma membrane (**Figure 1F**). Given the 1:1 peak ratio observed at $10\,\mu M$ protein concentration, a binding affinity of 10 to $20\,\mu M$ can be estimated for this interaction. The non-covalent α_3 homodimers were also detectable by native mass spectrometry (MS) in a concentration-dependent manner, and they easily separated into the monomers when the collision cell voltage was increased in an MS/MS experiment, suggesting a low-affinity interaction (**Figure 4E-G, Supplementary Figure S4A**). These biochemical results, together with the observations on the α_3 constructs above, suggest that FHC dimerization in the plasma membrane is at least partly based on the intrinsic affinity between the α_3 domains. To investigate whether the α_3 domain is indeed required for FHC dimerization, we performed a co-patterning experiment with an HA-K^b fusion protein that lacked the α_3 domain (HA-$K^b(\Delta\alpha_3)$-RFP) and K^b-GFP and found the interaction significantly reduced, but still measurable (**Supplementary Figure S4B**), which suggests that the α_1/α_2 domain is also involved in the interaction.

A molecular model of the K^b HC dimer

To image a possible arrangement of the K^b heavy chains in a dimer, we used molecular docking. A starting geometry was obtained by placing the α_3 domain of one K^b HC in the same position as the $\beta_2 m$ in the original K^b HC/$\beta_2 m$ heterodimer and adjusting the α_1/α_2 superdomain to avoid sterical overlap. The structure was then energy-minimized and further refined by Molecular Dynamics (MD) simulations (**Figure 5A**). During MD simulations of $400\,ns$, no signs of dissociation were observed, and a stable root-mean-square deviation (RMSD) from the start structure was reached (**Supplementary Figure S5**). In the model, the α_3 domain of one FHC substitutes for $\beta_2 m$, binding both the α_3 domain and the α_1/α_2 superdomain of another FHC with a buried interface area (BSA) of $2480\,\mathring{A}^2$, which is compa-

rable to the BSA of 2740 Å^2 between β_2m and the HC in heterodimers (**Figure 5B**, bottom),

and to other stable protein complexes (Bahadur and Zacharias, 2008). Similar results were

obtained with other arrangements of the two heavy chains (not shown).

Discussion

Interactions of MHC-I FHC with FHC of the same allotype, of different allotypes, and

even with other cell surface proteins have been proposed to play an important role in regulat-

ing adaptive and innate immune responses (Arosa et al., 2021, 2007; Campbell et al., 2012),

but the molecular principles that govern FHC interactions have remained unclear. By com-

bining live cell interaction and diffusion analysis using cell micropatterning and FRAP as

well as SMT and SMCT, we have shown here that, after losing β_2m, murine H-2K^b MHC-I

molecules at the cell surface interact in a homotypic and heterotypic manner to form dimers,

which are transient with a stability on the second timescale. The α_3 domains of the FHCs

alone are sufficient for such interactions, but it is conceivable that the α_1/α_2 domain is also

involved: in addition to the α_3-α_3 interaction, the α_3 domain might also bind to the α_1/α_2 do-

main of the other protein, or the α_1/α_2 domains of the two proteins might interact with each

other. We cannot currently distinguish these possibilities.

While we observed a somewhat increased tendency of FHCs to form higher oligomers, our

data surprisingly identify non-covalent FHC dimers as the prevalent species at the cell sur-

face. We analyzed the diffusion properties of FHCs and HC/β_2m/peptide trimers at different

cell surface expression levels by FRAP and SMT. FRAP experiments were carried out at high

surface densities, which probably exceeded the typical endogenous cell surface expression

levels of 10^5 copies/cell (Spack and Edidin, 1986). By contrast, SMT was performed at densi-

ties of ≈1 molecule/μm², *i.e.*, <5000 molecules/cell. Despite these differences in expression

1. levels, we found highly consistent diffusion coefficients D by both FRAP and SMT, which

2. revealed moderately decreased mobility upon dissociation of β_2m (**Table 1**). However, under

3. both conditions, the decrease in D was ≈40-50%, which perfectly agrees with the change in D

4. upon dimerizing intact MHC-I by the tandem nanobody (**Figure 3E**). Likewise, we and oth-

5. ers have previously observed very similar decreases of 30-50% upon dimerization of cell

6. surface receptors (Ho et al., 2017; Low-Nam et al., 2011; Moraga et al., 2015; Richter et al.,

7. 2017; Váradi et al., 2019; Wilmes et al., 2015a, 2020a), corroborating that the mobile HC/HC

8. complexes are mostly dimers.

9. Direct detection of FHC interaction by SMCT corroborated transient, non-covalent dimer-

10. ization and only minor clustering into higher oligomers. Estimated lifetime of the non-

11. covalent FHC/FHC dimer ($t_{1/2} = 220 \pm 150$ ms, **Figure 3G, H**) were well within the measura-

12. ble range, *i.e.*, substantially below the apparent half-life obtained for the quasi-irreversibly,

13. nanobody-crosslinked control ($t_{1/2} > 10$ s), which defined the limit of co-tracking fidelity. To-

14. gether with the immunoprecipitation experiments (**Figure 1F**) and micropatterning of FHC

15. lacking free cysteines (**Figure 1E**), these observations clearly establish that the FHC mole-

16. cules at the plasma membrane are not covalently linked. The short half-life and the non-

17. covalent monomer-dimer equilibrium observed by size exclusion chromatography at higher

18. micromolar concentration for the isolated α_3 domain (**Figure 4CD**), as well as the mass spec-

19. trometry data (**Figure 4E-G**), point to a low-affinity interaction with a dimerization K_d of

20. perhaps 10 to 20 µM. Previous quantitative studies on affinity-dimerization correlation of

21. heterodimeric cytokine receptors (Wilmes et al., 2015a) would predict efficient FHC dimeri-

22. zation at physiological densities above 10 molecules/µm², which is in line with the endoge-

23. nous MHC-I expression level. The relatively low level of dimerization we observed by

24. SMCT can be rationalized with the high background of endogenous MHC-I that are unla-

beled and invisible to us. In line with this observation, the decrease of the diffusion coefficient in SMT was much more prominent than the dimer fraction identified by SMCT. This interpretation is in line with the observation that more efficient interaction was observed at the elevated cell surface expression levels used in micropatterning experiments.

The weak and transient nature of HC/HC dimerization seen by SMT is similar to the nature of cell surface protein-protein association measured in other systems (Lin et al., 2014). It does not conflict with the clear and stringent patterning of the GFP fusion in the micropattern/FRAP experiments. In the latter, the concentration of HA-tagged bait HCs is considerably higher due to their immobilization by the antibodies, which prevents their endocytosis, and this probably creates an affinity matrix for the GFP fusions that can retain them for many seconds due to rebinding events. This also suggests that in endosomes, whose internal volume is very small, HC/HC interaction might be potentiated due to the increase in the concentration of the monomers compared to the plasma membrane.

In line with the observation of non-covalent dimerization in the plasma membrane, we obtained a robust structural model of a self-contained K^b FHC homodimer. The atomistic model (**Figure 5**) was derived from the experimental findings that dissociation of β_2m is required, that the α_3 domains are sufficient, and that a direct α_3/α_3 interaction exists (**Figures 1E, 4A-C**). We propose that β_2m dissociation exposes a binding site on the FHC for the α_3 domain of another FHC. However, other arrangements of the two FHCs in a dimer are theoretically possible, and only experimental data will give a definitive answer.

Several findings in the literature are consistent with the formation of complexes of MHC class I HCs in the absence of β_2m and peptide; this applies both to the 'classical', or class Ia, proteins HLA-A/B/C (and in mouse: H-2D/K/L) as well as to the 'non-classical', or class Ib, protein HLA-F (Armony et al., 2021; Bodnar et al., 2003; Chakrabarti et al., 1992; Matko et

al., 1994; Triantafilou et al., 2000). Still, it is important to differentiate these HC/HC dimers from class I associations described elsewhere (partially reviewed in (Arosa et al., 2021, 2007; Campbell et al., 2012)), namely homo- and heterotypic HC/β_2m/peptide trimers that are covalently dimerized via disulfide bonds in their cytosolic tails (Capps et al., 1993; Makhadiyeva et al., 2012) and that may play a role in binding the LILRB NK cell receptor (Baia et al., 2016); the non-covalent nano- and microscale clusters of HC/β_2m/peptide trimers (not detected in our system) that may stem from the fusion of exocytic vesicles with the plasma membrane and that may play a role in TCR recognition (Blumenthal et al., 2016; Ferez et al., 2014; Fooksman et al., 2006; Lu et al., 2012); the macroscopic 'clusters' of class I molecules at the signaling interface between cells (Fassett et al., 2001) and the covalent dimers of the HLA- B*27:05 heavy chain that are linked by disulfide bonds through Cys-67 (Chen et al., 2017), though our non-covalent HC/HC dimers may be a precursor to the formation of the latter.

The unexpected discovery that FHCs non-covalently associate into defined dimers allows exciting hypotheses of their distinct functional properties. FHC dimers might be responsible for the immunomodulatory functions of cell surface heavy chains, *i.e.,* the stabilization of MHC-I trimers to assist T cell activation (Geng et al., 2018; Schell, 2002) and direct binding of FHCs to receptors on other cells (for example, FHCs of HLA-F binding to activating receptors on NK cells (Dulberger et al., 2017, 2017; Goodridge et al., 2013)). Furthermore, dimerization of FHCs might enhance endocytosis in order to remove the non-functional FHCs, which themselves cannot activate T cells and are known to be short-lived (Mahmutefendic et al., 2011; S. Montealegre et al., 2015); alternatively or additionally, such associated HCs may bind to other proteins *in cis* and promote their removal from the plasma membrane. Such endocytic removal might be achieved by altered endosomal routing, since

the local density of membrane proteins in endosomes is higher than at the plasma membrane,

and thus, efficient dimerization of MHC-I FHCs is expected. In such a scenario, even transi-

ent oligomerization in endosomes might prevent the return of internalized MHC-I FHCs to

the cell surface (S. Montealegre et al., 2015). Taken together, the presence of non-covalent,

transient FHC dimers points to exciting new aspects in the regulation of MHC-I functions

with much potential for further investigation.

Materials and Methods

Cells and cell lines

TAP-deficient human STF1 fibroblasts (kindly provided by Henri de la Salle, Etablissement de Transfusion Sanguine de Strasbourg, Strasbourg, France) were cultivated at 37 °C and 5% CO_2 in Earle's minimum Essential Medium (MEM) with stable glutamine supplemented with 10% fetal bovine serum (FBS), non-essential amino acids and HEPES buffer without addition of antibiotics.

Genes, vectors, and gene expression

HA-K^b and K^b-GFP constructs were described previously (Dirscherl et al., 2018). HA-K^b carries an influenza hemagglutinin (HA) tag at the N terminus of the full-length murine H-2K^b, whereas K^b-GFP carries a GFP domain at the C terminus of H-2K^b. The D^b-GFP construct is analogous to the K^b-GFP construct. The α_3-GFP construct consists of the H-2K^b signal sequence and residues 204-369 of H-2K^b, including transmembrane and cytosolic domains. The HA-K^b($\Delta\alpha_3$)-RFP construct, compared to HA-K^b, lacks residues 204-294 and carries an additional RFP (red fluorescent protein) taken from pcDNA3-mRFP (see the key resources table). The GFP-K^b construct used in single-molecule imaging consists of a signal sequence and GFP fused to the N terminus of H-2K^b itself lacking a signal sequence. Amino acid numbering of K^b in this paragraph includes the signal sequence.

Stable cell lines were generated by lentiviral transduction as described (Hein et al., 2014), and transient transfection was achieved by electroporation (Garstka et al., 2007) or by calcium phosphate precipitation (Graham and van der Eb, 1973) as described.

Micropattern assay

Photolithography. Silicon master molds were prepared by semiconductor photolithogra-

phy as described previously (Dirscherl et al., 2017).

PDMS stamps and Antibody Patterns. PDMS stamps were generated from basic elasto-

mer and curing agent (Sylgard 184 Silicone Elastomer Kit) as described previously (Dirscherl

et al., 2017).

Patterning cell surface proteins. Coverslips with antibody pattern were placed into sterile

6-well plates. Cells were immediately seeded as indicated at a concentration of ca. 50 000

cells per well. Usually, cells were incubated for 4-6 hours at 37 °C for adhesion and then

shifted to 25 °C to accumulate MHC-I molecules at the cell surface. Samples were then kept

at 25 °C to increase cell surface heterodimer levels or shifted back to 37 °C for 3-4 hours to

induce FHCs by dissociation of β_2m.

Dyes. Purified antibodies were labeled with Alexa Fluor 647 NHS ester (Thermo Fisher

Scientific, Darmstadt, Germany) according to the manufacturer's protocol.

Peptides. The K^b-specific peptide SL8 (SIINFEKL in the single-letter amino acid code)

was synthesized by GeneCust (Ellange, Luxemburg) and emc microcollections (Tübingen,

Germany) and purified by HPLC (90% purity).) Peptides were added to the cells at a final

concentration of 2 µM for 15-30 min at 37 °C to induce peptide binding (Dirscherl et al.,

2018).

Washing and fixation. Cells were washed with phosphate buffered saline (PBS), 10 mM

phosphate pH 7.5, 150 mM NaCl), fixed with 3% paraformaldehyde (PFA), and observed by

confocal laser scanning microscopy (cLSM).

Microscopy. We used a confocal laser scanning microscope (LSM 510 Meta, Carl Zeiss

Jena GmbH, Germany) equipped with argon and helium-neon lasers at 488, 543, and 633 nm.

Images were recorded with a 63× Plan Apochromat oil objective (numerical aperture 1.4) at a

resolution of 1596 × 1596 pixels. Data acquisition was performed with the LSM 510 META

software, release 3.2 (Carl Zeiss Jena). During image acquisition, patterns and cells were im-

aged in the same focal plane at a pinhole of 1 Airy unit. Image analysis and processing were

performed using ImageJ (National Institutes of Health, Bethesda, USA). Image processing

comprised cropping, rotation and adjustment of brightness and contrast levels. Experiments

in Figure 1 were repeated at least three times each.

Recombinant α_3 domain of H-2K^b

The α_3 domain of H-2K^b (residues 205-295) was cloned into pET3a (preceded by the resi-

dues MAIQR and followed by DRDM) and expressed in *E. coli* BL21(*DE3*) *pLysS*, refolded

in vitro as described, and isolated by size exclusion chromatography (SEC) on a Cytiva

Hiload Superdex 200 16/600 column (Anjanappa et al., 2020). Molecular weights of the

peaks were determined by comparison to SEC protein standards (Cytiva), namely bovine

thyroglobulin (670 kDa), bovine gamma globulin (158 kDa), chicken ovalbumin (44 kDa),

horse myoglobin (17 kDa), and vitamin B12 (1.35 kDa). The D5 fraction, corresponding to

the elution peak at approximately 20-30 kDa, was boiled with or without DTT (0.6 M final

concentration) in sample buffer (LSB) (350 mM Tris-Cl pH 6.8, 10.28% sodium dodecyl

sulfate (SDS), 36% glycerol 0.012% bromophenol blue). Inclusion body extract boiled with-

out DTT (non-reducing) served as positive control for the formation of covalent oligomers.

Protein quality control after refolding and SEC was performed by nanoscale differential

scanning fluorimetry (nanoDSF) runs (see **Figure 4S2**) acquired with a Nanotemper Prome-

theus NT.48 fluorimeter (Nanotemper, Munich) controlled by PR.ThermControl (version

2.1.2).

Precipitation of surface class I

TAP2-deficient STF1 cells expressing either CE3-HA-K^b or CE3-HA-B*27:05 were kept overnight at 25 °C and then pretreated with tris(2-carboxyethyl)phosphine (TCEP; 1 mM, 10 min), labeled with 400 nm of Bio-MPAA-K3 (5 min) at room temperature (RT) (Reinhardt et al., 2014). Lysis was performed in native lysis buffer (50 mM Tris Cl (pH7.4), 150 mM NaCl, 5 mM EDTA, and 1% Triton X100) for 1 hour at 4 °C. Biotinylated surface proteins were then isolated with neutravidin-coated agarose beads (Thermo Fisher Scientific, Darmstadt Germany). The isolates were boiled at 95 °C for 7 min in the presence (reducing) or absence (non-reducing) of 10 mM dithiothreitol (DTT) in sample buffer as described above. Samples were separated by SDS-PAGE and transferred onto polyvinylidene fluoride (PVDF) membranes. MHC molecules were visualized on the membranes with polyclonal rabbit anti-HA antibody as primary antibody (ab9110, Abcam, Cambridge, United Kingdom) and alkaline phosphatase-conjugated anti-rabbit serum from goat as secondary antibody (1706518, Biorad, Munich, Germany). The signals were visualized by treating the blot with BCIP/NBT substrate (B1911, Sigma Aldrich, St. Louis, Missouri, United States).

Flow cytometry

For verification of cell surface levels of MHC-I, flow cytometry was performed with anti-HLA class I antibody W6/32 (Barnstable, 1978) and anti-HA antibody (12CA5, described in key resources table). Antibody-antigen complexes were labeled with goat secondary antibody against mouse IgG conjugated with allophycocyanin (APC) (115-135-164, Dianova, Hamburg, Germany). Fluorescent signal was recorded by a CyFlow1Space flow cytometer (Sysmex- Partec, Norderstedt, Germany) and analyzed by Flowjo, LLC software.

FRAP

Microcontact printing and antibody patterning for total internal reflection fluores-cence (TIRF) microscopy was performed as described previously (Lanzerstorfer et al., 2020). In short, a field of a large-area PFPE elastomeric stamp (1 µM grid size), obtained by the EV-Group (St. Florian am Inn, Upper Austria, Austria), was cut out, and washed by flushing with ethanol (100%) and distilled water. After drying with nitrogen, the stamp was incubated in 50 mL bovine serum albumin (BSA) solution (1 mg/mL) for 30 min. This step was followed by washing the stamp again with PBS and distilled water. After drying with nitrogen, the stamp was placed with homogeneous pressure onto the clean epoxy-coated glass bottom of a 96-well plate and incubated overnight at 4 °C. The next day, the stamp was stripped from the glass with forceps, and the glass bottom was bonded to a 96-well plastic casting with adhesive tape (3M) and closed with an appropriate lid. For the live cell experi-ments, a reaction chamber was incubated with 100 µL streptavidin solution (50 µg/mL) and incubated for 30 min at room temperature. After washing two times with PBS, 100 µL biotinylated antibody solution (10 µg/mL) was added for 30 min at room temperature result-ing in an antibody surface coverage of > 85% (**Supplementary Figure S1AB**). Finally, the incubation wells were washed twice with PBS, and cells were seeded at defined cell density for the live cell microscopy analysis. The cells were allowed to attach to the surface for at least 3-4 h prior to imaging to ensure a homogeneous cell membrane/substrate interface, which is a prerequisite for quantitative TIRF microscopy. Deformation of the plasma mem-brane on top of the pattern elements was excluded, since control cells that were stained with the lipophilic tracer DiD (1,1′-dioctadecyl-3,3,3′,3′-tetramethylindodicarbocyanine), which stains the plasma membrane uniformly, showed no patterning (**Supplementary Figure S1C**).

For negative control to test the adhesion of the antibodies, anti-HA antibody (Abcam,

ab26228) was labeled with a Zenon Alexa Fluor 488 IgG Labeling Kit (Thermofisher,

Z25102), printed, bleached, and fluorescence recovery was quantified as described below

(**Supplementary Figure S2AB**). In a second control experiment, binding and dissociation of

a construct with both tags (HA-K^b-GFP) from the antibody micropattern was tested, which

only accounted for less than 20% of the mobility of the K^b fraction (**Supplementary Figure

S2C**). Dissociation of Kb-GFP from micropattern elements was negligible (**Supplementary

Figure S2DE**). Thus, in our experiments, k_{slow} was determined by the binding events between

K^b-GFP and other K^b molecules.

Live-cell TIRF microscopy. The detection system was set up on an epi-fluorescence mi-

croscope (Nikon Eclipse Ti2). A multi-laser engine (Toptica Photonics, Munich, Germany)

was used for selective fluorescence excitation of GFP at 488 nm and RFP at 568 nm. The

samples were illuminated in total internal reflection (TIR) configuration (Nikon Ti-LAPP)

using a 60x oil immersion objective (NA = 1.49, APON 60XO TIRF). After appropriate fil-

tering with standard filter sets, the fluorescence was imaged onto a sCMOS camera (Zyla 4.2,

Andor, Northern Ireland). The samples were mounted on an x-y-stage (CMR-STG-MHIX2-

motorized table, Märzhäuser, Germany), and scanning of the larger areas was supported by a

laser-guided automated Perfect Focus System (Nikon PFS).

TIR-FRAP experiments and calculation of diffusion coefficients. FRAP experiments

were carried out on an epi-fluorescence microscope as described above. Single patterns (or

equivalent ROIs with 1 µM in diameter in unpatterned cells) were photobleached (Andor

FRAPPA) with a high-intensity laser pulse (488 nm) applied for 500 ms. Recovery images

were recorded at indicated time intervals. Normalization of data was done by pre-bleach im-

ages, and first data analysis was carried out using NIS Elements software package (Nikon).

Further data processing was done in Graphpad Prism. Resulting FRAP curves were plotted

based on the standard error of the mean and fitted using a bi-exponential equation. Kinetic

FRAP parameters were directly obtained from curve fitting with the following model as-

sumptions: Since unhindered MHC-I diffusion (e.g., off pattern and/or after peptide treat-

ment; Fig. 2E) was found to be much faster than the recovery within our pattern elements, we

applied a diffusion-uncoupled recovery scheme (Sprague and Mcnally, 2005). Here, after the

photobleaching step, fluorescent molecules rapidly diffuse throughout the bleached

micropattern, and only bound bleached molecules remain inside the spot. The bleached mole-

cules then gradually dissociate from their binding sites. Unbleached molecules can then re-

place the bleached molecules at the binding sites as they become vacant. The diffusion-

uncoupled FRAP recovery curve consists of two separable components: the early recovery

due to diffusion and the slower recovery due to exchange at binding sites, and can be fitted

using a diffusion-uncoupled two-component fit:

15
$$Y = Y_{t=0} + A_1(1 - e^{-kfast*x}) + A_2(1 - e^{-kslow*x}),$$

where A_1 is the amplitude of the fast-diffusing population, A_2 the amplitude of the slow

diffusing population (binding reaction), and k_{fast} and k_{slow} are the rate constants of A_1 and A_2,

respectively.

Diffusion coefficients were obtained using the initial image recordings and the simFRAP

plugin for ImageJ (Blumenthal *et al*., 2015).

Temperature-induced K^b associations. Temperature-dependent experiments were car-

ried out on an epi-fluorescence microscope as described above further equipped with a cage

incubator (Okolab, Shanghai, China). Cells were grown at 25 °C overnight on antibody-

patterned surfaces and treated with SIINFEKL peptide as indicated. For induction of K^b FHC

association, cells were mounted on pre-warmed microscopy stage, and imaging of the

GFP signal was started when the medium reached 37 °C.

Fluorescence contrast quantitation. Contrast analysis was performed as described previ-

ously (Lanzerstorfer et al., 2014) In short, initial imaging recording was supported by the

Nikon NIS Elements software. Images were exported as TIFF frames and fluorescence con-

trast analysis was performed with the Spotty framework (Borgmann et al., 2012). The fluo-

rescence contrast <c> was calculated as

$$\langle c \rangle = \frac{(F^+ - F^-)}{\left(F^+ - F_{bg}\right)}$$

, where F^+ is the intensity of the inner pixels of the pattern, F^- the intensity of the sur-

rounding pixels of the micropattern, and F_{bg} the intensity of the global background.

Single molecule microscopy

Imaging. Cells were transferred 48 h post transfection onto glass coverslips coated with a

poly-L-lysine-graft-(polyethylene glycol) copolymer functionalized with RGD tripeptide to

minimize non-specific binding of fluorescent nanobodies (You et al., 2014). Cells imaged in

presence of SL8 peptide (sequence SIINFEKL) were pre-incubated with 1 µM of SL8 pep-

tide 12 h before imaging. Single-molecule imaging experiments were conducted by total in-

ternal reflection fluorescence (TIRF) microscopy with an inverted microscope (Olympus

IX83) equipped with a motorized 4-Line TIR illumination condenser (Olympus) and a back-

illuminated electron multiplying (EM) CCD camera (iXon Ultra 897, Andor Technology). A

100× magnification objective with a numerical aperture of 1.45 (UPLAPO 100× HR, NA 1.5,

Olympus) together with a 1.6 × magnification changer was used for TIR illumination of the

sample. Imaging was conducted with or without 1 µM of fresh SL8 peptide. The sample was

preincubated with 10 µg/mL of Brefeldin A (BFA) for 15 min in order to inhibit protein

transport of GFP-K^b/β_2m heterodimers to the plasma membrane. All experiments were car-

ried out at 37 °C in medium without phenol red supplemented with an oxygen scavenger and

a redox-active photoprotectant to minimize photobleaching (Vogelsang et al., 2008) and pen-

icillin and streptomycin (PAA). For cell surface labeling of GFP-K^b, a 1:1 mixture of anti-

GFP nanobodies (2 nM each) site-specifically conjugated with ATTO 643 and ATTO Rho11

(Wilmes et al., 2020b), respectively, were added to the medium, thus ensuring >90 % binding

given the 0.3 nM binding affinity (Kirchhofer et al., 2010). After incubation for at least

5 min, image acquisition was started with the labeled nanobodies kept in the bulk solution

during the whole experiment in order to ensure high equilibrium binding. Dimerization of the

positive control was induced by applying 0.3 nM of a tandem nanobody crosslinker

(LaG16V2) binding to an orthogonal epitope (Fridy et al., 2014). For single molecule co-

localization and co-tracking experiments, orange (ATTO Rho11) and red (ATTO 643) emit-

ting fluorophores were simultaneously excited by illumination with a 561 nm laser (MPB

Communications) and a 642 nm laser (MPB Communications). Fluorescence was detected

with a spectral image splitter (QuadView QV2, Photometrics) with a dichroic beam splitter

(Chroma) combined with the bandpass filter 600/37 (BrightLine HC) for detection of

ATTO Rho11 and 685/40 (Brightline HC) for detection of ATTO 643 dividing each emission

channel into 256×256 pixels. Image stacks of 150 frames were recorded for each cell at a

time resolution of 32 ms/frame. Diffusion constants were determined by mean square dis-

placement analysis within a time window of 320 ms (10 frames).

Single molecule analysis. MHC homodimerization was quantified based on sequential co-

localization and co-tracking analysis using self-written Matlab code "SlimFAST " as de-

scribed in detail previously (Roder et al., 2014; Wilmes et al., 2015b). After aligning ATTO

643 and ATTO Rho11 channels with sub-pixel precision through spatial transformation based

on a calibration measurement with multicolor fluorescent beads (TetraSpeck microspheres

0.1 μm, Invitrogen), individual molecules detected in both spectral channels of the same

frame within a distance threshold of 100 nm were considered to be co-localized. For SMCT

analysis, the multiple-target tracing (MTT) algorithm was applied to this dataset of co-

localized molecules to reconstruct co-locomotion trajectories (co-trajectories) from the identi-

fied population of co-localizations. For the co-tracking analysis, only co-trajectories with

a minimum of 10 consecutive steps (320 ms) were considered. This cut-off was determined

based on systematic analysis of a negative control experiment with non-interacting model

transmembrane proteins (Wilmes et al., 2020b) in order to minimize background from ran-

dom co-localization. The relative fraction of co-tracked molecules was determined with re-

spect to the absolute number of trajectories from both channels and corrected for homodimers

stochastically double-labeled with the same fluorophore species as follows:

$$AB^* = \frac{AB}{2 \times \left[\left(\frac{A}{A+B}\right) \times \left(\frac{B}{A+B}\right)\right]}$$

$$rel.\,co-locomotion = \frac{2 \times AB^*}{(A+B)}$$

where A, B, AB and AB* are the numbers of trajectories observed for ATTO Rho11,

ATTO 643, co-trajectories and corrected co-trajectories, respectively.

For **Figures 3F and 3G**, the relative fraction of co-localized mobile and immobile molecules

was determined with respect to the absolute number of mobile or immobile molecules from

both channels and corrected for homodimers stochastically double-labeled with the same

fluorophore species as follows:

$$AB^* = \frac{AB/A + AB/B}{2}$$

$$rel.\,fraction = \frac{AB^*}{2 \times \left[\left(\dfrac{A}{A+B}\right) \times \left(\dfrac{B}{A+B}\right)\right]}$$

where A, B, AB and AB* are the numbers of molecules part of a trajectory observed for

ATTO Rho11, ATTO 643, co-localized molecules and corrected co-localized molecules, re-

spectively.

Immobile molecules were classified by their appearance within a radius described by the lo-

calization precision and molecular observation probability.

Statistics and statistical analysis. For **Figure 3D-H**, each data point represents the analy-

sis from one cell, with ≥14 cells measured per experiment and many trajectories analyzed per

cell (**3D**, 238-2513; **3E**, 238-2513; **3F**, 314-2732; **3G**, 41-354 tracked immobile particles and

362-3141 mobile particles; **3H**, 2-301 co-trajectories). Statistical significances were calculat-

ed by two-sample Kolmogorov-Smirnov test, Kruskal-Wallis test with multiple comparisons,

unpaired *t* test and two-way ANOVA with multiple comparisons as indicated in the figure

legends, using version Prism 8.4.0 for MacOS (GraphPad, San Diego, USA). Box plots were

used for visualization and indicate the data distribution of second and third quartile (box),

median (line), mean (square) and data range (whiskers).

Native mass spectrometry

In advance of native MS measurements, a small amount (<1%) of covalent dimers of the

α_3 domain, which had formed as side product during refolding, were removed by size exclu-

sion chromatography on Superdex 75 10/300 GL (Cytiva). Amicon Ultra 0.5 mL centrifugal

filter units (molecular weight cut-off 3 kDa; Merck Millipore) were used at 14,000 x g and

4 °C to exchange purified protein samples to 150 mM ammonium acetate (99.99 %; Sigma-

Aldrich), pH 7.2. The final concentration of the α_3 domain monomer was 5 µM, 10 µM or

20 µM. Native MS analysis was implemented on a Q-Tof II mass spectrometer in positive

electrospray ionization mode. The instrument was modified to enable high mass experiments

(Waters and MS Vision, (van den Heuvel et al., 2006)). Sample ions were introduced into the

vacuum using homemade capillaries via a nano-electrospray ionization source in positive ion

mode (source pressure: 10 mbar). Borosilicate glass tubes (inner diameter: 0.68 mm, outer

diameter: 1.2 mm; World Precision Instruments) were pulled into closed capillaries in a two-

step program using a squared box filament (2.5 mm × 2.5 mm) within a micropipette puller

(P-1000, Sutter Instruments). The capillaries were then gold-coated using a sputter coater

(5.0×10^{-2} mbar, 30.0 mA, 100 s, 3 runs to vacuum limit 3.0×10^{-2} mbar argon, distance of

plate holder: 5 cm; CCU-010, safematic). Capillaries were opened directly on the sample

cone of the mass spectrometer. In regular MS mode, spectra were recorded at a capillary

voltage of 1.45 kV and a cone voltage of 100 V to 150 V. Protein species with quaternary

structure were assigned by MS/MS analysis. These experiments were carried out using argon

as collision gas (1.2×10^{-2} mbar). The acceleration voltage ranged from 10 V to 100 V.

Comparability of results was ensured as MS quadrupole profiles and pusher settings were

kept constant in all measurements. A spectrum of cesium iodide (25 g/L) was recorded on the

same day of the particular measurement to calibrate the data.

All spectra were evaluated regarding experimental mass (MassLynx V4.1, Waters) and ar-

ea under the curve (AUC; UniDec (Marty et al., 2015)) of the detected mass species. The

values of the shown averaged masses and AUC of the different species as well as the corre-

sponding standard deviation result from at least three independent measurements. Exact ex-

perimental masses are presented in **Supplementary Figure S4A**.

Molecular model of the K^b heavy chain dimer

The crystal structure of K^b in complex with a chicken ovalbumin epitope (PDB 3p9m)

served as template to generate a model of the heavy chain dimer. Using the Pymol program

(DeLano, 2002), it was possible to superimpose a copy of the 3p9m heavy chain on the β_2m

subunit of a second 3p9m structure (FHC dimer). The superposition involved only the match-

ing of the α_3 domain backbone of the K^b molecule (residues 181-277) onto the β_2m subunit

resulting in a small root-mean-square deviation (RMSD) of 1.25 Å. Very little steric overlap

of the superimposed α_3 domain with the second heavy chain was detected. This minimal

overlap of the α_1/α_2 domain of the superimposed heavy chain was removed by adjusting

backbone dihedral angles of residues 179-181 in the linker between α_1/α_2 domain and the α_3

domain. The initial structural model was energy-minimized to remove any residual steric

overlap and was prepared for molecular dynamics (MD) simulations using the Amber18

package (Case et al., 2018).

Molecular docking

For comparison of FHC homodimers and K^b HC/β_2m heterodimers, MD simulations were

performed starting from the coordinates of the K^b structure in PDB 3p9m. Proteins were

solvated in octahedral boxes including explicit sodium and chloride ions (0.15 M) and explic-

it TIP3P water molecules keeping a minimum distance of 10 Å between protein atoms and

box boundaries (Jorgensen et al., 1983). The parm14SB force field was used for the proteins

and peptides (Maier et al., 2015). The simulation systems were again energy minimized

(5000 steps) after solvation followed by heating up to 300 K in steps of 100 K with position

restraints on all heavy atoms of the proteins. Subsequently, positional restraints were gradual-

ly removed from an initial 25 kcal·mol^{-1}·Å^{-2} to 0.5 kcal·mol^{-1}·Å^{-2} within 0.5 ns followed by a

ns unrestrained equilibration at 300 K. All production simulations were performed at a temperature of 300 K and a pressure of 1 bar. The hydrogen mass repartition option of Amber was used to allow a time step of 4 fs (Hopkins et al., 2015). Unrestrained production simulations for up to 400 ns were performed. The interface packing was analyzed by calculation of the buried surface area using the Shrake method, (Shrake and Rupley, 1973); analysis of root-mean-square deviations (RMSD) was performed using the cpptraj module of the Amber18 package.

High-speed atomic force measurements for antibody density estimation

We employed high-speed atomic force microscopy (HS-AFM), capable of resolving individual antibodies on surfaces (Preiner et al., 2014; Strasser et al., 2020) to estimate the antibody surface density in our pattern elements. Antibody micropatterns were prepared as described above. Antibody solution (10 µg/mL) was finally incubated for 15 – 1200 s, followed by washing and imaging in PBS. HS-AFM (RIBM, Japan) was operated in tapping mode at room temperature with free amplitudes of 1.5-2.5 nm and an amplitude set point of larger than 90%. Silicon nitride cantilevers (USC-F1.2-k0.15, Nanoworld AG, Neuchâtel, Switzerland) with nominal spring constants of 0.1-0.15 N/m, resonance frequencies of ~ 500 kHz, and a quality factor of ~2 in liquids were used.

Since the antibody coverage after 20 min incubation was too high to discriminate and count individual molecules, the images of the shortest incubation time (15 s) were used and thus lowest coverage and largest distance between individual molecules to estimate the area coverage per antibody (neglecting potential tip convolution effects). A height threshold of 2.7 nm above the streptavidin layer was applied for detection of particle density and mean area per particle (~190 nm^2), which were then used to convert the total area above the threshold in

the other images (30 s – 20 min) into a surface density. The resulting surface density Γ of antibodies as a function of incubation time is reasonably described by a mono-exponential function as typical for a pseudo-first order reaction (Artelsmair et al., 2008), i.e.

$$\Gamma(t) = A\left(1 - e^{-t/\tau}\right)$$

with A = 7646 ± 1828 IgGs/µm^2 (and τ = 77 ± 43 s). Image analysis was performed using threshold algorithms in Gwyddion (Nečas and Klapetek, 2012).

Acknowledgements

We thank the donors of reagents as mentioned in the Materials, Venkat Raman Ramnarayan for comments on the manuscript, Christian P. Richter for support with SMT/SMCT evaluation, Ankur Saikia and Christian Guenther for performing protein chromatography, the iBiOs staff for technical support with single molecule microscopy, the SPC facility at EMBL Hamburg for technical support, and Uschi Wellbrock for excellent technical assistance.

Funding

Deutsche Forschungsgemeinschaft (DFG SP583/7-2), Bundesministerium für Bildung und Forschung (BMBF, 031A153A); Tönjes Vagt Foundation (XXXII), iNEXT-Discovery (11911), Jacobs University (all to SSp); Deutsche Forschungsgemeinschaft (SFB 944, projects P8 and Z, Facility iBiOs, PI 405/14-1) to JP. PL and JW acknowledge funding from the province of Upper Austria as a part of the FH Upper Austria Center of Excellence for Technological Innovation in Medicine (TIMed Center) and the Christian Doppler Forschungsgesellschaft (Josef Ressel Center for Phytogenic Drug Research). CU acknowledges

funding from the Leibniz Association through SAW-2014-HPI-4 grant. The Heinrich-Pette-Institute, Leibniz Institute for Experimental Virology is supported by the Free and Hanseatic City Hamburg and the German Federal Ministry of Health.

Author Contributions

The experimental work was performed as follows: CD, Fig. 1CDEG; ZH, Fig. 1F; PL, Fig. 2, 4AB, S1, S2, S4B; SL, Fig. 3, S3, supplementary movies; ARH, Fig. 4CD and S4B; JDK, Fig. 4E-G, S4A; MZ, Fig. 5, S5; AK and JPr, Figure S1AB. MGA, CU, JW and JPi supervised and interpreted experiments. MZ performed all MD simulations (Fig. 5, 5S1). SSp conceived the work, supervised and coordinated experiments, and wrote the manuscript in cooperation with JPi and PL and assistance by NL with input from all authors. Funding was acquired by CU, JPi, JW, MGA, MZ, PL, and SSp.

Competing Interests:

The authors declare that no competing interests exist.

References

Allen RL, O'Callaghan CA, McMichael AJ, Bowness P. 1999. Cutting edge: HLA-B27 can form a novel beta 2-microglobulin-free heavy chain homodimer structure. *J Immunol Baltim Md 1950* **162**:5045–5048.

Anjanappa R, Garcia-Alai M, Kopicki J-D, Lockhauserbäumer J, Aboelmagd M, Hinrichs J, Nemtanu IM, Uetrecht C, Zacharias M, Springer S, Meijers R. 2020. Structures of peptide-free and partially loaded MHC class I molecules reveal mechanisms of peptide selection. *Nat Commun* **11**:1314. doi:10.1038/s41467-020-14862-4

Antoniou AN, Guiliano DB, Lenart I, Burn G, Powis SJ. 2011. The oxidative folding and misfolding of human leukocyte antigen-b27. *Antioxid Redox Signal* **15**:669–684. doi:10.1089/ars.2010.3692

Armony G, Heck AJR, Wu W. 2021. Extracellular crosslinking mass spectrometry reveals HLA class I – HLA class II interactions on the cell surface. *Mol Immunol* **136**:16–25. doi:10.1016/j.molimm.2021.05.010

Arosa FA, Esgalhado AJ, Reste-Ferreira D, Cardoso EM. 2021. Open MHC Class I Conformers: A Look through the Looking Glass. *Int J Mol Sci* **22**:9738. doi:10.3390/ijms22189738

Arosa FA, Santos SG, Powis SJ. 2007. Open conformers: the hidden face of MHC-I molecules. *Trends Immunol* **28**:115–123. doi:10.1016/j.it.2007.01.002

Artelsmair H, Kienberger F, Tinazli A, Schlapak R, Zhu R, Preiner J, Wruss J, Kastner M, Saucedo-Zeni N, Hoelzl M, Rankl C, Baumgartner W, Howorka S, Blaas D, Gruber HJ, Tampé R, Hinterdorfer P. 2008. Atomic force microscopy-derived nanoscale chip for the detection of human pathogenic viruses. *Small Weinh Bergstr Ger* **4**:847–854. doi:10.1002/smll.200700691

Bahadur RP, Zacharias M. 2008. The interface of protein-protein complexes: Analysis of contacts and prediction of interactions. *Cell Mol Life Sci* **65**:1059–1072. doi:10.1007/s00018-007-7451-x

Baia D, Pou J, Jones D, Mandelboim O, Trowsdale J, Muntasell A, Lopez-Botet M. 2016. Interaction of the LILRB1 inhibitory receptor with HLA class Ia dimers. *Eur J Immunol* **46**:1681–90. doi:10.1002/eji.201546149

Baía D, Pou J, Jones D, Mandelboim O, Trowsdale J, Muntasell A, López-Botet M. 2016. Interaction of the LILRB1 inhibitory receptor with HLA class Ia dimers. *Eur J Immunol* **46**:1681–1690. doi:10.1002/eji.201546149

Barnstable C. 1978. Production of monoclonal antibodies to group A erythrocytes, HLA and other human cell surface antigens-new tools for genetic analysis. *Cell* **14**:9–20. doi:10.1016/0092-8674(78)90296-9

Blumenthal D, Edidin M, Gheber LA. 2016. Trafficking of MHC molecules to the cell surface creates dynamic protein patches. *J Cell Sci* **129**:3342–3350. doi:10.1242/jcs.187112

Bodnar A, Bacso Z, Jenei A, Jovin TM, Edidin M, Damjanovich S, Matko J. 2003. Class I HLA oligomerization at the surface of B cells is controlled by exogenous beta(2)-microglobulin: implications in activation of cytotoxic T lymphocytes. *Int Immunol* **15**:331–9.

Bondar A, Jang W, Sviridova E, Lambert NA. 2020. Components of the Gs signaling cascade exhibit distinct changes in mobility and membrane domain localization upon β2 - adrenergic receptor activation. *Traffic Cph Den* **21**:324–332. doi:10.1111/tra.12724

Borgmann DM, Weghuber J, Schaller S, Jacak J, Winkler SM. 2012. Identification of patterns in microscopy images of biological samples using evolution strategies.

Campbell EC, Antoniou AN, Powis SJ. 2012. The multi-faceted nature of HLA class I dimer molecules. *Immunology* **136**:380–384. doi:10.1111/j.1365-2567.2012.03593.x

Capps GG, Robinson BE, Lewis KD, Zúñiga MC. 1993. In vivo dimeric association of class I MHC heavy chains. Possible relationship to class I MHC heavy chain-beta 2-microglobulin dissociation. *J Immunol* **151**:159–169.

Case DA, Ben-Shalom IY, Brozell SR, Cerutti DS, Cheatham TE III, Cruzeiro VWD, Darde TA, Duke RE, Ghoreishi D, Gilson MK, Gohlke H, Goetz AW, Greene D, Harris R, Homeyer N, Izadi S, Kovalenko A, Kurtzman T, Lee TS, LeGrand S, Li P, Lin C, Liu J, Luchko T, Luo R, Mermelstein DJ, Merz KM, Miao Y, Monard G, Nguyen C, Nguyen H, Omelyan I, Onufriev A, Pan F, Qi R, Roe DR, Roitberg A, Sagui C, Schott-Verdugo S, Shen J, Simmerling CL, Smith J, Salomon-Ferrer R, Swails J, Walker RC, Wang J, Wei H, Wolf RM, Wu X, Xiao L, York DM, Kollman PA. 2018. Amber18. San Fransisco: University of California.

Chakrabarti A, Matko J, Rahman NA, Barisas BG, Edidin M. 1992. Self-association of class I major histocompatibility complex molecules in liposome and cell surface membranes. *Biochemistry* **31**:7182–9.

Chen B, Li J, He C, Li D, Tong W, Zou Y, Xu W. 2017. Role of HLA-B27 in the pathogenesis of ankylosing spondylitis. *Mol Med Rep* **15**:1943–1951. doi:10.3892/mmr.2017.6248

Comber JD, Philip R. 2014. MHC class I antigen presentation and implications for developing a new generation of therapeutic vaccines. *Ther Adv Vaccines* **2**:77–89. doi:10.1177/2051013614525375

Dangoria NS, DeLay ML, Kingsbury DJ, Mear JP, Uchanska-Ziegler B, Ziegler A, Colbert RA. 2002. HLA-B27 Misfolding Is Associated with Aberrant Intermolecular Disulfide Bond Formation (Dimerization) in the Endoplasmic Reticulum. *J Biol Chem* **277**:23459–23468. doi:10.1074/jbc.M110336200

Day PM, Esquivel F, Lukszo J, Bennink JR, Yewdell JW. 1995. Effect of TAP on the generation and intracellular trafficking of peptide-receptive major histocompatibility complex class I molecules. *Immunity* **2**:137–147. doi:10.1016/S1074-7613(95)80014-X

DeLano WL. 2002. Pymol: An open-source molecular graphics tool. *CCP4 Newsl Protein Crystallogr* 82–92.

Dirscherl C, Hein Z, Ramnarayan VR, Jacob-Dolan C, Springer S. 2018. A two-hybrid antibody micropattern assay reveals specific in cis interactions of MHC I heavy chains at the cell surface. *eLife* **7**:e34150. doi:10.7554/eLife.34150

Dirscherl C, Palankar R, Delcea M, Kolesnikova TA, Springer S. 2017. Specific Capture of Peptide-Receptive Major Histocompatibility Complex Class I Molecules by Antibody Micropatterns Allows for a Novel Peptide-Binding Assay in Live Cells. *Small* **13**:1602974. doi:10.1002/smll.201602974

Dulberger CL, McMurtrey CP, Hölzemer A, Neu KE, Liu V, Steinbach AM, Garcia-Beltran WF, Sulak M, Jabri B, Lynch VJ, Altfeld M, Hildebrand WH, Adams EJ. 2017. Human Leukocyte Antigen F Presents Peptides and Regulates Immunity through Interac-

tions with NK Cell Receptors. *Immunity* **46**:1018-1029.e7. doi:10.1016/j.immuni.2017.06.002

Edidin M, Achilles S, Zeff R, Wei T. 1997. Probing the stability of class I major histocompatibility complex (MHC) molecules on the surface of human cells. *Immunogenetics* **46**:41–45. doi:10.1007/s002510050240

Elliott T, Cerundolo V, Elvin J, Townsend A. 1991. Peptide-induced conformational change of the class I heavy chain. *Nature* **351**:402–406. doi:10.1038/351402a0

Fassett MS, Davis DM, Valter MM, Cohen GB, Strominger JL. 2001. Signaling at the inhibitory natural killer cell immune synapse regulates lipid raft polarization but not class I MHC clustering. *Proc Natl Acad Sci U S A* **98**:14547–14552. doi:10.1073/pnas.211563598

Ferez M, Castro M, Alarcon B, van Santen HM. 2014. Cognate peptide-MHC complexes are expressed as tightly apposed nanoclusters in virus-infected cells to allow TCR crosslinking. *J Immunol* **192**:52–8. doi:10.4049/jimmunol.1301224

Fooksman DR, Grönvall GK, Tang Q, Edidin M. 2006. Clustering Class I MHC Modulates Sensitivity of T Cell Recognition. *J Immunol* **176**:6673–6680. doi:10.4049/jimmunol.176.11.6673

Fridy PC, Li Y, Keegan S, Thompson MK, Nudelman I, Scheid JF, Oeffinger M, Nussenzweig MC, Fenyö D, Chait BT, Rout MP. 2014. A robust pipeline for rapid production of versatile nanobody repertoires. *Nat Methods* **11**:1253–1260. doi:10.1038/nmeth.3170

Gakamsky DM, Bjorkman PJ, Pecht I. 1996. Peptide Interaction with a Class I Major Histocompatibility Complex-Encoded Molecule: Allosteric Control of the Ternary Complex Stability [†]. *Biochemistry* **35**:14841–14848. doi:10.1021/bi961707u

Gambin Y, Lopez-Esparza R, Reffay M, Sierecki E, Gov NS, Genest M, Hodges RS, Urbach W. 2006. Lateral mobility of proteins in liquid membranes revisited. *Proc Natl Acad Sci U S A* **103**:2098–2102. doi:10.1073/pnas.0511026103

Garstka M, Borchert B, Al-Balushi M, Praveen P, Kühl N, Majoul I, Duden R, Springer S. 2007. Peptide-receptive Major Histocompatibility Complex Class I Molecules Cycle between Endoplasmic Reticulum and *cis* -Golgi in Wild-type Lymphocytes. *J Biol Chem* **282**:30680–30690. doi:10.1074/jbc.M701721200

Geng J, Altman JD, Krishnakumar S, Raghavan M. 2018. Empty conformers of HLA-B preferentially bind CD8 and regulate CD8+ T cell function. *eLife* **7**:e36341. doi:10.7554/eLife.36341

Goodridge JP, Lee N, Burian A, Pyo C-W, Tykodi SS, Warren EH, Yee C, Riddell SR, Geraghty DE. 2013. HLA-F and MHC-I Open Conformers Cooperate in a MHC-I Antigen Cross-Presentation Pathway. *J Immunol* **191**:1567–1577. doi:10.4049/jimmunol.1300080

Graham FL, van der Eb AJ. 1973. A new technique for the assay of infectivity of human adenovirus 5 DNA. *Virology* **52**:456–467. doi:10.1016/0042-6822(73)90341-3

Hein Z, Uchtenhagen H, Abualrous ET, Saini SK, Janssen L, Van Hateren A, Wiek C, Hanenberg H, Momburg F, Achour A, Elliott T, Springer S, Boulanger D. 2014. Peptide-independent stabilization of MHC class I molecules breaches cellular quality control. *J Cell Sci* **127**:2885–2897. doi:10.1242/jcs.145334

Ho CCM, Chhabra A, Starkl P, Schnorr P-J, Wilmes S, Moraga I, Kwon H-S, Gaudenzio N, Sibilano R, Wehrman TS, Gakovic M, Sockolosky JT, Tiffany MR, Ring AM, Piehler

J, Weissman IL, Galli SJ, Shizuru JA, Garcia KC. 2017. Decoupling the Functional Pleiotropy of Stem Cell Factor by Tuning c-Kit Signaling. *Cell* **168**:1041-1052.e18. doi:10.1016/j.cell.2017.02.011

Hopkins CW, Le Grand S, Walker RC, Roitberg AE. 2015. Long-Time-Step Molecular Dynamics through Hydrogen Mass Repartitioning. *J Chem Theory Comput* **11**:1864–1874. doi:10.1021/ct5010406

Ibach J, Radon Y, Gelléri M, Sonntag MH, Brunsveld L, Bastiaens PIH, Verveer PJ. 2015. Single Particle Tracking Reveals that EGFR Signaling Activity Is Amplified in Clathrin-Coated Pits. *PloS One* **10**:e0143162. doi:10.1371/journal.pone.0143162

Jorgensen WL, Chandrasekhar J, Madura JD, Impey RW, Klein ML. 1983. Comparison of simple potential functions for simulating liquid water. *J Chem Phys* **79**:926–935. doi:10.1063/1.445869

Kaufman J. 2018. Generalists and Specialists: A New View of How MHC Class I Molecules Fight Infectious Pathogens. *Trends Immunol* **39**:367–379. doi:10.1016/j.it.2018.01.001

Kirchhofer A, Helma J, Schmidthals K, Frauer C, Cui S, Karcher A, Pellis M, Muyldermans S, Casas-Delucchi CS, Cardoso MC, Leonhardt H, Hopfner K-P, Rothbauer U. 2010. Modulation of protein properties in living cells using nanobodies. *Nat Struct Mol Biol* **17**:133–138. doi:10.1038/nsmb.1727

Lanzerstorfer P, Borgmann D, Schütz G, Winkler SM, Höglinger O, Weghuber J. 2014. Quantification and kinetic analysis of Grb2-EGFR interaction on micro-patterned surfaces for the characterization of EGFR-modulating substances. *PloS One* **9**:e92151. doi:10.1371/journal.pone.0092151

Lanzerstorfer P, Müller U, Gordiyenko K, Weghuber J, Niemeyer CM. 2020. Highly Modular Protein Micropatterning Sheds Light on the Role of Clathrin-Mediated Endocytosis for the Quantitative Analysis of Protein-Protein Interactions in Live Cells. *Biomolecules* **10**:540. doi:10.3390/biom10040540

Lin W-C, Iversen L, Tu H-L, Rhodes C, Christensen SM, Iwig JS, Hansen SD, Huang WYC, Groves JT. 2014. H-Ras forms dimers on membrane surfaces via a protein-protein interface. *Proc Natl Acad Sci U S A* **111**:2996–3001. doi:10.1073/pnas.1321155111

Ljunggren H-G, Stam NJ, Öhlén C, Neefjes JJ, Höglund P, Heemels M-T, Bastin J, Schumacher TNM, Townsend A, Kärre K, Ploegh HL. 1990. Empty MHC class I molecules come out in the cold. *Nature* **346**:476–480. doi:10.1038/346476a0

Low-Nam ST, Lidke KA, Cutler PJ, Roovers RC, van Bergen en Henegouwen PMP, Wilson BS, Lidke DS. 2011. ErbB1 dimerization is promoted by domain co-confinement and stabilized by ligand binding. *Nat Struct Mol Biol* **18**:1244–1249. doi:10.1038/nsmb.2135

Lu X, Gibbs JS, Hickman HD, David A, Dolan BP, Jin Y, Kranz DM, Bennink JR, Yewdell JW, Varma R. 2012. Endogenous viral antigen processing generates peptide-specific MHC class I cell-surface clusters. *Proc Natl Acad Sci U S A* **109**:15407–15412. doi:10.1073/pnas.1208696109

Mahmutefendic H, Blagojevic G, Tomas MI, Kucic N, Lucin P. 2011. Segregation of open Major Histocompatibility Class I conformers at the plasma membrane and during endosomal trafficking reveals conformation-based sorting in the endosomal system. *Int J Biochem Cell Biol* **43**:504–15. doi:10.1016/j.biocel.2010.12.002

Maier JA, Martinez C, Kasavajhala K, Wickstrom L, Hauser KE, Simmerling C. 2015. ff14SB: Improving the Accuracy of Protein Side Chain and Backbone Parameters from ff99SB. *J Chem Theory Comput* **11**:3696–3713. doi:10.1021/acs.jctc.5b00255

Makhadiyeva D, Lam L, Moatari M, Vallance J, Zheng Y, Campbell EC, Powis SJ. 2012. MHC class I dimer formation by alteration of the cellular redox environment and induction of apoptosis: Redox control of MHC class I dimer formation. *Immunology* **135**:133–139. doi:10.1111/j.1365-2567.2011.03518.x

Marty MT, Baldwin AJ, Marklund EG, Hochberg GKA, Benesch JLP, Robinson CV. 2015. Bayesian Deconvolution of Mass and Ion Mobility Spectra: From Binary Interactions to Polydisperse Ensembles. *Anal Chem* **87**:4370–4376. doi:10.1021/acs.analchem.5b00140

Matko J, Bushkin Y, Wei T, Edidin M. 1994. Clustering of class I HLA molecules on the surfaces of activated and transformed human cells. *J Immunol* **152**:3353–60.

Montealegre Sebastián, Venugopalan V, Fritzsche S, Kulicke C, Hein Z, Springer S. 2015. Dissociation of β $_2$ ☐microglobulin determines the surface quality control of major histocompatibility complex class I molecules. *FASEB J* **29**:2780–2788. doi:10.1096/fj.14-268094

Montealegre S., Venugopalan V, Fritzsche S, Kulicke C, Hein Z, Springer S. 2015. Dissociation of beta2-microglobulin determines the surface quality control of major histocompatibility complex class I molecules. *Faseb J* **29**:2780–8. doi:10.1096/fj.14-268094

Moraga I, Wernig G, Wilmes S, Gryshkova V, Richter CP, Hong W-J, Sinha R, Guo F, Fabionar H, Wehrman TS, Krutzik P, Demharter S, Plo I, Weissman IL, Minary P, Majeti R, Constantinescu SN, Piehler J, Garcia KC. 2015. Tuning cytokine receptor signaling by re-orienting dimer geometry with surrogate ligands. *Cell* **160**:1196–1208. doi:10.1016/j.cell.2015.02.011

Mylvaganam SM, Grinstein S, Freeman SA. 2018. Picket-fences in the plasma membrane: functions in immune cells and phagocytosis. *Semin Immunopathol* **40**:605–615. doi:10.1007/s00281-018-0705-x

Nečas D, Klapetek P. 2012. Gwyddion: an open-source software for SPM data analysis. *Open Phys* **10**. doi:10.2478/s11534-011-0096-2

Niman HL, Houghten RA, Walker LE, Reisfeld RA, Wilson IA, Hogle JM, Lerner RA. 1983. Generation of protein-reactive antibodies by short peptides is an event of high frequency: implications for the structural basis of immune recognition. *Proc Natl Acad Sci* **80**:4949–4953. doi:10.1073/pnas.80.16.4949

Ortiz-Navarrete V, Hammerling GJ. 1991. Surface appearance and instability of empty H-2 class I molecules under physiological conditions. *Proc Natl Acad Sci* **88**:3594–3597. doi:10.1073/pnas.88.9.3594

Preiner J, Kodera N, Tang J, Ebner A, Brameshuber M, Blaas D, Gelbmann N, Gruber HJ, Ando T, Hinterdorfer P. 2014. IgGs are made for walking on bacterial and viral surfaces. *Nat Commun* **5**:4394. doi:10.1038/ncomms5394

Reinhardt U, Lotze J, Zernia S, Mörl K, Beck-Sickinger AG, Seitz O. 2014. Peptide-Templated Acyl Transfer: A Chemical Method for the Labeling of Membrane Proteins on Live Cells. *Angew Chem Int Ed* **53**:10237–10241. doi:10.1002/anie.201403214

Richter D, Moraga I, Winkelmann H, Birkholz O, Wilmes S, Schulte M, Kraich M, Kenneweg H, Beutel O, Selenschik P, Paterok D, Gavutis M, Schmidt T, Garcia KC,

Müller TD, Piehler J. 2017. Ligand-induced type II interleukin-4 receptor dimers are sustained by rapid re-association within plasma membrane microcompartments. *Nat Commun* **8**:15976. doi:10.1038/ncomms15976

Roder F, Wilmes S, Richter CP, Piehler J. 2014. Rapid Transfer of Transmembrane Proteins for Single Molecule Dimerization Assays in Polymer-Supported Membranes. *ACS Chem Biol* **9**:2479–2484. doi:10.1021/cb5005806

Schell TD. 2002. The assembly of functional beta2-microglobulin-free MHC class I molecules that interact with peptides and CD8+ T lymphocytes. *Int Immunol* **14**:775–782. doi:10.1093/intimm/dxf041

Schwarzenbacher M, Kaltenbrunner M, Brameshuber M, Hesch C, Paster W, Weghuber J, Heise B, Sonnleitner A, Stockinger H, Schutz GJ. 2008. Micropatterning for quantitative analysis of protein-protein interactions in living cells. *Nat Methods* **5**:1053–60. doi:10.1038/nmeth.1268

Sevcsik E, Brameshuber M, Folser M, Weghuber J, Honigmann A, Schutz GJ. 2015. GPI-anchored proteins do not reside in ordered domains in the live cell plasma membrane. *Nat Commun* **6**:6969. doi:10.1038/ncomms7969

Shrake A, Rupley JA. 1973. Environment and exposure to solvent of protein atoms. Lysozyme and insulin. *J Mol Biol* **79**:351–371. doi:10.1016/0022-2836(73)90011-9

Spack E, Edidin M. 1986. The class I MHC antigens of erythrocytes: a serologic and biochemical study. *J Immunol Baltim Md 1950* **136**:2943–2952.

Sprague B, Mcnally J. 2005. FRAP analysis of binding: proper and fitting. *Trends Cell Biol* **15**:84–91. doi:10.1016/j.tcb.2004.12.001

Strasser J, de Jong RN, Beurskens FJ, Schuurman J, Parren PWHI, Hinterdorfer P, Preiner J. 2020. Weak Fragment Crystallizable (Fc) Domain Interactions Drive the Dynamic Assembly of IgG Oligomers upon Antigen Recognition. *ACS Nano* **14**:2739–2750. doi:10.1021/acsnano.9b08347

Townsend A, Bodmer H. 1989. Antigen recognition by class I -restricted cytotoxic T lymphocytes. *Annu Rev Immunol* 601–24.

Townsend A, Elliott T, Cerundolo V, Foster L, Barber B, Tse A. 1990. Assembly of MHC class I molecules analyzed in vitro. *Cell* **62**:285–295. doi:10.1016/0092-8674(90)90366-M

Townsend A, Öhlén C, Bastin J, Ljunggren H-G, Foster L, Kärre K. 1989. Association of class I major histocompatibility heavy and light chains induced by viral peptides. *Nature* **340**:443–448. doi:10.1038/340443a0

Triantafilou K, Triantafilou M, Wilson KM, Fernandez N. 2000. Human major histocompatibility molecules have the intrinsic ability to form homotypic associations. *Hum Immunol* **61**:585–598. doi:10.1016/S0198-8859(00)00112-9

Vámosi G, Friedländer-Brock E, Ibrahim SM, Brock R, Szöllősi J, Vereb G. 2019. EGF Receptor Stalls upon Activation as Evidenced by Complementary Fluorescence Correlation Spectroscopy and Fluorescence Recovery after Photobleaching Measurements. *Int J Mol Sci* **20**:E3370. doi:10.3390/ijms20133370

van den Heuvel RHH, van Duijn E, Mazon H, Synowsky SA, Lorenzen K, Versluis C, Brouns SJJ, Langridge D, van der Oost J, Hoyes J, Heck AJR. 2006. Improving the Performance of a Quadrupole Time-of-Flight Instrument for Macromolecular Mass Spectrometry. *Anal Chem* **78**:7473–7483. doi:10.1021/ac061039a

Váradi T, Schneider M, Sevcsik E, Kiesenhofer D, Baumgart F, Batta G, Kovács T, Platzer R, Huppa JB, Szöllősi J, Schütz GJ, Brameshuber M, Nagy P. 2019. Homo- and Heteroassociations Drive Activation of ErbB3. *Biophys J* **117**:1935–1947. doi:10.1016/j.bpj.2019.10.001

Vogelsang J, Kasper R, Steinhauer C, Person B, Heilemann M, Sauer M, Tinnefeld P. 2008. A Reducing and Oxidizing System Minimizes Photobleaching and Blinking of Fluorescent Dyes. *Angew Chem Int Ed* **47**:5465–5469. doi:10.1002/anie.200801518

Wilmes S, Beutel O, Li Z, Francois-Newton V, Richter CP, Janning D, Kroll C, Hanhart P, Hötte K, You C, Uzé G, Pellegrini S, Piehler J. 2015a. Receptor dimerization dynamics as a regulatory valve for plasticity of type I interferon signaling. *J Cell Biol* **209**:579–593. doi:10.1083/jcb.201412049

Wilmes S, Beutel O, Li Z, Francois-Newton V, Richter CP, Janning D, Kroll C, Hanhart P, Hötte K, You C, Uzé G, Pellegrini S, Piehler J. 2015b. Receptor dimerization dynamics as a regulatory valve for plasticity of type I interferon signaling. *J Cell Biol* **209**:579–593. doi:10.1083/jcb.201412049

Wilmes S, Hafer M, Vuorio J, Tucker JA, Winkelmann H, Löchte S, Stanly TA, Pulgar Prieto KD, Poojari C, Sharma V, Richter CP, Kurre R, Hubbard SR, Garcia KC, Moraga I, Vattulainen I, Hitchcock IS, Piehler J. 2020a. Mechanism of homodimeric cytokine receptor activation and dysregulation by oncogenic mutations. *Science* **367**:643–652. doi:10.1126/science.aaw3242

Wilmes S, Hafer M, Vuorio J, Tucker JA, Winkelmann H, Löchte S, Stanly TA, Pulgar Prieto KD, Poojari C, Sharma V, Richter CP, Kurre R, Hubbard SR, Garcia KC, Moraga I, Vattulainen I, Hitchcock IS, Piehler J. 2020b. Mechanism of homodimeric cytokine receptor activation and dysregulation by oncogenic mutations. *Science* **367**:643–652. doi:10.1126/science.aaw3242

You C, Richter CP, Löchte S, Wilmes S, Piehler J. 2014. Dynamic Submicroscopic Signaling Zones Revealed by Pair Correlation Tracking and Localization Microscopy. *Anal Chem* **86**:8593–8602. doi:10.1021/ac501127r

Figure legends

Figure 1: MHC I HC/HC association requires dissociation of β_2m but no disulfide
bonds.

A, Schematic of MHC class I states at the plasma membrane. Dissociation of peptide from
the HC/β_2m/peptide trimer results in an 'empty' HC/β_2m heterodimer. Dissociation of β_2m
then produces free HCs (FHCs), which can form FHC associations (HC/HC dimer and oli-
gomer shown). Other forms such as disulfide-linked dimers are known depending on the
allotype (see the text).

B, Schematic representation of the two-hybrid antibody micropattern assay. Cells expressing
a class I GFP fusion (green) and an N-terminally HA-tagged class I (gray) are seeded onto
glass slides that are printed with micrometer-sized patterns of fluorescently labelled anti-HA
antibodies. Dissociation of β_2m generates FHCs of both constructs, which diffuse freely in
the plasma membrane and eventually associate with each other to form HC/HC dimers (cen-
ter) or oligomers (not shown). The HC/HC associations, which contain both HA-tagged and
GFP-fused FHCs, localize in the pattern elements and are visible as pattern-shaped GFP fluo-
rescence on the plasma membrane.

C, Representative fluorescence micrograph showing one single STF1 cell expressing both
HA-tagged and GFP-fused H-2K^b in phase contrast (left), the purple anti-HA antibody pat-
tern on the glass slide (middle), and green K^b-GFP colocalizing with the antibody pattern
(right). The arrows point to the fluorophore observed in the respective panel and emphasize
the plane of the image. Scale bar, 20 μm.

D, Interaction occurs between HA-tagged K^b and D^b-GFP FHCs (37 °C) but not between
HC/β_2m heterodimers (25 °C). Scale bar, 20 μm.

E, HC/HC interaction does not involve intracellular disulfide bond formation, since the FHCs
(37 °C) of HA–K^b and K^b(C332S) GFP, which lacks the cytosolic cysteine, interact in the
micropattern assay. Scale bar, 20 μm.

F, B27, but not K^b, forms covalent dimers. HA-K^b (K^b in the label) and HA-B*27:05 (B27)
molecules were immunoprecipitated from the lysate of transduced STF1 cells with an anti-
HA monoclonal antibody, separated by reducing (+) or nonreducing (–) SDS-PAGE, and
monomers (heavy chain) and covalent homodimers as indicated were detected by Western
blotting with an anti-HA antiserum. * denotes a background band.

G, No interaction of K^b–GFP with F◻pocket-stabilized HA–K^b(Y84C/A139C), a disulfide-
stabilized K^b variant with increased β_2m affinity. Scale bar, 20 μm.

Figure 2: FHCs interact in TAP-proficient cells and in the absence of patterns.

A, Schematic of cell surface interaction between FHCs in TAP2-deficient (left) and TAP2-
proficient cells.

B, Time course of the interaction between HA-K^b and K^b-GFP in the anti-HA antibody
micropattern assay upon temperature shift from 25 °C to 37 °C in TAP2-proficient

(STF1/TAP2, right) and TAP2-deficient cells (STF1, left). Representative TIRF microscopy images are shown. 2 µM of cognate SIINFEKL peptide were added as indicated. Scale bar, 8 µm. Insets show enlarged intensity-adjusted regions within the selected cells.

C, Quantification of fluorescence contrast from B. Error bars are standard error of the mean (SEM) of >10 cells measured in ≥2 independent experiments.

D, FRAP analysis of K^b cell surface dynamics. Cells expressing K^b constructs as in B were monitored in the absence of patterns (top) or on anti-HA antibody patterns (bottom) with or without SIINFEKL peptide as indicated. Regions in the red circle were bleached, and recovery of fluorescence was followed over time. Scale bar, 5 µm. Insets show the enlarged intensity-adjusted region of bleached spot.

E, Normalized mean fluorescence recovery curves from FRAP experiments, error bars represent SEM of >20 cells for each condition measured in >2 independent experiments.

F, Diffusion coefficients calculated from E. Significant increase in diffusion coefficients upon peptide addition, both in the absence and presence of antibody patterns. ****, $p \leq 0.0001$ by two-way ANOVA. ns, not significant. Error bars are SEM of >20 cells for each condition measured in ≥2 independent experiments.

G, Immobile fraction of K^b–GFP molecules, i.e. molecules that remain in the bleached area during the time of the experiment. With peptide vs. without peptide ****, $p \leq 0.0001$ and on pattern vs. in the absence of pattern ****, $p \leq 0.0001$ by two-way ANOVA. Error bars are SEM of >20 cells for each condition measured in ≥2 independent experiments.

H, Exchange rates of K^b molecules calculated from fluorescence recovery curves. k_{fast} (left Y axis, hatched bars) is the rate of diffusion of K^b–GFP molecules. k_{slow} (right Y axis, solid-color bars) is the rate of K^b–GFP exchange on (= binding to and dissociation from) HA–K^b. With peptide vs. without peptide ****, $p \leq 0.0001$ on anti-HA pattern by two-way ANOVA. Error bars are SEM of >20 cells for each condition measured in ≥2 independent experiments.

Figure 3: Single-molecule microscopy shows K^b dimer formation.

A, Schematic on the left: analyzing FHC association by single-molecule microscopy. GFP-K^b☐molecules at the plasma membrane were labeled with two different fluorophores (ATTO Rho11, magenta, and ATTO 643, green) using anti-GFP nanobodies, with one GFP molecule binding one nanobody. Diffusion and interaction of individual molecules was quantified by tracking and co-tracking of individual molecules. Right: Top view of the plasma membrane with diffusing and co-diffusing nanobody-labeled GFP-K^b☐molecules (corresponding data are shown in **C**).

B, Formation and dissociation of an individual GFP-K^b dimer detected by co-tracking analysis. Co-locomoting molecules are highlighted by a white square. Scale bar, 400 nm.

C, Single molecule trajectories (green, magenta) and co-trajectories (black) observed for GFP-K^b FHCs (no peptide, top) and HC/β_2m/peptide trimers (With peptide, middle). As a control, the same experiment was carried out after crosslinking peptide-loaded K^b with a tandem anti-GFP nanobody (With crosslinker, bottom). Scale bar, 2 µm.

D-H, statistical analyses by two-sample Kolmogorov-Smirnov test (****, p ≤0.0001; ***, p ≤0.001; **, p ≤0.01). Sample numbers see the methods section.

D, Fraction of detected co-trajectories out of the total number of observed trajectories for K^b in the absence and in the presence of peptide and crosslinker as indicated.

E, Diffusion coefficients of K^b molecules determined by single-molecule tracking in the absence and presence of peptide and crosslinker.

F, The fraction of immobile molecules out of the total GFP-K^b surface molecules determined by single-molecule tracking. SIINFEKL peptide and crosslinker were added as indicated.

G, Comparison of the fractions of mobile and immobile GFP-K^b molecules that are associated. The fraction of (associated immobile K^b)/(total immobile K^b) is shown vs. the fraction of (associated mobile K^b)/(total mobile K^b).

H, Quantification of co-trajectory half-life for GFP-K^b as obtained from single molecule co-trajectories with SIINFEKL peptide and crosslinker as indicated; the dissociation rates calculated from the means are 3.1 (no peptide), 17 (peptide, no crosslinker), and 1.3 s^{-1} (with peptide and crosslinker).

Figure 4: The K^b α_3 domains form dimers.

A, HC/HC association between HA-K^b-RFP and α_3-GFP. Distance between pattern elements, 2 µm.

B, HC/HC association between HA-α_3 and α_3-GFP. Distance between pattern elements, 2 µm.

C, Size exclusion chromatography of the K^b α_3 domain expressed in E.coli and folded in vitro.

D, Nonreducing SDS-PAGE of the α_3 domain homodimer fraction from C shows absence of disulfide linkage between the monomers. The positive control for disulfide oligomer formation is a_3 inclusion bodies, isolated under oxidizing conditions, where some molecules are linked by disulfide bonds to form covalent oligomers as indicated on the left.

E, MS/MS analysis of the α_3 domain at 10 µM. The 9+ peak at 2551 m/z corresponding to the a_3 dimer was selected for MS/MS analysis at 10 V, 50 V and 100 V in the collision cell. The dimer (double spheres) easily dissociates in the gas-phase indicating a non-covalent rather low-affinity binding event. At 50 V, no dimer is detectable anymore displaced by monomeric signal (single spheres).

F, Overall area under the curve (AUC) for the detected α_3 domain dimers. Native mass spectra were recorded using 5 µM, 10 µM or 20 µM α_3 domain at 10 V (dark red) and 25 V (light red), respectively. The AUC was determined over the entire spectrum for the dimeric mass species and is shown as mean (n=3) ±SD. The dimeric fraction is concentration-dependent.

G, Representative mass spectra (20 µM α_3 domain) show the charge distributions of monomer (single sphere) and dimer (double spheres) at 10 V (dark red) and 25 V (light red) in the collision cell.

Figure 5: One hypothetical model of a K^b FHC dimer.

In this hypothetical structure of the extracellular portion of an FHC dimer (**A**), one of several
structures predicted by computational molecular docking and molecular dynamics simulation,
the α_3 domain of one FHC (orange) binds to another FHC (yellow/purple/lavender) in a man-
ner resembling the binding of β_2m in the crystal structure of peptide-loaded K^b (**B**, PDB
3P9L).

Tables

Table 1: Comparison of diffusion coefficients (mean $\pm$ SD in μm^2/s). NA, not applicable.

	FRAP	SMT
HC/β_2m/peptide trimers (+peptide)	0.35 ±0.12	0.35±0.06
FHCs (–peptide)	0.19 ±0.05	0.25±0.04
FHCs (–peptide)[a]	0.19 ±0.05	0.18±0.10
Dimerized HC/β_2m/peptide trimers (+peptide, + CL)	NA	0.20±0.05
Dimerized HC/β_2m/peptide trimers (+peptide, + CL)[a]	NA	0.15±0.04

[a]only dimers identified by SMCT

A heavy chain/β2m/peptide **trimer** | heavy chain/β2m **heterodimer** | free heavy chain (**FHC**) | FHC **dimer** | FHC **oligomer**

B Cell | HC/β2m/ peptide | HC/ β2m | FHC | FHC dimers on anti-HA micropatterns | HC-GFP fusion constructs | Glass slide | HA-tagged HC constructs

C HA-Kᵇ + Kᵇ-GFP | Phase contrast | Anti-HA pattern | GFP | 37 °C

D HA-Kᵇ + Dᵇ-GFP | 25 °C | 37 °C

E HA-Kᵇ + Kᵇ(C332S)-GFP | 25 °C | 37 °C

F Kᵇ | B27 | DTT | Homodimer | * | Heavy chain | 100 kDa | 46 | 32

G HA-Kᵇ + Kᵇ(Y84C/A139C)-GFP | 25 °C | 37 °C

Dirscherl *et* al., Figure 5

A Kᵇ–Kᵇ FHC dimer

α₁/α₂ domains

α₃ domain

α₃ domain

B Kᵇ HC–β₂m heterodimer

α₁/α₂ domain

β₂m

α₃ domain