Title

Megafaunal extinctions—not climate change—seem to explain Holocene genetic diversity declines in *Numenius* shorebirds

Author names and affiliations

Hui Zhen Tan[1], Justin J.F.J. Jansen[2], Gary A. Allport[3], Kritika M. Garg[1,4,5], Balaji Chattopadhyay[1,6], Martin Irestedt[7], Sean E.H. Pang[1], Glen Chilton[8], Chyi Yin Gwee[1,9], Frank E. Rheindt[1*]

[1]Department of Biological Sciences, National University of Singapore, 16 Science Drive 4, Singapore 117558, Singapore

[2]Naturalis Biodiversity Center, Leiden, P.O. Box 9517, 2300 RA Leiden, Netherlands

[3]BirdLife International, David Attenborough Building, Pembroke Street, Cambridge, UK CB2 3QZ, UK

[4](Present address) Centre for Interdisciplinary Archaeological Research, Ashoka University, Sonipat, Haryana, 131029, India

[5](Present address) Department of Biology, Ashoka University, Sonipat, Haryana, 131029, India

[6](Present address) Trivedi School of Biosciences, Ashoka University, Sonipat, Haryana, 131029, India

[7]Department of Bioinformatics and Genetics, Swedish Museum of Natural History, P. O. Box 50007, SE-104 05 Stockholm, Sweden

[8]Department of Biology, St. Mary's University, Calgary, Alberta, T2X 1Z4, Canada

[9](Present address) Division of Evolutionary Biology, Faculty of Biology, Ludwig Maximilian University of Munich, 82152 Planegg-Martinsried, Germany

*Corresponding Author; e-mail: dbsrfe@nus.edu.sg

Abstract

Understanding the relative contributions of historical and anthropogenic factors to declines in

genetic diversity is important for informing conservation action. Using genome-wide DNA of

fresh and historic specimens, including that of two species widely thought to be extinct, we

investigated fluctuations in genetic diversity and present the first complete phylogenomic tree for

all nine species of the threatened shorebird genus *Numenius*, known as whimbrels and curlews.

Most species faced sharp declines in effective population size, a proxy for genetic diversity, soon

after the Last Glacial Maximum (around 20,000 years ago). These declines occurred prior to the

Anthropocene and in spite of an increase in breeding area predicted by environmental niche

modelling, suggesting that they were not caused by climatic or recent anthropogenic factors.

Crucially, these genetic diversity declines coincide with mass extinctions of mammalian

megafauna in the Northern Hemisphere. Demise of ecosystem-engineering megafauna which

maintained open habitats would have been detrimental for grassland and tundra-breeding

Numenius shorebirds. Our work suggests that the impact of historical factors such as megafaunal

extinction may have had wider repercussions on present-day population dynamics of open

habitat biota than previously appreciated.

Introduction

Rates of population decline and extinction have risen sharply during the ongoing sixth mass extinction crisis (Ceballos, Ehrlich, & Raven, 2020; Dirzo & Raven, 2003; Sánchez-Bayo & Wyckhuys, 2019; S. N. Stuart et al., 2004). Species distribution models based on future climate scenarios forecast that rates of endangerment will further accelerate, underscoring the need for conservation action (Thomas et al., 2004). In this era of increasing biodiversity loss, the maintenance of genetic diversity within species has become a focus of conservation as it is thought to predict evolutionary adaptability and extinction risk (Frankham, 2005; Hoban et al., 2020; Jetz et al., 2014). Modern declines in genetic diversity have been documented for a handful of species (Allentoft & O'Brien, 2010; Chattopadhyay, Garg, Mendenhall, & Rheindt, 2019; S. R. Evans & Sheldon, 2008; Garner, Rachlow, & Hicks, 2005), but we continue to know little about the global mechanisms of genetic diversity loss.

Anthropogenic climate change is widely recognised for its pervasive impact on biodiversity and genetic diversity (C. N. Johnson et al., 2017; Miraldo et al., 2016; Turvey & Crees, 2019). However, historical events have equally left their signature in the genetic profiles of present-day species (Hewitt, 2000). Comparative genomics of extinct versus extant species could add an important perspective to elucidating such trends in faunal endangerment (Frankham, 2005).

We used a museomic approach to investigate fluctuations in effective population size in all nine species of the migratory shorebird genus *Numenius*, known as whimbrels and curlews, including two species, the slender-billed curlew (*N. tenuirostris*) and Eskimo curlew (*N. borealis*), that are presumed to be extinct (Buchanan et al., 2018; Butchart et al., 2018; Kirwan, Porter, & Scott, 2015; Pearce-Higgins et al., 2017; Roberts, Elphick, & Reed, 2010; Roberts & Jarić, 2016). Members of the genus *Numenius* breed across the Northern Hemisphere's tundras

and temperate grasslands, and are particularly vulnerable to endangerment due to comparatively

long generation times (Pearce-Higgins et al., 2017).

Our objective was to characterise genetic diversity fluctuations in *Numenius* shorebirds,

assess the relative impact of historical and anthropogenic factors on these fluctuations, and

determine the mechanisms that may have had the biggest impact on their populations. Because of

their dependence on open habitats, we expected the genetic diversity trends of whimbrels and

curlews to track the availability of such habitats across the late Quaternary. We also expected

significant declines in genetic diversity during the late Holocene when global human activity

intensified, not least because the demise of the two extinct species has been attributed to habitat

loss and hunting (Committee on the Status of Endangered Wildlife in Canada, 2009; Gallo-Orsi

& Boere, 2001). By testing the timing of genetic diversity fluctuations against that of important

ecological events, we could home in on the factors that influenced the evolutionary trajectory of

this threatened shorebird lineage over the last ~20,000 years.

Results and discussion

We sequenced 67 ancient and fresh samples across all nine *Numenius* species for target

enrichment (Figure 1A; Table S1). After filtering for quality, a final alignment of 514,771bp

across 524 sequence loci was retained for each of 62 remaining samples at a mean coverage of

118X. Phylogenomic analyses using MP-EST (L. Liu, Yu, & Edwards, 2010) revealed two

separate groups, here called the "whimbrel clade" and the "curlew clade", that diverged

approximately 5 million years ago (Figure 1B; Figure S1A). This is the first phylogenomic tree

to include all members of the genus *Numenius*. The use of degraded DNA from toepads of

museum specimens allowed us to include the two presumably extinct taxa. Of these, the slender-

billed curlew emerged as sister to the Eurasian curlew (*N. arquata*), a phenotypically similar

species that occurs in sympatry in Central Asia (Sharko et al., 2019). On the other hand, the

Eskimo curlew emerged as a distinct member of the curlew clade with no close relatives (Figure

1B). Our phylogenomic dating analyses demonstrated that 40.6% of the evolutionary distinctness

(Jetz et al., 2014) of the curlew clade has been lost with the presumable extinction of the two

species, and that another 15% is endangered (Figure 1B; Table S2).

To characterise the differential impacts of extinction pressures, we reconstructed the

demographic history of *Numenius* shorebirds. For five species with a sufficiently high sample

size, we employed stairway plots (X. Liu & Fu, 2020) to infer fluctuations in effective

population size (N_e), a proxy for genetic diversity, given that this method works well for reduced

representation genomic datasets such as ours, and has a relatively high accuracy for

reconstructions of diversity changes in the Late Quaternary (X. Liu & Fu, 2020). Fluctuations in

N_e were compared against key biotic and anthropogenic events of the Late Quaternary. We also

accounted for climatic changes by modelling the extent of suitable breeding areas of each species

under present day (1960–1990), mid-Holocene (6,000 years ago), and LGM (22,000 years ago)

climate conditions using the MaxEnt algorithm (Phillips, Anderson, & Schapire, 2006).

The Last Glacial Period preceding the LGM saw ice sheets at their maximum extent

(Hughes, Gibbard, & Ehlers, 2013). During this time, tundra habitats dominated the northern

latitudes and an increase in N_e in the tundra-inhabiting Eurasian whimbrel (*N. phaeopus*) was

observed (Binney et al., 2017; Y. Wang et al., 2021; Zimov et al., 1995; Figure 1C). Soon after,

during the Pleistocene-Holocene transition, our stairway plots revealed generally sharp declines

of N_e in most species despite an increase in area of suitable breeding habitat predicted (Figure

1C). The extent of breeding habitat predicted by our ecological niche models relied on

bioclimatic variables, suggesting that favourable conditions for *Numenius* shorebirds in the lead-

up to the Holocene did not trigger an increase in genetic diversity, but instead coincided with

precipitous declines in N_e. To understand the drivers of N_e declines in *Numenius* shorebirds,

factors other than climate change would need to be considered.

During the Pleistocene-Holocene transition (starting at roughly 20,000 years ago), a mass

extinction of megafaunal mammals (≥ 44 kg) was underway, known as the Late Quaternary

Extinctions (Hedberg, Lyons, & Smith, 2022; C. N. Johnson, 2009), with most becoming extinct

by 10kya (Figure 1C) (Koch & Barnosky, 2006; A. J. Stuart, 2015). Megafaunal mammals are

ecosystem-engineers that maintain open landscapes such as temperate grasslands and steppes

through grazing, browsing and physical impacts (Bakker et al., 2016; Goheen et al., 2018).

During the intervening period between their extinction and the spread of ungulate domestication,

there would have been no functional replacements for these ecosystem services (Hedberg et al.,

2022; Lundgren et al., 2020). Open grasslands would have been subject to increasing forest

succession (C. N. Johnson, 2009) and the amount of suitable habitat for *Numenius* shorebirds

might have been less than predicted by forecasts relying only on bioclimatic variables.

Therefore, genetic diversity fluctuations in *Numenius* shorebirds run counter to expectations

based on natural climate change and seem to be best explained by the demise of the ecosystem-

engineers that would have helped maintain shorebird breeding habitats.

By the late Holocene, the genetic diversity of most *Numenius* shorebirds stabilised at a time

when anthropogenic impact was only starting to expand across the Northern Hemisphere with a

steep rise in human population and land conversion for agriculture (Figure 1C). The timing of

these events is inconsistent with the hypothesis that direct anthropogenic activity has been the

main cause of genetic diversity declines in *Numenius* (Crisp, Trewick, & Cook, 2011). Events

unrelated to modern anthropogenic pressure seem to have played a bigger role in the declines

observed in *Numenius* shorebirds (Lucena-Perez et al., 2020; Nadachowska-Brzyska, Li, Smeds,

Zhang, & Ellegren, 2015). It is possible that additional adverse effects caused by more recent

anthropogenic impacts are not yet reflected in the genomes investigated, perhaps exacerbated by

shorebirds' long generation times.

At present, members of the curlew clade, which predominantly breed in temperate grasslands

at lower latitudes, generally exhibit levels of N_e that are lower than those of the higher latitude-

breeding whimbrels (Figure 1C). Temperate grasslands face far greater anthropogenic pressures

from land use than the northerly tundra (Pimm et al., 2014), contributing to further declines in

curlews more so than in whimbrels. Genetic diversity estimates were lowest in the presumably

extinct slender-billed curlew *N. tenuirostris* (Figure 1C). Low genetic diversity may contribute to

a species' extinction risk (Frankham, 2005; Spielman, Brook, & Frankham, 2004), although such

links must be examined for each species independently and could possibly be conflated with

other factors such as total population size (S. R. Evans & Sheldon, 2008; Teixeira & Huber,

2021).

Our study uncovers substantial declines in genetic diversity in curlews and whimbrels across

the Late Quaternary. Analysing N_e fluctuations over time allowed us to test which factors may

have coincided with genetic diversity declines. Of the factors investigated, megafaunal

extinctions—not natural climate change in the post-glacial period—best explain these declines

and may have had cascading effects on species' evolutionary trajectories that continue to impact

them to the present day. Our results underscore that grassland biomes and their biota face unique

challenges that warrant more conservation attention (Ceballos et al., 2010; Chan, Lacey, Pearson,

& Hadly, 2005; Helm et al., 2009; Nakahama, Uchida, Ushimaru, & Isagi, 2018; Török,

Ambarlı, Kamp, Wesche, & Dengler, 2016; Wesche et al., 2016). Our work demonstrates that

relatively brief evolutionary events, such as the Late Quaternary Extinctions of megafauna, can

have long-lasting evolutionary effects on populations, in our case for roughly ~10,000 years. The

plight of *Numenius* shorebirds is a sobering reminder of the importance of conserving remaining

genetic diversity to ensure the resilience of our planet's biota.

Figure 1. (A) Breeding distribution map and sampling localities of each *Numenius* species (BirdLife International and Handbook of the Birds of the World, 2017; Lappo, Tomkovich, & Syroeckovskiy, 2012); wintering and migratory ranges are not shown. Colours correspond to species identities in (B). Diagonal lines denote regions with co-distributed species. Each circle represents one sample unless otherwise specified by an adjacent number. The only known breeding records of *N. tenuirostris* were from near the village of Krasnoperova c.10 km south of Tara, Omsk (Russia), which is denoted by a black star (★), although this might not have been the core breeding area. (B) Phylogenomic tree constructed from an alignment of 514,771bp across 524 sequence loci. Tree topology (including bootstrap support values) and divergence times were estimated with MP-EST and MCMCTree respectively. Only bootstrap <100 is displayed. Sample sizes for each species are given in brackets. IUCN Red List status of critically endangered (CR) and endangered (EN) species are indicated. Illustrations of *Numenius* birds were reproduced with

permission from Lynx Edicions. (C) Results of demographic history reconstruction using stairway plot for selected
species displayed with key climatic, biotic and anthropogenic events. Effective population size: Line colours
correspond to species identities in the tree in (B) and numbers at present time represent present-day effective
population sizes. Thick lines represent the median effective population size while thin lines represent the 2.5 and
97.5 percentile estimation. The vertical grey line denotes the Last Glacial Maximum (LGM) and panels are shaded
to aid reference to the time axis. Suitable breeding area: predicted suitable breeding area at LGM (22,000 years ago),
mid-Holocene (6,000 years ago) and present day (1960–1990) using Maxent. Dot colours correspond to species
identities in the tree in (B). Dotted lines connecting the dots are for visualisation purposes and do not represent
fluctuations in breeding area. The following panels display the timings of key climatic, biotic and anthropogenic
events including megafaunal extinction (in terms of number of extinct genera with dotted shading denoting
uncertainty in estimates; Koch & Barnosky, 2006), agricultural land use and human population size (HYDE 3.2;
Goldewijk, Beusen, Doelman, & Stehfest, 2017; Goldewijk, Beusen, & Janssen, 2010). Line type corresponds to
geographical area (Nearctic versus Palaearctic) as denoted in the 'Human population size' panel.

Materials and methods

Taxon sampling

We acquired samples for all nine species in the genus *Numenius*, encompassing most of

the known subspecies. Species and subspecies identities are as provided by the source museum

or institution (Table S1) or assigned in reference to known breeding and wintering locations

(Birds of the World, 2022). We also included one common redshank *Tringa totanus* as an

outgroup for phylogenetic rooting. All samples were acquired through museum loans except for

an individual of the endangered subspecies *N. phaeopus alboaxillaris* that was sampled during

fieldwork by GAA (Table S1). Where possible, we acquired fresh samples (tissue or blood)

because of their higher DNA quality. To represent rarely-sampled or presently-rare taxa for

which no fresh samples were available, we acquired toepad material from historic museum

specimens and applied ancient DNA methods.

Baits design for target capture

We used the *Calidris pugnax* genome (accession no. GCA_001458055.1) (Küpper et al.,

2015) to design baits to capture selected exons. We used EvolMarkers (C. Li, Riethoven, &

Naylor, 2012) to identify single-copy exons conserved between *C. pugnax, Taeniopygia guttata*

(accession no. GCF_003957565.1; released by the Vertebrate Genomes Project) and *Ficedula*

albicollis (accession no. GCA_000247815.1). Exons longer than 500 bp with a minimum

identity of 55% and an e-value $< 10e^{-15}$ were isolated with bedtools 2.28.0 (Quinlan & Hall,

2010), forming our target loci. Only target loci with 40–60% GC content were retained and any

overlapping loci were merged (Quinlan & Hall, 2010). Target loci with repeat elements were

then filtered out in RepeatMasker 4.0.6 (Smit, Hubley, & Green, 2015). We arrived at a final set

of 565 unique target loci with a mean length of 970bp. These target loci were used to design

19,003 100 bp-long biotinylated RNA baits at 4X tiling density (MYcoarray/Arbor Biosciences,

USA).

<u>Laboratory methods</u>

Both fresh and historic samples were subjected to DNA extraction, followed by library

preparation and target enrichment, with slight modifications for various sample types to optimise

yield. DNA extractions of fresh samples were performed using the DNEasy Blood & Tissue Kit

(Qiagen, Germany) with an additional incubation step with heat-treated RNase. Extractions for

historic samples were performed using the same kit but with modifications (Chattopadhyay et al.,

2019). Historic samples were washed with nuclease-free molecular grade water before extraction

and dithiothreitol was added to the digestion mix. DNA precipitation was performed for at least

12 hours and MinElute Spin Columns were used for elution (Qiagen, Germany). Historic

samples were processed in a dedicated facility for highly degraded specimens.

DNA extracted from fresh samples was sheared via sonification using Bioruptor Pico

(Diagenode, Belgium) to a target size of 250 bp. DNA extracts from historic samples were

generally smaller than the target size; hence no further shearing was performed. Whole-genome

libraries were prepared using the NEBNext Ultra™ II DNA Library Prep Kit for Illumina (New

England Biolabs, Ipswich, USA) with modifications for subsequent target enrichment. For fresh

samples, adaptor concentrations were kept constant regardless of DNA input amount. Size

selection with AMPure XP beads (Beckman Coulter, USA) was performed for 250 bp insert

sizes. The reaction was split into two equal parts before polymerase chain reaction (PCR)

amplification and combined afterwards for subsequent steps. For historic samples, a formalin-

fixed, paraffin-embedded (FFPE) DNA repair step was first performed using NEBNext FFPE

DNA Repair Mix (New England BioLabs). A 10-fold dilution of adaptors was used, and no size

selection was performed. For both types of samples, twelve cycles of PCR amplification were

performed.

Target enrichment was carried out following the MYbaits manual (Arbor Biosciences,

USA) with modifications (Chattopadhyay et al., 2019). We used 1.1 uL of baits per fresh sample

(~5X dilution) and 2.46 uL of baits per ancient sample (~2X dilution). For fresh samples,

hybridization of baits and target loci was performed at 65°C for 20 hours and 15 cycles of

amplification were performed. For historic samples, hybridization was performed at 60°C for 40

hours, and 20 cycles of amplification were performed. For both fresh and historic samples, one

negative control sample was added for each batch of extraction, library preparation and target

enrichment. Extracts, whole-genome libraries, final enriched libraries, and all negatives were

checked for DNA concentration on a Qubit 2.0 Fluorometer using the Qubit dsDNA HS assay kit

(ThermoFisher Scientific, USA), and for fragment size on a Fragment Analyzer using the HS

NGS Fragment kit (1–6000 bp) (Agilent Technologies Inc., USA). Final enriched libraries were

pooled at equimolar quantities. A total of 67 enriched libraries were sequenced, with fresh and

historic samples sequenced separately on two Illumina HiSeq 150 bp paired-end lanes

(NovogeneAIT, Singapore).

<u>Reference genome assembly</u>

We obtained a sample of *N. phaeopus* (ZMUC 112728) from the Natural History

Museum of Denmark, Copenhagen, for reference genome assembly. Its genomic DNA was

extracted using the KingFisher™ Duo Prime Magnetic Particle Processor (ThermoFisher

Scientific, USA) and the KingFisher Cell and Tissue DNA Kit (Thermo Fisher Scientific). A

linked-read sequencing library was prepared using the Chromium Genome library kits (10X

Genomics) and sequenced on one Illumina Hiseq X lane at SciLifeLab Stockholm (Sweden). The

de novo assembly analysis was performed using 10X Chromium Supernova (v. 2.1.1). Reads

were filtered for low quality and duplication, while assemblies were checked for accuracy and

coverage and the best assembly was selected based on the highest genome coverage with the

fewest errors. The final genome had a size of 1.12 Gb at a coverage of 50X with N50 =

3504.2kbp.

<u>Raw reads processing</u>

Raw reads were checked for sequence quality in FastQC 0.11.8 (Babraham

Bioinformatics) and trimmed to remove low-quality termini and adaptors in fastp 0.20.0 (Chen,

Zhou, Chen, & Gu, 2018). We retained reads with a minimum length of 36bp and set a phred

quality threshold of 20. Retained reads started at the first base satisfying minimum quality

criteria at the 5'-end and were truncated wherever the average quality fell below the threshold in

a sliding window of 5bp. Duplicates were removed using FastUniq 1.1 (Xu et al., 2012) before

sequence quality, duplication rate and adaptor content were checked again in FastQC. We

employed FastQ Screen 0.14.0 (Wingett & Andrews, 2018) to assign the source of DNA against

a list of potential contaminants. We aligned reads to our assembled *Numenius phaeopus* genome,

Homo sapiens (accession no. GCF_000001405.39), and a concatenated database of all bacterial

genomes available on GenBank (National Center for Biotechnology Information (NCBI), 1988).

Only reads that mapped uniquely to the *N. phaeopus* genome were retained. Reads were sorted

and re-paired using BBtools 37.96 (Bushnell, 2014). Downstream bioinformatic procedures were

split into single nucleotide polymorphism (SNP)-based and sequence-based analyses.

SNP calling

For SNP-based analyses, reads were aligned to the target sequences used for bait design

with bwa-mem 0.7.17 (Li, 2013). The output alignment files were converted to bam files (view)

and sorted by coordinates (sort) using SAMtools 1.9 (H. Li et al., 2009). Alignments were

processed in Picard 2.20.0 (Picard tools, Broad Institute, Massachusetts, USA) to add read group

information (AddOrReplaceReadGroups), and another round of duplicate identification was

performed (MarkDuplicates) before alignment files were indexed (BuildBamIndex). The

reference file of target sequences was indexed in SAMtools (faidx) and a sequence dictionary

was created in Picard (CreateSequenceDictionary). To improve SNP calling accuracy, indel

realignment was performed in GATK 3.8 (McKenna et al., 2010) (RealignerTargetCreator,

IndelRealigner). We inspected historic DNA alignments in mapDamage 2.0.9 (Jónsson,

Ginolhac, Schubert, Johnson, & Orlando, 2013) and trimmed up to 5bp from the 3' ends of both

reads to minimise frequencies of G to A misincorporation (<0.1) and soft clipping (<0.2).

Finally, alignments were checked for quality and coverage in QualiMap 2.2.1 (Okonechnikov,

Conesa, & García-Alcalde, 2016).

We first generated likelihoods for alignment files in BCFtools 1.9 (Li, 2011) (mpileup),

skipping indels. Using the same program, we then called SNPs (call) for all *Numenius* samples

using the multiallelic and rare-variant calling model. Called SNPs were filtered in VCFtools

0.1.16 (Danecek et al., 2011) to retain sites with quality values >30, mean depth 30–150, minor

allele frequency ≥0.02 and missing data <5%, in this order. Missingness and depth of sites and

individuals, respectively, were quantified for SNPs called. We removed eight individuals from

downstream analyses due to a combination of high missing data (>0.4%) and low coverage

(<36X), yielding a SNP set representing 58 samples. A Perl script (rand_var_per_chr.pl) was

used to call one SNP per locus to avoid calling linked SNPs (Caballero, 2018). SNPs were

further screened for linkage disequilibrium in PLINK 1.9 (Purcell et al., 2007) using a sliding

window of 50 SNPs with a step size of 10 and an r^2 correlation threshold of 0.9. We also

screened for neutrality of SNPs in BayeScan 2.1 (Foll & Gaggiotti, 2008) using default settings.

We additionally created a dedicated SNP set per species for input into demographic history

reconstruction using the method described above, but without minor allele frequency cut-offs

and with all SNPs at each locus retained.

<u>Population genomic analyses</u>

We conducted principal component analysis (PCA) for all *Numenius* samples using the R

package SNPRelate 1.16.0 (R Core Team, 2022; Zheng et al., 2012) (Figure S1A). We did not

detect any considerable genomic differentiation along subspecific delimitations within *N.*

phaeopus and *N. arquata*, whose population-genetic structure had been resolved with thousands

of genome-wide markers in a previous study (Tan et al., 2019) (Figure S1B, C). Samples of *N. p.*

alboaxillaris and *N. a. suschkini,* two Central Asian taxa that are described in the literature as

phenotypically differentiated (Allport, 2017; Engelmoer & Roselaar, 1998b, 1998a; Morozov,

2000), did not emerge as genomically distinct from other conspecific populations and are likely

to represent ecomorphological adaptations controlled by few genes. Sample NBME 1039630,

which had been labeled as *N. tenuirostris*, and sample MCZR 15733, which was initially

identified as an *N. arquata* that shares many morphological features with *N. tenuirostris*,

clustered with *N. arquata* samples (Table S1, Figure S1D). Both samples were assigned to *N.*

arquata in subsequent phylogenetic analyses.

Sequence assembly

For sequence-based analyses, reads were assembled using HybPiper 1.3.1 (M. G. Johnson

et al., 2016) (reads_first) to yield sequence loci. Firstly, reads were mapped to the target

sequences using BWA 0.7.17 (H. Li & Durbin, 2009) and sorted by gene. Contigs were then

assembled from the reads mapped to respective loci using SPAdes 3.13 (Bankevich et al., 2012)

with a coverage cutoff value of 20. Using Exonerate 2.4.0 (Slater & Birney, 2005), these contigs

were then aligned to the target sequences and sorted before one contig per locus was chosen to

yield the final sequences. We inspected locus lengths (get_seq_lengths) and recovery efficiency

(hybpiper_stats) across all loci. We then investigated potentially paralogous loci

(paralog_investigator) by building gene trees using FastTree 2.1.11 (Price, Dehal, & Arkin,

2010) (paralog_retriever), leading to the removal of 10 loci. All loci retained were present in at

least 80% of individuals and constituted at least 60% of the length of total target loci. In

summary, a total of 525 loci with a mean length of 969 bp (492–6,054 bp) were recovered from

62 samples.

Phylogenomic analyses using sequence data

Multisequence alignment was performed for each locus using MAFFT 7.470 (Katoh &

Standley, 2013), allowing for reverse complement sequences as necessary. Alignments were

checked for gaps using a custom script, and loci with >35% gaps were removed from

downstream analyses. A total alignment length of 514,771 bp was obtained.

Phylogenomic analyses were performed on a concatenated dataset as well as on

individual gene trees. Concatenation was performed with abioscript 0.9.4 (Larsson, 2010)

(seqConCat). For the concatenated dataset, we constructed maximum-likelihood (ML) trees

using RAxML 8.2.12 (Stamatakis, 2014) with 100 alternative runs on distinct starting trees. We

applied the general time reversible substitution model with gamma distributed rate variation

among sites and with estimation of proportion of invariable sites (GTR+I+G) (Abadi, Azouri,

Pupko, & Mayrose, 2019; Arenas, 2015).

For individual gene trees, the best substitution model for each locus was determined

using jModelTest 2.1.10 (Darriba, Taboada, Doallo, & Posada, 2012) by virtue of the corrected

Akaike information criterion value. We then constructed ML trees in PhyML 3.1 with the subtree

pruning and regrafting algorithm, using 20 initial random trees. We performed 100 bootstrap

replicates with ML estimates for both proportion of invariable sites and value of the gamma

shape parameter. Individual gene trees were then rooted with Newick Utilities 1.3.0 (Junier &

Zdobnov, 2010). We removed one locus from downstream analyses due to the absence of an

outgroup sequence such that 524 loci were retained across 62 samples.

Species tree analyses were performed using the rooted gene trees in MP-EST 1.6 (L. Liu

et al., 2010), without calculation of triple distance among trees. We grouped samples by species

and performed three runs of 10 independent tree searches per dataset (Cloutier et al., 2019). To

calculate bootstrap values of the species tree, we performed multi-locus, site-only resampling

(Mirarab, 2014) from the bootstrap trees' (100 per gene) output from PhyML. The resulting 100

files, each with 100 bootstrap trees, were rooted and species tree analyses were performed in the

same manner for each file in MP-EST. The best tree from each run was identified by the best ML

score and compiled. Finally, we used the majority rule in PHYLIP 3.695 (Felsenstein, 2009) to

count the number of times a group descending from each node occurred so as to derive the

bootstrap value (consense).

362   For estimation of divergence times, we applied MCMCtree and BASEML (dos Reis &

Yang, 2011), a package in PAML 4.9e (Yang, 2007). To prepare the molecular data from 62

samples and 524 loci, we compiled the DNA sequence of each sample and combined all samples

onto separate rows of the same file. We then obtained consensus sequences for each species

using Geneious Prime 2020.2 (Kearse et al., 2012), with a majority support threshold of 50% and

ignoring gaps. We visually checked the resulting consensus sequences to ensure that ambiguous

bases remained infrequent. Consensus sequences were organised by loci as per the input format

for MCMCtree. We then prepared the input phylogenetic tree using the topology estimated in

MP-EST with calibrations of the two most basal nodes, namely between our outgroup (*Tringa*

totanus) and all *Numenius* species, as well as that between the whimbrel and curlew clades

within *Numenius*. Due to a lack of known fossils within the genus *Numenius,* we were unable to

perform fossil node calibrations. Instead, we utilised p-distance values calculated from the COI

sequences of *Numenius* species. Specifically, we applied the bird COI mutation rate of 1.8% per

million years (Lavinia, Kerr, Tubaro, Hebert, & Lijtmaer, 2016) and converted mean, maximum

and minimum p-distance values of both nodes to time (100 million years ago (MYA)). We

maintained a conservative position and scaled the COI-based timings by a factor of two to obtain

the final lower and upper bounds of node timings. We used the default probability of 0.025 that

the true node age is outside the calibration provided.

380   To run MCMCtree, we first calculated the gradient and Hessian matrix of the branch

lengths with the GTR substitution model applied, using default values of gamma rates and

numbers of categories (mcmctree-outBV.ctl). We then performed two independent Markov chain

Monte Carlo (MCMC) samplings of the posterior distribution of divergence times and rates

(mcmctree.ctl). All default values were used except that a constraint on the root age was set to

<0.3 (100 MYA). We also varied the prior for the birth-death process with species sampling and

ensured that time estimates are not affected by the priors applied (dos Reis & Yang, 2019). We

then performed convergence diagnostics for both runs in R to ensure that posterior means are

similar among multiple runs, while checking that the parameter space has been explored

thoroughly by the MCMC chain. Finally, we conducted MCMC sampling from the prior with no

data to check the validity of priors used by comparing them with posterior times estimated.

Again, two independent MCMC samplings were performed with convergence diagnostics.

Phylogenetic trees were visualised in FigTree 1.4.4 (Rambaut, 2018) with bootstrap

values and node ages (MYA) including the 95% credibility intervals. Evolutionary distinctness

and phylogenetic diversity was calculated for each branch (Jetz et al., 2014) using the divergence

times estimated in MCMCTree.

<u>Demographic history reconstruction</u>

We derived trends in effective population size using stairway plot 2.1.1, which uses the

SNP frequency spectrum and is suitable for reduced representation datasets (X. Liu & Fu, 2020;

Patton et al., 2019). From the dedicated SNP sets that were created without minor allele

frequency cut-off, we calculated a folded site frequency spectrum using vcf2sfs.py 1.1 (Marques,

Lucek, Sousa, Excoffier, & Seehausen, 2019). We assumed a mutation rate per site per

generation of $8.11e^{-8}$, as estimated for shorebirds in the same order as *Numenius*

(Charadriiformes) (X. Wang et al., 2019), and applied the following generation times

respectively: *N. americanus* 7 years, *N. arquata* 10 years, *N. hudsonicus* 6 years, *N. phaeopus* 6

years, *N. tenuirostris* 5 years (Bird et al., 2020; IUCN, 2020). We ran stairway plot on all

species, applying the recommended parameters.

Stairway plot is expected to perform at its highest accuracy in the reconstruction of

demographic history in the recent rather than distant past. However, the definition of recent past

varies from anywhere between 30 generations to ~40,000 generations before present (X. Liu &

Fu, 2015, 2020; Patton et al., 2019). We did not set a cutoff for the time period investigated but

let it be determined by the program itself. Additionally, we omitted reconstructions of the last 10

steps to avoid overinterpretation of the distant past (X. Liu & Fu, 2015). We only displayed the

results from the time period for which there was data across all species, and only for four species

represented by five or more samples (stairway_plot_es Stairbuilder), as recommended for

accurate inference (X. Liu, personal communication, October 14, 2020). We later also included

N. americanus, for which we had four samples, as its sample size did not appear to affect the

reliability of results (Figure 1). We were unable to include the remaining species (*N. borealis*, *N.*

tahitiensis, and *N. minutus*) as their demographic history reconstructions were clearly affected by

a lack of sufficient sample size. For *N. borealis,* two out of the five samples showed high

missingness, with adverse effects on stairway plot analyses, both in runs including all five

samples and those that excluded the two samples of high missingness (Figure S2). Our ability to

trial a large number of samples for laboratory work was also limited by the availability of target

enrichment baits.

We attempted to infer demographic history using sequentially Markovian coalescent-

based methods, which are more reliable for older timescales, to corroborate our stairway plot

results (Patton et al., 2019). In particular, we used the Pairwise Sequentially Markovian

Coalescent (PSMC) model (H. Li & Durbin, 2011) as it has been successfully applied to

reduced-representation datasets (S. Liu & Hansen, 2017). This method allows for analyses of all

species as only one sample per species is required as input. However, given the constraints

created by the sampling density of our target enrichment dataset, we were unable to run PSMC

successfully.

<u>Ecological Niche Modelling</u>

We performed ecological niche modelling (Peterson et al., 2011) to predict the extent of

suitable breeding area for species across the duration of our demographic history reconstruction.

We were able to do so for each species in the stairway plot except *Numenius tenuirostris* due to

the paucity of confirmed breeding records. We obtained species occurrence data from eBird

(2021) and the Global Biodiversity Information Facility (GBIF; using only records with

coordinate uncertainty < 1,000 m) (GBIF.org, 2022a, 2022b, 2022c, 2022d, 2022e, 2022f). For

N. phaeopus, we also included confirmed breeding localities from Lappo et al. (2012) to improve

sample size. Species occurrence data from various sources were combined and further filtered

(Table S3). Occurrence points were filtered by month to retain only records in peak breeding

months of respective species (Birds of the World, 2022). For species with sufficient occurrence

points, occurrence points were also filtered by year to match the time range of the climatic

variables, i.e., 1960–1990. Otherwise, occurrence records from all years were used to maximise

sample size. For species that span the entire Palaearctic (*N. phaeopus* and *N. arquata*), sampling

density was much higher in Europe. To account for the extreme sampling bias, in addition to

generating a kernel density estimate (see next paragraph), occurrence records within Europe for

these two species were randomly down-sampled to match sampling density across the rest of the

Palearctic. Occurrence records outside of the known breeding area of each species were removed

(BirdLife International and Handbook of the Birds of the World, 2017; Lappo et al., 2012).

Finally, to reduce spatial autocorrelation, occurrence records were thinned using a 50 km buffer

(Aiello-Lammens, Boria, Radosavljevic, Vilela, & Anderson, 2015).

To account for sampling bias specific to shorebirds, such as those of this study, we

generated a kernel density estimate using the R package spatialEco 1.3-7 (J. Evans, 2021) based

on the occurrences of species within Scolopacidae. The kernel density estimates was then used to

inform background point selection (i.e., matching sampling bias) (Kramer-Schadt et al., 2013).

For each species, we further limited the sampling of background points to areas outside a 10 km

buffer around occurrence points and within a 500 km buffer around the known breeding area

using the R packages terra 1.5-21 and raster 3.5-15 (Hijmans, 2022b, 2022a). A total of 10,000

background points were then sampled without replacement for each species.

All 19 bioclimatic variables (raster; 2.5 arcmin resolution of ~4.5 km) from WorldClim

1.4 (Hijmans, Cameron, Parra, Jones, & Jarvis, 2005) were obtained for the present day (1960–

1990), mid-Holocene (6,000 years ago), and LGM (22,000 years ago). Bioclimatic variables

were then prepared for input into Maxent 3.4.4 using QGIS 3.4 (QGIS.org, 2022) following De

Alban (2022). Polygon shapefiles were first created for each species, which included the present-

day breeding distribution as well as areas south of that to accommodate for potential shifts in

distribution around the LGM. These polygons were then used to crop the bioclimatic variable

raster for each respective species (Conrad et al., 2015).

We applied Maxent 3.4.4, which makes use of presence-only data and environmental

data to model species' geographical distributions (Phillips et al., 2006). Species-specific Maxent

analyses were performed using the respective breeding occurrence records, background points

and present-day bioclimatic variables of each species. To reduce collinearity among predictors,

we removed predictors with a high variance inflation factor (>3) for each species. To facilitate

parameter tuning, 20 candidate models were built for each species and evaluated using the R

package ENMeval 2.0.3, testing combinations of feature classes (L, LQ, LQH, LQPH) and

regularisation multipliers (0.5, 1, 2, 3, 4) (Kass et al., 2021; Merow, Smith, & Silander, 2013).

To test for model overfitting and transferability, candidate models cross-validated using the

'block' partitioning technique (i.e., occurrences and background points were partitioned into four

spatial blocks, where occurrence numbers among partitions are equal) (Fourcade, Besnard, &

Secondi, 2018; Muscarella et al., 2014). Candidate models with omission rates (minimum

training presence threshold) exceeding 0.2 were rejected. The candidate model with the highest

area under the receiver-operator curve (AUC) was selected as the final model (Table S4) and

used to predict suitable breeding area under present-day, mid-Holocene, and LGM climate

conditions (Table S5).

Predicted species distributions were visualised in R (R Core Team, 2022). We performed

a binary classification of predicted occurrence probability using the maximum sum of sensitivity

plus specificity threshold (C. Liu, White, & Newell, 2013) and calculated suitable breeding area

using the R package raster 3.5-15.

Data availability

DNA reads generated in this study are available on Sequence Read Archive under

BioProject PRJNA742889. Reference genome generated in this study is available on GenBank

under [accession number] (in submission). Pipelines and analysis codes are available upon

request.

Acknowledgements

We thank the following personnel and institutions for their generous contribution of samples (Table S1): Paul Sweet and Thomas Trombone at the American Museum of Natural History (AMNH, New York); Robert Palmer and Leo Joseph at the Australian National Wildlife Collection (ANWC, Canberra); Molly Hagemann at the Bernice Pauahi Bishop Museum (BPBM, Hawaii); David Allan at the Durban Natural Science Museum (DNSM, Durban) and Celine Santillan who assisted in sample transport; Ben Marks at the Field Museum of Natural History (FMNH, Chicago); Foo Maosheng at the Lee Kong Chian Natural History Museum (LKCNHM, Singapore); Carla Marangoni and Gloria Svampa at the Museo Civico di Zoologia (MCZR, Rome); Henry McGhie at the University of Manchester, Manchester Museum (MMUM, Manchester); Robert Prŷs-Jones, Mark Adams, Alex Bond, Ari Benucci and Douglas Russell at the Natural History Museum, London (NHMUK, Tring); Manuel Schweizer at the Naturhistorisches Museum der Bürgergemeinde Bern (NMBE, Bern); Bob McGowan at the Natural Museum of Scotland (NMS, Edinburgh); Joanna Sumner at Museums Victoria (NMV, Melbourne); Angela Ross at the National Museums NI (NMNI, Northern Ireland) and David Allen and Graeme Buchanan who assisted in sample transport; Jan Bolding Kristensen at the Natural History Museum of Denmark (SNM, Copenhagen); José Alves and Camilo Carneiro at the University of Iceland (UOI, Reykjavik); Sharon Birks at the Burke Museum, University of Washington (UWBM, Seattle); Pavel S. Tomkovich, Dmitry Shitikov and Vladimir Sotnikov at the Zoological Museum of Moscow State University (ZMMU, Moscow); and Fyodor Kondrashov and Lisa Chilton who assisted in sample transport. Fletcher Smith assisted in Mozambique (with permission from Lucilia Chuquela, Museu de História Natural and Universidade Eduardo Mondlane, Maputo) with further assistance from Rebecca and Cyril

Kormos, Patricia Zurita and Vinayagan Dhamarajah. HZT acknowledges Elize Ying Xin Ng,

Pratibha Baveja, Yong Chee Keita Sin, Shivaram Rasu, Dominic Yong Jie Ng, Meng Yue Wu,

Liu Xiaoming, and Jose Don De Alban for assistance with laboratory procedures and analyses.

The authors acknowledge support from the National Genomics Infrastructure in Stockholm

funded by Science for Life Laboratory, the Knut and Alice Wallenberg Foundation and the

Swedish Research Council, and SNIC/Uppsala Multidisciplinary Center for Advanced

Computational Science for assistance with massively parallel sequencing and access to the

UPPMAX computational infrastructure. MI acknowledges support from the Swedish Research

Council (2019-03900). KMG acknowledges support from the DBT-Ramalingaswami Fellowship

(BT/HRD/35/02/2006). BC acknowledges funding from the South East Asian Biodiversity

Genomics (SEABIG) Grant (WBS R-154-000-648-646 and WBS R-154-000-648-733) and

startup funding from Trivedi School of Biosciences, Ashoka University. Finally, FER

acknowledges a Singapore Ministry of Education Tier 2 grant (R-154-000-C41-112), which

funded most labwork.

Competing interests

The authors declare no competing interests.

Author Contributions

FER, JJFJJ, GAA, HZT conceptualised the research aims. JJFJJ, GAA, GC, and HZT collected

samples, KMG and BC designed the probe set and MI constructed the reference genome. HZT

performed all laboratory procedures with guidance from CYG, KMG, and BC. HZT performed

all bioinformatic analyses with guidance from CYG, SEHP, and FER. HZT and FER produced

the initial draft of the manuscript which was reviewed by all co-authors.

References

Abadi, S., Azouri, D., Pupko, T., & Mayrose, I. (2019). Model selection may not be a mandatory step for phylogeny reconstruction. *Nature Communications*, *10*(1), 1–11. https://doi.org/10.1038/s41467-019-08822-w

Aiello-Lammens, M. E., Boria, R. A., Radosavljevic, A., Vilela, B., & Anderson, R. P. (2015). spThin: An R package for spatial thinning of species occurrence records for use in ecological niche models. *Ecography*, *38*(5), 541–545. https://doi.org/10.1111/ecog.01132

Allentoft, M. E., & O'Brien, J. (2010). Global amphibian declines, loss of genetic diversity and fitness: A review. *Diversity*, *2*(1), 47–71. https://doi.org/10.3390/d2010047

Allport, G. (2017). Steppe Whimbrels *Numenius phaeopus alboaxillaris* at Maputo, Mozambique, in February–March 2016, with a review of the status of the taxon. *Bulletin of the African Bird Club*, *24*(1), 1–12.

Arenas, M. (2015). Trends in substitution models of molecular evolution. *Frontiers in Genetics*, *6*(OCT), 319. https://doi.org/10.3389/fgene.2015.00319

Bakker, E. S., Gill, J. L., Johnson, C. N., Vera, F. W. M., Sandom, C. J., Asner, G. P., & Svenning, J. C. (2016). Combining paleo-data and modern exclosure experiments to assess the impact of megafauna extinctions on woody vegetation. *Proceedings of the National Academy of Sciences of the United States of America*, *113*(4), 847–855. https://doi.org/10.1073/pnas.1502545112

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., … Pevzner, P. A. (2012). SPAdes: A new genome assembly algorithm and its applications to single-cell sequencing. *Journal of Computational Biology*, *19*(5), 455–477. https://doi.org/10.1089/cmb.2012.0021

Binney, H., Edwards, M., Macias-Fauria, M., Lozhkin, A., Anderson, P., Kaplan, J. O., … Zernitskaya, V. (2017). Vegetation of Eurasia from the last glacial maximum to present: Key biogeographic patterns. *Quaternary Science Reviews*, *157*, 80–97. https://doi.org/10.1016/j.quascirev.2016.11.022

Bird, J. P., Martin, R., Re, H., Gilroy, J., Burfield, I. J., Garnett, S. T., … Butchart, S. H. M. (2020). Generation lengths of the world's birds and their implications for extinction risk. *Conservation Biology*, *34*(5), 1252–1261. https://doi.org/10.1111/cobi.13486

BirdLife International and Handbook of the Birds of the World. (2017). Bird species distribution maps of the world. Retrieved from http://datazone.birdlife.org/species/requestdis

Birds of the World. (2022). *Birds of the World*. (S. M. Billerman, B. K. Keeney, P. G. Rodewald, & T. S. Schulenberg, Eds.). Ithaca, NY, USA: Cornell Laboratory of Ornithology. Retrieved from https://birdsoftheworld.org/bow/home

Buchanan, G. M., Bond, A. L., Pearce-Higgins, J. W., Hilton, G. M., Crockford, N. J., & Kamp, J. (2018). The potential breeding range of Slender-billed Curlew *Numenius tenuirostris* identified from stable-isotope analysis. *Bird Conservation International*, *28*(2), 228–237. https://doi.org/10.1017/S0959270916000551

Bushnell, B. (2014). BBMap. Retrieved from https://sourceforge.net/projects/bbmap/

Butchart, S. H. M., Lowe, S., Martin, R. W., Symes, A., Westrip, J. R. S., & Wheatley, H.
(2018). Which bird species have gone extinct? A novel quantitative classification approach.
Biological Conservation, *227*, 9–18. https://doi.org/10.1016/j.biocon.2018.08.014

Caballero, J. (2018). Scripts. Retrieved November 5, 2021, from
https://github.com/caballero/Scripts/blob/master/rand_var_per_chr.pl

Ceballos, G., Davidson, A., List, R., Pacheco, J., Manzano-Fischer, P., Santos-Barrera, G., &
Cruzado, J. (2010). Rapid decline of a grassland system and its ecological and conservation
implications. *PLoS ONE*, *5*(1), e8562. https://doi.org/10.1371/journal.pone.0008562

Ceballos, G., Ehrlich, P. R., & Raven, P. H. (2020). Vertebrates on the brink as indicators of
biological annihilation and the sixth mass extinction. *Proceedings of the National Academy*
of Sciences of the United States of America, *117*(24), 13596–13602.
https://doi.org/10.1073/pnas.1922686117

Chan, Y. L., Lacey, E. A., Pearson, O. P., & Hadly, E. A. (2005). Ancient DNA reveals
Holocene loss of genetic diversity in a South American rodent. *Biology Letters*, *1*(4), 423–
426. https://doi.org/10.1098/rsbl.2005.0354

Chattopadhyay, B., Garg, K. M., Mendenhall, I. H., & Rheindt, F. E. (2019). Historic DNA
reveals Anthropocene threat to a tropical urban fruit bat. *Current Biology*, *29*, R1269–
R1300. https://doi.org/10.1016/j

Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). Fastp: An ultra-fast all-in-one FASTQ
preprocessor. *Bioinformatics*, *34*(17), i884–i890.
https://doi.org/10.1093/bioinformatics/bty560

Cloutier, A., Sackton, T. B., Grayson, P., Clamp, M., Baker, A. J., & Edwards, S. V. (2019).
Whole-genome analyses resolve the phylogeny of flightless birds (Palaeognathae) in the
presence of an empirical anomaly zone. *Systematic Biology*, *68*(6), 937–955.
https://doi.org/10.1093/sysbio/syz019

Committee on the Status of Endangered Wildlife in Canada. (2009). *COSEWIC Assessment and*
Status Report on the Eskimo Curlew Numenius borealis in Canada. Ottawa. Retrieved from
https://species-registry.canada.ca/index-en.html#/species/21-22#threats

Conrad, O., Bechtel, B., Bock, M., Dietrich, H., Fischer, E., Gerlitz, L., … Böhner, J. (2015).
System for Automated Geoscientific Analyses (SAGA) v. 2.1.4. *Geoscientific Model*
Development, *8*(7), 1991–2007. https://doi.org/10.5194/gmd-8-1991-2015

Crisp, M. D., Trewick, S. A., & Cook, L. G. (2011). Hypothesis testing in biogeography. *Trends*
in Ecology and Evolution, *26*(2), 66–72. https://doi.org/10.1016/j.tree.2010.11.005

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., … Durbin, R.
(2011). The variant call format and VCFtools. *Bioinformatics*, *27*(15), 2156–2158.
https://doi.org/10.1093/bioinformatics/btr330

Darriba, D., Taboada, G. L., Doallo, R., & Posada, D. (2012, August 30). JModelTest 2: More
models, new heuristics and parallel computing. *Nature Methods*. Nature Publishing Group.
https://doi.org/10.1038/nmeth.2109

De Alban, J. D. T. (2022). A short species distribution modeling tutorial. Retrieved February 16, 2022, from https://github.com/dondealban/tutorial-qgis-maxent

Dirzo, R., & Raven, P. H. (2003). Global state of biodiversity and loss. *Annual Review of Environment and Resources*, *28*, 137–167. https://doi.org/10.1146/annurev.energy.28.050302.105532

dos Reis, M., & Yang, Z. (2011). Approximate likelihood calculation on a phylogeny for Bayesian estimation of divergence times. *Molecular Biology and Evolution*, *28*(7), 2161–2172. https://doi.org/10.1093/molbev/msr045

dos Reis, M., & Yang, Z. (2019). Bayesian molecular clock dating using genome-scale datasets. In *Methods in Molecular Biology* (Vol. 1910, pp. 309–330). Humana Press Inc. https://doi.org/10.1007/978-1-4939-9074-0_10

eBird. (2021). eBird Basic Dataset. Version: EBD_relNov-2021. Ithaca, New York: Cornell Lab of Ornithology.

Engelmoer, M., & Roselaar, C. S. (1998a). Eurasian curlew *Numenius arquata*. In M. Engelmoer & C. S. Roselaar (Eds.), *Geographical variation in waders* (pp. 213–223). Dordrecht, Netherlands: Springer Science+Business Media B.V. https://doi.org/10.1007/978-94-011-5016-3

Engelmoer, M., & Roselaar, C. S. (1998b). Whimbrel *Numenius phaeopus*. In M. Engelmoer & C. S. Roselaar (Eds.), *Geographical variation in waders* (pp. 199–212). Dordrecht, Netherlands: Springer Science+Business Media B.V. https://doi.org/10.1007/978-94-011-5016-3

Evans, J. (2021). spatialEco. R package version 1.3-7. Retrieved from https://github.com/jeffreyevans/spatialEco

Evans, S. R., & Sheldon, B. C. (2008). Interspecific patterns of genetic diversity in birds: Correlations with extinction risk. *Conservation Biology*, *22*(4), 1016–1025. https://doi.org/10.1111/j.1523-1739.2008.00972.x

Felsenstein, J. (2009). PHYLIP (PHYLogeny Inference Package). Retrieved from https://evolution.genetics.washington.edu/phylip.html

Foll, M., & Gaggiotti, O. (2008). A genome-scan method to identify selected loci appropriate for both dominant and codominant markers: A Bayesian perspective. *Genetics*, *180*(2), 977–993. https://doi.org/10.1534/genetics.108.092221

Fourcade, Y., Besnard, A. G., & Secondi, J. (2018). Paintings predict the distribution of species, or the challenge of selecting environmental predictors and evaluation statistics. *Global Ecology and Biogeography*, *27*(2), 245–256. https://doi.org/10.1111/geb.12684

Frankham, R. (2005). Genetics and extinction. *Biological Conservation*, *126*(2), 131–140. https://doi.org/10.1016/j.biocon.2005.05.002

Gallo-Orsi, U., & Boere, G. C. (2001). The slender-billed curlew *Numenius tenuirostris*: Threats and conservation. *Acta Ornithologica*, *36*(1), 73–77. https://doi.org/10.3161/068.036.0102

Garner, A., Rachlow, J. L., & Hicks, J. F. (2005). Patterns of genetic diversity and its loss in mammalian populations. *Conservation Biology*, *19*(4), 1215–1221. https://doi.org/10.1111/j.1523-1739.2005.00105.x

GBIF.org. (2022a). GBIF Occurrence Download. Retrieved April 27, 2022, from https://doi.org/10.15468/dl.c8xvcq

GBIF.org. (2022b). GBIF Occurrence Download. Retrieved May 11, 2022, from https://doi.org/10.15468/dl.7fs5mn

GBIF.org. (2022c). GBIF Occurrence Download. Retrieved April 27, 2022, from https://doi.org/10.15468/dl.sybhg8

GBIF.org. (2022d). GBIF Occurrence Download. Retrieved April 27, 2022, from https://doi.org/10.15468/dl.e8jkmw

GBIF.org. (2022e). GBIF Occurrence Download. Retrieved April 27, 2022, from https://doi.org/10.15468/dl.s8g354

GBIF.org. (2022f). GBIF Occurrence Download. Retrieved May 19, 2022, from https://doi.org/10.15468/dl.xuxpvq

Goheen, J. R., Augustine, D. J., Veblen, K. E., Kimuyu, D. M., Palmer, T. M., Porensky, L. M., … Young, T. P. (2018). Conservation lessons from large-mammal manipulations in East African savannas: the KLEE, UHURU, and GLADE experiments. *Annals of the New York Academy of Sciences*, *1429*(1), 31–49. https://doi.org/10.1111/nyas.13848

Goldewijk, K. K., Beusen, A., Doelman, J., & Stehfest, E. (2017). Anthropogenic land use estimates for the Holocene - HYDE 3.2. *Earth System Science Data*, *9*(2), 927–953. https://doi.org/10.5194/essd-9-927-2017

Goldewijk, K. K., Beusen, A., & Janssen, P. (2010). Long-term dynamic modeling of global population and built-up area in a spatially explicit way: HYDE 3.1. *Holocene*, *20*(4), 565–573. https://doi.org/10.1177/0959683609356587

Hedberg, C. P., Lyons, S. K., & Smith, F. A. (2022). The hidden legacy of megafaunal extinction: Loss of functional diversity and resilience over the Late Quaternary at Hall's Cave. *Global Ecology and Biogeography*, *31*(2), 294–307. https://doi.org/10.1111/geb.13428

Helm, A., Oja, T., Saar, L., Takkis, K., Talve, T., & Pärtel, M. (2009). Human influence lowers plant genetic diversity in communities with extinction debt. *Journal of Ecology*, *97*(6), 1329–1336. https://doi.org/10.1111/j.1365-2745.2009.01572.x

Hewitt, G. M. (2000). The genetic legacy of the Quaternary ice ages. *Nature*, *405*, 907–913. https://doi.org/10.1038/35016000

Hijmans, R. J. (2022a). raster: Geographic Data Analysis and Modeling. R package version 3.5-15. Retrieved from https://cran.r-project.org/package=raster

Hijmans, R. J. (2022b). terra: Spatial Data Analysis. R package version 1.5-21. Retrieved from https://cran.r-project.org/package=terra

Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology*, *25*(15), 1965–1978. https://doi.org/10.1002/joc.1276

Hoban, S., Bruford, M., D'Urban Jackson, J., Lopes-Fernandes, M., Heuertz, M., Hohenlohe, P. A., … Laikre, L. (2020). Genetic diversity targets and indicators in the CBD post-2020 Global Biodiversity Framework must be improved. *Biological Conservation*, *248*(July), 108654. https://doi.org/10.1016/j.biocon.2020.108654

Hughes, P. D., Gibbard, P. L., & Ehlers, J. (2013). Timing of glaciation during the last glacial cycle: Evaluating the concept of a global "Last Glacial Maximum" (LGM). *Earth-Science Reviews*, *125*, 171–198. https://doi.org/10.1016/j.earscirev.2013.07.003

IUCN. (2020). The IUCN Red List of Threatened Species. Version 2020-3. Retrieved from https://www.iucnredlist.org

Jetz, W., Thomas, G. H., Joy, J. B., Redding, D. W., Hartmann, K., & Mooers, A. O. (2014). Global distribution and conservation of evolutionary distinctness in birds. *Current Biology*, *24*(9), 919–930. https://doi.org/10.1016/j.cub.2014.03.011

Johnson, C. N. (2009). Ecological consequences of late quaternary extinctions of megafauna. *Proceedings of the Royal Society B: Biological Sciences*, *276*(1667), 2509–2519. https://doi.org/10.1098/rspb.2008.1921

Johnson, C. N., Balmford, A., Brook, B. W., Buettel, J. C., Galetti, M., Guangchun, L., & Wilmshurst, J. M. (2017). Biodiversity losses and conservation responses in the Anthropocene. *Science*, *356*(6335), 270–275. https://doi.org/10.1126/science.aam9317

Johnson, M. G., Gardner, E. M., Liu, Y., Medina, R., Goffinet, B., Shaw, A. J., … Wickett, N. J. (2016). HybPiper: Extracting coding sequence and introns for phylogenetics from high-throughput sequencing reads using target enrichment. *Applications in Plant Sciences*, *4*(7), 1600016. https://doi.org/10.3732/apps.1600016

Jónsson, H., Ginolhac, A., Schubert, M., Johnson, P. L. F., & Orlando, L. (2013). mapDamage2.0: Fast approximate Bayesian estimates of ancient DNA damage parameters. In *Bioinformatics* (Vol. 29, pp. 1682–1684). Oxford Academic. https://doi.org/10.1093/bioinformatics/btt193

Junier, T., & Zdobnov, E. M. (2010). The Newick utilities: high-throughput phylogenetic tree processing in the UNIX shell. *Bioinformatics*, *26*(13), 1669–1670. https://doi.org/10.1093/bioinformatics/btq243

Kass, J. M., Muscarella, R., Galante, P. J., Bohl, C. L., Pinilla-Buitrago, G. E., Boria, R. A., … Anderson, R. P. (2021). ENMeval 2.0: Redesigned for customizable and reproducible modeling of species' niches and distributions. *Methods in Ecology and Evolution*, *12*(9), 1602–1608. https://doi.org/10.1111/2041-210X.13628

Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molecular Biology and Evolution*, *30*(4), 772–780. https://doi.org/10.1093/molbev/mst010

Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., … Drummond, A.

(2012). Geneious Basic: An integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics*, *28*(12), 1647–1649. https://doi.org/10.1093/bioinformatics/bts199

Kirwan, G., Porter, R., & Scott, D. (2015). Chronicle of an extinction? A review of slender-billed curlew records in the Middle East. *British Birds*, *108*(11), 669–682.

Koch, P. L., & Barnosky, A. D. (2006). Late quaternary extinctions: State of the debate. *Annual Review of Ecology, Evolution, and Systematics*, *37*(1), 215–250. https://doi.org/10.1146/annurev.ecolsys.34.011802.132415

Kramer-Schadt, S., Niedballa, J., Pilgrim, J. D., Schröder, B., Lindenborn, J., Reinfelder, V., … Wilting, A. (2013). The importance of correcting for sampling bias in MaxEnt species distribution models. *Diversity and Distributions*, *19*(11), 1366–1379. https://doi.org/10.1111/ddi.12096

Küpper, C., Stocks, M., Risse, J. E., Dos Remedios, N., Farrell, L. L., McRae, S. B., … Burke, T. (2015). A supergene determines highly divergent male reproductive morphs in the ruff. *Nature Genetics*, *48*(1), 79–83. https://doi.org/10.1038/ng.3443

Lappo, E. G., Tomkovich, P. S., & Syroeckovskiy, E. E. (2012). *Atlas of breeding waders in the Russian Arctic*. Moscow: Institute of Geography, Russian Academy of Sciences.

Larsson, A. (2010). abioscript. Retrieved from http://www.ormbunkar.se/phylogeny/abioscripts/

Lavinia, P. D., Kerr, K. C. R., Tubaro, P. L., Hebert, P. D. N., & Lijtmaer, D. A. (2016). Calibrating the molecular clock beyond cytochrome b: Assessing the evolutionary rate of COI in birds. *Journal of Avian Biology*, *47*(1), 84–91. https://doi.org/10.1111/jav.00766

Li, C., Riethoven, J. J. M., & Naylor, G. J. P. (2012). EvolMarkers: A database for mining exon and intron markers for evolution, ecology and conservation studies. *Molecular Ecology Resources*, *12*(5), 967–971. https://doi.org/10.1111/j.1755-0998.2012.03167.x

Li, H. (2011). A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. *Bioinformatics*, *27*(21), 2987–2993. https://doi.org/10.1093/bioinformatics/btr509

Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *ArXiv Preprint*.

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics*, *25*(14), 1754–1760. https://doi.org/10.1093/bioinformatics/btp324

Li, H., & Durbin, R. (2011). Inference of human population history from individual whole-genome sequences. *Nature*, *475*(7357), 493–496. https://doi.org/10.1038/nature10231

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., … Durbin, R. (2009). The sequence alignment/map format and SAMtools. *Bioinformatics*, *25*(16), 2078–2079. https://doi.org/10.1093/bioinformatics/btp352

Liu, C., White, M., & Newell, G. (2013). Selecting thresholds for the prediction of species

occurrence with presence-only data. *Journal of Biogeography*, *40*(4), 778–789. https://doi.org/10.1111/jbi.12058

Liu, L., Yu, L., & Edwards, S. V. (2010). A maximum pseudo-likelihood approach for estimating species trees under the coalescent model. *BMC Evolutionary Biology*, *10*(1), 302. https://doi.org/10.1186/1471-2148-10-302

Liu, S., & Hansen, M. M. (2017). PSMC (pairwise sequentially Markovian coalescent) analysis of RAD (restriction site associated DNA) sequencing data. *Molecular Ecology Resources*, *17*(4), 631–641. https://doi.org/10.1111/1755-0998.12606

Liu, X., & Fu, Y. X. (2015). Exploring population size changes using SNP frequency spectra. *Nature Genetics*, *47*(5), 555–559. https://doi.org/10.1038/ng.3254

Liu, X., & Fu, Y. X. (2020). Stairway Plot 2: demographic history inference with folded SNP frequency spectra. *Genome Biology*, *21*(1), 1–9. https://doi.org/10.1186/s13059-020-02196-9

Lucena-Perez, M., Marmesat, E., Kleinman-Ruiz, D., Martínez-Cruz, B., Węcek, K., Saveljev, A. P., … Godoy, J. A. (2020). Genomic patterns in the widespread Eurasian lynx shaped by Late Quaternary climatic fluctuations and anthropogenic impacts. *Molecular Ecology*, *29*(4), 812–828. https://doi.org/10.1111/mec.15366

Lundgren, E. J., Ramp, D., Rowan, J., Middleton, O., Schowanek, S. D., Sanisidro, O., … Wallach, A. D. (2020). Introduced herbivores restore late pleistocene ecological functions. *Proceedings of the National Academy of Sciences of the United States of America*, *117*(14), 7871–7878. https://doi.org/10.1073/pnas.1915769117

Marques, D. A., Lucek, K., Sousa, V. C., Excoffier, L., & Seehausen, O. (2019). Admixture between old lineages facilitated contemporary ecological speciation in Lake Constance stickleback. *Nature Communications*, *10*(1), 1–14. https://doi.org/10.1038/s41467-019-12182-w

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., … DePristo, M. A. (2010). The genome analysis toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Research*, *20*(9), 1297–1303. https://doi.org/10.1101/gr.107524.110

Merow, C., Smith, M. J., & Silander, J. A. (2013). A practical guide to MaxEnt for modeling species' distributions: What it does, and why inputs and settings matter. *Ecography*, *36*(10), 1058–1069. https://doi.org/10.1111/j.1600-0587.2013.07872.x

Miraldo, A., Li, S., Borregaard, M. K., Flórez-rodríguez, A., Gopalakrishnan, S., Rizvanovic, M., … Nogués-bravo, D. (2016). An Anthropocene map of genetic diversity. *Science*, *353*(6307), 1532–1535. https://doi.org/10.1126/science.aaf4381

Mirarab, S. (2014). Multi-locus-bootstrapping. Retrieved February 25, 2021, from https://github.com/smirarab/multi-locus-bootstrapping

Morozov, V. V. (2000). Current status of the southern subspecies of the whimbrel *Numenius phaeopus alboaxillaris* (Lowe 1921) in Russia and Kazakstan. *Wader Study Group Bulletin*, *92*(August), 30–37.

Muscarella, R., Galante, P. J., Soley-Guardia, M., Boria, R. A., Kass, J. M., Uriarte, M., & Anderson, R. P. (2014). ENMeval: An R package for conducting spatially independent evaluations and estimating optimal model complexity for Maxent ecological niche models . *Methods in Ecology and Evolution*, *5*(11), 1198–1205. https://doi.org/10.1111/2041-210x.12261

Nadachowska-Brzyska, K., Li, C., Smeds, L., Zhang, G., & Ellegren, H. (2015). Temporal dynamics of avian populations during pleistocene revealed by whole-genome sequences. *Current Biology*, *25*(10), 1375–1380. https://doi.org/10.1016/j.cub.2015.03.047

Nakahama, N., Uchida, K., Ushimaru, A., & Isagi, Y. (2018). Historical changes in grassland area determined the demography of semi-natural grassland butterflies in Japan. *Heredity*, *121*(2), 155–168. https://doi.org/10.1038/s41437-018-0057-2

National Center for Biotechnology Information (NCBI). (1988). Retrieved from https://www.ncbi.nlm.nih.gov/

Okonechnikov, K., Conesa, A., & García-Alcalde, F. (2016). Qualimap 2: Advanced multi-sample quality control for high-throughput sequencing data. *Bioinformatics*, *32*(2), 292–294. https://doi.org/10.1093/bioinformatics/btv566

Patton, A. H., Margres, M. J., Stahlke, A. R., Hendricks, S., Lewallen, K., Hamede, R. K., … Storfer, A. (2019). Contemporary demographic reconstruction methods are robust to genome assembly quality: A case study in Tasmanian devils. *Molecular Biology and Evolution*, *36*(12), 2906–2921. https://doi.org/10.1093/molbev/msz191

Pearce-Higgins, J. W., Brown, D. J., Douglas, D. J. T., Alves, J. A., Bellio, M., Bocher, P., … Soloviev, S. A. (2017). A global threats overview for Numeniini populations: synthesising expert knowledge for a group of declining migratory birds. *Bird Conservation International*, *27*, 6–34. https://doi.org/10.1017/S0959270916000678

Peterson, A. T., Soberón, J., Pearson, R. G., Anderson, R. P., Martínez-Meyer, E., Nakamura, M., & Araújo, M. B. (2011). *Ecological Niches and Geographic Distributions*. (S. A. Levin & H. S. Horn, Eds.). New Jersey, USA: Princeton University Press. https://doi.org/10.1515/9781400840670

Phillips, S. J., Anderson, R. P., & Schapire, R. E. (2006). Maximum entropy modeling of species geographic distributions. *Ecological Modelling*, *190*, 231–259. https://doi.org/10.1016/j.ecolmodel.2005.03.026

Pimm, S. L., Jenkins, C. N., Abell, R., Brooks, T. M., Gittleman, J. L., Joppa, L. N., … Sexton, J. O. (2014). The biodiversity of species and their rates of extinction, distribution, and protection. *Science*, *344*(6187). https://doi.org/10.1126/science.1246752

Price, M. N., Dehal, P. S., & Arkin, A. P. (2010). FastTree 2 - Approximately maximum-likelihood trees for large alignments. *PLoS ONE*, *5*(3), e9490. https://doi.org/10.1371/journal.pone.0009490

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender, D., … Sham, P. C. (2007). PLINK: A tool set for whole-genome association and population-based linkage analyses. *American Journal of Human Genetics*, *81*(3), 559–575.

https://doi.org/10.1086/519795

QGIS.org. (2022). QGIS Geographic Information System. QGIS Association. Retrieved from
http://www.qgis.org

Quinlan, A. R., & Hall, I. M. (2010). BEDTools: A flexible suite of utilities for comparing
genomic features. *Bioinformatics*, *26*(6), 841–842.
https://doi.org/10.1093/bioinformatics/btq033

R Core Team. (2022). R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. Retrieved from https://www.r-project.org/

Rambaut, A. (2018). FigTree 1.4.4. Retrieved from https://github.com/rambaut/figtree

Roberts, D. L., Elphick, C. S., & Reed, J. M. (2010). Identifying anomalous reports of putatively
extinct species and why it matters: Contributed paper. *Conservation Biology*, *24*(1), 189–
196. https://doi.org/10.1111/j.1523-1739.2009.01292.x

Roberts, D. L., & Jarić, I. (2016). Inferring extinction in North American and Hawaiian birds in
the presence of sighting uncertainty. *PeerJ*, *2016*(9), e2426.
https://doi.org/10.7717/peerj.2426

Sánchez-Bayo, F., & Wyckhuys, K. A. G. (2019). Worldwide decline of the entomofauna: A
review of its drivers. *Biological Conservation*, *232*(January), 8–27.
https://doi.org/10.1016/j.biocon.2019.01.020

Sharko, F. S., Boulygina, E. S., Rastorguev, S. M., Tsygankova, S. V, Tomkovich, P. S., &
Nedoluzhko, A. V. (2019). Phylogenetic position of the presumably extinct slender-billed
curlew, *Numenius tenuirostris. Mitochondrial DNA Part A: DNA Mapping, Sequencing,*
and Analysis, *30*(4), 626–631. https://doi.org/10.1080/24701394.2019.1597862

Slater, G. S. C., & Birney, E. (2005). Automated generation of heuristics for biological sequence
comparison. *BMC Bioinformatics*, *6*(1), 1–11. https://doi.org/10.1186/1471-2105-6-31

Smit, A., Hubley, R., & Green, P. (2015). RepeatMasker Open-4.0.6. Retrieved from
http://www.repeatmasker.org/

Spielman, D., Brook, B. W., & Frankham, R. (2004). Most species are not driven to extinction
before genetic factors impact them. *Proceedings of the National Academy of Sciences of the*
United States of America, *101*(42), 15261–15264. https://doi.org/10.1073/pnas.0403809101

Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics*, *30*(9), 1312–1313.
https://doi.org/10.1093/bioinformatics/btu033

Stuart, A. J. (2015). Late Quaternary megafaunal extinctions on the continents: a short review.
Geological Journal, *50*, 338–363. https://doi.org/10.1002/gj.2633

Stuart, S. N., Chanson, J. S., Cox, N. A., Young, B. E., Rodrigues, A. S. L., Fischman, D. L., &
Waller, R. W. (2004). Status and trends of amphibian declines and extinctions worldwide.
Science, *306*(5702), 1783–1786. https://doi.org/10.1126/science.1112996

Tan, H. Z., Ng, E. Y. X., Tang, Q., Allport, G. A., Jansen, J. J. F. J., Tomkovich, pavel S., &

Rheindt, F. E. (2019). Population genomics of two congeneric Palaearctic shorebirds reveals differential impacts of Quaternary climate oscillations across habitats types. *Scientific Reports*, *9*(1), 18172. https://doi.org/10.1038/s41598-019-54715-9

Teixeira, J. C., & Huber, C. D. (2021). The inflated significance of neutral genetic diversity in conservation genetics. *Proceedings of the National Academy of Sciences of the United States of America*, *118*(10), 1–10. https://doi.org/10.1073/pnas.2015096118

Thomas, C. D., Cameron, A., Green, R. E., Bakkenes, M., Beaumont, L. J., Collingham, Y. C., … Williams, S. E. (2004). Extinction risk from climate change. *Nature*, *427*, 145–148. https://doi.org/https://doi.org/10.1038/nature02121

Török, P., Ambarlı, D., Kamp, J., Wesche, K., & Dengler, J. (2016). Step(pe) up! Raising the profile of the Palaearctic natural grasslands. *Biodiversity and Conservation*, *25*(12), 2187–2195. https://doi.org/10.1007/s10531-016-1187-6

Turvey, S. T., & Crees, J. J. (2019). Extinction in the Anthropocene. *Current Biology*, *29*(19), R982–R986. https://doi.org/10.1016/j.cub.2019.07.040

Wang, X., Maher, K. H., Zhang, N., Que, P., Zheng, C., Liu, S., … Liu, Y. (2019). Demographic histories and genome-wide patterns of divergence in incipient species of shorebirds. *Frontiers in Genetics*, *10*, 919. https://doi.org/10.3389/fgene.2019.00919

Wang, Y., Pedersen, M. W., Alsos, I. G., De Sanctis, B., Racimo, F., Prohaska, A., … Willerslev, E. (2021). Late Quaternary dynamics of Arctic biota from ancient environmental genomics. *Nature*, *600*(7887), 86–92. https://doi.org/10.1038/s41586-021-04016-x

Wesche, K., Ambarlı, D., Kamp, J., Török, P., Treiber, J., & Dengler, J. (2016). The Palaearctic steppe biome: a new synthesis. *Biodiversity and Conservation*, *25*(12), 2197–2231. https://doi.org/10.1007/s10531-016-1214-7

Wingett, S. W., & Andrews, S. (2018). FastQ Screen: A tool for multi-genome mapping and quality control. *F1000Research*, *7*, 1338. https://doi.org/10.12688/f1000research.15931.2

Xu, H., Luo, X., Qian, J., Pang, X., Song, J., Qian, G., … Chen, S. (2012). FastUniq: A fast de novo duplicates removal tool for paired short reads. *PLoS ONE*, *7*(12), e52249. https://doi.org/10.1371/journal.pone.0052249

Yang, Z. (2007). PAML 4: Phylogenetic analysis by maximum likelihood. *Molecular Biology and Evolution*, *24*(8), 1586–1591. https://doi.org/10.1093/molbev/msm088

Zheng, X., Levine, D., Shen, J., Gogarten, S. M., Laurie, C., & Weir, B. S. (2012). A high-performance computing toolset for relatedness and principal component analysis of SNP data. *Bioinformatics*, *28*(24), 3326–3328. https://doi.org/10.1093/bioinformatics/bts606

Zimov, S. A., Chuprynin, V. I., Oreshko, A. P., Chapin, F. S., Reynolds, J. F., & Chapin, M. C. (1995). Steppe-tundra transition: A herbivore-driven biome shift at the end of the Pleistocene. *American Naturalist*, *146*(4), 765–794. https://doi.org/10.1086/285824