An fMRI meta-analysis of the role of the striatum in everyday-life vs laboratory-developed habits

Pasqualina Guida[1,2,6]*, Mario Michiels[1,2,6]*, Peter Redgrave[5], David Luque[3,4], Ignacio Obeso[1,2]

* Co-first authors

[1] *HM CINAC, Centro Integral de Neurociencias AC. Hospital Universitario HM Puerta del Sur, HM Hospitales, Madrid, Spain*

[2] *CIBERNED, Instituto de Salud Carlos III, Madrid, Spain*

[3] *Departamento de Psicología Básica, Autonoma de Madrid University, Madrid, Spain*

[4] *Departamento de Psicología Básica, Universidad de Málaga, Madrid, Spain*

[5] *Department of Psychology, University of Sheffield, Sheffield, S10 2TN, UK*

[6] *PhD Program in Neuroscience, Autonoma de Madrid University-Cajal Institute, Madrid, Spain 28029*

Correspondence to Dr. Ignacio Obeso

Avda. Carlos V, 70

Móstoles, 28938

Spain

i.obesomartin@gmail.com

www.controlandhabit.com

Abstract

The dorsolateral striatum plays a critical role in the acquisition and expression of

stimulus-response habits that are learned in experimental laboratories. Here, we use meta-

analytic procedures to contrast the neural circuits activated by laboratory-acquired habits

with those activated by stimulus-response behaviours acquired in everyday-life. We

confirmed that newly learned habits rely more on the anterior putamen with activation

extending into caudate and nucleus accumbens. Motor and associative components of

everyday-life habits were identified. We found that motor-dominant stimulus-response

associations developed outside the laboratory primarily engaged posterior dorsal putamen,

supplementary motor area (SMA) and cerebellum. Importantly, associative components

were also represented in the posterior putamen. Thus, common neural representations for

both naturalistic and laboratory-based habits were found in the left posterior and right

anterior putamen. These findings suggest a partial common striatal substrate for habitual

actions that are performed predominantly by stimulus-response associations represented in

the posterior striatum. The overlapping neural substrates for laboratory and everyday-life

habits supports the use of both methods for the analysis of habitual behaviour.

Keywords: habits, everyday-life, probabilistic learning, cortex, striatum, meta-analysis, fMRI

-

Introduction

Repetition gradually enables human behaviour to become automated to the point

where it can be executed without conscious thought and attention. Thus, over a lifetime, a

vast array of sometimes sophisticated movements, thoughts, and motivations are assumed

under automatic stimulus-response control. Typically, we refer to such behaviour as *habits*.

Repeated associations between representations of specific stimuli and particular responses

enable stimulus-evoked reactions to be enacted without conscious intervention (Haith and

Krakauer, 2018). This allows us to perform established stimulus-response associations

automatically, thereby allowing conscious attention to be directed to less predictable aspects

of the world. An obvious example would be not having to think *how* to walk when consciously

thinking about *where* to walk. Hence, the development of habitual routines enable us to

perform predictable and less predictable tasks simultaneously – walking and talking (Sun,

2004).

A defining feature of habitual behaviour is once a stimulus-response association has

been established, it proceeds independently of any change in the outcome value of the

response (Dickinson et al., 1985; Miller et al., 2019). An example would be pressing the

lift/elevator button taking you to the floor of your old office, (the previous stimulus-response

outcome), rather than the new office. Here the lift is the stimulus while the habitual response

would be pressing the old floor button. The move to the new office has devalued the

outcome value of this response yet, when conscious attention is directed elsewhere, it is still

performed. This so-called action slip example satisfies the outcome-devaluation criteria and

would confirm its status as a habit. Automatic habitual control is often contrasted with

conscious goal-directed processing where the action selected is determined by the

value/appropriateness of the predicted outcome (Daw et al., 2005; Dickinson et al., 1985;

Gläscher et al., 2010). Typically, before statistical regularities of a task are determined,

adaptive goal-directed control is used flexibly to achieve the desired outcome at the expense

of speed and efficiency. Consequently, early in instrumental learning, and in uncertain

situations, flexible goal-directed control is specifically deployed for aspects of the task that

cannot easily be predicted.

Human neuroimaging studies have revealed activation of the rostro-medial

(associative) striatum during goal-directed instrumental learning, a pattern that gradually

shifts to caudo-lateral (sensorimotor) regions when habitual control takes over (Brovelli et al.,

2011a; de Wit et al., 2012a; Jankowski et al., 2009; Lehéricy et al., 2005a; Patterson and

Knowlton, 2018; Tanaka et al., 2008; Tricomi et al., 2009). These findings concur with those

described in non-human animals that demonstrate a similar involvement of the rostro-medial

striatal territories early, and caudo-lateral regions later in the acquisition of instrumental

tasks (Balleine and O'Doherty, 2010). Several neuropsychological studies including patients

with striatal abnormalities provide converging evidence for a critical involvement of the

caudal striatum in automatic habitual behaviour. The loss of dopaminergic input from the

midbrain to the striatum in Parkinson's disease is initially concentrated in the posterior

putamen (Pineda-Pardo et al., 2022; Redgrave et al., 2010a). This selective loss seems to

alter automatic moves such as arm swinging during walking or facial expressions in social

interactions. More recently habitual components of typing (Bannard et al., 2019) and

associative learning (Mi et al., 2021) seem to be reduced in Parkinson's disease. In contrast,

Huntington disease induces neural atrophy specifically in the head of the caudate (Vonsattel

and DiFiglia, 1998) preventing the initial associations needed to integrate the repetition of

stimulus-response associations into habits (Holl et al., 2012; Willingham and Koroshetz,

1993). Thus, evidence from experimental animal studies, functional imaging in healthy

humans, and behavioural studies in human patients with selective dysfunction in different

sub-regions of the basal ganglia, all point to the conclusion that conscious goal-directed and

automatic habitual control are exercised by regionally segregated regions in the striatum.

A complication in the study of habits is the likely overlap with a wide range of other

psychological concepts, including skills, automatic, routines, autonomous, inflexible,

unconscious behaviour (West and Brown, 2013). We have already established that the

formal definition of a habit is a stimulus-response association that proceeds automatically

and is not modulated by outcome expectations (Dickinson et al., 1985). This perspective

focuses on the stimulus evoked nature of habitual responses. Other definitions point more to

the acquisition of habitual behaviours (Gardner, 2015), where the role of statistical

regularities within tasks and the necessity of repetition are emphasised. Further, more recent

analyses consider the way ultimate goals and motivation can influence habitual performance

(Wood et al., 2021; Wood and Neal, 2007). However, due to the descriptive terminology

used in different literatures, where the extent of conceptual overlap is unclear,

inconsistencies and discrepancies between the previous studies of habitual behaviour

should be expected (De Houwer, 2019). The purpose of the present meta-analysis was

therefore, to determine the extent to which different forms of automatic stimulus-response

behaviour engage the same or different neural substrates.

Until recently, most studies of habits have been in experimental settings where new

stimulus-response relationships are learned in the laboratory. However, the relationship

between these newly acquired stimulus-response associations and those acquired naturally

over a lifetime when conducting everyday activities, is uncertain. It is equally important to

understand how these important functional systems might malfunction, as reflected in brain

disorders such as Parkinson's disease (Bannard et al., 2019; Redgrave et al., 2010b),

obsessive-compulsive disorder (Gillan, 2021), and drug addictions (Sjoerds et al., 2013),

where dysfunctional patterns of habits are evident. Despite the obvious importance of

habitual control, formal investigations of stimulus-response associations established in

everyday-life are notably absent. An important step to remedy this would be to ask whether

the neural circuits engaged by stimulus-response behaviour acquired in daily life are the

same or different to those activated by newly learned habits in the laboratory. The specific

purpose of the present investigation was therefore to compare the patterns of neural

activation evoked by long-acquired habits brought into the laboratory with those newly

acquired under formal experimental conditions.

Experimental research with animals has shown how instrumental learning occurs

initially through goal-directed computations where behaviour is influenced by predicted

outcome values and then later transitions into outcome-independent stimulus-response

mappings (Balleine and O'Doherty, 2010). Specific tests have been devised to demonstrate

the insensitivity of habitual responding to changes in outcome value or the stimulus-

response relationship. For example, when instrumental performance is seen to persist after

the outcome has been devalued, or the association between stimulus and response

outcome degraded (see Box 1 for a summary), then the observed behaviour is considered

under habitual rather than goal-directed control (Adams, 1982; Balleine and O'Doherty,

2010; Dickinson et al., 1985; Perez and Dickinson, 2020; for a review see Foerde, 2018).

The formal procedures developed in animal studies have been imported into experiments

investigating habits in humans (Luque et al., 2020; Tricomi et al., 2009; Valentin et al., 2007;

Watson et al., 2014; Watson and de Wit, 2018).

To date, most of the literature investigating the neural basis of human habits has

relied on subjects developing new stimulus-response associations under formal laboratory

conditions (Box 1B). This approach has provided important information about the transitions

between goal-directed and habitual neural systems (Balleine and O'Doherty, 2010), and the

factors (repetition, rewards or schedules of reinforcement) that promote the formation of

habitual control (Dickinson et al., 1985; Omar D. Perez and Dickinson, 2020; Smith and

Graybiel, 2016). However, this approach faces several challenges. Most pressing is the

assumed equivalence of stimulus-response habits learned in the laboratory with those

developed naturally in everyday-life. For example, one of the basic tenets of habitual

behaviour is that the potency of stimulus-response associations should become stronger

with repetition. It is therefore, not inspiring that several studies have failed to report a positive

relationship between the amount of training and the strength of habitual responses

measured in their experimental settings (de Wit et al., 2018; Pool et al., 2022.).

Everyday-life	Main habitual outcomes	Task example
READING Real words are read aloud and compared to pseudo-words reading. Depending of the nature of the study, content of words may vary (neutral, emotional, language, non-words, etc) as well as naturality (own language vs new unknown words).	Reading own language with highly familiar/frequent words

(1) |
| **WRITING**
Participants are asked to handwrite by free natural writing, copying sentences/words, air-writing (sensorimotor control), creative writing, naming or drawing different figures. | Writing own language, familiar words (compared to baseline or control conditions) |

(2) (3) |
| **WALKING**
While lying down inside the scanner, participants generate walking programmes on different block conditions, including voluntary alternating strides of the lower limbs, ankle moves, stepping, rapid walking or self-paced vs externally-paced. | Walking naturally at self-paced conditions |

(4) |
| **DRIVING**
MR-compatible driving simulator is often used where participants are asked to naturally drive with different environmental conditions (driving only, stopping at traffic lights, turning curves, avoidance of particular stimuli) and control conditions (driving with sub-task, sub-task only, passive viewing, resting). | Driving in a natural environment (without distractors or dual tasks) |

(5) |

- Cummine et al., 2016; (2) Yang et al., 2018; (3) Karimpoor et al., 2015; (4) Martínez et al., 2016; (5) Choi et al., 2017.

B. Experimental learning	Main habitual outcomes	Task example
PROBABILISTIC CLASSIFICATION TASK (PCT) Participants are shown visual cues that are probabilistically related to outcomes. Participants gradually learn the cue–outcome relationships through trial-by-trial feedback	Optimal responses to stimuli

 (1) |
| **TWO-STAGE REINFORCEMENT LEARNING TASK** Choices at stage 1 determine which state is more probable to occur in Stage 2 that leads to reward. At each Stage 2, participants choose between two stimuli and discover whether they receive a reward or not. | Reward sensitivity without sensitivity to transition probabilities |

 (1) |
| **OUTCOME DEVALUATION TASK** During the discrimination training, participants learn to discriminate cues related to right-left key presses with rewarded outcomes. During the test phase, some of these outcomes are devalued and no longer worth any points. Participants are asked to press the correct key when a stimulus signals the still-valuable outcome or withhold responding if outcome is no longer valuable | Continued responding for devalued outcomes |

 (1) |
| **MOTOR SEQUENCE LEARNING** Motor finger movements are repeated or alternated in high-frequent vs low-frequent sequences. Through repeated practice, participants learn the repeating sequences, as demonstrated by improvements in performance (speed and accuracy). | Accuracy differences between high-frequency and low frequency sequences |

 (2) |

- *Images obtained from Foerde et al., 2018; (2) Cellini, 2017*

***Box 1.** Description of most used paradigms to assess everyday-life **(A)** behaviours and*

*experimental learning **(B)** in fMRI contexts.*

An alternative and increasingly important way forward to study habits in humans

would be to have everyday habitual behaviour learned over a subject's lifetime brought into

the laboratory for investigation. Many behaviours in normal life including aspects of driving,

eating, dancing, reading, talking or walking have significant stimulus-response components

that can be performed automatically without thought while the person's conscious attention

is directed elsewhere (Wechsler et al., 2018; Worringer et al., 2019). Such components have

been acquired through frequent repetition throughout a life-time of everyday trials. Such

associations come to the laboratory fully formed and are independent of any new learning.

The principal challenge in studying everyday-life habits is getting subjects to express

their long-established stimulus-response associations in laboratory settings. This is

necessary so both the automatic behaviour and associated neural activity can be measured

quantitatively. Consequently, investigators have chosen behaviours that have significant

automatic stimulus-response components (Ceceli et al., 2020) and can be performed in the

laboratory while both behaviour and fMRI BOLD signals are measured simultaneously.

Examples of such tasks include reading, where comparisons are made between real words

of different familiarity and emotional content, foreign words and pseudo-words (Cummine et

al., 2016); writing and drawing (Lin et al., 2021; Yang et al., 2018) walking on a special

apparatus (Martínez et al., 2018); and driving an MR-compatible driving simulator (Box 1A;

Choi et al., 2017; Cummine et al., 2016; Huth et al., 2016; Karimpoor et al., 2015; Martínez

et al., 2016, 2018; Oberhuber et al., 2013; Varotto et al., 2020; Yang et al., 2018). The

specific question we addressed in the current study is whether the BOLD activity patterns

reported for stimulus-response associations acquired in everyday life are the same, partially

overlapping or separate from those described for recently established laboratory habits

(Figure 1).

To answer this question, we conducted a quantitative meta-analysis to investigate the

neural substrates of everyday-life and experimental habits. This novel approach allowed a

direct comparison of the respective neural signatures. Special interest was focused on the

system-level circuits involving the basal ganglia (Figure 1). The resulting information

provides important clues into the organisation of life-time habits against which pathologies of

habit and the results of therapeutic interventions can be referenced.

Figure 1. *Hypothesis diagram on striatal role in both everyday-life and experimental habits. Activities*
part of daily life such as writing, reading, walking or driving were selected as everyday-life habits (see
Box 1 for task measurements and details) to expect a critical striatal role in executing these habits
forms. Similar striatal activities can be expected compared to experimental paradigms commonly used
in the cognitive science literature.

Methods

Selection of studies. We conducted 2 independent searches in PubMed and Web

of Science databases to identify fMRI studies investigating stimulus-response habits

acquired: (1) in everyday-life; and (2) in experimental laboratories. The search of articles

related to everyday-life habits was focused on natural behaviours expected to have

significant automatic stimulus-response components: walking, driving, reading and writing.

The selection of these activities was in part based on their potential link to fundamental basal

ganglia functions (Figure 1) and were based on 3 additional criteria: (i) behaviours that are

testable within an MR scanner; (ii) entail a strong component of stimulus-response learning;

and (iii) acquired through lifetime repetition. A summary of typical measures used in these

studies is presented in Box 1. In the search for laboratory based habits we included articles

referenced by Patterson & Knowlton (2018) plus all new relevant articles published since

their last search (June 22, 2017) and June 1, 2022. To enable direct comparison with, and

the extension of the Patterson and Knowlton review (2018), we conducted literature searchs

using the same key search terms and included the task categories – probabilistic learning,

discriminative learning and sequence learning (see Box 1). We also included relevant MeSH

terms available in Pubmed. This allowed us to find articles that do not include targeted

keywords in the title or abstract but have proper MeSH terms linked to their metadata. The

complete list of our search terms can be found in Supplementary Material. To set limits on

inclusion, the articles were filtered further according to the following criteria:

(1)   Neuroimaging modality and analyses: spatial coordinates from human brain

fMRI reported in standardized stereotaxic space (MNI or Talairach space). Thus, other

functional imaging methods (e.g., PET, EEG source imaging, etc) were excluded;

(2)   Sample selection and signal sensitivity: healthy subjects over 18 years old

were included, as were the healthy control participants within clinical studies with

independent analyses. Since brain activity in sensorimotor-related regions of the basal

ganglia can deteriorate with neurological conditions (Redgrave et al., 2010), we included

studies only with healthy adult controls. Moreover, following other fMRI meta-analysis

(Murphy and Garavan, 2004; Sescousse et al., 2013), we did not include studies with

potential low sensitivity (less than 6 subjects);

(3)   Whole-brain analysis: ROI analyses were excluded as they can be biased by

the study hypothesis and may not report all significant regions;

(4)   Computational models: for studies using computational model parameters

(i.e., model-free and/or model-based parameters), we only selected those corresponding to

the reward prediction error (RPE) and the 2-step probabilistic task (a task validated to

assess the duality between habitual/goal-directed modelling (Daw et al., 2011). Model-based

related parameters of any kind (e.g., state prediction error, SPE) were not considered in the

analysis, as they are characteristics of goal-directed behavior;

(5)   Type of studies and language: studies reporting original data and published in

English peer-reviewed journals (other reviews and meta-analysis were excluded).

We then performed a second-step control using reference lists from already

included articles. From the list of references, we first excluded articles whose title had no

direct link to any of our inclusion criteria. From the remaining articles that met our inclusion

criteria we found 17 that had investigated everyday-life habits and 14 articles that had

investigated laboratory-developed habits. Since our second exclusion step involved manual,

rather than online searching, we were able to include articles published before 2017 not

included in Patterson and Knowlton (2018). These additional references were selected

because: (1) we checked all references from the list of accepted papers, not only those in

major review articles; and (2) our criteria allowed us to include studies that did not report

putative habit-related activation of the dorsal striatum (caudate, putamen, or both). Figure S1

depicts the identification, screening, eligibility and selection stages we used to select studies

for inclusion. Foci, scripts and statistical maps can be accessed in the Open Science

Framework (https://osf.io/w5ftm).

Contrasts of interest. The selected contrasts were based on the task condition

reflecting the least difficulty level and less goal-directed task demands as well as

comparisons linked to stimulus-response behaviours (see Table S1 and S2). Of interest, and

most relevant to the present work, was the use of activity maps that were not initially

designed to measure motor or cognitive habits, but rather, general human behaviour. Hence,

one strength of the current work is the study of common hubs for different measures not

initially intended to be directly compared. Ultimately, some contrasts are based upon

validated mechanisms part of habitual behaviour, such as experimental situations (stress or

time pressure) known to largely engage habits and were therefore selected from certain

studies;

256   ***Data analysis.***   The latest version of the GingerALE software v3.0.2 (Eickhoff et al.,

2009) was used to compute activation likelihood estimations (ALE) (Turkeltaub et al., 2002).

From each of the accepted articles, the coordinates of peak activations were manually

extracted and those in Talairach space were transformed to Montreal Neurological Institute

(MNI) space using the inverse transform of icbm2tal from GingerALE (Lancaster et al.,

2007). The ALE method was implemented as follows: (1) for each study a 3D Gaussian

distribution was created around every peak activation (variance proportional to the sample

size of the study). This allowed us to use the higher statistical power to reduce peak

uncertainty in studies with larger sample sizes. This step was done for every peak to

produce one modelled activation (MA) map per study. The value for each voxel in a MA-map

represented the probability of that voxel containing an activation foci. (2) Voxel probabilities

in all MA-maps were merged to produce an ALE map. Following recommendations of

Eickhoff et al., (2016), the ALE map was thresholded using a cluster-level Family-Wise Error

(FWE), with a cluster-forming threshold of .001 and a cluster-level FWE of .05 (as stated in

the GingerALE manual; Fox et al., 2013).

271   The first step in applying a cluster-level FWE was to threshold the ALE map at the

voxel-level (cluster-forming threshold). To do this, all possible combinations between all

peaks from each MA-map were tested. The ALE values provided a null distribution which

assumed the peaks would be randomly distributed following all potential combinations

amongst them. Here, an uncorrected *p*-value was used since in the next step, a FWE

correction was applied to correct for the possibility of multiple comparison errors.

Then, an additional threshold (cluster-level FWE) was applied to select only the

largest clusters. Peaks in every MA-map were distributed randomly and then combined into

a single ALE map, following the same union method described above for the voxel-level

threshold. The ALE maps were also thresholded with the same method used at the voxel-

level threshold. This procedure was repeated (1000 times for this meta-analysis) using

Monte Carlo permutations method, selecting on every run the largest cluster in the

thresholded ALE map (null distribution). Finally, we applied the selected threshold (0.05) to

obtain the FWE-corrected results. Using a 0.05 cluster-level threshold resulted in a

thresholded map where only 5% of the surviving clusters could have been introduced by

chance (false positives), following guidelines to discard non-significant clusters (Eickhoff et

al., 2009).

Contrasts between thresholded ALE maps were computed to compare between our

conditions of interest (e.g. Everyday-life > Experimental maps). All foci in the selected

studies were pooled into a single dataset, which we then split in two datasets by randomly

assigning to each one the same number of foci in their original file. The ALE-scores of these

two random maps were then subtracted in a voxel-wise manner generating the map of their

ALE-scores differences (repeated 10000 times to record ALE-scores and generate a null

distribution of randomly spatially distributed foci). Finally, the actual two ALE maps

corresponding to the contrast were combined voxel-wise by subtracting the ALE-score of

one map from the other. The resulting map of ALE-scores were then thresholded voxel-by-

voxel by comparing them with the null distribution previously obtained. We used the default

p-value suggested in GingerALE (0.01 uncorrected). All computations were run via in-house

python scripts to automate analyses in the GingerALE's interface. Thresholded maps are

reported in MNI152 space (Mazziotta et al., 2001).

Complementary analysis. To ensure the results were not driven by a small number

of studies from specific categories (i.e. heterogeneity), several additional tests were

conducted (as indicated by Müller et al., 2018). We categorized possible differences by

dividing each category (experimental, everyday-life) into subgroups and analysed

independently. Studies of experimental habits were separated into two subcategories:

probabilistic learning vs other experimental tasks (Table S2), on the ground that probabilistic

classification typically activates more anterior portions of the caudate and the putamen,

compared with other tasks (Patterson and Knowlton, 2018). Meanwhile, we included

categorical variables that separated motor (walking, driving) and cognitive (reading, writing)

habits acquired in everyday-life, a further sub-division motivated by the anatomo-functional

gradient along fronto-striatal circuits (Haynes and Haber, 2013). To best isolate the

sensorimotor properties of habitual driving as part of the motor category, we selected

contrasts including less cognitive demands. We also ensured to include > 20 studies in the

experimental and everyday-life categories in order to fulfil the expected effect (Eickhoff et al.,

2016). Finally, we also performed an additional check for heterogeneity and individual study

bias which consists in counting the number of studies that contribute to a specific ROI which

was significant in the ALE analysis (Müller et al., 2018). To do so, we applied an FWHM

smoothing, whose value depended on the number of subjects per study which was

configured according to recommendations in the GingerALE manual. Then, we reviewed the

clusters to check if the foci of each study had any influence for each ROI that resulted

significant in the group analysis.

Results

Study selection

First, to select a cohort of studies investigating everyday-life habits, we analysed data

from 57 studies (a total of 1536 subjects) that used diverse stimulus-response paradigms

(walking, reading, writing and driving). Imaging models that included automatic parameters

on each task were chosen (Tables S1-S2). Second, we sought to confirm the neural basis of

newly learned experimental habits in the 38 studies (a total of 938 subjects) that had used

probabilistic or discriminative learning, 2-step learning or sequential tasks (Box 1B).

Striatal role in everyday-life and experimental habits

To find the neural signatures of everyday-life habits, we obtained cluster activity

associated with stimulus-response behaviours learned throughout life, and compared these

with the signatures of habits newly learnt in laboratory settings. The main effects of

naturalistic habits revealed significant bilateral activity in the posterior putamen (Table 1;

Figure 2A). Specifically, this was sustained by activity in dorsal sections and left putamen

activation that expanded to its posterior boundary (Table 1; Figure 2A-B). For these tasks,

other active regions were seen in the cerebellum and cortical areas including the premotor,

SMA and inferior frontal gyrus pars opercularis (Table 1; Figure 2A-B).

Newly learned habits from laboratory settings were also associated with increased

levels of activation in putaminal sections, but with larger representations in anterior striatum

(Figure 2C). A gradient was seen along the rostro-ventral section of the putamen, right

caudate and the nucleus accumbens bilaterally (Table 2; Figure 2C). The right insula was

one of the extrastriatal hubs of activity associated with habits acquired in the laboratory

(Table 2; Figure 2A).

Of particular interest was the possibility that despite the differences between neural

activation patterns associated with everyday and experimental habits, there may be a

common neural substrate accessed by both categories of stimulus-response behaviour.

Specifically, we found common activations in anterior right putamen and posterior left

putamen (Figure 2A). Although both categories of habits recruited dorsal sub-regions of the

putamen bilaterally, the activation by everyday-life habits was stronger (Figure 2C). As

expected, no activation in the caudate nucleus survived FWE-corrected thresholds with

everyday-life habits. In contrast, habits acquired in the laboratory showed a differential

recruitment of the nucleus accumbens, and to a lesser extent, the most antero-ventral

section of the right caudate nucleus and putamen (Figure 2C). Hence, while the striatum is a

common hub for both categories of habitual behaviour, antero-posterior differences were

reflected in the activation patterns of habits acquired in everyday-life and the experimental

laboratory.

Figure 2. *Everyday-life and experimental thresholded activation maps.* **a)** *Axial views for the main*
regions. Overlap regions at the right column are shown in blue. Note that there exists activation in the
cerebellar cortex for the case of everyday-life studies but it is omitted here for brevity. Z=52 view for
the experimental studies is shown as an unthresholded map for visualization purposes (ALE value ≈
0.01). **b)** *3D striatum reconstruction showing all the activation that fall inside it. Overlap regions at the*
right column are shown in blue. **c)** *3D striatum reconstruction showing the differential activation of the*
Everyday -life > Experimental contrast (in green) and the Experimental > Everyday-life contrast (in
red).

Cluster ID	Volume (mm³)	Brain regions (%)	MNI peak coordinates			ALE value
			x	y	z	
1	9232	Left Putamen (54 %)	-24	-4	6	0.078
		Left Thalamus (28 %)	-14	-22	10	0.04
		Left Pallidum (16 %)	-23	-4	2	0.058
2	6616	Right Putamen (57 %)	24	4	6	0.056
		Right Thalamus (18 %)	16	-16	8	0.035
		Right Insula (12 %)	32	14	8	0.037
		Right Pallidum (11 %)	22	-8	8	0.04
3	4840	Left Supplementary Motor Area (40 %)	-6	0	52	0.035
		Right Paracingulate Gyrus (16 %)	6	14	50	0.031
		Left Paracingulate Gyrus (16 %)	0	8	53	0.031
		Right Supplementary Motor Area (14 %)	4	8	56	0.03
4	3088	Left Precentral Gyrus (97 %)	-54	-2	40	0.038
5	2288	Right-Cerebellum-Cortex (100 %)	8	-64	-20	0.037
6	2288	Left Central Opercular Cortex (54 %)	-46	-2	6	0.037
		Left Inferior Frontal Gyrus pars opercularis (19 %)	-54	8	24	0.027
		Left Precentral Gyrus (17 %)	-56	8	4	0.033
		Left Insula (7 %)	-44	-1	4	0.03
7	2152	Right Precentral Gyrus (76 %)	56	12	32	0.041
		Right Inferior Frontal Gyrus pars opercularis (19 %)	54	13	31	0.038
8	1072	Right Inferior Frontal Gyrus pars opercularis (34 %)	58	12	10	0.027
		Right Central Opercular Cortex (27%)	52	6	2	0.029
		Right Temporal Pole (13 %)	54	10	-2	0.034
		Right Precentral Gyrus (12 %)	3	-10	53	0.022
		Right Planum Polare (9 %)	53	5	-3	0.027
9	1016	Left Precentral Gyrus (43 %)	-26	-8	56	0.04
		Left Superior Frontal Gyrus (35 %)	-4	14	53	0.029
		Left Middle Frontal Gyrus (21 %)	-51	4	38	0.027

373 **Table 1.** *Everyday-life habits peaks coordinates in MNI152 space with region names from Harvard-*
374 *Oxford atlas. Percentages of each brain region indicate how much activation from a cluster fall into*
375 *such region. Coordinates of any activation comprising < 5% of its volume in a region are not shown*
376 *for conciseness.*

Cluster ID	Volume (mm³)	Brain regions (%)	MNI peak coordinates			ALE value
			x	y	z	
1	3000	Right Caudate (48 %)	12	6	10	0.028
		Right Pallidum (23 %)	14	4	0	0.023
		Right Accumbens (12 %)	10	8	-6	0.03
		Right Putamen (8 %)	14	9	-6	0.02
		Right Thalamus (7 %)	12	-1	10	0.018
2	2136	Left Putamen (30 %)	-14	5	-9	0.027
		Left Accumbens (28 %)	-12	6	-10	0.033
		Left Caudate (21 %)	-10	4	4	0.017
3	784	Left Putamen (99 %)	-28	-10	6	0.02
4	752	Right Insula (90 %)	32	22	-4	0.023
		Right Orbito-Frontal Cortex (10 %)	34	22	-7	0.016
5	736	Left Insula (90 %)	-34	16	2	0.025
		Left Frontal-Operculum Cortex (10 %)	-36	19	2	0.02

Table 2. Experimental habits peaks coordinates in MNI152 space with region names from Harvard-
Oxford atlas. Percentages of each brain region indicate how much activation from a cluster fall into
such region. Coordinates of any activation comprising < 5% of its volume in a region are not shown
for conciseness.

Motor and associative segregation in life-long habits

We next tested whether the neural correlates of everyday-life habits varied between

sensorimotor routines (walking or driving) and cognitive-associative ones (reading or

writing). Clustered activation in motor tasks revealed a significant presence along the dorsal

putamen compared with the associative tasks (Figure 3A). In addition, motor tasks extended

the pattern of activation into more posterior sections of the right putamen, but only dorsally

for the left putamen. This cluster was present across all the antero-posterior axes (Figure

3B).

In the case of habits acquired in everyday-life, we observed an activation pattern in

ventral sections of the putamen bilaterally (Table 1). This effect showed significant bilateral

asymmetry favoring the left putamen where activation extended into more posterior regions.

This was absent in the walking and driving tasks (Figure 3A).

Outside the striatum at the cortical level, the associative tasks acquired in everyday-

life recruited premotor area and SMA. Activation was also observed in the cerebellum

(Figure 3A). The same cortical regions were recruited during the walking and driving tasks,

but none survived FWE-corrected thresholds.

To investigate further the failure of our thresholding procedures to detect activation in

the caudate nucleus for any of the naturalistic habit categories, we included the distribution

of all the foci that fell into the striatum in both categories of everyday-life habits (Figure 3C).

This allowed activity in the right caudate to be observed (Figure 3C) in some associative

studies, but not for the motor ones. For the left caudate nucleus, no foci were reported in any

study, which confirms the absence of activity reported in the thresholded maps above.

Figure 3. Everyday-life cognitive-motor subcategories thresholded activation maps. a) Axial views for the main regions. Overlap regions at the right column are shown in blue. b) 3D striatum reconstruction showing all the activation that falls inside it. Overlap regions at the right column are shown in blue. c) Foci distribution of all the studies that falls in the striatum. Note that some foci may correspond to the same study.

Probabilistic learning and other tasks segregation in experimental habits

Finally, we aimed to replicate previous findings from laboratory conditions that linked

the striatum to the learning and execution of new habits. Consistent striatal activity has been

reported when evaluating probabilistic or discriminative learning, 2-step learning and

sequential tasks (Patterson and Knowlton, 2018). Here, the intention was to confirm and

update these results with more recent findings by separating the studies involving trial-and-

error probabilistic reward learning from those using different methodologies for stimulus

learning. As predicted, both probabilistic learning and the other tasks showed common

regions of striatal activation, with largest clusters in the nucleus accumbens and rostro-

ventral sections of the caudate and putamen (Figure 4A). However, only the probabilistic

tasks were associated with bilateral recruitment of the most anterior region of the putamen

(Figure 4B). These results confirm the previous findings reported by Patterson and

Knowlton, (2018). Similarly, the other experimental tasks differed with respect to the

probabilistic ones by demonstrating unilateral recruitment of the left rostral caudate nucleus

(Figure 4B). Outside the striatum, left insular cortex was activated in other tasks but not by

the probabilistic ones (Figure 4A).

Figure 4. *Experimental subcategories thresholded activation maps. **a)** Axial views for*
*the main regions. In this case there are no overlap regions at the right column. **b)** 3D*
striatum reconstruction showing all the activation that fall inside it. In this case there are no
*overlap regions at the right column. **c)** Foci distribution of all the studies that fall in the*
striatum. Note that some foci may correspond to the same study.

Between-study heterogeneity

The degree of heterogeneity was not strong (that is, a result not based on a reduced

number of studies), as was also seen in the subgroup analyses for everyday-life and

experimental results (Tables S3-S4).

Discussion

By comparing long-established habits acquired in everyday-life with newly learned

habits acquired under laboratory-controlled conditions, our meta-analysis identified two

distinct types of habit-related mechanisms in the human striatum. Both common and distinct

functional links between different sub-regions of the brain's habit-associated circuitry were

identified. Differences in neural substrate were found with life-time habits showing enhanced

activity in the dorsal posterior putamen, together with activity in the cerebellum and SMA,

while laboratory habits engaged anterior sections of the putamen with activation extending

into caudate nucleus and nucleus accumbens. Importantly, common regions of activation

were found in posterior left and anterior right putamen. Ultimately, delineation of specific

striatal contributions to motor-associative variables embedded in habits were responsible for

shared anatomical patterns in everyday-life associative habits, such as reading and writing,

that engaged rostral regions of the putamen.

The putamen function in everyday-life habits

The current findings provide direct evidence for bilateral putamen engagement in

habits, long acquired and executed in everyday-life. Typically, we observed larger bilateral

activity in studies of life-time habits compared with newly acquired experimental ones. This

may implicate a broad putaminal role for the stimuli-rich sensorimotor computations required

during complex everyday stimulus-response tasks. Consistent with this view, neuroimaging,

lesioning and animal electrophysiological investigations all converge to pinpoint a role for the

putamen in the expression of associations between specific stimuli and movement patterns

to produce habitual behaviours with predicted outcomes (Bellebaum et al., 2008; Cromwell

and Schultz, 2003; Haruno and Kawato, 2006; Sommer and Pollmann, 2016; Yamada et al.,

2004). Indeed, the neurophysiological properties of the putamen are supported by

subpopulations of neurons that respond to sensory stimuli (Gardiner and Nelson, 1992),

unifying actions sets for movement sequences (Cellini, 2017; Wymbs et al., 2012) and the

integration of elementary units such as individual finger movements (Andersen et al., 2020).

Moreover, putaminal activity is not solely dedicated to movement parameters, but is also

present while learning without motor plans (Romero et al., 2008), increasing neural response

magnitude to reinforced choices (Cohen et al., 2021), and for predicted well-learned and

contextually-driven actions (Kimura, 1986; Kimura et al., 1990, 1984; Kunimatsu et al., 2019;

Peters et al., 2021; Tunik et al., 2009). Hence, the diversity of sensorimotor-related and high-

order reinforcement neurons in the putamen support it having a pivotal role at several levels

in context-rich scenarios (sensorimotor control, predicted actions, object values) in the

acquisition and expression of stimulus-response habits.

The role of the putamen in habitual behaviour is directly influenced by ascending

dopaminergic system acting on cortical and thalamic inputs to this region (Haber et al.,

2000). Afferent dopamine signals provide critical modulatory influences on striatal

subregions whereby projections from substantia nigra pars compacta and the ventral

tegmental area differentially target terminals in dorsal and ventral striatum respectively

(Björklund and Dunnett, 2007). The main nigrostriatal projection is topographically organised

with a medial to lateral gradient (Matsuda et al., 2009). Tonic firing within this system

sustains, motivational, cognitive and action-specific decision making (Hikosaka et al., 2000;

Redgrave et al., 1999), while sensory-evoked phasic patterns of dopaminergic activity

provide a general mechanism for reinforcement learning associated with the acquisition of

novel actions (Redgrave et al., 2008; Schultz, 2006). The large bilateral activity found in the

striatum for the routines of everyday-life should be influenced by ascending dopaminergic

fibers facilitating the selection of appropriate low-level stimulus-response associations

required for writing or walking (proprioceptive and muscle sensorimotor control). Of interest,

the dorso-lateral caudal putamen was active for both cognitive (reading, writing) and motor

operations (driving, walking), which corresponds with our understanding of parallel inter-

related cortico-striatal functions (McFarland and Haber, 2002; Morris and Stein, 2017). The

stimulus-response selection role of the putamen may well represent the low level

sensorimotor selections necessary in all forms of habitual motor behaviour (Bromberg-Martin

et al., 2010; Romero et al., 2008). Hence, the putamen will likely respond preferentially to the

predictable sensorimotor contingences present in seemingly sophisticated behaviours that

rely, in part, on low-level automatic stimulus-response associations for their execution,

driving for example.

Cortical control over everyday-life habits

The substantial input that the striatum receives from cortical areas (Peters et al.,

2021) permits this region to receive contextual, motivational and/or motor cues relevant to

action selection (Haber and Knutson, 2009; Redgrave et al., 1999). Several cortical areas

represent potentially competing behavioural options that can be expressed under habitual

control, including ventral premotor (de Wit et al., 2012b; Ohbayashi et al., 2016),

extraestriate visual cortex (Luque et al., 2017) and bilateral insula or precentral gyrus

(Eryilmaz et al., 2017). In addition, premotor cortex has been shown to interact directly with

the posterior putamen during the acquisition of habits (de Wit et al., 2012b).

In the present study we found that SMA activity was an essential component of the

habitual circuitry engaged by the everyday-life tasks. The SMA represents one of the

important junctions between cortical-subcortical motor and cognitive circuits (Nachev et al.,

2008), with projections to the dorso-lateral striatum and posterior putamen (Leh et al., 2007).

The SMA has been shown to be involved in learning stimulus-response contingencies (Chen

and Wise, 1995; Nachev et al., 2008) and model-free (habitual) tasks (Morris et al., 2016).

Yet, the SMA is not represented in most studies involving new learning of actions. Probably,

the multi-dimensional components required while learning to walk or drive (somatosensory,

visuo-spatial, prediction, motor planning, etc) will be mediated via other cortical inputs (likely

SMA integration function) to the putamen (Haber and Knutson, 2009; Leisman et al., 2016).

When bolted together, serial selections in such complicated but eventually predictable tasks

can be viewed as coherent sequences of habitual behaviour. For example, the sophisticated

set of movements required to change gear when engine revs go above a certain level, and

the precise flow of hand movements required to write individual letters, with repetition, can

be expressed under automatic stimulus-response control. The absence of SMA activity in

experimental tasks included in the present, and previous investigations, attests to the

importance of multi-dimensional computations when cognitive and motor functions are

integrated to develop habitual sequences of behaviour in everyday-life. Such automatic

sequences are likely to recruit medial cortical motor areas including the SMA as a

mechanism to boost expected motor and associative computations that, with subcortical

structures, including putamen (Cunnington et al., 2002; Smittenaar et al., 2013) represent a

looped architecture that can select an appropriate sequences of sensory-driven movements

that implement predictable task components.

Our results also showed a consistent activation of the inferior frontal gyrus. but only

by everyday-life habits. It is known that this region is part of the mirror neuron system related

to action observation and execution (Kilner et al., 2009; Molenberghs et al., 2012). A

possible interpretation of this activation would be that mirror neurons encode behaviour at

the level of actions, rather than as individual movements comprising complex behaviour.

Consequently, only when an action representation is established and well-practiced would a

mirror system be expected to respond accordingly. Perhaps recent, laboratory-acquired

stimulus-response associations are not sufficiently well embedded.

Everyday-life vs experimental scenarios to study habits

Our analyses may also have implications for understanding previous studies that

investigated novel stimulus-response habits learned in experimental laboratories

(probabilistic or discriminative learning, 2-step learning or sequential tasks), and which also

report significant striatal activations (de Wit et al., 2012b; Eryilmaz et al., 2017; Huang et al.,

2020; Jankowski et al., 2009; Lehéricy et al., 2005b; Liljeholm et al., 2015a; Morris and

Stein, 2017; Patterson and Knowlton, 2018; Tricomi et al., 2009). Specifically, our analyses

show stimulus-responses behaviour acquired both in everyday-life and under laboratory

conditions showed common striatal activity in the right anterior putamen and left posterior

putamen. The everyday-life tasks included long established sensorimotor responses

triggered by sensory cues in the absence of new associative learning. In contrast, laboratory

learned habits typically involved relatively minor motor components (finger key presses), but

novel cue-response associations driven by reinforcement learning. In line with a previous

meta-analysis on basal ganglia activation across multiple motor disciplines (Arsalidou et al.,

2013), left lateralized putaminal activity was prominent in motor operations (such as eye

movements and body motion), has a larger volume in right-handed participants (Peterson et

al., 1993), and is critical in behaviours guided by stimulus-response mappings (Liljeholm et

al., 2015a; Patterson and Knowlton, 2018). Hence, our observation of left posterior putamen

activation by both long-established and novel habits suggests this region is critical for

aspects of automatic stimulus-response processing that are present in a wide range of

habitual behaviours.

In contrast, activation of the anterior putamen would accord well with neural patterns

associated with initial learning of stimulus-response associations in experimental settings

(Jankowski et al., 2009). Activity in key regions of the circuitry associated with goal-directed

behavioural control (caudate and nucleus accumbens) were also present in laboratory

studies of habits. These findings match those of several fMRI studies using various

reinforcement learning tasks. When encoding the value of predicted reward outcome linked

to new actions activation of the ventro-medial prefrontal cortex, insula and anterior striatum

has been widely reported (Grabenhorst and Rolls, 2011; Parkes et al., 2015; Tanaka et al.,

2008). An issue here could be that in some of these studies behaviours established in the

laboratory that are presumed to be habits fail to meet formal requirements. For example,

several recent human experimental studies failed to demonstrate an effect of training

duration on insensitivity to outcome-devaluation test (de Wit et al., 2018; Pool et al., n.d.;

Watson and de Wit, 2018). In such studies it is possible that more trials may be needed to

establish stimulus-response associations that are fully automatic and can persist after

devalued outcome challenges. In the case of every-day habits, independence of outcome

value is typically demonstrated in cases of action slips. That is, when a long established

habitual response produces an undesired outcome, such as taking the elevator to the floor of

an old office. Last, there is the possibility that some differences between everyday-life and

laboratory-developed habits, such as activation of the nucleus accumbens, may be due to

the explicit rewards component present in some laboratory paradigms rather than any

fundamental difference in the habitual network.

Dysfunctional habits: inability to exploit stimulus-response habits

Insofar as the caudal putamen has been identified as a critical node in the neural

substrate responsible for automatic habitual behaviour, malfunctioning of this region should

be associated with deficits in habitual performance. A notable instance of this would be the

differential loss of dopamine neurotransmission from this region in Parkinson's disease

(Redgrave et al., 2010a). The causes of this selective loss is not known, however given the

high metabolic cost of dopaminergic neurotransmission Hernandez et al., (2019) suggested

it could reflect an over-reliance throughout life on automatic habitual control. The particular

problems Parkinson's disease patients have with stimulus-response behaviours e.g. walking

and writing (Bannard et al., 2019; Peterson and Horak, 2016; Wu et al., 2016), and their

inability to demonstrate novel associations in the laboratory (de Wit et al., 2010; Wu et al.,

2015), can be seen as an inability of Parkinson's patients to enact established habits and

acquire new ones. In contrast, other neuropsychiatric conditions, such as the addictions,

patients exhibit an excessive cue-dependent use of certain rewards linked to increased

posterior putamen activity (Everitt et al., 2008; Sjoerds et al., 2013). In the light of this work,

an important future question would be whether the limbic and associative territories of the

basal ganglia (Alexander and Crutcher, 1990) can operate in a similar habitual mode as has

been established for stimulus-response motor habits in the sensorimotor territories. If so,

this would move towards a reconceptualization of stimulus evoked motivations (e.g. drug

craving), stimulus-evoked cognitions (e.g. prejudices), and stimulus-evoked motor responses

(motor habits) as part of a general mode of model-free stimulus-response behavioural

control. Were this to be the case, one might speculate there would likely be a general

mechanism of neural implementation across all territories of the basal ganglia.

Limitations & strengths

Despite the clear positive findings of the present investigation, certain limitations

must be acknowledged. First, the ALE analysis can cause some parameters, such as voxel

peaks, to be overlooked. Ideally, exploring the statistical activity maps of each study

individually would be of great value. Unfortunately, most studies did not have full imaging

datasets so this would be precluded.

Second, some studies using RPE contrasts combined mixed regressors of choice

and outcome thereby rendering it difficult to make separations between habitual stimulus-

response components. To avoid this, we explored our principal findings without these

studies and found similar patterns of striatal activation (anterior putamen; Figure 4A).

Moreover, our heterogeneity analysis showed a reduced number of RPE studies (6/25

studies for left putamen; 4/20 studies for right putamen) that contributed to the significant

activity of such clusters (as displayed in Table S4 for experimental studies). Consequently,

the putamen clusters we report are not mainly driven by the RPE studies

Third, as stimulus-response habits are driven by specific conjunctions of sensory

stimuli, the heterogeneity of every-day and laboratory tasks included in our analysis means

the stimuli driving brain responses necessarily will have differed. Moreover, the laboratory

studies included have minor motor components compared with the typical motor-

sophistication of habits established over a lifetime. To control for this possibility, and

following imaging meta-analysis best practice (Müller et al., 2018), as different stimuli and

task-related activities were collected, general and specific sub-analysis were conducted.

Last, due to the difficulty of performing some behaviours in the MRI scanner, necessarily we

had to restrict our selection to everyday-life behaviours where scanning was possible. This

could restrict the generalisation of our findings to other activities prevalent in every-day life.

The main strengths of the work point to the possible generalizability and external

validity of lab-developed habits, as well as proposing novel methodological procedures to

investigate human habitual behaviour. The heterogeneity of tasks necessarily means there

was a diverse range of both stimuli and behavioural responses deployed in the studies

included in our analyses. It is therefore remarkable we were able to pull out common

patterns of neural activation associated with stimulus-response control.

A further strength of our investigation was that we were able to find significant striatal

activation in tasks with high levels of stimulus-response behaviour in studies that were not

originally designed to study habits. This was despite the behaviours of interest not having

been formally identified as habitual using outcome devaluation or other tests. That we find

common neural maps with widely different means for testing habits lends support to this

idea.

Future research and conclusions

Recently, a range of procedures have been developed to establish and test the

expression of newly acquired habits in the experimental laboratory, including reversal of

learned actions after overtraining (Luque et al., 2020), overloading goal-directed top-down

control while measuring execution of learned stimulus-response associations (Hardwick et

al., 2019) or biasing movement kinematics (Wong et al., 2017). Interestingly, pre-existing

categorical associations established in everyday-life (i.e. color associations or prejudice)

have a clear advantage on measuring automatic processing (Amodio, 2014; Ceceli et al.,

2020). Further methodological options for studying long established habits acquired in

everyday-life include assessing the motor performance of expert musicians (Vaquero et al.,

2016), tennis players (Cacioppo et al., 2014) and expert shooters (Kim et al., 2014). Hence,

getting subjects to bring their everyday-life stimulus-response associations into the

laboratory under controlled conditions is an important option for studying the neural

substrates underlying habitual behaviour in humans. Although there is less control of the

independent variables in such studies, by selecting subjects with different amounts of

everyday-life experience it is possible to relate the amount of practice with habit strength.

In conclusion, the present study points to a fundamental functional role for the

posterior putamen in the expression of habits acquired in everyday-life. Importantly, a critical

dissociation was established between different brain regions whose contributions to motor-

cognitive representations of habits have, thus far, been largely indistinguishable. Conversely,

the engagement of the anterior putamen was shown to be associated more with habits newly

learned in experimental laboratories. Careful experimental protocols must be designed to

identify those chunks of complex behaviour that are under automatic stimulus-response

control and can occur independently of outcome value or response contingency. This is true

both for newly acquired associations in the laboratory and long-standing habits established

in everyday-life. Finally, the present study highlights the importance and value of having

subjects bring life-long habits into the laboratory to be investigated and compared with

recently acquired stimulus-response associations. This meta-analysis provides strong

evidence for a striatal role in everyday-life and lab-developed habits giving a novel

contribution to the literature on the general neural footprint of human habits.

References

Adams, C.D., 1982. Variations in the Sensitivity of Instrumental Responding to Reinforcer

Devaluation. Q. J. Exp. Psychol. Sect. B 34, 77–98.

https://doi.org/10.1080/14640748208400878

Alexander, G.E., Crutcher, M.D., 1990. Functional architecture of basal ganglia circuits:

neural substrates of parallel processing. Trends Neurosci. https://doi.org/10.1016/0166-

2236(90)90107-L

Allali, G., Montembeault, M., Brambati, S.M., Bherer, L., Blumen, H.M., Launay, C.P., Liu-

Ambrose, T., Helbostad, J.L., Verghese, J., Beauchet, O., 2019. Brain structure

covariance associated with gait control in aging. Journals Gerontol. - Ser. A Biol. Sci.

Med. Sci. 74, 705–713. https://doi.org/10.1093/gerona/gly123

Amodio, D.M., 2014. The neuroscience of prejudice and stereotyping. Nat. Rev. Neurosci.

15, 670–682. https://doi.org/10.1038/nrn3800

Andersen, K.W., Madsen, K.H., Siebner, H.R., 2020. Discrete finger sequences are widely

represented in human striatum. Sci. Rep. 10, 1–12. https://doi.org/10.1038/s41598-020-

69923-x

Anggraini, D., Glasauer, S., Wunderlich, K., 2018. Neural signatures of reinforcement

learning correlate with strategy adoption during spatial navigation. Sci. Reports 2018 81

8, 1–14. https://doi.org/10.1038/s41598-018-28241-z

Aron, A.R., Gluck, M.A., Poldrack, R.A., 2006. Long-term test-retest reliability of functional

MRI in a classification learning task. Neuroimage 29, 1000–1006.

https://doi.org/10.1016/J.NEUROIMAGE.2005.08.010

Aron, A.R., Shohamy, D., Clark, J., Myers, C., Gluck, M.A., Poldrack, R.A., 2004. Human

midbrain sensitivity to cognitive feedback and uncertainty during classification learning.

J. Neurophysiol. 92, 1144–1152. https://doi.org/10.1152/JN.01209.2003

Arsalidou, M., Duerden, E.G., Taylor, M.J., 2013. The Centre of the Brain : Topographical

Model of Motor , Cognitive , Affective , and Somatosensory Functions of the Basal

Ganglia 3054, 3031–3054. https://doi.org/10.1002/hbm.22124

Balleine, B.W., O'Doherty, J.P., 2010. Human and rodent homologies in action control:

Corticostriatal determinants of goal-directed and habitual action.

Neuropsychopharmacology 35, 48–69. https://doi.org/10.1038/npp.2009.131

Bannard, C., Leriche, M., Bandmann, O., Brown, C.H., Ferracane, E., Sánchez-Ferro, Á.,

Obeso, J., Redgrave, P., Stafford, T., 2019. Reduced habit-driven errors in Parkinson's

Disease. Sci. Rep. 9, 1–8. https://doi.org/10.1038/s41598-019-39294-z

Beeson, P.M., Rapcsak, S.Z., Plante, E., Chargualaf, J., Chung, A., Johnson, S.C., Trouard,

T.P., 2003. The neural substrates of writing: A functional magnetic resonance imaging

study. Aphasiology 17, 647–665. https://doi.org/10.1080/02687030344000067

Beierholm, U.R., Anen, C., Quartz, S., Bossaerts, P., 2011. Separate encoding of model-

based and model-free valuations in the human brain. Neuroimage 58, 955–962.

https://doi.org/10.1016/j.neuroimage.2011.06.071

Bellebaum, C., Koch, B., Schwarz, M., Daum, I., 2008. Focal basal ganglia lesions are

associated with impairments in reward-based reversal learning. Brain 131, 829–841.

https://doi.org/10.1093/brain/awn011

Binder, J.R., Medler, D.A., Westbury, C.F., Liebenthal, E., Buchanan, L., 2006. Tuning of the

human left fusiform gyrus to sublexical orthographic structure. Neuroimage 33, 739–

748. https://doi.org/10.1016/j.neuroimage.2006.06.053

Bisio, A., Pedullà, L., Bonzano, L., Tacchino, A., Brichetto, G., Bove, M., 2017. The

kinematics of handwriting movements as expression of cognitive and sensorimotor

impairments in people with multiple sclerosis. Sci. Rep. 7, 1–10.

https://doi.org/10.1038/s41598-017-18066-7

Björklund, A., Dunnett, S.B., 2007. Dopamine neuron systems in the brain: an update.

Trends Neurosci. https://doi.org/10.1016/j.tins.2007.03.006

Bookheimer, S.Y., Zeffiro, T.A., Blaxton, T., Gaillard, W., Theodore, W., 1995. Regional

cerebral blood flow during object naming and word reading. Hum. Brain Mapp. 3, 93–

106. https://doi.org/10.1002/hbm.460030206

Bromberg-Martin, E.S., Matsumoto, M., Hikosaka, O., 2010. Dopamine in Motivational

Control: Rewarding, Aversive, and Alerting. Neuron.

https://doi.org/10.1016/j.neuron.2010.11.022

Brovelli, A., Nazarian, B., Meunier, M., Boussaoud, D., 2011a. Differential roles of caudate

nucleus and putamen during instrumental learning. Neuroimage 57, 1580–90.

https://doi.org/10.1016/j.neuroimage.2011.05.059

Brovelli, A., Nazarian, B., Meunier, M., Boussaoud, D., 2011b. Differential roles of caudate

nucleus and putamen during instrumental learning. Neuroimage 57, 1580–1590.

https://doi.org/10.1016/j.neuroimage.2011.05.059

Buchsbaum, B.R., Olsen, R.K., Koch, P.F., Kohn, P., Kippenhan, J.S., Berman, K.F., 2005.

Reading, hearing, and the planum temporale. Neuroimage 24, 444–454.

https://doi.org/10.1016/j.neuroimage.2004.08.025

Cacioppo, S., Fontang, F., Patel, N., Decety, J., Monteleone, G., Cacioppo, J.T., 2014.

Intention understanding over T: A neuroimaging study on shared representations and

tennis return predictions. Front. Hum. Neurosci. 8, 781.

https://doi.org/10.3389/fnhum.2014.00781

Callan, A.M., Osu, R., Yamagishi, Y., Callan, D.E., Inoue, N., 2009. Neural correlates of

resolving uncertainty in driver's decision making. Hum. Brain Mapp. 30, 2804–2812.

https://doi.org/10.1002/hbm.20710

Carreiras, M., Mechelli, A., Estévez, A., Price, C.J., 2007. Brain activation for lexical decision

and reading aloud: Two sides of the same coin? J. Cogn. Neurosci. 19, 433–444.

https://doi.org/10.1162/jocn.2007.19.3.433

Ceceli, A.O., Myers, C.E., Tricomi, E., 2020. Demonstrating and disrupting well-learned

habits. PLoS One 15, 1–28. https://doi.org/10.1371/journal.pone.0234424

Cellini, N., 2017. Memory consolidation in sleep disorders. Sleep Med. Rev. 35, 101–112.

https://doi.org/10.1016/j.smrv.2016.09.003

Celone, K.A., Thompson-Brenner, H., Ross, R.S., Pratt, E.M., Stern, C.E., 2011. An fMRI

investigation of the fronto-striatal learning system in women who exhibit eating disorder

behaviors. Neuroimage 56, 1749–1757.

https://doi.org/10.1016/J.NEUROIMAGE.2011.03.026

Cheema, K., Lantz, N., Cummine, J., 2018. Exploring the role of subcortical structures in

developmental reading impairments: Evidence for subgroups differentiated by caudate

activity. Neuroreport 29, 271–279. https://doi.org/10.1097/WNR.0000000000000938

Chein, J.M., Albert, D., O'Brien, L., Uckert, K., Steinberg, L., 2011. Peers increase

adolescent risk taking by enhancing activity in the brain's reward circuitry. Dev. Sci. 14.

https://doi.org/10.1111/j.1467-7687.2010.01035.x

Chen, L.L., Wise, S.P., 1995. Neuronal activity in the supplementary eye field during

acquisition of conditional oculomotor associations. J. Neurophysiol. 73, 1101–1121.

https://doi.org/10.1152/jn.1995.73.3.1101

Choi, M.H., Kim, H.S., Yoon, H.J., Lee, J.C., Baek, J.H., Choi, J.S., Tack, G.R., Min, B.C.,

Lim, D.W., Chung, S.C., 2017. Increase in brain activation due to subtasks during

driving: FMRI study using new MR-compatible driving simulator. J. Physiol. Anthropol.

36, 1–12. https://doi.org/10.1186/s40101-017-0128-8

Christensen, M.S., Lundbye-Jensen, J., Petersen, N., Geertsen, S.S., Paulson, O.B.,

Nielsen, J.B., 2007. Watching your foot move - An fMRI study of visuomotor

interactions during foot movement. Cereb. Cortex 17, 1906–1917.

https://doi.org/10.1093/cercor/bhl101

Chung, S.C., Choi, M.H., Kim, H.S., You, N.R., Hong, S.P., Lee, J.C., Park, S.J., Baek, J.H.,

Jeong, U.H., You, J.H., Lim, D.W., Kim, H.J., 2014. Effects of distraction task on

driving: A functional magnetic resonance imaging study, in: Bio-Medical Materials and

Engineering. IOS Press, pp. 2971–2977. https://doi.org/10.3233/BME-141117

Church, J.A., Coalson, R.S., Lugar, H.M., Petersen, S.E., Schlaggar, B.L., 2008. A

developmental fMRI study of reading and repetition reveals changes in phonological

and visual mechanisms over age. Cereb. Cortex 18, 2054–2065.

https://doi.org/10.1093/cercor/bhm228

Ciccarelli, O., Toosy, A.T., Marsden, J.F., Wheeler-Kingshott, C.M., Sahyoun, C., Matthews,

P.M., Miller, D.H., Thompson, A.J., 2005. Identifying brain regions for integrative

sensorimotor processing with ankle movements. Exp. Brain Res. 166, 31–42.

https://doi.org/10.1007/s00221-005-2335-5

Cohen, Y., Schneidman, E., Paz, R., 2021. The geometry of neuronal representations during

rule learning reveals complementary roles of cingulate cortex and putamen. Neuron

109, 839-851.e9. https://doi.org/10.1016/j.neuron.2020.12.027

Cromwell, H.C., Schultz, W., 2003. Effects of expectations for different reward magnitudes

on neuronal activity in primate striatum. J. Neurophysiol. 89, 2823–2838.

https://doi.org/10.1152/jn.01014.2002

Cummine, J., Cribben, I., Luu, C., Kim, E., Bakhtiari, R., Georgiou, G., Boliek, C.A., 2016.

Understanding the Role of Speech Production in Reading: Evidence for a Print-to-

Speech Neural Network Using Graphical Analysis. Neuropsychology.

https://doi.org/10.1037/neu0000236

Cunnington, R., Windischberger, C., Deecke, L., Moser, E., 2002. The preparation and

execution of self-initiated and externally-triggered movement: A study of event-related

fMRI. Neuroimage 15, 373–385. https://doi.org/10.1006/nimg.2001.0976

Daw, N.D., Gershman, S.J., Seymour, B., Dayan, P., Dolan, R.J., 2011. Model-based

influences on humans' choices and striatal prediction errors. Neuron 69, 1204–1215.

https://doi.org/10.1016/j.neuron.2011.02.027

Daw, N.D., Niv, Y., Dayan, P., 2005. Uncertainty-based competition between prefrontal and

dorsolateral striatal systems for behavioral control. Nat. Neurosci. 8, 1704–1711.

https://doi.org/10.1038/nn1560

De Houwer, J., 2019. On How Definitions of Habits Can Complicate Habit Research. Front.

Psychol. 10, 1–9. https://doi.org/10.3389/fpsyg.2019.02642

de Wit, S., Barker, R. a, Dickinson, A.D., Cools, R., 2010. Habitual versus goal-directed

action control in Parkinson disease. J Cogn Neurosci 23, 1218–1229.

https://doi.org/10.1162/jocn.2010.21514 [doi]

de Wit, S., Kindt, M., Knot, S.L., Verhoeven, A.A.C., Robbins, T.W., Gasull-Camos, J.,

Evans, M., Mirza, H., Gillan, C.M., 2018. Shifting the balance between goals and habits:

Five failures in experimental habit induction. J. Exp. Psychol. Gen. 147, 1043–1065.

https://doi.org/10.1037/xge0000402

de Wit, S., Watson, P., Harsay, H.A., Cohen, M.X., van de Vijver, I., Ridderinkhof, K.R.,

2012a. Corticostriatal connectivity underlies individual differences in the balance

between habitual and goal-directed action control. J. Neurosci. 32, 12066–12075.

https://doi.org/10.1523/JNEUROSCI.1088-12.2012

de Wit, S., Watson, P., Harsay, H.A., Cohen, M.X., van de Vijver, I., Ridderinkhof, K.R.,

2012b. Corticostriatal connectivity underlies individual differences in the balance

between habitual and goal-directed action control. J. Neurosci. 32, 12066–12075.

https://doi.org/10.1523/JNEUROSCI.1088-12.2012

Delgado, M.R., Miller, M.M., Inati, S., Phelps, E.A., 2005. An fMRI study of reward-related

probability learning. Neuroimage 24, 862–873.

https://doi.org/10.1016/j.neuroimage.2004.10.002

Deserno, L., Huys, Q.J.M., Boehme, R., Buchert, R., Heinze, H.J., Grace, A.A., Dolan, R.J.,

Heinz, A., Schlagenhauf, F., 2015. Ventral striatal dopamine reflects behavioral and

neural signatures of model-based control during sequential decision making. Proc. Natl.

Acad. Sci. U. S. A. 112, 1595–1600. https://doi.org/10.1073/pnas.1417219112

Dickinson, A., Trans, P., Lond, R.S., 1985. Actions and habits: the development of

behavioural autonomy. Philos. Trans. R. Soc. London. B, Biol. Sci. 308, 67–78.

https://doi.org/10.1098/rstb.1985.0010

Doll, B.B., Duncan, K.D., Simon, D.A., Shohamy, D., Daw, N.D., 2015. Model-based choices

involve prospective neural activity. Nat. Neurosci. 18, 767–772.

https://doi.org/10.1038/NN.3981

Eickhoff, S.B., Laird, A.R., Grefkes, C., Wang, L.E., Zilles, K., Fox, P.T., 2009. Coordinate-

based activation likelihood estimation meta-analysis of neuroimaging data: A random-

effects approach based on empirical estimates of spatial uncertainty. Hum. Brain Mapp.

30, 2907–2926. https://doi.org/10.1002/hbm.20718

Eickhoff, S.B., Nichols, T.E., Laird, A.R., Hoffstaedter, F., Amunts, K., Fox, P.T., Bzdok, D.,

Eickhoff, C.R., 2016a. Behavior, sensitivity, and power of activation likelihood

estimation characterized by massive empirical simulation. Neuroimage 137, 70–85.

https://doi.org/10.1016/j.neuroimage.2016.04.072

Eickhoff, S.B., Nichols, T.E., Laird, A.R., Hoffstaedter, F., Amunts, K., Fox, P.T., Bzdok, D.,

Eickhoff, C.R., 2016b. Behavior, sensitivity, and power of activation likelihood

estimation characterized by massive empirical simulation. Neuroimage 137, 70–85.

https://doi.org/10.1016/J.NEUROIMAGE.2016.04.072

Erdeniz, B., Done, J., 2019. Common and distinct functional brain networks for intuitive and

deliberate decision making. Brain Sci. 9. https://doi.org/10.3390/brainsci9070174

Erhard, K., Kessler, F., Neumann, N., Ortheil, H.J., Lotze, M., 2014. Professional training in

creative writing is associated with enhanced fronto-striatal activity in a literary text

continuation task. Neuroimage 100, 15–23.

https://doi.org/10.1016/j.neuroimage.2014.05.076

Eryilmaz, H., Rodriguez-Thompson, A., Tanner, A.S., Giegold, M., Huntington, F.C.,

Roffman, J.L., 2017. Neural determinants of human goal-directed vs. habitual action

control and their relation to trait motivation. Sci. Rep. 7, 1–11.

https://doi.org/10.1038/s41598-017-06284-y

Etchamendy, N., Konishi, K., Pike, G.B., Marighetto, A., Bohbot, V.D., 2012. Evidence for a

virtual human analog of a rodent relational memory task: A study of aging and fMRI in

young adults. Hippocampus 22, 869–880. https://doi.org/10.1002/HIPO.20948

Everitt, B.J., Belin, D., Economidou, D., Pelloux, Y., Dalley, J.W., Robbins, T.W., 2008.

Neural mechanisms underlying the vulnerability to develop compulsive drug-seeking

habits and addiction. Philos. Trans. R. Soc. B Biol. Sci. 363, 3125–3135.

https://doi.org/10.1098/rstb.2008.0089

Fera, F., Weickert, T.W., Goldberg, T.E., Tessitore, A., Hariri, A., Das, S., Lee, S., Zoltick,

B., Meeter, M., Myers, C.E., Gluck, M.A., Weinberger, D.R., Mattay, V.S., 2005. Neural

mechanisms underlying probabilistic category learning in normal aging. J. Neurosci. 25,

11340–11348. https://doi.org/10.1523/JNEUROSCI.2736-05.2005

Fermin, A.S.R., Yoshida, T., Yoshimoto, J., Ito, M., Tanaka, S.C., Doya, K., 2016. Model-
869   based action planning involves cortico-cerebellar and basal ganglia networks. Sci. Rep.
870   6. https://doi.org/10.1038/SREP31378

871 Fernández-Seara, M.A., Aznárez-Sanado, M., Mengual, E., Loayza, F.R., Pastor, M.A.,
872   2009. Continuous performance of a novel motor sequence leads to highly correlated
873   striatal and hippocampal perfusion increases. Neuroimage 47, 1797–1808.
874   https://doi.org/10.1016/J.NEUROIMAGE.2009.05.061

875 Foerde, K., 2018. What are habits and do they depend on the striatum? A view from the
876   study of neuropsychological populations. Curr. Opin. Behav. Sci. 20, 17–24.
877   https://doi.org/10.1016/j.cobeha.2017.08.011

878 Foerde, K., Knowlton, B.J., Poldrack, R.A., 2006. Modulation of competing memory systems
879   by distraction. Proc. Natl. Acad. Sci. U. S. A. 103, 11778–11783.
880   https://doi.org/10.1073/PNAS.0602659103

881 Forstmann, B.U., Dutilh, G., Brown, S., Neumann, J., Von Cramon, D.Y., Ridderinkhof, K.R.,
882   Wagenmakers, E.J., 2008. Striatum and pre-SMA facilitate decision-making under time
883   pressure. Proc. Natl. Acad. Sci. U. S. A. 105, 17538–17542.
884   https://doi.org/10.1073/pnas.0805903105

885 Fox, P.T., Laird, A.R., Eickhoff, S.B., Lancaster, J.L., Fox, M., Uecker, A.M., Robertson, M.,
886   Ray, K.L., 2013. User manual for GingerALE 2.3, Retrieved from
887   http://brainmap.org/ale/manual.pdf.

888 Gardiner, T.W., Nelson, R.J., 1992. Striatal neuronal activity during the initiation and
889   execution of hand movements made in response to visual and vibratory cues. Exp.
890   Brain Res. 92, 15–26. https://doi.org/10.1007/BF00230379

891 Gardner, B., 2015. A review and analysis of the use of "habit" in understanding, predicting
892   and influencing health-related behaviour. Health Psychol. Rev. 9, 277–295.

https://doi.org/10.1080/17437199.2013.876238

Gillan, C.M., 2021. Recent Developments in the Habit Hypothesis of OCD and Compulsive

Disorders. Springer, Berlin, Heidelberg, pp. 1–21.

https://doi.org/10.1007/7854_2020_199

Gläscher, J., Daw, N., Dayan, P., O'Doherty, J.P., 2010. States versus rewards: Dissociable

neural prediction error signals underlying model-based and model-free reinforcement

learning. Neuron 66, 585–595. https://doi.org/10.1016/j.neuron.2010.04.016

Gottfried, J.A., O'Doherty, J., Dolan, R.J., 2003. Encoding predictive reward value in human

amygdala and orbitofrontal cortex. Science (80-.). 301, 1104–1107.

https://doi.org/10.1126/SCIENCE.1087919

Gould, L., Mickleborough, M.J.S., Lorentz, E., Ekstrand, C., Borowsky, R., 2018. A

behavioral and fMRI examination of the effect of rhythm on reading noun-verb

homographs aloud. https://doi.org/10.1080/23273798.2018.1442012 33, 829–849.

https://doi.org/10.1080/23273798.2018.1442012

Grabenhorst, F., Rolls, E.T., 2011. Value, pleasure and choice in the ventral prefrontal

cortex. Trends Cogn. Sci. 15, 56–67. https://doi.org/10.1016/j.tics.2010.12.004

Graydon, F.X., Young, R., Benton, M.D., Genik, R.J., Posse, S., Hsieh, L., Green, C., 2004.

Visual event detection during simulated driving: Identifying the neural correlates with

functional neuroimaging. Transp. Res. Part F Traffic Psychol. Behav. 7, 271–286.

https://doi.org/10.1016/j.trf.2004.09.006

Haber, S.N., Fudge, J.L., McFarland, N.R., 2000. Striatonigrostriatal Pathways in Primates

Form an Ascending Spiral from the Shell to the Dorsolateral Striatum. J. Neurosci. 20,

2369–2382. https://doi.org/10.1523/JNEUROSCI.20-06-02369.2000

Haber, S.N., Knutson, B., 2009. The Reward Circuit: Linking Primate Anatomy and Human

917   Imaging. Neuropsychopharmacology 35, 4–26. https://doi.org/10.1038/npp.2009.129

918 Haith, A.M., Krakauer, J.W., 2018. The multiple effects of practice: skill, habit and reduced

919   cognitive load. Curr. Opin. Behav. Sci. 20, 196–201.

920   https://doi.org/10.1016/j.cobeha.2018.01.015

921 Hardwick, R.M., Forrence, A.D., Krakauer, J.W., Haith, A.M., 2019. Time-dependent

922   competition between goal-directed and habitual response preparation. Nat. Hum.

923   Behav. 3, 1252–1262. https://doi.org/10.1038/s41562-019-0725-0

924 Haruno, M., Kawato, M., 2006. Different neural correlates of reward expectation and reward

925   expectation error in the putamen and caudate nucleus during stimulus-action-reward

926   association learning. J. Neurophysiol. 95, 948–959.

927   https://doi.org/10.1152/jn.00382.2005

928 Haynes, W.I.A., Haber, S.N., 2013. The organization of prefrontal-subthalamic inputs in

929   primates provides an anatomical substrate for both functional specificity and integration:

930   Implications for basal ganglia models and deep brain stimulation. J. Neurosci. 33,

931   4804–4814. https://doi.org/10.1523/JNEUROSCI.4674-12.2013

932 Hernandez, L.F., Obeso, I., Costa, R.M., Redgrave, P., Obeso, J.A., 2019. Dopaminergic

933   Vulnerability in Parkinson Disease: The Cost of Humans' Habitual Performance. Trends

934   Neurosci. 42, 375–383. https://doi.org/10.1016/j.tins.2019.03.007

935 Hikosaka, O., Takikawa, Y., Kawagoe, R., Kojima, S., Doupe, A.J., Silver, R., Boahen, K.,

936   Grillner, S., Kopell, N., Olsen, K.L., Neurosci F-W Zhou, J., Matta, S.G., Zhou, F.,

937   Neurophysiol, J., Bargas Carrillo-Reid, J.L., Tecuapetla, F., Tapia, D., Hernandez-Cruz,

938   A., Galarraga, E., Drucker-Colin, R., 2000. Tutor Song Exposure Song Selectivity in the

939   Pallial-Basal Ganglia Song Circuit of Zebra Finches Raised Without Role of the Basal

940   Ganglia in the Control of Purposive Saccadic Eye Movements. Physiol Rev 80, 953–

941   978.

Holl, A.K., Wilkinson, L., Tabrizi, S.J., Painold, A., Jahanshahi, M., 2012. Probabilistic

classification learning with corrective feedback is selectively impaired in early

Huntington's disease--evidence for the role of the striatum in learning with feedback.

Neuropsychologia 50, 2176–2186.

https://doi.org/10.1016/J.NEUROPSYCHOLOGIA.2012.05.021

Horovitz, S.G., Gallea, C., Najee-ullah, M.A., Hallett, M., 2013. Functional Anatomy of

Writing with the Dominant Hand. PLoS One 8, 1–10.

https://doi.org/10.1371/journal.pone.0067931

Hsieh, L., Young, R.A., Bowyer, S.M., Moran, J.E., Genik, R.J., Green, C.C., Chiang, Y.R.,

Yu, Y.J., Liao, C.C., Seaman, S., 2009. Conversation effects on neural mechanisms

underlying reaction time to visual events while viewing a driving scene: fMRI analysis

and asynchrony model. Brain Res. https://doi.org/10.1016/j.brainres.2008.10.002

Hsu, C.T., Jacobs, A.M., Citron, F.M.M., Conrad, M., 2015. The emotion potential of words

and passages in reading Harry Potter - An fMRI study. Brain Lang. 142, 96–114.

https://doi.org/10.1016/j.bandl.2015.01.011

Hu, X.Y., Wang, L., Liu, H., Zhang, S.Z., 2006. Functional magnetic resonance imaging

study of writer's cramp. Chin. Med. J. (Engl). 119, 1263–1271.

https://doi.org/10.1097/00029330-200608010-00006

Huang, Y., Yaple, Z.A., Yu, R., 2020. Goal-oriented and habitual decisions: Neural

signatures of model-based and model-free learning. Neuroimage 215, 116834.

https://doi.org/10.1016/j.neuroimage.2020.116834

Huth, A.G., De Heer, W.A., Griffiths, T.L., Theunissen, F.E., Gallant, J.L., 2016. Natural

speech reveals the semantic maps that tile human cerebral cortex. Nature 532, 453–

458. https://doi.org/10.1038/nature17637

Iaria, G., Petrides, M., Dagher, A., Pike, B., Bohbot, V.D., 2003. Cognitive strategies

dependent on the hippocampus and caudate nucleus in human navigation: variability

and change with practice. J. Neurosci. Off. J. Soc. Neurosci. 23, 5945–5952.

https://doi.org/10.1523/jneurosci.23-13-05945.2003

Jaeger L, Marchal-Crespo L, Wolf P, Riener R, Michels L, K.S., 2014. Brain activation

associated with active and passive lower limb stepping. Front. Hum. Neurosci. 8.

https://doi.org/10.3389/fnhum.2014.00828

Jankowski, J., Scheef, L., Hüppe, C., Boecker, H., 2009. Distinct striatal regions for planning

and executing novel and automated movement sequences. Neuroimage 44, 1369–

1379. https://doi.org/10.1016/j.neuroimage.2008.10.059

Jeanvoine, H., Labriffe, M., Tannou, T., Navasiolava, N., Ter Minassian, A., Girot, J.B.,

Leiber, L.M., Custaud, M.A., Annweiler, C., Dinomais, M., 2022. Specific age-correlated

activation of top hierarchical motor control areas during gait-like plantar stimulation: An

fMRI study. Hum. Brain Mapp. 43, 833–843. https://doi.org/10.1002/HBM.25691

Karimpoor, M., Churchill, N.W., Tam, F., Fischer, C.E., Schweizer, T.A., Graham, S.J., 2018.

Functional MRI of handwriting tasks: A study of healthy young adults interacting with a

novel touch-sensitive tablet. Front. Hum. Neurosci. 12, 1–14.

https://doi.org/10.3389/fnhum.2018.00030

Karimpoor, M., Tam, F., Strother, S.C., Fischer, C.E., Schweizer, T.A., Graham, S.J., 2015.

A computerized tablet with visual feedback of hand position for functional magnetic

resonance imaging. Front. Hum. Neurosci. 9, 150.

https://doi.org/10.3389/fnhum.2015.00150

Katanoda, K., Yoshikawa, K., Sugishita, M., 2001. A functional MRI study on the neural

substrates for writing. Hum. Brain Mapp. 13, 34–42. https://doi.org/10.1002/hbm.1023

Kilner, J.M., Neal, A., Weiskopf, N., Friston, K.J., Frith, C.D., 2009. Evidence of Mirror

Neurons in Human Inferior Frontal Gyrus. J. Neurosci. 29, 10153.

992  https://doi.org/10.1523/JNEUROSCI.2668-09.2009

993 Kim, W., Chang, Y., Kim, J., Seo, J., Ryu, K., Lee, E., Woo, M., Janelle, C.M., 2014. An fMRI

994  study of differences in brain activity among elite, expert, and novice archers at the

995  moment of optimal aiming. Cogn. Behav. Neurol. 27, 173–182.

996  https://doi.org/10.1097/WNN.0000000000000042

997 Kimura, M., 1986. The role of primate putamen neurons in the association of sensory stimuli

998  with movement. Neurosci. Res. 3, 436–443. https://doi.org/10.1016/0168-

999  0102(86)90035-0

1000 Kimura, M., Kato, M., Shimazaki, H., 1990. Physiological properties of projection neurons in

1001  the monkey striatum to the globus pallidus. Exp. Brain Res. 82, 672–676.

1002  https://doi.org/10.1007/BF00228811

1003 Kimura, M., Rajkowski, J., Evarts, E., 1984. Tonically discharging putamen neurons exhibit

1004  set-dependent responses. Proc. Natl. Acad. Sci. U. S. A. 81, 4998–5001.

1005  https://doi.org/10.1073/pnas.81.15.4998

1006 Kunimatsu, J., Maeda, K., Hikosaka, O., 2019. The caudal part of putamen represents the

1007  historical object value information. J. Neurosci. 39, 1709–1719.

1008  https://doi.org/10.1523/JNEUROSCI.2534-18.2018

1009 la Fougère, C., Zwergal, A., Rominger, A., Förster, S., Fesl, G., Dieterich, M., Brandt, T.,

1010  Strupp, M., Bartenstein, P., Jahn, K., 2010. Real versus imagined locomotion: A [18F]-

1011  FDG PET-fMRI comparison. Neuroimage 50, 1589–1598.

1012  https://doi.org/10.1016/j.neuroimage.2009.12.060

1013 Lancaster, J.L., Tordesillas-Gutiérrez, D., Martinez, M., Salinas, F., Evans, A., Zilles, K.,

1014  Mazziotta, J.C., Fox, P.T., 2007. Bias between MNI and talairach coordinates analyzed

1015  using the ICBM-152 brain template. Hum. Brain Mapp. 28, 1194–1205.

1016  https://doi.org/10.1002/hbm.20345

Leh, S.E., Ptito, A., Chakravarty, M.M., Strafella, A.P., 2007. Fronto-striatal connections in

the human brain: A probabilistic diffusion tractography study. Neurosci. Lett. 419, 113–

118. https://doi.org/10.1016/j.neulet.2007.04.049

Lehéricy, S., Benali, H., Van De Moortele, P.F., Pélégrini-Issac, M., Waechter, T., Ugurbil,

K., Doyon, J., 2005a. Distinct basal ganglia territories are engaged in early and

advanced motor sequence learning. Proc. Natl. Acad. Sci. U. S. A. 102, 12566–12571.

https://doi.org/10.1073/pnas.0502762102

Lehéricy, S., Benali, H., Van De Moortele, P.F., Pélégrini-Issac, M., Waechter, T., Ugurbil,

K., Doyon, J., 2005b. Distinct basal ganglia territories are engaged in early and

advanced motor sequence learning. Proc. Natl. Acad. Sci. U. S. A. 102, 12566–12571.

https://doi.org/10.1073/pnas.0502762102

Leisman, G., Moustafa, A., Shafir, T., 2016. Thinking, Walking, Talking: Integratory Motor

and Cognitive Brain Function. Front. Public Heal. 4, 1.

https://doi.org/10.3389/fpubh.2016.00094

Liljeholm, M., Dunne, S., O'Doherty, J.P., 2015a. Differentiating neural systems mediating

the acquisition vs. expression of goal-directed and habitual behavioral control. Eur. J.

Neurosci. 41, 1358–1371. https://doi.org/10.1111/ejn.12897

Liljeholm, M., Dunne, S., O'Doherty, J.P., 2015b. Differentiating neural systems mediating

the acquisition vs. expression of goal-directed and habitual behavioral control. Eur. J.

Neurosci. 41, 1358–1371. https://doi.org/10.1111/ejn.12897

Liljeholm, M., O'doherty, J.P., 2012. Contributions of the striatum to learning, motivation, and

performance: an associative account. Trends Cogn Sci 16, 467–475.

https://doi.org/10.1016/j.tics.2012.07.007

Lin, Z., Tam, F., Churchill, N.W., Schweizer, T.A., Graham, S.J., 2021. Tablet technology for

writing and drawing during functional magnetic resonance imaging: A review. Sensors

(Switzerland). https://doi.org/10.3390/s21020401

Longcamp, M., Lagarrigue, A., Nazarian, B., Roth, M., Anton, J.L., Alario, F.X., Velay, J.L.,

2014. Functional specificity in the motor system: Evidence from coupled fMRI and

kinematic recordings during letter and digit writing. Hum. Brain Mapp. 35, 6077–6087.

https://doi.org/10.1002/hbm.22606

Luque, D., Beesley, T., Morris, R.W., Jack, B.N., Griffiths, O., Whitford, T.J., Le Pelley, M.E.,

2017. Goal-Directed and Habit-Like Modulations of Stimulus Processing during

Reinforcement Learning. https://doi.org/10.1523/JNEUROSCI.3205-16.2017

Luque, D., Molinero, S., Watson, P., López, F.J., Le Pelley, M.E., 2020. Measuring habit

formation through goal-directed response switching. J. Exp. Psychol. Gen. 149, 1449–

1459. https://doi.org/10.1037/xge0000722

Marchal, V., Sellers, J., Pélégrini-Issac, M., Galléa, C., Bertasi, E., Valabrègue, R., Lau, B.,

Leboucher, P., Bardinet, E., Welter, M.L., Karachi, C., 2019. Deep brain activation

patterns involved in virtual gait without and with a doorway: An fMRI study. PLoS One

14. https://doi.org/10.1371/journal.pone.0223494

Martínez, M., Valencia, M., Vidorreta, M., Luis, E.O., Castellanos, G., Villagra, F., Fern??

ndez-Seara, M.A., Pastor, M.A., 2016a. Trade-off between frequency and precision

during stepping movements: Kinematic and BOLD brain activation patterns. Hum. Brain

Mapp. 37, 1722–1737. https://doi.org/10.1002/hbm.23131

Martínez, M., Valencia, M., Vidorreta, M., Luis, E.O., Castellanos, G., Villagra, F.,

Fernández-Seara, M.A., Pastor, M.A., 2016b. Trade-off between frequency and

precision during stepping movements: Kinematic and BOLD brain activation patterns.

Hum. Brain Mapp. 37, 1722–1737. https://doi.org/10.1002/hbm.23131

Martínez, M., Villagra, F., Castellote, J.M., Pastor, M.A., 2018. Kinematic and kinetic

patterns related to free-walking in parkinson's disease. Sensors (Switzerland) 18.

https://doi.org/10.3390/s18124224

Martinez, M., Villagra, F., Loayza, F., Vidorreta, M., Arrondo, G., Luis, E., Diaz, J.,

Echeverria, M., Fernandez-Seara, M.A., Pastor, M.A., 2014. MRI-compatible device for

examining brain activation related to stepping. IEEE Trans. Med. Imaging 33, 1044–

1053. https://doi.org/10.1109/TMI.2014.2301493

Mas-Herrero, E., Sescousse, G., Cools, R., Marco-Pallarés, J., 2019. The contribution of

striatal pseudo-reward prediction errors to value-based decision-making. Neuroimage

193, 67–74. https://doi.org/10.1016/j.neuroimage.2019.02.052

Matsuda, W., Furuta, T., Nakamura, K.C., Hioki, H., Fujiyama, F., Arai, R., Kaneko, T., 2009.

Single nigrostriatal dopaminergic neurons form widely spread and highly dense axonal

arborizations in the neostriatum. J. Neurosci. 29, 444–453.

https://doi.org/10.1523/JNEUROSCI.4029-08.2009

Mazziotta, J., Toga, A., Evans, A., Fox, P., Lancaster, J., Zilles, K., Woods, R., Paus, T.,

Simpson, G., Pike, B., Holmes, C., Collins, L., Thompson, P., MacDonald, D., Iacoboni,

M., Schormann, T., Amunts, K., Palomero-Gallagher, N., Geyer, S., Parsons, L., Narr,

K., Kabani, N., Le Goualher, G., Boomsma, D., Cannon, T., Kawashima, R., Mazoyer,

B., 2001. A probabilistic atlas and reference system for the human brain: International

Consortium for Brain Mapping (ICBM). Philos. Trans. R. Soc. B Biol. Sci.

https://doi.org/10.1098/rstb.2001.0915

McFarland, N.R., Haber, S.N., 2002. Thalamic relay nuclei of the basal ganglia form both

reciprocal and nonreciprocal cortical connections, linking multiple frontal cortical areas.

J. Neurosci. 22, 8117–8132. https://doi.org/10.1523/jneurosci.22-18-08117.2002

Mechelli, A., Gorno-Tempini, M.L., Price, C.J., 2003. Neuroimaging studies of word and

pseudoword reading: Consistencies, inconsistencies, and limitations. J. Cogn.

Neurosci. 15, 260–271. https://doi.org/10.1162/089892903321208196

Meschyan, G., Hernandez, A.E., 2006. Impact of language proficiency and orthographic

transparency on bilingual word reading: An fMRI investigation. Neuroimage 29, 1135–

1140. https://doi.org/10.1016/j.neuroimage.2005.08.055

Mi, T.M., Zhang, W., McKeown, M.J., Chan, P., 2021. Impaired Formation and Expression of

Goal-Directed and Habitual Control in Parkinson's Disease. Front. Aging Neurosci. 13.

https://doi.org/10.3389/FNAGI.2021.734807

Miller, K.J., Shenhav, A., Ludvig, E.A., 2019. Habits without values. Psychol. Rev.

https://doi.org/10.1037/rev0000120

Molenberghs, P., Cunnington, R., Mattingley, J.B., 2012. Brain regions with mirror

properties: A meta-analysis of 125 human fMRI studies. Neurosci. Biobehav. Rev. 36,

341–349. https://doi.org/10.1016/J.NEUBIOREV.2011.07.004

Moore, C.J., Price, C.J., 1999. Three distinct ventral occipitotemporal regions for reading

and object naming. Neuroimage 10, 181–192. https://doi.org/10.1006/nimg.1999.0450

Morris, L.S., Kundu, P., Dowell, N., Mechelmans, D.J., Favre, P., Irvine, M.A., Robbins,

T.W., Daw, N., Bullmore, E.T., Harrison, N.A., Voon, V., 2016. Fronto-striatal

organization: Defining functional and microstructural substrates of behavioural flexibility.

Cortex 74, 118–133. https://doi.org/10.1016/j.cortex.2015.11.004

Morris, N.J., Stein, C.M., 2017. Model-free linkage analysis of a quantitative trait, in:

Methods in Molecular Biology. Humana Press Inc., pp. 327–342.

https://doi.org/10.1007/978-1-4939-7274-6_16

Müller, V.I., Cieslik, E.C., Laird, A.R., Fox, P.T., Radua, J., Mataix-Cols, D., Tench, C.R.,

Yarkoni, T., Nichols, T.E., Turkeltaub, P.E., Wager, T.D., Eickhoff, S.B., 2018. Ten

simple rules for neuroimaging meta-analysis. Neurosci. Biobehav. Rev. 84, 151.

https://doi.org/10.1016/J.NEUBIOREV.2017.11.012

Murphy, K., Garavan, H., 2004. An empirical investigation into the number of subjects

required for an event-related fMRI study. Neuroimage 22, 879–885.

https://doi.org/10.1016/J.NEUROIMAGE.2004.02.005

Nachev, P., Kennard, C., Husain, M., 2008. Functional role of the supplementary and pre-

supplementary motor areas. Nat. Rev. Neurosci. https://doi.org/10.1038/nrn2478

Nakamura, K., Honda, M., Hirano, S., Oga, T., Sawamoto, N., Hanakawa, T., Inoue, H., Ito,

J., Matsuda, T., Fukuyama, H., Shibasaki, H., 2002. Modulation of the visual word

retrieval system in writing: A functional MRI study on the Japanese orthographies. J.

Cogn. Neurosci. 14, 104–115. https://doi.org/10.1162/089892902317205366

Nebe, S., Kroemer, N.B., Schad, D.J., Bernhardt, N., Sebold, M., Müller, D.K., Scholl, L.,

Kuitunen-Paul, S., Heinz, A., Rapp, M.A., Huys, Q.J.M., Smolka, M.N., 2018. No

association of goal-directed and habitual control with alcohol consumption in young

adults. Addict. Biol. 23, 379–393. https://doi.org/10.1111/ADB.12490

Noble, J.W., Eng, J.J., Boyd, L.A., 2014. Bilateral motor tasks involve more brain regions

and higher neural activation than unilateral tasks: An fMRI study. Exp. Brain Res. 232,

2785–2795. https://doi.org/10.1007/s00221-014-3963-4

Oberhuber, M., Hope, T.M.H., Seghier, M.L., Parker Jones, O., Prejawa, S., Green, D.W.,

Price, C.J., 2016. Four Functionally Distinct Regions in the Left Supramarginal Gyrus

Support Word Processing. Cereb. Cortex 26, 4212–4226.

https://doi.org/10.1093/cercor/bhw251

Oberhuber, M., Jones, O.P., Hope, T.M.H., Prejawa, S., Seghier, M.L., Green, D.W., Price,

C.J., 2013. Functionally distinct contributions of the anterior and posterior putamen

during sublexical and lexical reading. Front. Hum. Neurosci.

https://doi.org/10.3389/fnhum.2013.00787

Oh-Descher, H., Beck, J.M., Ferrari, S., Sommer, M.A., Egner, T., 2017. Probabilistic

inference under time pressure leads to a cortical-to-subcortical shift in decision

evidence integration. Neuroimage 162, 138–150.

https://doi.org/10.1016/j.neuroimage.2017.08.069

Ohbayashi, M., Picard, N., Strick, P.L., 2016. Inactivation of the dorsal premotor area

disrupts internally generated, but not visually guided, sequential movements. J.

Neurosci. 36, 1971–1976. https://doi.org/10.1523/JNEUROSCI.2356-15.2016

Parkes, S.L., Bradfield, L.A., Balleine, B.W., 2015. Interaction of insular cortex and ventral

striatum mediates the effect of incentive memory on choice between goal-directed

actions. J. Neurosci. 35, 6464–6471. https://doi.org/10.1523/JNEUROSCI.4153-

14.2015

Patterson, T.K., Knowlton, B.J., 2018. Subregional specificity in human striatal habit

learning: a meta-analytic review of the fMRI literature. Curr. Opin. Behav. Sci. 20, 75–

82. https://doi.org/10.1016/j.cobeha.2017.10.005

Perez, Omar D, Dickinson, A., 2020. A theory of actions and habits: The interaction of rate

correlation and contiguity systems in free-operant behavior.

https://doi.org/10.1037/rev0000201

Perez, Omar D., Dickinson, A., 2020. A theory of actions and habits: The interaction of rate

correlation and contiguity systems in free-operant behavior. Psychol. Rev. 127, 945–

971. https://doi.org/10.1037/REV0000201

Peters, A.J., Fabre, J.M.J., Steinmetz, N.A., Harris, K.D., Carandini, M., 2021. Striatal

activity topographically reflects cortical activity. Nature 591, 420–425.

https://doi.org/10.1038/s41586-020-03166-8

Peters, S., Eng, J.J., Liu-Ambrose, T., Borich, M.R., Dao, E., Amanian, A., Boyd, L.A., 2019.

Brain activity associated with Dual-task performance of Ankle motor control during

cognitive challenge. Brain Behav. 9. https://doi.org/10.1002/brb3.1349

Peterson, B.S., Riddle, M.A., Cohen, D.J., Katz, L.D., Smith, J.C., Leckman, J.F., 1993.

Human basal ganglia volume asymmetries on magnetic resonance images. Magn.

Reson. Imaging 11, 493–498. https://doi.org/10.1016/0730-725X(93)90468-S

Peterson, D.S., Horak, F.B., 2016. Neural control of walking in people with parkinsonism.

Physiology. https://doi.org/10.1152/physiol.00034.2015

Pineda-Pardo, J.D.S.A., Sánchez-Ferro, Á., Monje, M.H.G., Pavese, N., Obeso, J.A., 2022.

Onset pattern of nigrostriatal denervation in early Parkinson's disease. Brain 145,

1018–1028. https://doi.org/10.1093/BRAIN/AWAB378

Poldrack, R.A., Clark, J., Paré-Blagoev, E.J., Shohamy, D., Creso Moyano, J., Myers, C.,

Gluck, M.A., 2001. Interactive memory systems in the human brain. Nature 414, 546–

550. https://doi.org/10.1038/35107080

Poldrack, R.A., Prabhakaran, V., Seger, C.A., Gabrieli, J.D.E., 1999. Striatal activation

during acquisition of a cognitive skill. Neuropsychology 13, 564–574.

https://doi.org/10.1037//0894-4105.13.4.564

Pool, E., Gera, R., Fransen, A., Perez, O.D., Cremer, A., Aleksic, M., Tanwisuth, S., Quail,

S., Ceceli, A.O., Manfredi, D., Nave, G., Tricomi, E., Balleine, B., Schonberg, T.,

Schwabe, L., O'Doherty, J.P., n.d. Determining the effects of training duration on the

behavioral expression of habitual control in humans: a multi-laboratory investigation.

https://doi.org/10.31234/OSF.IO/Z756H

Potgieser, A.R.E., Van Der Hoorn, A., De Jong, B.M., 2015. Cerebral Activations Related to

Writing and Drawing with each hand. PLoS One 10, 1–25.

https://doi.org/10.1371/journal.pone.0126723

Redgrave, P., Gurney, K., Reynolds, J., 2008. What is reinforced by phasic dopamine

signals? Brain Res. Rev. https://doi.org/10.1016/j.brainresrev.2007.10.007

Redgrave, P., Prescott, T.J., Gurney, K., 1999. The basal ganglia: a vertebrate solution to

the selection problem? Neuroscience 89, 1009–1023.

https://doi.org/S0306452298003194 [pii]

Redgrave, P., Rodriguez, M., Smith, Y., Rodriguez-Oroz, M.C., Lehericy, S., Bergman, H.,

Agid, Y., Delong, M.R., Obeso, J.A., 2010a. Goal-directed and habitual control in the

basal ganglia: Implications for Parkinson's disease. Nat. Rev. Neurosci. 11, 760–772.

https://doi.org/10.1038/nrn2915

Redgrave, P., Rodriguez, M., Smith, Y., Rodriguez-Oroz, M.C., Lehericy, S., Bergman, H.,

Agid, Y., Delong, M.R., Obeso, J.A., 2010b. Goal-directed and habitual coRedgrave, P.,

Rodriguez, M., Smith, Y., Rodriguez-Oroz, M. C., Lehericy, S., Bergman, H., Agid, Y.,

Delong, M. R., & Obeso, J. A. (2010). Goal-directed and habitual control in the basal

ganglia: Implications for Parkinson's di. Nat. Rev. Neurosci. 11, 760–772.

https://doi.org/10.1038/nrn2915

Romero, M.C., Bermudez, M.A., Vicente, A.F., Perez, R., Gonzalez, F., 2008. Activity of

neurons in the caudate and putamen during a visuomotor task. Neuroreport 19, 1141–

1145. https://doi.org/10.1097/WNR.0b013e328307c3fc

Rueckl, J.G., Paz-Alonso, P.M., Molfese, P.J., Kuo, W.J., Bick, A., Frost, S.J., Hancock, R.,

Wu, D.H., Einar Mencl, W., Duñabeitia, J.A., Lee, J.R., Oliver, M., Zevin, J.D., Hoeft, F.,

Carreiras, M., Tzeng, O.J.L., Pugh, K.R., Frost, R., 2015. Universal brain signature of

proficient reading: Evidence from four contrasting languages. Proc. Natl. Acad. Sci. U.

S. A. 112, 15510–15515. https://doi.org/10.1073/pnas.1509321112

Sauvage, C., Jissendi, P., Seignan, S., Manto, M., Habas, C., 2013. Brain areas involved in

the control of speed during a motor sequence of the foot: Real movement versus

mental imagery. J. Neuroradiol. 40, 267–280.

https://doi.org/10.1016/j.neurad.2012.10.001

Schultz, W., 2006. Behavioral theories and the neurophysiology of reward. Annu. Rev.

Psychol. 57, 87–115. https://doi.org/10.1146/annurev.psych.56.091103.070229

Schwabe, L., Tegenthoff, M., Höffken, O., Wolf, O.T., 2013. Mineralocorticoid Receptor

Blockade Prevents Stress-Induced Modulation of Multiple Memory Systems in the

Human Brain. Biol. Psychiatry 74, 801–808.

https://doi.org/10.1016/J.BIOPSYCH.2013.06.001

Segal, E., Petrides, M., 2012. The anterior superior parietal lobule and its interactions with

language and motor areas during writing. Eur. J. Neurosci. 35, 309–322.

https://doi.org/10.1111/j.1460-9568.2011.07937.x

Seghier, M.L., Lee, H.L., Schofield, T., Ellis, C.L., Price, C.J., 2008. Inter-subject variability in

the use of two different neuronal networks for reading aloud familiar words. Neuroimage

42, 1226–1236. https://doi.org/10.1016/j.neuroimage.2008.05.029

Seghier, M.L., Price, C.J., 2010. Reading Aloud Boosts Connectivity through the Putamen.

Cereb. Cortex 20, 570–582. https://doi.org/10.1093/cercor/bhp123

Sescousse, G., Caldú, X., Segura, B., Dreher, J.C., 2013. Processing of primary and

secondary rewards: A quantitative meta-analysis and review of human functional

neuroimaging studies. Neurosci. Biobehav. Rev. 37, 681–696.

https://doi.org/10.1016/j.neubiorev.2013.02.002

Sjoerds, Z., De Wit, S., Van Den Brink, W., Robbins, T.W., Beekman, A.T.F., Penninx,

B.W.J.H., Veltman, D.J., 2013. Behavioral and neuroimaging evidence for overreliance

on habit learning in alcohol-dependent patients. Transl. Psychiatry 3, e337-8.

https://doi.org/10.1038/tp.2013.107

Smith, K.S., Graybiel, A.M., 2016. Habit formation. Dialogues Clin. Neurosci. 18, 33–43.

https://doi.org/10.31887/dcns.2016.18.1/ksmith

Smittenaar, P., Guitart-Masip, M., Lutti, A., Dolan, R.J., 2013. Preparing for selective

inhibition within frontostriatal loops. J. Neurosci. 33, 18087–18097.

https://doi.org/10.1523/JNEUROSCI.2167-13.2013

Soares, J.M., Sampaio, A., Ferreira, L.M., Santos, N.C., Marques, F., Palha, J.A., Cerqueira,

J.J., Sousa, N., 2012. Stress-induced changes in human decision-making are

reversible. Transl. Psychiatry 2. https://doi.org/10.1038/TP.2012.59

Sommer, S., Pollmann, S., 2016. Putamen Activation Represents an Intrinsic Positive

Prediction Error Signal for Visual Search in Repeated Configurations. Open Neuroimag.

J. 10, 126–138. https://doi.org/10.2174/1874440001610010126

Spiers, H.J., Maguire, E.A., 2007. Neural substrates of driving behaviour. Neuroimage 36,

245–255. https://doi.org/10.1016/j.neuroimage.2007.02.032

Steele, C.J., Penhune, V.B., 2010. Specific Increases within Global Decreases: A Functional

Magnetic Resonance Imaging Investigation of Five Days of Motor Sequence Learning.

J. Neurosci. 30, 8332–8341. https://doi.org/10.1523/JNEUROSCI.5569-09.2010

Sun, R., 2004. Desiderata for cognitive architectures. Philos. Psychol. 17, 341–373.

https://doi.org/10.1080/0951508042000286721

Swett, B.A., Contreras-Vidal, J.L., Birn, R., Braun, A., 2010. Neural Substrates of

Graphomotor Sequence Learning: A Combined fMRI and Kinematic Study. J.

Neurophysiol. 103, 3366. https://doi.org/10.1152/JN.00449.2009

Swinnen, S.P., Vangheluwe, S., Wagemans, J., Coxon, J.P., Goble, D.J., Van Impe, A.,

Sunaert, S., Peeters, R., Wenderoth, N., 2010. Shared neural resources between left

and right interlimb coordination skills: The neural substrate of abstract motor

representations. Neuroimage 49, 2570–2580.

https://doi.org/10.1016/j.neuroimage.2009.10.052

Tanaka, S.C., Balleine, B.W., O'Doherty, J.P., 2008. Calculating consequences: Brain

systems that encode the causal effects of actions. J. Neurosci. 28, 6750–6755.

https://doi.org/10.1523/JNEUROSCI.1808-08.2008

Toyomura, A., Shibata, M., Kuriki, S., 2012. Self-paced and externally triggered rhythmical

lower limb movements: A functional MRI study. Neurosci. Lett. 516, 39–44.

https://doi.org/10.1016/j.neulet.2012.03.049

Tricomi, E., Balleine, B.W., O'Doherty, J.P., 2009. A specific role for posterior dorsolateral

striatum in human habit learning. Eur. J. Neurosci. 29, 2225–2232.

https://doi.org/10.1111/j.1460-9568.2009.06796.x

Trinastic, J.P., Kautz, S.A., McGregor, K., Gregory, C., Bowden, M., Benjamin, M.B.,

Kurtzman, M., Chang, Y.L., Conway, T., Crosson, B., 2010. An fMRI study of the

differences in brain activity during active ankle dorsiflexion and plantarflexion. Brain

Imaging Behav. 4, 121–131. https://doi.org/10.1007/s11682-010-9091-2

Tunik, E., Houk, J.C., Grafton, S.T., 2009. Basal ganglia contribution to the initiation of

corrective submovements. Neuroimage 47, 1757–1766.

https://doi.org/10.1016/j.neuroimage.2009.04.077

Turkeltaub, P.E., Eden, G.F., Jones, K.M., Zeffiro, T.A., 2002. Meta-analysis of the functional

neuroanatomy of single-word reading: Method and validation. Neuroimage 16, 765–

780. https://doi.org/10.1006/nimg.2002.1131

Uchiyama, Y., Ebe, K., Kozato, A., Okada, T., Sadato, N., 2003. The neural substrates of

driving at a safe distance: A functional MRI study. Neurosci. Lett. 352, 199–202.

https://doi.org/10.1016/j.neulet.2003.08.072

Uchiyama, Y., Toyoda, H., Sakai, H., Shin, D., Ebe, K., Sadato, N., 2012. Suppression of

brain activity related to a car-following task with an auditory task: An fMRI study.

Transp. Res. Part F Traffic Psychol. Behav. 15, 25–37.

1288 https://doi.org/10.1016/j.trf.2011.11.002

1289 Valentin, V. V., Dickinson, A., O'Doherty, J.P., 2007. Determining the neural substrates of

1290  goal-directed learning in the human brain. J. Neurosci. 27, 4019–4026.

1291  https://doi.org/10.1523/JNEUROSCI.0564-07.2007

1292 van Steenbergen, H., Watson, P., Wiers, R.W., Hommel, B., de Wit, S., 2017. Dissociable

1293  corticostriatal circuits underlie goal-directed vs. cue-elicited habitual food seeking after

1294  satiation: evidence from a multimodal MRI study. Eur. J. Neurosci. 46, 1815–1827.

1295  https://doi.org/10.1111/ejn.13586

1296 Vannest, J., Eaton, K.P., Henkel, D., Siegel, M., Tsevat, R.K., Allendorfer, J.B., Schefft, B.K.,

1297  Banks, C., Szaflarski, J.P., 2012. Cortical correlates of self-generation in verbal paired

1298  associate learning. Brain Res. 1437, 104–114.

1299  https://doi.org/10.1016/j.brainres.2011.12.020

1300 Vaquero, L., Hartmann, K., Ripollés, P., Rojo, N., Sierpowska, J., François, C., Càmara, E.,

1301  van Vugt, F.T., Mohammadi, B., Samii, A., Münte, T.F., Rodríguez-Fornells, A.,

1302  Altenmüller, E., 2016. Structural neuroplasticity in expert pianists depends on the age of

1303  musical training onset. Neuroimage 126, 106–119.

1304  https://doi.org/10.1016/j.neuroimage.2015.11.008

1305 Varotto, S.F., Farah, H., Bogenberger, K., van Arem, B., Hoogendoorn, S.P., 2020.

1306  Adaptations in driver behaviour characteristics during control transitions from full-range

1307  Adaptive Cruise Control to manual driving: an on-road study. Transp. A Transp. Sci. 16,

1308  776–806. https://doi.org/10.1080/23249935.2020.1720856

1309 Vigneau, M., Jobard, G., Mazoyer, B., Tzourio-Mazoyer, N., 2005. Word and non-word

1310  reading: What role for the Visual Word Form Area? Neuroimage 27, 694–705.

1311  https://doi.org/10.1016/j.neuroimage.2005.04.038

1312 Vonsattel, J.P.G., DiFiglia, M., 1998. Huntington disease. J. Neuropathol. Exp. Neurol. 57,

369–384. https://doi.org/10.1097/00005072-199805000-00001

Wan Lee, S., Shimojo, S., O'Doherty, J.P., 2014. Neural computations underlying arbitration

between model-based and model-free learning. Neuron 81, 687.

https://doi.org/10.1016/J.NEURON.2013.11.028

Watson, P., de Wit, S., 2018. Current limits of experimental research into habits and future

directions. Curr. Opin. Behav. Sci. 20, 33–39.

https://doi.org/10.1016/j.cobeha.2017.09.012

Watson, P., van Wingen, G., de Wit, S., 2018. Conflicted between goal-directed and habitual

control, an fMRI investigation. eNeuro 5. https://doi.org/10.1523/ENEURO.0240-

18.2018

Watson, P., Wiers, R.W., Hommel, B., De Wit, S., 2014. Working for food you don't desire.

Cues interfere with goal-directed food-seeking. Appetite 79, 139–148.

https://doi.org/10.1016/j.appet.2014.04.005

Wechsler, K., Drescher, U., Janouch, C., Haeger, M., Voelcker-Rehage, C., Bock, O., 2018.

Multitasking during simulated car driving: A comparison of young and older persons.

Front. Psychol. 9, 910. https://doi.org/10.3389/fpsyg.2018.00910

West, R., Brown, J., 2013. Addiction, habit and instrumental learning. Theory Addict. 114–

135. https://doi.org/10.1002/9781118484890.CH5

Willingham, D.B., Koroshetz, W.J., 1993. Evidence for dissociable motor skills Huntington's

disease patients. In. Psychobiology 21.

Wong, A.L., Goldsmith, J., Forrence, A.D., Haith, A.M., Krakauer, J.W., 2017. Reaction times

can reflect habits rather than computations. Elife 6. https://doi.org/10.7554/eLife.28075

Wood, W., Mazar, A., Neal, D., 2021. Habits and Goals in Human Behavior: Separate but

Interacting Systems. Perspect. Psychol. Sci. https://doi.org/10.31234/OSF.IO/QVRBY

Wood, W., Neal, D.T., 2007. A New Look at Habits and the Habit-Goal Interface. Psychol.

Rev. 114, 843–863. https://doi.org/10.1037/0033-295X.114.4.843

Worringer, B., Langner, R., Koch, I., Eickhoff, S.B., Eickhoff, C.R., Binkofski, F.C., 2019.

Common and distinct neural correlates of dual-tasking and task-switching: a meta-

analytic review and a neuro-cognitive processing model of human multitasking. Brain

Struct. Funct. 224, 1845–1869. https://doi.org/10.1007/s00429-019-01870-4

Wu, T., Hallett, M., Chan, P., 2015. Motor automaticity in Parkinson's disease. Neurobiol.

Dis. 82, 226–234. https://doi.org/10.1016/j.nbd.2015.06.014

Wu, T., Zhang, J., Hallett, M., Feng, T., Hou, Y., Chan, P., 2016. Neural correlates

underlying Micrographia in Parkinson's disease. Brain 139, 144–160.

https://doi.org/10.1093/brain/awv319

Wunderlich, K., Dayan, P., Dolan, R.J., 2012. Mapping value based planning and extensively

trained choice in the human brain. Nat. Neurosci. 15, 786–791.

https://doi.org/10.1038/nn.3068

Wymbs, N.F., Bassett, D.S., Mucha, P.J., Porter, M.A., Grafton, S.T., 2012. Differential

Recruitment of the Sensorimotor Putamen and Frontoparietal Cortex during Motor

Chunking in Humans. Neuron 74, 936–946.

https://doi.org/10.1016/j.neuron.2012.03.038

Yamada, H., Matsumoto, N., Kimura, M., 2004. Tonically Active Neurons in the Primate

Caudate Nucleus and Putamen Differentially Encode Instructed Motivational Outcomes

of Action. J. Neurosci. 24, 3500–3510. https://doi.org/10.1523/JNEUROSCI.0068-

04.2004

Yang, Y., Zhang, J., Meng, Z.L., Qin, L., Liu, Y.F., Bi, H.Y., 2018. Neural correlates of

orthographic access in Mandarin Chinese writing: An fMRI study of the word-frequency

effect. Front. Behav. Neurosci. 12, 1–12. https://doi.org/10.3389/fnbeh.2018.00288

Yarkoni, T., Speer, N.K., Balota, D.A., McAvoy, M.P., Zacks, J.M., 2008. Pictures of a

thousand words: Investigating the neural mechanisms of reading with extremely rapid

event-related fMRI. Neuroimage 42, 973–987.

https://doi.org/10.1016/j.neuroimage.2008.04.258

Zwosta, K., Ruge, H., Goschke, T., Wolfensteller, U., 2018. Habit strength is predicted by

activity dynamics in goal-directed brain systems during training. Neuroimage 165, 125–

137. https://doi.org/10.1016/j.neuroimage.2017.09.062

1369

1370

Acknowledgments

We are thankful to Blanca Caijao and Cristina Rubio (Universidad Rey Juan Carlos)

for their help in article search. MM is funded by a PhD Fellowship from Instituto de Salud

Carlos III (PFIS-FI720/00068). IO is funded by Instituto de Salud Carlos III (Miguel Servet,

CP18/00038) and AES-ISCIII (PI19/00298) from Ministry of Science and Innovation in Spain.

The funders played no role in the idea, design, data collection or analysis, decision to

publish or manuscript editing and writing.

Author Contributions

P.G. design, data collection, writing; M.M. design, data collection, analysis, writing;

P.R. writing; D.L. design, data collection, writing; I.O. design, data collection, writing;

Competing Interests statement

The authors declare no competing interest.

Figure Legends and Tables

Figure 1. Hypothetical diagram on striatal role in both everyday-life and experimental habits.
Activities with strong stimulus-response components such as writing, reading, walking or
driving were selected as everyday-life habits (see Box 1 for task measurements and details).
Striatal activations elicited by these activities can be compared to those evoked by stimulus-
response paradigms commonly used to investigate habits in the cognitive science literature.

Figure 2. Everyday-life and experimental thresholded activation maps. **a)** Axial views for the
main regions. Overlap regions at the right column are shown in blue. Note activation in the
cerebellar cortex for everyday-life studies, but it is omitted here for brevity. Z=52 view for the
experimental studies is shown as an unthresholded map for comparison (ALE value ≈ 0.01)
b) 3D striatum reconstruction showing all the activation that fall inside it. Overlap regions at
the right column are shown in blue. **c)** 3D striatum reconstruction showing the differential
activation of the Everyday -life > Experimental contrast (in green) and the Experimental >
Everyday-life contrast (in red).

Figure 3. Everyday-life cognitive-motor subcategories thresholded activation maps. **a)** Axial
views for the main regions. Overlap regions at the right column are shown in blue. **b)** 3D
striatum reconstruction showing all the activation that fall inside it. Overlap regions at the
right column are shown in blue. **c)** Foci distribution of all the studies that fall in the striatum.
Note that some foci may come from the same study.

Figure 4. Experimental subcategories thresholded activation maps. **a)** Axial views for the
main regions. In this case there are no overlap regions in the right column. **b)** 3D striatum
reconstruction showing all the activation that fall inside it. In this case there are no overlap
regions in the right column. **c)** Foci distribution of all the studies that fall in the striatum. Note
that some foci may come from the same study.

Table 1. Activation by everyday-life habits. Percentages of each brain region indicate how
much activation from a cluster fall into that region. Coordinates of any activation comprising
< 5% of its volume in a region are not shown for brevity. Activity peak coordinates reported
based on MNI152 space with region names from Harvard-Oxford atlas.

Table 2. Activation by experimental. Percentages of each brain region indicate how much
activation from a cluster fall into that region. Coordinates of any activation comprising < 5%
of its volume in a region are not shown for brevity. Activity peak coordinates reported based
on MNI152 space with region names from Harvard-Oxford atlas.

1415

1416

Supplementary material

1418

An fMRI meta-analysis of the role of the striatum in everyday-life vs

laboratory-developed habits

1421 Pasqualina Guida[1,2,6]*, Mario Michiels[1,2,6]*, Peter Redgrave[5], David Luque[3,4], Ignacio Obeso[1,2]

1422 * Co-first authors

1423

[1] *HM CINAC, Centro Integral de Neurociencias AC. Hospital Universitario HM Puerta del Sur,*
HM Hospitales, Madrid, Spain

[2]*CIBERNED, Instituto de Salud Carlos III, Madrid, Spain*

[3]*Departamento de Psicología Básica, Universidad Autónoma de Madrid, Madrid, Spain*

[4]*Departamento de Psicología Básica, Universidad de Málaga, Madrid, Spain*

[5]*Department of Psychology, University of Sheffield, Sheffield, S10 2TN, UK*

[6]*PhD Program in Neuroscience, Autonoma de Madrid University-Cajal Institute, Madrid,*
Spain 28029

1432

1433

Methods

The complete set of search terms were as follows:

1. Search on PubMed

Everyday-life habits:

(walking[MeSH Terms] OR walking[Title/Abstract]) OR (Driving behaviour[MeSH

Terms] OR Driving behaviour[Title/Abstract]) OR (Driving behavior[MeSH Terms] OR

Driving behavior[Title/Abstract]) OR (car driving[MeSH Terms] OR Car

driving[Title/Abstract]) OR (writing[MeSH Terms] OR writing[Title/Abstract]) OR

(handwriting[MeSH Terms] OR handwriting[Title/Abstract]) OR

1443   (reading[Title/Abstract] OR reading[MeSH Terms) AND (functional magnetic

1444   resonance imaging[MeSH Terms] OR fMRI[Title/Abstract]) AND ((basal

1445   ganglia[Title/Abstract] OR caudate[Title/Abstract] OR putamen[Title/Abstract] OR

1446   striatum[Title/Abstract]) OR (basal ganglia[MeSH Terms] OR caudate nucleus[MeSH

1447   Terms] OR putamen[MeSH Terms] OR striatum[MeSH Terms]))

1448

1449   Experimental habits:

1450   ("habit" OR "habits" OR "probabilistic classification" OR "weather prediction" OR

1451   "response learning" OR "instrumental conditioning" OR "instrumental learning" OR

1452   "reinforcement learning" OR "outcome devaluation" OR "sequential decision" OR "two

1453   step" OR "2 step") AND ("basal ganglia" OR "caudate" OR "putamen" OR "striatum")

1454   AND ("fMRI" OR "functional magnetic resonance imaging" OR "functional MRI") AND

1455   ("2017/06/22"[PDAT] : "3000/12/31"[PDAT])

1456

1457  2. Search on Web of Science

1458   Everyday-life habits:

1459   TI=(((driving) OR (car driving) OR (walking) OR (writing) OR (handwriting) OR

1460   (reading)) AND ((functional magnetic resonance imaging OR fMRI)) AND (((basal

1461   ganglia OR caudate OR putamen OR striatum)))) OR AB=(((driving) OR (car driving)

1462   OR (walking) OR (writing) OR (handwriting) OR (reading)) AND ((functional magnetic

1463   resonance imaging OR fMRI)) AND (((basal ganglia OR caudate OR putamen OR

1464   striatum))))

1465

1466   Experimental habits:

1467   (TI=((habit OR habits OR probabilistic classification OR weather prediction OR

1468   response learning OR instrumental conditioning OR instrumental learning OR

1469   reinforcement learning OR outcome devaluation OR sequential decision OR two step

1470   OR 2 step) AND (basal ganglia OR caudate OR putamen OR striatum) AND (fMRI

OR functional magnetic resonance imaging OR functional MRI))) OR AB=((habit OR

habits OR probabilistic classification OR weather prediction OR response learning

OR instrumental conditioning OR instrumental learning OR reinforcement learning

OR outcome devaluation OR sequential decision OR two step OR 2 step) AND

(basal ganglia OR caudate OR putamen OR striatum) AND (fMRI OR functional

magnetic resonance imaging OR functional MRI))

1477

Supplementary Material Figure Legends and Tables

Figure S1. PRISMA chart describing the steps followed for the selection of studies.

Table S1. Studies with everyday-life habits included in the meta-analysis.

Table S2. Studies with experimental habits included in the meta-analysis.

Table S3. Additional diagnostics used to check the heterogeneity for everyday-life studies.
The list details which studies have any foci contributing to the significant clusters in the ALE
meta-analysis for everyday-life studies.

(IN CSV)

Table S4. Additional diagnostics to check the heterogeneity for experimental studies. The list
details which studies have any foci contributing to the significant clusters in the ALE meta-
analysis for experimental studies.

(IN CSV)

1490

1491

1492

1493

1494

1495

1496

1497

1498

1499

1500

1501

1502

1503

1504

1505

1506

1507 **Figure S1.** PRISMA chart describing the steps followed for the selection of studies.

Table S1. Studies with everyday-life habits included in the meta-analysis.

Study by domain	n = 1536	Foci	Task	Contrast	Statistical threshold
Writing					
Katanoda et al., (2001)	17	19	Write a name of an object or naming the object	Writing > naming	p < 0.001 voxel-level & p < 0.05 cluster-level corrected
Nakamura et al., (2002)	9	10	Upon auditory presentation subjects wrote kanji word.	Writing > Rest	p < 0.05 corrected for multiple comparison
Beeson et al., (2003)	12	29	Writing words or drawing circles	Writing words > drawing circles	p < 0.001 uncorrected
Hu et al., (2006)	10	11	Writing with a pencil	Main Effect of Writing with a pencil	p < 0.0001 uncorrected
Segal & Petrides, (2012)	9	19	Writing or naming pictures of objects and drawing one loop per syllable of the object's name	Writing > naming plus loops	p < 0.05 corrected
Horovitz et al., (2013)	13	5	Writing, tapping, and zigzagging with each limb	Right-handwriting > other tasks	p < 0.001 FWE corrected
Erhard et al., (2014)	20	13	Creative writing	Main effects of creative writing in expert writers	p < 0.05 FWE corrected
Longcamp et al., (2014)	18	13	Writing of letters or digits	Writing > holding the pen still	p < 0.05 voxel-level FWE corrected
Potgieser et al., (2015)	16	21	Write a sentence or hand tapping	Right-handwriting > right-hand tapping	p < 0.001 voxel-level & p < 0.05 cluster-level FWE corrected & k≥8
Bisio et al., (2017)	7	25	Writing sentences	Writing > resting	p < 0.05 FWE corrected
Karimpoor et al., (2018)	12	38	Copying grocery lists, phone numbers or sentences	All handwriting tasks > resting	p < 0.05 FDR corrected
Yang et al., (2018)	34	29	Copying chinese characters	Writing high frequency > writing low frequency	p < 0.05 FDR corrected
Swett et al., (2010)	12	11	Learning Chinese word characters	Late learning > baseline	p < 0.05 uncorrected
Reading					
Bookheimer et al., (1995)	16	33	Reading words or naming objects	Main effect of read aloud	p < 0.001 corrected
Moore & Price, (1999)	8	7	Reading/naming	Main effect of reading words	p < 0.001 uncorrected
Mechelli et al., (2003)	20	10	Early/late reading processing	Early reading > Fixation	p < 0.05 corrected
Buchsbaum et al., (2005)	17	15	Reading/Hearing	Reading > Control	z > 2.33 & p < 0.01 cluster-corrected
Vigneau et al., (2005)	23	39	Reading words/ non reading letters	Read > Cross fixation	p < 0.001 uncorrected
Meschyan & Hernandez, (2006)	12	12	Reading different languages	Reading > Resting	p < 0.001 uncorrected & p < 0.05 corrected spatial extent threshold
Binder et al., (2006)	30	31	Non-word reading	Reading > Fixation	p < 0.00001 uncorrected & p < 0.05 corrected
Carreiras et al., (2007)	36	32	Reading/Lexical decision	Reading > Baseline	p < 0.05 corrected
Church et al., (2008)	50	37	Reading/Repeat	Reading > Repeat	p < 0.05 corrected with k≥24 voxels

1508

1509

1510

Table S2. Studies with experimental habits included in the meta-analysis.

Study by domain	n = 938	Foci	Task	Contrast (short)	Statistical threshold
Probabilistic learning					
Poldrack et al., (1999)	8	15	Probabilistic learning task	Task > control (perceptual-motor)	p < 0.05 corrected with Gaussian Random Field Theory
Poldrack et al., (2001)	13	25	Probabilistic learning task	Task > control (perceptual-motor)	p < 0.005 uncorrected & k≥5 voxels
Aron et al., (2004)	15	9	Probabilistic learning task	Task > baseline	p < 0.05 FDR corrected
Delgado et al., (2005)	17	2	Probabilistic learning task	Condition and time interaction	p < 0.0001 uncorrected & k≥5 voxels
Fera et al., (2005)	18	5	Probabilistic learning task	Task > control (perceptual-motor)	p < 0.05 uncorrected
Aron et al., (2006)	8	9	Probabilistic learning task	Task > control (perceptual-motor)	z > 2.3 voxel-level & p < 0.01 cluster-level corrected with Gaussian Random Field Theory
Foerde et al., (2006)	14	2	Probabilistic learning task	Modulation by performance (accuracy)	p < 0.05 corrected with Gaussian Random Field Theory
Gläscher et al., (2010)	18	1	2-step probabilistic learning task	RPE	p < 0.006 corrected & k≥10
Celone et al., (2011)	19	8	Probabilistic learning task	Task > control (perceptual-motor)	p < 0.01 voxel-level & p < 0.05 cluster-level corrected
Wunderlich et al., (2012)	21	17	2-step probabilistic learning task	Trained > Planning (RPE)	p < 0.05 FWE corrected & k≥5
Soares et al., (2012)	12	10	Probabilistic learning task	Stress > control	p < 0.05 FWE corrected
Schwabe et al., (2013)	75	14	Probabilistic learning task	Task > control (perceptual-motor)	p < 0.05 FWE corrected & k≥5
Wan Lee et al., (2014)	22	5	2-step probabilistic learning task	RPE	p < 0.05 FWE corrected
Deserno et al., (2015)	29	7	2-step probabilistic learning task	Model-free component	p < 0.05 corrected
Doll et al., (2015)	20	5	2-step probabilistic learning task	RPE	p ≤ 0.0005 cluster-level FWE corrected
Oh-Descher et al., (2017)	25	3	Probabilistic learning task	High time pressure > low time pressure Positive subjective sum of evidence)	p < 0.005 voxel-level & p < 0.05 cluster-level corrected
Nebe et al., (2018)	146	35	2-step probabilistic learning task	RPE	p < 0.05 FWE corrected

Erdeniz & Done, (2019)	19	15	Probabilistic learning task	CS familiar > CS novel	p < 0.005 uncorrected
Mas-Herrero et al., (2019)	20	2	2-step probabilistic learning task	RPE pseudofeedback)	p < 0.001 corrected

Others

Gottfried et al., (2003)	13	7	Associative learning	Outcome devaluation	p < 0.001 uncorrected
Iaria et al., (2003)	7	5	Maze navigation	Task > control perceptual-motor)	p < 0.05 corrected
Lehéricy et al., (2005)	14	3	Motor sequence learning	Modulation by task experience	p < 0.0001 voxel-level & p < 0.05 cluster-level corrected
Forstmann et al., (2008)	19	3	Moving dots task	Speed > accuracy	p < 0.001 corrected
Fernández-Seara et al., (2009)	14	30	Motor sequence learning	Late > early	p < 0.01 uncorrected
Tricomi et al., (2009)	15	4	Discriminative learning	Late > early	p < 0.001 uncorrected & k≥5
Steele & Penhune, (2010)	15	14	Motor sequence learning	Late > early	p < 0.001 uncorrected & k≥100
Brovelli et al., (2011)	14	3	Discriminative learning	P correct | past observations) > P correct by chance)	P < 0.05 FWE corrected
Beierholm et al., (2011)	23	11	Sequential choice	Model-free component	p < 0.001 uncorrected & k≥5
de Wit et al., (2012)	24	3	Discriminative learning	Negative predictors of goal-directed action outcome devaluation)	p < 0.005 uncorrected & k≥20
Etchamendy et al., (2012)	15	3	Maze navigation	Q-signal, learners	p < 0.05 corrected
Liljeholm & O'doherty, (2012)	15	4	Sequential choice	Decreased activity in the R-O but not int the S-R	p < 0.05 corrected with cluster size thresholding
Liljeholm et al., (2015)	19	13	Sequential choice	Modulation by devaluation insensitivity	p < 0.05 cluster-level corrected
Fermin et al., (2016)	18	9	Grid-Sailing	Previous S-R learning – Model-based conditions	p < 0.0001 uncorrected
Eryilmaz et al., (2017)	72	5	Discriminative learning	Incongruent > standard cue	p < 0.001 voxel-level & p < 0.05 cluster-level corrected
van Steenbergen et al., (2017)	19	5	Discriminative learning	Cue-driven > Value-driven action control	p < 0.001 uncorrected & k≥15
Zwosta et al., (2018)	53	29	Discriminative learning	Decrease reward and no reward Contingency degradation)	p < 0.001 voxel-level & p < 0.05 cluster-level corrected
Watson et al., (2018)	23	8	Discriminative learning	Slip of action trials > still-valued	p < 0.05 corrected

Everyday-life & Experimental Habits

				responses	
Anggraini et al., (2018)	27	16	Maze navigation	Model-free component	p < 0.001 voxel-level & p < 0.05 cluster-level corrected

1511

1512

1513

Supplementary References

Adams, C.D., 1982. Variations in the Sensitivity of Instrumental Responding to Reinforcer Devaluation. Q. J. Exp. Psychol. Sect. B 34, 77–98. https://doi.org/10.1080/14640748208400878

Alexander, G.E., Crutcher, M.D., 1990. Functional architecture of basal ganglia circuits: neural substrates of parallel processing. Trends Neurosci. https://doi.org/10.1016/0166-2236(90)90107-L

Allali, G., Montembeault, M., Brambati, S.M., Bherer, L., Blumen, H.M., Launay, C.P., Liu-Ambrose, T., Helbostad, J.L., Verghese, J., Beauchet, O., 2019. Brain structure covariance associated with gait control in aging. Journals Gerontol. - Ser. A Biol. Sci. Med. Sci. 74, 705–713. https://doi.org/10.1093/gerona/gly123

Amodio, D.M., 2014. The neuroscience of prejudice and stereotyping. Nat. Rev. Neurosci. 15, 670–682. https://doi.org/10.1038/nrn3800

Andersen, K.W., Madsen, K.H., Siebner, H.R., 2020. Discrete finger sequences are widely represented in human striatum. Sci. Rep. 10, 1–12. https://doi.org/10.1038/s41598-020-69923-x

Anggraini, D., Glasauer, S., Wunderlich, K., 2018. Neural signatures of reinforcement learning correlate with strategy adoption during spatial navigation. Sci. Reports 2018 81 8, 1–14. https://doi.org/10.1038/s41598-018-28241-z

Aron, A.R., Gluck, M.A., Poldrack, R.A., 2006. Long-term test-retest reliability of functional MRI in a classification learning task. Neuroimage 29, 1000–1006. https://doi.org/10.1016/J.NEUROIMAGE.2005.08.010

Aron, A.R., Shohamy, D., Clark, J., Myers, C., Gluck, M.A., Poldrack, R.A., 2004. Human midbrain sensitivity to cognitive feedback and uncertainty during classification learning.

J. Neurophysiol. 92, 1144–1152. https://doi.org/10.1152/JN.01209.2003

Arsalidou, M., Duerden, E.G., Taylor, M.J., 2013. The Centre of the Brain : Topographical

Model of Motor , Cognitive , Affective , and Somatosensory Functions of the Basal

Ganglia 3054, 3031–3054. https://doi.org/10.1002/hbm.22124

Balleine, B.W., O'Doherty, J.P., 2010. Human and rodent homologies in action control:

Corticostriatal determinants of goal-directed and habitual action.

Neuropsychopharmacology 35, 48–69. https://doi.org/10.1038/npp.2009.131

Bannard, C., Leriche, M., Bandmann, O., Brown, C.H., Ferracane, E., Sánchez-Ferro, Á.,

Obeso, J., Redgrave, P., Stafford, T., 2019. Reduced habit-driven errors in Parkinson's

Disease. Sci. Rep. 9, 1–8. https://doi.org/10.1038/s41598-019-39294-z

Beeson, P.M., Rapcsak, S.Z., Plante, E., Chargualaf, J., Chung, A., Johnson, S.C., Trouard,

T.P., 2003. The neural substrates of writing: A functional magnetic resonance imaging

study. Aphasiology 17, 647–665. https://doi.org/10.1080/02687030344000067

Beierholm, U.R., Anen, C., Quartz, S., Bossaerts, P., 2011. Separate encoding of model-

based and model-free valuations in the human brain. Neuroimage 58, 955–962.

https://doi.org/10.1016/j.neuroimage.2011.06.071

Bellebaum, C., Koch, B., Schwarz, M., Daum, I., 2008. Focal basal ganglia lesions are

associated with impairments in reward-based reversal learning. Brain 131, 829–841.

https://doi.org/10.1093/brain/awn011

Binder, J.R., Medler, D.A., Westbury, C.F., Liebenthal, E., Buchanan, L., 2006. Tuning of the

human left fusiform gyrus to sublexical orthographic structure. Neuroimage 33, 739–

748. https://doi.org/10.1016/j.neuroimage.2006.06.053

Bisio, A., Pedullà, L., Bonzano, L., Tacchino, A., Brichetto, G., Bove, M., 2017. The

kinematics of handwriting movements as expression of cognitive and sensorimotor

impairments in people with multiple sclerosis. Sci. Rep. 7, 1–10.

https://doi.org/10.1038/s41598-017-18066-7

Björklund, A., Dunnett, S.B., 2007. Dopamine neuron systems in the brain: an update.

Trends Neurosci. https://doi.org/10.1016/j.tins.2007.03.006

Bookheimer, S.Y., Zeffiro, T.A., Blaxton, T., Gaillard, W., Theodore, W., 1995. Regional

cerebral blood flow during object naming and word reading. Hum. Brain Mapp. 3, 93–

106. https://doi.org/10.1002/hbm.460030206

Bromberg-Martin, E.S., Matsumoto, M., Hikosaka, O., 2010. Dopamine in Motivational

Control: Rewarding, Aversive, and Alerting. Neuron.

https://doi.org/10.1016/j.neuron.2010.11.022

Brovelli, A., Nazarian, B., Meunier, M., Boussaoud, D., 2011a. Differential roles of caudate

nucleus and putamen during instrumental learning. Neuroimage 57, 1580–90.

https://doi.org/10.1016/j.neuroimage.2011.05.059

Brovelli, A., Nazarian, B., Meunier, M., Boussaoud, D., 2011b. Differential roles of caudate

nucleus and putamen during instrumental learning. Neuroimage 57, 1580–1590.

https://doi.org/10.1016/j.neuroimage.2011.05.059

Buchsbaum, B.R., Olsen, R.K., Koch, P.F., Kohn, P., Kippenhan, J.S., Berman, K.F., 2005.

Reading, hearing, and the planum temporale. Neuroimage 24, 444–454.

https://doi.org/10.1016/j.neuroimage.2004.08.025

Cacioppo, S., Fontang, F., Patel, N., Decety, J., Monteleone, G., Cacioppo, J.T., 2014.

Intention understanding over T: A neuroimaging study on shared representations and

tennis return predictions. Front. Hum. Neurosci. 8, 781.

https://doi.org/10.3389/fnhum.2014.00781

Callan, A.M., Osu, R., Yamagishi, Y., Callan, D.E., Inoue, N., 2009. Neural correlates of

resolving uncertainty in driver's decision making. Hum. Brain Mapp. 30, 2804–2812.

https://doi.org/10.1002/hbm.20710

Carreiras, M., Mechelli, A., Estévez, A., Price, C.J., 2007. Brain activation for lexical decision

and reading aloud: Two sides of the same coin? J. Cogn. Neurosci. 19, 433–444.

https://doi.org/10.1162/jocn.2007.19.3.433

Ceceli, A.O., Myers, C.E., Tricomi, E., 2020. Demonstrating and disrupting well-learned

habits. PLoS One 15, 1–28. https://doi.org/10.1371/journal.pone.0234424

Cellini, N., 2017. Memory consolidation in sleep disorders. Sleep Med. Rev. 35, 101–112.

https://doi.org/10.1016/j.smrv.2016.09.003

Celone, K.A., Thompson-Brenner, H., Ross, R.S., Pratt, E.M., Stern, C.E., 2011. An fMRI

investigation of the fronto-striatal learning system in women who exhibit eating disorder

behaviors. Neuroimage 56, 1749–1757.

https://doi.org/10.1016/J.NEUROIMAGE.2011.03.026

Cheema, K., Lantz, N., Cummine, J., 2018. Exploring the role of subcortical structures in

developmental reading impairments: Evidence for subgroups differentiated by caudate

activity. Neuroreport 29, 271–279. https://doi.org/10.1097/WNR.0000000000000938

Chein, J.M., Albert, D., O'Brien, L., Uckert, K., Steinberg, L., 2011. Peers increase

adolescent risk taking by enhancing activity in the brain's reward circuitry. Dev. Sci. 14.

https://doi.org/10.1111/j.1467-7687.2010.01035.x

Chen, L.L., Wise, S.P., 1995. Neuronal activity in the supplementary eye field during

acquisition of conditional oculomotor associations. J. Neurophysiol. 73, 1101–1121.

https://doi.org/10.1152/jn.1995.73.3.1101

Choi, M.H., Kim, H.S., Yoon, H.J., Lee, J.C., Baek, J.H., Choi, J.S., Tack, G.R., Min, B.C.,

Lim, D.W., Chung, S.C., 2017. Increase in brain activation due to subtasks during

driving: FMRI study using new MR-compatible driving simulator. J. Physiol. Anthropol.

36, 1–12. https://doi.org/10.1186/s40101-017-0128-8

Christensen, M.S., Lundbye-Jensen, J., Petersen, N., Geertsen, S.S., Paulson, O.B.,

Nielsen, J.B., 2007. Watching your foot move - An fMRI study of visuomotor

interactions during foot movement. Cereb. Cortex 17, 1906–1917.

https://doi.org/10.1093/cercor/bhl101

Chung, S.C., Choi, M.H., Kim, H.S., You, N.R., Hong, S.P., Lee, J.C., Park, S.J., Baek, J.H.,

Jeong, U.H., You, J.H., Lim, D.W., Kim, H.J., 2014. Effects of distraction task on

driving: A functional magnetic resonance imaging study, in: Bio-Medical Materials and

Engineering. IOS Press, pp. 2971–2977. https://doi.org/10.3233/BME-141117

Church, J.A., Coalson, R.S., Lugar, H.M., Petersen, S.E., Schlaggar, B.L., 2008. A

developmental fMRI study of reading and repetition reveals changes in phonological

and visual mechanisms over age. Cereb. Cortex 18, 2054–2065.

https://doi.org/10.1093/cercor/bhm228

Ciccarelli, O., Toosy, A.T., Marsden, J.F., Wheeler-Kingshott, C.M., Sahyoun, C., Matthews,

P.M., Miller, D.H., Thompson, A.J., 2005. Identifying brain regions for integrative

sensorimotor processing with ankle movements. Exp. Brain Res. 166, 31–42.

https://doi.org/10.1007/s00221-005-2335-5

Cohen, Y., Schneidman, E., Paz, R., 2021. The geometry of neuronal representations during

rule learning reveals complementary roles of cingulate cortex and putamen. Neuron

109, 839-851.e9. https://doi.org/10.1016/j.neuron.2020.12.027

Cromwell, H.C., Schultz, W., 2003. Effects of expectations for different reward magnitudes

on neuronal activity in primate striatum. J. Neurophysiol. 89, 2823–2838.

https://doi.org/10.1152/jn.01014.2002

Cummine, J., Cribben, I., Luu, C., Kim, E., Bakhtiari, R., Georgiou, G., Boliek, C.A., 2016.

Understanding the Role of Speech Production in Reading: Evidence for a Print-to-

Speech Neural Network Using Graphical Analysis. Neuropsychology.

https://doi.org/10.1037/neu0000236

Cunnington, R., Windischberger, C., Deecke, L., Moser, E., 2002. The preparation and

execution of self-initiated and externally-triggered movement: A study of event-related

fMRI. Neuroimage 15, 373–385. https://doi.org/10.1006/nimg.2001.0976

Daw, N.D., Gershman, S.J., Seymour, B., Dayan, P., Dolan, R.J., 2011. Model-based

influences on humans' choices and striatal prediction errors. Neuron 69, 1204–1215.

https://doi.org/10.1016/j.neuron.2011.02.027

Daw, N.D., Niv, Y., Dayan, P., 2005. Uncertainty-based competition between prefrontal and

dorsolateral striatal systems for behavioral control. Nat. Neurosci. 8, 1704–1711.

https://doi.org/10.1038/nn1560

De Houwer, J., 2019. On How Definitions of Habits Can Complicate Habit Research. Front.

Psychol. 10, 1–9. https://doi.org/10.3389/fpsyg.2019.02642

de Wit, S., Barker, R. a, Dickinson, A.D., Cools, R., 2010. Habitual versus goal-directed

action control in Parkinson disease. J Cogn Neurosci 23, 1218–1229.

https://doi.org/10.1162/jocn.2010.21514 [doi]

de Wit, S., Kindt, M., Knot, S.L., Verhoeven, A.A.C., Robbins, T.W., Gasull-Camos, J.,

Evans, M., Mirza, H., Gillan, C.M., 2018. Shifting the balance between goals and habits:

Five failures in experimental habit induction. J. Exp. Psychol. Gen. 147, 1043–1065.

https://doi.org/10.1037/xge0000402

de Wit, S., Watson, P., Harsay, H.A., Cohen, M.X., van de Vijver, I., Ridderinkhof, K.R.,

2012a. Corticostriatal connectivity underlies individual differences in the balance

between habitual and goal-directed action control. J. Neurosci. 32, 12066–12075.

https://doi.org/10.1523/JNEUROSCI.1088-12.2012

de Wit, S., Watson, P., Harsay, H.A., Cohen, M.X., van de Vijver, I., Ridderinkhof, K.R.,

2012b. Corticostriatal connectivity underlies individual differences in the balance

between habitual and goal-directed action control. J. Neurosci. 32, 12066–12075.

https://doi.org/10.1523/JNEUROSCI.1088-12.2012

Delgado, M.R., Miller, M.M., Inati, S., Phelps, E.A., 2005. An fMRI study of reward-related

probability learning. Neuroimage 24, 862–873.

https://doi.org/10.1016/j.neuroimage.2004.10.002

Deserno, L., Huys, Q.J.M., Boehme, R., Buchert, R., Heinze, H.J., Grace, A.A., Dolan, R.J.,

Heinz, A., Schlagenhauf, F., 2015. Ventral striatal dopamine reflects behavioral and

neural signatures of model-based control during sequential decision making. Proc. Natl.

Acad. Sci. U. S. A. 112, 1595–1600. https://doi.org/10.1073/pnas.1417219112

Dickinson, A., Trans, P., Lond, R.S., 1985. Actions and habits: the development of

behavioural autonomy. Philos. Trans. R. Soc. London. B, Biol. Sci. 308, 67–78.

https://doi.org/10.1098/rstb.1985.0010

Doll, B.B., Duncan, K.D., Simon, D.A., Shohamy, D., Daw, N.D., 2015. Model-based choices

involve prospective neural activity. Nat. Neurosci. 18, 767–772.

https://doi.org/10.1038/NN.3981

Eickhoff, S.B., Laird, A.R., Grefkes, C., Wang, L.E., Zilles, K., Fox, P.T., 2009. Coordinate-

based activation likelihood estimation meta-analysis of neuroimaging data: A random-

effects approach based on empirical estimates of spatial uncertainty. Hum. Brain Mapp.

30, 2907–2926. https://doi.org/10.1002/hbm.20718

Eickhoff, S.B., Nichols, T.E., Laird, A.R., Hoffstaedter, F., Amunts, K., Fox, P.T., Bzdok, D.,

Eickhoff, C.R., 2016a. Behavior, sensitivity, and power of activation likelihood

estimation characterized by massive empirical simulation. Neuroimage 137, 70–85.

https://doi.org/10.1016/j.neuroimage.2016.04.072

Eickhoff, S.B., Nichols, T.E., Laird, A.R., Hoffstaedter, F., Amunts, K., Fox, P.T., Bzdok, D.,

Eickhoff, C.R., 2016b. Behavior, sensitivity, and power of activation likelihood

estimation characterized by massive empirical simulation. Neuroimage 137, 70–85.

https://doi.org/10.1016/J.NEUROIMAGE.2016.04.072

Erdeniz, B., Done, J., 2019. Common and distinct functional brain networks for intuitive and

deliberate decision making. Brain Sci. 9. https://doi.org/10.3390/brainsci9070174

Erhard, K., Kessler, F., Neumann, N., Ortheil, H.J., Lotze, M., 2014. Professional training in

creative writing is associated with enhanced fronto-striatal activity in a literary text

continuation task. Neuroimage 100, 15–23.

https://doi.org/10.1016/j.neuroimage.2014.05.076

Eryilmaz, H., Rodriguez-Thompson, A., Tanner, A.S., Giegold, M., Huntington, F.C.,

Roffman, J.L., 2017. Neural determinants of human goal-directed vs. habitual action

control and their relation to trait motivation. Sci. Rep. 7, 1–11.

https://doi.org/10.1038/s41598-017-06284-y

Etchamendy, N., Konishi, K., Pike, G.B., Marighetto, A., Bohbot, V.D., 2012. Evidence for a

virtual human analog of a rodent relational memory task: A study of aging and fMRI in

young adults. Hippocampus 22, 869–880. https://doi.org/10.1002/HIPO.20948

Everitt, B.J., Belin, D., Economidou, D., Pelloux, Y., Dalley, J.W., Robbins, T.W., 2008.

Neural mechanisms underlying the vulnerability to develop compulsive drug-seeking

habits and addiction. Philos. Trans. R. Soc. B Biol. Sci. 363, 3125–3135.

https://doi.org/10.1098/rstb.2008.0089

Fera, F., Weickert, T.W., Goldberg, T.E., Tessitore, A., Hariri, A., Das, S., Lee, S., Zoltick,

B., Meeter, M., Myers, C.E., Gluck, M.A., Weinberger, D.R., Mattay, V.S., 2005. Neural

mechanisms underlying probabilistic category learning in normal aging. J. Neurosci. 25,

11340–11348. https://doi.org/10.1523/JNEUROSCI.2736-05.2005

Fermin, A.S.R., Yoshida, T., Yoshimoto, J., Ito, M., Tanaka, S.C., Doya, K., 2016. Model-

based action planning involves cortico-cerebellar and basal ganglia networks. Sci. Rep.

6. https://doi.org/10.1038/SREP31378

Fernández-Seara, M.A., Aznárez-Sanado, M., Mengual, E., Loayza, F.R., Pastor, M.A.,

2009. Continuous performance of a novel motor sequence leads to highly correlated

striatal and hippocampal perfusion increases. Neuroimage 47, 1797–1808.

https://doi.org/10.1016/J.NEUROIMAGE.2009.05.061

Foerde, K., 2018. What are habits and do they depend on the striatum? A view from the

study of neuropsychological populations. Curr. Opin. Behav. Sci. 20, 17–24.

https://doi.org/10.1016/j.cobeha.2017.08.011

Foerde, K., Knowlton, B.J., Poldrack, R.A., 2006. Modulation of competing memory systems

by distraction. Proc. Natl. Acad. Sci. U. S. A. 103, 11778–11783.

https://doi.org/10.1073/PNAS.0602659103

Forstmann, B.U., Dutilh, G., Brown, S., Neumann, J., Von Cramon, D.Y., Ridderinkhof, K.R.,

Wagenmakers, E.J., 2008. Striatum and pre-SMA facilitate decision-making under time

pressure. Proc. Natl. Acad. Sci. U. S. A. 105, 17538–17542.

https://doi.org/10.1073/pnas.0805903105

Fox, P.T., Laird, A.R., Eickhoff, S.B., Lancaster, J.L., Fox, M., Uecker, A.M., Robertson, M.,

Ray, K.L., 2013. User manual for GingerALE 2.3, Retrieved from

http://brainmap.org/ale/manual.pdf.

Gardiner, T.W., Nelson, R.J., 1992. Striatal neuronal activity during the initiation and

execution of hand movements made in response to visual and vibratory cues. Exp.

Brain Res. 92, 15–26. https://doi.org/10.1007/BF00230379

Gardner, B., 2015. A review and analysis of the use of "habit" in understanding, predicting

and influencing health-related behaviour. Health Psychol. Rev. 9, 277–295.

https://doi.org/10.1080/17437199.2013.876238

Gillan, C.M., 2021. Recent Developments in the Habit Hypothesis of OCD and Compulsive

Disorders. Springer, Berlin, Heidelberg, pp. 1–21.

https://doi.org/10.1007/7854_2020_199

Gläscher, J., Daw, N., Dayan, P., O'Doherty, J.P., 2010. States versus rewards: Dissociable

neural prediction error signals underlying model-based and model-free reinforcement

learning. Neuron 66, 585–595. https://doi.org/10.1016/j.neuron.2010.04.016

Gottfried, J.A., O'Doherty, J., Dolan, R.J., 2003. Encoding predictive reward value in human

amygdala and orbitofrontal cortex. Science (80-.). 301, 1104–1107.

https://doi.org/10.1126/SCIENCE.1087919

Gould, L., Mickleborough, M.J.S., Lorentz, E., Ekstrand, C., Borowsky, R., 2018. A

behavioral and fMRI examination of the effect of rhythm on reading noun-verb

homographs aloud. https://doi.org/10.1080/23273798.2018.1442012 33, 829–849.

https://doi.org/10.1080/23273798.2018.1442012

Grabenhorst, F., Rolls, E.T., 2011. Value, pleasure and choice in the ventral prefrontal

cortex. Trends Cogn. Sci. 15, 56–67. https://doi.org/10.1016/j.tics.2010.12.004

Graydon, F.X., Young, R., Benton, M.D., Genik, R.J., Posse, S., Hsieh, L., Green, C., 2004.

Visual event detection during simulated driving: Identifying the neural correlates with

functional neuroimaging. Transp. Res. Part F Traffic Psychol. Behav. 7, 271–286.

https://doi.org/10.1016/j.trf.2004.09.006

Haber, S.N., Fudge, J.L., McFarland, N.R., 2000. Striatonigrostriatal Pathways in Primates

Form an Ascending Spiral from the Shell to the Dorsolateral Striatum. J. Neurosci. 20,

2369–2382. https://doi.org/10.1523/JNEUROSCI.20-06-02369.2000

Haber, S.N., Knutson, B., 2009. The Reward Circuit: Linking Primate Anatomy and Human

Imaging. Neuropsychopharmacology 35, 4–26. https://doi.org/10.1038/npp.2009.129

Haith, A.M., Krakauer, J.W., 2018. The multiple effects of practice: skill, habit and reduced

cognitive load. Curr. Opin. Behav. Sci. 20, 196–201.

https://doi.org/10.1016/j.cobeha.2018.01.015

Hardwick, R.M., Forrence, A.D., Krakauer, J.W., Haith, A.M., 2019. Time-dependent

competition between goal-directed and habitual response preparation. Nat. Hum.

Behav. 3, 1252–1262. https://doi.org/10.1038/s41562-019-0725-0

Haruno, M., Kawato, M., 2006. Different neural correlates of reward expectation and reward

expectation error in the putamen and caudate nucleus during stimulus-action-reward

association learning. J. Neurophysiol. 95, 948–959.

https://doi.org/10.1152/jn.00382.2005

Haynes, W.I.A., Haber, S.N., 2013. The organization of prefrontal-subthalamic inputs in

primates provides an anatomical substrate for both functional specificity and integration:

Implications for basal ganglia models and deep brain stimulation. J. Neurosci. 33,

4804–4814. https://doi.org/10.1523/JNEUROSCI.4674-12.2013

Hernandez, L.F., Obeso, I., Costa, R.M., Redgrave, P., Obeso, J.A., 2019. Dopaminergic

Vulnerability in Parkinson Disease: The Cost of Humans' Habitual Performance. Trends

Neurosci. 42, 375–383. https://doi.org/10.1016/j.tins.2019.03.007

Hikosaka, O., Takikawa, Y., Kawagoe, R., Kojima, S., Doupe, A.J., Silver, R., Boahen, K.,

Grillner, S., Kopell, N., Olsen, K.L., Neurosci F-W Zhou, J., Matta, S.G., Zhou, F.,

Neurophysiol, J., Bargas Carrillo-Reid, J.L., Tecuapetla, F., Tapia, D., Hernandez-Cruz,

A., Galarraga, E., Drucker-Colin, R., 2000. Tutor Song Exposure Song Selectivity in the

Pallial-Basal Ganglia Song Circuit of Zebra Finches Raised Without Role of the Basal

Ganglia in the Control of Purposive Saccadic Eye Movements. Physiol Rev 80, 953–

978.

Holl, A.K., Wilkinson, L., Tabrizi, S.J., Painold, A., Jahanshahi, M., 2012. Probabilistic

classification learning with corrective feedback is selectively impaired in early

Huntington's disease--evidence for the role of the striatum in learning with feedback.

Neuropsychologia 50, 2176–2186.

https://doi.org/10.1016/J.NEUROPSYCHOLOGIA.2012.05.021

Horovitz, S.G., Gallea, C., Najee-ullah, M.A., Hallett, M., 2013. Functional Anatomy of

Writing with the Dominant Hand. PLoS One 8, 1–10.

https://doi.org/10.1371/journal.pone.0067931

Hsieh, L., Young, R.A., Bowyer, S.M., Moran, J.E., Genik, R.J., Green, C.C., Chiang, Y.R.,

Yu, Y.J., Liao, C.C., Seaman, S., 2009. Conversation effects on neural mechanisms

underlying reaction time to visual events while viewing a driving scene: fMRI analysis

and asynchrony model. Brain Res. https://doi.org/10.1016/j.brainres.2008.10.002

Hsu, C.T., Jacobs, A.M., Citron, F.M.M., Conrad, M., 2015. The emotion potential of words

and passages in reading Harry Potter - An fMRI study. Brain Lang. 142, 96–114.

https://doi.org/10.1016/j.bandl.2015.01.011

Hu, X.Y., Wang, L., Liu, H., Zhang, S.Z., 2006. Functional magnetic resonance imaging

study of writer's cramp. Chin. Med. J. (Engl). 119, 1263–1271.

https://doi.org/10.1097/00029330-200608010-00006

Huang, Y., Yaple, Z.A., Yu, R., 2020. Goal-oriented and habitual decisions: Neural

signatures of model-based and model-free learning. Neuroimage 215, 116834.

https://doi.org/10.1016/j.neuroimage.2020.116834

Huth, A.G., De Heer, W.A., Griffiths, T.L., Theunissen, F.E., Gallant, J.L., 2016. Natural

speech reveals the semantic maps that tile human cerebral cortex. Nature 532, 453–

458. https://doi.org/10.1038/nature17637

Iaria, G., Petrides, M., Dagher, A., Pike, B., Bohbot, V.D., 2003. Cognitive strategies

dependent on the hippocampus and caudate nucleus in human navigation: variability

and change with practice. J. Neurosci. Off. J. Soc. Neurosci. 23, 5945–5952.

https://doi.org/10.1523/jneurosci.23-13-05945.2003

Jaeger L, Marchal-Crespo L, Wolf P, Riener R, Michels L, K.S., 2014. Brain activation

associated with active and passive lower limb stepping. Front. Hum. Neurosci. 8.

https://doi.org/10.3389/fnhum.2014.00828

Jankowski, J., Scheef, L., Hüppe, C., Boecker, H., 2009. Distinct striatal regions for planning

and executing novel and automated movement sequences. Neuroimage 44, 1369–

1379. https://doi.org/10.1016/j.neuroimage.2008.10.059

Jeanvoine, H., Labriffe, M., Tannou, T., Navasiolava, N., Ter Minassian, A., Girot, J.B.,

Leiber, L.M., Custaud, M.A., Annweiler, C., Dinomais, M., 2022. Specific age-correlated

activation of top hierarchical motor control areas during gait-like plantar stimulation: An

fMRI study. Hum. Brain Mapp. 43, 833–843. https://doi.org/10.1002/HBM.25691

Karimpoor, M., Churchill, N.W., Tam, F., Fischer, C.E., Schweizer, T.A., Graham, S.J., 2018.

Functional MRI of handwriting tasks: A study of healthy young adults interacting with a

novel touch-sensitive tablet. Front. Hum. Neurosci. 12, 1–14.

https://doi.org/10.3389/fnhum.2018.00030

Karimpoor, M., Tam, F., Strother, S.C., Fischer, C.E., Schweizer, T.A., Graham, S.J., 2015.

A computerized tablet with visual feedback of hand position for functional magnetic

resonance imaging. Front. Hum. Neurosci. 9, 150.

https://doi.org/10.3389/fnhum.2015.00150

Katanoda, K., Yoshikawa, K., Sugishita, M., 2001. A functional MRI study on the neural

substrates for writing. Hum. Brain Mapp. 13, 34–42. https://doi.org/10.1002/hbm.1023

Kilner, J.M., Neal, A., Weiskopf, N., Friston, K.J., Frith, C.D., 2009. Evidence of Mirror

Neurons in Human Inferior Frontal Gyrus. J. Neurosci. 29, 10153.

https://doi.org/10.1523/JNEUROSCI.2668-09.2009

Kim, W., Chang, Y., Kim, J., Seo, J., Ryu, K., Lee, E., Woo, M., Janelle, C.M., 2014. An fMRI

study of differences in brain activity among elite, expert, and novice archers at the

moment of optimal aiming. Cogn. Behav. Neurol. 27, 173–182.

https://doi.org/10.1097/WNN.0000000000000042

Kimura, M., 1986. The role of primate putamen neurons in the association of sensory stimuli

with movement. Neurosci. Res. 3, 436–443. https://doi.org/10.1016/0168-

0102(86)90035-0

Kimura, M., Kato, M., Shimazaki, H., 1990. Physiological properties of projection neurons in

the monkey striatum to the globus pallidus. Exp. Brain Res. 82, 672–676.

https://doi.org/10.1007/BF00228811

Kimura, M., Rajkowski, J., Evarts, E., 1984. Tonically discharging putamen neurons exhibit

set-dependent responses. Proc. Natl. Acad. Sci. U. S. A. 81, 4998–5001.

https://doi.org/10.1073/pnas.81.15.4998

Kunimatsu, J., Maeda, K., Hikosaka, O., 2019. The caudal part of putamen represents the

historical object value information. J. Neurosci. 39, 1709–1719.

https://doi.org/10.1523/JNEUROSCI.2534-18.2018

la Fougère, C., Zwergal, A., Rominger, A., Förster, S., Fesl, G., Dieterich, M., Brandt, T.,

Strupp, M., Bartenstein, P., Jahn, K., 2010. Real versus imagined locomotion: A [18F]-

FDG PET-fMRI comparison. Neuroimage 50, 1589–1598.

https://doi.org/10.1016/j.neuroimage.2009.12.060

Lancaster, J.L., Tordesillas-Gutiérrez, D., Martinez, M., Salinas, F., Evans, A., Zilles, K.,

Mazziotta, J.C., Fox, P.T., 2007. Bias between MNI and talairach coordinates analyzed

using the ICBM-152 brain template. Hum. Brain Mapp. 28, 1194–1205.

https://doi.org/10.1002/hbm.20345

Leh, S.E., Ptito, A., Chakravarty, M.M., Strafella, A.P., 2007. Fronto-striatal connections in
the human brain: A probabilistic diffusion tractography study. Neurosci. Lett. 419, 113–
118. https://doi.org/10.1016/j.neulet.2007.04.049

Lehéricy, S., Benali, H., Van De Moortele, P.F., Pélégrini-Issac, M., Waechter, T., Ugurbil,
K., Doyon, J., 2005a. Distinct basal ganglia territories are engaged in early and
advanced motor sequence learning. Proc. Natl. Acad. Sci. U. S. A. 102, 12566–12571.
https://doi.org/10.1073/pnas.0502762102

Lehéricy, S., Benali, H., Van De Moortele, P.F., Pélégrini-Issac, M., Waechter, T., Ugurbil,
K., Doyon, J., 2005b. Distinct basal ganglia territories are engaged in early and
advanced motor sequence learning. Proc. Natl. Acad. Sci. U. S. A. 102, 12566–12571.
https://doi.org/10.1073/pnas.0502762102

Leisman, G., Moustafa, A., Shafir, T., 2016. Thinking, Walking, Talking: Integratory Motor
and Cognitive Brain Function. Front. Public Heal. 4, 1.
https://doi.org/10.3389/fpubh.2016.00094

Liljeholm, M., Dunne, S., O'Doherty, J.P., 2015a. Differentiating neural systems mediating
the acquisition vs. expression of goal-directed and habitual behavioral control. Eur. J.
Neurosci. 41, 1358–1371. https://doi.org/10.1111/ejn.12897

Liljeholm, M., Dunne, S., O'Doherty, J.P., 2015b. Differentiating neural systems mediating
the acquisition vs. expression of goal-directed and habitual behavioral control. Eur. J.
Neurosci. 41, 1358–1371. https://doi.org/10.1111/ejn.12897

Liljeholm, M., O'doherty, J.P., 2012. Contributions of the striatum to learning, motivation, and
performance: an associative account. Trends Cogn Sci 16, 467–475.
https://doi.org/10.1016/j.tics.2012.07.007

Lin, Z., Tam, F., Churchill, N.W., Schweizer, T.A., Graham, S.J., 2021. Tablet technology for
writing and drawing during functional magnetic resonance imaging: A review. Sensors

(Switzerland). https://doi.org/10.3390/s21020401

Longcamp, M., Lagarrigue, A., Nazarian, B., Roth, M., Anton, J.L., Alario, F.X., Velay, J.L.,

2014. Functional specificity in the motor system: Evidence from coupled fMRI and

kinematic recordings during letter and digit writing. Hum. Brain Mapp. 35, 6077–6087.

https://doi.org/10.1002/hbm.22606

Luque, D., Beesley, T., Morris, R.W., Jack, B.N., Griffiths, O., Whitford, T.J., Le Pelley, M.E.,

2017. Goal-Directed and Habit-Like Modulations of Stimulus Processing during

Reinforcement Learning. https://doi.org/10.1523/JNEUROSCI.3205-16.2017

Luque, D., Molinero, S., Watson, P., López, F.J., Le Pelley, M.E., 2020. Measuring habit

formation through goal-directed response switching. J. Exp. Psychol. Gen. 149, 1449–

1459. https://doi.org/10.1037/xge0000722

Marchal, V., Sellers, J., Pélégrini-Issac, M., Galléa, C., Bertasi, E., Valabrègue, R., Lau, B.,

Leboucher, P., Bardinet, E., Welter, M.L., Karachi, C., 2019. Deep brain activation

patterns involved in virtual gait without and with a doorway: An fMRI study. PLoS One

14. https://doi.org/10.1371/journal.pone.0223494

Martínez, M., Valencia, M., Vidorreta, M., Luis, E.O., Castellanos, G., Villagra, F., Fern??

ndez-Seara, M.A., Pastor, M.A., 2016a. Trade-off between frequency and precision

during stepping movements: Kinematic and BOLD brain activation patterns. Hum. Brain

Mapp. 37, 1722–1737. https://doi.org/10.1002/hbm.23131

Martínez, M., Valencia, M., Vidorreta, M., Luis, E.O., Castellanos, G., Villagra, F.,

Fernández-Seara, M.A., Pastor, M.A., 2016b. Trade-off between frequency and

precision during stepping movements: Kinematic and BOLD brain activation patterns.

Hum. Brain Mapp. 37, 1722–1737. https://doi.org/10.1002/hbm.23131

Martínez, M., Villagra, F., Castellote, J.M., Pastor, M.A., 2018. Kinematic and kinetic

patterns related to free-walking in parkinson's disease. Sensors (Switzerland) 18.

https://doi.org/10.3390/s18124224

Martinez, M., Villagra, F., Loayza, F., Vidorreta, M., Arrondo, G., Luis, E., Diaz, J.,

Echeverria, M., Fernandez-Seara, M.A., Pastor, M.A., 2014. MRI-compatible device for

examining brain activation related to stepping. IEEE Trans. Med. Imaging 33, 1044–

1053. https://doi.org/10.1109/TMI.2014.2301493

Mas-Herrero, E., Sescousse, G., Cools, R., Marco-Pallarés, J., 2019. The contribution of

striatal pseudo-reward prediction errors to value-based decision-making. Neuroimage

193, 67–74. https://doi.org/10.1016/j.neuroimage.2019.02.052

Matsuda, W., Furuta, T., Nakamura, K.C., Hioki, H., Fujiyama, F., Arai, R., Kaneko, T., 2009.

Single nigrostriatal dopaminergic neurons form widely spread and highly dense axonal

arborizations in the neostriatum. J. Neurosci. 29, 444–453.

https://doi.org/10.1523/JNEUROSCI.4029-08.2009

Mazziotta, J., Toga, A., Evans, A., Fox, P., Lancaster, J., Zilles, K., Woods, R., Paus, T.,

Simpson, G., Pike, B., Holmes, C., Collins, L., Thompson, P., MacDonald, D., Iacoboni,

M., Schormann, T., Amunts, K., Palomero-Gallagher, N., Geyer, S., Parsons, L., Narr,

K., Kabani, N., Le Goualher, G., Boomsma, D., Cannon, T., Kawashima, R., Mazoyer,

B., 2001. A probabilistic atlas and reference system for the human brain: International

Consortium for Brain Mapping (ICBM). Philos. Trans. R. Soc. B Biol. Sci.

https://doi.org/10.1098/rstb.2001.0915

McFarland, N.R., Haber, S.N., 2002. Thalamic relay nuclei of the basal ganglia form both

reciprocal and nonreciprocal cortical connections, linking multiple frontal cortical areas.

J. Neurosci. 22, 8117–8132. https://doi.org/10.1523/jneurosci.22-18-08117.2002

Mechelli, A., Gorno-Tempini, M.L., Price, C.J., 2003. Neuroimaging studies of word and

pseudoword reading: Consistencies, inconsistencies, and limitations. J. Cogn.

Neurosci. 15, 260–271. https://doi.org/10.1162/089892903321208196

Meschyan, G., Hernandez, A.E., 2006. Impact of language proficiency and orthographic

transparency on bilingual word reading: An fMRI investigation. Neuroimage 29, 1135–

1140. https://doi.org/10.1016/j.neuroimage.2005.08.055

Mi, T.M., Zhang, W., McKeown, M.J., Chan, P., 2021. Impaired Formation and Expression of

Goal-Directed and Habitual Control in Parkinson's Disease. Front. Aging Neurosci. 13.

https://doi.org/10.3389/FNAGI.2021.734807

Miller, K.J., Shenhav, A., Ludvig, E.A., 2019. Habits without values. Psychol. Rev.

https://doi.org/10.1037/rev0000120

Molenberghs, P., Cunnington, R., Mattingley, J.B., 2012. Brain regions with mirror

properties: A meta-analysis of 125 human fMRI studies. Neurosci. Biobehav. Rev. 36,

341–349. https://doi.org/10.1016/J.NEUBIOREV.2011.07.004

Moore, C.J., Price, C.J., 1999. Three distinct ventral occipitotemporal regions for reading

and object naming. Neuroimage 10, 181–192. https://doi.org/10.1006/nimg.1999.0450

Morris, L.S., Kundu, P., Dowell, N., Mechelmans, D.J., Favre, P., Irvine, M.A., Robbins,

T.W., Daw, N., Bullmore, E.T., Harrison, N.A., Voon, V., 2016. Fronto-striatal

organization: Defining functional and microstructural substrates of behavioural flexibility.

Cortex 74, 118–133. https://doi.org/10.1016/j.cortex.2015.11.004

Morris, N.J., Stein, C.M., 2017. Model-free linkage analysis of a quantitative trait, in:

Methods in Molecular Biology. Humana Press Inc., pp. 327–342.

https://doi.org/10.1007/978-1-4939-7274-6_16

Müller, V.I., Cieslik, E.C., Laird, A.R., Fox, P.T., Radua, J., Mataix-Cols, D., Tench, C.R.,

Yarkoni, T., Nichols, T.E., Turkeltaub, P.E., Wager, T.D., Eickhoff, S.B., 2018a. Ten

simple rules for neuroimaging meta-analysis. Neurosci. Biobehav. Rev. 84, 151.

https://doi.org/10.1016/J.NEUBIOREV.2017.11.012

Müller, V.I., Cieslik, E.C., Laird, A.R., Fox, P.T., Radua, J., Mataix-Cols, D., Tench, C.R.,

Yarkoni, T., Nichols, T.E., Turkeltaub, P.E., Wager, T.D., Eickhoff, S.B., 2018b. Ten

simple rules for neuroimaging meta-analysis. Neurosci. Biobehav. Rev.

https://doi.org/10.1016/j.neubiorev.2017.11.012

Murphy, K., Garavan, H., 2004. An empirical investigation into the number of subjects

required for an event-related fMRI study. Neuroimage 22, 879–885.

https://doi.org/10.1016/J.NEUROIMAGE.2004.02.005

Nachev, P., Kennard, C., Husain, M., 2008. Functional role of the supplementary and pre-

supplementary motor areas. Nat. Rev. Neurosci. https://doi.org/10.1038/nrn2478

Nakamura, K., Honda, M., Hirano, S., Oga, T., Sawamoto, N., Hanakawa, T., Inoue, H., Ito,

J., Matsuda, T., Fukuyama, H., Shibasaki, H., 2002. Modulation of the visual word

retrieval system in writing: A functional MRI study on the Japanese orthographies. J.

Cogn. Neurosci. 14, 104–115. https://doi.org/10.1162/089892902317205366

Nebe, S., Kroemer, N.B., Schad, D.J., Bernhardt, N., Sebold, M., Müller, D.K., Scholl, L.,

Kuitunen-Paul, S., Heinz, A., Rapp, M.A., Huys, Q.J.M., Smolka, M.N., 2018. No

association of goal-directed and habitual control with alcohol consumption in young

adults. Addict. Biol. 23, 379–393. https://doi.org/10.1111/ADB.12490

Noble, J.W., Eng, J.J., Boyd, L.A., 2014. Bilateral motor tasks involve more brain regions

and higher neural activation than unilateral tasks: An fMRI study. Exp. Brain Res. 232,

2785–2795. https://doi.org/10.1007/s00221-014-3963-4

Oberhuber, M., Hope, T.M.H., Seghier, M.L., Parker Jones, O., Prejawa, S., Green, D.W.,

Price, C.J., 2016. Four Functionally Distinct Regions in the Left Supramarginal Gyrus

Support Word Processing. Cereb. Cortex 26, 4212–4226.

https://doi.org/10.1093/cercor/bhw251

Oberhuber, M., Jones, O.P., Hope, T.M.H., Prejawa, S., Seghier, M.L., Green, D.W., Price,

C.J., 2013. Functionally distinct contributions of the anterior and posterior putamen

during sublexical and lexical reading. Front. Hum. Neurosci.

https://doi.org/10.3389/fnhum.2013.00787

Oh-Descher, H., Beck, J.M., Ferrari, S., Sommer, M.A., Egner, T., 2017. Probabilistic

inference under time pressure leads to a cortical-to-subcortical shift in decision

evidence integration. Neuroimage 162, 138–150.

https://doi.org/10.1016/j.neuroimage.2017.08.069

Ohbayashi, M., Picard, N., Strick, P.L., 2016. Inactivation of the dorsal premotor area

disrupts internally generated, but not visually guided, sequential movements. J.

Neurosci. 36, 1971–1976. https://doi.org/10.1523/JNEUROSCI.2356-15.2016

Parkes, S.L., Bradfield, L.A., Balleine, B.W., 2015. Interaction of insular cortex and ventral

striatum mediates the effect of incentive memory on choice between goal-directed

actions. J. Neurosci. 35, 6464–6471. https://doi.org/10.1523/JNEUROSCI.4153-

14.2015

Patterson, T.K., Knowlton, B.J., 2018. Subregional specificity in human striatal habit

learning: a meta-analytic review of the fMRI literature. Curr. Opin. Behav. Sci. 20, 75–

82. https://doi.org/10.1016/j.cobeha.2017.10.005

Perez, Omar D, Dickinson, A., 2020. A theory of actions and habits: The interaction of rate

correlation and contiguity systems in free-operant behavior.

https://doi.org/10.1037/rev0000201

Perez, Omar D., Dickinson, A., 2020. A theory of actions and habits: The interaction of rate

correlation and contiguity systems in free-operant behavior. Psychol. Rev. 127, 945–

971. https://doi.org/10.1037/REV0000201

Peters, A.J., Fabre, J.M.J., Steinmetz, N.A., Harris, K.D., Carandini, M., 2021. Striatal

activity topographically reflects cortical activity. Nature 591, 420–425.

https://doi.org/10.1038/s41586-020-03166-8

Peters, S., Eng, J.J., Liu-Ambrose, T., Borich, M.R., Dao, E., Amanian, A., Boyd, L.A., 2019.

Brain activity associated with Dual-task performance of Ankle motor control during

cognitive challenge. Brain Behav. 9. https://doi.org/10.1002/brb3.1349

Peterson, B.S., Riddle, M.A., Cohen, D.J., Katz, L.D., Smith, J.C., Leckman, J.F., 1993.

Human basal ganglia volume asymmetries on magnetic resonance images. Magn.

Reson. Imaging 11, 493–498. https://doi.org/10.1016/0730-725X(93)90468-S

Peterson, D.S., Horak, F.B., 2016. Neural control of walking in people with parkinsonism.

Physiology. https://doi.org/10.1152/physiol.00034.2015

Pineda-Pardo, J.D.S.A., Sánchez-Ferro, Á., Monje, M.H.G., Pavese, N., Obeso, J.A., 2022.

Onset pattern of nigrostriatal denervation in early Parkinson's disease. Brain 145,

1018–1028. https://doi.org/10.1093/BRAIN/AWAB378

Poldrack, R.A., Clark, J., Paré-Blagoev, E.J., Shohamy, D., Creso Moyano, J., Myers, C.,

Gluck, M.A., 2001. Interactive memory systems in the human brain. Nature 414, 546–

550. https://doi.org/10.1038/35107080

Poldrack, R.A., Prabhakaran, V., Seger, C.A., Gabrieli, J.D.E., 1999. Striatal activation

during acquisition of a cognitive skill. Neuropsychology 13, 564–574.

https://doi.org/10.1037//0894-4105.13.4.564

Pool, E., Gera, R., Fransen, A., Perez, O.D., Cremer, A., Aleksic, M., Tanwisuth, S., Quail,

S., Ceceli, A.O., Manfredi, D., Nave, G., Tricomi, E., Balleine, B., Schonberg, T.,

Schwabe, L., O'Doherty, J.P., n.d. Determining the effects of training duration on the

behavioral expression of habitual control in humans: a multi-laboratory investigation.

https://doi.org/10.31234/OSF.IO/Z756H

Potgieser, A.R.E., Van Der Hoorn, A., De Jong, B.M., 2015. Cerebral Activations Related to

Writing and Drawing with each hand. PLoS One 10, 1–25.

https://doi.org/10.1371/journal.pone.0126723

Redgrave, P., Gurney, K., Reynolds, J., 2008. What is reinforced by phasic dopamine

signals? Brain Res. Rev. https://doi.org/10.1016/j.brainresrev.2007.10.007

Redgrave, P., Prescott, T.J., Gurney, K., 1999. The basal ganglia: a vertebrate solution to

the selection problem? Neuroscience 89, 1009–1023.

https://doi.org/S0306452298003194 [pii]

Redgrave, P., Rodriguez, M., Smith, Y., Rodriguez-Oroz, M.C., Lehericy, S., Bergman, H.,

Agid, Y., Delong, M.R., Obeso, J.A., 2010a. Goal-directed and habitual control in the

basal ganglia: Implications for Parkinson's disease. Nat. Rev. Neurosci. 11, 760–772.

https://doi.org/10.1038/nrn2915

Redgrave, P., Rodriguez, M., Smith, Y., Rodriguez-Oroz, M.C., Lehericy, S., Bergman, H.,

Agid, Y., Delong, M.R., Obeso, J.A., 2010b. Goal-directed and habitual coRedgrave, P.,

Rodriguez, M., Smith, Y., Rodriguez-Oroz, M. C., Lehericy, S., Bergman, H., Agid, Y.,

Delong, M. R., & Obeso, J. A. (2010). Goal-directed and habitual control in the basal

ganglia: Implications for Parkinson's di. Nat. Rev. Neurosci. 11, 760–772.

https://doi.org/10.1038/nrn2915

Romero, M.C., Bermudez, M.A., Vicente, A.F., Perez, R., Gonzalez, F., 2008. Activity of

neurons in the caudate and putamen during a visuomotor task. Neuroreport 19, 1141–

1145. https://doi.org/10.1097/WNR.0b013e328307c3fc

Rueckl, J.G., Paz-Alonso, P.M., Molfese, P.J., Kuo, W.J., Bick, A., Frost, S.J., Hancock, R.,

Wu, D.H., Einar Mencl, W., Duñabeitia, J.A., Lee, J.R., Oliver, M., Zevin, J.D., Hoeft, F.,

Carreiras, M., Tzeng, O.J.L., Pugh, K.R., Frost, R., 2015. Universal brain signature of

proficient reading: Evidence from four contrasting languages. Proc. Natl. Acad. Sci. U.

S. A. 112, 15510–15515. https://doi.org/10.1073/pnas.1509321112

Sauvage, C., Jissendi, P., Seignan, S., Manto, M., Habas, C., 2013. Brain areas involved in

the control of speed during a motor sequence of the foot: Real movement versus

mental imagery. J. Neuroradiol. 40, 267–280.

https://doi.org/10.1016/j.neurad.2012.10.001

Schultz, W., 2006. Behavioral theories and the neurophysiology of reward. Annu. Rev.

Psychol. 57, 87–115. https://doi.org/10.1146/annurev.psych.56.091103.070229

Schwabe, L., Tegenthoff, M., Höffken, O., Wolf, O.T., 2013. Mineralocorticoid Receptor

Blockade Prevents Stress-Induced Modulation of Multiple Memory Systems in the

Human Brain. Biol. Psychiatry 74, 801–808.

https://doi.org/10.1016/J.BIOPSYCH.2013.06.001

Segal, E., Petrides, M., 2012. The anterior superior parietal lobule and its interactions with

language and motor areas during writing. Eur. J. Neurosci. 35, 309–322.

https://doi.org/10.1111/j.1460-9568.2011.07937.x

Seghier, M.L., Lee, H.L., Schofield, T., Ellis, C.L., Price, C.J., 2008. Inter-subject variability in

the use of two different neuronal networks for reading aloud familiar words. Neuroimage

42, 1226–1236. https://doi.org/10.1016/j.neuroimage.2008.05.029

Seghier, M.L., Price, C.J., 2010. Reading Aloud Boosts Connectivity through the Putamen.

Cereb. Cortex 20, 570–582. https://doi.org/10.1093/cercor/bhp123

Sescousse, G., Caldú, X., Segura, B., Dreher, J.C., 2013. Processing of primary and

secondary rewards: A quantitative meta-analysis and review of human functional

neuroimaging studies. Neurosci. Biobehav. Rev. 37, 681–696.

https://doi.org/10.1016/j.neubiorev.2013.02.002

Sjoerds, Z., De Wit, S., Van Den Brink, W., Robbins, T.W., Beekman, A.T.F., Penninx,

B.W.J.H., Veltman, D.J., 2013. Behavioral and neuroimaging evidence for overreliance

on habit learning in alcohol-dependent patients. Transl. Psychiatry 3, e337-8.

https://doi.org/10.1038/tp.2013.107

Smith, K.S., Graybiel, A.M., 2016. Habit formation. Dialogues Clin. Neurosci. 18, 33–43.

https://doi.org/10.31887/dcns.2016.18.1/ksmith

Smittenaar, P., Guitart-Masip, M., Lutti, A., Dolan, R.J., 2013. Preparing for selective

inhibition within frontostriatal loops. J. Neurosci. 33, 18087–18097.

https://doi.org/10.1523/JNEUROSCI.2167-13.2013

Soares, J.M., Sampaio, A., Ferreira, L.M., Santos, N.C., Marques, F., Palha, J.A., Cerqueira,

J.J., Sousa, N., 2012. Stress-induced changes in human decision-making are

reversible. Transl. Psychiatry 2. https://doi.org/10.1038/TP.2012.59

Sommer, S., Pollmann, S., 2016. Putamen Activation Represents an Intrinsic Positive

Prediction Error Signal for Visual Search in Repeated Configurations. Open Neuroimag.

J. 10, 126–138. https://doi.org/10.2174/1874440001610010126

Spiers, H.J., Maguire, E.A., 2007. Neural substrates of driving behaviour. Neuroimage 36,

245–255. https://doi.org/10.1016/j.neuroimage.2007.02.032

Steele, C.J., Penhune, V.B., 2010. Specific Increases within Global Decreases: A Functional

Magnetic Resonance Imaging Investigation of Five Days of Motor Sequence Learning.

J. Neurosci. 30, 8332–8341. https://doi.org/10.1523/JNEUROSCI.5569-09.2010

Sun, R., 2004. Desiderata for cognitive architectures. Philos. Psychol. 17, 341–373.

https://doi.org/10.1080/0951508042000286721

Swett, B.A., Contreras-Vidal, J.L., Birn, R., Braun, A., 2010. Neural Substrates of

Graphomotor Sequence Learning: A Combined fMRI and Kinematic Study. J.

Neurophysiol. 103, 3366. https://doi.org/10.1152/JN.00449.2009

Swinnen, S.P., Vangheluwe, S., Wagemans, J., Coxon, J.P., Goble, D.J., Van Impe, A.,

Sunaert, S., Peeters, R., Wenderoth, N., 2010. Shared neural resources between left

and right interlimb coordination skills: The neural substrate of abstract motor

representations. Neuroimage 49, 2570–2580.

https://doi.org/10.1016/j.neuroimage.2009.10.052

Tanaka, S.C., Balleine, B.W., O'Doherty, J.P., 2008. Calculating consequences: Brain

systems that encode the causal effects of actions. J. Neurosci. 28, 6750–6755.

https://doi.org/10.1523/JNEUROSCI.1808-08.2008

Toyomura, A., Shibata, M., Kuriki, S., 2012. Self-paced and externally triggered rhythmical

lower limb movements: A functional MRI study. Neurosci. Lett. 516, 39–44.

https://doi.org/10.1016/j.neulet.2012.03.049

Tricomi, E., Balleine, B.W., O'Doherty, J.P., 2009. A specific role for posterior dorsolateral

striatum in human habit learning. Eur. J. Neurosci. 29, 2225–2232.

https://doi.org/10.1111/j.1460-9568.2009.06796.x

Trinastic, J.P., Kautz, S.A., McGregor, K., Gregory, C., Bowden, M., Benjamin, M.B.,

Kurtzman, M., Chang, Y.L., Conway, T., Crosson, B., 2010. An fMRI study of the

differences in brain activity during active ankle dorsiflexion and plantarflexion. Brain

Imaging Behav. 4, 121–131. https://doi.org/10.1007/s11682-010-9091-2

Tunik, E., Houk, J.C., Grafton, S.T., 2009. Basal ganglia contribution to the initiation of

corrective submovements. Neuroimage 47, 1757–1766.

https://doi.org/10.1016/j.neuroimage.2009.04.077

Turkeltaub, P.E., Eden, G.F., Jones, K.M., Zeffiro, T.A., 2002. Meta-analysis of the functional

neuroanatomy of single-word reading: Method and validation. Neuroimage 16, 765–

780. https://doi.org/10.1006/nimg.2002.1131

Uchiyama, Y., Ebe, K., Kozato, A., Okada, T., Sadato, N., 2003. The neural substrates of

driving at a safe distance: A functional MRI study. Neurosci. Lett. 352, 199–202.

https://doi.org/10.1016/j.neulet.2003.08.072

Uchiyama, Y., Toyoda, H., Sakai, H., Shin, D., Ebe, K., Sadato, N., 2012. Suppression of

brain activity related to a car-following task with an auditory task: An fMRI study.

Transp. Res. Part F Traffic Psychol. Behav. 15, 25–37.

https://doi.org/10.1016/j.trf.2011.11.002

Valentin, V. V., Dickinson, A., O'Doherty, J.P., 2007. Determining the neural substrates of

goal-directed learning in the human brain. J. Neurosci. 27, 4019–4026.

https://doi.org/10.1523/JNEUROSCI.0564-07.2007

van Steenbergen, H., Watson, P., Wiers, R.W., Hommel, B., de Wit, S., 2017. Dissociable

corticostriatal circuits underlie goal-directed vs. cue-elicited habitual food seeking after

satiation: evidence from a multimodal MRI study. Eur. J. Neurosci. 46, 1815–1827.

https://doi.org/10.1111/ejn.13586

Vannest, J., Eaton, K.P., Henkel, D., Siegel, M., Tsevat, R.K., Allendorfer, J.B., Schefft, B.K.,

Banks, C., Szaflarski, J.P., 2012. Cortical correlates of self-generation in verbal paired

associate learning. Brain Res. 1437, 104–114.

https://doi.org/10.1016/j.brainres.2011.12.020

Vaquero, L., Hartmann, K., Ripollés, P., Rojo, N., Sierpowska, J., François, C., Càmara, E.,

van Vugt, F.T., Mohammadi, B., Samii, A., Münte, T.F., Rodríguez-Fornells, A.,

Altenmüller, E., 2016. Structural neuroplasticity in expert pianists depends on the age of

musical training onset. Neuroimage 126, 106–119.

https://doi.org/10.1016/j.neuroimage.2015.11.008

Varotto, S.F., Farah, H., Bogenberger, K., van Arem, B., Hoogendoorn, S.P., 2020.

Adaptations in driver behaviour characteristics during control transitions from full-range

Adaptive Cruise Control to manual driving: an on-road study. Transp. A Transp. Sci. 16,

776–806. https://doi.org/10.1080/23249935.2020.1720856

Vigneau, M., Jobard, G., Mazoyer, B., Tzourio-Mazoyer, N., 2005. Word and non-word

reading: What role for the Visual Word Form Area? Neuroimage 27, 694–705.

https://doi.org/10.1016/j.neuroimage.2005.04.038

Vonsattel, J.P.G., DiFiglia, M., 1998. Huntington disease. J. Neuropathol. Exp. Neurol. 57,

369–384. https://doi.org/10.1097/00005072-199805000-00001

Wan Lee, S., Shimojo, S., O'Doherty, J.P., 2014. Neural computations underlying arbitration

between model-based and model-free learning. Neuron 81, 687.

https://doi.org/10.1016/J.NEURON.2013.11.028

Watson, P., de Wit, S., 2018. Current limits of experimental research into habits and future

directions. Curr. Opin. Behav. Sci. 20, 33–39.

https://doi.org/10.1016/j.cobeha.2017.09.012

Watson, P., van Wingen, G., de Wit, S., 2018. Conflicted between goal-directed and habitual

control, an fMRI investigation. eNeuro 5. https://doi.org/10.1523/ENEURO.0240-

18.2018

Watson, P., Wiers, R.W., Hommel, B., De Wit, S., 2014. Working for food you don't desire.

Cues interfere with goal-directed food-seeking. Appetite 79, 139–148.

https://doi.org/10.1016/j.appet.2014.04.005

Wechsler, K., Drescher, U., Janouch, C., Haeger, M., Voelcker-Rehage, C., Bock, O., 2018.

Multitasking during simulated car driving: A comparison of young and older persons.

Front. Psychol. 9, 910. https://doi.org/10.3389/fpsyg.2018.00910

West, R., Brown, J., 2013. Addiction, habit and instrumental learning. Theory Addict. 114–

135. https://doi.org/10.1002/9781118484890.CH5

Willingham, D.B., Koroshetz, W.J., 1993. Evidence for dissociable motor skills Huntington's

disease patients. In. Psychobiology 21.

Wong, A.L., Goldsmith, J., Forrence, A.D., Haith, A.M., Krakauer, J.W., 2017. Reaction times

can reflect habits rather than computations. Elife 6. https://doi.org/10.7554/eLife.28075

Wood, W., Mazar, A., Neal, D., 2021. Habits and Goals in Human Behavior: Separate but

Interacting Systems. Perspect. Psychol. Sci. https://doi.org/10.31234/OSF.IO/QVRBY

Wood, W., Neal, D.T., 2007. A New Look at Habits and the Habit-Goal Interface. Psychol.

Rev. 114, 843–863. https://doi.org/10.1037/0033-295X.114.4.843

Worringer, B., Langner, R., Koch, I., Eickhoff, S.B., Eickhoff, C.R., Binkofski, F.C., 2019.

Common and distinct neural correlates of dual-tasking and task-switching: a meta-

analytic review and a neuro-cognitive processing model of human multitasking. Brain

Struct. Funct. 224, 1845–1869. https://doi.org/10.1007/s00429-019-01870-4

Wu, T., Hallett, M., Chan, P., 2015. Motor automaticity in Parkinson's disease. Neurobiol.

Dis. 82, 226–234. https://doi.org/10.1016/j.nbd.2015.06.014

Wu, T., Zhang, J., Hallett, M., Feng, T., Hou, Y., Chan, P., 2016. Neural correlates

underlying Micrographia in Parkinson's disease. Brain 139, 144–160.

https://doi.org/10.1093/brain/awv319

Wunderlich, K., Dayan, P., Dolan, R.J., 2012. Mapping value based planning and extensively

trained choice in the human brain. Nat. Neurosci. 15, 786–791.

https://doi.org/10.1038/nn.3068

Wymbs, N.F., Bassett, D.S., Mucha, P.J., Porter, M.A., Grafton, S.T., 2012. Differential

Recruitment of the Sensorimotor Putamen and Frontoparietal Cortex during Motor

Chunking in Humans. Neuron 74, 936–946.

https://doi.org/10.1016/j.neuron.2012.03.038

Yamada, H., Matsumoto, N., Kimura, M., 2004. Tonically Active Neurons in the Primate

Caudate Nucleus and Putamen Differentially Encode Instructed Motivational Outcomes

of Action. J. Neurosci. 24, 3500–3510. https://doi.org/10.1523/JNEUROSCI.0068-

04.2004

Yang, Y., Zhang, J., Meng, Z.L., Qin, L., Liu, Y.F., Bi, H.Y., 2018. Neural correlates of

orthographic access in Mandarin Chinese writing: An fMRI study of the word-frequency

effect. Front. Behav. Neurosci. 12, 1–12. https://doi.org/10.3389/fnbeh.2018.00288

Yarkoni, T., Speer, N.K., Balota, D.A., McAvoy, M.P., Zacks, J.M., 2008. Pictures of a

thousand words: Investigating the neural mechanisms of reading with extremely rapid

event-related fMRI. Neuroimage 42, 973–987.

https://doi.org/10.1016/j.neuroimage.2008.04.258

Zwosta, K., Ruge, H., Goschke, T., Wolfensteller, U., 2018. Habit strength is predicted by

activity dynamics in goal-directed brain systems during training. Neuroimage 165, 125–

137. https://doi.org/10.1016/j.neuroimage.2017.09.062

2215

2216