Striatal role in everyday-life and laboratory-developed habits

Pasqualina Guida[1,2,6]*, Mario Michiels[1,2,6]*, Peter Redgrave[5], David Luque[3,4], Ignacio

Obeso[1,2]

* Co-first authors

[1]CINAC, Hospital Universitario HM Puerta del Sur, Móstoles, Madrid, Spain

[2]CIBERNED, Instituto de Salud Carlos III, Madrid, Spain

[3]Departamento de Psicología Básica, Autonoma de Madrid University, Madrid, Spain

[4]Departamento de Psicología Básica, Universidad de Málaga, Madrid, Spain

[5]Department of Psychology, University of Sheffield, Sheffield, S10 2TN, UK

[6]PhD program in Neuroscience, Autonoma de Madrid University, 28029 Madrid, Spain

Correspondence to Dr. Ignacio Obeso

Avda. Carlos V, 70

Móstoles, 28938

Spain

iobeso.hmcinac@hmhospitales.com

26   **Abstract**

The dorsolateral striatum plays a major role in stimulus-response habits that are

learned in the experimental laboratory. Here, we use meta-analytic procedures to identify the

neural circuits activated during the execution of stimulus-response behaviours acquired in

everyday-life and those activated by habits acquired in the laboratory. In the case of

everyday-life habits we dissociated motor and associative components. We found that

motor-dominant stimulus-response associations developed outside the laboratory engaged

posterior dorsal putamen, supplementary motor area (SMA) and cerebellum. Associative

components were also represented in the posterior putamen. Meanwhile, newly learned

habits relied more on the anterior putamen with activation expanding to caudate and nucleus

accumbens. Importantly, common neural representations for both naturalistic and laboratory

based habits were found in posterior left and anterior right putamen. Our findings suggest a

common striatal substrate for behaviours with significant stimulus-response associations,

independently of whether they were acquired in the laboratory or everyday-life.

Keywords: habits, everyday-life, probabilistic learning, cortex, striatum, meta-analysis, fMRI

Introduction

Much of human behaviour can become automated and executed without conscious thought and attention. Thus, over time, a vast array of sometimes sophisticated movements, thoughts, attitudes and motivations can be performed under automatic stimulus-response control. Typically, we refer to such behaviour as *habits*. Throughout life, repeated associations between representations of specific stimuli and particular responses enable stimulus-evoked behaviour to be enacted without conscious intervention [1]. This allows us to perform tasks with reliable stimulus-response components without thought, while at the same time, allowing conscious attention to be directed to less predictable aspects of the world. An obvious example would be not having to think about putting one foot in front of the other when walking to an interview. Hence, numerous habitual routines are constructed by humans to manage the simultaneous enactment of multiple tasks – walking and talking [2].

A significant feature of habitual behaviour is it occurs independently of how valuable or appropriate the outcome is [3,4]. An example would be habitually pressing the lift/elevator button taking you to the floor of your old office, rather than the new one. Automatic habitual control is often contrasted with conscious goal-directed processing where action selection is determined by the value/appropriateness of the predicted outcome [3,5,6]. Typically, before statistical regularities in the task are established, adaptive goal-directed control is slower but more flexible. Thus, early in instrumental learning, and in uncertain situations, flexible goal-directed control is normally deployed.

Until recently, habits have been studied in experimental settings where new stimulus-response relationships are learned in the laboratory. However, it is of great interest to understand how automatic habitual control operates in the normal circumstances of everyday life. It is also important to understand how brain disorders such as Parkinson's disease [7,8], obsessive-compulsive disorder [9], and drug addictions [10] lead to clinically dysfunctional patterns of habitual use of behaviour. Despite the importance of habitual

control in daily life and clinical pathology, formal investigation of the inherent stimulus-

response associations established in everyday life is in its infancy. However, even at this

early stage, an important question is whether the neural circuits engaged by habits acquired

in daily life are the same or different to those activated when newly acquired habits are

performed in the laboratory. The purpose of the present investigation was therefore to

compare the patterns of neural activation evoked by long-acquired habits brought into the

laboratory with those established under formal experimental conditions.

Experimental research with animals has shown how instrumental learning occurs

through initial goal-directed computations that later transition into stimulus-response

mappings [11]. The formal experimental paradigms (see Box 1 for a summary) used to

distinguish habits from goal-directed actions include (amongst others) outcome devaluation

or contingency degradation tests (Adams, 1982; Balleine and O'Doherty, 2010; Dickinson et

al., 1985; for a review see Foerde, 2018). In both cases, if the behaviour persists after the

outcome has been devalued or the outcome is no longer related to responding, the

behaviour is deemed to be under habitual control. The formal procedures developed in

animal studies have been imported into human experiments [14–18]. Significantly, human

neuroimaging studies have revealed activation of the rostro-medial (associative) striatum

during the initial stages of instrumental learning, which gradually shifted to caudo-lateral

(sensorimotor) regions of the striatum when habitual control of the instrumental task became

evident [14,19–24]. These findings concur with those in non-human animals that demonstrate a

similar involvement of the rostro-medial striatal territories early, and caudo-lateral regions

later in the acquisition of instrumental tasks [11].

To date, most of the literature investigating the neural basis of habits in humans has

relied on subjects developing new experimental habits in the laboratory under formal

experimental conditions (Box 1B). This approach, has provided important information about

the relationship between habitual and goal-directed neural systems [11] and the factors

(repetition, rewards or environmental cues) that promote the formation of habitual control [25].

However, this approach faces several challenges. The most pressing is the difficulty of

creating habits in an experimental situation that are equivalent to those developed in normal

everyday life. For example, a basic tenet of habit theory is that stimulus-response

associations typically get stronger with repetition. It is therefore, not exactly inspiring that

several studies have failed to report a direct positive relationship between the amount of

training and the strength of habitual responses measured in their experimental settings [26,27].

An alternative and increasingly important way forward to study habits in humans is to

have everyday habitual behaviour learned during a subjects lifetime brought into the

laboratory for investigation. Certain behaviours in normal life including driving, eating,

dancing, reading, talking or walking have significant stimulus-response components that can

be performed automatically while the person's conscious attention is directed elsewhere [28,29].

It is likely these components have been acquired through a life-time of everyday trials.

Therefore, such associations come to the laboratory fully formed and do not depend upon on

the multiple trial learning that is required to develop new experimental habitual behaviour.

The principal challenge in studying naturalistic habits is getting subjects to express long-

established stimulus-response behaviour in a laboratory setting. This is necessary so that

both the automatic behaviour and associated neural activity can be measured quantitatively.

Investigators of everyday habits have chosen behaviours that have critical automatic

stimulus-response components [30] and versions that can be performed in the laboratory while

BOLD signals are measured by fMRI imaging. Examples of such tasks include reading,

where comparisons are made between real words of different familiarity and emotional

content, foreign words and pseudo-words [31]; writing and drawing [32,33] walking on a special

apparatus [34]; and driving an MR-compatible driving simulator (Box 1A; Choi et al., 2017;

Cummine et al., 2016; Huth et al., 2016; Karimpoor et al., 2015; Martínez et al., 2016, 2018;

Oberhuber et al., 2013; Varotto et al., 2020; Yang et al., 2018). The specific question we

were interested in is whether recently established laboratory habits and those developed in

everyday life engaged the same, partially overlapping or separate neural circuits in the brain.

The purpose of the present investigation was therefore to directly compare the neural

signatures evoked by novel, laboratory-developed habits and those acquired over a life-time

(Figure 1). To answer this question, we conducted a quantitative meta-analysis to

investigate the neural substrates of everyday life and experimental habits. First, to select a

cohort of studies investigating naturalistic habits, we took data from 54 studies (a total of

1441 subjects) that had used diverse stimulus-response paradigms (walking, reading, writing

and driving). Imaging models that included automatic parameters on each task were chosen

(Table S1). Meanwhile, we included categorical variables that separated motor (walking,

driving) and cognitive (reading, writing) habits acquired in daily life, a further sub-division

motivated by the anatomo-functional gradient along fronto-striatal circuits [41]. Second, we

sought to confirm the neural basis of experimental habits in the 40 studies (a total of 973

subjects) that had used probabilistic or discriminative learning, 2-step learning or sequential

tasks to test for the laboratory development of novel habits (Box 1B). Studies of

experimental habits were separated into two subcategories: probabilistic learning vs other

experimental tasks (Table S2), on the ground that probabilistic classification typically

activates more anterior portions of the caudate and the putamen, compared with other tasks

[22]. This novel approach allowed the neural signatures of stimulus-response behaviour

developed in the laboratory and in everyday life to be compared directly. Special interest

was focused on the system-level circuits involving the basal ganglia (Figure 1). The resulting

information provides important clues into the organisation of normal habits against which

pathologies of habit and the results of therapeutic interventions can be referenced.

Figure 1. *Hypothesis diagram on striatal role in both everyday-life and experimental habits. Activities*
part of daily life such as writing, reading, walking or driving were selected as everyday-life habits (see
Box 1 for task measurements and details) to expect a critical striatal role in executing these habits
forms. Similar striatal activities can be expected compared to experimental paradigms commonly
used in the cognitive science literature.

Methods

Selection of studies. We conducted 2 independent searches in PubMed to

identify fMRI studies investigating stimulus-response habits acquired (1) in everyday life and

(2) in experimental laboratories. The search of articles related to everyday habits was

focused on natural behaviours where an important automatic stimulus-response component

would be expected (e.g. walking, speech, reading, etc). Selection of these activities was

based on their potential link to fundamental basal ganglia functions (Figure 1). A summary of

typical measures used in these studies is presented in Box 1. In the search for laboratory

based habits we included articles referenced by Patterson & Knowlton (2018) plus all new

relevant articles published since their last search (June 22, 2017) and March 26, 2021. To

enable direct comparison with, and extension of, the Patterson and Knowlton review (2018),

we conducted our literature search using the same key search terms and included the task

categories – probabilistic learning, discriminative learning and sequence learning (see Box

1). We also included relevant MeSH terms available in Pubmed. This allowed us to find

articles that do not include targeted keywords in the title or abstract but have proper MeSH

terms linked to their metadata. The complete list of our search terms can be found in

Supplementary Material. To set limits on inclusion, the articles were filtered according to the

following criteria:

(1)   Spatial coordinates from human brain fMRI reported in standardized

stereotaxic space (MNI or Talairach space). Other functional imaging methods (e.g., PET,

EEG source imaging, etc) were excluded;

(2)   Healthy subjects over 18 years old were included, as were the healthy

participants within clinical studies with independent analyses for the healthy controls;

(3)   At least 6 subjects;

(4)   Whole-brain analysis. ROI analyses were excluded as they can be biased by

the study hypothesis and may not report all significant regions;

(5)   For studies using the two-step probabilistic tasks and analyzing prediction

errors from computational models, we included results associated with model-free (reward

prediction error, RPE), but not from model-based errors (state prediction error, SPE);

(6)   When a study contained multiple experiments and/or contrasts, the contrasts

selected were based on the task condition and comparisons linked to stimulus-response

behaviours (see Table S1 and S2);

(7)   Studies reporting original data and published in English peer-reviewed

journals (reviews and meta-analysis were excluded).

We then performed a second-step control using reference lists from already

included articles. From the list of references, we first excluded articles whose title had no

direct link to any of our inclusion criteria. From the remaining articles that met our inclusion

criteria we found 17 that had investigated everyday-life habits and 14 articles that had

investigated laboratory-developed habits. Since our second exclusion step involved manual,

rather than online searching, we were able to include articles published before 2017 not

included in Patterson and Knowlton (2018). These additional references were selected

because: (1) we checked all references from the list of accepted papers, not only those in

major review articles; and (2) our criteria allowed us to include studies that did not report

putative habit-related activation of the dorsal striatum (caudate, putamen, or both). Figure S1

depicts the identification, screening, eligibility and selection stages we used to select studies

for inclusion.

We included a total of 54 studies investigating everyday-life habits (31 involved

cognitive tasks: writing, reading; and 23 studied motor tasks: walking, driving; Table S1). A

total of 40 studies involved reported experimental habits (20 involved probabilistic learning;

and 20 studied other experimental tasks; Table S2). Foci, scripts and statistical maps can be

accessed in the Open Science Framework (https://osf.io/w5ftm).

Data analysis. The latest version of the GingerALE software v3.0.2 [42] was used to

compute activation likelihood estimations (ALE) [43]. From each of the accepted articles, the

coordinates of peak activations were manually extracted and those in Talairach space were

transformed to Montreal Neurological Institute (MNI) space using the inverse transform of

icbm2tal from GingerALE [44]. The ALE method was implemented as follows: (1) for each

study a 3D Gaussian distribution was created around every peak activation (variance

proportional to the sample size of the study). This allowed us to use the higher statistical

power to reduce peak uncertainty in studies with larger sample sizes. This step was done for

every peak to produce one modelled activation (MA) map per study. The value for each

voxel in a MA-map represented the probability of that voxel containing an activation foci. (2)

Voxel probabilities in all MA-maps were merged to produce an ALE map. Following

recommendations of Eickhoff et al., (2016), the ALE map was thresholded using a cluster-

level Family-Wise Error (FWE), with a cluster-forming threshold of .001 and a cluster-level

FWE of .05 (as stated in the GingerALE manual; Fox et al., 2013).

The first step in applying a cluster-level FWE was to threshold the ALE map at the

voxel-level (cluster-forming threshold). To do this, all possible combinations between all

peaks from each MA-map were tested. The ALE values provided a null distribution which

assumed the peaks would be randomly distributed following all potential combinations

amongst them. Here, an uncorrected *p*-value was used since in the next step, a FWE

correction was applied to correct for the possibility of multiple comparison errors.

Then, an additional threshold (cluster-level FWE) was applied to select only the

largest clusters. Peaks in every MA-map were distributed randomly and then combined into

a single ALE map, following the same union method described above for the voxel-level

threshold. The ALE maps were also thresholded with the same method used at the voxel-

level threshold. This procedure was repeated (1000 times for this meta-analysis) using

Monte Carlo permutations method, selecting on every run the largest cluster in the

thresholded ALE map (null distribution). Finally, we applied the selected threshold (0.05) to

obtain the FWE-corrected results. Using a 0.05 cluster-level threshold resulted in a

thresholded map where only 5% of the surviving clusters could have been introduced by

chance (false positives), following guidelines to discard non-significant clusters [42].

Contrasts between thresholded ALE maps were computed to compare between our

conditions of interest (e.g. Everyday-life > Experimental maps). All foci in the selected

studies were pooled into a single dataset, which we then split in two datasets by randomly

assigning to each one the same number of foci in their original file. The ALE-scores of these

two random maps were then subtracted in a voxel-wise manner generating the map of their

ALE-scores differences (repeated 10000 times to record ALE-scores and generate a null

distribution of randomly spatially distributed foci). Finally, the actual two ALE maps

corresponding to the contrast were combined voxel-wise by subtracting the ALE-score of

one map from the other. The resulting map of ALE-scores were then thresholded voxel-by-

voxel by comparing them with the null distribution previously obtained. We used the default

p-value suggested in GingerALE (0.01 uncorrected). All computations were run via in-house

python scripts to automate analyses in the GingerALE's interface. Thresholded maps are

reported in MNI152 space [47].

Results

Striatal role in everyday and experimental habits

To find the neural signatures of everyday habits, we obtained cluster activity

associated with stimulus-response behaviours learned throughout life, compared with those

newly learnt in an experimental laboratory. The main effects of naturalistic habits revealed

significant bilateral activity in the posterior putamen (Table 1; Figure 2A). Specifically, this

was sustained by activity in dorsal sections and left putamen activation that expanded to its

posterior boundary (Table 1; Figure 2A-B). For these tasks, other active regions were seen

in the cerebellum and cortical areas including the premotor and SMA (Table 1; Figure 2A-B).

Habits established in laboratory settings were also associated with increased levels

of activation in putaminal sections, but with larger representations in anterior striatum (Figure

2C). A gradient was seen along the rostro-ventral section of the putamen, right caudate and

the nucleus accumbens bilaterally (Table 2; Figure 2C). These results confirm the striatum

as the most significant region for newly acquired experimental habits (Table 2; Figure 2C) [22].

They were also consistent with the idea that stimulus-response associations learned in the

laboratory would depend more on rostral striatal activity. The right insula was one of the

extrastriatal hubs of activity associated with habits acquired in the laboratory (Table 2; Figure

2A). These findings align with the evidence that sensorimotor territories of the striatum are a

critical for the expression of habitual behaviour, but with more rostral patterns of activity

observed when experimental habits were developed in laboratory setting.

266    Of particular interest in this investigation was the possibility that despite the

267 differences between neural activation patterns associated with everyday and experimental

268 habits, there may be a common neural substrate accessed by both forms of stimulus-

269 response behaviour. Common activations were seen in anterior right putamen and posterior

270 left putamen (Figure 2A). Although both categories of habits recruited dorsal sub-regions of

271 the putamen bilaterally, the activation by everyday habits was stronger (Figure 2C).

272 Unexpectedly, no activation in the caudate nucleus survived FWE-corrected thresholds with

273 everyday habits. In contrast, habits acquired in the laboratory showed a differential

274 recruitment of the nucleus accumbens, and to a lesser extent, the most antero-ventral

275 section of the right caudate nucleus and putamen (Figure 2C). Hence, while the striatum is a

276 common hub for both categories of habitual behaviour, antero-posterior differences were

277 reflected in the activation patterns of habits acquired in everyday life and the experimental

278 laboratory.

Figure 2. *Everyday-life and experimental thresholded activation maps.* **a)** *Axial views for the main regions. Overlap regions at the right column are shown in blue. Note that there exists activation in the cerebellar cortex for the case of everyday-life studies but it is omitted here for brevity. Z=52 view for the experimental studies is shown as an unthresholded map for visualization purposes (ALE value ≈ 0.01)* **(b)** *3D striatum reconstruction showing all the activation that fall inside it. Overlap regions at the right column are shown in blue.* **c)** *3D striatum reconstruction showing the differential activation of the Everyday -life > Experimental contrast (in green) and the Experimental > Everyday-life contrast (in red).*

Motor and associative segregation in life-long habits

We next tested whether the neural correlates of everyday habits varied between

sensorimotor routines (walking or driving) and cognitive-associative ones (reading or

writing). While all representations of habitual behaviour are characterized by stimulus-

response associations, for different tasks these may be represented in different sensorimotor

networks with differential connections to the basal ganglia. Clustered activation in motor

tasks revealed a significant presence along the dorsal putamen compared with the

associative tasks (Figure 3A). In addition, motor tasks extended the pattern of activation into

more posterior sections of the right putamen, but only dorsally for the left putamen. This

cluster was present across all the antero-posterior axes (Figure 3B).

In the case of associative habits acquired in everyday life, we observed an activation

pattern in ventral sections of the putamen bilaterally (Figure 3A). This effect showed

significant bilateral asymmetry favouring the left putamen where activation extended into

more posterior regions. This was absent in the walking and driving tasks (Figure 3A).

Outside the striatum at the cortical level, the associative tasks acquired in everyday

life recruited premotor area and SMA. Activation was also observed in the cerebellum

(Figure 3A). The same cortical regions were recruited during the walking and driving tasks,

but none survived FWE-corrected thresholds. Greater activation for the associative cognitive

tasks was expected given the more complex computational requirements when writing or

reading [48]. Such tasks are known to engage larger regions of cortical tissue [49].

To further investigate the failure of our thresholding procedures to detect activation in

the caudate nucleus for any of the naturalistic habit categories, we included the distribution

of all the foci that fell into the striatum in both categories of everyday habits (Figure 3C). This

allowed activity in the right caudate to be observed (Figure 3C) in some associative studies,

but not for the motor ones. For the left caudate nucleus, no foci were reported in any study,

which confirms the absence of activity reported in the thresholded maps above.

These findings provide insights into possible parallel habitual mechanisms in motor-

associative domains that may share regions of the posterior putamen for motor activities, but

a broader network for actions that require higher-order cognition. Given the stimulus-

response associations involved in everyday habits will simultaneously engage multiple

mechanisms, including visuo-spatial attention, movement planning and execution, the neural

substrate of cognitive and sensorimotor habits should be segregated.

Figure 3. *Everyday-life cognitive-motor subcategories thresholded activation maps. **a)** Axial views for the main regions. Overlap regions at the right column are shown in blue. **b)** 3D striatum reconstruction showing all the activation that fall inside it. Overlap regions at the right column are shown in blue. **c)** Foci distribution of all the studies that fall in the striatum. Note that some foci may correspond to the same study.*

Probabilistic learning and other tasks segregation in experimental habits

329   Finally, we aimed to replicate previous findings on laboratory conditions that have

330 linked the striatum to the learning and execution of new habits. Consistent striatal activity

331 has been reported when evaluating probabilistic or discriminative learning, 2-step learning

332 and sequential tasks [22]. Here, we intended to confirm and update these results with more

333 recent findings by separating studies that involve trial-and-error probabilistic reward learning

334 from those that used different methodologies for stimulus learning. As predicted, both

335 probabilistic learning and the other tasks showed common regions of striatal activation, with

336 largest clusters in the nucleus accumbens and rostro-ventral sections of the caudate and

337 putamen (Figure 4A). However, only the probabilistic tasks were associated with bilateral

338 recruitment of the most anterior region of the putamen (Figure 4B). This results confirms

339 previous findings reported by Patterson and Knowlton, (2018). Similarly, the other

340 experimental tasks differed with respect to the probabilistic ones by demonstrating unilateral

341 recruitment of the left rostral caudate nucleus (Figure 4B). Outside the striatum, left insular

342 cortex was activated in other tasks but not by the probabilistic ones (Figure 4A). This could

343 reflect a form of residual goal-directed activity due to the insula's complementary role in

344 evaluating future rewards in the decision phase [50].

Figure 4. Experimental subcategories thresholded activation maps. *a) Axial views for the main regions. In this case there are no overlap regions at the right column. b) 3D striatum reconstruction showing all the activation that fall inside it. In this case there are no overlap regions at the right column. c) Foci distribution of all the studies that fall in the striatum. Note that some foci may correspond to the same study.*

Discussion

Our meta-analysis has identified two distinct types of habit-related mechanisms in the

human striatum, by comparing long-established habits acquired in everyday life with newly

learned habits acquired under laboratory-controlled conditions. The results of our

investigation revealed both common and diverse functional links between different sub-

regions of the brain's habitual circuitry. Naturalistic habits showed enhanced activity in the

dorsal posterior putamen, together with activity in the cerebellum and SMA. In contrast,

laboratory acquired habits engaged anterior sections of the putamen with activation

expanding to caudate nucleus and nucleus accumbens. However, common regions of

activation were found in posterior left and anterior right putamen. Ultimately, delineation of

specific striatal contributions to motor-associative variables embedded in habits were

responsible for shared anatomical patterns in everyday-life associative habits, such as

reading and writing, that engaged rostral regions of the putamen.

The current findings provide direct evidence for bilateral putamen engagement in

habits acquired and executed in everyday life. Typically, we observed larger bilateral activity

in studies of long-established habits compared with newly acquired experimental ones. This

may implicate a broad putaminal role for the stimuli-rich sensorimotor computations required

during complex stimulus-response tasks enacted in everyday life. Consistent with this view,

neuroimaging, lesion and animal electrophysiological data all converge to pinpoint a role for

the putamen in the integration of movement units and stimulus associations to produce

behaviour with a predicted outcome [51–55]. Indeed, the neurophysiological properties of the

putamen are supported by a subpopulation of neurons that respond to sensory stimuli [56],

unifying actions sets for movement sequences [57,58] or integrating elementary movement

units such as individual finger moves [59]. Moreover, its activity is not solely dedicated to

movement parameters, but also in the absence of motor plans [60], increasing response

magnitude to the reinforced choice [61] and for predicted well-learned and contextually-driven

actions [62–67]. Hence, the diversity of sensory-related and high-order reinforcement neurons in

the putamen support a pivotal role in context-rich scenarios at several levels (sensorimotor,

predicted actions, object value, habitual action execution). We interpret the association of

putamen activity with habits in everyday life being due to its physiological and reinforcement

properties contributing to action sequence-specific and context-specific behaviour.

The role of the putamen in habitual behaviour is directly influenced by ascending

dopaminergic system acting on cortical and thalamic inputs [68]. Dopamine provides critical

modulatory influences on striatal subregions whereby projections from substantia nigra pars

compacta and the ventral tegmental area differentially target terminals in dorsal and ventral

striatum respectively [69]. The main nigrostriatal projection is topographically organised with a

medial to lateral gradient [70]. Tonic firing within this system sustains, motivational, cognitive

and action-specific decision making [71,72], while sensory-evoked phasic patterns of

dopaminergic activity provide a general mechanism for reinforcement learning [73,74]. The

large bilateral activity found in the striatum for the routines of everyday life should be

influenced by ascending dopaminergic fibers with the abovementioned functional properties

that likely contribute to sensorimotor control while writing or walking (proprioceptive and

muscle control). As well, dopaminergic projections will fire phasically to unpredicted

associative cues present in everyday life [75]. Hence, the putamen will likely respond

preferentially to the sensorimotor contingences present in highly-organised actions that rely

on automatic stimulus-response associations that characterise habitual responding.

A second key contributor comes from particular cortical structures that regulate the

expression of habits [23,67,76,77]. In the present study we found that SMA activity was an

essential component of the habitual circuitry present for the everyday tasks. The SMA

represents one of the important junctions between cortical-subcortical motor and cognitive

circuits [78]. It has projections to the dorso-lateral striatum and posterior putamen [79], a

pathway subject to neuromodulation by non-invasive brain stimulation [80]. The SMA has

been shown to be involved in learning stimulus-response contingencies [78,81] and model-free

tasks [82]. Multi-dimensional components required while driving or even walking

(somatosensory, visuo-spatial, prediction, motor planning, etc) will be mediated via cortical

inputs to the putamen [83,84]. How the putamen might operate on this range of input would be

to perform sequential selections of stimuli that trigger previously acquired stimulus-response

associations. When bolted together such serial selections can be viewed as coherent

sequences of habitual behaviour, such as changing gear while driving, or pen movements

associated with particular letters when writing. These automatic moves are likely to recruit

medial cortical motor areas including the SMA, that collaborate with subcortical structures,

including putamen (Cunnington et al., 2002; Smittenaar et al., 2013). Recurrent cortico-

striatal loops would therefore combine to mediate sequential cognitive and motor

components of habitual behaviour, both triggered by the specific sensory events they have

been repeatedly associated within everyday life.

To investigate the possible commonalities and differences in the neural networks

supporting categories of automatic behaviours acquired in everyday life, we subdivided

those activities with greater motor component (walking, driving) compared to those with

stronger cognitive requirements (reading and writing). This analysis was motivated by

possible differences in the contribution from motor, cognitive and/or emotional circuits in the

different types of everyday habitual activity [88]. The dorso-lateral caudal putamen was active

for both cognitive (reading, writing) and motor operations (driving, walking), which

corresponds with our understanding of parallel inter-related cortico-striatal functions [82,89].

The stimulus-response selection role of the putamen may well represent the low level

sensorimotor selections necessary in all forms of habitual motor behaviour [60,75]. Parallel

circuits have been suggested to integrate the different environmental signals that trigger

motivational, cognitive and motor responses [90,91]. Sequential selections in different territories

of the basal ganglia could be integrated by cortico-basal ganglia loops into coherent goal-

directed behaviour [92]. The question that remains is whether each of the limbic, associative

and sensorimotor territories [93] can all operate in stimulus-response habitual mode to elicit

respectively, stimulus-evoked motivations (e.g. drug cravings), stimulus-evoked cognitions

(prejudices) as well as the well-known motor habits.

Critically, our findings also have implications for previous studies investigating new

stimulus-response habits learned in experimental laboratories (probabilistic or discriminative

learning, 2-step learning or sequential tasks), which also report significant striatal activations

[14,19,21–23,82,94–96]. Specifically, our analyses show stimulus-responses behaviour acquired both

in everyday-life and under experimental conditions showed common striatal activity in the

right anterior putamen and left posterior putamen. The everyday-life tasks included long

established sensorimotor responses triggered by sensory cues in the absence of new

associative learning. In contrast, laboratory learned habits typically involved relatively minor

motor components (finger key presses), but novel cue-response associations driven by

reinforcement learning. In line with a previous meta-analysis on basal ganglia activation

across multiple motor disciplines (Arsalidou et al., 2013), left lateralized putaminal activity

was prominent in motor operations (such as eye movements and body motion), has a larger

volume in right-handed participants (Peterson et al., 1993) and is critical in behaviours

guided by stimulus-response mappings [22,95]. Hence, we interpret left posterior putamen

activation being present in both long-established and novel habit forms reflecting a critical

sensorimotor association embedded within every habitual response.

However, activation of the anterior putamen would accord well with neural patterns

associated with initial learning of stimulus-response associations in experimental settings [21].

Activity in key regions of the circuitry associated with goal-directed behavioural control

(caudate and nucleus accumbens) were also present in laboratory studies of habits. These

findings match those of several fMRI studies using various reinforcement learning tasks,

which report activation of the ventro-medial prefrontal cortex, insula or anterior striatum

when encoding the value of predicted reward outcome linked to new actions [20,50,98]. Yet, the

results with learning new habits may be explained by the theory and methodological

procedures often used when studying habit acquisition in the laboratory. For example, part of

the problem is that some of the presumed habits established in the laboratory fail to meet the

formal requirement of automatic stimulus-response behaviour. Thus, several recent human

experimental studies failed to demonstrate the expected effect of training duration on the

outcome-devaluation test [17,26,27]. This suggests rather more trials may be needed to establish

stimulus-response associations that can survive devalued outcome challenges and can be

enacted automatically without thought.

Recently, alternative procedures have been developed to establish and test new

habits in the experimental laboratory, including the sudden reversal of learned actions after

overtraining [18], overloading goal-directed top-down control while measuring execution of

learned stimulus-response associations [99] or biasing movement kinematics [100]. Interestingly,

pre-existing categorical associations established in everyday life (i.e. color associations or

prejudice) have a clear advantage on measuring automatic processing [30,101]. Further

methodological options for studying long established habits acquired in everyday life include

assessing expert musicians [102], tennis players [103] and expert shooters [104]. Hence, getting

subjects to bring their everyday-life stimulus-response associations into the laboratory under

controlled conditions is an important option for studying the neural substrates underlying

habitual behaviour in humans. Although there is less control of the independent variables in

such studies, by selecting subjects with different amounts of everyday life experience it is

possible to relate the amount of practice with habit strength. The dependent consequences

of life-long habits can be measured subsequently in the laboratory with traditional outcome

devaluation, contingency degradation, or dual processing procedures.

Insofar as the caudal putamen has been identified as a critical node in the neural

substrate responsible for automatic habitual behaviour, malfunctioning of this region has

been associated with deficits in habitual performance. One notable instance is the differential

loss of dopamine neurotransmission from this region in Parkinson's disease [7] a putative

pathophysiological condition linked to the cost of life-long use of habits [105]. The problems

Parkinson's disease patients have with walking and writing [8,106,107] and the new learning of

experimental habits [108,109], have been interpreted as an inability to express stimulus-

response habits. In contrast, other neuropsychiatric complications such as addictions exhibit

an excessive cue-dependent use of certain rewards linked to increased posterior putamen

activity [10,110]. Hence, depending on the nature of the neurobiological disruption, the habitual

system seems to be underused or overused in different clinical conditions.

Despite the clear positive results of our analysis certain limitations of meta-analytic

procedures must be acknowledged. First, the ALE analysis can cause some parameters,

such as voxel peaks, to be overlooked. Ideally, to explore the statistical activity maps of each

study individually would be of great value. However, the fact that most studies did not have

full imaging datasets would preclude this. Second, we included results from prior studies

whose experimental focus was not on habitual behaviours acquired in everyday life.

Specifically, in these studies behaviours were not formally identified as habitual using

outcome devaluation or contingency degradation tests. However, it should be noted that a

life-time repetition of everyday trials builds strong associations that do not depend upon new

stimulus learning required in experimental studies [111,112]. Last, our meta-analysis included

tasks with significant heterogeneity and regional foci were selected from studies with varying

contrasts. To overcome these issues, we selected activity maps only from studies that

reported contrasts measuring automatic components of behaviour. This suggests that our

dataset has enabled us to identify activity patterns that are shared by a diverse range of

human behaviours that contain a significant stimulus-response element. Consequently, it is

possible that the bilateral posterior putamen acts as a critical node when practiced behaviour

can be performed automatically and without thought in situations of everyday life.

Conclusions

The present study points to a fundamental functional role for the posterior putamen in

the expression of habits acquired in everyday life. Importantly, a critical dissociation was

established between different brain regions whose contributions to motor-cognitive

representations of habits have, thus far, been largely indistinguishable. Conversely, the

engagement of the anterior putamen is associated more with habits newly learned in

experimental laboratories. Careful experimental protocols must be designed to identify those

chunks of behaviour that are under stimulus-response control and can occur independently

of outcome value or response contingency. This is true both for newly acquired associations

in the laboratory and habits established in everyday life. Finally, the present study highlights

the importance and value of having subjects bring life-long habits into the laboratory to be

investigated and compared with recently acquired stimulus-response associations.

References

1. Haith, A. M. & Krakauer, J. W. The multiple effects of practice: skill, habit and reduced cognitive load. *Curr. Opin. Behav. Sci.* **20**, 196–201 (2018).

2. Sun, R. Desiderata for cognitive architectures. *Philos. Psychol.* **17**, 341–373 (2004).

3. Dickinson, A., Trans, P. & Lond, R. S. Actions and habits: the development of behavioural autonomy. *Philos. Trans. R. Soc. London. B, Biol. Sci.* **308**, 67–78 (1985).

4. Miller, K. J., Shenhav, A. & Ludvig, E. A. Habits without values. *Psychological Review* **126**, 292–311 (2019).

5. Daw, N. D., Niv, Y. & Dayan, P. Uncertainty-based competition between prefrontal and dorsolateral striatal systems for behavioral control. *Nat. Neurosci.* **8**, 1704–11 (2005).

6. Gläscher, J., Daw, N., Dayan, P. & O'Doherty, J. P. States versus rewards: Dissociable neural prediction error signals underlying model-based and model-free reinforcement learning. *Neuron* **66**, 585–595 (2010).

7. Redgrave, P. *et al.* Goal-directed and habitual control in the basal ganglia: Implications for Parkinson's disease. *Nature Reviews Neuroscience* **11**, 760–772 (2010).

8. Bannard, C. *et al.* Reduced habit-driven errors in Parkinson's Disease. *Sci. Rep.* **9**, 1–8 (2019).

9. Gillan, C. M. Recent Developments in the Habit Hypothesis of OCD and Compulsive Disorders. in 1–21 (Springer, Berlin, Heidelberg, 2021). doi:10.1007/7854_2020_199

10. Sjoerds, Z. *et al.* Behavioral and neuroimaging evidence for overreliance on habit learning in alcohol-dependent patients. *Transl. Psychiatry* **3**, e337-8 (2013).

11.   Balleine, B. W. & O'Doherty, J. P. Human and rodent homologies in action control:

Corticostriatal determinants of goal-directed and habitual action.

Neuropsychopharmacology **35**, 48–69 (2010).

12.   Adams, C. D. Variations in the Sensitivity of Instrumental Responding to Reinforcer

Devaluation. *Q. J. Exp. Psychol. Sect. B* **34**, 77–98 (1982).

13.   Foerde, K. What are habits and do they depend on the striatum? A view from the

study of neuropsychological populations. *Curr. Opin. Behav. Sci.* **20**, 17–24 (2018).

14.   Tricomi, E., Balleine, B. W. & O'Doherty, J. P. A specific role for posterior dorsolateral

striatum in human habit learning. *Eur. J. Neurosci.* **29**, 2225–32 (2009).

15.   Valentin, V. V., Dickinson, A. & O'Doherty, J. P. Determining the neural substrates of

goal-directed learning in the human brain. *J. Neurosci.* **27**, 4019–4026 (2007).

16.   Watson, P., Wiers, R. W., Hommel, B. & De Wit, S. Working for food you don't desire.

Cues interfere with goal-directed food-seeking. *Appetite* **79**, 139–148 (2014).

17.   Watson, P. & de Wit, S. Current limits of experimental research into habits and future

directions. *Curr. Opin. Behav. Sci.* **20**, 33–39 (2018).

18.   Luque, D., Molinero, S., Watson, P., López, F. J. & Le Pelley, M. E. Measuring habit

formation through goal-directed response switching. *J. Exp. Psychol. Gen.* **149**, 1449–

1459 (2020).

19.   Lehéricy, S. *et al.* Distinct basal ganglia territories are engaged in early and advanced

motor sequence learning. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 12566–12571 (2005).

20.   Tanaka, S. C., Balleine, B. W. & O'Doherty, J. P. Calculating consequences: Brain

systems that encode the causal effects of actions. *J. Neurosci.* **28**, 6750–6755 (2008).

21.   Jankowski, J., Scheef, L., Hüppe, C. & Boecker, H. Distinct striatal regions for

planning and executing novel and automated movement sequences. *Neuroimage* **44**,

1369–1379 (2009).

22.   Patterson, T. K. & Knowlton, B. J. Subregional specificity in human striatal habit

learning: a meta-analytic review of the fMRI literature. *Curr. Opin. Behav. Sci.* **20**, 75–

82 (2018).

23.   de Wit, S. *et al.* Corticostriatal connectivity underlies individual differences in the

balance between habitual and goal-directed action control. *J. Neurosci.* **32**, 12066–

12075 (2012).

24.   Brovelli, A., Nazarian, B., Meunier, M. & Boussaoud, D. Differential roles of caudate

nucleus and putamen during instrumental learning. *Neuroimage* **57**, 1580–90 (2011).

25.   Smith, K. S. & Graybiel, A. M. Habit formation. *Dialogues Clin. Neurosci.* **18**, 33–43

(2016).

26.   de Wit, S. *et al.* Shifting the balance between goals and habits: Five failures in

experimental habit induction. *J. Exp. Psychol. Gen.* **147**, 1043–1065 (2018).

27.   Pool, E. *et al.* Determining the effects of training duration on the behavioral

expression of habitual control in humans: a multi-laboratory investigation. 48 (2021).

doi:10.31234/OSF.IO/Z756H

28.   Wechsler, K. *et al.* Multitasking during simulated car driving: A comparison of young

and older persons. *Front. Psychol.* **9**, 910 (2018).

29.   Worringer, B. *et al.* Common and distinct neural correlates of dual-tasking and task-

switching: a meta-analytic review and a neuro-cognitive processing model of human

multitasking. *Brain Struct. Funct.* **224**, 1845–1869 (2019).

30.   Ceceli, A. O., Myers, C. E. & Tricomi, E. Demonstrating and disrupting well-learned

habits. *PLoS One* **15**, 1–28 (2020).

31.   Cummine, J. *et al.* Understanding the Role of Speech Production in Reading:

Evidence for a Print-to-Speech Neural Network Using Graphical Analysis.

Neuropsychology **30**, 385–97 (2016).

32. Yang, Y. *et al.* Neural correlates of orthographic access in Mandarin Chinese writing:

An fMRI study of the word-frequency effect. *Front. Behav. Neurosci.* **12**, 1–12 (2018).

33. Lin, Z., Tam, F., Churchill, N. W., Schweizer, T. A. & Graham, S. J. Tablet technology

for writing and drawing during functional magnetic resonance imaging: A review.

Sensors (Switzerland) **21**, 1–22 (2021).

34. Martínez, M., Villagra, F., Castellote, J. M. & Pastor, M. A. Kinematic and kinetic

patterns related to free-walking in parkinson's disease. *Sensors (Switzerland)* **18**,

(2018).

35. Huth, A. G., De Heer, W. A., Griffiths, T. L., Theunissen, F. E. & Gallant, J. L. Natural

speech reveals the semantic maps that tile human cerebral cortex. *Nature* **532**, 453–

458 (2016).

36. Martínez, M. *et al.* Trade-off between frequency and precision during stepping

movements: Kinematic and BOLD brain activation patterns. *Hum. Brain Mapp.* **37**,

1722–1737 (2016).

37. Oberhuber, M. *et al.* Functionally distinct contributions of the anterior and posterior

putamen during sublexical and lexical reading. *Front. Hum. Neurosci.* (2013).

doi:10.3389/fnhum.2013.00787

38. Varotto, S. F., Farah, H., Bogenberger, K., van Arem, B. & Hoogendoorn, S. P.

Adaptations in driver behaviour characteristics during control transitions from full-

range Adaptive Cruise Control to manual driving: an on-road study. *Transp. A Transp.*

Sci. **16**, 776–806 (2020).

39. Karimpoor, M. *et al.* A computerized tablet with visual feedback of hand position for

functional magnetic resonance imaging. *Front. Hum. Neurosci.* **9**, 1–13 (2015).

40.   Choi, M. H. *et al.* Increase in brain activation due to subtasks during driving: FMRI

study using new MR-compatible driving simulator. *J. Physiol. Anthropol.* **36**, 1–12

(2017).

41.   Haynes, W. I. A. & Haber, S. N. The organization of prefrontal-subthalamic inputs in

primates provides an anatomical substrate for both functional specificity and

integration: Implications for basal ganglia models and deep brain stimulation. *J.*

Neurosci. **33**, 4804–4814 (2013).

42.   Eickhoff, S. B. *et al.* Coordinate-based activation likelihood estimation meta-analysis

of neuroimaging data: A random-effects approach based on empirical estimates of

spatial uncertainty. *Hum. Brain Mapp.* **30**, 2907–2926 (2009).

43.   Turkeltaub, P. E., Eden, G. F., Jones, K. M. & Zeffiro, T. A. Meta-analysis of the

functional neuroanatomy of single-word reading: Method and validation. *Neuroimage*

16, 765–780 (2002).

44.   Lancaster, J. L. *et al.* Bias between MNI and talairach coordinates analyzed using the

ICBM-152 brain template. *Hum. Brain Mapp.* **28**, 1194–1205 (2007).

45.   Eickhoff, S. B. *et al.* Behavior, sensitivity, and power of activation likelihood estimation

characterized by massive empirical simulation. *Neuroimage* **137**, 70–85 (2016).

46.   Fox, P. T. *et al. User manual for GingerALE 2.3. Retrieved from*

http://brainmap.org/ale/manual.pdf (2013).

47.   Mazziotta, J. *et al.* A probabilistic atlas and reference system for the human brain:

International Consortium for Brain Mapping (ICBM). *Philos. Trans. R. Soc. Lond. B.*

Biol. Sci. **356**, 1293–322 (2001).

48.   Higashiyama, Y., Takeda, K., Someya, Y., Kuroiwa, Y. & Tanaka, F. The neural basis

of typewriting: A functional MRI study. *PLoS One* **10**, 1–20 (2015).

49.   Gould, R. L., Brown, R. G., Owen, A. M., Ffytche, D. H. & Howard, R. J. fMRI BOLD

response to increasing task difficulty during successful paired associates learning.

Neuroimage **20**, 1006–1019 (2003).

50.   Parkes, S. L., Bradfield, L. A. & Balleine, B. W. Interaction of insular cortex and

ventral striatum mediates the effect of incentive memory on choice between goal-

directed actions. *J. Neurosci.* **35**, 6464–6471 (2015).

51.   Sommer, S. & Pollmann, S. Putamen Activation Represents an Intrinsic Positive

Prediction Error Signal for Visual Search in Repeated Configurations. *Open*

Neuroimag. J. **10**, 126–138 (2016).

52.   Bellebaum, C., Koch, B., Schwarz, M. & Daum, I. Focal basal ganglia lesions are

associated with impairments in reward-based reversal learning. *Brain* **131**, 829–841

(2008).

53.   Cromwell, H. C. & Schultz, W. Effects of expectations for different reward magnitudes

on neuronal activity in primate striatum. *J. Neurophysiol.* **89**, 2823–2838 (2003).

54.   Haruno, M. & Kawato, M. Different neural correlates of reward expectation and reward

expectation error in the putamen and caudate nucleus during stimulus-action-reward

association learning. *J. Neurophysiol.* **95**, 948–959 (2006).

55.   Yamada, H., Matsumoto, N. & Kimura, M. Tonically Active Neurons in the Primate

Caudate Nucleus and Putamen Differentially Encode Instructed Motivational

Outcomes of Action. *J. Neurosci.* **24**, 3500–3510 (2004).

56.   Gardiner, T. W. & Nelson, R. J. Striatal neuronal activity during the initiation and

execution of hand movements made in response to visual and vibratory cues. *Exp.*

Brain Res. **92**, 15–26 (1992).

57.   Wymbs, N. F., Bassett, D. S., Mucha, P. J., Porter, M. A. & Grafton, S. T. Differential

665   Recruitment of the Sensorimotor Putamen and Frontoparietal Cortex during Motor

666   Chunking in Humans. *Neuron* **74**, 936–946 (2012).

667   58.   Cellini, N. Memory consolidation in sleep disorders. *Sleep Med. Rev.* **35**, 101–112

668   (2017).

669   59.   Andersen, K. W., Madsen, K. H. & Siebner, H. R. Discrete finger sequences are

670   widely represented in human striatum. *Sci. Rep.* **10**, 1–12 (2020).

671   60.   Romero, M. C., Bermudez, M. A., Vicente, A. F., Perez, R. & Gonzalez, F. Activity of

672   neurons in the caudate and putamen during a visuomotor task. *Neuroreport* **19**, 1141–

673   1145 (2008).

674   61.   Cohen, Y., Schneidman, E. & Paz, R. The geometry of neuronal representations

675   during rule learning reveals complementary roles of cingulate cortex and putamen.

676   *Neuron* **109**, 839-851.e9 (2021).

677   62.   Kimura, M., Kato, M. & Shimazaki, H. Physiological properties of projection neurons in

678   the monkey striatum to the globus pallidus. *Exp. Brain Res.* **82**, 672–676 (1990).

679   63.   Kimura, M., Rajkowski, J. & Evarts, E. Tonically discharging putamen neurons exhibit

680   set-dependent responses. *Proc. Natl. Acad. Sci. U. S. A.* **81**, 4998–5001 (1984).

681   64.   Kimura, M. The role of primate putamen neurons in the association of sensory stimuli

682   with movement. *Neurosci. Res.* **3**, 436–443 (1986).

683   65.   Tunik, E., Houk, J. C. & Grafton, S. T. Basal ganglia contribution to the initiation of

684   corrective submovements. *Neuroimage* **47**, 1757–1766 (2009).

685   66.   Kunimatsu, J., Maeda, K. & Hikosaka, O. The caudal part of putamen represents the

686   historical object value information. *J. Neurosci.* **39**, 1709–1719 (2019).

67.   Peters, A. J., Fabre, J. M. J., Steinmetz, N. A., Harris, K. D. & Carandini, M. Striatal

activity topographically reflects cortical activity. *Nature* **591**, 420–425 (2021).

68.   Haber, S. N., Fudge, J. L. & McFarland, N. R. Striatonigrostriatal Pathways in

Primates Form an Ascending Spiral from the Shell to the Dorsolateral Striatum. *J.*

Neurosci. **20**, 2369–2382 (2000).

69.   Björklund, A. & Dunnett, S. B. Dopamine neuron systems in the brain: an update.

Trends Neurosci. **30**, 194–202 (2007).

70.   Matsuda, W. *et al.* Single nigrostriatal dopaminergic neurons form widely spread and

highly dense axonal arborizations in the neostriatum. *J. Neurosci.* **29**, 444–53 (2009).

71.   Redgrave, P., Prescott, T. J. & Gurney, K. The basal ganglia: a vertebrate solution to

the selection problem? *Neuroscience* **89**, 1009–1023 (1999).

72.   Hikosaka, O. *et al.* Tutor Song Exposure Song Selectivity in the Pallial-Basal Ganglia

Song Circuit of Zebra Finches Raised Without Role of the Basal Ganglia in the

Control of Purposive Saccadic Eye Movements. *Physiol Rev* **80**, 953–978 (2000).

73.   Schultz, W. Behavioral theories and the neurophysiology of reward. *Annu. Rev.*

Psychol. **57**, 87–115 (2006).

74.   Redgrave, P., Gurney, K. & Reynolds, J. What is reinforced by phasic dopamine

signals? *Brain Research Reviews* **58**, 322–339 (2008).

75.   Bromberg-Martin, E. S., Matsumoto, M. & Hikosaka, O. Dopamine in Motivational

Control: Rewarding, Aversive, and Alerting. *Neuron* **68**, 815–834 (2010).

76.   Ashby, F. G., Turner, B. O. & Horvitz, J. C. Cortical and basal ganglia contributions to

habit learning and automaticity. *Trends in Cognitive Sciences* **14**, 208–215 (2010).

77.   Ohbayashi, M., Picard, N. & Strick, P. L. Inactivation of the dorsal premotor area

disrupts internally generated, but not visually guided, sequential movements. *J.*

Neurosci. **36**, 1971–1976 (2016).

712 78. Nachev, P., Kennard, C. & Husain, M. Functional role of the supplementary and pre-
713   supplementary motor areas. *Nature Reviews Neuroscience* **9**, 856–869 (2008).

714 79. Nambu, A., Tokuno, H. & Takada, M. Functional significance of the cortico-
715   subthalamo-pallidal 'hyperdirect' pathway. *Neurosci Res* **43**, 111–117 (2002).

716 80. Pineda-Pardo, J. A. *et al.* Static magnetic field stimulation of the supplementary motor
717   area modulates resting-state activity and motor behavior. *Commun. Biol.* **2**, 397
718   (2019).

719 81. Chen, L. L. & Wise, S. P. Neuronal activity in the supplementary eye field during
720   acquisition of conditional oculomotor associations. *J. Neurophysiol.* **73**, 1101–1121
721   (1995).

722 82. Morris, N. J. & Stein, C. M. Model-free linkage analysis of a quantitative trait. in
723   *Methods in Molecular Biology* **1666**, 327–342 (Humana Press Inc., 2017).

724 83. Haber, S. N. & Knutson, B. The Reward Circuit: Linking Primate Anatomy and Human
725   Imaging. *Neuropsychopharmacology* **35**, 4–26 (2009).

726 84. Leisman, G., Moustafa, A. & Shafir, T. Thinking, Walking, Talking: Integratory Motor
727   and Cognitive Brain Function. *Front. Public Heal.* **4**, 1 (2016).

728 85. Smittenaar, P., FitzGerald, T. H. B., Romei, V., Wright, N. D. & Dolan, R. J. Disruption
729   of Dorsolateral Prefrontal Cortex Decreases Model-Based in Favor of Model-free
730   Control in Humans. *Neuron* **80**, 914–919 (2013).

731 86. Cunnington, R., Windischberger, C., Deecke, L. & Moser, E. The preparation and
732   execution of self-initiated and externally-triggered movement: a study of event-related
733   fMRI. *Neuroimage* **15**, 373–385 (2002).

734 87. Aron, A. R. & Poldrack, R. A. Cortical and Subcortical Contributions to Stop Signal
735   Response Inhibition: Role of the Subthalamic Nucleus. *J. Neurosci.* **26**, 2424–2433

(2006).

88.   Arsalidou, M., Duerden, E. G. & Taylor, M. J. The Centre of the Brain : Topographical

Model of Motor , Cognitive , Affective , and Somatosensory Functions of the Basal

Ganglia. **3054**, 3031–3054 (2013).

89.   McFarland, N. R. & Haber, S. N. Thalamic relay nuclei of the basal ganglia form both

reciprocal and nonreciprocal cortical connections, linking multiple frontal cortical

areas. *J. Neurosci.* **22**, 8117–8132 (2002).

90.   Kim, H. F. & Hikosaka, O. Parallel basal ganglia circuits for voluntary and automatic

behaviour to reach rewards. *Brain* **138**, 1776–1800 (2015).

91.   Haber, S. N. Corticostriatal circuitry. *Dialogues Clin. Neurosci.* **18**, 7–21 (2016).

92.   Gurney, K., Humphries, M. D. & Redgrave, P. A New Framework for Cortico-Striatal

Plasticity: Behavioural Theory Meets In Vitro Data at the Reinforcement-Action

Interface. *PLoS Biol.* **13**, e1002034 (2015).

93.   Alexander, G. E. & Crutcher, M. D. Functional architecture of basal ganglia circuits:

neural substrates of parallel processing. *Trends in Neurosciences* **13**, 266–271

(1990).

94.   Huang, Y., Yaple, Z. A. & Yu, R. Goal-oriented and habitual decisions: Neural

signatures of model-based and model-free learning. *Neuroimage* **215**, 116834 (2020).

95.   Liljeholm, M., Dunne, S. & O'Doherty, J. P. Differentiating neural systems mediating

the acquisition vs. expression of goal-directed and habitual behavioral control. *Eur. J.*

Neurosci. **41**, 1358–1371 (2015).

96.   Eryilmaz, H. *et al.* Neural determinants of human goal-directed vs. habitual action

control and their relation to trait motivation. *Sci. Rep.* **7**, 1–11 (2017).

97.   Peterson, B. S. *et al.* Human basal ganglia volume asymmetries on magnetic

resonance images. *Magn. Reson. Imaging* **11**, 493–498 (1993).

98.   Grabenhorst, F. & Rolls, E. T. Value, pleasure and choice in the ventral prefrontal

cortex. *Trends Cogn. Sci.* **15**, 56–67 (2011).

99.   Hardwick, R. M., Forrence, A. D., Krakauer, J. W. & Haith, A. M. Time-dependent

competition between goal-directed and habitual response preparation. *Nat. Hum.*

Behav. **3**, 1252–1262 (2019).

100. Wong, A. L., Goldsmith, J., Forrence, A. D., Haith, A. M. & Krakauer, J. W. Reaction

times can reflect habits rather than computations. *Elife* **6**, (2017).

101. Amodio, D. M. The neuroscience of prejudice and stereotyping. *Nat. Rev. Neurosci.*

15, 670–682 (2014).

102. Vaquero, L. *et al.* Structural neuroplasticity in expert pianists depends on the age of

musical training onset. *Neuroimage* **126**, 106–119 (2016).

103. Cacioppo, S. *et al.* Intention understanding over T: A neuroimaging study on shared

representations and tennis return predictions. *Front. Hum. Neurosci.* **8**, 781 (2014).

104. Kim, W. *et al.* An fMRI study of differences in brain activity among elite, expert, and

novice archers at the moment of optimal aiming. *Cogn. Behav. Neurol.* **27**, 173–182

(2014).

105. Hernandez, L. F., Obeso, I., Costa, R. M., Redgrave, P. & Obeso, J. A. Dopaminergic

Vulnerability in Parkinson Disease: The Cost of Humans' Habitual Performance.

Trends Neurosci. **42**, 375–383 (2019).

106. Peterson, D. S. & Horak, F. B. Neural control of walking in people with parkinsonism.

Physiology **31**, 95–107 (2016).

107. Wu, T. *et al.* Neural correlates underlying Micrographia in Parkinson's disease. *Brain*

139, 144–160 (2016).

108.   de Wit, S., Barker, R. a, Dickinson, A. D. & Cools, R. Habitual versus goal-directed

785   action control in Parkinson disease. *J Cogn Neurosci* **23**, 1218–1229 (2010).

786   109.   Wu, T., Hallett, M. & Chan, P. Motor automaticity in Parkinson's disease. *Neurobiol.*

787   *Dis.* **82**, 226–234 (2015).

788   110.   Everitt, B. J. *et al.* Neural mechanisms underlying the vulnerability to develop

789   compulsive drug-seeking habits and addiction. *Philos. Trans. R. Soc. B Biol. Sci.* **363**,

790   3125–3135 (2008).

791   111.   Neal, D. T., Wood, W. & Quinn, J. M. Habits: A Repeat Performance. *Curr. Dir.*

792   *Psychol. Sci.* **15**, 198–202 (2006).

793   112.   Lally, P., van Jaarsveld, C. H. M., Potts, H. W. W. & Wardle, J. How are habits

794   formed: Modelling habit formation in the real world. *Eur. J. Soc. Psychol.* **40**, 998–

795   1009 (2010).

Acknowledgments

We are thankful to Blanca Caijao and Cristina Rubio (Universidad Rey Juan Carlos)

for their help in article search. MM is funded by a PhD Fellowship from Instituto de Salud

Carlos III (PFIS-FI720/00068). IO is funded by Instituto de Salud Carlos III (Miguel Servet,

CP18/00038) and AES-ISCIII (PI19/00298) from Ministry of Science and Innovation in Spain.

The funders played no role in the idea, design, data collection or analysis, decision to

publish or manuscript editing and writing.

Author Contributions

P.G. design, data collection, writing; M.M. design, data collection, analysis, writing;

P.R. writing; D.L. design, data collection, writing; I.O. design, data collection, writing;

Competing Interests statement

The authors declare no competing interest.

Figure Legends and Tables

Figure 1. Hypothesis diagram on striatal role in both everyday-life and experimental habits.
Activities part of daily life such as writing, reading, walking or driving were selected as
everyday-life habits (see Box 1 for task measurements and details) to expect a critical
striatal role in executing these habits forms. Similar striatal activities can be expected
compared to experimental paradigms commonly used in the cognitive science literature.

Figure 2. Everyday-life and experimental thresholded activation maps. **a)** Axial views for the
main regions. Overlap regions at the right column are shown in blue. Note that there exists
activation in the cerebellar cortex for the case of everyday-life studies but it is omitted here
for brevity. Z=52 view for the experimental studies is shown as an unthresholded map for
visualization purposes (ALE value ≈ 0.01) **(b)** 3D striatum reconstruction showing all the
activation that fall inside it. Overlap regions at the right column are shown in blue. **c)** 3D
striatum reconstruction showing the differential activation of the Everyday -life >
Experimental contrast (in green) and the Experimental > Everyday-life contrast (in red).

Figure 3. Everyday-life cognitive-motor subcategories thresholded activation maps. **a)** Axial
views for the main regions. Overlap regions at the right column are shown in blue. **b)** 3D
striatum reconstruction showing all the activation that fall inside it. Overlap regions at the
right column are shown in blue. **c)** Foci distribution of all the studies that fall in the striatum.
Note that some foci may correspond to the same study.

Figure 4. Experimental subcategories thresholded activation maps. **a)** Axial views for the
main regions. In this case there are no overlap regions at the right column. **b)** 3D striatum

reconstruction showing all the activation that fall inside it. In this case there are no overlap
regions at the right column. **c)** Foci distribution of all the studies that fall in the striatum. Note
that some foci may correspond to the same study.

Table 1. Everyday-life habits peaks coordinates in MNI152 space with region names from Harvard-Oxford atlas. Percentages of each brain region indicate how much activation from a cluster fall into such region. Coordinates of any activation comprising < 5% of its volume in a region are not shown for conciseness.

Cluster ID	Volume (mm³)	Brain regions (%)	MNI peak coordinates			ALE value
			x	y	z	
1	8792	Left Putamen (53.14%)	-24	-4	6	0.074
		Left Thalamus (28.84%)	-14	-22	10	0.04
		Left Pallidum (15.74%)	-23	-4	2	0.054
2	5320	Right Putamen (53.38%)	24	4	6	0.054
		Right Thalamus (20.45%)	16	-16	8	0.035
		Right Insula (13.08%)	32	14	8	0.037
		Right Pallidum (9.47%)	20	2	5	0.037
3	4840	Left Supplementary Motor Area (42.64%)	-6	0	52	0.035
		Right Paracingulate Gyrus (15.87%)	6	14	50	0.031
		Right Supplementary Motor Area (15.37%)	4	8	56	0.03
		Left Paracingulate Gyrus (13.06%)	0	8	53	0.027
4	3232	Left Precentral Gyrus (97.03%)	-54	-2	40	0.038
5	2416	Right-Cerebellum-Cortex (100%)	8	-64	-20	0.037
6	2368	Left Central Opercular Cortex (53.38%)	-46	-2	6	0.037
		Left Precentral Gyrus (18.58%)	-56	8	4	0.033
		Left Inferior Frontal Gyrus pars opercularis (18.58%)	-56	10	24	0.024
		Left Insula (7.09%)	-44	-1	4	0.03
7	2208	Right Precentral Gyrus (76.09%)	56	12	32	0.041
		Right Inferior Frontal Gyrus pars opercularis (18.48%)	54	13	31	0.036
		Right Middle Frontal Gyrus (5.43%)	46	13	30	0.013
8	1152	Right Inferior Frontal Gyrus pars opercularis (34.03%)	54	10	0	0.035
		Right Central Opercular Cortex (27.08%)	52	6	2	0.029
		Right Temporal Pole (13.89%)	54	10	-2	0.034
		Right Precentral Gyrus (11.81%)	-3	-14	54	0.022
		Right Planum Polare (9.03%)	53	5	-2	0.028
9	1032	Left Precentral Gyrus (42.64%)	-26	-8	56	0.04
		Left Superior Frontal Gyrus (36.43%)	6	11	54	0.029

Cluster ID	Volume (mm³)	Brain regions (%)	MNI peak coordinates			ALE value
			x	y	z	
		Left Middle Frontal Gyrus (20.93%)	47	12	31	0.025

Table 2. Experimental habits peaks coordinates in MNI152 space with region names from Harvard-Oxford atlas. Percentages of each brain region indicate how much activation from a cluster fall into such region. Coordinates of any activation comprising < 5% of its volume in a region are not shown for conciseness.

Cluster ID	Volume (mm³)	Brain regions (%)	MNI peak coordinates			ALE value
			x	y	z	
1	3344	Right Caudate (44.74%)	12	6	10	0.028
		Right Pallidum (20.57%)	-14	5	-6	0.025
		Right Accumbens (15.31%)	10	8	-6	0.03
		Right Putamen (10.05%)	14	9	-6	0.025
		Right Thalamus (6.46%)	11	-1	11	0.02
2	2088	Left Putamen (30.27%)	-14	5	-9	0.029
		Left Accumbens (27.97%)	-12	6	-10	0.033
		Left Caudate (20.69%)	-10	4	4	0.017
		Left Pallidum (12.26%)	14	5	-6	0.025
3	776	Left Putamen (98.97%)	-30	-8	-2	0.02
4	744	Right Insula (90.32%)	32	22	-4	0.023
		Right Orbito-Frontal Cortex (9.68%)	29	22	-7	0.019

Box 1. Description of most used paradigms to assess everyday-life (A) behaviours and experimental learning (B) in fMRI contexts.

Everyday-life	Main habitual outcomes	Task example
READING Real words are read aloud and compared to pseudo-words reading. Depending of the nature of the study, content of words may vary (neutral, emotional, language, non-words, etc) as well as naturality (own language vs new unknown words).	Reading own language with highly familiar/frequent words	 (1)
WRITING Participants are asked to handwrite by free natural writing, copying sentences/words, air-writing (sensorimotor control), creative writing, naming or drawing different figures.	Writing own language, familiar words (compared to baseline or control conditions)	 (2) (3)
WALKING While lying down inside the scanner, participants generate walking programmes on different block conditions, including voluntary alternating strides of the lower limbs, ankle moves, stepping, rapid walking or self-paced vs externally-paced.	Walking naturally at self-paced conditions	 (4)
DRIVING MR-compatible driving simulator is often used where participants are asked to naturally drive with different environmental conditions (driving only, stopping at traffic lights, turning curves, avoidance of particular stimuli) and control conditions (driving with sub-task, sub-task only, passive viewing, resting).	Driving in a natural environment (without distractors or dual tasks)	 (5)

(1) Cummine et al., 2016; (2) Yang et al., 2018; (3) Karimpoor et al., 2015; (4) Martínez et al., 2016; (5) Choi et al., 2017.

B. Experimental learning	Main habitual outcomes	Task example
PROBABILISTIC CLASSIFICATION TASK (PCT) Participants are shown visual cues that are probabilistically related to outcomes. Participants gradually learn the cue–outcome relationships through trial-by-trial feedback	Optimal responses to stimuli	
TWO-STAGE REINFORCEMENT LEARNING TASK Choices at stage 1 determines which state is more probable to occur in Stage 2 that leads to reward. At each Stage 2, participants choose between two stimuli and discover whether they receive a reward or not.	Reward sensitivity without sensitivity to transition probabilities	
OUTCOME DEVALUATION TASK During the discrimination training, participants learn to discriminative cues related to right-left key presses with rewarded outcomes. During the test phase, some of these outcomes are devaluated and no longer worth any points. Participants are asked to press the correct key when a stimulus signals the still-valuable outcome or withhold responding if outcome is no longer valuable	Continued responding for devalued outcomes	
MOTOR SEQUENCE LEARNING Motor finger movements are repeated or alternated in high-frequent vs low-frequent sequences. Through repeated practice, participants learn the repeating sequences, as demonstrated by improvements in performance (speed and accuracy).	Accuracy differences between high-frequency and low frequency sequences	

(1) *Images obtained from Foerde et al., 2018; (2) Cellini, 2017*

Supplementary material

Striatal role in everyday-life and laboratory-developed habits

Pasqualina Guida[1,2,6]*, Mario Michiels[1,2,6]*, Peter Redgrave[5], David Luque[3,4], Ignacio Obeso[1,2]

* Co-first authors

[1]*CINAC, Hospital Universitario HM Puerta del Sur, Móstoles, Madrid, Spain*
[2]*CIBERNED, Instituto de Salud Carlos III, Madrid, Spain*
[3]*Departamento de Psicología Básica, Universidad Autónoma de Madrid, Madrid, Spain*
[4]*Departamento de Psicología Básica, Universidad de Málaga, Madrid, Spain*
[5]*Department of Psychology, University of Sheffield, Sheffield, S10 2TN, UK*
[6]*PhD program in Neuroscience, Universidad Autónoma de Madrid, 28029 Madrid, Spain*

Methods

The complete set of search terms were as follows:

(1) Everyday life:

(walking[MeSH Terms] OR walking[Title/Abstract]) OR (Driving behaviour[MeSH Terms] OR Driving behaviour[Title/Abstract]) OR (Driving behavior[MeSH Terms] OR Driving behavior[Title/Abstract]) OR (car driving[MeSH Terms] OR Car driving[Title/Abstract]) OR (writing[MeSH Terms] OR writing[Title/Abstract]) OR (handwriting[MeSH Terms] OR handwriting[Title/Abstract]) OR (reading[Title/Abstract] OR reading[MeSH Terms) AND (functional magnetic resonance imaging[MeSH Terms] OR fMRI[Title/Abstract]) AND ((basal ganglia[Title/Abstract] OR caudate[Title/Abstract]

OR putamen[Title/Abstract] OR striatum[Title/Abstract]) OR (basal ganglia[MeSH Terms] OR caudate nucleus[MeSH Terms] OR putamen[MeSH Terms] OR striatum[MeSH Terms]))

(2) Experimental:

("habit" OR "habits" OR "probabilistic classification" OR "weather prediction" OR "response learning" OR "instrumental conditioning" OR "instrumental learning" OR "reinforcement learning" OR "outcome devaluation" OR "sequential decision" OR "two step" OR "2 step") AND ("basal ganglia" OR "caudate" OR "putamen" OR "striatum") AND ("fMRI" OR "functional magnetic resonance imaging" OR "functional MRI") AND ("2017/06/22"[PDAT] : "3000/12/31"[PDAT])

Table S1. Studies with everyday-life habits included in the meta-analysis.

Study by domain	n = 1441	Foci	Task	Contrast	Statistical threshold
Writing					
Katanoda et al. (2001)	17	19	Write a name of an object or naming the object	Writing > naming	p < 0.001 voxel-level & p < 0.05 cluster-level corrected
Nakamura et al. (2002)	9	10	Upon auditory presentation subjects wrote kanji word.	Writing > Rest	p < 0.05 corrected for multiple comparison
Beeson et al. (2003)	12	29	Writing words or drawing circles	Writing words > drawing circles	p < 0.001 uncorrected
Hu Xing-yue et al. (2006)	10	11	Writing with a pencil	Main Effect of Writing with a pencil	p < 0.0001 uncorrected
Segal et al. (2011)	9	19	Writing or naming pictures of objects and drawing one loop per syllable of the object's name	Writing > naming plus loops	p < 0.05 corrected
Horovitz et al. (2013)	13	5	Writing, tapping, and zigzagging with each limb	Right-handwriting > other tasks	p < 0.001 FWE corrected
Erhard et al. (2014)	20	13	Creative writing	Main effects of creative writing in expert writers	p < 0.05 FWE corrected
Longcamp et al. (2014)	18	13	Writing of letters or digits	Writing > holding the pen still	p < 0.05 voxel-level FWE corrected
Potgieser et al. (2015)	16	21	Write a sentence or hand tapping	Right-handwriting > right-hand tapping	p < 0.001 voxel-level & p < 0.05 cluster-level FWE corrected & k≥8
Bisio et al. (2017)	7	25	Writing sentences	Writing > resting	p < 0.05 FWE corrected
Karimpoor et al. (2018)	12	38	Copying grocery lists, phone numbers or sentences	All handwriting tasks > resting	p < 0.05 FDR corrected
Yang et al. (2018)	34	29	Copying chinese characters	Writing high frequency > writing low frequency	p < 0.05 FDR corrected
Reading					
Bookheimer et al. (1995)	16	33	Reading words or naming objects	Main effect of read aloud	p < 0.001 corrected
Moore et al. (1999)	8	7	Reading/naming	Main effect of reading words	p < 0.001 uncorrected
Mechelli et al. (2003)	20	10	Early/late reading processing	Early reading > Fixation	p < 0.05 corrected
Buchsbaum et al. (2005)	17	15	Reading/Hearing	Reading > Control	z > 2.33 & p < 0.01 cluster-corrected
Vigneau et al. (2005)	23	39	Reading words/ non reading letters	Read > Cross fixation	p < 0.001 uncorrected
Meschyan et al. (2006)	12	12	Reading different languages	Reading > Resting	p < 0.001 uncorrected & p < 0.05 corrected spatial extent threshold
Binder et al. (2006)	30	31	Non-word reading	Reading > Fixation	p < 0.00001 uncorrected & p < 0.05 corrected
Carreiras et al. (2007)	36	32	Reading/Lexical decision	Reading > Baseline	p < 0.05 corrected
Church et al. (2008)	50	37	Reading/Repeat	Reading > Repeat	p < 0.05 corrected with k≥24 voxels
Yarkoni et al. (2008)	28	21	Reading/Comprehension/lexical decision	Reading > Fixation	p < 0.05 uncorrected
Seghier et al. (2008)	43	5	Reading aloud/fixation	Reading > Fixation	p < 0.05 corrected
Seghier et al. (2010)	28	5	Reading aloud/naming	Reading > Fixation	p < 0.05 corrected
Oberhuber et al. (2013)	25	6	Reading/naming	Reading > Naming	p < 0.05, cluster level FWE-corrected
Vannest et al. (2013)	49	11	Reading/Generate words	Reading > Generate words	p ≤ 0.01 voxel-level corrected & t ≥ 7.5, cluster size 30

Hsu et al. (2015)	24	35	Reading emotional or neutral sentences	Reading > Resting	p < 0.005 voxel-level & p < 0.05 cluster-level FDR corrected
Rueckl et al. (2015)	84	18	Reading/Hearing	Reading > Resting	p < 0.001 FDR corrected
Cummine et al. (2016)	15	19	Reading words/pseudowords	Main effect of reading real words	p < 0.001 uncorrected
Oberhuber et al. (2016)	26	22	Reading words/ pseudowords or naming object/ colors	Reading words > other conditions	p < 0.05 FWE corrected
Cheema et al. (2018)	19	3	Reading words / non words	Reading > Control	p < 0.0001 uncorrected

Walking

Ciccarelliet al. (2005)	16	9	Right-left foot passive movement	Passive > Active movement	p < 0.05 corrected
Christensen et al. (2007)	18	10	Externally generated movement with or without visual feedback	Conjunctions of externally generated movements (regardless of feedback)	p < 0.05 FDR corrected
la Fougère et al. (2010)	16	14	Real locomotion	Walking > Resting condition	p < 0.05 FDR corrected
Trinastic et al (2010)	8	24	Active ankle dorsiflexion	Ankle dorsiflexion > Plantarflexion	p < 0.05
Swinnen et al. (2010)	14	49	90° out-of-phase versus iso task	90° Left > Iso directional Left	p < 0.001 FDR corrected
Toyomura et al. (2012)	12	9	Self-paced condition	Self > Externally paced	p < 0.0001 & k≥10 uncorrected; p < 0.05 FWE corrected
Sauvagea et al. (2013)	12	29	Speed execution task	Fast execution > low execution	p < 0.05 corrected
Martinez et al. (2014)	19	20	Stepping condition	Main effect of walking	p < 0.001 FDR corrected
Lukas Jaeger et al. (2014)	20	24	Active stepping	Main effect active walking	p < 0.001 cluster-level corrected & k≥42 voxels
Noble et al. (2014)	11	24	Bilateral plantarflexion exertion against physical resistance	Bilateral > Unilateral	p < 0.01 FWE corrected
Martín et al. (2016)	19	16	Pseudo-gait	Main effect of stability of stepping frequencies	p < 0.05 FWE corrected
Marchal et al. (2019)	20	7	Virtual gait task with doorway	Gait doorway > Walkway doorway	p < 0.001 uncorrected
Peters et al. (2019)	22	13	Ankle Task	Ankle main effect	p < 0.005 uncorrected
Allali G et al. (2019)	326	6	Varying walking speeds	Rapid walking > Normal walking speed	p < 0.05 cluster-level corrected

Driving

Uchiyama et al. (2003)	21	18	Driving simulator task	Active > Passive driving	p < 0.05 cluster-level FWE corrected
Graydon et al. (2004)	6	70	Driving simulator task	Simulated driving > Fixation	p < 0.05 corrected
Spiers et al. (2007)	20	8	The getaway game	Turning L > Turning R	p < 0.001 uncorrected & k≥5
Callan et al. (2009)	14	18	In-car video assist system	Driver's perspective with truck blocking viewing > driver's perspective occluded with a video from the perspective of the camera	p < 0.05 FDR corrected
Hsieh et al. (2009)	28	23	Static Load Paradigm	Driving video no distractor > fixation	p < 0.0001 corrected
Chein et al. (2011)	40	2	Stoplight driving game	Drive peer > alone	p < 0.05 FWE corrected
Uchiyama et al. (2012)	18	17	Driving simulator task (with dual task conditions)	Driving main effect	p < 0.001 voxel level & p < 0.05 cluster level corrected
Chung et al. (2014)	16	22	Driving simulator task	Driving only > Driving with task	p < 0.05 corrected

| Choi et al. (2017) | 15 | 25 | Driving simulator task | Driving with subtask condition > Control | p < 0.05 FDR corrected |

FWE: family-wise error; FDR: false-discovery rate

Table S2. Studies with experimental habits included in the meta-analysis.

Study by domain	n = 973	Foci	Task	Contrast (short)	Statistical threshold
Probabilistic learning					
Poldrack et al. (1999)	8	15	Probabilistic learning task	Task > control (perceptual-motor)	p < 0.05 corrected with Gaussian Random Field Theory
Poldrack et al. (2001)	13	25	Probabilistic learning task	Task > control (perceptual-motor)	p < 0.005 uncorrected & k≥5 voxels
Aron et al. (2004)	15	9	Probabilistic learning task	Task > baseline	p < 0.05 FDR corrected
Delgado et al. (2005)	17	2	Probabilistic learning task	Condition and time interaction	p < 0.0001 uncorrected & k≥5 voxels
Fera et al. (2005)	18	5	Probabilistic learning task	Task > control (perceptual-motor)	p < 0.05 uncorrected
Aron et al. (2006)	8	9	Probabilistic learning task	Task > control (perceptual-motor)	z > 2.3 voxel-level & p < 0.01 cluster-level corrected with Gaussian Random Field Theory
Foerde et al. (2006)	14	2	Probabilistic learning task	Modulation by performance (accuracy)	p < 0.05 corrected with Gaussian Random Field Theory
Glascher et al. (2010)	18	1	2-step probabilistic learning task	RPE	p < 0.006 corrected & k≥10
Celone et al. (2011)	19	8	Probabilistic learning task	Task > control (perceptual-motor)	p < 0.01 voxel-level & p < 0.05 cluster-level corrected
Wunderlich et al. (2012)	21	17	2-step probabilistic learning task	Trained > Planning (RPE)	p < 0.05 FWE corrected & k≥5
Soares et al. (2012)	12	10	Probabilistic learning task	Stress > control	p < 0.05 FWE corrected
Schwabe et al. (2013)	75	14	Probabilistic learning task	Task > control (perceptual-motor)	p < 0.05 FWE corrected & k≥5
Lee et al. (2014)	22	5	2-step probabilistic learning task	RPE	p < 0.05 FWE corrected
Deserno et al. (2015)	29	7	2-step probabilistic learning task	Model-free component	p < 0.05 corrected
Doll et al. (2015)	20	5	2-step probabilistic learning task	RPE	p ≤ 0.0005 cluster-level FWE corrected
Dunne et al. (2016)	17	3	Probabilistic learning task	RPE	p < 0.05 corrected
Oh-Descher et al. (2017)	25	3	Probabilistic learning task	High time pressure > low time pressure (Positive subjective sum of evidence)	p < 0.005 voxel-level & p < 0.05 cluster-level corrected
Nebe et al. (2017)	146	35	2-step probabilistic learning task	RPE	p < 0.05 FWE corrected
Erdeniz et al. (2019)	19	15	Probabilistic learning task	CS familiar > CS novel	p < 0.005 uncorrected
Mas-Herrero et al. (2019)	20	2	2-step probabilistic learning task	RPE (pseudofeedback)	p < 0.001 corrected
Others					
Gottfried et al. (2003)	13	7	Associative learning	Outcome devaluation	p < 0.001 uncorrected
Iaria et al. (2003)	7	5	Maze navigation	Task > control (perceptual-motor)	p < 0.05 corrected
Lehéricy et al. (2005)	14	3	Motor sequence learning	Modulation by task experience	p < 0.0001 voxel-level & p < 0.05 cluster-level corrected
Forstmann et al. (2008)	19	3	Moving dots task	Speed > accuracy	p < 0.001 corrected

Fernández-Seara et al. (2009)	14	30	Motor sequence learning	Late > early	p < 0.01 uncorrected
Tricomi et al. (2009)	15	4	Discriminative learning	Late > early	p < 0.001 uncorrected & k≥5
Worthy et al. (2010)	18	1	State maximizing	State-change uncertainty	p < 0.05 corrected
Steele et al. (2010)	15	14	Motor sequence learning	Late > early	p < 0.001 uncorrected & k≥100
Brovelli et al. (2011)	14	3	Discriminative learning	P (correct | past observations) > P (correct by chance)	P < 0.05 FWE corrected
Beierholm et al. (2011)	23	11	Sequential choice	Model-free component	p < 0.001 uncorrected & k≥5
De Wit et al. (2012)	24	3	Discriminative learning	Negative predictors of goal-directed action (outcome devaluation)	p < 0.005 uncorrected & k≥20
Etchamendy et al. (2012)	15	3	Maze navigation	Q-signal, learners	p < 0.05 corrected
Liljeholm et al. (2012)	15	4	Sequential choice	Decreased activity in the R-O but not int the S-R	p < 0.05 corrected with cluster size thresholding
Liljeholm et al. (2015)	19	13	Sequential choice	Modulation by devaluation insensitivity	p < 0.05 cluster-level corrected
Fermin et al. (2016)	18	9	Grid-Sailing	Previous S-R learning – Model-based conditions	p < 0.0001 uncorrected
Eryilmaz et al. (2017)	72	5	Discriminative learning	Incongruent > standard cue	p < 0.001 voxel-level & p < 0.05 cluster-level corrected
Van Steenbergen et al. (2017)	19	5	Discriminative learning	Cue-driven > Value-driven action control	p < 0.001 uncorrected & k≥15
Zwosta et al. (2018)	53	29	Discriminative learning	Decrease reward and no reward (Contingency degradation)	p < 0.001 voxel-level & p < 0.05 cluster-level corrected
Watson et al. (2018)	23	8	Discriminative learning	Slip of action trials > still-valued responses	p < 0.05 corrected
Anggraini et al. (2018)	27	16	Maze navigation	Model-free component	p < 0.001 voxel-level & p < 0.05 cluster-level corrected

RPE: reward prediction error; CS: conditioned stimuli; Q-signal: quality of state-action pairs; S-R: stimulus-response; R-O: response-outcome; FWE: family-wise error; FDR: false-discovery rate

Figure S1. PRISMA chart describing the steps followed in the selection of studies.