Truncation of IFT80 causes early embryonic loss in cattle

M. Sofía Ortega*, Derek M. Bickhart*, Kelsey N. Clark, Daniel J. Null, Jana L. Hutchison,

Jennifer C. McClure, John B. Cole[†]

* = co-first author

[†] = corresponding author

[1]Cell Wall Biology and Utilization Research Laboratory, U.S. Dairy Forage Research Center,

Agricultural Research Service, United States Department of Agriculture, Madison, WI, 53706

[2]Division of Animal Sciences, College of Agriculture, Food, and Natural Resources, University

of Missouri, Columbia, 65211

[3]Animal Genomics and Improvement Laboratory, Henry A. Wallace Beltsville Agricultural

Research Center, Agricultural Research Service, United States Department of Agriculture,

Beltsville, MD, 20705-2350

ABSTRACT

Recessive alleles represent a substantial risk in populations that have undergone significant

bottleneck events. Analyzing millions of genotypes and hundreds of sequenced bulls, we

identified one pernicious recessive variant that achieved 1.66% frequency in the US Holstein

population and confirmed loss of embryonic development via CRISPR-Cas9 editing. In the

process, we identified the developmental significance of two protein domains of the *IFT80* gene

in the embryonic development of cattle.

MAIN

Known lethal recessive alleles account for substantial economic losses to cattle breeders[1], and 20

such defects[2] are routinely tracked in the US population of 9.4 million dairy cows. Economic

losses resulting from embryonic death and stillbirths caused by these alleles have been estimated

to cost farmers at least $11 million per year[1] in the US and approximately £1.8 million annually

in the UK. The advent of artificial insemination techniques has exacerbated the problem by

increasing the frequency of recessive lethal alleles in commercial herds. The HH2 allele, which

accounts for annual fertility losses of more than $2 million in the US alone, is present in 1.66%

of the domestic Holstein population[3] and has a significant effect on cow fertility in the form of

early embryonic losses. The patrilineal nature of cattle reproduction and the availability of large

numbers of genotyped animals provides an opportunity to identify and validate such recessive

mutations and measure their potential impacts on other species.

We first sought to refine the HH2 genomic locus to narrow the window for variant discovery in

subsequent sequencing analysis (Fig 1a). Our analysis was aided by the recent release of a high-

quality reference assembly for taurine cattle[4] which had over 100-fold more contiguity than the

previously used UMD3.1 reference[5]. Using imputed SNP genotypes[6] derived from more than 3.7

million SNP genotypes of commercial cattle mapped to the new reference assembly, we were

able to reduce the size of the locus from 8.9 Mbp to 1.5 Mbp. This reduced the number of

candidate genes that were previously hypothesized to be causal for the lethality of the haplotype

by 72%, leaving only seven genes in the refined window out of 25 original candidates.

Separate, non-overlapping datasets were used for causal variant discovery and validation. The

discovery dataset included whole-genome sequence data for 9 carriers and 449 non-carriers from

the Animal Genomics and Improvement Laboratory (Beltsville, MD), the Collaborative Dairy

DNA Repository (Madison, WI) and the 1000 Bull Genomes Project[7]. A concordance analysis

was performed in order to identify candidate causal variants within the haplotype locus, and one

INDEL (Fig 1b) that was > 99% concordant with known carrier status was identified. One carrier

animal was predicted to not have a heterozygous copy of the INDEL, but manual review of read

pileups on the variant site identified the INDEL at a frequency of 25% which was below the

threshold of initial detection (data not shown). We interpreted this result as complete

concordance and attributed the lower proportion of variant reads to a bias in sampling the variant

haplotype during sequencing library preparation. The concordant INDEL was located on

btau1:107,172,615 and was found to induce a frameshift in the *IFT80* gene's 11th exon that led

to early truncation of the protein product (Supplementary Figure 1). *In silic*o translation of the

expected protein product of the edited transcript suggested an early truncation of the IFT80

protein, resulting in a protein of 385 amino acids (49.5 % of total) compared to the wild type.

The early termination of the gene interrupts one of seven WD 40-repeat containing domains

(Pfam: SSF50978) and a predicted weak polyampholyte domain

(https://mobidb.bio.unipd.it/A0A4W2EVR7), both of which likely impact the intra-flagellar

transport of materials in the cell[8]. Analysis of potential founders identified the bull Willowholme

Mark Anthony (73HO0219, born 1975) as a likely candidate. Pedigree, SNP and DNA sequence

data show that he is the founder with respect to the SNP haplotype, but he is not a carrier of the

actual *IFT80* variant, which appears to have arisen in one of his daughters (Elysa Anthony Lea,

HOCANF000003628269, born 1981) or grand-daughters (Comestar Laurie Sheik-ET,

HOCANF000004425038, born 1986). Neither descendant is genotyped or sequenced and DNA

is not available, so the true founder cannot be identified.

We hypothesized that the early termination of the *IFT80* gene was the causal variant of the

phenotype given its perfect concordance and the potential for the mutation to have a high impact

on downstream protein function. To induce the phenotype, we used a CRISPR-Cas9 gene editing

system (see Methods) to induce double-strand breaks at different exons at or upstream of the

candidate causal variant site (Fig 2a). In the first model, *IFT80* was truncated at exon 2, and in a

second model, *IFT80* was truncated at exon 11 (variant site). In both cases, treated embryos

failed to progress past the 8- to 16-cell transition period (Fig 2b, 2c; $P < 0.0001$), and

degenerated after this developmental stage. As alternative splicing could potentially recover

IFT80 protein function, we quantified the expression at different exons of *IFT80* transcript. For

embryos edited at exon 2, expression was measured at exons 1, 3–4 and 5–7 (before and after

edit site; Fig 2d). For the embryos edited at exon 11 (exon with causal mutation), *IFT80*

expression was determined at exons 9–10 and 12 (before and after the edited exon; Fig 2e) by

qPCR assays. Upstream regions of edited sites had no difference in *IFT80* expression (Exon 1, P

= 0.44; Exon 9-10, P = 0.62), between edited and non-edited embryos. Conversely, in both

models (truncation at exon 2 or 11) downstream exons of edited site, had significantly lower

IFT80 expression (P < 0.0001) in edited embryos than non-edited embryos, suggesting early

termination of the transcript due to the mutation or the altered splicing of these exons from the

transcript (Fig 2d). Differential expression was most pronounced in exon 12, in which the edited

cell line had 62.5% lower fold expression of the exon than the wildtype.

Mutations to the *IFT80* gene are linked to asphyxiating thoracic dystrophy 2 (ATD2) disease in

human[9] and hypomorphic expression of the gene has been shown to result in low postnatal

survival in gene-trap mice[10]. Previous surveys have speculated as to when *IFT80* gene expression

is essential during development[10], and our results suggest that expression is needed to progress

past the 8- to 16-cell developmental phase. This is consistent with previous RNA surveys of

cattle preimplantation embryonic development[11], and with the known fertility effects of HH2 in

cattle[1,3]. The results of editing experiments suggest that the carboxy-terminal WD 40-repeat or

the polyampholyte domains are essential for embryonic development. Recent studies suggest that

deletion of *IFT80* disrupt FGF2 signaling pathway, affecting cell proliferation, and self-renewal

of odontoblasts[12]. In the preimplantation mammalian embryo, FGF2 signaling is essential for

lineage commitment and blastocyst formation[13,14], in which disruption of FGF2 signaling

coincides with the developmental arrest observed in this study.

Using genotypes from millions of Holstein cattle and CRISPR-Cas9 gene editing, we were able

to track and validate a recessive lethal allele in the *IFT80* gene, respectively. This approach

conclusively identified the target mutation in such a way that it provided further insight into the

function of the gene in development of other mammalian species. There is a further economic

benefit to this discovery, as identification of the exact causal mutation underlying this haplotype

will benefit dairy producers by allowing them to avoid carrier-to-carrier matings which result in

pregnancy losses responsible for ~$437,000/year in US Holsteins. Concordance analysis and

molecular validation on naturally occurring genetic variants represent a potent tool for ongoing

efforts to translate genotype to phenotype in the field of Quantitative Genetics.

ONLINE METHODS

Haplotype discovery

Possible autosomal recessive mutations are routinely tracked during the imputation step of the

national genomic evaluations calculated by the Council on Dairy Cattle Breeding (CDCB;

Bowie, MD). Genotypes from 47 different arrays are currently stored in the CDCB database,

with SNP counts ranging from 2,900 to 777,962, and are imputed to a set of 79,276 (80k) for

genomic evaluation. No single chip includes all these markers, regardless of density, and

genomic calculations require that genotypes are available for all 80k SNP. Chromosomes are

phased into maternal and paternal haplotypes and the genome is divided into 100-SNP segments.

Inheritance of those segments is traced through the population and missing SNP are filled in

using a segment from the pedigree when available, and using the most common segment in the

population when pedigree information is not available[3]. Most males have complete pedigrees

available for many generations, but many females do not. For this study, haplotyping was

performed using the program Findhap version 3[15]. Haplotypes that either never appear in the

homozygous state, or which appear with far lower frequency than expected, are flagged as

possible recessives[3]. Such haplotypes are then tested for effects on fertility and stillbirth.

Founder animals are identified from the pedigree when possible. Currently, 26 such conditions

are tracked in the U.S. dairy cattle population

(https://www.ars.usda.gov/ARSUserFiles/80420530/Publications/ARR/Haplotype tests_ARR-

Genomic5.pdf), and causal mutations are known for all but the HH2 haplotype.

In April of 2011 a haplotype affecting fertility was identified on *Bos taurus* autosome (BTA) 1 in

the region of 92−97 Mbp and designated as HH2 (Figure 1a). At that time, a second haplotype on

BTA 1 at 101−106 Mbp with a similar carrier list, lack of homozygotes and negative effects on

fertility also was identified. When a new SNP list was introduced in August 2011 the haplotype

in the 101−106 Mbp region was no longer flagged as a potential lethal. A new 61,000 SNP list

was introduced in December 2013, and the haplotype segment upstream from HH2 again

appeared on the list of possible lethal haplotypes. In April 2020 the location of HH2 was

changed to correspond with the upstream segment based on results from fine-mapping described

below. Haplotype blocks are formed using fixed numbers of SNP, rather than LD boundaries, so

it can sometimes be difficult to narrow the candidates down to a single haplotype. In this case,

both segments appear to track the causal variant with high concordance.

Whole-genome sequencing, variant calling, and annotation

The dataset used for causal variant discovery and validation included whole-genome sequence

data for 9 carriers and 763 non-carriers, including animals from the Ayrshire, Holstein, and

Jersey breeds sequenced by the Animal Genomics and Improvement Laboratory (Beltsville,

MD), the Collaborative Dairy DNA Repository (Madison, WI), and the 1000 Bull Genomes

Project[7]. Sequencing depth for these animals averaged 14.7x. Sequence data were aligned to the

ARS-UCD1.2 reference genome[4] using BWA-MEM as implemented in bwa version 0.7.15-

r1140[16] and converted to the binary Sequence Alignment/Map format by samtools version 1.3[17].

1000 Bull Genomes run8 data were processed using version 4 of the Genome Analysis Toolkit[18],

and included genotyping, variant recalibration for SNP and INDEL, removal of individuals that

failed QC, and filtering of monomorphic alleles. The resulting VCF file was filtered to remove

reads with QS < 10 and those identified as "LOWQUAL". Sequence data variants were

identified using the Samtools v1.9 and BCFtools v1.9[19] and reads passing edits were aligned

against version ARS-UCD1.2 of the bovine genome[4]. The list of filtered variants was annotated

with the SNPEff[20] utility using the Ensembl annotation of the cattle reference genome (version

101). Variants listed as "High", "Medium" and "Low" priority within the haplotype region by

SNPEff were selected for further analysis. Impacts on predicted protein translation were

identified using custom scripts.

Production of IFT80$^{-/-}$ embryos

Guide RNA design

Guide RNAs (gRNAs) were designed to target exon 2 (Supplementary Table 1), or exon 11

(Supplementary Table 2) using the GPP portal from the Broad Institute

(https://portals.broadinstitute.org/gpp/public/analysis-tools/sgrna-design). To select gRNAs, two

criteria were used: (1) those with the highest on-target score after design based on the GPP portal

and the guide RNA checker portal (IDT technologies, San José, CA), and (2) those having the

least similarity with other bovine sequences containing a protospacer adjacent motif (PAM).

Selected gRNAs, as well as the universal 67mer tracrRNA® for form guide complexes were

ordered from IDT technologies.

The gRNAs and tracrRNA were resuspended to a concentration of 500 ng/μl. Each guide was

then annealed with equal amounts of tracrRNA by heating at 95 °C for 5 min and then cooled

down from 95 to 25 °C using a gradient of -1 °C/12 s in a Veriti 96-well gradient thermal cycler

(ThermoFisher, CA) to obtain a concentration of 250 ng/µl gRNA-tracrRNA complex. Prior to

zygote injection, gRNA-tracrRNA complexes were annealed to CAS9 mRNA (IDT

Technologies) by incubation at room temperature for 15 min. The final injection solution was 50

ng/µl each gRNAs complex and 20 ng/µl Cas9 mRNA. For each exon targeted, guides were

injected in pairs, to create a sizeable cut in the sequence and facilitate rapid genotyping on an

agarose gel. The expected cuts for guides targeting exon 2 was 82 bp and for those targeting

exon 11 were expected to generate a cut of 331 bp.

Production of embryos in vitro and zygote microinjection

Embryos were produced *in vitro* with a single sire known to be of high fertility *in vitro* as

previously described[21]. At the end of fertilization, putative zygotes (oocytes exposed to sperm)

were denuded from the surrounding cumulus cells and split in two groups: injected only with

CAS9 mRNA and injected with gRNA/CAS9 mRNA against *IFT80*. During the injection

procedure zygotes were maintained manipulation medium[22], and then transferred to four-well

dishes in groups of up to 50 zygotes in 500 µl of SOF-BE2[23], covered with 300 µl of mineral oil

per well at 38.5 °C in a humidified atmosphere of 5% (v/v) O_2 and 5% (v/v) CO_2. Percentage of

putative zygotes that cleaved was determined at day 3 of development (day 0 = day of

insemination) and blastocyst rate was estimated at day 8 of development. In addition, to

determine any developmental arrest, cell stage of all embryos was recorded at days 3, 5 and 8 of

development (Supplementary Figure 2).

Editing efficiency

To determine editing efficiency of the guides, embryos subjected to microinjection were

collected individually at day 8 of culture, placed in 6 µL of Embryo Lysis Buffer[22], and

subjected to 30 min at 60ºC, followed by 10 min at 85 ºC in a Veriti 96-well gradient thermal

cycler (ThermoFisher, CA). Lysed embryo solution was used as template DNA.

For genotyping, the region surrounding the target sequenced of the gRNAs was amplified by

endpoint PCR using the following primers (5'→3'): exon 2 F-

GGTTTCTTATCCTGCTTTCCATTC; R- GAAATTGAGTGTGAACCTTGGG, and for Exon

11 F- CACTGTTTAGGACTCTGCCT; R- CTCTCTGAGTAATGATACCATAGCA. The PCR

products were 639- and 620 bp for exon 2 (Supplementary Figure 3) and 11 (Supplementary

Figure 4), respectively. The PCR reaction, amplification (annealing temperature 53.4 °C), and

visualization of bands was performed as previously described[22]. Editing efficiency was

calculated from 3 separate *in vitro* embryo production runs, with 100 embryos per treatment

group per replicate. Editing efficiency for exon 2 was 26% bi-allelic, 32 % mono-allelic, and

42% non-edited, and for exon 11 was 50% bi-allelic, 23% mono-allelic, and 27% non-edited.

Nucleic acids isolation and RT-qPCR expression analysis

DNA and RNA were isolated from individual 8-cell embryos (65–72 h post insemination)

injected against *IFT80* or injected only with CAS9 mRNA. Embryos were collected and lysed

individually in 25 µl of RLT Lysis buffer (Qiagen, Valencia, CA), lightly vortexed, transferred

to a nano column (PuroSPIN™ Luna NANOTECH, Toronto, ON, CA), and centrifuged for 30 s

(all centrifugations were performed at 16,000 g). At this point, DNA is trapped in the column

and the flow-through contained the RNA. The column was placed in a new collection tube and

stored at room temp for later DNA purification. The flow-through containing RNA was mixed

with one volume of 70% ethanol, transferred to a new nano column, and centrifuged for 30 s.

Immediately after, 600 µl of buffer RW1 (Qiagen) were added to the column, and centrifuged for

30 s. Flow-through was discarded from the collection tube and 500 µl of buffer RPE (Qiagen)

were added to the column and centrifuged for 30 s. After discarding the flow-through, 500 µl of

80% ethanol were added to the column, and centrifuged for 2 min. To elute RNA, the column

was placed in a new microcentrifuge tube, and 15 µl of RNAse-free water were added directly to

the membrane of the column, incubated at room temperature for one minute and finally

centrifuged for 1 min. To complete DNA isolation, 500 µl of 70% ethanol were added to the

nano column containing DNA, followed by a 1-min centrifugation. Then, the column was placed

in a new microcentrifuge tube and 15 µl of EB buffer (Qiagen) preheated at 70 ºC were added

directly to the membrane of the column, incubated at room temperature for 10 min, followed by

1 min centrifugation to elute DNA. Embryos edit status was first assessed using PCR across the

site of the edit as described above, and the RNA from homozygous edited embryo cells were

selected for further analysis. Due to the small amount of total RNA in each cell, homozygous

embryos were pooled in groups of four prior to cDNA creation. Three pools of edited and control

embryos were analyzed in subsequent analyses. Two strand cDNA synthesis was conducted

using the high-capacity cDNA Reverse transcription kit (ThermoFisher) reverse transcriptase

following manufacturer instructions. Primers sequences for each locus were designed using the

PrimerQuest™ (IDT technologies). All primers were validated following procedures previously

described[24]. QPCR was conducted using the CFX384 Touch Real-Time System (Bio-Rad) using

10 µl of reaction mix as previously described[22]. The gene *SDHA* was chosen as a suitable

housekeeping gene for relative expression calculation in bovine embryos[25]. Fold changes were

calculated relative to the housekeeping gene ($2^{\Delta CT}$) as previously described[22,24].

Statistics

Differences in gene expression and development were analyzed by least-squares analysis of

variance of the ΔCT values using the GLM procedure of the Statistical Analysis System version

9.4 (SAS Institute Inc., Cary, NC, USA). Replicate and treatment were included as main effects

in the model.

Acknowledgments

Bickhart was supported by appropriated projects 8042-31000-001-00-D, "Enhancing Genetic

Merit of Ruminants Through Improved Genome Assembly, Annotation, and Selection" and

5090-31000-026-00-D, "Investigating Microbial, Digestive, and Animal Factors to Increase

Dairy Cow Performance and Nutrient Use Efficiency", of the Agricultural Research Service

(ARS) of the United States Department of Agriculture (USDA). Cole, Hutchison and Null were

supported by appropriated project 8042-31000-002-00-D, "Improving Dairy Animals by

Increasing Accuracy of Genomic Prediction, Evaluating New Traits, and Redefining Selection

Goals", of ARS, USDA. McClure was supported by appropriated project 5090-31000-026-00-D,

"Investigating Microbial, Digestive, and Animal Factors to Increase Dairy Cow Performance and

Nutrient Use Efficiency" of ARS, USDA. Clark was supported by USDA NIFA Grant 2019-

38420-28972.

The authors thank US dairy producers for providing phenotypic, genomic, and pedigree data

through the Council on Dairy Cattle Breeding under USDA Agricultural Research Service (ARS)

Material Transfer Research Agreement 58-8042-8-007. Access to whole-genome sequence data

from the Collaborative Dairy DNA Repository (Madison, WI) was provided under USDA-ARS

Material Transfer Research Agreement 58-8042-9-0010F. Access to whole-genome sequence

data from the 1000 Bull Genomes Project was provided under USDA-ARS Material Transfer

Agreement 14358. Authors also thank Bethany Redel and Joshua Benne from the University of

Missouri for their assistance with microinjections of bovine embryos. Mention of trade names or

commercial products in this article is solely for the purpose of providing specific information and

does not imply recommendation or endorsement by the US Department of Agriculture. The

USDA is an equal opportunity provider and employer.

Author Contributions

Cole and Ortega conceived the project. Null and Hutchison identified and annotated variant sites

within the haplotype region. Ortega and Clark designed CRISPR-Cas9 gRNAs, generated edited

embryos, and performed analysis on resulting embryos. Null and Bickhart ran analysis on

predicted variant sites and prepared figures for publication. McClure assisted with image

preparation. Ortega and Bickhart wrote the manuscript. All authors read and edited the

manuscript.

References

1. Cole, J. B., Null, D. J. & VanRaden, P. M. Phenotypic and genetic effects of recessive

haplotypes on yield, longevity, and fertility. *J. Dairy Sci.* **99**, 7274–7288 (2016).

2. Cole, J. B. *et al.* AIP Research Report Genomic5: Haplotype tests for economically

important traits of dairy cattle (update).

https://www.ars.usda.gov/ARSUserFiles/80420530/Publications/ARR/Haplotype%20tests_A

RR-Genomic5.pdf (2020).

3. VanRaden, P. M., Olson, K. M., Null, D. J. & Hutchison, J. L. Harmful recessive effects on

fertility detected by absence of homozygous haplotypes. *J. Dairy Sci.* **94**, 6153–6161 (2011).

4. Rosen, B. D. *et al.* De novo assembly of the cattle reference genome with single-molecule

sequencing. *Gigascience* **9**, (2020).

5. Zimin, A. V. *et al.* A whole-genome assembly of the domestic cow, Bos taurus. *Genome*

Biol. **10**, R42 (2009).

6. VanRaden, P. M. *et al.* Genomic imputation and evaluation using high-density Holstein
genotypes. *J. Dairy Sci.* **96**, 668–678 (2013).

7. Hayes, B. J. & Daetwyler, H. D. 1000 Bull Genomes Project to map simple and complex
genetic traits in cattle: Applications and outcomes. *Ann. Rev. Anim. Biosci.* **7**, 89–102 (2019).

8. Huang, D. W., Sherman, B. T. & Lempicki, R. A. Bioinformatics enrichment tools: paths
toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res.* **37**, 1–13
(2009).

9. Beales, P. L. *et al.* IFT80, which encodes a conserved intraflagellar transport protein, is
mutated in Jeune asphyxiating thoracic dystrophy. *Nat. Genet.* **39**, 727–729 (2007).

10. Rix, S., Calmont, A., Scambler, P. J. & Beales, P. L. An Ift80 mouse model of short rib
polydactyly syndromes shows defects in hedgehog signalling without loss or malformation
of cilia. *Hum. Mol. Genet.* **20**, 1306–1314 (2011).

11. Graf, A. *et al.* Fine mapping of genome activation in bovine embryos by RNA sequencing.
Proc. Natl. Acad. Sci. USA **111**, 4139–4144 (2014).

12. Yuan, X., Cao, X. & Yang, S. IFT80 is required for stem cell proliferation, differentiation,
and odontoblast polarization during tooth development. *Cell Death & Disease* **10**, 1–11
(2019).

13. Lanner, F. & Rossant, J. The role of FGF/Erk signaling in pluripotent cells. *Development*
137, 3351–3360 (2010).

14. Kuijk, E. W. *et al.* The roles of FGF and MAP kinase signaling in the segregation of the
epiblast and hypoblast cell lineages in bovine and human embryos. *Development* **139**, 871–
882 (2012).

15. VanRaden, Paul M., O'Connell, Jeffrey R., Wiggans, G. R. & Weigel, K. A. Genomic

evaluations with many more genotypes. *Genet. Sel. Evol.* **43**, 10 (2011).

16. Li, H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

arXiv:1303.3997 [q-bio] (2013).

17. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–

2079 (2009).

18. McKenna, A. *et al.* The Genome Analysis Toolkit: a MapReduce framework for analyzing

next-generation DNA sequencing data. *Genome Res.* **20**, 1297–1303 (2010).

19. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *GigaScience* **10**, giab008

(2021).

20. Cingolani, P. *et al.* A program for annotating and predicting the effects of single nucleotide

polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-

2; iso-3. *Fly (Austin)* **6**, 80–92 (2012).

21. Ortega, M. S. *et al.* Influences of sire conception rate on pregnancy establishment in dairy

cattle. *Biol. Reprod.* **99**, 1244–1254 (2018).

22. Ortega, M. S. *et al.* NANOG is required to form the epiblast and maintain pluripotency in the

bovine embryo. *Mol. Reprod. Devel.* **87**, 152–160 (2019).

23. Tríbulo, P., Rivera, R. M., Ortega Obando, M. S., Jannaman, E. A. & Hansen, P. J.

Production and Culture of the Bovine Embryo. in *Comparative Embryo Culture* (ed. Herrick,

J. R.) vol. 2006 115–129 (Springer New York, 2019).

24. Ortega, M. S., Kurian, J. J., McKenna, R. & Hansen, P. J. Characteristics of candidate genes

associated with embryonic development in the cow: Evidence for a role for WBP1 in

development to the blastocyst stage. *PLOS ONE* **12**, e0178041 (2017).

25. Goossens, K. *et al.* Selection of reference genes for quantitative real-time PCR in bovine

preimplantation embryos. *BMC Devel. Biol.* **5**, 27 (2005).

Figure Legends

Figure 1: A UCSC genome browser plot (a) shows the locations of prior Holstein Haplotype 2

(HH2) loci as determined in December 2013, April 2020 and August 2020. The *IFT80* gene is

highlighted by red dashed lines. Concordance analysis of variant sites against expected carrier

status (b) identified only one variant with > 99% concordance (highlighted in red). For genomic

context, the *IFT80* gene is drawn with exons (straight lines) and introns (diagonal lines) above

the concordance plot. Only variants predicted to have high functional effect by SNPEff are

highlighted by labels showing coordinate numbers.

Figure 2: A diagram of the *IFT80* gene (a) shows CRISPR-Cas9 edit sites (red triangles) and

qPCR primer design sites (blue boxes). Gene exons are represented by vertical lines, whereas

introns are represented by diagonal lines. Embryo counts for each developmental stage for exon

2 (b) and exon 3 (c) demonstrate that homozygous edit embryos (red) do not typically progress

past the 8-cell stage compared to vector Cas9 controls (blue). Single (*) and double (**)

asterisks indicate Student's T test p-values that are less than 0.001 and 0.0001, respectively.

Relative fold change of exons downstream of the exon 2 (d) and exon 11 (e) edit sites showed

significant decreases in edited embryos (red) compared to the vector Cas9 controls (blue).

 Figure 2