Truncation of IFT80 causes early embryonic loss in cattle

M. Sofía Ortega[1,†], Derek M. Bickhart[2,†], Kelsey N. Lockhart[1], Daniel J. Null[3], Jana L. Hutchison[3], Jennifer C. McClure[3], John B. Cole[3,4,*]

[1]Division of Animal Sciences, College of Agriculture, Food, and Natural Resources, University of Missouri, Columbia, 65211

[2]Cell Wall Biology and Utilization Research Laboratory, U.S. Dairy Forage Research Center, Agricultural Research Service, United States Department of Agriculture, Madison, WI, 53706

[3]Animal Genomics and Improvement Laboratory, Henry A. Wallace Beltsville Agricultural Research Center, Agricultural Research Service, United States Department of Agriculture, Beltsville, MD, 20705-2350

[4]Current address: URUS Group LP, Madison, WI, 53718

*John B. Cole

Email: john.cole@urus.org

Author Contributions: Cole and Ortega conceived the project. Null and Hutchison identified and annotated variant sites within the haplotype region. Ortega and Lockhart designed CRISPR-Cas9 gRNAs, generated edited embryos, and performed analysis on resulting embryos. Null and Bickhart ran analysis on predicted variant sites and prepared figures for publication. McClure assisted with image preparation. Ortega, Bickhart, and Cole wrote the manuscript. All authors read and edited the manuscript.

Competing Interest Statement: The authors declare no competing interests.

Keywords: dairy cattle, embryonic lethal, mammalian fertility, recessive allele.

Abstract

Recessive alleles represent a risk in populations that have undergone bottleneck events. We present a comprehensive framework for identification and validation of these genetic defects, including haplotype-based detection, variant selection from sequence data, and validation using knockout embryos. Holstein haplotype 2 (HH2), which causes embryonic death, was used to demonstrate the approach. HH2 was identified using a deficiency-of-homozygotes approach and confirmed to negatively affect conception rate and stillbirths. Five carriers were present in a group of 183 sequenced Holstein bulls selected to maximize the coverage of unique haplotypes. Three variants concordant with haplotype calls were found in HH2: a high-priority frameshift mutation resulting in a deletion, and two low-priority variants (1 synonymous variant, 1 premature stop codon). The frameshift in intraflagellar protein 80 (*IFT80*) was confirmed in a separate group of Holsteins from the 1000 Bull Genomes Project that shared no animals with the discovery set. *IFT80*-null embryos were generated by truncating the *IFT80* transcript at exon 2 or 11 using a CRISPR-Cas9 system. Abattoir-derived oocytes were fertilized *in vitro* and embryos were injected at the one-cell stage either with CRISPR-Cas9 complex (n=100) or Cas9 mRNA (control, n=100) before return to culture, and replicated 3 times. *IFT80* is activated at the 8-cell stage, and IFT80-null embryos arrested at this stage of development, which is consistent with data from mouse hypomorphs and HH2 carrier-to-carrier matings. This frameshift in IFT80 on chromosome 1 at

40   107,172,615 bp (p.Leu381fs) disrupts WNT and hedgehog signaling, and is responsible for the
41   death of homozygous embryos.

Significance Statement

Holstein haplotype 2 is an embryonic lethal present in 1.21% of the US Holstein cattle population,
and unrecognized carrier-to-carrier matings are responsible for >$2 million/year in additional
breeding expenses. A high-impact frameshift mutation in exon 11 of intraflagellar protein 80
(IFT80) was identified as the putative causal variant. Biallelic IFT80 knockout embryos were
produced *in vitro* and compared to wild-type embryos. IFT80-null embryos consistently arrested
at the 8-cell stage of development. The IFT80 protein expressed in knockout embryos had
substantially altered protein structure, resulting in a loss of functional domains. These results
validate the putative causal mutation observed in Holsteins. This system is a good model for
investigating possible causal variants that affect livestock fertility early in development.

52
Main Text

54
Introduction

56
Known lethal recessive alleles account for substantial economic losses to cattle breeders (1), and
20 such defects (2) are routinely tracked in the US population of 9.4 million dairy cows. Economic
losses resulting from embryonic death and stillbirths caused by these alleles have been estimated
to cost farmers at least $11 million per year in the US (1). The increased use of artificial
insemination on an already reduced genetic pool has exacerbated the problem by increasing the
rate at which recessive lethal alleles can be spread in commercial herds. The HH2 allele, which
accounts for annual fertility losses of more than $2 million in the US alone, is present in 1.66% of
the domestic Holstein population (3) and has a significant negative effect on fertility in the form of
early embryonic losses. The patrilineal nature of cattle reproduction and the availability of large
numbers of genotyped animals provides an opportunity to identify and validate such recessive
mutations and measure their potential impacts on other species.

68
Results and Discussion

70
We first refined the HH2 genomic locus to narrow the window for variant discovery in subsequent
sequencing analysis (Figure 1A). Our analysis was aided by the recent release of a high-quality
reference assembly for taurine cattle (4) which had over 100-fold more contiguity than the
previously used UMD3.1 reference (5). Using imputed SNP genotypes (6) derived from more than
3.7 million SNP genotypes of commercial cattle mapped to the new reference assembly, we were
able to reduce the size of the locus from 8.9 Mbp to 1.5 Mbp. This reduced the number of
candidate genes that were previously hypothesized to be causal for the lethality of the haplotype
by 72%, leaving only seven genes in the refined window out of 25 original candidates.

79
Separate, non-overlapping datasets were used for causal variant discovery and validation. The
discovery dataset included whole-genome sequence data for 9 carriers and 449 non-carriers from
the Animal Genomics and Improvement Laboratory (Beltsville, MD), the Collaborative Dairy DNA
Repository (Madison, WI) and the 1000 Bull Genomes Project (7). A concordance analysis was
used to identify candidate causal variants within the haplotype locus, and one INDEL (Figure 1B)
that was > 99% concordant with known carrier status was identified. One carrier animal was
predicted to not have a heterozygous copy of the INDEL, but manual review of read pileups on
the variant site identified the INDEL at a frequency of 25% which was below the threshold of initial
detection (data not shown). We interpreted this result as complete concordance and attributed the
lower proportion of variant reads to a bias in sampling the haplotype during sequencing library
preparation. The concordant INDEL was located on btau1:107,172,615 and was found to induce
a frameshift in the IFT80 gene's 11th exon that led to early truncation of the protein product

(Figure 2A). *In silico* translation of the expected protein product of the edited transcript suggested
an early truncation of the IFT80 protein, resulting in a protein of 385 amino acids (49.5% of total)
compared to the wild type (Figure 3D). The early termination of the gene interrupts one of seven
WD 40-repeat containing domains (Pfam: SSF50978) and a predicted weak polyampholyte
domain (https://mobidb.bio.unipd.it/A0A4W2EVR7), both of which likely impact the intra-flagellar
transport of materials in the cell (8). Analysis of potential founders identified the bull Willowholme
Mark Anthony (73HO0219, born 1975) as a likely candidate. Pedigree, SNP, and DNA sequence
data show that he is the founder with respect to the SNP haplotype, but he is not a carrier of the
actual IFT80 variant, which appears to have arisen in one of his daughters (Elysa Anthony Lea,
HOCANF000003628269, born 1981) or grand-daughters (Comestar Laurie Sheik-ET,
HOCANF000004425038, born 1986). Neither descendant is genotyped or sequenced and DNA is
not available, so the true founder cannot be identified.
104
We hypothesized that the early termination of the *IFT80* gene was the causal variant of the
phenotype given its perfect concordance and the potential for the mutation to have a high impact
on downstream protein function. To induce the phenotype, we used a CRISPR-Cas9 gene editing
system (see Methods) to induce double-strand breaks at different exons at or upstream of the
candidate causal variant site (Figure 2A). In the first model, IFT80 was truncated at exon 2, and in
a second model, IFT80 was truncated at exon 11 (variant site). In both cases, treated embryos
failed to progress past the 8- to 16-cell transition period (Figure 2A, 2c; P < 0.0001), and
degenerated after this developmental stage (Figure 3A). As alternative splicing could potentially
recover IFT80 protein function, we quantified the expression at different exons of IFT80 transcript.
For embryos edited at exon 2, expression was measured at exons 1, 3–4 and 5–7 (before and
after edit site; Figure 2D). For the embryos edited at exon 11 (exon with causal mutation), IFT80
expression was determined at exons 9–10 and 12 (before and after the edited exon; Figure 2E)
by qPCR assays. Upstream regions of edited sites had no difference in IFT80 expression (Exon
1, P = 0.44; Exon 9-10, P = 0.62), between edited and non-edited embryos. Conversely, in both
models (truncation at exon 2 or 11) downstream exons of edited site, had significantly lower
IFT80 expression (P < 0.0001) in edited embryos than non-edited embryos, suggesting early
termination of the transcript due to the mutation or the altered splicing of these exons from the
transcript (Figure 2D). Differential expression was most pronounced in exon 12, in which the
edited cell line had 62.5% lower fold expression of the exon than the wildtype (Figure 2E).
124
Mutations to the *IFT80* gene are linked to asphyxiating thoracic dystrophy 2 (ATD2) disease in
human (9) and hypomorphic expression of the gene has been shown to result in low postnatal
survival in gene-trap mice (10). Previous surveys have speculated as to when IFT80 gene
expression is essential during development (10), and our results suggest that expression is
needed to progress past the 8- to 16-cell developmental phase. This is consistent with previous
RNA surveys of cattle preimplantation embryonic development (11), and with the known fertility
effects of HH2 in cattle (1, 3). The results of editing experiments suggest that the carboxy-
terminal WD 40-repeat or the polyampholyte domains are essential for embryonic development.
Recent studies suggest that deletion of IFT80 disrupt FGF2 signaling pathway, affecting cell
proliferation, and self-renewal of odontoblasts (12). In the preimplantation mammalian embryo,
FGF2 signaling is essential for lineage commitment and blastocyst formation (13, 14), in which
disruption of FGF2 signaling coincides with the developmental arrest observed in this study.
137
Using genotypes from millions of Holstein cattle and CRISPR-Cas9 gene editing, we were able to
track and validate a recessive lethal allele in the IFT80 gene, respectively. This approach
conclusively identified the target mutation in such a way that it provided further insight into the
function of the gene in development of other mammalian species. There is a further economic
benefit to this discovery, as identification of the exact causal mutation underlying this haplotype
will benefit dairy producers by allowing them to avoid carrier-to-carrier matings which result in
pregnancy losses responsible for ~$437,000/year in US Holsteins. Concordance analysis and

molecular validation on naturally occurring genetic variants represent a potent tool for ongoing
efforts to translate genotype to phenotype in the field of quantitative genetics.
147
148 **Materials and Methods**
149
150 **Haplotype discovery**
151
Possible autosomal recessive mutations are routinely tracked during the imputation step of the
national genomic evaluations calculated by the Council on Dairy Cattle Breeding (CDCB; Bowie,
MD). Genotypes from 48 different arrays are currently stored in the CDCB database, with SNP
counts ranging from 2,900 to 777,962, and are imputed to a set of 78,964 SNP (80k) for genomic
evaluation. No single chip includes all these markers, regardless of density, and genomic
calculations require that genotypes are available for all 80k SNP. Chromosomes are phased into
maternal and paternal haplotypes and the genome is divided into 100-SNP segments. Inheritance
of those segments is traced through the population and missing SNP are filled in using a segment
from the pedigree when available, and using the most common segment in the population when
pedigree information is not available (3). Most males have complete pedigrees available for many
generations, but many females do not. For this study, haplotyping was performed using the
program Findhap version 3 (15). Haplotypes that either never appear in the homozygous state, or
which appear with far lower frequency than expected, are flagged as possible recessives (3).
Such haplotypes are then tested for effects on fertility and stillbirth. Founder animals are identified
from the pedigree when possible. Currently, 26 such conditions are tracked in the U.S. dairy
cattle population (20 genetic diseases and 6 coat-color and polled variants), and causal mutations
are known for all but the HH2 haplotype.
169
In April of 2011 a haplotype affecting fertility was identified on *Bos taurus* autosome (BTA) 1 in
the region of 92–97 Mbp and designated as HH2 (Figure 1A). At that time, a second haplotype on
BTA 1 at 101–106 Mbp with a similar carrier list, lack of homozygotes and negative effects on
fertility also was identified. When a new SNP list was introduced in August 2011 the haplotype in
the 101–106 Mbp region was no longer flagged as a potential lethal. A new 61,000 SNP list was
introduced in December 2013, and the haplotype segment upstream from HH2 again appeared
on the list of possible lethal haplotypes. In April 2020 the location of HH2 was changed to
correspond with the upstream segment based on results from fine-mapping described below.
Haplotype blocks are formed using fixed numbers of SNP, rather than LD boundaries, so it can
sometimes be difficult to narrow the candidates down to a single haplotype. In this case, both
segments appear to track the causal variant with high concordance.
181
182 **Whole-genome sequencing, variant calling, and annotation**
183
The dataset used for causal variant discovery and validation included whole-genome sequence
data for 9 carriers and 763 non-carriers, including animals from the Ayrshire, Holstein, and Jersey
breeds sequenced by the Animal Genomics and Improvement Laboratory (Beltsville, MD), the
Collaborative Dairy DNA Repository (Madison, WI), and the 1000 Bull Genomes Project (7).
Sequencing depth for these animals averaged 14.7x. Sequence data were aligned to the ARS-
UCD1.2 reference genome (4) using BWA-MEM as implemented in bwa version 0.7.15-r1140
(16) and converted to the binary Sequence Alignment/Map format by samtools version 1.3 (17).
1000 Bull Genomes run8 data were processed using version 4 of the Genome Analysis Toolkit
(18), and included genotyping, variant recalibration for SNP and INDEL, removal of individuals
that failed QC, and filtering of monomorphic alleles. The resulting VCF file was filtered to remove
reads with QS < 10 and those identified as "LOWQUAL". Sequence data variants were identified
using the Samtools v1.9 and BCFtools v1.9 (19) and reads passing edits were aligned against
version ARS-UCD1.2 of the bovine genome (4). The list of filtered variants was annotated with
the SNPEff (20) utility using the Ensembl annotation of the cattle reference genome (version
101). Variants listed as "High", "Medium" and "Low" priority within the haplotype region by SNPEff

were selected for further analysis. Impacts on predicted protein translation were identified using
custom scripts.
201
202 **Production of IFT80-/- embryos**
203
204 *Guide RNA design*
205
Guide RNAs (gRNAs) were designed to target exon 2 or exon 11 (Table 1), using the GPP portal
from the Broad Institute (https://portals.broadinstitute.org/gpp/public/analysis-tools/sgrna-design).
To select gRNAs, two criteria were used: (1) those with the highest on-target score after design
based on the GPP portal and the guide RNA checker portal (IDT technologies, San José, CA),
and (2) those having the least similarity with other bovine sequences containing a protospacer
adjacent motif (PAM). Selected gRNAs, as well as the universal 67mer tracrRNA® for form guide
complexes were ordered from IDT technologies.
213
The gRNAs and tracrRNA were resuspended to a concentration of 500 ng/µl. Each guide was
then annealed with equal amounts of tracrRNA by heating at 95 °C for 5 min and then cooled
down from 95 to 25 °C using a gradient of -1 °C/12 s in a Veriti 96-well gradient thermal cycler
(ThermoFisher, CA) to obtain a concentration of 250 ng/µl gRNA-tracrRNA complex. Prior to
zygote injection, gRNA-tracrRNA complexes were annealed to CAS9 mRNA (IDT Technologies)
by incubation at room temperature for 15 min. The final injection solution was 50 ng/µl each
gRNAs complex and 20 ng/µl Cas9 mRNA. For each exon targeted, guides were injected in pairs,
to create a sizeable cut in the sequence and facilitate rapid genotyping on an agarose gel. The
expected cuts for guides targeting exon 2 was 82 bp and for those targeting exon 11 were
expected to generate a cut of 331 bp.
224
225
226
227
228 *Production of embryos in vitro and zygote microinjection*
229
Embryos were produced in vitro with a single sire known to be of high fertility in vitro as previously
described (21). At the end of fertilization, putative zygotes (oocytes exposed to sperm) were
denuded from the surrounding cumulus cells and split in two groups: injected only with Cas9
mRNA and injected with gRNA/Cas9 mRNA against IFT80. During the injection procedure
zygotes were maintained manipulation medium (22), and then transferred to four-well dishes in
groups of up to 50 zygotes in 500 µl of SOF-BE2 (23), covered with 300 µl of mineral oil per well
at 38.5 °C in a humidified atmosphere of 5% (v/v) O_2 and 5% (v/v) CO_2. Percentage of putative
zygotes that cleaved was determined at day 3 of development (day 0 = day of insemination) and
blastocyst rate was estimated at day 8 of development. In addition, to determine any
developmental arrest, cell stage of all embryos was recorded at days 64.5, 71.5, 120 and 180 h of
development (Figure 2B, 2C and 3A).
241
242 *Editing efficiency*
243
244 To determine editing efficiency of the guides, embryos subjected to microinjection were collected
245 individually at day 8 of culture, placed in 6 µL of Embryo Lysis Buffer (22), and subjected to 30
246 min at 60ºC, followed by 10 min at 85 ºC in a Veriti 96-well gradient thermal cycler
247 (ThermoFisher, CA). Lysed embryo solution was used as template DNA.
248
For genotyping, the region surrounding the target sequenced of the gRNAs was amplified by
endpoint PCR using the following primers (5'→3'): exon 2 F-GGTTTCTTATCCTGCTTTCCATTC;
R- GAAATTGAGTGTGAACCTTGGG, and for Exon 11 F- CACTGTTTAGGACTCTGCCT; R-
CTCTCTGAGTAATGATACCATAGCA. The PCR products were 639- and 620 bp for exon 2

(Figure 3B) and 11 (Figure 3C), respectively. The PCR reaction, amplification (annealing
temperature 53.4 °C), and visualization of bands was performed as previously described (22).
Editing efficiency was calculated from 3 separate in vitro embryo production runs, with 100
embryos per treatment group per replicate. Editing efficiency for exon 2 was 26% bi-allelic, 32 %
mono-allelic, and 42% non-edited, and for exon 11 was 50% bi-allelic, 23% mono-allelic, and
27% non-edited.
259
260 ***Nucleic acids isolation and RT-qPCR expression analysis***
261
DNA and RNA were isolated from individual 8-cell embryos (65–72 h post insemination) injected
against IFT80 or injected only with Cas9 mRNA. Embryos were collected and lysed individually in
25 µl of RLT Lysis buffer (Qiagen, Valencia, CA), lightly vortexed, transferred to a nano column
(PuroSPIN™ Luna NANOTECH, Toronto, ON, CA), and centrifuged for 30 s (all centrifugations
were performed at 16,000 g). At this point, DNA is trapped in the column and the flow-through
contained the RNA. The column was placed in a new collection tube and stored at room temp for
later DNA purification. The flow-through containing RNA was mixed with one volume of 70%
ethanol, transferred to a new nano column, and centrifuged for 30 s. Immediately after, 600 µl of
buffer RW1 (Qiagen) were added to the column, and centrifuged for 30 s. Flow-through was
discarded from the collection tube and 500 µl of buffer RPE (Qiagen) were added to the column
and centrifuged for 30 s. After discarding the flow-through, 500 µl of 80% ethanol were added to
the column, and centrifuged for 2 min. To elute RNA, the column was placed in a new
microcentrifuge tube, and 15 µl of RNAse-free water were added directly to the membrane of the
column, incubated at room temperature for one minute and finally centrifuged for 1 min. To
complete DNA isolation, 500 µl of 70% ethanol were added to the nano column containing DNA,
followed by a 1-min centrifugation. Then, the column was placed in a new microcentrifuge tube
and 15 µl of EB buffer (Qiagen) preheated at 70 ºC were added directly to the membrane of the
column, incubated at room temperature for 10 min, followed by 1 min centrifugation to elute DNA.
Embryos edit status was first assessed using PCR across the site of the edit as described above,
and the RNA from homozygous edited embryo cells were selected for further analysis. Due to the
small amount of total RNA in each cell, homozygous embryos were pooled in groups of four prior
to cDNA creation. Three pools of edited and control embryos were analyzed in subsequent
analyses. Two strand cDNA synthesis was conducted using the high-capacity cDNA Reverse
transcription kit (ThermoFisher) reverse transcriptase following manufacturer instructions. Primers
sequences for each locus were designed using the PrimerQuest™ (IDT technologies). All primers
(Table 2) were validated following procedures previously described (24). qPCR was conducted
using the CFX384 Touch Real-Time System (Bio-Rad) using 10 µl of reaction mix as previously
described (22). The gene SDHA was chosen as a suitable housekeeping gene for relative
expression calculation in bovine embryos (25). Fold changes were calculated relative to the
housekeeping gene (2ΔCT) as previously described (22, 24).
292
293 **Statistics**
294
295 Differences in gene expression and development were analyzed by least-squares analysis of
296 variance of the ΔCT values using the GLM procedure of the Statistical Analysis System version
297 9.4 (SAS Institute Inc., Cary, NC, USA). Replicate and treatment were included as main effects in
298 the model.
299
300 **Data Availability**
301
302 **Performance and Pedigree Data.** The performance and pedigree data used to compute HH2
303 effects are controlled by the Council on Dairy Cattle Breeding (CDCB; Bowie, MD). Requests to
304 access CDCB data may be sent to João Dürr, CDCB Chief Executive Officer
305 (joao.durr@cdcb.us).
306

Genotype and Whole-Genome DNA Sequence Data. Access to SNP genotype data and whole-genome DNA sequence data for bulls held by the Collaborative Dairy DNA Repository (CDDR; Verona, WI) must be requested from Jay Weiker, CDDR Administrator (jweiker@naab-css.org).

1000 Bull Genomes Data. Access to whole-genome DNA sequence data from bulls distributed in Run7 is restricted to project members. Inquiries should be directed to Dr. Ben J. Hayes (b.hayes@uq.edu.au).

Acknowledgments

Bickhart was supported by appropriated projects 8042-31000-001-00-D, "Enhancing Genetic Merit of Ruminants Through Improved Genome Assembly, Annotation, and Selection" and 5090-31000-026-00-D, "Investigating Microbial, Digestive, and Animal Factors to Increase Dairy Cow Performance and Nutrient Use Efficiency", of the Agricultural Research Service (ARS) of the United States Department of Agriculture (USDA). Cole, Hutchison, and Null were supported by appropriated project 8042-31000-002-00-D, "Improving Dairy Animals by Increasing Accuracy of Genomic Prediction, Evaluating New Traits, and Redefining Selection Goals", of ARS, USDA. McClure was supported by appropriated project 5090-31000-026-00-D, "Investigating Microbial, Digestive, and Animal Factors to Increase Dairy Cow Performance and Nutrient Use Efficiency" of ARS, USDA. Clark was supported by USDA NIFA Grant 2019-38420-28972.

The authors thank US dairy producers for providing phenotypic, genomic, and pedigree data through the Council on Dairy Cattle Breeding under USDA Agricultural Research Service (ARS) Material Transfer Research Agreement 58-8042-8-007. Access to whole-genome sequence data from the Collaborative Dairy DNA Repository (Madison, WI) was provided under USDA-ARS Material Transfer Research Agreement 58-8042-9-0010F. Access to whole-genome sequence data from the 1000 Bull Genomes Project was provided under USDA-ARS Material Transfer Agreement 14358. The authors also thank Bethany Bauer and Joshua Benne from the University of Missouri for their assistance with microinjections of bovine embryos.

Mention of trade names or commercial products in this article is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the US Department of Agriculture. The USDA is an equal opportunity provider and employer.

References

1. J. B. Cole, D. J. Null, P. M. VanRaden, Phenotypic and genetic effects of recessive haplotypes on yield, longevity, and fertility. *Journal of Dairy Science* **99**, 7274–7288 (2016).

2. J. B. Cole, P. M. VanRaden, D. J. Null, J. L. Hutchison, S. M. Hubbard, "AIP Research Report GENOMIC5: Haplotype tests for recessive disorders that affect fertility and other traits" (2021).

3. P. M. VanRaden, K. M. Olson, D. J. Null, J. L. Hutchison, Harmful recessive effects on fertility detected by absence of homozygous haplotypes. *Journal of Dairy Science* **94**, 6153–6161 (2011).

4. B. D. Rosen, *et al.*, De novo assembly of the cattle reference genome with single-molecule sequencing. *Gigascience* **9** (2020).

5. A. V. Zimin, *et al.*, A whole-genome assembly of the domestic cow, Bos taurus. *Genome Biology* **10**, R42 (2009).

6. P. M. VanRaden, *et al.*, Genomic imputation and evaluation using high-density Holstein
genotypes. *Journal of Dairy Science* **96**, 668–678 (2013).

7. B. J. Hayes, H. D. Daetwyler, 1000 Bull Genomes Project to map simple and complex
genetic traits in cattle: applications and outcomes. *Annual Review of Animal Biosciences* **7**,
89–102 (2019).

8. D. W. Huang, B. T. Sherman, R. A. Lempicki, Bioinformatics enrichment tools: paths toward
the comprehensive functional analysis of large gene lists. *Nucleic Acids Res.* **37**, 1–13
(2009).

9. P. L. Beales, *et al.*, IFT80, which encodes a conserved intraflagellar transport protein, is
mutated in Jeune asphyxiating thoracic dystrophy. *Nat Genet* **39**, 727–729 (2007).

10. S. Rix, A. Calmont, P. J. Scambler, P. L. Beales, An Ift80 mouse model of short rib
polydactyly syndromes shows defects in hedgehog signalling without loss or malformation of
cilia. *Hum Mol Genet* **20**, 1306–1314 (2011).

11. A. Graf, *et al.*, Fine mapping of genome activation in bovine embryos by RNA sequencing.
PNAS **111**, 4139–4144 (2014).

12. X. Yuan, X. Cao, S. Yang, IFT80 is required for stem cell proliferation, differentiation, and
odontoblast polarization during tooth development. *Cell Death Dis* **10**, 1–11 (2019).

13. F. Lanner, J. Rossant, The role of FGF/Erk signaling in pluripotent cells. *Development* **137**,
3351–3360 (2010).

14. E. W. Kuijk, *et al.*, The roles of FGF and MAP kinase signaling in the segregation of the
epiblast and hypoblast cell lineages in bovine and human embryos. *Development* **139**, 871–
882 (2012).

15. P. M. VanRaden, J. R. O'Connell, G. R. Wiggans, K. A. Weigel, Genomic evaluations with
many more genotypes. *Genetics Selection Evolution* **43**, 10 (2011).

16. H. Li, Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv:1303.3997 [q-bio] (2013) (October 29, 2021).

17. H. Li, *et al.*, The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–
2079 (2009).

18. A. McKenna, *et al.*, The Genome Analysis Toolkit: A MapReduce framework for analyzing
next-generation DNA sequencing data. *Genome Research* **20**, 1297–1303 (2010).

19. P. Danecek, *et al.*, Twelve years of SAMtools and BCFtools. *GigaScience* **10**, giab008
(2021).

20. P. Cingolani, *et al.*, A program for annotating and predicting the effects of single nucleotide
polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-
2; iso-3. *Fly* **6**, 80–92 (2012).

21. M. S. Ortega, *et al.*, Influences of sire conception rate on pregnancy establishment in dairy
cattle†. *Biology of Reproduction* **99**, 1244–1254 (2018).

22. M. S. Ortega, *et al.*, NANOG is required to form the epiblast and maintain pluripotency in the
bovine embryo. *Molecular Reproduction and Development* **87,** 152–160 (2019).

23. P. Tribulo, R. M. Rivera, M. S. Ortega, E. A. Jannaman, P. J. Hansen, "Production and
Culture of the Bovine Embryo" in *Comparative Embryo Culture*, (Springer, 2019), pp. 115–
129.

24. M. S. Ortega, J. J. Kurian, R. McKenna, P. J. Hansen, Characteristics of candidate genes
associated with embryonic development in the cow: Evidence for a role for WBP1 in
development to the blastocyst stage. *PLOS ONE* **12**, e0178041 (2017).

25. K. Goossens, *et al.*, Selection of reference genes for quantitative real-time PCR in bovine
preimplantation embryos. *BMC Developmental Biology* **5**, 27 (2005).

Figures and Tables

403
Figure 1. A UCSC genome browser plot (**A**) shows the locations of prior Holstein Haplotype 2
(HH2) loci as determined in December 2013, April 2020 and August 2020. The IFT80 gene is
highlighted by red dashed lines. Concordance analysis of variant sites against expected carrier
status (**B**) identified only one variant with > 99% concordance (highlighted in red). For genomic
context, the IFT80 gene is drawn with exons (straight lines) and introns (diagonal lines) above the
concordance plot. Only variants predicted to have high functional effect by SNPEff are highlighted
by labels showing coordinate numbers.

412
Figure 2. A diagram of the IFT80 gene (**A**) shows CRISPR-Cas9 edit sites (red triangles) and
qPCR primer design sites (blue boxes). Gene exons are represented by vertical lines, whereas
introns are represented by diagonal lines. Embryo counts for each developmental stage for exon
2 (**B**) and exon 3 (**C**) demonstrate that homozygous edit embryos (red) do not typically progress
past the 8-cell stage compared to vector Cas9 controls (blue). Single (*) and double (**) asterisks
indicate Student's T test p-values that are less than 0.001 and 0.0001, respectively. Relative fold
change of exons downstream of the exon 2 (**D**) and exon 11 (**E**) edit sites showed significant
decreases in edited embryos (red) compared to the vector Cas9 controls (blue)

421
422
Figure 3. A. Development of control versus edited embryos. IFT80 injected group have increased
number of degenerating embryos (white arrows,120 and 80 hours post insemination), after the 8-
cell stage. In control embryos, development progressed normally with formation of blastocysts
with compact inner cell mass and defined blastocoel (white arrows,180 hours post insemination).
B. Embryo genotyping for exon 2 edits. Each numbered lane represents one embryo. Lanes 1, 5,
and 6 show non-edited embryos (wild type); lanes 2, 3, and 9 show biallelic edits (absence of wild
type band), and lanes 4, 7, 8, and 10 illustrate mono-allelic edits (both wild type, and edited band
present). **C**. Embryo genotyping for exon 11 edits. Each numbered lane represents one embryo.
Lanes 9, 12, 15, 26, 27, 28, and 29 show non-edited embryos (wild type); lanes 3 – 8, 10, 11, 16,
19, 20, 22, and 24 show biallelic edits (absence of wild type band), and lanes 2, 13, 14, 21, and
23 illustrate mono-allelic edits (both wild type, and edited band present). **D**. Protein model for
IFT80 truncation at exon 11. The pink model represents the 777-amino acid wild-type protein, the
blue model represents the 385-amino acid gene-edited protein (after truncation surrounding
mutation), and the yellow globular regions indicate putative active binding sites.

437　**Table 1.** IFT80 guide RNA design table for exons 2 and 11.

Exon	Guide	Guide name	Position Genomic location	Strand	Sequence – PAM[1] (5' -> 3')	On target Score[2] BI	IDT
	1	BI283	107,097,118 - 107,097,140	+	TGTTTTATGTCTCCTTAATA-TGG	33.75	68.0
2	2	BI313	107,097,148 - 107,097,170	+	TCAAGAATTAGTGAGCTGTG-TGG	62.53	65.0
	3	BI423	107,097,269 - 107,097,291	-	TCTATTGGGTAAATATCATC-AGG	47.26	60.0
11	1	Exon11-1	107,172,343 - 107,172,365	+	GGAATCAAAACTTTATAGGG-TGG	62.05	64.0
	2	Exon 11-3	107,172,670 - 107,172,692	-	CAGGAAATTTTGGAGAGGAA-AGG	53.38	32.0

438　[1]PAM: protospacer adjacent motif.

439　[2]On target score: predicted guide efficiency from two different design tools: BI (Broad Institute) and IDT (Integrated DNA Technologies).

440 **Table 2.** Primer sequences for gene expression analysis.

Exon	Primer Forward (5' → 3')	Primer Reverse (5' → 3')
1	TGAATTCAATCAGGTGGCATTT	AGACATGAGACTGAAGATATCTCTT
3–4	TGGCAGGACGATGGAATTATG	AAAGACGGGAATGCTAAGATCG
5–7	CCGCTTCAACCAAATGCTAAA	GTATGGGATAGTTATGGCCGTC
9–10	GCCTCTTTGAACTATGCACACTTAG	CTACAAAGAACTGGAATACAC
12	CTCTTTGAGGCATCAACTGGA	TTGGAAATTGCCCTGGATCA