Identification of novel translated small ORFs in *Escherichia coli* using complementary

ribosome profiling approaches

Anne Stringer[1†], Carol Smith[1†], Kyle Mangano[2] and Joseph T. Wade[1,3*]

[1]Wadsworth Center, New York State Department of Health, Albany, New York, USA.

[2]Center for Molecular Biosciences, University of Illinois, Chicago, USA.

[3]Department of Biomedical Sciences, School of Public Health, University at Albany, Albany, New

York, USA.

[†]Equal author contribution.

[*]Corresponding author: joseph.wade@health.ny.gov

ABSTRACT

Small proteins of <51 amino acids are abundant across all domains of life but are often overlooked

because their small size makes them difficult to predict computationally, and they are refractory

to standard proteomic approaches. Ribosome profiling has been used to infer the existence of small

proteins by detecting the translation of the corresponding open reading frames (ORFs). Detection

of translated short ORFs by ribosome profiling can be improved by treating cells with drugs that

stall ribosomes at specific codons. Here, we combine the analysis of ribosome profiling data for

Escherichia coli cells treated with antibiotics that stall ribosomes at either start or stop codons.

Thus, we identify ribosome-occupied start and stop codons for ~400 novel putative ORFs with

high sensitivity. The newly discovered ORFs are mostly short, with 365 encoding proteins of <51

amino acids. We validate translation of several selected short ORFs, and show that many likely

encode unstable proteins. Moreover, we present evidence that most of the newly identified short

ORFs are not under purifying selection, suggesting they do not impact cell fitness, although a small

subset have the hallmarks of functional ORFs.

IMPORTANCE

Small proteins of <51 amino acids are abundant across all domains of life but are often overlooked

because their small size makes them difficult to predict computationally, and they are refractory

to standard proteomic approaches. Recent studies have discovered small proteins by mapping the

location of translating ribosomes on RNA using a technique known as ribosome profiling.

Discovery of translated sORFs using ribosome profiling can be improved by treating cells with

drugs that trap initiating ribosomes. Here, we show that combining these data with equivalent data

for cells treated with a drug that stalls terminating ribosomes facilitates the discovery of small

proteins. We use this approach to discover 365 putative genes that encode small proteins in

Escherichia coli.

INTRODUCTION

Tens of thousands of bacterial genomes have been fully sequenced. These genome sequences are annotated using computational pipelines that predict the location of open reading frames (ORFs). One of the main criteria used by ORF prediction pipelines is the length of the putative ORFs; ORFs encoding proteins with >50 amino acids (aa) are unlikely to occur by chance; thus, there is a strong bias towards identifying longer ORFs. However, there is strong evidence for large numbers of short ORFs (sORFs; encoding proteins with <50 aa): phylogenetic analysis has identified many conserved sORFs (1), and transcriptomic and proteomic experimental approaches provide strong evidence for the translation of many sORFs (2–15). As more sORFs and their encoded small proteins are identified, it is becoming increasingly clear that sORFs and small proteins play important functional roles in bacteria. In particular, sORFs can function as *cis*-acting regulators of downstream operonic genes, and small proteins can function to modulate the activity of protein complexes (16, 17).

There are 118 described sORFs encoded by *Escherichia coli* K-12, most of which have not been functionally characterized. Recent studies have used genome-scale approaches to identify translated sORFs. Ribosome profiling is a method that experimentally maps ribosome association with the transcriptome. Two studies applied ribosome profiling to *E. coli* BL21 (closely related to *E. coli* K-12) cells treated with antibiotics, retapamulin ("Ribo-RET") and Onc112, that trap ribosomes at start codons (2, 3). Thus, sites of translation initiation were identified, leading to the identification of many putative translated sORFs. In one study, 38 of the encoded small proteins were successfully validated by western blot (3).

Ribo-RET has also been applied to *Mycobacterium tuberculosis*, revealing an even larger number

of putative sORFs than described for *E. coli* (4). Moreover, the majority of the sORFs identified

in *M. tuberculosis* do not appear to have been subject to purifying selection, suggesting that they

represent the products of "pervasive translation", whereby ribosomes initiate translation at large

numbers of locations across the transcriptome, with many of the translated proteins being non-

functional, i.e. not contributing to cell fitness.

Ribosome profiling has also been applied to *E. coli* cells treated with the antibiotic apidaecin,

which traps ribosomes at stop codons by binding in the nascent peptide exit tunnel and trapping

release factor (18). This method, "Ribo-Api", also leads to ribosome accumulation upstream and

downstream of stop codons, and at start codons (19); this additional signal can be partially reduced

by addition of puromycin to cell lysates ("Ribo-Api/Pmn"), but even with this reduction in signal

away from stop codons, Ribo-Api/Pmn data cannot be used to identify stop codons with high

specificity. Nonetheless, we reasoned that combining Ribo-RET and Ribo-Api/Pmn datasets for

E. coli would facilitate the identification of translated sORFs with high sensitivity, since the

likelihood of detecting Ribo-RET signal at a start codon and Ribo-Api/Pmn signal at the

corresponding stop codon by chance would be very low. Our analysis of Ribo-RET and Ribo-

Api/Pmn data led to the identification of 397 novel ORFs with high confidence, most of which are

sORFs. We validated expression of the associated small proteins for 9 of 18 novel sORFs tested,

and we showed that the ribosome-binding sites of all 18 selected sORFs could drive translation of

a luciferase reporter assay. Thus, our data suggest that most or all of the 365 putative sORFs

identified by combining Ribo-RET and Ribo-Api/Pmn data are robustly translated, but that many

of these sORFs encode scarce or unstable small proteins. We speculate that pervasive translation

leads to the production of many unstable, non-functional proteins in *E. coli*.

RESULTS

Reanalysis of Ribo-RET data to identify putative start codons

Using a previously described Ribo-RET dataset for *E. coli* BL21 (2) (Figure 1A), we identified 12,756 sites of ribosome occupancy that exceed the local background; we refer to these positions as "Initiation-Enriched Ribosome Footprints" (IERFs; Table S1). We then determined the enrichment of every possible trinucleotide sequence in the regions surrounding IERFs. Consistent with most IERFs representing sites of translation initiation, ATG, GTG and TTG trinucleotide sequences were enriched over the local background at positions 15-18 nt upstream of IERFs (Figure 1B), with the highest enrichment observed for ATG sequences, and the lowest enrichment observed for TTG sequences; we did not observe enrichment of CTG, ATT, or ATC sequences, consistent with these sequences being used infrequently as start codons (20). Based on the degree of enrichment for ATG, GTG and TTG sequences at different positions 14-18 nt upstream of IERFs, we reanalyzed the Ribo-RET data to identify putative start codons (see Methods). Thus, we identified 7,936 putative start codons with an estimated false discovery rate (FDR) of 11.7% (see Methods). 2,545 of these putative start codons match those of annotated ORFs. We compared the putative start codons to those predicted previously using the same Ribo-RET data using a higher read coverage threshold, but with a considerably more relaxed range of potential start codons (see methods for details). 116 of the start codons identified in the current study were identified previously in reference (2).

Figure 1

A

B

Position Relative to IERF (nt)

C

Position Relative to TERF (nt)

D

Figure 1. Identification of ORFs by combining Ribo-RET and Ribo-ApiPmn data. (A)

Sequence read coverage is shown across a selected genomic region for Ribo-RET (retapamulin

treatment; stalls initiating ribosomes), Ribo-Api/Pmn (apidaecin and puromycin treatment; stalls

terminating ribosomes), and Ribo-Pmn (puromycin treatment; control) data. Blue and orange

triangles indicate the positions of start codons paired with Initiation-Enriched Ribosome Footprints

(IERFs) and stop codons paired with Termination-Enriched Ribosome Footprints (TERFs),

respectively. Triangles joined by lines indicate start and stop codons from the same putative ORF.

The positions of the annotated *yjtD* and novel *ytjE* ORFs are shown by horizontal arrows above

the data. **(B)** Heatmap showing enrichment of possible start codon sequences at positions upstream

of IERFs. **(C)** Heatmap showing enrichment of possible stop codon sequences at positions

upstream of TERFs. **(D)** Classification of ORFs identified from Ribo-RET and Ribo-Api/Pmn

data, with categories defined based on the overlap of start/stop codons with annotated ORFs, as

shown in the schematic.

Reanalysis of Ribo-Api data to identify putative stop codons

Using a previously described Ribo-Api dataset, including a puromycin treatment ("Ribo-

Api/Pmn"), for *E. coli* BL21 (19) (Figure 1A), we identified 12,756 sites of ribosome occupancy

that exceed the local background; we refer to these positions as "Termination-Enriched Ribosome

Footprints" (TERFs; Table S2). We then determined the enrichment of every possible trinucleotide

sequence in the regions surrounding TERFs. Consistent with most TERFs representing sites of

translation termination, TAA, TGA and TAA trinucleotide sequences were enriched over the local

background at positions 12-14 nt upstream of TERFs (Figure 1C). Based on these enrichments, we

reanalyzed the Ribo-Api/Pmn data to identify putative stop codons by selecting TERFs with a

TAA/TGA/TAG trinucleotide sequence in a specified distance range upstream. Thus, we identified

6,877 putative stop codons with an estimated FDR of 33.7% (see Methods). 1,377 of these putative

stop codons match those of annotated genes.

Combining putative start and stop codons to identify putative novel ORFs

We reasoned that by selecting putative start codons inferred from Ribo-RET data that are paired

with putative stop codons inferred from Ribo-Api/Pmn data, we could identify ORFs with higher

confidence than by using Ribo-RET or Ribo-Api/Pmn data alone. Moreover, evidence of ribosome

occupancy at both start and stop codons of these putative ORFs implies they are actively translated

from beginning to end (Figure 1A). Using this approach, we detected 1,839 putative ORFs (Table

S3), with a conservative FDR estimate of 2.2% (see Methods). 1,005 of the ORFs we detected

perfectly match annotated genes; we refer to these as "annotated" ORFs. 437 of the ORFs we

detected share a stop codon with an annotated gene, but have a different start codon. These could

represent mis-annotated genes, or one of a pair of ORFs where the corresponding proteins share

the same C-terminal region. We refer to these as "isoform" ORFs. 397 of the ORFs we detected

have a stop codon that does not match any annotated gene. We refer to these as "novel" ORFs

(Figure 1D). The likelihood of detecting an annotated ORF by chance is exceedingly low (we

conservatively estimate the FDR to be ~0.07%; see Methods). We conservatively estimate that the

FDRs for identifying isoform and novel ORFs are 2.3% and 7.5%, respectively (see Methods).

Putative isoform and novel ORF start and stop codons are associated with RNA secondary

structure features expected of actively translated ORFs

Actively translated start codons in *E. coli* tend to be associated with regions of reduced RNA

secondary structure relative to the surrounding sequence (21). We examined the sequences around

the 1,005 annotated, 437 isoform, and 397 novel ORF start codons. For all three classes of ORF,

we observed significantly reduced predicted RNA secondary structure in the regions from -25 to

+15 nt from start codons, relative to that of 500 randomly selected genomic sequences of the same

length (Figure 2; Mann-Whitney U test $p < 2.2\mathrm{e}^{-16}$ for each of annotated and novel ORFs; $p = 3.2\mathrm{e}^{-}$

^{6} for isoform ORFs). We conclude that the start codons of isoform and novel ORFs are associated

with RNA secondary structure features expected of actively translated ORFs. Predicted RNA

secondary structure around the start codons of the annotated ORFs was slightly lower than that for

novel ORFs (Mann-Whitney U test $p = 1.4\mathrm{e}^{-8}$), and substantially lower than that for isoform ORFs

(Mann-Whitney U test $p < 2.2\mathrm{e}^{-16}$).

Figure 2

Figure 2. Reduced RNA secondary structure for sequences around identified start codons.
Strip plot showing the ΔG for the predicted minimum free energy structures for the regions from
-25 to +15 nt relative to putative start codons for annotated, isoform, and novel ORFs identified
by analyzing Ribo-RET and Ribo-Api/Pmn data, and for a set of 500 random sequences.

Classification of novel ORFs based on local gene context

We compared the positions of the 397 putative novel ORFs with the positions of annotated genes

(Figure 3). 27% of the novel ORFs are located entirely in intergenic regions; of these, roughly two

thirds are immediately upstream of an annotated gene on the same strand, suggesting that they are

co-transcribed. 44% of the novel ORFs overlap completely with an annotated gene in the sense

orientation, whereas only 12% overlap completely with an annotated gene in the antisense

orientation. 17% of the novel ORFs partially overlap an annotated gene.

Strikingly, the sizes of novel ORFs are typically much smaller than those of annotated ORFs; the

median length of novel ORFs is 15 codons, and 365 of the novel ORFs encode proteins of ≤ 50

amino acids, and hence are classified as sORFs. We compared the 365 putative novel sORFs found

in *E.* coli BL21 with sORFs described for *E. coli* K-12 (strain MG1655); 15 of the putative sORFs

perfectly match those described previously for MG1655, and 2 of the putative sORFs match

previously described ORFs with one or two nucleotide mismatches (3, 14, 15).

Experimental validation of putative novel sORFs

We selected 18 sORFs for experimental validation, covering a range of expression values based

on Ribo-RET and Ribo-Api/Pmn coverage at start and stop codons, respectively. We C-terminally

epitope tagged 17 of these ORFs at their native loci with SPA tags. We then used western blotting

to detect stably expressed tagged proteins for cells grown in rich medium. As a positive control,

we used a strain expressing SPA-tagged YhgP, an sORF that was previously detected by western

blotting (3). We observed single bands for the positive control protein and for 9 of the 17 proteins

encoded by putative sORFs. We observed no bands for an untagged strain grown under the same

Figure 3

Figure 3. Classification of novel ORFs by position relative to annotated genes. The pie chart

shows the distribution of novel ORFs across different categories defined by the type of overlap

with an annotated gene, as shown in the schematic.

conditions (Figure 4A). We conclude that approximately half of the putative sORFs are stably

expressed. We assigned new gene names to the ORFs for which a protein was detectable by

western blot (Figure 4; Table S3).

We reasoned that failure to detect a protein by western blot could either be due to the corresponding

sORF being a false positive, or instability of the protein. To distinguish between these possibilities

for the eight sORFs where the corresponding protein was not detected by western blot, for one

sORF not tested by western blotting, and two of the sORFs where the corresponding protein was

detected by western blot, we replaced the sORFs with a nano-luciferase gene at their native loci.

We constructed equivalent strains with mutations in the putative start codons (ATG → ACG). For

all ORFs tested, we detected robust luciferase expression; in all but one case, luciferase activity

was significantly lower in the start codon mutant strain (Figure 4B). We conclude that these sORFs

are actively translated, but the encoded proteins are not easily detected by western blot, likely due

to instability. In some cases, mutation of the start codon did not reduce luciferase activity to

background levels, suggesting that other in-frame trinucleotide sequences upstream of the

predicted start codon can serve as alternative start codons. In the case of sORF 4195023- (named

based on the genome coordinate and strand of the start codon), luciferase activity was the same for

both wild-type and mutant constructs, suggesting that the major start codon is upstream of the

predicted start codon.

Evidence for function of novel sORFs

Our data strongly suggest that the large majority of the 397 novel sORFs are actively translated in

E. coli BL21, but this does not indicate whether the sORFs or their encoded proteins are functional,

Figure 4

A

B

Figure 4. Experimental validation of sORFs by western blot and luciferase fusions. (A)

Western blotting of SPA-tagged novel small proteins. Each lane was loaded with whole cell extract

from cells expressing a SPA-tagged small protein, except for lanes labeled "Untagged", which

were loaded with wild-type MG1655. Underlined sample names indicate that the cells were Δ*thyA*

and grown in medium supplemented with thymine. Asterisks indicate non-specific bands. The

arrow indicates the expected position of SPA-tagged small proteins. YhgP is a positive control,

previously confirmed by western blot. Untagged strains are negative controls. Empty triangles

indicate the positive control lanes. Filled triangles indicate novel small proteins with bands of the

expected size. **(B)** Luciferase assays for nano-luciferase (nLuc) fusions immediately downstream

of the start codons of putative sORFs. Gray bars indicate normalized luciferase activity for nLuc

fusions with wild-type start codons; orange bars indicate normalized luciferase activity for nLuc

fusions with mutant (NCG) start codons. sORF names with asterisks indicate those for which we

detected a corresponding protein by western blot. "nLuc⁻" (purple bar) indicates control data for

cells without nLuc. Data represent the average of three independent biological replicates; error

bars represent ± one standard deviation from the mean.

with "function" defined as providing a positive contribution to cell fitness (22). Indeed, recent

studies suggest that bacteria and eukaryotes can express hundreds of sORFs that are likely non-

functional (4, 23, 24). We used three approaches to identify likely functional sORFs. First, we

searched the encoded proteins for functional domains in the PFAM database (25). Most of the

proteins are too short to include a functional domain; nonetheless, 6 of the novel ORFs have

significant matches to functional domains listed in the PFAM database (Table S3), with the

shortest of these ORFs encoding a protein of 61 amino acids. Second, we examined the codon bias

of the sORFs; codon sequences that are non-random compared to the overall nucleotide content of

the ORFs are indicative of purifying selection, a feature observed for annotated ORFs. We

reasoned that novel sORFs with codon usage similar to annotated ORFs are likely under purifying

selection. We analyzed the codon usage of 105 novel sORFs in regions that have at least 30 nt

non-overlapping with annotated genes (including annotated genes not identified by Ribo-RET or

Ribo-Api/Pmn). Specifically, we calculated the Relative Codon Deoptimization Index (RCDI), a

measure of how closely the codon usage of an ORF matches that of the annotated ORFs from the

same genome (26, 27). 11 of the 105 selected sORFs have codon usage similar to that of annotated

genes (RCDI score <1.9, $p < 0.01$; RCDI scores closer to 1 indicate greater codon optimality; see

Methods; Table S3), suggesting that these sORFs are under purifying selection, and hence are

likely to be functional. Third, we searched for homologues of proteins encoded by novel sORFs,

excluding those that have previously been described in *E. coli* MG1655, and limiting the analysis

to proteins encoded entirely in intergenic regions. Thus, we identified likely homologues for seven

of the encoded proteins (Table S3; Figure 5). Overall, our data identify a subset of sORFs that are

likely to be functional; however, we found no evidence of function for the majority of sORFs.

Figure 5

```
                  4087684+
Escherichia       MLNADVYPRYSGCGKAATE*
Salmonella        LLVADVYPRYSGCGKAATE*

                  YpeD
Escherichia       MNLIIILVWHSMKHSKGYVLEAQDDELSR*
Salmonella        MHLIIIINKHSMKHSKGYVFEAKDD*QSR-
Citrobacter       MNLIIILNKHSIKHSKGYVSEAKDDKRSR*

                  YnhH
Escherichia       LDCRTQRLKNRHSRWAETQAHIPLHAFSMSPILRARHHHFRNTGFAFRQCAQKASFSHPSQLKD*
Salmonella        LDCKTQRLKLSPSRWAETRAHIPLHALSMSPILRARHHHFSNTGLASGPCAPEARFPYPPQLKD*
Citrobacter       LDCKTQRLKNTPSRWAETRAHIPLHALSMSPILRARQHHFRNTGLASRPCAPEARFPYPSQLKD*

                  4195023+
Escherichia       MHLQLIKDNIHSVVICYT*
Salmonella        IHLQLIKDNILSVVICYT*

                  YtgB
Escherichia       VLILTLLRHWHVRTKSER*
Salmonella        VLILTLLRHWHVRTSSER*
```

Figure 5. Conservation of selected sORFs. Sequence alignments of five selected novel proteins

and their putative homologues in *Salmonella enterica* and (where found) *Citrobacter koseri*.

Alignments were extracted from tBLASTn output files. Shaded amino acids match the protein

sequence from *E. coli*. The boxed region for YpeD and putative homologues indicates the region

of overlap in the corresponding genes with the downstream gene (*mntH*) and its homologues.

Representative *S. enterica* and *C. koseri* genome sequences are shown (*Salmonella enterica* subsp.

enterica serovar Worthington, strain CFSAN051295, CP029041.1; *Citrobacter koseri*, strain

FDAARGOS_1029, CP066089.1).

DISCUSSION

Combining Ribo-RET and Ribo-Api datasets is a highly specific approach to identify translated

ORFs

Previous studies in prokaryotes and eukaryotes have identified start codons by combining

Ribosome Profiling coupled with antibiotics such as retapamulin or homoharringtonin that trap

initiating ribosomes (2, 3, 28–31). While this approach has been effective at mapping sites of

translation initiation, addition of the antibiotics does not completely remove signal from elongating

ribosomes. Hence, Ribo-RET and related approaches have uncertainty that can make it difficult to

identify start codons with high confidence, especially for start codons internal to highly expressed

ORFs (29). Moreover, Ribo-RET signal at a putative start codon does not indicate whether

ribosomes actively translate the corresponding ORF to completion in the absence of antibiotic.

Similar to Ribo-RET and related approaches, Ribo-Api/Pmn can be used to identify sites of

translation termination, i.e. active stop codons. However, apidaecin treatment does not cause

ribosomes to be exclusively associated with stop codons; ribosomes are also enriched in the

regions immediately upstream and downstream of stop codons, and at start codons (19), such that

stop codon identification from Ribo-Api/Pmn data is associated with a high FDR (estimated at

33.7% for our analysis). Moreover, identification of a stop codon position rarely indicates the

position of the associated start codon(s).

Since IERFs in Ribo-RET data and TERFs in Ribo-Api/Pmn data are independent measurements,

but start codons can be unambiguously associated with a single stop codon, combining data from

both approaches allows ORF identification with a greatly reduced FDR compared to Ribo-RET or

Ribo-Api/Pmn alone (estimated at 2.2% for our analysis). Moreover, detecting ribosomes at both

start and stop codons for the same ORF strongly suggests that translation runs from start to stop

codon.

The high sensitivity of an approach combining Ribo-RET and Ribo-Api/Pmn is evidenced by the

fact that 35% of novel ORFs and 9% of annotated ORFs we identified have only two Ribo-

Api/Pmn sequence reads at their stop codons. Nonetheless, the low FDR, and the propensity for

novel ORFs to have low RNA secondary structure around their start codons strongly suggest that

the majority of novel ORFs are actively translated. We anticipate that Ribo-RET and Ribo-

Api/Pmn or related approaches will be combined to identify translated ORFs with high confidence

in other bacterial and non-bacterial species. One potential barrier to these methods is the ability of

the drugs to accumulate within cells; this could be prevented due to low permeability, or active

efflux. While Ribo-RET has proved effective in *E. coli* (2) and *M. tuberculosis* (4), deletion of the

tolC gene encoding a component of the drug efflux pump was needed for successful Ribo-RET

with *E. coli*, and most Gram-negative species have similarly low sensitivity to retapamulin.

Apidaecin uptake requires the *sbmA* gene, which encodes an inner-membrane transport protein

(32); *sbmA* is missing in some Gram-negative and all Gram-positive bacterial species. If apidaecin

uptake is low, an alternative approach would be to encode and express apidaecin, a small protein,

inside the bacteria (33, 34). Similarly, Onc112, which functions similarly to retapamulin, is a small

protein that could be encoded and expressed within cells (35).

E. coli expresses large numbers of small proteins, many of which are likely unstable

Most of the novel ORFs we identified in this study are sORFs, i.e. they encode proteins of <51

amino acids. sORFs and small proteins have been challenging to identify by other methods; thus,

we have greatly expanded the number of novel ORFs known for *E. coli*, with the caveat that not

all have been experimentally validated using an independent approach. Luciferase assays

measuring the capacity of the translation initiation region of selected novel ORFs at their native

loci to direct translation indicate that expression levels vary greatly between sORFs, although it is

important to note that the native sequences immediately downstream of the start codons of these

sORFs were replaced in these assays by introduction of the nLuc gene; differences in expression

caused by these downstream sequences will have been lost. Western blotting also indicates large

differences in expression of small proteins encoded by selected novel sORFs. While differences

in protein levels detected by western blot roughly mirrored differential expression levels measured

using luciferase reporters, there were notable differences. In particular, protein levels of YtjE were

moderately higher than those of YtiE, whereas luciferase activity for the *ytiE-nLuc* reporter was

>100-fold higher than that for the *ytjE-nLuc* reporter. These differences are likely attributable to

differences in stability between small proteins.

Nine of the proteins encoded by selected sORFs were not detectable by western blot, yet we

detected robust luciferase activity for all the sORF-*nLuc* fusions. These data strongly suggest that

(i) the large majority of novel sORFs identified by combining Ribo-RET and Ribo-Api/Pmn are

actively translated, and (ii) many of the proteins encoded by identified sORFs are unstable.

Evidence that most small ORFs in E. coli *are not functional*

Several lines of evidence suggest that most of the *E. coli* sORFs we identified are not functional,

with "function" defined as providing a positive contribution to cell fitness, and hence being under

selective pressure (22): (i) the codon usage of most analyzed novel sORFs did not match that of

annotated ORFs; (ii) we detected sequence conservation at the amino acid level for only a few of

the novel sORFs in genera beyond *Escherichia* and *Shigella* (*Shigella* species were excluded due

to very high similarity between *Shigella* and *Escherichia* sequences). These analyses, however,

have important limitations: (i) codon usage patterns are only informative for sORFs that do not

overlap annotated ORFs; (ii) the statistical significance of both codon usage and sequence

conservation is limited by the small size of sORFs; (iii) regulatory sORFs can be functional even

if the corresponding proteins are not, and these sORFs often have non-standard codon usage. Based

on our study of sORFs in *M. tuberculosis*, we proposed that the *M. tuberculosis* transcriptome is

pervasively translated, with most translated proteins being non-functional. We propose that the *E.*

coli is also subject to pervasive translation, although seemingly at a level below that of *M.*

tuberculosis based on the relative number of novel and annotated ORFs identified in the two

studies. A lack of function for most sORFs is consistent with half of the small proteins tested

appearing to be weakly expressed and/or unstable, features expected of spurious proteins.

Annotated genes that contain start codons for novel ORFs tend to have suboptimal codon usage.

The position of novel ORFs with respect to annotated genes is non-random (Figure 3). Specifically,

we detected many more novel ORFs on the same strand as annotated overlapping or downstream

genes. We speculate that this bias is because novel ORFs require a transcript, and mRNAs for

annotated genes tend to be more abundant than antisense RNAs. Nonetheless, we were surprised

to find that 44% of all the novel ORFs we detected overlap completely with an annotated gene on

the same strand; this suggests that overlapping translation of the annotated ORFs does not always

prevent translation initiation at ORF-internal start codons, as suggested previously (2).

We speculated that novel ORFs whose start codons are within annotated genes reflect inefficient

translation of the overlapping annotated ORF. To test this idea, we assessed the codon usage of

annotated genes that encompass novel ORF start codons. The codon usage of annotated genes that

encompass novel ORFs (mean RCDI score = 1.41) is significantly less optimal than that for the

set of all other annotated ORFs (mean RCDI score = 1.33; Mann-Whitney U-test comparing RCDI

scores, $p = 1.7e^{-5}$). This difference was more pronounced when we considered only highly

expressed novel ORFs (mean RCDI score of 1.59 for annotated genes that encompass novel ORF

starts with Ribo-RET signal >5 RPM; $p = 2.1e^{-12}$). We conclude that suboptimal codon usage of

an annotated ORF increases the likelihood of internal initiation sites, likely due to a reduced

efficiency of translation elongation. Suboptimal codon usage is a feature of horizontally acquired

genes in *E. coli* (36), suggesting that horizontally acquired genes often have an increased

abundance of internal translation. Another feature of horizontally acquired genes is low G/C

content (36). Strikingly, annotated genes that encompass highly expressed novel ORF starts have

a much lower G/C content (mean of 42.2%) than the set of all annotated ORFs (mean of 51.1%).

Given that genes with internal start codons often have suboptimal codon usage, and often have low

G/C content, we speculate that horizontally acquired genes are hotspots for internal novel ORFs

due to inefficient translation elongation.

Identification of E. coli *sORFs that are likely to be functional*

A few of the novel sORFs we identified are associated with features expected of functional

proteins. Specifically, we detected stably expressed protein by western blot, we observed codon

usage patterns similar to those of annotated ORFs, and/or we identified putative homologues in

other genera. While the functions of these sORFs or their corresponding small proteins are

unknown, they represent promising candidates for further study. In the case of sORFs that are

immediately upstream or partially overlapping with annotated genes on the same strand, a potential

function is regulation of the downstream gene by translation the sORF, as has been described for

other sORFs in bacteria (17). Our data also suggest exercising caution when selecting sORFs/small

proteins for further study, since many sORFs/small proteins may be non-functional; relying on

codon usage patterns and/or sequence conservation is a simple way to prioritize likely functional

sORFs/small proteins.

MATERIALS AND METHODS

Strains and plasmids

All strains and plasmids used in this work are listed in Table 1. All oligonucleotides used in this

work are listed in Table S4. All strains used in this work are derivatives of *Escherichia coli* K-12

MG1655 (37). Strains AMD783-AMD800 were constructed using λ Red recombineering as

described previously (38). Specifically, PCR products used for recombineering were generated

with oligonucleotides JW10796-JW10803 and JW10883-JW10912, using DY330 *allR*-SPA::*kanR*

(39) as a template. The PCR products generated to make AMD783 and AMD784 were

electroporated into AMD052 (*Escherichia coli* K-12 MG1655 Δ*thyA*) harboring pKD46 (38). The

PCR products generated to make AMD785-AMD801 were electroporated into MG1655 harboring

pKD46.

Strains AMD827-AMD848 were constructed using the previously described FRUIT method of

recombineering (40). Oligonucleotides JW10919-JW10924, JW10934-JW10936 and JW11010-

JW11033 were used to generate PCR products for recombineering using pAMD375 (see below)

as a template. The PCR products generated were electroporated into AMD052 harboring pSIM6

(41).

Plasmid pAMD375 was constructed using oligonucleotides JW10881 and JW10882 to amplify the

NanoLuc luciferase gene from pNL1.1 (Promega). The resulting DNA fragment was cloned into

pAMD001 (40) digested with *Apa*I and *Nco*I, using the In-Fusion cloning system (Takara).

Identification of IEFRs from Ribo-RET data

Sequence reads for the Ribo-RET data (NCBI SRA accession number SRR8156054) (2) were

aligned to the *E. coli* BL21 reference genome and to a reverse complemented reference genome

using Rockhopper (version 2.03). The genome positions and strands of sequence read 3' ends were

inferred from the resultant .sam file. Sequence read coverage across known non-coding RNAs

(Table S5) was set to zero, and coverage was normalized to reads per million (RPM). Every

genome position on both strands was considered as a possible IEFR. IEFRs were selected if the

corresponding position/strand met three criteria: (i) normalized sequence read coverage was ≥0.5

RPM; (ii) normalized sequence read coverage was ≥10-fold higher than the mean sequence read

depth at all positions in a 101 nt window centered on the position being considered; (iii) normalized

sequence read depth was at least as high as that for any other position in a 21 nt window centered

on the position being considered.

Identification of putative start codons from IEFRs

The frequency of all trinucleotide sequences was determined for positions -50 to -1 nt relative to

the position of each IEFR. We reasoned that trinucleotide sequences used as start codons would

be enriched relative to the local background in the -14 to -18 nt range relative to IEFR positions,

with the strongest enrichment at position -15 (2); we considered the local background frequency

for each trinucleotide sequence to be that for positions -50 to -41 nt relative to the IEFRs. To be

considered as a possible start codon, we first selected candidate start codon sequences by requiring

that a trinucleotide sequence be enriched at least 1.4-fold above background at position -15 relative

to IEFRs, and that the highest level of enrichment in the -14 to -18 nt range be at position -15.

These criteria were met by only three trinucleotide sequences: ATG, GTG, and TTG; hence, only

these sequences were considered as possible start codons. ATG is enriched at least 1.4-fold above

background at positions -14 to -18 nt relative to IEFRs; GTG is enriched at least 1.4-fold above

background at positions -14 to -16 nt relative to IEFRs; TTG is enriched at least 1.4-fold above

background at positions -14 to -15 nt relative to IEFRs. We therefore identified putative start

codons as ATG sequences positioned 14-18 nt upstream of IEFRs, GTG sequences positioned 14-

16 nt upstream of IEFRs, and TTG sequences positioned 15-16 nt upstream of IEFRs.

Calculating the False Discovery Rate (FDR) for putative start codons

The likelihood of randomly selecting a genome coordinate with an associated start codon sequence

(as defined above for IERFs) was estimated by selecting 100,000 random genome coordinates and

determining the fraction, "R", that would be associated with a start codon. The set of IERFs

contains a number of true positives (i.e. corresponding to a genuine start codon), and a number of

false positives. We assume that true positive IERFs are all associated with a start codon using the

parameters described above for calling ORFs. We assume that false positive IERFs are associated

with a start codon at the same frequency as random genome coordinates, i.e. R. Since we know

how many IERFs were not associated with a start codon, we can use this number to estimate how

many false positive IERFs were associated with a start codon by chance. With the total number of

IERFs as "I" and the total number of identified ORFs as "O", the FDR for ORF calls is estimated

by:

$(100 * (I - O) * (R / (1 - R))) / O$

Identification of TERFs from Ribo-Api/Pmn data

TERFs were identified as described for IEFRs, but using Ribo-Api/Pmn data (NCBI SRA

accession number SRR11728142) (19).

Control Ribo-seq data for puromycin-treated cells

Sequence read coverage for a control dataset (puromycin-treated cells) were generated as

described above (NCBI SRA accession number SRR11728143) (19).

Identification of putative stop codons from TERFs

Putative stop codons were identified from TEFRs using the same approach as described above for

identifying putative start codons from IEFRs, except that only TAA, TGA and TAG trinucleotide

sequences were considered. Based on the enrichment of TAA, TGA and TAG sequences above

the local background in the regions upstream of TEFRs, we identified putative stop codons as

TAA, TGA or TAG sequences positioned 12-14 nt upstream of TEFRs. The FDR for putative stop

codons was determined as described for start codons.

Combining IERFs and TERFs to identify putative ORFs

Stop codons identified from TERFs, were discarded if the TERF was located within 3 nt of an

IERF on the same strand, since apidaecin stalls ribosomes at start codons in addition to stalling

ribosomes at stop codons. Putative start codons identified from IERFs were translated *in silico* to

identify the corresponding stop codon. These stop codons were compared to stop codons identified

from TERFs; for shared stop codons, the ORF defined by the IERF was selected as a putative

ORF.

To conservatively estimate an FDR for ORF identification, we rotated the normalized sequence

read coverage data for Ribo-RET and Ribo-Api/Pmn relative to the genome sequence by 1,000 nt,

each of 10 times. After each cycle of rotation, we repeated IERF, TERF, and ORF identification,

as described above. The average number of ORFs identified after each of the 10 rotations was

considered to be a conservative estimate of the number of falsely discovered ORFs in the non-

rotated data. FDRs for annotated, isoform and novel ORFs were estimated by determining how

many of the ORFs identified from rotated data were found in each category, and dividing the

overall FDR equivalently.

RNA folding prediction

The sequence from -25 to +15 nt relative to each start codon, or for 500 x 41 nt sequences randomly

selected from the *E. coli* BL21 genome, were selected for prediction of the free energy of the

predicted minimum free energy structure with a local installation of ViennaRNA Package tool

RNAfold, version 2.4.14, using default settings (42).

Western blotting

MG1655 (untagged) and strains AMD785-AMD800 were grown overnight at 37 °C in LB,

subcultured 1:100 in LB, and grown with shaking at 37 °C to an OD600 of 0.6–0.7. AMD052,

AMD783 and AMD784 were grown overnight at 37 °C in LB supplemented with 100 µg/mL

thymine, subcultured 1:100 in LB supplemented with 100 µg/mL thymine, and grown with shaking

at 37°C to an OD_{600} of 0.6–0.7. After growth, 500 µL of each culture was pelleted and resuspended

in 25 µL Laemmli sample buffer with a 19:1 ratio of buffer:β-mercaptoethanol. Samples were

separated on a 10% SDS-PAGE gel. Proteins were transferred to PVDF membranes and probed

with M2 anti-FLAG antibody (Sigma; 1 in 5,000 dilution) and HRP-conjugated goat anti-mouse

antibody (1 in 250,000 dilution). Tagged proteins were visualized using the SuperSignal™ West

Femto detection kit (ThermoFisher).

Luciferase reporter assays

Strains AMD825-AMD848 were grown overnight in LB at 37°C, subcultured 1:100 in LB and

grown with shaking at 37°C to an OD_{600} of 0.4-0.7. After recording the OD_{600}, 10 µL of each

culture was aliquoted into a 96-well opaque plate with three technical replicates each. All aliquots

were then mixed with 10 µL Nano-Glo (Promega). Luminescence readings (relative light units;

RLUs) were taken using a Turner Biosystems Veritas microplate luminometer. Relative

luminescence values were reported as RLU/OD_{600}.

Analysis of conserved protein domains

A FASTA file containing all proteins encoded by novel ORFs with >20 codons was submitted to

the PFAM webserver (version 34.0) (25) for a batch sequence search using default parameters.

Codon usage analysis

FASTA files containing the nucleotide sequences of regions of novel ORFs not overlapping with

annotated genes were generated and submitted to the RCDI/eRCDI web server (27). The *E. coli*

BL21 genome codon usage table (https://www.kazusa.or.jp/codon) was used, and the following

parameters were used: the upper confidence limit was set to 99% confidence/99% population, the

length of random sequence of 300 nt was selected, and the standard genetic code was used.

Sequence conservation analysis

The amino acid sequences of all proteins encoded by novel, fully intergenic ORFs were submitted

to the NCBI tBLASTn webserver (version 2.11.0) (43). The *Escherichia* (taxid:561) and *Shigella*

(taxid:620) genera were excluded from the search. The following parameters were adjusted: 250

max target sequences were selected, the expect threshold was set to 100, and the low complexity

region filter was unselected. The search results were further filtered by selecting only ORFs with

100% query coverage and a ≥70% sequence identity. Groups of conserved ORFs were realigned

using the Clustal Omega multiple sequence alignment program with default settings (44).

ACKNOWLEDGEMENTS

We thank the Wadsworth Center Applied Genomic Technologies Core Facility for DNA sequencing. We thank the Wadsworth Center Tissue Culture and Media Core Facility and Glassware Facility for technical support. We thank Alexander Mankin, Nora Vazquez-Laslop, and Gisela Storz for helpful suggestions, and comments on the manuscript. We thank Don Court for the recombineering plasmid pSIM6. KM was supported by a National Institutes of Health Training Grant (5T32AT007533). JTW was supported by National Institutes of Health grant R01GM139277.

REFERENCES

1. Sberro H, Fremin BJ, Zlitni S, Edfors F, Greenfield N, Snyder MP, Pavlopoulos GA, Kyrpides
NC, Bhatt AS. 2019. Large-Scale Analyses of Human Microbiomes Reveal Thousands of
Small, Novel Genes. Cell 178:1245-1259.e14.

2. Meydan S, Marks J, Klepacki D, Sharma V, Baranov PV, Firth AE, Margus T, Kefi A,
Vázquez-Laslop N, Mankin AS. 2019. Retapamulin-Assisted Ribosome Profiling Reveals the
Alternative Bacterial Proteome. Mol Cell https://doi.org/10.1016/j.molcel.2019.02.017.

3. Weaver J, Mohammad F, Buskirk AR, Storz G. 2019. Identifying Small Proteins by Ribosome
Profiling with Stalled Initiation Complexes. MBio 10.

4. Smith C, Canestrari JG, Wang J, Derbyshire KM, Gray TA, Wade JT. 2019. Pervasive
Translation in Mycobacterium tuberculosis. bioRxiv 665208.

5. Fuchs S, Kucklick M, Lehmann E, Beckmann A, Wilkens M, Kolte B, Mustafayeva A, Ludwig
T, Diwo M, Wissing J, Jänsch L, Ahrens CH, Ignatova Z, Engelmann S. 2021. Towards the
characterization of the hidden world of small proteins in Staphylococcus aureus, a
proteogenomics approach. PLoS Genet 17:e1009585.

6. Miravet-Verde S, Ferrar T, Espadas-García G, Mazzolini R, Gharrab A, Sabido E, Serrano L,
Lluch-Senar M. 2019. Unraveling the hidden universe of small proteins in bacterial genomes.
Mol Syst Biol 15:e8290.

7. Omasits U, Varadarajan AR, Schmid M, Goetze S, Melidis D, Bourqui M, Nikolayeva O,

Québatte M, Patrignani A, Dehio C, Frey JE, Robinson MD, Wollscheid B, Ahrens CH. 2017.

An integrative strategy to identify the entire protein coding potential of prokaryotic genomes

by proteogenomics. Genome Res 27:2083–2095.

8. Petruschke H, Anders J, Stadler PF, Jehmlich N, von Bergen M. 2020. Enrichment and

identification of small proteins in a simplified human gut microbiome. J Proteomics

213:103604.

9. Venturini E, Svensson SL, Maaß S, Gelhausen R, Eggenhofer F, Li L, Cain AK, Parkhill J,

Becher D, Backofen R, Barquist L, Sharma CM, Westermann AJ, Vogel J. 2020. A global

data-driven census of Salmonella small proteins and their potential functions in bacterial

virulence. microLife 1.

10. Baek J, Lee J, Yoon K, Lee H. 2017. Identification of Unannotated Small Genes in Salmonella.

G3: Genes, Genomes, Genetics 7:983–989.

11. Ndah E, Jonckheere V, Giess A, Valen E, Menschaert G, Van Damme P. 2017.

REPARATION: ribosome profiling assisted (re-)annotation of bacterial genomes. Nucleic

Acids Res 45:e168.

12. Shell SS, Wang J, Lapierre P, Mir M, Chase MR, Pyle MM, Gawande R, Ahmad R, Sarracino

DA, Ioerger TR, Fortune SM, Derbyshire KM, Wade JT, Gray TA. 2015. Leaderless

Transcripts and Small Proteins Are Common Features of the Mycobacterial Translational

Landscape. PLoS genetics 11:e1005641–e1005641.

13. Miranda-CasoLuengo AA, Staunton PM, Dinan AM, Lohan AJ, Loftus BJ. 2016. Functional

characterization of the Mycobacterium abscessus genome coupled with condition specific

transcriptomics reveals conserved molecular strategies for host adaptation and persistence.

BMC Genomics 17:553.

14. Hemm MR, Paul BJ, Schneider TD, Storz G, Rudd KE. 2008. Small membrane proteins found

by comparative genomics and ribosome binding site models. Mol Microbiol 70:1487–1501.

15. VanOrsdel CE, Kelly JP, Burke BN, Lein CD, Oufiero CE, Sanchez JF, Wimmers LE, Hearn

DJ, Abuikhdair FJ, Barnhart KR, Duley ML, Ernst SEG, Kenerson BA, Serafin AJ, Hemm

MR. 2018. Identifying New Small Proteins in Escherichia coli. Proteomics 18:e1700064.

16. Storz G, Wolf YI, Ramamurthi KS. 2014. Small proteins can no longer be ignored. Annu Rev

Biochem 83:753–777.

17. Dever TE, Ivanov IP, Sachs MS. 2020. Conserved Upstream Open Reading Frame Nascent

Peptides That Control Translation. Annu Rev Genet 54:237–264.

18. Florin T, Maracci C, Graf M, Karki P, Klepacki D, Berninghausen O, Beckmann R, Vázquez-

Laslop N, Wilson DN, Rodnina MV, Mankin AS. 2017. An antimicrobial peptide that inhibits

translation by trapping release factors on the ribosome. Nat Struct Mol Biol 24:752–757.

19. Mangano K, Florin T, Shao X, Klepacki D, Chelysheva I, Ignatova Z, Gao Y, Mankin AS,

Vázquez-Laslop N. 2020. Genome-wide effects of the antimicrobial peptide apidaecin on

translation termination in bacteria. Elife 9.

20. Gvozdjak A, Samanta MP. 2020. Genes Preferring Non-AUG Start Codons in Bacteria.
arXiv:200810758 [q-bio].

21. Del Campo C, Bartholomäus A, Fedyunin I, Ignatova Z. 2015. Secondary Structure across the
Bacterial Transcriptome Reveals Versatile Roles in mRNA Regulation and Function. PLoS
Genet 11:e1005613.

22. Keeling DM, Garza P, Nartey CM, Carvunis A-R. 2019. The meanings of "function" in
biology and the problematic case of de novo gene emergence. Elife 8.

23. Carvunis A-R, Rolland T, Wapinski I, Calderwood MA, Yildirim MA, Simonis N,
Charloteaux B, Hidalgo CA, Barbette J, Santhanam B, Brar GA, Weissman JS, Regev A,
Thierry-Mieg N, Cusick ME, Vidal M. 2012. Proto-genes and de novo gene birth. Nature
487:370–374.

24. Ruiz-Orera J, Verdaguer-Grau P, Villanueva-Cañas JL, Messeguer X, Albà MM. 2018.
Translation of neutrally evolving peptides provides a basis for de novo gene evolution. Nat
Ecol Evol 2:890–896.

25. Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, Tosatto
SCE, Paladin L, Raj S, Richardson LJ, Finn RD, Bateman A. 2021. Pfam: The protein families
database in 2021. Nucleic Acids Res 49:D412–D419.

26. Mueller S, Papamichail D, Coleman JR, Skiena S, Wimmer E. 2006. Reduction of the rate of
poliovirus protein synthesis through large-scale codon deoptimization causes attenuation of
viral virulence by lowering specific infectivity. J Virol 80:9687–9696.

27. Puigbò P, Aragonès L, Garcia-Vallvé S. 2010. RCDI/eRCDI: a web-server to estimate codon

usage deoptimization. BMC Res Notes 3:87.

28. Gelsinger DR, Dallon E, Reddy R, Mohammad F, Buskirk AR, DiRuggiero J. 2020. Ribosome

profiling in archaea reveals leaderless translation, novel translational initiation sites, and

ribosome pausing at single codon resolution. Nucleic Acids Res 48:5201–5216.

29. Eisenberg AR, Higdon AL, Hollerer I, Fields AP, Jungreis I, Diamond PD, Kellis M,

Jovanovic M, Brar GA. 2020. Translation Initiation Site Profiling Reveals Widespread

Synthesis of Non-AUG-Initiated Protein Isoforms in Yeast. Cell Syst 11:145-160.e5.

30. Ingolia NT, Lareau LF, Weissman JS. 2011. Ribosome profiling of mouse embryonic stem

cells reveals the complexity and dynamics of mammalian proteomes. Cell 147:789–802.

31. Lee S, Liu B, Lee S, Huang S-X, Shen B, Qian S-B. 2012. Global mapping of translation

initiation sites in mammalian cells at single-nucleotide resolution. Proc Natl Acad Sci USA

109:E2424-2432.

32. Mattiuzzo M, Bandiera A, Gennaro R, Benincasa M, Pacor S, Antcheva N, Scocchi M. 2007.

Role of the Escherichia coli SbmA in the antimicrobial activity of proline-rich peptides.

Molecular Microbiology 66:151–163.

33. Kuru E, Määttälä R-M, Noguera K, Stork DA, Narasimhan K, Rittichier J, Wiegand D, Church

GM. 2020. Release Factor Inhibiting Antimicrobial Peptides Improve Nonstandard Amino

Acid Incorporation in Wild-type Bacterial Cells. ACS Chem Biol 15:1852–1861.

34. Baliga C, Brown TJ, Florin T, Colon S, Shah V, Skowron KJ, Kefi A, Szal T, Klepacki D,

Moore TW, Vázquez-Laslop N, Mankin AS. 2021. Charting the sequence-activity landscape

of peptide inhibitors of translation termination. Proc Natl Acad Sci U S A 118:e2026465118.

35. Muthunayake NS, Islam R, Inutan ED, Colangelo W, Trimpin S, Cunningham PR, Chow CS.

2020. Expression and In Vivo Characterization of the Antimicrobial Peptide Oncocin and

Variants Binding to Ribosomes. Biochemistry 59:3380–3391.

36. Riley M. 1993. Functions of the gene products of Escherichia coli. Microbiol Rev 57:862–

952.

37. Blattner FR, Plunkett G, Bloch CA, Perna NT, Burland V, Riley M, Collado-Vides J, Glasner

JD, Rode CK, Mayhew GF, Gregor J, Davis NW, Kirkpatrick HA, Goeden MA, Rose DJ, Mau

B, Shao Y. 1997. The complete genome sequence of Escherichia coli K-12. Science (New

York, NY) 277:1453–1474.

38. Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in Escherichia

coli K-12 using PCR products. Proceedings of the National Academy of Sciences of the United

States of America 97:6640–5.

39. Butland G, Peregrín-Alvarez JM, Li J, Yang W, Yang X, Canadien V, Starostine A, Richards

D, Beattie B, Krogan N, Davey M, Parkinson J, Greenblatt J, Emili A. 2005. Interaction

network containing conserved and essential protein complexes in Escherichia coli. Nature

433:531–7.

40. Stringer AM, Singh N, Yermakova A, Petrone BL, Amarasinghe JJ, Reyes-Diaz L, Mantis NJ,

Wade JT. 2012. FRUIT, a scar-free system for targeted chromosomal mutagenesis, epitope

tagging, and promoter replacement in Escherichia coli and Salmonella enterica. PloS one

7:e44841–e44841.

41. Datta S, Costantino N, Court DL. 2006. A set of recombineering plasmids for gram-negative

bacteria. Gene 379:109–115.

42. Lorenz R, Bernhart SH, Höner Zu Siederdissen C, Tafer H, Flamm C, Stadler PF, Hofacker

IL. 2011. ViennaRNA Package 2.0. Algorithms Mol Biol 6:26.

43. Gertz EM, Yu Y-K, Agarwala R, Schäffer AA, Altschul SF. 2006. Composition-based

statistics and translated nucleotide searches: improving the TBLASTN module of BLAST.

BMC Biol 4:41.

44. Sievers F, Higgins DG. 2014. Clustal Omega, accurate alignment of very large numbers of

sequences. Methods Mol Biol 1079:105–116.

 Table 1. List of strains and plasmids used in this study.

Strain	Genotype/description	Source
MG1655	Wild-type	(37)
AMD052	MG1655 ΔthyA	(40)
N/A	DY330 *allR*-SPA::*kan*	(39)
AMD783	MG1655 Δ*thyA yhgP*-SPA::*kan*, constructed using JW10796 +JW10797	This study
AMD784	MG1655 Δ*thyA ytjE*-SPA::*kan*, constructed using JW10802 + JW10803	This study
AMD785	MG1655 [1995451_-]-SPA::*kan*, constructed using JW10887 + JW10888	This study
AMD786	MG1655 *ytiE*-SPA::*kan*, constructed using JW10798 + JW10799	This study
AMD787	MG1655 *yrfJ*-SPA::*kan*, constructed using JW10889 + JW10890	This study
AMD788	MG1655 [2179003_+]-SPA::*kan*, constructed using JW10800 + JW10801	This study
AMD789	MG1655 [1882207_-]-SPA::*kan*, constructed using JW10883 + JW10884	This study
AMD790	MG1655 *ykfO*-SPA::*kan*, constructed using JW10893 + JW10894	This study
AMD791	MG1655 *yoeJ*-SPA::*kan*, constructed using JW10901 + JW10902	This study
AMD792	MG1655 [2983454_+]-SPA::*kan*, constructed using JW10897 + JW10898	This study
AMD793	MG1655 *ytgB*-SPA::*kan*, constructed using JW10909 + JW10910	This study
AMD794	MG1655 [905143_-]-SPA::*kan*, constructed using JW10891 + JW10892	This study
AMD795	MG1655 *ymgN*-SPA::*kan*, constructed using JW10899 + JW10900	This study
AMD796	MG1655 [3953439_-]-SPA::*kan*, constructed using JW10905 + JW10906	This study
AMD797	MG1655 *ypeD*-SPA::*kan*, constructed using JW10907 + JW10908	This study
AMD798	MG1655 [3510920_-]-SPA::*kan*, constructed using JW10895 + JW10896	This study
AMD799	MG1655 *ynhH*-SPA:: *kan*, constructed using JW10911 + JW10912	This study
AMD800	MG1655 [4195023_+]-SPA::*kan*, constructed using JW10903 _+ JW10904	This study
AMD827	MG1655 Δ*thyA* wild-type start *ytjE-nLuc-thyA*, constructed using JW10923 + JW10924	This study
AMD828	MG1655 Δ*thyA* mutant start *ytjE-nLuc-thyA*, constructed using JW10936 + JW10924	This study
AMD829	MG1655 Δ*thyA* wild-type start *ytiE-nLuc-thyA*, constructed using JW10921 + JW10922	This study
AMD830	MG1655 Δ*thyA* mutant start *ytiE-nLuc-thyA*, constructed using JW10935 + JW10922	This study
AMD831	MG1655 Δ*thyA* wild-type start [2179003_+]-*nLuc-thyA*, constructed using JW10919 + JW10920	This study
AMD832	MG1655 Δ*thyA* mutant start [2179003_+]-*nLuc-thyA*, constructed using JW10934 + JW10920	This study
AMD833	MG1655 Δ*thyA* wild-type start [1995451_-]-*nLuc-thyA*, constructed using JW11016 + JW11018	This study
AMD834	MG1655 Δ*thyA* mutant start [1995451_-]-*nLuc-thyA*, constructed using JW11017 + JW11018	This study
AMD835	MG1655 Δ*thyA* wild-type start [1882207_-]-*nLuc-thyA*, constructed using JW11010 + JW11012	This study
AMD836	MG1655 Δ*thyA* mutant start [1882207_-]-*nLuc-thyA*, constructed using JW11011 + JW11012	This study

	MG1655 ΔthyA wild-type start [4440515_-]-*nLuc-thyA*, constructed using	
AMD837	JW11013 + JW11015	This study
	MG1655 ΔthyA mutant start [4440515_-]-*nLuc-thyA*, constructed using	
AMD838	JW11014 + JW11015	This study
	MG1655 ΔthyA wild-type start [3510920_-]-*nLuc-thyA*, constructed using	
AMD839	JW11022 + JW11024	This study
	MG1655 ΔthyA mutant start [3510920_-]-*nLuc-thyA*, constructed using	
AMD840	JW11023 + JW11024	This study
	MG1655 ΔthyA wild-type start [4195023_+]-*nLuc-thyA*, constructed using	
AMD841	JW11028 + JW11030	This study
	MG1655 ΔthyA mutant start [4195023_+]-*nLuc-thyA*, constructed using	
AMD842	JW11029 + JW11030	This study
	MG1655 ΔthyA wild-type start [2983454_+]-*nLuc-thyA*, constructed using	
AMD843	JW11025 + JW11027	This study
	MG1655 ΔthyA mutant start [2983454_+]-*nLuc-thyA*, constructed using	
AMD844	JW11026 + JW11027	This study
	MG1655 ΔthyA wild-type start [905143_-]-*nLuc-thyA*, constructed using	
AMD845	JW11019 + JW11021	This study
	MG1655 ΔthyA mutant start [905143_-]-*nLuc-thyA*, constructed using	
AMD846	JW11020 + JW11021	This study
	MG1655 ΔthyA wild-type start [3953439_-]-*nLuc-thyA*, constructed using	
AMD847	JW11031 + JW11033	This study
	MG1655 ΔthyA mutant start [3953439_-]-*nLuc-thyA*, constructed using	
AMD848	JW11032 + JW11033	This study

Plasmid	Description	Source
pKD46	p_{araBAD}-*gam-bet-exo* under the control of AraC, AmpR (Ts origin)	(38)
pSIM6	p_L-*gam-bet-exo* genes under the control of CI857 repressor, AmpR (Ts origin)	(41)
pAMD001	pGEMT-*thyA*	(40)
pNL1.1	Nanoluc (nLuc) luficerase reporter vector	Promega
pAMD375	pGEMT-nLuc-*thyA*	This study

SUPPLEMENTARY TABLES

Table S1. List of IERFs.

Table S2. List of TERFs.

Table S3. List of ORFs identified by combining Ribo-RET and Ribo-Api/Pmn.

Table S4. List of oligonucleotides used in this study.

Table S5. List of genomic regions containing non-coding genes that were excluded from the

Ribo-RET and Ribo-Api/Pmn analyses.