Bark beetles locate fungal symbionts by detecting volatile fungal metabolites of host tree resin monoterpenes

Dineshkumar Kandasamy[1,2,*$], Rashaduz Zaman[1], Yoko Nakamura[3,4], Tao Zhao[5], Henrik Hartmann[6], Martin N Andersson[2,7], Almuth Hammerbacher[8] and Jonathan Gershenzon[1*]

[1] Department of Biochemistry, Max Planck Institute for Chemical Ecology, Jena, Germany

[2] Max Planck Center for next Generation Insect Chemical Ecology (nGICE), Department of Biology, Lund University, Lund, Sweden

[3] Department of Natural Product Biosynthesis, Max Planck Institute for Chemical Ecology, Jena, Germany

[4] Reseacrch Group Biosynthesis/NMR, Max Planck Institute for Chemical Ecology, Jena, Germany

[5] School of Science and Technology, Örebro University, 701 82, Örebro, Sweden

[6] Department of Biogeochemical Processes, Max Planck Institute for Biogeochemistry, Jena, Germany

[7] Department of Biology, Lund University, Lund, Sweden

[8] Department of Zoology and Entomology, Forestry and Agricultural Biotechnology Institute, University of Pretoria, Pretoria, South Africa

[$] Current address: Department of Biology, Lund University, Lund, Sweden

*Corresponding authors

22   E-mail: dineshkumar.kandasamy@biol.lu.se (DK) and gershenzon@ice.mpg.de (JG)

Abstract

Outbreaks of the Eurasian spruce bark beetle (*Ips typographus*) have decimated millions of hectares of conifer forests in Europe in recent years. The ability of these 3-6 mm long insects to kill mature trees over a short period has been ascribed to two main factors: (1) mass attacks on the host tree to overcome tree defenses and (2) the presence of fungal symbionts that support successful beetle development in the tree. While the role of pheromones in coordinating mass attacks has been well studied, the role of chemical communication in maintaining the fungal symbiosis is poorly understood. We previously demonstrated that *I. typographus* can recognize beneficial fungal symbionts of the genera *Grosmannia*, *Endoconidiophora* and *Ophiostoma* by their *de novo* synthesized volatile compounds. Here, we hypothesize that the fungal symbionts of the bark beetles might metabolize spruce resin monoterpenes of the beetle's host tree, Norway spruce (*Picea abies*), and that the volatile products could be used as cues by beetles for locating breeding sites with beneficial symbionts. We show that *Grosmannia penicillata* and other fungal symbionts altered the profile of spruce bark volatiles by converting the major monoterpenes into an attractive blend of oxygenated derivatives. Bornyl acetate was metabolized to camphor, and α- and β-pinene to *trans*-4-thujanol and other oxygenated products. Other co-occurring non-beneficial fungi (*Trichoderma sp.*) also produce oxygenated monoterpenes, but in non-attractive ratios. Extensive electrophysiological measurements showed that the bark beetle possesses dedicated olfactory sensory neurons for oxygenated metabolites. Compounds such as camphor and *trans*-4-thujanol attracted beetles at specific doses in olfactory experiments, and the presence of symbiotic fungi enhanced attraction of

females to pheromones. Finally, we show that colonization of fungal symbionts on

spruce bark diet stimulated beetles to make tunnels into the diet. Collectively, our study

suggests that the blends of oxygenated metabolites of conifer monoterpenes produced

by fungal symbionts are used by bark beetles as attractive cues to find breeding or

feeding sites containing their essential microbial symbionts. The oxygenated

metabolites may aid beetles in assessing the presence of the fungus, the defense

status of the host tree and the density of conspecifics at potential feeding and breeding

sites.

Introduction

Many interactions between insects and their host plants are known to be mediated by volatile organic compounds [1–4]. In contrast, volatile signals between herbivorous insects and their symbiotic microbes have been less studied, aside from a few well-known examples including ambrosia beetles, termites, and the vinegar fly *Drosophila melanogaster* [1–4]. Yet, such signals could be as critical for insect fitness as their response to host plant cues. In some insect-microbe symbioses, microbes transform host plant metabolites creating volatile signals that are used by insects for food or brood site selection [3–6]. For example, yeasts vectored by *D. melanogaster* metabolize dietary phenolic antioxidants and release volatile phenolics that attract both larvae and adults to feed on antioxidant-rich foods [7]. Nevertheless, there is still little information about how microbial transformation of host plant chemicals influences insect-microbe symbioses, and whether the resulting metabolites represent honest signals of partner benefits.

Microbial symbioses are especially characteristic of wood-boring insects such as bark and ambrosia beetles. Bark beetles have captured much attention recently because of their large-scale outbreaks in many parts of the world. In Europe, for example, the Eurasian spruce bark beetle (*Ips typographus*) has killed millions of hectares of spruce stands as a result of major population bursts, likely in response to global warming and management practices, which increase forest vulnerability to epidemic outbreaks [8–12]. This bark beetle feeds and raises broods in the phloem tissue of trees, which contains high levels of terpene and phenolic defense chemicals [8,13]. This insect

overcomes its unfavorable environment by mass attacks and by introducing a suite of

microbes into the host tree, including ectosymbiotic ophiostomatoid fungi, primarily

Grosmannia penicillata, *Leptographium europhioides*, *Endoconidiophora polonica* and

Ophiostoma bicolor that cause blue staining of infected wood [14–18]. It is not yet well

understood how these free-living fungal symbionts benefit *I. typographus*. Though they

grow outside the insect, as necrotrophic fungi they may exhaust host tree defenses,

metabolize host defense compounds, and provide nutritional benefits to larvae and

adults [19–22].

Conifer oleoresins are a formidable defense against insects and pathogens, as they can

poison and physically entrap invaders [23–26]. However, the volatile fraction of the

resin, especially the monoterpenes, also plays a central role in the colonization of host

trees by bark beetles [24,27,28]. After locating a suitable tree, pioneer male *I.*

typographus oxidize the dominant host monoterpene α-pinene to *cis*-verbenol, which is

used as an aggregation pheromone in combination with the *de novo* produced 2-methyl-

3-buten-2-ol to attract conspecifics for a mass attack [29–31]. In addition to bark beetle

pheromones, several other oxygenated monoterpenes such as terpinen-4-ol, camphor,

trans-4-thujanol and borneol have also been detected at the entrance holes of *I.*

typographus galleries [32–35]. Interestingly, the phloem colonized by ophiostomatoid

fungi around these galleries also produces large amounts of oxygenated monoterpenes

compared to galleries without evident fungal growth [33]. However, the ecological

functions of these oxygenated monoterpenes remain poorly understood. In our previous

work, we showed that *I. typographus* utilizes *de novo* synthesized fungal volatiles to

maintain its association with specific beneficial symbionts and also to avoid saprophytes

[36]. However, it is unknown which volatiles are produced by these fungi when they

colonize their native substrate i.e., the phloem and sapwood of the tree.

In this study, we investigated the volatile compounds emitted when fungal symbionts of

I. typographus infect the bark of their Norway spruce (*Picea abies*) host trees. We show

that these fungi dramatically alter the volatile monoterpene composition of spruce bark

and demonstrate, using single sensillum recordings, that adult *I. typographus* can

perceive the fungal-produced monoterpenes and are attracted to these compounds in

behavioral bioassays. Our results indicate that bark beetles respond to symbiont

biotransformation products of host tree metabolites and employ them to identify suitable

sites for feeding and breeding.

Results

Ips typographus is attracted to volatiles from fungal symbionts on a spruce bark medium

Adult *I. typographus* beetles were strongly attracted to volatiles emitted by their

symbiotic fungus *G. penicillata* grown on both potato dextrose agar (PDA) or spruce

bark agar (SBA) compared to their respective fungus-free agar controls in laboratory

trap bioassays (Fig 1) (PDA, $z = 3.34$, $p = 0.001$; SBA, $z = 2.83$, $p = 0.005$, Wilcoxon's

test). Further, bark beetles showed a much stronger attraction towards *G. penicillata*

grown on SBA over the same fungus grown on PDA (Fig 1D) ($z = 4.28$, $p < 0.001$,

Wilcoxon's test). Volatiles from several other bark beetle primary and secondary fungal

symbionts, such as *E. polonica* and *L. europhioides*, grown on SBA were also highly

attractive to adult beetles, although not all bark beetle-associated fungi, nor a non-

beneficial saprophyte (*Trichoderma sp.*), tested in this way emitted attractive volatile

blends (S1 Fig).

Symbiotic fungi alter the volatile profile of the bark

The headspace volatiles of *P. abies* bark inoculated with *I. typographus* fungal

symbionts were distinct from those of fungus-free bark 4 d after inoculation when

analyzed using gas chromatography with flame ionization detection (GC-FID) and gas

chromatography-mass spectrometry (GC-MS) (S2-S7 Tables). For *G. penicillata*-

inoculated bark, PCA analysis revealed that nearly 68% of the variation in the volatile

profiles was explained by the first two principal components (Fig 2A).

Over a time course of 4, 8, 12 and 18 d post inoculation with *G. penicillata* on

detached bark plugs (Fig 2B, S3 Fig, S2, S3, and S5 Tables) 79 compounds comprising

host tree and fungal volatiles were detected and classified into different groups, namely

aliphatic hydrocarbons (17 compounds), aromatics (2 compounds), monoterpene

hydrocarbons (15 compounds), oxygenated monoterpenes) (26 compounds),

sesquiterpenes (17 compounds) and spiroketals (3 compounds). The proportion of total

oxygenated monoterpenes gradually increased to dominate the volatile profile of *G.*

penicillata-infected bark reaching a maximum at 18 d post inoculation ($F_{(3,14)}$ = 3.54, *p* =

0.04, ANOVA), while the proportion of total oxygenated monoterpenes was unchanged

in mock-inoculated controls (Fig 2B). Camphor was the major contributor to the overall

increase of oxygenated monoterpenes, with the highest relative abundance after 18 d

and a significant difference between time points ($F_{(1,16)}$ = 13.06, *p* = 0.002, ANOVA,

Tukey's test) (S5 Table). The proportion of monoterpene hydrocarbons gradually

decreased over time in both treatments (control, $F_{(3,7)}$ = 11.6 , p = 0.004; *G. penicillata*,

$F_{(3,15)}$ = 21.2, p < 0.001, ANOVA). The proportion of total sesquiterpenes also

decreased significantly over time in *G. penicillata* infected bark ($F_{(3,15)}$ = 4.4, p = 0.02,

ANOVA), but not in mock-inoculated control bark. When measuring emission rate (ng

mg DW^{-1} h^{-1}), the emission of total monoterpene hydrocarbons in the control and the *G.*

penicillata treated bark plugs was not significantly different at 4 d post inoculation (Fig

2C, S2 Table). However, there was a dramatic increase in the emission rate of total

oxygenated monoterpenes at this time point in spruce bark inoculated with *G. penicillata*

compared to the mock-inoculated control (Fig 2D) (9-fold increase, t(3.2) = 7.38, p =

0.004, Welch's t-test). Nineteen oxygenated monoterpenes were identified in the bark

inoculated with *G. penicillata*, and a total of 15 compounds significantly increased in

fungus infected bark compared to the control including camphor (Fig 2E) (S2 Table)

(51-fold increase, t(3) = 7.7, p = 0.004, Welch's t-test), *endo*-borneol (18-fold increase,

t(6) = 6.5, p < 0.001), isopinocamphone (3-fold increase, t(6) = 6.8, p < 0.001),

verbenone (2-fold increase, t(6) = 4.3, p = 0.005) and bornyl acetate (3-fold increase,

t(6) = 3.7, p = 0.01). Measurements conducted on four other *I. typographus* fungal

symbionts also showed differences in the volatile composition of fungus-inoculated

versus control bark (S2 Fig), with increases in the proportion of oxygenated

monoterpenes in bark inoculated with *L. europhioides* and *O. bicolor* over time, but not

for *E. polonica* (S3 Fig, S4, S6 and S7 Tables).

Symbiotic fungi produce oxygenated monoterpenes from spruce

monoterpene hydrocarbons and bornyl acetate

The symbiotic fungus, *G. penicillata*, significantly decreased the amount of (-)-bornyl

acetate added to PDA medium after 4 d compared to a fungus-free control (Fig 3A) (t =

-3.38, p = 0.003). However, the amounts of two other monoterpenes, (-)-β-pinene and (-

)-α-pinene, did not differ between *G. penicillata*-inoculated and fungus-free PDA.

The monoterpene metabolites of *G. penicillata* grown on either (-)-α-pinene- or (-

)-β-pinene-enriched agar were similar with the oxygenated monoterpene terpinen-4-ol

being the major product from both spruce precursors (Fig. 3B). Other oxygenated

monoterpenes formed by *G. penicillata* from (-)-α-pinene and (-)-β-pinene include (+)-

isopinocamphone, (+)-*trans*-4-thujanol and α-terpineol, with (-)-verbenone being

detected only from (-)-α-pinene (S5, S6 Figs). From (-)-bornyl acetate, the dominant

oxygenated monoterpenes of *G. penicillata* were camphor and *endo*-borneol (Fig 3B,

bottom panel; S8 Fig), and their production coincided with the decrease of the precursor

(Fig 3A). Similar results were obtained for the three other fungal symbionts tested (S4-

S8 Figs).

The volatile blend of the co-occurring, but a non-beneficial fungus, *Trichoderma*

sp., produced from a synthetic spruce monoterpene mixture (S8 Table) was distinct

from that of *G. penicillata* (S9 Fig). *Trichoderma* sp. produced mainly the oxygenated

monoterpenes, (-)-borneol and (-)-verbenone, while *G. penicillata* produced mainly (-)-

camphor and terpinen-4-ol. These results collectively show that symbiotic fungi can

dramatically alter the volatile profile of spruce bark by increasing the emission of

oxygenated monoterpenes, and the emission profile of symbiotic fungi is distinct from

that of a co-occurring fungal saprophyte.

Elimination of symbiotic ophiostomatoid fungi from *I. typographus* reduces the production of oxygenated monoterpenes in bark beetle galleries

To determine the primary source of oxygenated monoterpenes in beetle galleries, unsterilized pupae were reared into fungus-free mature adults using a diet fortified with the fungistatic agent sodium benzoate. Sterility tests and scanning electron microscopy of beetle body parts revealed that beetles reared on this diet were free from fungi and the elytral pits were devoid of fungal spores and yeasts (Fig. 4A-iii, S9 Table), though bacteria were abundant. In comparison, elytral pits of beetles reared on diet without sodium benzoate contained abundant spore masses, with structural similarity to *O. bicolor* and yeast (Fig. 4A-ii, S9 Table) (Solheim 1996). Re-inoculation of fungus-free beetles with *G. penicillata* resulted in acquisition of *G. penicillata* spores in the elytral pits (Fig. 4A-iv).

Next, we quantified the amount of oxygenated monoterpenes present in the bark galleries constructed by untreated, fungus-free, and *G. penicillata*-reinoculated beetles separately using GC-MS analyses of gallery extracts. The concentrations of (-)-camphor and terpinen-4-ol were significantly higher in the galleries of reinoculated beetles than fungus-free beetles or in bark samples without beetles at all ((-)-camphor: $F (3,23) =$ 15.6, $p < 0.001$, ANOVA; terpinen-4-ol: $H (3) = 18.6$, $p < 0.001$, Kruskal-Wallis) (Fig. 4B). On the other hand, the concentrations of these two oxygenated monoterpenes were similar when comparing galleries of *G. penicillata*-reinoculated beetles to those of untreated beetles. For (+)-isopinocamphone, the concentration in galleries constructed

by untreated beetles was significantly higher compared to bark samples without beetle

galleries, but similar to that in galleries constructed by fungus-free and *G. penicillata*-

reinoculated beetles (*F (3,23)* = 5.7, *p* = 0.005, ANOVA) (Fig. 4B). On the other hand,

the concentration of (-)-borneol in fungus-free galleries was significantly higher

compared to that in bark samples without galleries and was similar to that in the

galleries of *G. penicillata*-reinoculated and untreated beetles (*F (3,23)* = 5.3, *p* = 0.006,

ANOVA) (Fig. 4B). There were no significant differences in the concentrations of

pinocamphone, α-terpineol and (-)-verbenone among treatments.

Analysis of the microbes present in each treatment revealed that galleries constructed

by fungus-free beetles were indeed free from symbiotic ophiostomatoid fungi and yeasts

but contained many saprophytes and (or) endophytes and bacteria (S10 Table).

Microbes present in the galleries of untreated and *G. penicillata*-reinoculated beetles

were similar to those found on beetles that were used to infest the bark (S9-10 Table).

Ips typographus detects oxygenated monoterpenes through specialized olfactory sensory neurons (OSN) in their antennae

To test if *I. typographus* antennal olfactory sensilla contain OSNs that detect the

biotransformation products of monoterpenes, we challenged 231 olfactory sensilla with

a test panel comprising 92 ecologically relevant compounds diluted in paraffin oil (1 µg

µl^{-1}) using single cell recordings (S1 Table). Only 23 (~10%) of the sensilla housed

neurons that did not respond to any of the compounds from the odor panel although

their OSNs showed spontaneous firing. We obtained odor-evoked responses with

strong excitation (>120 Hz) from 198 OSNs and weak excitation (<50 Hz) from 10

OSNs, allowing the grouping of these neurons into different classes based on their

response profile. From initial screening experiments at a 10 µg dose on filter paper (to

determine the maximum receptive range of OSNs), we identified 20 classes of OSNs.

Three OSN classes responded primarily to fungal-produced oxygenated monoterpenes.

We also identified neurons belonging to previously described OSN classes tuned to

pheromones, host tree volatiles and non-host odorants [37] that are not further

considered here.

OSN classes tuned to fungal-produced oxygenated monoterpenes were

identified in both the A_m and B_m regions on the antennae (Fig 5A). One of these OSN

classes responded most strongly to (+)-isopinocamphone, and this class was highly

specific for oxygenated monoterpenes, especially ketones (Fig 5C, left panel). Apart

from (+)-isopinocamphone, relatively strong responses were also elicited by (+)-

pinocamphone, (-)-isopinocamphone, (±)-pinocarvone, (±)-camphor, and (-)-

pinocamphone. (Fig 5C, left panel). Dose-response tests showed that this OSN class

was the most sensitive to (+)-isopinocamphone of all the compounds tested with

responses evident at a dose of 100 pg. The responses to (+)-pinocamphone, (-)-

isopinocamphone, (±)-pinocarvone and (±)-camphor all appeared between 1 ng and 10

ng doses (Fig 5D, left panel). Another OSN class with specific responses to fungal-

derived compounds responded most strongly to (+)-*trans*-4-thujanol and weakly to (±)-3-

octanol, (±)-1-octen-3-ol, (+)- and (-)-terpinen-4-ol, and (+)- and (-)-α-terpineol (Fig 5C,

middle panel). This OSN showed a 1000-fold lower response threshold to (+)-*trans*-4-

thujanol compared to the next best ligands, the C_8 alcohols (Fig 5D, middle panel).

Finally, an OSN class responding strongly to verbenone, α-isophorone and β-

isophorone, followed by weaker responses to (-)- and (+)-*trans*-verbenol, pinocarvone,

and (-)-*cis*-verbenol (Fig 5C, right panel) was also found. Dose- response tests revealed

that this neuron class responded the strongest to α-isophorone across most tested

doses, followed by slightly weaker and similarly strong responses to both verbenone

and β-isophorone (Fig 5D, right panel).

A few previously characterized OSN classes for host tree monoterpenes,

including the classes with primary responses to α-pinene, *p*-cymene, and Δ3-carene,

respectively [37], showed varying secondary responses to some of the fungal-derived

compounds tested here for the first time. For example, the α-pinene OSN class

responded also to (+)-isopinocamphone, (-)-isopinocamphone and (±)-pinocarvone and

weakly to (±)-camphor, (-)-myrtenol, *trans*-pinocarveol, carvone, borneol and (-)-

fenchone (S10A Fig). The *p*-cymene OSN class showed intermediate responses to (+)-

trans-4-thujanol and carvone (S10B Fig). Although the Δ3-carene OSN class showed

high specificity towards Δ3-carene, ligands such as camphor and (-)-isopinocamphone

also elicited weak responses from this neuron class (S10C Fig).

Oxygenated monoterpenes produced by fungal symbionts attract *I.

typographus*

Grosmannia penicillata growing on spruce bark agar with added 0.1 mg g^{-1} or 0.5 mg g^{-1}

(-)-β-pinene (within the range of natural concentrations in *P. abies* bark) was much more

attractive to adult *I. typographus* than a fungus-free control medium supplemented with

(-)-β-pinene (Fig 6A, left) (0.1 mg g^{-1}, $z = 2.22$, $p = 0.02$; 0.5 mg g^{-1}, $z = 2.54$, $p = 0.01$,

Wilcoxon's test). However, addition of 1 mg g^{-1} (-)-β-pinene completely abolished the

attraction to *G. penicillata* and adult beetles were unresponsive in the binary choice

between fungus and fungus-free (-)-β-pinene-containing medium (Fig 6A, left). To

further understand these concentration-dependent responses, individual

electrophysiologically-active (-)-β-pinene biotransformation products were used in trap

bioassays against a mineral oil control. At the highest dose tested (1 mg), both *trans*-4-

thujanol and terpinen-4-ol were avoided by adult beetles (Fig 6B *trans*-4-thujanol, left

panel; terpinene-4-ol, right panel) (1 mg *trans*-4-thujanol, $z = -1.9$, $p = 0.05$; 1 mg

terpinen-4-ol, $z = -1.7$, $p = 0.08$, Wilcoxon's test). On the other hand, 100 µg *trans*-4-

thujanol significantly attracted adult beetles (Fig 6B, left panel) ($z = 2.78$, $p = 0.005$,

Wilcoxon's test). Adult beetles did not discriminate between *G. penicillata* grown on (-)-

β-pinene enriched medium and *G. penicillata* grown on non-enriched medium (Fig 6E).

Based on these results, we concluded that *I. typographus* shows concentration-specific

responses to some fungal metabolites of (-)-β-pinene, but not to (-)-β-pinene itself.

Addition of another host tree monoterpene, (-)-bornyl acetate, to fungal growth

medium at 0.05 mg g^{-1} (in the range of natural concentrations in *P. abies* bark) and 0.5

mg g^{-1} resulted in strong attraction of *I. typographus* adults towards *G. penicillata* when

tested against a fungus-free control after 4 d incubation (Fig 6C, left panel) (0.05 mg g^{-1},

$z = 3.31$, $p = 0.001$; 0.5 mg g^{-1}, $z = 3.21$, $p = 0.001$, Wilcoxon's test). The major

biotransformation product of (-)-bornyl acetate that formed in this period (Fig 6C),

camphor, was significantly more attractive to adult beetles at a 100 µg dose than the

mineral oil control (Fig 6D) ($z = 2.58$, $p = 0.01$, Wilcoxon's test). However, adult beetles

preferred *G. penicillata* grown on unenriched medium against *G. penicillata* grown on

high amount of (-)-bornyl acetate (0.5 mg g^{-1}) enriched medium (Fig 6F) ($z = 2.12$, $p =$

0.03, Wilcoxon's test) consistent with their preference for lower amounts of camphor. By

contrast, in the absence of fungus beetles did not discriminate between diet enriched

with monoterpenes and diet without monoterpenes (S11 Fig).

Volatiles of symbiotic fungi increase *I. typographus* attraction to pheromones

Ips typographus has been shown to display sex-specific responses to its aggregation

pheromone [38,39], so the sexes were tested separately with pheromones and volatiles

of their symbiotic fungi. Female beetles were significantly more attracted towards the

individual aggregation pheromone components *cis*-verbenol and 2-methyl-3-buten-2-ol

(*cis*-verbenol, $z = 2.98$, $p = 0.003$; 2-methyl-3-buten-2-ol, $z = 2$, $p = 0.046$, Wilcoxon's

test), and towards a mixture of the two pheromone components ($z = 5.19$, $p < 0.001$,

Wilcoxon's test) compared to the mineral oil control (Fig 7A). By contrast, adult males

did not make a significant choice between these options in accordance with previous

studies [42, 61]. However, when beetles had to choose between the pheromone mixture

with or without *G. penicillata* volatiles, females significantly preferred the pheromone

mixture together with *G. penicillata* volatiles (Fig 7B) ($z = 3.41$, $p = 0.001$, Wilcoxon's

test).

Symbiotic fungi increase the tunneling of adult *I. typographus*

The presence of a symbiotic fungus increased the tunneling success of adult *I.*

typographus beetles in medium after 48 h. Multiple logistic regression analysis revealed

that successful tunneling odds in (-)-bornyl acetate-amended medium were significantly

influenced by the fungus when the beetle sex and the monoterpene treatment remained

constant (Table 1) (β = 4.98, x^2 = 20.99, p < 0.001). The presence of the fungus

increased the tunneling odds by 145 units (percentage probability increased by 99%)

compared to in the absence of the fungus. Additionally, males had significantly lower

tunneling success compared to females (Table 1) (β = -1.72, x^2 = 8.78, p = 0.003) with

the tunneling odds for males being 0.18 units lower (percentage probability decreased

by 15.2%) than for females. In (-)-β-pinene-amended medium, only the presence of the

fungus significantly influenced the tunneling odds of beetles (Table 1) (β = 2.65, x^2 =

20.45, p < 0.001), with an increase of 14.06 units (percentage probability increased by

93%) compared to in the absence of the fungus. Similarly, in (-)-α-pinene-amended

medium only the presence of the fungus significantly influenced the tunneling odds of

beetles (Table 1) (β = 2.02, x^2 = 9.39, p = 0.002), with an increase of 7.51 units

(percentage probability increased by 88%) compared to the absence of the fungus.

Addition of the three monoterpenes without fungus did not have any effect on the

tunneling behavior of adult beetles.

The growth of *G. penicillata* on monoterpene-enriched media resulted in

significantly longer beetle tunnels than in the other treatments (Fig 8B-D) ((-)-β-pinene,

$F_{(3,99)}$ = 4.95, p = 0.003; (-)-α-pinene, $F_{(3,92)}$ = 14.8, p < 0.001; (-)-bornyl acetate, $F_{(3,106)}$

= 6.6, p < 0.001, ANOVA, Tukey's test). However, there was no significant difference in

tunnel lengths in treatments with fungus alone versus fungus plus monoterpenes except

for the treatment with (-)-β-pinene (Fig 8B-D, Tukey's test). The sex of the beetle and

the interaction of sex with other treatments had no effect on the tunnel length.

Discussion

The successful attack of bark beetles on their host trees is invariably associated with free-living fungal symbionts. These ascomycete symbionts may detoxify the terpene-rich defensive resin of the host tree, hasten host tree death, provide nutritional benefits, and increase resistance to pathogens, [16,17,22]. Here, we document the ability of *I. typographus* fungal symbionts to metabolize host tree monoterpenes to oxygenated derivatives that may assist adult beetles in locating suitable breeding and feeding sites [34,40]. Several oxygenated monoterpenes have been previously identified as volatiles released from trees that were attacked by *I. typographus* [32,33,35]. We showed, by removing symbiotic fungi from *I. typographus*, that these compounds were likely fungal metabolites that become dominant components of the volatile profile bark within 12 days following infection. Bark beetle olfactory sensory neurons can detect various oxygenated monoterpenes produced by symbiotic fungi, and these compounds attract adult beetles and stimulate tunneling on diets inoculated with fungal symbionts.

Oxygenated monoterpenes are widespread volatile cues in tree-feeding insects

Various forest insects that are associated with fungal symbionts, such as bark beetles, ambrosia beetles and wood wasps live in host trees producing large quantities of monoterpene volatiles. These insects are often attracted to their fungal symbionts through volatiles [2,36,41] and hence, fungus-produced monoterpene metabolites could be critical components of the attractive volatile blends. For bark beetles, oxygenated

monoterpenes derived from either the beetles, fungi or host trees are already known to
play important roles in their life history [42]. During mass attacks, the host-derived
oxygenated monoterpene *cis*-verbenol acts together with 2-methyl-3-buten-2-ol as an
aggregation pheromone for *I. typographus* to promote mass attack on individual trees
[28,43]. Beetles also utilize oxygenated monoterpenes to restrict the density of attack.
Microbes lining the gallery walls or living in the beetle gut oxidize *cis-* and *trans-*
verbenol into verbenone, which inhibits the attraction of both sexes to fully colonized
trees [44–46]. In addition, mated male *I. typographus* produce ipsenol and ipsdienol, of
which ipsenol acts as an anti-attractant [46]. Furthermore, an oxygenated monoterpene
from host trees, 1,8-cineole, which is produced in higher amounts in resistant or MeJA-
primed trees, inhibits attraction of beetles to their pheromones [47–49]. Oxygenated
monoterpenes are also used as reliable cues by parasitoids of bark beetles to locate
their prey [35,50].

Here we discovered that the oxygenated monoterpene blend emitted by bark beetle-
associated fungi growing on agar amended with spruce bark is distinct from the blend of
spruce monoterpene metabolites emitted by beetles alone [51] (S12 Fig) and attracted
adult *I. typographus*. Previously, we demonstrated that fungi grown on potato dextrose
agar without any spruce bark produced mixtures of non-terpene aliphatic and aromatic
volatiles that attract newly emerged (callow) adult beetles [36]. These compounds were
also detected here as major components of the volatile blend at later phases of fungal
growth (S3-7 Table). Although we focused principally on the symbiont *G. penicillata* in
this study, volatiles from other fungal symbionts were also investigated. *Leptographium*
europhioides and *E. polonica* also emitted volatile blends attractive to adult *I.*

typographus when growing on spruce bark agar (Fig. S1), but the volatiles of *O. bicolor*

and saprophytes such as *O. piceae* and *Trichoderma sp.* were not attractive. The

production of oxygenated monoterpenes by other co-inhabiting microbes including

fungal saprophytes, yeasts and bacteria has been reported before, but these

microorganisms produce different metabolites or the same metabolites in very different

ratios than produced by symbiotic fungi [52–54]. Since all these species produce

oxygenated derivatives of spruce bark monoterpenes (Tables S2-7) (S9 Fig), it is not

the simple presence of oxygenated monoterpenes, but the composition of the entire

volatile profile that determines its attractiveness. Notably, cross-attraction of *I.*

typographus to fungi associated with the North American spruce bark beetle *D.*

rufipennis attacking a different species of spruce has been reported recently. This

attraction could be attributed to similar oxygenated monoterpene profiles as reported

here, but it is unclear if an association with these fungi would be beneficial to *I.*

typographus [55].

411 Oxygenated monoterpenes signal the presence of fungi to *I.*
412 *typographus* and may modulate beetle colonization

The first chemical signals reported to mediate bark beetle colonization of their hosts

were aggregation pheromones. Yet even in the presence of these pheromones, a large

proportion of aggregating beetles that land on trees leave without tunneling into the bark

[40,56,57]. This suggests that other cues may be needed to induce beetles to stay and

bore into the bark. Indeed, bark beetles have been shown to respond to signals from

host and non-host species when selecting trees for colonization [28,36,58–61]. Based

on our results, fungus-produced oxygenated monoterpenes might also belong to the list

of colonization cues. For bark beetles, fungal metabolites can serve as indicators of

host tree sectors where their fungal symbionts are already established. These

compounds also provide evidence for the ongoing metabolism of host tree defenses,

which could improve the success of bark beetle colonization.

Fungal volatiles also enhance the attraction of *I. typographus* to aggregation

pheromones. Female *I. typographus* is known to use the aggregation pheromone (2-

methyl-3-buten-2-ol and *cis*-verbenol) to locate trees suitable for mating and oviposition

[40,62]. Here we showed that female *I. typographus* were more attracted at short range

to a combination of pheromone plus fungal volatiles than to pheromone alone. Similarly,

a recent study reported that the presence of fungal volatiles increased the attraction of

dispersing beetles to their pheromones in the field compared to traps with pheromones

only [63]. Oxygenated monoterpenes and other fungal volatiles provide information

about the presence of fungal symbionts, which promote the successful development of

their offspring. Similarly, the pheromone component *cis*-verbenol, itself an oxygenated

monoterpene produced by *I. typographus* from the host tree precursor α-pinene [64],

provides information about the presence of other beetles, especially males. The lack of

response of males to pheromones in our experiments is not unexpected, as male *I.*

typographus have been reported to be less responsive than females to high doses of

pheromones in walking bioassays [39,62]. This behavior may help them avoid dense

colonies of male conspecifics within a tree to reduce competition for mates and food.

Depending on how fast symbiotic fungi colonize the tree bark, fungal cues could benefit

both epidemic and endemic beetle populations. During the endemic phase, beetles

generally colonize stressed trees in a gradual fashion over months [56] and in this case,

volatiles emitted from fungal symbionts introduced into the tree by early colonizers, in

combination with beetle pheromones, could increase the probability for conspecifics to

locate attacked trees, present at low densities in spruce stands. In contrast, during

epidemics vigorous trees are typically colonized by large numbers of bark beetles within

a few days to overwhelm their defenses. Since ophiostomatoid fungi have been

reported to emit volatiles within one day following the infection of detached bark plugs in

the laboratory [65,66] these chemicals might function in attraction during natural attacks

upon epidemics as well. However, during rapid natural attacks it is unclear if fungi can

colonize a tree within few days and emit behaviorally active volatiles.

The oxygenated metabolites of host monoterpenes produced by fungal symbionts not

only attracted bark beetles, but also stimulated them to tunnel in a fungus-colonized

medium. Interestingly, both sexes showed increased tunneling in contrast to their

response to pheromones. The lack of sex-specific differences in tunneling could be due

to the fact that the nutritional advantage of feeding on fungus-colonized spruce bark is

beneficial to both sexes [17,18].

The proportion of oxygenated monoterpenes to total monoterpenes in the volatile blend

of *G. penicillata* increased over the time course studied to nearly 50% at 12 days and

nearly 70% at 18 days, a trend also observed for *L. europhioides* and *O. bicolor*. Thus,

higher proportions of oxygenated monoterpenes may indicate older fungal infection

sites and hence older beetle invasion sites that may be less attractive to newly arriving

beetles due to crowding. The lack of attraction and even repellency of higher

concentrations of individual oxygenated monoterpenes at higher concentrations, as

seen in laboratory bioassays in our own and in previous studies [67] is consistent with

this interpretation. Among the oxygenated monoterpenes, already reported to inhibit *I.*

typographus attraction, verbenone produced by microbial oxidation or auto-oxidation of

the pheromone *cis*-verbenol, reduces aggregation of *I. typographus* in later phases of

the attack on the tree [46,54,68]. The attractiveness or repellency of individual

oxygenated monoterpenes to *I. typographus* depends on their structures as well as the

stage of beetle attack cycle.

Oxygenated monoterpenes are signals not only for bark beetles, but also for their

enemies. Both beetle predators and parasitoids employ these compounds and other

volatiles to locate bark beetle larvae hidden under the bark [69]. Specifically, a three-

component blend comprising camphor, isopinocamphone and terpinen-4-ol, all fungal

metabolites of host tree monoterpene hydrocarbons, was reported to attract a

coleopteran predator and several hymenopteran parasitoids of *I. typographus* in the

presence of host tree background signals [50,70,71]. A similar mechanism is used by

parasitoids of the wood wasp, *Sirex noctilio*, which locate their concealed host insect

under the bark via the volatiles from the wood wasp fungal symbiont *Amylostereum*

areolatum [72]. Furthermore, the bark beetle predator, *Thanasimus formicarius*

possesses OSNs to detect oxygenated monoterpenes such as camphor and

pinocamphone as well as several additional OSN classes detecting various oxygenated

bark beetle pheromone compounds [73]. Thus, any benefit to the beetle arising from

oxygenated monoterpene production by its symbiotic fungi may come at the cost of

revealing its presence to natural enemies that employ these same volatiles to locate

bark beetles.

Formation of oxygenated derivatives may reduce monoterpene toxicity for *I. typographus*

The conversion of host tree monoterpene defenses by symbiotic fungi to oxygenated products may alleviate toxicity to bark beetles. Terpene-rich resins are a general defense of *P. abies* and other conifers against herbivores and pathogens [74–76]. Thus, it is not surprising that monoterpenes have exhibited toxicity to bark beetles in many studies [77–79]. Monoterpene hydrocarbons, such as α-pinene, are typically more toxic to beetles than host tree-produced oxygenated monoterpenes, such as bornyl acetate [80]. Hence the oxidative transformations carried out by fungal symbionts described in this study could reduce toxicity to *I. typographus* through conversion to less poisonous derivatives. Such detoxification of host tree defenses could represent a significant benefit of fungal symbionts [81].

By contrast, oxygenated monoterpenes may be more toxic to fungi compared to monoterpene hydrocarbons [82,83]. Thus, the initial oxidation of monoterpene hydrocarbons may not constitute a detoxification unless it is a step towards further metabolism. The potential toxicity of oxygenated monoterpenes may explain why these substances are degraded further after initial oxidation by fungi specialized on conifers such as *G. penicillata*, *Heterobasidion parviporum* and *Seridium cardinale* [82,83]. The fungus *G. clavigera*, a symbiont of the bark beetle *Dendroctonus ponderosae,* possesses genes encoding cytochromes P450 and other oxidative enzymes that are up-regulated by dietary monoterpenes [84,85]. Another class of terpenes present in *P. abies* bark and other conifers host trees of bark beetles are diterpene resin acids, which

510 are reported to be toxic to associated fungi in another bark beetle system [86]. Whether

511 bark beetles, their associated fungi or other microbes can detoxify diterpenes is not yet

512 known.

Other sources of oxygenated monoterpenes in spruce-bark beetle interactions

Oxygenated monoterpenes emitted from trees attacked by *I. typographu*s may arise

from sources other than fungal symbionts. The host tree *P. abies* itself synthesizes

large amounts of bornyl acetate [87] and small amounts of 1,8-cineole [88]. In these

compounds, the oxygen functions are incorporated during biosynthesis from basic

precursors, whereas the products from fungal symbionts are formed by oxidative

modification of a previously formed monoterpene hydrocarbon skeleton. The compound

trans-4-thujanol belongs to the latter group. We identified it as a *G. penicillata*

metabolite of α- and β-pinene, but *trans*-4-thujanol may also be synthesized by the tree,

although at low levels in *P. abies* bark [67]. As another alternative, this and other

oxygenated monoterpenes could be produced via autoxidation of monoterpene

hydrocarbons. The oxidation of monoterpenes upon exposure to air could explain the

release of low but readily detectible amounts of these compounds from uninfected

control bark plugs in our and other studies. In the field, oxygenated monoterpenes other

than bornyl acetate and 1,8-cineole have been detected from damaged *P. abies* trees

when monoterpenes were exposed to air [89,90]. However, in the present study the

emission rate from uninfected bark plugs was much lower than from fungus-infected

plugs, suggesting that microbial metabolism is a much more significant source of

oxygenated monoterpenes than autoxidation (S9 Fig) [91]. Nevertheless, the tree itself

cannot be ruled out as a source of any of the detected oxygenated monoterpenes, since

P. abies cell suspension cultures have been reported to oxidize added monoterpenes

[91,92]. If *P. abies* produced monoterpenes as a defense response only when infected

by fungi, it might be difficult to determine which organism was producing these

compounds- the tree or the fungus- without identifying specific biosynthetic genes in

one of the two genomes. However, our comparison of the volatiles produced by fungi

infecting *P. abies* bark showed large differences among fungal species (Figs. S2, S8,

S9). These seem more likely to occur if the compounds are of fungal origin. If the

compounds were produced by the tree, one might expect a very similar response to

each of these fungi, all ascomycetes.

Among microbial sources of oxygenated monoterpenes are several yeast species

including *Hansenula holstii, H. capsulata* and *Candida nitratophila*, which were isolated

from *I. typographus,* and produce terpinen-4-ol, α-terpineol, borneol and *trans-*

pinocarveol when grown in phloem medium or in α-pinene-supplemented medium [54].

In addition, several bacterial symbionts of bark beetles are capable of metabolizing

monoterpenes into oxygenated derivatives, such as verbenols and verbenone [53].

Some bacteria co-inhabit the beetle gallery together with ophiostomatoid fungi, but it is

not known if these bacteria induce the production of oxygenated monoterpenes, either

by producing them themselves or by interacting with fungi to form these compounds [92]

. Another bark beetle species, *Polygraphus poligraphus*, which is sometimes found

together with *I. typographus*, was shown to emit large amounts of terpinen-4-ol [93,94].

Intermediate amounts of α-terpineol, *cis-* and *trans*-4-thujanol were also identified from

the hindgut as well as the entrance holes of this beetle's gallery and could be formed by

microorganisms associated with this beetle species from host tree monoterpenes, or by

the beetle itself. Our experiments with fungus-free *I. typographus*, however,

demonstrated that live beetles themselves did not produce high concentrations of any of

the fungal biotransformation products identified in this study. Instead, beetles formed a

range of other oxidation products (Fig. S12) that have been identified in previous

studies [51,95].

High selectivity of *I. typographus* olfactory neurons to oxygenated monoterpenes suggest their role in detecting symbiotic fungi

Ips typographus possesses several classes of OSNs that were shown to detect the

oxygenated monoterpenes produced by their fungal symbionts with notable specificity.

For example, the isopinocamphone OSN class showed high specificity towards several

monoterpene ketones produced by fungal symbionts, including (+)- and (-)-

isopinocamphone, (+)- and (-)-pinocamphone, camphor, and pinocarvone, but not to

monoterpene alcohols such as borneol and *trans*-pinocarveol. The absence of any

response to monoterpene hydrocarbons indicates that this OSN is not tuned to detect

the host tree itself, but rather organisms metabolizing the major host monoterpenes.

The isopinocamphone OSN is similar to one recently reported OSN class from *I.*

typographus that responded best to pinocarvone and camphor (OSN class named Pcn;

isopinocamphone and pinocamphone were not tested) [90]. Our work shows that (+)-

isopinocamphone is the primary ligand of this OSN class, based on its greater activity

than the other active compounds. In addition, the response profile of this OSN class

matches very well with that of the odorant receptor (OR) ItypOR29, which recently was

characterized in *Xenopus laevis* oocytes [96].

Likewise, we showed that a previously described verbenone-sensitive OSN class [37]

also responds to *cis*- and *trans*-verbenol as well as α- and β-isophorone, compounds

that are believed to arise from bark beetle metabolism of host tree terpenes [51].

Verbenone is produced from the verbenols by microbes that colonize gallery walls and

beetle guts. Therefore, this OSN class appears to be tuned to signals from various

ecological sources providing information on bark beetle density as well as microbial

establishment [40,45,89]. Since, as discussed above, the individual oxygenated

monoterpenes arise from different organisms, including fungi, other microbes, the host

tree and the beetles themselves, the presence of OSNs responding to different

oxygenated monoterpenes may help the beetle identify the various life forms it

encounters.

Another OSN class responded most sensitively to the monoterpene alcohol *trans*-4-

thujanol, a fungal symbiont metabolite of α- and β-pinene. This OSN also responded to

the fungal metabolites terpinen-4-ol and α-terpineol, as well as C_8 alcohols, but only at

the highest doses tested, which extends prior results for this OSN [90] to other

compounds from our greatly expanded test odor panel. Strong electroantennographic

activity in *I. typographus* in response to these oxygenated monoterpenes has also been

reported [89,90], and the response spectrum of this OSN class matches well with that of

the receptor ItypOR23, which is evolutionarily related to the receptor (ItypOR29) that

detects isopinocamphone [96]. The specific responses of *I. typographus* bark beetles to

oxygenated monoterpenes may arise not only from the selectivity of the OSN classes,

but also due to further processing of the odor signal in the higher centers of the brain

[97]. Moreover, since some of the oxygenated monoterpenes elicited secondary

responses from other OSN classes, bark beetles like other insects may process

olfactory information through combinatorial codes [98], which could lead to very specific

responses to different compounds or mixtures.

Conclusion

We have shown that fungal ectosymbionts vectored by *I. typographus* metabolize host

tree monoterpene hydrocarbons to oxygenated derivatives as they infect the bark.

These oxygenated volatiles are detected by several classes of *I. typographus* olfactory

sensory neurons and lead to beetle attraction to fungi and stimulation of tunneling at

certain concentrations, but often to repulsion at higher concentrations. These

compounds appear to signal the presence of established fungi, but their effects on

beetles are context-dependent and could vary with the physiological status of the fungi,

the age, sex and density of beetles and the stage of attack [36,66]. Moreover, the

ecological roles we have proposed for these oxygenated monoterpenes are based on

laboratory assays with walking beetles, so studies under natural conditions are

necessary to confirm our findings. These compounds may also be useful in integrated

pest management strategies as attractants or repellents of bark beetles perhaps in

combination with pheromones [99–101]. In this way, oxygenated monoterpenes and

other microbial volatiles represent a rich source of untapped insect semiochemicals that

can be exploited for protecting forests from devastating pest species such as *I.*

typographus.

1 Materials and methods

1.1 Fungal strains and growth medium

The fungal strains used in this study have been previously described [36] (listed in *Table S1*). In order to obtain spores from fungi, freshly inoculated PDA plates were incubated at room temperature for 15-20 d until the mycelium was old and dark. After 20 d, plates were kept briefly at 4°C to induce sporulation. Four to six 1 cm diameter mycelium plugs were removed from each plate and inoculated into 20 mL potato dextrose broth and incubated at 25°C at 150 rpm for 4 days. Once the broth was turbid, the spores were filtered using a 40 µm EASYstrainer™ (Greiner Bio-One, Frickenhausen, Germany), and the filtrate was spun down at 4200 rcf for 10 min to precipitate the spores. The supernatant was discarded and the spore suspension was washed three times with autoclaved water and then stored at 4°C until used. The spore suspension prepared using this method was viable for several months when stored at 4°C.

1.2 Bark beetle rearing

Bark beetles were reared and stored in the laboratory as described [36]. The starting beetle culture was obtained from an infested tree in October 2017 near Jena, Thuringia, Germany. Beetles were reared throughout the year in the laboratory in freshly cut spruce logs (ca. 30 cm diameter x 50 cm height) placed in an environmental chamber set at 25°C throughout the day, 65% relative humidity and a photoperiod of 20 h per day. Beetles emerged from breeding logs after ca. 35 days and were collected

643   manually. Emerged adults were sexed based on the bristle density on their pronotum

644   [102] and stored separately in Falcon tubes lined with moist paper at 4°C at least for a

645   week before using them in bioassays. Adult beetles were used only once in bioassays.

1.3 Spruce bark diet

The outer bark of a freshly cut mature tree was scraped off gently using a drawing knife

and the inner bark (phloem) was carefully peeled off using a chisel. The bark was cut

into small pieces and ground to a fine powder in vibratory micro mill (Pulverisette 0,

Fritsch GmbH, Idar-Oberstein, Germany). The instrument was pre-cooled with liquid

nitrogen and bark pieces were pulverized at an amplitude of 2.0 for ca. 10 minutes with

addition of liquid nitrogen every two minutes to prevent thawing. The ground powder

was stored in Falcon tubes at -80°C until used for diet preparation. For preparing spruce

bark diet, 7% powdered inner spruce phloem (w/v) was added to 4% Bactoagar (Roth)

and heat sterilized at 121°C for 20 minutes.

1.4 Identification and quantification of headspace volatiles of fungal symbionts

Norway spruce bark plugs of approximately 28 mm diameter were removed from a

freshly felled tree in July, 2017 and a single bark plug was placed inside a 250 mL

volatile collection glass bottle. Before removing the bark plugs, the surface of the bark

and the cork borer were sterilized by thorough spraying with 70% ethanol in a laminar

hood. An 100 µL quantity of spore suspension ($1*10^6$ cells mL^{-1}), prepared as described

above, was added to the exposed section of the bark, and autoclaved water was added

to the control treatment. Each treatment was replicated four times including the control.

The glass bottle was secured tightly and incubated at 25°C for 4 days. After 4 d,

activated charcoal-filtered air was passed into the bottle inlet at the rate of 50 mL min^{-1}

and the outlet air was funneled through a SuperQ adsorbent filter (150 mg) for 4 hours.

Afterwards, the filters were eluted with 200 μL dichloromethane spiked with 10 ng μL^{-1}

nonyl acetate (Sigma Aldrich) as an internal standard and stored at -20°C. The spruce

bark plugs were oven dried at 80°C for 6 hours after the experiment and the dry weight

was measured.

The eluted volatile samples were subjected to GC-MS and GC-FID analysis using an

Agilent 6890 series GC (Agilent, Santa Clara, CA, USA) (injection, 1 μl splitless; flow, 2

ml min^{-1}; temperature, 45 to 180°C at 6°C min^{-1} and then to 300°C at 100°C min^{-1} for 10

min) coupled either to an Agilent 5973 quadrupole mass selective detector (interface

temperature 270 °C, quadrupole temperature 150°C, source temperature 230 °C;

electron energy 70 eV) or a flame ionization detector (FID, temp. 300 °C). The

constituents were separated on a DB-5MS column (Agilent (30 m x 0.25 mm x 0.25

μm)), with He (MS) or H_2 (FID) as carrier gas. Peaks arising from the solvent and

collection containers were identified by blank runs and excluded from the analysis. The

identity of each peak was determined by comparing its mass spectra and retention

times to those of reference libraries (NIST98 and Wiley275) and authentic standards.

The amount of each compound was calculated from the peak area obtained from the

FID detector relative to the internal standard and standardized to the spruce bark dry

weight.

1.5 Time series headspace volatile collection

For time series volatile analysis, spruce bark plugs (10 mm diameter) were removed

using a cork borer from a freshly felled spruce tree in October, 2016. Each spruce bark

plug was placed in a 15 mL clear glass vial (Supelco-Sigma-Aldrich) and 50 µL spore

suspension ($1*10^6$ cells mL^{-1}), prepared as described above, was added to treatment

plugs while control plugs received sterile water. The headspace volatiles were captured

on three polydimethylsiloxane (PDMS) sorbent silicone tubes (0.5 cm), which were hung

in each glass vial using a manually crafted metal hook attached to the bottom of

PTFE/silicone septa in the screw cap [103]. The headspace volatiles were collected

from each treatment for 2 h at 4, 8, 12 and 18 d after inoculation. After sampling,

silicone tubes were placed in 1.5 mL brown glass vials and stored at -20°C until

analysis.

Volatiles collected on PDMS tubes were analyzed using a GC-2010 plus gas

chromatograph coupled to a MS-QP2010 quadrupole mass spectrometer equipped with

a TD-20 thermal desorption unit (Shimadzu, Japan) and a GC Cryo-Trap filled with

Tenax. A single tube was placed in a 89 mm glass thermal desorption tube and

desorbed at a flow rate of 60 mL min^{-1} for 8 min at 200 °C under a stream of N_2

gas. The desorbed substances were focused on a cryogenic trap at -20 °C. The Tenax

adsorbent was heated to 230 °C and the analytes were injected using split mode

(1:100) onto a Rtx-5MS GC column (30 m x 0.25 mm x 0.25 µm) with helium as

carrier gas. Compounds were identified as above (1.5) from authentic standards and

libraries, and quantified from the area of each peak obtained using GC-MS post run

analysis software from Shimadzu. Peaks arising from contaminants from the solvent,

medium or bark plug container were identified by blank runs and excluded from the

analysis. The PLS-DA plot in Fig 2A was generated by using MetaboAnalyst 3.0

software with normalized GC-MS data (both log transformed, and range scaled) [104].

1.6 Biotransformation of host tree compounds

Experiments were conducted in 9 cm Petri dishes containing 2% potato dextrose agar

(PDA) supplemented with test solutions. The tested compounds were (-)-α-pinene, (+)-

α-pinene, (-)-β-pinene, myrcene, γ-terpinene, terpinolene, sabinene, camphene, *p*-

cymene, and (-)-bornyl acetate. Sources of these compounds are given in Table S1.

Compounds were added after dissolving in dimethyl sulfoxide (DMSO). These were

then added into PDA to reach a final concentration of 0.5 mg mL^{-1} before pouring into

Petri dishes. A 5 mm agar plug containing a fungal colony was placed in the center of

each dish and incubated at 25°C in darkness for 6 d. Each treatment was replicated four

times and for the control the PDA contained only DMSO plus monoterpene. The

headspace volatiles were collected after 4d using three PDMS tubes, which were

mounted on sterile metal wires and imbedded in PDA for one hour and stored at -20°C.

The identification and quantification of compounds were conducted in the same way as

reported for the time series (section 1.5). The headspace volatiles from fungus grown

on PDA enriched with myrcene, γ-terpinene, terpinolene, camphene and *p*-cymene did

not yield detectable amounts of monoterpene transformation products on analysis.

To identify if symbiotic fungi can reduce the amount of monoterpenes in their substrate,

fungi were grown on PDA enriched with 0.5 mg g^{-1} (-)-α-pinene, (+)-α-pinene, (-)-β-

pinene, and (-)-bornyl acetate as described above. Control plates contained only DMSO

and the tested monoterpene. After 4 d, three plugs of 6 mm diameter were removed,

weighed and transferred to 1.5 ml sterile glass vials. Agar plugs were homogenized

using sterile plastic pestles and 1 ml hexane (extraction solvent) spiked with 10 ng μL^{-1}

nonyl acetate was added and samples were vortexed for 30 s. Supernatants were

transferred to new vials and stored at -20°C until identification and quantification by GC-

MS and GC-FID (1.4). Data analysis was identical to that reported above.

1.7 Chemical synthesis of (+)-isopinocamphone and β-isophorone

(+)-isopinocamphone ((1*R*,2*R*,5*S*)-2,6,6-Trimethylbicyclo[3.1.1]heptan-3-one) was

synthesized as described in [105,106]. A mixture of (1*R*,2*R*,3*R*,5*S*)-(-)-isopinocampheol

(200 mg, 1.30 mmol, Sigma-Aldrich) and Dess-Martin-periodinane (825 mg, 1.95 mmol)

in anhydrous CH_2Cl_2 (15 mL) was stirred at room temperature for 1 hour, followed by

the addition of water and sat. aq. $NaHCO_3$ solution. The mixture was extracted twice

with methyl *t*-butyl ether. The organic phase was washed with brine, dried over

anhydrous Na_2SO_4 and concentrated *in vacuo*. The residue was purified by short-path

chromatography using an SPE cartridge (Chromabond SiOH, 6 mL, 500 mg, Macherey-

Nagel, *n*-hexane:EtOAc = 10:1) to yield (+)-isopinocamphone (159 mg, 1.04 mmol,

80%). NMR measurements were carried out on a Bruker Avance AV-500HD

spectrometer, equipped with a TCI cryoprobe using standard pulse sequences as

implemented in Bruker Topspin ver. 3.6.1. (Bruker Biospin GmbH, Rheinstetten,

Germany). Chemical shifts were referenced to the residual solvent signals of acetone-*d₆*

($δ_H$ 2.05/ $δ_C$ 29.84) or $CDCl_3$ ($δ_H$ 7.26/ $δ_C$ 77.16), respectively.

^{1}H-NMR (500 MHz, $CDCl_3$) *δ* ppm: 2.64 (*ddd*, *J*=18.6/3.0/3.0 Hz, 1H), 2.62 (*dddd*,

J=10.1/6.2/6.1/3.0 Hz, 1H), 2.52 (*bd*, *J*=18.6 Hz, 1H), 2.46 (*dq*, *J*=7.3/1.9 Hz, 1H), 2.12

(*ddd*, *J*=9.1/6.1/3.0 Hz, 1H), 2.06 (*ddd*, *J*=6.2/6.2/1.9 Hz, 1H), 1.31 (*s*, 3H), 1.21 (*d*,

J=7.3 Hz, 3H), 1.19 (*bd*, *J*=10.1 Hz, 1H), 0.88 (*s*, 3H). ^{13}C-NMR (125 MHz, CDCl$_3$) *δ*

ppm: 215.9, 51.7, 45.3, 45.1, 39.4, 39.1, 34.8, 27.4, 22.2, 17.3. GC-MS t$_R$: 13.6 min. EI-

MS (70 eV): *m/z* (%) 152 (14), 110 (15), 95 (44), 83 (89), 69 (97), 55 (100), 41(64) (S13

Fig).

β-isophorone (3,5,5-Trimethyl-3-cyclohexen-1-one). β-Isophorone was synthesized

from α-isophorone (Acros Organics, Fair Lawn, NJ, USA) following published methods

[107]. Since β-isophorone was very unstable during column chromatography [108], the

compound was used for the bioassay without purification. The purity of β-Isophorone

was 91% with 6% of α-isophorone as assessed by NMR analysis. ^{1}H-NMR (500 MHz,

acetone-*d$_6$*) *δ* ppm: 5.44 (*s*, 1H), 2.69 (*bs*, 2H), 2.27 (*s*, 2H), 1.70 (*bs*, 3H), 1.00 (*s*, 6H).

^{13}C-NMR (125 MHz, acetone-*d$_6$*) *δ* ppm: 209.3, 133.0, 130.6, 53.6, 44.2, 36.9, 29.8,

22.8. GC-MS t$_R$: 10.3 min. EI-MS (70 eV): *m/z* (%) 138 (69), 123 (68), 96 (99), 95 (84),

81 (100) (S14 Fig).

Purity of the synthesized compounds was also determined using the following GC-MS

program: injection, 1 μl splitless; flow, 2 ml min^{-1}; temperature, 45 °C (held for 2 min) to

250 °C with 6 °C min^{-1}.

1.8   Electrophysiology

Laboratory reared adult beetles from the same German culture that were used in

bioassays were used for electrophysiological single sensillum recordings (SSR) using

tungsten microelectrodes according to established methodology [36,37], using the SSR

set-up (Syntech, Kirchzarten, Germany) and odor delivery system previously described

[109]. The odor panel comprising 92 compounds consisted of beetle pheromones, host

tree, non-host tree and fungal compounds ([36], Table S2). Both major and minor

fungal volatiles identified during the chemical analysis were included in the odor panel.

All odors were dissolved in odorless paraffin oil (w/v). SSR traces were analyzed as

described [100] using Autospike 3.0 (Syntech). Males and females were initially

screened for responses to the odor panel using a high stimulus dose (10 µg on filter

paper placed inside capped standard Pasteur pipette odor cartridges [109]. OSN

classes shown to primarily respond to fungus-derived oxygenated monoterpenes were

subsequently studied in dose-response experiments with active stimuli diluted in ten-

fold steps and tested from lowest to highest dose with the least active ligands tested

first at each dose. To reduce variation due to odor depletion, stimulus cartridges were

used for a maximum of 8 stimulations during screening and 2 stimulations during dose-

response tests [110].

1.9 Trap bioassay

The arena [36] was designed so that adult beetles had to make their choice through

olfaction and not by contact cues. Fungi were inoculated on spruce bark agar-based

diet and incubated at 25°C for 4 days. With the help of a cork borer (10 mm diameter),

bark plugs with or without fungus were inserted into circular cups (1.8 cm height * 1.8

cm diameter) facing each other. Two beetles were placed inside each arena and the

olfactometer was placed inside a laminar flow cabinet in darkness. Each experiment

was replicated at least 25 times with 2 beetles per replicate. The choice of beetles was

determined periodically for up to six hours by counting the number of beetles trapped

inside the cups and represented as percentage choice (percentage of insects

responding to either control traps or treatment traps or no response). When two spruce

bark agar control plugs were tested simultaneously, adult beetles showed no preference

for traps on one side of the arena versus the other side. Preliminary experiments

showed that the sex of the beetle did not influence the olfactory response towards

fungus grown either alone or in the diet enriched with monoterpenes. Therefore, two

beetles were randomly chosen for trap bioassays.

For bioassays using terpenes, stock solutions were prepared by dissolving the

compounds in DMSO, which were then added to 7% spruce bark agar to a final

concentration of 0.05 to 1 mg g^{-1}. To determine the response of adult beetles to (-)-β-

pinene and (-)-bornyl acetate amended diet containing the fungus, *G. penicillata* was

used as this species emitted higher amounts of biotransformation products compared to

other fungi. Controls were treated with DMSO plus monoterpene (no fungus) or DMSO

plus *G. penicillata* (no monoterpene). 7% spruce bark agar plugs (10 mm)

supplemented with monoterpenes or plugs containing *G. penicillata* were placed in the

control cups, and *G. penicillata* colonized plugs from monoterpene-enriched medium

were placed in the treatment cups. The volatile emission from each control and

treatment plug used in the bioassays was determined using PDMS tubes as adsorbents

and analyzed as described previously (section 1.5). For bioassays with synthetic

compounds, stock solutions of authentic standards were prepared by dissolving them in

mineral oil (w/v) and further diluted in $\log_{10}$ steps by dissolving in mineral oil. 10 μL was

applied to 10 mm Whatmann filter paper laid on the top of spruce bark agar plugs

placed inside the cups. Control traps were treated with 10 μL paraffin oil. For the

experiment with pheromone blend in the presence of *G. penicillata* volatiles, *G.*

penicillata colonized spruce bark plugs were placed in treatment cups and 10 µL of a

pheromone mixture (*cis*-verbenol:2-methyl-3-buten-2-ol in the ratio of 1:50 diluted 1:100

in paraffin oil) was applied to filter paper as described above. Control cups were treated

with 10 µL of the pheromone mixture. Males and females were tested separately in this

experiment as *I. typographus* bark beetles show sex-specific responses to their

pheromones.

1.10 Tunneling behavior bioassay

Ips typographus tunneling behavior was studied using a protocol modified from [86]

We tested beetles in 35 ×10 mm Petri dishes (Greiner Bio-one, Frickenhausen,

Germany) filled with ca. 3 ml of spruce bark diet. The spruce bark diet was prepared

as before with some modifications: 7% (w/v) spruce inner bark powder was mixed

with 1% fibrous cellulose (Sigma), 2% glucose (Roth), and 4% Bactoagar (Roth) in

water and autoclaved for 20 minutes at 121 ºC. Before pouring the medium into the

Petri dishes, the medium was mixed with 2% solvent (DMSO: ethanol, 1:1) with 1 mg

g^{-1} of various monoterpenes ((-)-α-pinene, (-)-β-pinene and (-)-bornyl acetate) and

solvent only as a control. For treatment with fungus, 5 µl spore suspension of *G.*

penicillata (1×10^6 cells mL^{-1}) was added to the center of Petri dishes containing

monoterpene-enriched media or solvent controls and incubated at 25°C for 4 days. A

single beetle was introduced per plate, and the plates were sealed with Parafilm and

kept in the environmental chamber for 48 h under conditions described above

(section 1.2). The beetles were monitored for their tunneling activity after 2, 4, 6, 24

and 48 h with tunneling recorded as a binary event. If beetles were inside the media,

it was noted as 1 and if outside, noted as 0. After 48 h, tunnel lengths made by

beetles in each plate were measured using Image J software. Each treatment was

replicated with 15 male and female beetles.

1.11 Data analysis

IBM SPSS Statistics V25.0 was used to analyze the volatile differences between

treatments (*E. polonica*-, *G. penicillata*-, *L. europhioides*-, *O. bicolor*-, and *O. piceae*

treated bark samples and untreated control). Data were log-transformed to meet the

assumptions of normal distribution, as needed. All individual compounds assigned to

monoterpenes (MTs) or oxygenated monoterpenes were combined and their

concentrations (in dry weight (ng h^{-1} mg^{-1})) were subjected to a t-test or Welch's for

estimating differences between control and *G. penicillata* (Fig. 2). Additionally,

separate ANOVAs for all individual compounds in each group were also performed

(Table S2). For volatile time course samples, a separate ANOVA test was performed

for all individual compounds and compound groups from each fungus with time

intervals as an independent factor (Table S3-7). All ANOVA tests were followed by

Tukey's *post-hoc* tests to test for differences among treatment combinations. For

behavioral bioassays, the CI values from each experimental group were analyzed by

Wilcoxon's singed ranked test to compare the differences between control and

treatment samples. Binary data from bark beetle tunneling assays were subjected to

multiple logistic regression to analyze independent variables such as monoterpene,

sex, and fungus that influence the tunneling activity of beetles (dependent variable)

in the medium. During data analysis, the male was coded as 1 and female as 0, the

presence of fungus coded as 1 and absence of fungus as 0, tunneling inside the

medium coded as 1 and not tunneling or staying outside the medium as 0. After

testing all possible independent variables and their interactions among them, the following best-fitted logistic regression model was created to predict the odds of beetles tunneling in the different media.

Ln [odds] (tunneling odds) = β0 + β1*compound+ β2*sex + β3*fungus

Here, β0 is constant whereas β1, β2, and β3 are logistic coefficients or estimates for the parameters for compound, sex and fungus respectively. The strength of association between beetle tunneling odds and effect of monoterpenes or sex or fungus is expressed as odds ratios (OR=exp$^{\beta\cdot}$) where OR<1 indicates a negative relationship between the two events, i.e., the tunneling event is less likely to happen in response to a selected independent variable (coded as 1) in comparison with its base group (coded as 0), OR=1 indicates no relationship between two events, OR>1 shows positive relationship between two events.

Acknowledgements

We thank Dr. Hong-Lei Wang for assistance in identification of terpene metabolites of bark beetles and Erling Jirle for his help to obtain materials for rearing bark beetles in Lund. We thank Bettina Raguschke, and Daniel Veit for their assistance in the laboratory. We are grateful for the support of Stefan Engeter from ThüringenForst for permitting us to harvest spruce trees for the purpose of maintaining beetle culture in Jena. We also thank Dr. Jothi Kumar Yuvaraj for his assistance in single sensillum recordings, and Dr. Christian Schiebe, Prof. Rikard Unelius, and Prof. Fredrik Schlyter

888   for providing some of the stimuli used in SSR. We thank Ola Gustafsson for help with

889   scanning electron microscopy.

Funding Disclosure

891   Financial support was provided by the Max Planck society to DK, RZ, YN, HH, AH and

892   JG. MNA acknowledges funding from the Swedish Research Council FORMAS (grants

893   #217-2014-689 and #2018-01444) the Crafoord foundation, the Royal Physiographic

894   Society in Lund, and the Foundation in Memory of Oscar and Lily Lamm. The funders

895   had no role in study design, data collection and analysis, decision to publish and writing.

Competing Interests

897   I have read the journal's policy and I declare that the authors of this manuscript have no

898   competing interests.

Author Contributions

900   **Conceptualization:** Dineshkumar Kandasamy, Martin N Andersson, Almuth

901   Hammerbacher, Jonathan Gershenzon

902   **Data Curation:** Dineshkumar Kandasamy

903   **Formal Analysis:** Dineshkumar Kandasamy, Rashaduz Zaman

904   **Funding Acquisition:** Jonathan Gershenzon, Martin N Andersson

Investigation: Dineshkumar Kandasamy, Rashaduz Zaman, Yoko Nakamura, Tao

Zhao

Methodology: Dineshkumar Kandasamy, Martin N Andersson

Project Administration: Dineshkumar Kandasamy, Martin N Andersson, Almuth

Hammerbacher, Jonathan Gershenzon

Resources: Henrik Hartmann, Martin N Andersson, Jonathan Gershenzon

Supervision: Dineshkumar Kandasamy, Martin N Andersson, Almuth Hammerbacher,

Jonathan Gershenzon

Validation: Dineshkumar Kandasamy

Visualization: Dineshkumar Kandasamy

Writing – Original Draft Preparation: Dineshkumar Kandasamy

Writing – Review & Editing:, Jonathan Gershenzon, Almuth Hammerbacher, Martin N

Andersson, Henrik Hartmann

References

1. Becher PG, Flick G, Rozpedowska E, Schmidt A, Hagman A, Lebreton S, et al. Yeast, not fruit volatiles mediate *Drosophila melanogaster* attraction, oviposition and development. Thompson K, editor. Funct Ecol. 2012;26: 822–828. doi:10.1111/j.1365-2435.2012.02006.x

2. Hulcr J, Mann R, Stelinski LL. The scent of a partner: ambrosia beetles are attracted to volatiles from their fungal symbionts. J Chem Ecol. 2011;37: 1374–1377. doi:10.1007/s10886-011-0046-x

3.   Buser CC, Newcomb RD, Gaskett AC, Goddard MR. Niche construction initiates
the evolution of mutualistic interactions. Bonsall M, editor. Ecol Lett. 2014;17:
1257–1264. doi:10.1111/ele.12331

4.   Biedermann PHW, Kaltenpoth M. New Synthesis: The chemistry of partner choice
in insect-microbe mutualisms. J Chem Ecol. 5 Feb 201440: 99.
doi:10.1007/s10886-014-0382-8

5.   Archetti M, Scheuring I, Hoffman M, Frederickson ME, Pierce NE, Yu DW.
Economic game theory for mutualism and cooperation. Ecol Lett. Wiley/Blackwell
(10.1111); 2011. pp. 1300–1312. doi:10.1111/j.1461-0248.2011.01697.x

6.   Six DL. Niche construction theory can link bark beetle-fungus symbiosis type and
colonization behavior to large scale causal chain-effects. Curr Opin Insect Sci.
2020;39: 27–34. doi:10.1016/j.cois.2019.12.005

7.   Dweck HKM, Ebrahim SAM, Farhan A, Hansson BS, Stensmyr MC. Olfactory
proxy detection of dietary antioxidants in drosophila. Curr Biol. 2015;25: 455–466.
doi:10.1016/j.cub.2014.11.062

8.   Wermelinger B. Ecology and management of the spruce bark beetle *Ips
typographu*s—a review of recent research. For Ecol Manage. 2004;202: 67–82.
doi:10.1016/j.foreco.2004.07.018

9.   Raffa KF, Aukema BH, Bentz BJ, Carroll AL, Hicke JA, Turner MG, et al. Cross-
scale drivers of natural disturbances prone to anthropogenic amplification: the
dynamics of bark beetle eruptions. Bioscience. 2008;58: 501–517.
doi:10.1641/B580607

10.   Huang J, Kautz M, Trowbridge AM, Hammerbacher A, Raffa KF, Adams HD, et
al. Tree defence and bark beetles in a drying world: carbon partitioning, functioning
and modelling. New Phytol. 2020;225: 26–36. doi:10.1111/nph.16173

11.   Biedermann PHW, Müller J, Grégoire JC, Gruppe A, Hagge J, Hammerbacher A, et
al. Bark Beetle Population Dynamics in the Anthropocene: Challenges and
Solutions. Trends Ecol Evol. Elsevier Current Trends; 2019. pp. 914–924.
doi:10.1016/j.tree.2019.06.002

12.   Netherer S, Kandasamy D, Jirosová A, Kalinová B, Schebeck M, Schlyter F.
Interactions among Norway spruce, the bark beetle *Ips typographus* and its fungal
symbionts in times of drought. J Pest Sci (2004). 2021;94: 591–614.
doi:10.1007/s10340-021-01341-y

13.   Raffa KF, Grégoire JC, Lindgren BS. Natural history and ecology of bark beetles.
bark beetles: biology and ecology of native and invasive species. Academic Press;
2015. pp. 1–40. doi:10.1016/B978-0-12-417156-5.00001-0

14.   Kirisits T. Fungal associates of European bark beetles with special emphasis on the
963   ophiostomatoid fungi. Bark and wood boring insects in living trees in Europe, a
964   synthesis. Springer; 2007. pp. 181–236.

965   15.   Linnakoski R, Wilhelm de Beer ZB, Niemelä P, Wingfield MJ. Associations of
966   conifer-infesting bark beetles and fungi in Fennoscandia. Insects. 2012. pp. 200–
967   227. doi:10.3390/insects3010200

968   16.   Six DL, Wingfield MJ. The role of phytopathogenicity in bark beetle–fungus
969   symbioses: a challenge to the classic paradigm. Annu Rev Entomol. 2011;56: 255–
970   272. doi:10.1146/annurev-ento-120709-144839

971   17.   Six DL. Ecological and evolutionary determinants of bark beetle - Fungus
972   symbioses. Insects. 2012. doi:10.3390/insects3010339

973   18.   Six DL. The Bark Beetle Holobiont: Why Microbes Matter. J Chem Ecol. 2013;39:
974   989–1002. doi:10.1007/s10886-013-0318-8

975   19.   Wadke N, Kandasamy D, Vogel H, Lah L, Wingfield BD, Paetz C, et al. Catechol
976   dioxygenases catalyzing the first step in Norway spruce phenolic degradation are
977   key virulence factors in the bark beetle-vectored fungus *Endoconidiophora*
978   *polonica*. Plant Physiol. 2016;171: pp.01916.2015. doi:10.1104/pp.15.01916

979   20.   Boone CK, Keefover-Ring K, Mapes AC, Adams AS, Bohlmann J, Raffa KF.
980   Bacteria associated with a tree-killing insect reduce concentrations of plant defense
981   compounds. J Chem Ecol. 2013;39: 1003–1006. doi:10.1007/s10886-013-0313-0

982   21.   Six DL, Elser JJ. Extreme ecological stoichiometry of a bark beetle–fungus
983   mutualism. Ecol Entomol. 2019;44: 543–551. doi:10.1111/een.12731

984   22.   François L, Annie Y, Aurélien S. Stimulation of defenses by Ophiostomatoid fungi
985   can explain attack success of bark beetles on conifers. Ann For Sci. 2009;66: 1–22.
986   doi:10.1051/forest/2009066

987   23.   Krokene P. Conifer defense and resistance to bark beetles. Bark Beetles. Elsevier;
988   2015. pp. 177–207. doi:10.1016/B978-0-12-417156-5.00005-8

989   24.   Raffa KF, Berryman AA. The role of host plant resistance in the colonization
990   behavior and ecology of bark beetles (Coleoptera: Scolytidae). Ecol Monogr.
991   1983;53: 27–49. doi:10.2307/1942586

992   25.   Phillips MA, Croteau RB. Resin-based defenses in conifers. Trends Plant Sci.
993   1999;4: 184–190. doi:10.1016/S1360-1385(99)01401-6

994   26.   Klepzig KD, Smalley EB, Raffa KF. Combined chemical defenses against an
995   insect-fungal complex. J Chem Ecol. 1996;22: 1367–1388.
996   doi:10.1007/BF02027719

27.   Raffa KF. Terpenes tell different tales at different scales: glimpses into the
chemical ecology of conifer - bark beetle - microbial interactions. J Chem Ecol.
2014;40: 1–20. doi:10.1007/s10886-013-0368-y

28.   Raffa KF, Andersson MN, Schlyter F. Host selection by bark beetles: Playing the
odds in a high-stakes game. Adv In Insect Phys. 2016. pp. 1–74.
doi:10.1016/bs.aiip.2016.02.001

29.   Klimetzek D, Francke W. Relationship between the enantiomeric composition of α-
pinene in host trees and the production of verbenols in Ips species. Experientia.
1980;36: 1343–1345. doi:10.1007/BF01960087

30.   Lanne BS, Ivarsson P, Johnsson P, Bergström G, Wassgren AB. Biosynthesis of 2-
methyl-3-buten-2-ol, a pheromone component of *Ips typographus* (Coleoptera:
Scolytidae). Insect Biochem. 1989;19: 163–167. doi:10.1016/0020-1790(89)90087-
5

31.   Bakke A, Frøyen P, Skattebøl L. Field response to a new pheromonal compound
isolated from *Ips typographus*. Naturwissenschaften. 1977;64: 98–99.
doi:10.1007/BF00437364

32.   Leufvén A, Birgersson G. Quantitative variation of different monoterpenes around
galleries of *Ips typographus* (Colleoptera: Scolytidae) attacking Norway spruce.
Can J Bot. 1987;65: 1038–1044. doi:10.1139/b87-144

33.   Birgersson G, Leufvén A. The influence of host tree response to *Ips typographus*
and fungal attack on production of semiochemicals. Insect Biochem. 1988;18: 761–
770. doi:10.1016/0020-1790(88)90098-4

34.   Birgersson G, Bergström G. Volatiles released from individual spruce bark beetle
entrance holes Quantitative variations during the first week of attack. J Chem Ecol.
1989;15: 2465–2483. doi:10.1007/BF01020377

35.   Pettersson EM, Boland W. Potential parasitoid attractants, volatile composition
throughout a bark beetle attack. Chemoecology. 2003;13: 27–37.
doi:10.1007/s000490300003

36.   Kandasamy D, Gershenzon J, Andersson MN, Hammerbacher A. Volatile organic
compounds influence the interaction of the Eurasian spruce bark beetle (*Ips
typographus*) with its fungal symbionts. ISME J. 2019;13: 1788–1800.
doi:10.1038/s41396-019-0390-3

37.   Andersson MN, Larsson MC, Schlyter F. Specificity and redundancy in the
olfactory system of the bark beetle *Ips typographus*: Single-cell responses to
ecologically relevant odors. J Insect Physiol. 2009;55: 556–567.
doi:10.1016/j.jinsphys.2009.01.018

38.   Schlyter F, Byers JA, Löfqvist J. Attraction to pheromone sources of different
1034   quantity, quality, and spacing: Density-regulation mechanisms in bark beetle *Ips*
1035   *typographus*. J Chem Ecol. 1987;13: 1503–1523. doi:10.1007/BF01012294

1036   39.   Schlyter F, Löfqvist J. Response of walking spruce bark beetles *Ips typographus* to
1037   pheromone produced in different attack phases. Entomol Exp Appl. 1986;41: 219–
1038   230. doi:10.1111/j.1570-7458.1986.tb00532.x

1039   40.   Byers JA. Behavioral mechanisms involved in reducing competition in bark beetles.
1040   Ecography. 1989;12: 466–476. doi:10.1111/j.1600-0587.1989.tb00924.x

1041   41.   Fernández Ajó AA, Martínez AS, Villacide JM, Corley JC. Behavioural response
1042   of the woodwasp *Sirex noctilio* to volatile emissions of its fungal symbiont. J Appl
1043   Entomol. 2015;139: 654–659. doi:10.1111/jen.12211

1044   42.   Francke W, P. Vité J. Oxygenated terpenes in pheromone systems of bark beetles.
1045   Zeitschrift für Angewandte Entomologie. 1983;96: 146–156. doi:10.1111/j.1439-
1046   0418.1983.tb03655.x

1047   43.   Schlyter F, Birgersson G, Byers JA, Löfqvist J, Bergström G. Field response of
1048   spruce bark beetle, *Ips typographus*, to aggregation pheromone candidates. J Chem
1049   Ecol. 1987;13: 701–716. doi:10.1007/BF01020153

1050   44.   Leufvén A, Bergström G, Falsen E. Interconversion of verbenols and verbenone by
1051   identified yeasts isolated from the spruce bark beetle *Ips typographus*. J Chem Ecol.
1052   1984;10: 1349–1361. doi:10.1007/BF00988116

1053   45.   Cale JA, Ding R, Wang F, Rajabzadeh R, Erbilgin N. Ophiostomatoid fungi can
1054   emit the bark beetle pheromone verbenone and other semiochemicals in media
1055   amended with various pine chemicals and beetle-released compounds. Fungal Ecol.
1056   2019;39: 285–295. doi:10.1016/j.funeco.2019.01.003

1057   46.   Schlyter F, Birgersson G, Leufvén A. Inhibition of attraction to aggregation
1058   pheromone by verbenone and ipsenol - Density regulation mechanisms in bark
1059   beetle *Ips typographus*. J Chem Ecol. 1989;15: 2263–2277.
1060   doi:10.1007/BF01014114

1061   47.   Schiebe C, Hammerbacher A, Birgersson G, Witzell J, Brodelius PE, Gershenzon J,
1062   et al. Inducibility of chemical defenses in Norway spruce bark is correlated with
1063   unsuccessful mass attacks by the spruce bark beetle. Oecologia. 2012;170: 183–
1064   198. doi:10.1007/s00442-012-2298-8

1065   48.   Andersson MN, Larsson MC, Blazenec M, Jakus R, Zhang Q-H, Schlyter F.
1066   Peripheral modulation of pheromone response by inhibitory host compound in a
1067   beetle. J Exp Biol. 2010;213: 3332–3339. doi:10.1242/jeb.044396

1068   49.   Binyameen M, Jankuvová J, Blaženec M, Jakuš R, Song L, Schlyter F, et al. Co-
1069   localization of insect olfactory sensory cells improves the discrimination of closely

separated odour sources. Fox C, editor. Funct Ecol. 2014;28: 1216–1223.
doi:10.1111/1365-2435.12252

50.   Pettersson EM. Volatile attractants for three Pteromalid parasitoids attacking
concealed spruce bark beetles. Chemoecology. 2001;11: 89–95.
doi:10.1007/PL00001837

51.   Birgersson G, Schlyter F, Löfqvist J, Bergström G. Quantitative variation of
pheromone components in the spruce bark beetle *Ips typographus* from different
attack phases. J Chem Ecol. 1984;10: 1029–1055. doi:10.1007/BF00987511

52.   Grabarczyk M, Mączka W, Żołnierczyk AK, Wińska K. Transformations of
monoterpenes with the p☐menthane skeleton in the enzymatic system of bacteria,
fungi and insects. Molecules. Multidisciplinary Digital Publishing Institute; 2020.
p. 4840. doi:10.3390/molecules25204840

53.   Xu L, Lou Q, Cheng C, Lu M, Sun J. Gut-Associated Bacteria of *Dendroctonus*
valens and their involvement in verbenone production. Microb Ecol. 2015;70:
1012–1023. doi:10.1007/s00248-015-0625-4

54.   Leufvén A, Bergström G, Falsen E. Oxygenated monoterpenes produced by yeasts,
isolated from *Ips typographus* (Coleoptera: Scolytidae) and grown in phloem
medium. J Chem Ecol. 1988;14: 353–362. doi:10.1007/BF01022551

55.   Tanin SM, Kandasamy D, Krokene P. Fungal interactions and host tree preferences
in the spruce bark beetle *Ips typographus*. Front Microbiol. 2021;12: 695167.
doi:10.3389/fmicb.2021.695167

56.   Toffin E, Gabriel E, Louis M, Deneubourg JL, Grégoire JC. Colonization of
weakened trees by mass-attacking bark beetles: No penalty for pioneers, scattered
initial distributions and final regular patterns. R Soc Open Sci. 2018;5.
doi:10.1098/rsos.170454

57.   Paynter QE, Anderbrant O, Schlyter F. Behavior of male and female spruce bark
beetles, Ips typographus, on the bark of host trees during mass attack. J Insect
Behav. 1990;3: 529–543. doi:10.1007/BF01052016

58.   Erbilgin N, Krokene P, Kvamme T, Christiansen E. A host monoterpene influences
Ips typographus (Coleoptera: Curculionidae, Scolytinae) responses to its
aggregation pheromone. Agric For Entomol. 2007;9: 135–140. doi:10.1111/j.1461-
9563.2007.00329.x

59.   Zhang Q. Interruption of aggregation pheromone in *Ips typographus* (L.)(Col.
Scolytidae) by non☐host bark volatiles. Agric For Entomol. 2003;5: 145–153.

60.   Andersson MN. Mechanisms of odor coding in coniferous bark beetles: From
neuron to behavior and application. Psyche J Entomol. 2012.
doi:10.1155/2012/149572

61. Andersson MN, Löfstedt C, Newcomb RD. Insect olfaction and the evolution of
1108   receptor tuning. Front Ecol Evol. 2015;3: 53. doi:10.3389/fevo.2015.00053

1109   62. Byers JA. Chemical ecology of bark beetles. Experientia (Basel). 1989;45: 271–
1110   283.

1111   63. Jirošová A, Modlinger R, Hradecký J, Ramakrishnan R, Beránková K, Kandasamy
1112   D. Ophiostomatoid fungi synergize attraction of the Eurasian spruce bark beetle,
1113   *Ips typographus* to its aggregation pheromone in field traps. Front Microbiol.
1114   2022;0: 3652. doi:10.3389/FMICB.2022.980251

1115   64. Renwick JA, Hughes PR, Krull IS. Selective production of cis- and trans-verbenol
1116   from (-)-and (+)-alpha by a bark beetle. Science (1979). 1976;191: 199 LP – 201.
1117   doi:10.1126/science.1246609

1118   65. Zhao T, Axelsson K, Krokene P, Borg-Karlson AK. fungal symbionts of the spruce
1119   bark beetle synthesize the beetle aggregation pheromone 2-Methyl-3-buten-2-ol. J
1120   Chem Ecol. 2015;41: 848–852. doi:10.1007/s10886-015-0617-3

1121   66. Zhao T, Ganji S, Schiebe C, Bohman B, Weinstein P, Krokene P, et al. Convergent
1122   evolution of semiochemicals across Kingdoms: bark beetles and their fungal
1123   symbionts. ISME J. 2019;13: 1535–1545. doi:10.1038/s41396-019-0370-7

1124   67. Blažytė-Čereškienė L, Apšegaitė V, Radžiutė S, Mozūraitis R, Būda V, Pečiulytė
1125   D. Electrophysiological and behavioural responses of *Ips typographus* (L.) to *trans-*
1126   4-thujanol—a host tree volatile compound. Ann For Sci. 2016;73: 247–256.
1127   doi:10.1007/s13595-015-0494-5

1128   68. Kopper BJ, Klepzig KD, Raffa KF. Components of antagonism and mutualism in
1129   *Ips pini*–fungal interactions: relationship to a life history of colonizing highly
1130   stressed and dead trees. Environ Entomol. 2004;33: 28–34. doi:10.1603/0046-
1131   225X-33.1.28

1132   69. Boone CK, Six DL, Zheng Y, Raffa KF. Parasitoids and dipteran predators exploit
1133   volatiles from microbial symbionts to locate bark beetles. Environ Entomol.
1134   2008;37: 150–161. doi:10.1603/0046-225X(2008)37[150:PADPEV]2.0.CO;2

1135   70. Grégoire JC, Couillien D, Krebber R, König WA, Meyer H, Francke W.
1136   Orientation of *Rhizophagus grandis* (Coleoptera: Rhizophagidae) to oxygenated
1137   monoterpenes in a species-specific predator-prey relationship. Chemoecology.
1138   1992;3: 14–18. doi:10.1007/BF01261451

1139   71. Pettersson EM, Birgersson G, Witzgall P. Synthetic attractants for the bark beetle
1140   parasitoid *Coeloides bostrichorum* Giraud (Hymenoptera: Braconidae).
1141   Naturwissenschaften. 2001;88: 88–91. doi:10.1007/s001140100209

1142   72. Martínez AS, Fernández-Arhex V, Corley JC. Chemical information from the
1143   fungus *Amylostereum areolatum* and host-foraging behaviour in the parasitoid

Ibalia leucospoides. Physiol Entomol. 2006;31: 336–340. doi:10.1111/j.1365-3032.2006.00523.x

73. Tømmerås BÅ. Specialization of the olfactory receptor cells in the bark beetle Ips typographus and its predator *Thanasimus formicarius* to bark beetle pheromones and host tree volatiles. JComp Physiol. 1985;157: 335–341. doi:10.1007/BF00618123

74. Keeling CI, Bohlmann J. Genes, enzymes and chemicals of terpenoid diversity in the constitutive and induced defence of conifers against insects and pathogens. New Phytol. 2006;170: 657–675. doi:10.1111/j.1469-8137.2006.01716.x

75. Gershenzon J, Dudareva N. The function of terpene natural products in the natural world. Nat Chem Biol. 2007;3: 408–414. doi:10.1038/nchembio.2007.5

76. Pichersky E, Raguso RA. Why do plants produce so many terpenoid compounds? New Phytol. 2018;220: 692–702. doi:10.1111/nph.14178

77. Raffa KF, Smalley EB. Interaction of pre-attack and induced monoterpene concentrations in host conifer defense against bark beetle-fungal complexes. Oecologia. 1995;102: 285–295. doi:10.1007/BF00329795

78. Franceschi VR, Krokene P, Christiansen E, Krekling T. Anatomical and chemical defenses of conifer bark against bark beetles and other pests. New Phytol. 2005. pp. 353–376. doi:10.1111/j.1469-8137.2005.01436.x

79. Erbilgin N, Krokene P, Christiansen E, Zeneli G, Gershenzon J. Exogenous application of methyl jasmonate elicits defenses in Norway spruce (*Picea abies*) and reduces host colonization by the bark beetle *Ips typographus*. Oecologia. 2006;148: 426–436. doi:10.1007/s00442-006-0394-3

80. Chiu CC, Keeling CI, Bohlmann J. Toxicity of pine monoterpenes to mountain pine beetle. Sci Rep. 2017;7. doi:10.1038/s41598-017-08983-y

81. Hammerbacher A, Schmidt A, Wadke N, Wright LP, Schneider B, Bohlmann J, et al. A common fungal associate of the spruce bark beetle metabolizes the stilbene defenses of Norway spruce. Plant Physiol. 2013;162: 1324–1336. doi:10.1104/pp.113.218610

82. Achotegui-Castells A, della Rocca G, Llusià J, Danti R, Barberini S, Bouneb M, et al. Terpene arms race in the *Seiridium cardinale - Cupressus sempervirens* pathosystem. Sci Rep. 2016;6: 1–13. doi:10.1038/srep18954

83. Kusumoto N, Zhao T, Swedjemark G, Ashitani T, Takahashi K, Borg-Karlson AK. Antifungal properties of terpenoids in *Picea abies* against *Heterobasidion parviporum*. For Pathol. 2014;44: 353–361. doi:10.1111/efp.12106

1179   84.   Wang Y, Lim L, Madilao L, Lah L, Bohlmann J, Breuil C. Gene discovery for
1180   enzymes involved in limonene modification or utilization by the mountain pine
1181   beetle-associated pathogen *Grosmannia clavigera*. Appl Environ Microbiol.
1182   2014;80: 4566–4576. doi:10.1128/AEM.00670-14

1183   85.   Lah L, Haridas S, Bohlmann J, Breuil C. The cytochromes P450 of *Grosmannia*
1184   *clavigera*: Genome organization, phylogeny, and expression in response to pine
1185   host chemicals. Fungal GenetBiol. 2013;50: 72–81. doi:10.1016/j.fgb.2012.10.002

1186   86.   Kopper BJ, Illman BL, Kersten PJ, Klepzig KD, Raffa KF. Effects of diterpene
1187   acids on components of a conifer bark beetle–fungal interaction: Tolerance by *Ips*
1188   *pini* and sensitivity by its associate *Ophiostoma ips*. Environ Entomol. 2005;34:
1189   486–493. doi:10.1603/0046-225X-34.2.486

1190   87.   Martin DM, Tholl D, Gershenzon J, Bohlmann J. Methyl jasmonate induces
1191   traumatic resin ducts, terpenoid resin biosynthesis, and terpenoid accumulation in
1192   developing xylem of norway spruce stems. Plant Physiol. 2002;129: 1003 LP –
1193   1018. Available: http://www.plantphysiol.org/content/129/3/1003.abstract

1194   88.   Keeling CI, Weisshaar S, Ralph SG, Jancsik S, Hamberger B, Dullat HK, et al.
1195   Transcriptome mining, functional characterization, and phylogeny of a large
1196   terpene synthase gene family in spruce (*Picea spp.*). BMC Plant Biol. 2011;11.
1197   doi:10.1186/1471-2229-11-43

1198   89.   Kalinová B, Břízová R, Knížek M, Turčáni M, Hoskovec M. Volatiles from spruce
1199   trap-trees detected by *Ips typographus* bark beetles: Chemical and
1200   electrophysiological analyses. Arthropod Plant Interact. 2014;8: 305–316.
1201   doi:10.1007/s11829-014-9310-7

1202   90.   Schiebe C, Unelius CR, Ganji S, Binyameen M, Birgersson G, Schlyter F. Styrene,
1203   (+)-trans-(1R,4S,5S)-4-Thujanol and oxygenated monoterpenes related to host
1204   stress elicit strong electrophysiological responses in the bark beetle *Ips*
1205   *typographus*. J Chem Ecol. 2019; 1–16. doi:10.1007/s10886-019-01070-8

1206   91.   Hunt DWA, Borden JH, Lindgren BS, Gries G. The role of autoxidation of α-
1207   pinene in the production of pheromones of *Dendroctonus ponderosae* (Coleoptera:
1208   Scolytidae). Can J For Res. 1989;19: 1275–1282. doi:10.1139/x89-194

1209   92.   Adams AS, Currie CR, Cardoza Y, Klepzig KD, Raffa KF. Effects of symbiotic
1210   bacteria and tree chemistry on the growth and reproduction of bark beetle fungal
1211   symbionts. Can J For Rese. 2009;39: 1133–1147. doi:10.1139/X09-
1212   034/ASSET/IMAGES/X09-034T4H.GIF

1213   93.   Rahmani R, Hedenström E, Schroeder M. SPME collection and GC-MS analysis of
1214   volatiles emitted during the attack of male *Polygraphus poligraphus* (Coleoptera,
1215   Curcolionidae) on Norway spruce. Zeitschrift fur Naturforschung - Section C J
1216   Biosci. 2015;70: 265–273. doi:10.1515/znc-2015-5035

94. Rahmani R, Wallin EA, Viklund L, Schroeder M, Hedenström E. Identification and
field assay of two aggregation pheromone components emitted by males of the bark
beetle *Polygraphus punctifrons* (Coleoptera: Curculionidae). J Chem Ecol.
2019;45: 356–365. doi:10.1007/s10886-019-01056-6

95. Ramakrishnan R, Hradecký J, Roy A, Kalinová B, Mendezes RC, Synek J, et al.
Metabolomics and transcriptomics of pheromone biosynthesis in an aggressive
forest pest *Ips typographus*. Insect Biochem Mol Biol. 2022;140: 103680.
doi:10.1016/J.IBMB.2021.103680

96. Hou XQ, Yuvaraj JK, Roberts RE, Zhang DD, Unelius CR, Löfstedt C, et al.
Functional evolution of a bark beetle odorant receptor clade detecting
monoterpenoids of different ecological origins. Mol Biol Evol. 2021;38: 4934–
4947. doi:10.1093/MOLBEV/MSAB218

97. Clifford MR, Riffell JA. Mixture and odorant processing in the olfactory systems of
insects: A comparative perspective. J Comp Physiol A Neuroethol Sens Neural
Behav Physiol. 2013;199: 911–928. doi:10.1007/S00359-013-0818-6/FIGURES/3

98. Andersson MN, Löfstedt C, Newcomb RD. Insect olfaction and the evolution of
receptor tuning. Front Ecol Evol. 2015;3: 53. doi:10.3389/fevo.2015.00053

99. Davis TS, Crippen TL, Hofstetter RW, Tomberlin JK. Microbial volatile emissions
as insect semiochemicals. J Chem Ecol. 2013;39: 840–859. doi:10.1007/s10886-
013-0306-z

100. Kandasamy D, Gershenzon J, Hammerbacher A. volatile organic compounds
emitted by fungal associates of conifer bark beetles and their potential in bark
beetle control. J Chem Ecol. 2016;42: 952–969. doi:10.1007/s10886-016-0768-x

101. Beck JJ, Vannette RL. Harnessing insect-microbe chemical communications to
control insect pests of agricultural systems. J Agric Food Chem. 2017;65: 23–28.
doi:10.1021/acs.jafc.6b04298

102. Schlyter F, Cederholm I. Separation of the sexes of living spruce bark beetles, *Ips
typographus* (L.), (Coleoptera: Scolytidae). Zeitschrift für Angewandte
Entomologie. 1981;92: 42–47. doi:10.1111/j.1439-0418.1981.tb01650.x

103. Kallenbach M, Oh Y, Eilers EJ, Veit D, Baldwin IT, Schuman MC. A robust,
simple, high-throughput technique for time-resolved plant volatile analysis in field
experiments. Plant Journal. 2014;78: 1060–1072. doi:10.1111/tpj.12523

104. Chong J, Wishart DS, Xia J. Using MetaboAnalyst 4.0 for Comprehensive and
Integrative Metabolomics Data Analysis. Curr Protoc Bioinformatics. 2019;68.
doi:10.1002/cpbi.86

105. Kilbas B, Azizoglu A, Balci M. Incorporation of an Allene Unit into α-Pinene via
β-Elimination. Helv Chim Acta. 2006;89: 1449–1456.
doi:10.1002/HLCA.200690145

106. Buchbauer G, Shafii-Tabatabai A. Enones of (+)- and (−)-3-pinanone: Influence of
chirality on flavour. Flavour Fragr J. 2003;18: 441–445. doi:10.1002/FFJ.1247

107. Babler JH, Malek NC, Coghlan MJ. Selective hydrolysis of α,β-and β,γ-unsaturated
ketals: A method for deconjugation of β,β-disubstituted α,β-unsaturated ketones. J
Org Chem. 1978;43: 1821–1823. doi:10.1021/jo00403a047

108. Tückmantel W, Oshima K, Nozaki H. 1,4-Addition of triorganozincates and
silyldiorganozincates to α,β-unsaturated ketones. Chem Ber. 1986;119: 1581–1593.
doi:10.1002/CBER.19861190513

109. Andersson MN, Schlyter F, Hill SR, Dekker T. What reaches the antenna? How to
calibrate odor flux and ligand-receptor affinities. Chem Senses. 2012;37: 403–420.
doi:10.1093/chemse/bjs009

110. Andersson MN, Larsson MC, Svensson GP, Birgersson G, Rundlöf M, Lundin O,
et al. Characterization of olfactory sensory neurons in the white clover seed weevil,
Apion fulvipes (Coleoptera: Apionidae). J Insect Physiol. 2012;58: 1325–1333.
doi:10.1016/j.jinsphys.2012.07.006

1270

1271

Figure 1: *Ips typographus* **uses volatile cues from spruce bark inoculated with *G. penicillata (Gp)* to detect this symbiotic fungus.**

(A) Arena used for trap bioassays to study the behavior of *I. typographus* to volatiles emitted by a symbiotic fungus. Cups containing agar (with and without fungus) were offered to beetles as odor samples. Holes on each side of the cup allowed the beetles to smell, but not touch the agar until they entered the cup and then could no longer escape. **(B), (C)** Adult beetles chose *Gp*-colonized agar medium over fungus-free medium (*n* = 25, with two beetles per replicate). **(D)** Adult beetles chose *Gp*-inoculated spruce bark agar over *Gp*-inoculated agar without spruce bark (*n* = 25). **(B), (C), (D)** Deviation of response indices against zero was tested using Wilcoxon's test. Asterisks denotes significant differences, *P < 0.01, ***P < 0.001.

Figure 2: **Growth of *Ips typographus* symbiotic fungi on spruce bark induces increased emission of oxygenated monoterpenes.**

(A) Volatile emission pattern differed between spruce bark inoculated with *G. penicillata* (*Gp*) and uninfected bark 4 days after inoculation, as depicted in a partial least squares discriminant analysis (PLS-DA). Principal components (PC1 and PC2) explain 57.2% and 10.7% of total variation, respectively, and ellipses denote 95% confidence intervals around each treatment. Complete volatile emission data by compound for *G. penicillata* and other *I. typographus* fungal symbionts are given in Table S2. **(B)** Changes in volatile emission profiles of spruce bark due to *G. penicillata* infection over an 18 d time course. Compounds are classified into six groups by their chemical structures (*n* = 5). Complete volatile emission data by compound and time point for *G. penicillata* and other symbionts are given in Tables S3-S6. **(C)**, **(D)** Emission of specific monoterpenes from fresh spruce bark inoculated with *G. penicillata* at four days post inoculation. Identified compounds were classified into monoterpene hydrocarbons **(C)** and oxygenated monoterpenes **(D)**. The individual compounds are stacked within a single bar representing the total emission. Significant differences in the total emission levels induced by *G. penicillata* are denoted by asterisks

above the bars (*n* = 4, Welch's t-test with ns= not significant *P* < 0.05, **P* < 0.01, ***P* < 0.001). The numbers denote the identities of the compounds in the stacked bars. Complete volatile emission data are given in Table S2. **(E)** Emission rate of major oxygenated monoterpenes from spruce bark inoculated with *G. penicillata* four days post inoculation. Asterisks indicate significant differences between the spruce bark-inoculated *G. penicillata* and the fungus-free control (Welch's t-test).

Figure 3: **Metabolism of major spruce monoterpenes by a symbiotic fungus produces a variety of oxygenated monoterpenes.**

(A-B) Metabolism of major spruce monoterpenes by *G. penicillata* after fungal-infected vs. uninfected PDA was supplemented with 0.5 mg/g of (-)-β-pinene, (-)-α-pinene, and (-)-bornyl acetate. **(A)** The amounts of starting monoterpenes remaining after four days. Error bars represent SEM (n = 5). Asterisks indicate a significant difference between the PDA-inoculated *G. penicillata* and the fungus-free control (Welch's t-test) with ns = not significant, **P* < 0.01. **(B)** The most abundant metabolites of administered monoterpenes are depicted with their percentages relative to the total amounts of metabolites detected for each compound (derived from (-)-β-pinene in red; derived from (-)-α-pinene in purple; derived from (-)-bornyl acetate in green) (*n* = 4). Amounts were determined by headspace collection of volatiles from the agar (*n* = 3).

Figure 4: Oxygenated monoterpenes identified in *I. typographus*-infested spruce bark are produced by fungi associated with bark beetles

(**A**) Scanning electron micrographs of: i) an elytron of an untreated bark beetle showing ii) spores of an ophiostomatoid fungus in the elytral pit, iii) an empty elytral pit of a fungus-free beetle and iv) spore mass of *G. penicillata* in the elytral pit of a fungus-free beetle reinoculated with this fungal species. (**B**) Concentration of total oxygenated monoterpenes (i), camphor (ii), terpienen-4-ol (iii), isopinocamphone (iv), pinocamphone (v), borneol (vi), α-

terpineol (vii), and verbenone (viii) in bark without beetles, infested with fungus-free beetles, infested with fungus-free beetles reinoculated with *G. penicillata,* and untreated beetles. Different lowercase letters indicate significant differences between treatments (ANOVA, Tukey's test, *P*< 0.05).

Figure 5: **Oxygenated monoterpenes derived from fungal metabolism of host tree monoterpene hydrocarbons are detected by specialized olfactory neurons in the *I. typographus* antenna**

(A) Mapping of three classes of olfactory sensory neurons (OSN) selective for oxygenated monoterpenes (isopinocamphone; verbenone; (-)-*trans*-4-thujanol) on the antenna. Olfactory sensilla housing these OSN classes are distributed in medial (subscript "m") and lateral (subscript "l") regions of sensillum areas a and b (but not in area c) of the antenna. **(B)** Representative spike trains from an isopinocamphone-responsive neuron stimulated with 1 µg of (+)-isopinocamphone (top left) and (±)-camphor (bottom left); a verbenone-responsive neuron stimulated with 1 µg (-)-verbenone (top right); a (+)-*trans*-4-thujanol-responsive neuron stimulated with 1 µg (+)-*trans*-4-thujanol (bottom right). Black horizontal bars indicate the 0.5 s odor puffs. **(C)** Response spectra of OSN classes responding predominantly to

oxygenated monoterpenes produced by fungi at the 10 µg screening dose. The average number of spikes/second was recorded from the isopinocamphone-tuned OSN class (left) (n = 6 except (+) and (-)-pinocamphone (n = 3)), *trans*-4-thujanol-tuned OSN class (middle) (n = 5) and verbenone and isophorone-tuned OSN class (right) (n = 4) after neurons were stimulated with a panel of 97 odors. Error bars represent SEM. **(D)** Dose-response curves of the OSNs stimulated with their most active ligands: isopinocamphone-tuned OSN class (left) ((n = 9) except for (+) and (-)-pinocamphone (n = 3)), *trans*-4-thujanol-tuned OSN class (middle) (n = 3), and the OSN class tuned to isophorone and verbenone (right) (n = 3). Error bars represent SEM.

Figure 6: *Ips typographus* **is attracted to oxygenated monoterpenes produced by symbiotic fungi.**

(A) Adult beetles preferred spruce bark agar enriched with 0.1 mg/g, and 0.5 mg/g of (-)-β-pinene inoculated with *G. penicillata* for 4 days over spruce bark agar enriched with 0.1 mg/g, and 0.5 mg/g (-)-β-pinene but without fungus (left). GC-MS traces of the headspace volatiles of (-)-β-pinene-enriched agar with and without *G. penicillata* for 4 days (right) showing the oxygenated monoterpenes produced by the fungus from (-)-β-

pinene. Numbers refer to the identities of compounds. **(B)** Adult beetles chose *trans*-4-thujanol (left) at a 100 µg dose but avoided *trans*-4-thujanol (left) at a 1 mg dose diluted in mineral oil, when tested against a mineral oil control. Adult beetles showed indifferent responses to (+)-isopinocamphone (middle), and terpinen-4-ol (right), applied in various doses in mineral oil. **(C)** Adult beetles preferred spruce bark agar enriched with various amounts of (-)-bornyl acetate inoculated with *G. penicillata* for 4 days over spruce bark agar enriched with (-)-bornyl acetate but without fungus (left). GC-MS traces of the headspace volatiles of 0.5 mg/g (-)-bornyl acetate-enriched agar with and without *G. penicillata* (right) showing the oxygenated monoterpenes produced by the fungus from (-)-bornyl acetate. Numbers refer to the identities of compounds. **(D)** Adult beetles preferred ($\pm$)-camphor at a 100 µg dose against a mineral oil control, but not at other doses. **(E)** Adult beetles did not discriminate between *G. penicillata* on agar with three different concentrations of (-)-β-pinene and *G. penicillata* without (-)-β-pinene. **(F)** Adult beetles chose *G. penicillata* on agar without any administrated (-)-bornyl acetate vs. *G. penicillata* on agar with 0.5 mg/g (-)-bornyl acetate. **(A), (B), (C), (D), (E), (F)** Deviation of response indices against zero was tested using Wilcoxon's test. Asterisks denotes significant differences, *$P<0.05$, **$P<0.01$, ***$P< 0.001$. $n = 25$ for each trial.

Figure 7: Female adult *I. typographus* are attracted towards a pheromone mixture in the presence of *G. penicillata* volatiles

(A) Adult females chose traps containing 2-methyl-3-buten-2-ol (MB) and *cis*-verbenol at 10^{-2} concentration diluted in mineral oil over control traps containing mineral oil. Females strongly preferred traps containing a binary pheromone blend (*cis*-verbenol: 2-methyl-3-buten-2-ol, 50:1) diluted in mineral oil over the mineral oil control (bottom). Adult males were unresponsive to these concentrations of individual pheromones and their blend. **(B)** Adult females preferred pheromone blend in the presence of *G. penicillata* volatiles over pheromone blend without fungus. **(A), (B)**. Deviation of response indices against zero was tested using Wilcoxon's test. Asterisks denotes significant differences, *P<0.05, **P<0.01, ***P< 0.001 (*n* = 28 for each experiment).

Figure 8: *Ips typographus* **tunnels more in monoterpene-enriched diet in the presence of a symbiotic fungus.**

(A) Schematic drawing of a Petri dish arena used for the no-choice tunneling assay. The dish was filled with monoterpene-enriched spruce bark agar and inoculated with *G. penicillata* (Gp) (top). Example of the tunneling pattern of an adult beetle within the fungus-colonized diet as pictured from the bottom side of Petri dish (bottom). **(B), (C), (D)** Tunnel lengths (cm) made by adult beetles after 48 hours in diet containing *G. penicillata* only, monoterpenes only, *G. penicillata* and monoterpenes, or controls with neither *G. penicillata* nor monoterpenes. Error bars represent SEM (n = 30 (15 ♂, 15 ♀) for each trial). Monoterpenes: (-)-β-pinene **(B)**, (-)-α-pinene **(C)**, (-)-bornyl acetate **(D)**. Different lowercase letters indicate significant differences between treatments (ANOVA, Tukey's test, $P< 0.05$).

Table 1: Multiple logistic regression analysis predicting the odds of adult *I. typographus* tunneling into media enriched in different monoterpenes with and without *G. penicillata* in a no-choice assay (see also Fig 6A).

Successful tunneling in diets with[a]	Predictors	β	SE[b]	Wald χ^2	*P* value	Exp(β)	95% CI for Exp(β) Lower	Upper
(-)-Bornyl acetate	Monoterpene	-0.31	0.56	0.31	ns	0.73	0.25	2.19
	Sex	-1.72	0.58	8.78	**0.003**	0.18	0.06	0.56
	Fungus	4.98	1.09	20.99	**<0.001**	145.30	17.27	1222.52
(-)-β-Pinene	Monoterpene	-0.87	0.54	2.57	ns	0.42	0.14	1.22
	Sex	0.25	0.52	0.24	ns	1.29	0.47	3.55
	Fungus	2.65	0.58	20.45	**<0.001**	14.09	4.48	44.36
(-)-α-Pinene	Monoterpene	0.00	0.52	0.00	ns	1.00	0.36	2.75
	Sex	0.00	0.52	0.00	ns	1.00	0.36	2.75
	Fungus	2.02	0.66	9.39	**0.002**	7.51	2.07	27.28

[a]The reference category is unsuccessful tunneling
[b]Standard error of *β*
ns=not significant