Title: Multivalent, asymmetric IL-2-Fc fusions provide optimally enhanced regulatory T cell selectivity

Authors: Brian Orcutt-Jahns[1], Peter C. Emmel[1], Eli M. Snyder[1], Scott D. Taylor[1], Aaron S. Meyer[1,2,3,*]

Affiliations:

[1] Department of Bioengineering, University of California, Los Angeles, United States of America
[2] Jonsson Comprehensive Cancer Center, University of California, Los Angeles, United States of America
[3] Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research, University of California, Los Angeles, United States of America
[*]Corresponding author. Email: ameyer@asmlab.org

Classification: Research report

Keywords: Cytokines, IL-2, Immunity, Multivalency

Author Emails/Contact Information:

Brian Orcutt-Jahns: orcuttjahnsbrian@gmail.com
Peter C. Emmel: petercemmel@gmail.com
Eli M. Snyder: emsnyder@g.ucla.edu
Scott D. Taylor: sco.d.taylor@gmail.com
Aaron S. Meyer: ameyer@ucla.edu

Abstract: The common γ-chain receptor cytokines coordinate the proliferation and function of immune cell populations. One of these cytokines, interleukin (IL)-2, has potential as a therapy in autoimmune disease but is limited in effectiveness by its modest specificity toward regulatory T cells (T_{reg}s). Engineering T_{reg}-selective IL-2 has primarily focused on retaining binding to the high-affinity receptor, expressed more highly on T_{reg}s, while reducing binding to the lower affinity receptor with broader expression. However, other parameters, such as the orientation and valency of Fc fusion, have signaling effects that have never been systematically explored. Here, we systematically profiled the signaling responses to a panel of wild type and mutein IL-2-Fc fusions across time, cell types, and concentrations. Exploring these responses, we found that dimeric muteins have unique specificity for T_{reg}s through binding avidity. A mechanistic model of receptor interactions could capture these effects and directed the design of tetravalent IL-2-Fc fusions with greater T_{reg} specificity than possible with current design strategies. Exploration of other surface targets on T_{reg}s revealed that there are no other binding moieties that could be fused to IL-2 for greater selectivity. Instead, IL2Rα itself is a maximally unique surface target for T_{reg}s, and so avidity is likely the only route to more selective T_{reg} interaction. However, the binding model revealed that asymmetrical, multivalent IL-2 fusions can bias avidity effects toward IL2Rα for even further enhanced T_{reg} selectivity. These findings present a comprehensive analysis of how ligand properties and their effects on surface receptor-ligand interactions translate to selective activation of immune cell populations, and consequently reveals two new routes toward therapeutic cytokines with superior T_{reg} selectivity that can be exploited for designing selective therapies in many other contexts.

Significance Statement: Signaling in off-target immune cells has hindered the effectiveness of IL-2 as an immunotherapy. We show that IL-2-Fc fusions with higher valency can exhibit enhanced regulatory T cell selectivity. This altered selectivity is explained by the kinetics of surface receptor-ligand binding and can be quantitatively predicted using a multivalent binding model. Using these insights, we successfully develop two new strategies for IL-2 therapies with unprecedented selectivity.

Highlights:

- Current IL-2 therapies are limited by a selectivity/target potency tradeoff.
- Multivalency enhances selectivity for T_{reg}s through IL2Rα avidity.
- T_{reg} selectivity cannot be enhanced by targeting other surface protein markers.
- Multivalency can decouple selectivity from signaling using asymmetric cytokine fusions.

Main Text:

INTRODUCTION

Cytokines that bind to the common γ-chain ($γ_c$) receptor, interleukin (IL)-2, 4, 7, 9, 15, and 21, are a critical hub in modulating both innate and adaptive immune responses[1]. Each cytokine in this family binds to the common $γ_c$ receptor alongside a private receptor specific for each ligand to induce signaling. These cytokines are important control mechanisms for the activity of both effector and suppressor immune populations. For example, IL-2 can increase the effector functions of CD8[+] T cells through the upregulation of cytotoxic protein expression, as well as the suppressive capacity of T regulatory cells (T_{reg}s) through increased expression of suppressive cytokines and checkpoint proteins[2–4]. Signaling through the $γ_c$ family receptors also commonly results in lymphoproliferation across both suppressor and effector cell types; consequently, the $γ_c$ cytokines are an important

endogenous and exogenous mechanism for altering the balance of immune populations. The importance of these cytokines is observed most extremely from loss-of-function or reduced activity mutations in γ_c, which subvert T and NK cell maturation[5]. Disruptive mutations in the private receptors can lead to more selective reductions in cell types, such as T_{reg}s in the case of IL2Rα or T cells with IL-7Rα[1]. Conversely, activating mutations in these receptors, such as IL-7Rα, promote cancers such as B and T cell leukemias[6].

Due to γ_c cytokines' ability to control immune activity and abundance, they have been explored as immunotherapies in a diverse array of disease indications[7]. The most studied member of the family, IL-2, acts as both as an immunostimulant and immunosuppressant, and has been explored as a treatment for diseases ranging from cancer to autoimmunity[8–11]. Its ability to expand T_{reg} populations, particularly at low doses, has great promise as an effective treatment for autoimmune diseases such as graft vs. host disease and hepatitis-induced vasculitis[12,13]. The efficacy of these IL-2 therapies has been hindered, however, by IL-2's pleiotropic and non-specific activation of off-target immune populations, which simultaneously reduce therapeutic effectiveness and drive toxicities[14]. Enabling more selective activation of T_{reg}s is desired to reduce these detrimental effects. However, this goal has remained elusive; effector and suppressor immune populations have only subtly-differing abundances of each IL-2 receptor subunit, and no truly T_{reg}-specific marker has been discovered for targeting purposes[15]. Reducing IL-2's affinity for IL2Rβ moderately increases T_{reg} selectivity by increasing IL-2's reliance on IL2Rα which is found in greater abundance on T_{reg}s[15,16]. However, this increase in selectivity comes at the cost of potency, as IL2Rβ is necessary for signal transduction[17]. Thus, when targeting T_{reg}s, IL-2 therapies have faced a persistent tradeoff between selectivity and potency[18].

The challenges involved in engineering simultaneously cell-selective and potent forms of IL-2 have inspired varied therapeutic designs. As mentioned, the most common approach has been to alter the receptor affinities of IL-2 to weaken its interaction with IL2Rα, IL2Rβ, or both receptors[19–23]. In most cases, the wild-type cytokine or mutein is fused to an IgG Fc to take advantage of FcRn-mediated recycling for extended half-life. Fc fusion has taken many forms such as fusion to the cytokine at the N- or C-terminus, including one or two cytokines per IgG, and including or excluding Fc effector functions[15]. So-called immunocytokines have also been employed which bind to IL-2 and block certain receptor interactions to bias signaling responses[18]. More recently, cis-targeted IL-2 fusions, such as bispecific antibodies in which a non-fused Fab binds to CD8, have been designed to deliver cytokines to cytotoxic T cells[24]. However, it is unclear whether this approach can be applied to targeting T_{reg}s. These many potential therapeutic design choices necessitate design principals to direct therapeutic engineering.

Here, we systematically profile the signaling specificity effects of engineered cytokine alterations, including affinity-altering mutations and Fc-fusion formats, to map the current landscape of cell-selective IL-2 designs. Through this systematic evaluation, we identify Fc fusion valency to be an important factor in cell type selectivity. The signaling specificity of all muteins and Fc formats quantitatively matches a multivalent binding model, both between cell types and across cell-to-cell variation within a cell type, indicating that the effect of cytokine multivalency is derived from altered surface receptor binding avidity. We then use this model to demonstrate that cytokines engineered in higher valency formats are predicted to confer greater specificity towards a variety of immune cell types. These insights were experimentally validated by designing and testing two novel tetravalent IL-2 muteins in both symmetric and asymmetric forms which displayed superior T_{reg} signaling

selectivity. The superior performance of asymmetric tetravalent IL-2 fusion proteins also demonstrates how bitargeting/asymmetry can decouple targeting from signaling, enabling new therapeutic opportunities. In total, our analysis and experimental findings demonstrate that cytokine valency is an unexplored direction for further enhancing selective signaling responses and that many opportunities for using multivalency engineering exist within the γ_c cytokine family and beyond.

RESULTS

Systematic IL-2 variant profiling reveals multiple determinants of response

To explore how IL-2 mutations affect signaling across immune populations, we stimulated peripheral blood mononuclear cells (PBMCs), collected from a single donor, with 13 IL-2 muteins (Fig. 1a/b, Table S1). Our panel principally included IL-2 mutants previously developed to confer enhanced-T_{reg} selective signaling[15,22,23]. In addition to changes in receptor affinities, we included variation in two structural features: Fc fusion at either the C- or N-terminus, which has been shown to alter receptor-interaction kinetics[15], and fusion in both monomeric and dimeric formats. Stimulated cells were collected at four time points using 12 treatment concentrations. The PBMCs were then stained for canonical cell type markers and phosphorylated STAT5, a commonly used read-out of IL-2 signaling response, allowing us to separate signaling response by cell type. Four different cell types (T_{reg}, T_{helper}, CD8+, NK) were gated and quantified (Fig. S1a–d). T_{reg} and T_{helper} cells were further dissected into low, average, and high IL2Rα abundance by isolating subpopulations using three logarithmically spaced bins (Fig. S1j). For a surface-level visualization of the effects of time, cell type, receptor abundance, ligand format, ligand affinity, and concentration, we organized our signaling data into a heatmap (Fig. 1c). The complexity of the data demanded closer examination.

We selectively highlighted several dose-response curves to demonstrate the importance of our comprehensive characterization (Fig. 1d–o). First, as expected, we found that the affinity with which each IL-2 interacted with receptors divided responses (Fig. 1d–g); for example, wild-type (WT) IL-2 most potently activated all cell types, as expected given it bound to IL2Rβ with the greatest affinity (Fig. 1b, d–g). Valency played also had a prominent effect on signaling response; the bivalent Fc fusion form increased sensitivity and potency of response across all cell types.

We also found that temporal dynamics affected response characteristics (Fig. 1h–k). For example, we found that C- or N-terminus Fc-fused IL-2 demonstrated distinct responses in T_{reg}s at 1 hour of treatment but shared responses after 4 hours of treatment (Fig. 1h–k). Temporal effects are likely influenced by receptor mediated endocytosis of IL-2 receptor subunits and transcriptional changes enacted by IL-2 signaling and STAT5 phosphorylation[25,26].

Finally, we found that receptor abundance interacted with cell identity to modulate response (Fig. 1l–o). T_{reg} populations with high amounts of IL2Rα strongly responded to monovalent H16N N-term, and the bivalent form moderately enhanced this response. However, in the IL2Rα^{lo} T_{reg} cells, the effect of bivalency was even greater—only bivalent H16N induced a significant response. IL2Rα^{high} T_{helper} cells also showed a moderate increase in potency with bivalency, like the IL2Rα^{high} T_{reg} cells, but the IL2Rα^{lo} population showed no distinction between the monovalent and bivalent fusions. Thus, immune populations are further subdivided by receptor abundance into subpopulations with distinct cellular responses.

In total, the dynamics of response, cell type, concentration, ligand affinity, Fc fusion valency, and Fc fusion orientation all play roles in determining cellular response. These determinants interact in unique and often unintuitive manners, requiring a more comprehensive accounting of their effects.

Ligand valency and affinity interact to form unique cell type selectivity profiles

Given the coordinated importance of time, ligand valency, ligand affinity, cell type, and receptor expression, we next sought to focus on how ligand format affected T_{reg} selectivity. The selectivity of IL-2 for specific cell types corresponds closely to its therapeutic potency and potential toxicities[19,21,22,27].

To better understand the relationship between T_{reg} selectivity and ligand properties, we plotted the ratio of STAT5 phosphorylation (fit by a Hill curve) in T_{reg}s to that of off-target cells for each ligand across our concentration range (Fig. 2a/b/e/h). We saw that shape of each selectivity curve varied widely for each ligand and off-target cell type considered (Fig. 2b/d/f). T_{reg} selectivity quantified against CD8$^+$ and NK cells most prominently separated bivalent from monovalent ligands. Since IL2Rα affinity widely varied between our IL-2 mutants and is a known regulator of T_{reg} selectivity, we sought to understand how affinity differences contribute to T_{reg} selectivity. We plotted ligand IL2Rα affinity against the peak T_{reg} selectivity observed across concentrations (Fig. 2c/e/g). Due to the high abundance of IL2Rα displayed by T_{reg}s (Fig. S1i), we expected to see a positive correlation between IL2Rα affinity and peak T_{reg} selectivity. However, we saw that this relationship varied in a cell type- and valency-dependent manner. When considering either CD8$^+$ or NK cells, decreasing IL2Rα affinity led to decreases in the peak T_{reg} selectivity of monovalent muteins, but also, somewhat surprisingly, little relationship with the bivalent selectivity peaks (Fig. 2c/e). When considering T_{helper} populations, which have greater amounts of IL2Rα than CD8$^+$ and NK cells, we observed that decreases in IL2Rα affinity led to increases in maximum selectivity for both monovalent and bivalent muteins (Fig. 2g). Unsurprisingly, the mutein concentration at which the maximum T_{reg} selectivity occurred was higher for ligands with weaker IL2Rα affinity across all cell types (Fig. S2). In total, we show here that affinity and valency affect the selectivity profiles across ligand doses in distinct yet intertwined manners; to understand these relationships, a method mapping each of these factors simultaneously is needed.

The protein expression and abundance profile of T_{reg}s reveals limits to enhancing IL-2 T_{reg} selectivity through cis targeting

While IL2Rα is more abundant in T_{reg}s, the difference is subtle compared to some off-target cells, making selectively targeted activation more challenging[15,18]. Consequently, we wondered whether a cis targeting strategy, in which IL-2 is fused to a domain binding some other T_{reg}-specific surface marker, would provide even greater selectivity. To explore this possibility, we used a CITE-seq data set in which >211,000 human PBMCs were simultaneously analyzed for 228 surface markers coupled with single-cell RNA-seq[28]. Our previous work shows that specificity is conferred by markers expressed at a high ratio between target and off-target cells[29]. As a measure of difference, we calculated the Wasserstein distance and Kullback-Leibler divergence of each surface marker abundance and expression between T_{reg} populations and all off-target PBMCs. These complimentary distance metrics were chosen to reflect two different measures of difference: the Wasserstein distance is maximized when transforming one distribution to another would require changing the cells to a large degree, while the Kullback-Leibler distance is

maximized when the overlap between two distributions is minimized. We were surprised to find that IL2Rα was the most consistently and uniquely expressed marker on T_{reg}s by both proteomic and transcriptomic analysis (Fig 2l-o). These results were reinforced by using both a linear and non-linear classifier to identify which surface markers and transcripts were most informative for T_{reg} classification; this analysis again found that IL2Rα was optimal (Fig S3). In total, these analyses show that binding alternative surface markers cannot improve IL-2 selectivity for T_{reg}s.

Bivalent Fc-cytokine fusions have distinct cell specificity but shared dynamics

Understanding that selectivity for T_{reg}s must be derived through engineering binding to its cognate receptors, we sought to develop a more complete view of the various structural choices for IL-2 fusion design. Exploring variation in response across cell types and ligand treatments is challenging due to its multidimensional nature. Restricting ones' view to a single time point, cell type, or ligand concentration provides only a slice of the picture (Figs. 1 & 2)[15,30]. Dimensionality reduction is a generally effective tool for exploring multidimensional data. However, flattening our signaling data to two dimensions and using principal components analysis failed to help isolate the effects of concentration, ligand properties, time, and cell type (Fig. 3a). Therefore, to better resolve our data, we organized our profiling experiments into a four-dimensional tensor organized according to the ligand used, concentration, treatment duration, and cell type in the profiling. We then factored this data using non-negative canonical polyadic (CP) decomposition, a technique that represents n-dimensional tensors as additively separable patterns, themselves approximated by the outer product of dimension-specific vectors[31]. We used CP decomposition to derive factors summarizing the influence of each dimension (Fig. 3b). Three components explained roughly 90% of the variance within the dataset (Fig. 3c).

Factorization separated distinct response profiles into separate components, and the effect of each dimension (e.g., time, concentration) into separate factors. For instance, component 1 almost exclusively represented responses to wild-type cytokines (Fig. 3d), which were the only ligands which were not Fc-fused, showing a response primarily at high concentrations (Fig. 3e), with broad specificity (Fig. 3f) and a signaling profile that peaking 30 minutes and then more rapidly decreasing (Fig. 3g). An alternative way to interpret the factorization results is to compare profiles within a single factor. For example, component 1 led to a less sustained profile of signaling response as compared to the other signaling patterns (Fig. 3g).

Remarkably, components 2 and 3 cleanly separated ligands conjugated in bivalent or monovalent forms, respectively (Fig. 3d/h). In fact, ligand valency was represented more prominently than differences in receptor affinity between muteins. Component 2 had uniquely high T_{reg} specificity (Fig. 3f) and was most represented at intermediate concentrations (Fig. 3e). Component 2 was also highly correlated with IL2Rα abundance in subsets of T_{reg} and T_{helper} cells, suggesting that bivalent molecule's specificity for T_{reg}s is mediated by their higher abundance of IL2Rα. Component 3 had a broader cell response (Fig. 3f) and increased monotonically with concentration (Fig. 3e). Despite these strong differences in specificities, both components had nearly identical time dynamics (Fig. 3g). While other ligand variation influenced the potency and selectivity of each ligand, only the bivalent Fc fusions, regardless of their receptor affinities, more highly weighted component 2, where T_{reg} response was highly weighted, over components 1 and 3, which represented effector cell response (Fig. 3h). In total, these results indicated that mono- and multivalent

cytokines shared identical dynamics and that, while Fc fusion and affinity modulation affect
response, ligand valency was a critical and prominent determinant of specificity.

Variation in IL-2 responses is explained by a simple multivalent binding model

Having observed that T_{reg} selectivity is prominently enhanced by multivalency, we sought to
determine whether cell surface binding on its own could explain these selectivity
differences. To do so, we applied a two-step, equilibrium, multivalent binding model to
predict IL-2 response, assuming that signaling response was proportional to the amount of
active receptor-ligand complexes[32]. Within the model, ligand binding first occurs with
kinetics equivalent to the single binding site, and then subsequent interactions occur
proportionally to affinity, adjusted by K_x^*, a cross-lining constant that corrects for differences
between monovalent and multivalent interactions. We fit this model to our signaling
experiments and evaluated its concordance with the data. The model is very simple, with
the cross-linking parameter being the only non-scaling fit parameter; this parameter had an
optimum at 1.2×10^{-11} #/cell, consistent with that seen for other receptor families[33–35].
Overall, we observed remarkable consistency between predicted and observed responses
($R^2 = 0.85$), and accuracy was maintained when examining data subsets including
individual cell types and ligands (Fig. 4b–c).

To ensure that our model was not simply capturing a trend towards higher signaling with
increasing concentration, we examined our model's accuracy within specific cytokine
concentrations (Fig. 4d). Our model did not predict response at the lowest concentrations
as there was little to no response in the data itself but increased in accuracy at
concentrations where responses were observed. Finally, we examined how the model's
accuracy varied within each timepoint; initial responses (30 mins) were most accurately
predicted, with a slight decrease in accuracy over longer timescales (Fig. 4e). This is
expected, given that longer treatments likely involve various compensatory mechanisms
such as the degradation or increased transcription of IL-2 receptor subunits[26,36]. In total,
multivalent cell surface binding showed quantitative agreement with the pattern of cell-type-
specific responses to IL-2 muteins, supporting that the specificity enhancement of bivalency
is derived from receptor avidity effects and is explained by a simple model of cell surface
binding.

Upon finding that our model was broadly predictive of cell-type specific signaling
responses, we sought to use our model to understand and visualize how valency and
affinity interact to determine T_{reg} selectivity. Here, our model showed that T_{reg} response is
strongly governed by IL2Rα affinities and that these effects have an exceptionally strong
relationship with valency, particularly at intermediate cytokine doses, while NK signaling
barely varied across ligands of varying affinities (Fig. 4f/g). We then used the model to
explore how receptor abundance affects multivalent ligand binding (Fig. 4h/i). Here,
theoretical cell populations expressing 10^4 IL2Rα and 10^3 IL2Rβ molecules varied widely in
their response to multivalent IL-2 muteins (Fig. 4h), while cells expressing very few IL2Rα
receptors and the same abundance of IL2Rβ barely altered in their response (Fig. 4i).
These results demonstrate how multivalent cytokines with high IL2Rα affinities uniquely
and selectively target T_{reg}s through IL2Rα-mediated avidity effects.

Multivalency provides a general strategy for enhanced signaling selectivity and guides the development of superior IL-2 muteins

Given that a simple binding model accurately predicted cell type-specific responses to IL-2 and that bivalent, Fc-fused IL-2 muteins have favorable specificity properties, we computationally explored to what extent multivalency might be a generally useful strategy. While monovalent ligand binding scales linearly with receptor abundance, multivalent ligands bind nonlinearly depending upon receptor abundance[37]. Thus, multivalent ligands should be able to selectively target cells with uniquely high expression of certain γ_c family receptors.

Valency enhancements are only apparent with coordinated changes in receptor-ligand binding affinities[29]. Therefore, we optimized the receptor affinities of simulated ligands while varying valency. We first designed IL-2 muteins of varying valency to obtain optimal T_{reg} specificity (Fig. 5a). As expected, ligand valency increased achievable selectivity past that possible using a monovalent cytokine format at any receptor affinities. Muteins of higher valency required reduced IL2Rα affinity to achieve optimal T_{reg} selectivity (Fig. 5b).

We then explored whether IL-2 muteins lacking IL2Rα binding could selectively target NK cells, based on their uniquely high expression of IL2Rβ[15], with similar results; IL-2 muteins of higher valency were predicted to be increasingly selective for activation of NK cells, so long as IL2Rβ/γ_c affinity was coordinately decreased (Fig. 5c,d). Finally, we explored whether multivalent IL-7 could be used to target T_{helper}s, as they express high amounts of IL-7Rα (Fig. S1i). We again found that ligands of higher valency should achieve higher selectivity for these cells, but that the benefits of valency were less than the targeting of T_{reg}s or NK cells using IL-2 mutants because CD8+ T cells have similar IL7Rα amounts (Fig. 5e). These benefits were again contingent on decreasing IL-7Rα affinity at higher valency (Fig. 5f).

To experimentally show that muteins of higher valency could be engineered to increase T_{reg} selectivity, we expressed and purified three Fc fusions of R38Q/H16N IL-2 in monovalent, bivalent, and tetravalent formats (Fig. S6). PBMCs from three donors were stimulated for 30 minutes and stained for cell type markers as well as pSTAT5. Tetravalent IL-2 was designed by Fc-fusing IL-2 muteins at both the C- and N-terminus and allowing the Fc to dimerize (Fig. 5g). R38Q/H16N was selected as the mutant closest to optimal binding affinities in tetravalent form, though further optimization is possible (Figs. 1b & 5b). As predicted, valency increased the responsiveness of both T_{reg}s and off-target immune cells at each concentration (Fig. S7). However, the T_{reg} response increase far exceeded that in off-target cells; consequently, tetravalent R38Q/H16N was able to achieve much greater T_{reg} selectivity than bivalent and monovalent formats, two IL-2 fusions with already superior selectivity (Fig. 5h–j).

In total, these results show that valency beyond bivalency has unexplored potential for engineered cytokines with enhanced therapeutic potency alongside reduced toxicity. They also show the benefit of mechanistic modeling to guide ligand design, particularly when ligand affinity must be considered alongside other parameters such as valency.

Bitargeted IL-2–Fc fusions demonstrate even greater T_{reg} selectivity

Through the CITE-seq data analysis, we found that IL2Rα was the optimal surface target for T_{reg} selectivity (Fig. 2i–m). This result was further strengthened when we integrated

335 these data with our binding model and ligand optimization approach. We used the model to consider whether WT IL-2 fused to a selective binder for any surface markers could increase T_{reg} selectivity. WT IL-2 fusion to an IL2Rα binder was predicted to enhance T_{reg} selectivity over off-target immune cells (Fig. 6a/b). This was initially surprising because IL-2 itself binds IL2Rα but indicated to us that multivalent complexes provide the potential

340 opportunity to decouple T_{reg} selectivity (by binding IL2Rα) from cytokine potency.

To decouple cell-selective binding from signaling response, we expressed an asymmetric Fc fusion including both a signaling-competent R38Q/H16N IL-2 and signaling-deficient V91K/D20A/M104V IL-2 with only IL2Rα binding (Fig. 6c, Fig. S6). V91K/D20A/M104V IL-2 is reported to effectively eliminate IL2Rβ binding while maintaining IL2Rα interaction[23]. We

345 henceforth refer to this asymmetric construct as "bitargeted" IL-2 to reflect its inclusion of two IL-2 muteins with separate signaling and targeting roles. Bivalent bitargeted ligand was designed by introducing Fc mutations preventing Fc dimerization (Fig. 6c). Tetravalent bitargeted constructs were predicted to have greater T_{reg} specificity than their non-bitargeted counterparts for any IL2Rβ affinity, using either the IL2Rα affinity of WT or

350 R38Q/H16N (Fig. 6d). We tested our bivalent and tetravalent bitargeted constructs by again stimulating PBMCs and quantifying their pSTAT5 responses. We found that both bivalent and tetravalent bitargeted ligand increased or maintained high potency in T_{reg} cells (Fig. S8). This potency translated to greater T_{reg} selectivity for both bivalent and tetravalent forms of bitargeted, both of which outperformed any previously characterized monovalent

355 and bivalent IL-2 fusions, and modestly outperformed tetravalent R38Q/H16N (Fig. 6e–f).

We again applied non-negative CP decomposition of the R38Q/H16N and bitargeted ligand signaling responses to summarize our ligand engineering efforts (Fig. 6h–k). Three components captured >90% of the variation in the data. Here, T_{reg} responses were primarily represented by component 2, effector T cell responses by component 1, and NK

360 cell responses by component 3 (Fig. 6i). Valency again determined ligand component 2 weight most potently, with both tetravalent constructs, bitargeted or R38Q/H16N, demonstrating highest component 2 weight. However, while the bivalent bitargeted construct demonstrated no NK cell activity (component 3), both tetravalent constructs had some NK activity (Fig. 6j). Thus, our experimental results demonstrate that tradeoffs still

365 exist between T_{reg} potency and selectivity. As before, T_{reg} selectivity was maximized at low and intermediate dosages (Fig. 6k). Despite these tradeoffs, the selectivity demonstrated by both model-guided "cis"-targeting or higher valency fusion formats vastly improved upon the selectivity possible with existing approaches.

DISCUSSION

370 Here, we systematically explored how ligand properties determine signaling response and specificity across 13 engineered IL-2 variants (Fig. 1, 2). Our study is unique in its inclusion of clinically relevant muteins alongside variation in Fc fusion format. Dimensionality reduction in tensor form identified how ligand properties modulate response, revealing that multivalent cytokines have unique specificity advantages (Fig. 3). Using a multivalent

375 binding model, we uncovered that this unique specificity arises from surface binding avidity effects (Fig. 4). This model indicated that cytokines of higher valency can offer even greater cell type selectivity (Fig. 5), and we demonstrated this strategy experimentally by expressing tetravalent IL-2 fusions with greater T_{reg} selectivity than current state-of-the-art monovalent or bivalent affinity muteins. Finally, we uncovered that IL2Rα itself is the

380 optimal target for designing T_{reg}-selective binding, and that cis-targeting can be designed

into multivalent IL-2 fusions through asymmetric tetravalent IL-2 fusions, again improving on signaling selectivity (Fig. 6). In total, our results show that not only do multivalency and cis targeting of IL2Rα improve T_{reg} selectivity, but that these paired strategies represent the only known method for overcoming the selectivity-potency tradeoff faced by T_{reg}-selective muteins[18].

Our results have clear implications for the design of T_{reg}-directed IL-2 therapies, an area of enormous interest for the treatment and management of autoimmune diseases[22,38]. We showed computationally and experimentally that multivalency and bitargeting can enhance IL-2 T_{reg} selectivity for potential use in clinical settings, where IL-2 based therapies have traditionally struggled[39]. Engineering valency requires precise compensatory adjustments in the ligand affinity (Fig. 5); given that we limited our experiments to pre-existing muteins, we expect that our selectivity gains might be improved even further by identifying muteins with optimal affinities. Our multivalent and bitargeted designs will additionally need to be tested *in vivo* to see that these selectivity gains translate to the *in vivo* setting[23]. T_{reg} selectivity is central to the mechanism of action for these therapies, and so we expect that these benefits to selectivity will improve therapeutic properties in several ways: more potent activation of signaling in T_{reg}s without off-target effects may improve the potency of these therapies and the breadth of applications[18,40,41]; reduced toxicity may allow for more routine use with minimal patient monitoring[14]. The superior selectivity offered by engineered multivalent ligands will likely further increase their *in vivo* pharmacokinetic lifetimes, in turn requiring less frequent dosing, as most drug clearance occurs via receptor-mediated endocytosis in off-target populations[21,42–44]. However, known differences exist in IL-2 receptor expression between humans and mice, so we do not expect that results from murine models would be a reliable indication for the comparative advantages of these molecules[15,16], and primate studies are outside the scope of the present work.

Heterospecificity, in our case exploited through bitargeting, opens a whole range of new possibilities through its ability to decouple the targeting and signaling properties of cytokine therapy and/or combine synergistic signals. This capability has been demonstrated through bispecific antibodies previously, and through the design of cis-targeted cytokine-antibody fusions. However, we showed that, unlike other immune cells, T_{reg}s do not express any surface marker more selective than IL2Rα (Fig. 2i-l). Consequently, there are no alternative targeting options to our approach of using the IL-2 receptors themselves (Fig. 6). Beyond our results, heterospecificity creates opportunities for synergistic receptor agonism. For example, PD-1 cis-targeting with IL-2 increases the stemness of CD8[+] T cells and consequently their tumor killing capacity[45,46]. While IL-2 has been employed as a therapy because of its T_{reg} selectivity, there is no reason to believe that the cytokine's signaling effects are optimal for enhancing T_{reg} suppressive activities. In fact, with cell therapies, where selectivity is not a concern and non-natural cytokine receptors can be introduced, other cytokine signaling such as IL-9 is qualitatively more effective than IL-2 at promoting cytotoxic T cell function[47,48]. Thus, one possibility enabled by bitargeting is potentially plug-and-play combinations of one or more cytokines that are more capable than IL-2 of driving desirable T_{reg} properties, made T_{reg}-selective through their fusion to multivalent IL2Rα-targeting complexes[49–51]. More systems-level research into the signaling regulation of T_{reg} proliferation and suppressive activities, and comparisons to other cytokines beyond IL-2, is needed to develop these possibilities.

Most generally, our results demonstrate the value of computationally directed biologics design, particularly for fusion constructs incorporating more than one binding moiety. The

design of these modular ligands leads to a combinatorial explosion of ligand configurations[52]. Furthermore, multivalent ligands have several documented effects, including altered signal transduction[53,54], binding avidity, pharmacokinetics[55], and intracellular trafficking[56]. While valency has been extensively applied as a means to introduce binding selectivity based on receptor density[57,58], multiple receptors, ligands subunits with varied targeting, and differences in signaling effects lead to additional complexity. For instance, while we found that the contribution of multivalency was explained primarily through avidity effects (Fig. 4), bitargeting as a strategy arises through differences in the signaling capacity of IL2Rα versus IL2Rβ. The approach here—computationally designed, multivalent, bitargeted ligands for enhanced therapeutic selectivity—has widespread application to other receptor-ligand pathways, including IL-4/IL-13, bone morphogenic proteins, and the TNF cytokines[52,59,60]. There are likely still other design strategies to be found across these many signaling pathway structures.

MATERIALS AND METHODS

EXPERIMENTAL METHODS

Receptor abundance quantitation, octet binding assays, expression of recombinant bivalent and monovalent IL-2 muteins (Fig. 1–4), and measurement of those muteins' signaling in PBMCs were performed as described in Farhat *et al.*[15].

Receptor abundance quantitation

Receptor quantitation data was gathered as described previously in Farhat et al.[15]; the preprocessing of fluorescence measurements, population gating, and receptor abundance calculations were performed using these data. To quantify the number of antibodies bound to cells and to standard beads, the fluorescence intensity of isotype controls was subtracted from the signal from matched receptor stains and then calibrated using the two lowest quantitation standards. Cell gating was conducted as shown in Fig. S1. The geometric means of replicates were calculated to summarize the results.

pSTAT5 Measurement of IL-2 and -15 Signaling in PBMCs

Cryopreserved PBMCs (UCLA Virology Core) were thawed to room temperature and slowly diluted with 9 mL pre-warmed RPMI-1640 (Corning, 10040CV) supplemented with 10% FBS (VWR, 97068-091, lot#029K20) and Penicillin/Streptomycin (Gibco, 15140122). Media was removed, and cells were brought to 3×10^6 cells/mL, distributed at 300,000 cells per well in a 96-well V-bottom plate, and allowed to recover 2 hrs at 37℃ in an incubator at 5% CO_2. IL-2 (R&D Systems, 202-IL-010) or IL-15 (R&D Systems, 247-ILB-025) were diluted in RPMI-1640 in the absence of FBS. These dilutions were then added to the concentrations indicated. To quantify STAT5 phosphorylation, the media was taken away, and cells were fixed using 100 μL of 10% formalin (Fisher Scientific, SF100-4) for 15 mins at room temperature. Formalin was removed from the cells, and the PBMCs were placed on ice. They were then suspended in 50 μL of cold methanol (-30℃). PBMCs were then kept at -30℃ overnight. PBSA was used to wash the cells twice. The cells were then split into two identical plates and stained with fluorescent antibodies for 1 hr at room temperature in darkness using 50 μL of antibody panels 4 and 5 per well. Cells were suspended in 100 μL PBSA per well, and beads to 50 μL, and analyzed on an IntelliCyt iQue Screener PLUS with VBR configuration (Sartorius) using a sip time of 35 secs and beads 30 secs.

Compensation of measured fluorescent values was calculated as detailed above. Gating of cell populations was performed as shown in Fig. S1, and the median pSTAT5 level was calculated for each population in each well.

Recombinant proteins

The Expi293 expression system was used to express IL-2/Fc fusion proteins. Expression was conducted as prescribed by the manufacturer instructions (Thermo Scientific). Proteins were formulated as the Fc of human IgG1 fused at its N- or C-terminus to human IL-2 using a $(G_4S)_4$ linker. C-terminal lysine residues of human IgG1 were not included in C-terminal fusions. The AviTag sequence GLNDIFEAQKIEWHE was added to the Fc terminus which did not contain IL-2. Fc mutations which prevented dimerization were introduced into the Fc sequence for monovalent muteins[61]. MabSelect resin (GE Healthcare) was used to purify protein. Biotinylation of proteins was conducted using BirA enzyme (BPS Biosciences) according to manufacturer instructions. Extensive buffer-exchanging into phosphate buffered saline (PBS) was conducted using Amicon 10 kDa spin concentrators (EMD Millipore). The sequence which was used to express the IL2Rβ/γ Fc heterodimer was the same as that of a reported, active heterodimeric molecule (patent application US20150218260A1); a (G4S)2 linker was added between the Fc portion and each receptor ectodomain. The Expi293 system was used to express the protein, which was subsequently purified on MabSelect resin as above. The IL2Rα ectodomain was generated to include a C-terminal 6xHis tag and then purified on Nickel-NTA spin columns (Qiagen) according to manufacturer instructions.

pSTAT5 Measurement of Tetravalent IL-2 Signaling in PBMCs

Cryopreserved PBMCs (UCLA Virology Core) were thawed to room temperature and slowly diluted with 9 mL pre-warmed RPMI-1640 (Corning, 10040CV) supplemented with 10% FBS (VWR, 97068-091, lot#029K20) and Penicillin/Streptomycin (Gibco, 15140122). Media was removed, and cells were brought to 3×10^6 cells/mL, distributed at 300,000 cells per well in a 96-well V-bottom plate, and allowed to recover 2 hrs at 37℃ in an incubator at 5% CO_2. IL-2 (Peprotech, 200-02-50µg) and tetravalent IL-2 (expressed and purified as described below) were diluted in RPMI-1640 without FBS and added to the indicated concentrations. To measure pSTAT5, media was removed, and cells fixed in 100 µL of 4% paraformaldehyde (PFA, Election Microscopy Sciences, 15714) diluted in PBS for 15 mins at room temperature.

PFA was removed, cells were gently suspended in 100 µL of cold methanol (-30℃). Cells were stored overnight at -30℃, and then washed twice with 0.1% bovine serum albumin (BSA, Sigma-Aldrich, B4287-25G) in PBS (PBSA), and stained 1 hr at room temperature in darkness using antibody panel X with 40 µL per well. Cells were then washed twice with 0.1% PBSA and resuspended in 150 µL PBSA per well. Cells were analyzed on a BD FACSCelesta flow cytometer. Populations were gated as shown in **Fig. S1**, and the median pSTAT5 level was extracted for each population in each well.

Tetravalent IL-2 Expression

Proteins were expressed as human IgG1 Fc-fused at the N- or C- terminus to mutant human IL-2 through a flexible (G4S)4 linker. C-terminal fusions omitted the C-terminal lysine residue of human IgG1. In bivalent variants, Fc mutations to prevent dimerization were introduced into the Fc sequence. Each IL-2 fused via the 20 amino acid long linker to the Fc domain contained R38Q and H16N mutations to reduce the IL-2's affinity with which

it binds IL2Rβ. Plasmid DNA prepared by maxi-prep (Qiagen, 12162) were transfected into adherent HEK293T cells using Lipofectamine 3000 (Thermo-Fisher, L3000008) in 15 cm dishes in DMEM (Corning, 15017CV) supplemented with GlutaMax (Gibco, 35050061) and 10% FBS. Media was exchanged after 24 hrs with fresh DMEM supplemented with GlutaMax and 5% ultra-low IgG FBS (Thermo-Fisher, A3381901). Media was harvested after an additional 72 hrs. Media was incubated in the presence of Protein A/G Plus Agarose resin (Santa Cruz Biotechnology, sc-2003) overnight. The following day, the media-resin mixture was centrifuged, and the supernatant discarded. Resin was washed with PBS five times or until protein was no longer detected in supernatant by UV-Vis using a NanoDrop One Spectrophotometer (Thermo-Fisher, ND-ONE-W). IL-2 was eluted from resin using 0.1M glycine, pH 2.3, into 2M Tris-HCl, pH 8. IL-2 was then buffer exchanged into PBS for storage at -80℃. Concentration was determined by BCA assay and confirmed using an IgG1 ELISA.

Octet binding assays

An Octet RED384 (ForteBio) was used to measure the binding affinity of each IL-2 mutein. Monomeric, biotintylated IL-2/Fc fusion proteins were loaded to Streptavidin biosensors (ForteBio) at roughly 10% of saturation point and allowed to equilibrate for 10 min in PBS + 0.1% bovine serum albumin (BSA). Up to 40 min of association time in IL2Rβ/γ titrated in 2x steps from 400 nM to 6.25 nM, or IL2Rα from 25 nM to 20 pM, which was followed by dissociation in PBS + 0.1% BSA. A zero-concentration was included in each measurement and served as a negative control/reference signal. The affinity quantification experiments were performed in quadruplicate across two days. Binding of IL-2 to IL2Rα on its own did not fit to a simple binding model; K_D was calculated using equilibrium binding within each assay for this case. IL2Rβ/γ binding data fit a 1:1 binding model; thus, in these cases on-rate (k_{on}), off-rate (k_{off}) and K_D were determined by fitting to the entire binding curve. The average of each kinetic parameter across all concentrations with detectable binding (typically 12.5 nM and above) was used to calculate K_D.

Statistical analysis

The number of replicates performed for each experimental measurement, and the values of confidence intervals are described in corresponding figure captions. N is used to describe the number of times a particular experiment was performed. Flow cytometry experiments performed using initial panel of monovalent and bivalent cytokines (Figs. 1–4) were performed on hPBMCs were conducted using separate experimental replicates on cells gathered from a single donor. Each replicate of the flow cytometry signaling experiments in figures 5 and 6 were conducted using hPBMCs from different donors. To quantify population-level flow cytometry measurements for both signaling and receptor quantitation experiments, the mean fluorescent intensity (MFI) of a gated population was measured. Compensation to remove fluorescent spectral overlap was performed for each experimental measurement. Subtraction of either negative controls or cells treated with isotype antibodies was performed on signaling and receptor quantitation data respectively to remove background signal. Cells which were measured to display fluorescent intensities above 1,000,000 were excluded from analysis during signaling experiments. Pearson correlation coefficients (R^2) values were used to describe model accuracy when predicting signaling response to IL-2 and IL-2 muteins. The K_x^* parameter was fit with least-squares fitting using the Broyden–Fletcher–Goldfarb–Shanno minimization algorithm as implemented in SciPy.

MODELING

Binding model

565 The model was formulated as described in Tan *et al*[32]. The monomer composition of a ligand complex was represented by a vector $\boldsymbol{\theta} = (\theta_1, \theta_2, \ldots, \theta_{N_L})$, where each θ_i was the fraction of monomer ligand type i out of all monomers on that complex. Let $C_{\boldsymbol{\theta}}$ be the proportion of the $\boldsymbol{\theta}$ complexes in all ligand complexes, and Θ be the set of all possible $\boldsymbol{\theta}$'s. We have $\sum_{\boldsymbol{\theta} \in \Theta} C_{\boldsymbol{\theta}} = 1$.

570 The binding between a ligand complex and a cell expressing several types of receptors can be represented by a series of q_{ij}. The relationship between q_{ij}'s and θ_i is given by $\theta_i = q_{i0} + q_{i1} + \ldots + q_{iN_R}$. Let the vector $\mathbf{q}_i = (q_{i0}, q_{i1}, \ldots, q_{iN_R})$, and the corresponding $\boldsymbol{\theta}$ of a binding configuration $\mathbf{q}$ be $\boldsymbol{\theta}(\mathbf{q})$. For all i in $\{1, 2, \ldots, N_L\}$, we define $\psi_{ij} = R_{\text{eq},j} K_{a,ij} K_x^*$ where $j = \{1, 2, \ldots, N_R\}$ and $\psi_{i0} = 1$. The relative number of complexes bound to a cell with 575 configuration $\mathbf{q}$ at equilibrium is

$$v_{q,eq} = \frac{L_0 C_{\boldsymbol{\theta}(\boldsymbol{q})}}{K_x^*} \prod_{\substack{i=1 \\ j=0}}^{\substack{i=N_L \\ j=N_R}} \psi_{ij}^{q_{ij}} \prod_{i=1}^{N_L} \binom{\theta_i}{\boldsymbol{q}_i}.$$

Then we can calculate the relative amount of bound receptor n as

$$R_{\text{bound},n} = \frac{L_0}{K_x^*} \sum_{\boldsymbol{\theta} \in \Theta} C_{\boldsymbol{\theta}} \left[\sum_{i=1}^{N_L} \frac{\psi_{in} \theta_i}{\sum_{j=0}^{N_R} \psi_{ij}} \right] \prod_{i=1}^{N_L} \left(\sum_{j=0}^{N_R} \psi_{ij} \right)^{\theta_i}.$$

By mass conservation, ($R_{\text{tot},n} = R_{\text{eq},n} + R_{\text{bound},n}$), we can solve $R_{\text{eq},n}$ numerically for each 580 type of receptor.

Application of multivalent binding model to IL-2 signaling pathway

Each IL-2 molecule was allowed to bind to one free IL2Rα and one IL2Rβ/γc receptor. Initial IL-2-receptor association proceeds with the known kinetics of monomeric ligand-receptor interaction (Table S1). Subsequent ligand-receptor binding interactions then proceed with 585 an association constant proportional to available receptor abundance and affinity multiplied by the scaling constant, K_x^*, as described above. To predict pSTAT5 response to IL-2 stimulation, we assumed that pSTAT5 is proportional to the amount of IL-2-bound IL2Rβ/γc, as complexes which contain these species actively signal through the JAK/STAT pathway. Scaling factors converting from predicted active signaling species to pSTAT5 abundance 590 were fit to experimental data on a per-experiment and cell type basis. A single K_x^* value was fit for all experiments and cell types.

CITE-seq marker selectivity analysis

To assist in identifying possible methods through which to increase IL-2 selectivity towards T_regs, a publicly available Cellular Indexing of Transcriptomes and Epitopes by sequencing 595 (CITE-seq) dataset containing data gathered from human PBMCs was analyzed[28]. Only RNA transcripts encoding cell membrane extracellular-facing proteins were included. We first analyzed the data by determining the Wasserstein distance and Kullback-Leibler

divergence of markers and RNA measured in T_{reg}s against the distribution of these markers displayed by all other cells. We also analyzed the data using a ridge classification model, where all markers and RNA sequences were used by the model to distinguish between T_{reg}s and all other cell types.

Markers of interest were then used in conjunction with the binding model to determine whether they could confer selectivity, using the CITE-seq data to inform the number of markers per cell. Conversion factors for calculating marker abundance from CITE-seq marker and mRNA reads were estimated using proportional conversions from the data to previously experimentally determined marker abundances[15]. Single cell marker abundances were calculated for 1000 cells at a time, and the ratio of T_{reg} signaling to off-target signaling was calculated. To simulate bispecific binding, two distinct binding domains for each ligand were modeled, one for IL2, with affinity for IL2Ra and IL2RB, and the other for the marker of interest. An optimization function was used to calculate the binding affinity which would yield the best selectivity per marker. After finding IL2Rα to be the optimal epitope for increasing selectivity, we sought to explore the effects of increasing valency by doubling the number of binding domains per ligand.

Tensor Factorization

Before decomposition, the signaling response data was background subtracted and variance scaled across each cell population. Non-negative canonical polyadic decomposition was performed using the Python package TensorLy, using the HALS algorithm with non-negative SVD initialization[62].

References

1. Leonard, W. J., Lin, J.-X. & O'Shea, J. J. The γc Family of Cytokines: Basic Biology to Therapeutic Ramifications. *Immunity* **50**, 832–850 (2019).
2. Janas, M. L., Groves, P., Kienzle, N. & Kelso, A. IL-2 Regulates Perforin and Granzyme Gene Expression in CD8+ T Cells Independently of Its Effects on Survival and Proliferation. *The Journal of Immunology* **175**, 8003–8010 (2005).
3. Chinen, T. *et al.* An essential role for the IL-2 receptor in Treg cell function. *Nat Immunol* **17**, 1322–1333 (2016).
4. Fan, M. Y. *et al.* Differential Roles of IL-2 Signaling in Developing versus Mature Tregs. *Cell Rep* **25**, 1204-1213.e4 (2018).
5. Noguchi, M. *et al.* Interleukin-2 receptor gamma chain mutation results in X-linked severe combined immunodeficiency in humans. *Cell* **73**, 147–157 (1993).
6. Zenatti, P. P. *et al.* Oncogenic IL7R gain-of-function mutations in childhood T-cell acute lymphoblastic leukemia. *Nat Genet* **43**, 932–939 (2011).
7. Dwyer, C. J. *et al.* Fueling Cancer Immunotherapy With Common Gamma Chain Cytokines. *Frontiers in Immunology* **10**, (2019).
8. Abbas, A. K., Trotta, E., Simeonov, D. R., Marson, A. & Bluestone, J. A. Revisiting IL-2: Biology and therapeutic prospects. *Science Immunology* **3**, (2018).
9. Choudhry, H. *et al.* Prospects of IL-2 in Cancer Immunotherapy. *BioMed Research International* **2018**, e9056173 (2018).

10. Tahvildari, M. & Dana, R. Low-Dose IL-2 Therapy in Transplantation, Autoimmunity, and Inflammatory Diseases. *The Journal of Immunology* **203**, 2749–2755 (2019).

11. Spolski, R., Li, P. & Leonard, W. J. Biology and regulation of IL-2: from molecular mechanisms to human therapy. *Nat Rev Immunol* **18**, 648–659 (2018).

12. Koreth, J. *et al.* Interleukin-2 and regulatory T cells in graft-versus-host disease. *N Engl J Med* **365**, 2055–2066 (2011).

13. Saadoun, D. *et al.* Regulatory T-cell responses to low-dose interleukin-2 in HCV-induced vasculitis. *N Engl J Med* **365**, 2067–2077 (2011).

14. Krieg, C., Létourneau, S., Pantaleo, G. & Boyman, O. Improved IL-2 immunotherapy by selective stimulation of IL-2 receptors on lymphocytes and endothelial cells. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 11906–11911 (2010).

15. Farhat, A. M. *et al.* Modeling cell-specific dynamics and regulation of the common gamma chain cytokines. *Cell Rep* **35**, 109044 (2021).

16. Gonnord, P. *et al.* A hierarchy of affinities between cytokine receptors and the common gamma chain leads to pathway cross-talk. *Sci Signal* **11**, (2018).

17. Mitra, S. & Leonard, W. J. Biology of IL-2 and its therapeutic modulation: Mechanisms and strategies. *J. Leukoc. Biol.* **103**, 643–655 (2018).

18. VanDyke, D. *et al.* Engineered human cytokine/antibody fusion proteins expand regulatory T cells and confer autoimmune disease protection. *Cell Reports* **41**, (2022).

19. Silva, D.-A. *et al.* De novo design of potent and selective mimics of IL-2 and IL-15. *Nature* **565**, 186–191 (2019).

20. Khoryati, L. *et al.* An IL-2 mutein engineered to promote expansion of regulatory T cells arrests ongoing autoimmunity in mice. *Sci Immunol* **5**, eaba5264 (2020).

21. Bell, C. J. M. *et al.* Sustained in vivo signaling by long-lived IL-2 induces prolonged increases of regulatory T cells. *J Autoimmun* **56**, 66–80 (2015).

22. Peterson, L. B. *et al.* A long-lived IL-2 mutein that selectively activates and expands regulatory T cells as a therapy for autoimmune disease. *J. Autoimmun.* **95**, 1–14 (2018).

23. Ghelani, A. *et al.* Defining the Threshold IL-2 Signal Required for Induction of Selective Treg Cell Responses Using Engineered IL-2 Muteins. *Front Immunol* **11**, 1106 (2020).

24. Moynihan, K. *et al.* 717 Selective activation of CD8+ T cells by a CD8-targeted IL-2 results in enhanced anti-tumor efficacy and safety. *J Immunother Cancer* **9**, (2021).

25. Feinerman, O. *et al.* Single-cell quantification of IL-2 response by effector and regulatory T cells reveals critical plasticity in immune response. *Mol. Syst. Biol.* **6**, 437 (2010).

26. Lowenthal, J. W., Robson MacDonald, H. & Iacopetta, B. J. Intracellular pathway of interleukin 2 following receptor-mediated endocytosis. *European Journal of Immunology* **16**, 1461–1463 (1986).

27. Klatzmann, D. & Abbas, A. K. The promise of low-dose interleukin-2 therapy for autoimmune and inflammatory diseases. *Nature Reviews Immunology* **15**, 283–294 (2015).

28. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29 (2021).

29. Tan, Z. C., Orcutt-Jahns, B. T. & Meyer, A. S. A quantitative view of strategies to engineer cell-selective ligand binding. *Integrative Biology* **13**, 269–282 (2021).

30. Tan, Z. C., Murphy, M. C., Alpay, H. S., Taylor, S. D. & Meyer, A. S. Tensor-structured decomposition improves systems serology analysis. *Mol Syst Biol* **17**, e10243 (2021).

31. Kolda, T. G. & Bader, B. W. Tensor Decompositions and Applications. *SIAM Rev.* **51**, 455–500 (2009).

32. Tan, Z. C. & Meyer, A. S. A general model of multivalent binding with ligands of heterotypic subunits and multiple surface receptors. *Math Biosci* 108714 (2021) doi:10.1016/j.mbs.2021.108714.

33. Robinett, R. A. *et al.* Dissecting FcγR Regulation through a Multivalent Binding Model. *Cell Syst* **7**, 41-48.e5 (2018).

34. Perelson, A. S. & DeLisi, C. Receptor clustering on a cell surface. I. theory of receptor cross-linking by ligands bearing two chemically identical functional groups. *Mathematical Biosciences* **48**, 71–110 (1980).

35. Stone, J. D., Cochran, J. R. & Stern, L. J. T-cell activation by soluble MHC oligomers can be described by a two-parameter binding model. *Biophys J* **81**, 2547–2557 (2001).

36. Voss, S. D. *et al.* Increased expression of the interleukin 2 (IL-2) receptor beta chain (p70) on CD56+ natural killer cells after in vivo IL-2 therapy: p70 expression does not alone predict the level of intermediate affinity IL-2 binding. *J Exp Med* **172**, 1101–1114 (1990).

37. Mazor, Y. *et al.* Enhanced tumor-targeting selectivity by modulating bispecific antibody binding affinity and format valence. *Sci Rep* **7**, 40098 (2017).

38. Ye, C., Brand, D. & Zheng, S. G. Targeting IL-2: an unexpected effect in treating immunological diseases. *Signal Transduct Target Ther* **3**, 2 (2018).

39. Siegel, J. P. & Puri, R. K. Interleukin-2 toxicity. *J Clin Oncol* **9**, 694–704 (1991).

40. Hirai, T. *et al.* Selective expansion of regulatory T cells using an orthogonal IL-2/IL-2 receptor system facilitates transplantation tolerance. *J Clin Invest* **131**, (2021).

41. Buitrago-Molina, L. E. *et al.* Treg-specific IL-2 therapy can reestablish intrahepatic immune regulation in autoimmune hepatitis. *Journal of Autoimmunity* **117**, 102591 (2021).

42. Carr, A. *et al.* Outpatient continuous intravenous interleukin-2 or subcutaneous, polyethylene glycol-modified interleukin-2 in human immunodeficiency virus-infected patients: a randomized, controlled, multicenter study. Australian IL-2 Study Group. *J Infect Dis* **178**, 992–999 (1998).

43. de Picciotto, S. *et al.* Selective activation and expansion of regulatory T cells using lipid encapsulated mRNA encoding a long-acting IL-2 mutein. *Nat Commun* **13**, 3866 (2022).

44. Bernett, M. J. *et al.* An IL15/IL15Ra heterodimeric Fc-fusion engineered for reduced potency demonstrates an optimal balance of in vivo activity and exposure. (2018).

45. Hashimoto, M. *et al.* PD-1 combination therapy with IL-2 modifies CD8+ T cell exhaustion program. *Nature* **610**, 173–181 (2022).

46. Codarri Deak, L. *et al.* PD-1-cis IL-2R agonism yields better effectors from stem-like CD8+ T cells. *Nature* **610**, 161–172 (2022).

725 47. Kalbasi, A. *et al.* Potentiating adoptive cell therapy using synthetic IL-9 receptors. *Nature* **607**, 360–365 (2022).

48. Mo, F. *et al.* An engineered IL-2 partial agonist promotes CD8+ T cell stemness. *Nature* **597**, 544–548 (2021).

49. Elyaman, W. *et al.* IL-9 induces differentiation of TH17 cells and enhances function of
730   FoxP3+ natural regulatory T cells. *Proceedings of the National Academy of Sciences* **106**, 12885–12890 (2009).

50. Zaiss, D. M. W. *et al.* Amphiregulin enhances regulatory T cell suppressive function via the epidermal growth factor receptor. *Immunity* **38**, 275–284 (2013).

51. Marie, J. C., Letterio, J. J., Gavin, M. & Rudensky, A. Y. TGF-β1 maintains suppressor
735   function and Foxp3 expression in CD4+CD25+ regulatory T cells. *Journal of Experimental Medicine* **201**, 1061–1067 (2005).

52. Antebi, Y. E. *et al.* Combinatorial Signal Perception in the BMP Pathway. *Cell* **170**, 1184-1196.e24 (2017).

53. Kiessling, L. L., Gestwicki, J. E. & Strong, L. E. Synthetic Multivalent Ligands as Probes
740   of Signal Transduction. *Angewandte Chemie International Edition* **45**, 2348–2368 (2006).

54. Moraga, I. *et al.* Synthekines are surrogate cytokine and growth factor agonists that compel signaling through non-natural receptor dimers. *Elife* **6**, e22882 (2017).

55. Cuesta, A. M., Sainz-Pastor, N., Bonet, J., Oliva, B. & Alvarez-Vallina, L. Multivalent
745   antibodies: when design surpasses evolution. *Trends Biotechnol* **28**, 355–362 (2010).

56. York, S. J., Arneson, L. S., Gregory, W. T., Dahms, N. M. & Kornfeld, S. The rate of internalization of the mannose 6-phosphate/insulin-like growth factor II receptor is enhanced by multivalent ligand binding. *J Biol Chem* **274**, 1164–1171 (1999).

57. Csizmar, C. M. *et al.* Multivalent Ligand Binding to Cell Membrane Antigens: Defining
750   the Interplay of Affinity, Valency, and Expression Density. *J Am Chem Soc* **141**, 251–261 (2019).

58. Cb, C., P, M., Rm, O., Ec, D. & Ll, K. Selective tumor cell targeting using low-affinity, multivalent interactions. *ACS chemical biology* **2**, (2007).

59. Kelly-Welch, A. E., Hanson, E. M., Boothby, M. R. & Keegan, A. D. Interleukin-4 and
755   interleukin-13 signaling connections maps. *Science* **300**, 1527–1528 (2003).

60. Efimov, G. A. *et al.* Cell-type-restricted anti-cytokine therapy: TNF inhibition from one pathogenic source. *Proc Natl Acad Sci U S A* **113**, 3006–3011 (2016).

61. Ishino, T. *et al.* Engineering a monomeric Fc domain modality by N-glycosylation for the half-life extension of biotherapeutics. *J Biol Chem* **288**, 16529–16537 (2013).

760 62. Kossaifi, J., Panagakis, Y., Anandkumar, A. & Pantic, M. TensorLy: Tensor Learning in Python. *arXiv:1610.09555 [cs]* (2018).

Acknowledgements: The authors thank Cori Posner for contributing to initial profiling studies that enabled the study.

Funding: This work was partly supported by startup funds from UCLA Engineering and by NIH U01-AI148119 to A.S.M.

Author contributions statement: A.S.M. conceived of the study. S.D.T. performed the PBMC experiments with the IL-2 fusion proteins. A.S.M, B.O.J., E.M.S., and P.C.E. performed the computational analysis. All authors helped to design experiments and/or analyze the data. All authors contributed to writing the paper.

Competing interests: A.S.M. has filed a patent PCT/US22/35711 on the use of multivalent cytokines to enhance cell type-selective responses. A.S.M., B.O.J., and P.C.E. have filed a provisional patent on the use of bitargeting for engineered cytokine responses.

Data Sharing Plan: All analysis was implemented in Python v3.10 and can be found at

https://github.com/meyer-lab/gc-valent, release 1.0, along with all the experimental data.

FIGURES

Fig. 1. Systematically profiling IL-2 muteins reveals determinants of response.
(a) Schematic of affinity and structural mutants explored. **(b)** IL2Rα and IL2Rβ affinities of each IL-2 variant. **(c)** Heatmap of phosphorylated STAT5 measurements for each cell type, time point, ligand, and concentration. pSTAT5 measurements are normalized for each cell type. **(d–o)** STAT5 phosphorylation response curves for immune cells stimulated with select IL-2 muteins. Time points and cell types are indicated in subplot titles.

Fig. 2. IL-2 muteins display structural- and affinity-dependent T_reg selectivity that cannot be overcome with cis-targeting strategies. (a) Schematic describing ratio of activation between target and off-target immune populations. **(b, d, f)** Ratios of T_reg pSTAT5 to NK (b), CD8$^+$ (d), and T_helper (f) pSTAT5 dose response curves at 4 hours. Ratio was defined as the ratio of Hill curves fit to experimental data for target and off-target populations, as in Fig. 1. **(c, e, g)** Maximum ratio of T_reg to off-target population signaling vs. IL2Rα affinity. Lines of best fit were fit to monovalent (solid) and bivalent (dashed). **(h)** Schematic depicting how useful markers for conferring selectivity are selected. **(I–o)** Most unique markers for T_regs in CITE-seq dataset identified using Wasserstein distance (i,k), and Kullback-Leibler Divergence metrics (j,l) using surface markers (i,j) and RNA data (k,l).

800 **Fig. 3. Tensor-based decomposition reveals unique selectivity defined by fusion valency. (a)** Scores and loadings plot of pSTAT5 signaling data as calculated using principal components analysis. **(b)** Schematic representation of non-negative canonical polyadic (CP) decomposition. Experimental pSTAT5 measurements are arranged in a tensor according to the duration of treatment, ligand used, cytokine concentration, and cell 805 type. CP decomposition then helps to visualize this space. **(c)** Percent variance reconstructed (R2X) versus the number of components used. **(d)** Component values for each IL-2 mutant. **(e)** Component values representing the effect of IL-2 concentration. **(f)** Component values representing cell type specificity. **(g)** Component values for the effect of treatment duration. **(h)** Sum of Component 1 and 3 weights (off-target cellular signaling) vs. 810 Component 2 (T_{reg} signaling) weight for each monovalent and bivalent ligand. Thin line is included for visualization purposes only.

Fig. 4. Responses are predicted by a simple multivalent binding model. (a) A
simplified cartoon of the model. Initial association of multivalent ligands proceeds according
to monovalent affinity, and subsequent binding events proceed with that affinity scaled by
the K_x^* parameter. **(b, c)** Model's accuracy subset by cell type **(b)** and ligand **(c)** for all
mono- and bivalent IL-2 muteins. **(d, e)** Model's accuracy subset by concentration **(d)** for all
ligands and time **(e)** for all ligands, concentrations, and cell types. All accuracies are
calculated as a Pearson's correlation R^2 score for experimental cytokine responses at 30
mins and one hr. **(f, g)** Model-predicted pSTAT for T_{reg}s (f) and NK cells (g) in response to
mono- and bivalent IL-2 ligands with 10 nM IL2Rβ K_D. **(h, i)** Predicted number of active
signaling complexes formed on cells with 1000 IL2Rβ receptors and varying numbers of
IL2Rα for ligands with affinities of 10 nM K_D for IL2Rβ and either 1 nM (f) or 10 nM (i) K_D for
IL2Rα.

Fig. 5. Multivalency enhances the selectivity of cytokine fusion proteins. (a,c,e)
Signaling response of T_{reg}, NK cells, and T_{helper} cells predicted for ligand of optimal
selectivity at different valencies. Response predictions were normalized to each
population's response for the monovalent case. Selectivity for T_{reg}s and NK cells were

830   derived from IL-2 muteins, and selectivity for T_{helper}s was calculated using IL-7 muteins.
(b,d,f) Optimal receptor-ligand dissociation constants for each ligand optimized for
selectivity. Mutein affinity for IL2Rα and IL2Rβ/γc was allowed to vary for IL-2 muteins, and
affinity for IL-7Rα was allowed to vary for IL-7 muteins. Affinities were allowed to vary
across K_Ds of 10 pM–1 μM while K_x^* was fixed at its fitting optimum. All optimizations were

835   performed using a concentration of 1 nM. Selectivity was calculated as the ratio of
predicted pSTAT5 in target cells to the mean pSTAT5 predicted in off-target cells. **(g)**
Schematic of multivalent IL-2 mutant design. **(h–j)** Ratio of STAT5 phosphorylation in T_{reg}s
to T_{helper}s (h), NK cells (i), and CD8+ (j) cells at varying dosages for R38Q/H16N in various
valency formats. Dots are representative of mean of experimental replicates (N=3).

Fig. 6. Asymmetric IL-2 mutants display even greater T$_{reg}$ selectivity. (a,b) Predicted enhancements to T$_{reg}$ selectivities for IL-2 mutants including a binding domain as calculated using CITE-seq surface marker data (a) or surface expressed RNA transcripts (b). Selectivity was calculated for T$_{reg}$s against all other PBMC cells as the increase in average

845   T$_{reg}$ to off-target cell binding against WT IL-2 at a concentration of 0.1 nM. **(c)** Schematic of asymmetric IL-2 mutant design. **(d)** Predicted normalized T$_{reg}$ selectivity displayed by bivalent WT IL-2, R38Q/H16N, and Bitargeted IL-2 across IL2Rβ affinities. **(e–g)** Ratio of STAT5 phosphorylation in T$_{reg}$s to T$_{helper}$s (e), NK cells (f), and CD8$^+$ (g) cells at varying dosages for R38Q/H16N in various valency formats. Dots are representative of mean of

850   experimental replicates (N=3). **(h–j)** Tensor factorization of signaling responses to R38Q/H16N and Bitargeted variants. **(h)** Percent variance reconstructed (R2X) versus the number of components used. **(i)** Component values representing cell type specificity. **(j)** Component values for each IL-2 mutant. **(k)** Component values representing the effect of concentration.