Convergent evolution of venom gland transcriptomes across Metazoa

Giulia Zancolli[1*], Maarten Reijnders[1], Robert M. Waterhouse[1], Marc Robinson-Rechavi[1]

[1]Department of Ecology and Evolution, University of Lausanne and Swiss Institute of Bioinformatics, 1015, Lausanne, Switzerland

*Corresponding author: Giulia Zancolli

Email: giulia.zancolli@gmail.com

Author Contributions: GZ and MRR conceived the study design. GZ collected all data and performed all analyses except where specified, with input from MRR and RMW; MR performed GO annotations and GO term semantic similarity analysis; RMW performed species phylogeny. GZ wrote the first draft of the manuscript, and GZ and MRR wrote the final version.

Keywords: convergent evolution, evolutionary novelties, gene expression, RNA-seq, stress response, venom.

Abstract

Animals have repeatedly evolved specialized organs and anatomical structures to produce and deliver a cocktail of potent bioactive molecules to subdue prey or predators – venom. This makes it one of the most widespread convergent functions in the animal kingdom. Whether animals have adopted the same genetic toolkit to evolved venom systems is a fascinating question that still eludes us. Here, we performed the first comparative analysis of venom gland transcriptomes from 20 venomous species spanning the main Metazoan lineages, to test whether different animals have independently adopted similar molecular mechanisms to perform the same function. We found a strong convergence in gene expression profiles, with venom glands being more similar to each other than to any other tissue from the same species, and their differences closely mirroring the species phylogeny. Although venom glands secrete some of the fastest evolving molecules (toxins), their gene expression does not evolve faster than evolutionarily older tissues. We found 15 venom gland specific gene modules enriched in endoplasmic reticulum stress and unfolded protein response pathways, indicating that animals have independently adopted stress response mechanisms to cope with mass production of toxins. This, in turn, activates regulatory networks for epithelial development, cell turnover and maintenance which seem composed of both convergent and lineage-specific factors, possibly reflecting the different developmental origins of venom glands. This study represents the first step towards an understanding of the molecular mechanisms underlying the repeated evolution of one of the most successful adaptive traits in the animal kingdom.

Main Text

Introduction
Organisms often evolve predictably similar features when presented separately with the same
environmental or biological challenge (1). A long-standing question is whether the repeated evolution
of adaptive traits in distinct lineages involves similar molecular changes, such as protein-coding
sequences, cis-regulatory DNA elements, or gene expression (1–6). Animal venom represents one of
the most remarkable examples of convergent evolution. On more than 100 occasions animal lineages
have independently evolved the ability to secrete potent molecules to subdue prey or predators. Despite
having the same biological role, the origin, anatomy and organization of the venom apparatus differ
dramatically among lineages (7). Venom systems therefore represent an exceptional opportunity to test
whether different animal lineages have repeatedly adopted similar molecular mechanisms to perform
the same function (8).
Recent advances in sequencing technologies have allowed the molecular characterization of hundreds
of proteomes and transcriptomes from the venom glands of several taxa, with a focus on medically
important ones such as snakes and spiders, but also more neglected lineages (9–11). In particular,
most venom gland transcriptome studies are focused on the identification of toxin transcript sequences,
although recently there has been an increased interest for the "non-toxin" transcriptome (12, 13). The
availability of genomes and of venom-gland RNA-seq datasets from various venomous lineages provide
the opportunity for a comparative analysis across the animal tree of life to answer a simple, yet
unexplored questions: did animals independently employ the same genetic toolkit to achieve the same
function?
Here we address this question by comparing, for the first time, gene expression profiles from 20
venomous species representing eight independent origins of venom, and spanning ~ 700 million years
of evolution (protostome / deuterostome divergence) (14). We tested whether convergence in the ability
to produce venom corresponds to the convergent evolution of gene expression levels in the venom
glands. As venomous animals have evolved specialized organs for the biosynthesis and secretion of
toxins, we expect enrichment of similar biological processes even among distantly-related taxa.
However, animal venom glands are non-homologous structures with diverse origins (8); therefore, we
hypothesize that convergence in biological function does not imply similarity in global gene expression
profiles and regulatory networks. To study this, we first used a set of conserved ortholog genes to
assess global evolutionary patterns of expression across all taxa. Then, we examined the whole
transcriptome in each species separately to determine lineage-specific or shared expression changes,
pathways and regulatory networks. We found a striking similarity of global gene expression patterns in
evolutionary distinct venom glands, especially in genes involved in secretory functions, which indicates
that complex trait evolution may sometimes be more constrained and predictable than expected. On
the other hand, lineage specific profiles suggest that the way in which cells are regulated and
communicates might reflect the diverse developmental origins of venom systems.
73
74
75 **Results and Discussion**
76
Does convergence in function correspond to convergence in gene expression profiles?
To analyze to what extent gene expression profiles of non-homologous venom glands are convergent,
we compared publicly available RNA-seq datasets of venom glands and other body tissues from 20
venomous species representing eight different origins of venom: five spiders, two scorpions, one bee,
three wasps, one fly, two mollusks, four snakes, one fish, and one mammal (Table 1 and *SI Appendix*,
Table S1). In total, we used 2,528 orthogroups, sets of ortholog genes across all species, to create an
expression matrix of log-transformed, quantile normalized Transcripts Per Million (TPM) values (*SI*
Appendix, Dataset S1).
First, we explored gene expression patterns between lineages and tissues with principal component
analysis (PCA), and found that the first three components clearly separated the venom gland samples
from the other body tissues (Fig. 1 and *SI Appendix*, Fig S1). We had hypothesized that the expression
levels of toxin genes, which are generally either restricted to or highly enriched in the venom gland,
might be the drivers of this strong pattern. However, after removing the 38 orthogroups corresponding
to known toxin genes from the expression matrix, the PCA did not substantially change (*SI Appendix*,
Fig. S2). The clustering of venom gland samples in the PCA is remarkable considering the diverse
origins of the datasets, not only because they are from different species and experiments, therefore
different RNA processing and sequencing techniques, but also in terms of sampling procedures (e.g.

time of dissection, pooling of samples, sex, age). Thus, it most likely represents a lower estimate of the
degree of molecular convergence of the venom glands.
Venom glands are mostly composed of epithelial secretory cells; therefore, we can expect their gene
expression profiles to be similar to that of other exocrine tissues. Indeed, non-venom glandular tissues
were positioned between the venom glands and the other tissues in the principal component space
(Fig. 1 and *SI Appendix*, S1), suggesting shared expression patterns between secretory tissues. The
rattlesnake's accessory venom gland and the Indian cobra's salivary gland clustered with venom
glands: the accessory venom gland, as the name suggests, contributes to the production of venom and
therefore it is expected to have a similar transcriptome (15). The pattern of clustering of the Indian
Cobra's salivary gland is more complex; one possible explanation is that the gland may have been mis-
identified or there might have been contamination during dissection. The variation and diversity of
secreting dental glands in snakes makes it difficult to clearly distinguish them, and confusion in the
terminology of these homologous structures has been previously noted (16). Nonetheless, snake
venom gland transcriptomes seem to rely on a conserved secretory gene regulatory network shared
with salivary tissues of other amniotes (13).
Considering the diverse, and often recent, evolutionary independent origins of venom glands, we
hypothesized that they would have higher transcriptome similarity to other tissues of the same species
rather than between venom glands of different species. Contrary to this expectation, venom glands
were more similar to each other than to any other tissue even from the same species (Spearman rank
correlation coefficient ρ, Fig. 2). This is consistent with their clustering in the PCA (Fig. 1), although the
comparisons were not statistically significant after Benjamini-Hochberg correction due to the low sample
size of the non-venom tissues. However, we observed some particularly low correlation values between
venom glands of the echidna and of other species. The echidna has a peculiar venom system compared
to other animals – the venom gland is only active during the breeding season, it is found only in males,
and it is thought to play a role in scent communication and to aid in competition. Furthermore, the loss
of ability to erect the spur for venom delivery is thought to be the result of gradual decay of venom
function (17). For these reasons, it might be that the echidna's venom gland is diverging from the shared
function of high-level secretory machinery in other animals, thus the observed low similarity values and
the separation from the other tissues in the PC plot (Fig. 1).
123
Do closely related lineages have more similar transcriptomes?
We hypothesized that the expression levels of genes from homologous venom glands, i.e. from species
that share a common venomous ancestor (e.g. snakes), follow a phylogenetic pattern, as regulatory
changes accumulate over time. On the contrary, when comparing non-homologous transcriptomes, we
might observe different, unpredictable patterns, with distantly related lineages clustering together due
to functional convergence. To test this hypothesis, we compared a venom gland expression tree with
the species phylogenetic tree. The expression tree (Neighbor-Joining) was constructed using two
distance metrics, 1-Spearman coefficient and Euclidean distances, and the species tree was based on
multisequence alignments of 1:1 orthogroups using RAxML (18). Surprisingly, the expression tree was
overall consistent with the phylogenetic tree (Fig. 3A). The expression tree correctly separated
protostomes and deuterostomes and the eight independent origins of venom. The branching patterns
within these clades also broadly reflected the known phylogeny, except for the tree based on Euclidean
distances which grouped Octopi with Ecdysozoa (*SI Appendix*, Fig. S3).
Then, we compared sequence-based phylogenetic distances with expression distances between all
species pairs to test whether closer lineages have more similar transcriptomes. Expression distances
were positively correlated with phylogenetic distances ($R = 0.19$), confirming that closer taxa have more
similar expression patterns (Fig. 3B). We noticed some particularly high expression distance values;
these outliers are all pairwise distances between the echidna and other animals. As mentioned above,
the echidna's transcriptome is particularly divergent from all the other animals. The correlation between
phylogenetic and expression distances was, as expected, much stronger excluding the echidna ($R = 0.44$, *SI Appendix*, Fig. S4).
0.44, *SI Appendix*, Fig. S4).
145
Do venom-gland transcriptomes evolve faster than those of other tissues?
Venom glands are derived traits that evolved from already differentiated tissues. Furthermore, the main
product of venom glands, toxins, are among the fastest evolving genes, and their genomic makeup is
highly variable and dynamic, with duplications and deletions between individuals of the same species
(19). For these reasons we hypothesized that venom gland transcriptomes diverge at faster rates than
tissues which are evolutionarily older. To test this hypothesis, we built gene expression trees based on
1- Spearman and Euclidean distances for other homologous tissues with at least seven species in our

dataset, i.e. ovary and brain, and compared them with the venom gland expression tree (*SI Appendix*,
Fig. S5).
Similarly to the venom gland tree, the brain tree separated deuterostomes and protostomes, and
broadly reflected the species tree. On the other hand, the ovary tree resolved the two clades only with
Euclidean distances, and even then, octopi clustered with the deuterostomes (*SI Appendix*, Fig. S6).
Then, we compared pairwise phylogenetic distances with expression distances, and tested whether
venom glands have higher evolutionary rates (i.e. higher slope values). Overall, at similar phylogenetic
distances, ovary and brain transcriptomes were more divergent than venom gland transcriptomes (*SI
Appendix*, Fig. S5); however, the venom gland slope was not steeper compared to that of the ovary (F
= 0.41, p = 0.52), nor the brain (F = 1.41, p = 0.24). These results suggest that, contrary to our
hypothesis, venom gland expression transcriptomes do not evolve faster than homologous,
evolutionary older tissues, but they have comparable evolutionary rates.
165
Are there venom-gland specific transcription modules?
The observed convergence between venom gland transcriptomes might be driven by a set of genes
that have similar expression patterns, therefore producing tissue identity. We identified 'modules' by isa
clustering (20) based on the expression matrix of all tissue types. In total, we found 62 modules (Fig.
4A and *SI Appendix*, Dataset S2 and S3), of which 36 included venom gland samples. Of these, 15
were exclusively venom-gland specific, 12 included venom glands and various glandular tissue
samples, and the remaining nine were a mix of venom glands and other tissues. The venom gland-
specific modules differed in species composition (5 to 17 species) and number of orthogroups (44 to
411 orthogroups); notably, modules 13, 24 and 40 included almost all species, thus representing a core
gene set of 209 orthogroups. Some modules had lineage-enriched expression patterns, particularly for
snakes; for instance, three modules included all snake species and one octopus, four modules included
snakes and insect species, and one module included snakes and echidna. However, the genes in these
snake-enriched modules were mostly also found in other venom-gland modules.
Next, we screened for significant enrichment of KEGG pathways and GO functional categories based
on annotation of human orthologs (Fig. 4B and *SI Appendix*, Dataset S4 and S5). Venom gland-specific
modules were particularly enriched in pathways and in GO categories related to protein processing in
endoplasmic reticulum (ER), secretion, transport, and particularly to stress response mechanisms (Fig.
4C). Similar results were obtained using *Drososophila melanogaster* orthologs (*SI Appendix*, Fig. S7).
Also, we used TopAnat (21) to identify the anatomical structures where these gene modules are
particularly expressed in an organism without venom glands. Genes belonging to venom gland specific
modules were enriched in type B pancreatic cells and epithelial cells of pancreas in human, salivary
glands and embryonic foregut in *D. melanogaster*. Modules which also included other glands were also
enriched in entities of the oral and digestive system such as "mucosa of the sigmoid colon", or "pylorus".
A link between venom glands and pancreas has been previously observed in snakes (8), and our results
give a further hint that venom glands may have co-opted components from multiple anatomical origins.
The orthogroups with the highest weight values in venom gland specific modules included genes
involved in protein processing in ER and in ER stress, such as SEL1, SEC63, ERP44, DNAJC3, ER
oxidoreductin 1, and SPCS3. Interestingly, the top orthogroups of the modules including venom glands
and other glandular tissues were also related to protein secretion, such as UBA5, GRASP55/65,
SRP54, and TM9SF, but they were not specifically related to ER or ER stress. The top orthogroups in
the snake-enriched modules were mostly toxins or proteins also found in other venom gland modules.
These findings emphasize the extreme secretory capacity of venom glands, and indicate that venom
production necessitates the activation of stress response mechanisms. Dedicated protein-producing
gland cells, such as those in venom-secreting tissue, have an exceptionally high secretory load relative
to most cells. Consequently, during the emergence of venom gland cells, supporting mechanisms must
have evolved to accommodate mass protein trafficking. One key mechanism is the unfolded protein
response (UPR), which ensure reliable folding of proteins in the ER (22, 23). Our results suggest a
central role of stress response mechanisms in enabling extreme cellular performance of venom glands,
and this same mechanism seems to have been repeatedly adopted across the animal kingdom.
205
Lineage-specific molecular mechanisms underlie convergence in general function of venom
transcriptomes
The results so far point to convergence in gene expression profiles and the presence of venom gland-
specific modules suggests concerted expression changes of genes involved in the secretory function.
However, our first approach restricts the analysis to relatively few conserved orthogroups shared among
all taxa, which might hinder the identification of lineage-specific patterns, or of patterns in fast-evolving
genes missed by ortholog detection. To have a better resolution of the molecular underpinnings of

venom gland activity, we performed differential expression and functional enrichment analyses for each species separately. For these species-level analyses, we retained only the 16 species for which we had multiple tissues and samples from the same study (Table 1), but we used all genes and enrichment analyses were based on each species' annotation.

Across all species, the most upregulated genes, besides, of course, toxins, were involved in protein secretion and metabolism pathways (*SI Appendix*, Dataset S6). High level regulators of the unfolded protein response (UPR), e.g. ATF6, PERK, and IRE1, were upregulated in most lineages, confirming the isa-clustering results on orthogroups. Pathways that were enriched only in specific lineages were related to communication, such as "ECM-receptor interaction", which was significantly upregulated in spiders, octopus and catfish, or "MAPK signaling pathway" and "GnRH signaling pathway" in all studied vertebrates (snakes, echidna, and catfish). A possible interpretation of these results is that the genes involved in the main function of the organ, i.e. protein secretion, are convergent, whereas the expression patterns of their regulators are inherited from developmental precursors, hence differences in signaling pathways might reflect the different origins of venom glands.

Enrichment analysis of GO biological process terms revealed that the most upregulated genes were mainly related to tissue development, regulation, signaling, transport, and metabolic and biosynthetic processes. A specific GO term was rarely found in more than one species, i.e. most GO terms were singletons; however, the enriched terms were semantically similar and grouped together (Fig. 5 and *SI Appendix*, Dataset S7). Additionally, some processes were found only in certain lineages, for instance reproduction and behavior were enriched only in scorpion, signaling pathways and nervous system in octopus, and the immune system in catfish. Membrane and ER were commonly enriched cellular components, while in some lineages extracellular region was also enriched.

The reasons why species were enriched with unique GO terms could be technical - GO annotations were done for each species separately, and this might have resulted in orthologous proteins being assigned to slightly different GO terms. Also, different sets of tissues were used in the differential expression analyses, and genome annotations vary in completeness between lineages. Nonetheless, our results suggest that there is an overlap in general biological processes, but that different lineages might have evolved different specific molecular mechanisms to perform the same general function.

Even though global expression profiles are similar, the way in which biological processes are regulated might differ between species, hence convergence in function might not necessarily correspond to convergence in gene regulatory networks. To test this hypothesis, we examined which transcription factors (TFs) were upregulated ($log_2FC > 1$; FC = fold-change venom/average other tissues) in venom glands compared to other tissues. TFs were identified using the KEGG database and all the information related to the TFs were retrieved from the Human Protein Atlas (24), UniProt (25), and Bgee databases (21). The most upregulated TFs within each taxon were involved in ER stress and UPR response, in agreement with our previous results, but they were also largely involved in pluripotency, cell differentiation, and tissue development (*SI Appendix*, Dataset S8). We found thirteen orthologous TFs shared across all lineages, some of which were also found in venom gland-specific modules. These included TFs involved in the UPR and response to ER stress pathways (XBP1, CREB3), or typically expressed in the epithelium (ETS, BHLHB8/MIST1, BNC), but the majority were involved in cell proliferation, differentiation and growth (TWIST, MXD-MAD, FOXP2_4, SOX9). Various homeobox gene families that play pivotal roles in tissue development and differentiation were among the most upregulated TFs, but different members were found in different lineages: the NKX-homeodomain factor family, with NKX2.5 expressed in octopus, NKX1 in scorpions, and NKX3-1 in echidna and catfish; the SIX family, with SIX1 found in snakes and spiders while SIX4 in octopus, PBX, with PBX1 highly expressed exclusively in spiders and scorpions (Arachnida) and PBX3 in spiders and snakes, as well as DLX, and LHX. Myogenic factors were also upregulated, with MYOF5 found exclusively in spiders and MYOD1 in snakes (*SI Appendix*, Dataset S8).

An interesting finding was the high upregulation of the abdominal-B homeobox ABDB exclusively in scorpions and wasps. This TF specifies the identity of the posterior abdominal segments, the external genitalia and gonads, and is involved in regulating post-mating response. Another TF found only in these two lineages was the krueppel (KR) factor which is involved in differentiation of the Malpighian tubules, a type of excretory system in the posterior region of the alimentary canal of some arthropods. Compared to the other animals investigated, scorpions and wasps have in common the location of their venom apparatus which is in the posterior part of their body, or metasoma. In scorpions, venom glands are located in the telson where also cuticular pits and dermal gland openings have been described (26). The function of these other glands is not really understood, but it is hypothesized that they might produce sex pheromones and play a role in courtship (26). In parasitoid wasps, like those included in our study, venom glands are in the posterior dorsal surface and are connected to the female reproductive system. The close interaction between these two systems complements their functioning,

since venom is injected through a modified ovipositor to ensure successful development of the offspring
in the host. Several genes annotated with GO terms related to reproduction were expressed in the
venom glands of both wasps and scorpions, and in the latter behavior and reproduction terms were
significantly enriched (*SI Appendix*, Dataset S7). The exact function of these genes within venom glands
is unknown. Nonetheless, these findings provide a first evidence that venom gland regulatory networks
have evolved, to some extent, from the co-option of pre-existing genetic regulatory circuits of the tissues
from which venom glands derive, or that are most closely related to.
The high number of genes and magnitude of enrichment related to cell cycle regulation in our results is
intriguing, and might be indicative of high epithelial cell turnover. In many lineages, e.g. in scorpions
(27), spiders (28), echidna (17), and catfish (29), venom is released by holocrine or apocrine modes of
secretion, which cause cellular damage or complete destruction of the cell. Moreover, the high levels
of cellular stress caused by massive toxin production might result in DNA damage, apoptosis, or cellular
dysfunction. As a consequence, the activation of a regulatory network for epithelial cell turnover and
maintenance might be necessary. Undifferentiated epithelial cells have been morphologically identified
in the venom gland epithelium of various organisms, e.g. in spiders (30), scorpions (27), snakes (31).
Furthermore, non-venom epithelial supporting cells and stromal cells, which express stem cell markers
and niche factors, have been observed in snake venom gland single cell sequencing (31), supporting
our finding of active cell growth and differentiation pathways in venom glands. These findings combined
with our results suggest that conserved as well as lineage-specific regulators involved in cell
differentiation and organ development have been repeatedly adopted during the evolution of venomous
animals and that, besides secretion, regulation of cell cycle is a central task of venom glands.
294
295
Conclusions
Many animal cell types possess the capacity for protein secretion, and a conserved molecular and
organellar pathway exists for routing translated proteins out of the cell (23). However, dedicated protein-
secreting cells, such as those producing venom, have an exceptionally high secretory load. During the
evolutionary assembly of venom glands, stress response mechanisms seem to have been repeatedly
adopted by different animals to cope with mass protein production. The resulting DNA damage,
apoptosis and even complete destruction of those venom-producing cells with holocrine secretion,
activate a regulatory network for epithelial development, cell turnover and maintenance. While sets of
genes directly involved in the secretory function have similar expression patterns across animal
lineages, and might thus be conserved, the way in which cells are regulated and communicates, are
different between lineages, and might reflect their diverse developmental origins. Our findings provide
a first evidence that venom gland regulatory networks have evolved, to some extent, from the co-option
of pre-existing genetic regulatory circuits from the tissue most closely related to each venom glands.
This study represents the first step towards an understanding of the molecular mechanisms underlying
the convergent evolution of one of the most successful adaptive traits in the animal kingdom.
311
312
313 **Material and methods**
314
Species selection
For the analysis, we selected only venomous species with either an annotated genome or high quality
de novo transcriptome available for the same species or a close relative, and with RNA-seq data of
venom glands and other body tissues (Table 1 and *SI Appendix*, Table S1).
Considering the mixed nature of our dataset, for each species, we reduced the proteomes (proteins
from annotated genomes and *de novo* transcriptomes) to a set of non-redundant sequences as follows:
first, we filtered within-species 100% identical amino acid sequences with cdhit 4.6 (32) to select one
representative sequence. Then, we compared with BlastP (33) the sequences against the NCBI non-
redundant database (downloaded on 09.04.2020), the Uniprot-Toxprot (34) or Arachnoserver (35)
databases, and retained only those with evalue < 1e-05. These processed proteomes were used for
subsequent analyses.
326
Orthogroup assignment
For each species, we assigned protein sequences to orthogroups using the mapping tool in OrthoDB
v.10.1 (36). For the mapping we selected up to five closest taxa to assign proteins to orthogroups at
the Metazoa node. In parallel, we compared the proteomes with BlastP against the same species
selected for OrthoDB mapping. For each species, all Blast outputs were combined, and we kept only
one hit per sequence (the one with the lowest evalue). Blast and OrthoDB mapping were then merged

and proteins assigned to orthogroups using the OG2genes file at the Metazoa node. Species used for
the orthogroup assignment are listed in *SI Appendix*, Table S2.
Orthogroups containing proteins reported as venom components in the reference
genome/transcriptome paper, or that had the best Blast hit against a sequence in the Uniprot-Toxprot
or Arachnoserver databases, were assigned as toxin-containing orthogroups.
338
Expression levels
RNA-seq data were obtained from the Sequence Read Archive (SRA) database. Only Illumina SRA
reads were selected; where possible, we selected at least three libraries for each tissue from each
taxon, and only data generated from healthy, adult tissues were used. Raw fastq files were filtered with
trimmomatic (37), their quality checked with fastQC (38), and quantified with kallisto (39) using default
parameters for paired-end reads, and parameters -l 55 -s 1e-08 for single-end reads. All species were
mapped to their own specific transcriptome with the exception of *Microplitis mediator*, which was
mapped to *M. demolitor*, and *Octopus vulgaris* which was mapped to *O. bimaculoides*. We used *tximport*
(40) to estimate transcript abundances as Transcript Per Million (TPM) for the metazoan-level analysis,
and to aggregate read counts at the gene-level for the species-level analysis (see below).
349
Orthogroup expression matrix
For the comparative analysis at the Metazoa node, orthogroup-level abundances were obtained as
follows. Since most orthogroups included more than one protein per species (i.e. one-to-many or many-
to-many orthologs), we selected one representative sequence for each orthogroup in each species as
the transcript with the highest expression in venom gland samples (*SI Appendix* Dataset S9), and used
the TPM value estimated for that transcript as the orthogroup expression value. All samples of all
species were then merged into a matrix of orthogroup TPM values. To validate our method, we
randomly selected one transcript per orthogroup, and we obtained similar results (see *SI Appendix*, Fig.
S1 and S8).
Because the samples are from different experiments, to allow for comparison across samples, first, we
minimized the effects of technical artifacts by quantile normalization on log_2 transformed TPM values
to which a pseudo count of 1 was added to prevent $log_2(0)$ scores. Then, we removed the batch effect
caused by using multiple species and multiple SRA studies, using an empirical Bayes method
implemented via the ComBat function in the sva R package (41) which has proven to be efficient with
these kinds of datasets (13). Finally, we calculated tissue-level expression as mean TPM.
365
Transcriptome similarity analysis
All analyses were performed in R version 3.6.2 (42). To obtain an overview of expression patterns we
performed principal component analysis (PCA) with the rda function in *vegan* (43). To understand
whether the observed pattern was biased by the shared expression levels of toxins in venom glands,
we re-ran the analysis excluding the orthogroups containing toxin sequences.
We quantified transcriptome similarity between tissues as Spearman rank correlation coefficients to test
whether venom glands were more similar to each other than to any other tissue of the same species.
Mean pairwise distances were compared with the Wilcoxon signed-rank test and p-values adjusted with
(44).
375
Phylogenetic tree
We constructed a phylogenetic species tree using 77 one-to-one orthologs. First, proteome redundancy
was reduced by selecting a representative protein for all sequences > 90% identical using cd-hit 4.6.
Then, we selected orthogroups that were single-copy in all species, or all except one. Orthogroup
sequences were aligned with mafft 7.310 (45) and trimmed with trimAl 1.4.1 (46). The trimmed
alignments were concatenated, trimmed and used to build the phylogenetic tree with raxml 8.2.12 (18),
and bootstrap values of the consensus tree were obtained based on 100 replicates. The tree was rooted
at the deuterostome-protostome split using the function root_in_edge of the R package *castor* (47).
384
Expression trees
Expression trees were constructed for venom gland, ovary, and brain tissues. These homologous
organs were chosen for the comparisons because they had the highest number of samples (minimum
seven species).
We used the R package *ape* (48) to construct Neighbor-Joining expression trees based on two distance
measures: 1 – Spearman rank correlation coefficient and Euclidean distances. The reliability of
branching patterns was assessed with bootstrap analysis using 1000 replicates. When possible, trees
were rooted at the deuterostome-protostome split. To verify whether closer taxa have more similar

transcriptomes, we obtained pairwise distances using the function get_all_pairwise_distances in the
package *castor*, and tested for correlation between phylogenetic and expression matrices with Mantel
tests in the R package *vegan*. Expression divergence rates were calculated as the slope of linear
regressions (lm) between pairwise expression distances and phylogenetic distances.
397
Transcription Modules
We identified orthogroups with similar expression patterns using the iterative signature algorithm (isa)
implemented in the *isa2* Bioconductor package (20) with default parameters. Briefly, isa identifies, in an
unsupervised manner, sets of genes that exhibit coherent expression patterns over subsets of samples
from large sets of expression data. It selects genes that are significantly under- or over-expressed in a
random seed of samples, and then all samples are scored by the weighted average expression levels
across these genes.
KEGG pathway and GO enrichment analyses of all isa modules were performed with *clusterProfiler*
(49) based on human gene annotation after converting the OrthoDB ClusterId to NCBI EntrezId using
the OG2gene file at the Metazoa node obtained from the OrthoDB data page. For both analyses, the
foreground genes were the orthogroups of a module and the background genes were all the 2528
shared orthogroups.
For enrichment of anatomical structures, we used the TopAnat tool in Bgee 14.2 (21). TopAnat is based
on TopGO (50) and works similar to a GO enrichment test except that it analyzes Uberon anatomical
identities (51) where genes are expressed instead of GO terms. To find the anatomical entities in which
tissue-specific gene modules were enriched, we run TopAnat based on human annotation and with the
weight algorithm.
415
Species-level differential expression analysis
To identify taxon-specific patterns, we performed differential expression analysis at the gene-level for
each species separately using *edgeR* (52). Genes with low expression were automatically excluded
with the function filterByExpr. Differences in library size were accounted for using TMM-normalization
factors. We fit a quasi-likelihood (QL) negative binomial generalized log-linear model (glmQLFit
function) to estimate empirical Bayes moderated dispersion; this method accounts for gene-specific
variability from both biological and technical sources (53). Empirical Bayes QL F-tests were used to
compare gene expression levels between venom glands and the average of the other tissue types.
Genes with false discovery rate < 0.05 and $\log_2$ fold change > 1 were considered significantly
upregulated.
KEGG pathway enrichment analysis of upregulated genes was performed with *clusterProfiler*. For the
species which were not in the list of supported organisms in the KEGG catalog, we used the annotation
of the closest species. Orthologs between the two species were identified by reciprocal BlastP.
As none of the species in this study have available GO annotations, we annotated all the proteomes
using the program CrowdGO (54). CrowdGO is a consensus-based GO term meta-predictor that
employs machine learning models combined with GO term semantic similarities and information
contents to leverage strengths of individual predictors and produce comprehensive and accurate gene
functional annotations. For the GO annotations in this study, we used the CrowdGOFull model which
utilizes annotations from DeepGOPlus (55), FunFams (56), InterProScan (57), and Wei2GO (58).
GO enrichment analyses for the biological processes, molecular functions and cellular compartments
were performed with TopGO (50) using the elim algorithm and fisher statistic test. We summarized the
most enriched processes (pvalue < 0.01) across all species with GO-Figure! which reduces redundancy
by grouping together GO terms with similar functions, and produces semantic similarity scatterplots
where representative terms are plotted (59).
440
Transcription factors analysis
Finally, we focused on expression patterns of transcription factors (TFs) to test for convergence in gene
regulatory networks. We examined which transcription factors were upregulated in venom glands
compared to other tissues, and whether these TFs were expressed across all species or only within
certain lineages. For each species we downloaded the annotations from the KEGG BRITE database
('03000 Transcription factors'). For species which were not included in the KEGG Organism catalog,
we used the annotation of the closest species (as we did for the KEGG enrichment analysis). Because
one KEGG orthology entry can contain multiple genes (similarly to the OrthoDB orthogroups), for the
comparative analysis we kept one representative gene per KEGG ortholog.
450
451
452   **Acknowledgments and funding sources**

This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 845674 to GZ, and the Swiss National Science Foundation grant to PP00P3_170664 (RMW). We thank Nadia Ayoub for providing the amino acid sequences of the spider de novo transcriptomes, Robert Haney Zong-ji Wang for providing toxin sequences, and MRR lab members for bioinformatics support.

References

1. D. L. Stern, The genetic causes of convergent evolution. *Nat. Rev. Genet.* **14**, 751–764 (2013).

2. J. Arendt, D. Reznick, Convergence and parallelism reconsidered: what have we learned about the genetics of adaptation? *Trends Ecol. Evol.* **23**, 26–32 (2008).

3. P.-A. Christin, D. M. Weinreich, G. Besnard, Causes and evolutionary significance of genetic convergence. *Trends Genet.* **26**, 400–405 (2010).

4. K. R. Elmer, A. Meyer, Adaptation in the age of ecological genomics: insights from parallelism and convergence. *Trends Ecol. Evol.* **26**, 298–306 (2011).

5. Z. Zou, J. Zhang, Are convergent and parallel amino acid substitutions in protein evolution more prevalent than neutral expectations? *Mol. Biol. Evol.* **32**, 2085–2096 (2015).

6. T. B. Sackton, N. Clark, Convergent evolution in the genomics era: new insights and directions. *Philos. Trans. R. Soc. B Biol. Sci.* **374**, 20190102 (2019).

7. V. Schendel, L. D. Rash, R. A. Jenner, E. A. B. Undheim, The diversity of venom: the importance of behavior and venom system morphology in understanding its ecology and evolution. *Toxins* **11**, 666 (2019).

8. G. Zancolli, N. R. Casewell, Venom systems as models for studying the origin and regulation of evolutionary novelties. *Mol. Biol. Evol.* **37**, 2777–2790 (2020).

9. M. Holford, M. Daly, G. F. King, R. S. Norton, Venoms to the rescue. *Science* **361**, 842–844 (2018).

10. V. Herzig, *et al.*, Animal toxins — Nature's evolutionary-refined toolkit for basic research and drug discovery. *Biochem. Pharmacol.* **181**, 114096 (2020).

11. M. V. Modica, *et al.*, The new COST Action European Venom Network (EUVEN)—synergy and future perspectives of modern venomics. *GigaScience* **10** (2021).

12. B. W. Perry, D. R. Schield, A. K. Westfall, S. P. Mackessy, T. A. Castoe, Physiological demands and signaling associated with snake venom production and storage illustrated by transcriptional analyses of venom glands. *Sci. Rep.* **10**, 18083 (2020).

13. A. Barua, A. S. Mikheyev, An ancient, conserved gene regulatory network led to the rise of oral venom systems. *Proc. Natl. Acad. Sci.* **118** (2021).

14. M. dos Reis, *et al.*, Uncertainty in the timing of origin of animals and the limits of precision in molecular timescales. *Curr. Biol.* **25**, 2939–2950 (2015).

15. D. R. Schield, *et al.*, The origins and evolution of chromosomes, dosage compensation, and mechanisms underlying venom regulation in snakes. *Genome Res.* **29**, 590–601 (2019).

16. T. N. W. Jackson, *et al.*, Endless forms most beautiful: the evolution of ophidian oral glands, including the venom system, and the use of appropriate terminology for homologous structures. *Zoomorphology* **136**, 107–130 (2017).

494   17.   W. J. Krause, Morphological and histochemical observations on the crural gland-spur apparatus of
495   the echidna (*Tachyglossus aculeatus*) together with comparative observations on the femoral
496   gland-spur apparatus of the duckbilled platypus (*Ornithorhyncus anatinus*). *Cells Tissues Organs*
497   **191**, 336–354 (2010).

498   18.   A. Stamatakis, RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
499   phylogenies. *Bioinformatics* **30**, 1312–1313 (2014).

500   19.   G. Zancolli, *et al.*, When one phenotype is not enough: divergent evolutionary trajectories govern
501   venom variation in a widespread rattlesnake species. *Proc. R. Soc. B Biol. Sci.* **286**, 20182735
502   (2019).

503   20.   G. Csárdi, Z. Kutalik, S. Bergmann, Modular analysis of gene expression data with R.
504   *Bioinformatics* **26**, 1376–1377 (2010).

505   21.   F. B. Bastian, *et al.*, The Bgee suite: integrated curated expression atlas and comparative
506   transcriptomics in animals. *Nucleic Acids Res.* **49**, D831–D847 (2021).

507   22.   C. Hetz, The unfolded protein response: controlling cell fate decisions under ER stress and beyond.
508   *Nat. Rev. Mol. Cell Biol.* **13**, 89–102 (2012).

509   23.   A. Brückner, J. Parker, Molecular evolution of gland cell types and chemical interactions in animals.
510   *J. Exp. Biol.* **223** (2020).

511   24.   M. Uhlén, *et al.*, Tissue-based map of the human proteome. *Science* **347** (2015).

512   25.   The UniProt Consortium, UniProt: the universal protein knowledgebase in 2021. *Nucleic Acids Res.*
513   **49**, D480–D489 (2021).

514   26.   R. Foelix, B. Erb, M. Braunwalder, Fine structure of the stinger (aculeus) in *Euscorpius*. *J. Arachnol.*
515   **42**, 119–122 (2014).

516   27.   B. A. Soliman, N. M. Shoukry, M. E. Mohallal, H. A. W. Fetaih, H. S. khaled, Fine structure of the
517   stinger, histology and histochemistry of the venom gland in the scorpion *Androctonus amoreuxi*
518   (Buthidae). *J. Basic Appl. Zool.* **66**, 41–46 (2013).

519   28.   H. Malli, L. Kuhn-Nentwig, H. Imboden, M.-J. Moon, T. Wyler, Immunocytochemical localization
520   and secretion process of the toxin CSTX-1 in the venom gland of the wandering spider *Cupiennius*
521   *salei* (Araneae: Ctenidae). *Cell Tissue Res.* **299**, 417–426 (2000).

522   29.   J. J. Wright, "Evolutionary History of Venom Glands in the Siluriformes" in *Evolution of Venomous*
523   *Animals and Their Toxins*, Toxinology., A. Malhotra, Ed. (Springer Netherlands, 2017), pp. 279–
524   301.

525   30.   L. D. Foil, L. B. Coons, B. R. Norment, Ultrastructure of the venom gland of the brown recluse
526   spider, *Loxosceles reclusa* gertsch and mulaik (Araneae: Loxoscelidae). *Int. J. Insect Morphol.*
527   *Embryol.* **8**, 325–334 (1979).

528   31.   Y. Post, *et al.*, Snake venom gland organoids. *Cell* **180**, 233-247.e21 (2020).

529   32.   W. Li, A. Godzik, Cd-hit: a fast program for clustering and comparing large sets of protein or
530   nucleotide sequences. *Bioinformatics* **22**, 1658–1659 (2006).

531   33.   S. F. Altschul, *et al.*, Gapped BLAST and PSI-BLAST: a new generation of protein database search
532   programs. *Nucleic Acids Res.* **25**, 3389–3402 (1997).

533   34.   F. Jungo, L. Bougueleret, I. Xenarios, S. Poux, The UniProtKB/Swiss-Prot Tox-Prot program: a
534   central hub of integrated venom protein data. *Toxicon* **60**, 551–557 (2012).

35. S. S. Pineda, *et al.*, ArachnoServer 3.0: an online resource for automated discovery, analysis and
annotation of spider toxins. *Bioinforma. Oxf. Engl.* **34**, 1074–1076 (2018).

36. E. V. Kriventseva, *et al.*, OrthoDB v10: sampling the diversity of animal, plant, fungal, protist,
bacterial and viral genomes for evolutionary and functional annotations of orthologs. *Nucleic Acids*
Res. **47**, D807–D811 (2019).

37. A. M. Bolger, M. Lohse, B. Usadel, Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics **30**, 2114–2120 (2014).

38. S. Andrews, FastQC: A quality control tool for high throughput sequence data. Available online at:
http://www.bioinformatics.babraham.ac.uk/projects/fastqc (2010) (June 4, 2021).

39. N. L. Bray, H. Pimentel, P. Melsted, L. Pachter, Near-optimal probabilistic RNA-seq quantification.
Nat. Biotechnol. **34**, 525–527 (2016).

40. C. Soneson, M. I. Love, M. D. Robinson, Differential analyses for RNA-seq: transcript-level
estimates improve gene-level inferences. *F1000Research* **4**, 1521 (2015).

41. J. T. Leek, W. E. Johnson, H. S. Parker, A. E. Jaffe, J. D. Storey, The sva package for removing
batch effects and other unwanted variation in high-throughput experiments. *Bioinformatics* **28**, 882–
883 (2012).

42. R Core Team, R: A language and environment for statistical computing. R Foundation for Statistical
Computing, Vienna, Austria. URL http://www.R-project.org/ (2019) https:/doi.org/URL
http://www.R-project.org/.

43. J. Oksanen, *et al.*, vegan: community ecology package. R package version 2.5-6. https://CRAN.R-
project.org/package=vegan (2019).

44. Y. Benjamini, Y. Hochberg, Controlling the false discovery rate: a practical and powerful approach
to multiple testing. *J. R. Stat. Soc. Ser. B Methodol.* **57**, 289–300 (1995).

45. K. Katoh, K. Misawa, K. Kuma, T. Miyata, MAFFT: a novel method for rapid multiple sequence
alignment based on fast Fourier transform. *Nucleic Acids Res.* **30**, 3059–3066 (2002).

46. S. Capella-Gutiérrez, J. M. Silla-Martínez, T. Gabaldón, trimAl: a tool for automated alignment
trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972–1973 (2009).

47. S. Louca, M. Doebeli, Efficient comparative phylogenetics on large trees. *Bioinformatics* **34**, 1053–
1055 (2018).

48. E. Paradis, K. Schliep, ape 5.0: an environment for modern phylogenetics and evolutionary
analyses in R. *Bioinformatics* **35**, 526–528 (2019).

49. G. Yu, L.-G. Wang, Y. Han, Q.-Y. He, clusterProfiler: an R package for comparing biological themes
among gene clusters. *OMICS J. Integr. Biol.* **16**, 284–287 (2012).

50. A. Alexa, J. Rahnenfuhrer, topGO: enrichment analysis for gene ontology. R package version
2.38.1 (2019).

51. M. A. Haendel, *et al.*, Unification of multi-species vertebrate anatomy ontologies for comparative
biology in Uberon. *J. Biomed. Semant.* **5**, 21 (2014).

52. M. D. Robinson, D. J. McCarthy, G. K. Smyth, edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140 (2010).

53. S. P. Lund, D. Nettleton, D. J. McCarthy, G. K. Smyth, Detecting differential expression in RNA-
sequence data using quasi-likelihood with Shrunken dispersion estimates. *Stat. Appl. Genet. Mol.*
Biol. **11** (2012).

54. M. J. M. F. Reijnders, R. M. Waterhouse, CrowdGO: machine learning and semantic similarity
guided consensus Gene Ontology annotation. *bioRxiv*, 731596 (2021).

55. M. Kulmanov, R. Hoehndorf, DeepGOPlus: improved protein function prediction from sequence.
Bioinformatics **36**, 422–429 (2020).

56. L. Scheibenreif, M. Littmann, C. Orengo, B. Rost, FunFam protein families improve residue level
molecular function prediction. *BMC Bioinformatics* **20**, 400 (2019).

57. M. Blum, *et al.*, The InterPro protein families and domains database: 20 years on. *Nucleic Acids*
Res. **49**, D344–D354 (2021).

58. M. J. M. F. Reijnders, Wei2GO: weighted sequence similarity-based protein function prediction.
bioRxiv, 2020.04.24.059501 (2020).

59. M. J. M. F. Reijnders, R. M. Waterhouse, Summary visualizations of gene ontology terms with GO-
Figure! *Front. Bioinforma.* **1** (2021).

Figures and Tables

Figure 1. Global patterns of gene expression differences between multiple tissues and lineages. Principal component analysis based on the normalized expression levels (TPM) of 2,528 shared orthogroups among all taxa. The proportions of variance explained by the components are in parenthesis. 'Body tissues' include: abdomen, body tissue, cephalothorax, and viscera; 'Other glands' include: accessory venom gland, hypopharyngeal gland, rictal gland, salivary gland and silk gland; 'Muscle tissues' include: muscle, proboscis, heart, and leg; 'Other organs' include: liver, kidney, and pancreas.

Figure 2. Transcriptome similarity between venom glands and other tissues. For each species, interspecific similarity (Spearman rank correlation coefficient ρ) between venom glands is compared to intraspecific coefficients between the venom gland and all other tissues for that species. Significant comparisons (Wilcoxon signed-rank test, $p < 0.05$ before correction) are indicated with an asterisk, although they were not significant after Benjamini-Hochberg correction. The low data points correspond to the pairwise correlations with echidna.

Figure 3. Comparison of venom transcriptomes and species phylogeny. A: Phylogenetic species tree
(left), with circles marking the independent origins of venom in relation to venom gland expression tree
(right) with 100% bootstrap support throughout (not shown). B: Sequence-based phylogenetic distances
vs. venom gland expression distances (1-Spearman coefficient). Pair distances between echidna and
the other species are marked with triangles, all the others are circles. The dotted line indicates the
positive correlation between expression and phylogenetic distances excluding the echidna data points;
the corresponding correlation test values are in parentheses.

618
Figure 4. Tissue-specific modules and GO enrichment results. A: Heatmap based on Spearman
correlation coefficients between the 62 modules; most modules are tissue-specific and cluster together.
Definitions of tissue groups as in Fig. 1. B: Enrichment of the top 3 biological process GO terms of the
tissue-specific modules. Color bar representing the tissues as in A. C: Visualization of biological process
GO terms enriched in the venom gland core gene set (modules 13, 24 and 40) produced using GO-
Figure! (59). Each bubble represents a cluster of similar GO terms summarized by a representative
term reported in the legend, and sorted by the average p-values of the representative GO term across
the three modules. Bubble size indicates the amount of GO terms in each cluster, and the color is the
average p-value of the representative GO term across the gene modules. Similar clusters plot closer to
each other.

Figure 5. Semantic similarity scatterplot of GO biological process terms enriched in venom glands.
Functional enrichment of upregulated genes was performed separately for each species, and all
significant GO terms (p-value < 0.01) were summarized using GO-Figure! (59). Each bubble represents
a cluster of similar GO terms summarized by a representative term reported in the legend and sorted
by the amount of species with at least one term in the cluster. Bubble size indicates the number of terms
in the cluster and the color corresponds to the number of species in the cluster. Similar clusters plot
closer to each other.

Table 1. List of 20 species included in the analysis representing eight independent origins of venom,
one per lineage. The table reports the number of venom gland samples, number of other tissues and
the reference assemblies used for the analyses. Transcript and protein sequences were obtained either
from the NCBI Genome (GCA and GCF) databases, the NCBI Transcriptome Shotgun Assembly (for
de novo transcriptome assemblies), or from other repositories as indicated. *Species not used in the
species-level differential expression analysis.

Lineage	Species	# Venom gland samples	# Other tissues	Reference assembly
Wasps	*Apis cerana**	1	2	GCF_001442555.1
	Nasonia vitripennis	3	1	GCF_009193385.2
	*Microplitis demolitor**	1	1	GCF_000572035.2
	Microplitis mediator	4	1	GCF_000572035.2
Flies	*Dasypogon diadema*	2	2	http://dx.doi.org/10.5524/100612
Spiders	*Parasteatoda tepidariorum*	3	3	GCF_000365465.2
	Stegodyphus dumicola	3	3	GCF_010614865.1
	Steatoda grossa	2	3	GBJQ00000000.1 (*de novo*)
	Latrodectus geometricus	2	3	GBJM00000000.1 (*de novo*)
	Latrodectus hesperus	4	3	GBJN00000000.1 (*de novo*)
Scorpions	*Centruroides hentzi**	2	0	GCF_000671375.1
	Mesobuthus martensii	6	3	GCA_000484575.1
Octopi	*Octopus bimaculoides*	1	3	GCF_001194135.1
	*Octopus vulgaris**	1	1	GCF_001194135.1
Fishes	*Tachysurus fulvidraco*	1	2	GCF_003724035.1
Mammals	*Tachyglossus aculeatus*	1	5	GCF_015852505.1
Snakes	*Naja naja*	4	6	GCA_009733165.1
	Deinagkistrodon acutus	2	3	https://doi.org/10.5524/100196
	Protobothrops flavoviridis	3	7	GCA_003402635.1
	Crotalus viridis	3	8	https://doi.org/10.6084/m9.figshare.9031643.v1

18