Construction of a new chromosome-scale, long-read reference genome assembly for the Syrian hamster, *Mesocricetus auratus*

R. Alan Harris[1]   rharris1@bcm.edu

Muthuswamy Raveendran[1]   raveendr@bcm.edu

Dustin T. Lyfoung[2]   lyfoung@wisc.edu

Fritz J Sedlazeck[1]   fritz.sedlazeck@bcm.edu

Medhat Mahmoud[1]   medhat.mahmoud@bcm.edu

Trent M. Prall[3]   prall@wisc.edu

Julie A. Karl[3]   jakarl@wisc.edu

Harshavardhan Doddapaneni[1]   doddapan@bcm.edu

Qingchang Meng[1]   qingchang.meng@bcm.edu

Yi Han[1]   yhan@bcm.edu

Donna Muzny[1]   donnam@bcm.edu

Roger W. Wiseman[2, 3]   rwwiseman@wisc.edu

David H. O'Connor[2, 3]   dhoconno@wisc.edu

Jeffrey Rogers[1]   jr13@bcm.edu

(Corresponding author: jr13@bcm.edu; 713-798-7783)

[1]Human Genome Sequencing Center and Department of Molecular and Human Genetics,

Baylor College of Medicine, Houston, TX 77030

[2]Wisconsin National Primate Research Center, University of Wisconsin, Madison, WI 53711

[3]Department of Pathology and Laboratory Medicine, University of Wisconsin, Madison, WI

53711

Abstract

Background

The Syrian hamster (*Mesocricetus auratus*) has been suggested as a useful

mammalian model for a variety of diseases and infections, including infection with

respiratory viruses such as SARS-CoV-2. The MesAur1.0 genome assembly was

generated in 2013 using whole-genome shotgun sequencing with short-read sequence

data. Current more advanced sequencing technologies and assembly methods now

permit the generation of near-complete genome assemblies with higher quality and

greater continuity.

Findings

Here, we report an improved assembly of the *M. auratus* genome (BCM_Maur_2.0)

using Oxford Nanopore Technologies long-read sequencing to produce a chromosome-

scale assembly. The total length of the new assembly is 2.46 Gbp, similar to the 2.50

Gbp length of a previous assembly of this genome, MesAur1.0. BCM_Maur_2.0 exhibits

significantly improved continuity with a scaffold N50 that is 6.7 times greater than

MesAur1.0. Furthermore, 21,616 protein coding genes and 10,459 noncoding genes are

annotated in BCM_Maur_2.0 compared to 20,495 protein coding genes and 4,168

noncoding genes in MesAur1.0. This new assembly also improves the unresolved

regions as measured by nucleotide ambiguities, where approximately 17.11% of bases

in MesAur1.0 were unresolved compared to BCM_Maur_2.0 in which the number of

unresolved bases is reduced to 3.00%.

45   Conclusions

46   Access to a more complete reference genome with improved accuracy and continuity

47   will facilitate more detailed, comprehensive, and meaningful research results for a wide

48   variety of future studies using Syrian hamsters as models.

49   Keywords

50   Syrian hamster, *Mesocricetus auratus*, genome, disease model, COVID-19

Data Description

Introduction

The Syrian hamster (*Mesocricetus auratus*, NCBI:txid10036) has been used in biomedical research for decades because it is a good model for studies of cancer [1], reproductive biology [2] and infectious diseases [3,4], including SARS-CoV-2, influenza virus, and Ebola virus [5–9]. The use of Syrian hamsters in research has declined [10], likely due to advances in the genetic and molecular tools available for other rodents, especially laboratory mice, and not to a reduction in the utility of hamsters in biomedical research [3].

Syrian hamsters are particularly important for COVID-19 research. They spontaneously develop more severe lung disease than other animal models, such as wild-type mice, macaques, marmosets, and ferrets [5,11–14]. After intranasal infection, Syrian hamsters consistently show signs of respiratory distress, including labored breathing, but typically recover after 2 weeks [15]. This is in stark contrast to wild-type laboratory mice that are minimally susceptible to most SARS-CoV-2 strains that were circulating in 2020, though laboratory mice may be more susceptible to certain variants of concern that began circulating in 2021 [8,16]. Furthermore, a recent analysis has suggested that Syrian hamsters fed a high-fat, high-sugar diet exhibit accelerated weight gain and pathological changes in lipid metabolism, as well as more severe disease outcomes when subsequently infected with SARS-CoV-2 [17]. This result has obvious parallels with observations of the effects of comorbidities in humans suffering from COVID-19.

COVID-19 pathology in Syrian hamsters appears to be due to a dysregulated innate

immune response involving signal transducer and activator of transcription factor 2

(STAT2)-dependent type I (IFN-I) and type III interferon (IFN-III) signaling [18]. IFN-I

signaling can limit virus replication and dissemination and it has been shown that

intranasal administration of IFN-I in Syrian hamsters reduces viral load and tissue

damage [19]. The human angiotensin-converting enzyme 2 (ACE2) was identified as

the cell entry receptor of SARS-CoV-2 [20]. In addition, upon the engagement of ACE2

with SARS-CoV2, cellular transmembrane protease 'serine 2' (TMPRSS2) mediates the

priming of viral spike (S) protein by cleaving at the S1/S2 site and inducing the fusion of

viral and host cellular membranes, thus facilitating viral entry into the cells [21]. Human

ACE2 and hamster ACE2 receptors had previously been shown to share substantial

sequence homology, which strongly points to interaction with SARS-CoV-2 receptor

binding domain (RBD) structures and similar binding affinity [22]. *In silico* interaction

prediction analysis suggests that human and hamster TMPRSS2 are structurally very

similar. Even with slight differences in amino acid residue interactions, human and

hamster TMPRSS2 activity are identical for residue interactions related to SARS-CoV-2

infectivity [22]. As COVID-19 causes systemic disease in people, precision modeling of

specific aspects of pathogenesis will require carefully evaluating similarities and

differences across various biological processes in humans and Syrian hamsters which,

in turn, will require extensive genomic comparisons between the two species.

The currently available reference genome sequence for the Syrian hamster was

produced in 2013 using a whole-genome shotgun sequencing approach implementing

short read sequencing technology. The resulting MesAur1.0 reference sequence

(Genbank accession number GCA_000349665.1) is typical of those produced at that

time, containing 237,699 separate contigs with contig N50 of 22,512 bp. The quality and

research potential of the existing Syrian hamster genome is limited by the technology

that was available at the time of its development; for example, the cluster of type I

interferon genes was not resolvable with this technology. In this Data Note, we report

the production of a new Syrian hamster reference genome that was sequenced using

long-read methods on the Oxford Nanopore Technologies (ONT) PromethION platform

and assembled into highly contiguous chromosomes using a combination of Flye [23]

and Pilon [24] assembly software. The final assembly, BCM_Maur_2.0, improves upon

quality and contiguity in comparison with MesAur1.0, with longer contigs and more

contiguous sequence, allowing for a more complete reference genome with improved

accuracy that will benefit a wide variety of future studies using the Syrian hamster

reference genome.

Methods

DNA isolation, library construction, and sequencing

All genomic DNAs for this study were isolated from a single female LVG Golden Syrian

hamster (SY011) that was purchased from Charles River, Inc. (Kingston, NY). All

procedures were performed in accordance with the guidelines set by the Institutional

Animal Care and Use Committee at the University of Wisconsin-Madison. The protocol

was approved by the Institutional Animal Care and Use Committee at the University of

Wisconsin-Madison (protocol number V00806). Data from this individual are available in

NCBI BioProject PRJNA705675, BioSamples SAMN18096087 and SAMN18096088.

Qiagen AllPrep DNA/RNA Mini kits were used to extract DNA from frozen liver while

Qiagen Blood and Cell Culture DNA Midi Kits were used for extractions from frozen

kidney. Ultra-high molecular weight DNA for optical mapping was purified from frozen

liver using an Animal Tissue DNA Isolation Kit from Bionano Genomics, Inc. (San

Diego, CA).

123 Oxford Nanopore long-read sequencing

We prepared three separate genomic DNA isolates from the same Syrian hamster

(BioSample SAMN18096087). These aliquots were sheared to distinct target fragment

lengths (10 kb, 20kb and 30kb) in order to assess the effect of fragment size on flowcell

yield and improve efficiency. The two smaller length fragment libraries were sheared

using Covaris gTube and the 30kb targeted size library was fragmented with Diagnode

Megarupter 3, all following manufacturer's recommendations. The Oxford Nanopore

sequencing libraries were prepared using the ONT 1D sequencing by ligation kit (SQK-

LSK109). Briefly, 1-1.5ug of fragmented DNA was repaired with the NEB FFPE repair

kit, followed by end repair and A-tailing with the NEB Ultra II end-prep kit. After a clean

up step using AMPure beads, the prepared fragments were ligated to ONT specific

adapters via the NEB blunt/TA master mix kit. Each library underwent a final clean up

and was loaded onto a PromethION flow cell per manufacturer's instructions. One

library was sequenced per flow cell with standard parameters for 72 hrs. Base-calling

was done onboard the PromethION instrument using neuronal network based software

(Oxford Nanopore Technologies, UK).

Illumina sequencing

500ng of input genomic DNA from a kidney sample (BioSample SAMN18096088) was used to generate standard PCR-free Illumina paired-end sequencing libraries. Libraries were prepared using KAPA Hyper PCR-free library reagents (KK8505, KAPA Biosystems) in Beckman robotic workstations (Biomek FX and FXp models). Total genomic DNA was sheared into fragments of approximately 200-600 bp in a Covaris E220 system (96-well format) followed by purification of the fragmented DNA using AMPure XP beads. A double size selection step was employed, with different ratios of AMPure XP beads, to select a narrow size band of sheared DNA molecules for library preparation. DNA end-repair and 3'-adenylation were then performed in the same reaction followed by ligation of the barcoded adaptors to create PCR-Free libraries. The resulting libraries were evaluated using the Fragment Analyzer (Advanced Analytical Technologies, Ames, Iowa) to assess library size and presence of remaining adaptor dimers. This was followed by qPCR assay using KAPA Library Quantification Kit and their SYBR FAST qPCR Master Mix to estimate the size and quantify fragment yield.

Sequencing was performed on the NovaSeq 6000 Sequencing System using the S4 reagent kit (300 cycles) to generate 2 x 150 bp paired-end reads. The final concentration of the libraries loaded on flowcells was 400-450 pM. Briefly, the libraries were diluted in an elution buffer and denatured in sodium hydroxide. The denatured libraries were loaded into each lane of the S4 flow cell using the NovaSeq Xp Flow Cell Dock. Each lane included ~1% of a PhiX control library for run quality control.

Genome Assembly

We generated 221 gigabases of sequence data using the ONT PromethION platform (NCBI BioProject PRJNA705675, SRA Experiment SRX11206953). This represents an anticipated 88X coverage of the expected 2.5 Gbp Syrian hamster genome. The raw sequencing reads exhibited an N50 length of 15,730 bp. We used the Flye assembler v2.8.1 [23] to generate an initial *de novo* genome assembly. Given the potential sequence error rate of PromethION reads, it is advisable to use higher quality Illumina short reads mapped to an assembly to correct sequence errors in initial contigs. Consequently, we used Pilon software v. 1.23 [24] with default settings and 30X genome coverage of Illumina data (SRX10928323) generated from a kidney sample (SAMN18096088) obtained from the same individual for this sequence polishing step. Pilon sequence polishing was performed one time prior to the optical mapping analyses.

Optical mapping for scaffold improvement

Ultra-high molecular weight (UHMW) DNA was extracted following manufacturer's guidelines (Bionano Prep SP Tissue and Tumor DNA Isolation protocol) from frozen liver tissues obtained from the same animal used for ONT PromethION sequencing (SAMN18096087). Briefly, a total of 15-20mg of liver tissue was homogenized in cell buffer and digested with Proteinase K. DNA was precipitated with isopropanol and bound with nanobind magnetic disk (Bionano Genomics, USA). Bound UHMW DNA was resuspended in the elution buffer and quantified with Qubit dsDNA assay kits (ThermoFisher Scientific). DNA labeling was performed following manufacturer's protocols (Bionano Prep Direct Label and Stain protocol). Direct Labeling Enzyme 1

(DLE-1) reactions were carried out using 750 ng of purified UHMW DNA. Labeled DNA

was loaded on Saphyr chips for imaging. The fluorescently labeled DNA molecules

were imaged sequentially across nanochannel arrays (Saphyr chip) on a Saphyr

instrument (Bionano Genomics Inc, USA). Effective genome coverage of greater than

100X was achieved for all samples. All samples also met the following QC metrics:

labelling density of ~15/100 kbp; filtered (>15kbp) N50 > 230 kbp; map rate > 70%.

Genome analysis of the resulting data was performed using software solutions provided

by Bionano Genomics Inc. Briefly, automated optical genome mapping specific

pipelines consisting of Bionano Access v1.4.3 and Bionano Solve v. 3.6.1 were used for

data processing (BioNano Access Software User Guide). Hybrid scaffolding was

performed using Bionano's custom software program implementing the following steps:

1) generate *in silico* maps for sequence assembly; 2) align *in silico* sequence maps

against Bionano genome maps to identify and resolve potential conflicts in either data

set; 3) merge the non-conflicting maps into hybrid scaffolds; 4) align sequence maps to

the hybrid scaffolds; and 5) generate AGP and FASTA files for the scaffolds. Pairwise

comparisons of all DNA molecules were made to generate the initial consensus genome

maps (*.cmap). Genome maps were further refined and extended with best matching

molecules. Optical map statistics were generated using Bionano software producing the

Bionano Molecule Quality Report (MQR).

The optical map N50 (including only maps >=150 kbp and minSites >= 9) was 0.2341

Mbp and the average label density (scaffolds >= 150 kbp) was 17.40/100 kbp. This

yielded an effective molecule coverage with optical mapping information of 125.38X.

The optical mapping analysis identified 84 conflicts with the prior Flye/Pilon scaffolds

and these initial scaffolds were broken at those 84 sites. The completed assembly was

submitted to NCBI and is available under accession GCA_017639785.1.

Gene annotation

210 NCBI performed gene annotation using RNA-Seq data from multiple tissues including

211 lung, trachea, brain, olfactory bulb and small intestine that are targets for SARS-CoV-2

212 infection (NCBI BioProject PRJNA675865) [19].

Quality assessment

To assess the quality of our assembly compared to the previous MesAur1.0 we used

Quast v5.0.2 [25] together with MUMmer v3.23 [26]. These tools provided a detailed

comparison between these assemblies. In addition, the Illumina reads from the original

reference (NCBI SRA SRR413408) were mapped to our assembly and the MesAur1.0

reference using BWA v0.7.17 [27]. Quast was used to obtain discordant pair statistics.

We next used the software Benchmarking Universal Single-Copy Orthologs (BUSCO)

v3.0.2 [28] to assess the quality of the genome assembly. BUSCO is based on the

concept that single-copy orthologs should be highly conserved among closely related

species. BUSCO performs gene annotation on an assembly and reports the number of

gene models generated. BUSCO was performed using the OrthoDB v9 (odb9) release

consisting of 6192 genes shared across the superorder Euarchontoglires [29], the

appropriate test for the Syrian hamster.

In addition, FRCbam [30] was used to compute Feature Response Curves (FRCurve)

from the alignment of Illumina reads to the assembled contigs. FRC v1.3.0 was

employed to evaluate both assemblies, using default parameters. BCM_Maur_2 was

further evaluated using paired end mappings of the Illumina reads that had been used

for Pilon polishing (SRX10928323). MesAur1.0 was then similarly evaluated using

paired end mappings of Illumina reads used for the MesAur1.0 assembly (SRR413408).

Results

The initial Flye assembly consisted of 2.38 Gbp of sequence across 6,741 scaffolds with

a scaffold N50 of 10.56 Mbp (**Table 1**). Pilon polishing of the Flye assembly had little

effect on these metrics, but significant improvements were obtained when Bionano

optical mapping results were used to improve scaffolding. As shown in **Table 1**, the

optical mapping step reduced the total number of scaffolds in the final assembly by 395

(5.9%) while increasing the N50 scaffold length by more than 8-fold to 85.18 Mbp. Our

experience in comparing read lengths and total yield per flow-cell indicates that the

optimal target size for fragmented DNA as input into Nanopore libraries and sequencing

is 15-20 kb, which regularly yields 80-90 Gb of sequence data.

Of the 6,192 BUSCO gene models, 88.91% were annotated as complete genes in the

initial Flye assembly (**Table 2**). Pilon polishing of this Flye-alone assembly added

another 295 genes annotated completely and increased this proportion to 93.67% of the

BUSCO gene model dataset. Improvements in assembly scaffolding resulting from the

Bionano optical mapping step together with Pilon error correction decreased the

proportions of fragmented and missing BUSCO gene models in the new assembly to

2.9% and 3.2% respectively, also improvements over the MesAur1.0 assembly. This

advance translates to an additional 305 complete BUSCO genes identified in the new

assembly.

Table 1. Assembly statistics for BCM_Maur_2.0 versus the MesAur1.0 Syrian hamster
assembly

Parameter	MesAur1.0	Flye	Flye + Pilon	Flye + Pilon + Bionano (BCM_Maur_2.0)
Assembly length (bp)	2,504,908,775	2,381,258,546	2,383,228,608	2,457,062,007
Ungapped length (bp)	2,076,159,990	2,381,254,546	2,383,226,373	2,383,228,883
Number of scaffolds	21,483	6,741	6,741	6,346
N50 scaffold length (bp)	12,753,307	10,564,357	10,573,641	85,184,847
Number of contigs	237,699	6,781	6,779	7,057

N50 contig length (bp)	22,512	10,022,145	10,097,207	9,471,653

261 **Table 2.** BUSCO statistics for BCM_Maur_2.0 versus the MesAur1.0 Syrian hamster

262 assembly

	MesAur1.0	Flye	Flye + Pilon	Flye + Pilon + Bionano (BCM_Maur_2.0)
Complete[a]	88.97%	88.91%	93.67%	93.90%
Complete and single-copy	88.47%	88.05%	92.80%	93.12%
Complete and duplicated	0.50%	0.86%	0.87%	0.78%
Fragmented	5.89%	5.99%	3.04%	2.87%
Missing	5.14%	5.10%	3.29%	3.23%

263 [a]6192 gene models were included in this analysis

265 **Assembly Comparisons**

We also performed additional comparisons between the two assemblies. As

background, the karyotype of *M. auratus* is diploid 2n = 44, including 14 pairs of

metacentric chromosomes, 3 pairs of telocentrics and 5 pairs of acrocentrics [31].

Illumina read k-mer analyses were performed to estimate the genome size using SGA

preqc [32] (2.57 Gbp) and Jellyfish [33] (2.90 Gbp). The total length of the

BCM_Maur_2.0 assembly is 2.46 Gbp compared to the previous version's 2.50 Gbp.

Despite having a similar total length, BCM_Maur_2.0 shows an improved continuity with

a scaffold N50 that is 6.7 times greater than MesAur1.0 (**Table 1**); the L50 (i.e. the

number of contigs longer than or equal to the N50 length) of BCM_Maur_2.0 is 22

compared to MesAur1.0's 121. The longest scaffold of BCM_Maur_2.0 (187 Mb) is 2.35

times larger than the longest scaffold from the previous assembly. N50 is calculated in

the context of the assembly size rather than the genome size, so the NG50 statistic was

used to directly compare the different assemblies. NG50 is the same as N50 except that

it reports the length of the contig at which the size-ordered contigs (longest to shortest)

collectively reaches 50% of the known or estimated genome size [34]. **Figure 1**

illustrates the improved cumulative contig sequence length for any given NG50 value

that is generated from the BCM_Maur_2.0 assembly as compared to MesAur1.0, based

on the estimated genome size of 2.57 Gb calculated using SGA-preqc. The

BCM_Maur_2.0 assembly further improves the unresolved regions as measured by

nucleotide ambiguities (i.e. number of N's included in the final contigs). Approximately

17.11% of bases in MesAur1.0 were unresolved. BCM_Maur_2.0 reduces the number

of unresolved bases to 3.00%, with only very small gaps throughout the entire genome.

Figure 2 displays the overall increase in continuity of the BCM_Maur_2.0 assembly with

longer contigs than the MesAur1.0 assembly and fewer short contigs. Finally, we

compared feature response curves for BCM_Maur_2.0 and MesAur1.0 using FRC[Bam]

[30]. FRC^{Bam} shows that our new assembly is substantially more accurate based on the

feature response approach (Supplementary Figure 1).

To establish the correctness of the structure and completeness of BCM_Maur_2.0, we

also leveraged the Illumina short-reads that were published as part of the MesAur1.0

assembly project. When mapping the MesAur1.0 Illumina reads back to the MesAur1.0

reference, only 92.19% reads mapped successfully. When the same Illumina reads

were instead mapped to BCM_Maur_2.0, 97.32% mapped successfully. When

considering only properly paired reads, 75.76% and 87.62% mapped to MesAur1.0 and

BCM_Maur_2.0, respectively.

Alignments between the current and previous Syrian hamster assemblies performed by

NCBI [35] show that BCM_Maur_2.0 covers 98.95% of MesAur1.0 while MesAur1.0

only covers 86.67% of BCM_Maur_2.0. This together with the additional 307 Mbp of

ungapped sequence in BCM_Maur_2.0 indicates that BCM_Maur_2.0 is a more

complete representation of the Syrian hamster genome. The percent identity in the

regions aligned between the two assemblies is 99.76%.

306 **Transcript and Protein Alignments and Annotation**
307 **Comparisons**
308

NCBI annotation of BCM_Maur_2.0 [35] with Syrian hamster transcript and protein data

show this assembly to be of high quality. Transcript alignments of Syrian hamster

RefSeq (n=273), Genbank (n=751), and EST (n=558) data to BCM_Maur_2.0 show

99.44% or more average percent identity and 98.88% or more average percent

coverage. Alignments of these same transcript datasets to MesAur1.0 show 99.13% or

more average percent identity and 93.49% or more average percent coverage.

Alignments of RefSeq transcripts showed a similar average percent indels in the

BCM_Maur_2.0 (0.10%) and MesAur1.0 (0.11%) assemblies. Protein alignments of

Syrian hamster RefSeq (n=261) and Genbank (n=485) data to BCM_Maur_2.0 show

80.95% or more average percent identity and 89.18% or more average percent

coverage. Alignments of these same protein datasets to MesAur1.0 show 80.57% or

more average percent identity and 84.87% or more average percent coverage.

NCBI annotated 21,616 protein coding genes and 10,459 noncoding genes in

BCM_Maur_2.0 compared to 20,495 protein coding genes and 4,168 noncoding genes

in MesAur1.0 [36]. Only 7% of gene annotations are identical between BCM_Maur_2.0

and MesAur1.0, suggesting that a number of previous errors have been corrected,

though some differences are likely to be real differences between the animals used for

the different assemblies. Minor changes between BCM_Maur_2.0 and MesAur1.0 were

made in 46% of gene annotations and major changes were made in 15% of gene

annotations. We further note that, based on NCBI annotation feature counts,

BCM_Maur_2.0 has only 33 RefSeq models that were filled using transcript sequence

to compensate for an assembly gap [35]. This is compared to 5,050 RefSeq models

similarly compensated in MesAur1.0.

Interferon type 1 alpha gene cluster

Given the importance of type I interferon responses during SARS-CoV-2 infection, we

next compared the interferon type I alpha gene cluster in the BCM_Maur_2.0 assembly

relative to this genomic region in the original MesAur1.0 assembly. The MesAur1.0

scaffold NW_004801649.1 includes annotations for four interferon type I alpha loci but

this genomic sequence is riddled with numerous gaps. Of these four candidate loci,

only LOC101824534 appears to contain a complete interferon alpha-12-like coding

sequence with the ability to encode a predicted protein (XP_005074343.1). The

LOC101824794 gene sequence can only encode a 162 amino acid protein due to a 5'

truncation. The remaining pair of candidate genes (LOC101836618 and

LOC101836898) appear to have aberrant transcript models that have fused putative

exons from neighboring loci. In mice and humans, the interferon alpha gene cluster is

flanked by single copy interferon beta 1 (*Ifnb1*) and interferon epsilon (*Ifne*) genes.

Although neither of these genes are present on the MesAur1.0 scaffold

NW_004801649.1, this assembly does contain a *Ifne* gene on a short 2,408 bp contig

that is predicted to code for a protein of 192 amino acids. These observations

emphasize the need for an improved genomic assembly for Syrian hamsters given that

the interferon alpha gene cluster includes more than a dozen tightly linked functional

genes plus multiple pseudogenes in a wide variety of species including mice and

humans.

In the BCM_Maur_2.0 assembly, the interferon type I alpha gene cluster is contained on

the NW_024429197.1 super scaffold that spans nearly 75 Mbp. **Figure 3** illustrates this

genomic region in comparison with the interferon type 1 alpha regions of MesAur1.0

(NW_004801649.1) and the well-characterized C57BL/6J mouse assembly

(NC_000070.7). Fourteen predicted interferon type I alpha genes as well as five

presumptive pseudogenes lie within a span of 196 Kbp of the new Syrian hamster

assembly (**Figure 3** and **Supplemental Table 1**). This genomic organization is quite

comparable to that observed in the mouse genome where there are also fourteen

functional interferon alpha genes and four pseudogenes. This hamster gene cluster is

flanked by *Ifnb1* and *Ifne* genes consistent with expectations from the mouse and other

species. The NCBI annotations characterize twelve of these genes as interferon alpha-

12-like (**Supplemental Table 1**). The remaining pair of functional genes

(LOC101824794 and LOC121144100) are listed as interferon alpha-9-like and they

encode shorter predicted proteins. The increased length of this genomic region in the

mouse assembly is largely due to the presence of the interferon zeta gene family (*Ifnz*,

Gm13271, Gm13272, etc.). This *Ifnz* gene family appears to be absent in Syrian

hamsters since the closest matches to predicted hamster protein sequences are only

28% identical at the amino acid level. The interferon type I alpha gene cluster in the

BCM_Maur_2.0 assembly lies within more than 12 Mbp of contiguous genomic

sequence with the nearest flanking gaps located 2.66 Mbp proximal and 9.07 Mbp distal

to the *Ifne* and *Ifnb1* genes, respectively. The availability of a contiguous hamster

genomic sequence and associated transcriptional regulatory elements for this complex

immune gene region may be helpful for investigators who are interested in unravelling

mechanisms that control interferon expression during infections with SARS-CoV-2 as

well as challenges with other viral pathogens.

377   **Conclusions**

The improved Syrian hamster assembly and annotation described here will facilitate

research into this important animal model for COVID-19. Specifically, reagents for

studying immune responses in hamsters have lagged behind those available for

laboratory mice. BCM_Maur_2.0 will facilitate the identification of cross-reactive

reagents originally developed to study immunity in other species. Additionally, a more

accurate genome assembly will improve the analyses of host responses to infection by

enabling more accurate interpretation of RNA-seq experiments.

Relative to other recent assemblies that use a combination of long-read sequencing and

short-read polishing, this genome assembly and annotation compares very favorably.

The scaffold N50 of >85 Mbp is quite consistent with other long read assemblies. The

contig N50 and total number of scaffolds or contigs are likewise reasonable and

consistent with other similar mammalian reference genomes. The number of protein

coding genes identified is within the expected range, although additional attention will

likely be needed to resolve duplicated, repetitive gene loci, potentially leveraging recent

advances in ultralong read sequencing.

What additional genomic resources would be needed to make hamsters a better model

for COVID-19? Deep long read transcriptome analysis of multiple tissues and ages

would be the best next step, in order to define not just the genes expressed but the

alternative splicing of genes across tissues and developmental stages. Also, long read

RNA-seq of tissues following experimental challenge with SARS-CoV-2 and other

viruses would facilitate improvements in the quality of antiviral gene models.

The availability of higher accuracy sequences should lead to the development of

specific reagents for monitoring immune responses. For example, epitopes that are

shared between hamsters and other rodents can be used to identify monoclonal

antibody reagents for flow cytometry that are predicted to be cross-reactive. Additional

reagent development will be enabled by creating synthetic versions of hamster proteins

that can be used as immunogens to make hamster-specific antibodies.

One surprising motivation for this study is that Syrian hamsters, which were quickly

identified as a high value model for COVID-19, did not have a higher quality reference

genome at the start of the pandemic. While we worked quickly to generate this data and

make it available to the scientific community, better preparedness will be critical for

future unexpected epidemics. To this end, we would encourage investment in continued

refinement and improvement of reference genomes for all of the rodent, bat and

nonhuman primate models that are commonly used to study viruses in order to prevent

this situation from recurring in the future. Such an investment would also yield improved

genomic resources that would provide broad benefit to the entire scientific community.

Availability of Supporting Data and Materials

416
417

The MesAur1.0 genome assembly is available in the NCBI database under BioProject

PRJNA77669 (GenBank accession GCA_000349665.1). The new BCM_Maur_2.0

genome assembly is available in the NCBI data repository under BioProject

PRJNA705675 (GenBank accession GCA_017639785.1). Oxford Nanopore

(SRX11206953) and Illumina (SRX10928323) sequencing data are available through

the NCBI SRA. The Illumina RNA-Seq data from multiple tissues including lung,

trachea, brain, olfactory bulb and small intestine are available under NCBI BioProject [PRJNA675865](#).

Additional Files

Supplementary Table 1. Predicted genes in the Interferon type 1 alpha cluster of the BCM_Maur_2.0 assembly.

Abbreviations

ACE2: angiotensin-converting enzyme 2; BCM: Baylor College of Medicine; bp: base pairs; BUSCO: Benchmarking Universal Single-Copy Orthologs; BWA: Burrows-Wheeler Aligner; COVID-19: coronavirus disease 2019; EST: expressed sequence tag; FFPE: formalin-fixed, paraffin-embedded; Gbp: gigabase pairs; GC: guanine-cytosine; IFN: interferon; kbp: kilobase pairs; Mbp: megabase pairs; MQR: Molecule Quality Report; NCBI: National Center for Biotechnology Information; NEB: New England BioLabs; ng: nanogram; ONT: Oxford Nanopore Technologies; PCR: polymerase chain reaction; RBD: receptor-binding domain; RNA-Seq: RNA-sequencing; SARS-CoV-2: severe acute respiratory syndrome coronavirus 2; STAT2: signal transducer and activator of transcription factor 2; TMPRSS2: transmembrane protease serine 2

Competing interests

The authors declare that they have no competing interests.

Funding

This research was supported by contract HHSN272201600007C awarded to DHO from the National Institute of Allergy and Infectious Diseases of the National Institutes of Health. The content of this publication is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Authors' Contributions

Acknowledgements

We are extremely grateful to Dr. Tadashi Maemura for collecting the Syrian hamster tissues that were used for the sequence analyses described here. We also thank Dr. Benjamin tenOever for sharing Syrian hamster RNA-Seq datasets generated by his group prior to publication. And we also wish to thank two reviewers for their helpful comments.

Figure 1: Cumulative length and continuity comparison of MesAur1.0 and BCM_Maur_2.0. This summarizes the length of contigs/scaffolds across the assemblies. Given the length of contigs, the NG50 (mid x-axis) summarizes the sequence length of the shortest contig/scaffold at 50% of the total genome length. For genome length, the SGA preqc estimate of 2.57 Gbp was used.

Figure 2: Contig length and count comparison between BCM_Maur_2.0 and MesAur1.0. Log length of contigs on the X axis and normalized count on the Y axis comparing BCM_Maur_2.0 assembly and the previous assembly. Contigs from BCM_Maur_2.0 are shown red and contigs for MesAur1.0 are shown in gray.

Figure 3: Comparison of interferon type 1 alpha gene cluster between MesAur1.0, BCM_Maur_2.0 and GCRm39 mouse genome assembly. The genomic intervals illustrated here are defined by the flanking interferon beta 1 and interferon epsilon genes except for MesAur1.0 which does not include an interferon epsilon or beta 1 gene in a continuous sequence with interferon type 1 alpha genes. White space within each scaffold represents gaps in the MesAur1.0 assembly. Accession numbers for each genomic sequence are indicated on the right with genomic coordinates for the extracted intervals shown below their respective accession numbers. Predicted interferon type 1 alpha genes are highlighted in blue while putative pseudogenes are depicted with open symbols and labelled below each assembly.

References

1. LaRocca CJ, Han J, Gavrikova T, Armstrong L, Oliveira AR, Shanley R, et al.. Oncolytic adenovirus expressing interferon alpha in a syngeneic Syrian hamster model for the treatment of pancreatic cancer. *Surgery*. 2015; doi: 10.1016/j.surg.2015.01.006.

2. Pal S, Haldar C, Verma R. Photoperiodic modulation of ovarian metabolic, survival, proliferation and gap junction markers in adult golden hamster, Mesocricetus auratus. *Comp Biochem Physiol A Mol Integr Physiol*. 2021; doi: 10.1016/j.cbpa.2021.111083.

3. McCann KE, Sinkiewicz DM, Norvelle A, Huhman KL. De novo assembly, annotation, and characterization of the whole brain transcriptome of male and female Syrian hamsters. *Sci Rep*. 2017; doi: 10.1038/srep40472.

4. Saini S, Rai AK. Hamster, a close model for visceral leishmaniasis: Opportunities and challenges. *Parasite Immunol*. 2020; doi: 10.1111/pim.12768.

5. Chan JF-W, Zhang AJ, Yuan S, Poon VK-M, Chan CC-S, Lee AC-Y, et al.. Simulation of the Clinical and Pathological Manifestations of Coronavirus Disease 2019 (COVID-19) in a Golden Syrian Hamster Model: Implications for Disease Pathogenesis and Transmissibility. *Clin Infect Dis*. 2020; doi: 10.1093/cid/ciaa325.

6. Prescott J, Falzarano D, Feldmann H. Natural Immunity to Ebola Virus in the Syrian Hamster Requires Antibody Responses. *J Infect Dis*. 2015; doi: 10.1093/infdis/jiv203.

7. Huo J, Mikolajek H, Le Bas A, Clark JJ, Sharma P, Kipar A, et al.. A potent SARS-CoV-2 neutralising nanobody shows therapeutic efficacy in the Syrian golden hamster model of COVID-19. *Nat Commun*. 2021; doi: 10.1038/s41467-021-25480-z.

8. Mohandas S, Yadav PD, Shete A, Nyayanit D, Sapkal G, Lole K, et al.. SARS-CoV-2 Delta Variant Pathogenesis and Host Response in Syrian Hamsters. *Viruses*. 2021; doi: 10.3390/v13091773.

9. Mifsud EJ, Tai CM, Hurt AC. Animal models used to assess influenza antivirals. *Expert Opin Drug Discov*. 2018; doi: 10.1080/17460441.2018.1540586.

10. Gao M, Zhang B, Liu J, Guo X, Li H, Wang T, et al.. Generation of transgenic golden Syrian hamsters. *Cell Res*. 2014; doi: 10.1038/cr.2014.2.

11. Imai M, Iwatsuki-Horimoto K, Hatta M, Loeber S, Halfmann PJ, Nakajima N, et al.. Syrian hamsters as a small animal model for SARS-CoV-2 infection and countermeasure development. *Proc Natl Acad Sci U S A*. 2020; doi: 10.1073/pnas.2009799117.

12. Rockx B, Kuiken T, Herfst S, Bestebroer T, Lamers MM, Oude Munnink BB, et al.. Comparative pathogenesis of COVID-19, MERS, and SARS in a nonhuman primate model. *Science*. 2020; doi: 10.1126/science.abb7314.

13. Rogers TF, Zhao F, Huang D, Beutler N, Burns A, He W-T, et al.. Isolation of potent SARS-CoV-2 neutralizing antibodies and protection from disease in a small animal model. *Science*. 2020; doi: 10.1126/science.abc7520.

14. Shi J, Wen Z, Zhong G, Yang H, Wang C, Huang B, et al.. Susceptibility of ferrets, cats,
dogs, and other domesticated animals to SARS-coronavirus 2. *Science*. 2020; doi:
10.1126/science.abb7015.

15. Muñoz-Fontela C, Dowling WE, Funnell SGP, Gsell P-S, Riveros-Balta AX, Albrecht RA, et
al.. Animal models for COVID-19. *Nature*. 2020; doi: 10.1038/s41586-020-2787-6.

16. Montagutelli X, Prot M, Levillayer L, Salazar EB, Jouvion G, Conquet L, et al.. The B1.351
and P.1 variants extend SARS-CoV-2 host range to mice. bioRxiv.

17. Port JR, Adney DR, Schwarz B, Schulz JE, Sturdevant DE, Smith BJ, et al.. Western diet
increases COVID-19 disease severity in the Syrian hamster. *bioRxiv*. 2021; doi:
10.1101/2021.06.17.448814.

18. Boudewijns R, Thibaut HJ, Kaptein SJF, Li R, Vergote V, Seldeslachts L, et al.. STAT2
signaling restricts viral dissemination but drives severe pneumonia in SARS-CoV-2 infected
hamsters. *Nat Commun*. 2020; doi: 10.1038/s41467-020-19684-y.

19. Hoagland DA, Møller R, Uhl SA, Oishi K, Frere J, Golynker I, et al.. Leveraging the antiviral
type I interferon system as a first line of defense against SARS-CoV-2 pathogenicity. *Immunity*.
2021; doi: 10.1016/j.immuni.2021.01.017.

20. Brooke GN, Prischi F. Structural and functional modelling of SARS-CoV-2 entry in animal
models. *Sci Rep*. 2020; doi: 10.1038/s41598-020-72528-z.

21. Hoffmann M, Kleine-Weber H, Schroeder S, Krüger N, Herrler T, Erichsen S, et al.. SARS-
CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven
Protease Inhibitor. *Cell*. 2020; doi: 10.1016/j.cell.2020.02.052.

22. Rizvi ZA, Dalal R, Sadhu S, Kumar Y, Shrivastava T, Gupta SK, et al.. Immunological and
cardio-vascular pathologies associated with SARS-CoV-2 infection in golden syrian hamster.
Cold Spring Harbor Laboratory.

23. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. Assembly of long, error-prone reads using
repeat graphs. *Nat Biotechnol*. 2019; doi: 10.1038/s41587-019-0072-8.

24. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, et al.. Pilon: an
integrated tool for comprehensive microbial variant detection and genome assembly
improvement. *PLoS One*. 2014; doi: 10.1371/journal.pone.0112963.

25. Gurevich A, Saveliev V, Vyahhi N, Tesler G. QUAST: quality assessment tool for genome
assemblies. *Bioinformatics*. 2013; doi: 10.1093/bioinformatics/btt086.

26. Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, et al.. Versatile and
open software for comparing large genomes. *Genome Biol*. 2004; doi: 10.1186/gb-2004-5-2-r12.

27. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv [q-bio.GN].

28. Seppey M, Manni M, Zdobnov EM. BUSCO: Assessing Genome Assembly and Annotation
Completeness. *Methods Mol Biol*. 2019; doi: 10.1007/978-1-4939-9173-0_14.

29. : BUSCO. https://busco-archive.ezlab.org/v3/datasets/euarchontoglires_odb9.tar.gz

Accessed 2021 Jun 4.

30. Vezzi F, Narzisi G, Mishra B. Reevaluating assembly evaluations with feature response
curves: GAGE and assemblathons. *PLoS One.* 2012; doi: 10.1371/journal.pone.0052210.

31. Lehman JM, Macpherson I, Moorhead PS. KARYOTYPE OF THE SYRIAN HAMSTER. *J
Natl Cancer Inst.* 31:639–501963;

32. Simpson JT. Exploring genome characteristics and sequence quality without a reference.
Bioinformatics. 2014; doi: 10.1093/bioinformatics/btu023.

33. Marçais G, Kingsford C. A fast, lock-free approach for efficient parallel counting of
occurrences of k-mers. *Bioinformatics.* 2011; doi: 10.1093/bioinformatics/btr011.

34. Alhakami H, Mirebrahim H, Lonardi S. A comparative evaluation of genome assembly
reconciliation tools. *Genome Biol.* 2017; doi: 10.1186/s13059-017-1213-3.

35. : Mesocricetus auratus Annotation Report.
https://www.ncbi.nlm.nih.gov/genome/annotation_euk/Mesocricetus_auratus/103/ Accessed
2021 Jun 4.

36. : Mesocricetus auratus Annotation Report.
https://www.ncbi.nlm.nih.gov/genome/annotation_euk/Mesocricetus_auratus/102/ Accessed
2021 Jun 4.