Meta-analysis of tree diversity effects on the abundance, diversity and
activity of herbivores' enemies

Stemmelen[1], A., Jactel[1], H., Brockerhoff[2], E.G., Castagneyrol[1], B.

Affiliations

1 – INRAE, University of Bordeaux, UMR Biogeco, F-33612 Cestas, France

2 – Swiss Federal Research Institute WSL, Zürcherstrasse 111, CH-8903 Birmensdorf, Switzerland

Abstract

• The natural enemies hypothesis predicts that the abundance and diversity of
antagonists such as predators and parasitoids of herbivores increases with the
diversity of plants, which can lead to more effective top-down control of insect
herbivores. However, although the hypothesis has received large support in
agricultural systems, fewer studies have been conducted in forest ecosystems and a
comprehensive synthesis of previous research is still lacking.

• We conducted a meta-analysis of 65 publications comparing the diversity, abundance
or activity of various groups of natural enemies (including birds, bats, spiders and
insect parasitoids) in pure vs. mixed forest stands. We tested the effects of forest
biome, natural enemy taxon and type of study (managed vs experimental forest).

• We found a significant positive effect of forest tree diversity on natural enemy
abundance and diversity but not on their activity. The effect of tree diversity on natural
enemies was stronger towards lower latitudes but was not contingent on the natural
enemy taxon.

• Overall, our study contributes substantially toward a better understanding of the
"natural enemies hypothesis" in forest systems and provides new insights about the
mechanisms involved. Furthermore, we outline potential avenues for strengthening
forest resistance to the growing threat of herbivorous insects.

Keywords: Associational resistance ; Herbivory ; Meta-analysis ; Natural enemies ; Top-down
control ; Tree diversity ; Pest control

Introduction

Forest ecosystems offer a wide variety of ecosystem services that benefit the economy and human wellbeing (Brockerhoff et al., 2017; de Groot et al., 2002). However, in the context of global change, forests are facing increasing threats including herbivory by native and invasive pest species (Grünig et al., 2020; Seidl et al., 2014). It is generally accepted that mixed forests are less vulnerable to insect herbivory than single-species forests (Jactel et al., 2021), a phenomenon commonly referred to as 'associational resistance' (Barbosa et al., 2009; Jactel et al., 2017; Letourneau et al., 2011) – though the opposite, 'associational susceptibility', may also occur (Schuldt et al., 2014; White & Whitham, 2000). Associational resistance is usually explained by two main mechanisms, driven by the presence of heterospecific neighboring trees: (i) a bottom-up effect (*sensu* Hunter & Price, 1992) on focal host accessibility, with insect herbivores being more likely to find and remain on hosts that are growing in dense single-species stands (the resource concentration hypothesis) and (ii) a top-down effect, corresponding to the control of herbivores population by their natural enemies, supposed to be more abundant in more diverse habitats *(the natural enemies hypothesis)*. The "natural enemies hypothesis" was first presented by Root (1973) and postulates that more diverse habitats provide predators and parasitoids with more abundant and predictable resources (*i.e. prey, breeding and nesting sites, refuge and shelter, etc.*) across spatial and temporal scales, increasing natural enemy abundance and diversity and thus leading to more efficient top-down control of herbivorous pest populations. This hypothesis is well-supported in agricultural systems where it has been shown that crops growing in mixture generally support higher level of predation of herbivores and suffer less loss from herbivory than crops growing in monoculture (Russell, 1989; Letourneau, 1987, Bianchi et al., 2006; Wan et al., 2020). On the other hand, the applicability of the natural enemies hypothesis in forests is still a matter of discussion (Staab & Schuldt, 2020), as studies are reporting positive (e.g. Esquivel-Gómez et al., 2017; Novais et al., 2017), neutral (e.g. Klapwijk & Björkman, 2018; Schuldt et al., 2008) or negative (e.g. Nixon & Roland, 2012) effects of tree diversity on insect parasitism or predation.

Inconsistencies in studies having tested the natural enemies in forest ecosystems may arise from

multiple sources. Firstly, the natural enemies hypothesis, as it was initially presented, proposed that

an increase in abundance and diversity of natural enemies would lead to greater top-down control of

herbivores populations. However, relationships between abundance, diversity and actual biological

control that natural enemies exert on insect herbivores in forests have not been examined thoroughly.

Furthermore, different studies assessing the relevance of the natural enemies hypothesis have used

different proxies for top-down control by natural enemies which could lead to different results due to

methodological issues. For example, Esquivel-Gómez et al. (2017) found that the *richness* of spiders

was significantly higher in mixed stands than in pure stands in a forest of south Mexico. On the

contrary, Schuldt et al. (2008) found no effect of tree species mixing on the *abundance* of spiders in a

deciduous forest of central Germany. Both studies assessed the response of natural enemies to tree

diversity, but used different metrics (i.e. richness vs. abundance) which may have led to the difference

in the results observed. In recent years, studies using direct measures of predation and parasitism

while testing for an effect of tree diversity have increased (Abdala-Roberts et al., 2016; Bellone et al.,

2020) but they cannot be directly compared to those measuring only predator abundance.

Methodological issues can also arise from the type of study used to test the effect of tree diversity on

natural enemies. Tree diversity experiments, where the number of tree species per area is manipulated

(e.g in TreeDivNet, Paquette et al., 2018), are used increasingly because they allow for better, more

controlled tests of the mechanisms behind natural enemy responses to tree diversity. However, these

mainly recent experiments often include mixtures of rather low levels of tree diversity and they

typically lack the features of mature forest such as deadwood and diverse microhabitats (Staab &

Schuldt, 2020). Therefore, it was hypothesized that these different sources of heterogeneity in studies

of natural enemies in mixed forests may make it difficult for clear patterns to emerge (Staab & Schuldt,

2020).

Second, experimental tests of the natural enemies hypothesis in specific locations fail to account for

the effect of biogeographical gradients such as latitude, which is known to influence the biodiversity

of natural enemies and the strength of various biological interactions (Pennings & Silliman, 2005; Roslin

et al., 2017). For example it has been shown that ectothermic arthropod predators such as spiders,

ants or carabids are more active under warm climate (Becerra, 2015; Björkman et al., 2011).

Third, it has been found previously that the effect of tree diversity on natural enemies differed

between taxa. An experiment conducted in Germany on 150 forest plots showed that the species

richness of birds and bats was not related to tree diversity, while that of forest-dwelling predatory

arthropods was (Penone et al., 2019). Therefore, discrepancies could exist between studies conducted

in different forest biomes or focusing on different natural enemy taxa.

To assess the relevance of the natural enemies hypothesis in forests, we conducted a meta-analysis of

65 published studies from 1993 to 2020 that involved more than 200 case studies. We addressed

methodological issues arising from the broad definition of the natural enemies hypothesis by testing

whether the effect of tree diversity on natural enemies varied with the metrics used *(abundance,*

richness and activity) and the type of study *(tree diversity experiment vs mature forest)*. We also

evaluated the importance of the type of natural enemy *(birds, bats, arthropod predators vs.*

parasitoids) as well as the forest biome *(tropical, temperate or boreal forest)* to account for the

importance of ecological features and biogeographical gradients. Specifically, and albeit the potential

effect of the metrics used to assess natural enemy responses to tree diversity, we predict that,

following the work of Roslin et al. (2017), the strength of trophic interactions would be higher towards

the tropics, which would result in stronger effect of tree diversity on natural enemies at lower

latitudes. Additionally, we expect that more generalist natural enemies would benefit more from the

diversity of forests, being able to exploit complementary host or prey resources.

Material & Methods

Data collection

We conducted an extensive literature search focusing on the effect of tree diversity on natural enemies
(hereafter referred to as "NE") of herbivores. We searched the Web of Science and Scopus databases
using the following combination of relevant keywords: [forest OR woodland OR "planted forest"] AND
[tree NEAR (diversity OR richness OR composition OR gradients) OR (monoculture OR polyculture OR
(pure OR mix*) NEAR stand)] AND [enem* OR predat* OR prey OR parasit* OR top-down OR control*
OR bird* OR bat* OR chirop* OR spider* OR ant* OR carab* OR hoverfly OR syrph* OR ladybird OR
ichneum* OR hymeno* OR dipter* OR "enemies hypothesis"] AND [diversity OR richness OR
abundance OR density OR activity]. We limited the search to papers published in English, between
1950 and 2021. We then screened the title, abstract and full text when appropriate of each previously
returned article and included them in the meta-analysis if they met the following criteria:

1. Studies focused on the activity (e.g., predation rate, predation attempts on prey models, percentage
of parasitism), abundance or species richness or diversity of NE as a response variable (hereafter "NE
response to tree diversity") in pure vs mixed forest stands or along a gradient of tree species richness
or diversity.

2. Studies had to report the mean of the response variable, any measure of variability around means
(standard deviation, standard error or 95% confident interval) and the sample size used in the study,
or a correlation coefficient between one of the response variables and a gradient of tree species
richness or diversity. Wherever necessary, we retrieved that information by digitizing the figures using
WebPlotDigitizer version 4.2 (Rohatgi, 2019).

Our literature search yielded a total of 4666 candidate primary studies (last accessed in February 2021),
of which we discarded 87% (4083 studies) that did not satisfy our inclusion criteria. We then completed
our dataset by checking for any relevant articles present in the cited references of each article retained

after the first initial search or in recent reviews. A case study corresponded to any type of NE response

to tree diversity, such that there could be more than one case study per primary study. We gave each

primary study and case study a single ID. We finally gathered a dataset including 245 case studies from

65 primary studies that were published between 1993 and 2020 across 26 scientific journals (Fig. 1).

Calculating effect sizes

For each case study, we calculated Hedges' d effect size (Hedges & Olkin, 1985) as the standardized

differences between NE responses to tree diversity in mixed vs. pure stands. Hedges' d was used

because it can account for small sample sizes and uneven variance between control and treatment

groups. A positive d value corresponds to a positive effect of tree diversity on NE, meaning that NE

were more abundant, diverse or active in mixed stands as compared to pure stands. Negative d value

indicate the opposite. When authors reported correlations between tree diversity and NE responses,

we extracted Pearson's coefficient correlation and sample size from which we calculated Fisher's z-

score. Positive value of Fisher's z mean that there is a positive correlation between the NE response

variable and tree diversity. We then converted Fisher's z to Hedges' d (Koricheva et al., 2013). All effect

sizes were calculated with the "metafor" package, using R version 3.6.1 (R Core Team, 2016;

Viechtbauer, 2010).

Moderators

We modeled heterogeneity among effect sizes using three moderators (covariates).

<u>Natural enemy response type</u>

We first explored whether the type of response of NE to tree diversity was consistent across several

methodological approaches. We categorized NE response as 'abundance', 'diversity' and 'activity'.

Abundance was assessed using visual identification, trapping or other devices such as ultrasound

detectors and included relative and absolute abundance of NE as well as mean number of NE per tree

or per trapping device. NE diversity included species richness or Shannon index values. NE activity

referred to proportion of real or model hosts or prey that were missing, killed, or had evidence of

parasitism or predation attempts.

<u>Forest biome</u>

For each primary article, we extracted the geographic coordinates of forest stands, from which we

classified each system as 'boreal', 'temperate' or 'tropical' forest ecosystem.

<u>Taxa of NE</u>

We distinguished between three categories of NE: Birds and bats (predatory vertebrates), arthropods

(generalist predatory invertebrates such as spiders, ants, carabids) and insect parasitoids (specialist

predatory invertebrates). When such a classification could not be made from data reported by the

authors of primary papers, we categorized NE as 'unspecified'. When multiple and undiscerned NE taxa

were involved, NE were categorized as 'Multi-taxon'.

<u>Type of study</u>

For each study, we classified the forest in which the study was conducted as either "experimental

forest", where the tree diversity gradient was experimentally manipulated (e.g. TreeDivNet), or

"natural forest" for other cases.

Statistical analysis

We first calculated a grand mean effect size (Gurevitch & Hedges, 1999) across all studies to assess the

effect of tree diversity on NE. This effect was considered significant if the 95% confidence interval

around the grand mean effect size did not include zero. Following Cohen's guidelines (Cohen, 1988),

an effect size of 0.2 or smaller is considered a 'small' effect size, 0.5 represents a 'medium' effect size

and 0.8 or higher a 'large' effect size. We estimated consistency among studies by calculating between-

study heterogeneity (τ^2 and associated Q statistics) as well as I^2, the standardized estimate of total

heterogeneity ranging from 0 to 1 (Koricheva et al., 2013; Nakagawa et al., 2017). As we detected a

large amount of residual heterogeneity in the grand mean effect size, we accounted for this using three

moderators: NE response type, NE taxa and the forest biome.

We first addressed methodological issues by considering NE response type as a moderator, in

interaction with NE taxa or forest biome. The interaction terms were not significant (NE response

type × Biome: k = 247, Q_M = 0.50; P = 0.97; NE response type × NE taxa: k = 247, Q_M = 3.14, P = 0.53).

We therefore tested each moderator separately. We reported the results of the omnibus test and

interpreted the model coefficients and confidence intervals of each moderator level separately. We

considered that model coefficient parameter estimates were significantly different from zero if the

95% CI around each effect size did not overlap zero.

We ran each model using the *rma.mv* function of the "metafor" package and accounted for potential

non-independence between case study by including Case ID nested within Study ID as random factors.

The use of multiple comparisons to the same control was controlled by using a variance-covariance

matrix among effect sizes.

Preliminary analyses testing for an effect of study type (experimental vs natural forest) showed that

this factor did not significantly affect NE responses to tree diversity (Q_M = 0.005, P = 0.94) and

therefore was not included in the following results section.

Finally, we used several complementary approaches (funnel plot, leave-one out meta-analysis,

Rosenberg fail-safe number and cumulative meta-analysis) to evaluate the robustness and sensitivity

of our results to several sources of bias (see supplementary material 2)

193   Results

Of the 245 cases studies we analyzed, 159 were associated with arthropods (56 of which were

associated with parasitoids), 56 with birds, 16 with bats and 14 with mixed or unidentified NE. NE

abundance and richness were assessed 112 and 85 times respectively, while only 48 cases studies

assessed NE activity. Studies were mostly conducted in temperate forest (n = 30), followed by tropical

(n = 25) and boreal forests (n = 10).

The grand mean effect size (± 95% CI) was significantly positive (0.41 ± [0.17, 0.65], sample size: k =

245, indicating that, overall, NE responded moderately and positively to tree diversity in forest (Fig.

2A. However, there was a large and significant amount of total heterogeneity (τ^2 = 1.76, Q_T = 2899.32,

P < 0.001), mainly attributed to between-study heterogeneity (I^2 = 0.91).

We found a significant effect of the metric (i.e. abundance, diversity or activity) used to characterize

NE response to tree diversity (Q_M = 8.34, P = 0.015, k = 247). Both NE abundance (0.46 ± [0.14, 0.79])

and diversity (0.68 ± [0.36, 1.01]) were significantly higher in mixed than in pure forests, while NE

activity did not significantly influence NE response to tree diversity (-0.16 ± [-0.61, 0.30]) (Fig. 2B).

We found a significant effect of the forest biome in which studies took place (Q_M = 19.64, P = 0.04, k =

247). Mean effect size was large, positive and significantly different from zero in tropical forests (1.43

± [0.72, 2.14]) but non-significant in temperate forests (0.08 ± [-0.09, 0.95]) and in boreal forests (-0.16

± [-0.84, 0.50]) (Fig. 2C).

Model coefficient parameter estimates were positive for birds/bats (0.77 ± [0.05, 1.49]) and

arthropods (excluding parasitoids) (0.68 ± [0.10, 1.25]), almost null for parasitoids (0.09 ± [-0.78,

0.97]) and negative for "mixed" taxa (-0.56 ± [-1.76, 0.64]). However, the identity of the NE taxon did

not influence NE response to tree diversity in forest (Q_M = 2.11, P = 0.54, k = 247).

Finally, we found no significant two or three-way interaction between NE response type, forest biome

and NE taxa.

Discussion

Our meta-analysis provides the first quantitative and up-to-date assessment of the effect of tree

species diversity on natural enemies of insect herbivores in forest ecosystems. Overall, NE abundance

and diversity were higher in mixed than in pure forests, but not their activity, thus providing positive

although partial support to the natural enemies' hypothesis in forest. This also suggest that increasing

forest tree diversity may not automatically lead to stronger top-down control of herbivores by natural

enemies and this is corroborated by the large amount of between-study heterogeneity found in our

quantitative meta-analysis. Those findings are also consistent with recent studies assessing the effect

of tree diversity on natural enemies in forest (reviewed by Staab & Schuldt, 2020) and indicates that

this hypothesis may not apply universally in forests. However, we were able to identify some key

factors explaining the variation in response of NE to tree diversity in forest.

Consistent with the predictions of the natural enemies hypothesis, we confirm that the abundance and

diversity of NE are higher in mixed species forests. Two main non-exclusive mechanisms may explain

this pattern. First, diverse forests are expected to host more stable populations of herbivores

throughout the year (Lawton & Strong, 1981; Siemann et al., 1998; Vehviläinen et al., 2007) and are

thus more likely to provide natural enemies with abundant and various prey and hosts, allowing them

to maintain stable populations. In addition, more diverse habitats can offers a better supply of

complementary food such as pollen or nectar which can improve the fitness of natural enemies and

lead to higher abundance (Cappuccino et al., 1999; Russell, 1989) as well as more microhabitats for

nesting or resting (Asbeck et al., 2019; Batáry et al., 2014), and shelter against adverse conditions. In

turn, tree diversity had no overall effect on NE activity. One possible ecological explanation could be

that intraguild competition, which arises when natural enemies filling the same ecological niche co-

occur and share the same prey population (Finke & Denno, 2003; Rosenheim, 1998), or when intraguild

predation (e.g. super predators or hyperparasitoids) weakens top-down control of host or prey, even

though the abundance or richness of natural enemies increases in mixed forests. For example, Sanders

et al. (2011) found that increasing spider richness did not influence prey control and that the overall

outcome strongly depends on the interference between the predators species involved. Another

explanation could be that, just as host trees are more difficult to locate and colonize by herbivorous

insects in mixed forests because they are physically or chemically hidden by their heterospecific

neighbours (Jactel et al. 2021), prey may be more difficult to find by their predators in more complex

forests (Nell et al., 2018; Tarbox et al., 2018). Lastly, it is technically much more difficult to properly

estimate predation or parasitism of forest insect herbivores than to estimate NE abundance. We could

retrieve only 15 papers on NE activity and most of them were based on the use of sentinel caterpillars

(38% of all methods used to assess NE activity). There is therefore a need for more studies assessing

concomitantly natural enemy abundance/richness and activity in order to properly understand the link

between NE abundance and richness and the actual biological control they exert on herbivores.

Natural enemy responses to tree diversity were stronger towards lower latitudes. In a recent review,

Staab & Schuldt (2020) already pointed out that latitude and climate could be an important factor

shaping NE response to tree diversity in forest and, because tree diversity is much higher in tropical

than in temperate and boreal forests, it could have stronger effects on natural enemies. Additionally,

previous work highlighted an increase in natural enemy activity or abundance with decreasing latitude

(Roslin et al., 2017; Zvereva et al., 2020). Many natural enemies of herbivores are ectothermic

arthropods such as spiders, wasps, carabids or ants, and temperature is known to be a key abiotic

factors affecting their metabolism and activity (Deutsch et al., 2008; Zvereva & Kozlov, 2006). As such,

higher activity of natural enemies, associated with higher overall tree species diversity and stronger

trophic interactions for all NE taxa in tropical forests (Gaston, 2000; Hargreaves et al., 2019; Hillebrand,

2004), may explain why the natural enemies responses to tree diversity was stronger in tropical forests

than in temperate or boreal forests.

Finally, we found no significant difference of natural enemy responses to tree diversity between NE

taxa. Several studies suggest that natural enemy responses to tree diversity are species-specific

(Ampoorter et al., 2020; Riihimäki et al., 2005; Staab & Schuldt, 2020; Wan et al., 2020), and some

species may not respond to tree diversity, even though the overall effect is positive. For example,

Kaitaniemi et al. (2007) found that abundance of predatory ants was higher in mixed forests stands

and lead to higher predation of the European pine sawfly, but the abundance of other predators such

as spiders and predatory heteropterans did not change between mixed and pure stands. Additionally,

the natural enemies hypothesis predicts that generalist natural enemies may benefit more from an

increase in tree diversity than specialists, as generalists would be able to make better use of the greater

number of alternative prey and host species found in mixed habitats. For example, Legault et al. (2018)

found that tree diversity had a positive effect on the generalist parasitoid of the spruce budworm

Apanteles fumiferanae but did not affect parasitism rate by the specialist parasitoid *Glypta*

fumiferanae. Consequentially, and given the fact that NE responses to tree diversity can be closely

linked to NE identity, it is not surprising that no overall pattern have been found. Finally, it is important

to note that studies using multi-taxon approaches to test the responses of different natural enemies

to tree diversity under standardized methods were rare (only 6 studies among the 65 included in the

meta-analysis). The availability of more such multi-taxon studies would help to unfold global patterns

that may otherwise remain unexplored.

283 Conclusion and future work

Our results indicate that natural enemy abundance and richness respond positively to an increase in

tree diversity in forests, providing at least partial support for the natural enemies hypothesis in forest

ecosystems. However, considering that tree diversity does not seem to influence natural enemy

activity consistently, it is not yet clear how this effect of tree diversity will affect the ability of natural

enemies to exert biological control of herbivore populations. While mixed forests might sustain more

abundant and diverse communities of natural enemies, more attention is needed to further

understand how this could lead to stronger biotic interactions with their hosts or prey. We therefore

recommend that studies assessing NE abundance or richness as a proxy of biological control includes

accurate measures of NE activity as well, in order to reach a better understanding of the mechanisms

involved in the natural enemies hypothesis in forests.

Finally, we suggest that more attention is given to studies involving multiple taxa of NE and, ideally

that more standardized methods are applied to different species of NE. Otherwise predicting global

responses of natural enemies' communities to tree diversity will remain challenging.

Declaration of Competing Interests

None.

Data accessibility

Acknowledgments

This study was conducted in the framework of the HOMED project, which received funding from the

European Union's Horizon 2020 research and innovation program under grant 771271. We thank

Michael Staab for his friendly review of an early draft of this paper and for his insightful comments.

References

Ampoorter, E., Barbaro, L., Jactel, H., Baeten, L., Boberg, J., Carnol, M., Castagneyrol, B., Charbonnier,

Y., Dawud, S. M., Deconchat, M., Smedt, P. D., Wandeler, H. D., Guyot, V., Hättenschwiler, S., Joly, F.-

X., Koricheva, J., Milligan, H., Muys, B., Nguyen, D. Allan, E. (2020). Tree diversity is key for promoting

the diversity and abundance of forest-associated taxa in Europe. *Oikos*, *129*(2), 133–146.

https://doi.org/10.1111/oik.06290

Asbeck, T., Pyttel, P., Frey, J., & Bauhus, J. (2019). Predicting abundance and diversity of tree-related

microhabitats in Central European montane forests from common forest attributes. *Forest Ecology

and Management*, *432*, 400–408. https://doi.org/10.1016/j.foreco.2018.09.043

Barbosa, P., Hines, J., Kaplan, I., Martinson, H., Szczepaniec, A., & Szendrei, Z. (2009). Associational

Resistance and Associational Susceptibility: Having Right or Wrong Neighbors. *Annual Review of

Ecology, Evolution, and Systematics*, *40*(1), 1–20.

https://doi.org/10.1146/annurev.ecolsys.110308.120242

Batáry, P., Fronczek, S., Normann, C., Scherber, C., & Tscharntke, T. (2014). How do edge effect and

tree species diversity change bird diversity and avian nest survival in Germany's largest deciduous

forest? *Forest Ecology and Management*, *319*, 44–50. https://doi.org/10.1016/j.foreco.2014.02.004

Becerra, J. X. (2015). On the factors that promote the diversity of herbivorous insects and plants in

tropical forests. *Proceedings of the National Academy of Sciences*, *112*(19), 6098–6103.

https://doi.org/10.1073/pnas.1418643112

Bellone, D., Björkman, C., & Klapwijk, M. J. (2020). Top-down pressure by generalist and specialist

natural enemies in relation to habitat heterogeneity and resource availability. *Basic and Applied*

Ecology, *43*, 16–26. https://doi.org/10.1016/j.baae.2019.10.005

Bianchi, F. J. J. A., Booij, C. J. H., & Tscharntke, T. (2006). Sustainable pest regulation in agricultural

landscapes: A review on landscape composition, biodiversity and natural pest control. *Proceedings.*

Biological Sciences, *273*(1595), 1715–1727. https://doi.org/10.1098/rspb.2006.3530

Björkman, C., Berggren, Å., & Bylund, H. (2011). Causes behind insect folivory patterns in latitudinal

gradients. *Journal of Ecology*, *99*(2), 367–369. https://doi.org/10.1111/j.1365-2745.2010.01707.x

Cappuccino, N., Houle, M.-J., & Stein, J. (1999). The influence of understory nectar resources on

parasitism of the spruce budworm Choristoneura fumiferana in the field. *Agricultural and Forest*

Entomology, *1*(1), 33–36. https://doi.org/10.1046/j.1461-9563.1999.00009.x

Cohen, J. (1988). *Statistical Power Analysis for the Behavioral Sciences*. Routledge.

https://doi.org/10.4324/9780203771587

Cooper, H., & Hedges, L. V. (Eds.). (1994). *The handbook of research synthesis* (pp. xvi, 573). Russell

Sage Foundation.

Daily, C. G., Alexander, S., Ehrlich, R. P., Goulder, L., Lubchenco, J., Matson, A. P., Mooney, A. H.,

Postel, S., Schneider, H. S., Tilman, D., & Woodwell, M. G. (1997). Ecosystem Services: Benefits

Suppliedto Human Societies by Natural Ecosystems. *Issues in Ecology*, *1*(2), 1–18.

de Groot, R. S., Wilson, M. A., & Boumans, R. M. J. (2002). A typology for the classification,

description and valuation of ecosystem functions, goods and services. *Ecological Economics*, *41*(3),

393–408. https://doi.org/10.1016/S0921-8009(02)00089-7

Deutsch, C. A., Tewksbury, J. J., Huey, R. B., Sheldon, K. S., Ghalambor, C. K., Haak, D. C., & Martin, P.

R. (2008). Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of the*

National Academy of Sciences, *105*(18), 6668–6672. https://doi.org/10.1073/pnas.0709472105

Esquivel-Gómez, L., Abdala-Roberts, L., Pinkus-Rendón, M., & Parra-Tabla, V. (2017). Effects of tree

species diversity on a community of weaver spiders in a tropical forest plantation. *Biotropica*, *49*(1),

63–70. https://doi.org/10.1111/btp.12352

Finke, D. L., & Denno, R. F. (2003). Intra-guild predation relaxes natural enemy impacts on herbivore

populations. *Ecological Entomology*, *28*(1), 67–73. https://doi.org/10.1046/j.1365-2311.2003.00475.x

Finke, D. L., & Denno, R. F. (2004). Predator diversity dampens trophic cascades. *Nature*, *429*(6990),

407–410. https://doi.org/10.1038/nature02554

Gaston, K. J. (2000). Global patterns in biodiversity. *Nature*, *405*(6783), 220–227.

https://doi.org/10.1038/35012228

Grünig, M., Mazzi, D., Calanca, P., Karger, D. N., & Pellissier, L. (2020). Crop and forest pest metawebs

shift towards increased linkage and suitability overlap under climate change. *Communications*

Biology, *3*(1), 1–10. https://doi.org/10.1038/s42003-020-0962-9

Gurevitch, J., & Hedges, L. V. (1999). Statistical Issues in Ecological Meta-Analyses. *Ecology*, *80*(4),

1142–1149. https://doi.org/10.1890/0012-9658(1999)080[1142:SIIEMA]2.0.CO;2

Guyot, V., Jactel, H., Imbaud, B., Burnel, L., Castagneyrol, B., Heinz, W., Deconchat, M., & Vialatte, A.

(2019). Tree diversity drives associational resistance to herbivory at both forest edge and interior.

Ecology and Evolution, *9*(16), 9040–9051. https://doi.org/10.1002/ece3.5450

Hargreaves, A. L., Suárez, E., Mehltreter, K., Myers-Smith, I., Vanderplank, S. E., Slinn, H. L., Vargas-

Rodriguez, Y. L., Haeussler, S., David, S., Muñoz, J., Almazán-Núñez, R. C., Loughnan, D., Benning, J.

W., Moeller, D. A., Brodie, J. F., Thomas, H. J. D., & M, P. A. M. (2019). Seed predation increases from

the Arctic to the Equator and from high to low elevations. *Science Advances*, *5*(2), eaau4403.

https://doi.org/10.1126/sciadv.aau4403

Hedges, L., V., & Olkin, I. (1985). *Statistical methods for meta-analysis*.

Hillebrand, H. (2004). On the generality of the latitudinal diversity gradient. *The American Naturalist*,

163(2), 192–211. https://doi.org/10.1086/381004

Hobson, K. A., & Bayne, E. (2000). Breeding Bird Communities in Boreal Forest of Western Canada:

Consequences of "Unmixing" the Mixedwoods. *The Condor*, *102*(4), 759–769.

https://doi.org/10.1093/condor/102.4.759

Jactel, H., Bauhus, J., Boberg, J., Bonal, D., Castagneyrol, B., Gardiner, B., Gonzalez-Olabarria, J. R.,

Koricheva, J., Meurisse, N., & Brockerhoff, E. G. (2017). Tree Diversity Drives Forest Stand Resistance

to Natural Disturbances. *Current Forestry Reports*, *3*(3), 223–243. https://doi.org/10.1007/s40725-

017-0064-1

Jactel, H., Birgersson, G., Andersson, S., & Schlyter, F. (2011). Non-host volatiles mediate
associational resistance to the pine processionary moth. *Oecologia*, *166*(3), 703–711.

Jactel, H., Moreira, X., & Castagneyrol, B. (2021). Tree Diversity and Forest Resistance to Insect Pests:
Patterns, Mechanisms, and Prospects. *Annual Review of Entomology*, *66*(1), 277–296.
https://doi.org/10.1146/annurev-ento-041720-075234

Kaitaniemi, P., Riihimäki, J., Koricheva, J., & Vehviläinen, H. (2007). Experimental evidence for
associational resistance against the European pine sawfly in mixed tree stands. *Silva Fennica*, *41*(2),
259–268.

Klapwijk, M. J., & Björkman, C. (2018). Mixed forests to mitigate risk of insect outbreaks.
Scandinavian Journal of Forest Research, *33*(8), 772–780.
https://doi.org/10.1080/02827581.2018.1502805

Klapwijk, M. J., Hopkins, A. J. M., Eriksson, L., Pettersson, M., Schroeder, M., Lindelöw, Å., Rönnberg,
J., Keskitalo, E. C. H., & Kenis, M. (2016). Reducing the risk of invasive forest pests and pathogens:
Combining legislation, targeted management and public awareness. *Ambio*, *45*(Suppl 2), 223–234.
https://doi.org/10.1007/s13280-015-0748-3

Koricheva, J., Gurevitch, J., & Mengersen, K. (2013). *Handbook of Meta-analysis in Ecology and*
Evolution. https://press.princeton.edu/books/hardcover/9780691137285/handbook-of-meta-
analysis-in-ecology-and-evolution

Lawton, J. H., & Strong, D. R. (1981). Community Patterns and Competition in Folivorous Insects. *The*
American Naturalist, *118*(3), 317–338.

Legault, S., & James, P. M. A. (2018). Parasitism Rates of Spruce Budworm Larvae: Testing the Enemy
Hypothesis Along a Gradient of Forest Diversity Measured at Different Spatial Scales. *Environmental*
Entomology, *47*(5), 1083–1095. https://doi.org/10.1093/ee/nvy113

Letourneau, D. K. (1987). The Enemies Hypothesis: Tritrophic Interactions and Vegetational Diversity
in Tropical Agroecosystems. *Ecology*, *68*(6), 1616–1622. https://doi.org/10.2307/1939853

Letourneau, D. K., Armbrecht, I., Rivera, B. S., Lerma, J. M., Carmona, E. J., Daza, M. C., Escobar, S.,
Galindo, V., Gutiérrez, C., López, S. D., Mejía, J. L., Rangel, A. M. A., Rangel, J. H., Rivera, L., Saavedra,
C. A., Torres, A. M., & Trujillo, A. R. (2011). Does plant diversity benefit agroecosystems? A synthetic
review. *Ecological Applications*, *21*(1), 9–21. https://doi.org/10.1890/09-2026.1

Nakagawa, S., Noble, D. W. A., Senior, A. M., & Lagisz, M. (2017). Meta-evaluation of meta-analysis:

Ten appraisal questions for biologists. *BMC Biology*, *15*(1), 18. https://doi.org/10.1186/s12915-017-

0357-7

Nell, C. S., Abdala-Roberts, L., Parra-Tabla, V., & Mooney, K. A. (2018). Tropical tree diversity

mediates foraging and predatory effects of insectivorous birds. *Proceedings of the Royal Society B:*

Biological Sciences, *285*(1890), 20181842. https://doi.org/10.1098/rspb.2018.1842

Nixon, A. E., & Roland, J. (2012). Generalist predation on forest tent caterpillar varies with forest

stand composition: An experimental study across multiple life stages. *Ecological Entomology*, *37*(1),

13–23. https://doi.org/10.1111/j.1365-2311.2011.01330.x

Novais, S. M. A., Macedo-Reis, L. E., & Neves, F. S. (2017). Predatory beetles in cacao agroforestry

systems in Brazilian Atlantic forest: A test of the natural enemy hypothesis. *Agroforestry Systems*,

91(1), 201–209. https://doi.org/10.1007/s10457-016-9917-z

Paquette, A., Hector, A., Castagneyrol, B., Vanhellemont, M., Koricheva, J., Scherer-Lorenzen, M., &

Verheyen, K. (2018). A million and more trees for science. *Nature Ecology & Evolution*, *2*(5), 763–766.

https://doi.org/10.1038/s41559-018-0544-0

Pennings, S. C., & Silliman, B. R. (2005). Linking Biogeography and Community Ecology: Latitudinal

Variation in Plant-Herbivore Interaction Strength. *Ecology*, *86*(9), 2310–2319.

Penone, C., Allan, E., Soliveres, S., Felipe-Lucia, M. R., Gossner, M. M., Seibold, S., Simons, N. K.,

Schall, P., Plas, F. van der, Manning, P., Manzanedo, R. D., Boch, S., Prati, D., Ammer, C., Bauhus, J.,

Buscot, F., Ehbrecht, M., Goldmann, K., Jung, K., … Fischer, M. (2019). Specialisation and diversity of

multiple trophic groups are promoted by different forest features. *Ecology Letters*, *22*(1), 170–180.

https://doi.org/10.1111/ele.13182

Poch, T. J., & Simonetti, J. A. (2013). Ecosystem services in human-dominated landscapes: Insectivory

in agroforestry systems. *Agroforestry Systems*, *87*(4), 871–879. https://doi.org/10.1007/s10457-013-

9603-3

Riihimäki, J., Kaitaniemi, P., Koricheva, J., & Vehviläinen, H. (2005). Testing the enemies hypothesis in

forest stands: The important role of tree species composition. *Oecologia*, *142*(1), 90–97.

https://doi.org/10.1007/s00442-004-1696-y

Root, R. B. (1973). Organization of a Plant-Arthropod Association in Simple and Diverse Habitats: The

Fauna of Collards (Brassica Oleracea). *Ecological Monographs*, *43*(1), 95–124.

https://doi.org/10.2307/1942161

Rosenberg, M. S. (2005). The File-Drawer Problem Revisited: A General Weighted Method for

Calculating Fail-Safe Numbers in Meta-Analysis. *Evolution*, *59*(2), 464–468. JSTOR.

Rosenheim, J. A. (1998). Higher-Order Predators and the Regulation of Insect Herbivore Populations.

Annual Review of Entomology, *43*(1), 421–447. https://doi.org/10.1146/annurev.ento.43.1.421

Rosenheim, J. A., & Harmon, J. P. (2006). The Influence of Intraguild Predation on the Suppression of

a Shared Prey Population: An Empirical Reassessment. In J. Brodeur & G. Boivin (Eds.), *Trophic and*

Guild in Biological Interactions Control (pp. 1–20). Springer Netherlands. https://doi.org/10.1007/1-

4020-4767-3_1

Roslin, T., Hardwick, B., Novotny, V., Petry, W. K., Andrew, N. R., Asmus, A., Barrio, I. C., Basset, Y.,

Boesing, A. L., Bonebrake, T. C., Cameron, E. K., Dáttilo, W., Donoso, D. A., Drozd, P., Gray, C. L., Hik,

D. S., Hill, S. J., Hopkins, T., Huang, S., Slade, E. M. (2017). Higher predation risk for insect prey at low

latitudes and elevations. *Science*, *356*(6339), 742–744. https://doi.org/10.1126/science.aaj1631

Rumble, M. A., Flake, L. D., Mills, T. R., & Dykstra, B. L. (2001). Do pine trees in aspen stands increase

bird diversity? *Proceedings (P)*, *18*, 185–192.

Russell, E. P. (1989). Enemies Hypothesis: A Review of the Effect of Vegetational Diversity on

Predatory Insects and Parasitoids. *Environmental Entomology*, *18*(4), 590–599.

https://doi.org/10.1093/ee/18.4.590

Sanders, D., Schaefer, M., Platner, C., & Griffiths, G. J. K. (2011). Intraguild interactions among

generalist predator functional groups drive impact on herbivore and decomposer prey. *Oikos*, *120*(3),

418–426. https://doi.org/10.1111/j.1600-0706.2010.18924.x

Schuldt, A., Baruffol, M., Böhnke, M., Bruelheide, H., Härdtle, W., Lang, A. C., Nadrowski, K., Oheimb,

G. V., Voigt, W., Zhou, H., & Assmann, T. (2010). Tree diversity promotes insect herbivory in

subtropical forests of south-east China. *Journal of Ecology*, *98*(4), 917–926.

https://doi.org/10.1111/j.1365-2745.2010.01659.x

Schuldt, A., Fahrenholz, N., Brauns, M., Migge-Kleian, S., Platner, C., & Schaefer, M. (2008).

Communities of ground-living spiders in deciduous forests: Does tree species diversity matter?

Biodiversity and Conservation, *17*(5), 1267–1284. https://doi.org/10.1007/s10531-008-9330-7

Seidl, R., Schelhaas, M.-J., Rammer, W., & Verkerk, P. J. (2014). Increasing forest disturbances in

Europe and their impact on carbon storage. *Nature Climate Change*, *4*(9), 806–810.

https://doi.org/10.1038/nclimate2318

Siemann, E., Tilman, D., Haarstad, J., & Ritchie, M. (1998). Experimental Tests of the Dependence of

Arthropod Diversity on Plant Diversity. *The American Naturalist*, *152*(5), 738–750.

https://doi.org/10.1086/286204

Staab, M., & Schuldt, A. (2020). The Influence of Tree Diversity on Natural Enemies—A Review of the

"Enemies" Hypothesis in Forests. *Current Forestry Reports*, *6*(4), 243–259.

https://doi.org/10.1007/s40725-020-00123-6

Tarbox, B. C., Robinson, S. K., Loiselle, B., & Flory, S. L. (2018). Foraging ecology and flocking behavior

of insectivorous forest birds inform management of Andean silvopastures for conservation. *The*

Condor, *120*(4), 787–802. https://doi.org/10.1650/CONDOR-18-1.1

Vehviläinen, H., Koricheva, J., & Ruohomäki, K. (2007). Tree Species Diversity Influences Herbivore

Abundance and Damage: Meta-Analysis of Long-Term Forest Experiments. *Oecologia*, *152*(2), 287–

298.

Viechtbauer, W. (2010). Conducting Meta-Analyses in R with the metafor Package. *Journal of*

Statistical Software, *36*(1), 1–48. https://doi.org/10.18637/jss.v036.i03

Wan, N.-F., Zheng, X.-R., Fu, L.-W., Kiær, L. P., Zhang, Z., Chaplin-Kramer, R., Dainese, M., Tan, J., Qiu,

S.-Y., Hu, Y.-Q., Tian, W.-D., Nie, M., Ju, R.-T., Deng, J.-Y., Jiang, J.-X., Cai, Y.-M., & Li, B. (2020). Global

synthesis of effects of plant species diversity on trophic groups and interactions. *Nature Plants*, *6*(5),

503–510. https://doi.org/10.1038/s41477-020-0654-y

White, J. A., & Whitham, T. G. (2000). Associational Susceptibility of Cottonwood to a Box Elder

Herbivore. *Ecology*, *81*(7), 1795–1803. https://doi.org/10.1890/0012-

9658(2000)081[1795:ASOCTA]2.0.CO;2

Yang, B., Li, B., He, Y., Zhang, L., Bruelheide, H., & Schuldt, A. (2018). Tree diversity has contrasting

effects on predation rates by birds and arthropods on three broadleaved, subtropical tree species.

Ecological Research, *33*(1), 205–212. https://doi.org/10.1007/s11284-017-1531-7

Zvereva, E. L., & Kozlov, M. V. (2006). Consequences of simultaneous elevation of carbon dioxide and

temperature for plant–herbivore interactions: A metaanalysis. *Global Change Biology*, *12*(1), 27–41.

https://doi.org/10.1111/j.1365-2486.2005.01086.x

Zvereva, Elena L., Zverev, V., & Kozlov, M. V. (2020). Predation and parasitism on herbivorous insects

change in opposite directions in a latitudinal gradient crossing a boreal forest zone. *Journal of Animal*

Ecology, *89*(12), 2946–2957. https://doi.org/10.1111/1365-2656.13350

Figures

Figure 1. Map of study sites of the 65 articles included in the meta-analysis. Dot sizes are

proportional to the number of studies conducted at the same study site.

Figure 2. Summary of tree diversity effects on natural enemies (NE). (a) Grand mean effect size. (b)

Effect of NE responses type. (c) Effect of the forest biome. Dots and error bars represent model

parameter estimates and corresponding 95% confidence intervals. Black circled dots represent the

mean effect size for each moderator. Vertical thick lines represent the 95% confidence interval. Thin

vertical lines represent the prediction interval, which is the expected range of true effects in similar

studies. K (n) represent the number of case studies (K) and the number of primary studies (n) for

each moderator level.