TITLE

Members of the CUGBP Elav-Like Family of RNA-Binding Proteins are Expressed in Distinct Populations of Primary Sensory Neurons

AUTHORS

Eliza Grlickova-Duzevik[1,2]   egrlickovaduzevik@une.edu

Thomas M Reimonn[4]   Thomas.Reimonn@umassmed.edu

Merilla Michael[1,2]   merilla.michael@gmail.com

Tina Tian[5,6,7]   tina.tian@emory.edu

Jordan Owyoung[7,8]   jordan.nicole.owyoung@emory.edu

Aidan McGrath-Conwell[2,3]   amcgrathconwell@une.edu

Peter Neufeld[2,3]   pneufeld@une.edu

Madison Mueth[2,9]   mmueth@une.edu

Derek C Molliver[1,2]   dmolliver@une.edu

Patricia Jillian Ward[6,7]   jill.ward@emory.edu

Benjamin J Harrison[1,2]bharrison2@une.edu

1.   Biomedical Sciences, College of Osteopathic Medicine, University of New England, Biddeford. ME 04005

2.   Center for Excellence in the Neurosciences, University of New England. ME 04005

3.   College of Arts and Sciences, University of New England, Biddeford. ME 04005

4.   University of Massachusetts Medical School, Program in Bioinformatics and Integrative Biology, Worcester, MA

5.   Medical Scientist Training Program, Emory University, Atlanta, GA

6.   Neuroscience Graduate Program, Emory University, Atlanta, GA

7.   Emory University School of Medicine, Department of Cell Biology, Atlanta, GA

8.   Genetics and Molecular Biology Graduate Program, Atlanta, GA

9.   Graduate School of Biomedical Science and Engineering, University of Maine, Orono, ME 04469

ACKNOWLEDGMENTS:

P20GM103643 MH116492 T32GM107000

ABSTRACT

Primary sensory Dorsal Root Ganglia (DRG) neurons are diverse, with distinct populations that respond to specific stimuli. Previously, we observed that functionally distinct populations of DRG neurons express mRNA transcript variants with different 3' untranslated regions (3'UTR's). 3'UTRs harbor binding sites for interaction with RNA-binding proteins (RBPs) for transporting mRNAs to subcellular domains, modulating transcript stability and regulating the rate of translation. In the current study, analysis of publicly available single-cell RNA-Sequencing (scRNA-Seq) data generated from adult mice revealed that 17 3'UTR-binding RBPs were enriched in specific populations of DRG neurons. This included 4 members of the CUGBP Elav-Like Family (CELF): CELF2 and CELF4 were enriched in peptidergic, CELF6 in both peptidergic and nonpeptidergic and CELF3 in tyrosine hydroxylase-expressing neurons. Immunofluorescence studies confirmed that 60% of CELF4+ neurons are small diameter C fibers and 33% medium diameter myelinated (likely Aδ) fibers and showed that CELF4 is distributed to peripheral termini. Co-expression analyses using transcriptomic data and immunofluorescence revealed that CELF4 is enriched in nociceptive neurons that express GFRA3, CGRP and the capsaicin receptor TRPV1. Re-analysis of published transcriptomic data from macaque DRG revealed a highly similar distribution of CELF members, and re-analysis of single-nucleus RNA sequencing data derived mouse and rat DRG after sciatic injury revealed differential expression of CELFs in specific populations of sensory neurons. We propose that CELF RNA-binding proteins may regulate the fate of mRNAs in populations of nociceptors, and may play a role in pain and/or neuronal regeneration following nerve injury.

INTRODUCTION

RNA binding proteins (RBPs) are critical for orchestrating the post transcriptional fate of transcripts, including transport to sub-cellular domains for local translation, storage and release from ribonucleoprotein complexes and recruitment of translational machinery (e.g., review:

(Lenzken et al., 2014)). The 3' untranslated region (3'UTR) of transcripts is a hub for mRNA-

protein interactions (e.g., review:(Szostak & Gebauer, 2013)). For example, RBP binding to

3'UTRs on nascent transcripts regulates poly(A) signal site selection, cleavage and

polyadenylation (e.g., review: (Tian & Manley, 2013). On mature transcripts, 3'UTRs contain

interaction sites (originally termed "zip-code" motifs (Kislauskis et al., 1994; Kislauskis & Singer,

1992; Ross et al., 1997), for subcellular trafficking (Arora et al., 2022; Kim et al., 2015; Ribeiro

et al., 2020; Yergert et al., 2021) anchoring points for scaffolding interactions (Sahoo et al.,

2018) and interaction sites for recruitment of miRNAs (Grimson et al., 2007; Majoros & Ohler,

2007) or inhibition by competition for binding sites (Srikantan et al., 2012).

3'UTR-protein interactions are highly dynamic and provide a regulatory layer for modulation of

diverse cellular functions. By interaction with alternative cleavage sites at the distal end of

nascent transcripts, RBP-3'UTR interactions generate alternate 3'UTR isoforms through a

mechanism termed Alternative Polyadenylation (APA) (Singh et al., 2009). mRNA isoforms with

long 3'UTRs are predominantly expressed during development (Ji & Tian, 2009), and in the

adult are restricted to certain cell types most notably neurons that express mRNAs with long

3'UTRs (Harrison et al., 2014). Differential expression of 3'UTR isoforms by APA is associated

with diverse biological functions, for example long 3'UTR isoforms upregulated during

neuroplasticity (An et al., 2008; Miura et al., 2013) and during responses to stress (Graber et al.,

2013; Zheng et al., 2018), etc. 3'UTR variants are associated with human disease (Griesemer et

al., 2021). For example, shortened 3'UTR sequences contain fewer interactions sites for micro-

RNAs, and are associated with neurological disease such as Huntington's (Romo et al., 2017),

and facilitate oncogene activation and tumor growth (Mayr & Bartel, 2009).

RBP-3'UTR interactions are especially important for remote cellular compartments that rely on a

supply of mRNA for local protein synthesis. Sensory neurons have extremely long axons

projecting from peripheral targets through their cell bodies (soma) housed in DRG terminating in

the spinal cord dorsal horn. These termini require a supply of raw materials and machinery that

is transported along neuronal processes. RBPs are required for mRNA transport to and

translation in these remote domains for maintenance of synapses and for plasticity (Hornberg &

Holt, 2013; Wagnon et al., 2012). Numerous "neuron-specific" RBPs have been identified that

are highly enriched in neurons and are required for mRNA transport and neuronal excitability

(Darnell, 2013; Hornberg & Holt, 2013).

Primary somatosensory neurons are classified according to their responsiveness to specific

stimuli (light touch, temperature, stretch, etc.), neurochemistry and neurophysiology (Le Pichon

& Chesler, 2014), and can be identified by population-specific histological markers. Sensory

neurons with unmyelinated axons (C-fibers) tend to have small-diameter cell bodies, and the

great majority of these neurons are considered to be nociceptors (reviewed (Light, A.R. 1992)),

grossly divided into two subpopulations: peptidergic neurons, which express Calcitonin Gene

Related Peptide (CGRP) and Ntrk1 (gene for TRKA) and are particularly important for the

transduction of noxious heat stimuli, and non-peptidergic neurons, which bind the plant lectin

IB4 and express the G protein-coupled receptor MAS Related GPR Family Member D

(MRGPRD) and are particularly important for mechanical nociception (Dong et al., 2001;

Silverman & Kruger, 1988). Most sensory neurons that express the heat- and acid-gated

channel TRPV1 in adulthood are peptidergic neurons (Zwick et al., 2002). Peptidergic neurons

also include a subset of thinly myelinated (A-delta) nociceptors and a minority of A-beta

afferents (Lawson et al., 1993, 1996). A third population of C-fibers is comprised of neurons that

express tyrosine hydroxylase (TH) and are not nociceptors; instead they are sensitive

mechanoreceptors that have been implicated in the transduction of pleasant "social touch"

(Lallemend & Ernfors, 2012; Li et al., 2011). Although markers of the peptidergic/non-

peptidergic nociceptor populations are highly segregated in the mouse, the selectivity of these

markers varies considerably across species; in mouse, IB4-binding is seen in less than 10% of

neurons expressing CGRP or TRPV1, whereas in the rat this overlap is 35% (Price & Flores,

2007; Zwick et al., 2002), and recent evidence shows that in humans CGRP and TRPV1 are

expressed in most nociceptors (IB4 does not bind sensory neurons in human) (Shiers et al.,

2021). Recent advances in single-cell RNA-sequencing (scRNA-Seq) allow sensory neurons to

be characterized by comprehensive mRNA expression profiles, rather than a handful of

neurochemical markers. These transcriptomic studies align closely with histological data and

are a powerful resource for identifying the genomic signatures of functionally-distinct neuronal

populations (Usoskin et al., 2015).

Previously, we reported that functionally distinct populations of DRG neurons express mRNA

isoforms with divergent 3'UTR sequences (3'UTR isoforms), and that these 3'UTR variants

contain different RBP interaction motifs (Harrison et al., 2019). This led us to theorize that

sensory neuron diversity is controlled by differential RBP-3'UTR interactions. In the current

study we examine the expression of 3'UTR-binding RBPs in DRG, providing further evidence

suggesting divergence of 3'UTR-protein interactions in distinct populations of sensory neurons.

METHODS

Animals

All animal procedures were approved by the Institutional Animal Care and Use Committee of the

University of New England, and Emory University consistent with federal regulations and

guidelines. All animals used for this study were 6-8 week old male and female C57/BL6 mice

purchased from Jackson Labs (Bar Harbor, Maine).

Dorsal Root Ganglion Single Cell RNA-seq Analysis

Trimming, aligning, and quantification: Previously published single-cell RNA-seq data from adult

mouse dorsal root ganglion neurons produced by Usoskin et al. was accessed from GEO using

accession code GSE59739 (Usoskin et al., 2015). Fastq files were downloaded and reads were

trimmed using Trimmomatic version 0.39 (Bolger et al., 2014) in single-end mode with the

following parameters: ILLUMINACLIP:TruSeq3-SE:2:30:10 LEADING:3 TRAILING:3

SLIDINGWINDOW:4:15 MINLEN:36. Trimmed reads were aligned to the mouse genome

(GRCm38, primary assembly) and transcripts mapping to genes were quantified (Gencode

vM25, primary assembly) using STAR version 2.7.5a (Dobin et al., 2013) with default

parameters. Transcript quantifications were summed by gene when one cell contained reads

from multiple fastq files, and gene counts were tabulated into a gene-by-cell expression matrix.

Clustering, Visualization, and Annotation: Downstream processing was performed using Seurat

version 3.2.0 (Stuart et al., 2019). Cells that expressed fewer than 500 genes, expressed more

than 10,000 genes, contained fewer than 1,000 transcripts, or had more than 10% of transcripts

mapping to mitochondrial genes were removed, resulting in 811 cells passing all metrics. Gene

counts were normalized using "LogNormalize" and counts per 10,000 normalization. 3,000

highly variable genes were selected using the "vst" method. Normalized transcript counts were

scaled to 0 mean and unit variance, and principal component analysis with 100 principal

components was performed. Cells were Louvain clustered with a resolution of 1 and k = 20

nearest neighbors resulting in six clusters. Cells were embedded using t-SNE and UMAP with

default parameters. A grid search of clustering parameters for k nearest neighbors in {10, 15,

20, 30} and resolution in {0.5, 1, 1.5, and 2} was performed with visual inspection of

concordance between UMAP embedding and clustering to verify that k = 20 and resolution = 1

were reasonable. Normalized gene expression for 12 marker genes (Th, Sst, Mrgprd, P2rx3,

Tac1, Ntrk1, Calca, Nefh, Pvalb, B2m, Vim, Col6a2) were plotted and cluster cell types were

manually annotated as Non-neuronal, Peptidergic (Pep), non-peptidergic (NP), tyrosine

hydroxylase-expressing (TH) and neurofilament heavy chain (Nefh)-expressing (NF) (Fig 1). To

identify cell subpopulations, each cell type was separately processed using 2000 variable

features, 30 principal components, k = 10 nearest neighbors, and a louvain resolution of 1.

Differential Expression Analysis: Differentially expressed genes were identified using a Wilcoxon

rank sum test for each cell type compared to all other cell types where the log2 fold change was

greater than 0.25 and the gene was expressed in at least 10% of cells. A list of all known 3-

UTR-binding proteins was obtained from the Gene Ontology consortium website (Ashburner et

al., 2000; Gene Ontology Consortium, 2021) to highlight differentially expressed cell type

specific 3'UTR-binding proteins.

mRNA co-expression analysis: Pearson's correlation was calculated by comparison of CELF4

mRNA expression values vs expression values for all other detected mRNAs. For reference,

also shown are values for CGRP vs NTRK1 (known to be co-expressed in the same population

of neurons) and CGRP vs MRGPRD (known to be expressed separately in distinct populations)

(Fig 6A).

Antibody characterization (Table 1)

Polyclonal rabbit anti-CELF4 antibody (Prestige Antibodies, Sigma-Aldrich, HPA037986):

Developed using CUGBP, Elav-like family member 4 recombinant protein epitope signature tag

(PrEST). This antibody has previously been extensively validated for use with immunostaining:

This antibody detected CELF4 protein in the wildtype mouse hippocampal and cortical neurons,

but not in the CELF4 knockout samples (Wagnon et al., 2011, 2012). In addition, we determined

that CELF4 gene deletion (CELF4-fl/fl (Wagnon et al., 2011) ; Avil-cre-ERT2 mice treated with

tamoxifen (Lau et al., 2011) resulted in loss of signals (Fig 4A), as did omission of primary

antibody (Fig 6C, 7), providing further evidence that this reagent is specific and robust.

Polyclonal guinea pig anti-TRPV1 (Neuromics, GP14100): The antigen was residues of the

carboxy-terminus of rat TRPV1, YTGSLKPEDAEVFKDSMVPGEK. This antibody has been

assessed with wild-type tissue, as well as TRPV1-deficient mice, and no apparent

immunoreactivity was observed in TRPV1-KO mouse DRG, although the antibody robustly

stained the expected population of wild-type neurons (Sand et al., 2015). In addition, omission

of primary antibody resulted in loss of signals (Fig 6C).

Polyclonal chicken anti NF-H (EnCor, CPCA-NF-H): NF-H antibody robustly stained the

expected population of medium and large diameter DRG neurons and large caliber axons. No

signal was detected in NF-H negative small diameter neurons, small caliber fibers or non-

neuronal cells (Fig 5), thus confirming the specificity of this reagent.

Polyclonal guinea pig anti-CGRP (20R-CP001): CGRP antibody was raised in guinea pig using

calcitonin gene-related peptide conjugated to BSA as the immunogen. This antibody has been

extensively used for immunostaining DRG tissue slices to identify peptidergic neurons, and

specifically stains this population (Harrison et al., 2014).

Immunofluorescence Imaging and Analysis

Immunofluorescent staining was used to characterize populations of DRG neurons that express

CELF4 using reagents detailed in Table 1: CGRP for peptidergic, TRPV1 for heat responsive

nociceptors, and NF-H for myelinated neurons. Additionally, the plant lectin IB4 conjugated to

fluorophore was used to stain the non-peptidergic population, e.g.: (Burnstock, 2000)

Sample Collection and Processing: Animals were exsanguinated under deep pentobarbital

anesthesia, before perfusion with cold 4% paraformaldehyde in PBS. DRG were immediately

dissected, fixed in 4% paraformaldehyde (PFA) in PBS for 10 minutes before cryoprotection by

incubating overnight in 30% sucrose. 15 µm sections were cut using a cryostat, mounted onto

adhesion microscope slides, and left to dry, uncovered at room temperature for at least 1 hour.

Immunostaining: Non-specific binding sites were blocked by incubating sections with 5% donkey

serum in 0.4% Triton X-100 PBS for 1 hour, covered. CELF4 antibody was diluted 1:800,

TRPV1 1:500, CGRP 1:1000 and/or NF-H 1:40000 in blocking buffer and left on sections

overnight (i.e. 18-24 hours) at room temperature. The sections were then washed with 0.4%

Triton X-100 PBS; the washing solution was left on for 5 minutes for the first wash, then 10

minutes for the following three washes. Secondary antibodies and counterstains were prepared

1:200 in blocking buffer, added to sections and for 2 hours at room temperature. Alexa 488-

conjugated IB4 was diluted at 1:50. The sections were washed a second time using the same

procedure as the first wash. Slides were coverslipped with an antifade mountant and left

overnight at room temperature before imaging.

Imaging: Multichannel fluorescent micrographs of entire longitudinal DRG sections were

obtained using a Leica DMi8 epifluorescent inverted microscope with 20X objective and

automated field stitching. Image analysis was performed as previously published (Harrison et

al., 2014): Two sections per animal containing more than 200 neurons with visible nuclei were

selected from four animals. Sections were spaced more than 50 µm apart to ensure that no

single neuron could be included in the analysis twice. Using Image J software (Schneider et al.,

2012), individual neurons with visible nuclei were then manually circled to generate separate

regions of interest (ROIs). Mean fluorescent intensities were recorded along with the cross-

sectional area from each ROI. All ROI data from both sections from individual animals were then

collated into a single dataset per animal, yielding data from over 400 neurons per animal.

Background was estimated as the mean signal intensity of non-expressing neurons. The

median staining intensity for each channel was then calculated for each of the four datasets

(i.e., the median intensity of each DRG) to allow for normalization of fluorescence intensities to

the mean median value (division by central tendency [median] method). First, to observe the

distribution of CELF4 expression according to neuron soma size, normalized signals from all

animals were collated into a single dataset and subsequently binned according to soma cross-

sectional area (50µm bin sizes) (Fig. 4C). Neurons were assigned to either small (<300µm),

medium (300-700µm) or large (>700µm) populations. These size divisions were chosen due to

the well-established somatosensory characteristics that vary according to soma size. To

identify peptidergic, non-peptidergic and myelinated populations, sections were co-stained with

CGRP antibody, IB4 lectin or neurofilament heavy chain (NF-H) antibody respectively

(Ruscheweyh et al., 2007).

Neuronal Retrograde Tracing: For retrograde labeling of muscle afferents, 8 adult wild type

C57BL/6J mice (Jackson Laboratories, strain #000664) at least 8-weeks of age were studied.

The right bilateral heads of the gastrocnemius and tibialis anterior muscles received 2µL

injections of 5.0% fast blue retrograde tracer (Polysciences, 17740) with a 26-gauge 10µL

Hamilton microliter syringe (Model #701) distributed in at least 3 areas within the muscle

bellies under 2% isoflurane (Piramal Enterprises Limited, Telangana, India, ANADA 200-237,

NDC 66794-013-25) in 1L/min O_2, totaling 6µL of tracer. All mice received 1 dose of 5mg/kg

1.5mg/ml meloxicam (Pivetal, ANADA 200-550, NDC 46066-170-13). Four to 14 days later, to

allow for retrograde transport, mice were euthanized and perfused with 0.4% paraformaldehyde

as described above. The right L4 dorsal root ganglion (DRG) was dissected and placed in 20%

sucrose for cryoprotection prior to being cryosectioned for immunohistochemistry as detailed

above.

Skin Imaging: For analysis of lanceolate endings, 5 additional wild type C57BL/6J mice were

used. Following euthanasia and paraformaldehyde perfusion, a quadrangular piece of hairy hind

paw skin was dissected, cryoprotected and sectioned transversely at 30 µm. Following

immunolabeling with NF-H and CELF4, the skin was imaged on a Nikon A1R with 60x objective,

0.5 µm step size.

RESULTS

To identify RBPs expressed in populations of sensory neurons, we obtained publicly available

single-cell RNA-Sequencing data generated from adult mouse DRG (GEO GSE59739) (Usoskin

et al., 2015). Since release of those data, analysis tools have evolved significantly. In addition,

gene annotations, mapping coordinates and transcript variant sequences are under constant

revision. We therefore reanalyzed the raw sequencing data using current software (SEURAT

3.2.0) and annotations (Gencode vM25). Cells clustered into functionally defined populations as

expected (Fig 1): Peptidergic (PEP) that are predominantly nociceptors that express

neuropeptides such as CGRP, non-peptidergic (NP) that are predominantly nociceptors that do

not express these neuropeptides, tyrosine hydroxylase-expressing (TH) small-diameter sensory

neurons implicated in the pleasurable, affective component of touch and neurofilament heavy

chain (NF-H)-expressing that are myelinated and respond to low threshold stimuli such as light

touch. Within those clusters, neurons grouped into expected sub populations, for example

TRPV1-expressing neurons (heat responsive nociceptors) subclustered within the PEP cluster,

and parvalbumin-expressing neurons (chiefly proprioceptors) subclustered within the NF

(myelinated) cluster (Fig 1) (Usoskin et al., 2015).

Gene expression profiles from each population (Pep, NP, TH, NF) were compared by differential

gene expression analysis using Wilcoxon rank sum test with SEURAT software. Of

approximately 1100 known RBPs annotated in the mouse genome, 100 are assigned the gene

ontology (GO) term "mRNA 3'-UTR binding" (Table 2). Of those, 84 3'UTR-binding proteins

were detectable in DRG neurons (expression > 1 CFM) and 17 were enriched in at least 1

population (Table 2). Therefore, the majority (80%) of 3'UTR-binding proteins detected in DRG

neurons were ubiquitously expressed throughout all populations, as exemplified by FXR1 (Fig 2

a,b). Interestingly, 4 members of the CUGBP Elav-Like Family (CELF) were differentially

expressed: CELFs 2, 3, 4 and 6 (Table 3, Fig 2). CELF2 was enriched in the PEP population,

CELF3 in the TH population, CELF4 in the PEP population and CELF6 was enriched in both NP

and PEP populations.

During these studies, numerous single-cell/nucleus sequencing studies were published and the

data made publicly available. We therefore sought to confirm our findings using single nucleus

sequencing data from adult mouse DRG (Renthal et al., 2020), and adult macaque (Kupari et

al., 2021). We observed that CELF4 is predominantly in peptidergic populations (Fig. 3 a,b)

confirming our previous results. Interestingly, we also observed that CELF1 and 2 are

expressed in satellite/glial cells in addition to neurons (Figure 3 a).

CELF4 had the highest mean expression value of all family members (Figs. 2,3) and the largest

differential expression value (28.2 fold) over the mean (Table 3). This family member negatively

regulates neuron excitability in excitatory CNS neurons, likely by limiting translation of mRNAs

encoding synaptic proteins and ion channels (Wagnon et al., 2011, 2012). However, expression

and function of CELF4 in sensory neurons has not been described. RNA concentration does not

necessarily correlate with protein concentration, e.g.:(Edfors et al., 2016). Therefore, the

expression of CELF4 protein was examined using quantitative immunofluorescence microscopy.

By measurement of fluorescent signals from individual somata from 3 animals (approx. 200

somata/animal), we observed that 58% of small diameter (<300 μm) and 38% of medium

diameter (300-700 μm) DRG neurons were CELF4 positive (Fig 4). Using established

histological markers for non-peptidergic (plant lectin IB4), peptidergic (CGRP antiserum) and

myelinated (NFH antiserum) neurons, we observed that CELF4 protein distribution mirrors that

of CELF4 mRNA. CELF4 was detected in 53% of CGRP+, 34% of IB4-binding and 54% NF-H+

neurons (Fig 5). However, CELF4 protein is expressed to significantly lower levels in IB4-

binding ($p<0.05$, n=3) and NF-H+ ($p<0.01$, n=3 mice) neurons.

By comparing CELF4 mRNA expression to that of all other mRNAs detected in the scRNA-Seq

dataset, we determined that CELF4 expression is most highly correlated with GFRA3, TRPV1

and Calca (gene name for CGRP) expression in adult mouse DRG neurons (Fig 6, a).

Given the prominent functional role of TRPV1+ afferents in muscle nociception and mechanical

hyperalgesia, we chose to further characterize CELF4 protein expression in peripheral targets

by taking advantage of the well-known central and peripheral innervation pattern of the sciatic

nerve. Lumbar DRGs 3-5 contribute sensory innervation to peripheral targets to the muscles,

vasculature, and skin of the leg. We first performed retrograde tracing from the tibialis anterior

and gastrocnemius muscles to specifically label the DRG afferents innervating these muscles.

In total, we characterized 2,490 (1 DRG each from 8 mice) retrogradely labeled primary sensory

neurons for their expression of CELF4 and TRPV1. Many DRG neurons were CELF4+

(n=4876), and approximately 30% of them were also retrogradely labeled. We found that 42% of

retrogradely labeled afferents also expressed the neuropeptide TRPV1 in L4 DRG, which

agrees with previous findings using femoral nerve tracing and L2 DRG (Christianson et al.,

2006). The majority of triple labeled (fast blue, CELF4, TRPV1) neurons was small in

diameter. By co-staining DRG sections with TRPV1 and CELF4 antisera, we confirmed that the

majority CELF4 immunofluorescence signal intensity is correlated with TRPV1 signal intensity

(Fig 6b, c). Furthermore, among all TRPV1+ muscle afferents in the L4 DRG, 76% were also

CELF4+.

Again, using our predicted CELF4 phenotype (primarily peptidergic – GFRA3, CGRP, TRPV1),

we investigated CELF4 expression in another peripheral sensory target. Lanceolate endings are

innervated by low threshold mechanoreceptors (LTMRs) whose cell bodies reside in the DRG.

These specialized sensory termini are multiply innervated by a combination of Aβ-, Aδ-, and C-

LTMRs. The specialized combinations of these sensory endings encode distinct sensory

properties based on their myelination status and adaptation rates to sustained stimuli with at

least 5 physiologically distinct LTMRs in hairy skin (Kuehn et al., 2019). Given that 90% of the

body is covered with hair in most mammals, lanceolate endings are an indispensable unit for the

perception of mechanosensory stimuli upon the skin. The majority of these lanceolate endings

are also innervated by CGRP+ peptidergic nociceptors, which respond rapidly to hair pull and

are developmentally dependent on NGF-TRKA signaling (Ghitani et al., 2017; Li & Ginty, 2014).

Together with our finding that CELF4 is expressed in the cell bodies of this population, and that

CELF4 is transported to neuropil in the CNS (Wagnon et al., 2012), we reasoned that a large

proportion of lanceolate sensory endings in hairy skin would also express CELF4. We identified

114 lanceolate endings ~equally sampled among 5 mice, and 49.2% expressed CELF4. The

pattern of CELF4 labeling suggests that CELF4 is present in both the circumferential and

longitudinal endings surrounding the hair follicle (Fig 7), suggesting that CELF4 is present in the

CGRP+ endings and/or at least one population of LTRMs

Previous work has shown that CELF4 regulates neuron excitability (Wagnon et al., 2011) and

that CELF members may play a role in axon growth (Chen et al., 2016). We therefore wanted to

examine the expression of CELF proteins in a model of nerve injury where hyperexcitable

neurons are regenerating. Re-analysis of data from sciatic crush (Renthal et al., 2020) strongly

indicated that CELF RNA-binding proteins are differentially expressed during the time course of

recovery from sciatic injury (Fig. 8a). This was confirmed by analysis of bulk RNA-seq data

derived from rat sensory ganglia post sciatic injury (Harrison et al., 2019) where CELFs are

down-regulated during the course of recovery (Fig. 8 b).

DISCUSSION

Our previous work has shown that populations of sensory neurons express different mRNA

isoforms with alternate 3'UTRs (Harrison et al., 2014). In the current study, we investigated the

expression of 3'UTR-binding proteins in DRG, leading to the discovery that members of the

CUGBP-binding ELAV-like family of RBPs are expressed in distinct populations of sensory

neurons. By analysis of publicly available scRNA-Seq data, we revealed that CELF2 is

expressed in small diameter neurons and non-neuronal cells, CELF4 is restricted to the

peptidergic (PEP) population, and CELF6 is present in both peptidergic and non-peptidergic

(NP) populations. These neuronal populations predominantly represent different classes of C-

fiber nociceptors, and in the case of PEP also a subset of A-delta fibers. Most CGRP-

expressing (PEP) C-fibers also express the pro-inflammatory neuropeptide substance P

(Juránek & Lembeck, 1997). CGRP and SP are pro-inflammatory peptides that play key roles in

neurogenic inflammation; CGRP is a potent vasodilator and substance P induces mast cell

degranulation (Lim et al., 2017). In contrast, NP neurons have been reported to suppress mast

cell reactivity (Zhang et al., 2021). Notably, several studies have demonstrated that cutaneous

IB4-binding neurons transduce mechanical stimuli in the non-noxious range, and optogenetic

stimulation of these neurons fails to evoke nocifensive behavior unless they are sensitized by

tissue injury, calling into question whether these neurons act as nociceptors in healthy tissue

(Warwick et al., 2021). CELF3 is restricted to the TH population of non-peptidergic, innocuous

low threshold C-fiber mechanoreceptors, which have been implicated in the pleasurable,

affective component of touch (Lallemend & Ernfors, 2012; Li et al., 2011). Immunostaining

revealed that the family member with the highest expression, CELF4 is enriched in neurons that

express the capsaicin receptor TRPV1 and are therefore responsive to noxious heat. CELF4 is

a negative regulator of mRNA translation and neural excitability (Wagnon et al., 2011). We

therefore propose that further study of CELF4 could uncover mechanisms that regulate protein

synthesis in sensory neurons that convey pain and are important for neurogenic inflammation.

Previous studies examining the expression profile and function of CELF members have

determined that these proteins are expressed in the central nervous system during development

and in the adult (Table 4) (Giudice et al., 2016; Krismer et al., 2020; Samaras et al., 2020;

Schmidt et al., 2018). The RNA-binding motifs for all members are similar, with UG repeats

(Table 4), suggesting that these proteins have the ability to bind similar transcripts. However,

given that expression of CELF members is restricted in a cell-type dependent manner, this

could limit functional redundancy between members. Focusing on CELF4, in the CNS this

member is expressed primarily in excitatory neurons, including large pyramidal cells of the

cerebral cortex and hippocampus, where it tonically limits excitatory neurotransmission

(Wagnon et al., 2012). In these neurons, RNA-protein interaction profiling (RIP-Seq) revealed

that CELF4 predominantly binds to the 3'UTR of transcripts, and CELF4 knockout reduced

mRNA stability (Wagnon et al., 2012), and resulted in increases in expression of sodium

channel Na(v)1.6 (Sun et al., 2013). Therefore, CELF4 is a negative regulator of excitability of

excitatory CNS neurons most likely though limiting translation in these neurons. In accordance,

electrophysiological experiments using brain slices revealed that loss of CELF4 function

lowered the action potential (AP) initiation threshold and increased AP gain (Sun et al., 2013).

This is the first report characterizing expression of CELF4 in populations of neurons in

the peripheral nervous system, and its function in those neurons remains unknown.

Extrapolating from studies performed with CNS neurons where CELF4 primarily binds to sites

within the 3'UTR of mRNAs, indicated molecular functions of this protein could include alternate

polyadenylation thereby coordinating the expression of 3'UTR isoforms. This would be an

enticing possibility due to our previous observations of differential 3'UTR isoform expression in

DRG neuron populations (Harrison et al., 2014), and mechanisms responsible for cell-type

specific expression of 3'UTR variants are currently unknown. However, in the current study we

did not detect CELF4 in the nucleus of DRG neurons, making this possibility less likely. Also, in

the CNS CELF4 mRNA co-sediments with polysomes purified from neuropil, and CELF4

knockout increases the concentration of key excitability proteins in this compartment (Wagnon

et al., 2012). Therefore, CELF4 limits local translation in CNS axons. Together with our

findings, this suggests that CELF4 could also limit protein expression in populations of sensory

neuron termini, and experiments to characterize CELF4 expression in central and peripheral

endings are currently underway.

CELF4 tonically limits protein translation in and excitability of neurons. Therefore,

pharmacological targeting of CELF4-3'UTR interactions could lead to the development of tools

to study mechanisms of protein translation in neurons. Targeting CELF4 could have wide-

ranging implications for diseases of the nervous system. For example in the CNS, CELF4

knockout mice develop a seizure phenotype closely resembling an epilepsy syndrome caused

by CELF4 locus (18q12) deletion in humans (Halgren et al., 2012). Therefore, enhancing

CELF4 function has been proposed as a strategy to treat epilepsy. We have determined that in

the PNS CELF4 is enriched in C/Aδ neurons, a population of neurons critical for the etiology of

pathological pain, and that this protein is regulated during the time course of recovery from

sciatic injury. Therefore, enhancing CELF4 function in nociceptors could be a novel approach to

preventing/alleviating chronic pain.

FIGURE LEGENDS

Figure 1: Utilization of publicly available single cell RNA sequencing (scRNA-Seq) data,

derived from adult mouse sensory neurons, to generate population-specific gene

expression profiles.

To generate gene expression profiles for specific populations of mouse adult sensory neurons,

publicly available scRNA-seq data (GSE59739) were downloaded and processed with the latest

release of SEURAT software using up to date gene annotations. Graphs are Uniform Manifold

Approximation and Projections (UMAPs) of gene expression profiles. Each data point is a cell in

UMAP space. Yellow shading = high expression, Blue shading = low expression. Expression

profiles distributed into clusters as expected: **A-D** Peptidergic neurons were identified by

CALCA (CGRP) expression (PEP, purple cluster), non-peptidergic by MRGPRD expression

(NP, green cluster), tyrosine hydroxylase–expressing neurons (TH, red cluster) and myelinated

neurons by neurofilament heavy chain expression (NF, blue cluster). **E** Sub-cluster of TRPV1-

expressing nociceptors. **F** Sub-cluster of parvalbumin (PVALB) -expressing proprioceptors.

Figure 2: The CELF family of RNA-binding proteins are expressed in distinct populations

of sensory neurons.

A Mean vs variance plot of mRNA expression values for 3'UTR-binding RNA-binding protein

genes in adult mouse sensory neurons. X's are individual genes. FXR1 and CELF4 are

highlighted for reference. **B** Expression of a representative RBP, FXR1 is broadly expressed in

all clusters of DRG neurons. Yellow circles are individual neurons that express FXR1 to high

levels, blue circles = cells with low FXR1 expression **C** Heatmap of expression values of 3'UTR-

binding RBPs significantly differentially expressed (FDR<0.01, top-10) in distinct populations of

sensory neurons. Each line shows expression values in a single cell. Yellow = high expression ,

Purple = low expression. **D** Expression of significantly differentially expressed CELF-family

RBPs in DRG neuron clusters. Yellow = high expression , blue = low expression. **E** Violin plots

showing the distribution of expression values for significantly differentially expressed CELF

family proteins. Yellow triangles denote significant enrichment, purple triangles denote

significant exclusion. TH = tyrosine hydroxylase–expressing, NP = non-peptidergic, NF =

neurofilament heavy chain-expressing and PEP = peptidergic neurons.

Figure 3: Re-analysis of publicly available single cell RNA-sequencing datasets confirm

that CELF4 members are expressed in distinct populations of cells in sensory ganglia of

adult mouse and macaque.

A Expression of CELF member mRNAs in populations of neuronal and non-neuronal adult

mouse sensory ganglia cells. Data from GSE154659 (Renthal et al., 2020). CELF1 and 2

mRNAs are detected in non-neuronal populations, whereas CELF4 is enriched in peptidergic

neurons. **B** In the adult macaque, neuronal expression of CELF members is highly similar to

mouse. Data from (Kupari et al., 2021).

Figure 4: **CELF4 protein is expressed in small and medium diameter sensory neurons.**

A CELF4 gene deletion abolishes CELF4 immunoreactivity, confirming the specificity of the

CELF4 antibody. **B** Representative immunofluorescent micrograph of a 14µm DRG section

stained with CELF4 antibody. **C** Soma size distribution of CELF4+ neurons. Data are collated

from 3 sections, from 3 DRG from 3 replicate animals (approx. 100 soma per section). Somata

were categorized according to cross-sectional area - small (<300µm), medium (300-700µm) or

large (>700µm). **C i** Proportion of small, medium or large cross-sectional area DRG neurons

that express CELF4. **ii** Proportion of CELF4-expressing neurons with small, medium or large

cross-sectional areas.

Figure 5: CELF4 protein expression in peptidergic (CGRP+), non-peptidergic (IB4-

binding) and myelinated (NF-H+) neurons.

A-C 14µm DRG sections stained with CELF4 antibody, co-stained with established histological

markers. Peptidergic neurons were identified with CGRP antibody, non-peptidergic with isolectin

IB4 and myelinated with NFH. Scatter plots show distribution of fluorescence intensities of

somata collated from DRG from 3 replicate animals, 2 non-overlapping cross sections per

animal (approx. 100 soma per section). Dashed lines indicate background signals levels. **D** Pie

charts showing the proportion of double positive somata from each of the 3 stains. **E** Mean

CELF4 fluorescence values. Stats = ANOVA with post hoc t-tests, n=3 animals, *=p<.05,

**=p<.01.

Figure 6: CELF4 is co-expressed with the TRPV1 capsaicin receptor in DRG neurons.

(A) Pearson's r correlation coefficients were calculated for CELF4 mRNA concentration vs all

other transcripts. The 3 most positively and negatively correlated genes are shown. Also shown

are reference values for a known co-expressed pair of mRNAs (CGRP vs NTRK1) and a pair of

mRNA's know to be expressed in different neurons (CGRP vs MRGPRD). **(B)** 14µm DRG

sections were stained with CELF4 and TRPV1 antibody, and fluorescence intensity measured in

soma of 2 sections from 3 animals. Representative micrographs are shown. **(C)** To support the

mRNA correlations, we identified muscle afferents that co-express TRPV1 and CELF4 protein

by using retrograde neuronal tracing from the gastrocnemius and tibialis anterior muscles

combined with immunohistochemistry. Data collected from 3 DRG from 8 replicate animals

(lumbar level 4). Representative micrographs are shown along with the no primary controls.

Figure 7: CELF4 is distributed to the peripheral termini of sensory neurons.

The location of lanceolate endings in mouse hind paw hairy skin sections was identified by NFH

immunoreactivity. CELF4 was detected in 49.2% of lanceolate sensory nerve endings in hair

follicles within both the longitudinal and circumferential portions of the sensory terminals (arrow

heads). Representative micrographs are shown along with no primary control. n=5 animals.

Figure 8: CELF transcripts are downregulated in sensory ganglia in rodent models of neuropathic pain.

Single nucleus sequencing data GSE154659 (Renthal et al., 2020) from mouse DRG (**A**) and bulk homogenate RNA-Seq from rat DRG (Harrison et al., 2019) (**B**) showed differential expression of CELF members post sciatic injury. * = q<0.05, ** = q<0.01 where q values are determined by Student's t-tests with Bonferroni multiple testing correction.

References

An, J. J., Gharami, K., Liao, G.-Y., Woo, N. H., Lau, A. G., Vanevski, F., Torre, E. R., Jones, K. R.,

Feng, Y., Lu, B., & Xu, B. (2008). Distinct role of long 3' UTR BDNF mRNA in spine

morphology and synaptic plasticity in hippocampal neurons. *Cell*, *134*(1), 175–187. PubMed.

https://doi.org/10.1016/j.cell.2008.05.045

Arora, A., Castro-Gutierrez, R., Moffatt, C., Eletto, D., Becker, R., Brown, M., Moor, A. E., Russ, H. A.,

& Taliaferro, J. M. (2022). High-throughput identification of RNA localization elements in

neuronal cells. *Nucleic Acids Research*, *50*(18), 10626–10642.

https://doi.org/10.1093/nar/gkac763

Ashburner, M., Ball, C. A., Blake, J. A., Botstein, D., Butler, H., Cherry, J. M., Davis, A. P., Dolinski, K.,

Dwight, S. S., Eppig, J. T., Harris, M. A., Hill, D. P., Issel-Tarver, L., Kasarskis, A., Lewis, S.,

Matese, J. C., Richardson, J. E., Ringwald, M., Rubin, G. M., & Sherlock, G. (2000). Gene

Ontology: Tool for the unification of biology. *Nature Genetics*, *25*(1), 1.

https://doi.org/10.1038/75556

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina sequence

data. *Bioinformatics*, *30*(15), 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Burnstock, G. (2000). P2X receptors in sensory neurones. *British Journal of Anaesthesia*, *84*(4), 476–488.

https://doi.org/10.1093/oxfordjournals.bja.a013473

Chen, L., Liu, Z., Zhou, B., Wei, C., Zhou, Y., Rosenfeld, M. G., Fu, X.-D., Chisholm, A. D., & Jin, Y.

(n.d.). CELF RNA binding proteins promote axon regeneration in C. elegans and mammals

through alternative splicing of Syntaxins. *ELife*, *5*, e16072. https://doi.org/10.7554/eLife.16072

Christianson, J. A., McIlwrath, S. L., Koerber, H. R., & Davis, B. M. (2006). Transient receptor potential

vanilloid 1-immunopositive neurons in the mouse are more prevalent within colon afferents

compared to skin and muscle afferents. *Neuroscience*, *140*(1), 247–257.

https://doi.org/10.1016/j.neuroscience.2006.02.015

Darnell, R. B. (2013). RNA Protein Interaction in Neurons. *Annual Review of Neuroscience*, *36*, 243–270.

https://doi.org/10.1146/annurev-neuro-062912-114322

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., &

Gingeras, T. R. (2013). STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics*, *29*(1), 15–

21. https://doi.org/10.1093/bioinformatics/bts635

Dong, X., Han, S., Zylka, M. J., Simon, M. I., & Anderson, D. J. (2001). A diverse family of GPCRs

expressed in specific subsets of nociceptive sensory neurons. *Cell*, *106*(5), 619–632.

https://doi.org/10.1016/s0092-8674(01)00483-4

Edfors, F., Danielsson, F., Hallström, B. M., Käll, L., Lundberg, E., Pontén, F., Forsström, B., & Uhlén,

M. (2016). Gene-specific correlation of RNA and protein levels in human cells and tissues.

Molecular Systems Biology, *12*(10). https://doi.org/10.15252/msb.20167144

Gene Ontology Consortium. (2021). The Gene Ontology resource: Enriching a GOld mine. *Nucleic Acids*

Research, *49*(D1), D325–D334. https://doi.org/10.1093/nar/gkaa1113

Gentleman, R. C., Carey, V. J., Bates, D. M., Bolstad, B., Dettling, M., Dudoit, S., Ellis, B., Gautier, L.,

Ge, Y., Gentry, J., Hornik, K., Hothorn, T., Huber, W., Iacus, S., Irizarry, R., Leisch, F., Li, C.,

Maechler, M., Rossini, A. J., … Zhang, J. (2004). Bioconductor: Open software development for

computational biology and bioinformatics. *Genome Biology*, *5*(10), R80.

https://doi.org/10.1186/gb-2004-5-10-r80

Ghitani, N., Barik, A., Szczot, M., Thompson, J. H., Li, C., Le Pichon, C. E., Krashes, M. J., & Chesler,

A. T. (2017). Specialized Mechanosensory Nociceptors Mediating Rapid Responses to Hair Pull.

Neuron, *95*(4), 944-954.e4. https://doi.org/10.1016/j.neuron.2017.07.024

Giudice, G., Sánchez-Cabo, F., Torroja, C., & Lara-Pezzi, E. (2016). ATtRACT-a database of RNA-

binding proteins and associated motifs. *Database: The Journal of Biological Databases and*

Curation, *2016*, baw035. https://doi.org/10.1093/database/baw035

Graber, J. H., Nazeer, F. I., Yeh, P., Kuehner, J. N., Borikar, S., Hoskinson, D., & Moore, C. L. (2013).

DNA damage induces targeted, genome-wide variation of poly(A) sites in budding yeast.

Genome Research, *23*(10), 1690–1703. https://doi.org/10.1101/gr.144964.112

Griesemer, D., Xue, J. R., Reilly, S. K., Ulirsch, J. C., Kukreja, K., Davis, J. R., Kanai, M., Yang, D. K.,

Butts, J. C., Guney, M. H., Luban, J., Montgomery, S. B., Finucane, H. K., Novina, C. D.,

Tewhey, R., & Sabeti, P. C. (2021). Genome-wide functional screen of 3′UTR variants uncovers

causal variants for human disease and evolution. *Cell*, *184*(20), 5247-5260.e19.

https://doi.org/10.1016/j.cell.2021.08.025

Grimson, A., Farh, K. K.-H., Johnston, W. K., Garrett-Engele, P., Lim, L. P., & Bartel, D. P. (2007).

MicroRNA targeting specificity in mammals: Determinants beyond seed pairing. *Molecular Cell*,

27(1), 91–105. https://doi.org/10.1016/j.molcel.2007.06.017

Halgren, C., Bache, I., Bak, M., Myatt, M. W., Anderson, C. M., Brøndum-Nielsen, K., & Tommerup, N.

(2012). Haploinsufficiency of CELF4 at 18q12.2 is associated with developmental and behavioral

disorders, seizures, eye manifestations, and obesity. *European Journal of Human Genetics*,

20(12), 12. https://doi.org/10.1038/ejhg.2012.92

Harrison, B. J., Flight, R. M., Gomes, C., Venkat, G., Ellis, S. R., Sankar, U., Twiss, J. L., Rouchka, E.

C., & Petruska, J. C. (2014). IB4-binding sensory neurons in the adult rat express a novel 3′ UTR-

extended isoform of CaMK4 that is associated with its localization to axons. *Journal of*

Comparative Neurology, *522*(2), 308–336. https://doi.org/10.1002/cne.23398

Harrison, B. J., Park, J. W., Gomes, C., Petruska, J. C., Sapio, M. R., Iadarola, M. J., Chariker, J. H., &

Rouchka, E. C. (2019). Detection of Differentially Expressed Cleavage Site Intervals Within 3'

Untranslated Regions Using CSI-UTR Reveals Regulated Interaction Motifs. *Frontiers in*

Genetics, *10*, 182. https://doi.org/10.3389/fgene.2019.00182

Hornberg, H. K., & Holt, C. (2013). RNA-binding proteins and translational regulation in axons and

growth cones. *Frontiers in Neuroscience*, *7*. https://doi.org/10.3389/fnins.2013.00081

Huber, W., Carey, V. J., Gentleman, R., Anders, S., Carlson, M., Carvalho, B. S., Bravo, H. C., Davis, S.,

Gatto, L., Girke, T., Gottardo, R., Hahne, F., Hansen, K. D., Irizarry, R. A., Lawrence, M., Love,

M. I., MacDonald, J., Obenchain, V., Oleś, A. K., … Morgan, M. (2015). Orchestrating high-

throughput genomic analysis with Bioconductor. *Nature Methods*, *12*(2), 115–121.

https://doi.org/10.1038/nmeth.3252

Ji, Z., & Tian, B. (2009). Reprogramming of 3′ Untranslated Regions of mRNAs by Alternative

Polyadenylation in Generation of Pluripotent Stem Cells from Different Cell Types. *PLoS ONE*,

4(12). https://doi.org/10.1371/journal.pone.0008419

Juránek, I., & Lembeck, F. (1997). Afferent C-fibres release substance P and glutamate. *Canadian*

Journal of Physiology and Pharmacology, *75*(6), 661–664.

Kim, H. H., Lee, S. J., Gardiner, A. S., Perrone-Bizzozero, N. I., & Yoo, S. (2015). Different motif

requirements for the localization zipcode element of β-actin mRNA binding by HuD and ZBP1.

Nucleic Acids Research, *43*(15), 7432–7446. https://doi.org/10.1093/nar/gkv699

Kislauskis, E. H., & Singer, R. H. (1992). Determinants of mRNA localization. *Current Opinion in Cell*

Biology, *4*(6), 975–978. https://doi.org/10.1016/0955-0674(92)90128-y

Kislauskis, E. H., Zhu, X., & Singer, R. H. (1994). Sequences responsible for intracellular localization of

beta-actin messenger RNA also affect cell phenotype. *The Journal of Cell Biology*, *127*(2), 441–

451. https://doi.org/10.1083/jcb.127.2.441

Krismer, K., Bird, M. A., Varmeh, S., Handly, E. D., Gattinger, A., Bernwinkler, T., Anderson, D. A.,

Heinzel, A., Joughin, B. A., Kong, Y. W., Cannell, I. G., & Yaffe, M. B. (2020a). Transite: A

Computational Motif-Based Analysis Platform That Identifies RNA-Binding Proteins Modulating

Changes in Gene Expression. *Cell Reports*, *32*(8). https://doi.org/10.1016/j.celrep.2020.108064

Krismer, K., Bird, M. A., Varmeh, S., Handly, E. D., Gattinger, A., Bernwinkler, T., Anderson, D. A.,

Heinzel, A., Joughin, B. A., Kong, Y. W., Cannell, I. G., & Yaffe, M. B. (2020b). Transite: A

Computational Motif-Based Analysis Platform That Identifies RNA-Binding Proteins Modulating

Changes in Gene Expression. *Cell Reports*, *32*(8), 108064.

https://doi.org/10.1016/j.celrep.2020.108064

Kuehn, E. D., Meltzer, S., Abraira, V. E., Ho, C.-Y., & Ginty, D. D. (2019). Tiling and somatotopic

alignment of mammalian low-threshold mechanoreceptors. *Proceedings of the National Academy*

of Sciences of the United States of America, *116*(19), 9168–9177.

https://doi.org/10.1073/pnas.1901378116

Kupari, J., Usoskin, D., Parisien, M., Lou, D., Hu, Y., Fatt, M., Lönnerberg, P., Spångberg, M., Eriksson,

B., Barkas, N., Kharchenko, P. V., Loré, K., Khoury, S., Diatchenko, L., & Ernfors, P. (2021).

Single cell transcriptomics of primate sensory neurons identifies cell types associated with

chronic pain. *Nature Communications*, *12*(1), 1510. https://doi.org/10.1038/s41467-021-21725-z

Lallemend, F., & Ernfors, P. (2012). Molecular interactions underlying the specification of sensory

neurons. *Trends in Neurosciences*, *35*(6), 373–381. https://doi.org/10.1016/j.tins.2012.03.006

Lau, J., Minett, M. S., Zhao, J., Dennehy, U., Wang, F., Wood, J. N., & Bogdanov, Y. D. (2011).

Temporal control of gene deletion in sensory ganglia using a tamoxifen-inducible Advillin-Cre-

ERT2 recombinase mouse. *Molecular Pain*, *7*, 100. https://doi.org/10.1186/1744-8069-7-100

Lawson, S. N., McCarthy, P. W., & Prabhakar, E. (1996). Electrophysiological properties of neurones

with CGRP-like immunoreactivity in rat dorsal root ganglia. *The Journal of Comparative*

Neurology, *365*(3), 355–366. https://doi.org/10.1002/(SICI)1096-

9861(19960212)365:3<355::AID-CNE2>3.0.CO;2-3

Lawson, S. N., Perry, M. J., Prabhakar, E., & McCarthy, P. W. (1993). Primary sensory neurones:

Neurofilament, neuropeptides, and conduction velocity. *Brain Research Bulletin*, *30*(3–4), 239–

243. https://doi.org/10.1016/0361-9230(93)90250-f

Le Pichon, C. E., & Chesler, A. T. (2014). The functional and anatomical dissection of somatosensory

subpopulations using mouse genetics. *Frontiers in Neuroanatomy*, *8*, 21–21. PubMed.

https://doi.org/10.3389/fnana.2014.00021

Lenzken, S. C., Achsel, T., Carrì, M. T., & Barabino, S. M. L. (2014). Neuronal RNA-binding proteins in

health and disease. *Wiley Interdisciplinary Reviews. RNA*, *5*(4), 565–576.

https://doi.org/10.1002/wrna.1231

Li, L., & Ginty, D. D. (2014). The structure and organization of lanceolate mechanosensory complexes at

mouse hair follicles. *ELife*, *3*, e01901. https://doi.org/10.7554/eLife.01901

Li, L., Rutlin, M., Abraira, V. E., Cassidy, C., Kus, L., Gong, S., Jankowski, M. P., Luo, W., Heintz, N.,

Koerber, H. R., Woodbury, C. J., & Ginty, D. D. (2011). The functional organization of

cutaneous low-threshold mechanosensory neurons. *Cell*, *147*(7), 1615–1627.

https://doi.org/10.1016/j.cell.2011.11.027

Light, A. R. (n.d.). *The Initial Processing of Pain and Its Descending Control: Spinal and Trigeminal*

Systems | Karger Book. Retrieved February 1, 2023, from

https://www.karger.com/Book/Home/222289

Lim, J. E., Chung, E., & Son, Y. (2017). A neuropeptide, Substance-P, directly induces tissue-repairing

M2 like macrophages by activating the PI3K/Akt/mTOR pathway even in the presence of IFNγ.

Scientific Reports, *7*(1), 9417. https://doi.org/10.1038/s41598-017-09639-7

Majoros, W. H., & Ohler, U. (2007). Spatial preferences of microRNA targets in 3' untranslated regions.

BMC Genomics, *8*(1), 152. https://doi.org/10.1186/1471-2164-8-152

Mayr, C., & Bartel, D. P. (2009). Widespread Shortening of 3′UTRs by Alternative Cleavage and

Polyadenylation Activates Oncogenes in Cancer Cells. *Cell*, *138*(4), 673–684.

https://doi.org/10.1016/j.cell.2009.06.016

Mi, H., Ebert, D., Muruganujan, A., Mills, C., Albou, L.-P., Mushayamaha, T., & Thomas, P. D. (2021).

PANTHER version 16: A revised family classification, tree-based classification tool, enhancer

regions and extensive API. *Nucleic Acids Research*, *49*(D1), D394–D403.

https://doi.org/10.1093/nar/gkaa1106

Miura, P., Shenker, S., Andreu-Agullo, C., Westholm, J. O., & Lai, E. C. (2013). Widespread and

extensive lengthening of 3' UTRs in the mammalian brain. *Genome Research*, *23*(5), 812–825.

https://doi.org/10.1101/gr.146886.112

Price, T. J., & Flores, C. M. (2007). Critical evaluation of the colocalization between calcitonin gene-

related peptide, substance P, transient receptor potential vanilloid subfamily type 1

immunoreactivities, and isolectin B4 binding in primary afferent neurons of the rat and mouse.

The Journal of Pain: Official Journal of the American Pain Society, *8*(3), 263–272.

https://doi.org/10.1016/j.jpain.2006.09.005

Renthal, W., Tochitsky, I., Yang, L., Cheng, Y.-C., Li, E., Kawaguchi, R., Geschwind, D. H., & Woolf,

C. J. (2020). Transcriptional Reprogramming of Distinct Peripheral Sensory Neuron Subtypes

after Axonal Injury. *Neuron*, *108*(1), 128-144.e9. https://doi.org/10.1016/j.neuron.2020.07.026

Ribeiro, D. M., Prod'homme, A., Teixeira, A., Zanzoni, A., & Brun, C. (2020). The role of 3'UTR-

protein complexes in the regulation of protein multifunctionality and subcellular localization.

Nucleic Acids Research, *48*(12), 6491–6502. https://doi.org/10.1093/nar/gkaa462

Romo, L., Ashar-Patel, A., Pfister, E., & Aronin, N. (2017). Alterations in mRNA 3′ UTR Isoform

Abundance Accompany Gene Expression Changes in Human Huntington's Disease Brains. *Cell*

Reports, *20*(13), 3057–3070. https://doi.org/10.1016/j.celrep.2017.09.009

Ross, A. F., Oleynikov, Y., Kislauskis, E. H., Taneja, K. L., & Singer, R. H. (1997). Characterization of a

beta-actin mRNA zipcode-binding protein. *Molecular and Cellular Biology*, *17*(4), 2158–2165.

Ruscheweyh, R., Forsthuber, L., Schoffnegger, D., & Sandkühler, J. (2007). Modification of classical

neurochemical markers in identified primary afferent neurons with Aβ-, Aδ-, and C-fibers after

chronic constriction injury in mice. *Journal of Comparative Neurology*, *502*(2), 325–336.

https://doi.org/10.1002/cne.21311

Sahoo, P. K., Lee, S. J., Jaiswal, P. B., Alber, S., Kar, A. N., Miller-Randolph, S., Taylor, E. E., Smith,

T., Singh, B., Ho, T. S.-Y., Urisman, A., Chand, S., Pena, E. A., Burlingame, A. L., Woolf, C. J.,

Fainzilber, M., English, A. W., & Twiss, J. L. (2018). Axonal G3BP1 stress granule protein limits

axonal mRNA translation and nerve regeneration. *Nature Communications*, *9*(1), 3358.

https://doi.org/10.1038/s41467-018-05647-x

Samaras, P., Schmidt, T., Frejno, M., Gessulat, S., Reinecke, M., Jarzab, A., Zecha, J., Mergner, J.,

Giansanti, P., Ehrlich, H.-C., Aiche, S., Rank, J., Kienegger, H., Krcmar, H., Kuster, B., &

Wilhelm, M. (2020). ProteomicsDB: A multi-omics and multi-organism resource for life science

research. *Nucleic Acids Research*, *48*(D1), D1153–D1163. https://doi.org/10.1093/nar/gkz974

Sand, C. A., Grant, A. D., & Nandi, M. (2015). Vascular Expression of Transient Receptor Potential

Vanilloid 1 (TRPV1). *Journal of Histochemistry and Cytochemistry*, *63*(6), 449–453.

https://doi.org/10.1369/0022155415581014

Schmidt, T., Samaras, P., Frejno, M., Gessulat, S., Barnert, M., Kienegger, H., Krcmar, H., Schlegl, J.,

Ehrlich, H.-C., Aiche, S., Kuster, B., & Wilhelm, M. (2018). ProteomicsDB. *Nucleic Acids*

Research, *46*(D1), D1271–D1281. https://doi.org/10.1093/nar/gkx1029

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image

analysis. *Nature Methods*, *9*(7), 7. https://doi.org/10.1038/nmeth.2089

Shiers, S. I., Sankaranarayanan, I., Jeevakumar, V., Cervantes, A., Reese, J. C., & Price, T. J. (2021).

Convergence of peptidergic and non-peptidergic protein markers in the human dorsal root

ganglion and spinal dorsal horn. *The Journal of Comparative Neurology*, *529*(10), 2771–2788.

https://doi.org/10.1002/cne.25122

Silverman, J. D., & Kruger, L. (1988). Lectin and neuropeptide labeling of separate populations of dorsal

root ganglion neurons and associated "nociceptor" thin axons in rat testis and cornea whole-

mount preparations. *Somatosensory Research*, *5*(3), 259–267.

https://doi.org/10.3109/07367228809144630

Singh, P., Alley, T. L., Wright, S. M., Kamdar, S., Schott, W., Wilpan, R. Y., Mills, K. D., & Graber, J.

H. (2009). Global changes in processing of mRNA 3' untranslated regions characterize clinically

distinct cancer subtypes. *Cancer Research*, *69*(24), 9422–9430. https://doi.org/10.1158/0008-

5472.CAN-09-2236

Srikantan, S., Tominaga, K., & Gorospe, M. (2012). Functional interplay between RNA-binding protein

HuR and microRNAs. *Current Protein & Peptide Science, 13*(4), 372–379.

Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W. M., Hao, Y., Stoeckius, M.,

Smibert, P., & Satija, R. (2019). Comprehensive Integration of Single-Cell Data. *Cell, 177*(7),

1888-1902.e21. https://doi.org/10.1016/j.cell.2019.05.031

Sun, W., Wagnon, J. L., Mahaffey, C. L., Briese, M., Ule, J., & Frankel, W. N. (2013). Aberrant sodium

channel activity in the complex seizure disorder of Celf4 mutant mice. *The Journal of Physiology,*

591(1), 241–255. https://doi.org/10.1113/jphysiol.2012.240168

Szostak, E., & Gebauer, F. (2013). Translational control by 3′-UTR-binding proteins. *Briefings in*

Functional Genomics, 12(1), 58–65. https://doi.org/10.1093/bfgp/els056

Tian, B., & Manley, J. L. (2013). Alternative cleavage and polyadenylation: The long and short of it.

Trends in Biochemical Sciences, 38(6), 312–320. https://doi.org/10.1016/j.tibs.2013.03.005

Usoskin, D., Furlan, A., Islam, S., Abdo, H., Lönnerberg, P., Lou, D., Hjerling-Leffler, J., Haeggström, J.,

Kharchenko, O., Kharchenko, P. V., Linnarsson, S., & Ernfors, P. (2015a). Unbiased

classification of sensory neuron types by large-scale single-cell RNA sequencing. *Nature*

Neuroscience, 18(1), 145–153. https://doi.org/10.1038/nn.3881

Usoskin, D., Furlan, A., Islam, S., Abdo, H., Lönnerberg, P., Lou, D., Hjerling-Leffler, J., Haeggström, J.,

Kharchenko, O., Kharchenko, P. V., Linnarsson, S., & Ernfors, P. (2015b). Unbiased

classification of sensory neuron types by large-scale single-cell RNA sequencing. *Nature*

Neuroscience, 18(1), 145–153. https://doi.org/10.1038/nn.3881

Wagnon, J. L., Briese, M., Sun, W., Mahaffey, C. L., Curk, T., Rot, G., Ule, J., & Frankel, W. N. (2012).

CELF4 Regulates Translation and Local Abundance of a Vast Set of mRNAs, Including Genes

Associated with Regulation of Synaptic Function. *PLoS Genetics, 8*(11).

https://doi.org/10.1371/journal.pgen.1003067

Wagnon, J. L., Mahaffey, C. L., Sun, W., Yang, Y., Chao, H.-T., & Frankel, W. N. (2011). Etiology of a

genetically complex seizure disorder in Celf4 mutant mice. *Genes, Brain, and Behavior*, *10*(7),

765–777. https://doi.org/10.1111/j.1601-183X.2011.00717.x

Warwick, C., Cassidy, C., Hachisuka, J., Wright, M. C., Baumbauer, K. M., Adelman, P. C., Lee, K. H.,

Smith, K. M., Sheahan, T. D., Ross, S. E., & Koerber, H. R. (2021). MrgprdCre lineage neurons

mediate optogenetic allodynia through an emergent polysynaptic circuit. *Pain*, *162*(7), 2120–

2131. https://doi.org/10.1097/j.pain.0000000000002227

Yergert, K. M., Doll, C. A., O'Rouke, R., Hines, J. H., & Appel, B. (2021). Identification of 3' UTR

motifs required for mRNA localization to myelin sheaths in vivo. *PLoS Biology*, *19*(1),

e3001053. https://doi.org/10.1371/journal.pbio.3001053

Zhang, S., Edwards, T. N., Chaudhri, V. K., Wu, J., Cohen, J. A., Hirai, T., Rittenhouse, N., Schmitz, E.

G., Zhou, P. Y., McNeil, B. D., Yang, Y., Koerber, H. R., Sumpter, T. L., Poholek, A. C., Davis,

B. M., Albers, K. M., Singh, H., & Kaplan, D. H. (2021). Nonpeptidergic neurons suppress mast

cells via glutamate to maintain skin homeostasis. *Cell*, *184*(8), 2151-2166.e16.

https://doi.org/10.1016/j.cell.2021.03.002

Zheng, D., Wang, R., Ding, Q., Wang, T., Xie, B., Wei, L., Zhong, Z., & Tian, B. (2018). Cellular stress

alters 3'UTR landscape through alternative polyadenylation and isoform-specific degradation.

Nature Communications, *9*(1), 2268. https://doi.org/10.1038/s41467-018-04730-7

Zwick, M., Davis, B. M., Woodbury, C. J., Burkett, J. N., Koerber, H. R., Simpson, J. F., & Albers, K. M.

(2002). Glial cell line-derived neurotrophic factor is a survival factor for isolectin B4-positive,

but not vanilloid receptor 1-positive, neurons in the mouse. *The Journal of Neuroscience: The*

Official Journal of the Society for Neuroscience, *22*(10), 4057–4065. https://doi.org/20026394

Figure 1 (Methods)

Figure 2

(a)

(b)

Figure 3

Figure 4

Figure 5

(a)

mRNAs		r
CGRP vs NTRK1		**0.41**
CELF4 vs GFRA3		0.42
"	TRPV1	0.39
"	CALCA	0.34
"	TH	-0.11
"	MRGPRD	-0.09
"	NTRK3	-0.06
CGRP vs MRGPRD		**-0.05**

(b)

CELF4   TRPV1   Merge   150µm

r = 0.41

CELF4 Fluorescence

(c)

CELF4   TRPV1   Fast Blue   Merge

No Primary   No Primary   Fast Blue   Merge   200µm

CELF4+ / FaBlu+

small
medium
large

4%
26%
70%

FaBlu+ that are TRPV1+: 41.9%
TRPV1+ that are CELF4+: 76.0%
CELF4+ that are TRPV1+: 49.8%

Figure 6

Figure 7

Figure 8

Target	Immunogen	RRID	Host	Supplier	Cat #	Dilution Factor
CELF4	MYIKMATLANGQA DNASLSTNGLGSSPG SAGHMNGLSHSPG NPSTIPMKDH	AB_10673035	Rabbit	Millipore Sigma	HPA037986	1:800
TRPV1	YTGSLKPEDAEVFKD SMVPGEK	AB_1624142	Guinea Pig	Neuromics	GP14100	1:500
NF-H	NF-H purified from bovine spinal cord	AB_2149761	Chicken	EnCor	CPCA-NF-H	1:40000
CGRP	ACDTATCVTHRLAGL LSRSGGVVKNNFVPT NVGSKAF	AB_1282813	Guinea Pig	Fitzgerald Industries International	20R-CP007	1:1500

Table 1: Primary antisera used for this study

Gene Ontology (GO) Term	3'UTR-binding proteins (100)	Detected in DRG neurons (84)	Enriched in DRG neuron population(s) (17)
posttranscriptional regulation of gene expression (GO:0010608)	8	8	1
regulation of mRNA metabolic process (GO:1903311)	33	32	8
regulation of translation (GO:0006417)	43	35	4
regulation of mRNA catabolic process (GO:0061013)	33	11	
regulation of mRNA stability (GO:0043488)	39	33	5
mRNA processing (GO:0006397)	45	43	10
RNA splicing (GO:0008380)	32	31	10
RNA localization (GO:0006403)	5	4	
regulation of gene silencing by miRNA (GO:0060964)	10	9	
RNA transport (GO:0050658)	10	7	1
cytoplasmic ribonucleoprotein granule (GO:0036464)	34	28	4
growth cone (GO:0030426)	6	6	2
dendritic spine (GO:0043197)	5	5	1
ribosome (GO:0005840)	6	6	2
distal axon (GO:0150034)	4	4	
nucleolus (GO:0005730)	16	16	1
dendrite (GO:0030425)	13	12	2
postsynaptic density (GO:0014069)	7	7	1
nuclear body (GO:0016604)	13	13	2
neuronal cell body (GO:0043025)	10	10	3
postsynapse (GO:0098794)	8	8	2

Table 2: Function of 3'UTR mRNA-binding proteins in populations of adult mouse sensory (DRG) neurons. 100 mouse genes are assigned the "3'UTR-binding" ontology term (Ashburner et al., 2000). mRNA's for 84 of these were detected in publicly available scRNA-Seq data derived from adult mouse DRG neurons (GSE59739, (Usoskin et al., 2015). 17 were significantly differentially expressed (FDR<0.05) between one or more of 4 DRG neuron populations (tyrosine hydroxylase–expressing, peptidergic, non-peptidergic, myelinated).

Gene	Population	Average Enrichment	p
CELF4	PEP	28.11	4.07E-12
CARHSP1	NP	15.76	8.95E-63
RBMS3	TH	5.52	4.68E-33
AUH	NF	3.64	1.67E-35
CELF6	PEP	2.98	2.25E-06
CELF6	NP	2.94	1.86E-20
CELF2	PEP	2.58	0.002704
CELF3	TH	2.53	1.35E-21
HNRNPA2B1	TH	2.21	1.65E-10
FXR2	NF	2.02	1.24E-14
ELAVL4	TH	2.02	5.72E-13
CSDC2	NF	1.93	1.37E-25
FXR2	NP	1.92	2.30E-05
RBMS1	NP	1.88	2.24E-13
RBFOX1	TH	1.78	0.000245
FXR2	PEP	0.56	0.006902
AUH	NP	0.46	0.003118
YBX3	NF	0.45	0.007494
TAF15	NF	0.40	2.32E-15
CELF6	NF	0.36	1.05E-07
CELF4	TH	0.31	1.38E-05
RBMS3	NP	0.29	3.77E-05
HNRNPL	NF	0.23	4.45E-05
RPS7	NF	0.20	0.000137
CARHSP1	NF	0.18	0.000383
CARHSP1	TH	0.17	3.98E-09

Table 3: 3'UTR mRNA-binding proteins significantly enriched (FDR<0.05) in DRG neuron populations, sorted by average fold enrichment. TH = tyrosine hydroxylase–expressing, NP = non-peptidergic, NF = neurofilament heavy chain-expressing and PEP = peptidergic neurons.

Family member	Tissue Expression (Adult)	Subcellular Location	RNA-binding motif	Molecular/biological function(s)
CELF1	Brain (m), spleen (m),liver(m), testis(m) ubiquitous(h)	Nuclear and cytoplasmic	UGUGUG,UGUU, CUGUCUG,UUGUG,	splicing,RNA stability, translation regulation (1-5)
CELF2	Brain (m), spleen (m),heart(m), lung(m),testis(m), ubiquitous(h)	Nucleoplasm, vesicles	UGUGU, UUGUU,UGUU,	RNA binding, mRNA processing,splicing
CELF3	Brain (m), adrenal gland (h),retina (h), pancreas (h)	Nucleus, cytoplasm	UGUUGUG (h)	RNA binding,mRNA splicing,spermatogenesis
CELF4	Brain (m,h), adrenal gland (h), retina (h)	Nucleus, cytoplasm, Axons/Dendrites	UGUGUKK(h), UGUGUGU*	RNA binding, mRNA processing, mRNA splicing
CELF5	Brain (m), N/A(h)	Nucleus, cytoplasm	UGUGUKK (h), UGUGUGU*	RNA binding
CELF6	Brain (m), N/A (h)	Nucleus, cytoplasm, synapse	UGUGKKG (h)	RNA binding, mRNA processing

Table 4: **CELF family RNA-binding protein tissue distribution, subcellular location, RNA recognition motifs and summarized function**. Tissue expression data from ProteomicsDB (Samaras et al., 2020; Schmidt et al., 2018). RNA-binding motifs were obtained from Transite (Krismer et al., 2020) and ATtRACT (Giudice et al., 2016) databases. * reported consensus sequence