1 **Grizzly bear toe amputation due to seasonal overlap with furbearer trapping in southeast**
2 **British Columbia**
3

4 Clayton Lamb, Laura Smit, Bruce McLellan, Luke Vander Vennen, Garth Mowat, Michael
5 Proctor

6
7
8
9 **Abstract**
10

Science and adaptive management form crucial components of the North American
model of wildlife management. Under this model, wildlife managers are encouraged to update
management approaches when new information arises whose implementation could improve the
stewardship and viability of wildlife populations and the well-being of animals. Here we detail a
troubling observation of toe amputations from several grizzly bears in southeast British
Columbia and assemble evidence to inform immediate action to remedy the issue. During the
capture of 59 grizzly bears, we noticed that four individuals (~7%) were missing some or all
their toes on one of their front feet. The wounds were all well healed and linear in nature. Further
opportunistic record collection revealed that this pattern of missing toes occurred beyond our
study area, and that furbearer traps were responsible for toe loss. We documented a problematic
seasonal overlap between the active season for grizzly bears and the fall trapping seasons for
small furbearers with body grip traps and for wolves with leghold traps. Instead of opening these
trapping seasons on or prior to November 1, when more than 50% of bears are still active, we
recommend delaying the start of these seasons until December 1, when most bears have denned.
Innovative solutions, such as narrowing trap entrances to exclude bear feet while still allowing
entrance of target furbearers, have the potential to minimize accidental capture of bears, but the
effectiveness of these approaches is unknown. Solutions that do not involve season changes will
require monitoring of efficacy and compliance to ensure success.

29
30
31 **Identifying an Issue**
32
Grizzly bears (*Ursus arctos*) are wide-ranging mammals that are highly motivated to

rapidly ingest high-energy foods in preparation for denning (McLellan 2011). This is especially

true for females who give birth and lactate in the den. In the Rocky Mountains, grizzly bears

occupy large home ranges (200-1300 km^2) where they search for mates and seasonal foods

(Graham and Stenhouse 2014). Key natural foods for grizzly bears include high calorie fruits,

abundant herbaceous vegetation, ungulates (killed or scavenged), ants, and roots (McLellan and

Hovey 1995, Munro et al. 2006). In addition to these natural foods, grizzly bears are well known

to take advantage of poorly managed human foods such as residential fruit trees, garbage,

roadkill, grain, and livestock (Craighead and Craighead 1972, Lamb et al. 2017, 2019,

Morehouse et al. 2020). This food-motivated behaviour is generally adaptive and allows grizzly

bears to increase their weight by about 35% in the 6-8 months of non-denning season (McLellan

2011), but coexistence with bears can be challenging when bears are attracted to, and allowed to

feed on, human-sourced foods (Lamb et al. 2020).

The challenges to human-bear coexistence are exacerbated when high-density grizzly

bear populations co-occur in human settled areas with abundant natural foods and accessible

human-sourced foods. One area that fits this description well is the southeast corner of British

Columbia (BC), Canada. This region is home to high-density grizzly bear populations,

productive habitat, and growing rural and urban communities. In response to observed grizzly

bear population declines, ecological trap dynamics, and high mortality rates in this area (Lamb et

al. 2017, 2019, 2020), we initiated a radio collaring project to better understand sources of

grizzly bear mortality and examine how animals' use of the landscape influenced population

dynamics. Between 2016 and 2020, we captured and radio collared 59 grizzly bears in the Elk

Valley (near Fernie, BC, see Fig 1A). Here we focus on a particular observation from this live

capture work that has important ramifications: several of the bears were missing toes. This

observation is important because ethics and values have a major role in wildlife management in

British Columbia and resulted in closing a decades-long sustainable grizzly bear hunt in 2018,

despite a large and viable population (Auditor General 2017, McLellan et al. 2017, Hatter et al.

2018).

Like humans, grizzly bears have twenty digits. While capturing bears, we noticed that

multiple bears were missing some or all the toes on one of their front feet (Fig 2). Of the 59

individual bears captured, 4 (~7%) were missing digits. In all cases, the toes lost were adjacent

and terminated in a linear plane, and the amputation had healed. At the time of their capture,

these animals were all 12 years old or younger (ages: 12, 11, 8, and 4). Three of the four bears

we captured with missing toes were male, while the larger collar sample had slightly more

females (n=31) than males (n=28). The female with missing toes was captured with a yearling

cub, suggesting that missing toes did not preclude reproduction. Based on radio collar telemetry

data, these animals moved in a similar fashion to bears with all their digits. While the loss of

digits did not appear to impact their locomotion, it may have influenced conflict behaviour,

which was common in these animals; three of the four animals that were missing toes were

involved in human-bear conflicts. One of these four bears was killed shortly after capture while

breaking into a rancher's calf pen, another was likely involved in an attack on a human, and a

third was captured by conservation officers in late fall after complaints of bear issues on a farm.

While the sample size for this conflict behaviour is small, less than 30% of the toe-bearing

sample (n=55) died or were involved in conflict during the time they were collared. While we

had identified a concerning trend of missing bear digits, the healed wounds left us little evidence

of the cause of the lost toes.

Searching for Cause

Southern British Columbia is fortunate to have a long history of people interacting with

grizzly bears through grizzly bear research, human-wildlife conflict work, hunting, guide

outfitting, and trapping. After identifying the missing toe issue, we spoke with First Nations,

scientists, conservation officers, wildlife managers, guide outfitters, and trappers, all with

lifetimes of experience on this land. A common topic in these discussions was grizzly bears

being incidentally caught in baited traps set for furbearers, particularly in traps set for marten

(*Martes americana*) and wolves (*Canis lupus*).

We knew that grizzly bears could become stuck in leghold traps set for wolves, given that

in recent years several bears had either been killed in, or required release from, wolf traps in

southern BC; however, we were less familiar with the role that body grip traps could play. Body

grip traps are killing traps designed for furbearers such as marten, skunk (*Mephitis mephitis*),

fisher (*Pekania pennanti*), weasel (*Mustela sp.*), muskrat (*Ondatra zibethicus*), beaver (*Castor*

canadensis), bobcat (*Lynx rufus*), and lynx (*Lynx canadensis*) (Fig 2). Trapping legislation in BC

allows for the use of a variety of trap models made by different manufacturers, provided the

devices reliably and humanely kill their target species (Province of British Columbia 2021). In

our region of interest (Fig 1), small body grip traps (commonly referred to as 120 conibear traps)

are most often used to capture marten and weasels, and they are typically set at the mouth of a

baited wooden box that is affixed to a tree at a height of approximately 1.5 meters from the

ground.

In the spring of 2018, we deployed four body grip traps, complete with their boxes and

bait (usually a small piece of meat), on trees in grizzly bear habitat in the Elk Valley. We wired

the traps in such a way that they could fire but not close and trap the bears. We monitored the

traps with remote cameras for approximately two weeks to see if bears would visit and spring the

traps. Grizzly bears visited all four traps and sprung two of them. Pictures and videos showed

bears investigating the traps and manipulating the boxes with their paws (Fig 3). Video showed

that one of the traps was set off with a bear's nose, but the cause of the other trap being set off

could not be determined because we only had pictures, not video, of bears investigating the trap.

Although this was a small trial with low sample sizes, younger bears seemed to investigate and

set off traps more often than older bears. Even with the small sample, it was clear that baited

traps attracted bears and that bears set off the traps as they tried to get the bait. We heard from

some trappers that they voluntarily delayed the start of their marten trapping season (which

legally opens November 1, Province of British Columbia 2021) to avoid having bears wreck

their trap sets, further confirming that bears are attracted to these common sets.

We became more convinced that furbearer traps were the cause of the missing toes as we

accumulated anecdotes and photo evidence from the area of grizzly bear feet stuck in traps (Fig

2). Photos came from hunters, remote cameras on ranches, and conservation officers, and they all

suggested the same mechanism – a body grip trap was affixed just behind the toes and

amputation of the toes was underway. Beyond identifying a likely cause for the missing toes, the

photo evidence and information from people out on the land revealed that bears with traps stuck

on their feet and missing toes was an issue beyond the extent of our Elk Valley study. In addition

to the records of body grip traps on bears' feet in southeast BC (also known as the "Kootenays"),

photos of a grizzly bear with a body grip trap on its foot were reported in Wyoming, USA in

2017 (McKim 2017), and brown bears with body grip traps on their feet are observed almost

every year in Finland (Kai-Eerik Nyholm pers. comm.).

We were also aware of multiple reports of grizzly bears being caught in leghold traps set

for wolves, and we believe this is another possible source of toe loss. Between 2010 and 2020, at

least five grizzly bears in southern BC were caught in wolf leghold traps (with the trap often

closing right behind the toes) and had to be released by conservation officers and scientists.

However, in the records we were able to collect, body grip traps were the most common trap on

bears' feet where it was clear that the trap was the cause of toe loss. In one case, it is suspected

that a bear's foot became stuck in a body grip trap set on the ground for skunks on private

property, but given the near ubiquity of marten trapping with box traps across our area, and the

broad occurrence of missing toes and traps on feet, we focus on marten and weasel trapping as

likely sources of the majority of the body grip traps that end up on bear feet.

We compiled data from collaring projects in adjacent areas to assess if a broader pattern

of missing toes could be revealed. Collaring projects in the Selkirk and Purcell Mountains (near

Nelson and Creston, Fig 1A) between 2004 and 2018, and in the Flathead Valley (southeast of

Fernie, Fig 1A) between 1978 and 2020, captured 72 and 206 grizzly bears respectively. Of the

20 bears captured in the Purcell Mountains, one adult male bear (5%) fit the pattern of having

well-healed missing toes, while none of the 52 bears captured in the Selkirk Mountains had

missing toes. However, grizzly bears were accidently captured by trappers in leghold traps set for

wolves on at least three occasions during the study in the Selkirk and Purcell Mountains, but no

evidence of toe loss due to incidental grizzly bear capture in legholds was reported. None of the

bears captured in the Flathead Valley were known to be missing toes; however, a hunter killed a

bear just north of the Flathead study area with a body grip trap on its foot (Fig 2E). Results from

these adjacent studies did not reveal any general insights into the observed spatial distribution of

missing toes. Such a pattern is not surprising given that traplines in British Columbia are distinct

areas trapped exclusively by registered trappers. Variation in trapping intensity, target species,

and timing creates variable exposure of bears to body grip traps across the landscape.

Additionally, unsecured attractants such as apple trees, garbage, roadkill, and livestock may

increase bear use of low elevation areas and delay den entry perhaps increasing the chance of

overlap with trapping activities in these areas.

Observations of bears stuck in body grip traps and bears with missing toes have mainly

been documented in the past 15 years and are not restricted to a single area or trapline, but rather

occur broadly at low frequency across the Kootenays and beyond. This increased prevalence

could be related to an increase in grizzly bear research and monitoring; however, opportunities to

identify these issues have been abundant for several decades as many bears were captured by

conservation officers, harvested by hunters, and monitored for research purposes in the Flathead.

It is possible that in recent years the increased use of stronger traps, as mandated by evolving

humane trapping standards, has created a situation where bears now have more difficulty

extracting themselves from body grip traps when captured.

We summarized the den entry and exit dates for 61 animal-years in the Elk Valley to

assess when bears were denning in the fall. We found the median den entry date (i.e. the date

when 50% of bears had denned) was November 6, and the 95th quantile of den entry was

November 22, revealing a critical 3-week period of overlap between active bears and the

trapping season. We found no overlap between den emergence and the end of the trapping

season; thus, we focus on fall trapping as the source of the issue. More support for the idea that

the concurrence of the fall active season for bears and the beginning of the trapping season is the

main issue, is that we have not seen similar problems of missing toes in black bears, who den

several weeks earlier than grizzly bears and would thus be less exposed to traps.

Honing a Solution

Most trapping of furbearers is done in winter when underfur is dense and guard hairs are

long. Given that grizzly bears hibernate for the winter, trappers should generally be able to avoid

accidently catching bears. However, some trapping seasons open in late fall, such as the marten

season in much of British Columbia which begins November 1 and extends to February 15.

Reducing the overlap between the bear active season and the trapping season is one way to

minimize the loss of bear toes and prevent trappers from losing their traps and having their sets

destroyed by bears. Shifting the start of most trapping that coincides with the active bear season

from November 1 to December 1 would essentially eliminate overlap between trapping and

bears. This solution has been previously used in southeast British Columbia to avoid catching

and killing bears in neck snares set for wolves, an issue first documented in the Flathead project.

Indeed, such a change to the regulations to address the missing toe issue was proposed in

2019 but was not implemented. Much of the trapped landscape in the Kootenays is rugged,

mountainous terrain, and many trappers, especially in the west Kootenays, reported that they

were effectively unable to access their traplines for much of the winter beginning in early

December due to high avalanche risk and associated safety concerns. The trappers stated they did

most of their trapping in November and early December, and that shortening the season by

delaying the opening date to December 1 would largely remove their ability to harvest marten

from their trapline. As a result, provincial biologists implemented a condition on all active

trapping licenses in the Kootenay region stipulating that, beginning in 2021, all body grip traps

set for marten during the month of November must be enclosed in a box with an opening no

larger than 3 ½ inches. This constricted entrance was thought to be narrower than most bear

paws. The updated licenses also recommended that, prior to December 1, trappers should use

these same boxes to enclose similarly sized killing traps set for other species on dry ground. The

efficacy of the modified enclosure entrance in eliminating bear toe loss has not yet been tested,

but these modified marten trap boxes are believed to be sufficient to ensure grizzly bears are not

able to access a set trap. Given there is more uncertainty in the effectiveness of the modified

enclosure approach than there is in delaying the start of the trapping season, we recommend that

compliance and efficacy is monitored to ensure this intervention is effective and uptake is high.

Such monitoring would allow for adaptive management and changes to the approach as needed

to ensure a successful outcome for bears and trappers alike.

Trappers have a long history of supporting conservation efforts and are sentinels of

changing landscapes and wildlife populations. However, trapping of furbearers in British

Columbia has become an increasingly vulnerable policy amidst shifting sociopolitical trends in

society. Therefore, bold and novel strategies will be required from trapping organizations to

maintain public support while also recruiting and retaining the next generation of trappers. A

tactful strategy to uphold public support is to proactively assess and react to emerging evidence

that threatens social acceptance of trapping (e.g., grizzly bears missing toes), and where changes

could help increase the well-being of animals. This approach is the backbone of the North

American Model, and while emerging evidence or science can sometimes conclude no direct

threat to wildlife populations — organizations must equally weigh social ramifications and

understand that the future of trapping may be predicated more on social support than consensus

over sustainability. Here we document a troubling issue where grizzly bears are facing toe

amputation after becoming stuck in baited furbearer traps. The timing of furbearer seasons is the

primary issue, with trapping seasons opening weeks before all bears have denned, which creates

a problematic period of overlap between active bears and baited traps. We provide evidence of

the issue, solutions to remedy it, and ultimately urge immediate action.

Data Availability

All analyses and supporting data can be found at https://github.com/ctlamb/Grizzly-MissingToes

Acknowledgements

This work was conducted within ?amak?is Ktunaxa, the homelands of Ktunaxa people. We thank

the Ktunaxa Nation for their support of our k☐aw☐a work. Thanks to the trappers, guide

outfitters, hunters, First Nations, and scientists that discussed this work with us and helped us

identify the issue and explore solutions. Thanks to veterinarians Steven Chapman and Cali Lewis

for X-rays of EVGM55's foot, interpretation of these images, and discussion on potential causes.

Thanks to Patrick Stent for support of our research project and many helpful discussions as we

worked towards a solution to this issue. Thanks to Adam Ford and Melanie Dickie for

suggestions on displaying results, and to Mateen Hessami for suggestions on improving the

manuscript text. These data were collected with support from the Government of British

Columbia, Habitat Conservation Trust Foundation, Vanier Canada Graduate Scholarship, Liber

Ero Fellowship Program, National Science Engineering and Research Council, Fish and Wildlife

Compensation Program, Forest Enhancement Society of British Columbia, Teck Coal, Columbia

Basin Trust, Counter Assault, Wildsight, Nature Conservancy of Canada, Wildlife Conservation

Society, Yellowstone to Yukon Conservation Initiative, Safari Club International, Sparwood and

District Fish and Wildlife Association, Elkford Rod and Gun Club, Ministry of Transport and

Infrastructure, Outdoor Research, Patagonia, and the British Columbia Conservation Officer

Service.

References

Auditor General, B. C. 2017. An Independent Audit of Grizzly Bear Management | Auditor
General of British Columbia. Victoria.
Craighead, J. J., and F. C. Craighead. 1972. Grizzly Bear-Man Relationships in Yellowstone
National Park. Bears: Their Biology and Management 2:304.
Graham, K., and G. Stenhouse. 2014. Home Range, Movements, and Denning Chronology of the
Grizzly Bear (Ursus arctos) in West-Central Alberta. The Canadian Field-Naturalist
128:223–234. <http://www.canadianfieldnaturalist.ca/cfn/index.php/cfn/article/view/1600>.
Accessed 11 Dec 2014.
Hatter, I. W., G. Mowat, and B. N. McLellan. 2018. Statistical population reconstruction to
evaluate grizzly bear trends in British Columbia, Canada. Ursus 29:1.
Lamb, C., A. Ford, B. McLellan, M. Proctor, G. Mowat, L. Ciarniello, S. Nielsen, and S. Boutin.
2020. The ecology of human-carnivore coexistence. Proceedings of the National Academy
of Science.
Lamb, C. T., A. T. Ford, M. F. Proctor, J. A. Royle, G. Mowat, and S. Boutin. 2019. Genetic
tagging in the Anthropocene: scaling ecology from alleles to ecosystems. Ecological
Applications 29:e01876. <https://onlinelibrary.wiley.com/doi/abs/10.1002/eap.1876>.
Lamb, C. T., G. Mowat, B. McLellan, S. Nielsen, and S. Boutin. 2017. Forbidden Fruit: Human
settlement and abundant fruit create an ecological trap for grizzly bears. Journal of Animal
Ecology 86:55–65.
McKim, C. 2017. Grizzly Bear With Trap On Its Foot Still Not Found. Wyoming Public
Radio2017:1. Wyoming. <https://www.wyomingpublicmedia.org/natural-resources-
energy/2017-06-26/grizzly-bear-with-trap-on-its-foot-still-not-found>.
McLellan, B. 2011. Implications of a high-energy and low-protein diet on the body composition,
fitness, and competitive abilities of black (Ursus americanus) and grizzly (Ursus arctos).
Canadian Journal of Zoology 558:546–558.
<http://www.nrcresearchpress.com/doi/abs/10.1139/z11-026>. Accessed 18 Feb 2014.
McLellan, B., and F. Hovey. 1995. The diet of grizzly bears in the Flathead River drainage of
southeastern British Columbia. Canadian Journal of Zoology 73:704–712.
<http://www.nrcresearchpress.com/doi/pdf/10.1139/z95-082>. Accessed 9 Apr 2013.
McLellan, B. N., G. Mowat, T. Hamilton, and I. Hatter. 2017. Sustainability of the Grizzly Bear
Hunt in British Columbia, Canada. Journal of Wildlife Management 81:218–229.
Morehouse, A. T., C. Hughes, N. Manners, J. Bectell, and T. Bruder. 2020. Carnivores and
Communities: A Case Study of Human-Carnivore Conflict Mitigation in Southwestern
Alberta. Frontiers in Ecology and Evolution 8:1–15.
Munro, R. H. M., S. E. Nielsen, M. H. Price, G. B. Stenhouse, and M. S. Boyce. 2006. Seasonal
and Diel Patterns of Grizzly Bear Diet and Activity in West-Central Alberta. Journal of
Mammalogy 87:1112–1121. <http://asmjournals.org/doi/abs/10.1644/05-MAMM-A-
410R3.1>. Accessed 19 Feb 2014.
Province of British Columbia. 2021. British Columbia Hunting and Trapping Regulations
Synopsis.
288

A) Location of documented missing toes and by-catch

Trap ● body grip ▲ foothold ■ unknown   Missing toes ● N ● Y

B) Timing of grizzly bear activity and trapping season

Figure 1. A) Locations of observations of bears with missing toes and bears with traps on their feet. Detections made through research are outlined in grey, while detections opportunistically gathered throughout southeast British Columbia are not outlined in grey. This area is part of the homelands for the Ktunaxa Nation, and is also referred to as the Kootenays. B) The grizzly bear active season, defined from 61 collared bear den emergence and entrance dates, with the marten trapping season (November 1 to February 15) shown in grey. The 95[th] percentile of den entrance dates was November 22. We suggest that December 1 represents an evidence-based start date for trapping to reduce conflicts between bears and trappers.

Figure 2. Animals detected with missing toes through research captures in the Elk Valley (A-D), and opportunistic evidence of grizzly bears with traps on their feet (E-G). Each letter in A-D represents one of the four unique individuals detected with missing toes. Ci shows an X-ray of the foot of EVGM55, who was killed in a conflict with a cattle rancher. E) a body grip trap on the foot of a hunter-killed grizzly bear from the Flathead Valley, F) the paw of a 1-year-old grizzly bear stuck in a body grip trap captured by conservation officers near Pass Creek due to conflict behaviour from a mom and offspring; significant infection was present, and the toes had nearly completely separated from the foot by the time of capture, and G) a grizzly bear with a body grip trap on its foot detected feeding on a cow carcass by a remote camera west of Invermere.

310
311 Figure 3. Grizzly bears detected at trial marten body grip traps. The traps were wired so they
312 could spring but not close on bears' feet, and they were set for two weeks in areas frequented by
313 bears. Investigatory behaviour was common, and two of the four deployments were set off.