1   22 Oct 2021
2   Clayton Lamb
3   University of British Columbia
4   Kelowna, British Columbia, Canada, *V1V 1V7*
5   ctlamb@ualberta.ca
6

**Considerations for furbearer trapping regulations to prevent grizzly bear toe amputation**

**and injury**

CLAYTON LAMB,[1] *University of British Columbia, Department of Biology, Kelowna, BC,*

*Canada V1V 1V7*

LAURA SMIT, *Ministry of Forests, Lands, Natural Resource Operations and Rural*

*Development, Nelson, BC, Canada V1L 4K3*

BRUCE MCLELLAN, *International Union for the Conservation of Nature Bear Specialist*

*Group, D'Arcy, BC, Canada V0N 1L0*

LUCAS M. VANDER VENNEN, *Ministry of Forests, Lands, and Natural Resource Operations*

*and Rural Development, Nelson, BC, Canada V1L 4K*

MICHAEL PROCTOR, *International Union for the Conservation of Nature Bear Specialist*

*Group, Kaslo, BC, Canada V0G 1M0*

[1] *Email:* ctlamb@ualberta.ca

**ABSTRACT** Science and adaptive management form crucial components of the North

American model of wildlife management. Under this model, wildlife managers are encouraged to

update management approaches when new information arises whose implementation could

improve the stewardship and viability of wildlife populations and the well-being of animals.

Here we detail a troubling observation of several grizzly bears with amputated toes in southeast

British Columbia and assemble evidence to inform management strategies to remedy the issue.

During the capture of 59 grizzly bears, we noticed that four individuals (~7%) had amputated

toes on one of their front feet. The wounds were all healed and linear in nature. Further

opportunistic record collection revealed that similar examples of amputated toes occurred

beyond our study area, and that furbearer traps were frequently responsible for toe loss. We

found evidence that seasonal overlap between the active season for grizzly bears and the fall

trapping seasons for small furbearers with body grip traps and for wolves with foothold traps

may explain the issue. Multiple options to reduce or eliminate this issue exist, but have varying

degrees of expected efficacy and require differing levels of monitoring. The most certain

approach to greatly reduce the issue is to delay the start of the marten trapping season until

December 1, when most bears have denned, instead of opening the season on or prior to

November 1, when more than 50% of bears are still active. Additionally, innovative solutions,

such as narrowing trap entrances to exclude bear feet while still allowing entrance of target

furbearers, have the potential to minimize accidental capture of bears, but the effectiveness of

these approaches is unknown. Experimental evidence suggested that better anchoring traps was

not a viable solution. Solutions that do not involve season changes will require monitoring of

efficacy and compliance to ensure success.

**KEY WORDS**: Amputation, adaptive management, body-gripping trap, grizzly bear, leghold

trap, foothold trap, furbearer, bycatch, incidental catch, animal welfare

**IDENTIFYING AN ISSUE**

Grizzly bears (*Ursus arctos*) are wide-ranging mammals that are highly motivated to

rapidly ingest high-energy foods in preparation for denning (McLellan 2011). In the Rocky

Mountains of southern Canada, grizzly bears occupy large home ranges (200-1300 km$^2$) where

they search for mates and seasonal foods (Graham and Stenhouse 2014). Key natural foods for

grizzly bears include high-calorie fruits, abundant herbaceous vegetation, ungulates (killed or

scavenged), ants, and roots (McLellan and Hovey 1995, Munro et al. 2006). In addition to these

natural foods, grizzly bears are well known to take advantage of poorly managed human foods

such as residential fruit trees, garbage, roadkill, grain, and livestock (Craighead and Craighead

1972, Lamb et al. 2017, 2019, Morehouse et al. 2020). This food-motivated behaviour is

generally adaptive and allows grizzly bears to increase their weight by about 35% in the 6-8

months of non-denning season (McLellan 2011), but bears can also be attracted to, and allowed

to feed on, human-sourced foods which can create human and bear safety issues (Lamb et al.

2020).

The challenges to human-bear coexistence are exacerbated when high-density grizzly

bear populations co-occur in human settled areas with abundant natural foods and accessible

human-sourced foods. One area that fits this description well is the southeast corner of British

Columbia (BC), Canada. This region is home to high-density grizzly bear populations,

productive habitat, and growing rural and urban communities. In response to recent grizzly bear

population declines, ecological trap dynamics, and high human-caused mortality rates in this area

(Lamb et al. 2017, 2019, 2020), we initiated a radio collaring project to better understand sources

of grizzly bear mortality and examine how animals' use of the landscape influenced population

dynamics. Between 2016 and 2020, we captured and radio collared 59 grizzly bears in the Elk

Valley (near Fernie, BC, see Fig 1A). Captures were in accordance with University of Alberta

Animal Ethics #AUP00002181 and Province of British Columbia Capture Permit #CB17-

264200. Here we focus on a particular observation from this live capture work that has important

ramifications: several of the bears had amputated toes. That is the toes appeared to have been cut

off and then healed. This observation is important because ethics and values have a major role in

wildlife management in British Columbia and resulted in closing a decades-long sustainable

grizzly bear hunt in 2018, despite a large and viable population (Auditor General 2017, McLellan

et al. 2017, Hatter et al. 2018). Identifying how these toes were amputated and mitigating the

source of amputation became one of the objectives of our study.

Like humans, grizzly bears have twenty digits. Of the 59 individual bears captured, 4

(~7%) were missing some of the toes on one of their front feet (Fig 2). In all cases, the toes lost

were adjacent and terminated in a linear plane, and the injury had healed. At the time of their

capture, these animals were all between the ages of 4 and 12 years old. Three of the four

captured bears with amputated toes were male, while the larger collar sample had slightly more

females (n=31) than males (n=28). The female with amputated toes was captured with a yearling

cub, suggesting that amputated toes did not preclude reproduction. Based on radio collar

telemetry data, these animals moved in a similar fashion to bears with all their digits. While toe

loss did not appear to impact their locomotion, it may have influenced conflict behaviour, which

was common in these animals; three of the four animals with amputated toes were involved in

human-bear conflicts. One of these four bears was killed shortly after capture while breaking into

a rancher's calf pen, another was likely involved in an attack on a human, and a third was

captured by conservation officers in late fall after complaints of bear issues on a farm. While the

sample size for this conflict behaviour is small, less than one third of the bears with all their toes

(n=55) died or were involved in conflict during the time they were collared. The reason for

elevated conflict behaviour observed in the bears with amputated toes could be related to

challenges feeding on natural foods at certain times of year, such as digging roots during spring

and fall (McLellan and Hovey 1995), or these bears could be naturally bolder individuals and

thus more likely to become both trapped and involved in future conflict. Alternatively, the

pattern of increased conflict could be an artefact of the small sample sizes we had available

(n=4). Nevertheless, while we had identified a concerning trend of amputated bear toes but the

healed wounds left us little evidence for the cause of the toe loss.

**SEARCHING FOR CAUSE**

We first consulted with veterinary professionals to determine if the amputated toes could

be congenital (i.e., lost before birth and due to natural causes). After review of multiple photos

and X-rays (Fig 2) there was little evidence to suggest that this was congenital. On the contrary,

strong evidence for injury post-birth included clear evidence of toe bone fracture and healing,

and consistent linear wounds suggesting amputation related to human sources. As a result, we

proceeded to search for a human-caused source of the issue.

Southern British Columbia is fortunate to have a long history of people interacting with

grizzly bears through research, human-wildlife conflict work, hunting, guide outfitting, trapping,

and simply living in bear country. After identifying the amputated toe issue, we spoke with First

Nations, scientists, conservation officers, wildlife managers, guide outfitters, and trappers, all

with lifetimes of experience on this land. A common topic in these discussions was grizzly bears

being incidentally caught in baited traps set for furbearers, particularly in traps set for marten

(*Martes americana*) and wolves (*Canis lupus*). Indeed, non-target captures stemming from legal

trapping activities have been reported for many species. Cougars (*Puma concolor*) have been

incidentally captured in foothold traps or snares set for wolves and bobcat (*Lynx rufus*) (Knopff

et al. 2010, Andreasen et al. 2018), and free-ranging cats and dogs and even birds have been

incidentally caught in furbearer traps (White et al. 2021). We knew that grizzly bears could be

caught in foothold traps set for wolves, given that in recent years several bears had either been

killed in, or required release from, wolf traps in southern BC; however, we were less familiar

with the role that body grip traps could play in the loss of toes. Body grip traps are killing traps

designed for furbearers such as marten, skunk (*Mephitis mephitis*), fisher (*Pekania pennanti*),

weasel (*Mustela sp.),* muskrat (*Ondatra zibethicus*), beaver (*Castor canadensis*), bobcat, and

lynx (*Lynx canadensis*) (Fig 2). Trapping legislation in BC allows for the use of a variety of trap

models made by different manufacturers, provided the devices reliably and humanely kill their

target species (Province of British Columbia 2021, White et al. 2021). In our region of interest

(Fig 1), small body grip traps (commonly referred to as 120 conibear traps) are most often used

to capture marten and weasels, and they are typically set at the mouth of a baited wooden box

that is affixed to a tree at a height of approximately 1.5 meters from the ground.

In the spring of 2018, we deployed four body grip traps, complete with their boxes and

bait (usually a small piece of meat), on trees to see if bears would investigate these traps. We

wired the traps in such a way that they could fire but not close and trap the bears. We monitored

the traps with remote cameras for approximately two weeks to see if bears would visit and spring

the traps. Grizzly bears visited all four traps and sprung two of them. Pictures and videos showed

bears investigating the traps and manipulating the boxes with their paws (Fig 3). Video showed

that one of the traps was set off with a bear's nose, but the cause of the other trap being set off

could not be determined because we only had pictures, not video, of bears investigating the trap.

Although this was a small trial with low sample sizes, younger bears seemed to investigate and

set off traps more often than older bears. Even with the small sample, it was clear that baited

traps attracted bears and that bears set off the traps as they tried to get the bait. We heard from

some trappers that they voluntarily delayed the start of their marten trapping season (which

legally opens November 1, Province of British Columbia 2021) to avoid having bears wreck

their trap sets, further confirming that bears are attracted to these common sets.

        We became more convinced that furbearer traps were the cause of the amputated toes as

we accumulated anecdotes and photo evidence of grizzly bear feet stuck in traps (Fig 2). Photos

came from hunters, remote cameras on ranches, and conservation officers, and they all suggested

the same mechanism. A body grip trap was affixed just behind the toes and amputation was

underway. Beyond identifying a likely cause for the amputated toes, the photo evidence and

information from people out on the land revealed that bears with traps stuck on their feet and

amputated toes was an issue beyond the extent of our study area. In addition to the records of

body grip traps on bears' feet in southeast BC (also known as the "Kootenays"), photos of a

grizzly bear with a body grip trap on its foot were reported in Wyoming, USA in 2017 (McKim

2017), and brown bears with body grip traps on their feet are observed almost every year in

Finland (Kai-Eerik Nyholm pers. comm.).

        We were also aware of multiple reports of grizzly bears being caught in foothold traps set

for wolves, and we believe this is another possible source of toe loss. Between 2010 and 2020, at

least five grizzly bears in southern BC were caught in wolf foothold traps (with the trap often

closing right behind the toes) and had to be released by conservation officers and scientists.

However, in the records we were able to collect, body grip traps were the most common trap on

bears' feet where it was clear the trap was the cause of toe loss. In one case, it is suspected that a

bear's foot became stuck in a body grip trap set on the ground for skunks on private property.

However, given the near ubiquity of marten trapping with box traps across our area (an average

of 2,082 marten were harvested annually in the Kootenays from 2013-2017), and the broad

occurrence of amputated toes and traps on feet, we focus on marten and weasel trapping as likely

sources of most of the body grip traps that end up on bear feet.

We compiled data from collaring projects in adjacent areas within British Columbia to

assess if a broader pattern of toe loss could be revealed. Collaring projects in the Selkirk and

Purcell Mountains (near Nelson and Creston, Fig 1A) between 2004 and 2018, and in the

Flathead Valley (southeast of Fernie, Fig 1A) between 1978 and 2020, captured 72 and 206

grizzly bears respectively. Of the 20 bears captured in the Purcell Mountains, one adult male

bear (5%) fit the pattern of having well-healed but amputated toes, while none of the 52 bears

captured in the Selkirk Mountains had amputated toes. However, grizzly bears were accidently

captured by trappers in foothold traps set for wolves on at least three occasions during the study

in the Selkirk and Purcell Mountains, but no evidence of toe loss due to incidental grizzly bear

capture in footholds was reported. Likely because bears were either released from the traps or

killed. None of the bears captured in the Flathead Valley had amputated toes; however, a hunter

killed a bear just north of the Flathead study area with a body grip trap on its foot (Fig 2E).

Results from these adjacent studies did not reveal any general insights into the observed spatial

distribution of bears with amputated toes. This is not surprising given that traplines in British

Columbia are distinct areas trapped exclusively by registered trappers, and variation in trapping

intensity, target species, and timing creates variable exposure of bears to body grip traps across

the landscape. Additionally, unsecured attractants such as apple trees, garbage, roadkill, and

livestock may increase bear use of low elevation areas and delay den entry, perhaps increasing

the chance of overlap with trapping activities in these areas.

Observations of bears stuck in body grip traps and bears with amputated toes have mainly

been documented in the past 15 years and are not restricted to a single area or trapline, but rather

occur broadly at low frequency across the Kootenays and beyond. This increased detection of

bear toe loss could be related to an increase in grizzly bear research and monitoring; however,

opportunities to identify these issues have been abundant for several decades as many bears were

captured by conservation officers, harvested by hunters, and monitored for research purposes in

the Flathead. It is possible that in recent years the increased use of stronger traps, as mandated by

evolving humane trapping standards, has created a situation where bears now have more

difficulty extracting themselves from body grip traps when captured.

We summarized the den entry and exit dates for 61 animal-years in the Elk Valley to

assess when bears were denning in the fall. We found the median den entry date (i.e. the date

when 50% of bears had denned) was November 6, and the 95th quantile of den entry was

November 22, revealing a critical 3-week period of overlap between active bears and the

trapping season. We found no overlap between den emergence and the end of the trapping

season; thus, we focus on fall trapping as the source of the issue. More support for the idea that

the concurrence of the fall active season for bears and the beginning of the trapping season is the

main issue is that we have not seen similar problems of amputated toes in black bears, who den

several weeks earlier than grizzly bears and would thus be less exposed to traps.

**HONING A SOLUTION**

Most trapping of furbearers is done in winter when fur is prime (i.e., underfur is dense

and guard hairs are long) and most valuable. Given that grizzly bears hibernate for the winter,

trappers should generally be able to avoid accidently catching bears. However, some trapping

seasons open in late fall, such as the marten season in much of British Columbia which begins

November 1 and extends to February 15. Reducing the overlap between the period when bears

are active and the trapping season is open is one way to minimize the amputation of bear toes

and prevent trappers from losing their traps and having their sets destroyed by bears. Shifting the

start of most trapping that coincides with the active bear season from November 1 to December 1

would essentially eliminate overlap between trapping and bears. This solution has been

previously used in southeast British Columbia to avoid catching and killing bears in neck snares

set for wolves, an issue first documented in the Flathead project (McLellan 2015, Province of

British Columbia 2021).

Indeed, such a change to the rapping regulations to address the amputated toe issue was

proposed in 2019 but was not implemented. Much of the trapped landscape in the Kootenays is

rugged, mountainous terrain, and many trappers, especially in the west Kootenays, reported that

they were effectively unable to access their traplines for much of the winter beginning in early

December due to high avalanche risk and associated safety concerns. The trappers stated they did

most of their trapping in November and early December, and that shortening the season by

delaying the opening date to December 1 would largely remove their ability to harvest marten

from their trapline. However, this was not a ubiquitous issue because many trappers in the east

Kootenay reported voluntarily delaying their trapping until December to avoid bears, and other

users of the landscape such as backcountry skiers and snowmobilers recreate in the mountains

throughout the winter. Nevertheless, provincial biologists instead implemented a condition on all

active trapping licenses in the Kootenay region stipulating that, beginning in 2021, all body grip

traps set for marten during the month of November must be enclosed in a box with an opening no

larger than 3 ½ inches. This constricted entrance was thought to be narrower than most bear

paws. The license condition also recommended that, prior to December 1, trappers should use

these same boxes to enclose similarly sized killing traps set for other species on dry ground. The

efficacy of the modified enclosure entrance in eliminating bear toe loss has not yet been tested,

but these modified marten trap boxes are believed to be sufficient to ensure grizzly bears are not

able to access a set trap. Because there is more uncertainty in the effectiveness of the modified

enclosure approach than there is in delaying the start of the trapping season, we recommend that

compliance and efficacy is monitored to ensure this intervention is effective and uptake is high.

Such monitoring would allow for adaptive management and changes to the approach as needed

to ensure a successful outcome for bears and trappers alike.

245            Additional approaches have been suggested to help resolve the issue, but we do not

recommend their implementation. For example, one option is to sufficiently anchor traps so a

bear that is incidentally caught can hopefully pull its foot free. To address the viability of this

solution we conducted a small experiment using front appendages of deceased bears and four

types of traps approved for marten trapping in British Columbia (Province of British Columbia

2021). Our experiment consisted of three phases: the first used before and after X-rays to assess

whether traps could immediately break bear toe bones, the second assessed whether bears could

easily pull their feet free from traps (which we simulated by solidly anchoring the trap and

having an 86 kg person attempt to free the feet by pulling on the leg with maximum human pull),

and the third assessed how much pull (kg) was required to free the feet. We gathered the front

appendages from one adult and two cub grizzly bears killed in the Elk Valley during fall 2021

and tested four approved body grip traps: 1) Bélisle Super X 120, 2) Northwoods 155, 3)

Sauvageau 2001-5, and 4) Sauvageau 2001-6.

Results from our experiment suggested that traps do not immediately break bones, but

bears were unable to easily extract themselves from these traps. X-ray results confirmed that

setting traps off on adult or cub feet did not break or fracture any bones. Despite using maximum

human effort and both sustained and jerking pulls, we were only able to free adult grizzly feet

from traps 20% (95% CI: 7-33%) of the time, and cub feet 63% (95% CI: 53-72%) of the time.

For 80% of our attempts with adult feet and 37% of our attempts with cub feet, we were unable

to free the feet from the traps despite repeatedly jerking with as much force as we could muster.

Considerable variation in release efficacy was observed between trap models (Table 1), where

the Sauvageau traps generally released bears the least frequently, and Northwoods 155 released

bears the most. Finally, we used a come-along or a truck, with a scale affixed inline, to determine

the steady pull required to extract an adult foot from the Sauvageau traps. The average pull

required to free a foot was 164 kg (range=91-232 kg). Even after maximum pull was applied, we

did not detect superficial damage to toe bones or joints, although clearly significant pain would

be sustained by the bears during this time, and we are not certain that bears could always muster

the >230 kg pull required to ensure reliable release.

Based on the results of our experiment, we believe that better anchoring traps would

present safety concerns for bears, trappers, and wildlife professionals that might be called to

release bears. Although there is little work assessing the pull of a bear, a strong animal such as a

horse provides some insight into the forces animals can generate. A horse can exert ~70% of its

body weight at max exertion (Smith 1896). The adult female grizzly bear whose appendages we

used weighed approximately 140 kg, which would equate to a dead pull of ~98 kg if we use the

weight to exertion relationship from horses. This would be well below the maximum of >230 kg

required to reliably extract feet from these traps, and barely above the minimum recorded of 91

kg. As a result, bears may often be unable to escape from well anchored traps. There is a greater

chance that bears could free themselves if a long leash (>2 m) is used so bears could run and

generate more pull. It is thought that a 140 kg bear could generate between 400-1000 kg of pull

with a running start (Flaa et al. 2009), however, significant damage to bears' feet and traps is

possible in this case and we do not recommend this solution. Overall, our experiment suggests

that bears cannot easily extract their feet from traps and better anchoring traps presents

significant risk to people and bears. Furthermore, current regulations for trap check intervals on

killing traps such as body grips can be as long as 14 days in British Columbia and would require

significant reduction (<24 hours) if bears could become permanently held in traps. Such a change

in trap check intervals from 14 days to 24 hours would render most large winter traplines

impractical for trappers.

The most viable solution to the amputated toe issue requires that bears' feet do not enter

traps at all. Changing the season length is the most reliable solution, but possibly reduces trapper

access to trapping opportunities in portions of the Kootenays. Constricting the entrance size of

the trap presents a possible simple solution but requires further testing to ensure the design is

effective at excluding bears while not dissuading marten from entering. Modified entrance size

requirements will also require compliance monitoring, adding additional responsibilities to

already overtasked Conservation Officers. The solutions we present here have various pros and

cons, and we hope that this work can help policy makers make an informed decision and choose

a solution that will resolve the amputated toe issue while ensuring trappers have sufficient

opportunity to trap furbearers.

Trappers have a long history of supporting conservation efforts and are sentinels of

changing landscapes and wildlife populations. However, trapping of furbearers in British

Columbia has received increased scrutiny amidst shifting sociopolitical trends in society. Bold

and novel strategies will be required from trapping organizations and wildlife management

agencies to maintain public support while also recruiting and retaining the next generation of

trappers. A tactful strategy to uphold public support is to proactively assess and react to

emerging evidence that threatens social acceptance of trapping (e.g., grizzly bears with

amputated toes), and where changes could help increase the well-being of animals. This

approach is the backbone of the North American Model, and while emerging evidence or science

can sometimes conclude no direct threat to wildlife populations, organizations must equally

weigh social ramifications and understand that the future of trapping may be predicated more on

social support than consensus over sustainability. Here we document a troubling issue where

grizzly bears are losing their toes after becoming stuck in baited furbearer traps. The timing of

furbearer seasons is the primary issue, with trapping seasons opening weeks before all bears have

denned, which creates a problematic period of overlap between active bears and baited traps. We

provide evidence of the issue, solutions to remedy it, and ultimately urge immediate action.

**DATA AVAILABILITY**

All analyses and supporting data can be found at https://github.com/ctlamb/Grizzly-MissingToes

**ACKNOWLEDGEMENTS**

This work was conducted within ?amak?is Ktunaxa, the homelands of Ktunaxa people. We thank

the Ktunaxa Nation for their support of our kⱡawⱡa (grizzly bear) work. Thanks to the trappers,

guide outfitters, hunters, First Nations, and scientists that discussed this work with us, helped us

identify the issue, and explore solutions. A special thanks to trapper Blair Chatterson who lent us

traps, trapping supplies, and discussed this issue with us. Thanks to veterinarian Steven Chapman

for X-rays of EVGM55's foot, interpretation of these images, and discussion on potential causes.

Thanks to veterinarian Cali Lewis and Tanglefoot Veterinary Services for before and after X-

rays of bear feet for our experiment, interpretation of images, and many thoughtful discussions

on the causes of, and potential solutions to, this issue. Thanks to Emily Chow and Stewart Clow

for help setting traps and staunch efforts to free bear toes from these traps during the experiment.

Thanks to Patrick Stent for support of our research project and many helpful discussions as we

worked towards a solution to this issue. Thanks to Adam Ford and Melanie Dickie for

suggestions on displaying results. Thanks to Garth Mowat, Mateen Hessami, and Tim Killey for

suggestions on improving the manuscript. These data were collected with support from the

Government of British Columbia, Habitat Conservation Trust Foundation, Vanier Canada

Graduate Scholarship, Liber Ero Fellowship Program, National Science Engineering and

Research Council, Fish and Wildlife Compensation Program, Forest Enhancement Society of

British Columbia, Teck Coal, Columbia Basin Trust, Counter Assault, Wildsight, Nature

Conservancy of Canada, Wildlife Conservation Society, Yellowstone to Yukon Conservation

Initiative, Safari Club International, Sparwood and District Fish and Wildlife Association,

Elkford Rod and Gun Club, Ministry of Transport and Infrastructure, Outdoor Research,

Patagonia, and the British Columbia Conservation Officer Service.

**REFERENCES**

Andreasen, A. M., K. M. Stewart, J. S. Sedinger, C. W. Lackey, and J. P. Beckmann. 2018.

Survival of cougars caught in non-target foothold traps and snares. Journal of Wildlife

Management 82:906–917.

Auditor General, B. C. 2017. An Independent Audit of Grizzly Bear Management. Victoria,

British Columbia,Canada.

Craighead, J. J., and F. C. Craighead. 1972. Grizzly Bear-Man Relationships in Yellowstone

National Park. Bears: Their Biology and Management 2:304.

Flaa, J. P., S. B. Michel, and C. Borstad. 2009. Building a reliable snare cable for capturing

grizzly and American black bears. Ursus 20:50–55.

Graham, K., and G. Stenhouse. 2014. Home Range, Movements, and Denning Chronology of the

Grizzly Bear (Ursus arctos) in West-Central Alberta. The Canadian Field-Naturalist

128:223–234. <http://www.canadianfieldnaturalist.ca/cfn/index.php/cfn/article/view/1600>.

Accessed 11 Dec 2014.

Hatter, I. W., G. Mowat, and B. N. McLellan. 2018. Statistical population reconstruction to

evaluate grizzly bear trends in British Columbia, Canada. Ursus 29:1.

Knopff, K. H., A. A. Knopff, and M. S. Boyce. 2010. Scavenging Makes Cougars Susceptible to

Snaring at Wolf Bait Stations. Journal of Wildlife Management 74:644–653.

Lamb, C., A. Ford, B. McLellan, M. Proctor, G. Mowat, L. Ciarniello, S. Nielsen, and S. Boutin.

2020. The ecology of human-carnivore coexistence. Proceedings of the National Academy

of Science 117:17876–17883.

Lamb, C. T., A. T. Ford, M. F. Proctor, J. A. Royle, G. Mowat, and S. Boutin. 2019. Genetic

tagging in the Anthropocene: scaling ecology from alleles to ecosystems. Ecological

Applications 29:e01876. <https://onlinelibrary.wiley.com/doi/abs/10.1002/eap.1876>.

Lamb, C. T., G. Mowat, B. McLellan, S. Nielsen, and S. Boutin. 2017. Forbidden Fruit: Human

settlement and abundant fruit create an ecological trap for grizzly bears. Journal of Animal

Ecology 86:55–65.

McKim, C. 2017. Grizzly Bear With Trap On Its Foot Still Not Found. Wyoming Public

Radio2017:1. Wyoming. <https://www.wyomingpublicmedia.org/natural-resources-

energy/2017-06-26/grizzly-bear-with-trap-on-its-foot-still-not-found>.

McLellan, B. 2011. Implications of a high-energy and low-protein diet on the body composition,

fitness, and competitive abilities of black (Ursus americanus) and grizzly (Ursus arctos).

Canadian Journal of Zoology 558:546–558.

<http://www.nrcresearchpress.com/doi/abs/10.1139/z11-026>. Accessed 18 Feb 2014.

McLellan, B., and F. Hovey. 1995. The diet of grizzly bears in the Flathead River drainage of

southeastern British Columbia. Canadian Journal of Zoology 73:704–712.

<http://www.nrcresearchpress.com/doi/pdf/10.1139/z95-082>. Accessed 9 Apr 2013.

McLellan, B. N. 2015. Some Mechanisms Underlying Variation in Vital Rates of Grizzly Bears

on a Multiple Use Landscape. Journal of Wildlife Management 79:749–765.

McLellan, B. N., G. Mowat, T. Hamilton, and I. Hatter. 2017. Sustainability of the Grizzly Bear

Hunt in British Columbia, Canada. Journal of Wildlife Management 81:218–229.

Morehouse, A. T., C. Hughes, N. Manners, J. Bectell, and T. Bruder. 2020. Carnivores and

Communities: A Case Study of Human-Carnivore Conflict Mitigation in Southwestern

Alberta. Frontiers in Ecology and Evolution 8:1–15.

Munro, R. H. M., S. E. Nielsen, M. H. Price, G. B. Stenhouse, and M. S. Boyce. 2006. Seasonal

and Diel Patterns of Grizzly Bear Diet and Activity in West-Central Alberta. Journal of

Mammalogy 87:1112–1121. <http://asmjournals.org/doi/abs/10.1644/05-MAMM-A-

410R3.1>. Accessed 19 Feb 2014.

Province of British Columbia. 2021. British Columbia Hunting and Trapping Regulations

Synopsis.

Smith, F. 1896. The Maximum Muscular Effort of the Horse. The Journal of Physiology 19:224–

226.

White, H. B., G. R. Batcheller, E. K. Boggess, C. L. Brown, J. W. Butfiloski, T. A. Decker, J. D.

Erb, M. W. Fall, D. A. Hamilton, T. L. Hiller, G. F. Hubert, M. J. Lovallo, J. F. Olson, and

N. M. Roberts. 2021. Best Management Practices for Trapping Furbearers in the United

States. Wildlife Monographs 207:3–59.

**Table Captions**

404
405  Table 1. Phase two results of the toe extraction experiment where maximum human pulling
406  ability was applied in an attempt to free grizzly bear feet from traps.
407

| Age | Trap | Trials | Percent released |
|---|---|---|---|
| Adult | Bel120 | 5 | 0 |
| Adult | North155 | 5 | 60 |
| Adult | Sauv160 | 5 | 0 |
| Cub | Bel120 | 5 | 100 |
| Cub | North155 | 5 | 100 |
| Cub | Sauv120 | 8 | 50 |
| Cub | Sauv160 | 6 | 0 |

408
409  Table 2. Phase three results of the toe extraction experiment where a steady pull from either a
410  come along or a truck was applied to extract feet from traps. The maximum pull (kg) recorded on
411  an inline scale prior to release was recorded for each trial.
412

| Age | Trap | Trials | mean | min | max |
|---|---|---|---|---|---|
| Adult | Sauv120 | 3 | 170 | 118 | 232 |
| Adult | Sauv160 | 2 | 155 | 91 | 218 |

**Figure Captions**

A) Location of documented missing toes and by-catch

B) Timing of grizzly bear activity and trapping season

Figure 1. A) Locations of observations of bears with amputated toes and bears with traps on their feet. Detections made through research are outlined in grey, while detections opportunistically gathered throughout southeast British Columbia are not outlined in grey. B) The grizzly bear active season, defined from 61 collared bear den emergence and entrance dates, with the marten trapping season (November 1 to February 15) shown in grey. The 95th percentile of den entrance dates was November 22. We suggest that December 1 represents an evidence-based start date for trapping to reduce conflicts between bears and trappers.

422
Figure 2. Animals detected with amputated toes through research captures in the Elk Valley (A-
D), and opportunistic evidence of grizzly bears with traps on their feet (E-G). Each letter in A-D
represents one of the four unique individuals detected with amputated toes. Ci) an X-ray of the
foot of EVGM55, who was killed in a conflict with a cattle rancher. E) a body grip trap on the
foot of a hunter-killed grizzly bear from the Flathead Valley. F) the paw of a 1-year-old grizzly
bear stuck in a body grip trap captured by conservation officers near Pass Creek due to conflict
behaviour from a mom and offspring; significant infection was present, and the toes had nearly
completely separated from the foot by the time of capture. G) a grizzly bear with a body grip trap
on its foot detected feeding on a cow carcass by a remote camera west of Invermere.

432
433 Figure 3. Grizzly bears detected at trial marten body grip traps. The traps were wired so they
434 could spring but not close on bears' feet, and they were set for two weeks in areas frequented by
435 bears. Investigatory behaviour was common, and two of the four deployments were set off.