Development of *Kaptive* databases for *Vibrio parahaemolyticus* O- and K-antigen serotyping

Authors

Linda van der Graaf – van Bloois[1,2], Hongyou Chen[3], Jaap A. Wagenaar[1,2,4] and Aldert L. Zomer[1,2]

[1] Department of Biomolecular Health Sciences, Faculty of Veterinary Medicine, Utrecht University,

the Netherlands

[2] WHO Collaborating Centre for Reference and Research on *Campylobacter* and Antimicrobial

Resistance from a One Health Perspective / OIE Reference Laboratory for Campylobacteriosis,

Utrecht, the Netherlands

[3] Shanghai Municipal Center for Disease Control and Prevention, No. 1380, Zhong Shan Xi Rd.,

Shanghai 200336, PR China

[4] Wageningen Bioveterinary Research, Lelystad, the Netherlands

Running title: Development of *V. parahaemolyticus* O- and K-antigen Kaptive databases

Corresponding author: Aldert Zomer, a.l.zomer@uu.nl

Keywords: Vibrio parahaemolyticus, serotyping, whole genome sequencing, O-locus, K-locus

Abbreviations: KL, K-locus; KUT, K-Untypable; KLUT, K-locus Untypable; LPS/CPSgc, LPS and CPS gene

clusters; MLST, Multi Locus Sequence Typing; OL, O-locus; *V. parahaemolyticus*, Vibrio

parahaemolyticus; WGS, whole genome sequencing;

Data statement: All supporting data, code and protocols have been provided within the article or

through supplementary data files. The supplementary tables and supplementary figures are

available with the online version of this article.

Data summary: The sequence data are deposed at the European Nucleotide Archive (ENA) under

projects PRJEB39490 and PRJNA483379, and accession numbers are listed in Supplemental Table 1.

Abstract

Vibrio parahaemolyticus is an important food-borne human pathogen and is divided in 16 O-

serotypes and 71 K-serotypes. Agglutination tests are still the gold standard for serotyping, but many

V. parahaemolyticus isolates are not typable by agglutination. An alternative for agglutination tests

is serotyping using whole genome sequencing data. In this study, *V. parahaemolyticus* isolates are

serotyped and sequenced, and all known and several novel O- and K-loci are identified. We

developed *Kaptive* databases for all O- and K-loci after manual curation of the loci. These *Kaptive*

databases with the identified *V. parahaemolyticus* O- and K -loci can be used to identify the O- and

K-serotypes of *V. parahaemolyticus* isolates from genome sequences.

Introduction

Vibrio parahaemolyticus (*V. parahaemolyticus*) is an important food-borne human pathogen that

naturally inhabits marine environments worldwide, and can cause acute gastroenteritis and

septicemia in human (1,2). *V. parahaemolyticus* is typically serotyped on the basis of its heat-stable

somatic antigen (O) and its capsular antigen (K) and classified into 16 O-serotypes and 71 K-

serotypes.

Serotyping of *V. parahaemolyticus* is important for pathogen detection and epidemiological

surveillance. Many *V. parahaemolyticus* serotypes have been identified as pandemic clones, and

certain serotypes, for example O3:K6, O1:KUT and O4:K68, are generally considered to be more

virulent than others (3,4). Agglutination tests are the gold standard for *V. parahaemolyticus*

serotyping but frequently, *V. parahaemolyticus* strains are non-typable with agglutination tests in

routine screening (3,5–7). For several pathogenic bacterial species, e.g. several pathogenic *E. coli*

serogroups (8) and *Vibrio cholerae* serogroup O1 and O139 (9,10), PCR methods are developed to

detect the specific serogroups, but no PCR methods are available for *V. parahaemolyticus*

serotyping.

The software tool *Kaptive* was developed for rapid O- and K-loci typing of *Klebsiella* strains from

whole genome sequences (11). This tool includes an option to use other, self-created databases. The

genomic regions associated with the somatic synthesis locus (O-locus) and capsule synthesis locus

(K-locus) for *V. parahaemolyticus* are generally found on the same genomic location (2,12,13), which

makes it possible to locate and extract the O- and K-loci nucleotide sequences from whole genome

sequencing data of *V. parahaemolyticus* strains. Recently, the algorithm VPsero was developed,

which identified serogroup-specific genes as markers *in silico*, but only marker genes for 12 O- and

43 K-serotypes were covered (14).

The aim of this study was to identify all 16 O- and 71 K-loci of *V. parahaemolyticus* by serotyping,

sequencing and analyzing in-house isolates and investigating public data of serotyped strains and

manually curate reference databases for O- and K-loci, which can be used to determine the O- and K-

serotype of *V. parahaemolyticus* strains from whole genome sequencing data.

Materials and Method

Datasets of V. parahaemolyticus

In this study, 56 *V. parahaemolyticus* isolates were sequenced and serotyped. For whole genome

sequencing (WGS), DNA was extracted with TIANamp Bacteria DNA Kit (Tiagen Biotech (Beijing) CO.,

LTD, Beijing, China) according to the manufacturer's recommendation and DNA quality control was

performed using agarose gel electrophoresis and the Qubit dsDNA HS Assay Kit. DNA libraries and

DNA nanoballs (DNB) were constructed on BGISP-100 platform (WuHan MGI Tech Co.,Ltd, Wuhan,

China) with input of 150ng DNA. 50 bp single-end reads were generated with BGISEQ-50 sequencer

(WuHan MGI Tech Co.,Ltd, Wuhan, China) and assembled with SPAdes v3.12 (15). All sequence reads

of strains sequenced in this study are available under accession PRJEB39490 and PRJNA483379 of

the ENA short read archive.

Two datasets of public available *V. parahaemolyticus* genomes were included, consisting of 776

genomes from NCBI Genbank (Accession date: 14 December 2017) and 1498 genomes from PATRIC

database (16) (Accession date: 23 July 2020) were downloaded.

Taxonomic classification of all genomes was performed by *in silico* 16S DNA analysis with SPINGO

v1.3 (17) and genomes with ambiguous 16S results were classified as *Vibrio parahaemolyticus* using

KmerFinder v3.2 (18).

Serotyping of V. parahaemolyticus *strains*

Serotyping of the 56 in this study sequenced strains was performed by slide agglutination according

to the manufacturer's instructions, using the 65 type-K serum set and 11 type-O serum set (Denka

Seiken CO., LTD, Tokyo, Japan).

Development Kaptive *databases*

I.   Identification of O-loci

V. parahaemolyticus is divided into 16 O-serotypes (1,2). The O-locus has been defined located

between *dgkA* and *gmhD* genes, which encode a diacylglylcerol-kinase and an epimerase

respectively (2). With BlastN (minimum coverage of 80% and minimum identity of 80%), the flanking

genes of the O-region, *dgkA* and *gmhD*, were searched in all genome sequences. The O-loci

sequences were extracted from these sequences manually.

II.   Identification of K-loci

V. parahaemolyticus is divided in 71 K-serotypes (2). The K-locus has been defined located between

gene *gmhD* which encodes an epimerase at one side, and at the other side gene *rjg* encoding a

metallo-hydrolase (13). With BlastN (minimum coverage of 80% and minimum identity of 80%), the

flanking genes of the K-region, *gmhD* and *rjg,* were located in all 1601 genome sequences. In some

cases where the *rjg* gene was not found, gene *ugd* or gene *gtaB* was selected as flanking gene. The

K-loci sequences were extracted from these sequences manually.

III.   Development of the *Kaptive* databases

The extracted O- and K-loci sequences were annotated using Prokka v1.13 (19). For annotating the

loci, a custom database was used which was built with the annotations of the previous described loci

of *V. parahaemolyticus* (1,2). Genes of the O- and K-loci respectively were clustered using Roary with

an 80% cutoff on amino acid identity (20). The Roary gene presence absence table was used to

create gene presence-absence clusters of the O- and K-loci respectively and the unique gene

presence-absence loci were linked to the known serotypes of the genomes.

For the identified O- and K-loci, the nomenclature of the *Klebsiella* capsule synthesis loci (11) was

used; each distinct O- and K-locus was designated as OL (O-locus) and KL (K-locus), followed by an

unique number. The O- and K-loci of known serotypes were assigned with the same number as the

corresponding O- and K-serotypes, e.g. O-serotype 1 is encoded by the OL1 locus etcetera. O- and K-

loci with unknown serotypes were assigned with numbers starting from 101 (e.g. OL101 and KL101).

Variants of loci with maximal two genes different by insertion or deletion were given the suffix -1.

Eight gene patterns of K-Untypable (KUT) strains were found and these patterns are assigned in the

k-loci database as KLUT (K-locus UnTypable) followed by subsequent numbers 1-8

Of each loci, one genome was selected as reference and the nucleotide sequences of the O- and K-

loci of the selected genomes were added to the *Kaptive* database file and curated manually.

The *V. parahaemolyticus* O-locus and K-locus *Kaptive* databases are available on the following

Github page: https://github.com/aldertzomer/vibrio_parahaemolyticus_genomoserotyping and will

be added to Kaptive web: https://kaptive-web.erc.monash.edu/

Comparison of genes in O- and K-loci

The reference genomes of the identified O- and K-loci were aligned using Parsnp v1.2 (21),

recombination regions were filtered using Gubbins v2.3.4 (22) and the tree was built with FastTree

v2.1.8 (23). The tree was visualized with iTol v6.5.4 (24) and Multi Locus Sequence Typing (MLST)

was performed in silico with the MLST software (Seemann) using PubMLST typing schemes (26).

Gene cluster comparisons of the O- and K-loci are made with Clinker v0.0.37 (27).

Performance of Kaptive *databases compared to VPSero*

Of the total set of 831 selected genomes from Genbank (n=775) and in-house sequenced genomes

(n=56), 436 genomes had a known O-serotype and 294 genomes had a known K-serotype, whereas

the PATRIC genomes had all unknown O- and K-loci serotypes. The O- and K-loci of the genomes with

the known serotypes were all serotyped by the newly developed *Kaptive* databases and VPSero (14).

Furthermore, performance of the *Kaptive* databases was tested and compared with VPSero by using

the VPSero sequence data collection, deposited into CNGB Sequence Archive project CNP0000343

described by Bian *et al* (14).

Results and Discussion

Development of Kaptive *databases for V. parahaemolyticus O- and K-loci*

The study started with a set of 2330 *V. parahaemolyticus* genomes, consisting of 56 in this study

sequenced genomes, 776 genomes downloaded from NCBI GenBank and 1498 from the PATRIC

database. The genomes that were *in silico* identified as *V. parahaemolyticus* with 16S and

KmerFinder analysis were included. For the genomes downloaded from NCBI GenBank and PATRIC

database, only genomes that contained the complete loci on one contig were included. This

selection ended-up with a total set of 1601 genomes, consisting of 56 in-house sequenced strains,

775 from Genbank and 770 genomes from PATRIC database. For the development of the O- and K-

loci databases, only genomes that contained both flanking genes of the loci on one contig were

used. No isolate or genome were available for K-serotypes 26 and 71, and these serotypes were

therefore not included in the database.

The selected genomes that were used as reference for each O- and K-loci are shown in Supplemental

Table 1. For each O-serotype, unique loci were identified, except for serotype O3 and O13 genomes,

since the loci of these serotypes contained the same genes (1). Therefore, loci of serotype O3 and

O13 could not be distinguished based on gene presence and are assigned in the *Kaptive* O-database

as "O3_or_O13". It is possible that a second cluster modifies the O-antigen, similar to what has been

described for *Shigella* (28), however an O-modifying locus has not been described in literature for

O13.

Eight gene patterns of K-UnTypable (KUT) strains previously described were found and these

patterns are assigned in the *Kaptive* database as K-Locus UnTypable (KLUT) followed by subsequent

numbers 1-8 (Supplemental Table 1).

For both O- and K-loci, not all gene patterns could be assigned to specific serotypes, because these

gene patterns were found in genomes with unknown serotypes from NCBI and PATRIC databases.

These unknown gene patterns were assigned for O-loci as OL101 and for K-loci as KL101-KL157

(Supplemental Table 1).

For some O- and K-serotypes, variants in the loci gene patterns by insertion of deletion of maximal

two genes were found, and therefore, these serotypes have two gene patterns included in the

database, distinguished with the addition of "-1" to the locus name (OL1, OL4, OL7 and KL20, KL30,

KL68, KLUT4) (Supplemental Figures 1 and 2).

Examination of the phylogenetic tree of the reference isolates showed the O- and K-loci are not

associated with MLST sequence types, e.g. six different K-loci and four different O-loci are found in

ST3 reference isolates (Figure 1).

Performance of Kaptive *databases*

-    Genome collection of this study

To investigate the performance of the newly developed *Kaptive* databases, 436 genomes with a

known O-serotype and 295 genomes with a known K-serotype were used from our genome

collection. Of the 436 known O-serotypes, 398 O-serotypes were correctly identified (Figure 2A,

Supplemental Table 1) and 265 of the 295 known K-serotypes were correctly identified (Figure 2B,

Supplemental Table 1).

The tool VPSero was developed for 43 K- and 12 O-serotypes (14), and covers part of the known 16

O-serotypes and 71 K-serotypes. The performance of VPSero was tested with a set of 420 genomes

for O-serotyping and 274 genomes for K-serotyping from our genome collection, consisting of

genomes with known serotypes of the 12 and 46 O- and K-serotypes respectively which are included

in VPSero. Of this set, 343 out of 420 genomes with known O-serotypes were correctly identified

(Supplemental Figure 2A, Supplemental Table 1) and 183 out of 274 genomes with known K-

serotypes (Supplemental Figure 2B, Supplemental Table 1). The newly developed *Kaptive* databases

outcompete VPSero with 395/420 correctly identified O-serotype genomes and 254/274 correctly

identified K-serotype genomes (Supplemental Figure 2AB, Supplemental Table 1).

-    Sequence collection of VPsero

In the publication of VPsero, intact LPS and CPS gene clusters (LPS/CPSgc) were identified and

extracted, and deposited in CNGB Sequence Archive project CNP0000343. These gene clusters did

not contain the full O-serotype loci of the newly developed O-serotype *Kaptive* database, and

therefore, these sequence data collection could not be used to test the performance of the O-

serotype *Kaptive* database, the K-serotype loci were available however. The LPS/CPSgcs were tested

with both *Kaptive* K-database and VPSero, and results are listed in Supplemental Table 2. One K58

gene cluster was identified as K6 by both tested tools, possibly containing a wrong serotype in the

CNGB database. Of the remaining 56 gene clusters with known K-serotypes, seven were not typable

by VPSero and two were mis-identified by *Kaptive* databases.

The mis-identified O- and K-serotypes from our collection were all genomes downloaded from NCBI.

Since we only have the genomes and not the strains, we cannot check if these strains were

serotyped correctly. O-serotype O14 has been recently described (2) and only one genome with this

O-serotype is available. In our dataset of 831 genomes, 14 NCBI genomes with serotype O5 are

identified with the *Kaptive* database as the new serotype O14. It is very likely that these genomes

are mis-serotyped, because the serum for serotype O14 was not available when these genomes

were serotyped.

For several K-serotypes, only one genome sequence was available and we could not determine if

there is variation in these K-loci gene content. Furthermore, several of the genomes sequenced in

this study were sequenced with a 50-bp single-end BGI sequencer, resulting in a higher number of

short contigs, therefore for several of these genomes, the K-locus was not assembled on one single

contig. For these genomes, the contigs with flanking genes were selected and concatenated

manually. It is possible that some genes are missing in these K-locus reference sequences, although

these likely represent repeat sequences. If more *V. parahaemolyticus* genome sequences with closed

K-loci become available, the *Kaptive* K-serotype database will be updated with the closed K-locus

sequence of these K-serotypes.

Conclusion

231 The in this study developed *Kaptive* databases with the identified 16 O- and 71 K -loci can be used to

232 identify the O- and K-serotypes of *V. parahaemolyticus* isolates from whole genome sequencing

233 data. The variation of K-antigen loci is much higher than expected as we identified 57 new K-locus

234 variants.

Author Statements

Author contributions

Designed the study; L.G., J.W., A.Z. Performed the experiments; L.G., H.C., A.Z. Analyzed the data;

L.G., A.Z. Writing the paper; all authors.

Conflicts of interest

The authors declare that they have no competing interests.

Funding information

Sequencing of isolates was supported by the National Science and Technology Major Project of

China (No. 2018ZX10305409-003).

Acknowledgements

We thank the Kaptive team (Thomas Stanton, Kelly Wyres, Ryan Wick and Kathryn Holt) for the

undertaking to include the *Vibio parahaemolyticus* databases to Kaptive web.

References

1. Chen M, Guo D, Wong HC, Zhang X, Liu F, Chen H, et al. Development of O-serogroup specific PCR assay for detection and identification of Vibrio parahaemolyticus. Int J Food Microbiol. 2012;159(2):122–9.

2. Guo X, Liu B, Chen M, Wang Y, Wang L, Chen H, et al. Genetic and serological identification of three Vibrio parahaemolyticus strains as candidates for novel provisional O serotypes. Int J Food Microbiol. 2017;245:53–8.

3. Jones JL, Lüdeke CHM, Bowers JC, Garrett N, Fischer M, Parsons MB, et al. Biochemical, serological, and virulence characterization of clinical and oyster Vibrio parahaemolyticus isolates. J Clin Microbiol. 2012;50(7):2343–52.

4. Nair GB, Ramamurthy T, Bhattacharya SK, Dutta B, Takeda Y, Sack DA. Global dissemination of Vibrio parahaemolyticus serotype O3:K6 and its serovariants. Clin Microbiol Rev. 2007;20(1):39–48.

5. Gavilan RG, Zamudio ML, Martinez-Urtaza J. Molecular Epidemiology and Genetic Variation of Pathogenic Vibrio parahaemolyticus in Peru. PLoS Negl Trop Dis. 2013;7(5).

6. Gil AI, Miranda H, Lanata CF, Prada A, Hall ER, Barreno CM, et al. O3:K6 Serotype of Vibrio parahaemolyticus identical to the global pandemic clone associated with diarrhea in Peru. Int J Infect Dis. 2007;11(4):324–8.

7. Hashii N, Kondo S, Iguchi T, Nishibuchi M, Hisatsune K. Chemical and serological properties of lipopolysaccharides from Vibrio parahaemolyticus O-untypeable strains isolated from patients. Microbiol Immunol. 2000;44(4):229–34.

8. Wang Q, Ruan X, Wei D, Hu Z, Wu L, Yu T, et al. Development of a serogroup-specific multiplex PCR assay to detect a set of Escherichia coli serogroups based on the identification of their O-antigen gene clusters. Mol Cell Probes. 2010;24(5):286–90.

9. Albert MJ, Islam D, Nahar S, Qadri F, Falklind S, Weintraub A. Rapid detection of Vibrio cholerae O139 bengal from stool specimens by PCR. J Clin Microbiol. 1997;35(6):1633–5.

10. Hoshino K, Yamasaki S, Mukhopadhyay AK, Chakraborty S, Basu A, Bhattacharya SK, et al. Development and evaluation of a multiplex PCR assay for rapid detection of toxigenic Vibrio cholerae O1 and O139. FEMS Immunol Med Microbiol. 1998;20(3):201–7.

11. Wyres KL, Wick RR, Gorrie C, Jenney A, Follador R, Thomson NR, et al. Identification of Klebsiella capsule synthesis loci from whole genome data. Microb genomics. 2016;2(12):e000102.

12. Okura M, Osawa R, Tokunaga A, Morita M, Arakawa E, Watanabe H. Genetic analyses of the putative O and K antigen gene clusters of pandemic Vibrio parahaemolyticus. Microbiol Immunol. 2008;52(5):251–64.

13. Chen Y, Dai J, Morris JG, Johnson JA. Genetic analysis of the capsule polysaccharide (K antigen) and exopolysaccharide genes in pandemic Vibrio parahaemolyticus O3:K6. BMC Microbiol. 2010;10(1):274.

14. Bian S, Jia Y, Zhan Q, Wong NK, Hu Q, Zhang W, et al. VPsero: Rapid Serotyping of Vibrio parahaemolyticus Using Serogroup-Specific Genes Based on Whole-Genome Sequencing Data. Front Microbiol. 2021;12:2446.

15. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: A new genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol. 2012;19(5):455–77.

16. Wattam AR, Abraham D, Dalay O, Disz TL, Driscoll T, Gabbard JL, et al. PATRIC, the bacterial bioinformatics database and analysis resource. Nucleic Acids Res. 2014;42(D1):581–91.

17. Allard G, Ryan FJ, Jeffery IB, Claesson MJ. SPINGO: A rapid species-classifier for microbial amplicon sequences. BMC Bioinformatics. 2015;16(1):1–8.

18. Hasman H, Saputra D, Sicheritz-Ponten T, Lund O, Svendsen CA, Frimodt-Moller N, et al. Rapid whole-genome sequencing for detection and characterization of microorganisms directly from clinical samples. J Clin Microbiol [Internet]. 2014;52(1):139–46.

19. Seemann T. Prokka: Rapid prokaryotic genome annotation. Bioinformatics.

2014;30(14):2068–9.

20.   Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MTG, et al. Roary: Rapid large-
scale prokaryote pan genome analysis. Bioinformatics. 2015;31(22):3691–3.

21.   Treangen TJ, Ondov BD, Koren S, Phillippy AM. The harvest suite for rapid core-genome
alignment and visualization of thousands of intraspecific microbial genomes. Genome Biol.
2014;15(11):1–15.

22.   Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, et al. Rapid phylogenetic
analysis of large samples of recombinant bacterial whole genome sequences using Gubbins.
Nucleic Acids Res. 2015;43(3):e15.

23.   Price MN, Dehal PS, Arkin AP. FastTree 2 - Approximately maximum-likelihood trees for large
alignments. PLoS One. 2010;5(3).

24.   Letunic I, Bork P. Interactive Tree Of Life (iTOL): An online tool for phylogenetic tree display
and annotation. Bioinformatics. 2007;23(1):127–8.

25.   Seemann T. MLST. Available from: https://github.com/tseemann/mlst.

26.   Jolley KA, Maiden MCJ. BIGSdb: Scalable analysis of bacterial genome variation at the
population level. BMC Bioinformatics. 2010;11(1):1–11.

27.   Gilchrist CLM, Chooi Y-H. clinker & clustermap.js: Automatic generation of gene cluster
comparison figures. Bioinformatics. 2021;37(16); 2473-2475.

28.   Lehane AM, Korres H, Verma NK. Bacteriophage-encoded glucosyltransferase GtrII of Shigella
flexneri: Membrane topology and identification of critical residues. Biochem J.
2005;389(1):137–43.

Figures and Tables

Figure 1. Phylogenetic tree based on core genome analysis of O- and K-representative isolates,

including the O- and K-loci, and MLST STs.

Figure 2. Performance of the _Kaptive_ database and VPSero on genomes with known (A) O- and (B)

K-serotypes. In green the percentages of correctly typed genomes with _Kaptive_ databases, in orange

the percentages of incorrectly typed genomes with _Kaptive_ databases, in blue the percentages of

correctly typed genomes with VPSero, in red the percentages of incorrectly typed genomes with

VPSero, in grey the percentages of the with VPSero not-typable genomes. The numbers in the bars

represent the number of genomes.

Supplemental Figure 1. Gene cluster comparisons of the identified O-loci. All similar gene homologs

have been assigned an unique color and are linked with similar color boxes. Hypothetical proteins

are assigned with an "1".

Supplemental Figure 2. Gene cluster comparisons of the identified K-loci. All similar gene homologs

have been assigned an unique color and are linked with similar color boxes. Hypothetical proteins

are assigned with an "1".

Supplemental Table 1. Selected reference genomes used for the O- and K-loci _Kaptive_ databases.

Included are the numbers of genomes with known O- and K-serotypes and numbers of correctly

identified genomes with _Kaptive_ databases and VPSero.

Supplemental Table 2. Performance of _Kaptive_ databases and VPSero on sequence collection of

VPSero.

Tree scale: 10000