1 **Hurricane-induced population decrease in a Critically Endangered long-lived reptile**
2

3 Matthijs P. van den Burg[1,2,*], Hannah Madden[1,3,4,*], Timothy P. van Wagensveld[1,5], and Erik
4 Boman[6]
5
[1] IUCN SSC Iguana Specialist Group, Gland, Switzerland.
[2] BioCoRe S. Coop. Calle Villagarcía 6, 28010, Madrid, Spain.
[3] Caribbean Netherlands Science Institute, P.O. Box 65, St. Eustatius, Caribbean Netherlands.
[4] NIOZ Royal Netherlands Institute for Sea Research, and Utrecht University, P.O. Box 59, 1790
AB Den Burg, Texel, the Netherlands.
[5] Reptile Amphibian Fish Research the Netherlands, Nijmegen, The Netherlands.
[6] St. Eustatius National Park Foundation, St. Eustatius, Caribbean Netherlands.
13
14 * Joint first authors, email: thijs.burg@gmail.com, hannah.madden@cnsi.nl
15

ABSTRACT

Catastrophic events, like hurricanes, bring lethal conditions that can have population-altering

effects. The threatened Caribbean dry forest occurs in a region known for its high-intensity

hurricane seasons and high species endemism, highlighting the necessity to better understand

hurricane impacts as fragmentation and clearing of natural habitat continues. However, such studies

remain rare, and for reptiles are mostly restricted to *Anolis*. Here we used single-season occupancy

modeling to infer the impact of the intense 2017 Atlantic hurricane season on the critically

endangered Lesser Antillean Iguana, *Iguana delicatissima*. We surveyed 30 transects across eight

habitats on St. Eustatius during 2017-2019, which resulted in 344 individual surveys and 98 iguana

observations. Analyses of abundance and site occupancy indicated both measures for 2018 and

2019 were strongly reduced compared to the pre-hurricane 2017 state. Iguanas at higher elevations

were affected more profoundly, likely due to higher wind speeds, tree damage and extensive

defoliation. Overall, our results indicate a decrease in population estimates (23.3-26.5%) and

abundance (22-23.8%) for 2018 and 2019, and a 75% reduction in the number of opportunistic

sightings of tagged iguanas between 2017-2018. As only small and isolated *I. delicatissima*

populations remain, our study further demonstrates their vulnerability to stochastic events.

Considering the frequency and intensity of hurricanes are projected to increase, our results stress the

urgent need for population-increasing conservation actions in order to secure the long-term survival

of *I. delicatissima* throughout its range.

Keywords
Abundance, Caribbean, Detection probability, Disturbance event, Habitat, Hurricane, *Iguana delicatissima*, Monitoring, Site occupancy, St. Eustatius

Introduction

Hurricanes bring lethal conditions to populations in their path. These events can either directly

affect natural populations, through e.g., mortality (Reagan 1991; Wiley and Wunderle 1993; Spiller

et al. 1998; Behie and Pavelka 2005; Morcilo et al. 2020; Marroquin-Paramo et al. 2021), or

indirectly e.g., when food presence and abundance is impacted (Cely 1991). Selective mortality

sweeps have been shown to drive both evolution (Donihue et al. 2018) and community diversity

(Johnson and Winker 2010; Meléndez-Vazquez et al. 2019). Additionally, catastrophic winds can

aid dispersal of species (e.g., Censky et al. 1998). Although studies on the effects of hurricanes are

rising given their projected increase (Webster et al. 2005), such studies are dependent on pre-

hurricane data and hurricane trajectory, and are thus lacking for many species.

From a conservation perspective, the population-diminishing threat of hurricanes is

especially of concern for closed island populations, which often represent restricted-range and

endemic species. As natural habitat disappears and fragmentation continues to worsen (Jacobson et

al. 2019; Powers and Jetz 2019), understanding how stochastic events affect remaining populations

is crucial. In a rare long-term study on an endangered Bahamian woodpecker (*Melanerpes*

superciliaris nyeanus), Akresh et al. (2020) demonstrate that the population experienced significant

declines as a result of various hurricanes but was able to recover after several years, however

increased future hurricane frequency could prevent this. Contrarily, a study on a small, 350-

individual, population of *Cyclura nubila* revealed that the passing of Hurricane Ivan (2004) had no

effect on its size (Beovides-Casas and Mancina 2006), demonstrating the resilience of large rock

iguanas to catastrophic events. Overall, studies concerning the effect of hurricanes on reptile species

have mainly focused on small-size species (e.g., *Anolis*: Spiller et al. 1998; Schoener et al. 2001;

Losos et al. 2003; Donihue et al. 2018, 2020; Dufour et al. 2019; Rabe et al. 2020). Insular iguanas

are highly endangered ecological keystone species, and generally are the largest native remaining

terrestrial species (IUCN 2020). As Caribbean hurricane frequency is projected to increase (Bender

et al. 2010), understanding the impacts of hurricanes on iguana populations is important and is

frequently highlighted but rarely quantified (Alberts 2004; Fogarty et al. 2004; Powell 2004;

Pasachnik et al. 2012; but see Hayes et al. 2004, 2016; Beovides-Casas and Mancina 2006).

Among Caribbean Iguanids, the conservation status of the Lesser Antillean iguana (*Iguana*

delicatissima) has experienced the most rapid deterioration (van den Burg et al. 2018a). Its pre-

Colombian range has to date decreased by >80%, predominantly due to hybridization with *Iguana*

iguana lineages (Vuillaume et al. 2015; Pounder et al. 2020), a process that was recently initiated

on the last remaining ≥ 2 km^2 islands where *I. delicatissima* occurs (van den Burg et al. 2018b,

2020; B. Angin pers. comm.). Left uncontrolled, the genetic loss of these populations will lead to a

>99% decrease in their native range (van den Burg et al. 2018a). Critically, the five remaining

populations that are free of nonnative iguanas occur on <2 km^2 islets with low estimated population

sizes (only Petite Terre >1,000 individuals; Angin 2017; Pounder et al. 2020). Beyond the spread of

I. iguana (Knapp et al. 2020), other threats come from habitat destruction, pets and free-roaming

livestock, as well as other anthropogenic causes (e.g., vehicle-caused mortality) (Debrot et al. 2013;

Knapp et al. 2014, 2016; van den Burg et al. 2018c). Although hurricanes are cited as a potential

threat to population survival (e.g., Fogarty et al. 2004), only two studies have addressed this topic.

Namely, Knapp and Valeria (2008) provide some insight through an observational account of nest

site degeneration and iguana mortality caused by Hurricane Dean (2007) on Dominica. Focusing on

Petite Terre (Guadeloupe), a second study provides insight into a likely decrease in population size

due to Hurricane Hugo (1995; Lorvelec et al. 2004). However, since the monitoring methodology

has been questioned (Breuil and Ibéné, 2008; B Angin pers. comm.), no conclusions can be drawn.

This study also predated new analytical tools and updates (e.g., Fiske and Chandler 2011).

Within the northern Lesser Antilles, St. Eustatius supports one of the last remaining *I.*

delicatissima populations. This population suffered a major decline during the 1990s, in part due to

consumptional demand on neighboring St. Maarten (Debrot et al. 2013), and is currently threatened

by ongoing hybridization with *I. iguana* (van den Burg et al. 2018b). Despite the occasional arrival

of stowaway iguanas, continuous monitoring has only resulted in 15 (Debrot et al. in review.)

identified nonnative (pure or hybrid) animals, thus hybridization is believed to be minimal

(unpublished data). The most recent estimates put the population size at below 500 individuals

(Debrot et al. 2013), which has presumably since decreased due to ongoing anthropogenic threats

(van den Burg et al. 2018c). Additionally to threats common for the species (see above), the

population on St. Eustatius faces death by entrapment in abandoned cisterns, invasive rats, and the

spread of the nonnative invasive Mexican creeper or Corallita (*Antigonon leptopus*) (Fogarty et al.

2004; Debrot et al. 2013; van den Burg et al. 2018c). Although St. Eustatius fell within the

trajectory of major 2017-season hurricanes (Fig. 1), an impact assessment for *I. delicatissima*

population is lacking.

The 2017 Atlantic hurricane season was categorized as hyperactive, with an accumulated

cyclone energy score within the top ten since registration started (Schultz et al. 2018). Of the six

2017 hurricane-force systems, two closely passed St. Eustatius. On 6 September, the center of

Hurricane Irma passed at ~54 km, and on 19 September, Hurricane Maria passed at roughly 110 km

distance (Fig. 1; NOAA 2020). The goal of our study was to utilize iguana-survey data collected

during 2017–2019 in order to assess the impact of the 2017 hurricane season on the St. Eustatius *I.*

delicatissima population. To that end, we used island-wide transect survey data to estimate

abundance, detection probability, and site occupancy per habitat type and year. Changes in

occupancy, defined as the probability of a site being occupied (MacKenzie et al. 2002), provide a

robust proxy for population declines, particularly for sparsely populated and cryptic species (Dénes

et al. 2015; Beaudrot et al. 2016). We hypothesized that if population estimates were stable over

this 3-year period, this would suggest that iguanas are resilient to catastrophic weather events.

However, if population estimates varied, this would suggest that external factors (directly and/or

indirectly) impacted iguana survival.

114
Fig. 1 Best trajectory paths and regions with hurricane force winds ($\geq$ 64kt) for Hurricane Maria and Irma in

the northern Greater Caribbean region. Data from the National Hurricane Center.

Materials & Methods

Study area

This study took place on St. Eustatius (17.49°N, -62.98°W), a small (21 km^2) Leeward island within

the northern Lesser Antilles (Fig. 1), with a human population of approximately 3,100 (Statistics

Netherlands 2020). Habitat falls within the Caribbean dry forest and local climate is defined as

Tropical Humid (Trewartha and Horn 1980), with over 90% of the island having a tropical savanna

climate (<400 m) (De Freitas et al. 2012). Annual temperature fluctuates around 25–33 °C, with

average annual rainfall around 1000 mm while the highest elevations receive <2,000 mm (Rojer

1997; van Andel et al. 2016). Geographically, the island consists of two mountainous areas that

originate from volcanic activity and uplift on the northern and southern end. These areas are

separated by a lower elevational plain where most anthropogenic activity occurs. Historic island-

wide agricultural practices have downgraded the habitat quality which has partially recovered due

to a significant decrease in such practices and conservation efforts, though habitats are affected still

by grazers, erosion and invasive species (van Andel et al. 2016).

132
Transects

Between 2017–2019, we conducted repeated count transect surveys to estimate site

occupancy, detection probability and abundance of the *I. delicatissima* population (Royle and

Nichols 2003). For this, we placed a total of 30 transects over the island, each 100 m in length, and

visually surveyed up to 25 m on both sides (Fig. 2). Surveys were performed by minimally one

observer with iguana-fieldwork experience, during hours of general iguana activity (Pasachnik et al.

2002), and were limited to non-rainy days given ectotherm activity. We recorded the date, survey

duration (min.), elevation (m), number of observers, and observed number of iguanas. Due to

limited capacity to conduct surveys, transects were surveyed a maximum of six times per year.

Transects included 47% of present habitat types representing 78.3% of the island's surface. Several

habitat types were not surveyed due to access restrictions (e.g., *Bothriochloa - Antigonon*) or

physical safety (e.g., *Antirhea - Coccoloba* mountains). Furthermore, the northern area was

surveyed less often than other habitats due to logistics and access difficulty of the area, and given an

oil transhipment company occupies 10–15% of the island's surface area. Following De Freitas et al.

(2012) nomenclature, surveys were performed in eight habitat types (see Fig. 2, Table 1): *Pisonia -*

Antirhea mountains; miscellaneous urban/agricultural or disturbed land; *Pisonia - Justicia* hills;

Capparis - Pisonia mountains; *Rauvolfia - Antigonon* mountains; *Coccoloba - Bothriochloa* cliffs;

Pisonia - Eugenia mountains; *Chionanthus - Nectandra* mountains.

151
152 **Fig. 2** Map of St. Eustatius showing transect locations and habitat distribution. Habitat names following De

153 Freitas et al. (2012).

154 **Table 1** Summary of transect surveys conducted on St. Eustatius between 2017–2019. Habitat name

155 following De Freitas et al. (2012). All transects constituted 100 m in length with a surveyed area of 500 m^2.

Habitat name	Transect nr.	Lat.	Lon.	Start elev. (m)
Pisonia - Antirhea mountains	(1)	17.4672	-62.9681	69
	(2)	17.4659	-62.9643	41
Misc. urban/agricultural or disturbed land	(3)	17.4691	-62.9808	12
	(4)	17.4947	-62.9867	59
	(5)	17.4985	-62.9835	53
	(7)	17.5024	-62.9825	25
	(15)	17.4869	-62.9759	93
	(16)	17.4868	-62.9573	117
	(19)	17.4847	-62.9728	141
	(20)	17.4831	-62.9879	14
	(21)	17.4797	-62.9860	25
	(23)	17.4794	-62.9801	87
	(25)	17.4945	-62.9742	48
	(28)	17.4859	-62.9783	91
	(29)	17.4856	-62.9749	114
	(30)	17.4862	-62.9721	123
Pisonia - Justicia hills	(6)	17.4994	-62.9844	66
	(8)	17.5098	-62.9900	70
Capparis - Pisonia mountains	(9)	17.4742	-62.9518	79
	(10)	17.4774	-62.9499	50
	(11)	17.4790	-62.9508	72
	(12)	17.4806	-62.9510	80
	(13)	17.4837	-62.9527	104
	(14)	17.4862	-62.9547	99
Rauvolfia - Antigonon mountains	(17)	17.4892	-62.9642	104
	(18)	17.4884	-62.9654	117
Coccoloba - Bothriochloa cliffs	(22)	17.4745	-62.9852	12
Pisonia - Eugenia mountains	(24)	17.4764	-62.9748	149
	(26)	17.4754	-62.9726	190

Chionanthus - Nectandra mountains	(27)	17.4780	-62.9670	386

Opportunistic surveys/sightings

Since 2015, subadult and adult iguanas have been tagged with colored glass beads to

facilitate subsequent noninvasive tracking and observations of individuals using unique color

combinations (Binns and Burton 2007). To date, 540 unique bead codes have been assigned, and an

additional 96 iguanas have been tagged only with PIT tags. This methodology has allowed for

opportunistic data collection of the geographic locations of bead-tagged iguanas. We utilized these

data as an additional indicator for variation in population size.

164
Data analyses

As our dataset contained a large number of non-detections and small counts of iguanas, we

chose an *N*-mixture model approach over distance sampling (Royle 2004). Models were based on

four assumptions: (i) the iguana population in each site was closed to colonization and extinction

during the sampling period; (ii) iguanas were either detected or not detected during surveys; (iii)

iguanas were not falsely detected during surveys; and (iv) iguana detections at each site were

independent (Murray and Sandercock 2020). We followed the methods described by Madden et al.

(2021). Briefly, we used likelihood-based single-season occupancy models (MacKenzie et al.

2017), using the R packages "wiqid" and "unmarked" (MacKenzie et al. 2002; Fiske and Chandler

2011), to estimate site occupancy (ψ) in relation to habitat and elevation, while accounting for

detectability (p). We modeled detection probability and occupancy probability to estimate iguana

abundance (λ), and also tested the influence of independent variables on abundance estimates,

including quadratic terms. Resulting models were ranked using Akaike's Information Criterion,

where the detection model with the lowest AICc is best-fitting [AICc; Burnham and Anderson

(2002)], while correcting for small sample size. Finally, we tested for overdispersion by assessing

the goodness-of-fit of the most parsimonious model in each year (MacKenzie and Bailey 2004)

with the "mb.gof.test" function in the package "AICcmodavg" using 1,000 simulations, which

calculates a Pearson's chi-square fit statistic from the observed and expected frequencies of

detection histories for a given model. All analyses were performed in the R environment v3.5.1 (R

Core Team 2019).

185
Results

Pre-hurricane survey occupancy, detection and abundance estimates

We conducted 180 surveys between February and April 2017, prior to hurricanes Irma and Maria,

during which we detected 58 individual iguanas. Average transect time was 32 ±14 minutes. As

results for both R packages provided identical patterns we only report "wiqid" results here, and

"unmarked" results are presented in the Electronic Supplementary Information.

192   Occupancy probability was 100% (range 0–100%), and iguana occupancy and abundance

increased with elevation (Figs. 3–4) as well as with survey effort. Occupancy estimates were

generally consistent across the habitats sampled (Table 2). Mean detection probability (p) per

transect was 0.19 (range 0.14–0.26), which increased to 0.27 when an iguana had been previously

detected. The effects of predictor variables 'survey time' and 'week' on detection and occupancy

estimates were negligible (**Δ**AIC$_c$ <2). We observed a lack of fit of the null model (without

covariates; X^2 = inf, p = 0, c-hat (ĉ) = inf). Estimated mean individual abundance (λ) was 5.91 (± SE

0.01). When habitat and elevation were included as response variables, λ estimates were 1.52 ± 0.65

(Fig. 5), which translates to a population estimate across the entire study area (13.5 ha) of 1,364

iguanas. Lowest iguana abundance was found in *Coccoloba - Bothriochloa* cliff habitat, and highest

in *Chionanthus - Nectandra* mountain habitat.

Post-hurricane survey occupancy, detection and abundance estimates

Post-2017, we surveyed identical transects between July–December 2018 and March–September

2019. Overall, 164 transects were performed during which we detected 13 iguanas in 2018 and 27

iguanas in 2019. Average transect time was 19 ± 8 minutes.

Occupancy probability was 32.5% (range 11.8–63.4%) in 2018, and 36.7% (19.2–58.5%) in

2019. Iguana occupancy decreased with increasing elevation in 2018 and 2019 (Fig. 3), but

increased with survey effort; up to four repeats in 2018 and three in 2019. Occupancy estimates

were lower than those from 2017 but generally consistent across habitats (Table 2). p was 0.41

(range 0.14–0.74) in 2018, and 0.67 (0.39–0.87) in 2019, decreasing to 0.28 (0.07–0.67; 2018) and

0.36 (0.14–0.66; 2019) when an iguana had been previously detected. λ was 5.89 ± 0.02 in both

2018 and 2019. Considering habitat and elevation, λ estimates were 1.16 ± 0.15 in 2018 and $1.19 \pm$

0.18 in 2019 (Fig. 5), translating to population estimates of 1,046 and 1,003 iguanas, respectively;

declines of 23.3% and 26.5% compared to 2017. The covariate 'survey time' from the most

parsimonious model had a positive effect on detection probability (0.80 ± 0.49) in 2018, however

we observed a lack of fit of this model ('survey time'; $X^2 = \text{inf}$, p = 0, ĉ = inf). The effect of

covariates on detection and occupancy estimates in 2019 were negligible ($\Delta AIC_c < 2$). There was no

evidence for lack of fit in 2019 ($X^2 = 33.86$, p = 0.66, ĉ = 0.62). Habitat abundances were lower in

2018 and 2019 compared to 2017, with only abundance range overlap in the *Pisonia - Antirhea*

mountain and miscellaneous urban/agricultural or disturbed land habitat types (Fig. 5); compared to

2017, average decline in iguana abundance among habitats was 23.8% and 22.0% for 2018 and

2019, respectively. Overall, a two-way additive ANOVA indicated no significant difference of

iguana abundance among years ($F_{2,14} = 3.49$, $p = 0.059$), and a significant difference between

habitats ($F_{7,14} = 3.76$, $p = 0.017$).

227
228 **Fig. 3** The effect of elevation on abundance (lambda) estimates of *Iguana delicatissima* based on transect

229 surveys conducted on St. Eustatius in 2017 (pre-hurricane), 2018 and 2019 (post-hurricane).

230
231 **Fig. 4** The effect of elevation on site occupancy estimates of *Iguana delicatissima* based on transect surveys

232 conducted on St. Eustatius in 2017 (pre-hurricane), 2018 and 2019 (post-hurricane).

Fig. 5 Mean iguana abundance estimates per habitat type for 2017–2019, with standard deviations bars.

Habitat following De Freitas et al. (2012): 1 = *Pisonia - Antirhea* mountains; 2 = miscellaneous

urban/agricultural or disturbed land; 3 = *Pisonia - Justicia* hills; 4 = *Capparis - Pisonia* mountains; 5 =

Rauvolfia - Antigonon mountains; 6 = *Coccoloba - Bothriochloa* cliffs; 7 = *Pisonia - Eugenia* mountains; 8 =

Chionanthus - Nectandra mountains.

Table 2 Overview of 2017–2019 site occupancy estimates per habitat of *Iguana delicatissima* on St.
Eustatius. Data represents the probability of a site to be occupied (ψ), as well as confidence intervals (CI).
Habitat following De Freitas et al. (2012): 1 = *Pisonia - Antirhea* mountains; 2 = miscellaneous
urban/agricultural or disturbed land; 3 = *Pisonia - Justicia* hills; 4 = *Capparis - Pisonia* mountains; 5 =
Rauvolfia - Antigonon mountains; 6 = *Coccoloba - Bothriochloa* cliffs; 7 = *Pisonia - Eugenia* mountains; 8 =
Chionanthus - Nectandra mountains.

Habitat	2017		2018		2019	
	ψ	CI	ψ	CI	ψ	CI
1	0.98	<0.001–1.00	0.37	0.07–0.81	0.37	0.12–0.71
2	0.98	<0.001–1.00	0.35	0.10–0.73	0.37	0.16–0.63
3	0.99	<0.001–1.00	0.34	0.11–0.66	0.37	0.19–0.59
4	0.99	<0.001–1.00	0.32	0.11–0.64	0.36	0.18–0.61
5	0.99	<0.001–1.00	0.31	0.09–0.66	0.36	0.13–0.67
6	0.99	<0.001–1.00	0.29	0.06–0.72	0.36	0.09–0.76
7	0.99	<0.001–1.00	0.28	0.04–0.78	0.36	0.06–0.83
8	1.00	<0.001–1.00	0.26	0.02–0.84	0.35	0.04–0.88

Bead-tag data

Opportunistic sighting data were collected during 76, 78 and 56 days in 2017, 2018 and 2019, respectively. Assessment of the geographic distribution of those data indicates both time and geographic effort was comparably equal for 2017 and 2018, allowing a direct comparison (see Electronic Supplementary Information), though not for 2019. Of the 145 uniquely bead-tagged iguanas observed in 2017, 36 (24.8%) were observed during 2018.

Discussion

Our study demonstrates that multiple *I. delicatissima* population parameters decreased following the high-intensity hurricane year of 2017, with a decrease in estimates of population size, abundance, site occupancy, and opportunistic sightings of tagged iguanas. We did not observe a clear boomerang effect, as 2018-2019 parameters were mostly stable. These data are important to consider from a conservation and long-term stability perspective, given the small sizes of remaining populations of this highly endangered species. Overall, we present a rare effect-analysis of hurricane impact on a wild population of a large, long-lived reptile.

Iguana delicatissima inhabits most vegetation types within its native range, e.g., mangroves, high evergreen forest, dry shrublands, and gardens (for an overview see Henderson and Powell [2009]). On St. Eustatius, earlier studies found that iguana abundance was highest on vegetated slopes/cliffs and on the northwest flank of the dormant volcano where large estates are built within the tall evergreen forest (Fogarty et al. 2004; Debrot et al. 2013). Similarly, we found that iguana abundance was high in habitats on the western slopes of the volcano, though transects were not placed in gardens (Fig. 5). Thus, besides gardens, also the periferic habitats of this residential area are important iguana habitats. Indeed, Knapp and Perez-Heydrich (2012) found that biometric variables did not differ between iguanas living in- and outside of gardens. Furthermore, highest iguana abundance was found in the *Chionanthus - Nectandra* mountain habitat (De Freitas et al. 2012), at high (386m) elevation on the volcano, however this habitat was covered by a single

transect, thus we lack a perspective of variation. Additionally, we found higher iguana abundances in the *Pisonia-Eugenia* mountain habitat, located on the lower/mid (150–200m) south-western slopes of the Quill. This habitat is described as a secondary forest and, according to van Andel et al. (2016), "has a much lower diversity, density and canopy height than the forest on the wetter slope". Nevertheless, it is likely suitable and heterogenous habitat for iguanas and supports a variety of tree species including *Bourreria baccata, Pisonia subcordata, Guettarda scabra* and *Bursera simaruba* (van Andel et al. 2016) of which several are a known dietary source (Angin and Questel in prep.).

In contrast, *Rauvolfia - Antigonon* habitat is described as disturbed, regenerating shrubland with few trees, low floristic diversity and is covered extensively by an invasive vine (van Andel et al. 2016). Our results show that iguana abundance is surprisingly relatively high in this habitat type. We note that the patch covered by our survey efforts appears of higher quality than the remainder of this habitat type on the south-western side of the island, which is heavily grazed by free-roaming herbivores (van Andel et al. 2016; Madden 2020). Namely, the included patch has larger trees, is less degraded, and lies adjacent to secondary roads and semi-urban development. Consequently, we believe that our abundance estimates only apply to the covered patch of *Rauvolfia - Antigonon* habitat.

Detection probability increased from 2017 to 2019 but was overall low (<0.5) across all years. Nevertheless, detection probability estimates were similar to/higher than those from comparable studies (Bock et al. 2016; Rivera-Milán and Haakonsson 2020). However, when an iguana was previously observed, detection probability was equal across years. We did not test for observer effort, which may account for unequal detection between years and should ideally be consistent (Burton and Rivera-Milán 2014).; for our surveys, average number of observers was constant over the years (between 2–2.4), and surveys were always led by an experienced iguana-fieldwork technician. Vegetation damage (e.g., broken branches) may have resulted in higher detection probability post-hurricane compared to pre-hurricane, though generally iguanas are harder to detect in forested areas than other habitats. We were unable to locate specific published studies

on iguanas to support this, but reptile species overall are cryptic and considered difficult to study

(e.g., Durso et al. 2011; Sewell et al. 2012).

Our study provides evidence that the St. Eustatius *I. delicatissima* population declined

during 2017, during which two major catastrophic events hit the island; Hurricanes Maria and Irma.

As behavioural data of iguanas during such events are absent, pinpointing the exact cause of decline

is more challenging; how did iguanas die, and was that during or after the events? Post-event

mortality, in the absence of sustained physical injury, could result from a lower abundance of food.

Indeed, the majority of trees were defoliated due to high wind speeds and likely salt spray (Eppinga

and Pucko 2018). However, as not all trees were defoliated, and iguanas are known to survive

extended periods of drought (van Buurt 2010), we anticipate that post-event mortality due to

starvation had a minor contribution to the observed declines. Instead, we hypothesize that hurricane

conditions, either led to deadly physical injuries or translocated iguanas into the Caribbean Sea (see

Censky et al. 1998). Whether non-deadly on-island translocation is possible is unknown, though as

both abundances declined for all habitat and opportunistic sightings declined drastically, this

possibility appears unlikely. Although challenging to collect, future data on iguana movement

during hurricane events could provide more insight.

We found evidence for overdispersion when calculating site occupancy measures for 2017

and 2018, which resulted in inflated standard errors (Table 2). However, given the lack of

significance of covariates overall (only 'survey time' increased detection probability in 2018), we

chose to retain AICc model weights than pursue a quasi-likelihood theory (MacKenzie and Bailey

2004). We recognize the limitations of our study with regard to the assumptions of occupancy

modeling (Murray and Sandercock 2020), however we are confident that we met most if not all of

the assumptions. We assumed that iguana detection was independent between sites, given the

species' limited home range (van Wagensveld et al. in prep.), and that the sites surveyed were

discrete patches of habitat (except one survey track in the *Chionanthus - Nectandra* mountains

habitat). Nevertheless, we recognize that there may be more variation in the observed site

occupancy data than expected by the models. Considering the already wide confidence intervals,

however, we are of the opinion that the models presented offer a realistic representation of the pre-

/post-hurricane population data.

Despite the small size of St. Eustatius, our results suggest that the 2017 hurricanes affected

iguanas disproportionately across the island. Namely, although abundance and occupancy decreased

island wide, iguanas at higher elevations suffered stronger declines (Fig. 3–4). This finding is

corroborated by data from Eppinga and Pucko (2018) who assessed the impact of both 2017

hurricanes on the forests of St. Eustatius and found that a higher percentage of trees was snapped at

higher elevations. Eppinga and Pucko (2018) further show that over 90% of all trees were defoliated

for >75% and over 75% of trees lost primary branches, suggesting that Irma and Maria heavily

affected arboreal species. Indeed, hurricane wind speeds are stronger at higher elevations

(Holthuijsen et al. 2012), whereby our findings could be explained by increased mortality through

physical injury during hurricane events, as well as subsequently through longer food unavailability

(Cely 1991).

Understanding the natural movements of study species is important to consider in

population assessments (Neilson et al. 2018). Both *Iguana* species are known to remain static

throughout most of their life cycle, with movement mainly restricted to the hatchling life stage until

settling in breeding populations, and by adult females while traveling to nesting sites (Burghardt

and Rand 1985; Bock and McCracken 1988; Knapp et al. 2016). For *I. delicatissima*, the

reproductive cycle of populations on arid islands is clearly defined with nesting mainly occurring in

June-August (van den Burg et al. 2018a). Reproductive data for the St. Eustatius population remains

limited but appears to occur in June-July (pers. obs. by authors), which was considered while

planning field surveys. As no large communal nesting sites occur on St. Eustatius (as opposed to

Dominica, Knapp et al. 2016), and small isolated nesting sites have been located around the island

(Debrot et al. 2013; pers. obs. by authors), we suspect gravid females to undertake little movement

and nest within or close to their home range. This is further strengthened by the restricted size and

highly fragmented status of the population (Debrot et al. 2013; van den Burg et al. 2018b), with

interaction observations being rare (pers. obs. by authors).

Despite over half of Iguanidae diversity occurring in the Greater Caribbean region,

hurricane impacts on their populations have been rarely addressed. Beovides-Casas and Mancina

(2006) performed a total of 60 surveys (12 each during 4 visits) of the *Cyclura nubila* population on

Cayo Sijú (Cuba) in 2004, with the final visit post-Hurricane Ivan. Although Ivan passed at ~160

km distance with sea level rising by 2.5 meter, their data indicated the population size remained

stable. In another study, Hayes et al. (2004) report local extinction of a *Cyclura rileyi* population

from a small Bahamian cay following the passing of Hurricane Floyd (1999). However, it is

important to note that the authors indicated only three animals were observed by themselves there,

with a personal comment from the previous decade about 18 animals. Thus, the subsequent

reference of population extirpation following a hurricane event (e.g., Hayes et al. 2016) without

mention of these numbers and overall poor knowledge of the population in question appears

unjustified. Additionally, Hayes et al. (2004) also report no decrease in adult iguanas from a nearby

cay, with a large population. In our study, we found that multiple population parameters decreased

after an intense hurricane season; a decrease in I) population size of the studied area (23.3–26.5%)

and II) abundance (22–23.8%), as well as III) a 75% reduction in the number of opportunistic

sightings of tagged iguanas between 2017–2018. However, as studies remain scarce, equally are

generalizations and comparisons. Compared to *I. delicatissima*, *Cyclura* sp. are larger and more

robust, mainly non-arboreal iguanas. Although pre-hurricane behavior for *I. delicatissima* is

unknown, presumably lighter iguanas that remain arboreal during catastrophic weather events are

more vulnerable than heavier, terrestrial iguanas. However, inundation might force *Cyclura* sp. to

ascend trees in order to avoid being swept away given their range mostly includes low-elevation

islets.

To evaluate the impact of catastrophic events on closed populations, consideration of these

events' timing compared to a species' reproductive cycle is essential (Schoener et al. 2004). For St.

Eustatius, both 2017 hurricanes passed as strong early storms, preceding the main hatching period of the local *I. delicatissima* population while most eggs were incubating and therefore presumably safe in the absence of inundation. Later-seasonal storms are expected to have a more severe impact, as hatchlings are vulnerable to high wind speeds, thus potentially losing most individuals of an entire generation. Indeed, Hayes et al. (2004) compared hatching percentage between subsequent years and reported an extremely small hatchling cohort after the passing of Hurricane Floyd, which hit the Bahamas during the hatching period of *C. rileyi*. No information is available about the percentage of hatched clutches. On St. Eustatius only some nesting occurs on the island's lower elevations with the majority of nest sites found at elevations far higher than sea-level inundation (Debrot et al. 2013). Although data on egg survival and hatching success after inundation is known for some reptilians (Hsu et al. 2021), such data is missing for Iguanids.

Effect assessments of the 2017 Atlantic hurricane season are demonstrating its broad impact on ecosystems and their trophic levels; e.g., beach, mangrove and forest ecosystems (Liu et al. 2018; Barreto-Orta et al. 2019; Walker et al. 2019; Hall et al. 2020), salt- and freshwater fish communities (Meléndez-Vazquez et al. 2019; Neal et al. 2020), insect species and communities (Cabrera-Asencio and Meléndez-Ackerman 2021), birds (Palmer et al. 2018; Lloyd et al. 2019), and seas slugs (Middlebrooks et al. 2020). For St. Eustatius specifically, our data adds to the understanding of an ecosystem-wide hurricane impact during 2017 which, besides iguanas, affected both forests (Eppinga and Pucko 2018) and the populations of Bridled Quail-dove (Rivera-Milán et al. 2021) and Red-bellied racer (Madden et al. 2021). Importantly, these insights and novel risk assessments are demonstrating the high extinction risk that endemic, island species will face under projected climate change (Manes et al. 2021).

Conservation

From a conservation perspective, the status of *I. delicatissima* is at its most benign, with only five populations not directly threatened by on-island hybridization, of which just a single

island has a >200 population size; Petite Terre (Guadeloupe). Our study further highlights this

vulnerability, as local extinction or population declines to unsustainable sizes could be the result of

major hurricanes. Especially the extremely small population of Anguilla is vulnerable (Pounder et

al. 2020), which currently occurs on a small low-level islet. There, the potential of temporarily

holding iguanas to increase hurricane survival should be assessed, given adequate facilities can be

utilized. On St. Eustatius, we found no post-hurricane population recovery (or boomerang effect),

which could be linked to the small size of the existing population. Therefore, the projected increase

in frequency and intensity of catastrophic hurricanes (Bender et al. 2010) is troubling as populations

might be unable to recover between subsequent catastrophic events (McCoid 1996; Schriever et al.

2009; Selman 2015). However, we note that a short-term boomerang effect was likely absent given

the apparent low fecundity on St. Eustatius, possibly due to the population's low genetic diversity

(van den Burg et al. 2018b), although a detailed study on fecundity still is absent. Overall, our study

strengthens the need for population-increasing measures to be taken.

Acknowledgments
We thank Adam Mitchell, Sarinda Westerhout, Joey Markx, Rupert Redan, and all STENAPA staff
and volunteers that helped with data collection. Funding was provided through the Mohamed bin
Zayed Species Conservation fund (172517158), International Iguana Foundation, and, the BEST 2.0
program.

421
422

423 **References**
424 Angin B (2017) Plan National d'Actions pour le rétablissement de l'iguane des petites Antilles,
425 *Iguana delicatissima*, 2018–2022. 69p. + annexes.

426
427 Alberts AC (2004) Conservation strategies for West Indian rock iguanas (genus *Cyclura*): Current
428 efforts and future directions. Iguana 11:213–223.

429
430 Akresh ME, Askins AA, King DI, Hayes FE, Barry PE, Hayes WK (2020) Resilience in the
431 aftermath of hurricanes: fluctuations in a Critically Endangered population of West Indian
432 Woodpeckers *Melanerpes superciliaris nyeanus* over two decades. Bird Conserv Int
433 doi:10.1017/S0959270920000386

434
435 Barreto-Orta M, Méndez-Tejeda R, Rodriguez E, Cabrera N, Diaz E, Pérez K (2019) State of the
436 beaches in Puerto Rico after Hurricane Maria (2017). Shore Beach 87:16–23.

437
438 Beaudrot L, et al. (2016) Standardized assessment of biodiversity trends in tropical forest protected
439 areas: the end is not in sight. PLoS Biol 14:e1002357.
440

441 Behie AM, Pavelka MSM (2005) The short-term effects of a hurricane on the diet and activity of
442 Black Howlers (Alouatta pigra) in Monkey River, Belize. Folia Primatol 76:1–9.
443
444 Bender MA, Knutson TR, Tuleya RE, Sirutis JJ, Vecchi GA, Garner ST, Held IM (2010) Modeled
445 impact of anthropogenic warming on the frequency of intense Atlantic Hurricanes. Science
446 327:454–458.
447
448 Beovides-Casas K, Mancina CA (2006) Natural history and morphometry of the Cuban iguana
449 (*Cyclura nubila* Gray, 1831) in Cayo Siju, Cuba. Anim Biodivers Conserv 29:1–8.
450
451 Binns J, Burton F (2007) Bead Tags. International Reptile Conservation Foundation and Blue
452 Iguana Recovery Program.
453
454 Bock BC, McCracken GF (1988) Genetic structure and variability in the Green iguana (*Iguana
455 iguana*). J Herpetol 22:316–322.
456
457 Bock BC, Páez VP, Rand AS, Burghardt GM, Iverson JB, Grant TD, Knapp CR, Pasachnik SA
458 (2016) Life table and stochastic matrix projection analysis for a population of Green Iguanas
459 (Iguana iguana): implications for conservation and control. *Iguanas: biology, systematics, and
460 conservation.* Herpetol Conserv Biol 11:47–60.
461
462 Breuil M, Ibéné B (2008) Réponse au droit de réponse de Lorvelec et al. (2008). Bulletin Société
463 Herpétologique de France 128:49–52.
464
465 Burghardt GM, Rand AS (1985) Group size and growth rate in hatchling green iguanas (*Iguana
466 iguana*). Behav Ecol Sociobiol 18:101–104.
467
468 Burnham KP, Anderson DR (2002) Model selection and multimodel inference: a practical
469 information-theoretic approach. Second edition. Springer-Verlag, New York, USA.
470
471 Burton FJ, Rivera-Milán FF (2014) Monitoring a population of translocated Grand Cayman blue
472 iguanas: assessing the accuracy and precision of distance sampling and repeated counts. Anim
473 Conserv 17:40–47.
474
475 Cabrera-Asencio I, Meléndez-Ackerman EJ (2021) Community and Species-Level Changes of
476 Insect Species Visiting *Mangifera indica* Flowers Following Hurricane María: "The Devil Is in the
477 Details". Front Ecol Evol 9:556821.
478
479 Cely JE (1991) Wildlife effects of hurricane Hugo. J Coast Res 8:319–326.
480
481 Censky EJ, Hodge K, Dudley J (1998) Over-water dispersal of lizards due to hurricanes. Nature
482 395:556.
483
484 De Freitas JA, Rojer AC, Nijhof BSJ, Debrot AO (2012) A landscape ecological vegetation map of
485 Sint Eustatius (Lesser Antilles). IMARES, CARMABI and Royal Netherlands Academy of Arts
486 and Sciences, Amsterdam.
487
488 Debrot AO, Boman E, Madden H (2013) The Lesser Antillean Iguana on St. Eustatius: A 2012
489 population status update and causes for concern. IRCF Reptiles & Amphibians 20:44–52.
490

Dénes FV, Silveira LF, Beissinger SR (2015) Estimating abundance of unmarked animal populations: accounting for imperfect detection and other sources of zero inflation. Methods Ecol & Evol 6:543–556.

Donihue CM, Herrel A, Fabre A-C, Kamath A, Geneva AJ, Schoener TW, Kolbe JJ, Losos JB (2018) Hurricane-induced selection on the morphology of an island lizard. Nature 560:88–91.

Donihue CM, Kowaleski AM, Losos JB, Algar AC, Baeckens S, Buchkowski RW, Fabre A-C, Frank HK, Geneva AJ, Reynolds RG, Stroud JT, Velasco JA, Kolbe JJ, Mahler DL, Herrel A (2020) Hurricane effects on Neotropical lizards span geographic and phylogenetic scales. PNAS 117:10429–10434.

Dufour CMS, Donihue CM, Losos JB, Herrel A (2019) Parallel increases in grip strength in two species of *Anolis* lizards after a major hurricane on Dominica. J Zool 309:77–83.

Durso AM, Willson JD, Winne CT (2011) Needles in haystacks: estimating detection probability and occupancy of rare and cryptic snakes. Biol Conserv 144:1508–1515.

Eppinga MB, Pucko CA (2018) The impact of hurricanes Irma and Maria on the forest ecosystems of Saba and St. Eustatius, northern Caribbean. Biotropica 50:723–728.

Fiske I, Chandler R (2011) unmarked: An R package for fitting hierarchical models of wildlife occurrence and abundance. J Stat Softw 43:1–23.

Fogarty SP, Zero VH, Powell R (2004) Revisiting St. Eustatius: estimating the population size of Lesser Antillean Iguanas, *Iguana delicatissima*. Iguana 11:139–146.

Hall J, Muscarella R, Quebbeman A, Arellano G, Thompson J, Zimmerman JK, Uriarte M (2020) Hurricane-Induced Rainfall is a Stronger Predictor of Tropical Forest Damage in Puerto Rico Than Maximum Wind Speeds. Sci Rep 10:4318.

Hayes WK, Carter RL, Cyril Jr S, Thornton B (2004) Conservation of an endangered Bahamian Rock Iguana, I. Population assessments, habitat restoration, and behavioral ecology. Pp. 232–257 In Iguanas: Biology and Conservation. Alberts AC, Carter RL, Hayes WK, and Martins EP (Eds.). University of California Press, Berkeley and Los Angeles, California, USA.

Hayes W, Cyril S, Crutchfield T, Wasilewski J, Rothfus TA, Carter RL (2016) Conservation of the Endangered San Salvador Rock iguanas (*Cyclura rileyi rileyi*): Population estimation, invasive species control, translocation, and headstarting. Herp Conser Biol 11:90–105.

Henderson RW, Powell R (2009) Natural history of West Indian reptiles and amphibians. University of Florida Press, Gainesville, Florida, USA.

Holthuijsen LH, Powell MD, Pietrzak JD (2012) Wind and waves in extreme hurricanes. J Geophys Res 117:C09003.

Hsu M-H, Lin J-W, Liao C-P, Hsu J-Y, Huang W-S (2021) Trans-marine dispersal inferred from the saltwater tolerance of lizards from Taiwan. PLoS ONE 16:e0247009.

540 Jacobson AP, Riggio J, Tait AM, Baillie JEM (2019) Global areas of low human impact ('Low
541 Impact Areas') and fragmentation of the natural world. Sci Rep 9:14179.
542
543 Johnson AB, Winker K (2010) Short-term hurricane impacts on a neotropical community of marked
544 birds and implications for early-stage community resilience. PLoS ONE 30:e15109.
545
546 Knapp CR, Valerie S (2008) *Iguana delicatissima*. Mortality. Herpetol Rev 39:227–228.
547
548 Knapp CR, Perez-Heydrich C (2012) Using non-conspicuous metrics to examine selected impacts
549 of disturbance on a long-lived reptile. Endanger Species Res 17:193–200.
550
551 Knapp CR, Prince L, James A (2016) Movements and nesting of the Lesser Antillean iguana (*Iguana
552 delicatissima*) from Dominica, West Indies: Implications for conservation. Pp. 154–167 In Iguanas:
553 Biology, Systematics, and Conservation. Iverson JB, Grant TD, Knapp CR, Pasachnik SA (Eds.).
554 Herpetol Conserv Biol 11 (Monograph 6).
555
556 Knapp CR, Grant TD, Pasachnik S, Angin B, Boman E, Brisbane J, Buckner SD, Haakonsson JE,
557 Harlow PS, Mukhida F, Thomas-Moko N, van den Burg MP, Wasilewski JA (2020) The global
558 need to address threats from invasive alien iguanas. Anim Conserv ACV12660.
559 doi:10.1111/acv.12660.
560
561 Liu X, Zeng X, Zou X, González G, Wang C, Yang S (2018) Litterfall Production Prior to and
562 during Hurricanes Irma and Maria in Four Puerto Rican Forests. Forests 9:367.
563
564 Lloyd JD, Rimmer CC, Salguero-Faría JA (2019) Short-term effects of hurricanes Maria and Irma
565 on forest birds of Puerto Rico. PLoS ONE 14:e0214432.
566
567 Lorvelec O, Levesque A, Barré N, Feldmann P, Leblond G, Jaffard M-E, Pascal M, Pavis C (2004)
568 Evolution de la densité de population de l'iguane des Petites Antilles (*Iguana delicatissima*) dans la
569 réserve naturelle des Îles de la Petite Terre (Guadeloupe) entre 1995 et 2002. Revue d'Écologie (La
570 Terre et la Via) 59:331–344.
571
572 Losos JB, Schoener TW, Spiller DA (2003) Effect of immersion in seawater on egg survival in the
573 lizard *Anolis sagrei*. Oecologia 137:360–362.
574
575 MacKenzie DI, Nichols JD, Lachman GB, Droege S, Royle JA, Langtimm CA (2002) Estimating
576 site occupancy rates when detection probabilities are less than one. Ecol 83:2248–2255.
577
578 MacKenzie DI, Bailey LL (2004) Assessing the fit of site-occupancy models. J Agric Biol Environ
579 Stat 9(3):300–318.
580
581 MacKenzie DI, Nichols JD, Royle JA, Pollock KH, Bailey L, Hines JE (2017) Occupancy
582 estimation and modeling: inferring patterns and dynamics of species occurrence. Elsevier,
583 Amsterdam.
584
585 Madden H, Fernández DS, Tremblay RL, Verdel K, Kaboord B (2021) Find me if you can: Pre- and
586 Post-hurricane Densities of the Red-bellied Racer (*Alsophis rufiventris*) on St. Eustatius, and a
587 review of the genus in the Caribbean. bioRxiv. https://doi.org/10.1101/2021.07.05.451169
588

589 Madden H (2020) Free-roaming livestock distribution, densities and population estimates on St.
590 Eustatius: 2020 update. Caribbean Netherlands Science Institute, unpublished report.
591
592 Manes S, Costello MJ, Beckett H, Debnath A, Devenish-Nelson E, Grey K-A, Jenkins R, Ming
593 Khan T, Kiessling W, Krause C, Maharaj SS, Midgley GF, Price J, Talukdar G, Vale MM (2021)
594 Endemism increases species' climate change risk in areas of global biodiversity importance,
595 Biol Conserv 109070, doi:10.1016/j.biocon.2021.109070.
596
597 Marroquin-Paramo JA, Suazo-Ortuno I, Urbina-Cardona N, Benitez-Malvido J (2021) Cumulative
598 effects of high intensity hurricanes on herpetofaunal assemblages along a tropical dry forest
599 chronosequence. For Ecol Manag 479:118505.
600
601 McCoid, MJ (1996) Effect of typhoons on the lizard community of a shelf atoll. Atoll Research
602 Bulletin 439:1–5.
603
604 Meléndez-Vazquez F, Olmeda-Saldana, Cruz J, Arcila D, Betancur R (2019) Effects of Hurricane
605 Maria in hamlet communities (Serranidae: *Hypoplectrrrus* spp.) in Puerto Rico. Ecol Indic
606 107:105591.
607
608 Meredith M (2020) wiqid: Quick and Dirty Estimates for Wildlife Populations. R package version
609 0.3.0. https://CRAN.R-project.org/package=wiqid
610
611 Middlebrooks ML, Curtis NE, Pierce SK (2020) The complete disappearance of a long standing
612 sacoglossan sea slug population following Hurricane Irma, despite recovery of the local algal
613 community. Symbiosis 80:231–237.
614
615 Morcilo DO, Steiner UK, Grayson KL, Ruiz-Lambides AV, Hernandez-Pacheco R (2020)
616 Hurricane-induced demographic changes in a nonhuman primate population. R Soc Open Sci
617 7:200173.
618
619 Murray DL, Sandercock BK (eds.) (2020) *Population ecology in practice*. John Wiley & Sons.
620
621 Neal JW, Haley BM, Moran ZS (2020) Hurricane Maria in Puerto Rico: Effects on Reservoir Water
622 Quality and Fish Community Structure and Resilience. J Southeastern Association Fish Wildlife
623 Agencies 7:123–133.
624
625 Neilson EW, Avgar T, Burton AC, Broadley K, Boutin S (2018) Animal movement affects
626 interpretation of occupancy models from camera-trap surveys of unmarked animals. Ecosphere
627 9:e02092.
628
629 NOAA (2020) available at https://www.nhc.noaa.gov/gis/archive_besttrack.php?year=2017
630
631 Palmer CA, Martin TE, Durand S, Lamont M (2018) First observations of the impacts of Hurricane
632 Maria on the endemic imperial amazon. Oryx 52:410–411.
633
634 Pasachnik SA, Shew JJ, Townsend JH, Powell R (2002) *Iguana delicatissima* (Lesser Antillean
635 Iguana). Activity. Herpetol Rev 33:51–52.
636

Pasachnik SA, Montgomery CE, Ruyle LE, Corneil JP, Antunez EE (2012) Morphological and demographic analyses of the Black-chested Spiny-tailed iguana, *Ctenosaura melanosterna*, across their range: Implications for population level management. Herpetol Conserv Biol 7:399–406.

Pounder KC, Mukhida F, Brown RP, Carter D, Daltry JC, Fleming T, Goetz M, Halsey LG, Hughes G, Questel K, Saccheri IJ, Williams R, Soanes LM (2020) Testing for hybridisation of the Critically Endangered *Iguana delicatissima* on Anguilla to inform conservation efforts. Conserv Genet 21:405–420.

Powell R (2004) Conservation of iguanas (*Iguana delicatissima* and *I. iguana*) in the Lesser Antilles. Iguana 11:239–246.

Powers RP, Jetz W (2019) Global habitat loss and extinction risk of terrestrial vertebrates under future land-use-change scenarios. Nat Clim Change 9:323–329.

R Core Team (2019) A Language and Environment for Statistical Computing. Version 3.5.1. Available at www.R-project.org/.

Rabe AM, Herrmann NC, Culbertson KA, Donihue CM, Prado-Irwin SR (2020) Post-hurricane shifts in the morphology of island lizards. Biol J Linnean Soc 130:156–165.

Reagan DP (1991) The response of *Anolis* lizards to hurricane-induced habitat changes in a Puerto Rican rain forest. Biotropica 23:468–474.

River-Milan FF, Madden H, Verdel K (2021) Bridled Quail-dove *Geotrygon mystacea* population assessment after hurricanes Irma and Maria, St. Eustatius, Caribbean Netherlands. Bird Conserv Int 1–12.

Rojer AC (1997) Biological inventory of Sint Eustatius. KNAP-Project 96-10. Report, CARMABI Foundation, Curaçao, Netherlands Antilles. 50 pp.

Royle JA, Nichols JD (2003) Estimating abundance from repeated presence–absence data or point counts. Ecology 84:777–790.

Royle JA (2004) N-mixture models for estimating population size from spatially replicated counts. Biometrics 60:108–115.

Schoener TW, Spiller DA, Losos JB (2001) Natural restoration of the species-area relation for a lizard after a hurricane. Science 294:1525–1528.

Schoener TW, Spiller DA, Losos JB (2004) Variable ecological effects of hurricanes: The importance of seasonal timing for survival of lizards on Bahamian islands. PNAS 101:177–181.

Schriever, TA, Ramspott, J, Crother, BI, Fontenot, CL (2009) Effects of hurricanes Ivan, Katrina, and Rita on a southeastern Louisiana herpetofauna. Wetlands 29(1):112–122.

Schultz JMM, Kossin JP, Shepherd JM, Ransdell JM, Walshe R, Kelman I, Galea S (2018) Risks,
health consequences, and response challenges for small-island-based populations: Observations
from the 2017 Atlantic hurricane season. Disaster Med Public Health Prep 13:5–17.

Selman W (2015) Herpetofaunal diversity and seasonality from a remnant coastal chenier forest in
southwestern Louisiana. Southeastern Naturalist 14(3):491–505.

Sewell D, Guillera-Arroita G, Griffiths RA, Beebee TJ (2012) When is a species declining?
Optimizing survey effort to detect population changes in reptiles. PLoS ONE 7:e43387.

Spiller DA, Losos JB, Schoener TW (1998) Impact of a catastrophic hurricane on island
populations. Science 281:695–697.

Statistics Netherlands (2020) Statistics Netherlands, available at https://www.cbs.nl/en-gb.

Trewartha GT, Horn LH (1980) An Introduction to climate. McGraw-Hill, New York.

van Andel T, van der Hoorn B, Stech M, Bantjes Arostegui S, Miller J (2016) A quantitative
assessment of the vegetation types on the island of St. Eustatius, Dutch Caribbean. Global Ecol
Conserv 7:59–69.

van Buurt G (2010) A Short Natural History of Curaçao. In: Crossing Shifting Boundaries,
Language and Changing Political status in Aruba, Bonaire and Curaçao. Proceedings of the ECICC-
conference, Dominica 2009, Volume I, pp. 229-256. Faraclas, N., Severing, R., Weijer, C. Echteld,
E. (Eds.), FPI and UNA, Curaçao.

van den Burg M, Breuil M, Knapp C (2018a) *Iguana delicatissima.* The IUCN Red List of
Threatened Species 2018: e.T10800A122936983. https://dx.doi.org/10.2305/IUCN.UK.2018-
1.RLTS.T10800A122936983.en

van den Burg MP, Meirmans PG, van Wagensveld TP, Kluskens B, Madden H, Welch ME,
Breeuwer JAJ (2018b) The Lesser Antillean Iguana (*Iguana delicatissima*) on St. Eustatius:
genetically depauperate and threatened by ongoing hybridization. J Hered 109:426–437.

van den Burg MP, Madden H, van Wagensveld TP, Buma C (2018c) Anthropogenic mortality in
the Critically Endangered Lesser Antillean iguana (*Iguana delicatissima*) onn St. Eustatius. IRCF
Reptiles & Amphibians 25:120–124.

van den Burg MP, Brisbane JLK, Knapp CR (2020) Post-hurricane relief facilitates invasion and
establishment of two invasive alien vertebrate species in the Commonwealth of Dominica, West
Indies. Biol Invasions 22:195–203.

van Wagensveld TP, van den Burg MP (2018) First record on fecundity of an *Iguana* hybrid and its
implications for conservation: evidence for genetic swamping by non-native iguanas. Herpetology
Notes 11:1079–1082.

Vuillaume B, Valette V, Lepais O, Grandjean F, Breuil M (2015) Genetic evidence of hybridization
between the endangered native species *Iguana delicatissima* and the invasive *Iguana iguana*
(Reptilia, Iguanidae) in the Lesser Antilles: management implications. PLoS ONE 10:e0127575.
732

733 Walcker R, Laplanche C, Herteman M, Lambs L, Fromard F (2019) Damages caused by hurricane
734 Irma in the human-degraded mangroves of Saint Martin (Caribbean). Sci Rep 9:18971.
735

736 Webster PJ, Holland GJ, Curry JA, Chang H-R (2005) Changes in tropical cyclone number,
737 duration, and intensity in a warming environment. Science 309:1844–1846.
738

739 Wiley JW, Wunderle JM (1993) The effects of hurricanes on birds, with special reference to
740 Caribbean islands. Bird Conserv Int 3:319–349.