Comparative transcriptomics of tropical woody plants supports fast and furious strategy along the leaf economics spectrum in lianas

U. Uzay Sezen [1*], Samantha J. Worthy[2], Maria N. Umaña[3], Stuart J. Davies[4,5], Sean M. McMahon[1], Nathan G. Swenson[6,2],

1 Smithsonian Environmental Research Center, 647 Contees Wharf Rd, Edgewater, Maryland 21037 USA.
2 Department of Biology, University of Maryland, College Park, Maryland, 20742 USA.
3 Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, Michigan 48109 USA.
4 Forest Global Earth Observatory, Smithsonian Tropical Research Institute, Panama.
5 Department of Botany, National Museum of Natural History, Smithsonian Institution, Washington DC, USA.
6 Department of Biological Sciences, University of Notre Dame, Notre Dame, Indiana 46556 USA.

* uzay.sezen@gmail.com

Abstract

Lianas, climbing woody plants, influence the structure and function of tropical forests. Climbing traits have evolved multiple times, including ancestral groups such as gymnosperms and pteridophytes, but the genetic basis of the liana strategy is largely unknown. Here, we use a comparative transcriptomic approach for 47 tropical plant species, including ten lianas of diverse taxonomic origins, to identify genes that are consistently expressed or downregulated only in lianas. Our comparative analysis of full-length transcripts enabled the identification of a core interactomic network common to lianas. Sets of transcripts identified from our analysis reveal features related to functional traits pertinent to leaf economics spectrum in lianas, include upregulation of genes controlling epidermal cuticular properties, cell wall remodeling, carbon concentrating mechanism, cell cycle progression, DNA repair and a large suit of downregulated transcription factors and enzymes involved in ABA-mediated stress response as well as lignin and suberin synthesis. All together, these genes are known to be significant in shaping plant morphologies through responses such as gravitropism, phyllotaxy and shade avoidance.

Introduction

Lianas are woody vines that have evolved multiple times since at least the Devonian period in vascular plant lineages, including in the Gnetales, Coniferales, and repeatedly in the Angiosperms with fascinating diversity in stem anatomy even within a single genus [1,2]. The Neotropics carry close to 11,000 liana species belonging to 977 genera

under 119 families [3], while the Old-world tropics contain about 12,000 species found in 143 families and 1415 genera [4]. In tropical forests, lianas may account for as much as 25% of the woody plant species, but can also represent 40% of canopy leaf cover [5,6]. As structural parasites, lianas affect host tree recruitment, growth, survival, and reproduction [7–9]. Lianas also play significant roles in the cycling of carbon, nitrogen and water, and can hold back recovery after gap-forming events [6,10–17]. Despite their exceptional diversity and ecological importance, little is known about the genetic characteristics that distinguish them from other vascular plants that help explain their convergence.

The basic advantage of the liana life form is clear: lianas harvest light from the treetops without investing in the woody structures necessary to support a canopy of leaves [18]. Several trade-offs emerge from this strategy of structural parasitism that might provide a path toward understanding convergent liana phenotypes. Specifically, in order to gain access to the canopy, lianas need to balance leaf area, length of vascular tissues, and tolerance to high temperature, excess light and dehydration which suggest potential underlying genetic mechanisms towards a liana growth form in multiple plant families [19,20]. In the following, we consider each of these challenges to the liana lifestyle as a means of generating expectations regarding where similarities in gene expression profiles among a phylogenetically diverse set of lianas may exist. The liana life form involves a relatively longer stem length compared to that of trees culminating with a large total leaf area that sprawls over the crowns of many trees would make lianas more vulnerable to xylem vessel embolism than trees and shrubs [21,22]. Liana leaves must have evolved adaptations that complement and compensate for hydraulic limitations and occupy a distinct range along the leaf economic spectrum (LES) [23,24]. Compared to trees, lianas tend to have thinner blades with higher specific leaf area (SLA) and nutrient content. However, their leaves can also exhibit higher photosynthetic efficiencies despite the fact that the general leaf qualities reflect inexpensive, disposable, short longevity structures with little or no investment on defense [25].

The direct solar radiation and high heat in the canopies, where most liana leaves are deployed can impose significant constraints for lianas. For instance, the photosystem II has been shown to be vulnerable to temperature and light stress in a liana as compared to its tree relative [26]. Canopy trees have the potential to mediate this damage through higher leaf area with thinner boundary layers for efficient cooling. There is evidence that liana leaves have lower leaf transmittance values, better for light-capture, but worse for damage to tissues [26,27]. It is highly possible that lianas have evolved certain epicuticular and parenchymal properties that has a set upper limit for safe light harvest and deflect excessive photosynthetically active radiation. When leaf temperature exceeds photosynthetic optimum, CO_2 assimilation rates and stomatal conductance decrease while respiration increases [28]. Seasonal observations indicate that carbon fixation and efficiencies of water and nitrogen use are especially high in lianas during drought giving them a growth advantage [29–31]. We, therefore, expect that leaf surface qualities and mesophyll properties should be under selective pressures for dealing with light, thermal and dehydration stress. Guard cell openings can be set to maintain a steady rate of water loss enough to last through the day and fron *Vitis vinifera* (Grapevine from here in) we know that lianas can employ a range of anisohydric and isohydric control of stomatal conductance as water use strategy. Through their cheap but efficient leaves, lianas may also be carrying out sufficient levels of photosynthesis rapidly before midday to escape from the Sun's pernicious rays.

There has been some research on stress responses of a few liana species [32,33]. However, most of the information comes from Grapevine due to the large amount of genomic and transcriptomic information generated for the species [34–38]. A lack of high throughput sequencing data for additional liana species has reduced our ability to

explore their potential genetic commonalities that distinguish them from non-liana plants.

Here, we provide a comparative transcriptomic analysis across a phylogenetically diverse group of trees, shrubs, and lianas. We use reference transcriptome assemblies of 37 tree and shrub species and 10 liana species (SI Fig. S1, SI Dataset S1). Using network analyses, we ask the following questions: (1) what genes are uniquely expressed and downregulated in some or all of the lianas? (2) are there unique genetic similarities among lianas related to their shared ecological challenges (e.g., long distance nutrient transport, stress including hydraulic, thermal and light, leaf tissue quality, unidirectional growth and light sensing)? and (3) are there unique genetic similarities in lianas that reflect specific metabolic pathways underpinning the building of structural biomass?

Materials and Methods

Sample set.

The Luquillo transcriptomic set included six lianas (DIO) *Dioscorea poligonoides* (Dioscoreaceae), (HET) *Heteropterys laurifolia* (Malpighiaceae), (PAU) *Paulinia pinnata* (Sapindaceae), (SEC) *Securidaca virgata* (Polygonaceae), (SMI) *Smilax coriaceae* (Smilaceae), (DOL) *Dolichandra unguis-cati* (Bignoniaceae). We also included four more liana species from National Center for Biotechnology Information's Sequence Read Archive (NCBI-SRA) and One Thousand Plant Genomes (1KP) dataset [39, 40] with the following accessions (GNE) *Gnetum montanum* (Gnetaceae) SRR5908685, (SMISIE) *Smilax sieboldii* (Smilacae) SRR5134200, (PASCAE) *Passiflora caerulea* (Passifloraceae) 1KP id:SIZE, (SCHPAR) *Schlegelia parasitica* (Schlegeliaceae) 1KP id:GAKQ. A full list including non-liana species can be found in the Supplementary Information (SI Dataset S1, SI Fig. S1). An NCBI-SRA project accession number for the Luquillo transcriptomic set will be released upon acceptance of the manuscript.

Sample collection and RNA library construction.

To analyze the transcriptomes, we chose healthy and fully developed leaves from seedlings of tree and liana species distributed between 350 and 450 m in elevation from the Luquillo Experimental Forest (LEF) in the north eastern part of Puerto Rico. For each species, approximately 5 grams of leaf tissue was collected and placed in a 50 mL polypropylene conical tube with RNAlater (Thermo Fisher Scientific, Waltham, MA, USA). Explants were cut with a razor blade prior to being placed in the tube to allow the RNAlater to penetrate the mesophyll quickly. Samples were then frozen at -80ºC within two days. Rneasy Plant Mini Kit (Qiagen, Valencia, CA, USA) was used for RNA extraction. RNA quantification and quality metrics were carried out using a NanoDrop 2000 spectrophotometer (NanoDrop Products, Wilmington, DE, USA) and an Agilent Bioanalyzer 2100 (Agilent Technologies, Santa Clara, CA, USA) RNAseq library preparations and sequencing were performed at the Beijing Genomics Institute, Shenzen, China on Illumina Hiseq 2000 sequencer generating 100 bp paired-end reads.

Bioinformatics.

Paired-end raw Illumina reads were trimmed, and quality filtered using Sickle (-q 35 -l 30 for minimum quality score and retaining sequences longer than 30 nt). Trimmed fastq files were assembled by Trinity v.2.6.6 with minimum contig length 300 [41]. Quality of the assemblies were assessed by QUAST (Quality Assessment Tool for Genome Assemblies), validated by Bowtie2, and analyzed by BUSCO for completeness

using the Embryophyta (embryophyta_odb9) benchmark gene set [42–44] (SI Dataset S1). Transcriptome assemblies were annotated by EnTAP using the Diamond high performance aligner interrogating four protein databases (Uniprot, RefSeq plant proteins 94, RefSeq complete protein 94, RefSeq non-redundant protein 94) [45, 46]. Bacterial, archaeal and non-plant eukaryotic contaminants were filtered using Opisthokonta as the taxonomical cut-off (SI Dataset S3). Transcriptomes were frame selected and translated into proteins through built-in GeneMarkS-T module within EnTAP [47] (SI Dataset S3). Full-length protein sequences were filtered from EnTAP results and were blasted against indexed *V. vinifera* proteome v29720. Top hits were selected through usearch (-ublast -evalue 1e-9 -query_cov 0.9) [48]. Blast results were matched into Grapevine UNIPROT IDs and used as input into STRING database for interactome network construction [49, 50]. Co-downregulated and co-expressed protein-protein interactome networks were imported into Cytoscape and merged into a single network [51] (SI Dataset S2). The resulting interactome network was annotated with colored borders and fills to highlight biologically informative nodes. Phylogenetic tree of lianas and non-lianas was constructed using the TimeTree resource compiling evolutionary divergence times derived from molecular sequence data [52]. Gene Ontology (GO) terms were interrogated using g:Profiler by providing gene lists corresponding to UNIPROT IDs [53]. Results generated from g:Profiler are accessible through permalinks in SI Text.

A note on co-downregulation.

Co-downregulation should not be confused with gene deletions observed in the genomes of many parasitic and carnivorous plants [54]. Absence of gene activity does not necessarily mean gene loss or loss of function. Therefore, here we prefer to use the term co-downregulation to define orthologous genes that are most likely present in lianas but not actively transcribed. Similarly, we define co-expression as orthologous genes from climbers demonstrating a sufficiently high expression pattern compared to non-liana counterparts that allows detection as full-length transcripts.

Results

Transcriptome assembly and annotation.

Using a phylogenetically diverse sample of 10 liana species, we identified major convergent gene expression patterns that contrast with 37 coexisting species of trees and shrubs (SI Fig. S1, SI Fig. S3). Quality metrics of transcriptomes generated by Quality Assessment Tools for Genome Assemblies (QUAST) showed the transcriptomes range in size from 49,040 (*Tetragastris balsamifera*) to 182,060 (*Ixora ferrea*) (SI Dataset S1). The overall alignment rates for the *de novo* assembled reference transcriptomes, validated by Bowtie2 alignments of trimmed and quality filtered reads, were between 83 and 94 percent (SI Fig. S2A). The assembled transcriptomes captured an informative fraction of the expressed genes from 47 species where Benchmarking Universal Single-Copy Orthologs (BUSCO) analyses were able to represent between 52 and 92 percent of the benchmark eukaryotic genes within the Embryophyta (SI Fig. S2B, SI Dataset S1). Annotation and filtering of non-plant transcripts through EnTAP (Eukaryotic non-model Transcriptome Annotation Pipeline) resulted in a full-length set of Grapevine orthologs ranging from 6001 (*Smilax coriaceae*) to 9488 (*Dolichandra unguis-cati*) for cross-comparison (Table 1, SI Dataset S3).

Table 1. Ortholog numbers, families, and growth forms of species with full-length transcripts BLASTed onto Grapevine protein models.

Species	Family	Form	Orthologs
Dioscorea poligonoides	Dioscoreaceae	Woody	7289
Dolichandra unguis-cati	Bignoniaceae	Woody	9488
Gnetum montanum	Gnetaceae	Woody	7591
Heteropterys laurifolia	Malpighiaceae	Woody	7764
Paulinia pinnata	Sapindaceae	Woody	8275
Passiflora caerulea	Passifloraceae	Herbaceous	7237
Schlegelia parasitica	Schlegeliaceae	Herbaceous	8827
Securidaca virgata	Polygonaceae	Woody	8591
Smilax coriacea	Smilaceae	Woody	6002
Smilax sieboldii	Smilaceae	Herbaceous	8141
Non-liana	–	–	13267
Overall	–	–	18806

Figure 1. Protein orthologs of full-length liana and non-liana transcripts mapped onto Grapevine interactome and Manhattan plots of enriched GO terms. A, Protein-protein interactions of transcripts solely expressed in 10 lianas (the two Smilax species are represented as combined into SMI). Nodes shared by more than two lianas (L2-L5) and co-downregulated transcripts (salmon-colored borders) including their first shell interactors (gray colored borders) occupy distinctly sectorized footprints. GO terms represented molecular functions (MF) pertaining to the phytohormone auxin, biological processes (BP) including positive gravitropism and auxin transport. KEGG pathways included nitrogen metabolism and ABC transporters. No enrichment observed in cellular compartment (CC). **B** Nodes pertaining to mRNA metabolism (red border and green fill) and transcription factors (red border and black fill) are observed to be downregulated in all lianas in the dataset. GO term analysis of co-downregulated transcripts revealed terms such as regulation of gene expression concentrated around transcription factors. Molecular functions were enriched in DNA-binding transcription factor activity and sequence-specific DNA binding. Enriched biological processes included regulation of DNA templated transcription and regulation of macromolecule biosynthesis all localized to the nucleus.

A protein-protein interactome overlaid upon transcriptomic baselayer.

A Venn diagram of Grapevine protein orthologs corresponding to our transcripts revealed 827 protein models common to all 47 plant species (SI Dataset S1). After the exclusion of common proteins, a protein-protein interactome network based on curated STRING database was constructed using transcripts exclusively expressed by lianas and non-lianas. The total size of the network including vertices with first and second shells of interaction consisted of 3835 nodes and 14574 edges. Grapevine has 33,568 protein-coding genes and the network occupies a little over 11% of its gene space. Close to half of the network (1687 nodes) comprised genes solely expressed by non-lianas (Fig. 1). The rest included transcripts exclusively expressed by one or more lianas (Fig. 1A). The most connected node (VIT_11s0016g03430) with 199 interactions represented the core of the interactome corresponded to a protein phosphatase 2C (PP2C) also known as cyclic nucleotide-binding/kinase domain-containing protein (SI Fig. S3). This protein-serine/threonine phosphatase is a key enzyme in cGMP-dependent signaling, abscisic acid perception, commitment to cell cycle and is found within the additional shells of interaction in a region of the network subtending nodes exclusively expressed in lianas (Fig. 1, Table 2, SI Fig. S3). The subsequent highly connected nodes are all found within the co-downregulated portion of the network (Fig. 1B, Table 2)

Transcripts exclusively expressed in lianas.

The transcripts expressed exclusively by more than one liana (L2-L5) species generated a network of 705 nodes with 1609 edges (Fig. 1, SI Fig. S3). We found no genes that are unanimously expressed by all lianas. Two nodes, shared by five lianas (L5), corresponded to the ethylene responsive transcription factor SHINE2 (SHN2) and the Xyloglucan Glycosyltransferase 4/Cellulose Synthase-like C family 4 (CSLC4) enzyme, representing the highest points of convergence via individual gene transcripts (Table 2). The next set of most nodes shared by four lianas (L4) in different combinations included cell wall remodeling enzymes Pectin Methylesterase 7 (PME7), Glycosyl Transferase (GT), Polygalacturonase Inhibitor-like (PGIP) as well as a carbon concentrating enzyme alpha Carbonic Anhydrase 4 (alpha-CAH4), and the plant-specific cell cycle regulator F-box-like 17 (FBL17) (Table 2, Fig. S3).

The combined controlled vocabulary Gene Ontology (GO) analysis of transcripts shared by more than two lianas (L2-L5) through gProfiler displayed enriched molecular functions pertaining to the phytohormone auxin, such as auxin efflux transmembrane transporter activity (GO:0010329) and auxin transmembrane transporter activity (GO:0080161) (Fig. 1A). Biological processes included auxin associated activities such

as positive gravitropism (GO:0009958), auxin polar transport (GO:0009926), hormone transport (GO:0009914), auxin transport (GO:0060918), tropism (GO:0009606), response to gravity (GO:0009629), gravitropism (GO:0009630) and regulation of hormone levels (GO:0010817) (Fig. 1A). Enriched Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways included nitrogen metabolism (KEGG:00910) and ABC transporters (KEGG:02010) (Fig. 1A).

The term auxin and gravitropic response identified by gProfiler were consolidated around five genes including chaperone protein dnaJ 15-like (VIT_01S0011G03790), ABC transporter B family member 1 (VIT_08S0007G05060), protein kinase PINOID-like (VIT_10S0003G04320), auxin efflux carrier component (VIT_11S0052G00440), SEC7 domain-containing protein (VIT_02S0012G01790), and auxin efflux carrier component (VIT_14S0108G00020) (SI Dataset S1).

Table 2. Description of nodes of interest including transcripts exclusively expressed by more than one liana species (L2-L5), most connected nodes, and transcription factors (TFs) found within the co-downregulated portion of the protein-protein interactome network.

Top nodes exclusive to lianas

Liana spp.	Transcript ID	Annotation
L5	VIT_04s0008g05440.t01	ethylene-responsive transcription factor SHN2
L5	VIT_14s0066g01730.t01	xyloglucan glycosyltransferase 4/CSLC4
L4	VIT_03s0038g00680.t01	probable glycosyltransferase GT
L4	VIT_07s0005g00720.t01	pectinesterase/pectinesterase inhibitor 7 PME7
L4	VIT_13s0064g01370.t01	polygalacturonase inhibitor-like PGIP
L4	VIT_00s0252g00120.t01	alpha carbonic anhydrase 4 $\alpha - CAH4$
L4	VIT_07s0141g00330.t01	F-Box/LRR-repeat protein 17 FBL17

Most connected nodes

Number of edges	Transcript ID	Annotation
199	VIT_11s0016g03430.t01	protein phosphatase 2C PP2C
91	VIT_16s0013g00940.t01	kinetochore protein ndc80
86	VIT_07s0005g01430.t01	DNA replication licensing factor MCM7
80	VIT_16s0022g01590.t01	40S ribosomal protein S13
76	VIT_18s0001g03680.t01	serine/threonine-protein kinase mph1
64	VIT_00s1404g00010.t01	abnormal spindle-like microcephaly-associated protein homolog
63	VIT_12s0035g01130.t01	elongation factor 1-gamma
61	VIT_08s0007g03030.t01	ubiquitin-60S ribosomal protein L40-like
59	VIT_10s0003g03080.t01	DNA replication licensing factor MCM3 homolog 2
53	VIT_10s0003g00010.t01	30S ribosomal protein S5, chloroplastic

Downregulated TFs

Gene family	Transcript ID	Annotation
MYB-R2R3	VIT_01s0010g00410.t01	MYB308, MYB-related protein 308
MYB-R2R3	VIT_18s0117g00200.t01	MYB308, MYB-related protein 308
MYB-R2R3	VIT_04s0008g01800.t01	MYB308, MYB-related protein 308
MYB-R2R3	VIT_01s0011g04760.t01	R2R3 MYB transcription factor C2 repressor motif protein MYBC2-L1
MYB-R2R3	VIT_02s0025g00320.t01	MYB86, transcription factor MYB86
MYB-R2R3	VIT_02s0033g00410.t01	MYBA1, transcription factor MybA
MYB-R2R3	VIT_04s0008g00620.t01	MYB37, regulator of axillary meristems RAX1

Table 2 continued from previous page

MYB-R2R3	VIT_04s0008g01830.t01	MYB305, MYB-related protein 305
MYB-R2R3	VIT_06s0004g02110.t01	MYB46, transcription factor MYB46
MYB-R2R3	VIT_06s0004g06280.t01	MYB4, transcription factor MYB4
MYB-R2R3	VIT_08s0056g01190.t01	MYB44, transcription factor MYB44
MYB-R2R3	VIT_09s0002g01380.t01	MYB114, transcription factor MYB114
MYB-R2R3	VIT_09s0002g01400.t01	MYB38, regulator of axial meristems RAX2
MYB-R2R3	VIT_11s0016g02780.t01	MYB36, transcription factor MYB36
MYB-R2R3	VIT_11s0052g01490.t01	MYB36, transcription factor MYB36
MYB-R2R3	VIT_11s0078g00480.t01	MYB4, transcription factor MYB4
MYB-R2R3	VIT_12s0059g00700.t01	GAMYB, transcription factor GAMYB
MYB-R2R3	VIT_13s0067g01360.t01	myb-related protein B
MYB-R2R3	VIT_14s0006g01280.t01	MYB75, transcription factor MYB75
MYB-R2R3	VIT_15s0021g02040.t01	MYB48, transcription factor MYB48
MYB-R2R3	VIT_16s0050g00070.t01	MYB6, transcription repressor MYB6
MYB-R2R3	VIT_17s0000g06410.t01	transcription factor MYB34
MYB-R2R3	VIT_17s0000g08480.t01	MYB-WER, transcription factor MYB-WEREWOLF
MYB-R2R3	VIT_17s0000g08550.t01	MYB98, transcription factor MYB98
MYB-R2R3	VIT_18s0001g05670.t01	MYB39, transcription factor MYB39
MYB-R2R3	VIT_18s0001g15260.t01	MYB-GL1, transcription factor MYB-GL1
MYB-R2R3	VIT_19s0085g00940.t01	MYB26, transcription factor MYB26
MYB	VIT_14s0219g00050.t01	MYB315, myb-related protein 315-like
MYB	VIT_15s0048g02120.t01	transcriptional activator Myb
MYB-TT2	VIT_11s0016g01300.t01	MYB-TT2, MYBPAR, transcription factor MYB
MYB-3R4	VIT_02s0025g04220.t01	MYB transcription factor
MYB-SNAP	VIT_17s0000g02730.t01	MYB-SNAP, snRNA-activating protein complex subunit 4
MADS Box	VIT_03s0088g00590.t01	agamous-like MADS-box protein AGL62
MADS Box	VIT_03s0088g00610.t01	agamous-like MADS-box protein AGL62
MADS Box	VIT_14s0060g00300.t01	agamous-like MADS-box protein AGL80
BLH	VIT_08s0105g00230.t01	BEL1-like homeodomain protein 1
BLH	VIT_13s0019g02450.t01	BEL1-like homeodomain protein 9
HSF	VIT_10s0597g00050.t01	heat stress transcription factor B-2a
HSF	VIT_16s0100g00720.t01	heat stress transcription factor B-2a
NFY	VIT_02s0025g03770.t01	nuclear transcription factor Y subunit C-3
NFY	VIT_05s0020g02790.t01	nuclear transcription factor Y subunit C-3
ILP1	VIT_01s0011g02980.t01	ILP1, negative regulation of transcription by RNA PolII
DEM	VIT_04s0069g01060.t01	[histone H3]-dimethyl-L-lysine9 demethylase
DREB	VIT_08s0007g08150.t01	DREB 2F
TF	VIT_06s0061g00820.t01	RNA polymerase II regulatory region DNA binding
TF	VIT_08s0007g04490.t01	PHD-type domain-containing protein
TF	VIT_11s0016g02020.t01	zinc finger CCCH domain-containing protein 65
TF	VIT_11s0016g03750.t01	SPEAR3
TF	VIT_14s0006g02730.t01	zinc finger protein 6

Transcripts not expressed in lianas. [208]

Comparative analysis revealed 2684 Grapevine orthologs that were not expressed in our [209] liana transcriptome set (SI Dataset S1). Of these, 1687 had hits in the STRING [210] interactome database (Fig. 1). The co-downregulated transcripts (i.e. transcripts that [211] probably have a corresponding gene copy in the liana genomes but were not detected in [212] our set of liana transcripts) enveloped a large portion of the network with 3751 edges. [213] GO term analysis through gProfiler revealed terms pertaining to regulation of gene [214]

expression highly concentrated around transcription factors. Molecular functions were 215
enriched in DNA-binding transcription factor activity (GO:0003700) and 216
sequence-specific DNA binding (GO:0043565). Enriched biological processes included 217
regulation of DNA templated transcription (GO:0006355) and regulation of 218
macromolecule biosynthesis (GO:0010556) which are localized to the nucleus 219
(GO:0005634) (Fig. 1B). Transcription factors such as Nuclear Factor Y (NFY), 220
Agamous-like (AGL), Dehydration Responsive Element Binding (DREB), BEL-like 221
Homeodomain (BLH), and Myeloblastosis (MYB) showed significant presence in the 222
co-downregulated sector of the interactome. The largest group of transcription factors 223
consisted of MYB family represented by 32 members where 27 of them belonged to 224
MYB-R2R3 sub-family (Table 2). The co-downregulated portion of the interactome is 225
comprised of nodes pertaining to mRNA metabolism Fig. 1B). 226

Comparison of three families having liana and non-liana pairs. 227

Three families belonging to Bignoniaceae, Sapindaceae, and Malpighiaceae included 228
pairs of liana and non-liana members in our dataset. We scrutinized these with the 229
anticipation of detecting the most informative gene expression differences (Fig. 2). 230
Enrichment analysis for transcripts expressed only in the liana members resulted in five 231
GO terms and a single KEGG pathway shared between Bignoniaceae and Sapindaceae. 232
These shared terms are involved in developmental process during reproduction 233
(GO:0003006), system development (GO:0048731), intracellular anatomical structure 234
(GO:0005622), chloroplast (GO:0009507), plastid (GO:0009536), and biosynthesis of 235
cofactors (KEGG:01240) (Fig. 2). Terms enriched in Malpighiaceae didn't show any 236
overlap with the other two families (SI Dataset S1). 237

Figure 2. Phylogenetic placement of lianas and non-lianas together with interactome and GO term enrichment of three families bearing liana and non-liana pairs: *Dolichandra unguis-cati* vs *Tabebuia heterophylla* (Bignoniaceae), *Paulinia pinnata* vs *Matayba domingensis* (Sapindaceae), *Heteropterys laurifolia* vs *Byrsonima spicata* (Malpighiaceae). Members of the lianas (yellow colored border with green fill) and non-lianas (salmon colored border with no fill) are highlighted inside interactome networks for each of three within-family comparisons. Co-expressed nodes shared by more than two lianas (L2-L5) are retained for visual guidance as in Fig. 1. A total of six enriched GO and KEGG terms were shared between *Dolichandra unguis-cati* and *Paullinia pinnata*. Terms enriched in *Heteropterys laurifolia* did not show overlap with the other two lianas.

Discussion

Lianas and trees represent two widely separate growth forms with distinct trade off in biomass allocation and growth. Instead of self-supporting stems, lianas allocate most of their resources on shoot and leaf production. Liana leaves must have evolved adaptations to reduce hydraulic failure due to vascular embolism, damage on light harvesting complexes due to intense light and thermal stress, dehydration due to insufficient control of stomatal conductance, inefficient carbon fixation due to mesophyllic carbon diffusion limitations. Here, we provide a wide snapshot of physiologically interpretable differences in gene expression between lianas and non-lianas within a subtropical plant community and make an attempt to learn about genotypic drivers of life-history phenotypes from transcriptional responses of co-habitants. [55–57] (SI Fig. S1, SI Dataset S1). At early developmental stages, transcriptomic differences between lianas and non-lianas are most likely minimal. However, despite the ontogenetic and tissue diversity limitations there exists a suite of differences consistent with expectations from liana biology and functional traits theorized in LES [23–25, 58, 59]. Top leaf level functional traits of lianas correspond to higher SLA and maximal photosynthetic rates. Enzymes we highlight orchestrate cuticular and parenchymal properties related to mesophyll conductance. Liana leaves may do more with less through selective endoreduplication. The resultant action of cell wall building enzymes and control of lignification appears to be in line with the predictions of the LES in lianas. Downregulation of a high number of MYB transcription factors belonging to the largest gene family in plants indicates a crosstalk with cell wall building enzymes since many R2R3-MYBs are known to be involved in lignin deposition. A few R2R3-MYBs are also involved in ABA-mediated control of transpiration which may explain growth advantage experienced by lianas observed in seasonally dry tropical forests [30, 31, 60].

Top Co-expressed transcripts in lianas include enzymes relevant to LES.

Top co-expressed genes in our analysis correspond to enzymes synthesizing or modifying cell wall components, cuticle properties, carbon concentrating mechanism, cell cycle progression through endoreduplication and double-stranded DNA damage response (Table 2, SI Fig. S3). Modifications of cell wall components and cuticle properties are beneficial for lianas in developing leaves with high SLA and optical properties since excessive light levels may damage light harvesting complexes. Reduced production cost due to altered secondary cell wall composition, increased rates of carbon fixation and fast tissue turnover in most liana species may be an outcome of series of mesophyllic adaptations. Facilitation of carbon capture and diffusion through an active carbon concentrating enzyme capable of scavenging low levels of CO_2 floating inside the leaf could be adaptive since leaves with high SLA can accommodate only a few mesophyll cell layers. Lianas could be achieving more with less by increasing gene dosage through selective endoreduplication. In lianas, under hydraulic constraints, machinery evolved

for the repair of damaged nuclear genetic material may be primed as early as the   286
seedling stage.   287

Cuticle properties controlled by SHN2.   288

We found co-expression of SHN2 transcription factor exclusively by five liana species   289
involved in regulation of lipid biosynthesis (Table 2, SI Fig. S3). SHNs together with   290
Wax inducers (WINs) stimulate biosynthesis of cuticular wax in leaf epidermis and   291
reduce stomatal density [61, 62]. SHNs can induce changes in the leaf cuticle to make it   292
more resistant to light stress and avoid excessive heating [63]. Overexpressed SHN2 may   293
indirectly suppress the biosynthesis of lignified secondary wall [64]. This is relevant for   294
lianas because the leaf economical spectrum of lianas predict forms with cheap,   295
disposable but efficient blades with reduced longevity and defense [25]. In our study,   296
many MYBs some of which involved in lignified secondary wall biosynthesis have been   297
observed to be downregulated in lianas (Fig. 1B, Table 2, SI Fig. S3). All 149 genes   298
belonging to AP2/ERF protein family in grapevine genome have been transcriptionally   299
characterized and the Arabidopsis SHN2 ortholog gene   300
(GSVIVP00032652001/VvERF044) also showed an upregulated pattern in the grapevine   301
leaves [65].   302

Cell wall remodeling enzymes.   303

We observed upregulation of a suite of cell wall remodeling enzymes such as CSLC4,   304
PME7, GT, PGIP (Fig. 1A, Table 2, SI Fig. S3) [66]. These cell wall modifying   305
enzymes not only determine leaf morphogenesis but also phyllotaxis [67]. Less lignified   306
leaves with altered cellulose and hemicellulose composition may be one way to achieve   307
set of traits defining liana leaves with fast growth, resource uptake and high productivity   308
end of the leaf economic spectrum. The rigidity of the cell wall is proportional to the   309
number of xyloglucan connections maintained through Golgi-located xyloglucan   310
synthesizing enzymes [68, 69]. Removal of xyloglucan ties loosens cell wall leading to   311
asymmetries in elasticity and resulting in highly convoluted structures (Park and   312
Cosgrove 2015). Overexpression of xyloglucanase in transgenic poplar resulted in droopy   313
leaves with petioles slanted downward [70]. This could be significant because lianas tend   314
to evolve longer petioles when exposed to full light and rigidity of petioles can determine   315
photosynthetic efficiency especially in cordate leaves [71, 72]. Microtubules laid on the   316
cell wall during development can become a reinforcing factor in anisotropic growth since   317
cellulose synthases move along those tubules altering orientations of cellulose   318
deposits [73]. Cellulose is the main load bearing component especially in secondary cell   319
walls [66, 74]. Quintuple mutant analysis of Arabidopsis has revealed that besides   320
CSLC4 four other CSLC genes (CSLC5, CSLC6, CSLC8, and CSLC12) also possess   321
xyloglucan synthase activity (Kim et al. 2020). Another cell wall remodeling enzyme   322
showing upregulation among lianas is the PME7 (Table 2, SI Fig. S3). A composition of   323
esterified and de-esterified pectin residues control calcium-mediated stiffness of gel   324
matrix within the cell wall and contributes to diversification of organ shapes through   325
directional growth. Arabidopsis mutants show that xyloglucans and pectin can together   326
influence shoot meristem and phyllotaxis [67]. Cell wall remodeling could be a   327
morphogenetic response resulting in shade avoidance in lianas [75]. For instance,   328
disrupted phyllotaxy observed in Arabidopsis mutants may have been reinforced in   329
lianas by coordinated expression of cell wall modifying enzymes which are also   330
responsible from gravitropic response such as the positive gravitropism (GO:0009958),   331
tropism (GO:0009606), response to gravity (GO:0009629), gravitropism (GO:0009630)   332
enriched among GO term analysis of upregulated transcripts (Fig. 1A) [67, 68].   333

An active component of carbon concentrating mechanism along the mesophyll may increase fixation efficiency of liana leaves.

Our results also show exclusive activity of alpha-CAH4 in lianas but not in trees and shrubs indicating enhanced carbon concentrating mechanism along the liana mesophyll (Table 2, SI Fig. S3). The carbon concentrating mechanism describes the enzymatic pathways the atmospheric CO_2 flows through the mesophyll cell wall and the plasma membrane into the liquid cytosolic phase imposing limitations on diffusion known as mesophyll conductance. Mesophyll conductance is a significant driving factor in LES [59]. Carbonic anhydrases are zinc metalloenzymes driving the reversible hydration of CO_2 significantly aiding downstream carboxylases such as pyruvate carboxylase, acetyl-coenzyme A carboxylase, PEP carboxylase and RUBISCO [76]. Plants have three classes (alpha-, beta-, gamma-) of carbonic anhydrases [77]. Alpha-carbonic anhydrases are the largest class in the plant kingdom and are highly compartmentalized [78]. Their expression has been detected in cytosol and chloroplast. Carbonic anhydrases can capture CO_2 generated by dark respiration as well as photorespiration enhancing carbon assimilation efficiency of RUBISCO. Stomatal metrics and morphology can determine water use efficiency [79]. Liana leaves with typically lower stomatal densities could be expressing alpha-CAH4 in higher levels increasing the efficiency of the spongy mesophyllic conductance along the CO_2 diffusion pathway (Table 2). An active carbonic anhydrase may contribute to the maintenance of lower stomatal densities. Fossil evidence and present-day observations confirm that plants grown in elevated CO_2 environments decrease their stomatal conductance and stomatal index [80, 81]. Plants over-expressing carbonic anhydrase in their guard cells have the potential to improve their water use efficiency. In guard cells, carbonic anhydrases serve as upstream regulators induced by CO_2 levels independent of leaf photosynthetic rates [82]. High carbonic anhydrase activity could be making lianas more resilient in hotter and drier conditions by reducing the need for evapotranspiration [31, 60].

Liana leaves express FBL17 transcription factor controlling cell cycle progression and double-stranded DNA damage response.

We have observed that four lianas exlusively expressed the plant-specific FBL17 transcript which serves as a major checkpoint in cell cycle progression from Gap1 (G1) into Synthesis (S) phases [83] (Table 2, SI Fig. S3). From the LES perspective, leaf construction is the sum of spatially controlled cytoplasmic growth and cell division. During leaf expansion, a plant can do more with less through a process called endoreduplication where distinct cells may increase their genomic content and gene dosage without cell division [84–86]. During G1 into S phase transition Cyclin Dependent Kinase type A (CDKA) and Cyclin D (CYCD) work together to trigger an expression cascade of DNA replication genes. FBL17 is an indispensable protein component of E3 ubiquitin ligase SCF (ASK-Culling-F-box) for ubiquitination of CDK inhibitor proteins KRPs (KIP-related protein/Inhibitor-Interactor of CDKs) for proteasomal degradation and forms a checkpoint in G1/S transition together with CDKA/CYD complex. Unregulated FBL17 expression causes disruptions in cell cycle genes, while at the same time it can be involved in double stranded DNA damage response leading to abnormalities in root and shoot meristems [87, 88]. In plants, FBL17 is involved in double stranded DNA break induced damage response which is crucial for cell cycle arrest at the first G1/S checkpoint before being committed to division [88, 89]. Loss of function in FBL17 shuts off endoreduplication in Arabidopsis thrichomes [87]. Liana leaves appear to be well protected especially against UV-induced damage with thickened secondary cell wall and waxy cuticle therefore, upregulation of FBL17 might be a deliberate strategy to avoid interruptions in cell cycle. Downregulation of MYB4

and MYB75 involved in biosynthesis of UV-protecting flavonoids and anthocyanins and MYB46 involved in upstream of endoreduplication regulators could be a similarly reconciliatory exertion [90–93] (Table 2, SI Fig. S3). Compatible with FBL17, some of the enriched GO terms for the top 10 most connected nodes include (GO:0022402) cell cycle, (GO:0007049), double-strand break repair via break-induced replication (GO:0000727) (SI Dataset S1).

Figure 3. A conceptual sketch of transcripts and their cellular compartments highlighted in this study. Lianas appear to invest in thickened cuticular layers through upregulation of SHN2. Cell wall remodeling enzymes CSLC4, GT, PGIP, PME7 similarly show upregulation where CSLC4, PME7 and GT are Golgi localized together with pectin, xyloglucan and wax. GT and CSLC4 move into plasma membrane (PM) and deposit xyloglucan, pectin, and cellulose onto the cell wall following microtubule orientation patterns. Upregulated FBL17 is involved in cell cycle control, endoreduplication and double-stranded DNA damage response. Alpha-CAH4 is localized to the cytosol and chloroplast and is a significant player in the CO_2 concentrating mechanism. A large number of R2R3-MYB transcription factors were co-downregulated in lianas playing significant roles as activators and repressors controlling diverse functions such as pectin, lignin and suberin mediated xylogenesis, branching and phyllotaxis and ABA mediated stress tolerance. Cell wall qualities are affected by lignin deposition and pectin crosslinking complexed with Ca^{2+}. PP2C is the most connected node forming the core of the interactome as a key player in ABA induced stress response and interacts with MYBs. Salient liana leaf transcripts collectively may form genetic features underlying the fast end of the LES.

Co-downregulated transcripts are dominated by MYB-R2R3 transcription factors.

Silencing of even a single transcription factor can lead to drastic changes in phenotype with many examples from domesticated plants [94,95]. In our comparisons, transcripts not expressed in lianas are overwhelmingly dominated by R2R3 subclass MYB

transcription factors (Fig. 1, Table 2, SI Fig. S3). Significance of MYB-R2R3 ₄₀₈
co-downregulation for lianas could include stomatal control, lignification and plant ₄₀₉
architecture including phyllotaxis. The MYB-R2R3 serve as major activators and ₄₁₀
repressors controlling diverse functions such as pectin, lignin and suberin mediated ₄₁₁
xylogenesis through control of the phenolic acid metabolism, modulation of ₄₁₂
developmental signaling, cell cycle, epidermal cell fate and patterning [96–99]. There ₄₁₃
may be a crosstalk among the silent MYB-R2R3 transcription factors and exclusively ₄₁₄
upregulated set of cell wall building enzymes together shaping the liana leaf economy. ₄₁₅

MYB genes are found in all eukaryotes and is the largest transcription factor family ₄₁₆
in Arabidopsis [99]. The MYB-R2R3 subfamily in land plants (Embryophyta) has ₄₁₇
expanded during the Silurian widening the structural complexity of tissues through ₄₁₈
secondary cell wall biosynthesis and vascularization [100–102]. Functional predictions of ₄₁₉
most MYBs have been sufficiently characterized both experimentally and ₄₂₀
computationally in myriad model plants. The grapevine and Arabidopsis have 134 and ₄₂₁
126 R2R3-MYB genes, respectively [99, 103]. Structurally, the MYB family harbors ₄₂₂
significant levels of intrinsically disordered regions outside their canonical DNA-binding ₄₂₃
domains which possess potential for other functions regulated by post-transcriptional ₄₂₄
modifications [104]. Here we will highlight a subset of the R2R3-MYBs (MYB44, ₄₂₅
MYB4/MYB308, MYB46, MYB39, MYB38, MYB37) showing no expression in lianas ₄₂₆
(Fig. 1, Table 2, SI Fig. S3). ₄₂₇

MYB44 and ABA-mediated stress tolerance. ₄₂₈

One particular R2R3-MYB downregulated in our analysis is MYB44 which deserves ₄₂₉
special attention since it is heavily involved with the core node of the interactome the ₄₃₀
protein phosphatase PP2C (Table 2, SI Fig. S3). Lianas adopted features such as ₄₃₁
smaller and low-density stomatal openings with sunken guard cells for reduction of ₄₃₂
water loss due to excessive transpiration as early as the Paleozoic based on cuticular ₄₃₃
analysis of pteridosperm lianas [105]. The sesquiterpenoid plant hormone ABA mediates ₄₃₄
crucial physiological processes optimizing water use, carbon uptake and light signaling. ₄₃₅
Although guard cells can endogenously synthesize ABA, the main source for the whole ₄₃₆
plant is vascular parenchyma. In the presence of ABA, ABA receptors ₄₃₇
(PYR/PYL/RCAR) aggregate with clade A PP2Cs reducing the ability of PP2Cs to ₄₃₈
inhibit Sugar Non-fermenting 1 related protein kinase 2 (SnRk2) kinases. Freed from ₄₃₉
PP2C control, SnRK2 kinases phosphorylate and arm transcription factor genes to ₄₄₀
collectively induce or repress ABA-responsive genes downstream elucidating ₄₄₁
ABA-dependent stress response [106, 107]. The behavior of MYB44 appears to have ₄₄₂
opposing modes. In one mode, under stress MYB44 exerts suppressive effect on PP2C. ₄₄₃
The over expressed MYB44 has been shown to have antagonistic interaction with PP2C ₄₄₄
enzymes which control ABA signal transduction cascade in concert with ABA ₄₄₅
receptors [106, 108, 109] (Table 2). Overexpressed MYB44 also appeared to have binding ₄₄₆
interaction with ABA receptors [110]. In the other mode, MYB44 binds to transcription ₄₄₇
start sites of PP2C genes and represses their expression in normal unstressed ₄₄₈
conditions [111]. Moreover, akin to the prokaryotic end-product repression, MYB44 can ₄₄₉
bind to its own promoter in an act of self-silencing [112]. Induction of PP2Cs is ₄₅₀
reported from drought challenged grapevine transcriptomes [113]. In lianas, absence of ₄₅₁
MYB44 activity may be releasing repressed PP2C and could be a desensitization ₄₅₂
mechanism for less aggressive stomatal control. For instance, transgenic gray poplar ₄₅₃
trees expressing mutant ABA insensitive 1 (abi1) belonging to PP2Cs led to large and ₄₅₄
unresponsive stomata with inhibited lateral bud growth [114]. Downregulation of ₄₅₅
MYB37 could be a part of this desensitization scheme [115] (Table 2, SI Fig. S3). ₄₅₆

One noteworthy observation is that the top connected node forming the core of our ₄₅₇
interactome is another PP2C with kinase activity belonging to clade L protein ₄₅₈

phosphatases (VIT_11s0016g03430.t01/AT2G20050) [109](Table 2, SI Fig. S3). The   459
secondary messenger cyclic GMP binds and inhibits the phosphatase activity of this   460
PP2C in favor of kinase activity. For this reason, this PP2C is known as   461
cGMP-dependent protein kinase (PKG) and it phosphorylates the transcription factor   462
GAMYB to upregulate Gibberellic Acid-responsive genes [116]. Leaf cell expansion   463
following cell division also revolves around PP2C through ATPase activity acidifying   464
cell walls [86]. In addition to PP2C, many of the rest of the highly connected nodes are   465
pertaining to chromatin modification (Table 2). Chromatin modelers modify histone   466
tails and contribute to the repressor activity of MYB44 ([111, 112].   467

MYB44 is central in transduction of multiple local abiotic stresses into whole plant   468
response. For instance, SHN2 and MYB86 are responsive to excessive light stress in   469
leaves but when plants experience heat stress simultaneously, incorporation of multiple   470
forcings into a systemic acclimation is carried out by mediators including MYB44 [63].   471

R2R3-MYBs as regulators of secondary cell wall biosynthesis.   472

Lignin biosynthesis is repressed and activated by several R2R3-MYB genes and it is   473
curious whether the downregulated R2R3-MYBs identified in our study show biological   474
associations with the set of upregulated cell wall remodeling and epicuticular enzymes   475
(Fig. 1, Table 2, SI Fig. S3). For instance, overexpressed SHN2 suppresses many MYBs   476
involved in lignified secondary wall biosynthesis as evidenced from Poplar [64].   477
Transcriptional regulation of secondary cell wall related lignin deposition in liana leaves   478
appears to be fine-tuned since they downregulate a set of R2R3-MYBs serving as   479
suppressors and activators of lignin biosynthesis [74] (Table 2). Lignin biosynthesis   480
contains many redundant enzymes and lianas may be regulating multiple check points   481
through silencing of multiple MYBs to achieve their functional leaf morphology   482
encapsulated by LES hypothesis [59, 93].   483

Downregulated R2R3-MYBs as repressors of lignin biosynthesis   484

We have identified MYB46, MYB4/MYB308, and MYB75 reported to be involved in   485
lignin repression downregulated in liana leaves [117] (Table 2, SI Fig. S3). MYB46   486
targets a set of 13 genes in lignin biosynthesis [74, 93, 117–119]. A repressor of   487
endoreduplication called E2Fc is also known to bind to the promoter of MYB46 and   488
suggests a potential crosstalk with the cell cycle regulator FBL-17 [93, 118].   489
Overexpressed MYB4, an ortholog of MYB308, leads to suppression of lignin   490
biosynthesis and reduced elongation of stem internodes, which is a characteristic   491
liana-associated trait [120]. MYB4 is induced by MYB46 but also (similar to MYB44) is   492
inhibited by its own in an end-product repression fashion [74, 117]. MYB75 is a   493
repressor of lignin and secondary wall biosynthetic genes but also an inducer of   494
anthocyanins and flavonoids. Loss of function mutants in Arabidopsis displayed   495
thickened secondary cell walls [121, 122]. MYB75 is a highly phosphorylation dependent   496
protein with elevated internode expression in maize transcriptome and its role in lianas   497
could be parallel to that of MYB4 [120, 122, 123].   498

Downregulated R2R3-MYBs as activators of secondary cell wall   499
synthesis and changers of epidermal properties.   500

Downregulated R2R3-MYBs involved as positive regulators of secondary cell wall   501
synthesis are MYB48, MYB26, MYB39, MYB98, (Table 2, SI Fig. S3). MYB48 appears   502
to play role in xylogenesis but its mechanism of control is unclear containing evidence of   503
alternative splicing through intron-retention [97, 103]. MYB26 positively regulates   504
secondary cell wall biosynthesis in anthers [119]. MYB39 (SUBERMAN) has been   505

shown to be a positive regulator of suberin biosynthesis in Arabidopsis root endodermal tissue [124]. Reduction of suberin in liana leaves maybe a factor in generally low-quality foliage with high turnover as predicted by the LES hypothesis [59, 125].

R2R3-MYB functions linked to epidermal morphology and plant architecture.

MYB98 can be expressed in trichomes and has 83 downstream target genes in synergids [126, 127]. Downregulation of MYB98 maybe a coordinated expression pattern favoring a particular leaf epidermal structure in liana seedlings. This maybe in concert with MYB114 repressor affecting epidermal cell fate and with MYB48 and MYB39 where in Grapevine their low expression leads to anthocyanin accumulation in fruit skin [128] (Table 2, SI Fig. S3). Branching of the shoots into axillary meristems is controlled by MYB37 and MYB38 also known as Regulator of Axillary Meristems (RAX1 and RAX2) which are positively controlled by a WRKY transcription factor Excessive Branches 1 (EXB1) [129]. Transcripts of MYB37 and MBY38 are mobile from cell-to-cell [130]. Their downregulation could be a means to control side branching towards unidirectional growth in lianas especially required during the early ontogenetic developmental stages. Shoot meristems can be influenced by a mutant ribosomal protein gene [131]. In our interactome network some of the highest connected nodes were ribosomal proteins such as 40S ribosomal protein S13, 60S ribosomal protein L40, and chloroplast 30S ribosomal protein S5 (Table 2).

Conclusions

This study explores transcriptomic signatures related to liana leaf properties. Compared to trees, liana leaves generally carry traits colloquially conceptualized as "fast and furious" type life-history strategy associated with quick growth, low cost, high turnover, high capacity for water movement, minimal investment on defense. Our results are in accordance with the LES encapsulating construction costs, rates of carbon fixation and tissue turnover. A set of uniquely expressed enzymes in charge of cell wall building, epicuticular wax synthesis, carbon capture, cell cycle appears to complement with another set of a large number of transcription factors not expressed in lianas. Through comparative transcriptomics of a diverse spectrum of families, we believe we were able to interrogate a wide set of orthologs to contribute towards understanding some of the genetic underpinnings of biology leading to the liana growth form from the leaf perspective. As further sequencing data from additional liana species with more diverse tissue types become available, the genetic basis of this fascinatingly convergent plant form will be more comprehensible.

Supporting Information

Supporting information for Sezen et al. 2021, Comparative Transcriptomics of Tropical Woody Plants Supports Fast and Furious Strategy along the Leaf Economics Spectrum in Lianas. Dryad accession permalink https://doi.org/10.5061/dryad.n5tb2rbw3.

SI Dataset S1

Spreadsheet. A spreadsheet (Dataset-Sezen_etal_Liana.xlsx) containing species list, QUAST, Bowtie2, BUSCO, gProfiler analysis results, enriched genes, domains and categories.

Network file. A Cytoscape generated network used in Figure 1. For reproducible results we recommend selecting Ripple as the visual style and Preset as layout to retain formatting used in the manuscript. Source Cytoscape file (Liana-L5-L2.DRYAD.cys).

Sequences. Four fasta formatted files for each species: (a) TRINITY assembled unfiltered transcripts (SPECIES.Trinity.fasta), (b) EnTAP filtered transcripts where non-plant contaminant sequences have been removed (SPECIES.decon.fasta), (c) EnTAP filtered translated/frame-selected full-length protein sequences (SPECIES-fl-protein.fasta), (d) EnTAP filtered full-length transcript nucleotide sequences (SPECIES-fl-nt.fasta).

SI Figure S1

Caption of Figure S1. A community representation of transcriptome sequenced trees, shrubs and lianas from Luquillo ForestGEO Forest Dynamics Plot, Puerto Rico. Plant heights reflect approximate mature stature for each species. Lianas are shown with horizontal curves. Non-Luquillo climbers are listed on the upper right.

SI Figure S2

Caption of Figure S2. (A) Validation of the de novo assembled transcriptomes via Bowtie2 alignment of trimmed and quality filtered reads. (B) Evaluation for transcriptome completeness using BUSCO analysis. Climbing plant species highlighted in bold. Percentages represent alignment and matching hits.

SI Figure S3

Caption of Figure S3. An annotated view of the highlighted nodes within the co-downregulated (light salmon pink boundaries) and co-expressed transcripts (cyan boundaries) with their first shell interactors using the Grapevine interactome (STRING DB v29290). The interactome spans 1107 nodes and 2994 edges. Labels and colors are the same as those in Fig. 1.

Competing Interests

None declared.

Acknowledgments

We would like to thank Dr. Jill Wegrzyn for being a valuable resource person in bioinformatics and providing access to University of Connecticut's Xanadu High-Performance Computation Cluster. We would like to thank participants of the 2019 Smithsonian Tropical Research Institute-CTFS ForestGEO annual data analysis workshop in Singapore especially to Dr. Marco Visser for his useful insight and discussion that has helped shape the final version of this manuscript. This work was funded by the US National Science Foundation (DEB-1638488; DEB-1643052).

Author Contributions

UUS analyzed the data, prepared the figures and wrote the manuscript with input from all authors. UUS NGS SMM conceived and articulated the idea. SJW MNU NGS collected the samples and carried out the molecular lab work. NGS coordinated the sequencing, data curation, analytical verification and provided bioinformatic support. SJW MNU SJD SMM NGS contributed to the conceptualization, writing, review and proofreading, SJD NGS helped with the project coordination, implementation and funding.

References

1. William E. Stein, Christopher M. Berry, Linda VanAller Hernick, and Frank Mannolini. Surprisingly complex community discovered in the mid-Devonian fossil forest at Gilboa. *Nature*, 483(7387):78–81, March 2012.

2. Joyce G. Chery, Marcelo R. Pace, Pedro Acevedo-Rodríguez, Chelsea D. Specht, and Carl J. Rothfels. Modifications during Early Plant Development Promote the Evolution of Nature's Most Complex Woods. *Current Biology*, 30(2):237–244.e2, January 2020.

3. Alwyn H. Gentry. *The distribution and evolution of climbing plants*, page 3–50. Cambridge University Press, 1992.

4. Liang Hu and Mingguang Li. Diversity and Distribution of Climbing Plants in Eurasia and North Africa. In N. Parthasarathy, editor, *Biodiversity of Lianas*, Sustainable Development and Biodiversity, pages 57–79. Springer International Publishing, Cham, 2015.

5. Stefan A. Schnitzer and Frans Bongers. The ecology of lianas and their role in forests. *Trends in Ecology & Evolution*, 17(5):223–230, May 2002.

6. Francis E. Putz. The Natural History of Lianas on Barro Colorado Island, Panama. *Ecology*, 65(6):1713–1724, 1984.

7. George C. Stevens. Lianas as Structural Parasites: The Bursera Simaruba Example. *Ecology*, 68(1):77–81, 1987.

8. Frank W. Ewers, Julieta A. Rosell, and Mark E. Olson. Lianas as Structural Parasites. In Uwe Hacke, editor, *Functional and Ecological Xylem Anatomy*, pages 163–188. Springer International Publishing, Cham, 2015.

9. Marco D. Visser, Helene C. Muller-Landau, Stefan A. Schnitzer, Hans de Kroon, Eelke Jongejans, and S. Joseph Wright. A host–parasite model explains variation in liana infestation among co-occurring tree species. *Journal of Ecology*, 106(6):2435–2445, 2018.

10. Stefan A. Schnitzer, James W. Dalling, and Walter P. Carson. The impact of lianas on tree regeneration in tropical forest canopy gaps: evidence for an alternative pathway of gap-phase regeneration. *Journal of Ecology*, 88(4):655–666, 2000.

11. Martin G. Barker and Diego Pérez-Salicrup. Comparative water relations of mature mahogany (Swietenia macrophylla) trees with and without lianas in a subhumid, seasonally dry forest in Bolivia. *Tree Physiology*, 20(17):1167–1174, November 2000.

12. Stefan A. Schnitzer. A Mechanistic Explanation for Global Patterns of Liana Abundance and Distribution. *The American Naturalist*, 166(2):262–276, August 2005.

13. S. Joseph Wright, M. Alejandra Jaramillo, Javier Pavon, Richard Condit, Stephen P. Hubbell, and Robin B. Foster. Reproductive size thresholds in tropical trees: variation among individuals, species and forests. *Journal of Tropical Ecology*, 21(3):307–315, May 2005.

14. Stefan A. Schnitzer and Walter P. Carson. Lianas suppress tree regeneration and diversity in treefall gaps. *Ecology Letters*, 13(7):849–857, 2010.

15. José Luis Andrade, Frederick C. Meinzer, Guillermo Goldstein, and Stefan A. Schnitzer. Water uptake and transport in lianas and co-occurring trees of a seasonally dry tropical forest. *Trees*, 19(3):282–289, May 2005.

16. Oliver L. Phillips, Rodolfo Vásquez Martínez, Luzmila Arroyo, Timothy R. Baker, Timothy Killeen, Simon L. Lewis, Yadvinder Malhi, Abel Monteagudo Mendoza, David Neill, Percy Núñez Vargas, Miguel Alexiades, Carlos Cerón, Anthony Di Fiore, Terry Erwin, Anthony Jardim, Walter Palacios, Mario Saldias, and Barbara Vinceti. Increasing dominance of large lianas in Amazonian forests. *Nature*, 418(6899):770, August 2002.

17. Stefan A. Schnitzer, Scott A. Mangan, James W. Dalling, Claire A. Baldeck, Stephen P. Hubbell, Alicia Ledo, Helene Muller-Landau, Michael F. Tobin, Salomon Aguilar, David Brassfield, Andres Hernandez, Suzanne Lao, Rolando Perez, Oldemar Valdes, and Suzanne Rutishauser Yorke. Liana Abundance, Diversity, and Distribution on Barro Colorado Island, Panama. *PLOS ONE*, 7(12):e52114, December 2012.

18. Charles Darwin. On the Movements and Habits of Climbing Plants. *Journal of the Linnean Society of London, Botany*, 9(33-34):1–118, 1865.

19. Francis E. Putz and Harold A. Mooney, editors. *The Biology of vines*. Cambridge University Press, Cambridge ; New York, 1991.

20. Lan Zhang, Yajun Chen, Keping Ma, Frans Bongers, and Frank J. Sterck. Fully exposed canopy tree and liana branches in a tropical forest differ in mechanical traits but are similar in hydraulic traits. *Tree Physiology*, 39(10):1713–1724, October 2019. Publisher: Oxford Academic.

21. L. S. Santiago and S. J. Wright. Leaf functional traits of tropical forest plants in relation to growth form. *Functional Ecology*, 21(1):19–27, 2007. _eprint: https://besjournals.onlinelibrary.wiley.com/doi/pdf/10.1111/j.1365-2435.2006.01218.x.

22. Ryuji Ichihashi and Masaki Tateno. Biomass allocation and long-term growth patterns of temperate lianas in comparison with trees. *New Phytologist*, 207(3):604–612, 2015. _eprint: https://nph.onlinelibrary.wiley.com/doi/pdf/10.1111/nph.13391.

23. Ian J. Wright, Peter B. Reich, Mark Westoby, David D. Ackerly, Zdravko Baruch, Frans Bongers, Jeannine Cavender-Bares, Terry Chapin, Johannes H. C. Cornelissen, Matthias Diemer, Jaume Flexas, Eric Garnier, Philip K. Groom, Javier Gulias, Kouki Hikosaka, Byron B. Lamont, Tali Lee, William Lee, Christopher Lusk, Jeremy J. Midgley, Marie-Laure Navas, Ülo Niinemets, Jacek

Oleksyn, Noriyuki Osada, Hendrik Poorter, Pieter Poot, Lynda Prior, Vladimir I. Pyankov, Catherine Roumet, Sean C. Thomas, Mark G. Tjoelker, Erik J. Veneklaas, and Rafael Villar. The worldwide leaf economics spectrum. *Nature*, 428(6985):821–827, April 2004. Number: 6985 Publisher: Nature Publishing Group.

24. Jeanne L. D. Osnas, Jeremy W. Lichstein, Peter B. Reich, and Stephen W. Pacala. Global Leaf Trait Relationships: Mass, Area, and the Leaf Economics Spectrum. *Science*, 340(6133):741–744, May 2013. Publisher: American Association for the Advancement of Science Section: Report.

25. Felipe N. A. Mello, Sergio Estrada-Villegas, David M. DeFilippis, and Stefan A. Schnitzer. Can Functional Traits Explain Plant Coexistence? A Case Study with Tropical Lianas and Trees. *Diversity*, 12(10):397, October 2020. Number: 10 Publisher: Multidisciplinary Digital Publishing Institute.

26. S. B. Zhang, Q. L. Deng, and Y. J. Hao. Effects of high temperature and high light on photosystem II activity in leaves of two Bauhinia species. *Photosynthetica*, 57(4):1094–1099, November 2019. Publisher: Photosynthetica.

27. Gerardo Avalos, Stephen S. Mulkey, and Kaoru Kitajima. Leaf Optical Properties of Trees and Lianas in the Outer Canopy of a Tropical Dry Forest. *Biotropica*, 31(3):517–520, 1999. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1744-7429.1999.tb00395.x.

28. J. Antonio Guzmán Q., G. Arturo Sánchez-Azofeifa, and Benoit Rivard. Differences in Leaf Temperature between Lianas and Trees in the Neotropical Canopy. *Forests*, 9(6):307, June 2018. Number: 6 Publisher: Multidisciplinary Digital Publishing Institute.

29. Zhi-Quan Cai, Stefan A. Schnitzer, and Frans Bongers. Seasonal differences in leaf-level physiology give lianas a competitive advantage over trees in a tropical seasonal forest. *Oecologia*, 161(1):25–33, August 2009.

30. Stefan A. Schnitzer. Testing ecological theory with lianas. *New Phytologist*, 220(2):366–380, 2018. _eprint: https://nph.onlinelibrary.wiley.com/doi/pdf/10.1111/nph.15431.

31. José A. Medina-Vega, Frans Bongers, Lourens Poorter, Stefan A. Schnitzer, and Frank J. Sterck. Lianas have more acquisitive traits than trees in a dry but not in a wet forest. *Journal of Ecology*, 109(6):2367–2384, 2021. _eprint: https://besjournals.onlinelibrary.wiley.com/doi/pdf/10.1111/1365-2745.13644.

32. Helena Augusto Gioppato, Mariana Bombardi da Silva, Scott Carrara, Bruna Rafaella Zanardi Palermo, Tatiana de Souza Moraes, and Marcelo Carnier Dornelas. Genomic and transcriptomic approaches to understand Passiflora physiology and to contribute to passionfruit breeding. *Theoretical and Experimental Plant Physiology*, 31(1):173–181, March 2019.

33. Tin Hang Hung, Thea So, Syneath Sreng, Bansa Thammavong, Chaloun Boounithiphonh, David H. Boshier, and John J. MacKay. Reference transcriptomes and comparative analyses of six species in the threatened rosewood genus Dalbergia. *Scientific Reports*, 10(1):17749, October 2020. Number: 1 Publisher: Nature Publishing Group.

34. The French-Italian Public Consortium for Grapevine Genome Characterization, Olivier Jaillon, Jean-Marc Aury, Benjamin Noel, Alberto Policriti, Christian Clepet, Alberto Casagrande, Nathalie Choisne, Sébastien Aubourg, Nicola Vitulo, Claire Jubin, Alessandro Vezzi, Fabrice Legeai, Philippe Hugueney, Corinne Dasilva, David Horner, Erica Mica, Delphine Jublot, Julie Poulain, Clémence Bruyère, Alain Billault, Béatrice Segurens, Michel Gouyvenoux, Edgardo Ugarte, Federica Cattonaro, Véronique Anthouard, Virginie Vico, Cristian Del Fabbro, Michaël Alaux, Gabriele Di Gaspero, Vincent Dumas, Nicoletta Felice, Sophie Paillard, Irena Juman, Marco Moroldo, Simone Scalabrin, Aurélie Canaguier, Isabelle Le Clainche, Giorgio Malacrida, Eléonore Durand, Graziano Pesole, Valérie Laucou, Philippe Chatelet, Didier Merdinoglu, Massimo Delledonne, Mario Pezzotti, Alain Lecharny, Claude Scarpelli, François Artiguenave, M. Enrico Pè, Giorgio Valle, Michele Morgante, Michel Caboche, Anne-Françoise Adam-Blondon, Jean Weissenbach, Francis Quétier, and Patrick Wincker. The grapevine genome sequence suggests ancestral hexaploidization in major angiosperm phyla. *Nature*, 449(7161):463–467, September 2007.

35. Jun Wen, Zhiqiang Xiong, Ze-Long Nie, Likai Mao, Yabing Zhu, Xian-Zhao Kan, Stefanie M. Ickert-Bond, Jean Gerrath, Elizabeth A. Zimmer, and Xiao-Dong Fang. Transcriptome Sequences Resolve Deep Relationships of the Grape Family. *PLOS ONE*, 8(9):e74394, September 2013.

36. Daniel H. Chitwood, Aashish Ranjan, Ciera C. Martinez, Lauren R. Headland, Thinh Thiem, Ravi Kumar, Michael F. Covington, Tommy Hatcher, Daniel T. Naylor, Sharon Zimmerman, Nora Downs, Nataly Raymundo, Edward S. Buckler, Julin N. Maloof, Mallikarjuna Aradhya, Bernard Prins, Lin Li, Sean Myles, and Neelima R. Sinha. A Modern Ampelography: A Genetic Basis for Leaf Shape and Venation Patterning in Grape. *Plant Physiology*, 164(1):259–272, January 2014.

37. Silvina Dayer, Idan Reingwirtz, Andrew J. McElrone, and Gregory A. Gambetta. Response and Recovery of Grapevine to Water Deficit: From Genes to Physiology. In Dario Cantu and M. Andrew Walker, editors, *The Grape Genome*, Compendium of Plant Genomes, pages 223–245. Springer International Publishing, Cham, 2019.

38. Samia Daldoul, Hatem Boubakri, Mahmoud Gargouri, and Ahmed Mliki. Recent advances in biotechnological studies on wild grapevines as valuable resistance sources for smart viticulture. *Molecular Biology Reports*, 47(4):3141–3153, April 2020.

39. Naim Matasci, Ling-Hong Hung, Zhixiang Yan, Eric J Carpenter, Norman J Wickett, Siavash Mirarab, Nam Nguyen, Tandy Warnow, Saravanaraj Ayyampalayam, Michael Barker, J Gordon Burleigh, Matthew A Gitzendanner, Eric Wafula, Joshua P Der, Claude W dePamphilis, Béatrice Roure, Hervé Philippe, Brad R Ruhfel, Nicholas W Miles, Sean W Graham, Sarah Mathews, Barbara Surek, Michael Melkonian, Douglas E Soltis, Pamela S Soltis, Carl Rothfels, Lisa Pokorny, Jonathan A Shaw, Lisa DeGironimo, Dennis W Stevenson, Juan Carlos Villarreal, Tao Chen, Toni M Kutchan, Megan Rolf, Regina S Baucom, Michael K Deyholos, Ram Samudrala, Zhijian Tian, Xiaolei Wu, Xiao Sun, Yong Zhang, Jun Wang, Jim Leebens-Mack, and Gane Ka-Shu Wong. Data access for the 1,000 Plants (1KP) project. *GigaScience*, 3:17, October 2014.

40. One thousand plant transcriptomes and the phylogenomics of green plants. *Nature*, 574(7780):679–685, October 2019.

41. Manfred G. Grabherr, Brian J. Haas, Moran Yassour, Joshua Z. Levin, Dawn A. Thompson, Ido Amit, Xian Adiconis, Lin Fan, Raktima Raychowdhury, Qiandong Zeng, Zehua Chen, Evan Mauceli, Nir Hacohen, Andreas Gnirke, Nicholas Rhind, Federica di Palma, Bruce W. Birren, Chad Nusbaum, Kerstin Lindblad-Toh, Nir Friedman, and Aviv Regev. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature Biotechnology*, 29(7):644–652, July 2011.

42. Alexey Gurevich, Vladislav Saveliev, Nikolay Vyahhi, and Glenn Tesler. QUAST: quality assessment tool for genome assemblies. *Bioinformatics*, page btt086, February 2013.

43. Ben Langmead and Steven L. Salzberg. Fast gapped-read alignment with Bowtie 2. *Nature Methods*, 9(4):357–359, April 2012.

44. Felipe A. Simão, Robert M. Waterhouse, Panagiotis Ioannidis, Evgenia V. Kriventseva, and Evgeny M. Zdobnov. BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics*, 31(19):3210–3212, October 2015.

45. Alexander J. Hart, Samuel Ginzburg, Muyang (Sam) Xu, Cera R. Fisher, Nasim Rahmatpour, Jeffry B. Mitton, Robin Paul, and Jill L. Wegrzyn. EnTAP: Bringing Faster and Smarter Functional Annotation to Non-Model Eukaryotic Transcriptomes. *Molecular Ecology Resources*, 0(ja), October 2019.

46. Benjamin Buchfink, Chao Xie, and Daniel H. Huson. Fast and sensitive protein alignment using DIAMOND. *Nature Methods*, 12(1):59–60, January 2015.

47. Shiyuyun Tang, Alexandre Lomsadze, and Mark Borodovsky. Identification of protein coding regions in RNA transcripts. *Nucleic Acids Research*, 43(12):e78–e78, July 2015.

48. Torbjørn Rognes, Tomáš Flouri, Ben Nichols, Christopher Quince, and Frédéric Mahé. VSEARCH: a versatile open source tool for metagenomics. *PeerJ*, 4, October 2016.

49. Christian von Mering, Lars J. Jensen, Berend Snel, Sean D. Hooper, Markus Krupp, Mathilde Foglierini, Nelly Jouffre, Martijn A. Huynen, and Peer Bork. STRING: known and predicted protein–protein associations, integrated and transferred across organisms. *Nucleic Acids Research*, 33(suppl_1):D433–D437, January 2005.

50. Andrea Franceschini, Damian Szklarczyk, Sune Frankild, Michael Kuhn, Milan Simonovic, Alexander Roth, Jianyi Lin, Pablo Minguez, Peer Bork, Christian von Mering, and Lars J. Jensen. STRING v9.1: protein-protein interaction networks, with increased coverage and integration. *Nucleic Acids Research*, 41(D1):D808–D815, January 2013.

51. Paul Shannon, Andrew Markiel, Owen Ozier, Nitin S. Baliga, Jonathan T. Wang, Daniel Ramage, Nada Amin, Benno Schwikowski, and Trey Ideker. Cytoscape: A Software Environment for Integrated Models of Biomolecular Interaction Networks. *Genome Research*, 13(11):2498–2504, November 2003.

52. Sudhir Kumar, Glen Stecher, Michael Suleski, and S. Blair Hedges. TimeTree: A Resource for Timelines, Timetrees, and Divergence Times. *Molecular Biology and Evolution*, 34(7):1812–1819, July 2017.

53. Uku Raudvere, Liis Kolberg, Ivan Kuzmin, Tambet Arak, Priit Adler, Hedi Peterson, and Jaak Vilo. g:Profiler: a web server for functional enrichment analysis and conversions of gene lists (2019 update). *Nucleic Acids Research*, 47(W1):W191–W198, July 2019.

54. Paul G. Nevill, Katharine A. Howell, Adam T. Cross, Anna V. Williams, Xiao Zhong, Julian Tonti-Filippini, Laura M. Boykin, Kingsley W. Dixon, and Ian Small. Plastome-Wide Rearrangements and Gene Losses in Carnivorous Droseraceae. *Genome Biology and Evolution*, 11(2):472–485, February 2019.

55. Jenny Zambrano, Yoshiko Iida, Robert Howe, Luxiang Lin, Maria Natalia Umana, Amy Wolf, Samantha J. Worthy, and Nathan G. Swenson. Neighbourhood defence gene similarity effects on tree performance: a community transcriptomic approach. *Journal of Ecology*, 105(3):616–626, 2017. _eprint: https://besjournals.onlinelibrary.wiley.com/doi/pdf/10.1111/1365-2745.12765.

56. Nathan G. Swenson and F. Andrew Jones. Community transcriptomics, genomics and the problem of species co-occurrence. *Journal of Ecology*, 105(3):563–568, 2017. _eprint: https://besjournals.onlinelibrary.wiley.com/doi/pdf/10.1111/1365-2745.12771.

57. Nathan G. Swenson, Yoshiko Iida, Robert Howe, Amy Wolf, María Natalia Umaña, Krittika Petprakob, Benjamin L. Turner, and Keping Ma. Tree co-occurrence and transcriptomic response to drought. *Nature Communications*, 8(1):1–9, December 2017. Number: 1 Publisher: Nature Publishing Group.

58. Tomasz P. Wyka, Jacek Oleksyn, Piotr Karolewski, and Stefan A. Schnitzer. Phenotypic correlates of the lianescent growth form: a review. *Annals of Botany*, 112(9):1667–1681, December 2013.

59. Yusuke Onoda, Ian J. Wright, John R. Evans, Kouki Hikosaka, Kaoru Kitajima, Ülo Niinemets, Hendrik Poorter, Tiina Tosens, and Mark Westoby. Physiological and structural tradeoffs underlying the leaf economics spectrum. *New Phytologist*, 214(4):1447–1463, 2017. _eprint: https://nph.onlinelibrary.wiley.com/doi/pdf/10.1111/nph.14496.

60. María Natalia Umaña, Jimena Forero-Montaña, Christopher J. Nytch, Jill Thompson, María Uriarte, Jess Zimmerman, and Nathan G. Swenson. Dry conditions and disturbance promote liana seedling survival and abundance. *Ecology*, 100(1):e02556, January 2019.

61. Asaph Aharoni, Shital Dixit, Reinhard Jetter, Eveline Thoenes, Gert van Arkel, and Andy Pereira. The SHINE Clade of AP2 Domain Transcription Factors Activates Wax Biosynthesis, Alters Cuticle Properties, and Confers Drought Tolerance when Overexpressed in Arabidopsis. *The Plant Cell*, 16(9):2463–2480, September 2004. Publisher: American Society of Plant Biologists Section: Research Article.

62. Pierre Broun, Patricia Poindexter, Erin Osborne, Cai-Zhong Jiang, and José Luis Riechmann. WIN1, a transcriptional activator of epidermal wax accumulation in Arabidopsis. *Proceedings of the National Academy of Sciences*, 101(13):4706–4711, March 2004. Publisher: National Academy of Sciences Section: Biological Sciences.

63. Sara I. Zandalinas, Yosef Fichman, Amith R. Devireddy, Soham Sengupta, Rajeev K. Azad, and Ron Mittler. Systemic signaling during abiotic stress combination in plants. *Proceedings of the National Academy of Sciences*, 117(24):13810–13820, June 2020. Publisher: National Academy of Sciences Section: Biological Sciences.

64. Yingying Liu, Minjing Wei, Cong Hou, Tingting Lu, Lulu Liu, Hairong Wei, Yuxiang Cheng, and Zhigang Wei. Functional Characterization of Populus PsnSHN2 in Coordinated Regulation of Secondary Wall Components in Tobacco. *Scientific Reports*, 7(1):42, March 2017. Number: 1 Publisher: Nature Publishing Group.

65. Francesco Licausi, Federico M. Giorgi, Sara Zenoni, Fabio Osti, Mario Pezzotti, and Pierdomenico Perata. Genomic and transcriptomic analysis of the AP2/ERF superfamily in Vitis vinifera. *BMC Genomics*, 11(1):719, December 2010.

66. Edwin R. Lampugnani, Ghazanfar Abbas Khan, Marc Somssich, and Staffan Persson. Building a plant cell wall at a glance. *Journal of Cell Science*, 131(2), January 2018. Publisher: The Company of Biologists Ltd Section: CELL SCIENCE AT A GLANCE.

67. Feng Zhao, Wenqian Chen, Julien Sechet, Marjolaine Martin, Simone Bovio, Claire Lionnet, Yuchen Long, Virginie Battu, Grégory Mouille, Françoise Monéger, and Jan Traas. Xyloglucans and Microtubules Synergistically Maintain Meristem Geometry and Phyllotaxis. *Plant Physiology*, 181(3):1191–1206, November 2019. Publisher: American Society of Plant Biologists Section: Research Article.

68. Takahisa Hayashi and Rumi Kaida. Functions of Xyloglucan in Plant Cells. *Molecular Plant*, 4(1):17–24, January 2011.

69. Yi-Hsiang Chou, Gennady Pogorelko, Zachary T. Young, and Olga A. Zabotina. Protein–Protein Interactions Among Xyloglucan-Synthesizing Enzymes and Formation of Golgi-Localized Multiprotein Complexes. *Plant and Cell Physiology*, 56(2):255–267, February 2015.

70. Kei'ichi Baba, Yong Woo Park, Tomomi Kaku, Rumi Kaida, Miyuki Takeuchi, Masato Yoshida, Yoshihiro Hosoo, Yasuhisa Ojio, Takashi Okuyama, Toru Taniguchi, Yasunori Ohmiya, Teiji Kondo, Ziv Shani, Oded Shoseyov, Tatsuya Awano, Satoshi Serada, Naoko Norioka, Shigemi Norioka, and Takahisa Hayashi. Xyloglucan for Generating Tensile Stress to Bend Tree Stem. *Molecular Plant*, 2(5):893–903, September 2009.

71. Thomas J. Givnish and Geerat J. Vermeij. Sizes and Shapes of Liane Leaves. *The American Naturalist*, 110(975):743–778, 1976. Publisher: [University of Chicago Press, American Society of Naturalists].

72. Akio Takenaka. Effects of leaf blade narrowness and petiole length on the light capture efficiency of a shoot. *Ecological Research*, 9(2):109–114, 1994. _eprint: https://esj-journals.onlinelibrary.wiley.com/doi/pdf/10.1007/BF02347485.

73. Martin Bringmann, Benoit Landrein, Christian Schudoma, Olivier Hamant, Marie-Theres Hauser, and Staffan Persson. Cracking the elusive alignment hypothesis: the microtubule–cellulose synthase nexus unraveled. *Trends in Plant Science*, 17(11):666–674, November 2012.

74. Ruiqin Zhong and Zheng-Hua Ye. Secondary Cell Walls: Biosynthesis, Patterned Deposition and Transcriptional Regulation. *Plant and Cell Physiology*, 56(2):195–214, February 2015.

75. Sang-Jin Kim, Balakumaran Chandrasekar, Anne C. Rea, Linda Danhof, Starla Zemelis-Durfee, Nicholas Thrower, Zachary S. Shepard, Markus Pauly, Federica Brandizzi, and Kenneth Keegstra. The synthesis of xyloglucan, an abundant plant cell wall polysaccharide, requires CSLC function. *Proceedings of the National Academy of Sciences*, 117(33):20316–20324, August 2020. Publisher: National Academy of Sciences Section: Biological Sciences.

76. Claudiu T. Supuran. Carbonic Anhydrases and Metabolism. *Metabolites*, 8(2), March 2018.

77. Robert J. DiMario, Harmony Clayton, Ananya Mukherjee, Martha Ludwig, and James V. Moroney. Plant Carbonic Anhydrases: Structures, Locations, Evolution, and Physiological Roles. *Molecular Plant*, 10(1):30–46, January 2017.

78. Mina Momayyezi, Athena D. McKown, Shannon C. S. Bell, and Robert D. Guy. Emerging roles for carbonic anhydrase in mesophyll conductance and photosynthesis. *The Plant Journal*, 101(4):831–844, 2020. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/tpj.14638.

79. Lígia T. Bertolino, Robert S. Caine, and Julie E. Gray. Impact of Stomatal Density and Morphology on Water-Use Efficiency in a Changing World. *Frontiers in Plant Science*, 10, 2019. Publisher: Frontiers.

80. Gregory J. Retallack. A 300-million-year record of atmospheric carbon dioxide from fossil plant cuticles. *Nature*, 411(6835):287–290, May 2001. Number: 6835 Publisher: Nature Publishing Group.

81. Emmy I. Lammertsma, Hugo Jan de Boer, Stefan C. Dekker, David L. Dilcher, André F. Lotter, and Friederike Wagner-Cremer. Global CO2 rise leads to reduced maximum stomatal conductance in Florida vegetation. *Proceedings of the National Academy of Sciences*, 108(10):4035–4040, March 2011. Publisher: National Academy of Sciences Section: Biological Sciences.

82. Honghong Hu, Aurélien Boisson-Dernier, Maria Israelsson-Nordström, Maik Böhmer, Shaowu Xue, Amber Ries, Jan Godoski, Josef M. Kuhn, and Julian I. Schroeder. Carbonic Anhydrases are Upstream Regulators in Guard Cells of CO2-Controlled Stomatal Movements. *Nature cell biology*, 12(1):87–18, January 2010.

83. Hirofumi Harashima, Nico Dissmeyer, and Arp Schnittger. Cell cycle control across the eukaryotic kingdom. *Trends in Cell Biology*, 23(7):345–356, July 2013. Publisher: Elsevier.

84. J. E. Melaragno, B. Mehrotra, and A. W. Coleman. Relationship between Endopolyploidy and Cell Size in Epidermal Tissue of Arabidopsis. *The Plant Cell*, 5(11):1661–1668, November 1993. Publisher: American Society of Plant Biologists.

85. Shweta Kalve, Dirk De Vos, and Gerrit T. S. Beemster. Leaf development: a cellular perspective. *Frontiers in Plant Science*, 5, 2014. Publisher: Frontiers.

86. Jasmien Vercruysse, Alexandra Baekelandt, Nathalie Gonzalez, and Dirk Inzé. Molecular networks regulating cell division during Arabidopsis leaf growth. *Journal of Experimental Botany*, 71(8):2365–2378, April 2020.

87. Sandra Noir, Katia Marrocco, Kinda Masoud, Alexis Thomann, Andi Gusti, Marta Bitrian, Arp Schnittger, and Pascal Genschik. The Control of Arabidopsis thaliana Growth by Cell Proliferation and Endoreplication Requires the F-Box Protein FBL17. *The Plant Cell*, 27(5):1461–1476, May 2015. Publisher: American Society of Plant Biologists Section: Research Article.

88. Naomie Gentric, Kinda Masoud, Robin P. Journot, Valérie Cognat, Marie-Edith Chabouté, Sandra Noir, and Pascal Genschik. The F-Box-Like Protein FBL17 Is a Regulator of DNA-Damage Response and Colocalizes with RETINOBLASTOMA RELATED1 at DNA Lesion Sites. *Plant Physiology*, 183(3):1295–1305, July 2020. Publisher: American Society of Plant Biologists Section: Research Article.

89. Xin'Ai Zhao, Hirofumi Harashima, Nico Dissmeyer, Stefan Pusch, Annika K. Weimer, Jonathan Bramsiepe, Daniel Bouyer, Svenja Rademacher, Moritz K. Nowack, Bela Novak, Stefanie Sprunck, and Arp Schnittger. A General G1/S-Phase Cell-Cycle Control Module in the Flowering Plant Arabidopsis thaliana. *PLoS Genetics*, 8(8), August 2012.

90. Matthew R Hemm, Klaus M Herrmann, and Clint Chapple. AtMYB4: a transcription factor general in the battle against UV. *Trends in Plant Science*, 6(4):135–136, April 2001.

91. Takayuki Tohge, Miyako Kusano, Atsushi Fukushima, Kazuki Saito, and Alisdair R. Fernie. Transcriptional and metabolic programs following exposure of plants to UV-B irradiation. *Plant Signaling & Behavior*, 6(12):1987–1992, December 2011.

92. Takayuki Tohge, Yasutaka Nishiyama, Masami Yokota Hirai, Mitsuru Yano, Jun-ichiro Nakajima, Motoko Awazuhara, Eri Inoue, Hideki Takahashi, Dayan B. Goodenowe, Masahiko Kitayama, Masaaki Noji, Mami Yamazaki, and Kazuki Saito. Functional genomics by integrated analysis of metabolome and transcriptome of Arabidopsis plants over-expressing an MYB transcription factor. *The Plant Journal*, 42(2):218–235, 2005. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1365-313X.2005.02371.x.

93. M. Taylor-Teeples, L. Lin, M. de Lucas, G. Turco, T. W. Toal, A. Gaudinier, N. F. Young, G. M. Trabucco, M. T. Veling, R. Lamothe, P. P. Handakumbura, G. Xiong, C. Wang, J. Corwin, A. Tsoukalas, L. Zhang, D. Ware, M. Pauly, D. J. Kliebenstein, K. Dehesh, I. Tagkopoulos, G. Breton, J. L. Pruneda-Paz, S. E. Ahnert, S. A. Kay, S. P. Hazen, and S. M. Brady. An Arabidopsis gene regulatory network for secondary cell wall synthesis. *Nature*, 517(7536):571–575, January 2015. Number: 7536 Publisher: Nature Publishing Group.

94. Haibao Tang, Uzay Sezen, and Andrew H Paterson. Domestication and plant genomes. *Current Opinion in Plant Biology*, 13(2):160–166, April 2010.

95. Haibao Tang, Hugo E. Cuevas, Sayan Das, Uzay U. Sezen, Chengbo Zhou, Hui Guo, Valorie H. Goff, Zhengxiang Ge, Thomas E. Clemente, and Andrew H. Paterson. Seed shattering in a wild sorghum is conferred by a locus unrelated to domestication. *Proceedings of the National Academy of Sciences*, 110(39):15824–15829, September 2013. Publisher: National Academy of Sciences Section: Biological Sciences.

96. Erika Cavallini, José Tomás Matus, Laura Finezzo, Sara Zenoni, Rodrigo Loyola, Flavia Guzzo, Rudolf Schlechter, Agnès Ageorges, Patricio Arce-Johnson, and

Giovanni Battista Tornielli. The Phenylpropanoid Pathway Is Controlled at Different Branches by a Set of R2R3-MYB C2 Repressors in Grapevine. *Plant Physiology*, 167(4):1448–1470, April 2015. Publisher: American Society of Plant Biologists Section: Articles.

97. Sookyung Oh, Sunchung Park, and Kyung-Hwan Han. Transcriptional regulation of secondary growth in Arabidopsis thaliana. *Journal of Experimental Botany*, 54(393):2709–2722, December 2003.

98. José Tomás Matus, Felipe Aquea, and Patricio Arce-Johnson. Analysis of the grape MYB R2R3 subfamily reveals expanded wine quality-related clades and conserved gene structure organization across Vitis and Arabidopsis genomes. *BMC Plant Biology*, 8(1):83, July 2008.

99. Christian Dubos, Ralf Stracke, Erich Grotewold, Bernd Weisshaar, Cathie Martin, and Loïc Lepiniec. MYB transcription factors in Arabidopsis. *Trends in Plant Science*, 15(10):573–581, October 2010.

100. Peter H. Raven, University Ray F. Evert, Ray F. Evert, Susan E. Eichhorn, and University Susan E. Eichhorn. *Biology of Plants*. Macmillan, 2005.

101. Chen-Kun Jiang and Guang-Yuan Rao. Insights into the Diversification and Evolution of R2R3-MYB Transcription Factors in Plants. *Plant Physiology*, 183(2):637–655, June 2020. Publisher: American Society of Plant Biologists Section: Research Article.

102. Xiaojun Chang, Shupeng Xie, Lanlan Wei, Zhaolian Lu, Zhong-Hua Chen, Fei Chen, Zhongxiong Lai, Zhenguo Lin, and Liangsheng Zhang. Origins and Stepwise Expansion of R2R3-MYB Transcription Factors for the Terrestrial Adaptation of Plants. *Frontiers in Plant Science*, 11, 2020. Publisher: Frontiers.

103. Darren Chern Jan Wong, Rudolf Schlechter, Alessandro Vannozzi, Janine Höll, Ibrahim Hmmam, Jochen Bogs, Giovanni Battista Tornielli, Simone Diego Castellarin, and José Tomás Matus. A systems-oriented analysis of the grapevine R2R3-MYB transcription factor family uncovers new insights into the regulation of stilbene accumulation. *DNA Research*, 23(5):451–466, October 2016.

104. Peter S. Millard, Birthe B. Kragelund, and Meike Burow. R2R3 MYB Transcription Factors – Functions outside the DNA-Binding Domain. *Trends in Plant Science*, 24(10):934–946, October 2019.

105. Michael Krings, Hans Kerp, Thomas N. Taylor, and Edith L. Taylor. How Paleozoic vines and lianas got off the ground: On scrambling and climbing carboniferous-early permian pteridosperms. *The Botanical Review*, 69(2):204–224, April 2003.

106. Choonkyun Jung, Nguyen Hoai Nguyen, and Jong-Joo Cheong. Transcriptional Regulation of Protein Phosphatase 2C Genes to Modulate Abscisic Acid Signaling. *International Journal of Molecular Sciences*, 21(24):9517, January 2020. Number: 24 Publisher: Multidisciplinary Digital Publishing Institute.

107. Fen-Fen Soon, Ley-Moy Ng, X. Edward Zhou, Graham M. West, Amanda Kovach, M. H. Eileen Tan, Kelly M. Suino-Powell, Yuanzheng He, Yong Xu, Michael J. Chalmers, Joseph S. Brunzelle, Huiming Zhang, Huaiyu Yang, Hualiang Jiang, Jun Li, Eu-Leong Yong, Sean Cutler, Jian-Kang Zhu, Patrick R. Griffin, Karsten Melcher, and H. Eric Xu. Molecular Mimicry Regulates ABA

Signaling by SnRK2 Kinases and PP2C Phosphatases. *Science*, 335(6064):85–88, January 2012. Publisher: American Association for the Advancement of Science Section: Report.

108. Choonkyun Jung, Jun Sung Seo, Sang Won Han, Yeon Jong Koo, Chung Ho Kim, Sang Ik Song, Baek Hie Nahm, Yang Do Choi, and Jong-Joo Cheong. Overexpression of AtMYB44 Enhances Stomatal Closure to Confer Abiotic Stress Tolerance in Transgenic Arabidopsis. *Plant Physiology*, 146(2):623–635, February 2008. Publisher: American Society of Plant Biologists Section: ENVIRONMENTAL STRESS AND ADAPTATION TO STRESS.

109. Tongtong Xue, Dong Wang, Shizhong Zhang, Juergen Ehlting, Fei Ni, Stephen Jakab, Chengchao Zheng, and Yuan Zhong. Genome-wide and expression analysis of protein phosphatase 2C in rice and Arabidopsis. *BMC genomics*, 9:550, November 2008.

110. Masrur R. Jaradat, J. Allan Feurtado, Daiqing Huang, Yongquan Lu, and Adrian J. Cutler. Multiple roles of the transcription factor AtMYBR1/AtMYB44 in ABA signaling, stress responses, and leaf senescence. *BMC Plant Biology*, 13(1):192, November 2013.

111. Nguyen Hoai Nguyen and Jong-Joo Cheong. AtMYB44 interacts with TOPLESS-RELATED corepressors to suppress protein phosphatase 2C gene transcription. *Biochemical and Biophysical Research Communications*, 507(1):437–442, December 2018.

112. Nguyen Hoai Nguyen and Jong-Joo Cheong. H2A.Z-containing nucleosomes are evicted to activate AtMYB44 transcription in response to salt stress. *Biochemical and Biophysical Research Communications*, 499(4):1039–1043, May 2018.

113. Muhammad Salman Haider, Cheng Zhang, Mahantesh M. Kurjogi, Tariq Pervaiz, Ting Zheng, Chaobo Zhang, Chen Lide, Lingfie Shangguan, and Jinggui Fang. Insights into grapevine defense response against drought as revealed by biochemical, physiological and RNA-Seq analysis. *Scientific Reports*, 7, October 2017.

114. Matthias Arend, Jörg-Peter Schnitzler, Barbara Ehlting, Robert Hänsch, Theo Lange, Heinz Rennenberg, Axel Himmelbach, Erwin Grill, and Jörg Fromm. Expression of the Arabidopsis Mutant abi1 Gene Alters Abscisic Acid Sensitivity, Stomatal Development, and Growth Morphology in Gray Poplars. *Plant Physiology*, 151(4):2110–2119, December 2009. Publisher: American Society of Plant Biologists Section: DEVELOPMENT AND HORMONE ACTION.

115. Yong-Tao Yu, Zhen Wu, Kai Lu, Chao Bi, Shan Liang, Xiao-Fang Wang, and Da-Peng Zhang. Overexpression of the MYB37 transcription factor enhances abscisic acid sensitivity, and improves both drought tolerance and seed productivity in Arabidopsis thaliana. *Plant Molecular Biology*, 90:267–279, 2016.

116. Qingwen Shen, Xinqiao Zhan, Pei Yang, Jing Li, Jie Chen, Bing Tang, Xuemin Wang, and Yueyun Hong. Dual Activities of Plant cGMP-Dependent Protein Kinase and Its Roles in Gibberellin Signaling and Salt Stress. *The Plant Cell*, 31(12):3073–3091, December 2019.

117. Marc Behr, Gea Guerriero, Jacqueline Grima-Pettenati, and Marie Baucher. A Molecular Blueprint of Lignin Repression. *Trends in Plant Science*, 24(11):1052–1064, November 2019.

118. Won-Chan Kim, Joo-Yeol Kim, Jae-Heung Ko, Hunseung Kang, and Kyung-Hwan Han. Identification of direct targets of transcription factor MYB46 provides insights into the transcriptional regulation of secondary wall biosynthesis. *Plant Molecular Biology*, 85(6):589–599, August 2014.

119. Yoshimi Nakano, Masatoshi Yamaguchi, Hitoshi Endo, Nur Ardiyana Rejab, and Misato Ohtani. NAC-MYB-based transcriptional regulation of secondary cell wall biosynthesis in land plants. *Frontiers in Plant Science*, 6, 2015. Publisher: Frontiers.

120. Lodovico Tamagnone, Angel Merida, Adrian Parr, Steve Mackay, Francisco A. Culianez-Macia, Keith Roberts, and Cathie Martin. The AmMYB308 and AmMYB330 Transcription Factors from Antirrhinum Regulate Phenylpropanoid and Lignin Biosynthesis in Transgenic Tobacco. *The Plant Cell*, 10(2):135–154, February 1998. Publisher: American Society of Plant Biologists Section: Research Article.

121. Apurva Bhargava, Shawn D. Mansfield, Hardy C. Hall, Carl J. Douglas, and Brian E. Ellis. MYB75 Functions in Regulation of Secondary Cell Wall Formation in the Arabidopsis Inflorescence Stem. *Plant Physiology*, 154(3):1428–1438, November 2010. Publisher: American Society of Plant Biologists Section: SYSTEMS BIOLOGY, MOLECULAR BIOLOGY, AND GENE REGULATION.

122. Anna E. Kreynes, Zhenhua Yong, and Brian E. Ellis. Developmental phenotypes of Arabidopsis plants expressing phosphovariants of AtMYB75. *Plant Signaling & Behavior*, 16(1):1836454, January 2021. Publisher: Taylor & Francis _eprint: https://doi.org/10.1080/15592324.2020.1836454.

123. Scott C. Stelpflug, Rajandeep S. Sekhon, Brieanne Vaillancourt, Candice N. Hirsch, C. Robin Buell, Natalia de Leon, and Shawn M. Kaeppler. An Expanded Maize Gene Expression Atlas based on RNA Sequencing and its Use to Explore Root Development. *The Plant Genome*, 9(1):plantgenome2015.04.0025, 2016. _eprint: https://acsess.onlinelibrary.wiley.com/doi/pdf/10.3835/plantgenome2015.04.0025.

124. Hagai Cohen, Vadim Fedyuk, Chunhua Wang, Shuang Wu, and Asaph Aharoni. SUBERMAN regulates developmental suberization of the Arabidopsis root endodermis. *The Plant Journal*, 102(3):431–447, 2020. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/tpj.14711.

125. José Graça. Suberin: the biopolyester at the frontier of plants. *Frontiers in Chemistry*, 3, 2015. Publisher: Frontiers.

126. Ryushiro D. Kasahara, Michael F. Portereiko, Linda Sandaklie-Nikolova, David S. Rabiger, and Gary N. Drews. MYB98 Is Required for Pollen Tube Guidance and Synergid Cell Differentiation in Arabidopsis. *The Plant Cell*, 17(11):2981–2992, November 2005. Publisher: American Society of Plant Biologists Section: Research Article.

127. Jayson A. Punwani, David S. Rabiger, Alan Lloyd, and Gary N. Drews. The MYB98 subcircuit of the synergid gene regulatory network includes genes directly and indirectly regulated by MYB98. *The Plant Journal*, 55(3):406–414, 2008. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1365-313X.2008.03514.x.

128. Varsha Tirumalai, Chenna Swetha, Ashwin Nair, Awadhesh Pandit, and Padubidri V Shivaprasad. miR828 and miR858 regulate VvMYB114 to promote anthocyanin and flavonol accumulation in grapes. *Journal of Experimental Botany*, 70(18):4775–4792, September 2019.

129. Dongshu Guo, Jinzhe Zhang, Xinlei Wang, Xiang Han, Baoye Wei, Jianqiao Wang, Boxun Li, Hao Yu, Qingpei Huang, Hongya Gu, Li-Jia Qu, and Genji Qin. The WRKY Transcription Factor WRKY71/EXB1 Controls Shoot Branching by Transcriptionally Regulating RAX Genes in Arabidopsis. *The Plant Cell*, 27(11):3112–3127, November 2015.

130. Christoph J. Thieme, Monica Rojas-Triana, Ewelina Stecyk, Christian Schudoma, Wenna Zhang, Lei Yang, Miguel Miñambres, Dirk Walther, Waltraud X. Schulze, Javier Paz-Ares, Wolf-Rüdiger Scheible, and Friedrich Kragler. Endogenous *Arabidopsis* messenger RNAs transported to distant tissues. *Nature Plants*, 1(4):15025, April 2015.

131. Petra Stirnberg, Jin-Ping Liu, Sally Ward, Sarah L Kendall, and Ottoline Leyser. Mutation of the cytosolic ribosomal protein-encoding RPS10B gene affects shoot meristematic function in Arabidopsis. *BMC Plant Biology*, 12:160, September 2012.