1 **Title:**
2 Predictability and parallelism in the contemporary evolution of hybrid genomes
3

4 **Authors:** Quinn K. Langdon[1,2*], Daniel L. Powell[1,2], Bernard Kim[1], Shreya M. Banerjee[1,2],
5 Cheyenne Payne[1,2], Tristram O. Dodge[1,2], Ben Moran[1,2], Paola Fascinetto-Zago[2,3], Molly
6 Schumer[1,2,4*]
7
8 [1]Department of Biology, Stanford University
9 [2]Centro de Investigaciones Científicas de las Huastecas "Aguazarca", A.C.
10 [3]Department of Biology, Texas A&M University
11 [4]Hanna H. Gray Fellow, Howard Hughes Medical Institutes
12
*Correspondence to: qlangdon@stanford.edu and schumer@stanford.edu

<u>Abstract</u>

Hybridization between species is widespread across the tree of life. As a result, many
species, including our own, harbor regions of their genome derived from hybridization. Despite
the recognition that this process is widespread, we understand little about how the genome
stabilizes following hybridization, and whether the mechanisms driving this stabilization tend to
be shared across species. Here, we dissect the drivers of variation in local ancestry across the
genome in replicated hybridization events between two species pairs of swordtail fish:
Xiphophorus birchmanni × *X. cortezi* and *X. birchmanni* × *X. malinche*. We find unexpectedly
high levels of repeatability in local ancestry across the two types of hybrid populations. This
repeatability is attributable in part to the fact that the recombination landscape and locations of
functionally important elements play a major role in driving variation in local ancestry in both
types of hybrid populations. Beyond these broad scale patterns, we identify dozens of regions of
the genome where minor parent ancestry is unusually low or high across species pairs. Analysis
of these regions points to shared sites under selection across species pairs, and in some cases,
shared mechanisms of selection. We show that one such region is a previously unknown hybrid
incompatibility that is shared across *X. birchmanni* × *X. cortezi* and *X. birchmanni* × *X. malinche*
hybrid populations.

Introduction

  In the past decade the number of species with sequenced genomes has risen exponentially [1–3]. This has allowed for a greater exploration of the evolutionary histories of the diversity of life on the planet. One consequence of this work has been a new understanding of the ubiquity of genetic exchange between species [4–6]. Studies have demonstrated the impact of both ancient and recent hybridization on the genomes of extant species from fruit flies to humans [7–14]. While it is now clear that hybridization is commonplace, what is less understood is how the genome changes after admixture and what genetic and evolutionary factors generally shape this process.

  An important piece of predicting how the genome will respond after hybridization is understanding the sources of selection that act on hybrids. A large number of studies have shown that hybrids between different species tend to experience fitness consequences; they are often less capable of surviving and reproducing than the species that formed them [15–20]. Several types of selection on hybrids can drive this pattern. Decades of empirical and theoretical work have indicated that Dobzhansky-Muller hybrid incompatibilities (DMIs), or incompatible interactions between mutations that are derived in each of the parental species, are a common cause of inviability and infertility in hybrids [21–24]. Hybrids can also suffer from the effects of intermediate or transgressive traits that place them far from the phenotypic optimum for either parental species [25–28]. In both of these scenarios, selection is generally expected to act against alleles derived from the "minor" parent species, or the species that contributed less to the genome of the hybrids. This is because removing incompatible alleles derived from the minor parent species (or variants that move a hybrid further away from the major parent's phenotypic optimum) is the fastest route to restoring fitness [6]. In addition, recent work has indicated that another mechanism of selection against hybrids can stem from differences in historical effective populations size between the parental species. Such differences can result in the disproportionate accumulation of deleterious mutations along the parental lineage with the smaller historical effective population size and drive the removal of haplotypes from this parental species after hybridization [19,29]. However, this mechanism is most likely to impact hybrids that form between species with dramatically different historical effective population sizes [19,29].

  Although it is well-established that hybrids between species frequently experience selection, it is less clear how predictable the outcomes of selection are at a genome-wide scale, both within and between species pairs. While it has been appreciated for decades that certain regions of the genome, such as sex chromosomes, tend to retain less minor parent ancestry [30–34], recent studies have highlighted other patterns that appear to be consistent across several independent hybridization events. These include broad selection against minor parent ancestry across the genome (presumably due to the mechanisms outlined above) and stronger depletion of minor parent ancestry in regions of the genome with a high density of conserved or coding elements [8,12,35,36]. Another key variable highlighted by empirical and theoretical studies is the importance of the interplay between recombination rate and selection, with several groups finding that selection on hybrids generates a positive correlation between local recombination rate and minor parent ancestry [8,35–41]. This correlation is thought to be driven by how the local recombination rate impacts the decoupling of neutral or adaptive alleles from alleles that are deleterious in hybrids. Neutral or adaptive minor parent alleles that occur in regions of the genome with high local recombination rates are less likely to remain linked to deleterious

neighbors, resulting in the retention of more minor parent haplotypes in such regions after
selection.
These emerging findings highlight how certain features of genome organization can
interact with selection after hybridization in a predictable way across diverse species [6]. This
suggests that we can expect some broad scale repeatability in the outcomes of hybridization
across species and the types of regions of the genome that are more or less permeable to
introgression. In hybridization events between pairs of closely related species we may expect
repeatability in the outcomes of genome evolution after hybridization to be even higher. This is
in part because close relatives are expected to share genome composition and conserved local
recombination maps (in some species; [42,43]). However, in close relatives, repeatability in the
outcomes of hybridization could also be driven by shared mechanisms of selection acting on
individual sites. For example, close relatives may share DMIs due to shared phylogenetic history
[44] or similarities in underlying genetic networks [45] and among some species groups, certain
alleles may be globally adaptive, leading to shared ancestry driven by adaptive introgression.
Thus, leveraging comparisons of independent natural hybridization events between closely
related species pairs is one promising approach for evaluating the factors that drive repeatable
evolution after hybridization [12].
Recently formed hybrid populations are an especially powerful tool for addressing these
questions because ancestry variation along the genome can reflect the action of strong selection
on hybrids. Here, we leverage swordtail fish species (genus *Xiphophorus*) to investigate genome
evolution after several recent, independent hybridization events. Natural replicate hybrid
populations formed between *X. birchmanni* and *X. malinche* are an emerging model system for
studying genome evolution after hybridization. Research on this sister species pair has shown
that variation in local ancestry along the genome is best explained by the presence of prevalent
hybrid incompatibilities (as opposed to historical effective populations size differences in the
parental species or ecological drivers of ancestry variation; [35]), and has documented strong
DMIs between them [23]. Here, we expand on our recent description of replicate hybrid
populations of the species pair *X. birchmanni* × *X. cortezi* [46] to explore repeatability in local
ancestry between these two types of hybrid populations.
X. birchmanni × *X. cortezi* are more deeply diverged than *X. birchmanni* × *X. malinche*
(~450k generations diverged and ~250k generations diverged respectively) and unlike *X.*
birchmanni and *X. malinche*, they do not differ in their time-averaged historical effective
population sizes (Fig. S1) [46]. Hybrid populations between both species pairs formed recently
(~100-150 generations ago [20,46]), likely due to human disturbance [47]. Comparisons across
these two types of hybridization events provide one of the first windows into the drivers of
evolution after hybridization in related species, allowing us to address questions about
predictability and parallelism in selection on hybrids at both a local and broad-scale along the
genome.
116

Results

Background

      Throughout the manuscript we explore repeatability in genome evolution across two types of hybrid populations, those formed between the *X. birchmanni* × *X. cortezi* species pair and those formed between the *X. birchmanni* × *X. malinche* species pair. These three species are closely related, with divergence between *X. birchmanni* × *X. cortezi* slightly higher than divergence between *X. birchmanni* × *X. malinche* (Fig. 1; sequence divergence 0.6% and 0.4% per basepair respectively). Nucleotide diversity and inferred long-term effective population sizes are similar between *X. birchmanni* and *X. cortezi* ($\pi \sim$ 0.1% per basepair; [46]) and lower in *X. malinche* ($\pi \sim$ 0.03% per basepair; [35]) (Fig. S1). *X. birchmanni* is distributed between *X. malinche* and *X. cortezi*, with hybridization occurring in areas of the rivers where the species are sympatric (Fig. 1); *X. malinche* and *X. cortezi* are allopatric throughout their ranges. We have previously shown that replicate hybrid populations in both species pairs formed recently, within the last ~100-150 generations [20,35,46,48], likely due to human-mediated habitat disturbance [47].

      For clarity, we refer to the hybridization events between each pair of species as the two "types" of hybridization events. In addition, we analyzed data from two *X. birchmanni* × *X. cortezi* (Huextetitla and Santa Cruz) and three *X. birchmanni* × *X. malinche* (Acuapa, Aguazarca, and Tlatemaco) hybrid populations, each of which occur in different rivers or tributaries (Fig. 1). When discussing comparisons within a hybridization type (e.g. *X. birchmanni* × *X. cortezi* or *X. birchmanni* × *X. malinche*) we refer to these geographically distinct samples as "replicate" hybrid populations. Past work in the *X. birchmanni* × *X. malinche* system has shown that the replicate hybrid populations formed independently [49]. Additionally, they also differ in their genome-wide ancestry: *X. birchmanni* is the minor parent in Tlatemaco, while *X. malinche* is the minor parent in Acuapa and Aguazarca (Fig. 1).

      Despite the fact that similar factors (i.e. population disturbance) have likely driven recent hybridization across the two species pairs [46,47], each replicate population occurs in a distinct ecological environment, and varies in the degree of assortative mating observed [35,46,48,49]. For a more detailed description of what is known about the ecological environments and mating dynamics of each population from past work see Supporting Information 1.

      The Santa Cruz and Huextetitla hybrid populations occur along geographically separated tributaries of the same river (Fig. 1) and have similar genome-wide admixture proportions and ancestry structure [46]. Hybrids in both populations derive 85-90% of their genomes from the *X. cortezi* parent species (Fig. 1). Using a hidden Markov model (HMM) based approach, we collected low-coverage whole genome sequence data and inferred local ancestry across the genome in 254 hybrids from the two populations (see Methods). While we initially planned to use the two sites to explore repeatability in genome evolution within the replicate populations, we found evidence that they were not completely independent hybridization events (Fig. S2-S3; Supporting Information 2). Thus, we focus our discussion on the Santa Cruz hybrid population but present analyses for the Huextetitla hybrid population (for which results are qualitatively similar) in parts of the main text and supplement.

Inference of demographic histories of hybrid populations

      Initial work investigating the demographic history of the *X. birchmanni* × *X. cortezi* hybrid populations at Santa Cruz and Huextetitla [46] suggested that they formed recently and

experience low levels of migration from the parental species [46]. In this paper we expanded our
sampling from previous work [46] and explored the demographic history of *X. birchmanni × X.*
cortezi and *X. birchmanni × X. malinche* hybrid populations in more detail. We used simulations
and an approximate Bayesian computation approach (ABC, Supporting Information 3) to infer
the demographic history of two populations: the Santa Cruz and Acuapa hybrid populations.
Drawing from uniform (or log-uniform) prior distributions for parameters of interest (initial
admixture proportions, time since initial admixture, population size, and migration rates), we
conducted simulations in SLiM and performed rejection sampling based on summary statistics
derived from the observed data (Supporting Information 3). These simulations yielded well-
resolved posterior distributions for most parameters in both hybrid populations (Fig. S4). We
discuss these results in more detail in Supporting Information 3. Briefly, our findings are
consistent with our previous work [46]; both hybrid zones are young and experience limited
migration from the parental species, but the *X. birchmanni × X. cortezi* hybrid population formed
less recently (~120 generations ago) than the *X. birchmanni × X. malinche* (~80 generations ago)
hybrid population (Fig. S4). Together, our ABC results allow us to model plausible demographic
histories for each type of hybrid population to explore expected patterns of genome evolution.
179
Predictable genomic features correlate with minor parent ancestry in X. birchmanni × X. cortezi
hybrid populations
Although admixture began recently between *X. birchmanni* and *X. cortezi*, we observe
substantial heterogeneity in local ancestry along the genome, with regions fixed for both major
and minor parent ancestry (e.g. Fig. S3). While severe bottlenecks could also generate fixation in
recently formed hybrid populations ([46]), this variance in local ancestry could also be indicative
of selection on hybrids. Indeed, simulations of variation in local ancestry using the demographic
parameters we infer with ABC (admixture time, admixture proportion, and hybrid population
size) suggest that the patterns of fixation in major and minor parent ancestry we observe are
unexpected under the most likely scenario of neutral admixture (Fig. S5; Supporting Information
3).
We next evaluated correlations between minor parent ancestry and genomic features
known to interact with selection in other hybrid systems. Studies in a number of systems have
found that selection on hybrids can drive a positive correlation between recombination rate and
minor parent ancestry [8,35,36,40]. Using averages of minor parent ancestry and recombination
rate summarized in non-overlapping windows we found strong evidence that recombination rate
is positively correlated with minor parent ancestry at both a local (Fig. 2; Santa Cruz – 100 kb
windows: $\rho = 0.44$, $p<10^{-300}$; Huextetitla – 100 kb windows: $\rho = 0.42$, $p<10^{-275}$) and
chromosome-wide scale (Supporting Information 4; Fig. S6-7) in replicate hybrid populations of
X. birchmanni × X. cortezi. This pattern is robust across spatial scales (Table S1) and when
controlling for a number of possible technical variables; e.g. thinning windows based on linkage
disequilibrium and use of recombination rate estimates from observed crossovers in F_2s (see
Table S2-S3).
We also investigated how ancestry in *X. birchmanni × X. cortezi* hybrid populations
covaries with coding and conserved basepairs across the genome. Based on our previous findings
in *X. birchmanni × X. malinche* hybrid populations [35], we predicted that regions with more
coding and conserved basepairs would be depleted in minor parent ancestry. We summarized
average minor parent ancestry and number of coding and conserved basepairs in non-overlapping
genetic windows, to account for variation in recombination rate, and found depletion of minor

parent ancestry in regions especially dense in coding and conserved basepairs (Fig. 2; Table S4;
Santa Cruz – 0.1 cM windows coding basepairs: $\rho = -0.15$, p<10^{-73}, conserved basepairs: $\rho = -$
0.22, p<10^{-161}; Huextetitla – 0.1 cM windows: coding basepairs: $\rho = -0.09$, p<10^{-27}, conserved
basepairs: $\rho = -0.16$, p<10^{-79}). This pattern is robust across a range of window sizes (Table S4),
and when controlling for a number of potential technical artifacts (Tables S5-S9). Interestingly,
we did not detect an impact of nonsynonymous substitutions between species on minor parent
ancestry after controlling for local substitution rates (Supporting Information 5; Fig. S8).
216
Sources of selection driving variation in minor parent ancestry
Above we confirmed via simulations that the observed variation in local ancestry is not
expected by chance or as a consequence of the demographic history of hybrid populations
(Supporting Information 3), motivating us to ask about the role different sources of selection
might play in driving the broad-scale ancestry patterns we observed in *X. birchmanni* × *X. cortezi*
type and *X. birchmanni* × *X. malinche* type hybrid populations. To do so, we used simulations,
grounded in the inferred demographic history of the hybrid populations (Supporting Information
3,6).
In our empirical data, minor parent ancestry is consistently depleted in regions of the
genome with low recombination rate (and higher coding/conserved density), regardless of *which*
species is the minor parent in a given population. Our simulations indicate that these positive
correlations between minor parent ancestry and recombination rate across replicate populations
and species pairs are best explained by a model where selection acts against minor parent
ancestry at many sites across the genome. In simulations we use a model of selection against
hybrid incompatibilities, which results in selection against minor parent ancestry at incompatible
sites and has been well-documented in swordtails [20,23,35,50]. We note, however, that a subset
of other models of selection on hybrids could generate similar results (see discussion in
Supporting Information 6; reviewed in [6]). However, the patterns we observe are inconsistent
with models of selection against hybridization load or global selection against one of the parental
species (Fig. S9; Supporting Information 6). This latter finding indicates that selection on the *X.*
birchmanni genome, which is a shared parent species in both hybrid population types, does not
underlie the patterns that we document.
239
Repeatability extends across hybrid population types
Given evidence for selection against minor parent ancestry in hybrid populations of both
types, the key question we wish to understand is how repeatable the outcomes of admixture are
across different species pairs, and what mechanisms contribute to this repeatability. Past work
has suggested that we expect repeatability in local ancestry in replicate hybrid populations
formed between the same species pairs [35,51]. This repeatability often exceeds expectations
from shared genetic architecture (e.g. recombination rate and functional basepair density),
indicating that selection on the same sites in replicate hybrid populations drives concordance in
local ancestry [35,51].
We wanted to evaluate whether there was evidence for a similar phenomenon across the
two distinct types of hybridization events studied here. The *X. birchmanni* × *X. cortezi*
hybridization events are independent from each of the three *X. birchmanni* × *X. malinche* hybrid
populations (Fig. 1). In addition, the three replicate *X. birchmanni* × *X. malinche* hybrid
populations we analyze (Acuapa, Aguazarca, and Tlatemaco) occur in geographically isolated
river systems (Fig. 1) and formed independently of each other [49].

We observe repeatability in local patterns of minor parent ancestry across the genome
between most pairs of hybrid populations formed between these distinct species pairs (in 100 kb
non-overlapping windows $\rho_{\text{Santa Cruz - Acuapa}} = 0.25$, p<$10^{-93}$; $\rho_{\text{Santa Cruz - Aguazarca}} = 0.17$, p<$10^{-42}$;
Fig. 2; Table S10), an intriguing observation given the demographic independence of these
hybrid populations. These results were robust to a number of factors, including accounting for
the locations of structural differences between the species (see Table S11; Supporting
Information 4, 6). One exception to this pattern is in comparisons involving the Tlatemaco
population. We speculate that this is driven by the fact that hybrids in this population derive the
majority of their genome from *X. malinche*, a species with low historical effective population
sizes (Fig. S1). We discuss this pattern in more detail in Supporting Information 6.
Because many features of the architecture of the genome, such as the local recombination
rate and the locations of coding and conserved basepairs, are largely conserved across *X.*
birchmanni, *X. malinche*, and *X. cortezi* (Supporting Information 4), we wanted to evaluate
whether cross-population repeatability exceeded what was expected from these features alone.
Partial correlation analysis of local ancestry across pairs of populations, including these features
as covariates, indicated that some of the signal of shared local ancestry across hybrid population
types can be explained by these features (Table S12). However, after accounting for this shared
genomic architecture, we still observed repeatability in patterns of minor parent ancestry
between most pairs of hybrid populations, including those formed between distinct species pairs
(in 100 kb non-overlapping windows $\rho_{\text{Santa Cruz - Acuapa}} = 0.15$, p<$10^{-33}$; $\rho_{\text{Santa Cruz - Aguazarca}} = 0.09$,
p<10^{-15}; Table S12). Simulations indicate that the observed correlations in local ancestry across
the two types of hybridization events are unlikely to be driven by demographic history but can
readily be generated by shared sites under selection (Fig. S10; Supporting Information 3,6).
278
Shared islands and deserts of minor parent ancestry across hybrid population types
The analyses described above indicate that not all cross-population repeatability in
ancestry can be explained by shared genomic architecture between these closely related species
pairs (Table S10, S12-S13; Supporting Information 6). We next explored how other factors, such
as shared sites where minor parent ancestry is favored or disfavored, contributed to repeatability
in local ancestry. Specifically, we asked whether there were certain regions of the genome with
shared patterns of minor parent ancestry across hybrid population types, and whether we could
identify the mechanisms of selection driving these shared patterns.
We first focused on the Santa Cruz *X. birchmanni* × *X. cortezi* population to identify
regions of the genome with unusual minor parent ancestry. We identified minor parent "deserts"
by scanning the genome for ancestry informative sites that fell in the lower 2.5% tail of genome-
wide ancestry. We then expanded out from these sites to define regions of reduced minor parent
ancestry (defined as those falling in the lower <5% tail of genome-wide ancestry; see Methods).
These regions were then filtered to exclude very short ancestry tracts, those supported by few
ancestry informative sites, and were merged with nearby deserts (see Methods; Fig. S11). This
procedure resulted in the identification of 81 minor parent deserts with an average length of 317
kb spread across 18 of the 24 swordtail chromosomes. Of these 81 minor parent ancestry deserts,
21 (26%) overlapped with regions of low minor parent ancestry in one or more replicate *X.*
birchmanni × *X. malinche* hybrid population (see Methods).
Using the same approach, we identified 76 regions of unusually high minor parent
ancestry (minor parent ancestry "islands"), in the Santa Cruz *X. birchmanni* × *X. cortezi*
population, where minor parent ancestry fell in the upper 2.5% tail of the genome wide ancestry

distribution. Approximately 25% of these minor parent islands were also identified in one or
more *X. birchmanni × X. malinche* hybrid population.
We next explored whether the amount of sharing of both deserts and islands exceeded
what would be expected by chance using two approaches. First, we randomly sampled 0.05 cM
windows from each focal population and counted the number of windows overlapping with
detected deserts and islands that had low or high minor parent ancestry. With this approach we
found that sharing of both low and high minor parent ancestry regions in the real data exceeded
sharing expected by chance (Fig. 3A; Methods; both deserts and islands p<0.001 by
permutation). Because this simple approach could introduce artifacts by disrupting
autocorrelation in local ancestry along the genome, we also evaluated expected ancestry sharing
by chance using an approach where we preserved the structure of local ancestry by shuffling the
genome in large chunks (Supporting Information 8, Fig. S12; deserts p<0.008, islands p=0.2, by
permutation). While both approaches indicated that the number of shared minor parent deserts in
the real data is unexpected by chance, support for shared islands is weaker.
We also evaluated shared ancestry deserts and islands in light of a number of technical
factors that could bias ancestry calls and potentially drive artifacts in the data, such as variation
in power to infer ancestry along the genome and coincidence of minor parent deserts and islands
with repetitive regions (Supporting Information 8). We do not find evidence for systematic
differences in power or expected error rates between shared and unshared regions (Fig. S13-14;
Supporting Information 8). We also found that shared minor parent deserts and islands are not
unusual in the number of conserved or coding basepairs they contain, suggesting that they are not
driven by regions with an especially high or low density of conserved or coding basepairs
(Supporting Information 8; Fig. S15). Moreover, minor parent deserts were still ancestry outliers
when compared to regions of the genome predicted to have especially high constraint and minor
parent islands remained ancestry outliers when compared to regions of the genome predicted to
have especially low constraint (Fig. S16; Supporting Information 8). Finally, simulations
indicated that the approach we use to identify shared minor parent deserts has good power even
at moderate selection coefficients (e.g. *s*~0.05; Supporting Information 7). Overall, our analyses
suggest that our approach to identifying regions of high and low minor parent ancestry between
hybrid population types is robust to technical artifacts.
331
Features coinciding with shared minor parent ancestry deserts and islands
The unexpectedly high number of shared minor parent ancestry deserts and islands is
consistent with a model where there are some shared loci under selection across species pairs.
These shared ancestry patterns could be driven by a number of shared sources of selection on
hybrids across the two species pairs, including selection against hybrid incompatibilities, shared
signatures of adaptive introgression, among other possible mechanisms (Supporting Information
6). We evaluated the locations of minor parent deserts and islands across the two types of hybrid
populations to begin to explore the mechanisms of selection driving them. Two of the 21 shared
minor parent deserts overlap with known segregation distorters in *X. birchmanni × X. malinche*
that are likely caused by hybrid incompatibilities (see next section), and three shared deserts
correspond to structural differences between species on chromosome 7 and chromosome 17.
Another shared desert was found in all replicate hybrid populations where *X. birchmanni* is the
minor parent. This region spans the first two megabases of chromosome 21, the putative sex
chromosome (Fig. S17), and is depleted in *X. birchmanni* ancestry. However, consistent with
previous work in swordtails, we do not detect an enrichment of shared minor parent deserts (or

islands) on chromosome 21 (Supporting Information 8), possibly because sex chromosomes are
young and not strongly differentiated between males and females [52].
Of the 19 regions with especially high minor parent ancestry across hybrid population
types, two correspond to an inversion that occurs on chromosome 17 and one corresponds to an
inversion on chromosome 19 (Fig. S18). Intriguingly, inversions that differed between species
had unusually high minor parent ancestry in all hybrid populations on average (Fig. S19), in
contrast to what is most frequently observed in other systems ([53–55] but see [40,56,57]).
The remaining minor parent deserts and islands do not coincide with known structural
differences between species [35,46], mapped hybrid incompatibilities between *X. birchmanni* ×
X. malinche [20,23,50], or mapped QTL underlying trait differences between the species [23,58].
We investigate other features associated with these regions below.
358
Ancestry at sites under selection in X. birchmanni × X. malinche hybrids
We observe some overlap between known hybrid incompatibilities and minor parent
ancestry deserts, suggesting that this is one mechanism that can drive cross-population
repeatability in ancestry (among several plausible mechanisms; Supporting Information 6). To
test the role of hybrid incompatibilities in driving cross-repeatability more formally, we took
advantage of sites likely to be involved in hybrid incompatibilities in replicate *X. birchmanni* ×
X. malinche hybrid populations and asked if they had particularly low minor parent ancestry in
X. birchmanni × *X. cortezi* hybrid populations.
Nearly a decade of work has aimed to identify hybrid incompatibilities between *X.*
birchmanni × *X. malinche* using natural hybrid populations and artificial crosses [20,23,35,50].
We focused on two available datasets: putative segregating DMIs identified in natural hybrid
populations between *X. birchmanni* × *X. malinche* [20,50] as well as six segregation distorters
identified in F_2 hybrids of lab crosses between *X. birchmanni* × *X. malinche* that show ancestry
patterns consistent with the action of strongly selected DMIs (e.g. $s>0.25$; see Methods; Fig.
S20). For the segregation distorter dataset, we found that sites under selection in *X. birchmanni* ×
X. malinche F_2 hybrids had lower minor parent ancestry on average in the replicate *X.*
birchmanni × *X. cortezi* hybrid populations (Fig. 3B). Moreover, we show that this result is
robust to excluding a segregation distorter that is also a shared minor parent ancestry desert
across all populations (Fig. S20-21). We do not see an effect of DMIs mapped in *X. birchmanni*
× *X. malinche* natural hybrid populations that were inferred to be under weaker selection, which
could indicate that they are not enriched for hybrid incompatibilities between *X. birchmanni* × *X.*
cortezi or that we have low power to detect incompatibilities under weak selection [20].
381
Mapping a newly identified shared hybrid incompatibility
In the previous section we showed that hybrid incompatibilities identified in one species
pair are enriched for low minor parent ancestry in natural hybrid populations formed between
another species pair, indicating that some hybrid incompatibilities are repeatable across the two
types of hybrid populations. One striking signal we observe in our data is the coincidence of an
ancestry desert on chromosome 6 shared across all populations where *X. birchmanni* is the minor
parent and a strong segregation distorter identified in F_2 hybrids between *X. birchmanni* × *X.*
malinche (Fig. 3C-D; see Methods). Further examination of this region indicated that it is
depleted for minor parent ancestry in all replicate *X. birchmanni* × *X. malinche* populations (Fig.
3C-D, though it did not reach the shared desert threshold in all populations).

Since minor parent ancestry in this region appears to be deleterious across all hybrid
populations in both species pairs, we predicted that it was likely to be a shared hybrid
incompatibility ([35]; Supporting Information 6). By definition, hybrid incompatibilities involve
two or more loci, so we set out to identify loci interacting with the chromosome 6 region in
hybrids. Because this region appears to be under selection in *X. birchmanni* × *X. malinche* F_2
hybrids, we leveraged our data from *X. birchmanni* × *X. malinche* F_2s to scan for interacting loci.
Using deviations from expected two locus genotypes (see Methods), we mapped a locus that
interacts with the chromosome 6 region to chromosome 13 (Fig. 4A). Several genotype
combinations are found in F_2 hybrids at lower than expected frequencies (Fig. 4).
Examining the chromosome 13 region in natural hybrid populations of both *X.*
birchmanni × *X. malinche* and *X. birchmanni* × *X. cortezi*, we found that it also tended to have
lower minor parent ancestry across populations than expected by chance, although it was not
extreme enough to have been identified as a shared desert in our initial scan. At the center of the
associated chromosome 13 region, minor parent ancestry in four out of five of the hybrid
populations fell into the lower 10% quantile genome-wide (Fig. S22; empirical p-value based on
ancestry in 10 kb non-overlapping windows – p<0.005).
The associated regions on chromosome 6 and chromosome 13 identified in F_2 hybrids
between *X. birchmanni* × *X. malinche* contain 13 and 29 genes respectively. We used the
STRING database [59] to evaluate whether any pairs of genes across the two regions are known
to interact, restricting interaction criteria to those supported with experimental evidence,
evidence of dimerization, or evidence of co-expression. Only one pair of genes was annotated as
interacting based on these criteria across the two regions on chromosomes 6 and 13, the
mitochondrial complex I genes *ndufs5* and *ndufa13*. Both these genes are strongly conserved in
other *Xiphophorus* species [60], however several nonsynonymous substitutions distinguish these
three species and appear to be derived in *X. birchmanni* (Table 1). Intriguingly, all of the hybrid
populations analyzed have fixed the mitochondrial haplotype of their major parent species (Fig.
S23; see also [60]), and our cross setup means that all F_2 hybrids had *X. malinche* mitochondria
since the reverse cross is rarely successful (see Methods). We explore the possibility that these
genes form a complex incompatibility with mitochondrial interactors in a companion study [60].
Similar analyses provided some evidence for the presence of additional shared hybrid
incompatibilities involving other shared minor parent deserts (see Methods; Supporting
Information 9). These included evidence for another pairwise hybrid incompatibility involving a
minor parent ancestry desert on chromosome 9 and an interacting locus on chromosome 4 (Fig. 4
C-D) and evidence for a complex hybrid incompatibility involving a minor parent ancestry desert
identified on chromosome 5 (Fig. S24; Supporting Information 9), both detected at FPR
threshold of 10%.
Evidence from simulations indicated that in contrast to good power to detect minor parent
deserts in the population data driven by moderate selection (s~0.05), we have poor power to
detect interactions in the F_2 dataset, with the exception of cases in which selection is extremely
strong (e.g. $s > 0.25$; Supporting Information 7, 9). Given that we likely lack power to detect
many interactions between shared minor parent deserts and other regions of the genome, we also
analyzed the data using a broader approach. We asked whether there was evidence that genes in
minor parent deserts (or islands) were enriched for known protein-protein interactions or high
expression network centrality compared to matched null datasets. We did not find clear evidence
that the interaction networks of genes in minor parent deserts (or islands) deviated from the
genome-wide background (Supporting Information 10).

Table 1. Nucleotide (Nt) and amino acid (AA) changes between species pairs in interacting
genes between the chromosome 6 and chromosome 13 regions. Ratios of rates of
nonsynonymous to synonymous substitutions (dN/dS) inferred by codeml [61] are also shown.
Comparisons where dN/dS is marked as > 99 are due to all nucleotide changes being
nonsynonymous.

Species Comparison	*ndufs5*			*ndufa13*		
	dN/dS	Nt changes	AA changes	dN/dS	Nt changes	AA changes
X. birchmanni - X. cortezi	> 99	5	5	> 99	3	3
X. birchmanni - X. malinche	> 99	5	5	1.2	4	3
X. cortezi – X. malinche	0	0	0	0.001	1	0

445
446
447

Discussion

We now know that species across the tree of life have hybridized with their close relatives and that many species derive substantial proportions of their genomes from past hybridization events. Based on these observations, it is likely that hybridization plays an important role in the evolution of genomes and species, but we still understand relatively little about how this process unfolds. There are major open questions about which mechanisms shape retention and loss of minor parent ancestry after hybridization and how repeatable this process is across species.

While the predictability of evolution has been a classic question in evolutionary biology for decades [62], such questions have been less central to the hybridization literature [6,10,63]. With the expanding number of known hybridization events between closely related species, we are now poised to tackle these questions. One general principle that has emerged is that ancestry from the minor parent species, or the species that contributed less to the genome of hybrids, is more frequently purged by selection after hybridization [6]. This purging can be explained by several mechanisms of selection on hybrids, including selection against hybrid incompatibilities [19,25,28,29,35], and tends to be stronger in regions of the genome with many linked coding or conserved basepairs [8,12,35]. We also uncover these patterns in the *X. birchmanni × X. cortezi* hybrid populations studied here, showing a substantial effect of local recombination rate and linked coding and conserved basepairs on minor parent ancestry in hybrids (Fig. 2). This represents at least the sixth system in which these effects have been detected [8,12,35,36,38,40], suggesting that these patterns of selection on minor parent ancestry are predictable outcomes of the process of genome stabilization after hybridization.

Here, we dramatically expand our understanding of how repeatable the outcomes of hybridization are by comparing patterns of local ancestry in hybridization events between *X. birchmanni × X. cortezi* and *X. birchmanni × X. malinche* species pairs. This allows us to evaluate, for the first time, some of the factors driving repeatability in the outcomes of hybridization across related species pairs with similar histories of admixture. Both hybridization events began recently (in the last ~80-150 generations [20,35]) but occur in different geographic regions between distinct species pairs (Fig. 1). Because we find that broad scale factors like genome structure are important in generating local ancestry variation in both species pairs, this is expected to drive some concordance in local ancestry across the two species pairs. Indeed, we detect repeatability in where in the genome minor parent ancestry is retained or purged across pairs of *X. birchmanni × X. cortezi* and *X. birchmanni × X. malinche* hybrid populations (Fig. 2; Table S10).

Beyond these broad patterns generated by shared genetic architecture, little is known about whether the same genes or regions of the genome are intolerant of hybridization in related species pairs. Intriguingly, we still observe correlations in local ancestry across the two types of hybrid populations after accounting for shared genomic architecture (Table S12). This indicates that although shared genome structure plays a role in some of the repeatability in local ancestry we observe, other factors are likely to contribute, such as shared sites under selection across the two types of hybridization events. Indeed, when we searched for local signals of shared ancestry across the two hybrid population types, we found evidence that some of the same regions of the genome are under selection in hybrid populations of both species pairs. Compared to expected overlap by chance, *X. birchmanni × X. cortezi* and *X. birchmanni × X. malinche* hybrid populations are >10X enriched in both shared minor parent deserts and islands.

What factors contribute to this high repeatability of sites under selection across the two
types of hybrid populations? From first principles, possible mechanisms driving repeatable local
ancestry include shared hybrid incompatibilities or shared regions of adaptive introgression.
Given that the genetic architecture of hybrid incompatibilities has been well-studied in *X.*
birchmanni × *X. malinche* hybrids, we can directly ask about the role of such sites in contributing
to repeatable ancestry across the two types of hybrid populations. In regions identified as likely
hybrid incompatibilities in F_2 hybrids between *X. birchmanni* × *X. malinche*, minor parent
ancestry was on average lower near these sites in *X. birchmanni* × *X. cortezi* hybrid populations.
Only two of these regions overlap with the minor parent deserts discussed above, suggesting that
they represent largely complementary signals of shared sites under selection in replicate *X.*
birchmanni × *X. cortezi* and *X. birchmanni* × *X. malinche* hybrid populations.
Further comparisons across hybrid population types reveal at least one and possibly
multiple shared hybrid incompatibilities between *X. birchmanni* × *X. cortezi* and *X. birchmanni* ×
X. malinche. We identified a single region on chromosome 6 that was a segregation distorter in
early generation hybrids and depleted in minor parent ancestry in all five replicate *X. birchmanni*
× *X. cortezi* and *X. birchmanni* × *X. malinche* hybrid populations. We found that this region on
chromosome 6 interacts with a locus on chromosome 13 (Fig. 4). Based on gene network
analysis, we identified two mitonuclear genes in the chromosome 6 and chromosome 13 regions
that physically interact to form part of mitochondrial complex I. Intriguingly, all hybrid
populations studied here of both *X. birchmanni* × *X. cortezi* and *X. birchmanni* × *X. malinche* are
fixed for mitochondrial haplotypes from the major parent species, and crosses only included
individuals with *X. malinche* mitochondria. This raises the possibility that the incompatibility
may be driven by complex interactions between the mitochondrial genome and its interacting
nuclear counterparts. Indeed, we find evidence that this is the case in *X. birchmanni* × *X.*
malinche hybrids in a companion study [60]. Similar analyses of other regions suggest that there
are multiple shared hybrid incompatibilities between *X. birchmanni* × *X. cortezi* and *X.*
birchmanni × *X. malinche* associated with shared minor parent ancestry deserts (Fig. 4).
While it is clear that hybrid incompatibilities play a role in driving some of the shared
minor parent deserts across the two types of hybrid populations, other mechanisms of selection
are likely important in generating shared minor parent islands (although we note that minor
parent islands are less strongly enriched compared to expectations by chance). In some cases,
they may reflect cases where the minor parent haplotype is globally beneficial (i.e. adaptive
introgression). For example, *X. birchmanni* is the minor parent in both the Santa Cruz and
Tlatemaco populations, hinting that *X. birchmanni* haplotypes may be globally favored within
the four shared minor parent islands identified in these two populations. In other pairs of
populations, however, when the same regions are enriched in minor parent ancestry this means
that different ancestries are favored in the two populations (i.e. in the seven islands shared
between Santa Cruz and Aguazarca, *X. birchmanni* ancestry is overrepresented in the former and
X. malinche in the latter). We speculate that some minor parent islands could correlate with
regions under balancing selection such that minor parent ancestry is generally advantageous or
other mechanisms that can favor minor parent ancestry such as masking of weakly deleterious
alleles [64]. Notably, three shared minor parent islands occur in inversions that differentiate
species, conflicting with the prevailing wisdom that inversions should resist introgression (Fig.
S18; see [40,56,57] for similar findings).
The shared hybrid incompatibility we identify here in *X. birchmanni* × *X. malinche* and
X. birchmanni × *X. cortezi* hybrids is one of the first described cases in any species group.

However, this finding should not be surprising since some hybrid incompatibilities are predicted
to be shared between related species from first principles. Hybrid incompatibilities are typically
described as interactions between derived mutations in each species but incompatibilities can
also arise between the ancestral and derived genotypes when multiple derived mutations have
accumulated along one lineage (Fig. S25; [44]). This type of ancestral-derived incompatibility
may be more likely in cases where there are biased patterns of lineage specific substitutions in
particular gene classes, as appears to be the case for the *X. birchmanni* mitochondrially
interacting proteins discussed above (Table 1). Our findings add to a handful of known cases
where the same or similar genetic interactions are under selection in hybridization events across
multiple species pairs [21,23,65,66]. Whether this phenomenon is widespread awaits progress
mapping additional hybrid incompatibilities in related species pairs [6]. The comparative
approach used here is a powerful strategy for evaluating this question on a genome-wide scale.
Together, our results highlight predictable outcomes of selection after hybridization. We
find that the outcomes of hybridization are repeatable across closely related species pairs, and are
driven in part by broad scale factors such as shared genetic architecture. Beyond shared genomic
architecture, we find that diverse mechanisms contribute to repeatability in local ancestry,
including shared hybrid incompatibilities between species and shared sites where minor parent
ancestry is favored. This parallelism at both the broad and local scale is an important first step
towards untangling the drivers of local variation in ancestry in hybrid genomes.
559
560

Methods

Sample Collection

Wild fish were caught using baited minnow traps in the states of Hidalgo and San Luis Potosí, Mexico [46]. Individuals from populations that contain a mixture of pure *X. birchmanni* individuals and *X. birchmanni × X. cortezi* hybrids were sampled from two distinct collection sites occurring in separate tributaries of the Río Santa Cruz in northern Hidalgo: Huextetitla (21°9'43.82"N 98°33'27.19"W) and Santa Cruz (21°9'27.63"N 98°31'13.79"W). We excluded pure *X. birchmanni* individuals based on genome-wide ancestry, resulting in 254 hybrid individuals included in this study. Ninety-eight of these individuals were originally sequenced as part of a previous study documenting the hybridization event between *X. cortezi* and *X. birchmanni*, and the remaining individuals were collected from the Santa Cruz hybrid population in July of 2020. Collected fish were anesthetized in a 100 mg/mL buffered solution of MS-222 and water (Stanford APLAC protocol #33071). Fish were then photographed and a small fin clip was taken from each individual's caudal fin. Fin clips were preserved in 95% ethanol for later DNA extraction.

Tn5 libraries for low-coverage whole genome sequencing of X. birchmanni × X. cortezi hybrids

We prepared libraries for low-coverage whole genome sequencing from DNA extracted from fin clips collected from fish caught at the Huextetitla and Santa Cruz populations as described in Powell et al. [46]. Briefly, we extracted DNA from fin tissue using the Agencourt DNAdvance kit (Beckman Coulter, Brea, California) using half the recommended reaction volume but otherwise as specified by the manufacturer. We quantified isolated DNA with a TECAN Infinite M1000 microplate reader (Tecan Trading AG, Switzerland). We then prepared tagmentation-based whole genome libraries for low coverage sequencing by enzymatically shearing DNA diluted to approximately 2.5 ng/ul using the Illumina Tagment DNA TDE1 Enzyme and Buffer Kits (Illumina, San Diego, CA) at 55°C for 5 minutes. We amplified the sheared DNA in dually-indexed PCR reactions for 12 cycles. We purified pooled amplified PCR reactions with 18% SPRI magnetic beads. Library concentrations were measured with a Qubit fluorometer (Thermo Scientific, Wilmington, DE) and library size distributions were measured using Agilent 4200 Tapestation (Agilent, Santa Clara, CA). Libraries were sent for sequencing on an Illumina HiSeq 4000 at Admera Health Services, South Plainfield, NJ.

Local ancestry inference in X. birchmanni × X. cortezi hybrids

We previously developed and validated local ancestry inference methods for *X. birchmanni × X. cortezi* hybrids using a combination of demographic inference, simulations, analysis of parental individuals, and analysis of a subset of the hybrid individuals described here [46]. Briefly, we used panels of parental individuals (N=37 for *X. cortezi* and N=55 for *X. birchmanni*) to identify 1.1 million ancestry informative sites genome wide that were nearly fixed between the two parental species (90% frequency difference or greater), yielding approximately 3 ancestry informative sites per 2 kb. Performance was tested on pure parental individuals not included in the reference panel and in a series of simulations [46]. We found that across scenarios tested, accuracy in local ancestry inference was remarkably high (≤0.3% errors per ancestry informative site in all tested scenarios), presumably due to the high density of ancestry informative markers.

We applied a hidden Markov model (HMM) to infer local ancestry based on low-
coverage whole genome sequencing data collected from Huextetitla and Santa Cruz hybrids,
using the pipeline described above [46,67]. Average coverage across individuals was 1.1X and
the minimum coverage cutoff used in this study was 0.15X. We set priors for the number of
generations since initial admixture to 150 and the admixture proportion to 85% *X. cortezi* based
on the results of previous work [46]. We set the error rate for the HMM to 0.02 based on
empirically inferred error and cross-contamination rates for our library preparation protocol [67].
After running the HMM and excluding ancestry informative sites that were not covered in any
individual, we recovered posterior probabilities for each ancestry state at 1,050,362 informative
sites genome wide in 254 individuals.
For many of our analyses, it was convenient to convert these posterior probabilities for a
given ancestry state to hard-calls. We used a posterior probability threshold of 0.9 or greater to
assign a site to a given ancestry state (homozygous *X. birchmanni*, heterozygous, or homozygous
X. cortezi). Sites with lower than 0.9 posterior probability were masked, as were sites that were
present in fewer than 25% of individuals. Minor parent ancestry was summarized across
individuals by averaging the ancestry hard-calls at each site across a range of non-overlapping
window sizes (e.g. 10 kb – 1 Mb; 0.1 – 1 cM).
Historical recombination events in admixed populations are detectable as ancestry
transitions in present day individuals [68]. We were also able to use this data to infer the
approximate locations of crossover events in a hybrid ancestor that resulted in a transition
between ancestry states. We identified the interval over which the posterior probability changes
from 0.9 for one ancestry state to 0.9 in support of a different ancestry state. We excluded
markers that were not in Hardy-Weinberg equilibrium at a Bonferroni corrected p-value of 0.1,
since errors in genotyping and ancestry inference could generate false ancestry switches
(excluded sites in this analysis: 84,366 – 8% of ancestry informative markers). This resulted in
the identification of 168,858 ancestry transitions genome-wide across the two populations with a
median resolution of 22 kb (~29 per individual per chromosome).
The density of ancestry informative sites in swordtails varies with features of interest
such as the local recombination rate and the number of linked functional basepairs [35], likely
due to processes such as background selection within the parental lineages. As a result, we must
be cautious of technical biases that could arise due to the distribution of ancestry informative
sites that could mislead us into inferring biological differences when there are in fact none. To
address this, we thinned the data with respect to the density of ancestry informative sites and re-
inferred local ancestry as described above. On average, an ancestry informative site between *X.*
birchmanni × *X. cortezi* occurs once every 395 basepairs. For the ~50% of the genome where
ancestry informative sites occurred more frequently, we thinned informative sites to retain only
one per 395 basepairs. We inferred ancestry using this thinned set of informative sites (N=
689,983) and performed all of the analyses described on both thinned and unthinned datasets to
ask if any results qualitatively change as a result of the input dataset. Results for the thinned
datasets are reported in Supporting Information 11.
646
Local ancestry inference in X. birchmanni × X. malinche hybrids
For the replicate *X. birchmanni* × *X. malinche* hybrid populations analyzed in our study,
we took advantage of data collected from previous studies of these replicate populations [35,58].
As previously described, we inferred local ancestry with an HMM and used population specific
priors for the admixture proportions and time since initial admixture [23,35]. We limited our

analysis to individuals collected over a single sampling year. Briefly, we included 95 individuals
collected from Tlatemaco in 2013 (resulting in 629,584 ancestry informative sites), 97
individuals collected from Acuapa in 2018 (613,171 sites), and 51 individuals collected in 2016
from Aguazarca (628,811 sites).
Previous work has explored the accuracy of these local ancestry inference approaches in
X. birchmanni × *X. malinche* hybrids in great detail [23,67]. Briefly, initial definition of ancestry
informative markers relied on 183 parental individuals [23], ancestry informative sites that were
not fixed or nearly fixed between the two parental species were excluded (<98% frequency
difference). Performance was evaluated in simulations, on pure parental individuals not used in
the definition of ancestry informative sites and on early generation F_1 and F_2 hybrids where
switches in ancestry over short genetic distances can confidently be inferred to be errors. Based
on these analyses we estimated error rates to be ≤0.2% errors per ancestry informative site in all
tested scenarios [23,67]. Thus, given high expected accuracy of ancestry inference in both *X.*
birchmanni × *X. cortezi* and *X. birchmanni* × *X. malinche* hybrids, we do not expect that errors
will drive to the global or local patterns that we observe in the empirical data.
667
Comparing minor parent ancestry as a function of local genetic architecture
For the *X. birchmanni* × *X. cortezi* hybrid populations, we wanted to evaluate whether
minor parent ancestry was correlated with recombination rate, as had previously been described
in other species [35,36,40]. We summarized average ancestry in non-overlapping 50 kb, 100 kb,
and 250 kb windows for each population. We previously created a linkage disequilibrium (LD)
based recombination map for *X. birchmanni* [35] and calculated recombination rate based on this
map over these same spatial scales. Available data suggests that recombination maps are likely to
be conserved between *X. birchmanni* and *X. cortezi* (Supporting Information 4; [35]), and
between parental species and hybrids [35].
Because the local recombination rate covaries with other genomic features in swordtails
[35], we also calculated the number of coding (or conserved) basepairs in each window so that
we could account for this in our analysis. We performed these calculations using annotations
developed for the *X. birchmanni* genome [23]. We then used a Spearman's partial correlation in
R to determine if ancestry was correlated with recombination rate, after accounting for the
number of coding and conserved basepairs in each window. To aid in interpreting these
empirical results, we also analyzed simulations that were performed under the inferred
demographic history of *X. birchmanni* × *X. cortezi* and *X. birchmanni* × *X. malinche* hybrid
populations, with and without selection (Supporting Information 3, 6).
After evaluating the relationship between recombination rate and ancestry in *X.*
birchmanni × *X. cortezi* populations, we were also interested in the relationship between minor
parent ancestry and the locations of coding and conserved basepairs along the genome. To
control for recombination rate, which is positively correlated with coding and conserved bases in
swordtails [35], we summarized average ancestry in 0.25 cM non-overlapping windows. As
noted above, we used the gene annotation file for *X. birchmanni* and a set of regions conserved
in percomorph fish that we had previously identified using phastCons. Comparisons of
alignments across species indicate that the locations of these elements are largely conserved
across *X. birchmanni*, *X. malinche*, and *X. cortezi* (Supporting Information 4). We counted the
number of coding or conserved basepairs in each window. Using this approach, we evaluated the
relationship between ancestry and the number of linked coding or conserved basepairs with a
Spearman's correlation (and partial correlation) in R. We chose to present results for 0.25 cM

windows in the main text because 50% of minor parent ancestry tracts in Santa Cruz are 0.25
cMs in length or longer (75% are 0.1 cMs in length or greater). However, we evaluated these
relationships across a range of non-overlapping window sizes (0.1-1 cM) and found the results
were qualitatively similar (Table S4).
Because power to accurately infer ancestry may be correlated with the number of nearby
conserved basepairs, we also repeated these analyses using ancestry probabilities generated from
a set of ancestry informative sites thinned to reduce power differences between different regions
of the genome (see above, *Local ancestry inference in X. birchmanni × X. cortezi hybrids*). We
found that our results were qualitatively unchanged (Table S8). We also repeated analyses of the
relationship between the number of conserved and coding basepairs and minor parent ancestry,
excluding windows that fell in the lower or upper 25% quantile of the number of ancestry
informative sites. We found that excluding these windows, where we expect to have particularly
low or high power to infer ancestry, did not impact our results (Table S9).
For analyses including recombination rate, we repeated tests using an LD-based map that
we inferred using the same methods after thinning SNPs in regions of the genome with high SNP
density in the *X. birchmanni* population sample it was originally generated with [35]. Because
our ability to reliably estimate recombination rate depends on the density of SNPs, we expect our
power to estimate recombination rate to vary along the genome. If power to estimate
recombination rate varies with other features of interest, such as density of coding or conserved
basepairs, this could mislead us into interpreting variation in power as variation in signal. To
evaluate this possibility, we calculated the median distance between SNPs in our dataset, which
was 184 basepairs. We thinned the SNPs input into LDhelmet so that when SNPs were denser
than one every 184 basepairs, a single SNP was retained per 184 basepairs. We then re-inferred
recombination rates using LDhelmet as described previously [35]. We repeated all analyses
described above that incorporate recombination rate with this thinned LD map and found the
results qualitatively unchanged (Supporting Information 11).
As a further validation of the relationship between minor parent ancestry and
recombination rate, we repeated analyses with a lower-resolution crossover-based map generated
from the 943 F_2 *X. birchmanni* x *X. malinche* hybrids. We observed a total of 34,939 ancestry
transitions genome-wide in our dataset, or an average of 1.5 per individual per chromosome. We
summarized the number of observed ancestry transitions, reflecting crossovers in the F_1 parents,
in windows. On average we observed only ~5 crossovers per 100 kb, so we chose to summarize
recombination rate in large windows of 1 Mb and 5 Mb. We repeated analyses of the relationship
between recombination rate and minor parent ancestry using this lower resolution crossover map
and found that our results were consistent with those recovered using the LD-based maps (Table
S3).
Finally, we tested the robustness of our results in several additional ways to control for a
number of potential technical artifacts (see full description in Supporting Information 11 and
Table S2). Since unexpectedly short ancestry tracts may represent errors, we repeated our
analyses masking minor parent ancestry tracts that were shorter than 0.004 cM and major parent
ancestry tracts that were shorter than 0.035 cM (tract length expectations given population
history were derived from [69]). To control for potential autocorrelation in nearby windows we
calculated correlations between minor parent ancestry and features of interest when we retained
one window per 500 kb (approximately the scale of admixture LD in these hybrid populations;
[35,46]). To control for the fact that recombination rates in hybrids are expected to differ within

species-specific structural rearrangements, we identified and removed windows spanning known
inversions and repeated our analyses.
745
Repeatability of local ancestry between cross types
To evaluate correlations in minor parent ancestry across *X. birchmanni* × *X. cortezi* and
X. birchmanni × *X. malinche* populations we used summaries of observed ancestry in non-
overlapping physical (50-250 kb) and genetic windows (0.1-1 cM), as described above. We first
calculated pairwise correlations between populations in minor parent ancestry without
accounting for other features (Fig. 2, Table S10). We then repeated these calculations using
partial correlation tests in physical windows (50-250 kb) that included the recombination rate,
the number of linked coding, and conserved basepairs as covariates (Table S12) and partial
correlation tests in genetic windows (0.1-1 cM) that included the number of linked coding and
conserved basepairs as covariates (Table S13). This allowed us to infer both whether broad-scale
correlations in local ancestry could be detected across populations and whether these correlations
could be largely explained by shared genetic architecture (i.e. locations of coding and conserved
basepairs and local recombination rate). We also explored this question using simulations. We
performed simulations under the inferred demographic histories for *X. birchmanni* × *X. cortezi*
and *X. birchmanni* × *X. malinche* hybrid populations with and without selection (Supporting
Information 3, 6) and evaluated expected cross-population correlations in ancestry under
different scenarios.
763
Identification of minor parent ancestry "deserts" and "islands"
Beyond shared correlations as a result of shared broad-scale architecture of the genome,
we were interested if we could identify shared minor parent ancestry "deserts," where minor
parent ancestry was unusually low, and shared minor parent ancestry "islands, where minor
parent ancestry was unusually high, that were shared across *X. birchmanni* × *X. cortezi* and *X.*
birchmanni × *X. malinche* hybrid populations.
To do so we identified regions of especially high and low minor parent ancestry in Santa
Cruz, the *X. birchmanni* × *X. cortezi* hybrid population for which we have the largest dataset (for
results of the same analysis using the Huextetitla populations see Supporting Information 12).
We first identified ancestry informative sites where the minor or major parent ancestry fell in the
lower 2.5% tail of genome-wide ancestry. We expanded out in the 5' and 3' direction from the
focal ancestry informative site until we reached a site on each edge that exceeded the 5%
ancestry quantile, and treated this interval as the boundary of the minor parent ancestry desert or
island. To control for possible technical artifacts, we then evaluated ancestry in these regions in
0.05 cM windows. We identified the window that contained the midpoint of the islands or desert
and checked that this entire window fell in the 10% minor or major ancestry tail, contained at
least 10 ancestry informative sites, and was more than 10 kb in length. We also merged any
detected regions that were closer together than 50 kb (deserts N=54, islands N=14) and removed
any regions shorter than 10 kb (deserts N=1). We also filtered regions where inferred ancestry
varied substantially between inference with unthinned and thinned ancestry informative markers
(Fig. S13). Specifically, we filtered islands where inferred ancestry differed by more than 10%
between thinned and unthinned data (N=58, based on 0.05 cM windows). This resulted in the
identification of 81 minor parent ancestry deserts with an average length of 317 kb, and 76
islands with an average length of 242 kb in the Santa Cruz population.

We next wanted to compare these regions across populations. Because each population
has a distinct recombination history and distribution of ancestry tract sizes, we took the midpoint
of the deserts and islands identified in the Santa Cruz hybrid population and overlapped these
regions with average ancestry in other populations, calculated at a fine spatial scale (0.05 cM, or
on average 11 kb). We then determined if these regions had unusually high or low minor parent
ancestry in other populations (defined as falling in the 10% minor or major ancestry tail in
another population). We identified several shared islands and deserts between Santa Cruz and *X.*
birchmanni × *X. malinche* populations (Fig. 2, 3). Simulations suggest that this two-step
approach has excellent power to detect shared sites under selection, even when selection
coefficients are relatively weak (Supporting Information 7).
Next, we asked whether the number of shared minor parent islands and deserts across *X.*
birchmanni × *X. malinche* and *X. birchmanni* × *X. cortezi* hybrid populations exceeded what
would be expected by chance. To investigate this, we used the same 0.05 cM windows we used
above to determine whether minor parent ancestry deserts and islands were shared. We permuted
the average ancestry values in *X. birchmanni* × *X. malinche* populations across windows and
asked whether any windows that were major or minor parent ancestry outliers in the permuted
data overlapped with the ancestry deserts identified in Santa Cruz (defined as falling in the 10%
tail of major or minor parent ancestry, as in the real data). We repeated this procedure 1000
times. Based on these permutations, we found that few minor (or major) parent ancestry outliers
in *X. birchmanni* × *X. malinche* hybrid populations overlap minor parent deserts (or islands)
identified in the Santa Cruz hybrid population by chance (Fig. 3A). For pairs of populations, the
expected false positive rate across all simulations was less than 10% (Fig. 3A). When we instead
asked about the probability that any minor parent desert or island identified in Santa Cruz
overlapped an ancestry outlier in simulations, the false positive rate was higher (~20%).
Our interest in performing permutations was to determine how much overlap in minor
parent ancestry deserts and islands is expected by chance, and how much we can attribute to
shared sources of selection on hybrids. While our permutations above suggest that little overlap
is expected by chance, one challenge of this type of analysis is that permuting the data disrupts
all correlations in ancestry across hybrid populations. This means that it will also disrupt
correlations driven by interactions between selection and features of the genome such as the local
recombination rate and coding/conserved basepair density. Since we are primarily interested in
minor parent deserts and islands driven by sites under shared selection rather than windows that
happen to be in low recombination rate or high functional density regions of the genome, we
attempted to control for this. First, our permutations use genetic windows, which account in part
for the relationship between minor parent ancestry and recombination rate variation. To evaluate
whether minor parent ancestry deserts and islands were likely driven by functional basepair
density, we compared the windows identified in our analysis to the genomic background. We
describe these methods in detail in Supporting Information 8 and the results of these analyses are
presented in Fig. S14-S16.
Given the importance of this analysis for the conclusions of our manuscript, we also took
two additional approaches to determine the expected false positive rates for desert and island
regions. In the first, we shuffled ancestry in blocks along the genome to preserve local ancestry
structure during permutations but otherwise evaluated false positive rates as described above. We
found the results of these permutations to be similar, but with a somewhat higher false positive
rate for minor parent deserts and a substantially higher false positive rate for minor parent islands
(Supporting Information 8, Fig. S12). In the second, we analyzed expected false positive rates for

each identified desert and island individually. Briefly, for each minor parent desert or island we
generated windows of matched cM length, and selected all windows in the genome that had a
similar content of coding basepairs. We next used permutations to determine the expected
overlap by chance of regions of extreme ancestry within these matched sets (Supporting
Information 8).
839
Identifying loci under selection using controlled crosses and natural populations
Hybridization between $X.\ birchmanni \times X.\ malinche$ has been studied for more than a
decade. As such, we have generated controlled crosses between $X.\ birchmanni \times X.\ malinche$ that
are not yet feasible between $X.\ birchmanni \times X.\ cortezi$. We have also mapped a number of
hybrid incompatibilities that occur between the two species, including a gene interaction that
causes melanoma in hybrids [23] and dozens of pairs of loci that show signatures consistent with
selection against DMIs in natural hybrid populations [20,50].
We also recently generated a large mapping panel of F_2 hybrids between $X.\ birchmanni \times$
$X.\ malinche$, which we used for QTL mapping of male sexually selected traits [70]. Here, we
reanalyzed this mapping panel, doubling our sample size by including females, to identify strong
segregation distorters. Segregation distortion can be generated by selfish genetic elements but in
hybrids segregation distortion is often a hallmark of regions of the genome that impact hybrid
viability or fertility [31]. Since segregation distorters that we have power to detect in our
mapping panel are likely to have a strong impact on viability or fertility of $X.\ birchmanni \times X.$
$malinche$, they also represent loci that we were likely unable to map previously since our initial
work on hybrid incompatibilities relied on segregating variation in $X.\ birchmanni \times X.\ malinche$
hybrid populations [20,23]. Briefly, we generated F_1 hybrids between $X.\ malinche$ females and $X.$
$birchmanni$ males, and produced F_2s by crossing this F_1 generation [70]. The reverse cross
direction is rarely successful [60]. We inferred local ancestry as described previously for this
cross [67,70]. As expected from the cross design, average ancestry across ancestry informative
sites in our dataset of 943 males and females indicated that on average 50% of the genome was
derived from each parental species. Our previously published dataset for QTL mapping included
568 males [70].
In reanalyzing this data, we next turned to identifying regions of the genome with
unusually high $X.\ birchmanni$ or $X.\ malinche$ ancestry, consistent with loci having a major
impact on viability or fertility in hybrids (or generating segregation distortion for other reasons).
We first thinned the F_2 data to retain one ancestry informative marker per 50 kb for
computational efficiency. We then used a binomial test to identify loci that deviated from the
expected segregation ratio of 50% $X.\ birchmanni$ and 50% $X.\ malinche$ ancestry. We used a p-
value threshold of $p<5^{-4}$, approximately corresponding to the 5% tail of ancestry outliers in our
dataset. We note that several of these regions were robust to Bonferroni correction for the
number of markers tested (N= 20,011; $p<5^{-6}$ corresponding to FDR of 10%). We consider this
analysis conservative due to the extent of ancestry linkage disequilibrium in F_2 hybrids. After
identifying regions of the genome with evidence for segregation distortion, we visualized
ancestry in each region to ensure that the pattern was consistent with an extended region of
segregation distortion rather than a short interval as might be expected from e.g. genotyping error
(see Fig. S20).
To explore the strength of selection that would be required to generate segregation
distortion at the threshold we impose on the real data, we performed simple simulations. We
simulated genotypes formed from 50-50 mixtures of the two parental gametes. Next, we

randomly drew a selection coefficient from 0-1 against one of the homozygous genotypes and a
dominance coefficient (0-1). We implemented selection on this pool of genotypes, treating 1-s
(or 1-hs) as the probability of survival. We drew 943 individuals from the surviving genotypes
and calculated average ancestry at the simulated sites. Simulations where ancestry fell in the 5%
tail of the real data were treated as accepted simulations; we repeated this procedure 10,000
times. Based on the distribution of accepted simulations (Fig. S26), we predict that very strong
selection ($s > 0.25$) is required to generate ancestry patterns as extreme as the loci we focus on in
our F_2 dataset. Distributions of the dominance coefficient from accepted simulations mirrored the
prior distribution.
889
890   *Role of loci under selection in ancestry variation in natural populations*
891   One of our major questions is whether loci under selection are shared not just across
hybrid populations of the same cross type but between hybrid populations of different cross
types. To tackle this question we took advantage of what is known about selection in the *X.*
birchmanni × *X. malinche* system, using the segregation distortion regions identified above, and
more weakly selected putative DMIs that have been previously identified in natural hybrid
populations between *X. birchmanni* × *X. malinche* [20,50]. Both sets of loci are associated with
lower minor parent ancestry in *X. birchmanni* × *X. malinche* hybrid populations [35], as expected
by theory.
899   We asked if these loci were also associated with lower minor parent ancestry in *X.*
birchmanni × *X. cortezi* hybrid populations, and the degree to which they coincided with
ancestry deserts that are shared across *X. birchmanni* × *X. cortezi* and *X. birchmanni* × *X.*
malinche hybrid populations. We summarized ancestry in 10 kb non-overlapping windows and
used bedtools [71] to determine the distance of each window to the closest selected site in each
of the two datasets and plotted average major parent ancestry as a function of that distance (Fig.
S27).
906
907   *Identifying potential partners of shared ancestry desert and segregation distortion region*
908   One of the regions that we identified as a significant segregation distorter on
chromosome 6 in F_2 hybrids between *X. birchmanni* × *X. malinche* was also a shared minor
ancestry desert, and had unusually low minor parent ancestry across all surveyed *X. birchmanni*
× *X. malinche* and *X. birchmanni* × *X. cortezi* hybrid populations. The observation that minor
parent ancestry is disfavored across populations of diverse ancestry (majority *malinche*, majority
birchmanni, and majority *cortezi*) indicates that this locus is likely to be involved in a hybrid
incompatibility [35], as opposed to being driven by other possible mechanisms of selection on
hybrids [6].
916   Based on these observations, we wanted to identify the interacting partner of the
chromosome 6 locus. Given the clear expectations for genotype frequencies in F_2 hybrids and
apparent shared selection on this locus across species pairs, we used the F_2 data generated
between *X. birchmanni* × *X. malinche* to perform this scan. We selected an ancestry informative
marker from the center of the shared ancestry desert on chromosome 6, which falls within the
peak segregation distortion region on chromosome 6 identified in F_2 hybrids (Fig. 3). Next, we
performed a scan against all other ancestry informative markers across the genome and
calculated χ^2 statistics for deviations from expected two-locus genotype frequencies (with four
degrees of freedom). Importantly, we used the observed genotype frequencies across individuals
at each of the two loci to generate this expectation, such that we identified loci with deviations

from expected two-locus genotype combinations rather than deviations in ancestry at one locus
or the other.
To determine the appropriate significance threshold for this scan, we performed
simulations. Using the F_2 ancestry data described in the previous paragraph, we simulated
genotypes at the focal locus based on each individual's genome-wide ancestry and performed the
χ^2 test as described above. We repeated this for every locus in the dataset (i.e. over 20,011
markers) and recorded the minimum p-value. We performed this procedure 500 times and took
the lower 5% quantile of minimum p-values across simulations. This value served as our false
positive rate (FPR) cutoff for analysis of the real data. We describe this approach in more detail
in Supporting Information 9. We initially explored an approach using permutations at the focal
locus to set the FPR but found that this approach was less conservative; these results are also
described in Supporting Information 9. Only one region in the genome (found on chromosome
13) passed the FPR threshold determined by permutation.
We also repeated this scan for interacting loci for all ancestry deserts shared between *X.*
birchmanni × *X. cortezi* and *X. birchmanni* × *X. malinche* populations. While we did not identify
any interacting loci that passed the 5% genome-wide FPR correction threshold for other regions,
we highlight two other cases where interactions are detected with minor parent deserts at a FPR
of 10% (Supporting Information 9). Simulations suggest that our power to detect interactions in
is relatively low (Supporting Information 9), leaving open the possibility that other minor parent
deserts interact with partner loci that we are unable to detect in the current data.
946
Pathway analysis of genes falling in a shared hybrid incompatibility
We wanted to ask whether genes in the identified chromosome 6 and chromosome 13
regions might be known to interact from past experimental, fusion, or co-expression data. To
evaluate this, we took advantage of the STRING database [59]. We used haltools [72] to liftover
coordinates from the *X. birchmanni* reference genome to the *X. maculatus* reference genome for
the shared ancestry desert on chromosome 6 and the interacting region on chromosome 13. This
allowed us to extract gene IDs for *X. maculatus* in this region, which is included in the STRING
database.
We input gene names from both chromosomes (chromosome 6 – 13 genes; chromosome
13 – 29 genes) into STRING using the protein batch search function. Two pairs of interacting
genes with supporting experimental, fusion, or co-expression data were identified. For one of
these pairs, both genes fell within the chromosome 6 region but for the other pair, one gene fell
within the chromosome 6 region and the other within the chromosome 13 region.
Given that our power to map interacting regions of shared minor parent deserts is
relatively low (Supporting Information 9), we wanted to perform a broad scan to ask whether
there was evidence for enrichment of genes that have many protein-protein interactions or dense
regulatory networks within shared minor parent ancestry deserts and islands. We thus performed
similar analyses for genes found in other shared minor parent ancestry deserts and islands. These
analyses included both STRING database searches similar to those described above, and
evaluation of enrichment of hub genes from co-expression analyses in minor parent deserts and
islands compared to matched null sets (Supporting Information 10).
968
Investigating other features associated with ancestry variation
Because theory predicts that many of the changes in genome-wide ancestry that occur in
hybrid populations happen rapidly after admixture [29,39], we were interested in investigating

correlations between minor parent ancestry and chromosome-level features. We evaluated
whether there were correlations between chromosome length, number of genes per chromosome,
and minor parent ancestry using a Spearman's rank correlation test.
We also evaluated minor parent ancestry as a function of structural differences detected
between the *X. birchmanni*, *X. malinche*, and *X. cortezi* assemblies. To identify chromosomal
inversions between species, we compared de-novo assemblies using MUMmer (mummer-
4.0.0beta2) [73]. After aligning the 24 largest linkage groups, which correspond to the 24
Xiphophorus chromosomes, we excluded alignments under 5 kb before proceeding with the
analysis. We focused our analysis on scanning for large inversions (> 100 kb). To identify the
putative inversion breakpoints, we used mummer's show-coords command. By aligning the *X.*
birchmanni [23] assembly to *X. cortezi* [46] and to *X. malinche* [23], and comparing to an
outgroup sequence (*X. maculatus;* [52]), we determined which inversions were derived in each
species. For simplicity, we assumed the inversion configuration detected in the reference
genomes was fixed at the species level (π in *X. birchmanni*, *X. malinche*, and *X. cortezi* is
<0.12%) but we note that some segregating inversions have been detected in *Xiphophorus* [23].
987
988 *Animal Ethics*
989   All animal use followed protocols that were approved by Stanford IACUC (Stanford
990 APLAC protocol number 33071). We are grateful to the Mexican federal government for
991 permission to collect samples (permit number: PFF/DGOPA-064/20).
992

Acknowledgements

      We thank Stepfanie Aguillon, Yaniv Brandvain, Erin Calfee, Yuki Haba, Priya Moorjani, Rachel Moran, Emilie Richards, Ken Thompson, Leslie Turner, and members of the Brandvain, Coop, Martin, Moorjani, Goldberg and Schumer labs for helpful discussion and/or feedback on earlier versions of this work. We also thank Baruc Zago-Mazzocco for field work support. We are grateful to the Mexican federal government for permission to collect samples. We thank Stanford University and the Stanford Research Computing Center for providing computational support for this project. This work was supported by a Knight-Hennessy Scholars fellowship and NSF GRFP 2019273798 to B. Moran, NRSA F32 GM135998 to B. Kim, a CEHG fellowship and NSF PRFB (2010950) to Q. Langdon, and a Hanna H. Gray fellowship and NIH 1R35GM133774 grant to M. Schumer

Figures

Figure 1. Description of focal species, study sites, and hybrid population ancestry structure. **A.** Phylogenetic relationships and contemporary hybridization events between focal species *X. birchmanni*, *X. malinche*, and *X. cortezi*. **B.** Distributions of *X. birchmanni* (*Xbir*), *X. malinche* (*Xmal*), and *X. cortezi* (*Xcor*), and their hybrids in Hidalgo and San Luis Potosí, Mexico. Blue shading shows the known range of *X. malinche*, red shading shows the known range of *X. birchmanni*, and green shading shows the known range of *X. cortezi*. **C**. Genome-wide admixture proportions of *X. birchmanni* × *X. cortezi* hybrids found in the Santa Cruz and Huextetitla replicate hybrid populations. **D**. Genome-wide admixture proportions of *X. birchmanni* × *X. malinche* hybrids found in the replicate populations at Tlatemaco (N=95), Aguazarca (N=51), and Acuapa (N=97).

1021
Figure 2. Relationship between minor parent ancestry, recombination rate and functional
basepairs in *X. birchmanni* × *X. cortezi* hybrid populations, as well as cross-population type
comparisons. **A.** Minor parent ancestry (*X. birchmanni* ancestry) in the Santa Cruz hybrid
population is higher in regions of the genome with higher recombination rates. Semi-transparent
points show results from individual 100 kb non-overlapping windows, black points and whiskers
show the mean ± two standard errors of the mean. After accounting for recombination rate by
averaging ancestry in 0.25 cM non-overlapping windows, we also observe a strong negative
correlation between minor parent ancestry and the number of linked coding (middle) and
conserved basepairs **(right)**. We find the same results when analyzing data from the Huextetitla
hybrid population and across a range of window sizes (Table S1, S4). Note that results for **A** are
shown in quintiles for visualization but statistical tests reported in the text were performed on
unbinned data. **B**. Correlations in local ancestry, plotted here in 1 Mb non-overlapping windows.
Cross-population ancestry correlations between Santa Cruz (or Huextetila – Table S10) and *X.*
birchmanni × *X. malinche* hybrid populations are significant in all comparisons except for the
majority *X. malinche* population Tlatemaco (far right; see main text & Supporting Information
6). Some of the ancestry covariance between *X. birchmanni* × *X. cortezi* and *X. birchmanni* × *X.*
malinche hybrid populations can be explained by shared features of genome architecture such as
the locations of coding and conserved basepairs and the local recombination rate, but substantial
covariance remains after accounting for these features (Table S12).

1041
1042
Figure 3. A. Shared minor parent deserts and islands are enriched compared to expectations by
chance between *X. birchmanni* × *X. cortezi* and *X. birchmanni* × *X. malinche* hybrid populations.
Results shown here indicate the number of shared minor parent deserts (or islands) between the
Santa Cruz *X. birchmanni* × *X. cortezi* hybrid population and each *X. birchmanni* × *X. malinche*
hybrid population (Acuapa, Aguazarca, and Tlatemaco). Black diamonds show the observed
number of shared minor parent deserts or islands and colored points show the results when the
data is jack-knife bootstrapped. In contrast, few shared minor parent deserts or islands are
expected by chance (results of permutations shown by gray points). The column labeled
Xmal/Xbir shows the number of shared deserts or islands between the Santa Cruz *X. birchmanni*
× *X. cortezi* hybrid population and any single *X. birchmanni* × *X. malinche* population. See
Supporting Information 8 and Fig. S12 for the results of alternative analysis approaches. **B.**
Minor parent ancestry in *X. birchmanni* × *X. cortezi* hybrid populations is depleted near 6
segregation distorters identified in artificial crosses between *X. birchmanni* × *X. malinche* (see
Results). Results plotted here are for the Santa Cruz hybrid population; gray lines show 500
replicates bootstrap resampling windows in the data, blue line indicates the mean across
simulations. **C.** One minor parent ancestry desert on chromosome 6 was detected across all *X.*
birchmanni × *X. cortezi* and *X. birchmanni* × *X. malinche* hybrid populations. Points represent
minor parent ancestry calculated in 10 kb non-overlapping windows divided by the proportion of
the genome derived from the minor parent species. Red – Tlatemaco (*malinche* × *birchmanni*),
Blue – Acuapa and Aguazarca (*birchmanni* × *malinche*), Green –Santa Cruz (*birchmanni* ×

cortezi). **D**. The shared minor parent ancestry desert on chromosome 6 coincides with a strong
segregation distorter in artificial crosses between *X. birchmanni* × *X. malinche*. Plotted here is
average ancestry at ancestry informative sites across 943 F_2 hybrids, where expected ancestry is
50% *X. birchmanni* and 50% *X. malinche*. The blue envelop indicates the 99% quantile of
variation in ancestry in this cross genome-wide. The dashed lines indicate the location of the
shared ancestry desert in hybrid populations plotted in **C**.

1069

Figure 4. Evidence for repeated hybrid incompatibilities between *X. birchmanni* × *X. malinche*
and *X. birchmanni* × *X. cortezi* hybrids. **A**. Interacting locus of the shared chromosome 6
ancestry desert identified in a genome-wide scan. This region was identified scanning for
deviations from expected two locus genotypes using a χ^2 test in F_2 hybrids, based on observed
genotypes at the shared chromosome 6 desert and genotypes at other loci throughout the genome.
Genome-wide significance threshold (FPR 5%; blue line) was determined using permutations,
see Methods for details. Inset shows STRING network with experimentally verified interactions
between *ndufa13* (contained in the chromosome 6 region) and *ndufs5* (contained in the
chromosome 13 region). **B**. Ratio of expected to observed two-locus genotype combinations in
F_2 hybrids between *X. birchmanni* × *X. malinche* at the shared chromosome 6 desert and
associated chromosome 13 region. Capital letters indicate genotype at chromosome 6 and
lowercase letters indicate genotype at the chromosome 13 region. For visualization, certain
genotype combinations are collapsed (i.e. homozygous parental – BBbb & MMmm;
heterozygous locus 1, homozygous locus 2 – MBbb & MBmm). Note that several genotype
combinations were expected to be found at low frequency due to low *X. birchmanni* ancestry on
chromosome 13 (MMbb, MBbb, BBbb). **C.** A second interaction between the minor parent
desert on chromosome 9 and a region on chromosome 4 is detected at a relaxed FPR of 10%. **D**.
This minor parent desert is shared between *X. birchmanni* × *X. cortezi* hybrid populations and the
Acuapa *X. birchmanni* × *X. malinche* hybrid population and occurs on chromosome 9 at 12.1
Mb. Points represent minor parent ancestry calculated in 10 kb non-overlapping windows
divided by the proportion of the genome derived from the minor parent species. Red – Tlatemaco
(*malinche* × *birchmanni*), Blue – Acuapa and Aguazarca (*birchmanni* × *malinche*), Green –Santa
Cruz (*birchmanni* × *cortezi*).

1093

References

1. Rhie A, McCarthy SA, Fedrigo O, Damas J, Formenti G, Koren S, et al. Towards complete and error-free genome assemblies of all vertebrate species. Nature. 2021;592:737-746. doi: 10.1038/s41586-021-03451-0

2. Genereux DP, Serres A, Armstrong J, Johnson J, Marinescu VD, Murén E, et al. A comparative genomics multitool for scientific discovery and conservation. Nature. 2020;587: 240–245. doi:10.1038/s41586-020-2876-6

3. The Global Invertebrate Genomics Alliance (GIGA): Developing Community Resources to Study Diverse Invertebrate Genomes. J Hered. 2014;105: 1–18. doi:10.1093/jhered/est084

4. Abbott R, Albach D, Ansell S, Arntzen JW, Baird SJE, Bierne N, et al. Hybridization and speciation. Journal of Evolutionary Biology. 2013;26: 229–246. doi:10.1111/j.1420-9101.2012.02599.x

5. Mallet J. Hybridization as an invasion of the genome. Trends Ecol Evol. 2005;20: 229–237. doi:10.1016/j.tree.2005.02.010

6. Moran BM, Payne C, Langdon Q, Powell DL, Brandvain Y, Schumer M. The genomic consequences of hybridization. Wittkopp PJ, editor. eLife. 2021;10: e69016. doi:10.7554/eLife.69016

7. Turissini DA, Matute DR. Fine scale mapping of genomic introgressions within the Drosophila yakuba clade. PLOS Genetics. 2017;13: e1006971. doi:10.1371/journal.pgen.1006971

8. Brandvain Y, Kenney AM, Flagel L, Coop G, Sweigart AL. Speciation and Introgression between Mimulus nasutus and Mimulus guttatus. PLOS Genetics. 2014;10: e1004410. doi:10.1371/journal.pgen.1004410

9. D'Angiolo M, De Chiara M, Yue J-X, Irizar A, Stenberg S, Persson K, et al. A yeast living ancestor reveals the origin of genomic introgressions. Nature. 2020;587: 420–425. doi:10.1038/s41586-020-2889-1

10. Rieseberg LH, Raymond O, Rosenthal DM, Lai Z, Livingstone K, Nakazato T, et al. Major ecological transitions in wild sunflowers facilitated by hybridization. Science. 2003;301: 1211–1216. doi:10.1126/science.1086949

11. Cui R, Schumer M, Kruesi K, Walter R, Andolfatto P, Rosenthal G. Phylogenomics reveals extensive reticulate evolution in Xiphophorus fishes. Evolution. 2013;67: 2166–2179.

12. Sankararaman S, Mallick S, Patterson N, Reich D. The Combined Landscape of Denisovan and Neanderthal Ancestry in Present-Day Humans. Current Biology. 2016;26: 1241–1247. doi:10.1016/j.cub.2016.03.037

1129   13.   Heliconius Genome C. Butterfly genome reveals promiscuous exchange of mimicry
1130   adaptations among species. Nature. 2012;487: 94–8.

1131   14.   Trier CN, Hermansen JS, Sætre G-P, Bailey RI. Evidence for Mito-Nuclear and Sex-Linked
1132   Reproductive Barriers between the Hybrid Italian Sparrow and Its Parent Species. PLOS
1133   Genetics. 2014;10: e1004075. doi:10.1371/journal.pgen.1004075

1134   15.   Orr HA. The population genetics of speciation: the evolution of hybrid incompatibilities.
1135   Genetics. 1995;139: 1805–1813.

1136   16.   Arnegard ME, McGee MD, Matthews B, Marchinko KB, Conte GL, Kabir S, et al.
1137   Genetics of ecological divergence during speciation. Nature. 2014;511: 307–311.
1138   doi:10.1038/nature13301

1139   17.   Svedin N, Wiley C, Veen T, Gustafsson L, Qvarnström A. Natural and sexual selection
1140   against hybrid flycatchers. Proceedings of the Royal Society B: Biological Sciences.
1141   2008;275: 735–744. doi:10.1098/rspb.2007.0967

1142   18.   Christie K, Strauss SY. Along the speciation continuum: Quantifying intrinsic and extrinsic
1143   isolating barriers across five million years of evolutionary divergence in California
1144   jewelflowers. Evolution. 2018;72: 1063–1079. doi:10.1111/evo.13477

1145   19.   Juric I, Aeschbacher S, Coop G. The Strength of Selection against Neanderthal
1146   Introgression. PLOS Genetics. 2016;12: e1006340. doi:10.1371/journal.pgen.1006340

1147   20.   Schumer M, Cui R, Powell DL, Dresner R, Rosenthal GG, Andolfatto P. High-resolution
1148   mapping reveals hundreds of genetic incompatibilities in hybridizing fish species. McVean
1149   G, editor. eLife. 2014;3: e02535. doi:10.7554/eLife.02535

1150   21.   Chae E, Bomblies K, Kim S-T, Karelina D, Zaidem M, Ossowski S, et al. Species-wide
1151   genetic incompatibility analysis identifies immune genes as hot spots of deleterious
1152   epistasis. Cell. 2014;159: 1341–1351. doi:10.1016/j.cell.2014.10.049

1153   22.   Larson EL, Vanderpool D, Sarver BAJ, Callahan C, Keeble S, Provencio LL, et al. The
1154   Evolution of Polymorphic Hybrid Incompatibilities in House Mice. Genetics. 2018;209:
1155   845–859. doi:10.1534/genetics.118.300840

1156   23.   Powell DL, García-Olazábal M, Keegan M, Reilly P, Du K, Díaz-Loyo AP, et al. Natural
1157   hybridization reveals incompatible alleles that cause melanoma in swordtail fish. Science.
1158   2020;368: 731–736. doi:10.1126/science.aba5216

1159   24.   Zuellig MP, Sweigart AL. A two-locus hybrid incompatibility is widespread, polymorphic,
1160   and active in natural populations of Mimulus*. Evolution. 2018;72: 2394–2405.
1161   doi:10.1111/evo.13596

1162   25.   Barton NH. The role of hybridization in evolution. Molecular Ecology. 2001;10: 551–568.
1163   doi:https://doi.org/10.1046/j.1365-294x.2001.01216.x

1164   26.   Slatkin M, Lande R. Segregation variance after hybridization of isolated populations. Genet
1165   Res. 1994;64: 51–56. doi:10.1017/s0016672300032547

1166   27.   Thompson KA. Experimental Hybridization Studies Suggest That Pleiotropic Alleles
1167   Commonly Underlie Adaptive Divergence between Natural Populations. The American
1168   Naturalist. 2020;196: E16–E22. doi:10.1086/708722

1169   28.   Simon A, Bierne N, Welch JJ. Coadapted genomes and selection on hybrids: Fisher's
1170   geometric model explains a variety of empirical patterns. Evolution Letters. 2018;2: 472–
1171   498. doi:10.1002/evl3.66

1172   29.   Harris K, Nielsen R. The Genetic Cost of Neanderthal Introgression. Genetics. 2016;203:
1173   881–891. doi:10.1534/genetics.116.186890

1174   30.   Presgraves DC. Sex chromosomes and speciation in Drosophila. Trends in Genetics.
1175   2008;24: 336–343. doi:10.1016/j.tig.2008.04.007

1176   31.   Maheshwari S, Barbash DA. The Genetics of Hybrid Incompatibilities. Annual Review of
1177   Genetics. 2011;45: 331–355. doi:10.1146/annurev-genet-110410-132514

1178   32.   Qvarnström A, Bailey RI. Speciation through evolution of sex-linked genes. Heredity.
1179   2009;102: 4–15. doi:10.1038/hdy.2008.93

1180   33.   Trier CN, Hermansen JS, Sætre G-P, Bailey RI. Evidence for Mito-Nuclear and Sex-Linked
1181   Reproductive Barriers between the Hybrid Italian Sparrow and Its Parent Species. PLoS
1182   Genet. 2014;10. doi:10.1371/journal.pgen.1004075

1183   34.   Burton RS, Barreto FS. A disproportionate role for mtDNA in Dobzhansky–Muller
1184   incompatibilities? Molecular Ecology. 2012;21: 4942–4957. doi:10.1111/mec.12006

1185   35.   Schumer M, Xu C, Powell DL, Durvasula A, Skov L, Holland C, et al. Natural selection
1186   interacts with recombination to shape the evolution of hybrid genomes. Science. 2018;360:
1187   656. doi:10.1126/science.aar3684

1188   36.   Martin SH, Davey JW, Salazar C, Jiggins CD. Recombination rate variation shapes barriers
1189   to introgression across butterfly genomes. PLOS Biology. 2019;17: e2006288.
1190   doi:10.1371/journal.pbio.2006288

1191   37.   Nachman MW, Payseur BA. Recombination rate variation and speciation: theoretical
1192   predictions and empirical results from rabbits and mice. Philos Trans R Soc Lond B Biol
1193   Sci. 2012;367: 409–421. doi:10.1098/rstb.2011.0249

1194   38.   Sankararaman S, Mallick S, Dannemann M, Prüfer K, Kelso J, Pääbo S, et al. The genomic
1195   landscape of Neanderthal ancestry in present-day humans. Nature. 2014;507: 354–357.
1196   doi:10.1038/nature12961

39. Veller C, Edelman NB, Muralidhar P, Nowak MA. Recombination, variance in genetic
relatedness, and selection against introgressed DNA. bioRxiv. 2019; 846147.
doi:10.1101/846147

40. Calfee E, Gates D, Lorant A, Perkins MT, Coop G, Ross-Ibarra J. Selective sorting of
ancestral introgression in maize and teosinte along an elevational cline. PLoS Genet.
2021;17:e1009810. doi: 10.1371/journal.pgen.1009810

41. Vilgalys TP, Fogel AS, Mututua RS, Warutere JK, Siodi L, Anderson JA, et al. Selection
against admixture and gene regulatory divergence in a long-term primate field study. 2021
Aug p. 2021.08.19.456711. doi:10.1101/2021.08.19.456711

42. Baker Z, Schumer M, Haba Y, Bashkirova L, Holland C, Rosenthal GG, et al. Repeated
losses of PRDM9-directed recombination despite the conservation of PRDM9 across
vertebrates. In: eLife [Internet]. 6 Jun 2017 [cited 23 Jul 2019]. doi:10.7554/eLife.24133

43. Singhal S, Leffler EM, Sannareddy K, Turner I, Venn O, Hooper DM, et al. Stable
recombination hotspots in birds. Science. 2015;350: 928–932. doi:10.1126/science.aad0843

44. Wang RJ, Ané C, Payseur BA. The Evolution of Hybrid Incompatibilities Along a
Phylogeny. Evolution. 2013;67: 2905–2922. doi:https://doi.org/10.1111/evo.12173

45. Yang C-H, Scarpino SV. Reproductive barriers as a byproduct of gene network evolution.
bioRxiv. 2020; 2020.06.12.147322. doi:10.1101/2020.06.12.147322

46. Powell DL, Moran BM, Kim BY, Banerjee SM, Aguillon SM, Fascinetto-Zago P, et al.
Two new hybrid populations expand the swordtail hybridization model system. Evolution.
2021;75: 2524–2539. doi:10.1111/evo.14337

47. Fisher HS, Wong BBM, Rosenthal GG. Alteration of the chemical environment disrupts
communication in a freshwater fish. Proc R Soc London Ser B. 2006;273: 1187–1193.
doi:10.1098/rspb.2005.3406

48. Schumer M, Powell DL, Delclós PJ, Squire M, Cui R, Andolfatto P, et al. Assortative
mating and persistent reproductive isolation in hybrids. Proc Natl Acad Sci USA. 2017;114:
10936. doi:10.1073/pnas.1711238114

49. Culumber ZW, Fisher HS, Tobler M, Mateos M, Barber PH, Sorenson MD, et al.
Replicated hybrid zones of Xiphophorus swordtails along an elevational gradient.
Molecular Ecology. 2011;20: 342–356. doi:10.1111/j.1365-294X.2010.04949.x

50. Schumer M, Brandvain Y. Determining epistatic selection in admixed populations.
Molecular Ecology. 2016;25: 2577–2591. doi:10.1111/mec.13641

51. Matute DR, Comeault AA, Earley E, Serrato-Capuchina A, Peede D, Monroy-Eklund A, et
al. Rapid and Predictable Evolution of Admixed Populations Between Two Drosophila
Species Pairs. Genetics. 2019 [cited 20 Apr 2020]. doi:10.1534/genetics.119.302685

52.   Schartl M, Walter RB, Shen Y, Garcia T, Catchen J, Amores A, et al. The genome of the
platyfish, Xiphophorus maculatus, provides insights into evolutionary adaptation and
several complex traits. Nature Genetics. 2013;45: 567.

53.   Noor MAF, Grams KL, Bertucci LA, Reiland J. Chromosomal inversions and the
reproductive isolation of species. PNAS. 2001;98: 12084–12088.
doi:10.1073/pnas.221274498

54.   Kirkpatrick M, Barton N. Chromosome Inversions, Local Adaptation and Speciation.
Genetics. 2006;173: 419–434. doi:10.1534/genetics.105.047985

55.   Wellenreuther M, Bernatchez L. Eco-Evolutionary Genomics of Chromosomal Inversions.
Trends in Ecology & Evolution. 2018;33: 427–440. doi:10.1016/j.tree.2018.04.002

56.   Edelman NB, Frandsen PB, Miyagi M, Clavijo B, Davey J, Dikow RB, et al. Genomic
architecture and introgression shape a butterfly radiation. Science. 2019;366: 594–599.
doi:10.1126/science.aaw2090

57.   Todesco M, Owens GL, Bercovich N, Légaré J-S, Soudi S, Burge DO, et al. Massive
haplotypes underlie ecotypic differentiation in sunflowers. Nature. 2020;584: 602–607.
doi:10.1038/s41586-020-2467-6

58.   Powell DL, Payne C, Banerjee SM, Keegan M, Bashkirova E, Cui R, et al. The Genetic
Architecture of Variation in the Sexually Selected Sword Ornament and Its Evolution in
Hybrid Populations. Current Biology. 2021 [cited 28 Jan 2021].
doi:10.1016/j.cub.2020.12.049

59.   Szklarczyk D, Morris JH, Cook H, Kuhn M, Wyder S, Simonovic M, et al. The STRING
database in 2017: quality-controlled protein–protein association networks, made broadly
accessible. Nucleic Acids Res. 2017;45: D362–D368. doi:10.1093/nar/gkw937

60.   Moran BM, Payne CY, Powell DL, Iverson ENK, Banerjee SM, Langdon QK, et al. A
Lethal Genetic Incompatibility between Naturally Hybridizing Species in Mitochondrial
Complex I. 2021 Jul p. 2021.07.13.452279. doi:10.1101/2021.07.13.452279

61.   Yang Z. PAML 4: Phylogenetic Analysis by Maximum Likelihood. Mol Biol Evol.
2007;24: 1586–1591. doi:10.1093/molbev/msm088

62.   Blount ZD, Lenski RE, Losos JB. Contingency and determinism in evolution: Replaying
life's tape. Science. 2018;362. doi:10.1126/science.aam5979

63.   Chaturvedi S, Lucas LK, Buerkle CA, Fordyce JA, Forister ML, Nice CC, et al. Recent
hybrids recapitulate ancient hybrid outcomes. Nat Commun. 2020;11: 2179.
doi:10.1038/s41467-020-15641-x

64.   Kim BY, Huber CD, Lohmueller K. Deleterious variation shapes the genomic landscape of
introgression. PLoS Genet. 2018;14:e1007741. doi: 10.1371/journal.pgen.1007741

65.   Lu Y, Sandoval A, Voss S, Lai Z, Kneitz S, Boswell W, et al. Oncogenic allelic interaction
1268   in Xiphophorus highlights hybrid incompatibility. PNAS. 2020;117: 29786–29794.
1269   doi:10.1073/pnas.2010133117

1270   66.   Alcázar R, von Reth M, Bautor J, Chae E, Weigel D, Koornneef M, et al. Analysis of a
1271   plant complex resistance gene locus underlying immune-related hybrid incompatibility and
1272   its occurrence in nature. PLoS Genet. 2014;10: e1004848.
1273   doi:10.1371/journal.pgen.1004848

1274   67.   Schumer M, Powell DL, Corbett-Detig R. Versatile simulations of admixture and accurate
1275   local ancestry inference with mixnmatch and ancestryinfer. Mol Ecol Resour. 2020;20:
1276   1141–1151. doi:10.1111/1755-0998.13175

1277   68.   Hinch AG, Tandon A, Patterson N, Song Y, Rohland N, Palmer CD, et al. The landscape of
1278   recombination in African Americans. Nature. 2011;476: 170–175. doi:10.1038/nature10336

1279   69.   Gravel S. Population Genetics Models of Local Ancestry. Genetics. 2012;191: 607.
1280   doi:10.1534/genetics.112.139808

1281   70.   Powell DL, Payne C, Banerjee SM, Keegan M, Bashkirova E, Cui R, et al. The Genetic
1282   Architecture of Variation in the Sexually Selected Sword Ornament and Its Evolution in
1283   Hybrid Populations. Current Biology. 2021;31: 923-935.e11.
1284   doi:10.1016/j.cub.2020.12.049

1285   71.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic
1286   features. Bioinformatics. 2010;26: 841–842. doi:10.1093/bioinformatics/btq033

1287   72.   Hickey G, Paten B, Earl D, Zerbino D, Haussler D. HAL: a hierarchical format for storing
1288   and analyzing multiple genome alignments. Bioinformatics. 2013;29: 1341–1342.
1289   doi:10.1093/bioinformatics/btt128

1290   73.   Marçais G, Delcher AL, Phillippy AM, Coston R, Salzberg SL, Zimin A. MUMmer4: A
1291   fast and versatile genome alignment system. PLOS Computational Biology. 2018;14:
1292   e1005944. doi:10.1371/journal.pcbi.1005944

1293