Defining the Immune Responses for SARS-CoV-2-Human Macrophage Interactions

Mai M. Abdelmoaty[1,2], Pravin Yeapuri[1], Jatin Machhi[3], Katherine E. Olson[3], Farah Shahjin[3],

You Zhou[4], Liang Jingjing[3], Kabita Pandey[3], Arpan Acharya[3], Siddappa N. Byrareddy[3],

R. Lee Mosley[3] and Howard E. Gendelman[3*]

[1]Department of Pharmaceutical Sciences, College of Pharmacy, University of Nebraska Medical

Center, NE 68198, USA

[2]Therapeutic Chemistry Department, Pharmaceutical and Drug Industries Research Division,

National Research Centre, Giza, Egypt

[3]Department of Pharmacology and Experimental Neuroscience, College of Medicine, University

of Nebraska Medical Center, NE 68198, USA

[4]Center for Biotechnology, University of Nebraska-Lincoln, Lincoln, NE, USA
14

Corresponding author: Howard E. Gendelman, Department of Pharmacology and Experimental

Neuroscience, University of Nebraska Medical Center, Omaha, NE 68198-5800; phone 402-559-

8920; fax 402-559-3744; email hegendel@unmc.edu

Abstract

Host innate immune response follows severe acute respiratory syndrome coronavirus 2 (SARS-

CoV-2) infection, and it is the driver of the acute respiratory distress syndrome (ARDS) amongst

other inflammatory end-organ morbidities. Such life-threatening coronavirus disease 2019

(COVID-19) is heralded by virus-induced activation of mononuclear phagocytes (MPs;

monocytes, macrophages, and dendritic cells). MPs play substantial roles in aberrant immune

secretory activities affecting profound systemic inflammation and end organ malfunctions. All

follow an abortive viral infection. To elucidate SARS-CoV-2-MP interactions we investigated

transcriptomic and proteomic profiles of human monocyte-derived macrophages. While

expression of the SARS-CoV-2 receptor, the angiotensin-converting enzyme 2, paralleled

monocyte-macrophage differentiation it failed to affect productive viral infection. In contrast,

simple macrophage viral exposure led to robust pro-inflammatory cytokine and chemokine

expression but attenuated type I interferon (IFN) activity. Both paralleled dysregulation of innate

immune signaling pathways specifically those linked to IFN. We conclude that the SARS-CoV-2-

infected host mounts a robust innate immune response characterized by a pro-inflammatory storm

heralding consequent end-organ tissue damage.

Keywords: macrophages, SARS-CoV-2, cytokine storm, interferon, end-organ disease,

inflammation, transcriptomics, proteomics

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), the causative agent of coronavirus disease 2019 (COVID-19), is an enveloped positive-stranded RNA virus belonging to the *Coronaviridae* family, *Betacoronaviruses* genus (Lu et al., 2020). COVID-19 has posed an unprecedented global threat to public health and in March of 2020, it was declared a pandemic by the World Health Organization (WHO) (World Health Organization, 2020). COVID-19 ranges from asymptomatic infection to mild pneumonia and, in its most severe form, progression to acute respiratory distress syndrome (ARDS). Such pulmonary compromise is associated with dyspnea and hypoxia that can progress to severely compromised lung dysfunction and multiorgan system failure and death (Gavriatopoulou et al., 2020). Disease mortality is linked to cytokine storm syndrome (CSS), heralded by innate immune activation with secretion of excessive pro-inflammatory cytokines (Ruan et al., 2020). Indeed, nearly 15% of reported COVID-19 disease cases progress to ARDS defined by widespread inflammatory-associated lung tissue damage and multiorgan failure (Guan et al., 2020) involving heart, liver, gastrointestinal tract, kidney, and brain (Salamanna et al., 2020). Viral persistence in the face of such end-organ disease is linked to cell expression of angiotensin-converting enzyme 2 (ACE2), the molecule that SARS-CoV-2 utilizes for receptor-mediated cell entry (Wan et al., 2020).

Mononuclear phagocytes (MPs; monocytes, macrophages, and dendritic cells) are the governors of innate immunity serving to contain microbial infection (Lugo-Villarino et al., 2019). Immediately following viral exposure, the process of intracellular microbial removal is initiated through recognition of viral pathogen-associated molecular patterns (PAMPs) by pattern recognition receptors (PRRs). This includes, but is not limited to, cytosolic retinoic acid-inducible

gene I (RIG-I)-like receptors (RLRs) and extracellular and endosomal toll-like receptors (TLRs).

During the initiation of virus-cell interactions, secretion of pro-inflammatory cytokines and

chemokines initiate intracellular killing, antigen presentation, and mobilization of adaptive

immunity. Factors that are engaged include, but are not limited to, interleukin-6 (IL-6), IL-1, tumor

necrosis factor-alpha (TNF-α), and C-X-C motif chemokine ligand 10 (CXCL10) (Nikitina et al.,

2018). Type I interferon (IFN) α and β responses occur in tandem to control both viral replication

and dissemination (Swiecki and Colonna, 2011). During infection with SARS-CoV-2, CSS

contributes to COVID-19-associated multiorgan failure (Huang et al., 2020; Xiong et al., 2020).

In particular, infiltrating inflammatory MPs in lungs, heart, kidney, spleen, and lymph nodes are

seen in post-mortem tissues of COVID-19-infected patients (Diao et al., 2021; Feng et al., 2020;

Merad and Martin, 2020; Xu et al., 2020). Despite the critical role of MPs to clear the infection,

these same cells underlie the pathobiology of COVID-19.

Mounting evidence confirms abortive MP infection by SARS-CoV-2 (Yang et al., 2020; Zheng et

al., 2021). Nonetheless, such viral cell engagements are sufficient to induce activation and pro-

inflammatory cytokine secretory responses. The nature of virus-MP interactions and their role in

persistent viral infection and associated tissue pathologies remains enigmatic. To such ends, we

pursued transcriptomic and proteomic analyses of immune cells following virus infection to

identify modulations of host immune responses. We employed MPs to assess immune responses

following SARS-CoV-2 cell engagements for cell activation using human immune response arrays

and mass spectrometry-based label-free proteomic quantification methods. These techniques serve

to define virus-induced innate immune responses linked to antiviral immunity. Based on these

tests, we have uncovered the dysregulation of a spectrum of viral-induced responses related to IFN

signaling pathways, complement activation, and linked adaptive immune responses. All provide

unique insight into how the inflammatory response occurs as a consequence of SARS-CoV-2

infection. Virion persistence for up to 14 days after viral exposure despite abortive infection

underlies the persistence of immune activation. Most importantly, the data provides a signature

for virus-induced disease pathobiology, including COVID-19-associated CSS and multiorgan

dysfunction. All affect the most severe disease morbidities and mortalities seen as a consequence

of viral exposure, transmission, and dissemination.

Materials and methods

Isolation and cultivation of human monocytes

Human monocytes were obtained by leukapheresis from hepatitis B and HIV-1/2 seronegative

donors and purified by counter-current centrifugal elutriation (Faradji et al., 1994). Monocytes

were seeded in 6-well plates (3×10^6 cells/well) in Dulbecco's Modified Eagle's Media (DMEM)

containing 4.5 g/l glucose, L-glutamine, and sodium pyruvate, and supplemented with 10% heat-

inactivated human serum, 50 μg/ml gentamicin, 10 μg/ml ciprofloxacin, and 1000 U/ml human

macrophage colony-stimulating factor (M-CSF) to facilitate differentiation of monocytes into

monocyte-derived macrophages (MDMs). Cells were incubated at 37°C with 5% CO_2 and the

culture medium was half-exchanged with fresh medium every other day (Gendelman et al., 1992).

Flow cytometry assays

Human monocytes were evaluated by flow cytometry for levels of SARS-CoV-2 cell entry

receptor ACE2 and monocyte-macrophage phenotypic surface markers, CD14 and CD16, during

macrophage differentiation with or without captopril (Sigma-Aldrich, C4042). Captopril was

added to the culture medium to increase ACE2 cell expression. On days 0, 1, 3, 5, and 7 during

differentiation, monocytes-macrophages were stained with fluorescently-conjugated antibodies to

detect human ACE2 (APC, LSBio, LS-C275129, polyclonal), CD14 (Alexa Fluor 488,

eBioscience, clone 61D3), and CD16 (PE, eBioscience, eBioCB16, clone CB16), and with isotype-

matched antibodies serving as negative controls. Stained cells were examined with an LSR II flow

cytometer (BD Biosciences) and analyzed using BD FACSDiva software.

SARS-CoV-2 infection

Experiments involving SARS-CoV-2 were performed in the University of Nebraska Medical

Center (UNMC) biosafety level 3 (BSL-3) core facility and approved by UNMC Institutional

Biosafety Committee (IBC) (protocol number 20-05-027-BL3). The SARS-CoV-2 strain used in

this study was isolate USA-WI1/2020 (BEI, NR-52384). The virus was passaged on Vero.STAT1

knockout (KO) cells (ATCC, CCL-81-VHG) and titer was determined by plaque assay in Vero E6

cells (ATCC, CRL-1586) (Mendoza et al., 2020). On day 5 following cell cultivation, MDMs were

exposed to the WI1/2020 viral strain at a multiplicity of infection (MOI) of 0.01. Exposure was

maintained with or without captopril, and the virus-cell mixtures incubated at 37°C in 5% CO_2

with shaking at 15 minute intervals for 1 hour. This was followed by incubation with the virus for

an additional 3 hours. At the termination of viral cell incubation, the virus inoculum was removed,

and cells were washed 3 times with phosphate-buffered saline (PBS). Mock-challenged cells were

treated with a culture medium alone. Culture supernatants were collected at defined time points

then used for viral quantitative reverse transcription-polymerase chain reaction (RT-qPCR) and

type I IFN activity assays. Vero.STAT1 KO cells were maintained for study based on their high

susceptibility to virus infection due to lack of Signal Transducer and Activator of Transcription 1

(STAT1) protein required for cellular antiviral responses (Durbin et al., 1996). Thus, the SARS-

CoV-2 replication kinetics in MDMs were compared against viral infection in Vero.STAT1 KO

cells that were monitored over 5 days post-viral exposure.

RT-qPCR assay

Culture supernatants were collected on days 1, 3, 5, 7, and 10 following viral exposure. Viral

RNA was extracted using QIAamp Viral RNA Mini Kit (Qiagen, 52906). SARS-CoV-2 genome

equivalents were quantified in culture supernatant by RT-qPCR using 2019-nCoV CDC probe and

Primer Kit for SARS-CoV-2 (Biosearch Technologies, KIT-nCoV-PP1-1000). Forward primer: 5'

GACCCCAAAATCAGCGAAAT 3', Reverse primer: 5' TCTGGTTACTGCCAGTTGAATCTG

3', and Probe: 5' FAM-ACCCCGCATTACGTTTGGTGGACC-BHQ-1 3'. RT-qPCR was

performed using Taqman Fast Virus 1-step Master Mix (Thermo Fisher Scientific, 4444434) in

StepOne Plus real-time PCR thermocycler (Applied Biosystems) using the following cycling

conditions: 50°C for 10 minutes, 95°C for 3 minutes, and 40 cycles of 95°C for 15 seconds,

followed by 60°C for 1 minute. The SARS-CoV-2 genome equivalent copies were calculated using

control RNA from heat-inactivated SARS-CoV-2 USA-WA1/2020 (BEI, NR-52347).

Transmission electron microscopy (TEM)

For negative staining analysis of viral particles used for cell infection experiment, purified SARS-

CoV-2 viruses were de-activated and fixed in 2% glutaraldehyde and 2% paraformaldehyde (PFA)

in 0.1 M Sorenson's phosphate buffer for 1 hour at room temperature. Briefly, for negative

staining, a drop of the virus sample in a fixative solution was placed onto a formvar and carbon-

coated grid for 30-40 seconds. After removing excess sample solution with filter paper, the grid

was placed with sample side down on a drop of 1% phosphotungstic acid (PTA) in water, stained

for 30 seconds, then excess PTA solution was blotted with filter paper. Stained samples were

examined and imaged using a Hitachi H7500 TEM (Hitachi High-Tech GLOBAL) and a bottom-

mount AMT camera (AMT Imaging). For ultrastructural analysis, mock and SARS-CoV-2-

challenged MDMs sampled at 1, 3, and 5 days post-viral inoculation were washed 2 times with

PBS and fixed in a solution of 2% glutaraldehyde and 2% PFA in 0.1 M Sorenson's phosphate

buffer for 24 hours at 4°C which have been washed 3 times with PBS to clear excess fixative

solution. TEM analysis was performed as previously described (Mukadam et al., 2020) in samples

post-fixed in a 1% aqueous solution of osmium tetroxide for 30 minutes that were dehydrated in

50, 70, 90, 95, and 100% graded ethanol. Spurr's resin was used as embedding medium after

solvent transition with ethanol and Spurr's resin (50:50 ethanol:resin, followed by twice immersion

in 100% Spurr's resin for 2-3 hours for each solution), and embedded samples were cured at 60 -

65°C for 24 hours. Ultrathin sections (100 nm) were cut with Leica UC7 ultramicrotome, placed

on 200-mesh copper grids, followed by staining with 2% uranyl acetate and Reynold's lead citrate,

and examined with a Hitachi H7500 TEM at 80 kV. Images were acquired digitally with an AMT

digital imaging system.

Transcriptomic analyses

SARS-CoV-2-challenged MDMs were collected on days 1, 3, and 5 post-viral exposure. Total

RNA was isolated using RNeasy Mini Kit (Qiagen, 74104), and cDNA was generated utilizing

RevertAid First Strand cDNA synthesis kit (Thermo Fisher Scientific, K1622) followed by

amplification and quantification using RT^2 Profiler Human Innate and Adaptive Immune Response

96-well Array (Qiagen, 330231) with RT^2 SYBR Green ROX qPCR Mastermix (Qiagen, 330523).

The qPCR cycling conditions were 95°C for 10 minutes for 1 cycle, followed by 40 cycles of 95°C

for 15 seconds and 60°C for 1 minute using Eppendorf Mastercycler ep realplex 2S. Fold changes

were determined via Qiagen's RT^2 Profiler analysis software (version 3.5). Ingenuity Pathway

Analysis (IPA) (Qiagen) was used to identify the pathways and networks affected post-viral

exposure. Functional and pathway enrichment analyses of screened genes in SARS-CoV-2-

challenged MDMs were compared to mock-challenged MDMs at different time intervals following

viral exposure. Gene ontology (GO) annotation was conducted using the GO Resource database

(http://geneontology.org). The Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway

enrichment analysis was conducted utilizing DAVID (http://david.abcc.ncfcrf.gov/) (Huang da et

al., 2009), an online tool providing a comprehensive set of functional annotations for

understanding biological meaning in large list of genes. The Reactome gene set enrichment

analysis (GSEA) was conducted using ReactomeFIViz (https://reactome.org/tools/reactome-fiviz)

(Wu et al., 2014), a Cytoscape application for pathway and network-based data analysis. The

Search Tool for the Retrieval of Interacting Genes/Proteins (STRING) local network cluster

enrichment analysis was conducted using STRING database (http://string-db.org), which provides

a critical assessment and integration of protein-protein interaction (PPI), including direct

(physical) and indirect (functional) associations in a given organism (Szklarczyk et al., 2019).

Measures of IFN activity

MDBK cells (ATCC, CCL-22) were cultured in Eagle's Minimum Essential Medium (EMEM)

containing 10% heat-inactivated fetal bovine serum (FBS) and 50 µg/ml gentamicin (Rubinstein

et al., 1981). Vesicular stomatitis virus (VSV Indiana laboratory strain) (V-520-001-522, ATCC,

VR-1238) was passaged on Vero cells (ATCC, CCL-81), and viral titer was determined using the

plaque assay in Vero cells. MDBK cells bind and respond to human type I IFNs, α and β, but not

IFN-gamma (IFN-γ) (Rubinstein et al., 1981). Culture supernatants were collected from virus-

exposed MDMs on days 1, 3, and 5 following infection with SARS-CoV-2 at MOI of 0.01, then

assessed for IFN-α/β activity by protection against the VSV-induced cytopathicity measured in

MDBK cells (Rubinstein et al., 1981). Polyinosinic-polycytidylic acid (poly(I:C); Sigma-Aldrich,

P9582) served as a positive control for IFN induction via TLR3 engagement (Field et al., 1967).

Recombinant human IFN-α (PBL Assay Science, 11200-2) was used as assay standard. We also

investigated IFN activity upon the challenge of Teflon flask-suspended monocytes with SARS-

CoV-2 (MOI=0.01), before and after treatment of cells with 100 µg/ml poly(I:C).

Western blot analysis

At days 1, 3, and 5 after infection, MDMs were collected, and total protein was extracted using

lysis buffer containing 0.1% SDS, 100 mM Tris-HCL, 150 mM NaCl, 1 mM EDTA, 1% Triton

X-100, supplemented with protease and phosphatase inhibitor cocktail (Sigma-Aldrich, PPC1010).

Protein concentration was determined utilizing Pierce 660 Protein Assay kit (Thermo Fisher

Scientific, 22662) with ionic detergent compatibility reagent (Thermo Fisher Scientific, 22663)

following the manufacturer's instructions. Protein lysates (25 µg) were resolved by SDS-PAGE

and transferred to Immobilon-P PVDF membrane (Sigma-Aldrich, IPVH00010). Membranes were

blocked in 5% nonfat milk in TBST buffer at room temperature for 1 hour, followed by incubation

with primary antibodies to IFN-α (1:500, Thermo Fisher Scientific, MA5-37518), IFN-β (1:1000,

Abcam, ab85803), and β-actin (1:3000, Sigma-Aldrich, A3854) at 4°C overnight, followed by 1

hour incubation in 3% nonfat milk in Tris-buffered saline with 0.1% Tween 20 detergent

(TBST) buffer with horseradish peroxidase-conjugated anti-rabbit (1:2000, R&D Systems,

HAF008) or mouse (1:2000, R&D Systems, HAF018) secondary antibody. Immunoreactive bands

were detected using SuperSignal West Pico Chemiluminescent substrate (Thermo Fisher

Scientific, 34080), and images were captured using an iBright CL750 Imager (Thermo Fisher

Scientific). Immunoblots were quantified using ImageJ software (NIH) relative to β-actin

expression.

Proteomic analysis

On days 1, 3, and 5 post-infection, MDMs were collected, lysed with 2% SDS in 100 mM Tris-

HCL and 100 mM dithiothreitol, pH 7.6, and supplemented with protease and phosphatase

inhibitors. Protein concentration was determined using Pierce 660 Protein Assay kit with ionic

detergent compatibility reagent according to the manufacturer's instructions. Afterward, samples

were processed as previously described (Arainga et al., 2015) using filter-aided sample preparation

(FASP, Pall Life Sciences, OD010C34) for digestion of 50 μg per sample. Following overnight

digestion, samples were cleaned using the Oasis MCX column (Waters, 186000252) and C18 Zip-

Tips (Sigma-Aldrich, ZTC18M960). Cleaned peptides were quantitated using NanoDrop2000 at

205 nm. Following resuspension in 0.1% formic acid, 2 μg of the sample was used for label-free

quantification (LFQ) in UNMC Mass Spectrometry and Proteomics Core Facility as previously

described (Gao et al., 2020). Proteins identified by mass spectrometry were quantified to determine

differentially expressed proteins between mock and SARS-CoV-2-challenged MDMs at different

time points post-infection. Statistical analysis for proteomic data was performed using ANOVA,

and false discovery rate (FDR) for pathway analysis was controlled using the Benjamini-Hochberg

(BH) method (Benjamini and Hochberg, 1995). A protein was considered to be differentially

expressed if p value ≤ 0.05 and the absolute value of fold change ≥ 2. Next, functional and pathway

enrichment analysis of differentially expressed proteins in SARS-CoV-2-challenged MDMs

compared to mock-challenged MDMs was conducted at different times after virus challenge. Gene

enrichment analysis to identify immune system processes affected after virus exposure was

performed using Cytoscape in conjunction with the plug-in ClueGO (Bindea et al., 2009). GO

annotation, KEGG, Reactome GSEA, and STRING analyses were conducted using GO Resource,

DAVID, ReactomeFIViz, and STRING databases, respectively.

Statistical analysis

Results are presented as the mean ± standard error of the mean (SEM). Student's t-test or one-way

ANOVA followed by Tukey's multiple comparison test was used to analyze differences in the

mean values between groups. P values ≤ 0.05 were considered statistically significant. Statistical

analysis was performed using GraphPad Prism 9.1.0 software (GraphPad Software, San Diego,

CA).

Results

ACE2 receptor expression during human monocyte-macrophage maturation

ACE inhibitors were used to upregulate the SARS-CoV-2 receptor ACE2 and potentiate cell entry

(Ferrario et al., 2005; Pedrosa et al., 2021; Soler et al., 2009). As ACE inhibitors are frequently

prescribed to treat hypertension, we posit that they can increase susceptibility to SARS-CoV-2.

Therefore, we measured ACE2 expression on freshly isolated monocytes throughout cell isolation

and differentiation. Days 1, 3, 5, and 7 were evaluated for cells cultured with or without captopril,

an ACE inhibitor which is thought to increases ACE2 expression. In these studies, ACE2

expression by monocytes was found to peak by day 5 after initiation of cell differentiation then

decreased after that (**Figure 1A**), which corresponds to the susceptibility of infection of

monocytes-macrophages by other viruses (Gendelman et al., 1992; Lum et al., 2018;

Schmidtmayerova et al., 1996). On day 5, ACE2 expression increased by 23% in captopril-treated

cells. In parallel, the expression of phenotypic monocyte-macrophage surface markers, CD14 and

CD16, were not changed (**Figure 1A**). Based on these observations, we challenged MDMs with

SARS-CoV-2 on day 5 after start of cell differentiation and assessed whether captopril affects

susceptibility to infection.

Abortive infection of MDMs is established following SARS-CoV-2 exposure

To investigate whether human MDMs are susceptible to SARS-CoV-2 infection, monocytes were

isolated from healthy donors and differentiated in the presence or absence of captopril and infected

at day 5 after cell culture. The kinetic growth of SARS-CoV-2 in MDMs was determined from

culture supernatant tests of challenged cells over 10 days post-infection. In all studied groups

(captopril-treated and controls), the SARS-CoV-2 genome equivalent copies in the supernatant

significantly decreased during 5 days of viral exposure, becoming undetectable after that (**Figure**

1B). Additionally, the number of genome copies did not change significantly with or without

captopril. These results demonstrated that SARS-CoV-2 infection of human MDMs was abortive

and differentiated monocytes could not generate progeny virus in contrast to productive infection

of SARS-CoV-2 in Vero.STAT1 KO cells (**Figure 1B**). Our data also indicated that captopril did

not alter the susceptibility of MDMs to SARS-CoV-2 infection.

Ultrastructural features of SARS-CoV-2-challenged MDMs

Viral particles and mock and SARS-CoV-2-challenged MDMs were fixed at different time points

A

% ACE2 positive cells / Days of differentiation

% CD14 positive cells / Days of differentiation

% CD16 positive cells / Days of differentiation

B

■ w/o captopril ■ with captopril

SARS-CoV-2 genome equivalents/ml — MDMs / Days post-inoculation

SARS-CoV-2 genome equivalents/ml — Vero.STAT1 KO cells

C

(i) 50 nm

(ii) 400 nm

(iii) 50 nm / 400 nm / Cyt

(iv) 50 nm / 400 nm / N / Cyt

(v) 50 nm / 400 nm / Cyt / IB

(vi) 50 nm / 400 nm

Figure 1. SARS-CoV-2 abortive infection of human MDMs. (A) Expression of SARS-CoV-2 cell entry receptor ACE2 and phenotypic surface markers CD14 and CD16, during differentiation of monocytes into macrophages, was analyzed by flow cytometry in absence or presence of captopril. (B) SARS-CoV-2 presence in MDMs. SARS-CoV-2 (MOI=0.01) was used to infect MDMs (with or without captopril). The number of virus genome equivalents per ml was measured in the culture supernatant by RT-qPCR. Vero.STAT1 KO cells served as a positive control. (C) Transmission electron micrographs of SARS-CoV-2-challenged MDMs. (C.i) A viral particle from a pool of SARS-CoV-2 used for the virus challenges. Mock (C.ii) and SARS-CoV-2-challenged MDMs (C.iii: day 1, C.iv: day 3, C.v: day 5, and C.vi: day 14 after viral exposure). Red arrows and boxes demonstrate virus particles inside inclusion bodies and within the macrophage cytoplasm. All experiments were done at least twice with representative images depicted here. (**A** and **B**) Data are represented as mean ± SEM (n=3-6). Statistical significance between groups was determined using unpaired Student's t-test or one-way ANOVA, and $p < 0.05$ was considered significant (*** $p < 0.001$). Abbreviations: w/o: without; Cyt: cytoplasm; N: nucleus; IB: inclusion body. Scale bars: 50 nm (i and left panels of iii, iv, v, and vi) and 400 nm (ii and right panels of iii, iv, v, and vi).

after viral exposure and then examined by TEM to determine ultrastructural changes in the virus-challenged cells (**Figure 1C.i-vi**). Negative stain TEM shows the ultrastructure of a viral particle from a pool of SARS-CoV-2 used for the virus challenges (**Figure 1C.i**). TEM analysis of mock-treated cells showed normal architecture (**Figure 1C.ii**). By days 1 and 3 after exposure, viral particles were observed inside the cytoplasm of many cells (**Figures 1C.iii** and **1C.iv**). Five days after viral exposure, an increased number of virions were observed inside inclusion bodies and cell cytoplasm (**Figure 1C.v**). Surprisingly, while no viral genome copies were detected after 5 days (**Figure 1B**), mature virions were observed 14 days after initial exposure (**Figure 1C.vi**). These underly the persistence of virus in macrophages despite evidence of abortive infection.

Abortive SARS-CoV-2 infection induces pro-inflammatory factors, but not IFN

To investigate alterations in macrophage-mediated innate immune responses after SARS-CoV-2 exposure, a total of 84 key genes were examined. Transcriptional changes linked to immunity were screened using RT^2 Profiler Human Innate and Adaptive Immune Response Array. Expression of

immune response genes in SARS-CoV-2-challenged MDMs were investigated against control

"mock-challenged cells" on days 1, 3, and 5 (**Supplementary Figure 1**). On day 1, more than two-

fold increase in expression of pro-inflammatory cytokine and chemokine genes were detected,

including *IL-6, TNF-α, IL-1α, IL17A, IL8*, colony-stimulating factor 2 (*CSF2*), chemokine (C-C

motif) ligand 2 (*CCL2*), and *CCL5*. Interestingly, on day 3, gene expression of the node-like

receptor family pyrin domain containing 3 (*NLRP3*) and *IL-1β* increased by 8.84- and 5.64-fold,

respectively. On day 5, expression for these pro-inflammatory cytokines and chemokines were at

baseline or decreased with the exception of *IL-6* and *CCL2*. Gene expression of anti-inflammatory

cytokine *IL-10* was elevated on days 1 and 5 supporting the attenuation of pro-inflammatory

responses (Lu et al., 2021). Importantly, expression of IFN-linked genes *IFN-α1, IFN-β1, IFN-γ*,

IFN regulatory factor 7 (*IRF7*), and tyrosine kinase 2 (*TYK2*) genes increased on day 1 but returned

to near baseline by days 3 and 5. These data demonstrate a complex pro- and anti-inflammatory

network interactions that are sustained to day 5 with attenuated IFN responses following SARS-

CoV-2 exposure (**Figure 2A**).

Functional and pathway enrichment of immune regulated genes

SARS-CoV-2 infection may induce dynamic changes of immune-based gene expression in specific

cellular biological processes and pathways in virus-challenged cells. To assess these changes,

functional and pathway enrichment analyses were performed on immune-regulated genes screened

in SARS-CoV-2-challenged MDMs compared to controls at days 1, 3, and 5 (**Supplementary file

1**). Multiple immune processes were enriched upon viral exposure, including defense response to

other organism, positive regulation of immune system processes, cell surface receptor signaling

pathways, and positive regulation of immune responses (**Figure 2B**).

Figure 2. Transcriptomic profile of SARS-CoV-2-challenged MDMs. Expression of 84 genes
specific for human innate and adaptive immune responses were screened in SARS-CoV-2-
challenged MDMs compared to mock-challenged MDMs at different times post-challenge, using
RT^2 Profiler Human Innate and Adaptive Immune Response 96-well Array. Fold changes in the
gene expression were determined via Qiagen's RT^2 Profiler analysis software (n=4). (A) IPA was
performed with upregulated or downregulated genes to identify putative network interactions
involved in SARS-CoV2-MDM interactions. Upregulated genes are shaded red with darker shade
indicating higher upregulation, while green shades denote downregulation in gene expression.
Solid grey lines indicate direct interactions, while dotted grey lines correspond to indirect
interactions. (B) Functional and pathway enrichment analysis of transcriptomic dataset was
performed using GO-term, KEGG, and Reactome analyses at different time points post-infection.
Different immune and inflammation-related biological processes and pathways affected in SARS-
CoV-2-challenged MDMs were plotted as a bar chart compared to mock-challenged MDMs.

All indicated they comprised activated immune responses in the virus-challenged MDMs. In

addition, a series of inflammation-related processes were enriched, including inflammatory

response, positive regulation of cytokine production, and cellular response to cytokine stimulus

(**Figure 2B**), highlighting inflammation responses induced after viral exposure. Similarly,

different immune and inflammation-related molecular functions such as cytokine activity,

chemokine activity, complement component C1q binding, type I IFN receptor binding, and IL-1

receptor binding were enriched after virus exposure (**Supplementary file 1**). Furthermore, KEGG

pathway analysis showed enrichment of TLR, NOD-like receptor, and RLR signaling pathways

(**Figure 2B**), suggesting the activation of PRRs on MDMs to recognize viral PAMPs initiating

innate immune responses. Moreover, KEGG pathway analysis depicted enrichment of IL-17, NF-

κB, TNF, and chemokine signaling pathways (**Figure 2B**), highlighting different inflammatory

pathways induced in the SARS-CoV-2-challenged MDMs. Interestingly, Reactome analysis

revealed enrichment of multiple IFN-related signaling pathways including IFN-α/β signaling, IFN-

γ signaling, TICAM1-dependent activation of IRF3/IRF7, TRAF3-dependent IRF activation

pathway, and TRAF6 mediated IRF7 activation (**Figure 2B**). In addition, immune and

inflammation-mediated molecular functions were enriched in virus-challenged cells

(**Supplementary file 1**). The interacting proteins involved in the screened innate and adaptive

immune-related genes were also identified using STRING analysis of their interaction networks

(**Supplementary file 1**).

SARS-CoV-2 monocyte-macrophage engagement and failed induction of IFN activity

To determine whether exposure to the SARS-CoV-2 can trigger antiviral IFN activities in human

macrophages, culture supernatants were collected at different times post-viral exposure and

assayed in VSV-challenged MDBK cells. Notably, no type I IFN activity was observed in SARS-

CoV-2-challenged MDM cultures at different time intervals following viral challenge (**Figure**

3A). These data stand in contrast to the transcriptomic results, which displayed increases in IFN

pathway-linked genes. Furthermore, no additive or synergistic IFN activity was detected in control

and SARS-CoV-2-challenged monocytes treated with poly(I:C) (**Figure 3B**). Taken together, the

data demonstrate that SARS-CoV-2-MDM interactions do not affect IFN activities. To affirm this,

we evaluated the protein expression level of IFN-α and IFN-β in SARS-CoV-2 exposed MDMs at

days 1, 3, and 5 post-exposure. Similar to IFN activity, no IFN-α or IFN-β proteins were produced

in the SARS-CoV-2-challenged MDMs (**Figure 3C**). Altogether, our data demonstrate that SARS-

CoV-2 exposure of MDM does not trigger IFN activity or protein production in human monocytes-

macrophages.

Proteomic profiles of SARS-CoV-2-challenged MDMs

In attempts to elucidate the mechanisms of the MDM response against the SARS-CoV-2 challenge,

we obtained the proteome profiles from lysates of viral exposed macrophages and compared those

to non-exposed counterparts. On defined days after viral exposure, the expression of more than

Figure 3. SARS-CoV-2 does not induce IFN activity in monocytes-macrophages. (A) MDMs were infected with SARS-CoV-2 (MOI=0.01) and the culture supernatants were collected at days 1, 3, and 5 post-infection. The collected culture supernatants were used to assess IFN activity in VSV-challenged bovine MDBK cells. Recombinant human IFN-α was used as standard. (A.i) Experimental timeline. (A.ii) Representative images of IFN activity assay plates. (B) No additive or synergistic responses in poly(I:C)-induced IFN activity between control and SARS-CoV-2-challenged monocytes. Suspended monocytes were treated with 100 μg poly(I:C) overnight before or after the infection with SARS-CoV-2 (MOI=0.01) for 4 h. (B.i) Experimental timeline. (B.ii) Representative images of IFN activity assay plates. (C) Western blot analysis was performed to determine expression of IFN-α and IFN-β in cell lysates at different times after viral exposure. Representative immunoblot and densitometric quantification are shown. All experiments were done at least twice, and one representative image is shown. (C) Data represent mean ± SEM (n=3). Statistical significance between the groups was determined with unpaired Student's t-test and $p < 0.05$ was considered significant.

4,000 proteins were identified and quantified using differential proteomic tests. Amongst total identified proteins, 1776, 1372, and 2448 proteins were significantly differentially expressed ($p \leq 0.05$ and fold change ≥ 2), on days 1, 3, and 5, respectively, after viral challenge (**Supplementary file 2**). Volcano plots depicting differentially expressed proteins in SARS-CoV-2-challenged MDMs compared to controls are shown in **Supplementary Figure 2**. To obtain a detailed understanding of changes in the MDM proteomic profile after viral challenge, we performed functional and pathway enrichment analyses of differentially regulated proteins in virus-challenged MDMs at different times. Amongst total proteins altered on day 1, only 5% were involved in negative regulation of inflammatory response to an antigenic stimulus. In contrast, on days 3 and 5, 38.1% and 23.53% of altered proteins, respectively, belonged to myeloid cell activation involved in immune responses (**Figure 4A**). Immune processes of each function indicated in pie charts include, but are not limited to, activation of the innate immune response, myeloid cell activation, innate immune response activating signal transduction, and innate immune response activating cell surface receptor signaling pathway (**Figure 4A** and **Supplementary file 3**). All indicated activated immune and inflammatory responses in MDMs upon viral challenge.

Figure 4. Proteomic profile of SARS-CoV-2-challenged MDMs. (A) Gene enrichment analysis was performed using Cytoscape in conjunction with the plug-in ClueGO at different times after viral exposure. Pie charts representing the distribution of identified differentially expressed proteins according to observed immune processes and bar charts demonstrating the specific processes that correspond to their classification. The same color key used in the pie charts applied in the bar charts. (B) Functional and pathway enrichment analyses of proteomic dataset were performed using GO annotation, KEGG, Reactome GSEA, and STRING analyses at different time points post-infection. Different immune-related biological processes and PPI networks affected in SARS-CoV-2-challenged MDMs were plotted as a bar chart compared to mock-challenged MDMs.

Additionally, viral exposure induced changes in biological processes related to the endoplasmic reticulum (ER) and mRNA on days 1 and 3 after challenge. With positive fold changes, the enriched biological processes included protein localization to ER, protein targeting to ER, regulation of mRNA metabolic process, nuclear-transcribed mRNA catabolic process, and mRNA splicing via spliceosome (**Supplementary file 4**). Similarly, on days 1 and 3 after viral challenge, Reactome analysis depicted enrichment of processes related to RNA processing, with positive fold change, including rRNA processing in the nucleus and cytosol and mRNA splicing (**Supplementary file 4**). Moreover, KEGG pathway analysis showed an enrichment of ribosome function as a pathway of genetic information processing and translation on day 1. Similarly, the molecular function of a structural constituent of the ribosome was enriched with positive fold change on day 1 (**Supplementary file 4**). Interestingly, GO-term analysis showed robust enrichment of different immune responses such as complement activation, humoral immune response, adaptive immune response, and regulation of inflammatory response with negative fold changes suggesting dysregulated immune responses to SARS-CoV-2 in MDMs (**Figure 4B**). Importantly, STRING analysis of PPIs showed enrichment of IFN-related interaction networks such as "IFN signaling and positive regulation of RIG-I signaling pathway" and "IFN-α/β signaling and IFN-γ signaling", with negative fold changes, suggesting dysregulated IFN

responses induced by SARS-CoV-2 (**Figure 4C** and **Supplementary file 4**). Overall, these results

support transcriptomic, IFN activity, and protein evaluations that demonstrated attenuated IFN

responses in SARS-CoV-2-challenged MDMs.

Discussion

MPs are the first line of host defense against viral infection (Nikitina et al., 2018). These sentinel

cells interact with SARS-CoV-2 to define COVID-19 pathogenesis. That includes, but is not

limited to, CSS, ARDS, and multiorgan dysfunction (Guan et al., 2020). As a first line defense

mechanism, the initial virus-MP interaction following the entry of the pathogenic virus into host

cells represents a critical determinant for infectivity and pathogenesis (Li, 2016; Perlman and

Netland, 2009; Shang et al., 2020). To this end, we have better defined such virus-host cell

interactions and determined relationships between ACE2 receptor expression and monocyte-

macrophage differentiation during abortive SARS-CoV-2 infection. Like other viral infections, we

reasoned that monocytes would respond with an elevated expression of pro-inflammatory

molecules and antiviral responses. Indeed, this has already been reported for influenza,

Chikungunya, herpes, Zika, and lentiviral infections that include human immunodeficiency virus

type one (HIV-1) and herpes viruses (Coates et al., 2018; Her et al., 2010; Krzyzowska et al., 2013;

Lum et al., 2018; Schmidtmayerova et al., 1996). Each elicits profound inflammation during early

viral infection. However, unlike other known viral infections, SARS-CoV-2 infection is abortive

and balances pro- and anti-inflammatory processes with a unique cell signature. The resulting

signature defines the pathways towards the major cause for morbidity and mortality in COVID-

19: the CSS leading to ARDS and other multi-organ failures. The specific lack of induction of IFN

also serves to define a specific monocyte-macrophage-virus phenotype during both early and

progressive infection (Dalskov et al., 2020; Yang et al., 2020).

Herein, we show increased expression of viral receptor ACE2 during monocyte-macrophage

differentiation. Interestingly, further induction of ACE2 expression by captopril, a well-known

ACE inhibitor, was observed. Captopril induces a reduction of angiotensin II, which increases

ACE2 expression and activity. This occurs through angiotensin type 1 receptor-dependent ACE2

internalization followed by lysosomal degradation (Deshotels et al., 2014). However, the observed

upregulation of ACE2 expression failed to affect infection with SARS-CoV-2. Despite infection

of MDMs with SARS-CoV-2 when ACE2 receptor expression peaked, the infection remained

abortive in agreement with prior reports (Boumaza et al., 2021; Yang et al., 2020; Zheng et al.,

2021). The inability of captopril to increase the susceptibility of MDM infection with SARS-CoV-

2 is in accordance with a recent study indicating that treatment of human alveolar type-II

pneumocytes with captopril induced upregulation of ACE2 expression and counteracted drug-

induced reduction of SARS-CoV-2 spike protein entry (Pedrosa et al., 2021). This occurred

through inhibition of A disintegrin and metalloprotease 17 (ADAM17), which has been shown to

play an essential role in ACE2 shedding and viral entry into the cells (Lambert et al., 2005).

Nonetheless, viral particles were present inside cytoplasmic structures of macrophages up to two

weeks after the viral challenge underscoring any continuity of immune responses.

SARS-CoV-2 abortive infection likely results from intracellular mechanisms induced upon MP

activation. Human coronaviruses can infect human peripheral blood mononuclear cells leading to

cell activation and aberrant production of pro-inflammatory mediators with increased

507 chemoattraction (Channappanavar and Perlman, 2017; Yilla et al., 2005; Zhou et al., 2014). In

508 particular, virus-exposed monocytes-macrophages can serve as perpetrators for virus-induced

509 inflammatory responses within different body organs, as seen by the abundance of pro-

510 inflammatory macrophages in bronchoalveolar lavage fluid obtained from severe COVID-19 cases

511 (Liao et al., 2020). The MP pro-inflammatory factors can contribute to local tissue inflammation

512 and systemic inflammatory responses that characterize cytokine storm (Nikitina et al., 2018). In

513 the current study, the transcriptomic profile of SARS-CoV-2-challenged MDMs demonstrated

514 increased mRNA expression of multiple pro-inflammatory molecules, including *IL-6, TNF-α, IL-*

515 *1α, IL17A, IL8, CSF2, CCL2, CCL5, NLRP3,* and *IL-1β* upon the virus exposure. Our data

516 confirmed previous reports which showed excessive production of pro-inflammatory cytokines

517 and chemokines in the SARS-CoV-2-exposed MPs (Boumaza et al., 2021; Yang et al., 2020;

518 Zheng et al., 2021). In parallel, our data displayed enrichment of different inflammation-related

519 pathways such as IL-17, NF-κB, and TNF signaling pathways, following viral exposure. IL-17 is

520 known for its pivotal role in inducing and mediating pro-inflammatory responses and its

521 involvement in different inflammatory autoimmune diseases (Kuwabara et al., 2017). NF-κB

522 serves as a central mediator of inflammation since the DNA binding site for NF-κB was found in

523 the promoter regions of multiple pro-inflammatory molecules (Hayden and Ghosh, 2004). Thus,

524 the activation of NF-κB with other pro-inflammatory transcription factors leads to the transcription

525 of several pro-inflammatory molecules such as IL-1β, inducible nitric oxide synthase (iNOS), and

526 TNF-α (Dasgupta et al., 2003; Liu et al., 2002). In addition, the transcriptome of SARS-CoV-2-

527 challenged MDMs showed enrichment of multiple immune-related biological processes,

528 molecular functions, and signaling pathways such as positive regulation of immune response, cell

529 surface receptor signaling pathway, cytokine activity, chemokine activity, TLR signaling pathway,

and NOD-like receptor signaling pathway, demonstrating cell activation of MDMs after viral

exposure. Notably, the proteome of virus-challenged macrophages revealed the dysregulation of

different immune processes such as regulation of inflammatory response, complement activation,

and linked adaptive and humoral immune responses. Overall, our findings demonstrate MDM

activation by SARS-CoV-2 is associated with exaggerated inflammatory responses and

dysregulated immune activities.

Significantly, SARS-CoV-2 induced an attenuated MP IFN response, as the transcript expression

of type I (IFN-α1 and IFN-β1) and type II (IFN-γ) IFNs in the virus-challenged cells increased

early after exposure but returned to or below normal levels at later times. Our findings are

consistent with previous reports depicting a dysregulated IFN response in SARS-CoV-2-

challenged MDMs and alveolar macrophages (Dalskov et al., 2020; Yang et al., 2020). This

attenuated IFN response provides a possible mechanism for delayed viral clearance from cells up

to two weeks after inoculation. Similarly, our study showed the inability of SARS-CoV-2 to induce

IFN activity or production in MDMs, as illustrated by the failure of culture supernatants of infected

cells to provide protection against the cytopathic effects induced by VSV in MDBK cells as well

as unchanged levels of IFN-α and IFN-β after the virus challenge. These findings are following a

recent study that demonstrated the absence of IFN induction in the alveolar macrophages

challenged with SARS-CoV-2 (Dalskov et al., 2020). A possible explanation is the presence of a

cap structure on the viral genome. The *Coronoviridae* family contains this structure enabling the

virus to evade recognition by PRRs and prevent the host innate immune response mediated by the

RIG-I/mitochondrial antiviral-signaling (MAVS) pathway that recognizes single-stranded RNAs

without a cap structure (Chen and Guo, 2016; Pichlmair et al., 2006). Moreover, different SARS-

CoV-2 proteins were found to antagonize type I IFN production through other mechanisms (Xia

et al., 2020). SARS-CoV-2 nonstructural protein 6 (nsp6) binds TANK binding kinase 1 (TBK1)

to suppress IRF3 phosphorylation, and nsp13 binds and blocks TBK1 phosphorylation. In addition,

open reading frame 6 (ORF6) binds importin Karyopherin α 2 to inhibit IRF3 nuclear

translocation, and ORF7b prevents STAT1 phosphorylation and nuclear translocation,

consequently inhibiting the transcription of multiple IFN stimulated genes (ISGs) which possess

antiviral functions (Xia et al., 2020). In parallel, the negative fold change of IFN-related interaction

networks: "IFN signaling and positive regulation of RIG-I signaling pathway" and "IFN-α/β

signaling and IFN-γ signaling" indicates dysregulated IFN responses in the SARS-CoV-2-

challenged MDMs. Thus far, our data suggest an aberrant IFN response induced by SARS-CoV-2

in human MDMs. Early IFN induction was halted at the transcriptional level one day post-infection

and did not proceed to the translational level. Thus, we postulate that aberrant IFN responses in

the face of a robust inflammatory environment presage a lack of viral infection control and

multiorgan damage in COVID-19 (Gavriatopoulou et al., 2020).

In summary, we demonstrate that abortive SARS-CoV-2 macrophage infection triggers a unique

signature of inflammatory responses. This includes combinations of abortive infection with

delayed viral clearance, excessive pro-inflammatory cytokine and chemokine productions, and

attenuated IFN responses. Each and all imply dysfunction of the innate immune system after viral

exposure. Transcriptomic and proteomic profiles of SARS-CoV-2-challenged MPs provide

signatures for understanding COVID-19 pathogenesis. The lack of adequate antiviral innate

immune responses in the midst of cytokine storm heralds an absence of viral infection control that

exacerbates clinical manifestations and contributes to end-organ damage for COVID-19-related

morbidities and mortalities.

Conflict of interest

Acknowledgments

The authors thank the UNMC Elutriation and Cell Separation Core Facility (Myhanh Che and Na

Ly) for providing human monocytes. The authors would also like to thank Dr. Pawel Ciborowski

and his research group within the Department of Pharmacology and Experimental Neuroscience

at UNMC for discussion and assistance with the proteomic sample preparation. The authors

acknowledge the UNMC Mass Spectrometry and Proteomics Core Facility for assistance with the

proteomic analysis. The authors would also like to acknowledge the UNMC Flow Cytometry

Research Core Facility for assistance with the flow cytometric analysis. The authors would like to

thank Dirk Anderson (The Biotech Microscopy Core Research Facility of University of Nebraska-

Lincoln) for his assistance with TEM sample preparations.

Funding

The work was supported, in part, by the University of Nebraska Foundation, which includes

donations from the Carol Swarts, M.D. Emerging Neuroscience Research Laboratory, the

Margaret R. Larson Professorship, the Frances and Louie Blumkin, and the Harriet Singer

Research Donations. We thank Dr. Bradley Britigan, Dean of the College of Medicine at UNMC,

for providing funds to support the SARS-CoV-2 studies. The research also received support from

National Institutes of Health grants PO1 DA028555, R01 NS36126, PO1 MH64570, P30

MH062261, P20 GM113126, R01 AG043540, and 2R01 NS034239. We also thank the INBRE

grant support from 2P20GM103427 for infrastructure research support.

Author contributions

M.M.A.: Investigation, Methodology, Validation, Visualization, Formal analysis, Data curation,

Writing-original draft, review and editing; P.Y.: Investigation, Methodology, Formal analysis,

Data curation, Writing-review and editing; J.M.: Investigation, Methodology, Data curation,

Writing-review and editing; K.E.O.: Validation, Visualization, Writing-review and editing; F.S.:

Investigation, Validation, Writing-review and editing; Y.Z.: Investigation, Validation, Data

curation, Writing-review and editing; L.J.: Investigation, Formal analysis, Writing-review and

editing; K.P.: Investigation, Validation, Writing-review and editing; A.A.: Investigation,

Validation, Visualization, Writing-review and editing; S.N.B.: Supervision, Visualization,

Writing-review and editing; R.L.M.: Conceptualization, Methodology, Supervision, Writing-

review and editing; H.E.G.: Conceptualization, Funding Acquisition, Methodology, Resources,

Supervision, Writing-original draft, review and editing; All authors have read and approved the

final version of the manuscript.

References

Arainga, M., Guo, D., Wiederin, J., Ciborowski, P., McMillan, J., and Gendelman, H. E. (2015). Opposing regulation of endolysosomal pathways by long-acting nanoformulated antiretroviral therapy and HIV-1 in human macrophages. *Retrovirology, 12*, 5. doi:10.1186/s12977-014-0133-5

Benjamini, Y., and Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B (Methodological), 57*(1), 289-300. doi:https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Bindea, G., Mlecnik, B., Hackl, H., Charoentong, P., Tosolini, M., Kirilovsky, A., Fridman, W. H., Pages, F., Trajanoski, Z., and Galon, J. (2009). ClueGO: a Cytoscape plug-in to decipher functionally grouped gene ontology and pathway annotation networks. *Bioinformatics, 25*(8), 1091-1093. doi:10.1093/bioinformatics/btp101

Boumaza, A., Gay, L., Mezouar, S., Bestion, E., Diallo, A. B., Michel, M., Desnues, B., Raoult, D., La Scola, B., Halfon, P., Vitte, J., Olive, D., and Mege, J. L. (2021). Monocytes and macrophages, targets of SARS-CoV-2: the clue for Covid-19 immunoparalysis. *J Infect Dis*. doi:10.1093/infdis/jiab044

Channappanavar, R., and Perlman, S. (2017). Pathogenic human coronavirus infections: causes and consequences of cytokine storm and immunopathology. *Semin Immunopathol, 39*(5), 529-539. doi:10.1007/s00281-017-0629-x

Chen, Y., and Guo, D. (2016). Molecular mechanisms of coronavirus RNA capping and methylation. *Virol Sin, 31*(1), 3-11. doi:10.1007/s12250-016-3726-4

Coates, B. M., Staricha, K. L., Koch, C. M., Cheng, Y., Shumaker, D. K., Budinger, G. R. S., Perlman, H., Misharin, A. V., and Ridge, K. M. (2018). Inflammatory monocytes drive influenza A virus-mediated lung injury in juvenile mice. *J Immunol, 200*(7), 2391-2404. doi:10.4049/jimmunol.1701543

Dalskov, L., Mohlenberg, M., Thyrsted, J., Blay-Cadanet, J., Poulsen, E. T., Folkersen, B. H., Skaarup, S. H., Olagnier, D., Reinert, L., Enghild, J. J., Hoffmann, H. J., Holm, C. K., and Hartmann, R. (2020). SARS-CoV-2 evades immune detection in alveolar macrophages. *EMBO Rep, 21*(12), e51252. doi:10.15252/embr.202051252

Dasgupta, S., Jana, M., Liu, X., and Pahan, K. (2003). Role of very-late antigen-4 (VLA-4) in
myelin basic protein-primed T cell contact-induced expression of proinflammatory
cytokines in microglial cells. *J Biol Chem, 278*(25), 22424-22431.
doi:10.1074/jbc.M301789200

Deshotels, M. R., Xia, H., Sriramula, S., Lazartigues, E., and Filipeanu, C. M. (2014). Angiotensin
II mediates angiotensin converting enzyme type 2 internalization and degradation through
an angiotensin II type I receptor-dependent mechanism. *Hypertension, 64*(6), 1368-1375.
doi:10.1161/HYPERTENSIONAHA.114.03743

Diao, B., Wang, C., Wang, R., Feng, Z., Zhang, J., Yang, H., Tan, Y., Wang, H., Wang, C., Liu,
L., Liu, Y., Liu, Y., Wang, G., Yuan, Z., Hou, X., Ren, L., Wu, Y., and Chen, Y. (2021).
Human kidney is a target for novel severe acute respiratory syndrome coronavirus 2
infection. *Nat Commun, 12*(1), 2506. doi:10.1038/s41467-021-22781-1

Durbin, J. E., Hackenmiller, R., Simon, M. C., and Levy, D. E. (1996). Targeted disruption of the
mouse Stat1 gene results in compromised innate immunity to viral disease. *Cell, 84*(3),
443-450. doi:10.1016/s0092-8674(00)81289-1

Faradji, A., Bohbot, A., Schmitt-Goguel, M., Siffert, J. C., Dumont, S., Wiesel, M. L., Piemont,
Y., Eischen, A., Bergerat, J. P., Bartholeyns, J., and et al. (1994). Large scale isolation of
human blood monocytes by continuous flow centrifugation leukapheresis and counterflow
centrifugation elutriation for adoptive cellular immunotherapy in cancer patients. *J
Immunol Methods, 174*(1-2), 297-309. doi:10.1016/0022-1759(94)90033-7

Feng, Z., Diao, B., Wang, R., Wang, G., Wang, C., Tan, Y., Liu, L., Wang, C., Liu, Y., Liu, Y.,
Yuan, Z., Ren, L., Wu, Y., and Chen, Y. (2020). The novel severe acute respiratory
syndrome coronavirus 2 (SARS-CoV-2) directly decimates human spleens and lymph
nodes. *medRxiv*, 2020.2003.2027.20045427. doi:10.1101/2020.03.27.20045427

Ferrario, C. M., Jessup, J., Chappell, M. C., Averill, D. B., Brosnihan, K. B., Tallant, E. A., Diz,
D. I., and Gallagher, P. E. (2005). Effect of angiotensin-converting enzyme inhibition and
angiotensin II receptor blockers on cardiac angiotensin-converting enzyme 2. *Circulation,
111*(20), 2605-2610. doi:10.1161/CIRCULATIONAHA.104.510461

Field, A. K., Tytell, A. A., Lampson, G. P., and Hilleman, M. R. (1967). Inducers of interferon
and host resistance. II. Multistranded synthetic polynucleotide complexes. *Proc Natl Acad
Sci U S A, 58*(3), 1004-1010. doi:10.1073/pnas.58.3.1004

Gao, L., Kumar, V., Vellichirammal, N. N., Park, S. Y., Rudebush, T. L., Yu, L., Son, W. M.,
Pekas, E. J., Wafi, A. M., Hong, J., Xiao, P., Guda, C., Wang, H. J., Schultz, H. D., and
Zucker, I. H. (2020). Functional, proteomic and bioinformatic analyses of Nrf2- and
Keap1- null skeletal muscle. *J Physiol, 598*(23), 5427-5451. doi:10.1113/JP280176

Gavriatopoulou, M., Korompoki, E., Fotiou, D., Ntanasis-Stathopoulos, I., Psaltopoulou, T.,
Kastritis, E., Terpos, E., and Dimopoulos, M. A. (2020). Organ-specific manifestations of
COVID-19 infection. *Clin Exp Med, 20*(4), 493-506. doi:10.1007/s10238-020-00648-x

Gendelman, H. E., Baca, L. M., Kubrak, C. A., Genis, P., Burrous, S., Friedman, R. M., Jacobs,
D., and Meltzer, M. S. (1992). Induction of IFN-alpha in peripheral blood mononuclear
cells by HIV-infected monocytes. Restricted antiviral activity of the HIV-induced IFN. *J*
Immunol,   *148*(2),   422-429.   Retrieved   from
https://www.ncbi.nlm.nih.gov/pubmed/1729362

Guan, W. J., Ni, Z. Y., Hu, Y., Liang, W. H., Ou, C. Q., He, J. X., Liu, L., Shan, H., Lei, C. L.,
Hui, D. S. C., Du, B., Li, L. J., Zeng, G., Yuen, K. Y., Chen, R. C., Tang, C. L., Wang, T.,
Chen, P. Y., Xiang, J., Li, S. Y., Wang, J. L., Liang, Z. J., Peng, Y. X., Wei, L., Liu, Y.,
Hu, Y. H., Peng, P., Wang, J. M., Liu, J. Y., Chen, Z., Li, G., Zheng, Z. J., Qiu, S. Q., Luo,
J., Ye, C. J., Zhu, S. Y., Zhong, N. S., and China Medical Treatment Expert Group for, C.
(2020). Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med,*
382(18), 1708-1720. doi:10.1056/NEJMoa2002032

Hayden, M. S., and Ghosh, S. (2004). Signaling to NF-kappaB. *Genes Dev, 18*(18), 2195-2224.
doi:10.1101/gad.1228704

Her, Z., Malleret, B., Chan, M., Ong, E. K., Wong, S. C., Kwek, D. J., Tolou, H., Lin, R. T.,
Tambyah, P. A., Renia, L., and Ng, L. F. (2010). Active infection of human blood
monocytes by Chikungunya virus triggers an innate immune response. *J Immunol, 184*(10),
5903-5913. doi:10.4049/jimmunol.0904181

Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., Zhang, L., Fan, G., Xu, J., Gu, X., Cheng,
Z., Yu, T., Xia, J., Wei, Y., Wu, W., Xie, X., Yin, W., Li, H., Liu, M., Xiao, Y., Gao, H.,
Guo, L., Xie, J., Wang, G., Jiang, R., Gao, Z., Jin, Q., Wang, J., and Cao, B. (2020). Clinical
features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet,*
395(10223), 497-506. doi:10.1016/S0140-6736(20)30183-5

Huang da, W., Sherman, B. T., and Lempicki, R. A. (2009). Systematic and integrative analysis
of large gene lists using DAVID bioinformatics resources. *Nat Protoc, 4*(1), 44-57.
doi:10.1038/nprot.2008.211

Krzyzowska, M., Baska, P., Orlowski, P., Zdanowski, R., Winnicka, A., Eriksson, K., and
Stankiewicz, W. (2013). HSV-2 regulates monocyte inflammatory response via the
Fas/FasL pathway. *PLoS One, 8*(7), e70308. doi:10.1371/journal.pone.0070308

Kuwabara, T., Ishikawa, F., Kondo, M., and Kakiuchi, T. (2017). The role of IL-17 and related
cytokines in inflammatory autoimmune diseases. *Mediators Inflamm, 2017*, 3908061.
doi:10.1155/2017/3908061

Lambert, D. W., Yarski, M., Warner, F. J., Thornhill, P., Parkin, E. T., Smith, A. I., Hooper, N.
M., and Turner, A. J. (2005). Tumor necrosis factor-alpha convertase (ADAM17) mediates
regulated ectodomain shedding of the severe-acute respiratory syndrome-coronavirus
(SARS-CoV) receptor, angiotensin-converting enzyme-2 (ACE2). *J Biol Chem, 280*(34),
30113-30119. doi:10.1074/jbc.M505111200

Li, F. (2016). Structure, Function, and Evolution of Coronavirus Spike Proteins. *Annu Rev Virol,*
3(1), 237-261. doi:10.1146/annurev-virology-110615-042301

Liao, M., Liu, Y., Yuan, J., Wen, Y., Xu, G., Zhao, J., Cheng, L., Li, J., Wang, X., Wang, F., Liu,
L., Amit, I., Zhang, S., and Zhang, Z. (2020). Single-cell landscape of bronchoalveolar
immune cells in patients with COVID-19. *Nat Med, 26*(6), 842-844. doi:10.1038/s41591-
020-0901-9

Liu, X., Jana, M., Dasgupta, S., Koka, S., He, J., Wood, C., and Pahan, K. (2002). Human
immunodeficiency virus type 1 (HIV-1) tat induces nitric-oxide synthase in human
astroglia. *J Biol Chem, 277*(42), 39312-39319. doi:10.1074/jbc.M205107200

Lu, L., Zhang, H., Dauphars, D. J., and He, Y. W. (2021). A potential role of interleukin 10 in
COVID-19 pathogenesis. *Trends Immunol, 42*(1), 3-5. doi:10.1016/j.it.2020.10.012

Lu, R., Zhao, X., Li, J., Niu, P., Yang, B., Wu, H., Wang, W., Song, H., Huang, B., Zhu, N., Bi,
Y., Ma, X., Zhan, F., Wang, L., Hu, T., Zhou, H., Hu, Z., Zhou, W., Zhao, L., Chen, J.,
Meng, Y., Wang, J., Lin, Y., Yuan, J., Xie, Z., Ma, J., Liu, W. J., Wang, D., Xu, W.,
Holmes, E. C., Gao, G. F., Wu, G., Chen, W., Shi, W., and Tan, W. (2020). Genomic
characterisation and epidemiology of 2019 novel coronavirus: implications for virus

origins and receptor binding. *Lancet, 395*(10224), 565-574. doi:10.1016/S0140-6736(20)30251-8

Lugo-Villarino, G., Cougoule, C., Meunier, E., Rombouts, Y., Verollet, C., and Balboa, L. (2019). Editorial: The mononuclear phagocyte system in infectious disease. *Front Immunol, 10*, 1443. doi:10.3389/fimmu.2019.01443

Lum, F. M., Lee, D., Chua, T. K., Tan, J. J. L., Lee, C. Y. P., Liu, X., Fang, Y., Lee, B., Yee, W. X., Rickett, N. Y., Chia, P. Y., Lim, V., Leo, Y. S., Matthews, D. A., Hiscox, J. A., and Ng, L. F. P. (2018). Zika virus infection preferentially counterbalances human peripheral monocyte and/or NK cell activity. *mSphere, 3*(2). doi:10.1128/mSphereDirect.00120-18

Mendoza, E. J., Manguiat, K., Wood, H., and Drebot, M. (2020). Two detailed plaque assay protocols for the quantification of infectious SARS-CoV-2. *Curr Protoc Microbiol, 57*(1), ecpmc105. doi:10.1002/cpmc.105

Merad, M., and Martin, J. C. (2020). Pathological inflammation in patients with COVID-19: a key role for monocytes and macrophages. *Nat Rev Immunol, 20*(6), 355-362. doi:10.1038/s41577-020-0331-4

Mukadam, I. Z., Machhi, J., Herskovitz, J., Hasan, M., Oleynikov, M. D., Blomberg, W. R., Svechkarev, D., Mohs, A. M., Zhou, Y., Dash, P., McMillan, J., Gorantla, S., Garrison, J., Gendelman, H. E., and Kevadiya, B. D. (2020). Rilpivirine-associated aggregation-induced emission enables cell-based nanoparticle tracking. *Biomaterials, 231*, 119669. doi:10.1016/j.biomaterials.2019.119669

Nikitina, E., Larionova, I., Choinzonov, E., and Kzhyshkowska, J. (2018). Monocytes and macrophages as viral targets and reservoirs. *Int J Mol Sci, 19*(9). doi:10.3390/ijms19092821

Pedrosa, M. A., Valenzuela, R., Garrido-Gil, P., Labandeira, C. M., Navarro, G., Franco, R., Labandeira-Garcia, J. L., and Rodriguez-Perez, A. I. (2021). Experimental data using candesartan and captopril indicate no double-edged sword effect in COVID-19. *Clin Sci (Lond), 135*(3), 465-481. doi:10.1042/CS20201511

Perlman, S., and Netland, J. (2009). Coronaviruses post-SARS: update on replication and pathogenesis. *Nat Rev Microbiol, 7*(6), 439-450. doi:10.1038/nrmicro2147

Pichlmair, A., Schulz, O., Tan, C. P., Naslund, T. I., Liljestrom, P., Weber, F., and Reis e Sousa,
C. (2006). RIG-I-mediated antiviral responses to single-stranded RNA bearing 5'-
phosphates. *Science, 314*(5801), 997-1001. doi:10.1126/science.1132998

Ruan, Q., Yang, K., Wang, W., Jiang, L., and Song, J. (2020). Clinical predictors of mortality due
to COVID-19 based on an analysis of data of 150 patients from Wuhan, China. *Intensive*
Care Med, 46(5), 846-848. doi:10.1007/s00134-020-05991-x

Rubinstein, S., Familletti, P. C., and Pestka, S. (1981). Convenient assay for interferons. *J Virol,*
37(2), 755-758. doi:10.1128/JVI.37.2.755-758.1981

Salamanna, F., Maglio, M., Landini, M. P., and Fini, M. (2020). Body Localization of ACE-2: On
the Trail of the Keyhole of SARS-CoV-2. *Front Med (Lausanne), 7*, 594495.
doi:10.3389/fmed.2020.594495

Schmidtmayerova, H., Nottet, H. S., Nuovo, G., Raabe, T., Flanagan, C. R., Dubrovsky, L.,
Gendelman, H. E., Cerami, A., Bukrinsky, M., and Sherry, B. (1996). Human
immunodeficiency virus type 1 infection alters chemokine beta peptide expression in
human monocytes: implications for recruitment of leukocytes into brain and lymph nodes.
Proc Natl Acad Sci U S A, 93(2), 700-704. doi:10.1073/pnas.93.2.700

Shang, J., Wan, Y., Luo, C., Ye, G., Geng, Q., Auerbach, A., and Li, F. (2020). Cell entry
mechanisms of SARS-CoV-2. *Proc Natl Acad Sci U S A, 117*(21), 11727-11734.
doi:10.1073/pnas.2003138117

Soler, M. J., Ye, M., Wysocki, J., William, J., Lloveras, J., and Batlle, D. (2009). Localization of
ACE2 in the renal vasculature: amplification by angiotensin II type 1 receptor blockade
using telmisartan. *Am J Physiol Renal Physiol, 296*(2), F398-405.
doi:10.1152/ajprenal.90488.2008

Swiecki, M., and Colonna, M. (2011). Type I interferons: diversity of sources, production
pathways and effects on immune responses. *Curr Opin Virol, 1*(6), 463-475.
doi:10.1016/j.coviro.2011.10.026

Szklarczyk, D., Gable, A. L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., Simonovic, M.,
Doncheva, N. T., Morris, J. H., Bork, P., Jensen, L. J., and Mering, C. V. (2019). STRING
v11: protein-protein association networks with increased coverage, supporting functional
discovery in genome-wide experimental datasets. *Nucleic Acids Res, 47*(D1), D607-D613.
doi:10.1093/nar/gky1131

Wan, Y., Shang, J., Graham, R., Baric, R. S., and Li, F. (2020). Receptor recognition by the novel coronavirus from Wuhan: An analysis based on decade-long structural studies of SARS coronavirus. *J Virol, 94*(7). doi:10.1128/JVI.00127-20

World Health Organization. (2020). Coronavirus disease 2019 (COVID-19). Situation report – 51. Retrieved from https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200311-sitrep-51-covid-19.pdf?sfvrsn=1ba62e57_10

Wu, G., Dawson, E., Duong, A., Haw, R., and Stein, L. (2014). ReactomeFIViz: a Cytoscape app for pathway and network-based data analysis. *F1000Res, 3*, 146. doi:10.12688/f1000research.4431.2

Xia, H., Cao, Z., Xie, X., Zhang, X., Chen, J. Y., Wang, H., Menachery, V. D., Rajsbaum, R., and Shi, P. Y. (2020). Evasion of type I interferon by SARS-CoV-2. *Cell Rep, 33*(1), 108234. doi:10.1016/j.celrep.2020.108234

Xiong, Y., Liu, Y., Cao, L., Wang, D., Guo, M., Jiang, A., Guo, D., Hu, W., Yang, J., Tang, Z., Wu, H., Lin, Y., Zhang, M., Zhang, Q., Shi, M., Liu, Y., Zhou, Y., Lan, K., and Chen, Y. (2020). Transcriptomic characteristics of bronchoalveolar lavage fluid and peripheral blood mononuclear cells in COVID-19 patients. *Emerg Microbes Infect, 9*(1), 761-770. doi:10.1080/22221751.2020.1747363

Xu, Z., Shi, L., Wang, Y., Zhang, J., Huang, L., Zhang, C., Liu, S., Zhao, P., Liu, H., Zhu, L., Tai, Y., Bai, C., Gao, T., Song, J., Xia, P., Dong, J., Zhao, J., and Wang, F. S. (2020). Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *Lancet Respir Med, 8*(4), 420-422. doi:10.1016/S2213-2600(20)30076-X

Yang, D., Chu, H., Hou, Y., Chai, Y., Shuai, H., Lee, A. C., Zhang, X., Wang, Y., Hu, B., Huang, X., Yuen, T. T., Cai, J. P., Zhou, J., Yuan, S., Zhang, A. J., Chan, J. F., and Yuen, K. Y. (2020). Attenuated interferon and proinflammatory response in SARS-CoV-2-infected human dendritic cells is associated with viral antagonism of STAT1 phosphorylation. *J Infect Dis, 222*(5), 734-745. doi:10.1093/infdis/jiaa356

Yilla, M., Harcourt, B. H., Hickman, C. J., McGrew, M., Tamin, A., Goldsmith, C. S., Bellini, W. J., and Anderson, L. J. (2005). SARS-coronavirus replication in human peripheral monocytes/macrophages. *Virus Res, 107*(1), 93-101. doi:10.1016/j.virusres.2004.09.004

Zheng, J., Wang, Y., Li, K., Meyerholz, D. K., Allamargot, C., and Perlman, S. (2021). Severe acute respiratory syndrome coronavirus 2-induced immune activation and death of

monocyte-derived human macrophages and dendritic cells. *J Infect Dis, 223*(5), 785-795.

doi:10.1093/infdis/jiaa753

Zhou, J., Chu, H., Li, C., Wong, B. H., Cheng, Z. S., Poon, V. K., Sun, T., Lau, C. C., Wong, K.

K., Chan, J. Y., Chan, J. F., To, K. K., Chan, K. H., Zheng, B. J., and Yuen, K. Y. (2014).

Active replication of Middle East respiratory syndrome coronavirus and aberrant induction

of inflammatory cytokines and chemokines in human macrophages: implications for

pathogenesis. *J Infect Dis, 209*(9), 1331-1342. doi:10.1093/infdis/jit504

Supplementary information

Defining the Immune Responses for SARS-CoV-2-Human Macrophage Interactions

Mai M. Abdelmoaty[1,2], Pravin Yeapuri[1], Jatin Machhi[3], Katherine E. Olson[3], Farah Shahjin[3], You Zhou[4], Liang Jingjing[3], Kabita Pandey[3], Arpan Acharya[3], Siddappa N. Byrareddy[3], R. Lee Mosley[3] and Howard E. Gendelman[3*]

[1]Department of Pharmaceutical Sciences, College of Pharmacy, University of Nebraska Medical Center, NE 68198, USA

[2]Therapeutic Chemistry Department, Pharmaceutical and Drug Industries Research Division, National Research Centre, Giza, Egypt

[3]Department of Pharmacology and Experimental Neuroscience, College of Medicine, University of Nebraska Medical Center, NE 68198, USA

[4]Center for Biotechnology, University of Nebraska-Lincoln, Lincoln, NE, USA

Supplementary Figure 1. Fold changes of immune response genes in SARS-CoV-2-challenged MDMs. Heat map of fold changes in the expression of 84 genes specific for human innate and adaptive immune responses in SARS-CoV-2-challenged MDMs compared to mock-challenged MDMs at different time points after the infection, determined using RT2 Profiler Human Innate and Adaptive Immune Response 96-well Array. Fold changes were determined via Qiagen's RT2 Profiler analysis software (n=4).

Supplementary Figure 2. Differential proteomic analysis of SARS-CoV-2-challenged MDMs.

Volcano plots showing the fold change plotted against the P value highlighting significantly changed proteins (red – upregulation and green – downregulation; $p \leq 0.05$ and an absolute fold change ≥ 2) in SARS-CoV-2-challenged MDMs compared to mock-challenged MDMs at different time points (n=4). The vertical lines correspond to the absolute fold change of 2, and the horizontal line represents a p value of 0.05.