Phylogenomic evidence for the Origin of Obligately Anaerobic Anammox Bacteria around the Great Oxidation Event

Tianhua Liao[1][^], Sishuo Wang[1][^], Eva E. Stüeken[2], Haiwei Luo[1][*]

[1]Simon F. S. Li Marine Science Laboratory, School of Life Sciences and State Key Laboratory of Agrobiotechnology, The Chinese University of Hong Kong, Shatin, Hong Kong SAR

[2]School of Earth and Environmental Sciences and Centre for Exoplanet Science, University of St Andrews, Bute Building, Queen's Terrace, KY16 9TS, UK

[^]These authors contribute equally to this work.

[*]**Corresponding author:**
Haiwei Luo
The Chinese University of Hong Kong
Shatin, Hong Kong SAR
Phone: (+852) 39436121
E-mail: hluo2006@gmail.com

Running title: The origin time and phylogeny of anammox bacteria

Keywords: anammox bacteria, molecular dating analysis, Planctomycetes

Abstract

The anaerobic ammonium oxidation (anammox) bacteria could transform ammonium and nitrite to dinitrogen gas, and this obligate anaerobic process accounts for up to half of the global nitrogen loss in surface environments. Yet its origin and evolution, which may give important insights into the biogeochemistry of early Earth, remains enigmatic. Here, we performed comprehensive phylogenomic analysis and showed a single origin of anammox bacteria within the phylum Planctomycetes. After accommodating the uncertainties and factors influencing time estimates, which includes implementing both a traditional cyanobacteria-based and a recently developed mitochondria-based approach, we estimated that anammox bacteria originated at early Proterozoic and most likely around the so-called Great Oxidation Event (GOE; 2.32 to 2.5 billion years ago [Ga]) which fundamentally changed global biogeochemical cycles. We further showed that during the origin of anammox bacteria, genes involved in oxidative stress, bioenergetics and anammox granules formation were recruited, which might have contributed to their survival on an increasingly oxic Earth. Our findings suggest the rising levels of atmospheric oxygen, which made nitrite increasingly available, was a potential driving force for the emergence of anammox bacteria. This is one of the first studies that link the GOE to the evolution of obligate anaerobic bacteria.

Introduction

Anaerobic ammonium oxidation (anammox, $NH_4^+ + NO_2^- \rightarrow N_2 + 2H_2O$) [1], which

usually occurs in anoxic marine, freshwater and wetland settings, accounts for up to 50% of the

removal of fixed nitrogen (N) in nature. Along with denitrification, it is recognized as an

important biological process that leads to N loss from the environment [2, 3]. In wastewater

treatment, anammox is more cost-effective and environmentally friendlier than denitrification

due to its lower oxygen requirement for aeration (nitrite [N(+III)], instead of nitrate [N(+V)] is

sufficient for the anammox metabolism), carbon-free cultivation (some denitrifying bacteria are

heterotrophic) and its negligible emissions of greenhouse gases like N_2O [4, 5]. Consequently,

anammox bacteria, the organisms that perform anammox, have been widely used in wastewater

treatment plants [3, 6-8]. Despite the environmental and industrial importance of these organisms,

their evolutionary history and antiquity are poorly known, which hinders accurate

reconstructions of the biogeochemical nitrogen cycle over geologic time.

Previous studies investigated the roles of key genes driving anammox, including *hzsABC*

(hydrazine synthase)[9], *hdh* (hydrazine dehydrogenase), *hao* (hydroxylamine oxidoreductase)

[10] and *nxr* (nitrite oxidoreductase) [11]. In general, the vital enzymes encoded by these genes

either are directly involved in anammox or participate in replenishing electrons to the cyclic

electron flow [12, 13]. Further, all known anammox bacteria are found within the phylum

Planctomycetes [14]. Six candidate genera of anammox bacteria, namely '*Candidatus* Brocadia',

'Candidatus Kuenenia', 'Candidatus Jettenia', 'Candidatus Scalindua', 'Candidatus

Anammoximicrobium' and 'Candidatus Anammoxoglobus', have been proposed based on 16S

ribosomal RNA (rRNA) gene sequences [15], but none of them have been successfully isolated

into pure cultures. The habitat of anammox bacteria requires the simultaneous presence of

reduced (ammonia) and oxidized (nitrite) inorganic N compounds. Such habitats are often found

at the aerobic-anaerobic interface in aquatic ecosystems, including the margins of oxygen

minimum zones (OMZs) in the ocean and sediment-water interfaces, where ammonium

originates from the anaerobic degradation of organic matter and nitrite can be produced by

aerobic ammonia oxidation [16]. Ammonium, one of the two substrates of the anammox

metabolism, was thermodynamically stable and likely present in moderate concentrations in the

deep ocean throughout the Archean [17] and probably extending well into the Proterozoic [18].

The availability of nitrite, the second important substrate, is less certain. Geochemical data from

sedimentary rocks suggest that the early Earth (before 3 Ga) was fully anoxic and deficient in

aerobic ecosystems that were able to generate nitrite/nitrate [19-21]. The first transient and/or

localized occurrences of aerobic nitrogen cycling appear in the Mid- to Neoarchean (2.7 Ga

[22-25]), but evidence of widespread nitrate availability does not appear until around 2.4 billion

years ago (Ga), the so-called Great Oxidation Event (GOE) [26-28]. The GOE marks the rise of

free O_2 in the atmosphere above 10^{-5} times modern levels, and was ultimately a result of the

emergence of oxygenic cyanobacteria [29]. Hence it is conceivable that the appearance of

anammox was linked to the appearance of aerobic ecosystems around the time of the GOE.

However, it is also possible that some nitrite was provided much earlier, through lightning

reactions in the Archean atmosphere [30]. Lightning, even in the absence of O_2, can generate NO

gas, which dissolves in water and converts into aqueous species, including nitrite [31]. Some

anammox bacteria are even capable of using NO rather than nitrite directly as a substrate [12]. If

these organisms capitalized on the lightning flux, then anammox might long pre-date the GOE.

Several tools exist for investigating the link between the evolution of metabolic pathways

and geo-environmental transformations. One way to explore the evolutionary history of a

specific metabolic pathway is based on organic biomarkers, which, however, are often affected

by the poor preservation over geologic timescales [32]. For instance, ladderanes, a type of lipids

that is unique to anammox bacteria [33], are rarely preserved to a level that can be used to date

their evolutionary origin. Another approach is based on nitrogen isotope ratios of sedimentary

records [17, 34, 35]. However, the isotopic fractionation factors for different metabolic pathways

overlap widely (denitrification: -5 to -30‰; anammox: -16 or -24‰ [18, 36]), making it difficult

to single out and elucidate the evolution of the different redox reactions within the N cycle by

this method [37]. Alternatively, molecular dating, which estimates the age of the last common

ancestor (LCA) of analyzed lineages by comparing their sequences based on the molecular clock

theory [38], provides an alternative strategy to investigate this issue. Briefly, it can estimate the

divergence timescale of organisms using genetic data while accounting for issues like

uncertainties in the calibrations and different evolutionary rates among lineages, thereby

bypassing the paucity and uncertainty of biomarkers and other biogeochemical records. On the

one hand, because anammox is an anaerobic reaction, it is intuitive to assume that anammox

should originate before the GOE, perhaps using nitrite or NO provided by lightning reactions in

the atmosphere [39]. On the other hand, it is because of GOE that O_2 on Earth rose permanently

to a concentration that is biologically meaningful, and as a result, (micro-)environments were

created in the surface ocean that were rich in nitrate/nitrite produced by aerobic organisms. These

redox-stratified conditions may have stimulated the anammox metabolism by providing both

nitrite and ammonium at the redoxcline. In this regard, it could be hypothesized that anammox

did not arise until the GOE. Here, we ask the question when anammox originated. To answer this

question and to obtain more insights into the evolution of anammox bacteria, we compiled an

up-to-date genomic data set of Planctomycetes (see Supplementary Text section 1; Dataset S1.1),

placed the evolution of anammox bacteria into the context of geological events, and investigated

genomic changes associated with the origin of this ecologically important bacterial lineage.

Results and discussion

Monophyletic origins of anammox bacteria and anammox genes

The anammox bacteria form a monophyletic group within the phylum Planctomycetes in

a comprehensive phylogenomic tree with 881 Planctomycetes genomes (Fig. S1). The anammox

bacteria clade in this phylogenomic tree (Fig. S1) comprises four known genera including the

early-branching *Ca.* Scalindua and *Ca.* Kuenenia, and the late-branching *Ca.* Jettenia and *Ca.*

Brocadia. It also includes two understudied lineages which we named 'basal lineage' [40] and

'*hzsCBA*-less lineage' (Fig. S1), both of which have relatively low genome completeness

compared to the other anammox bacteria and are represented by metagenome-assembled

genomes (MAGs) sampled from groundwater (see Supplementary Text section 2.1). Note that

the absence of *hzsCBA* in several genomes within the '*hzsCBA*-less lineage' might be ascribed to

the loss of these genes in evolution or their low genome completeness (Fig. S1). Because of their

shallow (late-branching) phylogenetic position, this uncertainty should not affect the inference of

the origin of anammox bacteria. Two described genera (*Ca.* Anammoximicrobium and *Ca.*

Anammoxoglobus) do not have genomes available by the time of the present study (last accessed:

April 2021), and are therefore not included in the phylogenomic analysis. Furthermore, using the

last updated set of 2,077 Planctomycetes genomes (retrieved in April 2021) from NCBI, we

obtained a consistent topology of anammox bacteria (Fig. S2).

To check whether important groups, particularly early-branching lineages, have been

encompassed in our analyzed genome data sets, we built a 16S rRNA gene tree (Fig. S3) using

the identified 16S rRNA genes from genomic sequences of anammox bacteria and the deposited

16S rRNA gene amplicons in SILVA from the class Brocadiae, to which anammox bacteria

belong (see Supplementary Texts section 2.2). Since the anammox bacteria with genome

sequences in this 16S rRNA gene tree (Fig. S3) showed a branching order congruent with the

topology of the phylogenomic tree (Fig. S1), and since the 16S rRNA gene tree included 913

(clustered from 20,142) sequences sampled from a wide array of habitats (marine, sediments,

man-made reactor, freshwater and terrestrial ecosystems), the 16S rRNA gene phylogeny, which

could better capture the diversity of anammox bacteria by including uncultured samples, likely

encompasses the early-branching lineages of anammox bacteria. Taken together, our genome set,

which encompasses the genomes of the early-branching lineages, is appropriate to be used to

study the evolutionary origin of anammox bacteria (Dataset S1.3). We further analyzed the

evolution (Fig. S4) of the genes (*hzsCBA*) critical to the anammox reaction [10, 41].

Reconciliation (see Supplementary Texts section 2.3) of the gene and the genome-based species

phylogenies pointed to a single origin of anammox metabolism at the LCA of anammox bacteria

(Fig. S4).

Taken together, the above analyses suggest that our genome data sets have encompassed

the earliest-branching lineages and deep phylogenetic diversity of anammox bacteria, and

indicate its monophyletic origin, which agrees with previous studies [42-44]. This should provide

a foundation for estimating the origin of anammox metabolism by estimating the origin time of

anammox bacteria.

<u>The evolutionary origin of anammox bacteria coincided with the rising O_2</u>

We employed MCMCTree to estimate the divergence times of the anammox bacteria with

i) soft bounds (time constraint allowing a small probability of violation) based on cyanobacteria

fossils, and ii) the relaxed molecular clock approaches allowing substitution rates to vary among

branches. We first employed a cyanobacteria fossil-based strategy, where cyanobacteria fossils

were used as the only calibrations, a widely used strategy in dating deep-time bacterial evolution

[45-47]. Using a best-practice dating scheme (Fig. 1; see Supplementary Text section 3), we

dated the origin of anammox bacteria to 2,117 million years ago (Ma) (95% highest posterior

density [HPD] interval, 2,002 - 2,226 Ma). Further, we repeated the analysis based on the

expanded Planctomycetes genomes (Genome set 2; Dataset S1.1) and the constraint topology

inferred with a profile mixture model (LG+C60+G; Genome set 1; Dataset S1.1). In general,

both analyses based on different genome sets resolved consistent topologies (Fig. S5) as shown

in Figure 1. Overall, they gave similar posterior times of the LCA of the anammox bacteria at

2,105 Ma (95% HPD: 1,961 - 2,235 Ma) or 2,005 Ma (95% HPD: 1,869 - 2,146 Ma).

Analyses with different combinations of calibrations and parameters broadly converge to

similar time estimates (Fig. S6; Dataset S2.1). Specifically, assigning a more ancient time

constraint to the total group of oxygenic cyanobacteria, which is based on the biogeochemical

evidence of the presence of O_2 at nearly 3.0 Ga [48] instead of GOE (Dataset S2.1), shifted the

posterior dates of the LCA of anammox bacteria into the past by ~300 million years (Myr) (e.g.,

C1 versus C10; Fig. S6). Varying the maximum bounds of the root by using 4.5 Ga, 3.8 Ga or 3.5

Ga had a minor impact (around 100 Myr) on the time estimates of anammox bacteria (e.g., C1

versus C3; Fig. S6). Though the auto-correlated rates (AR) model was less favored than the

independent rates (IR) model based on the model comparison (see Supplementary Text section

3.3; Dataset S2.2), it is worth noting that the posterior ages estimated by the AR model were 1-12%

younger than estimated by the IR model (Dataset S2.1).

While cyanobacteria fossils are the most widely used calibrations (and in many times the

only calibrations) in bacterial molecular dating studies [47, 49, 50], there are two important

issues: i) the available cyanobacterial fossils are very rare, ii) cyanobacteria fossils do not

provide appropriate maximum calibrations [51]. Hence, we also used a recently developed

mitochondria-based dating strategy [51] to estimate the origin time of anammox bacteria. This

approach, based on mitochondria endosymbiosis, integrates mitochondrial lineages as a sister

group to Alphaproteobacteria, thus taking the advantages of eukaryotic fossils, particularly

several maximum calibrations, to improve date estimates. We compiled two gene sets (see

Dataset S1.4), one according to the original 24 mitochondria-encoded genes used in [51] (mito24;

6295 sites), the other being six (mito6; 1238 sites) out of these 24 genes conserved across the

bacterial tree of life (see Fig. 2A and Supplementary Text section 3.5). These analyses showed

that the posterior time of the LCA of anammox bacteria shifted to the past by 90 to 500 Myr (Fig.

2), compared with only using the cyanobacteria-fossil-based approach (Fig. 1). The difference

between calibrations C1 and C8 resulted from the removal of the maximum of the Node2 in C8,

and accordingly, the posterior time estimated by C8 was greatly shifted to the past (Fig. 2B).

When all maximum time constraints except for the one imposed on the root were removed, the

posterior time was mainly affected by changes in the root maximum bound (C8 & C10; Fig. 2B).

Despite the large phylogenetic distance between mitochondrial lineages and anammox bacteria,

the mitochondria-based dating strategy may provide additional time constraints to the origin of

anammox bacteria. For example, the 'C8+Euk' scheme effectively constrained the time estimates

compared using a single maximum bound (C8; Fig. 2B). Furthermore, the posterior dates

obtained by incorporating eukaryotic fossils ('C8+euk' to 'C10+euk') displayed less variation

across calibration schemes (C8 to C10) based on cyanobacterial fossils (Fig. 2B). Nevertheless,

the broadly consistent time estimates between analyses using cyanobacterial and mitochondrial

calibrations highlight an origin of anammox bacteria roughly around or shortly after the GOE.

Besides, there remains a possibility that there are unsampled or even extinct lineages of

anammox bacteria that diverged earlier than all anammox bacteria analyzed in the present study.

This scenario, if true, hints that the first anammox bacterium could have originated before the

occurrence of the LCA of sequenced anammox bacteria but later than their split from the sister

non-anammox lineage (Node X in Fig. 1; up to 2,600 Ma). Actually, a pre-GOE origin of

anammox bacteria was also obtained by applying a larger minimum age (3.0 Ga) [52-54] to the

total group of oxygenic cyanobacteria based on only cyanobacteria fossils (Fig. S5), or by using

the mitochondria-based strategy (Fig. 2B). Accommodating the above uncertainties, our analyses

suggest that the origin of anammox bacteria, and hence the origin of anammox, most likely falls

into the 2.6-2.0 Ga interval, or more generally, around the early Proterozoic. Running

MCMCTree analysis with no sequence data showed different distributions of time estimates for

both the clade of anammox bacteria and the four calibration points (Fig. S7), suggesting that

sequence data are informative for our molecular clock analysis [55]. Besides, we estimated

similar origin times of anammox at around 2.1 Ga (the AR model) or 2.3 Ga (the IR model; see

Fig. S9) with the complete taxon sampling shown in Fig. S2 (Genome set 2; see Dataset S1.1).

For this analysis, we used a likelihood-based dating algorithm since the Bayesian molecular

clock analysis is known to be computationally intensive and not suitable to take a data set with a

broad taxon sampling [56]. Note that unlike Bayesian molecular dating software, the

likelihood-based method we used only makes point estimations for the node ages (no standard

deviation), thereby not allowing for fully appreciating the uncertainties in age estimates and any

inference based on that. In general, the above result suggests that taxon sampling may not affect

our dating analysis.

The geochemical context of the origin of anammox bacteria

We speculate that the timing of the origin of the anammox metabolism is linked to the

increasing availability of nitrite in surface environments, because ammonium was likely present

in the deep anoxic ocean throughout the Precambrian [17, 18] and therefore presumably not a

limiting substrate. The first abiotic source of nitrogen oxides on the early Earth, including NO

and nitrite, would have been lightning reactions between N_2 and CO_2 in the Archean atmosphere

[57, 58]. A prior study suggested that this process led to micromolar levels of nitrite in seawater

[57], and the supply of NO may have been even higher, considering that it is the initial reaction

product of lighting [57, 58]. This implies the possibility of an earlier origin of anammox bacteria.

In fact, an intriguing hypothesis is that NO-dependent anammox [12] is ancestral to all anammox

bacteria and arose before the appearance of significant nitrite levels with the GOE, perhaps

driven by the lightning-source of NO. However, the original estimates of the lightning-derived

flux of nitrogen oxides may have been based on perhaps unrealistically high amounts of CO_2

[59]. Furthermore, to our knowledge, there has been no isotopic evidence in the early Archean

rock record prior to 3 Ga for the presence of a significant nitrite or nitrate reservoir in the early

ocean [19-21], and it is relatively conceivable that any nitrogen oxides that were supplied to the

ocean by lightning were rapidly reduced to ammonium or N_2 by ferrous iron, possibly even

abiotically [60, 61]. Hence, although a background lightning flux of nitrogen oxides almost

certainly existed, to our knowledge, there seems little evidence that it could supply a reliable

metabolic substrate for early life. Furthermore, it is important to note that our phylogenetic

results are not supportive of this hypothesis since the NO-utilizing anammox bacteria

discovered to date are affiliated with phylogenetically shallow lineages from *Ca.* Kuenenia and

Ca. Brocadia (Fig. 1). To our knowledge, there have been no studies reporting the ability of

using NO as the sole electron acceptor in any other lineages of anammox bacteria. Hence, if

NO-dependent anammox would have been present in the LCA of anammox bacteria, it was

not clear why it was lost in all other anammox bacteria which resulted from multiple

independent losses of the trait, a scenario apparently not favored from an evolutionary

perspective (as shown by the arrows in Fig. S2). There are also anammox bacteria that are

capable of using hydroxylamine as an oxidant; however, this molecule is a highly reactive

compound and is usually considered as an intermediate in the nitrogen cycle, thereby rendering it

a less important compound from the perspective of geological timescales. We are not aware of

any non-biological sources of hydroxylamine. Taken together, our results are inconsistent with an

Archean emergence of the anammox metabolism driven by lightning-derived nitrogen oxides.

Instead, our results are most parsimoniously explained by the appearance of more significant

nitrite/nitrate reservoirs around the time of the GOE.

The first transient appearances of nitrite/nitrate-dependent metabolisms are captured by

the sedimentary nitrogen isotope record in late Mesoarchean soils at 3.2 Ga [23] and in

Neoarchean shallow-marine settings at 2.7 Ga and 2.5 Ga [22, 24, 25]. These observations may

reflect local and/or temporally restricted oxygen oases [62, 63]. Widespread nitrite/nitrate

availability is inferred for the Paleoproterozoic (2.4-1.8 Ga), i.e., in the immediate aftermath of

the GOE [26-28]. Importantly, O_2 is necessary, although a trace amount is feasible, to extant

nitrifying organisms for the oxidation of ammonium to nitrite and nitrate [64-66]. We noticed

that a recent study [67] reports the ability of the model ammonia-oxidizing archaeal species

(*Nitrosopumilus maritimus*) to continue ammonia oxidation after consuming all supplied O_2 by

nitric oxide disproportionation to generate O_2, but the underlying genetic pathway is not known,

making it difficult to extrapolate this mechanism to any other ammonia-oxidizing organisms.

Hence the rise of oxygen almost certainly triggered the growth of the nitrite/nitrate reservoir, and

therefore potentially provided one of the key substrates for anammox bacteria. We note that also

NO would have become more abundant after the GOE, because it is an intermediate product in

nitrification and denitrification [68]. This would put the possibility of NO-driven anammox back

on the table. However, NO levels in seawater are highly variable, and so nitrite would likely have

been a more reliable substrate. In any case, it would not violate our conclusion that the

appearance of anammox in the Paleoproterozoic was linked to the GOE and directly or indirectly

to the rise of marine nitrite. It has been shown that a low concentration of nitrite significantly

decreases the rate of N removal by anammox bacteria in reactors [69, 70]. In modern OMZ in the

Bay of Bengal, which may to some extent serve as analogues for the Paleoproterozoic

redox-stratified ocean, the concentration of nitrite, instead of ammonium, was proposed as the

rate-limiting factor to the anammox metabolism [71]. These imply that the rise of nitrite, which

is driven by the rise of O_2, could have facilitated the appearance of anammox bacteria in the

early Proterozoic.

In modern environments, the majority of nitrite used in anammox is derived from nitrate

reduction or aerobic ammonia oxidation [72], the latter performed by ammonia-oxidizing

archaea (AOA) or bacteria (AOB). As noted above, nitrification could have appeared by at least

2.7 Ga, as indicated by nitrogen-isotope evidence for local accumulation of nitrate in surface

ocean waters [18, 73, 74]. A recent study inferred that the LCA of AOA dates back to ~2.3 Ga

and first appeared on land, driven by increasing O_2 concentrations in the atmosphere at that time,

and that the expansion of AOA from land to the ocean did not occur until nearly 1.0 Ga [75]. If

true, this hints at a dominant role of AOB in early ocean, before 2.3 Ga. Consistent with this idea,

the vast majority of the early-branching lineages of anammox bacteria (*Ca.* Scalindua) and the

sister Planctomycetes lineages of the anammox clade in the phylogenomic tree (Fig. 1) are found

in marine or groundwater habitats. Furthermore, in the 16S rRNA gene tree which represents a

greater diversity than the phylogenomic tree, marine lineages still account for the majority of the

earliest-branching lineages of anammox bacteria (Fig. S3). Thus, it seems likely that the LCA of

anammox bacteria originated in the marine realm where AOB thrived first. The above arguments,

although speculative, tentatively suggest that the nitrite demand for anammox had been readily

met by GOE by taking advantage of the significant amount of O_2 newly available, which is

broadly consistent with our molecular dating results suggesting an origin of anammox shortly

before or after the GOE.

Genomic changes potentially related to anammox metabolism and habitat adaptation upon the origin of anammox bacteria

We further explored the genomic changes characterizing the origin of anammox bacteria (see Supplementary Text section 4). This includes the gains of the aforementioned genes that directly participate in anammox, viz., *hzsABC*, *hdh*, and *hao*. The current view of the anammox metabolism posits that electrons consumed by anammox for carbon fixation are replenished by the oxidation of nitrite to nitrate by nitrite oxidoreductase (NXR) [13], which was inferred to be acquired upon the origin of anammox bacteria (Fig. 3). Moreover, multiple auxiliary genes for N assimilatory pathways including N regulatory protein (*glnB*), assimilatory nitrite reductase large subunit (*nirB*), transporters like *amt* (ammonium), *nirC* (nitrite) and NRT family proteins (nitrate/nitrite), were also gained at the origin of anammox bacteria (Fig. 3). However, genes for nitrogen-related dissimilatory pathways encoding nitrite reduction to nitric oxide (*nirK* or *nirS*) and to ammonium (*nrfAH*), respectively, were likely acquired after the origin of anammox bacteria (Fig. 3). Anammox occurs at the membrane of anammoxosome, an organelle predominantly composed of a special type of lipid called ladderane. The ladderane membranes show low proton permeability, which helps maintain the proton motive force during the anammox metabolism [33]. Although the biosynthetic pathway of ladderane is yet to be characterized, a previous study [76] predicted 34 candidate genes responsible for the synthesis of ladderane lipids, four of which were potentially acquired during the origin of anammox bacteria

(see Supplementary Text section 4.2; Fig. 3), hinting at their potential roles in ladderane

synthesis and providing clues to future experimental investigations.

Anammox bacteria occur in anoxic habitats and exhibit low oxygen tolerance [15].

Specifically, the O_2-sensitive intermediates generated by anammox, e.g., hydrazine, a powerful

reductant, requires anoxic conditions. Accordingly, the likely acquired peroxidases include

cytochrome c peroxidase (Ccp), which could scavenge hydrogen peroxide in the periplasm [77],

and the most prevalent peroxidase [78], thioredoxin-dependent peroxiredoxin (AhpC). Besides

peroxide scavengers, the desulfoferrodoxin (Dfx), which functions as superoxide reductase (SOR)

to reduce superoxide to hydrogen peroxide and which is broadly distributed among anaerobic

bacteria [78], were likely acquired during the origin of anammox bacteria (Fig. 3). Another

acquired gene is *fprA*, which encodes flavo-diiron proteins that scavenge O_2.

Investigating other metabolisms that were also acquired upon the origin of anammox

bacteria allows reconstructing the coeval ecology. Our data show that *sat* (Fig. 3), which encodes

sulfate adenylyltransferase for assimilatory incorporation of sulfate into bioavailable adenylyl

sulfate, as well as *aprAB* and *fsr* (Fig. 3) for encoding dissimilatory sulfite reductase, were

acquired upon the origin of anammox bacteria (Fig. 3). This acquired capability of sulfate

metabolism is coincident with the increased concentrations of seawater sulfate from ~100 μM

or less throughout much of the Archean [79, 80] to over 1mM after the rise of atmospheric O_2

[81, 82]. Unlike other Planctomycetes, anammox bacteria are generally autotrophs that use the

Wood-Ljungdahl pathway for carbon assimilation [83]. Consequently, they potentially lost many

genes involved in carbohydrate utilization (Fig. 3). The metabolic loss is further strengthened by

the enrichment analysis where the genes predicted to be lost by the LCA of anammox bacteria

are enriched in pathways involving hydrolase activity, intramolecular oxidoreductase activity and

carbohydrate kinase activity (Fig. 3; see Data and Code availability), hinting at a decreased

ability of anammox bacteria to degrade organics.

Additionally, iron is a vital element for HZS [9] and HDH [84], and *in vitro* studies

revealed that increased iron concentrations can promote the growth of anammox bacteria [85].

The ancient oceans are thought to have been ferruginous (high iron availability) according to

sedimentary records of iron speciation across several coeval marine settings from the Archean

through most of the Proterozoic [86], meaning that large amounts of soluble iron would have

been available for anammox bacteria. Interestingly, a series of iron-related genes to make use of

the abundant iron from the environment were likely acquired upon the origin of anammox

bacteria. First, anammox bacteria likely acquired the gene *fur*, which encodes a ferric uptake

regulator for iron assimilation. Moreover, the iron-containing proteins, specifically cytochrome,

are encoded by the acquired genes *petBC*, which potentially encodes Rieske-heme b [13], and

CcsAB, which encodes a protein involved in cytochrome c maturation systems (Fig. 3). Notably,

ahbABCD, which encodes proteins-synthesizing *b*-type hemes in anammox bacteria [87], were

likely acquired before the origin of anammox bacteria (Fig. 3). Furthermore, we identified that

the gene *Bfr* that encodes the oligomeric protein, bacterioferritin involved in the uptake and

storage of iron [88] was likely acquired during the origin of anammox bacteria (Fig. 3), which

may help to hoard iron in settings where iron became locally sparse, such as along euxinic

(sulfide-rich) or oxic marine margins in the Paleoproterozoic [89]. A recent study highlights the

vital role of bacterioferritin to the anammox regulation [90]. It is thought that the appearance of

euxinic margins around the time of the GOE trapped iron that upwelled from the deep ocean,

leading to the demise of iron oxide deposits (i.e., banded iron formations) [91]. Our results

tentatively support this model, if the microenvironments that those primordial anammox bacteria

colonized were located near the euxinic-oxic interface where nitrite and ammonium were

available while upwelled iron was titrated out of the water column by freely dissolved hydrogen

sulfide.

Taken together, the timelines and corresponding genomic changes of anammox bacteria

provide implications for the physiological characteristics of descendant anammox bacteria and

the origin of other nitrogen-transforming pathways in the context of an estimated evolutionary

timeline of anammox bacteria. For example, the earlier branching lineage *Ca.* Scalindua exhibit

high nitrite affinity (0.45 μM) compared to other anammox genera (up to 370 μM) [15, 92],

which may reflect a progressive growth of the marine nitrite reservoir, linked to the protracted

oxygenation of the surface environments over geologic timescales [29]. Likewise, genomic data

reveal that also canonical denitrification radiated across the tree of life after the GOE [93], and

the gains (Fig. S10; see Supplementary Text section 4.4) of genes *nrfAH* for dissimilatory nitrate

reduction to ammonium (DNRA) at the late-branching lineages (*Ca.* Brocadia and *Ca.* Kuenenia)

at around 800 Ma are consistent with the time when the nitrite/nitrate availability was increased

by the Neoproterozoic Oxygenation Event [94]. The time estimate and genomic changes of the

anammox bacteria have gone some way towards enhancing the understanding of the

physiological characteristics of modern anammox bacteria and the historical N cycle.

Caveats and concluding remarks

In the present study, we link an estimated evolutionary timeline of anammox and

geological context of early Earth on a simplified view that nitrite-dependent anammox is

ancestral to anammox bacteria. However, there are several limitations. First, dating the

bacterial evolution has many challenges, which, among others, include the paucity of

appropriate fossils, and a long evolutionary distance between anammox bacteria and

cyanobacteria [51]. Hence, the date estimates should be read as potential ranges of time

(indicated by the 95% HPD interval), instead of accurate time points, under the molecular

clock model. Moreover, it is worth pointing out the metabolism called feammox, a process

where ferric instead of nitrite is taken as the electron acceptor for ammonia oxidation. Both

feammox and anammox can anaerobically convert ammonia into dinitrogen [95], but the former

process is supposed to be carried out by Acidimicrobiaceae from the Actinobacteria phylum [96,

97] and under anaerobic environments without the presence of anammox bacteria [98]. Likewise,

a recent experimental study [99] proposed extant anammox bacteria as electroactive organisms

and suggested the feasibility to utilize graphene oxides and man-made metallic electrodes

mimicking metal oxides as the electron acceptor. However, the lack of detailed molecular

mechanism hampers the inference of its evolutionary history. In any case, metal oxides such as

Fe_2O_3 or MnO_2 also became more abundant in the Neoarchean to Paleoproterozoic, i.e. around

the time of the GOE [91, 100], and hence even if these oxides were used as substrates for

anammox, it would not violate our overall conclusion that the GOE played a primary role in

driving the evolution of this metabolic pathway. However, it is also worth noting that metal

oxides are solids and thus require extra-cellular electron transfer [99]. An entirely intra-cellular

metabolism, using dissolved nitrite, may be more likely to evolve first, especially given that

nitrate became bioavailable in the ocean around the same time (see above).

Another important aspect to highlight about our approach is that, our analysis began with

inferring the LCA of anammox bacteria based on the presence of *hzsCBA*, a common strategy in

phylogenetic studies. It is important to note that such an enzyme complex encoded by multiple

genes might originate in different genomic backgrounds and were presented together in a suitable

genomic background later in evolution. This scenario, if true, indicates that the appearance of

any single subunit of *hzsCBA* genes, which could not assembly a functional hydrazine synthase,

should predate the LCA of anammox bacteria. Nevertheless, the alternative hypothesis would not

affect our inference and following analysis since we only focus on the inferred ancestor of extant

anammox bacteria, a strategy commonly used in modern phylogenetic analysis. These alternative

possibilities, although not affecting the molecular clock analysis, should be carefully considered

when making inference based on our estimated evolutionary timeline of anammox bacteria.

Here, using molecular dating and comparative genomics approaches, we link the

emergence of an obligate anaerobic bacterial group, which drives the loss of fixed N in many

environments, to the rise of O_2, and highlight their evolutionary responses to major

environmental disturbances. Apparently, the GOE opened novel niches for the origin and

subsequent expansion of diverse aerobic prokaryotes such as AOA and AOB, which in turn

facilitated the origin of other bacterial lineages, including some anaerobes like the anammox

bacteria, by providing the resources for their energy conservation. Our results open up the

possibility that a significant proportion of the Proterozoic nitrite budget was consumed by

anammox bacteria, and that the sedimentary nitrogen isotope record might be influenced by their

activity. Our findings may have implications for Proterozoic climate, because N_2O has

previously been invoked as an important greenhouse gas at that time [101], but a relatively

higher contribution of anammox bacteria to the marine nitrogen cycle may have hampered N_2O

production [6]. Lastly, our study suggests that the impacts of the GOE go well beyond aerobic

prokaryotes. We therefore conclude that molecular dating is a likely feasible approach to

complement isotopic evidence for resolving the timeline of biological evolution and for

providing additional constraints on climate models of the distant past.

Materials and Methods

Overall, we compiled two genome sets and one 16S rRNA gene set in our study (Dataset

S1.1). A phylogenomic tree (Fig. S1) was generated based on the concatenated alignment of 120

ubiquitous proteins (bac120; Dataset S1.3) proposed for tree inference by the Genome Taxonomy

Database [102]. All published genomic sequences (952 in total) affiliated with the phylum

Planctomycetes in NCBI Genbank up to December 2019 were retrieved (Dataset S1.2). To

further examine whether the patterns obtained with this Planctomycetes dataset hold the same,

we further constructed an expanded set of Planctomycetes genomes by retrieving a total of 2,077

Planctomycetes genomes released by April 2021 at Genbank (Fig. S2; see Data and Code

availability). Key anammox genes *hzsCBA* and *hdh* were identified against manually curated

reference sequences by BLASTP. For each gene, identified protein sequences were aligned using

MAFFT (v7.222) [103] and the alignments were refined by trimAl (v1.4) [104]. All phylogenies

were constructed by IQ-tree (v1.6.2) [105] with substitution models automatically selected by

ModelFinder [106] and branch support assessed with 1,000 ultrafast bootstrap replicates. Note

that the constraint topologies for dating analysis (Fig. 1; Fig. S5) were inferred with the profile

mixture model (LG+C60+G) which better accommodates across-site heterogeneity in deep-time

evolution. Following manual curation, four well-recognized genera and two separate lineages of

anammox bacteria were highlighted with different colours, and the presence of key genes were

annotated with filled symbols beside the labels (Fig. S1). Furthermore, the phylogenomic tree

was generated with similar methods using 16S rRNA genes identified from downloaded genomes

and those retrieved from the SILVA database (Fig. S3; Dataset S1.3). All trees (including

phylogenomic and 16S trees) in our study were visualized with iTOL [107]. More details are

shown in supplementary text, section 2.

Molecular dating analysis was carried out using the program MCMCTree from the PAML

package (4.9j) [108]. In our study, 13 calibration sets constructed with different time constraints

of the root and three calibration nodes within the cyanobacteria lineage were used (see

Supplemental Texts section 3.2). The topology constraint for dating analysis was generated using

85 genomes, which were sampled from the phylogenomic tree of phylum Planctomycetes (see

Supplemental Texts section 3.1), and the model LG+C20+F+G under posterior mean site

frequency (PMSF) approximation [109]. To perform dating analysis, clock models and different

calibration sets were compared in 26 schemes (see Supplemental Texts section 3; Dataset S2.1)

[110, 111]. For each scheme, the approximate likelihood method [112] of MCMCTree were

conducted in duplicate with identical iteration parameters (burn-in: 10,000; sample frequency: 20;

number of sample: 20,000). The convergence of each scheme was evaluated by comparing the

posterior dates of two independent runs (see Data and Code availability). With the updated

genome set 2 (Dataset S1.1), we repeated the taxon sampling process and generated an

alternative phylogeny for dating analysis (Fig. S5) using LG+C20+F+G model (C20: 20 classes

of site-specific amino acid profiles) under PMSF approximations. Similarly, we also generated

another phylogeny for dating analysis using the same genome set as used in Fig. 1 but with the

profile mixture model (LG+C60+G) which uses 60 classes of amino acid profiles [113]. Note

that the missing parameter +F would not significantly change the results since it only adds

another profile amino acid calculated from the original data. These two repeat dating analyses

were conducted with the best practice dating scheme (IR model and calibration C1). Moreover,

we performed likelihood based dating analyses [114] (Fig. S9) using the phylogenomic tree (Fig.

S2) and the calibration C1 (see Data and Code availability) implemented in ape [56].

The phylogenetic differences between gene trees and the species tree were reconciliated

by GeneRax [115] with unrooted gene tree as input and automatically optimized duplication,

transfer, loss (DTL) rates. For each gene identified from the genome set 2 (Dataset S1.1), we

used a species tree comprised by all anammox bacteria pruned from the phylogenomic tree (Fig.

S2) as the reference. We used recommended parameters including SPR strategy, undated DTL

model and a maximum radius of five.

For comparative genomics analysis, the protein-coding sequences were annotated against

KEGG [116], CDD [117], InterPro [118], Pfam [119], TIGRFAM [120] and TCDB [121],

individually (see Supplemental Texts section 4). Following annotations, the potentially

gained/lost genes between groups (anammox bacteria versus non-anammox bacteria; an

anammox bacterial genus versus all other anammox bacteria) were statistically tested using

two-sided fisher's exact tests. Finally, the resulting p-values were corrected with the

Benjamini-Hochberg FDR (false discovery rate) procedure. Moreover, genes with corrected

p-values smaller than 0.05 and with larger ratio in the study group were defined as gained genes,

and genes with corrected p-values smaller than 0.05 and with smaller ratio in the study group

were defined as potentially lost genes. Those potentially gained/lost genes were summarized for

their detailed functions (see Data and Code availability).

Acknowledgements

We thank Bite Pei for building the basic dataset for this project, Yang Qian for his help in

editing the earlier version of the manuscript, Jinjin Tao for her helpful discussion, and Mario dos

Reis for his suggestion on dating analysis. This work is funded by the National Science

Foundation of China (92051113), the Hong Kong Research Grants Council Area of Excellence

Scheme (AoE/M-403/16), the Direct Grant of CUHK (4053495), and The CUHK Impact

Postdoctoral Fellowship Scheme to (S.W.).

Data and Code availability

513   All used genomic sequences, generated phylogenetic trees, estimated divergence times

514 and the used python codes are deposited in the online repository

515 https://github.com/luolab-cuhk/anammox-origin.

517 **Conflict of interest**

518   The authors declare that they have no conflict of interest.

520 **References**

521 1 Broda, E. 1977 Two kinds of lithotrophs missing in nature. *Z Allg Mikrobiol*. **17**, 491-493.
522 (10.1002/jobm.3630170611)
523 2 Oshiki, M., Satoh, H., Okabe, S. 2016 Ecology and physiology of anaerobic ammonium
524 oxidizing bacteria. *Environ Microbiol*. **18**, 2784-2796. (10.1111/1462-2920.13134)
525 3 Stein, L. Y., Klotz, M. G. 2016 The nitrogen cycle. *Current Biology*. **26**, R94-R98. (DOI
526 10.1016/j.cub.2015.12.021)
527 4 Van Dongen, U., Jetten, M. S., Van Loosdrecht, M. 2001 The SHARON®-Anammox® process
528 for treatment of ammonium rich wastewater. *Water science and technology*. **44**, 153-160.
529 5 Jetten, M. S. M., Wagner, M., Fuerst, J., van Loosdrecht, M., Kuenen, G., Strous, M. 2001
530 Microbiology and application of the anaerobic ammonium oxidation ('anammox') process.
531 *Current Opinion in Biotechnology*. **12**, 283-288. (Doi 10.1016/S0958-1669(00)00211-1)
532 6 Li, L., Ling, Y., Wang, H. Y., Chu, Z. S., Yan, G. K., Li, Z. W., Wu, T. 2020 N_2O emission in
533 partial nitritation-anammox process. *Chinese Chemical Letters*. **31**, 28-38.
534 (10.1016/j.cclet.2019.06.035)
535 7 Okabe, S., Oshiki, M., Takahashi, Y., Satoh, H. 2011 N2O emission from a partial
536 nitrification-anammox process and identification of a key biological process of N2O emission
537 from anammox granules. *Water Research*. **45**, 6461-6470. (10.1016/j.watres.2011.09.040)
538 8 Gao, D. W., Tao, Y. 2011 Versatility and application of anaerobic ammonium-oxidizing bacteria.
539 *Applied Microbiology and Biotechnology*. **91**, 887-894. (10.1007/s00253-011-3411-3)

9 Dietl, A., Ferousi, C., Maalcke, W. J., Menzel, A., de Vries, S., Keltjens, J. T., Jetten, M. S.,

Kartal, B., Barends, T. R. 2015 The inner workings of the hydrazine synthase multiprotein

complex. *Nature*. **527**, 394-397.   (10.1038/nature15517)

10 Kartal, B., Maalcke, W. J., de Almeida, N. M., Cirpus, I., Gloerich, J., Geerts, W., Op den

Camp, H. J., Harhangi, H. R., Janssen-Megens, E. M., Francoijs, K. J.*, et al.* 2011 Molecular

mechanism of anaerobic ammonium oxidation. *Nature*. **479**, 127-130.   (10.1038/nature10453)

11 Chicano, T. M., Dietrich, L., de Almeida, N. M., Akram, M., Hartmann, E., Leidreiter, F.,

Leopoldus, D., Mueller, M., Sanchez, R., Nuijten, G. H. L.*., et al.* 2021 Structural and functional

characterization of the intracellular filament-forming nitrite oxidoreductase multiprotein complex.

Nature Microbiology.   (10.1038/s41564-021-00934-8)

12 Hu, Z., Wessels, H., van Alen, T., Jetten, M. S. M., Kartal, B. 2019 Nitric oxide-dependent

anaerobic ammonium oxidation. *Nat Commun*. **10**, 1244.   (10.1038/s41467-019-09268-w)

13 Kartal, B., Keltjens, J. T. 2016 Anammox Biochemistry: a Tale of Heme c Proteins. *Trends*

Biochem Sci. **41**, 998-1011.   (10.1016/j.tibs.2016.08.015)

14 Wang, Y., Xu, L. Y., Wang, S. Y., Ye, F., Zhu, G. B. 2019 Global Distribution of Anaerobic

Ammonia Oxidation (Anammox) Bacteria - Field Surveys in Wetland, Dryland, Groundwater

Aquifer and Snow. *Frontiers in Microbiology*. **10**,   (10.3389/fmicb.2019.02583)

15 Zhang, L., Okabe, S. 2020 Ecological niche differentiation among anammox bacteria. *Water*

Res. **171**, 115468.   (10.1016/j.watres.2020.115468)

16 Sonthiphand, P., Hall, M. W., Neufeld, J. D. 2014 Biogeography of anaerobic

ammonia-oxidizing (anammox) bacteria. *Front Microbiol*. **5**, 399.   (10.3389/fmicb.2014.00399)

17 Yang, J., Junium, C. K., Grassineau, N. V., Nisbet, E. G., Izon, G., Mettam, C., Martin, A.,

Zerkle, A. L. 2019 Ammonium availability in the Late Archaean nitrogen cycle. *Nature*

Geoscience. **12**, 553-557.   (10.1038/s41561-019-0371-1)

18 Stueken, E. E., Kipp, M. A., Koehler, M. C., Buick, R. 2016 The evolution of Earth's

biogeochemical nitrogen cycle. *Earth-Science Reviews*. **160**, 220-239.

(10.1016/j.earscirev.2016.07.007)

19 Ossa Ossa, F., Hofmann, A., Spangenberg, J. E., Poulton, S. W., Stueken, E. E., Schoenberg,

R., Eickmann, B., Wille, M., Butler, M., Bekker, A. 2019 Limited oxygen production in the

Mesoarchean ocean. *Proc Natl Acad Sci U S A*. **116**, 6647-6652.   (10.1073/pnas.1818762116)

20 Stueken, E. E., Buick, R., Guy, B. M., Koehler, M. C. 2015 Isotopic evidence for biological

nitrogen fixation by molybdenum-nitrogenase from 3.2 Gyr. *Nature*. **520**, 666-669.

(10.1038/nature14180)

21 Koehler, M. C., Buick, R., Barley, M. E. 2019 Nitrogen isotope evidence for anoxic deep

marine environments from the Mesoarchean Mosquito Creek Formation, Australia. *Precambrian*

Research. **320**, 281-290.   (10.1016/j.precamres.2018.11.008)

22 Koehler, M. C., Buick, R., Kipp, M. A., Stueken, E. E., Zaloumis, J. 2018 Transient surface
ocean oxygenation recorded in the similar to 2.66-Ga Jeerinah Formation, Australia. *Proceedings*
of the National Academy of Sciences of the United States of America. **115**, 7711-7716.
(10.1073/pnas.1720820115)
23 Homann, M., Sansjofre, P., Van Zuilen, M., Heubeck, C., Gong, J., Killingsworth, B., Foster, I.
S., Airo, A., Van Kranendonk, M. J., Ader, M.*, et al.* 2018 Microbial life and biogeochemical
cycling on land 3,220 million years ago. *Nature Geoscience*. **11**, 665-+.
(10.1038/s41561-018-0190-9)
24 Godfrey, L. V., Falkowski, P. G. 2009 The cycling and redox state of nitrogen in the Archaean
ocean. *Nature Geoscience*. **2**, 725-729.   (10.1038/Ngeo633)
25 Garvin, J., Buick, R., Anbar, A. D., Arnold, G. L., Kaufman, A. J. 2009 Isotopic Evidence for
an Aerobic Nitrogen Cycle in the Latest Archean. *Science*. **323**, 1045-1048.
(10.1126/science.1165675)
26 Zerkle, A. L., Poulton, S. W., Newton, R. J., Mettam, C., Claire, M. W., Bekker, A., Junium, C.
K. 2017 Onset of the aerobic nitrogen cycle during the Great Oxidation Event. *Nature*. **542**,
465-+.   (10.1038/nature20826)
27 Luo, G. M., Junium, C. K., Izon, G., Ono, S., Beukes, N. J., Algeo, T. J., Cui, Y., Xie, S. C.,
Summons, R. E. 2018 Nitrogen fixation sustained productivity in the wake of the
Palaeoproterozoic Great Oxygenation Event. *Nature Communications*. **9**,
(10.1038/s41467-018-03361-2)
28 Kipp, M. A., Stueken, E. E., Yun, M., Bekker, A., Buick, R. 2018 Pervasive aerobic nitrogen
cycling in the surface ocean across the Paleoproterozoic Era. *Earth and Planetary Science*
Letters. **500**, 117-126.   (10.1016/j.epsl.2018.08.007)
29 Lyons, T. W., Reinhard, C. T., Planavsky, N. J. 2014 The rise of oxygen in Earth's early ocean
and atmosphere. *Nature*. **506**, 307-315.   (10.1038/nature13068)
30 Navarro-González, R., Molina, M. J., Molina, L. T. 1998 Nitrogen fixation by volcanic
lightning in the early Earth. *Geophysical Research Letters*. **25**, 3123-3126.
(https://doi.org/10.1029/98GL02423)
31 Kasting, J. F., Walker, J. C. G. 1981 Limits on oxygen concentration in the prebiological
atmosphere and the rate of abiotic fixation of nitrogen. *Journal of Geophysical Research:*
Oceans. **86**, 1147-1158.   (https://doi.org/10.1029/JC086iC02p01147)
32 Hallmann, C., Kelly, A. E., Gupta, S. N., Summons, R. E. 2011 Reconstructing Deep-Time
Biology with Molecular Fossils. *Quantifying the Evolution of Early Life : Numerical Approaches*
to the Evaluation of Fossils and Ancient Ecosystems. **36**, 355-401.
(10.1007/978-94-007-0680-4_14)
33 Moss, F. R., Shuken, S. R., Mercer, J. A. M., Cohen, C. M., Weiss, T. M., Boxer, S. G., Burns,
N. Z. 2018 Ladderane phospholipids form a densely packed membrane with normal hydrazine

and anomalously low proton/hydroxide permeability. *Proceedings of the National Academy of Sciences of the United States of America*. **115**, 9098-9103.   (10.1073/pnas.1810706115)

Mettam, C., Zerkle, A. L., Claire, M. W., Prave, A. R., Poulton, S. W., Junium, C. K. 2019 Anaerobic nitrogen cycling on a Neoarchaean ocean margin. *Earth and Planetary Science Letters*. **527**,   (10.1016/j.epsl.2019.115800)

Ader, M., Thomazo, C., Sansjofre, P., Busigny, V., Papineau, D., Laffont, R., Cartigny, P., Halverson, G. P. 2016 Interpretation of the nitrogen isotopic composition of Precambrian sedimentary rocks: Assumptions and perspectives. *Chemical Geology*. **429**, 93-110. (10.1016/j.chemgeo.2016.02.010)

Brunner, B., Contreras, S., Lehmann, M. F., Matantseva, O., Rollog, M., Kalvelage, T., Klockgether, G., Lavik, G., Jetten, M. S., Kartal, B.*, et al.* 2013 Nitrogen isotope effects induced by anammox bacteria. *Proc Natl Acad Sci U S A*. **110**, 18994-18999. (10.1073/pnas.1310488110)

Fuchsman, C. A., Stueken, E. E. 2021 Using modern low-oxygen marine ecosystems to understand the nitrogen cycle of the Paleo- and Mesoproterozoic oceans. *Environ Microbiol*. **23**, 2801-2822.   (10.1111/1462-2920.15220)

Yang, Z., Rannala, B. 2006 Bayesian estimation of species divergence times under a molecular clock using multiple fossil calibrations with soft bounds. *Mol Biol Evol*. **23**, 212-226. (10.1093/molbev/msj024)

Ducluzeau, A. L., van Lis, R., Duval, S., Schoepp-Cothenet, B., Russell, M. J., Nitschke, W. 2009 Was nitric oxide the first deep electron sink? *Trends in Biochemical Sciences*. **34**, 9-15. (10.1016/j.tibs.2008.10.005)

Suarez, C., Dalcin Martins, P., Jetten, M., Karacic, S., Wilen, B. M., Modin, O., Hagelia, P., Hermansson, M., Persson, F. 2022 Metagenomic evidence of a novel family of anammox bacteria in a subsea environment. *Environ Microbiol*.   (10.1111/1462-2920.16006)

Harhangi, H. R., Le Roy, M., van Alen, T., Hu, B. L., Groen, J., Kartal, B., Tringe, S. G., Quan, Z. X., Jetten, M. S., Op den Camp, H. J. 2012 Hydrazine synthase, a unique phylomarker with which to study the presence and biodiversity of anammox bacteria. *Appl Environ Microbiol*. **78**, 752-758.   (10.1128/AEM.07113-11)

Hamasaki, K., Shishikura, R., Suzuki, S., Shiozaki, T., Ogawa, H., Nakamura, T., Suwa, Y. 2018 Distribution and phylogeny of anaerobic ammonium-oxidizing (anammox) bacteria in the water column of the central Pacific Ocean. *Deep-Sea Research Part Ii-Topical Studies in Oceanography*. **156**, 60-67.   (10.1016/j.dsr2.2017.11.013)

Terada, A., Zhou, S., Hosomi, M. 2011 Presence and detection of anaerobic ammonium-oxidizing (anammox) bacteria and appraisal of anammox process for high-strength nitrogenous wastewater treatment: a review. *Clean Technologies and Environmental Policy*. **13**, 759-781.   (10.1007/s10098-011-0355-3)

Schmid, M., Walsh, K., Webb, R., Rijpstra, W. I. C., van de Pas-Schoonen, K., Verbruggen, M. J., Hill, T., Moffett, B., Fuerst, J., Schouten, S., *et al.* 2003 Candidatus "Scalindua brodae", sp nov., Candidatus "Scalindua wagneri", sp nov., two new species of anaerobic ammonium oxidizing bacteria. *Systematic and Applied Microbiology*. **26**, 529-538.   (Doi 10.1078/072320203770865837)

Sanchez-Baracaldo, P., Raven, J. A., Pisani, D., Knoll, A. H. 2017 Early photosynthetic eukaryotes inhabited low-salinity habitats. *Proc Natl Acad Sci U S A*. **114**, E7737-E7745. (10.1073/pnas.1620089114)

Zhang, H., Sun, Y., Zeng, Q., Crowe, S. A., Luo, H. 2021 Snowball Earths, population bottlenecks, and the evolution of marine photosynthetic bacteria. *bioRxiv*. 2020.2011.2024.395392.   (10.1101/2020.11.24.395392)

Wang, S. S., Meade, A., Lam, H. M., Luo, H. W. 2020 Evolutionary Timeline and Genomic Plasticity Underlying the Lifestyle Diversity in Rhizobiales. *Msystems*. **5**,   (ARTN e00438-20 10.1128/mSystems.00438-20)

Betts, H. C., Puttick, M. N., Clark, J. W., Williams, T. A., Donoghue, P. C. J., Pisani, D. 2018 Integrated genomic and fossil evidence illuminates life's early evolution and eukaryote origin. *Nat Ecol Evol*. **2**, 1556-1562.   (10.1038/s41559-018-0644-x)

Battistuzzi, F. U., Hedges, S. B. 2009 A Major Clade of Prokaryotes with Ancient Adaptations to Life on Land. *Molecular Biology and Evolution*. **26**, 335-343.   (10.1093/molbev/msn247)

Lin, W., Paterson, G. A., Zhu, Q., Wang, Y., Kopylova, E., Li, Y., Knight, R., Bazylinski, D. A., Zhu, R., Kirschvink, J. L., *et al.* 2017 Origin of microbial biomineralization and magnetotaxis during the Archean. *Proc Natl Acad Sci U S A*. **114**, 2171-2176. (10.1073/pnas.1614654114)

Wang, S., Luo, H. 2021 Dating Alphaproteobacteria evolution with eukaryotic fossils. *Nature Communications*. **12**, 3324.   (10.1038/s41467-021-23645-4)

Fournier, G. P., Moore, K. R., Rangel, L. T., Payette, J. G., Momper, L., Bosak, T. 2021 The Archean origin of oxygenic photosynthesis and extant cyanobacterial lineages. *Proceedings of the Royal Society B-Biological Sciences*. **288**,   (ARTN 20210675 10.1098/rspb.2021.0675)

Cardona, T. 2019 Thinking twice about the evolution of photosynthesis. *Open biology*. **9**, 180246.

Crowe, S. A., Dossing, L. N., Beukes, N. J., Bau, M., Kruger, S. J., Frei, R., Canfield, D. E. 2013 Atmospheric oxygenation three billion years ago. *Nature*. **501**, 535-+. (10.1038/nature12426)

Roger, A. J., Susko, E. 2018 Molecular clocks provide little information to date methanogenic Archaea. *Nat Ecol Evol*. **2**, 1676-1677.   (10.1038/s41559-018-0687-z)

56 Paradis, E., Schliep, K. 2019 ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. *Bioinformatics*. **35**, 526-528.   (10.1093/bioinformatics/bty633)

57 Wong, M. L., Charnay, B. D., Gao, P., Yung, Y. L., Russell, M. J. 2017 Nitrogen Oxides in

Early Earth's Atmosphere as Electron Acceptors for Life's Emergence. *Astrobiology*. **17**, 975-983.

(10.1089/ast.2016.1473)

58 Kasting, J. F., Walker, J. C. G. 1981 Limits on Oxygen Concentration in the Prebiological

Atmosphere and the Rate of Abiotic Fixation of Nitrogen. *Journal of Geophysical*

Research-Oceans. **86**, 1147-1158.   (DOI 10.1029/JC086iC02p01147)

59 Charnay, B., Le Hir, G., Fluteau, F., Forget, F., Catling, D. C. 2017 A warm or a cold early

Earth? New insights from a 3-D climate-carbon model. *Earth and Planetary Science Letters*. **474**,

97-109.

60 Brandes, J. A., Boctor, N. Z., Cody, G. D., Cooper, B. A., Hazen, R. M., Yoder, H. S. 1998

Abiotic nitrogen reduction on the early Earth. *Nature*. **395**, 365-367.   (Doi 10.1038/26450)

61 Summers, D. P., Chang, S. 1993 Prebiotic Ammonia from Reduction of Nitrite by Iron(Ii) on

the Early Earth. *Nature*. **365**, 630-632.   (DOI 10.1038/365630a0)

62 Lalonde, S. V., Konhauser, K. O. 2015 Benthic perspective on Earth's oldest evidence for

oxygenic photosynthesis. *Proceedings of the National Academy of Sciences of the United States*

of America. **112**, 995-1000.   (10.1073/pnas.1415718112)

63 Anbar, A. D., Duan, Y., Lyons, T. W., Arnold, G. L., Kendall, B., Creaser, R. A., Kaufman, A.

J., Gordon, G. W., Scott, C., Garvin, J.*, et al.* 2007 A whiff of oxygen before the Great Oxidation

Event? *Science*. **317**, 1903-1906.   (10.1126/science.1140325)

64 Ranjan, S., Todd, Z. R., Rimmer, P. B., Sasselov, D. D., Babbin, A. R. 2019 Nitrogen Oxide

Concentrations in Natural Waters on Early Earth. *Geochemistry Geophysics Geosystems*. **20**,

2021-2039.   (10.1029/2018gc008082)

65 Cheng, C., Busigny, V., Ader, M., Thomazo, C., Chaduteau, C., Philippot, P. 2019 Nitrogen

isotope evidence for stepwise oxygenation of the ocean during the Great Oxidation Event.

Geochimica Et Cosmochimica Acta. **261**, 224-247.   (10.1016/j.gca.2019.07.011)

66 Lehtovirta-Morley, L. E. 2018 Ammonia oxidation: Ecology, physiology, biochemistry and

why they must all come together. *FEMS Microbiol Lett*. **365**,   (10.1093/femsle/fny058)

67 Kraft, B., Jehmlich, N., Larsen, M., Bristow, L. A., Könneke, M., Thamdrup, B., Canfield, D.

E. 2022 Oxygen and nitrogen production by an ammonia-oxidizing archaeon. *Science*. **375**,

97-100.   (doi:10.1126/science.abe6733)

68 Stanton, C. L., Reinhard, C. T., Kasting, J. F., Ostrom, N. E., Haslun, J. A., Lyons, T. W.,

Glass, J. B. 2018 Nitrous oxide from chemodenitrification: A possible missing link in the

Proterozoic greenhouse and the evolution of aerobic respiration. *Geobiology*. **16**, 597-609.

(10.1111/gbi.12311)

69 Li, W., Zhuang, J. L., Zhou, Y. Y., Meng, F. G., Kang, D., Zheng, P., Shapleigh, J. P., Liu, Y. D.
2020 Metagenomics reveals microbial community differences lead to differential nitrate
production in anammox reactors with differing nitrogen loading rates. *Water Res*. **169**, 115279.
(10.1016/j.watres.2019.115279)
70 Zhang, L., Narita, Y., Gao, L., Ali, M., Oshiki, M., Ishii, S., Okabe, S. 2017 Microbial
competition among anammox bacteria in nitrite-limited bioreactors. *Water Res*. **125**, 249-258.
(10.1016/j.watres.2017.08.052)
71 Bristow, L. A., Callbeck, C. M., Larsen, M., Altabet, M. A., Dekaezemacker, J., Forth, M.,
Gauns, M., Glud, R. N., Kuypers, M. M. M., Lavik, G.*, et al.* 2017 N-2 production rates limited
by nitrite availability in the Bay of Bengal oxygen minimum zone. *Nature Geoscience*. **10**, 24-29.
(10.1038/Ngeo2847)
72 Dalsgaard, T., Thamdrup, B., Farias, L., Revsbech, N. P. 2012 Anammox and denitrification
in the oxygen minimum zone of the eastern South Pacific. *Limnology and Oceanography*. **57**,
1331-1346. (10.4319/lo.2012.57.5.1331)
73 Canfield, D. E., Glazer, A. N., Falkowski, P. G. 2010 The evolution and future of Earth's
nitrogen cycle. *Science*. **330**, 192-196. (10.1126/science.1186120)
74 Zerkle, A. L., Poulton, S. W., Newton, R. J., Mettam, C., Claire, M. W., Bekker, A., Junium, C.
K. 2017 Onset of the aerobic nitrogen cycle during the Great Oxidation Event. *Nature*. **542**,
465-467. (10.1038/nature20826)
75 Ren, M., Feng, X., Huang, Y., Wang, H., Hu, Z., Clingenpeel, S., Swan, B. K., Fonseca, M.
M., Posada, D., Stepanauskas, R.*, et al.* 2019 Phylogenomics suggests oxygen availability as a
driving force in Thaumarchaeota evolution. *ISME J*. **13**, 2150-2161.
(10.1038/s41396-019-0418-8)
76 Rattray, J. E., Strous, M., Op den Camp, H. J., Schouten, S., Jetten, M. S., Damste, J. S. 2009
A comparative genomics study of genetic products potentially encoding ladderane lipid
biosynthesis. *Biol Direct*. **4**, 8. (10.1186/1745-6150-4-8)
77 Van Vliet, A. H., Ketley, J. M., Park, S. F., Penn, C. W. 2002 The role of iron in
Campylobacter gene regulation, metabolism and oxidative stress defense. *FEMS Microbiol Rev*.
26, 173-186. (10.1016/s0168-6445(02)00095-5)
78 Johnson, L. A., Hug, L. A. 2019 Distribution of reactive oxygen species defense mechanisms
across domain bacteria. *Free Radical Biology and Medicine*. **140**, 93-102.
(10.1016/j.freeradbiomed.2019.03.032)
79 Crowe, S. A., Paris, G., Katsev, S., Jones, C., Kim, S. T., Zerkle, A. L., Nomosatryo, S.,
Fowle, D. A., Adkins, J. F., Sessions, A. L.*, et al.* 2014 Sulfate was a trace constituent of Archean
seawater. *Science*. **346**, 735-739. (10.1126/science.1258966)

80 Jamieson, J. W., Wing, B. A., Farquhar, J., Hannington, M. D. 2013 Neoarchaean seawater
sulphate concentrations from sulphur isotopes in massive sulphide ore. *Nature Geoscience*. **6**,
61-64.   (10.1038/Ngeo1647)
81 Blattler, C. L., Claire, M. W., Prave, A. R., Kirsimae, K., Higgins, J. A., Medvedev, P. V.,
Romashkin, A. E., Rychanchik, D. V., Zerkle, A. L., Paiste, K*., et al.* 2018 Two-billion-year-old
evaporites capture Earth's great oxidation. *Science*. **360**, 320-+.   (10.1126/science.aar2687)
82 Fakhraee, M., Hancisse, O., Canfield, D. E., Crowe, S. A., Katsev, S. 2019 Proterozoic
seawater sulfate scarcity and the evolution of ocean-atmosphere chemistry. *Nature Geoscience*.
12, 375-380.   (10.1038/s41561-019-0351-5)
83 Kartal, B., Keltjens, J. T., Jetten, M. S. M. 2011 Metabolism and Genomics of Anammox
Bacteria. *Nitrification*. 181-200.
84 Akram, M., Dietl, A., Mersdorf, U., Prinz, S., Maalcke, W., Keltjens, J., Ferousi, C., de
Almeida, N. M., Reimann, J., Kartal, B*., et al.* 2019 A 192-heme electron transfer network in the
hydrazine dehydrogenase complex. *Sci Adv*. **5**, eaav4310.   (10.1126/sciadv.aav4310)
85 Bi, Z., Qiao, S., Zhou, J., Tang, X., Zhang, J. 2014 Fast start-up of Anammox process with
appropriate ferrous iron concentration. *Bioresour Technol*. **170**, 506-512.
(10.1016/j.biortech.2014.07.106)
86 Poulton, S. W., Canfield, D. E. 2011 Ferruginous Conditions: A Dominant Feature of the
Ocean through Earth's History. *Elements*. **7**, 107-112.   (10.2113/gselements.7.2.107)
87 Ferousi, C., Lindhoud, S., Baymann, F., Kartal, B., Jetten, M. S., Reimann, J. 2017 Iron
assimilation and utilization in anaerobic ammonium oxidizing bacteria. *Curr Opin Chem Biol*. **37**,
129-136.   (10.1016/j.cbpa.2017.03.009)
88 Keren, N., Aurora, R., Pakrasi, H. B. 2004 Critical roles of bacterioferritins in iron storage
and proliferation of cyanobacteria. *Plant Physiol*. **135**, 1666-1673.   (10.1104/pp.104.042770)
89 Poulton, S. W., Fralick, P. W., Canfield, D. E. 2010 Spatial variability in oceanic redox
structure 1.8 billion years ago. *Nature Geoscience*. **3**, 486-490.   (10.1038/Ngeo889)
90 Peng, M.-W., Qi, J., Yan, P., Guan, Y., Liu, Y.-Y., Sun, Z.-H., Zhang, L.-J., Weng, X., Shen, Y.,
Fang, F*., et al.* 2022 Insight into the structure and metabolic function of iron-rich nanoparticles in
anammox bacteria. *Science of The Total Environment*. **806**, 150879.
(https://doi.org/10.1016/j.scitotenv.2021.150879)
91 Konhauser, K. O., Planaysky, N. J., Hardisty, D. S., Robbins, L. J., Warchola, T. J., Haugaard,
R., Lalonde, S. V., Partin, C. A., Oonk, P. B. H., Tsikos, H*., et al.* 2017 Iron formations: A global
record of Neoarchaean to Palaeoproterozoic environmental history. *Earth-Science Reviews*. **172**,
140-177.   (10.1016/j.earscirev.2017.06.012)
92 Wu, Z., Meng, H., Huang, X., Wang, Q., Chen, W. H., Gu, J. D., Lee, P. H. 2019
Salinity-driven heterogeneity toward anammox distribution and growth kinetics. *Appl Microbiol*
Biotechnol. **103**, 1953-1960.   (10.1007/s00253-018-9521-4)

93 Parsons, C., Stueken, E. E., Rosen, C. J., Mateos, K., Anderson, R. E. 2021 Radiation of
nitrogen-metabolizing enzymes across the tree of life tracks environmental transitions in Earth
history. *Geobiology*. **19**, 18-34.   (10.1111/gbi.12419)
94 Ader, M., Sansjofre, P., Halverson, G. P., Busigny, V., Trindade, R. I. F., Kunzmann, M.,
Nogueira, A. C. R. 2014 Ocean redox structure across the Late Neoproterozoic Oxygenation
Event: A nitrogen isotope perspective. *Earth and Planetary Science Letters*. **396**, 1-13.
(10.1016/j.epsl.2014.03.042)
95 Li, X., Huang, Y., Liu, H. W., Wu, C., Bi, W., Yuan, Y., Liu, X. 2018 Simultaneous Fe(III)
reduction and ammonia oxidation process in Anammox sludge. *Journal of Environmental*
Sciences. **64**, 42-50.   (10.1016/j.jes.2017.01.002)
96 Zhu, J., Li, T., Liao, C., Li, N., Wang, X. 2021 A promising destiny for Feammox: From
biogeochemical ammonium oxidation to wastewater treatment. *Science of The Total Environment*.
790, 148038.   (https://doi.org/10.1016/j.scitotenv.2021.148038)
97 Wan, L., Liu, H., Wang, X. 2021 Anaerobic ammonium oxidation coupled to Fe(III) reduction:
Discovery, mechanism and application prospects in wastewater treatment. *Science of The Total*
Environment. 151687.   (https://doi.org/10.1016/j.scitotenv.2021.151687)
98 Yang, Y., Xiao, C., Yu, Q., Zhao, Z., Zhang, Y. 2021 Using Fe(II)/Fe(III) as catalyst to drive a
novel anammox process with no need of anammox bacteria. *Water Research*. **189**, 116626.
(https://doi.org/10.1016/j.watres.2020.116626)
99 Shaw, D. R., Ali, M., Katuri, K. P., Gralnick, J. A., Reimann, J., Mesman, R., van Niftrik, L.,
Jetten, M. S. M., Saikaly, P. E. 2020 Extracellular electron transfer-dependent anaerobic
oxidation of ammonium by anammox bacteria. *Nature Communications*. **11**,
(10.1038/s41467-020-16016-y)
100 Johnson, J. E., Webb, S. M., Ma, C., Fischer, W. W. 2016 Manganese mineralogy and
diagenesis in the sedimentary rock record. *Geochimica et Cosmochimica Acta*. **173**, 210-231.
(https://doi.org/10.1016/j.gca.2015.10.027)
101 Buick, R. 2007 Did the Proterozoic 'Canfield Ocean' cause a laughing gas greenhouse?
Geobiology. **5**, 97-100.   (10.1111/j.1472-4669.2007.00110.x)
102 Parks, D. H., Rinke, C., Chuvochina, M., Chaumeil, P. A., Woodcroft, B. J., Evans, P. N.,
Hugenholtz, P., Tyson, G. W. 2017 Recovery of nearly 8,000 metagenome-assembled genomes
substantially expands the tree of life. *Nat Microbiol*. **2**, 1533-1542.
(10.1038/s41564-017-0012-7)
103 Katoh, K., Standley, D. M. 2013 MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol Biol Evol*. **30**, 772-780.
(10.1093/molbev/mst010)

104 Capella-Gutierrez, S., Silla-Martinez, J. M., Gabaldon, T. 2009 trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*. **25**, 1972-1973.
(10.1093/bioinformatics/btp348)
105 Nguyen, L. T., Schmidt, H. A., von Haeseler, A., Minh, B. Q. 2015 IQ-TREE: a fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol*.
32, 268-274.   (10.1093/molbev/msu300)
106 Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A., Jermiin, L. S. 2017
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods*. **14**,
587-589.   (10.1038/nmeth.4285)
107 Letunic, I., Bork, P. 2019 Interactive Tree Of Life (iTOL) v4: recent updates and new
developments. *Nucleic Acids Research*. **47**, W256-W259.
108 Yang, Z. H. 2007 PAML 4: Phylogenetic analysis by maximum likelihood. *Molecular*
Biology and Evolution. **24**, 1586-1591.
109 Wang, H.-C., Minh, B. Q., Susko, E., Roger, A. J. 2018 Modeling site heterogeneity with
posterior mean site frequency profiles accelerates accurate phylogenomic estimation. *Systematic*
biology. **67**, 216-235.
110 Reis, M. D., Gunnell, G. F., Barba-Montoya, J., Wilkins, A., Yang, Z., Yoder, A. D. 2018
Using Phylogenomic Data to Explore the Effects of Relaxed Clocks and Calibration Strategies
on Divergence Time Estimation: Primates as a Test Case. *Syst Biol*. **67**, 594-615.
(10.1093/sysbio/syy001)
111 Dos Reis, M., Donoghue, P. C., Yang, Z. 2016 Bayesian molecular clock dating of species
divergences in the genomics era. *Nat Rev Genet*. **17**, 71-80.   (10.1038/nrg.2015.8)
112 Dos Reis, M., Yang, Z. 2011 Approximate likelihood calculation on a phylogeny for
Bayesian estimation of divergence times. *Mol Biol Evol*. **28**, 2161-2172.
(10.1093/molbev/msr045)
113 Quang, L. S., Gascuel, O., Lartillot, N. 2008 Empirical profile mixture models for
phylogenetic reconstruction. *Bioinformatics*. **24**, 2317-2323.   (10.1093/bioinformatics/btn445)
114 Sanderson, M. J. 2002 Estimating absolute rates of molecular evolution and divergence
times: A penalized likelihood approach. *Molecular Biology and Evolution*. **19**, 101-109.   (DOI
10.1093/oxfordjournals.molbev.a003974)
115 Morel, B., Kozlov, A. M., Stamatakis, A., Szollosi, G. J. 2020 GeneRax: A tool for
Species-Tree-Aware Maximum Likelihood-based gene family tree inference under gene
duplication, transfer, and loss. *Mol Biol Evol*. **37**, 2763-2774.   (10.1093/molbev/msaa141)
116 Kanehisa, M., Goto, S. 2000 KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic*
Acids Research. **28**, 27-30.

117 Lu, S., Wang, J., Chitsaz, F., Derbyshire, M. K., Geer, R. C., Gonzales, N. R., Gwadz, M.,
Hurwitz, D. I., Marchler, G. H., Song, J. S.*, et al.* 2020 CDD/SPARCLE: the conserved domain
database in 2020. *Nucleic Acids Res*. **48**, D265-D268.   (10.1093/nar/gkz991)
118 Mitchell, A. L., Attwood, T. K., Babbitt, P. C., Blum, M., Bork, P., Bridge, A., Brown, S. D.,
Chang, H. Y., El-Gebali, S., Fraser, M. I.*, et al.* 2019 InterPro in 2019: improving coverage,
classification and access to protein sequence annotations. *Nucleic Acids Res*. **47**, D351-D360.
(10.1093/nar/gky1100)
119 El-Gebali, S., Mistry, J., Bateman, A., Eddy, S. R., Luciani, A., Potter, S. C., Qureshi, M.,
Richardson, L. J., Salazar, G. A., Smart, A.*, et al.* 2019 The Pfam protein families database in
2019. *Nucleic Acids Res*. **47**, D427-D432.   (10.1093/nar/gky995)
120 Haft, D. H., Loftus, B. J., Richardson, D. L., Yang, F., Eisen, J. A., Paulsen, I. T., White, O.
2001 TIGRFAMs: a protein family resource for the functional identification of proteins. *Nucleic*
Acids Res. **29**, 41-43.   (10.1093/nar/29.1.41)
121 Saier, M. H., Jr., Reddy, V. S., Tsu, B. V., Ahmed, M. S., Li, C., Moreno-Hagelsieb, G. 2016
The Transporter Classification Database (TCDB): recent advances. *Nucleic Acids Res*. **44**,
D372-D379.   (10.1093/nar/gkv1103)

 Figure legends

 Figure 1. The evolutionary timeline of anammox bacteria using MCMCTree. The chronogram

 was estimated based on the calibration set C1 (see Supplementary Text section 3) and sequence

 alignments of the 25 orthologs conserved in bacteria (20040 sites). The blue bars on the four

 calibration nodes and the last common ancestor (LCA) of anammox bacteria represent the

 posterior 95% highest probability density (HPD) interval of the posterior time estimates. These

 alternative calibration sets were selected by choosing the three that accommodate different

 calibration constraints (see Supplementary Texts section 3.3). More alternative time estimates are

 provided in Figure S4. The vertical grey bar represents the period of the GOE from 2,500 to

 2,320 Ma. The calibration constraints used within the phylum Cyanobacteria are marked with

 orange texts: the LCA of Planctomycetes and Cyanobacteria (Root), the total group of oxygenic

 Cyanobacteria (Node 2), the total group of *Nostocales* (Node 3), and the total group of

 Pleurocapsales (Node 4). Planctomycetes from marine or groundwater habitats were labelled by

 a blue circle. The genome completeness estimated by CheckM is visualized with a gradient color

 strip. The right next color strip indicates the genome type of genomic sequences used in our

 study including metagenome-assembled genomes (MAGs) and whole-genome sequencing (WGS)

 from either enriched culture sample or isolate. The diagram below the dated tree illustrates the

 change of atmospheric partial pressure of O_2 (PO_2) and nitrogen isotope fractionations. The blue

 line shows the proposed model according to Lyons TW, Reinhard CTPlanavsky NJ [29]. The

green arrow suggests the earliest evidence for aerobic nitrogen cycling at around 2.7 Ga. PAL on

right axis means PO$_2$ relative to the present atmospheric level. The black dots denote the change

of nitrogen δ^{15}N isotope values according to Kipp MA, Stueken EE, Yun M, Bekker ABuick R

[28].

Figure 2. (A) The chronogram of the Anammox lineage estimated by mitochondria-based dating

analysis. The time tree was estimated based on the calibration set "C1+Euk" and sequence

alignments of the 24 mitochondria encoded proteins at the amino acid level (6295 sites). The

vertical grey bar represents the period of the GOE from 2,500 to 2,320 Ma. The calibration

constraints are marked with orange texts: the LCA of Planctomycetes and Cyanobacteria (Root),

the total group of oxygenic Cyanobacteria (Node 2), the total group of Nostocales (Node 3), the

total group of Pleurocapsales (Node 4), the total group of Bangiales (crown group of red algae)

(Node 5), the total group of Florideophyceae (Node 6), the total group of mosses (crown group

of Embryophyta) (Node 7) and the total group of eudicots (crown group of angiosperms) (Node

8). The color strip next to the label represent the major lineages of analyzed genomes. The filled

and empty squares, respectively, represent the presence and absence of particular genes used in

molecular dating analysis. The six genes selected to comprise the mito6 gene set are represented

by blue squares. (B) The posterior times of anammox bacteria estimated using

cyanobacteria-based and mitochondria-based dating analyses. The calibration sets starting with C

represent the calibration sets with cyanobacterial calibrations, while those starting with Euk

represent the calibration sets with eukaryotic calibrations. Calibration information: calibration

scheme C1 node1: <4.5 Ga, node2: 3.0-2.32 Ga; C3 node1: <3.8 Ga, node2: 3.0-2.32 Ga; C8

node1: <4.5 Ga, node2: >3.0 Ga; C10 node1: <3.8 Ga, node2: >3.0 Ga. All of these calibration

schemes share the same calibration for nodes 3 (>1.6 Ga) and 4 (>1.7 Ga). The horizontal grey

bar represents GOE from 2.5 to 2.32 Ga. The detailed constraints of calibrations and time

estimates are provided in Dataset S2.1.

Figure 3. The phyletic pattern of ecologically relevant genes in the comparison between

anammox bacteria and non-anammox bacteria. Solid circles at the nodes indicate the ultrafast

bootstrap values in 1,000 bootstrapped replicates. Note that copy number difference is not

indicated since most genes displayed here are present as a single copy in genome except a few

exceptions like *hao* (see Dataset S1.5 for the table summarizing the copy number of each gene).

The phylogenomic tree on the left was constructed with 887 genomic sequences described in

Supplementary Texts section 2.1. Those Planctomycetes genomes not used for comparative

genomics analyses are collapsed into grey triangles, and the numbers of collapsed genomes are

labelled next to the triangles. The target group and reference group for comparative genomic

analysis are within an orange or blue box, separately. For each genome, the genome

completeness estimated by CheckM is visualized with a color strip and labelled besides leaf

nodes. The right next color strip represents the type of genomic sequences used in our study

including metagenome-assembled genomes (MAGs) and whole-genome sequencing (WGS)

from either enriched culture sample or isolate. The filled and empty circles, respectively,

represent the presence and absence of particular genes in corresponding genomes. For gene

clusters, only genomes with at least half of the members of the gene cluster are indicated by a

filled circle. The classifications of annotated genes are labelled above the gene names. *hzsCBA*,

hydrazine synthase subunit C, B and A; *hdh*, hydrazine dehydrogenase; *hao*, hydroxylamine

dehydrogenase; *nxrAB*, nitrite oxidoreductase subunit A and B; *nirK*, copper-containing and

NO-forming nitrite reductase; *nirS*, cytochrome NO-forming nitrite reductase; *nrfAH*,

ammonia-forming nitrite reductase subunit A and H; *glnB*, nitrogen regulatory protein P-II; *nirC*,

nitrite transporter; NRT, nitrate/nitrite transporter; *amt*, ammonium transporter; kuste2805, 3603,

3605-3606, proposed genes relative to the synthetic pathways for ladderane at Rattray JE, Strous

M, Op den Camp HJ, Schouten S, Jetten MSDamste JS [76]; *cbiG*, cobalt-precorrin 5A hydrolase;

cbiD, cobalt-precorrin-5B(C1)-methyltransferase; *cbiOMQ*, cobalt/nickel transport system; *AhpC*,

peroxiredoxin; *Ccp*, cytochrome *c* peroxidase; *dfx*, superoxide reductase; *fprA*, H_2O-forming

enzyme flavoprotein; *CcsAB*, cytochrome c maturation systems; *petB*, Cytochrome b subunit of

the bc complex; *petC*, Rieske Fe-S protein; *ahbABCD*, heme biosynthesis; *sat*, sulfate

adenylyltransferase; *aprAB*, adenylylsulfate reductase, subunit A and B; *fsr*, sulfite reductase

(coenzyme F420); *higB-1*, toxin; *higA-1*, antitoxin; *mnhABCDEG*, multicomponent Na^+/H^+

antiporter; *fdhAB*, formate dehydrogenase subunit A and B; *nuo*(A-N), NADH-quinone

oxidoreductase; *ndh*(A-N), NAD(P)H-quinone oxidoreductase; *nqrABCDEF*, Na$^+$-transporting

NADH:ubiquinone oxidoreductase; *rnfABCDEG*, Na$^+$-translocating ferredoxin:NAD$^+$

oxidoreductase; *atp*(A-H), F-type H$^+$-transporting ATPase; *lacAZ*, beta-galactosidase; *melA*, galA,

alpha-galactosidase; *ebgA*, beta-galactosidase; *galK*, galactokinase; , fructokinase; *fruK*,

1-phosphofructokinase; *rhaB*, rhamnulokinase; rbsK, ribokinase; *araB*, L-ribulokinase; *xylB*,

xylulokinase; *kdgK*, 2-dehydro-3-deoxygluconokinase; GALK2, N-acetylgalactosamine kinase;

cah, cephalosporin-C deacetylase; *argE*, acetylornithine deacetylase; *nagA*;

N-acetylglucosamine-6-phosphate deacetylase; *HDAC11*, histone deacetylase 11.