Phylogenomic evidence for the Origin of Obligately Anaerobic Anammox Bacteria around the Great Oxidation Event

Tianhua Liao[1,^], Sishuo Wang[1,^], Eva E. Stüeken[2], Haiwei Luo[1,*]

[1]Simon F. S. Li Marine Science Laboratory, School of Life Sciences and State Key Laboratory of Agrobiotechnology, The Chinese University of Hong Kong, Shatin, Hong Kong SAR

[2]School of Earth and Environmental Sciences and Centre for Exoplanet Science, University of St Andrews, Bute Building, Queen's Terrace, KY16 9TS, UK

[^]These authors contribute equally to this work.

[*]**Corresponding author:**
Haiwei Luo
The Chinese University of Hong Kong
Shatin, Hong Kong SAR
Phone: (+852) 39436121
E-mail: hluo2006@gmail.com

Running title: The origin time and phylogeny of anammox bacteria

Keywords: anammox bacteria, molecular dating analysis, Planctomycetes

Abstract

The anaerobic ammonium oxidation (anammox) bacteria transform ammonium and nitrite to dinitrogen gas, and this obligate anaerobic process accounts for nearly half of the global nitrogen loss for surface environments. Yet its origin and evolution, which may give important insights into the biogeochemistry of early Earth, remains enigmatic. Here, we compile a comprehensive sequence data set of anammox bacteria and confirm their single origin within the phylum Planctomycetes. After accommodating the uncertainties and factors influencing time estimates with different statistical methods, we estimate that anammox bacteria originated at around the so-called Great Oxidation Event (GOE; 2.32 to 2.5 billion years ago [Ga]) which is thought to have fundamentally changed global biogeochemical cycles. We further show that during the origin of anammox bacteria, genes involved in oxidative stress, bioenergetics and anammox granules formation were recruited, which may have contributed to their survival on an increasingly oxic Earth. Our findings suggest the rising levels of atmospheric oxygen, which made nitrite increasingly available, was a potential driving force for the emergence of anammox bacteria. This is one of the first studies that link the GOE to the evolution of obligate anaerobic bacteria.

Introduction

Anaerobic ammonium oxidation (anammox, $NH_4^+ + NO_2^- \rightarrow N_2 + 2H_2O$), which usually

occurs in anoxic marine, freshwater and wetland settings, accounts for up to 50% of the removal

of fixed nitrogen (N) in nature. Along with denitrification, it is recognized as an important

biological process that leads to N loss from the environment [1,2]. In contrast to the

nitrification-denitrification process for wastewater treatment, coupling partial nitrification and

anammox is cost-effective and environmentally friendly due to its lower oxygen requirement for

aeration (nitrite [N(+III)], instead of nitrate [N(+V)] is sufficient for the anammox metabolism),

carbon free cultivation (some denitrifying bacteria are heterotrophic) and its negligible emissions

of greenhouse gases like N_2O. Consequently, anammox bacteria, the organisms that perform

anammox, have been widely used in wastewater treatments [2-5]. Nevertheless, their evolutionary

history and antiquity are poorly known, which hinders accurate reconstructions of the

biogeochemical nitrogen cycle over geologic time.

Previous studies investigated the roles of key genes driving anammox, including *hzsABC*

(hydrazine synthase)[6], *hdh* (hydrazine dehydrogenase), *hao* (hydroxylamine oxidoreductase) [7]

and *nxr* (nitrite oxidoreductase) [8]. In general, the vital enzymes encoded by these genes either are

directly involved in anammox or participate in replenishing electrons to the cyclic electron flow

which is consumed by the reduction of NAD^+ during the ATP synthesis [9,10]. Further, all members

of anammox bacteria are found within the phylum Planctomycetes [11]. Six candidate genera of

anammox bacteria, namely 'Candidatus Brocadia', 'Candidatus Kuenenia', 'Candidatus

Jettenia', 'Candidatus Scalindua', 'Candidatus Anammoximicrobium' and 'Candidatus

Anammoxoglobus', have been proposed based on 16S ribosomal RNA (rRNA) gene sequences

[12], but none of them have been successfully isolated into pure cultures. Apparently, the habitat of

anammox bacteria requires simultaneous presence of reduced (ammonia) and oxidized (nitrite)

inorganic N compounds. Such habitats are often found at the aerobic-anaerobic interface in

aquatic ecosystems, including the margins of oxygen minimum zones (OMZs) in the ocean and

sediment-water interfaces, where ammonium originates from the anaerobic degradation of

organic matter and nitrite can be produced by aerobic ammonia oxidation [13]. Although a recent

study [14] revealed the potential of anammox bacteria to utilize metal oxides as the electron

acceptor instead of nitrite, the feasibility of this metabolism in nature remains largely unresolved.

Thus, the availability of nitrite is presumably the determinant factor to the origin of anammox. It

is believed that the early Earth (before 3 Ga) was fully anoxic and deficient in aerobic

ecosystems that were able to generate nitrite/nitrate [15-17]. The first transient and/or localized

occurrences of aerobic nitrogen cycling appear in the Mid- to Neoarchean (2.7 Ga [18-21]), but

evidence of widespread nitrate availability does not appear until around 2.4 billion years ago

(Ga), the so-called Great Oxidation Event (GOE) [22-24]. The GOE marks the rise of free O_2 in the

atmosphere above 10^{-5} times modern levels, and was ultimately a result of the emergence of

oxygenic cyanobacteria [25]. In contrast to nitrite, ammonium, the other substrate of anammox

metabolism, was thermodynamically stable in the deep ocean throughout the Archean [26] and

probably extending well into the Proterozoic [27]. It is therefore tempting to hypothesize that the

origin of anammox coincided with Neoarchean oxygen oases or the GOE itself, when molecular

oxygen (O_2) rose permanently to a concentration that is biologically meaningful [28], thereby

creating the (micro-)environments satisfying anammox metabolism by providing enough nitrite.

Several tools exist for investigating the link between the evolution of metabolic pathways

and geo-environmental transformations. One way to explore the evolutionary history of a

specific metabolic pathway is based on organic biomarkers, which, however, are often affected

by the poor preservation over geologic timescales [29]. For instance, ladderanes, a type of lipids

that is unique to anammox bacteria [30], are rarely preserved to a level that can be used to date their

evolutionary origin. Another approach is based on nitrogen isotope ratios of sedimentary records

[26,31,32]. However, the isotopic fractionation factors for different metabolic pathways overlap

widely (denitrification: -5 to -30‰; anammox: -16 or -24‰ [27]), making it difficult to single out

and elucidate the evolution of the different redox reactions within the N cycle by this method [33].

Alternatively, molecular dating, which estimates the age of the last common ancestor (LCA) of

analyzed lineages by comparing their sequences based on the molecular clock theory [34], provides

a powerful strategy to investigate this issue. Here, we hypothesize that the rising level of O_2

during GOE, which made nitrite available to power anammox [27], was a driving force of the

origin of anammox bacteria. To test this hypothesis and to obtain more insights into the evolution

of anammox bacteria, we compiled an up-to-date genomic data set of Planctomycetes (see

Supplementary Text section 1; Dataset S1.1), placed the evolution of anammox bacteria into the

context of geological events, and investigated genomic changes underlying the origin of this

ecologically important bacterial lineage.

Results and discussion

<u>Monophyletic origins of anammox bacteria and anammox genes</u>

The anammox bacteria form a monophyletic group within the phylum Planctomycetes in

a comprehensive phylogenomic tree with 881 Planctomycetes genomes (Fig. S1). The anammox

bacteria clade in this phylogenomic tree (Fig. S1) comprises four known genera including the

early-branching *Ca.* Scalindua and *Ca.* Kuenenia, and the late-branching *Ca.* Jettenia and *Ca.*

Brocadia. It also includes two understudied lineages which we named 'basal lineage' and

'*hzsCBA*-less lineage' (Fig. S1), both of which have relatively low genome completeness

compared to the other anammox bacteria and are represented by metagenome-assembled

genomes (MAGs) sampled from groundwater (see Supplementary Text section 2.1). Note that

the absence of *hzsCBA* in several genomes within the '*hzsCBA*-less lineage' might be ascribed to

the loss of these genes in evolution or their low genome completeness (Fig. S1). Because of their

shallow (late-branching) phylogenetic position, this uncertainty should not affect the inference of

the origin of anammox bacteria. Two described genera (*Ca.* Anammoximicrobium and *Ca.*

Anammoxoglobus) do not have genomes available by the time of the present study (last accessed:

April 2021), and are therefore not included in the phylogenomic analysis. Furthermore, using the

last updated set of 2,077 Planctomycetes genomes (retrieved in April 2021) from NCBI, we

obtained a consistent topology of anammox bacteria (Fig. S2).

To check whether important groups, particularly early-branching lineages, have been

encompassed in our analyzed genome data sets, we built a 16S rRNA gene tree (Fig. S3) using

the identified 16S rRNA genes from genomic sequences of anammox bacteria and the deposited

16S rRNA gene amplicons in SILVA from the class Brocadiae, which comprises both anammox

and non-anammox bacteria (see Supplementary Texts section 2.2). Since the anammox bacteria

with genome sequences in this 16S rRNA gene tree (Fig. S3) showed a branching order

congruent with the topology in the phylogenomic tree (Fig. S1), and since the 16S rRNA gene

tree included 913 (clustered from 20,142) sequences sampled from a wide array of habitats

(marine, sediments, man-made reactor, freshwater and terrestrial ecosystems), the 16S rRNA

gene phylogeny, which could better capture the diversity of anammox bacteria by including

uncultured samples, likely encompasses the early-branching lineages of anammox bacteria.

Taken together, our genome set, which encompasses the genomes of the early-branching lineages,

is appropriate to be used to study the evolutionary origin of anammox bacteria (Dataset S1.3).

We further analyzed the evolution (Fig. S4) of the genes (*hzsCBA*) critical to the

anammox reaction [7,35]. Reconciliation (see Supplementary Texts section 2.3) of the gene and the

genome-based species phylogenies pointed to a single origin of anammox metabolism at the

LCA of anammox bacteria (Fig. S4). Taken together, the above analyses suggest that our genome

data sets have encompassed the earliest-branching lineages and deep phylogenetic diversity of

anammox bacteria, and indicate its monophyletic origin, thereby providing a foundation for

estimating the origin of anammox metabolism by estimating the origin time of anammox

bacteria.

<u>The evolutionary origin of anammox bacteria coincided with the rising O_2</u>

We employed MCMCTree to estimate the divergence times of the anammox bacteria with

i) soft bounds (time constraint allowing a small probability of violation) based on cyanobacteria

fossils, and ii) the relaxed molecular clock approaches allowing substitution rates to vary among

branches. Using a best-practiced dating scheme (Fig. 1; see Supplementary Text section 3), we

dated the origin of anammox bacteria to 2,117 million years ago (Ma) (95% highest posterior

density [HPD] interval, 2,002 - 2,226 Ma). Further, we performed repeat analysis based on the

expanded Planctomycetes genomes (Genome set 2; Dataset S1.1) or based on the alternative

constraint topology inferred with a better profile mixture model (LG+C60+G; Genome set 1;

Dataset S1.1). In general, both analyses resolved consistent topologies (Fig. S5) as shown in

Figure 1. Overall, they gave similar posterior times of the LCA of the anammox bacteria at 2,105

Ma (95% HPD: 1,961 - 2,235 Ma) or 2,005 Ma (95% HPD: 1,869 - 2,146 Ma).

Analyses with different combinations of calibrations and parameters broadly converge to

similar time estimates (Fig. S6; Dataset S2.1). Specifically, assigning a more ancient time

constraint to the total group of oxygenic cyanobacteria, which is based on the biogeochemical

evidence of the presence of O_2 at nearly 3.0 Ga [36] instead of GOE (Dataset S2.1), shifted the

posterior dates of the LCA of anammox bacteria to the past by ~300 million years (Myr) (e.g.,

C1 versus C10; Fig. S6). Varying the maximum bounds of the root of the tree of life between 4.5

Ga, 3.8 Ga or 3.5 Ga had a minor impact (around 100 Myr) on the time estimates of anammox

bacteria (e.g., C1 versus C3; Fig. S6). Though the auto-correlated rates (AR) model was less

favored than the independent rates (IR) model based on the model comparison (see

Supplementary Text section 3.3; Dataset S2.2), it is worth noting that the posterior ages

estimated by the AR model were 1-12% younger than estimated by the IR model (Dataset S2.1).

Besides, there remains a possibility that there are unsampled or even extinct lineages of

anammox bacteria that diverged earlier than all anammox bacteria analyzed in the present study.

This scenario, if true, hints that the first anammox bacterium could have originated before the

occurrence of the LCA of sequenced anammox bacteria but later than their split from the sister

lineage (Node X in Fig. 1; up to 2,600 Ma). Accommodating the above uncertainties, our

analyses suggest that the origin of anammox bacteria, hence the origin of anammox occurred

most likely falls into the 2.6-2.0 Ga interval. Running MCMCTree analysis with no sequence

data showed different distributions of time estimates for both the clade of anammox bacteria and

the four calibration points (Fig. S7), suggesting that sequence data are informative for our

molecular clock analysis [37].

<u>The geochemical context of the origin of anammox bacteria</u>

We speculate that the timing of the origin of the anammox metabolism is linked to the

increasing availability of nitrite in surface environments, because ammonium was likely present

in the deep anoxic ocean throughout the Precambrian [27] and therefore probably not a limiting

substrate. The first abiotic source of nitrite on the early Earth would have been lightning

reactions between N_2 and CO_2 in the Archean atmosphere [38,39]. A prior study suggested that this

process led to micromolar levels of nitrite in seawater [38], which implies the possibility of an

earlier origin of anammox bacteria. However, this estimate was based on perhaps unrealistically

high amounts of CO_2 [40]. Furthermore, there is no isotopic evidence in the early Archean rock

record prior to 3 Ga for the presence of a significant nitrite or nitrate reservoir in the early ocean

[15-17], and it is conceivable that any nitrite that was supplied to the ocean by lightning was rapidly

reduced to ammonium or N_2 by ferrous iron, possibly even abiotically [41,42]. Hence, although a

background lightning flux of nitrite almost certainly existed, there is no evidence that it supplied

a reliable metabolic substrate for early life.

The first transient appearances of nitrite/nitrate-dependent metabolisms are captured by

the sedimentary nitrogen isotope record in late Mesoarchean soils at 3.2 Ga [19] and in Neoarchean

shallow-marine settings at 2.7 Ga and 2.5 Ga [18,20,21]. These observations may reflect local and/or

temporally restricted oxygen oases [43,44]. Widespread nitrite/nitrate availability is inferred for the

Paleoproterozoic (2.4-1.8 Ga), i.e. in the immediate aftermath of the GOE [22-24]. Importantly, O_2

is necessary, although a trace amount is feasible, to extant nitrifying organisms for the oxidation

of ammonium to nitrite and nitrate [45-47]. Hence the rise of oxygen almost certainly triggered the

growth of the nitrite/nitrate reservoir, and therefore provided one of the key substrates for

anammox bacteria. It has been shown that a low concentration of nitrite significantly decreases

the N removal rate by anammox bacteria in reactors [48]. In modern OMZs, which may to some

extent serve as analogues for the Paleoproterozoic redox-stratified ocean, the concentration of

nitrite is the rate-limiting factor to the anammox metabolism [49]. Hence it was most likely that the

rise of nitrite, driven by the rise of O_2, facilitated the appearance of anammox bacteria at 2.6-2.0

Ga.

In modern environments, the majority of nitrite used in anammox is derived from nitrate

reduction or aerobic ammonia oxidation [50], the latter performed by ammonia-oxidizing archaea

(AOA) or bacteria (AOB). As noted above, the nitrification likely appeared by at least 2.7 Ga, as

indicated by nitrogen-isotope evidence for local accumulation of nitrate in surface ocean waters

[27,51,52]. A recent study inferred that the LCA of AOA dates back to ~2.3 Ga and first appeared on

land, driven by increasing O_2 concentrations in the atmosphere at that time, and that the

expansion of AOA from land to the ocean did not occur until nearly 1.0 Ga [53]. This hints at a

dominant role of AOB in early ocean, before 2.3 Ga. Consistent with this idea, the vast majority

of the early-branching lineages of anammox bacteria (*Ca.* Scalindua) and the sister

Planctomycetes lineages of the anammox clade in the phylogenomic tree (Fig. 1) are found in

marine environments. Furthermore, in the 16S rRNA gene tree which represents a greater

diversity than the phylogenomic tree, marine lineages still account for the majority of the

earliest-branching lineages of anammox bacteria (Fig. S3). Thus, it seems likely that the LCA of

anammox bacteria originated in the marine realm where AOB thrived first. The above arguments,

although speculative, tentatively suggest that the nitrite demand for anammox had been readily

met by GOE by taking advantage of the significant amount of O_2 newly available, which is

broadly consistent with our molecular dating results suggesting an origin of anammox shortly

before or after the GOE. Note that there are recently reported anammox bacteria lineages

showing less dependence on nitrite, which may use nitric oxide or hydroxylamine as the electron

receptor in anammox [12]. These lineages are affiliated with *Ca.* Kuenenia and *Ca.* Brocadia,

which evolved recently, implying that the ability to use alternative electron receptors are likely

not ancestral to all anammox bacteria. This aspect further indicates that lightning was likely not

an important source of metabolic substrates in the early Archean, because nitric oxide is the

major product of the lightning reaction between N_2 and CO_2 [38,39].

Genomic changes related to anammox metabolism upon the origin of anammox bacteria

We further explored the genomic changes characterizing the origin of anammox bacteria

(see Supplementary Text section 4). This includes the gains of the aforementioned genes that

directly participate in anammox, viz., *hzsABC*, *hdh*, and *hao*. The current view of the anammox

metabolism posits that electrons consumed by anammox for carbon fixation are replenished by

the oxidation of nitrite to nitrate by nitrite oxidoreductase (NXR) [10], which was inferred to be

acquired upon the origin of anammox bacteria (Fig. 2). Moreover, multiple auxiliary genes for N

assimilatory pathways including N regulatory protein (*glnB*), assimilatory nitrite reductase large

subunit (*nirB*), transporters like *amt* (ammonium), *nirC* (nitrite) and NRT family proteins

(nitrate/nitrite), were also gained at the origin of anammox bacteria (Fig. 2). However, genes for

nitrogen-related dissimilatory pathways encoding nitrite reduction to nitric oxide (*nirK* or *nirS*)

and to ammonium (*nrfAH*), respectively, were likely acquired after the origin of anammox

bacteria (Fig. 2). Anammox occurs at the membrane of anammoxosome, an organelle

predominantly composed of a special type of lipid called ladderane. The ladderane membranes

show low proton permeability, which helps maintain the proton motive force during the

anammox metabolism [30]. Although the biosynthetic pathway of ladderane is yet to be

characterized, a previous study [54] predicted 34 candidate genes responsible for the synthesis of

ladderane lipids, four of which were potentially acquired during the origin of anammox bacteria

(see Supplementary Text section 4.2; Fig. 2), hinting at their potential roles in ladderane

synthesis and providing clues to future experimental investigations.

Anammox bacteria occur in anoxic habitats and exhibit low oxygen tolerance [12].

Specifically, the O_2-sensitive intermediates generated by anammox, e.g., hydrazine, a powerful

reductant, requires anoxic conditions. Accordingly, the likely acquired peroxidases include

cytochrome c peroxidase (Ccp), which could scavenge hydrogen peroxide in the periplasm [55],

and the most prevalent peroxidase [56], thioredoxin-dependent peroxiredoxin (AhpC). Besides

peroxide scavengers, the desulfoferrodoxin (Dfx), which functions as superoxide reductase (SOR)

to reduce superoxide to hydrogen peroxide and which is broadly distributed among anaerobic

bacteria [56], were likely acquired during the origin of anammox bacteria (Fig. 2). Another

acquired gene is *fprA*, which encodes flavo-diiron proteins that scavenge O_2. Furthermore, *cbi*

genes, which are exclusively involved in the anaerobic biosynthesis of vitamin B_{12} [57], are widely

present in the genomes of anammox bacteria, suggesting their acquisition during the early

evolution of anammox bacteria (Fig. 2).

Investigating other metabolisms that were also acquired upon the origin of anammox

bacteria allow reconstructing the coeval ecology. Our data show that *sat* (Fig. 2), which encodes

sulfate adenylyltransferase for assimilatory incorporation of sulfate into bioavailable adenylyl

sulfate, as well as *aprAB* and *fsr* (Fig. 2) for encoding dissimilatory sulfite reductase, were

acquired upon the origin of anammox bacteria (Fig. 2). This acquired capability of sulfate

metabolism is coincident with the increased concentrations of seawater sulfate from ~100 μM

or less throughout much of the Archean [58,59] to over 1mM after the rise of atmospheric O_2 [60,61].

Unlike other Planctomycetes, anammox bacteria are generally autotrophs that use the

Wood-Ljungdahl pathway for carbon assimilation [62]. Consequently, they potentially lost many

genes involved in carbohydrate utilization (Fig. 2). The metabolic loss is further strengthened by

the enrichment analysis where the genes predicted to be lost by the LCA of anammox bacteria

are enriched in pathways involving hydrolase activity, intramolecular oxidoreductase activity and

carbohydrate kinase activity (Fig. 2; see Data and Code availability), hinting at a decreased

ability of anammox bacteria to degrade organics. Also lost are genes (*mnhABCDEG*) encoding

the Na^+/H^+ antiporter (Fig. 2), which usually regulates sodium tolerance and pH homeostasis in

marine environments [63]. Only members of the dominant marine genus *Ca.* Scalindua keep this

gene cluster. Although building up a proton motive force is vital to energy harvesting, the

mechanism for this process was far unexplored in anammox bacteria [64]. Our analysis shows that

nuo genes, which encode NADH-quinone oxidoreductase that can build up a proton motive force,

was likely acquired by the LCA of anammox bacteria (Fig. 2). Moreover, the presence of genes

coding for Na^+-coupled respiratory enzymes, for example *rnf* (ferredoxin-NADH oxidoreductase)

and *nqr* (ubiquinone-NADH oxidoreductase), suggests that anammox bacteria could also use the

sodium motive force for ATP synthesis (Fig. 2).

Additionally, iron is a vital element for HZS [6] and HDH [65], and *in vitro* studies revealed

that increased iron concentrations can promote the growth of anammox bacteria [66]. The ancient

oceans are thought to have been ferruginous (high iron availability) according to sedimentary

records of iron speciation across several coeval marine settings from the Archean through most

of the Proterozoic [67], meaning that large amounts of soluble iron would have been available for

anammox bacteria. Interestingly, a series of iron-related genes to make use of the abundant iron

from the environment were likely acquired upon the origin of anammox bacteria. First, anammox

bacteria likely acquired the gene *fur*, which encodes a ferric uptake regulator for iron

assimilation. Moreover, the iron-containing proteins, specifically cytochrome, are encoded by the

acquired genes *petBC*, which potentially encodes Rieske-heme b [10], and *CcsAB*, which encodes a

protein involved in cytochrome c maturation systems (Fig. 2). Notably, *ahbABCD*, which

encodes proteins-synthesizing *b*-type hemes in anammox bacteria [68], were likely acquired before

the origin of anammox bacteria (Fig. 2). Furthermore, we identified that the gene *Bfr* that

encodes the oligomeric protein, bacterioferritin involved in the uptake and storage of iron [69] was

likely acquired during the origin of anammox bacteria (Fig. 2), which may help to hoard iron in

settings where iron became locally sparse, such as along euxinic (sulfide-rich) or oxic marine

margins in the Paleoproterozoic [70]. It is thought that the appearance of euxinic margins trapped

iron that upwelled from the deep ocean, leading to the demise of iron oxide deposits (i.e., banded

iron formations) [71]. Our results tentatively support this model, if the microenvironments that

those primordial anammox bacteria colonized were located near the euxinic-oxic interface where

nitrite and ammonium were available while upwelled iron was titrated out of the water column

by freely dissolved hydrogen sulfide.

Taken together, the timelines and corresponding genomic changes of anammox bacteria

provide implications for the physiological characteristics of descendant anammox bacteria and

the origin of other nitrogen-transforming pathways in the context of an estimated evolutionary

timeline of anammox bacteria. For example, the earlier branching lineage *Ca.* Scalindua exhibit

high nitrite affinity (0.45 μM) compared to other anammox genera (up to 370 μM) [12,72], which

may reflect a progressive growth of the marine nitrite reservoir, linked to the protracted

oxygenation of the surface environments over geologic timescales [25]. Likewise, genomic data

reveal that also canonical denitrification radiated across the tree of life after the GOE [73], and the

gains (Fig. S8; see Supplementary Text section 4.4) of genes *nrfAH* for dissimilatory nitrate

reduction to ammonium (DNRA) at the late-branching lineages (*Ca.* Brocadia and *Ca.* Kuenenia)

at around 800 Ma are consistent with the time when the nitrite/nitrate availability was increased

by the Neoproterozoic Oxygenation Event [74]. The time estimate and genomic changes of the

anammox bacteria have gone some way towards enhancing the understanding of the

physiological characteristics of modern anammox bacteria and the historical N cycle.

Concluding remarks

Traditionally, the metabolic evolution of the biosphere across the GOE has mainly been

investigated in the phylum Cyanobacteria [75], likely because oxygenic photosynthesis is believed

as a major driver of the GOE. More recently, scientists have increasingly realized that the impact

of the GOE on the bacterial world may not be limited to cyanobacteria. This was well illustrated

by Ren, et al. [53], who showed that the GOE was likely a driving force of the origin of the aerobic

AOA and thus an aerobic nitrogen cycle on Earth. Furthermore, another molecular dating study

suggested expanded arsenic resistance systems as biological adaptations to the increased toxicity

of oxidized arsenic species after the GOE [76]. However, to our knowledge, little was so far known

about the relationships between the GOE and obligate anaerobic metabolisms.

Here, using molecular dating and comparative genomics approaches, we link the

emergence of an obligate anaerobic bacterial group, which drives the loss of fixed N in many

environments, to the rise of O_2, and highlight their evolutionary responses to major

environmental disturbances. Apparently, the GOE opened novel niches for the origin and

subsequent expansion of diverse aerobic prokaryotes such as AOA and AOB, which in turn

facilitated the origin of other bacterial lineages, including some anaerobes like the anammox

bacteria, by providing the resources for their energy conservation. Our results open up the

possibility that a significant proportion of the Proterozoic nitrite budget was consumed by

anammox bacteria, and that the sedimentary nitrogen isotope record may be influenced by their

activity. Given that anammox bacteria tend to produce less N_2O as a by-product compared to

canonical nitrification/denitrification [4,5], our findings may have implications for Proterozoic

climate, because N_2O has previously been invoked as an important greenhouse gas at that time [77].

Lastly, our study suggests that the impacts of the GOE go well beyond aerobic prokaryotes.

Specifically, the origin of anammox bacteria was likely driven by the expansion of nitrite/nitrate

availability, as supported by isotopic evidence from sedimentary records. We therefore conclude

that molecular dating is a powerful approach to complement isotopic evidence for resolving the

timeline of biological evolution and for providing additional constraints on climate models of the

distant past.

Materials and Methods

Overall, we compiled two genome sets and one 16S rRNA gene set in our study (Dataset

S1.1). A phylogenomic tree (Fig. S1) was generated based on the concatenated alignment of 120

ubiquitous proteins (bac120; Dataset S1.3) proposed for tree inference by the Genome Taxonomy

Database [78]. All published genomic sequences (952 in total) affiliated with the phylum

Planctomycetes in NCBI Genbank up to December 2019 were retrieved (Dataset S1.2). To

further examine whether the patterns obtained with this Planctomycetes dataset hold the same,

we further constructed an expanded set of Planctomycetes genomes by retrieving a total of 2,077

Planctomycetes genomes released by April 2021 at Genbank (Fig. S2; see Data and Code

availability). Key anammox genes *hzsCBA* and *hdh* were identified against manually curated

reference sequences by BLASTP. For each gene, identified protein sequences were aligned using

MAFFT (v7.222) [79] and the alignments were refined by trimAl (v1.4) [80]. All phylogenies were

constructed by IQ-tree (v1.6.2) [81] with substitution models automatically selected by

ModelFinder [82] and branch support assessed with 1,000 ultrafast bootstrap replicates. Note that

the constraint topologies for dating analysis (Fig. 1; Fig. S5) were inferred with the profile

mixture model (LG+C60+G) which better accommodates across-site heterogeneity in deep-time

evolution. Following manual curation, four well-recognized genera and two separate lineages of

anammox bacteria were highlighted with different colours, and the presence of key genes were

annotated with filled symbols beside the labels (Fig. S1). Furthermore, the phylogenomic tree

was generated with similar methods using 16S rRNA genes identified from downloaded genomes

and those retrieved from the SILVA database (Fig. S3; Dataset S1.3). All trees (including

phylogenomic and 16S trees) in our study were visualized with iTOL [83]. More details are shown

in supplementary text, section 2.

Molecular dating analysis was carried out using the program MCMCTree from the PAML

package (4.9j) [84]. In our study, 13 calibration sets constructed with different time constraints of

the root and three calibration nodes within the cyanobacteria lineage were used (see

Supplemental Texts section 3.2). The topology constraint for dating analysis was generated using

85 genomes, which were sampled from the phylogenomic tree of phylum Planctomycetes (see

Supplemental Texts section 3.1), and the model LG+C20+F+G under posterior mean site

frequency (PMSF) approximation [85]. To perform dating analysis, clock models and different

calibration sets were compared in 26 schemes (see Supplemental Texts section 3; Dataset S2.1)

[86,87]. For each scheme, the approximate likelihood method [88] of MCMCTree were conducted in

duplicate with identical iteration parameters (burn-in: 10,000; sample frequency: 20; number of

sample: 20,000). The convergence of each scheme was evaluated by comparing the posterior

dates of two independent runs. With the updated genome set 2 (Dataset S1.1), we repeated the

taxon sampling process and generated an alternative phylogeny for dating analysis (Fig. S5)

using LG+C20+F+G model (C20: 20 classes of site-specific amino acid profiles) under PMSF

approximations. Similarly, we also generated another phylogeny for dating analysis using the

same genome set as used in Fig. 1 but with the profile mixture model (LG+C60+G) which uses

60 classes of amino acid profiles [89]. Note that the missing parameter +F would not significantly

change the results since it only adds another profile amino acid calculated from the original data.

These two repeat dating analyses were conducted with the best practice dating scheme (IR model

and calibration C1).

The phylogenetic differences between gene trees and the species tree were reconciliated

by GeneRax [90] with unrooted gene tree as input and automatically optimized duplication, transfer,

loss (DTL) rates. For each gene identified from the genome set 2 (Dataset S1.1), we used a

species tree comprised by all anammox bacteria pruned from the phylogenomic tree (Fig. S2) as

the reference. We used recommended parameters including SPR strategy, undated DTL model

and a maximum radius of five.

For comparative genomics analysis, the protein-coding sequences were annotated against

KEGG [91], CDD [92], InterPro [93], Pfam [94], TIGRFAM [95] and TCDB [96], individually (see

Supplemental Texts section 4). Following annotations, the potentially gained/lost genes between

groups (anammox bacteria versus non-anammox bacteria; an anammox bacterial genus versus all

other anammox bacteria) were statistically tested using two-sided fisher's exact tests. Finally, the

resulting p-values were corrected with the Benjamini-Hochberg FDR (false discovery rate)

procedure. Moreover, genes with corrected p-values smaller than 0.05 and with larger ratio in the

study group were defined as gained genes, and genes with corrected p-values smaller than 0.05

and with smaller ratio in the study group were defined as potentially lost genes. Those potentially

gained/lost genes were summarized for their detailed functions (see Data and Code availability).

Acknowledgements

We thank Bite Pei for building the basic dataset for this project, Yang Qian for his help in

editing the earlier version of the manuscript, Jinjin Tao for her helpful discussion, and Mario dos

Reis for his suggestion on dating analysis. This work is funded by the National Science

Foundation of China (92051113), the Hong Kong Research Grants Council Area of Excellence

Scheme (AoE/M-403/16), the Direct Grant of CUHK (4053495), and The CUHK Impact

Postdoctoral Fellowship Scheme to (S.W.).

Data and Code availability

All used genomic sequences, generated phylogenetic trees, estimated divergence times

and the used python codes are deposited in the online repository

https://github.com/luolab-cuhk/anammox-origin.

Conflict of interest

The authors declare that they have no conflict of interest.

References

1   Oshiki, M., Satoh, H. & Okabe, S. Ecology and physiology of anaerobic ammonium
oxidizing bacteria. *Environ Microbiol* **18**, 2784-2796, doi:10.1111/1462-2920.13134
(2016).
2   Stein, L. Y. & Klotz, M. G. The nitrogen cycle. *Current Biology* **26**, R94-R98, doi:DOI
10.1016/j.cub.2015.12.021 (2016).
3   Li, L. *et al.* N_2O emission in partial nitritation-anammox process. *Chinese Chemical*
Letters **31**, 28-38, doi:10.1016/j.cclet.2019.06.035 (2020).
4   Okabe, S., Oshiki, M., Takahashi, Y. & Satoh, H. N2O emission from a partial
nitrification-anammox process and identification of a key biological process of N2O
emission from anammox granules. *Water Research* **45**, 6461-6470,
doi:10.1016/j.watres.2011.09.040 (2011).
5   Gao, D. W. & Tao, Y. Versatility and application of anaerobic ammonium-oxidizing
bacteria. *Applied Microbiology and Biotechnology* **91**, 887-894,
doi:10.1007/s00253-011-3411-3 (2011).
6   Dietl, A. *et al.* The inner workings of the hydrazine synthase multiprotein complex.
Nature **527**, 394-397, doi:10.1038/nature15517 (2015).
7   Kartal, B. *et al.* Molecular mechanism of anaerobic ammonium oxidation. *Nature* **479**,
127-130, doi:10.1038/nature10453 (2011).

8   Chicano, T. M. *et al.* Structural and functional characterization of the intracellular
filament-forming nitrite oxidoreductase multiprotein complex. *Nature Microbiology*,
doi:10.1038/s41564-021-00934-8 (2021).
9   Hu, Z., Wessels, H., van Alen, T., Jetten, M. S. M. & Kartal, B. Nitric oxide-dependent
anaerobic ammonium oxidation. *Nat Commun* **10**, 1244,
doi:10.1038/s41467-019-09268-w (2019).
10   Kartal, B. & Keltjens, J. T. Anammox Biochemistry: a Tale of Heme c Proteins. *Trends*
Biochem Sci **41**, 998-1011, doi:10.1016/j.tibs.2016.08.015 (2016).
11   Wang, Y., Xu, L. Y., Wang, S. Y., Ye, F. & Zhu, G. B. Global Distribution of Anaerobic
Ammonia Oxidation (Anammox) Bacteria - Field Surveys in Wetland, Dryland,
Groundwater Aquifer and Snow. *Frontiers in Microbiology* **10**,
doi:10.3389/fmicb.2019.02583 (2019).
12   Zhang, L. & Okabe, S. Ecological niche differentiation among anammox bacteria. *Water*
Res **171**, 115468, doi:10.1016/j.watres.2020.115468 (2020).
13   Sonthiphand, P., Hall, M. W. & Neufeld, J. D. Biogeography of anaerobic
ammonia-oxidizing (anammox) bacteria. *Front Microbiol* **5**, 399,
doi:10.3389/fmicb.2014.00399 (2014).
14   Shaw, D. R. *et al.* Extracellular electron transfer-dependent anaerobic oxidation of
ammonium by anammox bacteria. *Nature Communications* **11**,
doi:10.1038/s41467-020-16016-y (2020).
15   Ossa Ossa, F. *et al.* Limited oxygen production in the Mesoarchean ocean. *Proc Natl*
Acad Sci U S A **116**, 6647-6652, doi:10.1073/pnas.1818762116 (2019).
16   Stueken, E. E., Buick, R., Guy, B. M. & Koehler, M. C. Isotopic evidence for biological
nitrogen fixation by molybdenum-nitrogenase from 3.2 Gyr. *Nature* **520**, 666-669,
doi:10.1038/nature14180 (2015).
17   Koehler, M. C., Buick, R. & Barley, M. E. Nitrogen isotope evidence for anoxic deep
marine environments from the Mesoarchean Mosquito Creek Formation, Australia.
Precambrian Research **320**, 281-290, doi:10.1016/j.precamres.2018.11.008 (2019).
18   Koehler, M. C., Buick, R., Kipp, M. A., Stueken, E. E. & Zaloumis, J. Transient surface
ocean oxygenation recorded in the similar to 2.66-Ga Jeerinah Formation, Australia.
Proceedings of the National Academy of Sciences of the United States of America **115**,
7711-7716, doi:10.1073/pnas.1720820115 (2018).
19   Homann, M. *et al.* Microbial life and biogeochemical cycling on land 3,220 million years
ago. *Nature Geoscience* **11**, 665-+, doi:10.1038/s41561-018-0190-9 (2018).
20   Godfrey, L. V. & Falkowski, P. G. The cycling and redox state of nitrogen in the
Archaean ocean. *Nature Geoscience* **2**, 725-729, doi:10.1038/Ngeo633 (2009).

21   Garvin, J., Buick, R., Anbar, A. D., Arnold, G. L. & Kaufman, A. J. Isotopic Evidence for
an Aerobic Nitrogen Cycle in the Latest Archean. *Science* **323**, 1045-1048,
doi:10.1126/science.1165675 (2009).

22   Zerkle, A. L. *et al.* Onset of the aerobic nitrogen cycle during the Great Oxidation Event.
Nature **542**, 465-+, doi:10.1038/nature20826 (2017).

23   Luo, G. M. *et al.* Nitrogen fixation sustained productivity in the wake of the
Palaeoproterozoic Great Oxygenation Event. *Nature Communications* **9**,
doi:10.1038/s41467-018-03361-2 (2018).

24   Kipp, M. A., Stueken, E. E., Yun, M., Bekker, A. & Buick, R. Pervasive aerobic nitrogen
cycling in the surface ocean across the Paleoproterozoic Era. *Earth and Planetary*
Science Letters **500**, 117-126, doi:10.1016/j.epsl.2018.08.007 (2018).

25   Lyons, T. W., Reinhard, C. T. & Planavsky, N. J. The rise of oxygen in Earth's early ocean
and atmosphere. *Nature* **506**, 307-315, doi:10.1038/nature13068 (2014).

26   Yang, J. *et al.* Ammonium availability in the Late Archaean nitrogen cycle. *Nature*
Geoscience **12**, 553-557, doi:10.1038/s41561-019-0371-1 (2019).

27   Stueken, E. E., Kipp, M. A., Koehler, M. C. & Buick, R. The evolution of Earth's
biogeochemical nitrogen cycle. *Earth-Science Reviews* **160**, 220-239,
doi:10.1016/j.earscirev.2016.07.007 (2016).

28   Klotz, M. G. & Stein, L. Y. Nitrifier genomics and evolution of the nitrogen cycle. *Fems*
Microbiology Letters **278**, 146-156, doi:10.1111/j.1574-6968.2007.00970.x (2008).

29   Hallmann, C., Kelly, A. E., Gupta, S. N. & Summons, R. E. Reconstructing Deep-Time
Biology with Molecular Fossils. *Quantifying the Evolution of Early Life : Numerical*
Approaches to the Evaluation of Fossils and Ancient Ecosystems **36**, 355-401,
doi:10.1007/978-94-007-0680-4_14 (2011).

30   Moss, F. R. *et al.* Ladderane phospholipids form a densely packed membrane with normal
hydrazine and anomalously low proton/hydroxide permeability. *Proceedings of the*
National Academy of Sciences of the United States of America **115**, 9098-9103,
doi:10.1073/pnas.1810706115 (2018).

31   Mettam, C. *et al.* Anaerobic nitrogen cycling on a Neoarchaean ocean margin. *Earth and*
Planetary Science Letters **527**, doi:10.1016/j.epsl.2019.115800 (2019).

32   Ader, M. *et al.* Interpretation of the nitrogen isotopic composition of Precambrian
sedimentary rocks: Assumptions and perspectives. *Chemical Geology* **429**, 93-110,
doi:10.1016/j.chemgeo.2016.02.010 (2016).

33   Fuchsman, C. A. & Stueken, E. E. Using modern low-oxygen marine ecosystems to
understand the nitrogen cycle of the Paleo- and Mesoproterozoic oceans. *Environ*
Microbiol **23**, 2801-2822, doi:10.1111/1462-2920.15220 (2021).

34   Yang, Z. & Rannala, B. Bayesian estimation of species divergence times under a
molecular clock using multiple fossil calibrations with soft bounds. *Mol Biol Evol* **23**,
212-226, doi:10.1093/molbev/msj024 (2006).
35   Harhangi, H. R. *et al.* Hydrazine synthase, a unique phylomarker with which to study the
presence and biodiversity of anammox bacteria. *Appl Environ Microbiol* **78**, 752-758,
doi:10.1128/AEM.07113-11 (2012).
36   Betts, H. C. *et al.* Integrated genomic and fossil evidence illuminates life's early evolution
and eukaryote origin. *Nat Ecol Evol* **2**, 1556-1562, doi:10.1038/s41559-018-0644-x
(2018).
37   Roger, A. J. & Susko, E. Molecular clocks provide little information to date
methanogenic Archaea. *Nat Ecol Evol* **2**, 1676-1677, doi:10.1038/s41559-018-0687-z
(2018).
38   Wong, M. L., Charnay, B. D., Gao, P., Yung, Y. L. & Russell, M. J. Nitrogen Oxides in
Early Earth's Atmosphere as Electron Acceptors for Life's Emergence. *Astrobiology* **17**,
975-983, doi:10.1089/ast.2016.1473 (2017).
39   Kasting, J. F. & Walker, J. C. G. Limits on Oxygen Concentration in the Prebiological
Atmosphere and the Rate of Abiotic Fixation of Nitrogen. *Journal of Geophysical
Research-Oceans* **86**, 1147-1158, doi:DOI 10.1029/JC086iC02p01147 (1981).
40   Charnay, B., Le Hir, G., Fluteau, F., Forget, F. & Catling, D. C. A warm or a cold early
Earth? New insights from a 3-D climate-carbon model. *Earth and Planetary Science
Letters* **474**, 97-109 (2017).
41   Brandes, J. A. *et al.* Abiotic nitrogen reduction on the early Earth. *Nature* **395**, 365-367,
doi:Doi 10.1038/26450 (1998).
42   Summers, D. P. & Chang, S. Prebiotic Ammonia from Reduction of Nitrite by Iron(Ii) on
the Early Earth. *Nature* **365**, 630-632, doi:DOI 10.1038/365630a0 (1993).
43   Lalonde, S. V. & Konhauser, K. O. Benthic perspective on Earth's oldest evidence for
oxygenic photosynthesis. *Proceedings of the National Academy of Sciences of the United
States of America* **112**, 995-1000, doi:10.1073/pnas.1415718112 (2015).
44   Anbar, A. D. *et al.* A whiff of oxygen before the Great Oxidation Event? *Science* **317**,
1903-1906, doi:10.1126/science.1140325 (2007).
45   Ranjan, S., Todd, Z. R., Rimmer, P. B., Sasselov, D. D. & Babbin, A. R. Nitrogen Oxide
Concentrations in Natural Waters on Early Earth. *Geochemistry Geophysics Geosystems*
20, 2021-2039, doi:10.1029/2018gc008082 (2019).
46   Cheng, C. *et al.* Nitrogen isotope evidence for stepwise oxygenation of the ocean during
the Great Oxidation Event. *Geochimica Et Cosmochimica Acta* **261**, 224-247,
doi:10.1016/j.gca.2019.07.011 (2019).

47   Lehtovirta-Morley, L. E. Ammonia oxidation: Ecology, physiology, biochemistry and
554   why they must all come together. *FEMS Microbiol Lett* **365**, doi:10.1093/femsle/fny058
555   (2018).
556   48   Li, W. *et al.* Metagenomics reveals microbial community differences lead to differential
557   nitrate production in anammox reactors with differing nitrogen loading rates. *Water Res*
558   **169**, 115279, doi:10.1016/j.watres.2019.115279 (2020).
559   49   Zhang, L. *et al.* Microbial competition among anammox bacteria in nitrite-limited
560   bioreactors. *Water Res* **125**, 249-258, doi:10.1016/j.watres.2017.08.052 (2017).
561   50   Dalsgaard, T., Thamdrup, B., Farias, L. & Revsbech, N. P. Anammox and denitrification
562   in the oxygen minimum zone of the eastern South Pacific. *Limnology and Oceanography*
563   **57**, 1331-1346, doi:10.4319/lo.2012.57.5.1331 (2012).
564   51   Canfield, D. E., Glazer, A. N. & Falkowski, P. G. The evolution and future of Earth's
565   nitrogen cycle. *Science* **330**, 192-196, doi:10.1126/science.1186120 (2010).
566   52   Zerkle, A. L. *et al.* Onset of the aerobic nitrogen cycle during the Great Oxidation Event.
567   *Nature* **542**, 465-467, doi:10.1038/nature20826 (2017).
568   53   Ren, M. *et al.* Phylogenomics suggests oxygen availability as a driving force in
569   Thaumarchaeota evolution. *ISME J* **13**, 2150-2161, doi:10.1038/s41396-019-0418-8
570   (2019).
571   54   Rattray, J. E. *et al.* A comparative genomics study of genetic products potentially
572   encoding ladderane lipid biosynthesis. *Biol Direct* **4**, 8, doi:10.1186/1745-6150-4-8
573   (2009).
574   55   Van Vliet, A. H., Ketley, J. M., Park, S. F. & Penn, C. W. The role of iron in
575   Campylobacter gene regulation, metabolism and oxidative stress defense. *FEMS*
576   *Microbiol Rev* **26**, 173-186, doi:10.1016/s0168-6445(02)00095-5 (2002).
577   56   Johnson, L. A. & Hug, L. A. Distribution of reactive oxygen species defense mechanisms
578   across domain bacteria. *Free Radical Biology and Medicine* **140**, 93-102,
579   doi:10.1016/j.freeradbiomed.2019.03.032 (2019).
580   57   Moore, S. J. & Warren, M. J. The anaerobic biosynthesis of vitamin B12. *Biochem Soc*
581   *Trans* **40**, 581-586, doi:10.1042/BST20120066 (2012).
582   58   Crowe, S. A. *et al.* Sulfate was a trace constituent of Archean seawater. *Science* **346**,
583   735-739, doi:10.1126/science.1258966 (2014).
584   59   Jamieson, J. W., Wing, B. A., Farquhar, J. & Hannington, M. D. Neoarchaean seawater
585   sulphate concentrations from sulphur isotopes in massive sulphide ore. *Nature*
586   *Geoscience* **6**, 61-64, doi:10.1038/Ngeo1647 (2013).
587   60   Blattler, C. L. *et al.* Two-billion-year-old evaporites capture Earth's great oxidation.
588   *Science* **360**, 320-+, doi:10.1126/science.aar2687 (2018).

61 Fakhraee, M., Hancisse, O., Canfield, D. E., Crowe, S. A. & Katsev, S. Proterozoic
seawater sulfate scarcity and the evolution of ocean-atmosphere chemistry. *Nature*
Geoscience **12**, 375-380, doi:10.1038/s41561-019-0351-5 (2019).
62 Kartal, B., Keltjens, J. T. & Jetten, M. S. M. Metabolism and Genomics of Anammox
Bacteria. *Nitrification*, 181-200 (2011).
63 Ito, M., Morino, M. & Krulwich, T. A. Mrp Antiporters Have Important Roles in Diverse
Bacteria and Archaea. *Front Microbiol* **8**, 2325, doi:10.3389/fmicb.2017.02325 (2017).
64 Peeters, S. H. & van Niftrik, L. Trending topics and open questions in anaerobic
ammonium oxidation. *Curr Opin Chem Biol* **49**, 45-52, doi:10.1016/j.cbpa.2018.09.022
(2019).
65 Akram, M. *et al.* A 192-heme electron transfer network in the hydrazine dehydrogenase
complex. *Sci Adv* **5**, eaav4310, doi:10.1126/sciadv.aav4310 (2019).
66 Bi, Z., Qiao, S., Zhou, J., Tang, X. & Zhang, J. Fast start-up of Anammox process with
appropriate ferrous iron concentration. *Bioresour Technol* **170**, 506-512,
doi:10.1016/j.biortech.2014.07.106 (2014).
67 Poulton, S. W. & Canfield, D. E. Ferruginous Conditions: A Dominant Feature of the
Ocean through Earth's History. *Elements* **7**, 107-112, doi:10.2113/gselements.7.2.107
(2011).
68 Ferousi, C. *et al.* Iron assimilation and utilization in anaerobic ammonium oxidizing
bacteria. *Curr Opin Chem Biol* **37**, 129-136, doi:10.1016/j.cbpa.2017.03.009 (2017).
69 Keren, N., Aurora, R. & Pakrasi, H. B. Critical roles of bacterioferritins in iron storage
and proliferation of cyanobacteria. *Plant Physiol* **135**, 1666-1673,
doi:10.1104/pp.104.042770 (2004).
70 Poulton, S. W., Fralick, P. W. & Canfield, D. E. Spatial variability in oceanic redox
structure 1.8 billion years ago. *Nature Geoscience* **3**, 486-490, doi:10.1038/Ngeo889
(2010).
71 Konhauser, K. O. *et al.* Iron formations: A global record of Neoarchaean to
Palaeoproterozoic environmental history. *Earth-Science Reviews* **172**, 140-177,
doi:10.1016/j.earscirev.2017.06.012 (2017).
72 Wu, Z. *et al.* Salinity-driven heterogeneity toward anammox distribution and growth
kinetics. *Appl Microbiol Biotechnol* **103**, 1953-1960, doi:10.1007/s00253-018-9521-4
(2019).
73 Parsons, C., Stueken, E. E., Rosen, C. J., Mateos, K. & Anderson, R. E. Radiation of
nitrogen-metabolizing enzymes across the tree of life tracks environmental transitions in
Earth history. *Geobiology* **19**, 18-34, doi:10.1111/gbi.12419 (2021).

74   Ader, M. *et al.* Ocean redox structure across the Late Neoproterozoic Oxygenation Event:
A nitrogen isotope perspective. *Earth and Planetary Science Letters* **396**, 1-13,
doi:10.1016/j.epsl.2014.03.042 (2014).
75   Schopf, J. W. Geological evidence of oxygenic photosynthesis and the biotic response to
the 2400-2200 ma "great oxidation event". *Biochemistry (Mosc)* **79**, 165-177,
doi:10.1134/S0006297914030018 (2014).
76   Chen, S. C. *et al.* The Great Oxidation Event expanded the genetic repertoire of arsenic
metabolism and cycling. *Proc Natl Acad Sci U S A* **117**, 10414-10421,
doi:10.1073/pnas.2001063117 (2020).
77   Buick, R. Did the Proterozoic 'Canfield Ocean' cause a laughing gas greenhouse?
Geobiology **5**, 97-100, doi:10.1111/j.1472-4669.2007.00110.x (2007).
78   Parks, D. H. *et al.* Recovery of nearly 8,000 metagenome-assembled genomes
substantially expands the tree of life. *Nat Microbiol* **2**, 1533-1542,
doi:10.1038/s41564-017-0012-7 (2017).
79   Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol Biol Evol* **30**, 772-780,
doi:10.1093/molbev/mst010 (2013).
80   Capella-Gutierrez, S., Silla-Martinez, J. M. & Gabaldon, T. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972-1973,
doi:10.1093/bioinformatics/btp348 (2009).
81   Nguyen, L. T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: a fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol*
Evol **32**, 268-274, doi:10.1093/molbev/msu300 (2015).
82   Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S.
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods* **14**,
587-589, doi:10.1038/nmeth.4285 (2017).
83   Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v4: recent updates and new
developments. *Nucleic Acids Research* **47**, W256-W259 (2019).
84   Yang, Z. H. PAML 4: Phylogenetic analysis by maximum likelihood. *Molecular Biology*
and Evolution **24**, 1586-1591 (2007).
85   Wang, H.-C., Minh, B. Q., Susko, E. & Roger, A. J. Modeling site heterogeneity with
posterior mean site frequency profiles accelerates accurate phylogenomic estimation.
Systematic biology **67**, 216-235 (2018).
86   Reis, M. D. *et al.* Using Phylogenomic Data to Explore the Effects of Relaxed Clocks
and Calibration Strategies on Divergence Time Estimation: Primates as a Test Case. *Syst*
Biol **67**, 594-615, doi:10.1093/sysbio/syy001 (2018).

87   Dos Reis, M., Donoghue, P. C. & Yang, Z. Bayesian molecular clock dating of species
divergences in the genomics era. *Nat Rev Genet* **17**, 71-80, doi:10.1038/nrg.2015.8
(2016).
88   Dos Reis, M. & Yang, Z. Approximate likelihood calculation on a phylogeny for
Bayesian estimation of divergence times. *Mol Biol Evol* **28**, 2161-2172,
doi:10.1093/molbev/msr045 (2011).
89   Quang, L. S., Gascuel, O. & Lartillot, N. Empirical profile mixture models for
phylogenetic reconstruction. *Bioinformatics* **24**, 2317-2323,
doi:10.1093/bioinformatics/btn445 (2008).
90   Morel, B., Kozlov, A. M., Stamatakis, A. & Szollosi, G. J. GeneRax: A tool for
Species-Tree-Aware Maximum Likelihood-based gene family tree inference under gene
duplication, transfer, and loss. *Mol Biol Evol* **37**, 2763-2774,
doi:10.1093/molbev/msaa141 (2020).
91   Kanehisa, M. & Goto, S. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic
Acids Research* **28**, 27-30 (2000).
92   Lu, S. *et al.* CDD/SPARCLE: the conserved domain database in 2020. *Nucleic Acids Res*
48, D265-D268, doi:10.1093/nar/gkz991 (2020).
93   Mitchell, A. L. *et al.* InterPro in 2019: improving coverage, classification and access to
protein sequence annotations. *Nucleic Acids Res* **47**, D351-D360,
doi:10.1093/nar/gky1100 (2019).
94   El-Gebali, S. *et al.* The Pfam protein families database in 2019. *Nucleic Acids Res* **47**,
D427-D432, doi:10.1093/nar/gky995 (2019).
95   Haft, D. H. *et al.* TIGRFAMs: a protein family resource for the functional identification
of proteins. *Nucleic Acids Res* **29**, 41-43, doi:10.1093/nar/29.1.41 (2001).
96   Saier, M. H., Jr. *et al.* The Transporter Classification Database (TCDB): recent advances.
Nucleic Acids Res **44**, D372-D379, doi:10.1093/nar/gkv1103 (2016).

 Figure legends

Figure 1. The evolutionary timeline of anammox bacteria using MCMCTree. The chronogram

was estimated based on the calibration set C1 (see Supplementary Text section 3). The blue bars

on the four calibration nodes and the last common ancestor (LCA) of anammox bacteria

represent the posterior 95% highest probability density (HPD) interval of the posterior time

estimates. These alternative calibration sets were selected by choosing the three that

accommodate different calibration constraints (see Supplementary Texts section 3.3). More

alternative time estimates are provided in Figure S4. The vertical grey bar represents the period

of the GOE from 2,500 to 2,320 Ma. The calibration constraints used within the phylum

Cyanobacteria are marked with orange texts: the LCA of Planctomycetes and Cyanobacteria

(Root), the total group of oxygenic Cyanobacteria (Node 2), the total group of *Nostocales* (Node

3), and the total group of *Pleurocapsales* (Node 4). The genome completeness estimated by

checkM is visualized with a gradient color strip. The right next color strip indicates the genome

type of genomic sequences used in our study including metagenome-assembled genomes (MAGs)

and whole-genome sequencing (WGS) from either enriched culture sample or isolate. The

diagram below the dated tree illustrates the change of atmospheric partial pressure of O_2 (P_{O_2})

and nitrogen isotope fractionations. The blue line shows the proposed model according to Lyons,

et al. [25]. The green arrow suggests the earliest evidence for aerobic nitrogen cycling at around 2.7

Ga. PAL on right axis means P_{O_2} relative to the present atmospheric level. The black dots denote

the change of nitrogen δ^{15}N isotope values according to Kipp, et al. [24].

Figure 2. The phyletic pattern of ecologically relevant genes in the comparison between

anammox bacteria and non-anammox bacteria. The phylogenomic tree on the left was

constructed with 887 genomic sequences described in Supplementary Texts section 2.1. Solid

circles at the nodes indicate the ultrafast bootstrap values in 1,000 bootstrapped replicates. Those

Planctomycetes genomes not used for comparative genomics analyses are collapsed into grey

triangles, and the numbers of collapsed genomes are labelled next to the triangles. The target

group and reference group for comparative genomic analysis are within an orange or blue box,

separately. For each genome, the genome completeness estimated by CheckM is visualized with

a color strip and labelled besides leaf nodes. The right next color strip represents the type of

genomic sequences used in our study including metagenome-assembled genomes (MAGs) and

whole-genome sequencing (WGS) from either enriched culture sample or isolate. The filled and

empty circles, respectively, represent the presence and absence of particular genes in

corresponding genomes. For gene clusters, only genomes with at least half of the members of the

gene cluster are indicated by a filled circle. The classifications of annotated genes are labelled

above the gene names. *hzsCBA*, hydrazine synthase subunit C, B and A; *hdh*, hydrazine

dehydrogenase; *hao*, hydroxylamine dehydrogenase; *nxrAB*, nitrite oxidoreductase subunit A and

B; *nirK*, copper-containing and NO-forming nitrite reductase; *nirS*, cytochrome NO-forming

nitrite reductase; *nrfAH*, ammonia-forming nitrite reductase subunit A and H; *glnB*, nitrogen

regulatory protein P-II; *nirC*, nitrite transporter; NRT, nitrate/nitrite transporter; *amt*, ammonium

transporter; kuste2805, 3603, 3605-3606, proposed genes relative to the synthetic pathways for

ladderane at Rattray, et al. [54]; *cbiG*, cobalt-precorrin 5A hydrolase; *cbiD*,

cobalt-precorrin-5B(C1)-methyltransferase; *cbiOMQ*, cobalt/nickel transport system; *AhpC*,

peroxiredoxin; *Ccp*, cytochrome *c* peroxidase; *dfx*, superoxide reductase; *fprA*, H_2O-forming

enzyme flavoprotein; *CcsAB*, cytochrome c maturation systems; *petB*, Cytochrome b subunit of

the bc complex; *petC*, Rieske Fe-S protein; *ahbABCD*, heme biosynthesis; *sat*, sulfate

adenylyltransferase; *aprAB*, adenylylsulfate reductase, subunit A and B; *fsr*, sulfite reductase

(coenzyme F420); *higB-1*, toxin; *higA-1*, antitoxin; *mnhABCDEG*, multicomponent Na^+/H^+

antiporter; *fdhAB*, formate dehydrogenase subunit A and B; *nuo*(A-N), NADH-quinone

oxidoreductase; *ndh*(A-N), NAD(P)H-quinone oxidoreductase; *nqrABCDEF*, Na^+-transporting

NADH:ubiquinone oxidoreductase; *rnfABCDEG*, Na^+-translocating ferredoxin:NAD^+

oxidoreductase; *atp*(A-H), F-type H^+-transporting ATPase; *lacAZ*, beta-galactosidase; *melA*, galA,

alpha-galactosidase; *ebgA*, beta-galactosidase; *galK*, galactokinase; , fructokinase; *fruK*,

1-phosphofructokinase; *rhaB*, rhamnulokinase; rbsK, ribokinase; *araB*, L-ribulokinase; *xylB*,

xylulokinase; *kdgK*, 2-dehydro-3-deoxygluconokinase; GALK2, N-acetylgalactosamine kinase;

cah, cephalosporin-C deacetylase; *argE*, acetylornithine deacetylase; *nagA*;

N-acetylglucosamine-6-phosphate deacetylase; *HDAC11*, histone deacetylase 11.