# Phylogenomic evidence for the Origin of Obligately Anaerobic Anammox Bacteria around the Great Oxidation Event

Tianhua Liao[1^], Sishuo Wang[1^], Eva E. Stüeken[2], Haiwei Luo[1*]

[1]Simon F. S. Li Marine Science Laboratory, School of Life Sciences and State Key Laboratory of Agrobiotechnology, The Chinese University of Hong Kong, Shatin, Hong Kong SAR

[2]School of Earth and Environmental Sciences and Centre for Exoplanet Science, University of St Andrews, Bute Building, Queen's Terrace, KY16 9TS, UK

[^]These authors contribute equally to this work.

[*]**Corresponding author:**
Haiwei Luo
The Chinese University of Hong Kong
Shatin, Hong Kong SAR
Phone: (+852) 39436121
E-mail: hluo2006@gmail.com

**Running title:** The origin time and phylogeny of anammox bacteria

**Keywords**: anammox bacteria, molecular dating analysis, Planctomycetes

**Abstract**

The anaerobic ammonium oxidation (anammox) bacteria could transform ammonium and nitrite to dinitrogen gas, and this obligate anaerobic process accounts for up to half of the global nitrogen loss for surface environments. Yet its origin and evolution, which may give important insights into the biogeochemistry of early Earth, remains enigmatic. Here, we performed comprehensive phylogenomic analysis and showed a single origin of anammox bacteria within the phylum Planctomycetes. After accommodating the uncertainties and factors influencing time estimates with different statistical methods, we estimated that anammox bacteria originated at around the so-called Great Oxidation Event (GOE; 2.32 to 2.5 billion years ago [Ga]) which is thought to have fundamentally changed global biogeochemical cycles. We further showed that during the origin of anammox bacteria, genes involved in oxidative stress, bioenergetics and anammox granules formation were recruited, which might have contributed to their survival on an increasingly oxic Earth. Our findings suggest the rising levels of atmospheric oxygen, which made nitrite increasingly available, was a potential driving force for the emergence of anammox bacteria. This is one of the first studies that link the GOE to the evolution of obligate anaerobic bacteria.

**Introduction**

Anaerobic ammonium oxidation (anammox, $NH_4^+ + NO_2^- \rightarrow N_2 + 2H_2O$), which usually

occurs in anoxic marine, freshwater and wetland settings, accounts for up to 50% of the removal

of fixed nitrogen (N) in nature. Along with denitrification, it is recognized as an important

biological process that leads to N loss from the environment [1,2]. In wastewater treatment,

anammox is more cost-effective and environmentally friendlier than denitrification due to its

lower oxygen requirement for aeration (nitrite [N(+III)], instead of nitrate [N(+V)] is sufficient

for the anammox metabolism), carbon-free cultivation (some denitrifying bacteria are

heterotrophic) and its negligible emissions of greenhouse gases like $N_2O$. Consequently,

anammox bacteria, the organisms that perform anammox, have been widely used in wastewater

treatment plants [2-5]. Despite the environmental and industrial importance of these organisms,

their evolutionary history and antiquity are poorly known, which hinders accurate

reconstructions of the biogeochemical nitrogen cycle over geologic time.

Previous studies investigated the roles of key genes driving anammox, including *hzsABC*

(hydrazine synthase)[6], *hdh* (hydrazine dehydrogenase), *hao* (hydroxylamine oxidoreductase) [7]

and *nxr* (nitrite oxidoreductase) [8]. In general, the vital enzymes encoded by these genes either are

directly involved in anammox or participate in replenishing electrons to the cyclic electron flow

[9,10]. Further, all known anammox bacteria are found within the phylum Planctomycetes [11]. Six

candidate genera of anammox bacteria, namely '*Candidatus* Brocadia', '*Candidatus* Kuenenia',

'Candidatus Jettenia', 'Candidatus Scalindua', 'Candidatus Anammoximicrobium' and

'Candidatus Anammoxoglobus', have been proposed based on 16S ribosomal RNA (rRNA) gene

sequences [12], but none of them have been successfully isolated into pure cultures. The habitat of

anammox bacteria requires the simultaneous presence of reduced (ammonia) and oxidized

(nitrite) inorganic N compounds. Such habitats are often found at the aerobic-anaerobic interface

in aquatic ecosystems, including the margins of oxygen minimum zones (OMZs) in the ocean

and sediment-water interfaces, where ammonium originates from the anaerobic degradation of

organic matter and nitrite can be produced by aerobic ammonia oxidation [13]. Ammonium, one of

the two substrates of the anammox metabolism, was thermodynamically stable and likely present

in moderate concentrations in the deep ocean throughout the Archean [14] and probably extending

well into the Proterozoic [15]. The availability of nitrite, the second important substrate, is less

certain. Geochemical data from sedimentary rocks suggest that the early Earth (before 3 Ga) was

fully anoxic and deficient in aerobic ecosystems that were able to generate nitrite/nitrate [16-18].

The first transient and/or localized occurrences of aerobic nitrogen cycling appear in the Mid- to

Neoarchean (2.7 Ga [19-22]), but evidence of widespread nitrate availability does not appear until

around 2.4 billion years ago (Ga), the so-called Great Oxidation Event (GOE) [23-25]. The GOE

marks the rise of free $O_2$ in the atmosphere above $10^{-5}$ times modern levels, and was ultimately a

result of the emergence of oxygenic cyanobacteria [26]. Hence it is conceivable that the appearance

of anammox was linked to the appearance of aerobic ecosystems around the time of the GOE.

However, it is also possible that some nitrite was provided much earlier, through lightning

reactions in the Archean atmosphere [27]. Lightning, even in the absence of $O_2$, can generate NO

gas, which dissolves in water and converts into aqueous species, including nitrite [28]. Some

anammox bacteria are even capable of using NO rather than nitrite directly as a substrate [9]. If

these organisms capitalized on the lightning flux, then anammox might long pre-date the GOE.

Several tools exist for investigating the link between the evolution of metabolic pathways

and geo-environmental transformations. One way to explore the evolutionary history of a

specific metabolic pathway is based on organic biomarkers, which, however, are often affected

by the poor preservation over geologic timescales [29]. For instance, ladderanes, a type of lipids

that is unique to anammox bacteria [30], are rarely preserved to a level that can be used to date their

evolutionary origin. Another approach is based on nitrogen isotope ratios of sedimentary records

[14,31,32]. However, the isotopic fractionation factors for different metabolic pathways overlap

widely (denitrification: -5 to -30‰; anammox: -16 or -24‰ [15]), making it difficult to single out

and elucidate the evolution of the different redox reactions within the N cycle by this method [33].

Alternatively, molecular dating, which estimates the age of the last common ancestor (LCA) of

analyzed lineages by comparing their sequences based on the molecular clock theory [34], provides

an alternative strategy to investigate this issue. Briefly, it can estimate the divergence timescale

of organisms using genetic data while accounting for issues like uncertainties in the calibrations

and different evolutionary rates among lineages, thereby bypassing the paucity and uncertainty of

biomarkers and other biogeochemical records. On the one hand, because anammox is an

anaerobic reaction, it is intuitive to assume that anammox should originate before the GOE,

perhaps using nitrite or NO provided by lightning reactions in the atmosphere [35]. On the other

hand, it is because of GOE that $O_2$ on Earth rose permanently to a concentration that is

biologically meaningful, and as a result, (micro-)environments were created in the surface ocean

that were rich in nitrate/nitrite produced by aerobic organisms. These redox-stratified conditions

may have stimulated the anammox metabolism by providing both nitrite and ammonium at the

redoxcline. In this regard, it could be hypothesized that anammox did not arise until the GOE.

Here, we ask the question when anammox originated. To answer this question and to obtain more

insights into the evolution of anammox bacteria, we compiled an up-to-date genomic data set of

Planctomycetes (see Supplementary Text section 1; Dataset S1.1), placed the evolution of

anammox bacteria into the context of geological events, and investigated genomic changes

associated with the origin of this ecologically important bacterial lineage.

**Results and discussion**

<u>Monophyletic origins of anammox bacteria and anammox genes</u>

The anammox bacteria form a monophyletic group within the phylum Planctomycetes in

a comprehensive phylogenomic tree with 881 Planctomycetes genomes (Fig. S1). The anammox

bacteria clade in this phylogenomic tree (Fig. S1) comprises four known genera including the

early-branching *Ca.* Scalindua and *Ca.* Kuenenia, and the late-branching *Ca.* Jettenia and *Ca.*

Brocadia. It also includes two understudied lineages which we named 'basal lineage' and

'*hzsCBA*-less lineage' (Fig. S1), both of which have relatively low genome completeness

compared to the other anammox bacteria and are represented by metagenome-assembled

genomes (MAGs) sampled from groundwater (see Supplementary Text section 2.1). Note that

the absence of *hzsCBA* in several genomes within the '*hzsCBA*-less lineage' might be ascribed to

the loss of these genes in evolution or their low genome completeness (Fig. S1). Because of their

shallow (late-branching) phylogenetic position, this uncertainty should not affect the inference of

the origin of anammox bacteria. Two described genera (*Ca.* Anammoximicrobium and *Ca.*

Anammoxoglobus) do not have genomes available by the time of the present study (last accessed:

April 2021), and are therefore not included in the phylogenomic analysis. Furthermore, using the

last updated set of 2,077 Planctomycetes genomes (retrieved in April 2021) from NCBI, we

obtained a consistent topology of anammox bacteria (Fig. S2).

To check whether important groups, particularly early-branching lineages, have been

encompassed in our analyzed genome data sets, we built a 16S rRNA gene tree (Fig. S3) using

the identified 16S rRNA genes from genomic sequences of anammox bacteria and the deposited

16S rRNA gene amplicons in SILVA from the class Brocadiae, to which anammox bacteria

belong (see Supplementary Texts section 2.2). Since the anammox bacteria with genome

sequences in this 16S rRNA gene tree (Fig. S3) showed a branching order congruent with the

topology of the phylogenomic tree (Fig. S1), and since the 16S rRNA gene tree included 913

(clustered from 20,142) sequences sampled from a wide array of habitats (marine, sediments,

man-made reactor, freshwater and terrestrial ecosystems), the 16S rRNA gene phylogeny, which

could better capture the diversity of anammox bacteria by including uncultured samples, likely

encompasses the early-branching lineages of anammox bacteria. Taken together, our genome set,

which encompasses the genomes of the early-branching lineages, is appropriate to be used to

study the evolutionary origin of anammox bacteria (Dataset S1.3). We further analyzed the

evolution (Fig. S4) of the genes (*hzsCBA*) critical to the anammox reaction [7,36]. Reconciliation

(see Supplementary Texts section 2.3) of the gene and the genome-based species phylogenies

pointed to a single origin of anammox metabolism at the LCA of anammox bacteria (Fig. S4).

Taken together, the above analyses suggest that our genome data sets have encompassed

the earliest-branching lineages and deep phylogenetic diversity of anammox bacteria, and

indicate its monophyletic origin, which agrees with previous studies [37-39]. This should provide a

foundation for estimating the origin of anammox metabolism by estimating the origin time of

anammox bacteria.

The evolutionary origin of anammox bacteria coincided with the rising $O_2$

We employed MCMCTree to estimate the divergence times of the anammox bacteria with

i) soft bounds (time constraint allowing a small probability of violation) based on cyanobacteria

fossils, and ii) the relaxed molecular clock approaches allowing substitution rates to vary among

branches. Using a best-practice dating scheme (Fig. 1; see Supplementary Text section 3), we

dated the origin of anammox bacteria to 2,117 million years ago (Ma) (95% highest posterior

density [HPD] interval, 2,002 - 2,226 Ma). Further, we repeated the analysis based on the

expanded Planctomycetes genomes (Genome set 2; Dataset S1.1) and the constraint topology

inferred with a profile mixture model (LG+C60+G; Genome set 1; Dataset S1.1). In general,

both analyses based on different genome sets resolved consistent topologies (Fig. S5) as shown

in Figure 1. Overall, they gave similar posterior times of the LCA of the anammox bacteria at

2,105 Ma (95% HPD: 1,961 - 2,235 Ma) or 2,005 Ma (95% HPD: 1,869 - 2,146 Ma).

Analyses with different combinations of calibrations and parameters broadly converge to

similar time estimates (Fig. S6; Dataset S2.1). Specifically, assigning a more ancient time

constraint to the total group of oxygenic cyanobacteria, which is based on the biogeochemical

evidence of the presence of $O_2$ at nearly 3.0 Ga [40] instead of GOE (Dataset S2.1), shifted the

posterior dates of the LCA of anammox bacteria into the past by ~300 million years (Myr) (e.g.,

C1 versus C10; Fig. S6). Varying the maximum bounds of the root by using 4.5 Ga, 3.8 Ga or 3.5

Ga had a minor impact (around 100 Myr) on the time estimates of anammox bacteria (e.g., C1

versus C3; Fig. S6). Though the auto-correlated rates (AR) model was less favored than the

independent rates (IR) model based on the model comparison (see Supplementary Text section

3.3; Dataset S2.2), it is worth noting that the posterior ages estimated by the AR model were 1-12%

younger than estimated by the IR model (Dataset S2.1). Besides, there remains a possibility that

there are unsampled or even extinct lineages of anammox bacteria that diverged earlier than all

anammox bacteria analyzed in the present study. This scenario, if true, hints that the first

anammox bacterium could have originated before the occurrence of the LCA of sequenced

anammox bacteria but later than their split from the sister non-anammox lineage (Node X in Fig.

1; up to 2,600 Ma). Accommodating the above uncertainties, our analyses suggest that the origin

of anammox bacteria, and hence the origin of anammox, most likely falls into the 2.6-2.0 Ga

interval, or more generally, around the early Proterozoic. Running MCMCTree analysis with no

sequence data showed different distributions of time estimates for both the clade of anammox

bacteria and the four calibration points (Fig. S7), suggesting that sequence data are informative

for our molecular clock analysis [41]. Besides, we estimated similar origin times of anammox at

around 2.1 Ga (the AR model) or 2.3 Ga (the IR model; see Fig. S8) with the complete taxon

sampling shown in Fig. S2 (Genome set 2; see Dataset S1.1). For this analysis, we used a

likelihood-based dating algorithm since the Bayesian molecular clock analysis is known to be

computationally intensive and not suitable to take a data set with a broad taxon sampling [42]. Note

that unlike Bayesian molecular dating software, the likelihood-based method we used only

makes point estimations for the node ages (no standard deviation), thereby not allowing for fully

appreciating the uncertainties in age estimates and any inference based on that. In general, the

above result suggests that taxon sampling may not affect our dating analysis.

### The geochemical context of the origin of anammox bacteria

We speculate that the timing of the origin of the anammox metabolism is linked to the

increasing availability of nitrite in surface environments, because ammonium was likely present

in the deep anoxic ocean throughout the Precambrian [14,15] and therefore presumably not a

limiting substrate. The first abiotic source of nitrogen oxides on the early Earth, including NO

and nitrite, would have been lightning reactions between $N_2$ and $CO_2$ in the Archean atmosphere

[43,44]. A prior study suggested that this process led to micromolar levels of nitrite in seawater [43],

and the supply of NO may have been even higher, considering that it is the initial reaction

product of lighting [43,44]. This implies the possibility of an earlier origin of anammox bacteria. In

fact, an intriguing hypothesis is that NO-dependent anammox [12] is ancestral to all anammox

bacteria and arose before the appearance of significant nitrite levels with the GOE, perhaps

driven by the lightning-source of NO. However, the original estimates of the lightning-derived

flux of nitrogen oxides may have been based on perhaps unrealistically high amounts of $CO_2$ [45].

Furthermore, to our knowledge, there has been no isotopic evidence in the early Archean rock

record prior to 3 Ga for the presence of a significant nitrite or nitrate reservoir in the early ocean

[16-18], and it is relatively conceivable that any nitrogen oxides that were supplied to the ocean by

lightning were rapidly reduced to ammonium or $N_2$ by ferrous iron, possibly even abiotically [46,47].

Hence, although a background lightning flux of nitrogen oxides almost certainly existed, to our

knowledge, there seems little evidence that it could supply a reliable metabolic substrate for

early life. Furthermore, it is important to note that our phylogenetic results are not supportive

of this hypothesis since the NO-utilizing anammox bacteria discovered to date are affiliated

with phylogenetically shallow lineages from *Ca.* Kuenenia and *Ca.* Brocadia (Fig. 1). To our

knowledge, there have been no studies reporting the ability of using NO as the sole electron

acceptor in any other lineages of anammox bacteria. Hence, if NO-dependent anammox would

have been present in the LCA of anammox bacteria, it was not clear why it was lost in all

other anammox bacteria which resulted from multiple independent losses of the trait, a

scenario apparently not favored from an evolutionary perspective (as shown by the arrows in

Fig. S2). There are also anammox bacteria that are capable of using hydroxylamine as an

oxidant; however, this molecule is a highly reactive compound and is usually considered as an

intermediate in the nitrogen cycle, thereby rendering it a less important compound from the

perspective of geological timescales. We are not aware of any non-biological sources of

hydroxylamine. Taken together, our results are inconsistent with an Archean emergence of the

anammox metabolism driven by lightning-derived nitrogen oxides. Instead, our results are most

parsimoniously explained by the appearance of more significant nitrite/nitrate reservoirs around

the time of the GOE.

The first transient appearances of nitrite/nitrate-dependent metabolisms are captured by

the sedimentary nitrogen isotope record in late Mesoarchean soils at 3.2 Ga [20] and in Neoarchean

shallow-marine settings at 2.7 Ga and 2.5 Ga [19,21,22]. These observations may reflect local and/or

temporally restricted oxygen oases [48,49]. Widespread nitrite/nitrate availability is inferred for the

Paleoproterozoic (2.4-1.8 Ga), i.e., in the immediate aftermath of the GOE [23-25]. Importantly, $O_2$

is necessary, although a trace amount is feasible, to extant nitrifying organisms for the oxidation

of ammonium to nitrite and nitrate [50-52]. We noticed that a recent study [53] reports the ability of the

model ammonia-oxidizing archaeal species (*Nitrosopumilus maritimus*) to continue ammonia

oxidation after consuming all supplied $O_2$ by nitric oxide disproportionation to generate $O_2$, but

the underlying genetic pathway is not known, making it difficult to extrapolate this mechanism

to any other ammonia-oxidizing organisms. Hence the rise of oxygen almost certainly triggered

the growth of the nitrite/nitrate reservoir, and therefore potentially provided one of the key

substrates for anammox bacteria. We note that also NO would have become more abundant after

the GOE, because it is an intermediate product in nitrification and denitrification [54]. This would

put the possibility of NO-driven anammox back on the table. However, NO levels in seawater are

highly variable, and so nitrite would likely have been a more reliable substrate. In any case, it

would not violate our conclusion that the appearance of anammox in the Paleoproterozoic was

linked to the GOE and directly or indirectly to the rise of marine nitrite. It has been shown that a

low concentration of nitrite significantly decreases the rate of N removal by anammox bacteria in

reactors [55,56]. In modern OMZ in the Bay of Bengal, which may to some extent serve as

analogues for the Paleoproterozoic redox-stratified ocean, the concentration of nitrite, instead of

ammonium, was proposed as the rate-limiting factor to the anammox metabolism [57]. These imply

that the rise of nitrite, which is driven by the rise of $O_2$, could have facilitated the appearance of

anammox bacteria in the early Proterozoic.

In modern environments, the majority of nitrite used in anammox is derived from nitrate

reduction or aerobic ammonia oxidation [58], the latter performed by ammonia-oxidizing archaea

(AOA) or bacteria (AOB). As noted above, nitrification could have appeared by at least 2.7 Ga,

as indicated by nitrogen-isotope evidence for local accumulation of nitrate in surface ocean

waters [15,59,60]. A recent study inferred that the LCA of AOA dates back to ~2.3 Ga and first

appeared on land, driven by increasing $O_2$ concentrations in the atmosphere at that time, and that

the expansion of AOA from land to the ocean did not occur until nearly 1.0 Ga [61]. If true, this

hints at a dominant role of AOB in early ocean, before 2.3 Ga. Consistent with this idea, the vast

majority of the early-branching lineages of anammox bacteria (*Ca.* Scalindua) and the sister

Planctomycetes lineages of the anammox clade in the phylogenomic tree (Fig. 1) are found in

marine environments. Furthermore, in the 16S rRNA gene tree which represents a greater

diversity than the phylogenomic tree, marine lineages still account for the majority of the

earliest-branching lineages of anammox bacteria (Fig. S3). Thus, it seems likely that the LCA of

anammox bacteria originated in the marine realm where AOB thrived first. The above arguments,

although speculative, tentatively suggest that the nitrite demand for anammox had been readily

met by GOE by taking advantage of the significant amount of $O_2$ newly available, which is

broadly consistent with our molecular dating results suggesting an origin of anammox shortly

before or after the GOE.

<u>Genomic changes potentially related to anammox metabolism and habitat adaptation upon the</u>

<u>origin of anammox bacteria</u>

We further explored the genomic changes characterizing the origin of anammox bacteria

(see Supplementary Text section 4). This includes the gains of the aforementioned genes that

directly participate in anammox, viz., *hzsABC*, *hdh*, and *hao*. The current view of the anammox

metabolism posits that electrons consumed by anammox for carbon fixation are replenished by

the oxidation of nitrite to nitrate by nitrite oxidoreductase (NXR) [10], which was inferred to be

acquired upon the origin of anammox bacteria (Fig. 2). Moreover, multiple auxiliary genes for N

assimilatory pathways including N regulatory protein (*glnB*), assimilatory nitrite reductase large

subunit (*nirB*), transporters like *amt* (ammonium), *nirC* (nitrite) and NRT family proteins

(nitrate/nitrite), were also gained at the origin of anammox bacteria (Fig. 2). However, genes for

nitrogen-related dissimilatory pathways encoding nitrite reduction to nitric oxide (*nirK* or *nirS*)

and to ammonium (*nrfAH*), respectively, were likely acquired after the origin of anammox

bacteria (Fig. 2). Anammox occurs at the membrane of anammoxosome, an organelle

predominantly composed of a special type of lipid called ladderane. The ladderane membranes

show low proton permeability, which helps maintain the proton motive force during the

anammox metabolism [30]. Although the biosynthetic pathway of ladderane is yet to be

characterized, a previous study [62] predicted 34 candidate genes responsible for the synthesis of

ladderane lipids, four of which were potentially acquired during the origin of anammox bacteria

(see Supplementary Text section 4.2; Fig. 2), hinting at their potential roles in ladderane

synthesis and providing clues to future experimental investigations.

Anammox bacteria occur in anoxic habitats and exhibit low oxygen tolerance [12].

Specifically, the $O_2$-sensitive intermediates generated by anammox, e.g., hydrazine, a powerful

reductant, requires anoxic conditions. Accordingly, the likely acquired peroxidases include

cytochrome c peroxidase (Ccp), which could scavenge hydrogen peroxide in the periplasm [63],

and the most prevalent peroxidase [64], thioredoxin-dependent peroxiredoxin (AhpC). Besides

peroxide scavengers, the desulfoferrodoxin (Dfx), which functions as superoxide reductase (SOR)

to reduce superoxide to hydrogen peroxide and which is broadly distributed among anaerobic

bacteria [64], were likely acquired during the origin of anammox bacteria (Fig. 2). Another

acquired gene is *fprA*, which encodes flavo-diiron proteins that scavenge $O_2$.

Investigating other metabolisms that were also acquired upon the origin of anammox

bacteria allows reconstructing the coeval ecology. Our data show that *sat* (Fig. 2), which encodes

sulfate adenylyltransferase for assimilatory incorporation of sulfate into bioavailable adenylyl

sulfate, as well as *aprAB* and *fsr* (Fig. 2) for encoding dissimilatory sulfite reductase, were

acquired upon the origin of anammox bacteria (Fig. 2). This acquired capability of sulfate

metabolism is coincident with the increased concentrations of seawater sulfate from ~100 μM

or less throughout much of the Archean [65,66] to over 1mM after the rise of atmospheric $O_2$ [67,68].

Unlike other Planctomycetes, anammox bacteria are generally autotrophs that use the

Wood-Ljungdahl pathway for carbon assimilation [69]. Consequently, they potentially lost many

genes involved in carbohydrate utilization (Fig. 2). The metabolic loss is further strengthened by

the enrichment analysis where the genes predicted to be lost by the LCA of anammox bacteria

are enriched in pathways involving hydrolase activity, intramolecular oxidoreductase activity and

carbohydrate kinase activity (Fig. 2; see Data and Code availability), hinting at a decreased

ability of anammox bacteria to degrade organics.

Additionally, iron is a vital element for HZS [6] and HDH [70], and *in vitro* studies revealed

that increased iron concentrations can promote the growth of anammox bacteria [71]. The ancient

oceans are thought to have been ferruginous (high iron availability) according to sedimentary

records of iron speciation across several coeval marine settings from the Archean through most

of the Proterozoic [72], meaning that large amounts of soluble iron would have been available for

anammox bacteria. Interestingly, a series of iron-related genes to make use of the abundant iron

from the environment were likely acquired upon the origin of anammox bacteria. First, anammox

bacteria likely acquired the gene *fur*, which encodes a ferric uptake regulator for iron

assimilation. Moreover, the iron-containing proteins, specifically cytochrome, are encoded by the

acquired genes *petBC*, which potentially encodes Rieske-heme b [10], and *CcsAB*, which encodes a

protein involved in cytochrome c maturation systems (Fig. 2). Notably, *ahbABCD*, which

encodes proteins-synthesizing *b*-type hemes in anammox bacteria [73], were likely acquired before

the origin of anammox bacteria (Fig. 2). Furthermore, we identified that the gene *Bfr* that

encodes the oligomeric protein, bacterioferritin involved in the uptake and storage of iron [74] was

likely acquired during the origin of anammox bacteria (Fig. 2), which may help to hoard iron in

settings where iron became locally sparse, such as along euxinic (sulfide-rich) or oxic marine

margins in the Paleoproterozoic [75]. A recent study highlights the vital role of bacterioferritin to

the anammox regulation [76]. It is thought that the appearance of euxinic margins around the time

of the GOE trapped iron that upwelled from the deep ocean, leading to the demise of iron oxide

deposits (i.e., banded iron formations) [77]. Our results tentatively support this model, if the

microenvironments that those primordial anammox bacteria colonized were located near the

euxinic-oxic interface where nitrite and ammonium were available while upwelled iron was

titrated out of the water column by freely dissolved hydrogen sulfide.

Taken together, the timelines and corresponding genomic changes of anammox bacteria

provide implications for the physiological characteristics of descendant anammox bacteria and

the origin of other nitrogen-transforming pathways in the context of an estimated evolutionary

timeline of anammox bacteria. For example, the earlier branching lineage *Ca.* Scalindua exhibit

high nitrite affinity (0.45 μM) compared to other anammox genera (up to 370 μM) [12,78], which

may reflect a progressive growth of the marine nitrite reservoir, linked to the protracted

oxygenation of the surface environments over geologic timescales [26]. Likewise, genomic data

reveal that also canonical denitrification radiated across the tree of life after the GOE [79], and the

gains (Fig. S9; see Supplementary Text section 4.4) of genes *nrfAH* for dissimilatory nitrate

reduction to ammonium (DNRA) at the late-branching lineages (*Ca.* Brocadia and *Ca.* Kuenenia)

at around 800 Ma are consistent with the time when the nitrite/nitrate availability was increased

by the Neoproterozoic Oxygenation Event [80]. The time estimate and genomic changes of the

anammox bacteria have gone some way towards enhancing the understanding of the

physiological characteristics of modern anammox bacteria and the historical N cycle.

**Caveats and concluding remarks**

In the present study, we link an estimated evolutionary timeline of anammox and

geological context of early Earth on a simplified view that nitrite-dependent anammox is

ancestral to anammox bacteria. However, there are several limitations. First, dating the

bacterial evolution has many challenges, which, among others, include the paucity of

appropriate fossils, and a long evolutionary distance between anammox bacteria and

cyanobacteria [81]. Hence, the date estimates should be read as potential ranges of time

(indicated by the 95% HPD interval), instead of accurate time points, under the molecular

clock model. Moreover, it is worth pointing out the metabolism called feammox, a process

where ferric instead of nitrite is taken as the electron acceptor for ammonia oxidation. Both

feammox and anammox can anaerobically convert ammonia into dinitrogen [82], but the former

process is supposed to be carried out by Acidimicrobiaceae from the Actinobacteria phylum[83,84]

and under anaerobic environments without the presence of anammox bacteria [85]. Likewise, a

recent experimental study [86] proposed extant anammox bacteria as electroactive organisms and

suggested the feasibility to utilize graphene oxides and man-made metallic electrodes mimicking

metal oxides as the electron acceptor. However, the lack of detailed molecular mechanism

hampers the inference of its evolutionary history. In any case, metal oxides such as $Fe_2O_3$ or

$MnO_2$ also became more abundant in the Neoarchean to Paleoproterozoic, i.e. around the time of

the GOE [77,87], and hence even if these oxides were used as substrates for anammox, it would not

violate our overall conclusion that the GOE played a primary role in driving the evolution of this

metabolic pathway. However, it is also worth noting that metal oxides are solids and thus require

extra-cellular electron transfer [86]. An entirely intra-cellular metabolism, using dissolved nitrite,

may be more likely to evolve first, especially given that nitrate became bioavailable in the ocean

around the same time (see above).

Another important aspect to highlight about our approach is that, our analysis began with

inferring the LCA of anammox bacteria based on the presence of *hzsCBA*, a common strategy in

phylogenetic studies. It is important to note that such an enzyme complex encoded by multiple

genes might originate in different genomic backgrounds and were presented together in a suitable

genomic background later in evolution. This scenario, if true, indicates that the appearance of

any single subunit of *hzsCBA* genes, which could not assembly a functional hydrazine synthase,

should predate the LCA of anammox bacteria. Nevertheless, the alternative hypothesis would not

affect our inference and following analysis since we only focus on the inferred ancestor of extant

anammox bacteria, a strategy commonly used in modern phylogenetic analysis. These alternative

possibilities, although not affecting the molecular clock analysis, should be carefully considered

when making inference based on our estimated evolutionary timeline of anammox bacteria.

Here, using molecular dating and comparative genomics approaches, we link the

emergence of an obligate anaerobic bacterial group, which drives the loss of fixed N in many

environments, to the rise of $O_2$, and highlight their evolutionary responses to major

environmental disturbances. Apparently, the GOE opened novel niches for the origin and

subsequent expansion of diverse aerobic prokaryotes such as AOA and AOB, which in turn

facilitated the origin of other bacterial lineages, including some anaerobes like the anammox

bacteria, by providing the resources for their energy conservation. Our results open up the

possibility that a significant proportion of the Proterozoic nitrite budget was consumed by

anammox bacteria, and that the sedimentary nitrogen isotope record might be influenced by their

activity. Our findings may have implications for Proterozoic climate, because $N_2O$ has

previously been invoked as an important greenhouse gas at that time [88], but a relatively higher

contribution of anammox bacteria to the marine nitrogen cycle may have hampered $N_2O$

production [3]. Lastly, our study suggests that the impacts of the GOE go well beyond aerobic

prokaryotes. We therefore conclude that molecular dating is a likely feasible approach to

complement isotopic evidence for resolving the timeline of biological evolution and for

providing additional constraints on climate models of the distant past.

**Materials and Methods**

Overall, we compiled two genome sets and one 16S rRNA gene set in our study (Dataset

S1.1). A phylogenomic tree (Fig. S1) was generated based on the concatenated alignment of 120

ubiquitous proteins (bac120; Dataset S1.3) proposed for tree inference by the Genome Taxonomy

Database [89]. All published genomic sequences (952 in total) affiliated with the phylum

Planctomycetes in NCBI Genbank up to December 2019 were retrieved (Dataset S1.2). To

further examine whether the patterns obtained with this Planctomycetes dataset hold the same,

we further constructed an expanded set of Planctomycetes genomes by retrieving a total of 2,077

Planctomycetes genomes released by April 2021 at Genbank (Fig. S2; see Data and Code

availability). Key anammox genes *hzsCBA* and *hdh* were identified against manually curated

reference sequences by BLASTP. For each gene, identified protein sequences were aligned using

MAFFT (v7.222) [90] and the alignments were refined by trimAl (v1.4) [91]. All phylogenies were

constructed by IQ-tree (v1.6.2) [92] with substitution models automatically selected by

ModelFinder [93] and branch support assessed with 1,000 ultrafast bootstrap replicates. Note that

the constraint topologies for dating analysis (Fig. 1; Fig. S5) were inferred with the profile

mixture model (LG+C60+G) which better accommodates across-site heterogeneity in deep-time

evolution. Following manual curation, four well-recognized genera and two separate lineages of

anammox bacteria were highlighted with different colours, and the presence of key genes were

annotated with filled symbols beside the labels (Fig. S1). Furthermore, the phylogenomic tree

was generated with similar methods using 16S rRNA genes identified from downloaded genomes

and those retrieved from the SILVA database (Fig. S3; Dataset S1.3). All trees (including

phylogenomic and 16S trees) in our study were visualized with iTOL [94]. More details are shown

in supplementary text, section 2.

Molecular dating analysis was carried out using the program MCMCTree from the PAML

package (4.9j) [95]. In our study, 13 calibration sets constructed with different time constraints of

the root and three calibration nodes within the cyanobacteria lineage were used (see

Supplemental Texts section 3.2). The topology constraint for dating analysis was generated using

85 genomes, which were sampled from the phylogenomic tree of phylum Planctomycetes (see

Supplemental Texts section 3.1), and the model LG+C20+F+G under posterior mean site

frequency (PMSF) approximation [96]. To perform dating analysis, clock models and different

calibration sets were compared in 26 schemes (see Supplemental Texts section 3; Dataset S2.1)

[97,98]. For each scheme, the approximate likelihood method [99] of MCMCTree were conducted in

duplicate with identical iteration parameters (burn-in: 10,000; sample frequency: 20; number of

sample: 20,000). The convergence of each scheme was evaluated by comparing the posterior

dates of two independent runs (see Data and Code availability). With the updated genome set 2

(Dataset S1.1), we repeated the taxon sampling process and generated an alternative phylogeny

for dating analysis (Fig. S5) using LG+C20+F+G model (C20: 20 classes of site-specific amino

acid profiles) under PMSF approximations. Similarly, we also generated another phylogeny for

dating analysis using the same genome set as used in Fig. 1 but with the profile mixture model

(LG+C60+G) which uses 60 classes of amino acid profiles [100]. Note that the missing parameter

+F would not significantly change the results since it only adds another profile amino acid

calculated from the original data. These two repeat dating analyses were conducted with the best

practice dating scheme (IR model and calibration C1). Moreover, we performed likelihood based

dating analyses [101] (Fig. S8) using the phylogenomic tree (Fig. S2) and the calibration C1 (see

Data and Code availability) implemented in ape [42].

The phylogenetic differences between gene trees and the species tree were reconciliated

by GeneRax [102] with unrooted gene tree as input and automatically optimized duplication,

transfer, loss (DTL) rates. For each gene identified from the genome set 2 (Dataset S1.1), we

used a species tree comprised by all anammox bacteria pruned from the phylogenomic tree (Fig.

S2) as the reference. We used recommended parameters including SPR strategy, undated DTL

model and a maximum radius of five.

For comparative genomics analysis, the protein-coding sequences were annotated against

KEGG [103], CDD [104], InterPro [105], Pfam [106], TIGRFAM [107] and TCDB [108], individually (see

Supplemental Texts section 4). Following annotations, the potentially gained/lost genes between

groups (anammox bacteria versus non-anammox bacteria; an anammox bacterial genus versus all

other anammox bacteria) were statistically tested using two-sided fisher's exact tests. Finally, the

resulting $p$-values were corrected with the Benjamini-Hochberg FDR (false discovery rate)

procedure. Moreover, genes with corrected $p$-values smaller than 0.05 and with larger ratio in the

study group were defined as gained genes, and genes with corrected $p$-values smaller than 0.05

and with smaller ratio in the study group were defined as potentially lost genes. Those potentially

gained/lost genes were summarized for their detailed functions (see Data and Code availability).

**Acknowledgements**

We thank Bite Pei for building the basic dataset for this project, Yang Qian for his help in

editing the earlier version of the manuscript, Jinjin Tao for her helpful discussion, and Mario dos

Reis for his suggestion on dating analysis. This work is funded by the National Science

Foundation of China (92051113), the Hong Kong Research Grants Council Area of Excellence

Scheme (AoE/M-403/16), the Direct Grant of CUHK (4053495), and The CUHK Impact

Postdoctoral Fellowship Scheme to (S.W.).

**Data and Code availability**

All used genomic sequences, generated phylogenetic trees, estimated divergence times

and the used python codes are deposited in the online repository

https://github.com/luolab-cuhk/anammox-origin.

**Conflict of interest**

The authors declare that they have no conflict of interest.

**References**

1      Oshiki, M., Satoh, H. & Okabe, S. Ecology and physiology of anaerobic ammonium
oxidizing bacteria. *Environ Microbiol* **18**, 2784-2796, doi:10.1111/1462-2920.13134
(2016).
2      Stein, L. Y. & Klotz, M. G. The nitrogen cycle. *Current Biology* **26**, R94-R98, doi:DOI
10.1016/j.cub.2015.12.021 (2016).
3      Li, L. *et al.* $N_2O$ emission in partial nitritation-anammox process. *Chinese Chemical*
*Letters* **31**, 28-38, doi:10.1016/j.cclet.2019.06.035 (2020).
4      Okabe, S., Oshiki, M., Takahashi, Y. & Satoh, H. N2O emission from a partial
nitrification-anammox process and identification of a key biological process of N2O
emission from anammox granules. *Water Research* **45**, 6461-6470,
doi:10.1016/j.watres.2011.09.040 (2011).
5      Gao, D. W. & Tao, Y. Versatility and application of anaerobic ammonium-oxidizing
bacteria. *Applied Microbiology and Biotechnology* **91**, 887-894,
doi:10.1007/s00253-011-3411-3 (2011).
6      Dietl, A. *et al.* The inner workings of the hydrazine synthase multiprotein complex.
*Nature* **527**, 394-397, doi:10.1038/nature15517 (2015).
7      Kartal, B. *et al.* Molecular mechanism of anaerobic ammonium oxidation. *Nature* **479**,
127-130, doi:10.1038/nature10453 (2011).

8     Chicano, T. M. *et al.* Structural and functional characterization of the intracellular
filament-forming nitrite oxidoreductase multiprotein complex. *Nature Microbiology*,
doi:10.1038/s41564-021-00934-8 (2021).
9     Hu, Z., Wessels, H., van Alen, T., Jetten, M. S. M. & Kartal, B. Nitric oxide-dependent
anaerobic ammonium oxidation. *Nat Commun* **10**, 1244,
doi:10.1038/s41467-019-09268-w (2019).
10    Kartal, B. & Keltjens, J. T. Anammox Biochemistry: a Tale of Heme c Proteins. *Trends*
*Biochem Sci* **41**, 998-1011, doi:10.1016/j.tibs.2016.08.015 (2016).
11    Wang, Y., Xu, L. Y., Wang, S. Y., Ye, F. & Zhu, G. B. Global Distribution of Anaerobic
Ammonia Oxidation (Anammox) Bacteria - Field Surveys in Wetland, Dryland,
Groundwater Aquifer and Snow. *Frontiers in Microbiology* **10**,
doi:10.3389/fmicb.2019.02583 (2019).
12    Zhang, L. & Okabe, S. Ecological niche differentiation among anammox bacteria. *Water*
*Res* **171**, 115468, doi:10.1016/j.watres.2020.115468 (2020).
13    Sonthiphand, P., Hall, M. W. & Neufeld, J. D. Biogeography of anaerobic
ammonia-oxidizing (anammox) bacteria. *Front Microbiol* **5**, 399,
doi:10.3389/fmicb.2014.00399 (2014).
14    Yang, J. *et al.* Ammonium availability in the Late Archaean nitrogen cycle. *Nature*
*Geoscience* **12**, 553-557, doi:10.1038/s41561-019-0371-1 (2019).
15    Stueken, E. E., Kipp, M. A., Koehler, M. C. & Buick, R. The evolution of Earth's
biogeochemical nitrogen cycle. *Earth-Science Reviews* **160**, 220-239,
doi:10.1016/j.earscirev.2016.07.007 (2016).
16    Ossa Ossa, F. *et al.* Limited oxygen production in the Mesoarchean ocean. *Proc Natl*
*Acad Sci U S A* **116**, 6647-6652, doi:10.1073/pnas.1818762116 (2019).
17    Stueken, E. E., Buick, R., Guy, B. M. & Koehler, M. C. Isotopic evidence for biological
nitrogen fixation by molybdenum-nitrogenase from 3.2 Gyr. *Nature* **520**, 666-669,
doi:10.1038/nature14180 (2015).
18    Koehler, M. C., Buick, R. & Barley, M. E. Nitrogen isotope evidence for anoxic deep
marine environments from the Mesoarchean Mosquito Creek Formation, Australia.
*Precambrian Research* **320**, 281-290, doi:10.1016/j.precamres.2018.11.008 (2019).
19    Koehler, M. C., Buick, R., Kipp, M. A., Stueken, E. E. & Zaloumis, J. Transient surface
ocean oxygenation recorded in the similar to 2.66-Ga Jeerinah Formation, Australia.
*Proceedings of the National Academy of Sciences of the United States of America* **115**,
7711-7716, doi:10.1073/pnas.1720820115 (2018).
20    Homann, M. *et al.* Microbial life and biogeochemical cycling on land 3,220 million years
ago. *Nature Geoscience* **11**, 665-+, doi:10.1038/s41561-018-0190-9 (2018).

Godfrey, L. V. & Falkowski, P. G. The cycling and redox state of nitrogen in the Archaean ocean. *Nature Geoscience* **2**, 725-729, doi:10.1038/Ngeo633 (2009).

Garvin, J., Buick, R., Anbar, A. D., Arnold, G. L. & Kaufman, A. J. Isotopic Evidence for an Aerobic Nitrogen Cycle in the Latest Archean. *Science* **323**, 1045-1048, doi:10.1126/science.1165675 (2009).

Zerkle, A. L. *et al.* Onset of the aerobic nitrogen cycle during the Great Oxidation Event. *Nature* **542**, 465-+, doi:10.1038/nature20826 (2017).

Luo, G. M. *et al.* Nitrogen fixation sustained productivity in the wake of the Palaeoproterozoic Great Oxygenation Event. *Nature Communications* **9**, doi:10.1038/s41467-018-03361-2 (2018).

Kipp, M. A., Stueken, E. E., Yun, M., Bekker, A. & Buick, R. Pervasive aerobic nitrogen cycling in the surface ocean across the Paleoproterozoic Era. *Earth and Planetary Science Letters* **500**, 117-126, doi:10.1016/j.epsl.2018.08.007 (2018).

Lyons, T. W., Reinhard, C. T. & Planavsky, N. J. The rise of oxygen in Earth's early ocean and atmosphere. *Nature* **506**, 307-315, doi:10.1038/nature13068 (2014).

Navarro-González, R., Molina, M. J. & Molina, L. T. Nitrogen fixation by volcanic lightning in the early Earth. *Geophysical Research Letters* **25**, 3123-3126, doi:https://doi.org/10.1029/98GL02423 (1998).

Kasting, J. F. & Walker, J. C. G. Limits on oxygen concentration in the prebiological atmosphere and the rate of abiotic fixation of nitrogen. *Journal of Geophysical Research: Oceans* **86**, 1147-1158, doi:https://doi.org/10.1029/JC086iC02p01147 (1981).

Hallmann, C., Kelly, A. E., Gupta, S. N. & Summons, R. E. Reconstructing Deep-Time Biology with Molecular Fossils. *Quantifying the Evolution of Early Life : Numerical Approaches to the Evaluation of Fossils and Ancient Ecosystems* **36**, 355-401, doi:10.1007/978-94-007-0680-4_14 (2011).

Moss, F. R. *et al.* Ladderane phospholipids form a densely packed membrane with normal hydrazine and anomalously low proton/hydroxide permeability. *Proceedings of the National Academy of Sciences of the United States of America* **115**, 9098-9103, doi:10.1073/pnas.1810706115 (2018).

Mettam, C. *et al.* Anaerobic nitrogen cycling on a Neoarchaean ocean margin. *Earth and Planetary Science Letters* **527**, doi:10.1016/j.epsl.2019.115800 (2019).

Ader, M. *et al.* Interpretation of the nitrogen isotopic composition of Precambrian sedimentary rocks: Assumptions and perspectives. *Chemical Geology* **429**, 93-110, doi:10.1016/j.chemgeo.2016.02.010 (2016).

Fuchsman, C. A. & Stueken, E. E. Using modern low-oxygen marine ecosystems to understand the nitrogen cycle of the Paleo- and Mesoproterozoic oceans. *Environ Microbiol* **23**, 2801-2822, doi:10.1111/1462-2920.15220 (2021).

34    Yang, Z. & Rannala, B. Bayesian estimation of species divergence times under a
molecular clock using multiple fossil calibrations with soft bounds. *Mol Biol Evol* **23**,
212-226, doi:10.1093/molbev/msj024 (2006).
35    Ducluzeau, A. L. *et al.* Was nitric oxide the first deep electron sink? *Trends in*
*Biochemical Sciences* **34**, 9-15, doi:10.1016/j.tibs.2008.10.005 (2009).
36    Harhangi, H. R. *et al.* Hydrazine synthase, a unique phylomarker with which to study the
presence and biodiversity of anammox bacteria. *Appl Environ Microbiol* **78**, 752-758,
doi:10.1128/AEM.07113-11 (2012).
37    Hamasaki, K. *et al.* Distribution and phylogeny of anaerobic ammonium-oxidizing
(anammox) bacteria in the water column of the central Pacific Ocean. *Deep-Sea Research*
*Part Ii-Topical Studies in Oceanography* **156**, 60-67, doi:10.1016/j.dsr2.2017.11.013
(2018).
38    Terada, A., Zhou, S. & Hosomi, M. Presence and detection of anaerobic
ammonium-oxidizing (anammox) bacteria and appraisal of anammox process for
high-strength nitrogenous wastewater treatment: a review. *Clean Technologies and*
*Environmental Policy* **13**, 759-781, doi:10.1007/s10098-011-0355-3 (2011).
39    Schmid, M. *et al.* Candidatus "Scalindua brodae", sp nov., Candidatus "Scalindua
wagneri", sp nov., two new species of anaerobic ammonium oxidizing bacteria.
*Systematic and Applied Microbiology* **26**, 529-538, doi:Doi
10.1078/072320203770865837 (2003).
40    Betts, H. C. *et al.* Integrated genomic and fossil evidence illuminates life's early evolution
and eukaryote origin. *Nat Ecol Evol* **2**, 1556-1562, doi:10.1038/s41559-018-0644-x
(2018).
41    Roger, A. J. & Susko, E. Molecular clocks provide little information to date
methanogenic Archaea. *Nat Ecol Evol* **2**, 1676-1677, doi:10.1038/s41559-018-0687-z
(2018).
42    Paradis, E. & Schliep, K. ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics* **35**, 526-528,
doi:10.1093/bioinformatics/bty633 (2019).
43    Wong, M. L., Charnay, B. D., Gao, P., Yung, Y. L. & Russell, M. J. Nitrogen Oxides in
Early Earth's Atmosphere as Electron Acceptors for Life's Emergence. *Astrobiology* **17**,
975-983, doi:10.1089/ast.2016.1473 (2017).
44    Kasting, J. F. & Walker, J. C. G. Limits on Oxygen Concentration in the Prebiological
Atmosphere and the Rate of Abiotic Fixation of Nitrogen. *Journal of Geophysical*
*Research-Oceans* **86**, 1147-1158, doi:DOI 10.1029/JC086iC02p01147 (1981).

45    Charnay, B., Le Hir, G., Fluteau, F., Forget, F. & Catling, D. C. A warm or a cold early
Earth? New insights from a 3-D climate-carbon model. *Earth and Planetary Science*
*Letters* **474**, 97-109 (2017).
46    Brandes, J. A. *et al.* Abiotic nitrogen reduction on the early Earth. *Nature* **395**, 365-367,
doi:Doi 10.1038/26450 (1998).
47    Summers, D. P. & Chang, S. Prebiotic Ammonia from Reduction of Nitrite by Iron(Ii) on
the Early Earth. *Nature* **365**, 630-632, doi:DOI 10.1038/365630a0 (1993).
48    Lalonde, S. V. & Konhauser, K. O. Benthic perspective on Earth's oldest evidence for
oxygenic photosynthesis. *Proceedings of the National Academy of Sciences of the United*
*States of America* **112**, 995-1000, doi:10.1073/pnas.1415718112 (2015).
49    Anbar, A. D. *et al.* A whiff of oxygen before the Great Oxidation Event? *Science* **317**,
1903-1906, doi:10.1126/science.1140325 (2007).
50    Ranjan, S., Todd, Z. R., Rimmer, P. B., Sasselov, D. D. & Babbin, A. R. Nitrogen Oxide
Concentrations in Natural Waters on Early Earth. *Geochemistry Geophysics Geosystems*
**20**, 2021-2039, doi:10.1029/2018gc008082 (2019).
51    Cheng, C. *et al.* Nitrogen isotope evidence for stepwise oxygenation of the ocean during
the Great Oxidation Event. *Geochimica Et Cosmochimica Acta* **261**, 224-247,
doi:10.1016/j.gca.2019.07.011 (2019).
52    Lehtovirta-Morley, L. E. Ammonia oxidation: Ecology, physiology, biochemistry and
why they must all come together. *FEMS Microbiol Lett* **365**, doi:10.1093/femsle/fny058
(2018).
53    Kraft, B. *et al.* Oxygen and nitrogen production by an ammonia-oxidizing archaeon.
*Science* **375**, 97-100, doi:doi:10.1126/science.abe6733 (2022).
54    Stanton, C. L. *et al.* Nitrous oxide from chemodenitrification: A possible missing link in
the Proterozoic greenhouse and the evolution of aerobic respiration. *Geobiology* **16**,
597-609, doi:10.1111/gbi.12311 (2018).
55    Li, W. *et al.* Metagenomics reveals microbial community differences lead to differential
nitrate production in anammox reactors with differing nitrogen loading rates. *Water Res*
**169**, 115279, doi:10.1016/j.watres.2019.115279 (2020).
56    Zhang, L. *et al.* Microbial competition among anammox bacteria in nitrite-limited
bioreactors. *Water Res* **125**, 249-258, doi:10.1016/j.watres.2017.08.052 (2017).
57    Bristow, L. A. *et al.* N-2 production rates limited by nitrite availability in the Bay of
Bengal oxygen minimum zone. *Nature Geoscience* **10**, 24-29, doi:10.1038/Ngeo2847
(2017).
58    Dalsgaard, T., Thamdrup, B., Farias, L. & Revsbech, N. P. Anammox and denitrification
in the oxygen minimum zone of the eastern South Pacific. *Limnology and Oceanography*
**57**, 1331-1346, doi:10.4319/lo.2012.57.5.1331 (2012).

Canfield, D. E., Glazer, A. N. & Falkowski, P. G. The evolution and future of Earth's nitrogen cycle. *Science* **330**, 192-196, doi:10.1126/science.1186120 (2010).

Zerkle, A. L. *et al.* Onset of the aerobic nitrogen cycle during the Great Oxidation Event. *Nature* **542**, 465-467, doi:10.1038/nature20826 (2017).

Ren, M. *et al.* Phylogenomics suggests oxygen availability as a driving force in Thaumarchaeota evolution. *ISME J* **13**, 2150-2161, doi:10.1038/s41396-019-0418-8 (2019).

Rattray, J. E. *et al.* A comparative genomics study of genetic products potentially encoding ladderane lipid biosynthesis. *Biol Direct* **4**, 8, doi:10.1186/1745-6150-4-8 (2009).

Van Vliet, A. H., Ketley, J. M., Park, S. F. & Penn, C. W. The role of iron in Campylobacter gene regulation, metabolism and oxidative stress defense. *FEMS Microbiol Rev* **26**, 173-186, doi:10.1016/s0168-6445(02)00095-5 (2002).

Johnson, L. A. & Hug, L. A. Distribution of reactive oxygen species defense mechanisms across domain bacteria. *Free Radical Biology and Medicine* **140**, 93-102, doi:10.1016/j.freeradbiomed.2019.03.032 (2019).

Crowe, S. A. *et al.* Sulfate was a trace constituent of Archean seawater. *Science* **346**, 735-739, doi:10.1126/science.1258966 (2014).

Jamieson, J. W., Wing, B. A., Farquhar, J. & Hannington, M. D. Neoarchaean seawater sulphate concentrations from sulphur isotopes in massive sulphide ore. *Nature Geoscience* **6**, 61-64, doi:10.1038/Ngeo1647 (2013).

Blattler, C. L. *et al.* Two-billion-year-old evaporites capture Earth's great oxidation. *Science* **360**, 320-+, doi:10.1126/science.aar2687 (2018).

Fakhraee, M., Hancisse, O., Canfield, D. E., Crowe, S. A. & Katsev, S. Proterozoic seawater sulfate scarcity and the evolution of ocean-atmosphere chemistry. *Nature Geoscience* **12**, 375-380, doi:10.1038/s41561-019-0351-5 (2019).

Kartal, B., Keltjens, J. T. & Jetten, M. S. M. Metabolism and Genomics of Anammox Bacteria. *Nitrification*, 181-200 (2011).

Akram, M. *et al.* A 192-heme electron transfer network in the hydrazine dehydrogenase complex. *Sci Adv* **5**, eaav4310, doi:10.1126/sciadv.aav4310 (2019).

Bi, Z., Qiao, S., Zhou, J., Tang, X. & Zhang, J. Fast start-up of Anammox process with appropriate ferrous iron concentration. *Bioresour Technol* **170**, 506-512, doi:10.1016/j.biortech.2014.07.106 (2014).

Poulton, S. W. & Canfield, D. E. Ferruginous Conditions: A Dominant Feature of the Ocean through Earth's History. *Elements* **7**, 107-112, doi:10.2113/gselements.7.2.107 (2011).

73 Ferousi, C. *et al.* Iron assimilation and utilization in anaerobic ammonium oxidizing
bacteria. *Curr Opin Chem Biol* **37**, 129-136, doi:10.1016/j.cbpa.2017.03.009 (2017).
74 Keren, N., Aurora, R. & Pakrasi, H. B. Critical roles of bacterioferritins in iron storage
and proliferation of cyanobacteria. *Plant Physiol* **135**, 1666-1673,
doi:10.1104/pp.104.042770 (2004).
75 Poulton, S. W., Fralick, P. W. & Canfield, D. E. Spatial variability in oceanic redox
structure 1.8 billion years ago. *Nature Geoscience* **3**, 486-490, doi:10.1038/Ngeo889
(2010).
76 Peng, M.-W. *et al.* Insight into the structure and metabolic function of iron-rich
nanoparticles in anammox bacteria. *Science of The Total Environment* **806**, 150879,
doi:https://doi.org/10.1016/j.scitotenv.2021.150879 (2022).
77 Konhauser, K. O. *et al.* Iron formations: A global record of Neoarchaean to
Palaeoproterozoic environmental history. *Earth-Science Reviews* **172**, 140-177,
doi:10.1016/j.earscirev.2017.06.012 (2017).
78 Wu, Z. *et al.* Salinity-driven heterogeneity toward anammox distribution and growth
kinetics. *Appl Microbiol Biotechnol* **103**, 1953-1960, doi:10.1007/s00253-018-9521-4
(2019).
79 Parsons, C., Stueken, E. E., Rosen, C. J., Mateos, K. & Anderson, R. E. Radiation of
nitrogen-metabolizing enzymes across the tree of life tracks environmental transitions in
Earth history. *Geobiology* **19**, 18-34, doi:10.1111/gbi.12419 (2021).
80 Ader, M. *et al.* Ocean redox structure across the Late Neoproterozoic Oxygenation Event:
A nitrogen isotope perspective. *Earth and Planetary Science Letters* **396**, 1-13,
doi:10.1016/j.epsl.2014.03.042 (2014).
81 Wang, S. & Luo, H. Dating Alphaproteobacteria evolution with eukaryotic fossils. *Nature*
*Communications* **12**, 3324, doi:10.1038/s41467-021-23645-4 (2021).
82 Li, X. *et al.* Simultaneous Fe(III) reduction and ammonia oxidation process in Anammox
sludge. *Journal of Environmental Sciences* **64**, 42-50, doi:10.1016/j.jes.2017.01.002
(2018).
83 Zhu, J., Li, T., Liao, C., Li, N. & Wang, X. A promising destiny for Feammox: From
biogeochemical ammonium oxidation to wastewater treatment. *Science of The Total*
*Environment* **790**, 148038, doi:https://doi.org/10.1016/j.scitotenv.2021.148038 (2021).
84 Wan, L., Liu, H. & Wang, X. Anaerobic ammonium oxidation coupled to Fe(III)
reduction: Discovery, mechanism and application prospects in wastewater treatment.
*Science of The Total Environment*, 151687,
doi:https://doi.org/10.1016/j.scitotenv.2021.151687 (2021).

85    Yang, Y., Xiao, C., Yu, Q., Zhao, Z. & Zhang, Y. Using Fe(II)/Fe(III) as catalyst to drive a
novel anammox process with no need of anammox bacteria. *Water Research* **189**, 116626,
doi:https://doi.org/10.1016/j.watres.2020.116626 (2021).
86    Shaw, D. R. *et al.* Extracellular electron transfer-dependent anaerobic oxidation of
ammonium by anammox bacteria. *Nature Communications* **11**,
doi:10.1038/s41467-020-16016-y (2020).
87    Johnson, J. E., Webb, S. M., Ma, C. & Fischer, W. W. Manganese mineralogy and
diagenesis in the sedimentary rock record. *Geochimica et Cosmochimica Acta* **173**,
210-231, doi:https://doi.org/10.1016/j.gca.2015.10.027 (2016).
88    Buick, R. Did the Proterozoic 'Canfield Ocean' cause a laughing gas greenhouse?
*Geobiology* **5**, 97-100, doi:10.1111/j.1472-4669.2007.00110.x (2007).
89    Parks, D. H. *et al.* Recovery of nearly 8,000 metagenome-assembled genomes
substantially expands the tree of life. *Nat Microbiol* **2**, 1533-1542,
doi:10.1038/s41564-017-0012-7 (2017).
90    Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol Biol Evol* **30**, 772-780,
doi:10.1093/molbev/mst010 (2013).
91    Capella-Gutierrez, S., Silla-Martinez, J. M. & Gabaldon, T. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**, 1972-1973,
doi:10.1093/bioinformatics/btp348 (2009).
92    Nguyen, L. T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: a fast and
effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol*
*Evol* **32**, 268-274, doi:10.1093/molbev/msu300 (2015).
93    Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S.
ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat Methods* **14**,
587-589, doi:10.1038/nmeth.4285 (2017).
94    Letunic, I. & Bork, P. Interactive Tree Of Life (iTOL) v4: recent updates and new
developments. *Nucleic Acids Research* **47**, W256-W259 (2019).
95    Yang, Z. H. PAML 4: Phylogenetic analysis by maximum likelihood. *Molecular Biology*
*and Evolution* **24**, 1586-1591 (2007).
96    Wang, H.-C., Minh, B. Q., Susko, E. & Roger, A. J. Modeling site heterogeneity with
posterior mean site frequency profiles accelerates accurate phylogenomic estimation.
*Systematic biology* **67**, 216-235 (2018).
97    Reis, M. D. *et al.* Using Phylogenomic Data to Explore the Effects of Relaxed Clocks
and Calibration Strategies on Divergence Time Estimation: Primates as a Test Case. *Syst*
*Biol* **67**, 594-615, doi:10.1093/sysbio/syy001 (2018).

98   Dos Reis, M., Donoghue, P. C. & Yang, Z. Bayesian molecular clock dating of species
divergences in the genomics era. *Nat Rev Genet* **17**, 71-80, doi:10.1038/nrg.2015.8
(2016).
99   Dos Reis, M. & Yang, Z. Approximate likelihood calculation on a phylogeny for
Bayesian estimation of divergence times. *Mol Biol Evol* **28**, 2161-2172,
doi:10.1093/molbev/msr045 (2011).
100  Quang, L. S., Gascuel, O. & Lartillot, N. Empirical profile mixture models for
phylogenetic reconstruction. *Bioinformatics* **24**, 2317-2323,
doi:10.1093/bioinformatics/btn445 (2008).
101  Sanderson, M. J. Estimating absolute rates of molecular evolution and divergence times:
A penalized likelihood approach. *Molecular Biology and Evolution* **19**, 101-109, doi:DOI
10.1093/oxfordjournals.molbev.a003974 (2002).
102  Morel, B., Kozlov, A. M., Stamatakis, A. & Szollosi, G. J. GeneRax: A tool for
Species-Tree-Aware Maximum Likelihood-based gene family tree inference under gene
duplication, transfer, and loss. *Mol Biol Evol* **37**, 2763-2774,
doi:10.1093/molbev/msaa141 (2020).
103  Kanehisa, M. & Goto, S. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic
Acids Research* **28**, 27-30 (2000).
104  Lu, S. *et al.* CDD/SPARCLE: the conserved domain database in 2020. *Nucleic Acids Res*
**48**, D265-D268, doi:10.1093/nar/gkz991 (2020).
105  Mitchell, A. L. *et al.* InterPro in 2019: improving coverage, classification and access to
protein sequence annotations. *Nucleic Acids Res* **47**, D351-D360,
doi:10.1093/nar/gky1100 (2019).
106  El-Gebali, S. *et al.* The Pfam protein families database in 2019. *Nucleic Acids Res* **47**,
D427-D432, doi:10.1093/nar/gky995 (2019).
107  Haft, D. H. *et al.* TIGRFAMs: a protein family resource for the functional identification
of proteins. *Nucleic Acids Res* **29**, 41-43, doi:10.1093/nar/29.1.41 (2001).
108  Saier, M. H., Jr. *et al.* The Transporter Classification Database (TCDB): recent advances.
*Nucleic Acids Res* **44**, D372-D379, doi:10.1093/nar/gkv1103 (2016).

**Figure legends**

**Figure 1.** The evolutionary timeline of anammox bacteria using MCMCTree. The chronogram

was estimated based on the calibration set C1 (see Supplementary Text section 3) and sequence

alignments of the 25 orthologs conserved in bacteria (20040 sites). The blue bars on the four

calibration nodes and the last common ancestor (LCA) of anammox bacteria represent the

posterior 95% highest probability density (HPD) interval of the posterior time estimates. These

alternative calibration sets were selected by choosing the three that accommodate different

calibration constraints (see Supplementary Texts section 3.3). More alternative time estimates are

provided in Figure S4. The vertical grey bar represents the period of the GOE from 2,500 to

2,320 Ma. The calibration constraints used within the phylum Cyanobacteria are marked with

orange texts: the LCA of Planctomycetes and Cyanobacteria (Root), the total group of oxygenic

Cyanobacteria (Node 2), the total group of *Nostocales* (Node 3), and the total group of

*Pleurocapsales* (Node 4). The genome completeness estimated by checkM is visualized with a

gradient color strip. The right next color strip indicates the genome type of genomic sequences

used in our study including metagenome-assembled genomes (MAGs) and whole-genome

sequencing (WGS) from either enriched culture sample or isolate. The diagram below the dated

tree illustrates the change of atmospheric partial pressure of $O_2$ ($P_{O_2}$) and nitrogen isotope

fractionations. The blue line shows the proposed model according to Lyons, et al. [26]. The green

arrow suggests the earliest evidence for aerobic nitrogen cycling at around 2.7 Ga. PAL on right

axis means $P_{O_2}$ relative to the present atmospheric level. The black dots denote the change of

nitrogen $\delta^{15}N$ isotope values according to Kipp, et al. [25].

**Figure 2**. The phyletic pattern of ecologically relevant genes in the comparison between

anammox bacteria and non-anammox bacteria. Solid circles at the nodes indicate the ultrafast

bootstrap values in 1,000 bootstrapped replicates. Note that copy number difference is not

indicated since most genes displayed here are present as a single copy in genome except a few

exceptions like *hao* (see Dataset S1.5 for the table summarizing the copy number of each gene).

The phylogenomic tree on the left was constructed with 887 genomic sequences described in

Supplementary Texts section 2.1. Those Planctomycetes genomes not used for comparative

genomics analyses are collapsed into grey triangles, and the numbers of collapsed genomes are

labelled next to the triangles. The target group and reference group for comparative genomic

analysis are within an orange or blue box, separately. For each genome, the genome

completeness estimated by CheckM is visualized with a color strip and labelled besides leaf

nodes. The right next color strip represents the type of genomic sequences used in our study

including metagenome-assembled genomes (MAGs) and whole-genome sequencing (WGS)

from either enriched culture sample or isolate. The filled and empty circles, respectively,

represent the presence and absence of particular genes in corresponding genomes. For gene

clusters, only genomes with at least half of the members of the gene cluster are indicated by a

filled circle. The classifications of annotated genes are labelled above the gene names. *hzsCBA*,

hydrazine synthase subunit C, B and A; *hdh*, hydrazine dehydrogenase; *hao*, hydroxylamine

dehydrogenase; *nxrAB*, nitrite oxidoreductase subunit A and B; *nirK*, copper-containing and

NO-forming nitrite reductase; *nirS*, cytochrome NO-forming nitrite reductase; *nrfAH*,

ammonia-forming nitrite reductase subunit A and H; *glnB*, nitrogen regulatory protein P-II; *nirC*,

nitrite transporter; NRT, nitrate/nitrite transporter; *amt*, ammonium transporter; kuste2805, 3603,

3605-3606, proposed genes relative to the synthetic pathways for ladderane at Rattray, et al. [62];

*cbiG*, cobalt-precorrin 5A hydrolase; *cbiD*, cobalt-precorrin-5B(C1)-methyltransferase; *cbiOMQ*,

cobalt/nickel transport system; *AhpC*, peroxiredoxin; *Ccp*, cytochrome *c* peroxidase; *dfx*,

superoxide reductase; *fprA*, $H_2O$-forming enzyme flavoprotein; *CcsAB*, cytochrome c maturation

systems; *petB*, Cytochrome b subunit of the bc complex; *petC*, Rieske Fe-S protein; *ahbABCD*,

heme biosynthesis; *sat*, sulfate adenylyltransferase; *aprAB*, adenylylsulfate reductase, subunit A

and B; *fsr*, sulfite reductase (coenzyme F420); *higB-1*, toxin; *higA-1*, antitoxin; *mnhABCDEG*,

multicomponent $Na^+/H^+$ antiporter; *fdhAB*, formate dehydrogenase subunit A and B; *nuo*(A-N),

NADH-quinone oxidoreductase; *ndh*(A-N), NAD(P)H-quinone oxidoreductase; *nqrABCDEF*,

$Na^+$-transporting NADH:ubiquinone oxidoreductase; *rnfABCDEG*, $Na^+$-translocating

ferredoxin:$NAD^+$ oxidoreductase; *atp*(A-H), F-type $H^+$-transporting ATPase; *lacAZ*,

beta-galactosidase; *melA*, galA, alpha-galactosidase; *ebgA*, beta-galactosidase; *galK*,

galactokinase; , fructokinase; *fruK*, 1-phosphofructokinase; *rhaB*, rhamnulokinase; rbsK,

ribokinase; *araB*, L-ribulokinase; *xylB*, xylulokinase; *kdgK*, 2-dehydro-3-deoxygluconokinase;

GALK2, N-acetylgalactosamine kinase; *cah*, cephalosporin-C deacetylase; *argE*, acetylornithine

deacetylase; *nagA*; N-acetylglucosamine-6-phosphate deacetylase; *HDAC11*, histone deacetylase

11.