Reconstruction of nitrogenase predecessors suggests origin from maturase-like proteins

Amanda K. Garcia[1], Bryan Kolaczkowski[2], Betül Kaçar[1]

[1]Department of Bacteriology, University of Wisconsin – Madison, USA;

[2]Department of Microbiology and Cell Science, University of Florida, Gainesville, Florida, USA;

Corresponding author: Betül Kaçar

Email: betul@wisc.edu

ABSTRACT

The evolution of biological nitrogen fixation, uniquely catalyzed by nitrogenase enzymes, has been one of the most consequential biogeochemical innovations over life's history. Though understanding the early evolution of nitrogen fixation has been a longstanding goal from molecular, biogeochemical, and planetary perspectives, its origins remain enigmatic. In this study, we reconstructed the evolutionary histories of nitrogenases, as well as homologous maturase proteins that participate in the assembly of the nitrogenase active-site cofactor but are not able to fix nitrogen. We combined phylogenetic and ancestral sequence inference with an analysis of predicted functionally divergent sites between nitrogenases and maturases to infer the nitrogen-fixing capabilities of their shared ancestors. Our results provide phylogenetic constraints to the emergence of nitrogen fixation and are consistent with a model wherein nitrogenases emerged from maturase-like predecessors. Though the precise functional role of such a predecessor protein remains speculative, our results highlight evolutionary contingency as a significant factor shaping the evolution of a biogeochemically essential enzyme.

SIGNIFICANCE STATEMENT

The origin of nitrogenase-catalyzed nitrogen fixation was a transformative event in life's history, garnering long-term study from molecular, biogeochemical, and planetary perspectives. Reconstruction of ancestral nitrogenases suggests that the protein sequence space capable of yielding a nitrogen-fixing enzyme in the past was likely more constrained than previously thought. Specifically, here we show that nitrogenases likely evolved from ancestors that resemble maturases, homologs that today participate in nitrogenase cofactor assembly, contrary to the commonly accepted view that maturases evolved from a nitrogenase ancestor. We further submit that the molecular architecture that may have been required for nitrogenase origins was unlikely to have been shaped by the same environmental drivers often implicated in the evolution of nitrogen fixation. If this decoupling is found to be a recurring pattern in metabolic origins, then the presented results would undercut the common, systems-focused rationale of using ancient environmental conditions to explain the timing of critical and singular biogeochemical innovations in life's past.

KEYWORDS

nitrogenase, nitrogen fixation, maturase, ancestral sequence reconstruction, early life, historical contingency

INTRODUCTION

The modern biosphere is shaped by a variety of essential and ancient enzymes that have co-evolved with the Earth environment for billions of years. Though general mechanisms for the gain of novel enzymatic functions have been explored (Ohno 1970; Gerlt and Babbitt 2001; Copley 2015; Noda-Garcia et al. 2018; Copley 2021), the co-evolutionary steps toward the origins of many specific, key enzymes during Earth's early history remain unresolved. An unavoidable task in addressing this challenge is constraining the ancestral functions of early-evolved enzyme families and their precursors (Benner et al. 2007, Kacar and Garcia 2019).

Biological nitrogen fixation is a notable example of a critical metabolic process with ancient and enigmatic origins. All life requires fixed, or bioavailable, nitrogen. For much of Earth history, this biologically vital element has primarily been acquired by organisms via the activities of nitrogenase metalloenzymes, early evolved and conserved catalysts that uniquely reduce dinitrogen (N_2) to ammonia (NH_3) (Hoffman et al. 2014; Einsle and Rees 2020). The evolution of nitrogenases has constrained the long-term productivity of the biosphere and has itself been shaped by the co-evolving biogeochemistry of Earth's environment (Falkowski 1997; Glass et al. 2009; Stüeken et al. 2016; Luo et al. 2018; Allen et al. 2019; Mus et al. 2019; Garcia et al. 2020). Nitrogenases represent the only known biomolecular solution for the reduction of N_2, a remarkable innovation given that the N≡N bond is one of the most inert in nature. The answer to how biology converged on this solution billions of years ago remains elusive (Boyd and Peters 2013; Mus et al. 2019; Garcia et al. 2020).

Insights into the origins of biological nitrogen fixation can be gained by reconstructing the protein ancestors of nitrogenases and their close homologs. Nitrogenases do not operate alone, but instead exist within a larger protein network required for their assembly and function. The closest homologs to nitrogenases are themselves a key player in this larger assembly network, serving as maturases (also referred to as assembly scaffold proteins) for the final steps in nitrogenase cofactor biosynthesis (Fay et al. 2016; Buren et al. 2020) (fig. 1a). These maturases are

considered necessary in most nitrogenase assembly pathways to modify a complex metal cluster

precursor that, when matured, serves as the nitrogenase active site for N_2 reduction. (The only

forms of nitrogenases that are confirmed to assemble without maturases are those that only

incorporate iron into their active-site cofactors (Perez-Gonzalez et al. 2021)). Though not

themselves known to reduce N_2, these maturases reduce a variety of other substrates *in vitro*

under highly reducing conditions, including C_2H_2, CO, and CN^-, that also serve as alternative, non-

physiological substrates of nitrogenases (Hu et al. 2010; Fay et al. 2016; Seefeldt et al. 2020).

These findings establish maturases as catalytically similar to nitrogenases with the exception of

their inability to reduce N_2. The divergent protein features between nitrogenases and maturases

must therefore account for this difference in N_2-reduction capability.

To investigate the origins of biological nitrogen fixation, we reconstructed the evolutionary history

of nitrogenases and maturases. By comparing patterns of sequence conservation between

nitrogenases and maturases, we identified divergent residues that might account for their

functional differentiation—namely, their ability or inability to reduce N_2, respectively. These

sequence features were then leveraged to infer the N_2-reduction capability of reconstructed

ancestral proteins and phylogenetically map the origins of biological nitrogen fixation within the

nitrogenase family evolutionary history. The relative timing of the evolutionary relationship

between nitrogenases and maturases is debated, with some studies suggesting that maturases

are evolutionarily derived from nitrogenases (Boyd et al. 2011a). Our findings support an origin of

the canonical nitrogenase clade from predecessor proteins that were unlikely to have been

capable of N_2 reduction and more closely resemble extant maturases—proteins that are today

only ancillary in biological nitrogen fixation. Nitrogenases may therefore represent a case in

molecular evolution where a pre-existing but already complex protein architecture, adapted to an

alternative role, shaped the origins of one of the most consequential biomolecular innovations in

Earth history.

RESULTS AND DISCUSSION

Uncharacterized homologs root canonical nitrogenases within maturase protein clades

We reconstructed the phylogenetic history of nitrogenase and maturase homologs to explore

ancestral states for this protein family. There are three forms of nitrogenase—Nif, Vnf, and Anf—

that each differ in the composition of their active-site iron-sulfur cluster ("M-cluster"; incorporating

molybdenum, vanadium, or additional iron, respectively). The heterotetrameric ($\alpha_2\beta_2$) catalytic

protein component of nitrogenase (Nif/Vnf/AnfDK) has its counterpart in a homologous maturase

protein complex (Nif/VnfEN; the Anf nitrogenase system does not have dedicated maturase

proteins (Perez-Gonzalez et al. 2021)) (fig. 1a). We compiled a comprehensive dataset including

nitrogenase Nif/Vnf/AnfDK and maturase NifEN protein sequences (fig. 1b), as well as outgroup

dark-operative protochlorophyllide oxidoreductase homologs that share the $\alpha_2\beta_2$ subunit

arrangement (BchNB) (Fujita and Bauer 2000; Moser and Brocker 2011). Four maximum

likelihood phylogenies were built to test the robustness of tree topology and downstream ancestral

sequence inference to sequence sampling and alternate alignment methods (table 1): 1) 2,425

nitrogenase, maturase, and outgroup homologs aligned by MAFFT (Katoh and Standley 2013),

Tree-1; 2) removal of "uncharacterized" nitrogenase and maturase homologs, Tree-2 (see

definition and discussion of "uncharacterized" homologs below); 3) removal of β-subunit

nitrogenase, maturase, and outgroup homologs, Tree-3; and 4) alignment with MUSCLE (Edgar

2004) instead of MAFFT, Tree-4.

General shared features across the reconstructed phylogenies include clustering of α-subunit

sequences sister to β-subunit sequences (except in Tree-3, which lacks β-subunit sequences),

reproducing the $\alpha_2\beta_2$ structural distinction for nitrogenases and maturases (fig. 1c, supplementary

fig. S1). These α- and β-subunit clades themselves each segregate into nitrogenase and

maturase protein sequences. This topology is consistent with an initial gene duplication event that

resulted in separate α- and β-subunits, followed by a secondary duplication event that resulted in

functionally distinct nitrogenase and maturase proteins (Fani et al. 2000; Raymond et al. 2004;

Boyd et al. 2011a; Boyd and Peters 2013). Within the nitrogenase clade, vanadium- and iron-

nitrogenase sequences nest within molybdenum-nitrogenase clades, as has been previously

observed (Raymond et al. 2004; Boyd et al. 2011a; Garcia et al. 2020). By contrast, the

phylogenetic clustering of maturase sequences associated with different metal-dependent forms

of nitrogenases do not reproduce this nesting pattern. Vanadium-maturase sequences are each

split into two groups: one forms a small clade with a relatively long branch that diverges prior to

all other nitrogenase and maturase sequences, whereas another diverges relatively recently

within nitrogenase clades associated with aerobic or facultative bacteria. This topology suggests

that maturases for the vanadium nitrogenase system originated independently at least twice, with

one origin associated with an early divergence from ancestors of unknown function and another

from a recent nitrogenase ancestor.

For the phylogenies that include them (i.e., Tree-1, Tree-3, Tree-4), several maturase-like homologs diverge prior to nitrogenases and root the latter within the maturase clade (fig. 1c, supplementary fig. S1). These homologs belong to "uncharacterized" bacterial and archaeal taxa that lack extensive experimental characterization regarding the metal dependence and N_2-reducing capability of their nitrogenase-like proteins (McGlynn et al. 2012; Garcia et al. 2020). We obtained preliminary functional predictions for these uncharacterized, maturase-like sequences by KEGG's BlastKOALA (Kanehisa et al. 2016), including a control subset of sequences as well that branch within canonical maturase clades. BlastKOALA returned a mix of maturase and nitrogenase annotations, even for certain control sequences (supplementary table S1). Given this discrepancy and the absence of experimental data, we assign these uncharacterized maturase-like sequences as putative maturases (rather than nitrogenases) based on three lines of evidence. First, the genes that encode these homologs in uncharacterized taxa are located closely downstream of nitrogenase-like genes, as is frequently the case with *bona fide* maturases (fig. 1b). Second, certain uncharacterized taxa (including those that are missing a NifN-like maturase subunit gene) have been shown to fix nitrogen, evidencing a functioning nitrogenase and, by extension, maturase (Mehta and Baross 2006; Dekas et al. 2009; Chen et al. 2021). Third, these homologs conserve a Cys48 residue (numbering here and hereafter from aligned *Azotobacter vinelandii* nitrogenase NifD) present in most maturase proteins (with the exception of certain VnfE homologs) and considered important for binding the cluster precursor prior to maturation (fig. 2a) (Kaiser et al. 2011). At the same time, these sequences all lack the strictly conserved nitrogenase His442 residue that ligates the active-site M-cluster and is critical for N_2 reduction (Kim and Rees 1992; Lee et al. 1998; Li 2002; Jimenez-Vicente et al. 2018). Together, these observations suggest that the uncharacterized maturase-like homologs are unlikely to be functioning as nitrogenases and are more likely operating as canonical maturases.

Our alternate tree reconstructions demonstrate that the rooting of nitrogenases within maturase clades is primarily determined by the inclusion of uncharacterized homologs. Though the phylogenetic position of certain uncharacterized clades is ambiguous (e.g., one clade diverges immediately prior to nitrogenases in Tree-1, but prior to both nitrogenases and other maturases in Tree-3 and Tree-4), we do not observe rooting of maturases within nitrogenase sequences in any of these trees (fig. 1c, supplementary fig. S1). These topological features are also unaffected by trimming the Tree-1 alignment (supplementary fig. S2). This consistency suggests that the observed rooting pattern is robust to the exclusion of the β-subunit protein subtree (Tree-3) and

variation in tested alignment methodology (Tree-4). By contrast, the exclusion of uncharacterized

homologs from the sequence dataset results in reciprocal monophyly between α-subunit

nitrogenase and maturase clades and nesting of β-subunit maturase sequences within

nitrogenase sequences (Tree-2; supplementary fig. S1). The α-subunit topology is therefore more

stable and less affected by the presence of uncharacterized homologs than the β-subunit

topology, which is consistent with the comparatively low branch support values among β-subunit

sequences in Tree-2.

The rooting of nitrogenases within the broader maturase protein clade might parsimoniously

suggest that the functionality of the common ancestor of both protein groups more likely

resembled that of extant maturases. These phylogenetic observations present a hypothesis that

can be tested by evaluating sequence features of reconstructed ancestral proteins inferred to

contribute to the functional divergence between nitrogenases and maturases.

Phylogenetically divergent protein sites between extant nitrogenases and maturases map

to functionally important structural regions

We performed a comprehensive, comparative sequence-structure analysis to identify protein sites

that likely contribute to the functional divergence (i.e., (in)ability to reduce N_2) between

nitrogenases and maturases. Our goal was to subsequently leverage this analysis for

identification of similar sites in reconstructed protein ancestors and phenotypic inference. We

limited our analysis to α-subunit nitrogenase (NifD) and maturase (NifE) sequences in part due to

the greater topological uncertainty within β-subunit subtrees across our phylogenetic

reconstructions (fig. 1c, supplementary fig. S1). In addition, α-subunit sequences host the active-

site M-cluster or cluster precursor. We therefore expected that functional differences between

nitrogenases and maturases are more likely to be modulated by sequence- and structural-level

differences between α-subunit proteins.

To predict functionally divergent protein sites, we first calculated the amino acid frequency

distributions of each alignment column for nitrogenase and maturase sequences (Materials and

Methods). The Jensen-Shannon (J-S) distance between the two protein groups for every

alignment column was calculated, where larger distances indicate greater divergence for that site

between nitrogenases and maturases. We estimated the expected distribution of site-wise J-S

distances by randomly partitioning our protein sequences into 2 groups 10,000 times and

calculating site-wise J-S distances from each random partition. The p-value for each site's J-S

distance from the nitrogenase-maturase partition was calculated from the distribution of J-S

distances for that site across random sequence partitions. We defined functionally divergent sites

as those exceeding the 75[th]-percentile distance across all alignment columns, as well as having

an FDR-corrected *p*-value <0.0001. This analysis was repeated for all four alignments used to

build phylogenetic trees (table 1), yielding 117 (116 for the Tree-4 alignment) predicted sites

(supplementary table S2).

Putative functionally divergent sites identified from the sequence alignments cluster with known

functionally important structural regions of the nitrogenase subunit (fig. 2b). These include the M-

cluster active site for N_2 reduction, the interface between nitrogenase NifD and NifK subunits, and

the M-cluster insertion funnel (Hu et al. 2008). This observed correlation suggests that the set of

putative functionally divergent sites is likely enriched for positions contributing to the functional

divergence between nitrogenases and maturases. Six sites are of the most divergent across all

sequence alignments: 69, 189, 362, 383, 440, and 444 (supplementary table S2). Two additional

sites, 442 and 445, are highly divergent in alignments for Tree-1, Tree-2, and Tree-3, but not for

Tree-4, likely because Tree-4 was constructed by a different alignment method that would have

impacted downstream distance calculations (table 1). All are in or proximal to the nitrogenase

active site or M-cluster insertion funnel. Some have specific inferred or experimentally determined

functional roles in nitrogenases, including as an M-cluster ligand (site 442) (Kim and Rees 1992),

a "lock" to hold the M-cluster within the active site (site 444) (Hu et al. 2008; Solomon et al. 2020),

and a "lid" at the cluster insertion funnel opening (site 362) (Hu et al. 2008). Finally, our set of

divergent sites includes the maturase cluster precursor ligand, site 48 (Kaiser et al. 2011). The

clustering of highly divergent sites with these protein regions highlights the differential interaction

of nitrogenases or maturases with the M-cluster. Specifically, maturase function does not require

the conservation of residues that permit the insertion and stabilization of the cluster within the

active site (Hu et al. 2008), nor the fine-tuning of residues in the active site for catalysis.

In addition to divergent sites, we identified several conserved residues among nitrogenase and

maturase sequences (fig. 2a). His83, Cys88, Cys154, Gly160, Gly185, Gly194, and Cys275

residues are conserved in both nitrogenases and maturases. These sites are likely essential to

both groups and arose prior to their evolutionary divergence. 23 sites are conserved only in

nitrogenases, compared to just three sites that are conserved only in maturases: Cys62, Gly246,

and Gly455. However, these three residues are still present in most nitrogenase sequences. The

greater number of uniquely conserved residues in nitrogenases relative to maturases may reflect

the stronger selective constraint associated with N_2-reduction functionality.

Putative functionally divergent sites of oldest ancestors resemble extant maturases more

than nitrogenases

With a set of predicted functionally divergent protein sites between extant α-subunit nitrogenase

and maturase proteins, we used a probabilistic approach to compare features of ancestral

proteins inferred from all reconstructed phylogenies (table 1, supplementary fig. S1). For divergent

sites, J-S distances were calculated between ancestral amino acid posterior probability

distributions and extant amino acid frequency distributions for either extant nitrogenase or

maturase homologs. These distance scores were then normalized to yield a value between -1

and +1, here called the "D-score" (supplementary table S3). Positive D-scores indicate greater

similarity to α-subunit nitrogenase NifD homologs and negative scores indicate greater similarity

to NifE homologs (see Materials and Methods for additional details). D-scores were averaged

across divergent sites for each ancestral node.

Ancestral protein sequences inferred from oldest phylogenetic nodes on average resemble extant

maturase homologs more than nitrogenase homologs at predicted functionally divergent sites.

Mean D-scores across divergent sites for nodes ancestral to all nitrogenase and maturase

homologs (fig. 3, "nitrogenase/maturase last common ancestor 1") range between -0.08 and 0.00.

These values are low-magnitude relative to the full range of mean D-scores across all ancestral

nodes (~0.2 to +0.2), which is expected due to the mixing of both nitrogenase- and maturase-like

sequence features at oldest nodes. Nevertheless, the primarily negative mean D-score values at

these nodes indicate greater sequence-level similarity to maturase Nif/VnfE sequences than to

nitrogenase Nif/Vnf/AnfD sequences. The ambiguity of the Tree-3 node (D-score = 0.00) may

result from the removal of β-subunit sequences for this phylogeny that would otherwise form an

outgroup to the α-subunit clade and constrain ancestral sequence composition at this node. At

more recent ancestral nodes that exclude early diverged VnfE clades (which have long, less

highly supported branches and, thus, ambiguous evolutionary context; "nitrogenase/maturase last

common ancestor 2"), mean D-scores range between -0.10 and +0.02. The only tree that yields

positive D-scores for these more recent nodes is that reconstructed from the MUSCLE alignment

(Tree-4), indicating that this analysis is more sensitive to the tested alignment method than
sequence sampling. The sensitivity of ancestral sequence inference to alignment method has
been observed previously with simulated data, also finding that MUSCLE produces less accurate
inferences than MAFFT (Vialle et al. 2018). Finally, nodes associated with the "nitrogenase last
common ancestor" yield positive D-scores for all tree, and therefore resemble nitrogenases more
than maturases at predicted functionally divergent sites.

In addition to surveying mean ancestral similarity to extant nitrogenases or maturases across all
putative functionally divergent sites, we also investigated site-wise D-score values, some of which
have been experimentally determined to contribute to either nitrogenase or maturase function.
These site-wise D-scores were assessed specifically along the phylogenetic transect between the
last common ancestor of nitrogenases and the last common ancestor of all nitrogenase and
maturases (supplementary table S3). Certain divergent sites that become "nitrogenase-like" early
(i.e., prior to the nitrogenase ancestor) include site 195 (important for N_2 substrate binding (Kim
et al. 1995)), site 444 (locks the M-cluster in the nitrogenase active site (Hu et al. 2008; Solomon
et al. 2020)), and site 359 (helps form the cluster insertion funnel (Hu et al. 2006)). By contrast,
site 48 (involved with L-cluster binding in maturases (Kaiser et al. 2011)), as well as sites 361 and
362 (involved with M-cluster insertion at the nitrogenase active site (Hu et al. 2008)), remain
primarily "maturase-like" until the nitrogenase last common ancestor.

The site 442 histidine M-cluster ligand conserved across all extant canonical nitrogenase proteins
is considered critical for nitrogenase activity (Kim and Rees 1992; Lee et al. 1998; Li 2002;
Jimenez-Vicente et al. 2018) and, notably, remains maturase-like prior to the nitrogenase last
common ancestor (mean D-score ≈ -0.40 across alternate alignments, compared to a minimum
D-score of -0.76 for all divergent sites; supplementary table S3). Site 442 is only not predicted to
be functionally divergent for the tree reconstructed by a MUSCLE alignment (Tree-4). This
difference is likely due to an inferred homology, unique to the MUSCLE alignment, between the
nitrogenase His442 residue and a frequently observed histidine residue in maturases. However,
the other alignments, which instead infer homology with a neighboring arginine residue in many
maturases (fig. 2a), is supported by studies which indicate that this arginine residue is structurally
aligned to nitrogenase His442 (Kaiser et al. 2011). Thus, it is likely that the MUSCLE alignment
is erroneous at this site, consistent with the observed reduced accuracy of MUSCLE compared
to MAFFT for ancestral sequence inference (Vialle et al. 2018). Due to its functional significance

and conservation among extant nitrogenases, the appearance of a histidine residue at site 442
may have been critical for the origins of N_2 reduction.

Nitrogenases likely originated from a non-N_2-reducing maturase-like protein with possible
biosynthetic or alternate catalytic roles

Our exploration of nitrogenase and maturase ancestry, coupled with the investigation of global
sequence and structural features that account for their functional divergence, is generally
consistent with the hypothesis that the last common ancestor of nitrogenase and maturase
proteins was unlikely to have functioned as a nitrogenase. This inference is also supported by the
absence of residue-level similarity at divergent sites that have empirically been shown to be
critical for nitrogenase function. These results are robust to phylogenetic uncertainty stemming
fromsequence sampling of early diverged uncharacterized lineages and incorporates statistical
uncertainty associated with ancestral sequence inference. Though we find that an alternative
alignment method, MUSCLE, does modulate these sequence-based functional inferences at one
ancestral node (fig. 3), there is reason to suspect reduced alignment accuracy given probable
misalignment of at least one key nitrogenase residue and decreased performance in simulated
data (Vialle et al. 2018) (see above).

Conservatively, our results suggest that phylogenetic inference of ancient nitrogen fixation can
reliably extend only to within the canonical nitrogenase clade, and that inferences of earlier
nitrogen fixation activity require further evidence. More definitive assessment of the N_2-reduction
capability of the common ancestor of nitrogenases and maturases awaits experimental
investigation, which can be directly achieved through the laboratory resurrection of ancestral
proteins inferred in this study (Thornton 2004; Benner 2017; Garcia and Kacar 2019).
Nevertheless, a model that posits a maturase-like ancestry for nitrogenases deviates from existing
hypotheses regarding their early evolution (fig. 4). Previous models are based on parsimonious
interpretations of nitrogenase and maturase phylogenetic topology that is not observed in the
trees reconstructed here. For instance, previous studies root the maturase clade within
nitrogenase sequences, suggesting that the former evolved via gene duplication of nitrogenase
ancestors (Boyd et al. 2011a; Boyd and Peters 2013). An updated phylogenetic analysis
incorporating the Boyd et al. (2011a) dataset, lacking uncharacterized sequences, produces a
topology similar to Tree-2 where nitrogenase and maturase α-subunit clades are reciprocally

monophyletic (supplementary fig. S2). However, with the inclusion of uncharacterized nitrogenase
and maturase homologs, maturases instead root nitrogenases, and support a model wherein
nitrogen fixation is instead a derived feature of a maturase-like ancestor. Our reconstruction of
ancestral states within the nitrogenase and maturase phylogeny provides additional constraints
on ancestral phenotypes within a maximum likelihood framework, extending beyond inferences
drawn from phylogenetic topology alone. These results constrain the likely origin of nitrogen
fixation to a relatively well-resolved lineage within the nitrogenase/maturase topology, rather than
to a deeper history that bridges nitrogenases with more distantly related homologs (e.g.,
coenzyme F430 biosynthesis proteins (Mus et al. 2019)).

Our results suggest that ancestral maturase-like proteins may have provided the molecular
architecture for the origin of nitrogen fixation. However, the precise functional role of such an
ancestor is not clear. Candidate phenotypic attributes, shared between extant nitrogenases and
maturases, may shed light on how biological N_2 reduction evolved. For example, both are the only
proteins known to bind the nitrogenase active-site M-cluster (Fay et al. 2016). In addition, both
extant nitrogenases and maturases can reduce several non-physiological, alternative substrates
including C_2H_2, CO, and CN^-, albeit in highly reducing experimental conditions (Fay et al. 2016;
Seefeldt et al. 2020). It has previously been argued that the ability of nitrogenases to reduce
alternative substrates may simply be the byproduct of overcoming the significant activation barrier
required for N_2 reduction (Boyd and Peters 2013). However, combined with the evidence
presented here for a maturase-like ancestry, a plausible scenario is an ancestral protein capable
of reducing the shared substrates of extant nitrogenases and maturases at an M-cluster-like
active site (perhaps in a role as detoxyases, as has previously been proposed ((Silver and
Postgate 1973); fig. 4). This scenario would provide a stepwise path for the evolution of
nitrogenases from ancestral proteins capable of catalyzing less ATP-intensive reactions (Hu et
al. 2010), and requires only residue-level tuning of an already complex peptide environment to
achieve the earliest whiffs of N_2 reduction. Our reconstruction of ancestral residues at predicted
functionally divergent sites provides possible mutational trajectories toward the evolution of
nitrogenases.

The origin of nitrogenases from proteins involved with cofactor biosynthesis such as the
maturases studied here might be expected given the prevalence of biosynthetic proteins
associated with the broader family of nitrogenase-like homologs. These include chlorophyll
biosynthesis proteins used as phylogenetic outgroups in our study (Fujita and Bauer 2000; Moser

and Brocker 2011), as well as coenzyme F430 biosynthesis proteins that are conserved among
methanogens (Staples et al. 2007; Zheng et al. 2016). In addition, more distantly related and
poorly studied nitrogenase-like homologs may have roles in assembly of hydrogenase
metalloclusters and metal transport (reviewed in Ghebreamlak and Mansoorabadi (2020)). It is
not clear whether a hypothetical, maturase-like nitrogenase ancestor may have had a central
function in cofactor biosynthesis as its descendants and several related homologs do today. Other
distantly related nitrogenase-like homologs have putative catalytic roles, including a recently
reported homolog suggested to participate in a methionine salvage pathway that forms ethylene
(North et al. 2020). It is possible that a maturase-like predecessor would have been promiscuous,
a suggested general feature of early-evolved proteins (Copley 2015; De Tarafder et al. 2021),
and capable of both providing a scaffold for cluster maturation as well as catalysis using the same
matured cluster. Gene duplication and divergence might then have subsequently specialized
maturases to only function as a scaffold and evolve residues that permit the release of the
matured cluster. In parallel, nitrogenases would have specialized to function in a solely catalytic
role.

The possible capability of an ancestral maturase-like protein to bind an M-cluster-like cofactor
would have important implications for the coevolutionary trajectory of nitrogenases and the
biogeochemical environment (Anbar and Knoll 2002; Glass et al. 2009; Boyd et al. 2011b). A
defining role of extant maturases in the molybdenum-dependent nitrogenase system is to provide
a scaffold for the incorporation of molybdenum into the active-site M-cluster. Thus, the origin of
maturase proteins has previously been suggested to coincide with the origin of molybdenum
dependence in nitrogenases (Boyd et al. 2011a). However, if maturase-like proteins predate
nitrogenases, the molybdenum-containing M-cluster itself may predate nitrogenases as well (fig.
4). Inferences for the age of nitrogenases extend to more than 3 billion years ago (Stueken et al.
2015; Parsons et al. 2020) when molybdenum in Earth's oceans was likely exceedingly scarce
(Anbar 2008). A molybdenum-incorporating maturase-like protein existing prior to 3 billion years
ago would suggest that the bulk geochemistry of the early Earth environment may not necessarily
have provided strict constraints on enzyme evolution (Garcia et al. 2020), and would be in
agreement with geochemical evidence of early molybdenum-dependent nitrogenase activity
(Stueken et al. 2015). Rather, localized environments may have provided sufficient molybdenum
for the function of nitrogenases and their predecessors (Stueken et al. 2015), or molybdenum
may have simply been selected despite its scarcity due to its invaluable chemical features.

Another possibility is that earliest maturases-like ancestors did not incorporate molybdenum, but
rather the iron-only cluster precursor that is matured to the M-cluster today (Mus et al. 2019).
409
410

411 **The role of contingency and subsumed complexity in nitrogenase evolution**
412

The alternative model of nitrogenase origins that we propose exemplifies a case for molecular
evolution in which a novel and consequential metabolic function was built off a pre-existing,
complex molecular architecture. An open question is whether the pre-existing complexity and
functional role of a maturase-like protein was needed to evolve an enzyme capable of reducing
N_2. The role of evolutionary contingency in shaping biological diversity has long been examined
(Gould 1989; Vermeij 2006; Blount et al. 2018), particularly to envision future evolutionary
scenarios or alternate trajectories on other worlds characterized by distinct environmental
parameters (Kaçar and Gaucher 2012). Regarding the evolution of biological nitrogen fixation in
particular, it has been argued that necessity—i.e., the need for bioavailable sources of nitrogen—
or environmental geochemistry likely controlled the timing of early nitrogenase evolution and
diversification on Earth (Navarro-Gonzalez et al. 2001; Anbar and Knoll 2002; Mus et al. 2018).
However, another possibility is that the origin of biological nitrogen fixation required the subsumed
complexity of a protein predecessor (Adam et al. 2018), which was initially positively selected for
an entirely different metabolic role. The origin of nitrogenases thus may not have occurred (or
may have been significantly delayed) without a suitable protein on which to build, despite the
scarcity of bioavailable nitrogen, an abundance of possible metal cofactors, or both. Testing these
possibilities would require experimentally replaying the evolutionary path that led to the origin of
biological nitrogen fixation.
431

There is no evidence that Terran biology evolved nitrogen fixation more than once. Whether this
is the product of very exceptional circumstances of origination or of a survivorship bias so
pronounced that there is only one remaining functional example supported across all of biology,
nitrogenases are therefore on par with other singular molecular-level innovations such as the
ribosome (Fox 2010) and oxygenic photosynthesis (Blankenship and Hartman 1998). The existing
variation among different metal-dependent forms of nitrogenase enzymes do not constitute truly
independent evolutionary experiments, but variations on a theme that was determined and
uniquely constrained by the common ancestral form. Even within this narrow range of constraints,

the degree to which contingent amino-acid substitutions shaped the diversification of nitrogenase

metal preference and specificity remains unknown.

Identifying the singular circumstances that left Earth with a single common ancestor for all

nitrogenase function may be critical for understanding the pervasiveness of nitrogen fixation as a

universal biological capability. This is particularly important for assessing whether metal

availability significantly guides protein evolution or whether other internal biophysical constraints

lead to background-dependent, epistatic interactions (Williams and da Silva 1996; Anbar and

Knoll 2002; Moore et al. 2017; Smethurst and Shcherbik 2021). Efforts to generate artificial

nitrogenases and nitrogenase metalloclusters (Tanifuji et al. 2015; Sickerman et al. 2017) may

expand the suite of molecular structures capable of reducing N_2, but biotic experiments integrating

gene regulatory and protein-protein interaction constraints are needed to test different

macroevolutionary hypotheses of nitrogenase emergence. A survey of such functional constraints

on nitrogenase and maturase predecessors could reveal the sequence of biomolecular functions

conducive for the evolution of nitrogen fixation, which could then be integrated into a more

comprehensive accounting of internal selective forces, geochemical features and planetary

environments that can host similar evolutionary pathways (Kacar et al. 2020).

Perhaps most intriguingly, our results suggest that nitrogen fixation may have emerged from

natural selection acting on a maturase-like protein whose ancestral function was largely

decoupled from extracellular conditions frequently implicated as drivers for the origins of nitrogen

fixation. This scenario would thus cast the origin of nitrogen fixation as an act of extreme

contingency bordering on happenstance, betraying its utility as one of the most evolutionarily

significant and biologically limiting metabolic pathways on Earth. If borne out by further study or

found to be a recurring pattern for other critical molecular innovations, enzyme origins largely

decoupled from putative environmental drivers may severely compromise the soundness of

systems-focused hypotheses that tie organismal or ecological need to bulk geochemical substrate

or cofactor availabilities. The paleobiology of molecular innovations would require disciplinary

approaches and conceptual foundations quite distinct from the study of their more recent

evolution.

MATERIALS AND METHODS

Phylogenetic reconstruction and ancestral sequence inference

Protein homologs were identified from the National Center for Biotechnology Information non-redundant protein database by BLASTp (Camacho et al. 2009) with an expect value cutoff of <1e-5 (accessed January 2020). Query sequences from *Azotobacter vinelandii* (NifD: WP_01270336, NifK: WP_012698833; NifE: WP_012698838, NifN: WP_012698839) were used for nitrogenase and maturase homolog identification, and sequences from *Synechococcus elongatus* (BchN: WP_126148028, BchB: WP_126147769) for outgroup dark-operative protochlorophyllide oxidoreductase homolog identification. Sequences from this relatively permissive BLASTp search were aligned using MAFFT v7.450 (Katoh and Standley 2013) to build a preliminary phylogeny with FastTree v2.1.11 (Price et al. 2010). Putative nitrogenase homologs were identified based on previously published phylogenies (Boyd et al. 2011a; Garcia et al. 2020) as well as sequence features known to be critical for N_2 reduction (e.g., Cys275, His442). Putative maturase homologs were only retained if the encoding genes were co-localized with nitrogenase genes in the same genome. Finally, sequences in overrepresented clades were pruned to obtain a roughly equal number of nitrogenase and maturase versus outgroup sequences, so as not to bias subsequent ancestral sequence inference.

A final untrimmed MAFFT alignment was used as input for phylogenetic reconstruction by RAxML v8.2.11 (Stamatakis 2014) using 100 rapid bootstrap searches and the best-fit LG+G+F evolutionary model determined by ModelFinder (Kalyaanamoorthy et al. 2017) (an additional phylogeny was also built from an alignment trimmed by TrimAl (Capella-Gutierrez et al. 2009) (supplementary fig. S2)). The tree was further optimized by applying nearest-neighbor-interchanges before calculation of SH-like branch support values (Guindon et al. 2010), resulting in the final phylogeny, Tree-1. Additional trees incorporated in ancestral sequenc inference were generated by altering sequence sampling or alignment with MUSCLE v3.8.425 (Edgar 2004) instead of MAFFT (see Results and Discussion, table 1). Finally, the Bayesian phylogenetic analysis by Boyd et al. (Boyd et al. 2011a) was replicated using their reported methods (supplementary fig. S2).

Ancestral sequences were inferred by maximum likelihood marginal reconstruction in PAML v4.9j (Yang 2007) using the same evolutionary model parameters described above for RAxML. Sequence gaps were reconstructed in PAML using the binary encoding approach described in Aadland et al. (2019). Briefly, the protein sequence alignment was recoded as a 'presence-

absence' alignment matrix, and the posterior-probability of the presence (amino-acid residue) or

absence (gap) state at each position in each ancestral sequence was calculated using maximum-

likelihood reconstruction, assuming a binary character model with state frequencies inferred by

maximum likelihood. All phylogenetic data, including sequence alignments, trees, and ancestral

sequence inference outputs can be found at https://github.com/kacarlab/maturase2021.

Prediction of functionally divergent protein sites between nitrogenases and maturases

For each position in the sequence alignment, we calculated the Jensen-Shannon (J-S) distance

between the amino-acid frequency distribution estimated from extant nitrogenase Nif/Vnf/AnfD

sequences in the alignment and that estimated from extant maturase Nif/VnfE sequences, with

nitrogenase and maturase sequences being defined based on monophyly in the tree topology

and gene location within the *nif*, *vnf*, or *anf* loci (see Results and Discussion, fig. 1b). Briefly, the

J-S distance is calculated as the average Kullback-Leibler divergence, or "relative entropy," which

estimates the loss of information when one frequency distribution is used to represent another.

Intuitively, the site-wise J-S distance between nitrogenase and maturase amino-acid frequency

distributions describes how dissimilar the distribution of extant amino-acids is between

nitrogenase and maturase sequences for each alignment site.

We estimated the expected distribution of site-wise J-S distances, given our sequence alignment,

by randomly partitioning the alignment into two sequence groups of sizes equivalent to the sizes

of our actual nitrogenase and maturase groups, respectively, and calculating site-wise J-S

distances between these randomly partitioned groups. We performed 10,000 random partitions

and site-wise J-S distance calculations. For each alignment column i, we calculated the probability

of observing J-S(nitrogenase, maturase)$_i$, given the distribution of J-S-distances at column i in the

randomly partitioned dataset (i.e., p-value). We enriched for highly divergent positions using an

FDR-corrected p-value cutoff of 0.0001. We additionally excluded any sites with J-S distances in

the lower 75th percentile of the J-S distance distribution across all alignment positions.

Probabilistic assessment of nitrogenase-like ancestral sequence features

Nitrogenase-like ancestral sequence features were assessed by incorporating the statistical
uncertainty of ancestral sequence inference into comparisons between those of extant
nitrogenases and maturases. For putative functionally divergent protein sites identified as
described above, J-S distances were calculated between the ancestral amino acid posterior
probability distributions and the extant amino acid frequency distributions across either extant
nitrogenase (Nif/Vnf/AnfD) or maturase (Nif/VnfE) clades. These distance values were then
normalized to yield a value between -1 and +1 indicating the relative similarity of an ancestral
protein site to a homologous extant nitrogenase site, here called the "D-score" (i.e., similarity to
the nitrogenase D-subunit):

$$\text{D-score} = (\text{dist}_E - \text{dist}_D) / (\text{dist}_E + \text{dist}_D)$$

where dist_D is the J-S distance between ancestral and maturase E-subunit sites, and dist_E is the
J-S distance between ancestral and nitrogenase D-subunit sites. D-scores were analyzed on a
site-wise basis as well as averaged across the length of each ancestral sequence for all
constructed phylogenies. All data and scripts related to the prediction of functionally divergent
sites prediction and D-score calculations can be found at
https://github.com/kacarlab/maturase2021.

ACKNOWLEDGEMENTS

This work was supported by the National Aeronautics and Space Administration Postdoctoral
Program to A.K.G., the National Science Foundation Division of Molecular and Cellular
Biosciences (1817942 to B. Kolaczkowski), the National Science Foundation Emerging Frontiers
Program (1724090 to B. Kaçar), the National Aeronautics and Space Administration Early Career
Faculty Award (80NSSC19K1617 to B. Kaçar), and the Metal Utilization and Selection across
Eons (MUSE) Interdisciplinary Consortium for Astrobiology Research, sponsored by the National
Aeronautics and Space Administration Science Mission Directorate (19-ICAR19_2-0007 to B.
Kaçar). We thank Eric Boyd and Jennifer Glass for the helpful discussions.

FIGURES AND FIGURE LEGENDS

Fig. 1. Nitrogenase and maturase functionality, genetic organization, and evolutionary history. (*A*)

Simplified biosynthetic pathway for nitrogenase active-site cluster maturation and incorporation

(molybdenum-dependent nitrogenase system shown). Pathway steps are indicated by circled

numbers. Step 1). Maturase NifEN proteins (NifE, dark green; NifN, brown), which are $\alpha_2\beta_2$

heterotetramer homologs to nitrogenase NifDK proteins (NifD, teal; NifK, yellow), are scaffolds for

maturation of nitrogenase iron-sulfur cluster precursors (yellow diamond). Step 2) Cluster

precursors are matured to M-clusters (yellow diamond with teal circle) by incorporation of

molybdenum and homocitrate, delivered to the maturase complex by NifH proteins. Step 3)

Mature M-clusters are incorporated into the nitrogenase complex, where they serve as the active

sites for N_2 reduction to NH_3. During enzyme turnover, NifH proteins transiently interface with the

nitrogenase NifDK complex to deliver electrons to the M-cluster active site. (*B*) Representative

gene locus structures for molybdenum-dependent (Nif), vanadium-dependent (Vnf), and iron-

dependent (Anf) nitrogenase systems, as well as for uncharacterized nitrogenase homologs (see

text for discussion). Gene and intergenic region lengths are approximate. Hash marks indicate

significant distance between represented genes. (*C*) Maximum likelihood phylogenetic tree (Tree-

1; see text for details) built from nitrogenase and maturase protein sequences. α- or β-subunit

designations for protein sequences are indicated on the right. Homologs from uncharacterized

taxa are highlighted in red. Clade widths are not to scale. "LCA": Last common ancestor.

A

		96
A. vinelandii NifD	MTG-MSREEVESLIQEVLEVYPEKARKDRNKHLAVNDPAV-TQS-KKCIILSNKKSQPGLMTIRGCAYAGSKGVVWGPIKDMIHISHGPVGCGCYSRAGR	
A. vinelandii NifE	·KA-------K-----DIA---DLL--DEPACSH·K---KE-K--S--GCA·P-K---ADG·SFT·VQ·AQ-IALLVA·VA·V·IAAGS·WDN	
M. thermoacetica NifE	ELSVD----------MLPE·RQ·MQRVG-------·LPAC·RPTI··VVSQ·A·VLX·AR-WMLAT·AL·LI···AY·ETV	
Ancestor △	·SKSTT---------T·-·-VLPE·K·-ICLKG---EDEPL-S-V·ACT·TI·V·Q·S···AR-·LA···LV··I·AV·WGV	
Ancestor △	·EAA---------TI·--·-VLPE·K·ICLKG---EDEK--EAK·AC--·V··V·Q··F··AA-R·LAG·A·ILV··Q·AY·-V	
Ancestor ▲	·E----------TV·--·-LPE·K---K----E-------C·TA·SVNPQ··L·AA-R·LAG··AMILV·Q·AY·--V	

		190
	RNYY----IGTTGVNAFVTMNFTSDFQEKDIVFG-GDKKLAKLIDEVETLFPLNKGISVQSECFIGLIGDDIESVSKVKGAEL-SKTIVPVRCEGFRGVS	
	GTRS------S-ED-LYRIGMT·LT·N·VIM·RAE·R·FHA·RQAVESY-SPPAVE·YNT·VPA···VDA·C·AAAERF-GTPVI·DSA··-YGT	
	·SLS------S-E·K-RYEV-·STALD···I··I·-E-·YQA·L·AVEE··EP·A·YAT·TT···A·C·KAEE·I-GIPVI·N·P··-	
	TALSRH------EE-VYRPVASTNL···I·-·I-·E·K·A·R·VERY-RP·V·E·V·T·AT·I···G·CRKLEE·I-GVPVI·N·P··-	
	TALSRHFA-----·E-PVYRPVASTNL··--EE··K·A·R·VERY-RP·V·E·V·T·ASEI···GICRKLEE·IPGVPVI·NAP··-SG	

		280
	QSLGHHIANDAVRDWVLGKRDED-TT-------FASTPYDVAIIGDYNIC-GJAWSSRILLEEMGLRCVAQWSGDGSISEIELTPKVKLNLVHCYRSMN	
	KN--·NR·GE-·MLKY·I·T·E----PDPLPVGSERPGIRVH··NL·E··A-·DF·HVLP·D·L··VLCTLA·ARYR·VQTMHRAEV·MMV·SKA·L	
	·CD-·EA·AL·LLEHFI·RGG--EGS-------PLAK-S·N·L·D·Q--·LKEIEG··QRL·VEVLCTV··RATVAGLARAHRAG····I··Q·AT	
	·H·A·YDL·FE·LLEHLV·TKE---PE-------EP··IN··E··VA-·L·VIKP-··K··I·VISTFT··A·D·LAKAHRA···I··Q·AT	
	·H·E·YDL·FE·LLEHLAKAEP----E-------KPGK-·IN··Q·SPA·LREIKR-··K··I·V·SVPT·NTT··E·LKRAGRA-V·I·L·QDVAT	

		376
	YISRHMEEKYGIPWMEYNF-FGPTKTIESLRAIAAKF-DESIQKKCCEEVIAKYKPEWEAVVAKYRPRLEGKRVMLYIGGLRPRHVIGAYE-DLGMEVVG	
	NVA-·KLQ·T··T·F·GS-·-Y·I·D·SQA--·DF·RLLD·PDLTART·AL·REEAKVR·ALEPW·A-·-··L··T··V-KSWS·VS·LQ--··K·A	
	RLALE-·A·RF-··RKVS-·-·LAE·AAA··T··DV·H---CPE-KATWLEEEIESAREKASP·LE·-A·-··G·FF·AS-·MGSMAR·FR-E···-··F	
	·LAKM-----·-·YLKVE·-·IEN·SK··K·KF·G·ELEERA·-·EEEMAKIKPKLEQ··K····K·AI·V··S-·ANSMVK·FK-E·····V	
	·LAKK-----·-·YLKVE·P·IEN·SKW··K·KF·G·--LEERA··L·ERREREKIKDKLEQA·KY·Q····AI·G·DATKALSLVRMLKRE-····P·V	
	·LAKKL----·V·YLKVE·P·IEN·SRW··-KF·GE--LEGRA··L·ERERERVKDKLEEA·KY·R····AI·G-DATLASLVRFLKRE-····P·V	

		467
	TGYEFAHNDDYDRTM--KEMGDS-TLLYDDVTSYEFEEFVKRIKEDLIGSGLKEKFIFCKMG------IPFRQMHSWDYSGPYHGFDGFAIFARDMDMTL	
	·-TK·STEE·KA--IREL·-·D-VKML·EGNARVLLKT·DEYQA··ILIA·CRNMYTAL·GR------V·-··LDINQER-EEG·A·Y··MLELV·QLCI	
	C·SQ·GCRE·-AE--SRSRL·RE-·-·V·AHCR·L·-·LH·YRCH·LGA·SR·Q·MAH·L·------V·-··LVFFQE--TA··A··N·-·VNL·EVATLI	
	A·TQ·GSKE·-EK--MREQVK·G-·-·II·ANSM·L·LL·KY··FVG·T··YLAH·L-------V·-··LNF··-EE-TS··A·E··VN·--·V·KAI	
	V·TQ·GTKE·-EK-LKELVE·G-·-IIIN·ANA··L·LL·EA··LIG·AN··YLAH·L·ILTKVS·-·LLDLNAAR--NR··F·YE··ALNL·KTINKA	
	V·TQ·GTKKEFEKE-LKELVEEGL·II-N·ADA··L·LI·EA··LIG·AN··YLAH·L·ILTKVS·-·LLDRNAAR--NR··F·YE··AMNL·ETISNA	

		492
	N--NPCWKKLQA--PWEA--------SE--GAEKVA--ASA	
	E--C·V·EAVRRPA·DIPASQDAAP·APARS·N-------	
	G--A·V·RLA---AG------------------SP---APG-	
	--·S·V··LVRK-A-·KT----------------------	
	--·S·V··L·RKRA-·-RAA---------------------	
	KERS·A··L·KRRA-·-RAA---------------------	

◆ nitrogenase M-cluster active site
✳ conserved in nitrogenases
☐ conserved in maturases
▨ conserved in both

B

■ Sites at nitrogenase NifD-NifK interface

J-S distance
0.54 — 0.80

A. vinelandii nitrogenase (NifD)

A. vinelandii nitrogenase (NifD)

A. vinelandii maturase (NifE)

M-cluster active site

180°

180°

180°

NifD

M-cluster insertion funnel

NifK

NifD

M-cluster precursor binding site

NifN

NifE

Fig. 2. Structure and sequence maps of putative functionally divergent protein sites between nitrogenases and maturases. (*A-B*) Putative functionally divergent sites are defined as those above the 75th percentile J-S distance between nitrogenase NifD and maturase NifE proteins (see text for details). The J-S distance scale applies to both (*A*) and (*B*). (*A*) Maximum likelihood ancestors (inferred from Tree-1; fig. 1c) aligned to representative extant nitrogenase (NifD) and maturase (NifE) sequences. *Moorella thermoacetica (M. thermoacetica)* NifE is an uncharacterized, putative maturase sequence. Ancestor triangle symbols correspond to labeled nodes in the simplified phylogeny (*left*) and match those in fig. 3. Nitrogenase M-cluster active-site residues are defined as those within 5 Å of the M-cluster. Dots within the alignment indicate residue identity to *Azotobacter vinelandii* (*A. vinelandii*) NifD. Site numbering based on *A. vinelandii* NifD. (*B*) Divergent sites mapped to aligned nitrogenase (*center*; *A. vinelandii* NifD, PDB 3U7Q) and maturase (*right*; *A. vinelandii* NifE, PDB 3PDI) subunit structures. Protein sites at the nitrogenase NifD-NifK interface (*left*) are defined as those within 5 Å of the NifK subunit.

Fig. 3. Ancestral sequence similarity to extant nitrogenases or maturases, mapped across four alternate phylogenies: (*A*) Tree-1, (*B*) Tree-2, (*C*) Tree-3, and (*D*) Tree-4. Similarity is expressed as the "D-score" parameter, where a positive D-score (red) indicates greater similarity to extant nitrogenases, and a negative D-score (blue) indicates greater similarity to extant maturases (see Materials and Methods). D-scores were averaged across putative functionally divergent sites for each ancestral node. Nodes are labeled nitrogenase/maturase last common ancestor ("LCA") 1— including early diverged VnfE homologs (black triangle; see text for discussion); nitrogenase

621 /maturase LCA 2—excluding early diverged VnfE homologs (grey triangle); and nitrogenase LCA

622 (white triangle), along with their mean D-scores.

624
625

Fig. 4. Proposed model for the origins and functional divergence of maturase and nitrogenase

proteins. An ancestral maturase-like protein (~NifEN, grey and light brown), incapable of reducing

N_2, may have otherwise reduced various carbon-containing substrates and/or played a role in

cluster (yellow diamond) biosynthesis. The ancestor may have been capable of incorporating

molybdenum (teal circle) into the cluster. Duplication of the encoding ancestral genes and

functional divergence would then have yielded canonical maturase (NifEN, dark green and brown)

and nitrogenase (NifDK, teal and yellow) proteins. Maturases would have specialized to provide

a scaffold for the maturation of the nitrogenase cluster precursor (yellow diamond) to the

nitrogenase active-site M-cluster (yellow diamond with teal circle). In parallel, tuning of the

ancestral peptide environment along a divergent lineage would have spurred the origin of N_2

reduction and specialization of nitrogenases for a solely catalytic role in nitrogen fixation. Protein

components of the $\alpha_2\beta_2$ heterotetrameric nitrogenase and maturase structures are labeled.

TABLES

Table 1. Nitrogenase and maturase phylogenies built in this study.

Tree	Alignment method	Sequence Dataset

Tree-1	MAFFT	Nif/Vnf/AnfDK, Nif/VnfEN, BchNB
Tree-2	MAFFT	Tree-1 dataset without uncharacterized homologs
Tree-3	MAFFT	Tree-1 dataset without β-subunit sequences (Nif/Vnf/AnfK, Nif/VnfN, BchB)
Tree-4	MUSCLE	Same as Tree-1 dataset

DATA AVAILABILITY STATEMENT

All phylogenetic datasets and scripts are available at https://github.com/kacarlab/maturase2021.

REFERENCES

Aadland K, Pugh C, Kolaczkowski B. 2019. High-Throughput Reconstruction of Ancestral Protein Sequence, Structure, and Molecular Function. *Methods Mol Biol*. 1851:135-170. doi: 10.1007/978-1-4939-8736-8_8.

Adam ZR, Fahrenbach AC, Kacar B, Aono M. 2018. Prebiotic Geochemical Automata at the Intersection of Radiolytic Chemistry, Physical Complexity, and Systems Biology. *Complexity*. 2018:1-21. doi: 10.1155/2018/9376183.

Allen JF, Thake B, Martin WF. 2019. Nitrogenase Inhibition Limited Oxygenation of Earth's Proterozoic Atmosphere. *Trends Plant Sci*. 24:1022-1031. doi: 10.1016/j.tplants.2019.07.007.

Anbar AD. 2008. Oceans. Elements and evolution. *Science*. 322:1481-1483. doi: 10.1126/science.1163100.

Anbar AD, Knoll AH. 2002. Proterozoic ocean chemistry and evolution: a bioinorganic bridge? *Science*. 297:1137-1142. doi: 10.1126/science.1069651.

Benner S. 2017. Uniting natural history with the molecular sciences. The ultimate multidisciplinarity. *Acc Chem Res*. 50:498-502. doi: 10.1021/acs.accounts.6b00496.

Benner SA, Sassi SO, Gaucher EA. 2007. Molecular paleoscience: systems biology from the
past. *Adv Enzymol Relat Areas Mol Biol*. 75:1-132, xi.

Blankenship RE, Hartman H. 1998. The origin and evolution of oxygenic photosynthesis. *Trends*
Biochem Sci. 23:94-97. doi: 10.1016/S0968-0004(98)01186-4.

Blount ZD, Lenski RE, Losos JB. 2018. Contingency and determinism in evolution: Replaying
life's tape. *Science*. 362:eaam5979. doi: 10.1126/science.aam5979.

Boyd ES, Anbar AD, Miller S, Hamilton TL, Lavin M, Peters JW. 2011a. A late methanogen
origin for molybdenum-dependent nitrogenase. *Geobiology*. 9:221-232. doi: 10.1111/j.1472-
4669.2011.00278.x.

Boyd ES, Hamilton TL, Peters JW. 2011b. An alternative path for the evolution of biological
nitrogen fixation. *Front Microbiol*. 2:205. doi: 10.3389/fmicb.2011.00205.

Boyd ES, Peters JW. 2013. New insights into the evolutionary history of biological nitrogen
fixation. *Front Microbiol*. 4:201. doi: 10.3389/fmicb.2013.00201.

Buren S, Jimenez-Vicente E, Echavarri-Erasun C, Rubio LM. 2020. Biosynthesis of Nitrogenase
Cofactors. *Chem Rev*. 120:4921-4968. doi: 10.1021/acs.chemrev.9b00489.

Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. 2009.
BLAST+: architecture and applications. *BMC Bioinformatics*. 10:421. doi: 10.1186/1471-2105-
10-421.

Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T. 2009. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*. 25:1972-1973. doi:
10.1093/bioinformatics/btp348.

Chen Y, Nishihara A, Haruta S. 2021. Nitrogen-fixing Ability and Nitrogen Fixation-related
Genes of Thermophilic Fermentative Bacteria in the Genus *Caldicellulosiruptor*. *Microbes*
Environ. 36:ME21018. doi: 10.1264/jsme2.ME21018.

Copley SD. 2015. An evolutionary biochemist's perspective on promiscuity. *Trends Biochem*
Sci. 40:72-78. doi: 10.1016/j.tibs.2014.12.004.

Copley SD. 2021. Setting the stage for evolution of a new enzyme. *Curr Opin Struct Biol*. 69:41-
49. doi: 10.1016/j.sbi.2021.03.001.

De Tarafder A, Parajuli NP, Majumdar S, Kacar B, Sanyal S. 2021. Kinetic Analysis Suggests
Evolution of Ribosome Specificity in Modern Elongation Factor-Tus from 'Generalist' Ancestors.
Mol Biol Evol.msab114. doi: 10.1093/molbev/msab114.

Dekas AE, Poretsky RS, Orphan VJ. 2009. Deep-sea archaea fix and share nitrogen in
methane-consuming microbial consortia. *Science*. 326:422-426. doi: 10.1126/science.1178223.

Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res*. 32:1792-1797. doi: 10.1093/nar/gkh340.

Einsle O, Rees DC. 2020. Structural Enzymology of Nitrogenase Enzymes. *Chem Rev*.
120:4969-5004. doi: 10.1021/acs.chemrev.0c00067.

Falkowski PG. 1997. Evolution of the nitrogen cycle and its influence on the biological
sequestration of CO_2 in the ocean. *Nature*. 387:272-275. doi: 10.1038/387272a0.

Fani R, Gallo R, Lio P. 2000. Molecular evolution of nitrogen fixation: the evolutionary history of
the *nifD*, *nifK*, *nifE*, and *nifN* genes. *J Mol Evol*. 51:1-11. doi: 10.1007/s002390010061.

Fay AW, Blank MA, Rebelein JG, Lee CC, Ribbe MW, Hedman B, Hodgson KO, Hu Y. 2016.
Assembly scaffold NifEN: A structural and functional homolog of the nitrogenase catalytic
component. *Proc Natl Acad Sci USA*. 113:9504-9508. doi: 10.1073/pnas.1609574113.

Fox GE. 2010. Origin and Evolution of the Ribosome. *CSH Perspect Biol*. 2:a003483-a003483.
doi: 10.1101/cshperspect.a003483.

Fujita Y, Bauer CE. 2000. Reconstitution of Light-independent Protochlorophyllide Reductase
from Purified BchI and BchN-BchB Subunits. *J Biol Chem*. 275:23583-23588. doi:
10.1074/jbc.M002904200.

Garcia AK, Kacar B. 2019. How to resurrect ancestral proteins as proxies for ancient
biogeochemistry. *Free Radic Biol Med*. 140:260-269. doi: 10.1016/j.freeradbiomed.2019.03.033.

Garcia AK, McShea H, Kolaczkowski B, Kacar B. 2020. Reconstructing the evolutionary history
of nitrogenases: Evidence for ancestral molybdenum-cofactor utilization. *Geobiology*. 18:394-
411. doi: 10.1111/gbi.12381.

Gerlt JA, Babbitt PC. 2001. Divergent Evolution of Enzymatic Function: Mechanistically Diverse
Superfamilies and Functionally Distinct Suprafamilies. *Annu Rev Biochem*. 70:209-246. doi:
10.1146/annurev.biochem.70.1.209.

Ghebreamlak SM, Mansoorabadi SO. 2020. Divergent Members of the Nitrogenase
Superfamily: Tetrapyrrole Biosynthesis and Beyond. *ChemBioChem*. 21:1723-1728. doi:
10.1002/cbic.201900782.

Glass JB, Wolfe-Simon F, Anbar AD. 2009. Coevolution of metal availability and nitrogen
assimilation in cyanobacteria and algae. *Geobiology*. 7:100-123. doi: 10.1111/j.1472-
4669.2009.00190.x.

Gould SJ. 1989. Wonderful Life: The Burgess Shale and the Nature of History. New York, NY:
W. W. Norton & Company, Inc.

Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010. New algorithms
and methods to estimate maximum-likelihood phylogenies: assessing the performance of
PhyML 3.0. *Syst Biol*. 59:307-321. doi: 10.1093/sysbio/syq010.

Hoffman BM, Lukoyanov D, Yang ZY, Dean DR, Seefeldt LC. 2014. Mechanism of nitrogen
fixation by nitrogenase: the next stage. *Chem Rev*. 114:4041-4062. doi: 10.1021/cr400641x.

Hu Y, Fay AW, Lee CC, Wiig JA, Ribbe MW. 2010. Dual functions of NifEN: insights into the

evolution and mechanism of nitrogenase. *Dalton T*. 39. doi: 10.1039/b922555b.

Hu Y, Fay AW, Lee CC, Yoshizawa J, Ribbe MW. 2008. Assembly of Nitrogenase MoFe

Protein. *Biochemistry*. 47:3973-3981. doi: 10.1021/bi7025003.

Hu Y, Fay AW, Schmid B, Makar B, Ribbe MW. 2006. Molecular Insights into Nitrogenase

FeMoco Insertion. *J Biol Chem*. 281:30534-30541. doi: 10.1074/jbc.M605527200.

Jimenez-Vicente E, Yang ZY, Ray WK, Echavarri-Erasun C, Cash VL, Rubio LM, Seefeldt LC,

Dean DR. 2018. Sequential and differential interaction of assembly factors during nitrogenase

MoFe protein maturation. *J Biol Chem*. 293:9812-9823. doi: 10.1074/jbc.RA118.002994.

Kacar B, Garcia AK, Anbar AD. 2020. Evolutionary History of Bioessential Elements Can Guide

the Search for Life in the Universe. *ChemBioChem*. doi: 10.1002/cbic.202000500.

Kaçar B, Gaucher EA. 2012. Towards the recapitulation of ancient history in the laboratory:

Combining synthetic biology with experimental evolution. *Artif Life*. 13:11-18. doi: 10.1162/978-

0-262-31050-5-ch002.

Kaiser JT, Hu Y, Wiig JA, Rees DC, Ribbe MW. 2011. Structure of Precursor-Bound NifEN: A

Nitrogenase FeMo Cofactor Maturase/Insertase. *Science*. 331:91-94. doi:

10.1126/science.1196954.

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017. ModelFinder: fast

model selection for accurate phylogenetic estimates. *Nature Methods*. 14:587-589. doi:

10.1038/nmeth.4285.

Kanehisa M, Sato Y, Morishima K. 2016. BlastKOALA and GhostKOALA: KEGG Tools for

Functional Characterization of Genome and Metagenome Sequences. *J Mol Biol*. 428:726-731.

doi: 10.1016/j.jmb.2015.11.006.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. *Mol Biol Evol*. 30:772-780. doi:

10.1093/molbev/mst010.

Kim CH, Newton WE, Dean DR. 1995. Role of the MoFe protein alpha-subunit histidine-195

residue in FeMo-cofactor binding and nitrogenase catalysis. *Biochemistry*. 34:2798-2808.

Kim J, Rees D. 1992. Structural models for the metal centers in the nitrogenase molybdenum-

iron protein. *Science*. 257:1677-1682. doi: 10.1126/science.1529354.

Lee H-I, Thrasher KS, Dean DR, Newton WE, Hoffman BM. 1998. ^{14}N Electron Spin−Echo

Envelope Modulation of the S = 3/2 Spin System of the *Azotobacter vinelandii* Nitrogenase

Iron−Molybdenum Cofactor. *Biochemistry*. 37:13370-13378. doi: 10.1021/bi980956a.

Li H. 2002. *Azotobacter vinelandii* Nitrogenase: Multiple Substrate-Reduction Sites and Effects

of pH on Substrate Reduction and CO Inhibition. [Dissertation]. [Blacksburg, VA]: Virginia

Polytechnic Institute and State University.

Luo G, Junium CK, Izon G, Ono S, Beukes NJ, Algeo TJ, Cui Y, Xie S, Summons RE. 2018.

Nitrogen fixation sustained productivity in the wake of the Palaeoproterozoic Great Oxygenation

Event. *Nat Commun*. 9:978. doi: 10.1038/s41467-018-03361-2.

McGlynn SE, Boyd ES, Peters JW, Orphan VJ. 2012. Classifying the metal dependence of

uncharacterized nitrogenases. *Front Microbiol*. 3:419. doi: 10.3389/fmicb.2012.00419.

Mehta MP, Baross JA. 2006. Nitrogen fixation at 92 degrees C by a hydrothermal vent

archaeon. *Science*. 314:1783-1786. doi: 10.1126/science.1134772.

Moore EK, Jelen BI, Giovannelli D, Raanan H, Falkowski PG. 2017. Metal availability and the

expanding network of microbial metabolisms in the Archaean eon. *Nat Geosci*. 10:629-636. doi:

10.1038/ngeo3006.

Moser J, Brocker MJ. 2011. Enzymatic Systems with Homology to Nitrogenase. In: Ribbe M,

editor. Nitrogen Fixation. Methods in Molecular Biology (Methods and Protocols): Humana

Press. p. 67-77.

Mus F, Alleman AB, Pence N, Seefeldt LC, Peters JW. 2018. Exploring the alternatives of

biological nitrogen fixation. *Metallomics*. 10:523-538. doi: 10.1039/c8mt00038g.

Mus F, Colman DR, Peters JW, Boyd ES. 2019. Geobiological feedbacks, oxygen, and the

evolution of nitrogenase. *Free Radic Biol Med*. doi: 10.1016/j.freeradbiomed.2019.01.050.

Navarro-Gonzalez R, McKay CP, Mvondo DN. 2001. A possible nitrogen crisis for Archaean life

due to reduced nitrogen fixation by lightning. *Nature*. 412:61-64. doi: Doi 10.1038/35083537.

Noda-Garcia L, Liebermeister W, Tawfik DS. 2018. Metabolite-Enzyme Coevolution: From

Single Enzymes to Metabolic Pathways and Networks. *Annu Rev Biochem*. 87:187-216. doi:

10.1146/annurev-biochem-062917-012023.

North JA, Narrowe AB, Xiong W, Byerly KM, Zhao G, Young SJ, Murali S, Wildenthal JA,

Cannon WR, Wrighton KC, et al. 2020. A nitrogenase-like enzyme system catalyzes

methionine, ethylene, and methane biogenesis. *Science*. 369:1094-1098. doi:

10.1126/science.abb6310.

Ohno S. 1970. Evolution by gene duplication. Berlin: Springer-Verlag.

Parsons C, Stüeken EE, Rosen CJ, Mateos K, Anderson RE. 2020. Radiation of nitrogen-

metabolizing enzymes across the tree of life tracks environmental transitions in Earth history.

Geobiology. 19:18-34. doi: 10.1111/gbi.12419.

Perez-Gonzalez A, Jimenez-Vicente E, Gies-Elterlein J, Salinero-Lanzarote A, Yang ZY, Einsle

O, Seefeldt LC, Dean DR. 2021. Specificity of NifEN and VnfEN for the Assembly of

Nitrogenase Active Site Cofactors in Azotobacter vinelandii. *mBio*. 12:e0156821. doi:

10.1128/mBio.01568-21.

Price MN, Dehal PS, Arkin AP. 2010. FastTree 2--approximately maximum-likelihood trees for
large alignments. *PLoS One*. 5:e9490. doi: 10.1371/journal.pone.0009490.

Raymond J, Siefert JL, Staples CR, Blankenship RE. 2004. The natural history of nitrogen
fixation. *Mol Biol Evol*. 21:541-554. doi: 10.1093/molbev/msh047.

Seefeldt LC, Yang ZY, Lukoyanov DA, Harris DF, Dean DR, Raugei S, Hoffman BM. 2020.
Reduction of Substrates by Nitrogenases. *Chem Rev*. 120:5082-5106. doi:
10.1021/acs.chemrev.9b00556.

Sickerman NS, Tanifuji K, Hu Y, Ribbe MW. 2017. Synthetic Analogues of Nitrogenase
Metallocofactors: Challenges and Developments. *Chemistry*. 23:12425-12432. doi:
10.1002/chem.201702496.

Silver WS, Postgate JR. 1973. Evolution of asymbiotic nitrogen fixation. *J Theor Biol*. 40:1-10.
doi: 10.1016/0022-5193(73)90160-4.

Smethurst DGJ, Shcherbik N. 2021. Interchangeable utilization of metals: New perspectives on
the impacts of metal ions employed in ancient and extant biomolecules. *J Biol Chem*.
297:101374. doi: 10.1016/j.jbc.2021.101374.

Solomon JB, Lee CC, Jasniewski AJ, Rasekh MF, Ribbe MW, Hu Y. 2020. Heterologous
Expression and Engineering of the Nitrogenase Cofactor Biosynthesis Scaffold NifEN. *Angew*
Chem Int Ed Engl. 59:6887-6893. doi: 10.1002/anie.201916598.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics*. 30:1312-1313. doi: 10.1093/bioinformatics/btu033.

Staples CR, Lahiri S, Raymond J, Von Herbulis L, Mukhophadhyay B, Blankenship RE. 2007.
Expression and association of group IV nitrogenase NifD and NifH homologs in the non-
nitrogen-fixing archaeon *Methanocaldococcus jannaschii*. *J Bacteriol*. 189:7392-7398. doi:
10.1128/JB.00876-07.

Stueken EE, Buick R, Guy BM, Koehler MC. 2015. Isotopic evidence for biological nitrogen
fixation by molybdenum-nitrogenase from 3.2 Gyr. *Nature*. 520:666-669. doi:
10.1038/nature14180.

Stüeken EE, Kipp MA, Koehler MC, Buick R. 2016. The evolution of Earth's biogeochemical
nitrogen cycle. *Earth-Sci Rev*. 160:220-239. doi: 10.1016/j.earscirev.2016.07.007.

Tanifuji K, Lee CC, Ohki Y, Tatsumi K, Hu Y, Ribbe MW. 2015. Combining a Nitrogenase
Scaffold and a Synthetic Compound into an Artificial Enzyme. *Angew Chem Int Ed Engl*.
127:14228-14231. doi: 10.1002/ange.201507646.

Thornton JW. 2004. Resurrecting ancient genes: experimental analysis of extinct molecules.
Nat Rev Genet. 5:366-375. doi: 10.1038/nrg1324.

Vermeij GJ. 2006. Historical contingency and the purported uniqueness of evolutionary
innovations. *Proc Natl Acad Sci USA*. 103:1804-1809. doi: 10.1073/pnas.0508724103.

Vialle RA, Tamuri AU, Goldman N. 2018. Alignment Modulates Ancestral Sequence
Reconstruction Accuracy. *Mol Biol Evol*. 35:1783-1797. doi: 10.1093/molbev/msy055.

Williams RJP, da Silva JRRF. 1996. The natural selection of the chemical elements: the
environment and life's chemistry: Clarendon Press.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol*. 24:1586-
1591. doi: 10.1093/molbev/msm088.

Zheng K, Ngo PD, Owens VL, Yang X-p, Mansoorabadi SO. 2016. The biosynthetic pathway of
coenzyme F430 in methanogenic and methanotrophic archaea. *Science*. 354:339-342. doi:
10.1126/science.aag2947.