Causes of variability in estimates of mutational variance from mutation

accumulation experiments

Cara Conradsen[1*], Mark W. Blows[1] and Katrina McGuigan[1]

[1]School of Biological Sciences; The University of Queensland; St. Lucia, Queensland, Australia 4072

Running Title: Variability of mutational variance

Keywords: meta-analysis, Drosophila, life history, morphology, heritability, coefficient of variance,

wing, micro-environment, measurement error

[*]Corresponding Author: Cara Conradsen

School of Biological Sciences

The University of Queensland;

St. Lucia, QLD, 4072 AUSTRALIA

T: +61(0)450645318

E-mail: cara.conradsen@uqconnect.edu.au

ABSTRACT

Characteristics of the new phenotypic variation introduced via mutation have broad implications in

evolutionary and medical genetics. Estimates of this mutational variance, V_M, span two orders of

magnitude, but factors contributing to this range remain poorly resolved. We applied meta-analyses

to ~200 estimates from 62 studies to assess how previously implicated factors affect the magnitude of

V_M. While estimates are available from a range of taxa, and quantitative trait types over a range of

timescales, these factors are confounded with one another. We call for further directly comparable

empirical data to resolve these influences on variability of V_M. We also analysed data from

Drosophila serrata lines, derived from a mutation accumulation experiment, observing substantial

heterogeneity among replicated estimates of among-line variance for 11 morphological (wing) traits.

Micro-environmental variation in mutational effects was implicated as causing variability in

mutational variance for just two of 11 traits, suggesting estimation environment may contribute little

to variability of published estimates. Varying the opportunity for evolutionary processes to increase

within-line genetic variation did not affect among-line variance estimates, suggesting transient

segregation of mutations is also an unlikely general explanation of variability in published estimates.

We conclude that estimation error, expected to be relatively high due to the low differentiation

among lines diverging solely through new mutation, is likely to contribute strongly to variability of

estimates of mutational variance. Analyses of repeated measures of mutation accumulation lines can

improve the precision of estimates, consequently expanding our understanding of the dynamics of

mutations in natural populations.

INTRODUCTION

The magnitude of per-generational increase in genetic variance due to spontaneous mutations (V_M) is important for a wide range of genetic and evolutionary phenomena, including the maintenance of quantitative genetic variance (LYNCH 1988b; BARTON AND TURELLI 1989; JOHNSON AND BARTON 2005). Much of our understanding of V_M comes from mutation accumulation (MA) experiments, where populations diverge phenotypically due solely to the fixation of new mutations (MUKAI 1964; HALLIGAN AND KEIGHTLEY 2009). Estimates from MA experiments in a wide range of traits and taxa indicate that mutation increases the variance in quantitative traits by 10^{-4} to 10^{-2} times the environmental variance of the trait per generation (HOULE *et al.* 1996; LYNCH AND WALSH 1998; LYNCH *et al.* 1999; HALLIGAN AND KEIGHTLEY 2009). This range of values (two orders of magnitude) suggests variation in either the rate or the average magnitude of effects of new mutation, potentially leading to differences in the magnitude of standing quantitative genetic variation and, ultimately, rates of phenotypic evolution (HOULE 1998; LYNCH *et al.* 1999; HOULE *et al.* 2017; WALSH AND LYNCH 2018). However, the causes of variation in estimates of V_M, and thus the evolutionary interpretation of this variability, are not well resolved.

A relatively simple potential cause of variation in V_M estimates is scale. HOULE *et al.* (1996) showed that life history traits have lower mutational heritability ($h_M^2 = V_M / V_E$, where V_E is the environmental variance) than morphological traits, but the reverse pattern holds when comparing coefficients of mutational variance (CV_M; $100 \times \sqrt{V_M}/\bar{X}$, where $\bar{X}$ is the trait mean). While the pattern is reversed on the different scales, the variation in the magnitude of mutational variance between trait types nonetheless persists. Life-history traits have been hypothesised to be affected by more loci than morphological traits, with this larger target of mutation resulting in a higher mutation rate and greater mutational variance (HOULE 1991; HOULE 1992; HOULE *et al.* 1996; HOULE 1998; MERILÄ AND SHELDON 1999). However, the exact role mutational target size plays in generating

variability of V_M is difficult to assess, as a full catalogue of causal loci is unknown for most

quantitative traits (BARTON AND KEIGHTLEY 2002; MACKAY *et al.* 2009; YANG *et al.* 2010;

ROCKMAN 2012). Indeed, emerging evidence that diverse traits are all highly polygenic (YANG *et al.*

2010; BOYLE *et al.* 2017) suggests that differences in the distribution of mutational effect sizes

(SIMONS *et al.* 2018), rather than simply in number of contributing loci, might cause heterogeneity in

estimates of V_M.

Mutation rate is known to vary widely among taxa (reviewed in KATJU AND BERGTHORSSON

2019), which may reflect differences in ploidy levels, genome size, generation time, and/or effective

population size (LYNCH *et al.* 1999; LYNCH 2010; SUNG *et al.* 2012). Marked variation in mutation

rate has also been observed within species, both among replicated MA experiments (i.e., different

founder genotypes: NESS *et al.* 2012; SUNG *et al.* 2012; SCHRIDER *et al.* 2013) and among lines

within a single MA panel (HUANG *et al.* 2016). This heterogeneity may explain observed variation in

V_M estimates for the same trait in the same taxon. For example, h_M^2 of body size in *Caenorhabditis*

elegans ranged from 0.0035 to 0.0076 between two strains (genetic isolates) (OSTROW *et al.* 2007),

and from 0.0011 to 0.0044 among four sets of MA lines established from the same strain (AZEVEDO

et al. 2002; ESTES *et al.* 2005; OSTROW *et al.* 2007). Different combinations of rate and effect size

have been inferred in different experiments despite highly similar h_M^2 values (Table 1 in HALLIGAN

AND KEIGHTLEY 2009), again suggesting the need to disentangle the complex relationships between

allelic effect sizes and genomic mutation rates to understand the causes of the observed heterogeneity

in V_M.

Ultimately, the random nature of mutation-drift processes themselves might contribute to

variability of estimates, where it is the distribution of estimates obtained from multiple experiments

that is most informative of the expected input of mutational variance. The extent to which

experimental error or biologically relevant sampling variance contribute to variability in estimates

might be revealed by further consideration of the MA experimental design. Firstly, sampling the

mutation-drift process using a finite number of lines will yield variation in estimates, representing

variation in the realisation of the mutation-drift process. LYNCH (1988a) recommended that the

number of replicate lines (L) should be greater than 400 for selfing and 300 for full-sib lines (the two

most common modes in bottlenecking in MA designs) to ensure less than 10% error in the estimate

of among-line variance. However, maintaining a large number of lines simultaneously is often

logistically prohibitive, and this recommendation is rarely met, with experiments typically

considering far fewer (< 50) lines (see Methods).

Secondly, the timeframe over which mutations accumulate might influence estimates of

mutational variance. If the base population was homozygous (heterozygous) when MA lines were

established, estimates of V_M will be downwardly (upwardly) biased before $6N_e$ generations (LYNCH

AND HILL 1986). However, V_M is typically estimated after more than $6N_e$ generations (see Methods),

suggesting little potential for bias due to variation in ancestral segregating variation. Conversely,

long-running MA experiments might under-estimate mutational variance, as the cumulative effect of

low fitness alleles can cause line extinction, or within-line selection against further accumulation

(LYNCH et al. 1999; ESTES et al. 2004; MCGUIGAN AND BLOWS 2013). While a decline in the

magnitude of V_M over time has been observed (MACKAY et al. 1995), other studies have reported

consistent estimates through time (GARCÍA-DORADO et al. 2000). Apart from consistent temporal

trends, transient within-line segregation of mutations might generate variability in V_M estimates from

generation to generation (LYNCH AND HILL 1986).

Environmental context within which MA lines are assayed could also contribute variation to

estimates of V_M. Several studies have considered the effect of macro-environmental variation (i.e.,

known, replicable, experimenter-imposed, variation), including temperature (WAYNE AND MACKAY

1998), light (KAVANAUGH AND SHAW 2005), density (FRY et al. 1996) or food availability (LATTA et

al. 2015). Although the magnitude of mutational variance often varies among environments, there is

only weak evidence for predicable patterns, such as novel or stressful conditions increasing the

magnitude of variance (KONDRASHOV AND HOULE 1994; MARTIN AND LENORMAND 2006). Even in

carefully controlled laboratory experiments, micro-environmental conditions (due to uncontrolled,

un-replicated, and typically unknown sources) contribute to phenotypic variation (LYNCH 1988a;

FALCONER AND MACKAY 1996). If all MA lines respond similarly, micro-environmental variation

will not affect among-line variance; if lines differ in their response, then this genotype by micro-

environment variance could contribute undetected variability to estimates of V_M (LYNCH 1988a;

GARCÍA-DORADO *et al.* 2000). The magnitude of inter-generation variation of among-line estimates

has rarely been assessed (GARCÍA-DORADO *et al.* 2000) and the relative contributions to any such

variation from micro-environment, measurement error and segregation remain unknown.

Here, we combined two approaches to investigate causes of variability in estimates of

mutational variance. Given that it has been over twenty years since causes of variability in mutational

variance have been broadly investigated (HOULE *et al.* 1996; HOULE 1998; LYNCH *et al.* 1999), we

first conducted a meta-analysis to update tests of the previously implicated causal factors of taxon

(LYNCH AND WALSH 1998; LYNCH *et al.* 1999; HALLIGAN AND KEIGHTLEY 2009), trait type (HOULE

et al. 1996; HOULE 1998) and experimental duration (generation number) (LYNCH AND HILL 1986;

MACKAY *et al.* 1995). Multiple estimates of V_M were available for different values of these factors,

allowing us to test the hypothesised effects, although confounding across the factors resulted in little

insight into how these factors affect estimates of V_M. Second, we conducted a new empirical

experiment in *Drosophila serrata* to disentangle the effects of transient segregation, micro-

environment and sampling error, overcoming the limitation on meta-analysis of too few studies of the

same taxon, trait and macro-environment to address these potential causes of estimate variability. Our

results suggest that sampling error is likely to contribute markedly to heterogeneity of published

estimates, but that repeated measures can mitigate these effects.

METHODS

Meta-analysis of empirical estimates of mutational variance

Literature search

We extracted all papers from seven articles that have assembled and reviewed empirical estimates of mutational variance published between 1942 and 2014: LYNCH (1988b); KEIGHTLEY *et al.* (1993); HOULE *et al.* (1996); LYNCH AND WALSH (1998); LYNCH *et al.* (1999); HALLIGAN AND KEIGHTLEY (2009); WALSH AND LYNCH (2018). We then searched the Web of Science database on the 11/12/2019 at 4:38 PM AEST for *journal article* document types meeting the topic criteria of "mutation accumulation" and (varia* or "mutat* coefficient*") and published between 1998 and 2019. This time range overlapped HALLIGAN AND KEIGHTLEY (2009) (which considered fitness traits only) and WALSH AND LYNCH (2018) (which only gives a brief in-text update on their 1998 review), allowing us to capture papers potentially missed in previous reviews.

Further details on the papers identified and preliminary handling steps can be found in Figure S1. From the 473 unique papers identified, we screened titles and abstracts, then the full text, for relevance, using four strict criteria: 1) quantitative mutational variance was reported; 2) estimates were from spontaneous mutation accumulation experiments; 3) estimates were made under the same macro-environmental conditions as mutations were accumulated and; 4) estimates were not re-reporting of previously published estimates. We excluded six studies that analysed transcriptomic data due to the extremely large number of traits relative to other trait categories. We also excluded mitotic cell division traits, examined in only one study (FARHADIFAR *et al.* 2016), cases where V_M estimates were not reported on either a heritability or coefficient of variance scale nor the information to calculate these (i.e., the environmental variance or trait mean) provided and, finally, studies where

the number of mutation accumulation generations was not reported. After applying these criteria, 69

papers were retained.

Meta-analysis data collection

For each of the 69 papers, mutational parameter estimates were extracted, associated with the identity

of the taxon and trait (Figure 1), number of generations of mutation accumulation, and the number of

lines sampled (Figure S2). Where studies reported estimates from multiple environments, only

estimates from the MA maintenance environment were extracted. Papers reported estimates of

mutational variance for a median of three traits (range: 1–29). Where possible, we extracted (or

calculated from provided information) both the coefficient of variance (CV_M; $100 \times \sqrt{V_M}/\bar{X}$, where

$\bar{X}$ is the trait mean) and mutational heritability (h_M^2; V_M/V_E) for each trait. As detailed in Table S1,

after further inspection we excluded extreme estimates, and those reported with very low precision

(Table S1.B).

Before proceeding with the meta-analyses, we transformed estimates to improve normality,

applying a $\log_{10}$ transformation to h_M^2 estimates and a Box-Cox transformation (BOX AND COX 1964)

to CV_M estimates. For the Box-Cox transformation, using the MASS (VENABLES AND RIPLEY 2002)

package in R and maximum-likelihood, $\lambda = 0.2$ was determined as the best-fit power term for these

data. Both the number of generations and the number of lines sampled were log-transformed to a

normal distribution, and an extreme outlier (> 8 SD) excluded (Table S1B).

Predictor variables for the meta-analysis

Estimates came from 22 species, and based on the distribution of estimates, we defined six taxon

categories: Drosophila (*Drosophila melanogaster* [$n = 89$] and *D. serrata* [$n = 15$]), plant

(*Arabidopsis thaliana* [five studies, $n = 15$]; cumulatively three studies and $n = 15$ from *Amsinckia*

douglasiana, *A. gloriosa*, *Hordeum vulgare* and *Zea mays*), microbe (*Cryptococcus neoformans*,

Escherichia coli, Chlamydomonas reinhardtii, Saccharomyces cerevisiae, Cystovirus pseudomonas

virus φ6 *and Vesicular stomatitis virus*), nematode (*Caenorhabditis briggsae, C. brenneri, C.*

elegans, C. remanei and *Oscheius myriophila*), and Daphnia (*Daphnia pulex*) (Figure 1A). We

excluded from analyses estimates from two other insects (*Daktulosphaira vitifoliae* and *Nasonia*

vitripennis) for clarity of inference (insect versus *Drosophila*), and from *Mus musculus*, the only

vertebrate, due to low sample size (four estimates: Figure 1A).

HOULE *et al.* (1996; 1998) contrasted life-history, morphology and growth traits, which were

expected to differ in mutational target size and sensitivity to micro-environmental variation. We

differ from HOULE *et al.* (1996; 1998) in considering weight and height of juveniles as morphological

(not growth) traits. Within the morphology category (20% of all estimates), 76% of estimates were

for size-related traits. Reflecting more recent publications, we defined a physiology category (28% of

all estimates), which included locomotive, enzymatic and metabolic activity traits, which may differ

from life-history or morphological traits in mutational target size or micro-environmental sensitivity.

Life history traits accounted for 53% of estimates, prompting us to subdivide life history traits into

relatively evenly represented narrower sub-categories: total fitness (either measured directly, or

reported based on combined data from individual components), survival, productivity, development

time and a miscellaneous category (capturing traits such as phenology, longevity and mating success,

which were individually less well represented) (Figure 1B; Table S2).

Meta-analyses of mutational variance estimates

Our final meta-analysis data set consisted of 201 estimates of h_M^2 and 211 estimates of CV_M from 62

studies. We implemented the following model (separately for h_M^2 and CV_M data) via PROC MIXED

in SAS v.9.4 (SAS Institute Inc., Cary, NC.), using maximum likelihood:

$$y_{ijklmn} = \mu + tx_i + trt_j + g_k + l_l + s_m + \varepsilon_{ijklmn} \qquad (1)$$

where y is the vector of published estimates, and μ was the overall mean h_M^2 or CV_M. We fit the

categorical predictors taxon (*tx, i* =1, 2, ..., 5) and trait (*trt, j*=1, 2, ..., 7), and the continuous

predictors generation number (*g*) and number of lines sampled (*l*) as fixed effects. The number of

MA lines and generations spanned different (non-overlapping) ranges among taxa (Figure S2), and

interactions were therefore not fit. Study (*s*) was fit as a random effect, accounting for non-

independence among estimates within a paper. The same MA lines were sometimes analysed in

multiple studies, for different traits, or after further generations of accumulation (Table S3).

However, an additional random effect of shared lines (with study nested within this) did not account

for any variance, potentially a consequence of the unbalanced design, with only some studies sharing

lines. The residual error (ε) captured all unexplained variation among estimates. We used the

Satterthwaite approximation to correct the denominator degrees of freedom.

Variation in estimates of among-line variance within the same taxon, trait type and MA

duration: An experiment in *Drosophila serrata*

Quantitative genetic parameters are associated with large sampling errors (KLEIN *et al.* 1973; KLEIN

1974; LYNCH AND WALSH 1998), and the relatively low signal (i.e., few genetic differences) among

MA lines will make them particularly vulnerable to "noisy" estimation. We conducted a new

empirical experiment to quantify the contribution of this estimation error relative to the effects of

other sources of random variation, particularly the chance sampling of different genotypes within

inbred lines (i.e., segregating mutations in the process of becoming fixed or lost), and micro-

environmental variation due to the specific time and place of sampling. We conducted the experiment

in *Drosophila serrata*, a member of the *montium* species group, endemic to Australia and Papua New

Guinea, that has been extensively used in quantitative genetic research, including study of mutational

variance (e.g., MCGUIGAN AND BLOWS 2013; MCGUIGAN *et al.* 2014; HINE *et al.* 2018; DUGAND *et*

al. 2021).

 Experimental design

A panel of 200 MA lines were founded from one of the *D. serrata* Reference Genome Panel

(DsRGP) lines described in REDDIEX *et al.* (2018). The MA lines were each maintained by brother-

sister inbreeding for 20 generations, following husbandry protocols previously established by

MCGUIGAN *et al.* (2011) to minimise extinction and within-line selection. Given the relatively short

duration of mutation accumulation, among-line variation may be inflated by segregating variation in

the base population (LYNCH AND HILL 1986). Notably, genome-wide heterozygosity was very low

(0.3%) in the DsRGP line that founded the MA lines, and among-line variance for wing phenotypes

not statistically supported in the first generation of mutation accumulation (S. Chenoweth, pers.

comm.). This panel of lines are therefore representative of the relatively low among-line variance

characteristic of typical MA lines, and are well-suited to investigating the relative potential of

estimation error versus other factors to generate variability in estimates of mutational variance.

We randomly chose 42 of the MA lines for further experimentation to partition putative

causes of variation in estimates. Our experimental design had three main goals. Firstly, we aimed to

retard ongoing mutation accumulation to maintain the low among-line variance, and achieved this by

increased population size within each of the lines. The fate of a new mutation is determined by drift

when its fitness effect is less than $1/2N_e$ (WRIGHT 1931; KIMURA 1983). Empirical evidence suggests

that population sizes as low as ten may be sufficient to prevent fixation of mutations with deleterious

fitness effects (ESTES *et al.* 2004; KATJU *et al.* 2015; LUIJCKX *et al.* 2018). We imposed a minimum

population size of 12 males and 12 females (Figure 2A). This change in census population size

complicates calculation of a per-generation rate of mutational increase in phenotypic variance; here,

we instead focus on the among-line variance, V_L, the parameter estimated for calculating V_M (LYNCH

AND WALSH 1998).

253   The second aim of our experimental design was to vary the opportunity for mutation-

254 selection-drift operating within a line, and hence, the dynamics of segregating genetic variation; we

255 achieved this by imposing two, substantially different, population sizes on sublines of each of the 42

256 MA lines ($N = 24$ versus 288, referred to hereafter as small, S, and large, L, population size

257 treatments: Figure 2A). These treatments contrast deterministic evolution of mutations with relatively

258 strong ($s > 0.029$: $N_e \sim 17$), versus weak ($s > 0.003$: $N_e \sim 202$) fitness effects (effective population

259 size estimated as 70% of census size, CROW AND MORTON 1955), and thus a different opportunity of

260 within-line variance to increase during the six generations, and for this segregating variance to

261 influence estimates of among-line variance.

262   The final aim of the experimental design was to determine how variability in the effects of

263 new mutations under different random micro-environments impact among-line variance estimates.

264 Environmental variation contributes to phenotypes even under tightly controlled laboratory

265 conditions (LANDE 1976; FALCONER AND MACKAY 1996); we randomly sampled the environmental

266 conditions present within our laboratory by repeatedly sampling the lines (genotypes) over six

267 consecutive generations. Thus, our experiment consisted of applying two population size treatments

268 (S, L) to each of 42 lines (derived from a classical MA, with low among-line variation), where these

269 84 lines were maintained under the same micro-environment conditions (12 flies per sex per vial

270 founding each generation, with S and L differing in the number of vials) for six generations (Figure

271 2A).

272 *Data collection*

273 Each generation, 12 males per subline (six per rearing vial) were randomly sampled for wing

274 phenotypes (Figure 2). Wings were mounted on microscope slides and photographed using a Leica

275 MZ6 microscope camera and the software LAS EZ v2.0.0 (Leica Microsystems Ltd, Switzerland). A

276 total of 5,135 wings were landmarked for nine positions, defined by wing vein and margin

intersections (Figure 2B), using the software tpsDIG2 (ROHLF 2013). The number of wings were

evenly distributed across treatments (2,583 in S and 2,552 in L) and generations (~8.3% per

generation, per treatment). Landmarks were aligned using a General Procrustes fit in tpsRelw (ROHLF

2007). Centroid size (CS), the square root of the sum of squared deviations of the coordinates from

the centroid (ROHLF 1999), was recorded as a metric of wing size. The aligned X-Y coordinates for

each landmark were then used to calculate ten inter-landmark distances (ILDs) (Figure 2B). ILDs

scores were re-scaled prior to analysis (multiplied by 100) to aid model convergence. Outliers >3.0

SD from the mean were removed for each of the 11 traits (10 ILD and size) (329 measures across the

56,485 total measures).

Analyses of trait means

Micro-environmental variation can generate both temporal trends and idiosyncratic changes among

generations (LANDE 1976; FALCONER AND MACKAY 1996), but here is expected to have consistent

effects on the two synchronously reared population size treatments. In contrast, genetic divergence,

due to independent mutation and subsequent drift or selection within each of the 84 subpopulations

could lead to increasing divergence between the treatment means if mutational effects on either the

measured traits, or pleiotropic effects on fitness are biased toward a higher or lower value (JOHNSON

AND BARTON 2005; MCGUIGAN AND BLOWS 2013). To test the null hypothesis that wing trait mean

remained constant across generations and treatments we implemented the following model using

PROC MIXED in SAS v9.4 (SAS Institute Inc., Cary, NC.), within a maximum-likelihood

framework:

$$y_{ijklm} = \mu + G_i + T_j + GT_{ij} + Line_{kj} + Vial_{l(ijk)} + \varepsilon_{ijklm} \tag{2}$$

where y_{ijkm} was the trait value for the *m*th wing, from the *l*th vial, within the *k*th line, for the *j*th

treatment in the *i*th generation. The model fits the overall mean (μ) and the fixed effects of generation

(G), treatment (T), and their interaction (GT), as well as random effects due to MA line (Line),

replicate rearing vials (Vial) within each line, and residual error (ε), capturing unexplained variance

among individual flies within a rearing vial. Each line was sampled in all generations and we

accounted for this repeated measures design, fitting a reduced-rank factor analytic covariance

structure (specifying TYPE=FA(1)), which estimated generation-specific among-line variance (a

diagonal matrix) and one dimension of among-generation covariance (i.e., among-line variance

shared across repeated measures of generations) (MEYER 2009; MCGUIGAN AND BLOWS 2010).

Preliminary analyses indicated that this FA(1) model captured all among-line variance. Vial and

residual variances were estimated for each generation and treatment, accounting for potential

heterogeneity in micro-environmental effects, and segregating variation.

Analyses of among-line variance

To investigate the effects of evolutionary processes (captured by our two population size treatments)

and micro-environment (captured by the six generations) on estimates of among-line variance, we

first treated data from each generation and treatment as independent experimental estimates for each

trait, allowing us to characterise the variability in estimates (12 estimates per trait, 132 estimates in

total). This was done by reducing model 2 to:

$$y_{klm} = \mu + Line_k + Vial_{l(k)} + \varepsilon_{klm} \tag{3}$$

fit using restricted-maximum likelihood (REML) in PROC MIXED in SAS v9.4 (SAS Institute Inc.,

Cary, NC.). We used REML-MVN sampling to place confidence intervals about our estimates,

obtained by sampling from $N \sim (\hat{\theta}, \mathbf{V})$ 10,000 times, using the MASS package (VENABLES AND

RIPLEY 2002) in R [v. 3.6.1], where $\hat{\theta}$ was the vector of observed covariance parameter estimates,

and $\mathbf{V}$ was the asymptotic variance-covariance matrix, the inverse of the Fisher information matrix,

$I(\hat{\theta})^{-1}$ (MEYER AND HOULE 2013; HOULE AND MEYER 2015; SZTEPANACZ AND BLOWS 2017). G-scale

REML-MVN samples, which are not constrained to the parameter space (i.e., can be negative), allow

inference of statistical support for variance when the lower 5% CI does not encompass zero (a one-

tailed test); for individual variables, this approach provides the same conclusions as a Log Likelihood

Ratio test (LRT) (DUGAND *et al.* 2021), but has the advantage of also providing a robust visualisation

of the confidence in estimate values. We are interested here only in general patterns of variability

among estimates and do not correct for multiple testing.

Next, to formally test whether among-line variance differed among generations we again

modified model 2, and used PROC MIXED and REML to fit a model with the intercept (μ),

generation (G), Line, Vial and error, and included an additional random effect of the interaction

between line and generation (G × Line), estimating the variation in genetic effects among

generations (GARCÍA-DORADO *et al.* 2000):

$$y_{iklm} = \mu + G_i + Line_k + (G \times Line)_{ik} + Vial_{l(ik)} + \varepsilon_{iklm} \qquad (4)$$

This model was applied to each trait and population size treatment separately (22 models), with vial

and residual variances estimated separately for each generation. Statistical support for G × Line and

for Line was determined using log-likelihood ratio tests (LRT, 0.5 d.f.: SELF AND LIANG 1987;

LITTELL *et al.* 2006) to compare model 4 to a reduced models that did not fit G x Line, or did not fit

Line. We applied the Benjamini-Hochberg method (BENJAMINI AND HOCHBERG 1995) to correct for

multiple hypothesis testing (within each random effect), using a conservative 5% false discovery rate

(FDR).

To formally test the hypothesis that differences in the opportunity for mutation-drift-selection

between the Large and Small population size treatments affected the estimates of among-line

variance, or the among-line by generation variance, we ran the following model for each of the 11

traits, again using PROC MIXED and REML, where we fit the intercept (μ), generation (G),

treatment (T) and the interaction between generation and treatment (GT), as well as the random

effects of Line, G × Line, Vial and error:

$$y_{ijklm} = \mu + G_i + T_j + GT_{ij} + Line_k + (G \times Line)_{ik} + Vial_{l(ijk)} + \varepsilon_{ijklm} \qquad (5)$$

Although estimates of among-line by generation variance were typically not statistically

distinguishable from zero (see Results), we nonetheless modelled it here to ensure that small,

statistically undetectable, variation did not inflate the among-line variance of interest. Generation and

treatment-specific vial and residual variances were estimated. We used LRTs to determine whether

estimating among-line variance separately for the two population size treatments (modelled using a

Group statement in PROC MIXED) improved model fit over modelling a single, pooled, estimate,

again applying the Benjamini-Hochberg method (BENJAMINI AND HOCHBERG 1995) to allow a 5%

FDR. For the one trait with statistically supported G × Line variance in at least one population size

treatment (see Results), we again used a LRT to test the hypothesis that Large and Small population

treatments differed in the magnitude of this G × Line variance; a 5% FDR correction was applied

assuming 11 traits were tested.

Finally, to infer whether the magnitude of among-line (V_L) variance differed among the 10

different wing shape traits, we took the V_L estimates from model (5), estimated from the data from

both treatments, and obtained REML-MVN confidence intervals as described above. We regressed

the among-line variance on phenotypic variance (V_P: sum of among-line, among-vial and residual)

for the REML estimate and each of the 10,000 REML-MVNA samples. Similarly, we regressed the

REML V_L estimates on the trait mean (estimated over all generations and treatments), and the REML-

MVN samples of V_L on 10,000 samples of the trait mean taken from a MVN distribution with the

observed mean and standard deviation, sampled using the MASS package (VENABLES AND RIPLEY

2002). Where 95% CI of the regression slope did not include zero, we concluded the parameter (V_P or

mean) significantly predicted V_L.

RESULTS

Meta-analysis of published mutational variance estimates

Unexpectedly, we observed mutational heritability to have a similar magnitude across all trait types

($F_{6,161} = 2.01$, $P = 0.0670$; Figure 3A). In contrast, CV_M significantly varied among trait types ($F_{6,151}$

$= 15.89$, $P < 0.0001$), following the predicted pattern of morphological traits having smaller

magnitude CV_M than life history traits (Figure 3B). The number of lines sampled did not affect the

magnitude of mutational variance (h_M^2: $F_{1,68.6} = 0.83$, $P = 0.3658$; CV_M: $F_{1,70.6} = 0.27$, $P = 0.6076$).

Taxon significantly predicted the magnitude of mutational variance, on both scales (h_M^2: $F_{4,54.5} =$

3.12, $P = 0.0221$; CV_M: $F_{4,56.5} = 6.13$, $P = 0.0004$), with plants having the lowest model estimates of

mutational variance and Nematodes the highest (Figure 4). The magnitude of mutational variance

significantly decreased with increasing number of generations of mutation accumulation (h_M^2: $F_{1,58.7}$

$= 12.06$, $P = 0.0010$, $\beta = -0.622$; CV_M: $F_{1,65.6} = 21.05$, $P < 0.0001$, $\beta = -1.065$). However, as plants

and nematodes differed markedly in generation number (median of 17 and 250 generations,

respectively; Figure S2A), the effect of taxon versus generation number on the magnitude of V_M

cannot be clearly inferred. To further investigate this, we analysed V_M separately for each of the two

best represented taxonomic groups, Drosophila (19 – 300 generations; Figure S2A) and Nematode

(50 – 370 generations; Figure S2A). In Drosophila, there was a significant negative relationship

between number of generations and the magnitude of mutational variance (h_M^2: $F_{1,32} = 7.76$, $P =$

0.0089, $\beta = -0.486$, $N = 56$; CV_M: $F_{1,20.1} = 28.05$, $P < 0.0001$, $\beta = -1.299$, $N = 73$). In contrast,

 generation number did not explain the magnitude of mutational variance in Nematodes (h^2_M: $F_{1,14.1} =$

1.38, $P = 0.2600$, $\beta = -0.337$, $N = 86$; CV_M: $F_{1,18.9} < 0.00$, $P = 0.9708$, $\beta = -0.023$, $N = 76$).

Variation in estimates of among-line variance within the same taxon, trait type and duration: An experiment in *Drosophila serrata*

While there was statistically significant generation by treatment effects on most *D. serrata* wing traits, these appeared as idiosyncratic changes in direction and magnitude of differences between population size treatments (Figure 5), more consistent with effects of micro-environmental fluctuations than evolving phenotypic differentiation. One trait differed in mean between the treatments (ILD4.8: $F_{1, 711} = 25.93$, $P = 0.0001$), but again there was no evidence of increasing magnitude of divergence through time (Figure 5), expected under evolution.

Next, we investigated the variation in among-line variance, treating our data as representing 12 different "experiments" for each of our 11 traits. There was inconsistent statistical evidence for among-line variance within each trait, where 60 out of 132 estimates (45.45%) were not significantly different from zero (Figure 6). We could draw very different conclusions about the magnitude of among-line variance, depending on which "experiment" we had conducted (Figure 6). For example, among-line variance was estimated to be >2 times higher in the first than third generation in the analyses of wing (centroid) size (CS) in the large population size treatment (3.43×10^{-4} and 1.38×10^{-4}, respectively; Figure 6). Marked fluctuations in the magnitude of among-line variance persisted when accounting for heterogeneity among generations in the magnitude of phenotypic variance or trait mean (Supplementary Figures S3 and S4, respectively).

Among-line variance did not exhibit a directional trend over time for any trait (Figure 6), and variation in mutational effects among generations (among-line by generation variance: model 4) was statistically supported for only four of the 22 estimates (two treatments by eleven traits), with only two estimates remaining significant at a 5% FDR (LRT for CS in L: $\chi^2_1 = 13.9$, $P = 0.0001$; LRT for

ILD2.5 in L: $\chi_1^2 = 11.1$, $P = 0.0005$; Figure 7A). Consistent with the largely overlapping CI of the

two treatments (Figure 6), a difference between the Small and Large population size treatments in

among-line by generation variance was detected for only one trait (centroid size), and this did not

remain significant at the 5% FDR (Table S4B). Furthermore, there was no statistical support for an

effect of the population size treatment on the among-line variance for any trait (Figure 7B; Table

S4A). The variability in estimates (Figure 6) was thus typically within the magnitude of the error

associated with each estimate.

421   Given these results, we utilised all data (six generations for two treatments) to obtain a single,

statistically supported, estimate of among-line variance for each trait (Figure 8A). Utilising these

robust estimates (and excluding size), we found that the total phenotypic variance ($\beta = 0.075$, 95%

CI: 0.050, 0.106), or the magnitude of the trait mean ($\beta = 1.689 \times 10^{-6}$, 95% CI: 0.406×10^{-6}, $3.026 \times$

10^{-6}) significantly predicted the observed magnitude of among-line variance (Figure 8B).

Nonetheless, substantial heterogeneity in estimates remained unexplained (Figure 8B; for V_L on V_P,

$R^2 = 0.286$ [95% CI: 0.141-0.477]; for V_L on mean, $R^2 = 0.029$ [95% CI: 0.002-0.076]), suggesting

either inherent stochasticity of the mutation process, or differences in genetic architecture of the traits

affected estimates of among-line variance.

DISCUSSION

Although numerous estimates of mutational variance (V_M) have been published, it remains unclear

why they vary in magnitude. Our meta-analytic investigation revealed substantial confoundment

between potential causal factors, suggesting that further empirical data must be collected to fully

resolve the contribution of these general factors to variation in V_M. Mutation is an inherently random

process, but the expected variance about the true V_M due to stochastic sampling of newly segregating

mutations, from micro-environment specific effects of mutations, or from estimation error, is not

known. Analyses of data from a manipulative experiment in *D. serrata* suggests that estimation error

may contribute most generally to variation in estimates. Below we discuss the specific outcomes and

limitations of both approaches, and the implications our analyses have for future work characterising

mutational input to quantitative genetic variation.

Effects of taxon, trait type and mutation accumulation duration on the magnitude of mutational

variance

Among eight multicellular eukaryotes, the reported per nucleotide mutation rate is highest for

Arabidopsis and lowest for the nematode *C. briggsae* (KATJU AND BERGTHORSSON 2019). Consistent

with this, we observed a significant effect on both CV_M and h_M^2 of taxon category, with plant

(predominantly estimates from *Arabidopsis*) and Nematode (predominantly estimates from

Caenorhabditis species) differing most in the magnitude of estimates. However, experiments in

plants had the shortest duration of mutation accumulation (Figure S2A), where our analyses also

supported a significant decline in V_M as the number of MA generations increased. Thus, the apparent

elevation of V_M for plants could be a consequence of short duration, or the apparent decline in V_M

over time could be an artifact, (partly) caused by the high V_M in plants. We observed a significant

decline in the magnitude of V_M with number of MA generations in Drosophila, but not in Nematodes.

Decline in V_M over time has been reported previously in *D. melanogaster,* but only for some assayed

traits (MACKAY *et al.* 1995; GARCÍA-DORADO *et al.* 2000). Therefore, while the data suggest that

taxon and MA duration could account for some variation in the magnitude of V_M, further experiments

are required to decouple their effects. Advances in accessibility of genome data for a broad range of

taxa provide substantial scope for such experiments to explicitly estimate relevant genomic

parameters (e.g., frequency spectra for different types of mutations) alongside the phenotypic

variation generated by those mutations (i.e., V_M) (KATJU AND BERGTHORSSON 2019). Furthermore,

given evidence that epigenetic mutations arise more frequently than genetic mutations (e.g., VAN DER

GRAAF et al. 2015) the potential contribution of epimutation to patterns of heterogeneity of V_M

should be explicitly assessed in future studies.

HOULE et al. (1996); (see also LYNCH AND WALSH 1998; LYNCH et al. 1999) strongly

concluded that, relative to morphological traits, life history traits had lower h^2_M but higher CV_M. Our

results were consistent with this for CV_M, but not for h^2_M, where estimates did not significantly vary

among trait types. Notably, 76% of estimates in the morphology category for the meta-analysis were

the size of either part of (e.g., wing length; SANTIAGO et al. 1992) or the whole organism (e.g., plant

height; RUSSELL et al. 1963), reflecting bias in the traits studied. In contrast to other aspects of

morphology, size may share the same heightened sensitivity to environmental variance (and hence

depressed h^2_M) expected for life-history traits (SANTOS et al. 1994; HOULE et al. 1996). Size might

also be more similar to fitness traits in its underlying genetic complexity (and hence, mutational

target size; HOULE 1998). Investigation of diverse morphological traits, in conjunction with

assessment of environmental sensitivities, will help inform where the magnitude of h^2_M for size-

related traits lies in relationship to fitness versus morphological traits.

Notably, life-history categories varied in h^2_M and CV_M (Figure 3), which may reflect

differences in the genomic basis of different life history traits. Alternatively, the MA design may

allow less effective relaxation of selection on some fitness traits (e.g., survival) than others (e.g.,

productivity). Within the D. serrata experiment, variation among wing-shape traits in the magnitude

of among-line variance was not fully accounted for by differences in the magnitude of phenotypic

variance or trait mean (Figure 8). Establishing whether these differences are informative of the

inherent genetic architecture, or are a manifestation of the stochastic nature of mutation, requires

repeating the estimation using either the same or a different genetic background to determine if

consistent differences among traits persist. Again, combining phenotypic investigation with fine-

scale genomic analyses of MA lines could lead to further insights on why traits differ in mutational

variance (KATJU AND BERGTHORSSON 2019).

The contributions of mutation-drift-selection processes, micro-environmental variation, and

estimation error to variation in the magnitude of mutational variance

Individual point estimates of among-line variance, and thus of V_M, might deviate from theoretical

expectations due to factors including transient segregation of mutations, and interactions with micro-

environmental variation, as well as simple estimation error (LYNCH AND HILL 1986; LYNCH 1991).

Here, we observed substantial heterogeneity in the magnitude of, and statistical support for, estimates

of among-line variance of 11 wing traits measured in *D. serrata* (Figure 6). In the only other study to

investigate short-term (over two or three generations) variability of estimates, GARCÍA-DORADO *et al.*

(2000) found that V_M varied across consecutive generations for one (sternopleural bristle count) of

four traits investigated (no temporal variation in estimates of egg-to-adult viability, wing length or

abdominal bristle count). GARCÍA-DORADO *et al.* (2000) attributed the variation to (mutant)

genotype-by-environment (GxE) interactions, specifically to micro-environmental variation in

rearing conditions altering the phenotypic effects of mutations from generation to generation. We

similarly found statistical support for variation in mutational effects among generations for few (two

of 11) traits. Notably, wing (centroid) size exhibited significant generation by among-line variance;

as discussed above, size might be more dependent on micro-environmental variation in resources or

energy requirements than wing shape traits (CAVICCHI *et al.* 1985; BITNER-MATHÉ AND KLACZKO

1999). Thus, our results and those of GARCÍA-DORADO *et al.* (2000) suggest that changes in

mutational effects with micro-environment may contribute to heterogeneity among point estimates of

some traits, which may reflect differences in trait environmental sensitivity, or potentially covariation

of environmental sensitivity and mutational effect size (LYNCH *et al.* 1999; GARCÍA-DORADO *et al.*

2000). Further data on whether phenotypic traits differ in the frequency or effect size of epi-

mutations may also help to inform how and why traits differ in temporal heterogeneity or micro-

environmental effects on among-line variance.

The mutation-drift process itself can also cause heterogeneity in among-line variance due to

chance sampling of a newly-arisen mutation, segregating within lines (LYNCH AND HILL 1986).

Transient segregation of mutation could contribute to estimation error (increasing within relative to

among-line variance), as well as causing bounce in estimates if a line's breeding value is extreme due

to the presence of a segregating variant that is later lost (or not sampled in subsequent phenotype

assays). In the current study, statistical analyses did not support any effect of the population size

treatments on estimates of among-line variance. Given the different opportunity for mutation-drift-

selection afforded by the ~order of magnitude difference in census size in these paired sets of lines,

this suggests that segregating variation may have little effect on variability of among-line variance

estimates. Notably, populations as small as five might be sufficiently large for selection to prevent

declines in fitness due to new mutations (ESTES *et al.* 2004; KATJU *et al.* 2015; LUIJCKX *et al.* 2018).

Here, pleiotropic effects of mutations on wing traits and fitness (MCGUIGAN AND BLOWS

2013) may have resulted in little difference between Small and Large treatments in the segregating

frequencies of new mutation, or consequently within-line variance. Inter-generational changes in the

phenotype of a line, contributing to temporal heterogeneity in among-line variance, will depend on

dominance coefficients of mutations. Mutations influencing fitness may typically be at least partially

recessive, with the dominance coefficient positively correlated with the selection coefficient

(AGRAWAL AND WHITLOCK 2011). In contrast, the pleiotropic effects of these mutations on wing

traits (which may be typical of other non-fitness phenotypes) appear to typically be additive (HOULE

AND FIERST 2013; DUGAND *et al.* 2021). Thus, our focus on morphological traits may have under-

estimated the temporal heterogeneity that transient segregation of mutations may cause.

Rejecting micro-environmental variation and transient segregation of variants as general

explanations of estimate heterogeneity in our data, we conclude that the observed variability is

largely the consequence of estimation error, and suggest that estimation error is likely to contribute to

heterogeneity of published estimates. We aimed to capture typical levels of estimation error by

considering a low, but typical number of MA lines (42; see Supplementary Figure S2B), and

balancing the relatively low among line variance (due to relatively few generations of mutation

accumulation) by considering traits that were expected to have strong genetic signal relative to

residual error (ROFF AND MOUSSEAU 1987; HOULE *et al.* 1996).

The sequential repeated-measures experimental design provided statistical control over the

experimental noise, allowing us to consistently detect statistically significant mutational variance in

all traits (Figure 8A). Several previous studies have analysed repeated measures of MA lines to

estimate V_M as the slope of the regression of among-line variance on generation (VASSILIEVA *et al.*

2000; HOULE AND NUZHDIN 2004; MCGUIGAN *et al.* 2011), or included generation as a (fixed or

random) factor in the model to similarly obtain a single estimate of among-line variance

(MARTORELL *et al.* 1998; SHAW *et al.* 2000; BAER *et al.* 2005). Our analysis particularly highlights

the potential benefits of short-term repeated measures (sequential generations) to improve estimate

precision, and power to detect small effects.

Understanding the contribution that mutations make to evolutionary and genetic phenomena

relies on better estimates of the phenotypic variance generated by new mutation. Our meta-analysis

of empirical estimates of mutational variance was unsuccessful in determining the relative

contribution of trait type, taxon and experimental duration due to confoundment between these

factors, an issue requiring further data to resolve. Our manipulative experiment suggested that

estimation error may contribute substantially to estimate variability, and demonstrated that repeated

measures over few (e.g., sequential) generations provides a simple but effective approach to address

656 this and improve inference. Overall, further empirical studies are needed to fully assess how both

657 general and study specific factors affect V_M estimates, where improved precision and replicability in

658 estimates will consequently advance broader evolutionary questions such as those addressing the

659 maintenance of quantitative genetic variance (BARTON AND TURELLI 1989; JOHNSON AND BARTON

660 2005; WALSH AND LYNCH 2018).

Data Availability

663 All analysed data are available at: https://doi.org/10.6084/m9.figshare.14913051.v2

Acknowledgments

666 We thank Adam Reddiex, Nicholas Appleton, Jack Price and Derek Sun for their help with

667 maintaining, collecting, and digitising the mutation accumulation data. We also thank Stephen F.

668 Chenoweth for providing us with the *Drosophila serrata* MA lines, and Jan Engelstädter for

669 suggesting the combinational approach. This manuscript was improved by the input of two

670 anonymous reviewers and the AE.

References

Agrawal, A. F., and M. C. Whitlock, 2011 Inferences about the distribution of dominance drawn from yeast gene knockout data. Genetics 187**:** 553-566.

Azevedo, R. B., P. D. Keightley, C. Laurén-Määttä, L. L. Vassilieva, M. Lynch *et al.*, 2002 Spontaneous mutational variation for body size in *Caenorhabditis elegans*. Genetics 162**:** 755-765.

Baer, C. F., F. Shaw, C. Steding, M. Baumgartner, A. Hawkins *et al.*, 2005 Comparative evolutionary genetics of spontaneous mutations affecting fitness in rhabditid nematodes. Proceedings of the National Academy of Sciences, USA 102**:** 5785-5790.

Barton, N. H., and P. D. Keightley, 2002 Understanding quantitative genetic variation. Nat Rev Genet 3**:** 11-21.

Barton, N. H., and M. Turelli, 1989 Evolutionary quantitative genetics: how little do we know? Annual Review of Genetics 23**:** 337-370.

Benjamini, Y., and Y. Hochberg, 1995 Controlling the false discovery rate: a practical and powerful approach to multiple testing. Journal of the Royal Statistical Society: Series B (Statistical Methodology) 57**:** 289-300.

Bitner-Mathé, B. C., and L. B. Klaczko, 1999 Plasticity of *Drosophila melanogaster* wing morphology: effects of sex, temperature and density. Genetica 105**:** 203.

Box, G. E., and D. R. Cox, 1964 An analysis of transformations. Journal of the Royal Statistical Society: Series B (Statistical Methodology) 26**:** 211-243.

Boyle, E. A., Y. I. Li and J. K. Pritchard, 2017 An expanded view of complex traits: From polygenic to omnigenic. Cell 169**:** 1177-1186.

Cavicchi, S., D. Guerra, G. Giorgi and C. Pezzoli, 1985 Temperature-related divergence in experimental populations of *Drosophila melanogaster*. I. Genetic and developmental basis of wing size and shape variation. Genetics 109**:** 665-689.

Crow, J. F., and N. E. Morton, 1955 Measurement of gene frequency drift in small populations. Evolution 9**:** 202-214.

Dugand, R. J., J. D. Aguirre, E. Hine, M. W. Blows and K. McGuigan, 2021 The contribution of mutation and selection to multivariate quantitative genetic variance in an outbred population of *Drosophila serrata*. Proceedings of the National Academy of Sciences, USA 118**:** e2026217118.

Estes, S., B. C. Ajie, M. Lynch and P. C. Phillips, 2005 Spontaneous mutational correlations for life-history, morphological and behavioral characters in *Caenorhabditis elegans*. Genetics 170**:** 645-653.

Estes, S., P. C. Phillips, D. R. Denver, W. K. Thomas and M. Lynch, 2004 Mutation accumulation in populations of varying size: The distribution of mutational effects for fitness correlates in *Caenorhabditis elegans*. Genetics 166**:** 1269-1279.

Falconer, D. S., and T. F. C. Mackay, 1996 *Introduction to quantitative genetics*. Longman, Burnt Mill, England.

Farhadifar, R., J. M. Ponciano, E. C. Andersen, D. J. Needleman and C. F. Baer, 2016 Mutation is a sufficient and robust predictor of genetic variation for mitotic spindle traits in *Caenorhabditis elegans*. Genetics 203**:** 1859-1870.

Fry, J. D., S. L. Heinsohn and T. F. Mackay, 1996 The contribution of new mutations to genotype-environment interaction for fitness in *Drosophila melanogaster*. Evolution 50**:** 2316-2327.

García-Dorado, A., J. Fernández and C. López-Fanjul, 2000 Temporal uniformity of the spontaneous mutational variance of quantitative traits in *Drosophila melanogaster*. 75**:** 47-51.

Halligan, D. L., and P. D. Keightley, 2009 Spontaneous mutation accumulation studies in evolutionary genetics. Annual Review of Ecology, Evolution and Systematics 40**:** 151-172.

Hine, E., D. E. Runcie, K. McGuigan and M. W. Blows, 2018 Uneven distribution of mutational variance across the transcriptome of *Drosophila serrata* revealed by high-dimensional analysis of gene expression. Genetics 209**:** 1319-1328.

Houle, D., 1991 Genetic covariance of fitness correlates: what genetic correlations are made of and why it matters. Evolution 45**:** 630-648.

Houle, D., 1992 Comparing evolvability and variability of quantitative traits. Genetics 130**:** 195-204.

Houle, D., 1998 How should we explain variation in the genetic variance of traits? Genetica 102**:** 241-253.

Houle, D., G. H. Bolstad, K. van der Linde and T. F. Hansen, 2017 Mutation predicts 40 million years of fly wing evolution. Nature 548**:** 447-450.

Houle, D., and J. Fierst, 2013 Properties of spontaneous mutational variance and covariance for wing size and shape in *Drosophila melanogaster*. Evolution 67**:** 1116-1130.

Houle, D., and K. Meyer, 2015 Estimating sampling error of evolutionary statistics based on genetic covariance matrices using maximum likelihood. Journal of Evolutionary Biology 28**:** 1542-1549.

Houle, D., B. Morikawa and M. Lynch, 1996 Comparing mutational variabilities. Genetics 143**:** 1467-1483.

Houle, D., and S. V. Nuzhdin, 2004 Mutation accumulation and the effect of *copia* insertions in *Drosophila melanogaster*. Genetical Research 83**:** 7-18.

Huang, W., R. F. Lyman, R. A. Lyman, M. A. Carbone, S. T. Harbison *et al.*, 2016 Spontaneous mutations and the origin and maintenance of quantitative genetic variation. Elife 5**:** e14625.

Johnson, T., and N. Barton, 2005 Theoretical models of selection and mutation on quantitative traits. Philosophical Transactions of the Royal Society of London, Series B: Biological Sciences 360**:** 1411-1425.

Katju, V., and U. Bergthorsson, 2019 Old trade, new tricks: insights into the spontaneous mutation process from the partnering of classical mutation accumulation experiments with high-throughput genomic approaches. Genome Biology and Evolution 11**:** 136-165.

Katju, V., L. B. Packard, L. Bu, P. D. Keightley and U. Bergthorsson, 2015 Fitness decline in spontaneous mutation accumulation lines of *Caenorhabditis elegans* with varying effective population sizes. Evolution 69**:** 104-116.

Kavanaugh, C. M., and R. G. Shaw, 2005 The contribution of spontaneous mutation to variation in environmental responses of *Arabidopsis thaliana*: Responses to light. Evolution 59**:** 266-275.

Keightley, P. D., T. F. Mackay and A. Caballero, 1993 Accounting for bias in estimates of the rate of polygenic mutation. Proceedings of the Royal Society of London, Series B: Biological Sciences 253**:** 291-296.

Kimura, M., 1983 *The Neutral Theory of Molecular Evolution*. Cambridge University Press, New York.

Klein, T. W., 1974 Heritability and genetic correlation: Statistical power, population comparisons and sample size. Behavior Genetics 4**:** 171-189.

Klein, T. W., J. C. De Vries and C. T. Finkbeiner, 1973 Heritability and genetic correlation: Standard errors of estimates and sample size. Behavior Genetics 3**:** 355-364.

Kondrashov, A. S., and D. Houle, 1994 Genotype-environment interactions and the estimation of the genomic mutation rate in *Drosophila melanogaster*. Proceedings of the Royal Society of London, Series B: Biological Sciences 258**:** 221-227.

Lande, R., 1976 Natural selection and random genetic drift in phenotypic evolution. Evolution 30**:** 314-334.

Latta, L. C., M. Peacock, D. J. Civitello, J. L. Dudycha, J. M. Meik *et al.*, 2015 The phenotypic effects of spontaneous mutations in different environments. American Naturalist 185**:** 243-252.

Littell, R., G. Milliken, W. Stroup, R. Wolfinger and O. Schabenberger, 2006 *SAS for Mixed Models*. SAS Institute, Cary, NC.

Luijckx, P., E. K. H. Ho, A. Stanic and A. F. Agrawal, 2018 Mutation accumulation in populations of varying size: Large effect mutations cause most mutational decline in the rotifer *Brachionus calyciflorus* under UV-C radiation. Journal of Evolutionary Biology 31**:** 924-932.

Lynch, M., 1988a Design and analysis of experiments on random drift and inbreeding depression. Genetics 120**:** 791-807.

Lynch, M., 1988b The rate of polygenic mutation. Genetical Research 51**:** 137-148.

Lynch, M., 1991 Methods for the analysis of comparative data in evolutionary biology. Evolution 45**:** 1065-1080.

Lynch, M., 2010 Evolution of the mutation rate. Trends in Genetics 26**:** 345-352.

Lynch, M., J. Blanchard, D. Houle, T. Kibota, S. Schultz *et al.*, 1999 Perspective: spontaneous deleterious mutation. Evolution 53**:** 645-663.

Lynch, M., and W. G. Hill, 1986 Phenotypic evolution by neutral mutation. Evolution 40**:** 915-935.

Lynch, M., and B. Walsh, 1998 *Genetics and analysis of quantitative traits*. Sinauer, Sunderland, MA.

Mackay, T., R. F. Lyman and W. G. Hill, 1995 Polygenic mutation in *Drosophila melanogaster*: non-linear divergence among unselected strains. Genetics 139**:** 849-859.

Mackay, T. F. C., E. A. Stone and J. F. Ayroles, 2009 The genetics of quantitative traits: challenges and prospects. Nature Reviews Genetics 10**:** 565-577.

Martin, G., and T. J. E. Lenormand, 2006 The fitness effect of mutations across environments: a survey in light of fitness landscape models. 60**:** 2413-2427.

Martorell, C., M. Toro and C. Gallego, 1998 Spontaneous mutation for life-history traits in *Drosophila melanogaster*, pp. 315-324 in *Mutation and Evolution*. Springer.

McGuigan, K., and M. W. Blows, 2010 Evolvability of individual traits in a multivariate context: partitioning the additive genetic variance into common and specific components. Evolution 64**:** 1899-1911.

McGuigan, K., and M. W. Blows, 2013 Joint allelic effects on fitness and metric traits. Evolution 67**:** 1131-1142.

McGuigan, K., J. M. Collet, S. L. Allen, S. F. Chenoweth and M. W. Blows, 2014 Pleiotropic mutations are subject to strong stabilizing selection. Genetics 197**:** 1051-1062.

McGuigan, K., D. Petfield and M. W. Blows, 2011 Reducing mutation load through sexual selection on males. Evolution 65**:** 2816-2829.

Merilä, J., and B. C. Sheldon, 1999 Genetic architecture of fitness and nonfitness traits: empirical patterns and development of ideas. Heredity 83**:** 103-109.

Meyer, K., 2009 Factor-analytic models for genotype x environment type problems and structured covariance matrices.
Genetics Selection Evolution 41: 21.
Meyer, K., and D. Houle, 2013 Sampling based approximation of confidence intervals for functions of genetic covariance
matrices. Proc. Assoc. Adv. Anim. Breed 20: 523-526.
Moher, D., A. Liberati, J. Tetzlaff, D. G. Altman and P. Group, 2009 Preferred reporting items for systematic reviews and
meta-analyses: the PRISMA statement. PLoS Medicine 6: e1000097.
Mukai, T., 1964 Genetic structure of natural populations of *Drosophila melanogaster*. I. Spontaneous mutation rate of
polygenes controlling viability. Genetics 50: 1-19.
Ness, R. W., A. D. Morgan, N. Colegrave and P. D. Keightley, 2012 Estimate of the spontaneous mutation rate in
Chlamydomonas reinhardtii. Genetics 192: 1447-1454.
Ostrow, D., N. Phillips, A. Avalos, D. Blanton, A. Boggs *et al.*, 2007 Mutational bias for body size in Rhabditid nematodes.
176: 1653-1661.
Reddiex, A. J., S. L. Allen and S. F. Chenoweth, 2018 A genomic reference panel for *Drosophila serrata*. G3 8: 1335-1346.
Rockman, M. V., 2012 The QTN program and the alleles that matter for evolution: All that's gold does not glitter.
Evolution 66: 1-17.
Roff, D. A., and T. A. Mousseau, 1987 Quantitative genetics and fitness: Lessons from *Drosophila*. Heredity 58: 103-118.
Rohlf, F., 2007 tpsRelw version 1.45. Department of Ecology and Evolution, State University of New York, Stony Brook.
Rohlf, F. J., 1999 Shape statistics: Procrustes superimpositions and tangent spaces. J. Classif. 16: 197-223.
Rohlf, F. J., 2013 tpsDig, digitize landmarks and outlines, version 2.17. Department of Ecology and Evolution, State
University of New York at Stony Brook.
Russell, W., G. Sprague and L. Penny, 1963 Mutations affecting quantitative characters in long-time inbred lines of
maize. Crop Science 3: 175-178.
Santiago, E., J. Albornoz, A. Domínguez, M. A. Toro and C. López-Fanjul, 1992 The distribution of spontaneous mutations
on quantitative traits and fitness in *Drosophila melanogaster*. Genetics 132: 771-781.
Santos, M., K. Fowler and L. Partridge, 1994 Gene-environment interaction for body size and larval density in *Drosophila*
melanogaster: an investigation of effects on development time, thorax length and adult sex ratio. Heredity 72:
515-521.
Schrider, D. R., D. Houle, M. Lynch and M. W. J. G. Hahn, 2013 Rates and genomic consequences of spontaneous
mutational events in *Drosophila melanogaster*. 194: 937-954.
Self, S. G., and K.-Y. Liang, 1987 Asymptotic properties of maximum likelihood estimators and likelihood ratio tests
under nonstandard conditions. Journal of the American Statistical Association 82: 605-610.
Shaw, R. G., D. L. Byers and E. Darmo, 2000 Spontaneous mutational effects on reproductive traits of *Arabidopsis*
thaliana. Genetics 155: 369-378.
Simons, Y. B., K. Bullaughey, R. R. Hudson and G. Sella, 2018 A population genetic interpretation of GWAS findings for
human quantitative traits. Plos Biology 16: e2002985.
Sung, W., M. S. Ackerman, S. F. Miller, T. G. Doak and M. Lynch, 2012 Drift-barrier hypothesis and mutation-rate
evolution. Proceedings of the National Academy of Sciences, USA 109: 18488-18492.
Sztepanacz, J. L., and M. W. Blows, 2017 Accounting for sampling error in genetic eigenvalues using random matrix
theory. Genetics 206: 1271-1284.
van der Graaf, A., R. Wardenaar, D. A. Neumann, A. Taudt, R. G. Shaw *et al.*, 2015 Rate, spectrum, and evolutionary
dynamics of spontaneous epimutations. Proceedings of the National Academy of Sciences of the United States
of America 112: 6676-6681.
Vassilieva, L. L., A. M. Hook and M. Lynch, 2000 The fitness effects of spontaneous mutations in *Caenorhabditis elegans*.
Evolution 54: 1234-1246.
Venables, W., and B. Ripley, 2002 *Modern Applied Statistics with S*. Springer, New York.
Walsh, B., and M. Lynch, 2018 *Evolution and Selection of Quantitative Traits*. Oxford University Press.
Wayne, M. L., and T. F. Mackay, 1998 Quantitative genetics of ovariole number in *Drosophila melanogaster*. II.
Mutational variation and genotype-environment interaction. Genetics 148: 201-210.
Wright, S., 1931 Evolution in Mendelian populations. Genetics 16: 97-159.
Yang, J., B. Benyamin, B. P. McEvoy, S. Gordon, A. K. Henders *et al.*, 2010 Common SNPs explain a large proportion of
the heritability for human height. Nature genetics 42: 565-569.

FIGURES AND TABLES

A. Taxa

B. Trait Type

Figure 1 The distribution of published mutational variance estimates by taxon (A) and trait (B) categories. The number of studies per category are shown (first value in brackets), where many studies included multiple estimates (from different traits, life stages or sexes; second value in brackets). See Methods (and Table S2) for details of the categories.

Figure 2 Schematic of design (A) and phenotypes (B) from a manipulative experiment in *Drosophila serrata*. (A) 42 MA lines (evolved through 20 generations of brother-sister mating) each founded two sublines: Small (S; 24 flies) and Large (L; 288 flies distributed evenly among 12 vials). These 84 lines (S and L subline per 42 MA lines) were maintained at these census population sizes for six generations (only two shown here). Each generation, all emergent flies from the 12 vials per L subline were pooled prior to virgin collection. For S sublines, two vials were established each generation; the focal vial contributed offspring to the next generation, while the replicate vial (grey shaded) did not. Each generation, wings were randomly sampled for six males from each of two vials per line (focal and replicate vials for S; randomly chosen two for L). (B) Wing size and shape were characterised from landmarks recorded on each image: proximal (1) and distal intersections of the radial vein (2); distal intersections of medial (3), cubital vein (4), and distal veins (5) and; the posterior (6, 7) and anterior (8, 9) cross-veins. Inter-landmark distance traits were described by their end-point landmarks (e.g., ILD1.2 was the distance between landmark1 and landmark 2).

Figure 3 Variation in estimates of mutational heritability (A) and coefficient of mutational variance (B) across trait categories. For each trait type (x-axis, defined in Methods), least-squares means ($\pm$ SE) were estimated from the analysis of model 1 for: all meta-analyses data (solid circles), Nematodes (grey squares) and Drosophila (open circles). The number of estimates, *n*, and studies for all taxa are shown across the top of each panel. For both (A) and (B), the transformed (log10 or BoxCox) values, as input to the analysis, are shown on the left-hand y-axis, and the untransformed values on the right-hand y-axis. No estimates of mutational variance in development time were

available for Drosophila. Trait type significantly predicted the magnitude of CV_{M} in the total data (see main text), of h_{M}^2 and CV_{M} in Drosophila (h_{M}^2: $F_{5,38.9} = 7.25$, $P < 0.0001$; CV_{M}: $F_{5,26.9} = 11.87$, $P < 0.0001$), and CV_{M} in Nematodes (h_{M}^2: $F_{6,31.7} = 0.53$, $P = 0.7780$; CV_{M}: $F_{6,62.8} = 3.20$, $P = 0.0083$).

Figure 4 Variation in estimates of mutational heritability (A) and coefficient of mutational variance (B) across taxa categories. Point estimates are the least-squares mean estimate (± SE) from the analysis of model 1. The number of estimates (*n*) and of studies (*x*) are shown for each category. For both (A) and (B), the transformed (*log*$_{10}$ or Box-Cox) values input to the analysis are shown on the left-hand y-axis, and the untransformed values on the right-hand y-axis.

Figure 5 Variation in wing trait mean across six generations of an experiment in *Drosophila serrata.* For each of the 11 wing traits (panels), the least-squares mean (± SE) is plotted for each generation for the Small (solid circle, solid lines) and Large (open circles, dashed lines) population size treatments. Centroid size (CS) is measured in millimetres; the ten inter-landmark distances are in units of centroid size (multiplied by 100; see Figure 2B for trait definitions). The F-ratio (and degrees of freedom) for the generation by population size treatment effect for each trait, from model 2, is given, where the asterisk indicates P ≤ 0.05.

Figure 6 Among-line variance estimates across six generations in an experiment in *Drosophila serrata.* Variances were estimated independently for each trait (panel) in each generation (x-axis) for each of the two population size treatments (Small: solid circles; Large: open circles). Plotted are the REML point estimate, and the REML-MVN 90% confidence intervals (CI). The dashed horizontal line indicates zero; estimates for which the lower CI did not overlap zero were interpreted as statistically supported. Where REML estimates of among-line variance were zero, no CI are plotted.

Figure 7 Estimates of variance from an experiment in *Drosophila serrata*. (A) Among-line by generation variance and (B) among-line variance estimated for 11 *D. serrata* wing traits (x-axis), in two different population size treatments (Small: solid circles; Large: open circles). Plotted are the REML point estimates (from model 4) and the REML-MVN 90% confidence intervals (CI) around this. The dashed horizontal line indicates zero; statistical significance was inferred where the lower 5% CI did not overlap zero. After applying a conservative 5% FDR correction, only two estimates in (A) and 21 in (B) remained significant (asterisk above CI).

Figure 8 Among-line variances for eleven wing traits in *Drosophila serrata*. (A) Among-line variance, V_L, was estimate per trait (model 5), and 90% CI estimated from the 10,000 REML-MVN samples as described in the Methods. (B) REML estimates of among line variance (points in panel A) were plotted against the total phenotypic variance (black circles, bottom x-axis) or mean (grey squares, top x-axis) of the trait.

SUPPLEMENTARY INFORMATION

Supplementary Table S1 **Information on mutational variance estimates obtained from the literature.**

Supplementary Table S2 **Broad and narrow trait categories.**

Supplementary Table S3 **Mutation accumulation (MA) lines shared across studies.**

Supplementary Table S4 **Statistical support for the effect of population size treatment on (A) among-line and (B) among-line by generation variance estimates.**

Supplementary Figure S1 **Diagram of systematic literature search for the meta-analysis of mutational variance estimates from mutation accumulation (MA) experiments.**

Supplementary Figure S2. **Distribution of the (log-transformed) number of generations (A) and the (log-transformed) number of lines (B) of mutation accumulation experiments in different taxon categories.**

Supplementary Figure S3 **Contribution of variability in phenotypic variance to estimates of among-line variance in *D. serrata*.**

Supplementary Figure S4 **Contribution of variability in trait mean to variability of among-line variance estimates in *D. serrata*.**

Table S1. Information on mutational variance estimates obtained from the literature. We detail estimates that were included in the meta-analysis (A) and those that were disqualified (L). For each paper, the source (a previous review or Web of Science search) is indicated (see footnote for source key), along with information on where the standardised estimates of mutational variance (mutational heritability, h_M^2 and, coefficient of mutational variance, CV_M) were obtained from or, where relevant, the justification for exclusion from the meta-analyses. Note: some papers reported estimates that were included in the meta-analysis (A) and other estimates that were excluded (B). Estimates (papers) identified in B were not included in the summaries of data (Figure 1) or analyses. All estimates (A and B) are available at:

https://doi.org/10.6084/m9.figshare.14913051.v2.

Reference	Source	Description of where estimates were obtained from
(A) ESTIMATES USED IN THE META-ANALYSIS		
Ajie, L. C., S. Estes, M. Lynch, and P. C. Phillips, 2005 Behavioural degradation under mutation accumulation in *Caenorhabditis elegans*. Genetics **170**: 655–660	1	Reference from in-text in Halligan & Keightley (2009). We calculated CV_M using the V_M (taking the number of generations as 307) from Estes *et al.* (2005) and the mean in Ajie *et al.* (2005)
Ávila, V. and A. García-Dorado, 2002 The effects of spontaneous mutation on competitive fitness in *Drosophila melanogaster*. J. Evol. Biol. **15**: 561–566	1	Halligan & Keightley (2009)'s Table 1.
Azevedo, R. L., P. D. Keightley, C. Laurén-Määttä, L. L. Vassilieva, M. Lynch, *et al.*, 2002 Spontaneous mutational variation for body size in *Caenorhabditis elegans*. Genetics **162**: 755–765	1	Table 1 of paper.
Baer, C. F., F. Shaw, C. Steding, M. Baumgartner, A. Hawkins, *et al.*, 2005 Comparative evolutionary genetics of spontaneous mutations affecting fitness in rhabditid nematodes. Proc. Natl. Acad. Sci. **102**: 5785–5790	1	Halligan & Keightley (2009)'s Table 1.
Baer, C. F., N. Phillips, D. Ostrow, A. Avalos, D. Blanton, *et al.*, 2006 Cumulative effects of spontaneous mutations for fitness in Caenorhabditis: Role of genotype, environment and stress. Genetics **174**: 1387–1395	1	Table 1 of paper.
Baer, C. F., J. Joyner-Matos, D. Ostrow, V. Grigaltchik, M. P. Salomon, *et al.*, 2010 Rapid decline in fitness of mutation accumulation lines of gonochoristic (outcrossing) *Caenorhabditis nematodes*. Evolution **64**: 3242–3253	2	h_M^2 estimates were from Table 2 of paper, insufficient information provided to calculate CV_M.
Bailey, D. W., 1959 Rates of subline divergence in highly inbred strains of mice. J. Hered **50**: 26–30	3	Houle *et al.* (1996)'s Table 1.
Burch, C. L., S. Guyader, D. Samarov, and H. Shen, 2007 Experimental estimate of the abundance and effects of nearly neutral mutations in the RNA virus φ6. Genetics **176**: 467–476	1	Halligan & Keightley (2009)'s Table 1.
Caballero, A., E. Cusi, C. García, and A. García-Dorado, 2002 Accumulation of deleterious mutations: Additional *Drosophila melanogaster* estimates and a simulation of the effects of selection. Evolution **56**: 1150–1159	1	CV_M calculated from in-text information, taking V_B from $\Delta V = V_B/(mean_G \times F_G)$; where $F_G=203$ and $mean_G = 0.625$.
Casellas, J. and J. F. Medrano, 2008 Within-generation mutation variance for litter size in inbred mice. Genetics **179**: 2147–55	4	MA litter mean was taken for generation 46 in Table 1. The V_M and h_M^2 estimates were taken from text and Table 3. CV_M was calculated from reported information.
Chang, S. and R. G. Shaw, 2003 The contribution of spontaneous mutation to variation in environmental response in *Arabidopsis thaliana*: responses to nutrients. Evolution **57**: 984–994	4	Table 3 of paper.
Charlesworth, L., H. Borthwick, C. Bartolomé, and P. Pignatelli, 2004 Estimates of the genomic mutation rate for detrimental alleles in *Drosophila melanogaster*. Genetics **167**: 815–826	1	Halligan & Keightley (2009)'s Table 1.
Clark, A. G., L. Wang, and T. Hulleberg, 1995 Spontaneous mutation rate of modifiers of metabolism in drosophila. Genetics **139**: 767–779	5	h_M^2 and CV_M estimates taken from Lynch *et al.* (1999)'s Table 1, then CV_M converted to percentage scale.

Reference	Source	Description of where estimates were obtained from
Clarke, D. K., E. A. Duarte, A. Moya, S. F. Elena, E. Domingo, et al., 1993 Genetic bottlenecks and population passages cause profound fitness differences in RNA viruses. J. Virol. **67**: 222–228	1	Halligan & Keightley (2009)'s Table 1.
Cox, T., D. D. Cox, and K. Frey, 1987 Mutations for polygenic traits in barley under nutrient stress. Euphytica **36**: 823–829	3	Houle et al (1996)'s Table 1.
Davies, S. K., A. Leroi, A. Burt, J. G. Bundy, and C. F. Baer, 2016 The mutational structure of metabolism in Caenorhabditis elegans. Evolution **70**: 2239–2246	2	Table S1 of paper.
Downie, D., 2003 Effects of short-term spontaneous mutation accumulation for life history traits in grape phylloxera, Daktulosphaira vitifoliae. Genetica **119**: 237–251	4	V_M and h^2_m were taken from Table 1. Line means were estimated from figure 3 (L =0.288, C = 0.235 for G12, and G24 survival, respectively). CV_M calculated from information provided.
Estes, S., L. C., Ajie, M. Lynch, and P. C. Phillips, 2005 Spontaneous mutational correlations for life-history, morphological and behavioral characters in Caenorhabditis elegans. Genetics **170**: 645–653	1	We calculated CV_M using V_M (taking the number of generations as 307) from Estes et al (2005) and the mean from Ajie et al (2005)
Etienne,V.,E.C.Andersen,J.M.Ponciano,D.Blanton,A.Cadavid,et al.,2015 The red death meets the abdominal bristle: polygenic mutation for susceptibility to a bacterial pathogen in Caenorhabditis elegans. Evolution **69**: 508–519	2	Table 1 of paper.
Farhadifar, R., J. M. Ponciano, E. C. Andersen, D. J. Needleman, and C. F. Baer, 2016 Mutation is a sufficient and robust predictor of genetic variation for mitotic spindle traits in Caenorhabditis elegans. Genetics **203**: 1859–1870	2	Trait means from Table 1 of paper, and variances from Table S4.
Fernández, J and C. López-Fanjul, 1996 Spontaneous mutational variances and covariances for fitness-related traits in Drosophila melanogaster. Genetics **143**: 829–837	1	V_M and h^2_m taken from Table 3 & trait mean from Table 1 of paper; CV_M calculated from those estimates.
Festing, M., 1973 A multivariate analysis of subline divergence in the shape of the mandible in c57bl/gr mice. Genet. Res. **21**: 121–132	3	Houle et al. (1996)'s Table 1.
Fry, J. D. and S. L. Heinsohn, 2002 Environment dependence of mutational parameters for viability in Drosophila melanogaster. Genetics **161**: 1155–1167	1	We calculated CV_M by dividing $\sqrt{V_M}$/mean$_{MA}$×100, where V_M was from Table 7, the mean$_{MA}$ was from Table 1 of paper.
Fry, J. D., 2001 Rapid mutational declines of viability in Drosophila. Genet. Res. **77**: 53–60	1	Halligan & Keightley (2009)'s Table 1.
Fry, J. D., P. D. Keightley, S. L. Heinsohn, and S. V. Nuzhdin, 1999 New estimates of the rates and effects of mildly deleterious mutation in Drosophila melanogaster. Proc. Natl. Acad. Sci. **96**: 574–579	1	Halligan & Keightley (2009)'s Table 1.
García-Dorado, A., J. L. Monedero, and C. López-Fanjul, 1998 The mutation rate and the distribution of mutational effects of viability and fitness in Drosophila melanogaster. In Mutation and Evolution, pp. 255–265, Springer	1	Halligan & Keightley (2009)'s Table 1.
García-Dorado, A., J. Fernández, and C. López-Fanjul, 2000 Temporal uniformity of the spontaneous mutational variance of quantitative traits in Drosophila melanogaster. Genet. Res. **75**: 47–51	1	Table 2 of paper.
Hall, D. W. R. Mahmoudizad, A. W. Hurd, and S. L. Joseph, 2008 Spontaneous mutations in diploid Saccharomyces cerevisiae: Another thousand cell generations. Genet. Res. **90**: 229–241	4	Table 2 of paper.
Harada, K., 1995 A quantitative analysis of modifier mutations which occur in mutation accumulation lines in Drosophila melanogaster. Heredity **75**: 589–598	5	CV_M was calculated from information in Table 3 of paper; among line variance was back-calculated using 4V_G (reported) and then divided by 2× the number of generations. h^2_m could not be calculated as no V_E estimates were reported.
Houle, D., D. Hoffmaster, S. Assimacopoulos, and L. Charlesworth. 1992 The genomic mutation rate for fitness in Drosophila. Nature **359**: 58–60	1	Halligan & Keightley (2009)'s Table 1.
Houle, D., K. A. Hughes, D. K. Hoffmaster, J. Ihara, S. Assimacopoulos, et al., 1994 The effects of spontaneous mutation on quantitative traits. I. Variances and covariances of life history traits. Genetics **138**: 773–785	3	Houle et al. (1996)'s Table 1.

Reference	Source	Description of where estimates were obtained from
Huang, W., R. F. Lyman, R. A. Lyman, M. A. Carbone, S. T. Harbison, *et al.*, 2016 Spontaneous mutations and the origin and maintenance of quantitative genetic variation. Elife **5**: e14625	2	h²ₘ estimates were taken from supplementary information 1D. Estimates were averaged across sex. Could not calculate CV_M.
Joseph, S. L. and D. W. Hall, 2004 Spontaneous mutations in diploid *Saccharomyces cerevisiae* more beneficial than expected. Genetics **168**: 1817–1825	1	Halligan & Keightley (2009)'s Table 1.
Joyner-Matos, J., A. Upadhyay, M. P. Salomon, V. Grigalchik, and C. F. Baer, 2009 Genetic (co)variation for life span in rhabditid nematodes: Role of mutation, selection, and history. J. Gerontol. A Biol. Sci. Med. Sci. **64**: 1134–1145	2	Table 2 of paper.
Joyner-Matos, J., L. C. Bean, H. L. Richardson, T. Sammeli, and C. F. Baer, 2011 No evidence of elevated germline mutation accumulation under oxidative stress in *Caenorhabditis elegans*. Genetics **189**: 1439–1447	2	Estimates of h²ₘ(V_M/V_E) and CV_M(V_M/ mean for relative fitness) then taken as a percentage, were calculated using information from Table 1 of paper.
Kavanaugh, C. M. and R. G. Shaw, 2005 The contribution of spontaneous mutation to variation in environmental responses of *Arabidopsis thaliana*: Responses to light. Evolution **59**: 266–275	1	Estimates from information in Table 5 of paper.
Keightley, P. D. and A. Caballero, 1997 Genomic mutation rates for lifetime reproductive output and lifespan in *Caenorhabditis elegans*. Proc. Natl. Acad. Sci. **94**: 3823–3827	1	Estimates for productivity and r (intrinsic growth rate) from Halligan & Keightley (2009)'s Table 1. Lifespan estimates were taken from text on p. 3825 of paper.
Kibota, T. T. and M. Lynch, 1996 Estimate of the genomic mutation rate deleterious to overall fitness in *E. coli*. Nature **381**: 694–696	1	Halligan & Keightley (2009)'s Table 1.
Latimer, C. A. L., K. McGuigan, R. S. Wilson, M. W. Blows, and S. F. Chenoweth, 2014 The contribution of spontaneous mutations to thermal sensitivity curve variation in *Drosophila serrata*. Evolution **68**: 1824–1837	2	Table 1 of paper.
Latta, L. C., M. Peacock, D. J. Civitello, J. L. Dudycha, J. M. Meik, *et al.*, 2015 The phenotypic effects of spontaneous mutations in different environments. Am. Nat. **185**: 243–252	2	Authors did not specify exact generation of MA (taken as average of 93) and the reported estimates of V_M had not been divided by number of generations. V_M, h²ₘ and CV_M provided in supplementary have all been recalculated using V_M/(2*93).
López, M. A. and C. López-Fanjul, 1993 Spontaneous mutation for a quantitative trait in *Drosophila melanogaster*. II. Distribution of mutant effects on the trait and fitness. Genet. Res. **61**: 117–126	3	Houle *et al.* (1996)'s Table 1.
Lynch, M., 1985 Spontaneous mutations for life-history characters in an obligate parthenogen. Evolution pp. 804–818	3	Houle *et al.* (1996)'s Table 1.
Lynch, M., L. Latta, J. Hicks, and M. Giorgianni, 1998 Mutation, selection, and the maintenance of life-history variation in a natural population. Evolution **52**: 727–733	4	Reference taken from in-text citations in Walsh & Lynch (2018); Estimates of h²ₘ, V_M and V_E were taken from Table 1 of paper. MA mean calculated using $z_0 + R_m$ (Table 1), and used to calculate CV_M.
Mackay, T. F. and R. F. Lyman, 1998 Polygenic mutation in *Drosophila melanogaster*: Genotype × environment interaction for spontaneous mutations affecting bristle number. Genetica **102**: 199	2	Table 9 of paper. Only the unselected lines were considered here, and we averaged estimates across sex.
Martorell, C., M. Toro, and C. Gallego, 1998 Spontaneous mutation for life-history traits in *Drosophila melanogaster*. In Mutation and Evolution, pp. 315–324, Springer	5	Table 2 of paper.
McGuigan, K., D. Petfield, and M. W. Blows, 2011 Reducing mutation load through sexual selection on males. Evolution **65**: 2816–2829	2	h²ₘ was taken from text. CV_M was calculated by determining V_M as (mean(V_E) × h²ₘ), where V_E and MA means were taken from Tables S1 and S2 of paper.
Morgan, A. D., R. W. Ness, P. D. Keightley, and N. Colegrave, 2014 Spontaneous mutation accumulation in multiple strains of the green alga, *Chlamydomonas reinhardtii*. Evolution **68**: 2589–2602	2	Generation estimates from Table 2. Authors had repeated the fitness assays twice, where we report estimates of CV_M from the second attempt in Table 3. No estimates of V_E or h²ₘ were reported.
Mukai, T., 1964 The genetic structure of natural populations of *Drosophila melanogaster*. I. spontaneous mutation rate of polygenes controlling viability. Genetics **50**: 1–19	3	Houle *et al.* (1996)'s Table 1.

Reference	Source	Description of where estimates were obtained from
Mukai, T. and T. Yamazaki, 1968 The genetic structure of natural populations of *Drosophila melanogaster*. V. coupling-repulsion effect of spontaneous mutant polygenes controlling viability. Genetics **59**: 513–535.	1	Estimates taken from text of paper.
Mukai, T., S. I. Chigusa, J. F. Crow, and L. E. Mettler, 1972 Mutation rate and dominance of genes affecting viability in *Drosophila melanogaster*. Genetics **72**: 335–355	1	Halligan & Keightley (2009)'s Table 1.
Mukai,T., K. Harada,and H. Yoshimaru, 1984 Spontaneous mutations modifying the activity of alcohol dehydrogenase (adh) in *Drosophila melanogaster*. Genetics **106**: 73–84	3	Houle *et al.* (1996)'s Table 1.
Ohnishi, O., 1977 Spontaneous and ethyl methanesulfonate-induced mutations controlling viability in *Drosophila melanogaster*. II. homozygous effect of polygenic mutations. Genetics **87**: 529–545	1	Halligan & Keightley (2009)'s Table 1.
Ostrow, D., N. Phillips, A. Avalos, D. Blanton, A. Boggs, *et al.*, 2007 Mutational bias for body size in Rhabditid nematodes. Genetics **176**: 1653–1661	4	Table 2 of paper.
Paxman, G., 1957 A study of spontaneous mutation in *Drosophila melanogaster*. Genetica **29**: 39–57	3	Houle *et al.* (1996)'s Table 1.
Pannebakker, L. A., D. L. Halligan, K. T. Reynolds, G. A. Ballantyne, D. M. Shuker, *et al.*, 2008 Effects of spontaneous mutation accumulation on sex ratio traits in a parasitoid wasp. Evolution **62**: 1921–1935	1	MA mean and line numbers taken from Table 2 of paper. Variance estimates were taken from Table 3, where CV_M has been converted to a percentage scale.
Pletcher, S. D., D. Houle, and J. W. Curtsinger, 1998 Age-specific properties of spontaneous mutations affecting mortality in *Drosophila melanogaster*. Genetics **148**: 287–303	5	h^2_m, V_M and mean from Table 5 of paper and CV_M estimated from that information.
Roles, A. J., M. T. Rutter, I. Dworkin, C. L. Fenster, and J. K. Conner, 2016 Field measurements of genotype by environment interaction for fitness caused by spontaneous mutations in *Arabidopsis thaliana*. Evolution **70**: 1039–1050	2	Table 2 of paper. Count traits were identified as outliers in preliminary analyses and excluded.
Russell, W., G. Sprague, and L. Penny, 1963 Mutations affecting quantitative characters in long-time inbred lines of maize. Crop Sci. **3**: 175–178	3	Houle *et al.* (1996)'s Table 1.
Rutter, M. T., F. H. Shaw, and C. L. Fenster, 2010 Spontaneous mutation parameters for *Arabidopsis thaliana* measured in the wild. Evolution **64**: 1825–35	2	Table 2 of paper.
Santiago, E., J. Albornoz, A. Domínguez, M. Toro, and C. López-Fanjul, 1992 The distribution of spontaneous mutations on quantitative traits and fitness in *Drosophila melanogaster*. Genetics **132**: 771–781	3	Houle *et al.* (1996)'s Table 1.
Schoen, D. J., 2005 Deleterious mutation in related species of the plant genus *Amsinckia* with contrasting mating systems. Evolution **59**: 2370–2377	1	Halligan & Keightley (2009)'s Table 1.
Shaw, R. G., D. L. Byers, and E. Darmo, 2000 Spontaneous mutational effects on reproductive traits of *Arabidopsis thaliana*. Genetics **155**: 369–78	1	Estimates for number of fruits and seeds per fruit taken from Halligan & Keightley (2009)'s Table 1. Reproductive mass estimates were taken from Table 1 of paper.
Vassilieva, L. L., A. M. Hook, and M. Lynch, 2000 The fitness effects of spontaneous mutations in *Caenorhabditis elegans*. Evolution **54**: 1234–1246	1	h^2_M and V_M estimates were taken from Table 2. For CV_M, the mean of MA was calculated as the intercept plus the slope from Table 1, and V_M taken as the slope from Table 2, divided by 2 (diploid genome).
Vassilieva, L. L. and M. Lynch, 1999 The rate of spontaneous mutation for life-history traits in *Caenorhabditis elegans*. Genetics **151**: 119–129	1	h^2_M and V_M estimates were taken from Table 1 of paper. The MA mean calculated using $z_0 + R_n$ (Table 1), and used for calculating CV_M. We omitted CV_M estimate for Generation rate (1/T, the reciprocal of the mean age at reproduction) as estimates were an order of magnitude larger than any other estimate (MA$_{mean}$ = 0.23, CV_M=93.796).
Wayne, M. L. and T. F. Mackay, 1998 Quantitative genetics of ovariole number in *Drosophila melanogaster*. II. Mutational variation and genotype-environment interaction. Genetics **148**: 201–210	1	Table 3 of paper.
Xu, J., 2004 Genotype-environment interactions of spontaneous mutations for vegetative fitness in the human pathogenic fungus *Cryptococcus neoformans*. Genetics **168**: 1177–1188	1	h^2_m estimates from Table 2 of paper. Insufficient information to calculate CV_M values.
Zeyl, C. and J. A. G. DeVisser, 2001 Estimates of the rate and distribution of fitness effects of spontaneous mutation in *Saccharomyces cerevisiae*. Genetics **157**: 53–61	1	Halligan & Keightley (2009)'s Table 1.

Reference	Source Description of where estimates were obtained from

(B) ESTIMATES REMOVED FROM META-ANALYSIS

Reference	Source Description of where estimates were obtained from
Baer, C. F., J. Joyner-Matos, D. Ostrow, V. Grigaltchik, M. P. Salomon, et al., 2010 Rapid decline in fitness of mutation accumulation lines of gonochoristic (outcrossing) Caenorhabditis nematodes. Evolution 64: 3242–3253	As in (A): Removed Caenorhabditis brenneri heritability estimate for Productivity as the estimate was an outlier at 6.3 standard deviations from the mean.
Farhadifar, R., J. M. Ponciano, E. C. Andersen, D. J. Needleman, and C. F. Baer, 2016 Mutation is a sufficient and robust predictor of genetic variation for mitotic spindle traits in Caenorhabditis elegans. Genetics 203: 1859–1870	As in (A): One estimate (for posterior centrosome oscillation frequency) was removed as the estimate for V_M was negative.
Gong, Y., R. Woodruff, and J. Thompson Jr., 2005 Deleterious genomic mutation rate for viability in Drosophila melanogaster using concomitant sibling controls. Biol. Lett. 1: 492–495	CV_M estimates were taken from Halligan & Keightley (2009)'s Table 1. However, these were an order of magnitude larger than any other CV_M estimate. Halligan & Keightley (2009) re-scaled the estimates to whole genome ($\times$ 5.5) and expressed as a proportion of the mean control, but nonetheless omit these values from their summary. We also chose to omit them here.
Harada, K., 1995 A quantitative analysis of modifier mutations which occur in mutation accumulation lines in Drosophila melanogaster. Heredity 75: 589–598	As in (A): Could not calculate CV_M (negative mean) for one metabolite (Isocitrate-dehydrogenase activity) which was therefore excluded.
Houle, D. and S. V. Nuzhdin, 2004 Mutation accumulation and the effect of copia insertions in Drosophila melanogaster. Genet. Res. 83: 7–18	Excluded from the meta-analysis as the estimate of mutational variance was derived from the slope of regression on generation, and the number of generations of mutation accumulation therefore undefined.
Kavanaugh, C. M. and R. G. Shaw, 2005 The contribution of spontaneous mutation to variation in environmental responses of Arabidopsis thaliana: Responses to light. Evolution 59: 266–275	As in (A): Several estimates of mutational variance were negative, and these traits were excluded (Biomass, Bolting day, Fruit number, Height, Leaf length and Leaf number).
Latta, L. C., M. Peacock, D. J. Civitello, J. L. Dudycha, J. M. Meik, et al., 2015 The phenotypic effects of spontaneous mutations in different environments. Am. Nat. 185: 243–252	As in (A): Estimates for First instar duration, Spine length at maturity, Spine length at second adult instar, Third clutch size, and Total length at maturity excluded from analysis due to poor precision (all reported rounded to have the same value).
Pletcher, S. D., D. Houle, and J. W. Curtsinger, 1999 The evolution of age-specific mortality rates in Drosophila melanogaster: Genetic divergence among unselected lines. Genetics 153: 813–823	V_M and V_E estimates in Table 4 were pooled across generations of 10, 19 and 47, making the data unsuitable for meta-analysis of the effect of generation number. V_M was reported for each specific generation in Table 5, however corresponding estimates of V_E or trait mean were not reported.
Schultz, S. T., M. Lynch, and J. H. Willis, 1999 Spontaneous deleterious mutation in Arabidopsis thaliana. Proc. Natl. Acad. Sci. 96: 11393-11398	Estimates were taken from Table 3, where $\sqrt{CV_M}$ was calculated from $V_{B/2\sigma}{}^2$ reported in Table 3 as $(\sqrt{V_L}/z\bar{z})2$. Excluded from our analyses as the number of lines were a large outlier (8.7 SD above mean).

[1] Halligan, D. L. and P. D. Keightley, 2009 Spontaneous mutation accumulation studies in evolutionary genetics. Annu. Rev. Ecol. Evol. Syst. 40: 151-172

[2] Web of Science database. Journal article document types were searched for with the following search topics criteria: "mutation accumulation" and (varia* or "mutat* coefficient*"), on the 11/12/2019 at 4:38 PM AEST.

[3] Houle, D., B. Morikawa, and M. Lynch, 1996 Comparing mutational variabilities. Genetics 143: 1467–1483

[4] Walsh, L. and M. Lynch 2018 Evolution and Selection of Quantitative Traits. Oxford University Press

[5] Lynch, M., L. Walsh, 1998 Genetics and Analysis of Quantitative Traits. Sinauer Sunderland, MA

Table S2. Broad and narrow trait categories. The trait description is a list of traits included within each category.

Broad Category	Trait	Trait Description
Life-history	Development time	Age at maturity, Age at reproduction, Growth rate, Instar duration, The reciprocal of the mean age at reproduction, Time to maturity
	Fitness	Colony size, Fitness, Competitive mating success, Phi (rate of convergence to the stable age distribution), Population growth, r (intrinsic rate of increase)
	Miscellaneous	Bolting day, Female longevity, Flowering day, Flowering timing, Lifespan, Longevity, Male longevity, Male mating, Sex ratio, Silking date
	Productivity	Clutch size, Fecundity, Grain yield, Litter size, Number of flowers, Ovariole number, Productivity, Reproductive mass, Tassel branches
	Survival	Age-specific survivorship, Germination, Pathogen susceptibility, Survival, Viability
Morphology	Morphology	Abdominal bristles, Biomass, Body length, Body size, Body width, Ear diameter, Ear length, Fruit size, Height, Leaf length, Leaf number, Limb lengths, Mandible length, Skull length, Spine length, Sternopleural bristles, Weight, Wing length
Physiology	Physiology	Day sleeping characteristics, Enzyme activity, Locomotor activity, Metabolites, Night sleeping characteristics, Velocity, Waking activity

Table S3. Mutation accumulation (MA) lines shared across studies. The first published description of a set of MA lines is indicated in bold; the following non-bolded entries represent subsequent investigations of the same set of MA lines. See Table S1 for citation details.

Species name	Reference	Generation	Trait Description
Arabidopsis thaliana	**Shaw *et al.* (2000)**	17	Fecundity, Reproductive mass
	Chang & Shaw (2003)	17	Height, Leaf number, Fruit size
	Kavanaugh & Shaw (2005)	17	Bolting day, Flowering day, Fecundity, Height, Leaf length, Leaf number
	Rutter *et al.* (2010)	25	Fecundity
	Roles *et al.* (2016)	25	Fecundity, Germination, Age-specific survivorship, Fitness
	Rutter *et al.* (2018)	25	Biomass, Flowering timing, Fruit size, Germination, Survival
Caenorhabditis briggsae	**Baer *et al.* (2005)**	200	Productivity
	Baer *et al.* (2006)	220	Productivity
	Ostrow *et al.* (2007)	200	Body size
	Joyner-Matos *et al.* (2009)	250	Life span
Caenorhabditis elegans	**Baer *et al.* (2005)**	200	Productivity
	Baer *et al.* (2006)	220	Productivity
	Ostrow *et al.* (2007)	200	Body size
	Joyner-Matos *et al.* (2009)	250	Life span
	Etienne *et al.* (2015)	250	Pathogen susceptibility
	Davies *et al.* (2016)	250	Metabolites
	Farhadifar *et al.* (2016)	250	Mitotic cell division traits
	Keightley & Caballero (1997)	60	Lifespan, Productivity, Population growth
	Azevedo *et al.* (2002)	60	Body size
	Vassilieva *et al.* (1999)	49.4	Longevity, Productivity, Survivorship, Population growth
	Vassilieva *et al.* (2000)	214	Longevity, Productivity, Survivorship, Population growth
	Azevedo *et al.* (2002)	152	Body size
	Ajie *et al.* (2005)	370	Locomotor activity, Productivity, Velocity
	Estes *et al.* (2005)	370	Population growth parameter, Body width
Drosophila melanogaster	**Houle *et al.* (1994)**	44	Fecundity, Fitness, Longevity, Male mating, Productivity
	Pletcher *et al.* (1998)	19	Female longevity, Male longevity
	Pletcher *et al.* (1999)	47	Female longevity, Male longevity
	Mackay & Lyman (1998)	223	Abdominal bristles, Sternopleural bristles
	Wayne & Mackay (1998)	228.5	Body size, Ovariole number
	Mukai (1964)	25	Viability
	Mukai & Yamazaki (1968)	60	Viability
	Santiago *et al.* (1992)	46	Sternopleural bristles, Wing dimensions
	Fernandez & Lopez-Fanjul (1996)	106	Viability, Fecundity
	Garcia-Dorado *et al.* (1998)	183	Fitness
	Martorell *et al.* (1998)	82	Productivity, Longevity
	Garcia-Dorado *et al.* (2000)	170	Wing length
		204	Abdominal bristles, Sternopleural bristles
		209	Viability
	Chavarrias *et al.* (2001)	250	Viability
		255	Viability
	Caballero *et al.* (2002)	210	Viability
	Avila & Garca-Dorado (2002)	288	Fitness
Drosophila serrata	**McGuigan *et al.* (2011a)**	23	Mating Success
		26	Productivity
	Latimer *et al.* (2014)	32	Locomotor activity
Oscheius myriophila	**Baer *et al.* (2005)**	200	Productivity
	Ostrow *et al.* (2007)	200	Body size
	Joseph & Hall (2004)	1012	Fitness

Saccharomyces cerevisiae	Hall *et al.* (2008)	2062	Fitness

Table S4. Statistical support for the effect of population size treatment on (A) among-line and (B) among-line by generation variance estimates. Model-fit information for model 5 (see main text) is reported per trait (defined in Figure 2B), including: the number of parameters; the -2 log-likelihood score and; the Akaike information criterion (AIC). Model fit was assessed using a Likelihood Ratio Test (LRT) based on the difference in -2 log-likelihood scores between the model estimating treatment-specific variance, and the model estimating a single pooled estimate (i.e., the Chi-square statistic), where the degrees of freedom was the difference in the number of model parameters. Both the P-value from this LRT test (bold values <0.05), and the multiple-test corrected *P*-value (at a conservative 5% false discovery rate, FDR; significant tests in bold) are reported.

Trait	Model	Parameters	-2 Log likelihood	AIC	Chi-square statistic	*P*−value	5% FDR *P*−value
(A)			Model fit for estimating treatment-specific versus pooled *among-line* variance				
CS	Treatment-specific	28	-22247.9	-22191.9			
	Pooled	27	-22281.6	-22227.6	33.7	**0.000**	**0.000**
1.2	Treatment-specific	28	11381.7	11433.7			
	Pooled	27	11356	11406	25.7	**0.000**	**0.000**
1.5	Treatment-specific	28	9414.9	9466.9			
	Pooled	27	9338.7	9388.7	76.2	**0.000**	**0.000**
1.9	Treatment-specific	28	8626.1	8676.1			
	Pooled	27	8609.3	8657.3	16.8	**0.000**	**0.000**
2.5	Treatment-specific	28	10839.3	10887.3			
	Pooled	27	10818.2	10866.2	21.1	**0.000**	**0.000**
2.8	Treatment-specific	28	10598.6	10652.6			
	Pooled	27	10574.8	10626.8	23.8	**0.000**	**0.000**
3.7	Treatment-specific	28	9642.2	9694.2			
	Pooled	27	9603.5	9655.5	38.7	**0.000**	**0.000**
3.8	Treatment-specific	28	4803.1	4853.1			
	Pooled	27	4757.8	4805.8	45.3	**0.000**	**0.000**
4.5	Treatment-specific	28	9707.6	9755.6			
	Pooled	27	9668.3	9710.3	39.3	**0.000**	**0.000**
4.8	Treatment-specific	28	5102.4	5148.4			
	Pooled	27	5081	5125	21.4	**0.000**	**0.000**
5.6	Treatment-specific	28	7490.6	7540.6			
	Pooled	27	7480.9	7526.9	9.7	**0.002**	**0.002**
(B)			Model fit for estimating treatment-specific versus pooled *among-line by generation* variance				
CS	Treatment-specific	28	-22247.9	-22191.9			
	Pooled	27	-22242.4	-22188.4	5.5	**0.0190**	0.21
1.2	Treatment-specific	28	11381.7	11433.7			
	Pooled	27	11381.5	11431.5	0.2	0.6547	0.72
1.5	Treatment-specific	28	9414.9	9466.9			
	Pooled	27	9415.2	9465.2	0.3	0.5839	0.72
1.9	Treatment-specific	28	8626.1	8676.1			
	Pooled	27	8624.3	8672.3	1.8	0.1797	0.47
2.5	Treatment-specific	28	10839.3	10887.3			
	Pooled	27	10842.2	10890.2	2.9	0.0886	0.47
2.8	Treatment-specific	28	10598.6	10652.6			
	Pooled	27	10597.3	10649.3	1.3	0.2542	0.47
3.7	Treatment-specific	28	9642.2	9694.2			
	Pooled	27	9642.1	9694.1	0.1	0.7518	0.75
3.8	Treatment-specific	28	4803.1	4853.1			

	Pooled	27	4802.9	4852.9	0.2	0.6547	0.72
4.5	Treatment-specific	28	9707.6	9755.6			
	Pooled	27	9708.4	9754.4	0.8	0.3711	0.58
4.8	Treatment-specific	28	5102.4	5148.4			
	Pooled	27	5103.9	5149.9	1.5	0.2207	0.47
5.6	Treatment-specific	28	7490.6	7540.6			
	Pooled	27	7492.4	7538.4	1.8	0.1797	0.47

Figure S1. Diagram of systematic literature search for the meta-analysis of mutational variance estimates from mutation accumulation (MA) experiments. The below schematic follows the PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analysis, MOHER *et al.* 2009) protocol. Further information, including mutational parameter estimates, available at: https://doi.org/10.6084/m9.figshare.14913051.v2.

Figure S2. Distribution of the (log-transformed) number of generations (A) and the (log-transformed) number of lines (B) of mutation accumulation experiments in different taxon categories. For each box plot, the median of the distribution is shown as a black band, grey boxes represent the interquartile range, whiskers represent 1.5 times the interquartile range, and estimates that fell outside this range are designated with open circles. The data are plotted on the log scale, but values reported on the y-axes correspond to the raw scale.

Figure S3. Contribution of variability in phenotypic variance to estimates of among-line variance in *D. serrata*. The among-line estimate from the per trait per generation per treatment implementation of model (3) (Figure 6) was divided by the corresponding estimate of total phenotypic variance (summed among-line, among-vial and residual variance from model (3)). Plotted are these variance-scaled REML estimates (and REML-MVN 90% CI) for each trait (panel; trait descriptions in Figure 2B) in each generation (x-axis) for each of the two population size treatments (Small: solid circles; Large: open circles). The dashed horizontal line indicates zero.

Figure S4. Contribution of variability in trait mean to variability of among-line variance estimates in *D. serrata*. Among-line variance estimates (V_L, Figure 6) were placed on a coefficient of variance scale to account for the influence of trait mean on magnitude of estimates, using the following formula: $\sqrt{\frac{V_L}{\bar{x}}} \times 100$ where $\bar{x}$ is the trait least squares mean (Figure 5). Plotted are the REML point estimates (and 90% CI) for each trait (panel; trait descriptions in Figure 2B) in each generation (x-axis) for each of the two population size treatments (Small: solid circles; Large: open circles). The 90% confidence intervals were estimated using 10,000 REML-MVN samples of V_L and the trait mean as described in the Methods. Some MVN sample estimates of V_L were negative and the coefficient of variance was thus undefined for those samples. The dashed horizontal line indicates zero. Where the REML estimate of among-line variance was zero, CI were not estimated.