A behavioral architecture for realistic simulations of *Drosophila* larva locomotion and foraging

Panagiotis Sakagiannis[1*], **Anna-Maria Jürgensen**[1], **Martin Paul Nawrot**[1*]

[1]Computational Systems Neuroscience, University of Cologne

*For correspondence:
p.sakagiannis@uni-koeln.de (PS);
mnawrot@uni-koeln.de (MN)

Present address: Computational Systems Neuroscience, Institute of Zoology, 50674, University of Cologne, Germany

Abstract The *Drosophila* larva is extensively used as model organism in neuroethological studies where precise behavioral tracking enables the statistical analysis of individual and population-level behavioral metrics that can inform mathematical models of larval behavior. Here, we propose a hierarchical model architecture comprising three layers to facilitate modular model construction, closed-loop simulations, and direct comparisons between empirical and simulated data. At the basic layer, the autonomous locomotory model is capable of performing exploration. Based on novel kinematic analyses our model features intermittent forward crawling that is phasically coupled to lateral bending. At the second layer, navigation is achieved via active sensing in a simulated environment and top-down modulation of locomotion. At the top layer, behavioral adaptation entails associative learning. We evaluate virtual larval behavior across agent-based simulations of autonomous free exploration, chemotaxis, and odor preference testing. Our behavioral architecture is ideally suited for the modular combination of neuromechanical, neural or mere statistical model components, facilitating their evaluation, comparison, extension and integration into multifunctional control architectures.

Introduction

Drosophila larvae express a fairly tractable behavioral repertoire that is consistent across the 4-5 days of the larval life stage (*Almeida-Carvalho et al., 2017*). Most of the larval lifetime is dedicated to foraging for suitable nutrients while avoiding threats via stereotyped evasive behaviors. Behavioral control is achieved via neural circuits conserved throughout development (*Gerhard et al., 2017*), that span the entire tripartite CNS consisting of the brain, the subesophageal zone (SEZ) and the ventral nerve cord (VNC), making the larva a formidable system for studying control, execution and adaptation of behavior (*Zarin et al., 2019*; *Jovanic, 2020*; *Eschbach and Zlatic, 2020*; *Imambocus et al., 2022*). After reaching critical mass for pupation, homeostatic signals switch behavior towards food aversion, hypermobility and collaborative burrowing (*Wu et al., 2003*), terminating the feeding state and leading to pupation and metamorphosis.

High resolution methods for behavioral tracking (*Ohyama et al., 2013*; *Risse et al., 2017*; *Schumann and Triphan, 2020*; *Tadres and Louis, 2020*; *Thane et al., 2023*; *Laurent et al., 2024*), now routinely used in neuroethological experiments, have revealed detailed insight in the organisation of larval foraging behavior. In the absence of food resources, larvae engage in free exploration to locate food patches (*Sims et al., 2019*) with a characteristic alternation of locomotory activity and brief pauses (*Sakagiannis et al., 2020*), a property also reported for adult fly behavior (*Ueno et al., 2012*; *Reynolds et al., 2015*). Food consumption involves repetitive feeding motion and digging into the substrate (*Kim et al., 2017*). Statistical regularities that govern foraging behavior have been unveiled by analysis at the microscale of body kinematics and at the macroscale of larva trajectories

(*Denisov et al., 2013*; *Risse et al., 2017*; *Karagyozov et al., 2018*). Locomotion combines the basic sensorimotor primitives of crawling and turning, and has been in the main focus of recent studies (*Heckscher et al., 2012*; *Wystrach et al., 2016*; *Sims et al., 2019*), whereas studies of feeding behavior remain scarce (*Ruiz-Dubreuil et al., 1996*). Both, crawling and feeding consist of recurring sensorimotor cycles controlled by central pattern generating circuits (*Heckscher et al., 2012*; *Mantziaris et al., 2020*; *Miroschnikow et al., 2018a*).

Salient olfactory cues can trigger appetitive or aversive chemotaxis, during which larvae navigate up or down a chemical gradient (*Gomez-Marin et al., 2011*; *Slater et al., 2015*; *Schleyer et al., 2015a*; *Klein et al., 2017*). During appetitive chemotaxis, the detection of minor concentration changes during lateral bending causes a directional bias in the turning movement, establishing a mechanism of active sensing (*Gomez-Marin and Louis, 2012*; *Wystrach et al., 2016*; *Thane et al., 2019*). Encounters with novel odorants in the presence of a food reward or a negative reinforcement such as the bitter taste substance quinine can induce olfactory learning, enabling short- and long-term behavioral adaptations (*Schleyer et al., 2011*, *2015b*; *Gerber and Stocker, 2007*; *Diegelmann et al., 2013*; *Widmann et al., 2018*; *Weiglein et al., 2019*; *Jürgensen et al., 2024*). For quantification of choice behavior, widely-used standard group assays for behavioral preference testing have been established (*Gerber and Stocker, 2007*; *Gerber et al., 2014*; *Schleyer et al., 2015a*,b, *2011*).

The routine access to detailed behavioral data, and the broad approaches to neuroethological experiments makes the *Drosophila* larva an ideal model system for computational modeling studies on behavioral control principles. Existing generative models typically aim at specific aspects of larval behavior and can vary widely in model type and abstraction level, ranging from basic neuromechanics to the abstract mathematical description. We therefore propose the concept of the behavioral architecture, a three-layered hierarchical and modular framework, nested from behavioral primitives to high-level function, to integrate models of diverse types and different levels of abstraction (*Figure 1A*). At the basic layer we establish a refined coupled-oscillator model for larval locomotion capable of autonomous exploratory behavior. To this end we provide a detailed analysis of larval tracking data from experiments of free exploration. Introducing sensory modulation at the second layer allows us to reproduce several experimental findings of appetitive chemotaxis. The combination with a biologically realistic spiking neural network model for associative learning at the adaptive layer reproduces empirical population-level learning scores across a multiple trial training protocol. The architecture allows for agent-based, potentially closed-loop, simulations of virtual larvae in virtual environments and generates simulated data at the level of the experimentally observed behavior facilitating model calibration and evaluation.

Model

Behavioral architecture

Larval behavior is hierarchically structured, i.e. sensorimotor primitives such as crawling, bending and feeding can be integrated into more complex behaviors such as exploration, taxis or exploitation. It has been proposed that the hierarchy of animal behavior is reflected by the underlying neuroanatomy as a hierarchy of nested sensorimotor loops (*Prescott et al., 1999*; *Wilson and Prescott, 2022*; *Prescott and Wilson, 2023*). A functional modeling paradigm that exploits this idea regards the neural system as a layered control architecture (or subsumption architecture) (*Brooks, 1986*) where low-level stereotyped reflexive and repetitive behaviors are autonomously generated by localized peripheral motor circuitries, while multisynaptic loops involving more centralized neural circuits act as descending modulators on the local circuits in order to coordinate global and complex behavior. The central idea is that energy-efficient decentralized neural control is the rule, while higher centers are recruited only when more extensive integration is needed e.g. in order to react suitably to unexpected sensory stimulation. Furthermore, there are only limited degrees of freedom by which higher layers can influence local sensorimotor loops via descending pathways

Figure 1. Behavioral architecture for larva foraging. A: In the trilayer architecture the bottom layer consists of three basic sensorimotor effectors that constitute the locomotory model. The intermediate layer features innate reactive behavior in response to sensory input that reflects changes in the environment. The top layer allows for behavioral adaptation through experience. Framed areas denote more complex behaviors that require subsumption of subordinate behaviors. **B:** Suggested implementation of basic behavioral modules at the bottom layer. Initiation or cessation of a behavior is controlled by the intermittency module. The turner and crawler module are phasically coupled during forward locomotion, while crawling and feeding are implemented as mutually exclusive sensorimotor primitives.

⁹¹ e.g. by starting or stopping, or by accelerating, decelerating or inverting their autonomous function ⁹² (*Sen et al., 2017*; *Feng et al., 2020*; *Bidaye et al., 2020*). The complexity of inter-layer top-down mod-⁹³ ulation is thus predicted to be considerably lower than the complexity of signal integration within ⁹⁴ layers. Layered control architectures have been used mainly in behavior-based robotics (*Bicho,* ⁹⁵ *1999*), allowing sequential calibration and modular integration of partial neuroscientific models ⁹⁶ under a common framework (*Brooks, 1986*; *Prescott et al., 1999*).

⁹⁷ We here propose a three-layered behavioral architecture for *Drosophila* larva foraging as illus-⁹⁸ trated in *Figure 1A*. The bottom layer consists of three basic behaviors: crawling, turning and feed-⁹⁹ ing. Each is realized by a low-level local sensorimotor loop between motor effectors and sensory ¹⁰⁰ feedback. For exploration this involves proprioception and mechanoreception, for feeding it addi-¹⁰¹ tionally involves gustatory input. Integration of these basic behaviors within the layer gives rise to ¹⁰² composite behaviors. For example while exploration in stimulus-free conditions only entails crawl-¹⁰³ ing and turning, differences in the temporal microstructure between these two yield a spectrum ¹⁰⁴ from localized search to remote dispersal. The peripheral ventral nerve cord (VNC) circuits medi-¹⁰⁵ ating exploration have been shown to be capable of autonomous, decentralised function without ¹⁰⁶ the need for any top-down regulation (*Sims et al., 2019*).

¹⁰⁷ The intermediate layer introduces salient sensory stimulation of different modalities and there-¹⁰⁸ fore allows for reactive behavior in the face of presented risks and opportunities. Following the ¹⁰⁹ subsumption architecture paradigm, we only assume a single modulatory input from the inter-¹¹⁰ mediate to the bottom layer. This assumption is in line with the idea of pre-motor integration of ¹¹¹ signals from different sensory modalities into a final integrated descending modulatory pathway, ¹¹² which eventually affects behavior (*Wystrach et al., 2016*; *Eschbach and Zlatic, 2020*). Modulation ¹¹³ of exploratory behavior in the presence of salient stimulation enables coherent navigation along ¹¹⁴ sensory gradients. In the present study we consider odor-driven chemotaxis as a process of active ¹¹⁵ sensing in which the larva constantly samples the odor concentration during lateral bending mo-¹¹⁶ tions, enabling odorscape navigation and odor source discovery. In an alternative application, we ¹¹⁷ recently simulated larval behavior during active sensing in thermotaxis experiments across differ-¹¹⁸ ent *Drosophila* species (*Kafle et al., 2024*).

¹¹⁹ Finally, at the top layer, associative learning biases the evaluation of recurring sensory stimuli

and can modulate the innate valence of odors to allow for experience-dependent adaptation of
navigation. Again, the modulatory top-down signal to the underlying layers is limited to the mod-
ulation of approach versus avoidance behavior, in line with the research on the mushroom body
output signal in *Drosophila* (*Gerber and Stocker, 2007*; *Gerber et al., 2014*; *Schleyer et al., 2015a*,b,
2011; *Owald and Waddell, 2015*).

The proposed behavioral architecture is naive to the level of model abstraction of the underly-
ing neural mechanisms. Its role is to create a modular framework where diverse and potentially
competing models can be positioned, integrated and tested in behavioral simulations. It is there-
fore justified to use simple generative models to bridge the gaps that have not yet been studied
in detail until they can be substituted by more elaborate future versions. To ensure flexibility and
compatibility, every module has a defined basic input and output regardless of the internal struc-
ture, complexity, level of abstraction and spatiotemporal scale of its possible instantiations. For
the purpose of modeling under the behavioral architecture framework we provide the *larvaworld*
python package available at https://pypi.org/project/larvaworld/.

Locomotory model

We propose a model for locomotion in the two-dimensional plane that defines the momentary
body state by the instantaneous forward velocity v and angular velocity ω as generated through
crawling and bending, respectively. Extending on previous models (see Discussion) we incorporate
two new features based on our kinematic analysis of larval locomotion: (i) the peristaltic cycle
phase dependent attenuation of angular motion, and (ii) the intermittent crawling as transitions
between runs and pauses. We briefly describe the modular components of our model. A detailed
mathematical description is provided in the Materials and Methods section. The complete pipeline
for model calibration is described in *Appendix 2*.

We choose to simplify the larva to a two-segment body (*Figure 2F*). This abstraction allows to
describe the body state at any point in time by only three variables: (i) position of a selected spine
midpoint (*Appendix 1-Figure 1*), (ii) absolute orientation of the front segment θ, and (iii) bending
angle θ_b between the front and rear segments. This approach is in line with the common practice of
quantifying larva bending via a single angle between one anterior and one posterior vector (*Gomez-*
Marin et al., 2011; *Lahiri et al., 2011*; *Paisios et al., 2017*).

Locomotion is achieved via forward crawling and lateral body bending. Crawling moves the
midpoint along the orientation vector. Bending reorients the front segment by rotation around the
midpoint. Forward displacement restores θ_b back to zero by gradually aligning the rear segment to
the orientation axis. A demonstration of the bisegmental model simplification is shown in *Video 4*.

Based on our empirical results (*Figure 2* and *Figure 3*), we propose a coupled oscillator model
of locomotion that generates both, weathervaning and headcasts. To this end we implement an
oscillatory process (crawler) that generates the forward-velocity v during subsequent stride cycles
(*Figure 1B*) where a tonic input I_C modulates crawling frequency f_C. A second oscillator (turner)
generates alternating left-right bending as previously proposed by *Wystrach et al.* (*2016*) with oscil-
latory amplitude A_T. To incorporate the empirically observed crawl-bend interference, we couple
the two oscillators by imposing a crawler-phase dependent suppression on A_T where the phase-
dependent modulation is modelled as Gaussian function and fitted to the empirical data (red pro-
file in *Figure 3C*).

To accommodate the intermittent behavior with alternating crawling runs and crawling pauses,
we chose to implement a stochastic model approach where the number of strides per stridechain
and the pause durations each follows a log-normal distribution fitted to the empirical data (*Fig-*
ure 3B). For a demonstration of the locomotory model in different configurations see *Video 1*.

Under the constraints of the subsumption architecture paradigm the crawler and turner can
be influenced by higher-order circuits in a limited number of ways. In the case of the crawler,
frequency modulation and the initiation or cessation of the oscillation-cycle are the two available
top-down modulations. Likewise for the turner, top-down modulation of the tonic input affects

170 both, the frequency and the amplitude of oscillation.

Results

Kinematic analysis of larva locomotion

173 We start out with the kinematic analysis of experimental larva trajectories and body postures in
174 order to infer and parameterize several aspects of larva locomotion that will inform our modeling
175 approach. Using diverse metrics that capture spatial and temporal dynamics we specifically assess
176 the oscillation of forward velocity during consecutive peristaltic strides and its influence on lateral
177 bending, the less pronounced oscillation of angular velocity, the intermittent nature of crawling,
178 and the inter-individual variability of a number of locomotion-related parameters across different
179 larvae.

180 We analyzed the trajectory of each single larva by tracking the forward velocity v of a posterior
181 midline point (*Figure 2A*) and the head orientation angular velocity ω (*Figure 2E*). The former illus-
182 trates that forward locomotion indeed consists of consecutive steps (strides) that are characterized
183 by the alternation of increased (green) and decreased (red) values of v. The latter confirms that lat-
184 eral bending occurs both, during crawl-runs (weathervaning) and during stationary crawl-pauses
185 (headcasts). The individual trajectory depicted in *Figure 2* can be seen in full length in *Figure 2—*
186 *video 1*, while the short epoch depicted in the inset is shown in *Figure 2—video 2*.

187 To characterize both types of oscillations we first perform a Fourier analysis of v and ω on each
188 of 100 larva tracks (*Figure 3A*). Across larvae we observe a robust peristaltic rhythm with a crawling
189 frequency f_C normally distributed around 1.4 Hz in line with earlier results (*Heckscher et al., 2012*;
190 *Mantziaris et al., 2020*; *Thane et al., 2023*). We also confirm previous reports that lateral bending
191 manifests a slower, more variable rhythm f_T around 0.4 Hz (*Wystrach et al., 2016*). Then we detect
192 all strides performed by an individual animal and verify their stereotypical structure in terms of
193 stride duration, resulting body displacement, and the phase-dependence of v (*Figure 2D*).

194 In order to quantify the phase-dependence of angular motion we interpolate each detected
195 stride into a 64-bin oscillation cycle of period 2π. The progression of forward velocity from head to
196 tail during the peristaltic cycle is illustrated in *Figure 3C* (top). To analyze the angular metrics during
197 the peristaltic cycle we consider the absolute angular change ignoring the left or right direction of
198 a turn. This analysis reveals a robust phase dependence of the angular velocity ω and acceleration
199 $\dot{\omega}$, and of the bend angle θ_b (*Appendix 2-Figure 1A*). Specifically, the angular velocity ω exhibits a
200 smooth unimodal dependency on the stride period peaking during the early stride phase and just
201 before the head forward velocity reaches its maximum (*Figure 3C*). Nevertheless, ω during crawl-
202 ing is lower than during pauses as shown when we normalize it by dividing with the average $\bar{\omega}_P$
203 computed during the pause epochs (*Figure 3C*, bottom). This implies a phasic attenuating interfer-
204 ence of the peristaltic cycle on the lateral bending mechanism (*Figure 1B*). A plausible mechanistic
205 explanation featuring bodily interference of crawling and bending is suggested in the Discussion.

206 Larvae intermittently pause crawling before re-assuming it. This results in sequences of con-
207 catenated strides (runs or stridechains) that alternate with brief crawl-pauses (*Figure 2C*). We an-
208 alyzed the stridechain length with respect to the number of strides and the pause duration in the
209 experimental dataset. To this end we pooled both measures across 100 larvae to obtain the em-
210 pirical distributions. Testing power-law, exponential, log-normal and Levy distributions revealed
211 the highest quality of fit for the log-normal distribution for both parameters (*Figure 3B*).

Simulation of behavioral experiments

213 In this section we simulate increasingly complex behavioral experiments. Starting from stimulus-
214 free exploration we advance to chemotactic navigation and finally to adaptive odor preference
215 experiments. We simulate individual larvae using the model as calibrated in *Appendix 2* and all
216 model parameters are shown in *Appendix 2-Figure 1A*. Larval populations are constructed by pool-
217 ing individual virtual larvae that behave independently as they move through the spatial arena and

Figure 2. Kinematic analysis of a single Drosophila larva in locomotion. A: Individual larva trajectory tracking a posterior point along the midline of the animal . Trajectory color denotes the forward velocity v from 0 (red) to maximum (green). Inset focuses on a shorter track epoch analyzed in C and G. The full-length trajectory and the epoch in the inset are shown in *Figure 2—video 1* and *Figure 2—video 2* respectively. Dark green rectangle denotes the single stride described in B. **B:** Sketch of the single crawling stride indicated in A. The larva first stretches its head forward, anchors it to the substrate and then drags its body forward via peristaltic contraction. Scaled stride displacement $\mathring{d}_S$ is defined as the resulting displacement d divided by the body-length l. **C:** Scaled forward velocity v during the $40\,$s track epoch selected in A (inset). Green and red markers denote the local maxima and minima used for stride annotation. Individual strides are tiled by vertical dashed lines. Successive strides constitute uninterrupted stridechains (white). Epochs that do not show any strides are annotated as crawl-pauses (gray). **D:** Scaled forward velocity v of head, midpoint and tail as a function of the stride cycle phase Φ. All detected strides have been interpolated to a stride oscillation cycle of period 2π. Solid lines denote the median, shaded areas the lower and upper quartiles across strides. **E:** Same trajectory as in (A) now tracking the head segment. Color denotes the absolute orientation angular velocity ω from 0 (red) to maximum (green). The full-length trajectory and the epoch in the inset are shown in *Figure 2—video 1* and *Figure 2—video 2* respectively. **F:** Definition of bending angle θ_b and orientation angle θ for the original 12-segment (blue) and the simplified 2-segment (red) larvae. **G:** Three angular parameters during the same track epoch shown in (C). Bending angle θ_b, bend and orientation angular velocities ω_b, ω are shown. Background shadings denote left and right turning bouts. For illustration purposes only turns resulting in a change of orientation angle $\Delta\theta > 20°$ are shown. **H:** Absolute orientation angular velocity ω during the stride cycle, as shown for v in (D).

Figure 2—video 1. The full-length trajectory (A, E) colored according to angular and forward velocity.

http://computational-systems-neuroscience.de/wp-content/uploads/2024/10/3.mp4

Figure 2—video 2. The short track epoch depicted in the insets (A, E) colored according to forward and angular velocity

http://computational-systems-neuroscience.de/wp-content/uploads/2024/10/4.mp4

Figure 3. Population-level analysis. A: Fourier analysis of the forward v (red) and angular ω (blue) velocity across 100 larvae. Inset shows the respective dominant frequencies within suitable ranges $1 \leqslant f_C \leqslant 2.5$ and $0.1 \leqslant f_T \leqslant 0.8$ for v and ω, respectively. Crawling exhibits a dominant frequency of around 1.4 while lateral bending has a slower more variable rhythm of around 0.4. **B:** Epoch-duration distribution. Dots describe the cumulative probability density over logarithmic bins for the length of stridechains and the duration of crawl-pauses pooled across the larva population. Lines indicate the distribution with the lowest Kolmogorov-Smirnov distance among the best-fitting power-law, exponential, log-normal and Levy distributions. Stridechain length and pause duration are best approximated by log-normal distributions. **C-D:** Crawl-bend interference. The stride cycle kinematics are depicted for a single individual. All detected strides have been interpolated into a 64-bin oscillation cycle of period 2π. **C:** Forward velocity of 5 points along the larva midline. Velocity is scaled to the larva body-length. **D:** Absolute angular velocity ω (blue) normalized by the average value $\bar{\omega}_P$ computed during the pause epochs. Fitted Gaussian function (red) describes well the phase-dependent attenuation imposed on ω and is used for the implementation of the coupled-oscillator locomotory model. Solid lines denote the median, shaded areas the lower and upper quartiles. Vertical dashed lines denote the cycle phase where the respective velocity reaches its maximum value. Inset : Phase offset $\Delta\phi$ between the peak phase of each midline point's forward velocity $\phi_C^{v_i}$ and the peak phase of angular velocity ϕ_C^{ω} across a dataset of 100 tracked larvae. Notably, ω reaches its maximum just before the head forward velocity reaches its maximum.

Figure 4. Free exploration in simulation and experiment. A: Dispersal of 200 larvae in experiment (left) and simulation (right) during 40 seconds. Individual tracks have been transposed to originate from the center of the arena. The entire temporal course shown in *Figure 4—video 1*. **B:** Dispersal from origin. Line indicates the group median while shaded area denotes first and third quartiles. **C:** Histograms for total number of strides, time ratio allocated to crawling and pathlength. (arena dimensions = 500x500mm, N = 200 larvae, experiment duration = 3 minutes, simulation timestep =1/16).

Figure 4—video 1. The temporal course of dispersal for real (left) and virtual (right) larvae shown in (A).

http://computational-systems-neuroscience.de/wp-content/uploads/2024/10/5.mp4

---

odorscape (*Niewalda et al., 2014*).

Exploration

In the first virtual experiment, we consider free exploration (*Figure 4*) and evaluate our simulation
results against an empirical data set (see Materials and Methods). To this end we simulate a popu-
lation of 200 virtual larvae during three minutes while exploring a stimulus-free environment on a
non-nutritious substrate in a Petri-dish, mimicking the experimental lab conditions (*Figure 4*). Due
to the absence of sensory input and food, the proposed intermittent coupled-oscillator locomotory
model at the basic layer is sufficient to autonomously generate exploratory movement.

Larvae are initially placed at the center of the dish, each with a random body orientation. Over
the course of the experiment, both the simulated and the real larval population disperses in space
(*Figure 4A*, *Figure 4—video 1*).

The quantitative comparison shows a good agreement between simulated and empirical data
with respect to the radial dispersal from the initial position (*Figure 4B*). Importantly, the variability
across individual animals is well captured by the variability between individual model instances as
shown for the number of strides, the crawling time, and the total distance travelled by each larva
(*Figure 4C*).

Chemotaxis

Chemotaxis describes the process of exploiting an odor gradient in space to locate an attractive or
avoid a repelling odor source. An olfactory sensor (olfactor) placed at the front end of the virtual

237 body enables active sensing during body bending and allows detection of concentration changes
238 that modulate turning behavior accordingly. Sensory-driven behavior is enabled via modulation
239 of the frequency and amplitude of the lateral oscillator (turner) by sensory feedback, while the
240 peristaltic oscillator is not affected, as proposed in (*Wystrach et al., 2016*).

241 To assess the chemotactic efficiency of our coupled-oscillator model we reconstruct the arena
242 and odor landscape (odorscape) of two behavioral experiments described in (*Gomez-Marin et al.,*
243 *2011*). In the first, larvae are placed on the left side of the arena facing to the right. An appetitive
244 odor source is placed on the right side. The virtual larvae navigate up the odor gradient approach-
245 ing the source (*Figure 5A,C,E*), reproducing the experimental observation in Fig.1 C in (*Gomez-Marin*
246 *et al., 2011*). In the second, both the odor source and the virtual larvae are placed at the center
247 of the arena. The larvae perform localized exploration, generating trajectories across and around
248 the odor source. (*Figure 5B,D,F*), again replicating the observation in Fig.1 D in (*Gomez-Marin et al.,*
249 *2011*). Two sample simulations can be seen in *Figure 5—video 1*.

Odor preference test

251 We simulate the odor preference paradigm as described in the Maggot Learning Manual (*Michels*
252 *et al., 2017*). Larvae are placed at the center of a dish containing two odor sources in opposite
253 sides and left to freely explore. The odor concentrations are Gaussian-shaped and overlapping,
254 resulting in an odorscape of appetitive and/or aversive opposing gradients. After 3 minutes the
255 final situation is evaluated. The established population-level metric used is the olfactory preference
256 index (PI), computed for the left odor as $PI_l = \frac{N_l - N_r}{N}$ where N_l and N_r is the number of larvae on
257 the left and right side of the dish while N is the total number of larvae.

258 The extend of olfactory modulation on the turning behavior is determined by the odor-specific
259 gain G. As this is measured in arbitrary units, we first need to define a realistic value range that cor-
260 relates with the behaviorally measured PI. We perform a parameter-space search independently
261 varying the gain for left and right odors and measuring the resulting PI in simulations of 30 larvae.
262 The results for a total of 25^2 gain combinations within a suitable range of $G \in [-100, 100]$ are illus-
263 trated in *Figure 6A*. Simulation examples for one appetitive and one aversive odor are shown in
264 *Figure 6—video 1*.

265 In order to simulate larval group behavior in response to an associative learning paradigm we
266 interface our behavioral simulation with the spiking mushroom body (MB) model introduced in
267 (*Jürgensen et al., 2024*) (*Figure 6C*). It implements a biologically realistic neural network model of
268 the olfactory pathway according to detailed anatomical data using leaky integrate-and-fire neurons
269 (*Jürgensen et al., 2021*). The MB network undergoes associative plasticity at the synapses between
270 the Kenyon cells and two MB output neurons as a result of concurrent stimulation with an odor
271 and a reward signal. Both, odor and reward is simulated as spike train input to the receptor neu-
272 rons and the reinforcement signalling dopaminergic neuron, respectively. The model employs two
273 output neurons (MB_+, MB_-), representing a larger number of MB compartments associated with
274 approach/avoidance learning respectively (*Saumweber et al., 2018*). The initially balanced firing
275 rates between MB_+ and MB_- are skewed after learning and encode the acquired odor valence
276 (*Owald et al., 2015*; *Owald and Waddell, 2015*) here defined as:

$$MB_{out} = \frac{MB_+ - MB_-}{MB_+ + MB_-} \qquad | \qquad MB_{out} \epsilon [-1, 1]$$

278 We first trained the MB model via a classical conditioning experiment where, in each condition-
279 ing trial, it experiences an odor (conditioned stimulus, CS_+) in combination with a sugar reward
280 during 5 min, following the standard training protocol in (*Michels et al., 2017*). 5 groups of 30 MB
281 models undergo between 1 and 5 sequential conditioning trials (*Weiglein et al., 2019*). The result-
282 ing odor valence MB_{out} from each MB model was converted to an odor gain G via a simple linear
283 transformation and used to generate a virtual larva (*Figure 6B*). Each population of 30 larvae was
284 then tested in an odor preference simulation. The larvae were placed on a dish in presence of the
285 previously rewarded odor (CS+) and a neutral odor in opposite sides of the dish (*Figure 6—video 1*),

Figure 5. Simulation of chemotaxis. A: Experiment 1: A single odor source of $8.9\mu M$ peak concentration is placed on the right side of the rectangular arena creating a chemical gradient as indicated by the color scale. Larvae are placed on the left side facing to the right. Larvae are expected to navigate up the gradient approaching the source. A single larva trajectory is shown. This setup mimics the first experiment in (*Gomez-Marin et al., 2011*). **B:** Experiment 2: A single odor source of $2.0\mu M$ peak concentration is placed at the center of the rectangular arena. Larvae are placed in close proximity to the odor source. Larvae are expected to locally explore generating trajectories around and across the source. A single larva trajectory is shown. This setup mimics the second experiment in (*Gomez-Marin et al., 2011*). **C,D:** The trajectories of 25 virtual larvae during the two experiments. **E,F:** The odor concentration encountered by the virtual larvae as a function of time. Red curves refer to the single larva in A and B. Gray denotes the mean and quartiles of all 25 larvae in C and D. The simulation results fit well to the experimental estimates of concentration sensing during larval chemotaxis in (*Gomez-Marin et al., 2011*). (arena dimensions = 100x60mm, N = 30 larvae, experiment duration = 3 and 5 minutes respectively, simulation timestep =1/16).

Figure 5—video 1. Time course of the two simulated chemotaxis experiments.

http://computational-systems-neuroscience.de/wp-content/uploads/2024/10/6.mp4

Figure 6. Simulation of innate and learned odor preference. A: A total of 25^2 simulations are shown with the resulting Preference Index for different gains of the left and right odor. On the top left the initial state is shown with the larvae randomly generated at the center of the dish. The final state of three additional simulations is depicted on the top right and bottom left and right. See *Figure 6—video 1* for videos of two sample simulations. **B:** The pipeline used for coupling the Mushroom Body (MB) model with the behavioral simulation. First a MB model is trained via a classical conditioning experiment where olfactory input is combined with reward. The resulting odor valence MB_{out} is then converted to odor gain G via a simple linear transformation and used to generate a virtual larva. Finally the odor preference of a virtual larva population is evaluated in a behavioral simulation. **C:** The spiking neural network comprising the MB model. The number of neurons comprising each layer is indicated. **D:** The resulting PIs for 100 simulations per number of training trials. In each of the 100 simulations per condition a population of 30 virtual larvae was generated and evaluated using a different random seed, always bearing the exact same 30 odor gains derived from the respective group of 30 trained MB models (arena dimensions = $100x100$mm, N = 30 larvae, experiment duration = 3 minutes, simulation timestep =0.1.

Figure 6—video 1. Two odor preference simulations, one with an appetitive and one with an aversive odor source placed on the left side of the dish. A non-valenced odor source is placed on the right side.

http://computational-systems-neuroscience.de/wp-content/uploads/2024/10/7.mp4

---

again following standard experimental procedures (*Michels et al., 2017*). To obtain robust results we replicated the experiment 100 times per population with a different random seed for a total of 600 simulations. The obtained preference indexes (PI) are illustrated in *Figure 6D*. The PI increase with increasing number of trials as well as its saturation resembles empirical observations (*Wei-glein et al., 2019*). Note that the variability of the PI across the 100 simulations per condition is introduced solely by the behavioral simulation and resembles that seen across real experiments.

The current implementation only sequentially couples a trained MB model to be tested in a behavioral simulation. In the discussion we further elaborate on a possible extension featuring their closed-loop integration allowing for full behavioral simulations of both the training and the testing phase of the associative learning paradigm in a virtual environment.

Discussion

Experimental evidence for layered behavioral architecture

The proposed behavioral architecture is based on two underlying principles that justify our modular, hierarchically layered approach. Firstly, we attempt to horizontally structure it into behaviorally functional modules, each generating a well-defined behavior. Secondly, we vertically parse it into semi-independent layers, each under top-down modulation from the ones above but still capable of generating behavior independently of them. We summarize the neuroanatomical observations

and behavioral experiments that support this approach.

The neural mechanisms underlying the three basic larval behaviors (*Figure 1A*, basic layer) have been extensively studied. Crawling is characterized by fairly stereotypical repetitive strides. Initial contraction of the head and tail segments driven by a 'visceral piston' mechanism is followed by a laterally symmetric peristaltic wave traversing neighboring segments longitudinally from back to front (*Heckscher et al., 2012*) (*Figure 2B*). Segmental central pattern generators (CPGs) coupled via intersegmental short- and long-range connectivity motifs involving also premotor neurons constitute the underlying neural circuitry (*Kohsaka et al., 2019*; *Zarin et al., 2019*; *Mantziaris et al., 2020*). Lateral bending results from asymmetric contraction of body musculature initiated at the thoracic segments (*Berni, 2015*). Feeding is generated via a network of mono- and multi-synaptic sensorimotor loops from enteric, pharyngeal and external sensory organs to motor neurons controlling mouth-hook movement, head-tilt and pharyngeal pumping (*Miroschnikow et al., 2018b*; *Schoofs et al., 2024*). A noteworthy facet of these behaviors is their autonomous generation without the need for descending control. This has been demonstrated by continued exploration even after brain ablation (*Sims et al., 2019*). Exploitation of a nutritious substrate requires all three basic behaviors as the larva constantly consumes food and re-positions its body. According to the layer-independence principle, we postulate that peripheral consummatory circuits at the sub-esophageal zone (SEZ) are also capable of autonomous exploitation.

Higher brain centers play a pivotal role in behavior modulation, with neurotransmitters like dopamine, serotonin, acetylcholine, and octopamine being key players in this intricate process (*Berni et al., 2012*; *Zhang et al., 2013*; *Miroschnikow et al., 2018b*; *Malloy et al., 2019*; *Eschbach et al., 2020*; *Vogt et al., 2021*). It has been suggested that the transition between exploration and exploitation (feeding) is acutely induced via dopaminergic signaling (*Schleyer et al., 2020*) while their long-term balance is regulated via hugin-mediated homeostatic neuromodulation (*Schoofs et al., 2014*). Identification of sensory pathways towards motor effector neuropiles further elucidates the role of interoception in behavioral modulation (*Qian et al., 2018*).

The neural mechanisms that underlie olfactory modulation of the basic locomotory behavior are also under intense investigation. Chemotactic approach and avoidance to innately valenced odors has been attributed to a predominantly innate pathway involving the antennal lobe (AL) and its direct projection to the lateral horn (LH), both in the larva (*Schulze et al., 2015*; *Vogt et al., 2021*) and in the adult fly (*de Belle and Heisenberg, 1994*; *Strutz et al., 2014*; *Dolan et al., 2018*). Modulation of behavior by learned odors strongly involves descending control by the MB in juvenile (*Saumweber et al., 2018*; *Eschbach et al., 2020*) and adult (*Slater et al., 2015*) stages. Both pathways have similar modulating effects on foraging behavior and are likely integrated in a premotor network downstream of the AL (*Schleyer et al., 2015a*; *Eschbach et al., 2020*). In the sensorimotor loop, descending pathways involving the LH control cessation of crawling, possibly triggering sharper re-orientation when navigating down-gradient, facilitating chemotaxis (*Tastekin et al., 2018*). Finally, the internal homeostatic state (e.g. starvation vs. satiation) is implicated in behavioral regulation via neurotransmitter release at multiple levels, including AL, LH and MB and neuropeptide expression (*Vogt et al., 2021*; *de Tredern et al., 2024*).

Previous computational models of larva locomotion

Larva locomotion has been modeled extensively. Generative models of peristaltic crawling feature neural and/or neuromuscular dynamics. Sequential neural activity patterns across the segmental body can be generated by longitudinally repeated CPGs of paired excitatory and inhibitory population rate units, possibly involving proprioceptive feedback (*Gjorgjieva et al., 2013*; *Pehlevan et al., 2016*). This relatively abstract CPG model has been elaborated into a connectome-based circuitry of premotor and motor neurons. Under tonic activation the model exhibits two functionally distinct, though structurally overlapping, recurring patterns of neuromuscular activity, responsible for forward and backward peristalsis respectively (*Zarin et al., 2019*). At the other end of the modeling spectrum the contribution of the visceral-piston mechanism to the peristaltic cycle has been

assessed in a biomechanical model (*Ross et al., 2015*). A more elaborate neuromechanical model,
based on segmental localized reflexes and substrate frictional forces and assuming empirically
informed axial and transverse oscillatory frequencies, has been shown to generate forward and
backward crawling without the need for any neural activation pattern (*Loveless et al., 2019*).

Lateral bending has also been captured in statistical (*Davies et al., 2015*) or generative models
(*Wystrach et al., 2016*; *Loveless et al., 2019*; *Loveless and Webb, 2018*). In the context of free-
exploration it has been shown to be a byproduct of chaotic body neuromechanics underlying peri-
stalsis (*Loveless et al., 2019*). In the context of chemotaxis it has been attributed to a distinct oscil-
latory process, autonomous (*Wystrach et al., 2016*) or semi-autonomous to crawling (*Davies et al.,*
2015). In the former case oscillation is driven by mutual inhibition between excitatory-inhibitory
circuits. In the latter, bending behavior is dissected into low-amplitude weathervaning while crawl-
ing and high-amplitude headcasting during crawl-pauses, an approach essentially equivalent to
an attenuation of the lateral oscillation amplitude due to crawling. Autonomous exploration and
chemotaxis can be generated by an integration of the neuromechanical and the independent lat-
eral oscillator models (*Loveless and Webb, 2018*).

Apart from the rare occasion where such models implement mutually exclusive mechanistic
hypotheses and are indubitably incompatible to each other, in most cases they are indeed comple-
mentary, overlapping, nested or disconnected in terms of the generative mechanisms they aim to
capture and could potentially coexist under a broader control system. In this context our unifying
architecture for larval behavior in which any partial model can be positioned can be valuable for
modelers and roboticists, interested in a behavior-based synthetic approach.

374 Intermittent coupled-oscillator model for realistic locomotion

Each of the above models could be adjusted so that at minimum it generates the 2D translational
and angular motion of a virtual body and could therefore populate the basic locomotory layer
of the behavioral architecture. In this study we propose such a model, assembled in a synthetic
approach by distinct modules, which either extend previously proposed locomotory models or
integrate novel findings derived from our analysis of kinematic parameters. Given a simplified
bisegmental body, a simple oscillator under frequency-regulating tonic activation is adequate for
generating recurrent strides, efficiently summarizing the complex underlying neural and neurome-
chanical dynamics (*Heckscher et al., 2012*; *Mantziaris et al., 2020*). Concerning angular motion, the
previously introduced lateral oscillator model (*Wystrach et al., 2016*) meets the requirements and
can therefore be coupled to the forward oscillator.

Our proposed model contributes two novel features. First, the intermittent nature of crawling
as transitions between runs and pauses (*Figure 3B*). And, second, the peristaltic cycle-phase depen-
dent attenuation of angular motion (*Figure 3D*). By combining these two features, the two bending
behavioral modes termed weathervaning and head-casting are naturally generated via a phasic
coupling between the two oscillators.

Behavioral intermittency

Larval locomotion is intermittent meaning that crawling runs are transiently intermitted by brief
pauses. Transitions between these two behavioral states occur autonomously in stimulus-free
conditions as during free exploration. Traditionally, in the context of movement ecology, intermit-
tency has been studied in the spatial regime by characterizing the distributions of run distances
and turn amplitudes occurring during brief stationary reorientation events (pauses). Power-law
distributed runs in line with Levy-walk theoretical models (*Günther et al., 2016*; *Sims et al., 2019*)
and diffusion-like kinematics have been reported (*Klein et al., 2017*). On the other hand, the turn
amplitude distribution diverges from the original Levy-walk-predicted uniform distribution, as it is
highly skewed towards small amplitudes, even if only significant reorientation events are taken into
account (*Sims et al., 2019*). Moreover, the speed-curvature power-law relationship has been dis-
puted (*Zago et al., 2016*; *Marken and Shaffer, 2017*; *Zago et al., 2017*; *Marken and Shaffer, 2018*).

Regarding the temporal dynamics of intermittency, the duration distribution of these pauses is commonly neglected in traditional Levy-walk literature as they are usually characterised via the amplitude distribution of their resulting turning events. A recent analysis of larval tracks reported that the duration of pauses follows a power-law while that of activity bouts a log-normal distribution (*Sakagiannis et al., 2020*), partly in line with findings in adult-fly studies (*Ueno et al., 2012*; *Reynolds et al., 2015*). In our current study we found log-normal best fits for both pause duration and stridechain length. The disagreement over the pause duration might be attributed to the different timescale assessed in the two studies. Contrary to the high-resolution 180-sec tracks in this study, those analyzed previously lasted at minimum 1024 sec and up to 1 hour, allowing for the detection of longer pauses and imposing the necessity to fit over 4 orders of magnitude bypassing the apparent drop around 10 sec, also seen in Fig.3 of *Sakagiannis et al.* (*2020*).

Concerning computational modeling, the contribution of behavioral intermittency to locomotion has not been adequately appreciated. Candidate generative models can either simply sample statistical distributions (*Sims et al., 2019*) or feature a generative mechanism that yields state transitions. Stochastic state-transitions have been included in a model of larva exploration yielding exponentially distributed epochs of both runs and stationary headcasts (*Davies et al., 2015*). At a mechanistic level, a recent study presented a simple binary-neuron model exhibiting state transitions between power-law and non power-law regimes via self-limiting neuronal avalanches and proposed a plausible underlying mechanism that explains initiation/cessation of crawling (*Sakagiannis et al., 2020*). All these attempts can be considered instantiations of a behavioral intermittency module (*Figure 1B*) controlling cessation and re-initiation of crawling, central for generating epoch transitions. In the present study, we chose stochastic sampling from the empirically fitted parametric distribution models.

Crawl-bend interference

Crawling includes mouth hook motion. Specifically, the first phase of a crawling stride consists of concurrent forward motion of head and tail segments, aided by a 'visceral pistoning' mechanism that generates forward displacement of the gut. Subsequently, the mouth hooks anchor the head to the substrate so that the second phase of peristaltic motion can drag all other segments forward as well, completing the stride (*Heckscher et al., 2012*). With respect to turning, it is still debated whether individual turns should be considered as discrete reorientation events that are temporally non-overlapping with crawling bouts (*Sims et al., 2019*), or whether lateral bending occurs in an oscillatory fashion generating turns both during crawling (weathervaning) and during pauses (headcasts) (*Wystrach et al., 2016*; *Thane et al., 2019*). The latter is supported by detailed eigenshape analysis confirming that larvae rarely crawl straight, rather forward locomotion is always accompanied by continuous small amplitude lateral bending (*Szigeti et al., 2015*). It follows that crawling does not exclude bending, rather the two strictly co-occur.

Crawling and bending partially recruit the same effector neural circuitry and body musculature at least at the level of the thorax. Peristaltic motion during crawling includes sequential symmetric bilateral contraction of all segments while bending occurs due to asymmetric unilateral contraction of the thoracic segments. This partial effector overlap could result in interference between the two processes. Indeed here we report a phase-dependent attenuation of angular velocity (*Figure 3C*). Attenuation reaches a minimum at a specific phase of the cycle, closely preceding the increase of head forward velocity (*Figure 3C:inset*). This coincides with the stride phase when the head stops being anchored to the substrate and is therefore free to move laterally. When applying a phase-dependent Gaussian attenuation kernel on angular velocity we managed to accurately reproduce the empirical relation (*Appendix 2-Figure 1C*).

A reasonable hypothesis would then be that the asymmetric thoracic contraction generating lateral bending becomes easier when the head is not anchored to the substrate therefore during a specific phase interval of the stride cycle. We propose that crawling phasically interferes with lateral bending because of these bodily constraints. A consequence of this hypothesis is that the

amplitude of turns generated during crawl-pauses (headcasts) is larger in comparison to those generated during crawling (weathervaning) because during pauses the crawling interference to lateral bending is lifted. It is this phenomenon that dominates the description of larva exploration as a Levy-walk with non-overlapping straight runs and reorientation events, where weathervaning is neglected (*Günther et al., 2016*; *Sims et al., 2019*). Nevertheless, it has been included in a previous stochastic model of larva exploration, where it has been treated as entirely distinct to headcasts, occurring during crawl-pauses, via the application of differential constraints on both the angular velocity and the resulting turn amplitude (*Davies et al., 2015*). By implementing a coupled-oscillator locomotory model we avoid such dual treatment of headcasts and weathervaning.

Olfactory learning in closed loop behavioral simulations

We have reproduced the results of a basic associative learning experiment in the fruit fly larva (*Schleyer et al., 2018*; *Jürgensen et al., 2024*) by the open-loop simulation of classical conditioning trials and subsequent closed-loop behavioral simulation during the memory retention test for individual virtual larvae (*Figure 6B-D*). This modeling approach can be extended in multiple ways. First, the larva demonstrates a number of additional learning capabilities such as differential conditioning (*Schleyer et al., 2011*, *2015b*, *2018*), extinction learning (*Felsenberg and Waddell, 2019*; *Lesar et al., 2021*), and relief learning (*Saumweber et al., 2018*; *Weiglein et al., 2019*; *Gerber et al., 2014*; *König et al., 2018*). Interfacing neural network simulations with candidate circuit and synaptic mechanisms of plasticity with our behavioral model allows to directly compare virtual and empirical behavioral experiments, both at the level of individual and group assays. Second, while information about odor concentration is provided through active sensing, simultaneous input from a feeder module could activate the dopaminergic reward pathway required for synaptic plasticity in the mushroom body. At the same time, simulation of food uptake and energy expenditure will regulate the agent's energy homeostasis. This would further allow realistic foraging scenarios with food depletion and competition among animals.

Closing the loop from active sensing to associative memory formation and behavioral control requires to synchronize a (spiking) neural network at the adaptive layer with the sensory module (reactive layer), and the locomotory and feeding modules at the basic layer (*Figure 1A*). This will enable the simulation of virtual larvae experiencing spatial and temporal dynamics in a virtual environment or on a robotic platform (*Helgadottir et al., 2013*), and it will allow to test model hypotheses on sensory-motor integration and to infer predictions for experimental interventions such as optogenetic stimulation (*Saumweber et al., 2018*) or genetic manipulation (*Saumweber et al., 2011*; *Michels et al., 2011*; *Widmann et al., 2016*; *Springer and Nawrot, 2021*).

Materials and Methods

We first describe the experimental dataset and the software package used in this study. Then we explicitly describe each of the computational modules that comprise the proposed behavioral architecture. The metrics used throughout the analyses of both real and simulated datasets are described in *Appendix 1*.

Dataset description

The larva-tracking dataset was obtained by M.Schleyer and J. Thoener at the Leipzig Institute of Neurobiology. It consists of 31 experimental groups. Each group of ~ 30 third-instar larvae (Canton S) was placed on agarose-filled Petri dishes of 15 cm diameter with no particular stimuli and video-filmed from above at a framerate of $16\,Hz$ for 3 minutes. During video-tracking, 12 points are detected along the longitudinal axis of each larva. For the present study groups were pooled together in a single population and timepoints of detected collisions have been excluded. A subset consisting of the 200 most complete, uninterrupted tracks was selected. The x-y coordinate timeseries have been filtered with a first-order butterworth low-pass filter with a cutoff frequency

of $2\,Hz$ in order to decrease tracking-related noise but retain the behaviorally relevant crawling frequency of $f_C \simeq 1.4\,Hz$. The effect of inadequate and excessive filtering is illustrated in *Video 3*.

Software package and code availability

All data processing, data analyses and model simulations were performed using our freely available python package *Larvaworld* (https://pypi.org/project/larvaworld/), a behavioral analysis and simulation platform for *Drosophila* larva. In *Larvaworld* simulated and empirical data are treated indistinguishably, meaning that the exact same analysis pipeline and behavioral metrics are applied to both. Concerning modeling, the behavioral architecture we propose here comprises the backbone for the construction, extension, configuration and fitting of behavioral models in *Larvaworld*. Moreover, the intermittent coupled-oscillator model as introduced in the present manuscript is available for simulations alongside other preconfigured models. Extensive documentation can be found at https://larvaworld.readthedocs.io.

Inter-individual variability in virtual populations

The dominant methodological approach in animal behavioral modeling yields generative models that capture the average behavior of a group of animals. Each model therefore strives to generate behavior resembling that of an idealized average animal, neglecting inter-individual variability. To challenge this, we contrast it with a group-level modeling approach that preserves individuality within the population. Here, the generative model is instantiated by a group of non-identical animats, with parameter combinations drawn from a fitted multivariate normal distribution, preserving their pairwise correlations. We first measured a number of endpoint parameters across a population of 200 larvae and fitted a multivariate Gaussian distribution. A subset of three of these parameters, body-length l, crawling frequency f_C, and scaled stride displacement $\mathring{d}_S$ is shown in *Figure 7*. The univariate and bivariate empirical distributions are illustrated in blue, while the bivariate projections are shown in red.

Module definition

The building blocks comprising the behavioral architecture are described as separate modules of defined input and output. The modular architecture only specifies the required placeholders and remains agnostic to the specific module implementations. Nevertheless, in what follows, the specific modular implementation for the proposed coupled-oscillator locomotory model will be described alongside the general module-placeholder definition. Calibration of each module on the empirical dataset and parameter specification is described in detail in *Appendix 2*. The parameters of the final model configuration are shown in *Table 1*.

Bisegmental body

The virtual body is the architecture's interface with the simulated environment. It moves through space under the control of the motor effectors comprising the basic locomotory layer of the architecture, and thereby repositions the sensors introduced in the intermediate reactive layer. For the simplest case of 2D locomotion it is therefore imperative to define at least the forward velocity v and the angular orientation velocity ω. Any locomotory model dynamically generating these two parameters is adequate for an abstract point-body or a single-segment body implementation.

For the proposed coupled-oscillator model we choose a bisegmental body implementation, additionally featuring the bending angle (posture) θ_b between the front and rear segment. Following (*Wystrach et al., 2016*), the body is modeled as a torsional analog of the mass-spring damper model. Torque angularly accelerates an inertia I against angular damping that resists motion and viscoelastic forces that resist deformation (lateral bending). The original mass-spring damper model and its torsional analogue are defined by the equations:

$$m\dot{v} = F_{ext} - zv - kx$$

Figure 7. Individuality: empirical (blue) and fitted (red) parameter distributions. Diagonal: Histogram and kernel density estimates (KDE) for body-length l, crawling frequency f_C and mean scaled displacement per stride $\mathring{d}_S$ across a population of 200 larvae in the experimental dataset. Below: Bivariate projection of 3-dim. KDE outlined contours for each parameter pair. Above: Red ellipses represent the bivariate projections of the 3-dim. fitted Gaussian distributions at 0.5, 1, 2 and 3 standard deviations. In our model this Gaussian is used to sample a parameter set for each individual larva. The blue dots denote the empirically measured parameters.

	Parameter	Symbol	Value	Unit
PHYSICS	torque coefficient	c_T	0.5	sec^{-2}
	angular damping	z	1.0	sec^{-1}
	body spring constant	k	1.0	sec^{-2}
	bend correction coefficient	c_b	1.0	-
BODY	length	l	4.0	mm
	number of body segments	-	2	# segments
TURNER	tonic input	I_T^0	21.43	-
	input range	$[I_T^{min}, I_T^{max}]$	[10. 40.]	-
	time constant	tau_T	0.1	sec
	spike response steepness	n_T	2.37	-
CRAWLER	crawling frequency	f_C	1.42	Hz
	maximum scaled velocity	$\hat{v}_{max}$	0.51	$body-lengths/sec$
	stride distance mean	$\bar{d}_S$	0.24	$body-lengths$
	stride distance std	$\tilde{d}_S$	0.04	$body-lengths$
	max velocity phase	ϕ_C^v	3.49	$rads$
INTERFERENCE	suppression coefficient	c_{CT}^0	0.46	-
	suppression relief coefficient.	c_{CT}^1	0.54	-
	max relief phase	ϕ_C^ω	2.05	$rads$
INTERMITTER	run length distribution	N_R	Lognormal(m=1.4, s=1.15)	-
	run length range	$[N_R^{min}, N_R^{max}]$	[1, 142]	# strides
	pause duration distribution	t_P	Exp(b=1.0)	-
	pause duration range	$[t_P^{min}, t_P^{max}]$	[0.12, 16.0]	sec

Table 1. Locomotory model configuration. The parameters of the calibrated average locomotory model, organized per module.

$$I\dot{\omega} = T_{ext} - Z \cdot \omega - K \cdot \theta_b.$$

545 We introduce simplified coefficients for angular damping $z = {}^Z/_I$ in sec^{-1} and bend deformation
546 resistance $k = {}^K/_I$ in sec^{-2} as the dimensions of inertia $M L^2$ cancel out. Similarly, the external
547 torque-per-unit-of-inertia is now in angular acceleration units $\dot{\omega}_{ext} = {}^{T_{ext}}/_I$. This drive is generated by
548 the turner module and will be described below. Additionally we introduce a crawl-phase dependent
549 suppression of angular motion during crawling described by the coefficient $c_{CT}(t) = c_{CT}(\phi_C)$, which
550 will also be described below.

$$\dot{\omega} = \dot{\omega}_{ext} - k \cdot \theta_b - z \cdot \omega - (1 - c_{CT}) \cdot \omega \tag{1}$$

551 In its original implementation (*Wystrach et al., 2016*) the torsional body model deliberately neglects
552 two aspects of the real turning behavior of the larva. First, there is no distinction between bending
553 ω_b and orientation ω angular velocities. Second, there is no correction of the bending angle θ_b
554 due to forward motion. We tackle the first via the bisegmental body so that ω_b between the front
555 and rear vector is distinct from the front vector's ω. Regarding the second, we introduce a simple
556 linear bending-angle correction as the rear vector is aligned to the front vector's orientation during
557 forward motion, according to the equation:

$$\theta_b = \begin{cases} \check{\theta}_b \cdot (1 - c_b^{max}) & if \quad 0 \le c_b^{max} < 1 \\ 0 & if \quad c_b^{max} \ge 1 \end{cases}$$

$$c_b^{max} = \frac{2 d c_b}{l}$$

558 where d is the linear displacement during a timestep, $\check{\theta}_b$ and θ_b is the bending angle before and
559 after correction, l is the body length and $c_b = 1$ is the bend correction coefficient.
560 Concerning the forward motion, the simple kinematic implementation of the crawler module
561 directly generates forward velocity v without taking into account any biomechanical dynamics.

Crawler module
562
563 The crawler module generates forward velocity v, under a continuous (tonic) activation signal I_C
564 and an additional intermittent initiation/cessation signal generated by the intermittency module.
565 Velocity generation involves three parameters : the larval body-length l, the scaled displacement
566 per stride $\mathring{d}_S$ and the crawling frequency f_C. The latter linearly reflects the tonic input I_C, which is
567 kept constant during the short-duration simulations. For an individual virtual larva $l, \bar{\mathring{d}}_S, \tilde{\mathring{d}}_S$ and f_C
568 are set during initialization, where $\bar{\mathring{d}}_S, \tilde{\mathring{d}}_S$ are the mean and standard deviation of $\mathring{d}_S$.
569 In the present study crawling behavior is modeled as an oscillatory process. Each oscillation
570 generates a cycle of forward velocity v increase-decrease resulting in displacement $l \cdot \mathring{d}_S$ of the
571 larva along the axis of its front-segment, simplistically modeling the result of exactly one peristaltic
572 stride. After termination of a stride a new $\mathring{d}_S$ is sampled. An analytically tractable curve is fitted
573 to the average empirical velocity curve measured during strides by setting all 5 crawler-related
574 parameters to the median empirically measured value (*Appendix 2-Figure 1B*):

$$\mathring{d}_S = N\left(\bar{\mathring{d}}_S, \tilde{\mathring{d}}_S\right) \tag{2}$$

$$v = l \cdot \mathring{d}_S \cdot f_C \cdot \left(\mathring{v}_{max} \cos\left(\phi_C - \phi_C^v\right) + 1\right) \tag{3}$$

575 where ϕ_C is the instantaneous phase of the crawler oscillation iterating from 0 to 2π during an
576 oscillatory cycle, $\mathring{v}_{max}$ is the maximum scaled velocity during the cycle and ϕ_C^v is the phase where
577 the maximum velocity occurs.

 Turner module

 The turner module is defined as the generative model of a torque-like output $A_T(t)$ under a continu-

 uous activation signal $I_T(t)$. The (considered non-dimensional) output is scaled to angular acceler-

 ation by a coefficient c_T in sec^{-2} and applied to the body as external bending drive (**Equation 1**):

$$\dot{\omega}_{ext}(t) = c_T \cdot A_T(t), \tag{4}$$

 which is then applied to the body yielding the instantaneous angular velocity $\omega(t)$ and angular accel-

 eration $\dot{\omega}(t)$ bending the virtual body laterally. It has been previously suggested that an underlying

 oscillatory process drives alternating bending to the left and right side (**Wystrach et al., 2016**). We

 empirically confirmed this by detecting a slow,variable rhythm $f_T \approx 0.4$ on the angular velocity

 timeseries of individual larvae. Based on this observation we adopt an oscillatory approach for

 the alternating lateral bending and extend the lateral oscillator model described in **Wystrach et al.**

 (**2016**). It consists of two mutually inhibiting components each having an excitatory and inhibitory

 neuron (E_L&C_L vs E_R&C_R). The spike-response of a neuron to a cumulative input x is given by

 $R(x,h)$ where h is the half-response threshold. The neuromodulatory activation $I_T(t)$ excites both

 components but also affects the gain $g(I_T)$ and time constant $\tau_H(I_T)$ of an additional adaptation

 component H_x of each neuron. The left and right components quickly settle in antiphase, while

 adaptation ensures that periodic transitions occur. Under the baseline activation I_T^0 the oscillations

 occur at a dominant frequency f_T. Perturbations of this external drive cause transient changes in

 both amplitude and frequency because both , up to transient loss of oscillation. This feature is ex-

 ploited during olfactory modulation. The resulting oscillator activity is defined as the instantaneous

 difference in the firing rates $A_T = (E_L - E_R)$.

$$R(x, h) = \begin{cases} \frac{mx^n}{h^n + x^n} & \text{if } x \geq 0 \\ 0 & \text{if } x < 0 \end{cases} \tag{5}$$

$$g(I_T) = 6 + (0.09 \cdot I_T)^2$$
$$\tau_H(I_T) = \frac{3}{1 + (0.04 \cdot I_T^2)}$$

$$\tau_T \dot{E}_L = -E_L + R(I_T + W_{ee}E_L - W_{ec}E_R, 64 + g(I_T)H_{EL})$$

$$\tau_T \dot{C}_L = -C_L + R(I_T + W_{ce}E_L - W_{cc}C_R, 64 + g(I_T)H_{CL})$$

$$\dot{H}_{EL} = \frac{E_L - H_{EL}}{\tau_H(I_T)}$$

$$\dot{H}_{CL} = \frac{E_L - H_{CL}}{\tau_H(I_T)}.$$

 Calibration of the turner module parameters in order to achieve a resulting angular motion that

 is in agreement to the empirical observations requires concurrent specification of the angular-

 motion related angular damping z, bend deformation resistance k and torque scaling coefficient

 c_T. The calibration process is described in detail in **Appendix 2**.

 Crawler-turner coupling

 We couple the crawler and turner oscillators by imposing a crawler-phase dependent attenuation

 $c_{CT}(\phi_C)$ on the angular velocity ω (**Equation 1**). During the entire stride-cycle ω is scaled by a base-

 line suppression coefficient c_{CT}^0. Our kinematic analysis revealed that this baseline attenuation is

 partially lifted during a phase interval of the stride-cycle. The maximum additional relief is defined

 by the relief coefficient c_{CT}^1. Concerning the mode of transition from c_{CT}^0 to $c_{CT}^0 + c_{CT}^1$ and back

 to c_{CT}^0, we study two implementations, each defined by an additional parameter. In the "square"

 mode there is an acute transition to maximum relief and back to baseline for a specific interval of

the cycle $[\phi_C^{\omega_0}, \phi_C^{\omega_1}]$. In the "phase" mode the transition is described as a Gaussian kernel reaching maximum relief at phase ϕ_C^{ω} when crawl-induced angular attenuation is minimum and the anterior body is maximally free to bend laterally, as shown in the equation:

$$c_{CT} = \begin{cases} c_{CT}^1 \cdot A_{CT} + c_{CT}^0, & \text{during runs} \\ 1, & \text{during pauses} \end{cases} \tag{6}$$

$$A_{CT} = \begin{cases} \exp(-\frac{(\phi_C - \phi_C^{\omega})^2}{2}) & \text{phase mode} \\ \\ 1 \quad \text{if } \phi_C^{\omega_0} \leq \phi_C \leq \phi_C^{\omega_1} \\ & \text{square mode} \\ 0 \quad \text{otherwise} \end{cases} \tag{7}$$

The parameter set is optimized to match the average empirically measured angular motion observed during the stride cycle. The optimization process used is described in *Appendix 2*. The optimal Gaussian kernel is shown in *Figure 3*.

Intermittency module

We define the intermittency module as a placeholder for any model capable of generating transitions between runs and pauses (or equivalently runs and headcasts) (*Figure 1B*). During pauses, the input to the crawler module $I_C = 0$ and therefore forward velocity $v = 0$ while during runs this is unaffected. For the specific study we implement a statistical model where the duration for both the run and pause states is sampled at each initiation from a distribution that has been fitted to the experimental data. In the case of pauses the duration is normally measured in time units. For runs we employ an equivalent metric and therefore measure them as number of consecutive crawling strides (stridechains). The fitted distributions are shown in *Table 1*.

Olfactory sensor

Olfaction is introduced in the intermediate layer of the behavioral architecture allowing chemotactic behavior. The olfactory sensor is located at the front end of the virtual larva therefore any reorientation and/or displacement influences sensory input. As in (*Gomez-Marin et al., 2011*) we assume that olfactory perception A_O relates to changes in odor concentration C according to the Weber-Fechner law, meaning that $\Delta A_O \sim \ln \Delta C$. We further add a decay term that slowly resets A_O back to 0. The rate of change is given in *Equation 8* where $c_O = 1$ is the olfactory decay coefficient, G_i is the gain for odor i and C_i the respective odor concentration:

$$\dot{A}_O = -c_O A_O + \sum_i G_i \cdot \frac{\dot{C}_i}{C_i} \qquad with \qquad -1 \leq A_O \leq 1. \tag{8}$$

Concerning how the perceived olfactory stimulation A_O modulates exploratory behavior we adopt the mechanism proposed in (*Wystrach et al., 2016*) (*Figure 5*). According to this model, the turner activation I_T is perturbed from its baseline value I_T^0 within a suitable range $I_T^{range} = [I_T^{min}, I_T^{max}] = [10, 40]$:

$$I_T = I_T^0 + A_O \left(I_T^{lim} - I_T^0 \right) \qquad where \qquad I_T^{lim} = \begin{cases} I_T^{max} & if \quad A_O \geq 0 \\ I_T^{min} & if \quad A_O \leq 0. \end{cases} \tag{9}$$

Mushroom body module

The underlying spiking network model of the *Drosophila* larva olfactory pathway and mushroom body (MB) implements the model published by *Jürgensen et al.* (*2024*). It consists of conductance-based leaky integrate-and-fire neurons. It encompasses 21 olfactory receptor neurons ($ORNs$) and

the same number of projection neurons (PNs) and local interneurons (LNs) that receive input from the $ORNs$ in a one-to-one manner (*Figure 6C*). LNs form inhibitory connections with the PNs. In the MB, each of the 72 Kenyon cells (KC) receive excitatory input from 2-6 random PNs. Feedback inhibition to the KCs is provided via the single GABAergic anterior paired lateral neuron (APL). All KC provide excitatory input to two MB output neurons ($MBONs$). The two $MBONs$ represent two different types of output neurons that exist in the larval MB and either represent approach or avoidance behavior. Plasticity at the $KC :: MBON$ synapses facilitates the association of odors with reward or punishment (associative learning). $KC :: MBON$ synapses employ a two factor learning rule: Pre-synaptic activation by an odor stimulus (action potential of the KC) triggers an exponentially decaying eligibility trace $e(t)$, which determines the window of opportunity for synaptic change. Additional neuromodulatory input $r(t)$ from one of the two $DANs$ will lead to a reduction in synaptic strength (*Figure 6C*) proportional to $e(t) \cdot r(t)$. The acquired imbalance between the outputs of the two $MBONs$ (behavioral bias) represents the association of odors with rewards/punishments. An initially balanced interaction of excitatory and inhibitory feedback components from $MBONs$ onto both $DANs$ encodes the models' learning history in form of a prediction error. The concept of prediction error suggests that associative learning about stimulus A is proportional to the difference between the reinforcement currently received with stimulus A minus the reinforcement predicted by stimulus A (previously learned association) (*Rescorla, 1972*). In vertebrates an implementation of this error signal has been demonstrated in dopaminergic neurons (*Schultz, 2015*, *2016*) and a similar mechanism is proposed for learning in the insect MB (*Riemensperger et al., 2005*; *Springer and Nawrot, 2021*; *Bennett et al., 2021*; *Jürgensen et al., 2024*). The time-resolved behavioral bias is used to compute gains for odors, which bias motor output of the virtual larvae.

Videos

Here we provide the videos cited in the text. All videos are derived from real or simulated experiments as visualized via the *Larvaworld* software package, after some additional editing. Videos can be played by clicking on the image frame.

Video 1.

Locomotory model for Drosophila larva. The function of the locomotory model in *Figure 1B* is illustrated by gradually integrating its 4 modules (crawler, turner, oscillator-coupling, crawling-intermittency). In each of the 6 videos the implemented modules are shown in the inset.

Video 2.

Free exploration simulation. A population of 25 real (left) or virtual (right) larvae is placed on a dish and left to freely explore. The body of the real larvae has been simplified into 2 segments as described in *Video 4*.

Video 3.

Filter selection. The effect of inadequate or excessive filtering of the empirical larva recordings is illustrated. The left video shows the jittery original recording while the effect of lowpass filtering at cut-off frequencies of 4 Hz, 2 Hz and 0.5 Hz is shown on the rest. Selection of an intermediate 2 Hz cut-off frequency eliminates the unrealistic jitter while preserving the behaviorally relevant crawling frequency.

Video 4.

Bisegmental larva-body simplification. The first video shows the original larva body as recorded by the tracker. 12 points are tracked along its longitudinal axis defining 11 segments while 22 points constitute the body contour. In the second video the body contour is dropped. In the third video an artificial rectangular contour is added for each body segment. In the last video the body-midline is segmented into 2 segments. The absolute head orientation angle θ is preserved while the single bending angle between the 2 segments is defined as $\theta_b = \sum_{i=1}^{5} \theta_i$.

Acknowledgments

This project received funding from the Ministry of Culture and Science of the State of Northrhine Westphalia through the science network 'iBehave' (https://ibehave.nrw/) within the program "Netzwerke 2021", and from the German Research Foundation through the Research Unit 'Structure, Plasticity and Behavioral Function of the *Drosophila* mushroom body' (DFG-FOR 2705, grant no. 365082554, https://www.unigoettingen.de/en/601524.html). PS received a stipendship within the Research Training Group 'Neural Circuit Analysis' (DFG-RTG 1960, grant no. 233886668). We would like to thank Michael Schleyer and Juliane Thoener for providing the tracking dataset and for valuable discussions on the data analyses. We thank Bertram Gerber for valuable discussion and Anna Morozova for substantial support with the generation of the video and figure material.

References

Almeida-Carvalho MJ, Berh D, Braun A, Chen YC, Eichler K, Eschbach C, Fritsch PMJ, Gerber B, Hoyer N, Jiang X, Kleber J, Klämbt C, König C, Louis M, Michels B, Miroschnikow A, Mirth C, Miura D, Niewalda T, Otto N, et al. The Ol1mpiad: Concordance of behavioural faculties of stage 1 and stage 3 Drosophila larvae. J Exp Biol. 2017; 220(13):2452–2475. doi: 10.1242/jeb.156646.

de Belle JS, Heisenberg M. Associative Odor Learning in Drosophila Abolished by Chemical Ablation of Mushroom Bodies. Science (80-). 1994; 263(5147):692–695y. doi: 10.1126/science.8303280.

Bennett JE, Philippides A, Nowotny T. Learning with reinforcement prediction errors in a model of the Drosophila mushroom body. Nature communications. 2021; 12(1):2569. doi: 10.1038/s41467-021-22592-4.

Berni J. Genetic dissection of a regionally differentiated network for exploratory behavior in drosophila larvae. Curr Biol. 2015; 25(10):1319–1326. http://dx.doi.org/10.1016/j.cub.2015.03.023, doi: 10.1016/j.cub.2015.03.023.

Berni J, Pulver SR, Griffith LC, Bate M. Autonomous circuitry for substrate exploration in freely moving drosophila larvae. Curr Biol. 2012; 22(20):1861–1870. http://dx.doi.org/10.1016/j.cub.2012.07.048, doi: 10.1016/j.cub.2012.07.048.

Bicho E. Dynamic approach to behavior-based robotics : design, specification, analysis, simulation and implementation. PhD thesis, University of Minho; 1999.

Bidaye SS, Laturney M, Chang AK, Liu Y, Bockemühl T, Büschges A, Scott K. Two brain pathways initiate distinct forward walking programs in Drosophila. Neuron. 2020; 108(3):469–485. doi: 10.1016/j.neuron.2020.07.032.

Brooks R. A Robust Layered Control System For A Mobile Robot. IEEE J Robot Autom. 1986; RA-2(I):14–23.

Davies A, Louis M, Webb B. A Model of Drosophila Larva Chemotaxis. PLoS Comput Biol. 2015; 11(11):1–24. doi: 10.1371/journal.pcbi.1004606.

Denisov G, Ohyama T, Jovanic T, Zlatic M. Model-based detection and analysis of animal behaviors using signals extracted by automated tracking. BIOSIGNALS 2013 - Proc Int Conf Bio-Inspired Syst Signal Process. 2013; p. 175–181. doi: 10.5220/0004235101750181.

Diegelmann S, Klagges B, Michels B, Schleyer M, Gerber B. Maggot learning and synapsin function. J Exp Biol. 2013; 216(6):939–951. doi: 10.1242/jeb.076208.

Dolan MJ, Belliart-Guérin G, Bates AS, Frechter S, Lampin-Saint-Amaux A, Aso Y, Roberts RJV, Schlegel P, Wong A, Hammad A, Bock D, Rubin GM, Preat T, Plaçais PY, Jefferis GSXE. Communication from Learned to Innate Olfactory Processing Centers Is Required for Memory Retrieval in Drosophila. Neuron. 2018; 100(3):651–668.e8. doi: 10.1016/j.neuron.2018.08.037.

Eschbach C, Fushiki A, Winding M, Schneider-Mizell CM, Shao M, Arruda R, Eichler K, Valdes-Aleman J, Ohyama T, Thum AS, Gerber B, Fetter RD, Truman JW, Litwin-Kumar A, Cardona A, Zlatic M. Recurrent architecture for adaptive regulation of learning in the insect brain. Nat Neurosci. 2020; 23(4):544–555. doi: 10.1038/s41593-020-0607-9.

Eschbach C, Zlatic M. Useful road maps: studying Drosophila larva's central nervous system with the help of connectomics. Curr Opin Neurobiol. 2020; 65:129–137. https://doi.org/10.1016/j.conb.2020.09.008, doi: 10.1016/j.conb.2020.09.008.

Felsenberg J, Waddell S. Switching Gears, Structuring the Right Search Strategy. Neuron. 2019; 102(2):273–275. https://doi.org/10.1016/j.neuron.2019.03.040, doi: 10.1016/j.neuron.2019.03.040.

Feng K, Sen R, Minegishi R, Dübbert M, Bockemühl T, Büschges A, Dickson BJ. Distributed control of motor circuits for backward walking in Drosophila. Nature Communications. 2020 Dec; 11(1):6166. https://www.nature.com/articles/s41467-020-19936-x, doi: 10.1038/s41467-020-19936-x.

Gerber B, Stocker RF. The drosophila larva as a model for studying chemosensation and chemosensory learning: A review. Chem Senses. 2007; 32(1):65–89. doi: 10.1093/chemse/bjl030.

Gerber B, Yarali A, Diegelmann S, Wotjak CT, Pauli P, Fendt M. Pain-relief learning in flies, rats, and man: Basic research and applied perspectives. Learn Mem. 2014; 21(4):232–252. doi: 10.1101/lm.032995.113.

Gerhard S, Andrade I, Fetter RD, Cardona A, Schneider-Mizell CM. Conserved neural circuit structure across drosophila larval development revealed by comparative connectomics. Elife. 2017; 6:1–17. doi: 10.7554/eLife.29089.

Gjorgjieva J, Berni J, Evers JF, Eglen SJ. Neural circuits for peristaltic wave propagation in crawling drosophila larvae: Analysis and modeling. Front Comput Neurosci. 2013; 7(MAR):1–19. doi: 10.3389/fncom.2013.00024.

Gomez-Marin A, Louis M. Active sensation during orientation behavior in the Drosophila larva: More sense than luck. Curr Opin Neurobiol. 2012; 22(2):208–215. http://dx.doi.org/10.1016/j.conb.2011.11.008, doi: 10.1016/j.conb.2011.11.008.

Gomez-Marin A, Stephens GJ, Louis M. Active sampling and decision making in Drosophila chemotaxis. Nat Commun. 2011; 2(1):410–441. http://dx.doi.org/10.1038/ncomms1455, doi: 10.1038/ncomms1455.

Günther MN, Nettesheim G, Shubeita GT. Quantifying and predicting Drosophila larvae crawling phenotypes. Sci Rep. 2016; 6:1–10. http://dx.doi.org/10.1038/srep27972, doi: 10.1038/srep27972.

Heckscher ES, Lockery SR, Doe CQ. Characterization of Drosophila Larval Crawling at the Level of Organism, Segment, and Somatic Body Wall Musculature. J Neurosci. 2012; 32(36):12460–12471. doi: 10.1523/jneurosci.0222-12.2012.

Helgadottir LI, Haenicke J, Landgraf T, Rojas R, Nawrot MP. Conditioned behavior in a robot controlled by a spiking neural network. In: *Int. IEEE/EMBS Conf. Neural Eng. NER*; 2013. doi: 10.1109/NER.2013.6696078.

Imambocus BN, Zhou F, Formozov A, Wittich A, Tenedini FM, Hu C, Sauter K, Macarenhas Varela E, Herédia F, Casimiro AP, Macedo A, Schlegel P, Yang CH, Miguel-Aliaga I, Wiegert JS, Pankratz MJ, Gontijo AM, Cardona A, Soba P. A neuropeptidergic circuit gates selective escape behavior of Drosophila larvae. Current Biology. 2022; 32(1):149–163.e8. doi: 10.1016/j.cub.2021.10.069.

Jovanic T. Studying neural circuits of decision-making in Drosophila larva. J Neurogenet. 2020; 34(1):162–170. doi: 10.1080/01677063.2020.1719407.

Jürgensen AM, Khalili A, Chicca E, Indiveri G, Nawrot MP. A neuromorphic model of olfactory processing and sparse coding in the Drosophila larva brain. Neuromorphic Comput Eng. 2021; doi: https://doi.org/10.1088/2634- 4386/ac3ba6.

Jürgensen AM, Sakagiannis P, Schleyer M, Gerber B, Nawrot MP. Prediction error drives associative learning and conditioned behavior in a spiking model of Drosophila larva. iScience. 2024 Jan; 27(1):108640. https://linkinghub.elsevier.com/retrieve/pii/S2589004223027177, doi: 10.1016/j.isci.2023.108640.

Kafle T, Grub M, Sakagiannis P, Nawrot MP, Arguello R. Fast and recurrent evolution of temperature preference among drosophilids. bioRxiv. 2024 Mar; http://biorxiv.org/lookup/doi/10.1101/2024.03.15.585210, doi: 10.1101/2024.03.15.585210.

Karagyozov D, Mihovilovic Skanata M, Lesar A, Gershow M. Recording Neural Activity in Unrestrained Animals with Three-Dimensional Tracking Two-Photon Microscopy. Cell Reports. 2018 Oct; 25(5):1371–1383.e10. https://linkinghub.elsevier.com/retrieve/pii/S2211124718315766, doi: 10.1016/j.celrep.2018.10.013.

Kim D, Alvarez M, Lechuga LM, Louis M. Species-specific modulation of food-search behavior by respiration and chemosensation in Drosophila larvae. Elife. 2017; 6:1–23. doi: 10.7554/eLife.27057.

Klein M, Krivov SV, Ferrer AJ, Luo L, Samuel AD, Karplus M. Exploratory search during directed navigation in C. elegans and Drosophila larva. Elife. 2017; 6:1–14. doi: 10.7554/elife.30503.

Kohsaka H, Zwart MF, Fushiki A, Fetter RD, Truman JW, Cardona A, Nose A. Regulation of forward and backward locomotion through intersegmental feedback circuits in Drosophila larvae. Nat Commun. 2019; 10(1):2654. http://www.nature.com/articles/s41467-019-10695-y, doi: 10.1038/s41467-019-10695-y.

König C, Khalili A, Ganesan M, Nishu AP, Garza AP, Niewalda T, Gerber B, Aso Y, Yarali A. Reinforcement signaling of punishment versus relief in fruit flies. Learn Mem. 2018; 25(6):247–257. doi: 10.1101/lm.047308.118.

Lahiri S, Shen K, Klein M, Tang A, Kane E, Gershow M, Garrity P, Samuel ADT. Two alternating motor programs drive navigation in Drosophila larva. PLoS One. 2011; 6(8). doi: 10.1371/journal.pone.0023180.

Laurent F, Blanc A, May L, Gándara L, Cocanougher BT, Jones BMW, Hague P, Barré C, Vestergaard CL, Crocker J, Zlatic M, Jovanic T, Masson JB. LarvaTagger: manual and automatic tagging of *Drosophila* larval behaviour. Bioinformatics. 2024 Jul; 40(7):btae441. https://academic.oup.com/bioinformatics/article/doi/10.1093/bioinformatics/btae441/7708398, doi: 10.1093/bioinformatics/btae441.

Lesar A, Tahir J, Wolk J, Gershow M. Switch-like and persistent memory formation in individual Drosophila larvae. Elife. 2021; 10:e70317.

Loveless J, Lagogiannis K, Webb B. Modelling the neuromechanics of exploration in larval Drosophila. PLoS Comput Biol. 2019; 15:7. doi: 10.5281/zenodo.1432637.

Loveless J, Webb B. A Neuromechanical Model of Larval Chemotaxis. Integr Comp Biol. 2018; 58(5):906–914. doi: 10.1093/icb/icy094.

Malloy CA, Somasundaram E, Omar A, Bhutto U, Medley M, Dzubuk N, Cooper RL. Pharmacological identification of cholinergic receptor subtypes: modulation of locomotion and neural circuit excitability in Drosophila larvae. Neuroscience. 2019; 411:47–64. https://doi.org/10.1016/j.neuroscience.2019.05.016, doi: 10.1016/j.neuroscience.2019.05.016.

Mantziaris C, Bockemühl T, Büschges A. Central Pattern Generating Networks in Insect Locomotion. Dev Neurobiol. 2020; 00(October 2019):1–15. doi: 10.1002/dneu.22738.

Marken RS, Shaffer DM. The power law of movement : an example of a behavioral illusion. Exp Brain Res. 2017; 235(6):1835–1842. doi: 10.1007/s00221-017-4939-y.

Marken RS, Shaffer DM. The power law as behavioral illusion: reappraising the reappraisals. Exp Brain Res. 2018; 236(5):1537–1544. http://dx.doi.org/10.1007/s00221-018-5208-4, doi: 10.1007/s00221-018-5208-4.

Michels B, Chen YC, Saumweber T, Mishra D, Tanimoto H, Schmid B, Engmann O, Gerber B. Cellular site and molecular mode of synapsin action in associative learning. Learn Mem. 2011; 18(5):332–344. doi: 10.1101/lm.2101411.

Michels B, Saumweber T, Biernacki R, Thum J, Glasgow RDV, Schleyer M, Chen YC, Eschbach C, Stocker RF, Toshima N, Tanimura T, Louis M, Arias-Gil G, Marescotti M, Benfenati F, Gerber B. Pavlovian conditioning of larval Drosophila: An illustrated, multilingual, hands-on manual for odor-taste associative learning in maggots. Front Behav Neurosci. 2017; 11(April):1–6. doi: 10.3389/fnbeh.2017.00045.

Miroschnikow A, Schlegel P, Schoofs A, Hueckesfeld S, Li F, Schneider-Mizell CM, Fetter RD, Truman JW, Cardona A, Pankratz MJ. Convergence of monosynaptic and polysynaptic sensory paths onto common motor outputs in a *Drosophila* feeding connectome. eLife. 2018 dec; 7:e40247. https://doi.org/10.7554/eLife.40247, doi: 10.7554/eLife.40247.

Miroschnikow A, Schlegel P, Schoofs A, Hueckesfeld S, Li F, Schneider-Mizell CM, Fetter RD, Truman JW, Cardona A, Pankratz MJ. The Feeding Connectome: Convergence of Monosynaptic and Polysynaptic Sensory Paths onto Common Motor Outputs. Elife. 2018; 7:1–23. doi: 10.7554/elife.40247.

Niewalda T, Jeske I, Michels B, Gerber B. 'Peer pressure' in larval Drosophila? Biol Open. 2014; 3(7):575–582. doi: 10.1242/bio.20148458.

Ohyama T, Jovanic T, Denisov G, Dang TC, Hoffmann D, Kerr RA, Zlatic M. High-Throughput Analysis of Stimulus-Evoked Behaviors in Drosophila Larva Reveals Multiple Modality-Specific Escape Strategies. PLoS One. 2013; 8(8). doi: 10.1371/journal.pone.0071706.

Owald D, Felsenberg J, Talbot CB, Das G, Perisse E, Huetteroth W, Waddell S. Activity of defined mushroom body output neurons underlies learned olfactory behavior in Drosophila. Neuron. 2015; 86(2):417–427. http://dx.doi.org/10.1016/j.neuron.2015.03.025, doi: 10.1016/j.neuron.2015.03.025.

Owald D, Waddell S. Olfactory learning skews mushroom body output pathways to steer behavioral choice in Drosophila. Curr Opin Neurobiol. 2015; 35(Figure 1):178–184. http://dx.doi.org/10.1016/j.conb.2015.10.002, doi: 10.1016/j.conb.2015.10.002.

Paisios E, Rjosk A, Pamir E, Schleyer M. Common microbehavioral "footprint" of two distinct classes of conditioned aversion. Learn Mem. 2017; 24(5):191–198. doi: 10.1101/lm.045062.117.

Pehlevan C, Paoletti P, Mahadevan L. Integrative neuromechanics of crawling in D. melanogaster larvae. Elife. 2016; 5:1–23. doi: 10.7554/elife.11031.

Prescott TJ, Redgrave P, Gurney K. Layered control architectures in robots and vertebrates. Adapt Behav. 1999; 7(1):99–127. doi: 10.1177/105971239900700105.

Prescott TJ, Wilson SP. Understanding brain functional architecture through robotics. Science Robotics. 2023 May; 8(78). https://www.science.org/doi/10.1126/scirobotics.adg6014, doi: 10.1126/scirobotics.adg6014, publisher: American Association for the Advancement of Science (AAAS).

Qian CS, Kaplow M, Lee JK, Grueber WB. Diversity of internal sensory neuron axon projection patterns is controlled by the POU-domain protein pdm3 in drosophila larvae. J Neurosci. 2018; 38(8):2081–2093. doi: 10.1523/JNEUROSCI.2125-17.2018.

Rescorla RA. Informational Variables in Pavlovian Conditioning. Psychol Learn Motiv. 1972; 6:1–64.

Reynolds AM, Jones HBC, Hill JK, Pearson AJ, Wilson K, Wolf S, Lim KS, Reynolds DR, Chapman JW, Reynolds AM. Evidence for a pervasive 'idling-mode' activity template in flying and pedestrian insects. R Soc Open Sci. 2015; 2. doi: 10.1098/rsos.150085.

Riemensperger T, Völler T, Stock P, Buchner E, Fiala A. Punishment prediction by dopaminergic neurons in Drosophila. Curr Biol. 2005; 15(21):1953–1960. doi: 10.1016/j.cub.2005.09.042.

Risse B, Berh D, Otto N, Klämbt C, Jiang X. FIMTrack: An open source tracking and locomotion analysis software for small animals. PLoS Comput Biol. 2017; 13(5):1–15. doi: 10.1371/journal.pcbi.1005530.

Ross D, Lagogiannis K, Webb B. A Model of Larval Biomechanics Reveals Exploitable Passive Properties for Efficient Locomotion. In: Wilson SP, Verschure PFMJ, Mura A, Prescott TJ, editors. *Biomimetic and Biohybrid Systems* Cham: Springer International Publishing; 2015. p. 1–12.

Ruiz-Dubreuil G, Burnet B, Connolly K, Furness P. Larval foraging behaviour and competition in Drosophila melanogaster. Heredity (Edinb). 1996; 76(1):55–64. doi: 10.1038/hdy.1996.7.

Sakagiannis P, Aguilera M, Nawrot MP. A Plausible Mechanism for Drosophila Larva Intermittent Behavior. In: Vouloutsi V, Mura A, Tauber F, Speck T, Prescott TJ, Verschure PFMJ, editors. *Biomimetic and Biohybrid Systems* Cham: Springer International Publishing; 2020. p. 288–299.

Saumweber T, Rohwedder A, Schleyer M, Eichler K, Chen Yc, Aso Y, Cardona A, Eschbach C, Kobler O, Voigt A, Durairaja A, Mancini N, Zlatic M, Truman JW, Thum AS, Gerber B. Functional architecture of reward learning in mushroom body extrinsic neurons of larval Drosophila. Nat Commun. 2018; p. 1–19. http://dx.doi.org/10.1038/s41467-018-03130-1, doi: 10.1038/s41467-018-03130-1.

Saumweber T, Weyhersmüller A, Hallermann S, Diegelmann S, Michels B, Bucher D, Funk N, Reisch D, Krohne G, Wegener S, Buchner E, Gerber B. Behavioral and synaptic plasticity are impaired upon lack of the synaptic protein SAP47. J Neurosci. 2011; 31(9):3508–3518. doi: 10.1523/JNEUROSCI.2646-10.2011.

Schleyer M, Fendt M, Schuller S, Gerber B. Associative learning of stimuli paired and unpaired with reinforcement: Evaluating evidence from maggots, flies, bees, and rats. Front Psychol. 2018; 9(AUG):1–15. doi: 10.3389/fpsyg.2018.01494.

Schleyer M, Miura D, Tanimura T, Gerber B. Learning the specific quality of taste reinforcement in larval Drosophila. Elife. 2015; 4. doi: 10.7554/eLife.04711.

Schleyer M, Reid SF, Pamir E, Saumweber T, Paisios E, Davies A, Gerber B, Louis M. The impact of odor-reward memory on chemotaxis in larval Drosophila. Learn Mem. 2015; 22(5):267–277. doi: 10.1101/lm.037978.114.

Schleyer M, Saumweber T, Nahrendorf W, Fischer B, von Alpen D, Pauls D, Thum A, Gerber B. A behavior-based circuit model of how outcome expectations organize learned behavior in larval Drosophila. Learn Mem. 2011; 18(10):639–653. doi: 10.1101/lm.2163411.

Schleyer M, Weiglein A, Thoener J, Strauch M, Hartenstein V, Weigelt MK, Schuller S, Saumweber T, Eichler K, Rohwedder A, Merhof D, Zlatic M, Thum AS, Gerber B. Identification of dopaminergic neurons that can both establish associative memory and acutely terminate its behavioral expression. J Neurosci. 2020; 40(31):5990–6006. doi: 10.1523/JNEUROSCI.0290-20.2020.

Schoofs A, Hückesfeld S, Schlegel P, Miroschnikow A, Peters M, Zeymer M, Spieß R, Chiang AS, Pankratz MJ. Selection of Motor Programs for Suppressing Food Intake and Inducing Locomotion in the Drosophila Brain. PLoS Biol. 2014; 12(6). doi: 10.1371/journal.pbio.1001893.

Schoofs A, Miroschnikow A, Schlegel P, Zinke I, Schneider-Mizell CM, Cardona A, Pankratz MJ. Serotonergic modulation of swallowing in a complete fly vagus nerve connectome. Current Biology. 2024 Sep; p. S0960982224011370. https://linkinghub.elsevier.com/retrieve/pii/S0960982224011370, doi: 10.1016/j.cub.2024.08.025.

Schultz W. Neuronal reward and decision signals: From theories to data. Physiol Rev. 2015; 95(3):853–951. doi: 10.1152/physrev.00023.2014.

Schultz W. Dopamine reward prediction error coding. Dialogues Clin Neurosci. 2016; 18(1):23.

Schulze A, Gomez-Marin A, Rajendran VG, Lott G, Musy M, Ahammad P, Deogade A, Sharpe J, Riedl J, Jarriault D, Trautman ET, Werner C, Venkadesan M, Druckmann S, Jayaraman V, Louis M. Dynamical feature extraction at the sensory periphery guides chemotaxis. Elife. 2015; 4(JUNE):1–52. doi: 10.7554/eLife.06694.

Schumann I, Triphan T. The PEDtracker: An Automatic Staging Approach for Drosophila melanogaster Larvae. Front Behav Neurosci. 2020; 14. doi: 10.3389/fnbeh.2020.612313.

Sen R, Wu M, Branson K, Robie A, Rubin GM, Dickson BJ. Moonwalker Descending Neurons Mediate Visually Evoked Retreat in Drosophila. Current Biology. 2017; 27(5):766–771. http://dx.doi.org/10.1016/j.cub.2017.02.008, doi: 10.1016/j.cub.2017.02.008, publisher: Elsevier Ltd.

Sims DW, Humphries NE, Hu N, Medan V, Berni J. Optimal searching behaviour generated intrinsically by the central pattern generator for locomotion. Elife. 2019; 8:1–31. https://elifesciences.org/articles/50316, doi: 10.7554/eLife.50316.

Slater G, Levy P, Andrew Chan KL, Larsen C. A central neural pathway controlling odor tracking in drosophila. J Neurosci. 2015; 35(5):1831–1848. doi: 10.1523/JNEUROSCI.2331-14.2015.

Springer MA, Nawrot MP. A mechanistic model for reward prediction and extinction learning in the fruit fly. eNeuro - Press. 2021 mar; https://www.eneuro.org/content/early/2021/03/27/ENEURO.0549-20.2021, doi: 10.1523/ENEURO.0549-20.2021.

Strutz A, Soelter J, Baschwitz A, Farhan A, Grabe V, Rybak J, Knaden M, Schmuker M, Hansson BS, Sachse S. Decoding odor quality and intensity in the Drosophila brain. Elife. 2014; 3:1–21. doi: 10.7554/elife.04147.

Szigeti B, Deogade A, Webb B. Searching for motifs in the behaviour of larval Drosophila melanogaster and Caenorhabditis elegans reveals continuity between behavioural states. J R Soc Interface. 2015; 12(113). doi: 10.1098/rsif.2015.0899.

Tadres D, Louis M. PiVR: An affordable and versatile closed-loop platform to study unrestrained sensorimotor behavior. PLoS Biol. 2020; 18(7):1–25. http://dx.doi.org/10.1371/journal.pbio.3000712, doi: 10.1371/journal.pbio.3000712.

Tastekin I, Khandelwal A, Tadres D, Fessner ND, Truman JW, Zlatic M, Cardona A, Louis M. Sensorimotor pathway controlling stopping behavior during chemotaxis in the Drosophila melanogaster larva. Elife. 2018; 7:1–38. doi: 10.7554/elife.38740.

Thane M, Paisios E, Stöter T, Krüger AR, Gläß S, Dahse AK, Scholz N, Gerber B, Lehmann DJ, Schleyer M. High-resolution analysis of individual *Drosophila melanogaster* larvae uncovers individual variability in locomotion and its neurogenetic modulation. Open Biology. 2023 Apr; 13(4):220308. https://royalsocietypublishing.org/doi/10.1098/rsob.220308, doi: 10.1098/rsob.220308.

Thane M, Viswanathan V, Meyer TC, Paisios E, Schleyer M. Modulations of microbehaviour by associative memory strength in Drosophila larvae. PLoS One. 2019; 14(10):e0224154. doi: 10.1371/journal.pone.0224154.

de Tredern E, Manceau D, Blanc A, Sakagiannis P, Barre C, Sus V, Viscido F, Hasan MA, Autran S, Nawrot M, Mason JB, Jovanic T. Feeding-state dependent neuropeptidergic modulation of reciprocally interconnected inhibitory neurons biases sensorimotor decisions in Drosophila. bioRxiv. 2024 Apr; doi: https://doi.org/10.1101/2023.12.26.573306.

Ueno T, Masuda N, Kume S, Kume K. Dopamine Modulates the Rest Period Length without Perturbation of Its Power Law Distribution in Drosophila melanogaster. PLoS One. 2012; 7(2). doi: 10.1371/journal.pone.0032007.

Vogt K, Zimmerman DM, Schlichting M, Hernandez-Nunez L, Qin S, Malacon K, Rosbash M, Pehlevan C, Cardona A, Samuel ADT. Internal state configures olfactory behavior and early sensory processing in Drosophila larvae. Sci Adv. 2021; 7:1, eabd6900. doi: 10.1126/sciadv.abd6900.

Weiglein A, Gerstner F, Mancini N, Schleyer M, Gerber B. One-trial learning in larval Drosophila. Learn Mem. 2019; 26(4):109–120. doi: 10.1101/lm.049106.118.

Widmann A, Artinger M, Biesinger L, Boepple K, Peters C, Schlechter J, Selcho M, Thum AS. Genetic Dissection of Aversive Associative Olfactory Learning and Memory in Drosophila Larvae. PLoS Genet. 2016; 12(10):1–32. doi: 10.1371/journal.pgen.1006378.

Widmann A, Eichler K, Selcho M, Thum AS, Pauls D. Odor-taste learning in Drosophila larvae. J Insect Physiol. 2018; 106(August):47–54. http://dx.doi.org/10.1016/j.jinsphys.2017.08.004, doi: 10.1016/j.jinsphys.2017.08.004.

Wilson SP, Prescott TJ. Scaffolding layered control architectures through constraint closure: insights into brain evolution and development. Philosophical Transactions of the Royal Society B: Biological Sciences. 2022 Feb; 377(1844). https://royalsocietypublishing.org/doi/10.1098/rstb.2020.0519, doi: 10.1098/rstb.2020.0519, publisher: The Royal Society.

Wu Q, Wen T, Lee G, Park JH, Cai HN, Shen P. Developmental control of foraging and social behavior by the Drosophila neuropeptide Y-like system. Neuron. 2003; 39(1):147–161. doi: 10.1016/S0896-6273(03)00396-9.

Wystrach A, Lagogiannis K, Webb B. Continuous lateral oscillations as a core mechanism for taxis in Drosophila larvae. Elife. 2016; 5. doi: 10.7554/elife.15504.

Zago M, Lacquaniti F, Gomez-marin A, Gomez-marin A. The speed - curvature power law in Drosophila larval locomotion. Biol Lett. 2016; 12. doi: https://dx.doi.org/10.6084/m9.fig-share.c.3517350.

Zago M, Matic A, Flash T, Gomez-Marin A, Lacquaniti F. The speed - curvature power law of movements: a reappraisal. Exp Brain Res. 2017; 236:69–82. doi: 10.1007/s00221-017-5108-z.

Zarin AA, Mark B, Cardona A, Litwin-Kumar A, Doe CQ. A multilayer circuit architecture for the generation of
distinct locomotor behaviors in Drosophila. Elife. 2019; 8:1–34. doi: 10.7554/eLife.51781.

Zhang T, Branch A, Shen P. Octopamine-mediated circuit mechanism underlying controlled appetite for palat-
able food in Drosophila. Proc Natl Acad Sci. 2013; 110(38):15431–15436. doi: 10.1073/pnas.1308816110.

Metric definition

Body length

The instantaneous body-length of an individual larva fluctuates during crawling due to sub-
sequent stretching and contraction. Its histogram is well fitted by a Gaussian distribution
(data not shown). Therefore individual larva length l is defined as the median of the midline
length across time (total length of the line connecting all 12 midline points). All spatial pa-
rameters including displacement and velocity can be scaled to this body-length, converting
spatial units m or mm to dimensionless body-length units. Scaled spatial metrics are denoted
by an additional ° over the metric symbol.

Segmentation and angular metrics

To specify the body segmentation providing the most suitable contact/rotation point for
the definition of the bending ω_b and orientation ω angular velocities we analyse their rela-
tionship in a subset of 40 larvae. Tracking of 12 midline points allows computation of the
absolute orientation of 11 body-segments and the respective 10 angles θ_{1-10} between suc-
cessive body segments (**Figure 2F**). We define θ as the head-segment absolute orientation
in reference to the x-axis because this defines the movement orientation of the animal. We
ask how ω results from the bending of the body as this is captured by the 10 angular veloci-
ties ω_{1-10}. The regression analysis depicted in **Figure 1B** shows as expected that ω depends
primarily on the front angular velocities while this dependence decays as we move towards
the rear segments, in line with previous studies (**Lahiri et al., 2011**). Timeshift analysis also
shows that the front 3 angles change concurrently while angles further down the midline
are increasingly lagging behind (data not shown). The correlation analysis depicted in **Fig-**
ure 1C shows that the sum of the front 5 angular velocities best correlates to ω. In other
words the cumulative body bend of the front 5 segments best predicts head reorientation.
Therefore we define the reorientation-relevant bending angle as $\theta_b = \sum_{i=1}^{5} \theta_i$ (**Figure 2F**). The
remaining 5 angles between the rear body-segments can safely be neglected as they do not
contribute to reorientation. This analysis results in a segmentation of the body in a front
and a rear segments of length ratio 5 : 6. The segmentation process is demonstrated in
Video 4.

Forward velocity

To define forward velocity we need to choose which midline-point is most suitable to track
and which velocity metric to use for defining the start and end of a stride. To this end we
perform stride annotation of 3-minute tracks of a population of 20 larvae using each of 24
candidate instantaneous velocity metrics, namely the velocities of the 12 points, the compo-
nent velocities of the rearest 11 points parallel to their front segment's absolute orientation
and finally the centroid velocity. To compare the candidate metrics we compute the spatial
cv_s and temporal cv_t coefficient of variation of the annotated strides for each larva to assess
how variant their time duration and displacement is. We finally compute the mean $\overline{cv}_s$ and
$\overline{cv}_t$ across individuals. In **Figure 1A** the spatiotemporal stride variance is shown for each can-
didate metric. We choose the metric that provides the minimal spatial and temporal stride
variance, assuming that strides of an individual larva are more or less stereotypical in both
duration and displacement (**Heckscher et al., 2012**). Our study reveals that the centroid
velocity is the most suitable metric for stride annotation. All spatial metrics are therefore
computed via this point's displacement.

Appendix 1—figure 1. Segmentation and velocity definition. A: Forward velocity definition. 13 candidate velocity metrics are compared for use in stride annotation of 3-minute tracks of a population of 30 larvae. For each candidate the mean coefficient of variation of temporal duration $\overline{cv}_t$ and spatial displacement $\overline{cv}_s$ of the annotated strides is shown. Midline point 9 velocity provides the most temporally and spatially stereotypical strides, therefore it is selected as the reference forward velocity for stride annotation and model fitting. v_{cen} : centroid velocity, $v_1 - v_{12}$: 1^{st}-12^{th} point's velocity. **B:** Regression analysis of individual and cumulative angular velocities $\omega_{i=1-10}$ to orientation angular velocity ω. When considered individually, ω_2 best predicts reorientation with the ω_3 and ω_1 following. When considered cumulatively the anterior 4 ω_i allow optimal prediction of reorientation velocity. **C:** Correlation analysis of the sum of all possible ω_i combinations to ω. The sum $\sum_{i=1}^{4} \omega_i$ shows the highest correlation therefore we define $\theta_b = \sum_{i=1}^{4} \theta_i$ as shown in A. For illustration purposes only the 5 highest correlations are shown.

Track tortuosity

Track tortuosity is quantified by the straightness index (S.I), a metric previously used in larva-track analysis (**Sims et al., 2019**), computed by advancing a fixed time window along the track and calculating at each point the ratio of the straight line distance to the actual distance travelled. This index, which varies from 0 (no movement) to 1 (straight line movement), can capture very well the complexity of the movement at various scales (set by the window time frame) throughout the track. As the time window decreases the smoothing effect is also reduced revealing increasing track details. Different window sizes have been used ranging from 2 to 20 seconds in order to capture both large-scale spatial trajectories and small-scale local movements. Tortuosity was computed for each larva through time and revealed changes in movement as larva alternated between straight line relocation, changes of direction and different degrees of tortuosity. Hence, a change in the S.I. along a larva's track captures the magnitude of the change in movement pattern from intensive, area-restricted searching movements (higher tortuosity) to extensive, straighter line movements (lower tortuosity), and vice versa, across a wide range of spatial scales.

Epoch annotation

Strides (S) are annotated using the scaled forward velocity $\mathring{v}$. First we apply fourier analysis to detect the dominant crawling frequency within a suitable range $1 \leqslant f_C \leqslant 2.5$ (**Figure 3A**). From this we derive the reference stride duration $t_S = f_C^{-1}$. Then epochs are annotated under a number of constraints (**Figure 2C**):

- Each stride is contained between two $\mathring{v}$ local minima.
- The $\mathring{v}$ local maxima contained in the epoch needs to exceed a threshold : $\mathring{v}_{max} \geqslant 0.3$.
- The duration of the epoch t needs to range within $0.75 \leqslant {}^t S/_{\leqslant 2}$. This allows individual

strides to temporally vary without overlapping so that adjacent strides can be concatenated in stridechains.

After stride annotation the displacement due to each individual stride is computed for each larva and divided by the larva's body-length (*Figure 2A-C*). The individual distributions are well fitted by Gaussians (data not shown). Therefore $\bar{d}_S, \mathring{d}_S$ are defined as the average and standard deviation of the scaled displacement per stride for each larva.

Crawling runs (R) are defined as uninterrupted sequences of successive strides, also termed stridechains (*Figure 2C*). Stridechain length is the number of concatenated strides in a run and is a discrete metric equivalent to the continuous crawling run duration. If the locomotory model under evaluation does not generate v oscillations and therefore strides can not be detected, runs are annotated using the plain forward velocity v timeseries as in a previous study (*Sakagiannis et al., 2020*). We first define a suitable threshold v_{thr} by detecting the minima of the pooled v bimodal distribution. Then we define runs as epochs where constantly $v \geqslant v_{thr}$. Pauses are then defined as epochs containing no strides (or equivalently not overlapping with runs) and during which $\mathring{v} \leq 0.3$ (or $v \leq v_{thr}$).

Turn epochs (T) are contained between pairs of successive sign changes of orientation angular velocity ω. Annotated epochs of left (T_L) and right (T_R) turns yield the respective turn-angle amplitudes θ_{T_L} and θ_{T_R} as the absolute total change of orientation angle θ (*Figure 2G*), which can then be pooled into the overall absolute turn-angle amplitudes θ_T. Notably by this definition turns include both headcasts (H) occurring during crawl-pauses and weathervaning (W) occurring during runs. Turns happening exclusively during runs and pauses yield the turn-angle amplitudes of weathervaning θ_W and headcasts θ_H respectively.

Model calibration

Here we describe in detail the calibration procedure for a locomotory model of an average
idealized individual based on a reference dataset of 150 larvae tracked while freely exploring
a Petri-dish over 3 minutes. The pipeline consists of 3 initial steps, which can be performed
independently of each other and set the configuration parameters of the crawler, intermit-
tency and turner modules respectively. In a final step that builds upon the results of the
initial three, the parameters of the crawl-bend phase-locked interference are set. The first
two steps are directly defined by the kinematic analysis results. The third and the final steps
involve an optimization process to reach the parameter set best fitting the empirical data.
Notably, along with the intrinsic parameters of a module, a number of additional module-
related parameters might need to be defined, such as the body-length for the crawler mod-
ule and the angular-motion relevant physical parameters for the turner module.

1093
Appendix 2—figure 1. Average locomotory model summary. A: Distribution of stride-cycle related
parameters over the empirical dataset. Red dotted line denotes the median value used in the Crawler
module of the average locomotory model. **B:** Normalized average curves of angular metrics during
the stride cycle for each individual larva. Red line denotes the group median. **C:** Pooled distribution of
runs and pauses over the entire dataset (blue). Runs are detected as chains of concatenated strides
(stridechains). Stridechains and pauses generated by the fitted distributions are shown in red. These
distributions are used in the Intermitter module of the average locomotory model. **D:** Sample
simulation of the model with all modules active. The model features additionally bend correction due
to forward motion and crawling phase-coupled suppression of angular motion.

For the hereby described average-larva model no noise is introduced to any module

input or output and no parameter variability is allowed across the population of the generated virtual larvae. The sole source of stochasticity then is due to the random distribution sampling processes operated during the behavioral simulations. For the analysis of the generated datasets the exact same pipeline is used as for the empirical dataset. The configuration of the final locomotory model is illustrated in **Table 1**.

Forward motion

For the Crawler module a set of 5 parameters need to be defined, along with a default body-length. This parameter-set is adequate to dynamically generate realistic forward velocity v oscillations as defined in **Equation 3**. All parameters are set to the median value across the empirical larva group. In **Figure 1A** the empirical distributions of the 5 crawler-module parameters are shown along with the median value selected for the average module.

Crawl intermittency

For the Intermittency module the sampling distributions for the run and pause epochs need to be defined. For both a continuous time-duration distribution is define while for the former an additional discrete distribution of number of concatenated strides per stridechain (run) is computed. In all cases, the pooled distribution of the given epoch across the entire dataset is approximated by power-law, exponential, Levy, log-normal and combined log-normal/power-law distributions. All distributions are truncated within an empirically observed range and are fully defined by their usual arguments. The best-fitting candidates from each class are sorted according to their Kolmogorov-Smirnov distance D_{KS} from the empirical distribution and the one exhibiting the minimum D_{KS} is selected for use in the model's intermittency module (**Table 1**). A validation is shown in **Figure 1D** where the distributions of the generated epochs (blue) match the empirical ones (red).

Angular motion

Realistically calibrating the locomotory model in the angular domain is more demanding. The Turner module only generates a torque-equivalent output while the torsional-spring body model applies both angular damping and restorative force due to the body-bend angle (**Equation 1**). We need to calibrate the 3 involved physics parameters, namely the angular damping z in sec^{-1}, the bend deformation resistance k in sec^{-2} and the torque coefficient c_T in sec^{-2}. Additionally the parameters of the turner module itself need to be specified. Here we study two implementations. The first is the above described neural oscillator, for which we need to define the baseline tonic input I_T^0, the spike-response steepness coefficient n_T and the time constant τ_T. The second is a simple sinusoidal oscillator, for which we need to define the baseline amplitude A_T^0 and frequency f_T. In order to have an empirical reference for isolated angular-only motion while avoiding the effect of crawling on the angular motion and we select the detected pause epochs exceeding 3 seconds and derive the pooled distributions for 3 angular metrics, namely the bending angle θ_b, the angular velocity ω and the angular acceleration $\dot{\omega}$. We run an optimization algorithm aiming to minimize the Kolmogorov-Smirnov distance between the simulated and empirical distributions for these 3 angular metrics. In each iteration a Turner module of a given configuration is simulated for 3 minutes while **Equation 1** is applied to derive θ_b and ω. The obtained optimal turner and physics parameters are shown in **Table 1**.

Crawl-bend interference

For the crawl-phase dependent attenuation of angular motion we study two implementations, each having three parameters that define the attenuation coefficient $c_{CT}(\phi_C)$ during the stride-cycle (**Equation 6**). To achieve this we use a genetic algorithm (GA) optimization process to reach a best-fitting parameter-set for this module. Although optimization is only

allowed to search the parameter space for these three parameters, each configuration is evaluated using a complete virtual larva model, combined with the optimal configurations of all other modules defined in the previous three calibration steps. During each iteration of the GA process a group of 30 model configurations is simulated for 3 minutes in free exploration conditions. The best 6 configurations are then mutated and/or combined in a novel generation. The evaluation function minimized is two-fold.Firstly the distributions of the three angular metrics derived from the simulated track, namely the bending angle θ_b, the angular velocity ω and the angular acceleration $\dot{\omega}$ are compared to the empirically measured target distributions via the Kolmogorov-Smirnov distance as described above for the angular motion. Secondly, stride-cycle analysis is carried out for each simulated larva. Detected strides are interpolated in over $n = 64$ bins in a 2π cycle and direction-normalized by inverting the right-turning strides. The average stride-cycle curve $\hat{x}$ is then computed for each of the three metrics and compared to the average empirically measured curve x via the evaluation metric defined below. The metric has been selected so that the different scales of angles, angular velocities and accelerations are normalized before being summed. The final evaluation metric to be minimized is defined as the cumulative error across the three angular metrics for both the distributions and the average stride-cycle curves.

$$Error(x, \hat{x}) = \sqrt{\frac{1}{n} \sum_{i=1}^{n} \frac{(x_i - \hat{x}_i)^2}{|\max(x) - \min(x)|^2}} \tag{10}$$

Here we study two configurations for the crawl phase-dependent suppression of the angular motion, namely a smooth gaussian relief curve and an acute square transition as described in *Equation 6*.The target empirical and optimal model's average curve for the angular velocity ω are shown in *Figure 1B*.

A sample simulation of the complete model is shown in *Figure 1D*. The Crawler-generated forward velocity oscillations (1st row) attenuate the angular motion in a phase-locked manner (2nd row) resulting in low angular velocity during runs and acute headcasts during pauses (3rd row). The bending angle is additionally restored to 0 during forward motion (4th row).