Title: Mutualism-enhancing mutations dominate early adaptation in a microbial community

Authors: Sandeep Venkataram[1], Huan-Yu Kuo[1,2], Erik F. Y. Hom[3], Sergey Kryazhimskiy[1,*]

Affiliations:

[1]Division of Biological Sciences, University of California San Diego, La Jolla, CA 92093;

[2]Department of Physics, University of California San Diego, La Jolla, CA 92093;

[3]Department of Biology and Center for Biodiversity and Conservation Research, University of
Mississippi, University, MS 38677

[*]Corresponding author: skryazhi@ucsd.edu

Abstract:

Species interactions drive species evolution while evolution shapes these interactions. The
resulting eco-evolutionary dynamics, their outcomes and their repeatability depend on how
adaptive mutations available to community members affect fitness and ecologically relevant
traits. However, the diversity of adaptive mutations available to community members is not well
characterized, and we do not know how this diversity is affected by the ecological milieu. Here
we use barcode lineage tracking to address this gap in a competitive mutualism between the
yeast *Saccharomyces cerevisiae* and the alga *Chlamydomonas reinhardtii*. We find that yeast has
access to many adaptive mutations with diverse ecological consequences, in particular, those that
increase and reduce the yields of both species. The presence of alga does not change which
mutations are adaptive in yeast (i.e., there is no fitness trade-off for yeast between growing alone
or with alga), but rather shifts selection to favor yeast mutants that increase the yields of both
species and make the mutualism stronger. Thus, in the presence of alga, we find that yeast
repeatably evolves to become more cooperative, even though cooperativity is not directly
favored by natural selection in our system. Our results demonstrate that ecological interactions
not only alter the trajectory of evolution but also dictate its repeatability; in particular, weak
mutualisms can repeatably evolve to become stronger.

Introduction

Ecological communities are constantly subjected to perturbations by environmental shifts[1,2],
demographic noise[3] and species turnover[4,5]. Such perturbations can not only displace
communities from their ecological equilibria but also precipitate adaptive evolution[6–9]. When
evolution is rapid, ecological and evolutionary dynamics may become coupled, potentially
propelling communities towards otherwise inaccessible ecological attractors[10–12]. Although rapid
eco-evolutionary dynamics have been identified in many communities[7,9,10,13–26], the population
genetics of eco-evolutionary coupling is poorly understood. As a result, we do not know how and
under what conditions such coupling occurs, how repeatable the ensuing eco-evolutionary
dynamics are and what role these dynamics play in the evolution and maintenance of
biodiversity[27–29].

Previous studies have shown that the coupling between ecology and evolution is often driven by
interactions between species, in the sense that the evolutionary trajectory of one community-
member population is affected by its interactions with other members, which in turn feeds back
to alter the ecology of the community[10,15–17,20,23,30]. However, many empirical questions
regarding the population genetic mechanisms of such coupling remain unresolved[29]. How does
the spectrum of adaptive mutations available to a species depend on the composition of the
surrounding community? For example, how does it change when another species is lost from the
community or a third species invades it? How many and which of the adaptive mutations
available to a species affect its interactions with the rest of the community? Which of these
mutations are likely to spread and fix? And thus, how diverse and repeatable are the ecological
outcomes of evolution?

Empirical data supporting answers to these fundamental questions are important for a proper
theoretical understanding of eco-evolutionary dynamics. In particular, it is critical that models
are based on realistic assumptions, especially if the assumptions can qualitatively alter model
behavior. For example, many eco-evolutionary models assume that any combination of traits can
be produced by mutations so that the eco-evolutionary trajectories and outcome are determined
exclusively by natural selection[31]. But recent evidence shows that the availability of mutations
can significantly impact evolution[32–36]. Yet, we know very little about the distributions of
ecological and fitness effects of new mutations in multi-species communities and how these
distributions shift when the ecological milieu changes—for example, due to the addition or
extinction of community members.

Here, we set out to address these gaps in one of the simplest experimentally tractable microbial
communities. Our community consists of two species, the alga *Chlamydomonas reinhardtii* and
the yeast *Saccharomyces cerevisiae*[37], that interact in our environment via competition and
mutualism. Although communities in nature often contain more members, understanding eco-
evolutionary dynamics in simple model communities is helpful in developing an intuition and
expectations for the behaviors of more complex ecosystems[38–40]. We measure how adaptive
mutations in our community affect competitive fitness (a metric that determines the evolutionary
success of a mutant lineage) and species abundances (a metric that informs us about the type of
interactions between species and the stability of the community), depending on the ecological
milieu. Specifically, we focus on mutations available to one member of our community, the
yeast, and ask whether and how the presence of alga alters the spectrum of these mutations, their

effects on competitive fitness and on the abundances of both species in the community. To obtain
a statistical characterization of adaptive mutations available to yeast, we use barcode lineage
tracking (BLT) technology[41,42], which allows us to isolate hundreds of adaptive mutations that
arise in yeast when it evolves alone or in community with the alga. We characterize these two
sets of mutations at the genetic level and measure their effects on individual fitness, basic life-
history traits (growth rates and carrying capacities) and on the abundances of both species in the
yeast-alga community. These observations offer us a detailed view on how inter-species
interactions can affect the evolutionary dynamics of new mutations, and how these mutations in
turn alter the ecology of the community.

Results

Yeast and alga form a facultative competitive mutualism

In a previous study, Hom and Murray showed that in a sealed environment in which nitrite is
provided as a sole source of nitrogen and glucose is provided as a sole source of organic carbon,
the yeast *Saccharomyces cerevisiae* and the alga *Chlamydomonas reinhardtii* spontaneously
form an obligate mutualism[37]. Under such conditions, *C. reinhardtii* consumes nitrite and
produces ammonium that is secreted and utilized by *S. cerevisiae*, which consumes glucose and
produces CO_2 that is in turn utilized by the alga. If the environment is open to ambient gas
exchange and if some ammonium is supplied in the medium, this mutualism becomes facultative
in the sense that both yeast and alga can survive without each other although both can benefit
from each other's presence, as we demonstrate below.

We grow yeast and alga alone or together over multiple 5-day growth and dilution cycles in a
medium supplemented with 0.5 mM of ammonium under well-mixed and well-lit conditions
(Supplementary Information). Yeast grown alone exhausts the initial supply of ammonium after
about 48 hours and thereafter gradually loses viability due to starvation (Figure 1). When yeast
and alga grow together in the same environment, they initially compete, likely for ammonium,
which can be seen by the reduction of the maximum yeast cell density at around 48 hours (Figure
1). Upon the depletion of supplemented ammonium, the alga subsequently reduces nitrite to
ammonium that it secretes[37]; this nitrogen provisioning by the alga allows yeast to persist
without experiencing a detectable viability loss (Figure 1). The alga also grows more robustly in
the community compared to growth alone because of the dissolved CO_2 supplied by yeast. Thus,
the ecological interactions between yeast and alga are complex and multi-phasic, exhibiting clear
signatures of both competition and mutualism, as it is common for mutualistic communities[43].
We thus refer to this system as a facultative "competitive mutualism".

As a result of growth in the presence of both competitive and mutualistic interactions, the final
densities of both yeast and alga at the end of the 5-day cycle, which we refer to as "yields", are
greater in the community than when each species is grown alone (Figure 1). Specifically, yeast
yield in community (YYC) is 24% percent higher than its yield alone (YYA), and algal yield in
community (AYC) is 288% percent higher than its yield alone (AYA), although these differences
are not statistically significant (Figure 1; Student's t-test, YYA vs YYC $t = 1.26$, $df = 2$, $P =$
0.336, 95% Confidence Interval (CI) [–59%, 107%]; AYA vs AYC $t = 3.84$, $df = 2$, $P = 0.06$,
95% CI [–34%, 611%]). We quantify the relative strength of competition and cooperation in our
community by the ratios of YYC to YYA and AYC to AYA, so that both ratios exceeding unity

indicate that mutualism is stronger than competition and vice versa. By this metric, in our
conditions mutualism and competition between yeast and alga roughly balance each other out,
with a slight predominance of the mutualistic interaction.

The fact that the interaction between yeast and alga is non-obligate allows us to compare the
evolution of each species alone relative to their evolution in the community under otherwise
identical initial conditions, and to understand how species evolution is affected by ecological
interactions. We focus here on the adaptive evolution in one community member, the yeast,
because it is likely to adapt faster than the alga (see Supplementary Information). Specifically,
we characterize the distribution of ecological and fitness effects of individual adaptive mutations
available to yeast and examine how these distributions depend on the presence of alga in the
environment, as we describe next.

Figure 1. Growth of ancestral yeast and alga. Growth of ancestral yeast (yellow) and alga (green)
alone (dashed lines) and in a community (solid lines) over the 5-day growth cycle. Error bars show ±1
standard error of the mean calculated from measurements of two biological replicates. Numbers next to
the gray brackets represent *P*-values for the statistical comparison of the abundances alone versus in
community. These *P*-values are obtained from two-sample t-tests.

The presence of alga alters the fitness effects of beneficial mutations in yeast without apparent trade-offs

We first asked whether and how ecological interactions with alga change the distribution of
fitness effects of adaptive mutations available to yeast. To this end, we carried out five replicate
BLT experiments in yeast evolving alone (the "A-condition") and in yeast evolving in a
community with alga (the "C-condition"; Supplementary Information). In each population, we

tracked the frequencies of ~5×10^5 neutral DNA barcodes integrated into the yeast genome[44] for
17 growth and dilution cycles. We identified a total of 14,098 and 14,523 adapted barcode
lineages across all replicates in A- and C-conditions, respectively (Supplementary Information;
Figures S1–S4). The similarity of these numbers suggests that the presence of alga does not
dramatically change the rate at which beneficial mutations arise in yeast.

Based on prior work[41,42], each adapted lineage is expected to initially carry a single beneficial
driver mutation. Thus, by tracking barcode frequencies, we can estimate the competitive fitness
benefits of many simultaneously segregating driver mutations, relative to the ancestral yeast
strain (Supplementary Information). The estimated distributions of fitness effects of beneficial
mutations (bDFEs) were much broader than expected from measurement noise alone in both A-
and C-conditions (Supplementary Information), indicating that yeast has access to multiple
mutations with different fitness benefits, consistent with previous work measuring bDFEs[41,45].
The bDFEs in A- and C-conditions were different in both median effect and breadth (Figure 2A).
Specifically, ecological interactions with alga reduced the bDFE median (1.60 in A vs. 1.50 in C;
$P < 10^{-4}$, two-sided permutation test, see Supplementary Information) but increased its width
(interquartile range (IQR) = 0.31 in A vs. IQR = 0.37 in C; $P < 10^{-4}$, two-sided permutation test).
This increase in width was associated with the appearance of two peaks with higher relative
fitness values around 2.0 and 2.5 (Figure 2A). Since the dynamics of adaptation depend on the
shape of the bDFE[46,47], these results indicate that the presence of alga alters evolutionary
dynamics in the yeast population.

The presence of alga can alter the yeast bDFE by changing which mutations are beneficial (i.e.,
by imposing a fitness trade-off relative to the A-condition) or by changing the fitness benefits
provided by adaptive mutations, or both. To discriminate between these possibilities, we
randomly sampled 221 yeast clones from distinct adapted lineages in the A-condition ("A-
mutants") and 189 yeast clones from distinct adapted lineages in the C-condition ("C-mutants").
Clones were sampled at cycle nine, a time point at which most adapted lineages are expected to
still be driven by a single beneficial mutation (Supplementary Information). We then used
competition assays to measure the fitness of all A- and C-mutants in both A- and C-conditions
relative to the ancestral yeast background (Supplementary Information; Figures S5–S7; Data S1).
These direct fitness measurements are concordant with estimates from the BLT experiment (see
Supplementary Information; Figure S8). We found that C-mutants have significantly higher
fitness in their "home" C-condition than A-mutants (by on average 10.8%, 95% CI [7.7%,
14.0%], $P = 1\times10^{-11}$, ANOVA testing the effect of the evolution treatment and fitness
measurement replicate, $F = 46.98$, $df = 1$) and that A-mutants are more fit in their home A-
condition than C-mutants (by on average 14.4%, 95% CI [10.2%, 18.5%], $P = 2\times10^{-11}$; $F = $
46.25, $df = 1$), consistent with a signature of local adaptation. Interactions with alga significantly
altered the fitness of 88% (362/410) of A- and C-mutants (FDR = 11% by permutation).
However, fitness was positively correlated between the two conditions across all mutants (Figure
2B; Pearson $R = 0.70$, 95% CI [0.65, 0.75], two-sided $P < 10^{-10}$, $t = 19.9$, $df = 408$). Importantly,
none of the A- or C-mutations were deleterious in their "non-home" condition. Thus, the
presence of alga changes the fitness benefits provided by adaptive mutations but does not impose
a measurable fitness trade-off in the sense that it does not alter which mutations are beneficial.

Figure 2. Ecological interactions with alga alter adaptation in yeast at fitness and genetic levels.
(A) Yeast bDFE when evolving alone (top) and in the community with alga (bottom). Each histogram is constructed based on five replicates of the BLT experiment. Dashed blue lines show the outlines of the bDFE in the other condition. Each colored point indicates the average fitness of a mutant carrying a mutation at the indicated adaptive locus (same data as in panel B, see Supplementary Information for details). Large copy-number variants are referred to as chr$x$-yn, where x is the chromosome number and y is the number of copies. **(B)** Fitness of A- and C-mutants measured in competition assays in the A- and C-conditions. Mutants carrying mutations at the 7 most common driver loci are colored. Error bars show ±1 standard error. The solid line shows y=x. **(C)** Distribution of adaptive mutations among A- and C-mutants. The same 7 most common mutation classes are shown explicitly as in panels A and B, all other classes are grouped into "Other" (see Data S3 for full distributions). Adaptive mutations that are expected to be present in our mutants but are not identified in the genome data are labeled as "Unknown" (see Supplementary Information for details). The number of times each locus was independently mutated is given in parentheses.

The presence of alga alters the distribution of mutations contending for fixation in yeast

Given that all sampled yeast mutants are beneficial in the presence and absence of alga and that mutant fitness is correlated between the two conditions, we expected that the sets of A- and C-mutants would be genetically indistinguishable. To test this expectation, we sequenced the genomes of 181 out of 189 C-mutants, 215 out of 221 A-mutants, as well as 24 ancestral isolates as controls (Supplementary Information; Figures S9–S12; Data S2). We found 176 large chromosomal duplications awoulnd deletions across 14 loci in the genomes of A- and C-mutants and none in the ancestral isolates. All of these large mutations are likely adaptive (see Supplementary Information), with a typical A- and C-mutant carrying on average 0.39 ± 0.04 and 0.51 ± 0.04 of these mutations, respectively. We also discovered 185 small indels/point mutations at 63 loci for which mutations are found in A- and C-mutants significantly more often than expected by chance (see Supplementary Information), suggesting that these small mutations are also adaptive. A typical A- and C-mutant carries on average 0.35 ± 0.03 and 0.60 ± 0.06 of these small adaptive mutations, respectively. Overall, we identified 361 beneficial mutations across 77 loci in 250 out of 396 adapted mutants, with each A- and C-mutant carrying on average 0.74 ± 0.04 and 1.11 ± 0.07 such mutations, respectively (Table S1, Data S2).

We quantified the diversity of adaptive mutations carried by A- and C-mutants with the probability of genetic parallelism, P_g, which is the probability that two random clones share a mutation at the same driver locus (lower P_g values imply higher diversity). We found that P_g is slightly higher among A-mutants than among C-mutants ($P_g = 6.0 \pm 0.6\%$ and $8.5 \pm 0.8\%$, respectively), although this difference was not statistically significant ($P = 0.06$, two-sided permutation test; see Supplementary Information). Nevertheless, there were large differences in the frequency distribution of driver mutations among A- and C-mutants (Figures 2C and S13, Data S3; $P = 5 \times 10^{-8}$, two-sided χ^2-test, $\chi^2 = 160.51$, $df = 76$), suggesting that the chance for any given beneficial mutation to rise to a high enough frequency and be sampled varied dramatically between A- and C-conditions. The starkest difference was observed for chromosome XIV: 10% (18/181) of C-mutants carried a whole chromosome amplification (chrXIV-3n) but none of the 215 A-mutants did (Figures 2C and S13), despite chrXIV-3n mutations being beneficial in both conditions (Figure 2A and B).

This initially puzzling observation could be explained by the dynamics of rapid adaptation in populations with limited recombination. In such populations, clonal interference prevents weak or rare beneficial mutations from reaching even moderately high frequencies[36,48,49]. Thus, the same beneficial mutation can have dramatically different chances of being sampled in the two conditions if it is located in different parts of the respective bDFEs. Consistent with this explanation, we find that a typical chrXIV-3n mutant is 2.05 ± 0.04 times more fit per cycle than the ancestral wild-type and ranks in the top $11\% \pm 1.3\%$ of most fit mutants in the C-condition, while being only 1.59 ± 0.03 times more fit than the ancestor and ranking at $53\% \pm 5.3\%$ in the A-condition (Figure 2A). Other mutations with strong discrepancies in their representation among A- and C-mutants show similar shifts in their fitness effects and rank order (Figure 2A, Data S3). In general, interactions with the alga shifted the fitness ranks of mutants between conditions by $14.1\% \pm 0.8\%$ on average. Using evolutionary simulations, we confirmed that such differences in fitness effect and rank are sufficient to explain the observed genetic differences between A- and C-mutants (see Supplementary Information; Figures S14, S15, Table S2).

The fact that the sets of A- and C-mutants are genetically different implies that the yeast
populations in these two conditions embark on distinct evolutionary trajectories. Indeed, A- and
C-mutants primarily represent high-frequency lineages in the respective populations. Therefore,
the mutations that they carry are more likely to win the clonal competition towards fixation in the
condition from which they were sampled compared to other mutations. Then the fact that A- and
C-mutants carry statistically distinct sets of mutations implies that mutations contending for
fixation in yeast in the A- versus C-conditions are also different. In other words, by altering the
fitness benefits of mutations, the ecological interactions with alga change the evolutionary
trajectory of yeast, at least over the short-term.

Adaptive mutations in yeast have diverse ecological consequences for the community

A mutation that spreads and fixes in the yeast population could subsequently alter the ecological
dynamics of the yeast-algal community; and adaptive mutations at different loci may have
different ecological consequences. For example, some mutations could increase yeast's
competitive ability and ultimately lead to the exclusion of the alga from the community. Others
could increase yeast's cooperativity and thereby strengthen the mutualism. To assess the
prevalence and the magnitude of different ecological effects of adaptive mutations in yeast, we
selected 28 C-mutants and 31 A-mutants that are representative of the genetic diversity of
contending mutations (Supplementary Information). We formed 59 "mutant communities" by
culturing each of these yeast mutants with the ancestral strain of the alga. We then quantified the
ecological effect of each mutation by measuring YYC (yeast yield in community) and AYC
(algal yield in community), as well as YYA (yeast yield alone; Supplementary Information,
Figure S16). As before, we define yield as the species density at the end of the 5-day growth
cycle. We focused on yields for two reasons. Yield is a measure of absolute species abundance
which determines the robustness of our community to demographic fluctuations: communities
with higher yields of both species are more ecologically stable[3]. In addition, as discussed above,
the ratio of YYC to YYA as well as the ratio of AYC to AYA (algal yield alone) inform us about
the relative strength of competition versus cooperation in our communities.

We found that many of the adaptive mutations significantly affected both YYA and YYC (Figure
3A, S16). The majority of tested mutations, 58% (34/59), significantly decreased YYA while
15% (9/59) of them significantly increased it (two-sided FDR = 0.18, see Supplementary
Information). At the same time, we found no mutations that significantly decreased YYC, but
39% (23/59) of mutations significantly increased it (FDR = 0.07). Mutations had uncorrelated
effects on YYA and YYC (Figure 3A), suggesting yeast yield in the two conditions is
determined by different underlying traits. Mutations in yeast also significantly altered AYC, with
58% (34/59) mutations increasing it and 17% (10/59) decreasing it (Figures 3B and S16, FDR =
0.03).

The fact that mutations alter the yields of both species in the community suggests that they may
also alter the relative strength of cooperation and competition between yeast and alga. As before,
we measured this strength by the ratios of YYC to YYA and AYC to AYA. We found that 5%
(3/59) of the tested adaptive yeast mutants significantly decreased the AYC to AYA ratio while
increasing the YYC to YYA ratio; while 44% (26/59) of the mutants significantly increased both
the YYC to YYA and the AYC to AYA ratios (two-tailed FDR = 0.36, Figure 3C). The former

two mutants thus weaken the yeast-algal mutualism, at least for one of the partners, and the latter
26 mutants significantly enhance the mutualism.

These results show that yeast has access to beneficial mutations with ecologically diverse
consequences and suggest that our community has the potential to embark on a variety of eco-
evolutionary trajectories with possibly different ecological outcomes.

Figure 3. Adaptive mutations in yeast have diverse ecological consequences. (A) Pearson correlation between
yeast yield in community (YYC) and yeast yield alone (YYA), normalized by ancestral values, across sampled
mutants. **(B)** Pearson correlation between yeast yield in community (YYC) and algal yield in community (AYC),
normalized by ancestral values, across sampled mutants. AYC values were estimated using fluorescence
measurements and then adjusted to cell densities (see Supplementary Information). **(C)** Ratio of YYC to YYA
versus ratio of AYC to AYA (algal yield alone), across sampled mutants. Darker shades of green indicate stronger
mutualism. In all panels, error bars represent ±1 standard error of the mean. In panels A and B, *P*-values are
obtained by a permutation test (see Supplementary Information) and solid lines are fitted by linear regression.

Mutations favored by selection in the presence of alga strengthen the mutualism

Given that adaptive mutations in yeast have a variety of ecological consequences and that yeast
populations in the absence or presence of alga likely fix different sets of adaptive mutations, we
next asked whether mutations that contend for fixation in A- or C-conditions might have
systematically different ecological effects.

We first noticed that, in comparison to the A-mutants, the C-mutants clustered in the top right
corner of the YYC versus AYC plot in Figure 3B. A formal statistical test confirmed that YYC
and AYC values for the C-mutant communities were on average 103% (95% CI [62%, 143%]; *P*
= 3×10^{-4}, permutations of two-sided Welch's t-test, n_1 = 28 C-mutants and n_2 = 31 A-mutants)
and 57% (95% CI [26%, 90%]; *P* = 4×10^{-3}) higher than those for the A-mutant communities,
respectively. These differences remained large and significant even after accounting for the
frequencies with which different driver mutations were observed in our yeast populations (YYC
P = 10^{-5}; AYC *P* = 4×10^{-3}, permutations of weighted two-sided Welch's t-test, Figure S17, see
Supplementary Information), indicating that this trend is not an accidental byproduct of our
selection of A- and C-mutants.

To determine whether the observed differences in yield between A- and C-mutants were driven
by natural selection, we calculated the correlation between these yields and competitive fitness in
both A- and C-conditions (Figure 4 and S18). We found that both YYC and AYC correlated
strongly and significantly with fitness in the C-condition, but only weakly (and in case of YYC,
not significantly) with fitness in the A-condition. This observation implies that natural selection
in the C-condition, but not in the A-condition, favors yeast mutations that lead to yield gains in
community with the alga.

Figure 4. Relationship between yield and fitness. (A) Pearson correlation between yeast yield in community
(YYC), normalized to ancestral values, and competitive fitness in community, across sampled mutants. (**B**) Pearson
correlation between yeast yield in community (YYC), normalized to ancestral values, and competitive fitness alone,
across sampled mutants. In both panels, error bars show ±1 standard error. Solid lines are fitted by linear regression.

A mutation in yeast that increases YYC may concomitantly lead to either a gain or a loss of
YYA (Figure 3A). Therefore, such a mutation may either increase or decrease the mutualistic
benefit that yeast derives from its interactions with the alga. We found that the YYC to YYA
ratio was significantly increased relative to the ancestor for 86% (24/28) of C-mutants (FDR =
49%) but only for 58% (18/31) of A-mutants (FDR = 53%; Figure 3C). Likewise, the AYC to
AYA ratio was increased relative to the ancestor for 68% (19/28) of C-mutants (FDR = 61%) but
not for any A-mutants. As a result, 18 C-mutants and eight A-mutants significantly increased
both the YYC to YYA and AYC to AYA ratios (FDR = 35%). Meanwhile, three A-mutants, but
only one C-mutant significantly decreased the YYC to YYA ratio. Thus, C-mutants both benefit
more often from the presence of alga and reciprocally more often provide benefits to the alga
compared to the A-mutants. In other words, yeast is more likely to become a more cooperative
mutualist when it evolves in the presence of the alga than in its absence.

An interesting potential consequence of this shift in selection on yeast precipitated by the
presence of the alga is that it can change the repeatability of yeast evolution along the
competition-mutualism continuum. We quantified such ecological repeatability by the
probability that two randomly drawn yeast mutants both increase or both decrease the YYC to
YYA ratio and simultaneously both increase or both decrease the AYC to AYA ratio. The
probability of ecological parallelism would be 25% under a uniform null model. Instead, we find
that it is $33 \pm 2.8\%$ for A-mutants ($P = 0.21$, two-sided χ^2-test, $\chi^2 = 1.55$, $df = 1$) and $68 \pm 5.5\%$
for C-mutants ($P = 6 \times 10^{-8}$, $\chi^2 = 29.3$, $df = 1$), and this difference is statistically significant ($P =$

0.01, two-sided permutation test; see Supplementary Information). Thus, yeast evolves more
repeatably (towards stronger mutualism) in the presence of the alga than in its absence.

In summary, our results show that mutations contending for fixation in yeast populations
evolving alone have diverse effects on the ecology of the yeast-algal community, with some
strengthening and some weakening the mutualism. In contrast, mutations contending for fixation
in yeast evolving in the presence of the alga predominantly strengthen the yeast-algal mutualism
making evolution more repeatable at the ecological level.

Mutualism enhancement is likely a byproduct of selection for other yeast life-history traits

We next asked how stronger mutualism could possibly evolve in our community. Specifically,
does natural selection in the presence of alga favor increased cooperativity in yeast directly or is
cooperativity a byproduct of selection for other traits[26,50]? Natural selection can directly favor
rare mutualism-enhancing (i.e., more cooperative) yeast mutants only if such mutants
preferentially receive fitness benefits from their algal partners, that is, if there is a partner-fidelity
feedback[51]. Our system is well-mixed, so that all diffusible benefits are shared by the entire
culture, eliminating any potential fitness advantage of rare mutualism-enhancing mutants[26,52].
The only way to prevent such diffusion and ensure preferential benefit exchange with an algal
partner is for such a mutant to form a physical association with the partner[22]. However, we found
no evidence for such associations in any of the sampled mutants (Figure S19; Supplementary
Information). Given the absence of a plausible partner-fidelity feedback, the increased
cooperation of C-mutants must be a byproduct (pleiotropic effect) of selection for one or more
other traits.

We sought to identify traits under selection in the C-condition that could cause yields to increase.
Both competitive fitness and yield depend on fundamental physiological and life-history traits
embodied by yeast and alga, such as their growth rates, mortalities, nutrient consumption
efficiencies, etc[53]. Since measuring all potentially relevant traits was not feasible in this study,
we focused on two key traits that are known to be under selection in environments with variable
nutrient availability. The maximum population growth rate, r, is important for competitive
fitness when resources are abundant[54,55], a condition that takes place at the beginning of each
growth cycle in our cultures. The carrying capacity, K, is an indicator of nutrient utilization
efficiency, which is important for competitive fitness when resources are scarce[54,55], a condition
that takes place at the later phase of each growth cycle. We estimated these two traits for all 59
mutants from monoculture growth-curve measurements (Figures S20, S21; Supplementary
Information), reasoning that these intrinsic traits would be relevant for fitness in both A- and C-
conditions.

We found that the sampled adaptive mutations ($n = 59$) have diverse effects on r and K, with
many mutations increasing and decreasing both traits (Figure 5A). There is a negative correlation
between the effects of mutations on r and K (Pearson $R = -0.38$, 95% CI: [-0.59, -0.13], two-
sided permutation $P = 0.004$, Figure 5A), indicating a commonly observed trade-off between
growth rate and nutrient utilization efficiency[56-58]. To test whether r and K are under selection,
we examined the correlation between these traits and competitive fitness among all 59 assayed
mutants. We found weak or no correlation between either r or K and fitness in the A-condition

(permutation $P \geq 0.05$, Figure S22A, B). As a result, the A-mutants have r and K values that are
indistinguishable from the ancestor (average $\Delta r = 2\% \pm 1.9\%$, $P = 0.27$; average $\Delta K = 3\% \pm$
2.4%; $P = 0.29$; $n = 31$, permutations of two-sided Welch's t-test). In contrast, fitness in the C-
condition is positively correlated with K (Figure 5B, $R = 0.51$, 95% CI [0.29, 0.68], $P < 10^{-4}$)
and negatively correlated with r (Figure S22C, Pearson's $R = -0.51$, 95% CI [-0.68, -0.29], $P <$
10^{-4}), consistent with the observed r vs K trade-off. C-mutants reach K values on average 12.7%
higher than the ancestor ($P = 4 \times 10^{-4}$, permutations of two-sided Welch's t-test) and 9.5% higher
than A-mutants (95% CI [2.6%, 16.3%], $P = 0.03$). C-mutants also have a significantly lower r
than both the ancestor (average $\Delta r = -8.7\% \pm 2.3\%$, $P = 0.001$) and A-mutants ($\Delta r = -11\%$, 95%
CI [-17.0%, -5.1%], $P = 7 \times 10^{-4}$). We conclude that there is no detectable selection on r or K in
the absence of alga, while the presence of the alga induces selection that favors yeast mutants
with lower r and higher K.

Figure 5. The effects of adaptive mutations on life-history traits, fitness and yield. The effects of
mutations on r and K, Δr and ΔK, are reported as fractional differences relative to the ancestral values
(see Supplementary Information). **(A)** Pearson correlation between ΔK and Δr among adapted mutants.
(B) Pearson correlation between ΔK and their relative fitness in the original community among adapted
mutants. **(C)** Pearson correlation between ΔK and YYC (yeast yield in community), normalized to
ancestral values. Error bars show ±1 standard error of the mean. Solid lines are fitted by linear
regression. P-values are computed by permutation testing (see Supplementary Information).

For the r and K traits to mediate the correlation between competitive fitness and yield, mutants
with higher K would need to produce higher yield at the end of the growth cycle. Such
correlation is generally expected because, all else being equal, mutants that reach higher density
in the middle of the growth cycle due to their higher carrying capacity are more likely to
maintain higher density at the end of the growth cycle. We found that this is indeed the case
(Figures 5C and S22D). Taken together, our observations suggest a plausible model for how
adaptive evolution can favor mutualism enhancement in the absence of partner-fidelity
feedbacks: ecological interactions with the alga intensify selection for yeast mutants that use
resources more efficiently (i.e., those that reach higher K perhaps at the expense of reduced r);
once these mutants spread in the yeast population, they support higher yields of both members of
the community. Whether mutualistic partners generally induce selection for lower r and/or
higher K, and whether such selection consistently leads to increased yields of both species
remains an open question.

Similar to our analysis of ecological parallelism, we asked whether the presence of alga alters the
probability of parallelism at the level of life-history traits, which we define as the probability that
r would be affected in the same direction in two randomly sampled mutants and that K would
also be affected in the same direction in these mutants (see Supplementary Information). We find
that the probability of trait parallelism is $30.7 \pm 1.7\%$ for the A-mutants, not significantly
different from 25% expected under the uniform null model ($P = 0.47$, two-sided χ^2-test, $\chi^2 =$
0.53, $df = 1$). In contrast, the probability of trait parallelism is $46.3 \pm 4.8\%$ for the C-mutants,
which significantly exceeds 25% ($P = 0.009$, $\chi^2 = 6.7$, $df = 1$), suggesting that evolution in the
presence of alga becomes more repeatable not only at the ecological level, as shown in the
previous section, but also at the level of underlying life-history traits.

In summary, our results show that interactions with alga shift natural selection on yeast to favor
mutants that increase K and decrease r, which in turn leads to increasing yields of both species in
the community. The shift in selection imposed by the alga makes evolution more repeatable both
at the level of life-history traits and even more so at the ecological level.

427 Discussion

We characterized early adaptation in the experimental yeast-algal community and made three
main observations. First, we found that yeast have access to adaptive mutations that are not only
genetically diverse but also have diverse ecological effects. Second, even though there are no
measurable fitness trade-offs for yeast between growing alone or with alga, the presence of the
alga modifies the fitness benefits provided by many mutations. This relatively subtle shift in
selection pressure changes the set of mutations contending for fixation in yeast and thereby alters
the course of its evolution. Third, mutations that are strongly favored by selection in the presence
versus absence of alga have different ecological consequences. Specifically, the presence of alga
shifts selection on yeast to favor mutations that enhance the yeast-algal mutualism (as measured
by the yield of both species at the end of the growth cycle), making evolution at the ecological
level more repeatable.

Insofar as our yeast-algal community is representative of other ecological communities, our
results suggest that (i) organisms have access to a variety of adaptive mutations with diverse
ecological consequences and (ii) ecological perturbations, such as removal or addition of species,
can change the fitness effects of many of these mutations, thereby altering future outcomes of
evolution not only at the genetic but also at the ecological level. As a result, the eco-evolutionary
dynamics of multi-species communities are likely historically contingent on both prior
evolution[59] and ecology[60]. Thus, we might expect that ecological communities would generically
have the potential to embark on a variety of divergent eco-evolutionary trajectories and approach
different ecological attractors. For example, mutations that are beneficial to yeast in our
community can either increase or decrease the yields of both species suggesting that our
community has the potential to evolve either towards stronger mutualism or towards mutualism
breakdown, with probabilities of these outcomes being dependent on whether yeast previously
evolved in the presence or absence of alga.

Given this potential for ecological and evolutionary historical contingency, one might expect *a*
priori that replicate communities would often diverge towards different ecological states.
However, recent laboratory studies have found that replicates tend to evolve towards similar

ecological states with surprising repeatability[14,16,17,21,22,25,26,61–65]. Our results show that, despite
the diversity of available adaptive mutations, selection in the community preferentially favors
mutations with certain restricted ecological properties (namely, those that lead to higher yields of
both yeast and alga). Taken together, these prior observations and our findings suggest that eco-
evolutionary dynamics are subject to constraints that limit the space of likely trajectories. In our
system in particular, the presence of alga modifies the effects of mutations in yeast in such a way
that its evolution becomes more repeatable at the ecological level, at least over the short-term.
Thus, ecological interactions may canalize evolution. Whether such canalization is a general
feature of evolution in a community context remains to be determined.

In our competitive mutualistic community, canalization occurs in the direction of enhanced
mutualism in the sense that the presence of the alga shifts selection on yeast in favor of
mutations that benefit both species. While there are no known mechanisms in our community
that favor cooperativity directly, we have demonstrated another plausible ecological mechanism
by which increased cooperativity can evolve. Mutations favored in the presence of alga tend to
simultaneously increase yeast's carrying capacity in our medium and reduce its growth rate.
Increased carrying capacity could provide the competitive advantage necessary for such mutants
to spread. Increased K and/or decreased r could enhance the cooperation with the alga.
Specifically, increased K means that there are more yeast cells to generate CO_2, which stimulates
algal growth. Reduction in r could also benefit the alga via the "competitive restraint"
mechanism[66], because slower growing yeast compete less for the initial supply of ammonium
thereby offering the alga an opportunity to grow more and in turn supply more ammonium
during the latter portions of the growth cycle. We note, however, that competition for initial
ammonium can probably not be reduced to zero solely by mutations in yeast. This is because
yeast lacks the molecular machinery for metabolizing the only other nitrogen source, nitrite, so
that a single mutation or even a few mutations cannot alleviate yeast's basic requirement for
ammonium. Careful measurements will be needed to determine how adaptations discovered here
modify the competitive and cooperative phases of the growth cycle to provide an evolutionary
advantage for yeast and increase the yield of both species in the community.

Analysis of the genetic and biochemical basis of adaptation may further help us disentangle the
ecological mechanisms of mutualism enhancement and assess how general they might be.
However, one challenge is that many adaptive mutations that we discovered are large
chromosomal amplifications and deletions, and it is unclear which amplified/deleted genes
actually cause the fitness gains and changes in the ecologically relevant traits. At this point, we
can only speculate on this subject. For example, it is known that ChrXIV-3n amplifications are
adaptive under ammonium limitation, possibly driven by the copy number of the gene *MEP2* that
encodes a high affinity ammonium transporter[67]. We suspect that these adaptations are
particularly beneficial to yeast in the C-condition because the alga provides a continuous but low
flux of ammonia. Another interesting example are mutations in genes *HEM1*, *HEM2* and *HEM3*
which provide much larger fitness benefits in the C-condition compared to the A-condition (Data
S3) possibly because they shift the metabolic balance towards fermentation at higher
concentrations of dissolved oxygen produced by the alga (see Supplementary Information).
Elucidating these and other mechanisms of physiological adaptation in our competitive
mutualistic systems is the subject of future work.

Overall, our results suggest that microbial adaptation in the community context is driven by many mutations that are genetically and phenotypically diverse and have diverse ecological consequences. Changes in the ecological milieu, such as loss of some species or invasions by others, may not necessarily alter which mutations are beneficial to community members. Nevertheless, such ecological changes can quantitatively alter the benefits of mutations, so that evolutionary trajectories become canalized towards certain ecological outcomes.

References

1. Agostini, S. *et al.* Ocean acidification drives community shifts towards simplified non-calcified habitats in a subtropical-temperate transition zone. *Sci. Rep.* **8**, 11354 (2018).
2. Walther, G.-R. *et al.* Ecological responses to recent climate change. *Nature* **416**, 389–395 (2002).
3. Gilbert, B. & Levine, J. M. Ecological drift and the distribution of species diversity. *Proc. Biol. Sci.* **284**, (2017).
4. White, E. P. *et al.* A comparison of the species-time relationship across ecosystems and taxonomic groups. *Oikos* **112**, 185–195 (2006).
5. Sax, D. F. & Gaines, S. D. Species invasions and extinction: the future of native biodiversity on islands. *Proc. Natl. Acad. Sci. U. S. A.* **105 Suppl 1**, 11490–11497 (2008).
6. Thompson, J. N. Rapid evolution as an ecological process. *Trends Ecol. Evol.* **13**, 329–332 (1998).
7. Post, D. M. & Palkovacs, E. P. Eco-evolutionary feedbacks in community and ecosystem ecology: interactions between the ecological theatre and the evolutionary play. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **364**, 1629–1640 (2009).
8. Reznick, D. N. & Travis, J. Experimental Studies of Evolution and Eco-Evo Dynamics in Guppies (Poecilia reticulata). *Annu. Rev. Ecol. Evol. Syst.* (2019) doi:10.1146/annurev-ecolsys-110218-024926.
9. Hendry, A. P. *Eco-evolutionary Dynamics*. (Princeton University Press, 2020).
10. Yoshida, T., Jones, L. E., Ellner, S. P., Fussmann, G. F. & Hairston, N. G. Rapid evolution drives ecological dynamics in a predator–prey system. *Nature* vol. 424 303–306 (2003).
11. Bedford, T., Rambaut, A. & Pascual, M. Canalization of the evolutionary trajectory of the human influenza virus. *BMC Biol.* **10**, 38 (2012).
12. Kotil, S. E. & Vetsigian, K. Emergence of evolutionarily stable communities through eco-evolutionary tunnelling. *Nature Ecology & Evolution* **2**, 1644–1653 (2018).
13. Hansen, S. K., Rainey, P. B., Haagensen, J. A. J. & Molin, S. Evolution of species interactions in a biofilm community. *Nature* **445**, 533–536 (2007).
14. Hillesland, K. L. & Stahl, D. A. Rapid evolution of stability and productivity at the origin of a microbial mutualism. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 2124–2129 (2010).
15. Turcotte, M. M., Reznick, D. N. & Hare, J. D. The impact of rapid evolution on population dynamics in the wild: experimental test of eco-evolutionary dynamics. *Ecol. Lett.* **14**, 1084–1092 (2011).
16. Lawrence, D. *et al.* Species interactions alter evolutionary responses to a novel environment. *PLoS Biol.* **10**, e1001330 (2012).
17. Celiker, H. & Gore, J. Clustering in community structure across replicate ecosystems following a long-term bacterial evolution experiment. *Nat. Commun.* **5**, 4643 (2014).
18. Andrade-Domínguez, A., Salazar, E., Vargas-Lagunas, M. del C., Kolter, R. & Encarnación, S. Eco-evolutionary feedbacks drive species interactions. *ISME J.* **8**, 1041–1054 (2014).

19. Reznick, D. Hard and Soft Selection Revisited: How Evolution by Natural Selection Works in the
Real World. *J. Hered.* **107**, 3–14 (2016).

20. Matthews, B., Aebischer, T., Sullam, K. E., Lundsgaard-Hansen, B. & Seehausen, O. Experimental
Evidence of an Eco-evolutionary Feedback during Adaptive Divergence. *Curr. Biol.* **26**, 483–489
(2016).

21. Harcombe, W. R., Chacón, J. M., Adamowicz, E. M., Chubiz, L. M. & Marx, C. J. Evolution of
bidirectional costly mutualism from byproduct consumption. *Proc. Natl. Acad. Sci. U. S. A.* **115**,
12000–12004 (2018).

22. Preussger, D., Giri, S., Muhsal, L. K., Oña, L. & Kost, C. Reciprocal Fitness Feedbacks Promote the
Evolution of Mutualistic Cooperation. *Curr. Biol.* (2020) doi:10.1016/j.cub.2020.06.100.

23. Adamowicz, E. M., Muza, M., Chacón, J. M. & Harcombe, W. R. Cross-feeding modulates the rate
and mechanism of antibiotic resistance evolution in a model microbial community of *Escherichia
coli* and *Salmonella enterica*. *PLoS Pathog.* **16**, e1008700 (2020).

24. Rodríguez-Verdugo, A. & Ackermann, M. Rapid evolution destabilizes species interactions in a
fluctuating environment. *ISME J.* **15**, 450–460 (2021).

25. Barber, J. N. *et al.* The evolution of coexistence from competition in experimental co-cultures of
Escherichia coli and Saccharomyces cerevisiae. *ISME J.* **15**, 746–761 (2021).

26. Hart, S. F. M., Chen, C.-C. & Shou, W. Pleiotropic mutations can rapidly evolve to directly benefit
self and cooperative partner despite unfavorable conditions. *Elife* **10**, (2021).

27. Schoener, T. W. The newest synthesis: understanding the interplay of evolutionary and ecological
dynamics. *Science* **331**, 426–429 (2011).

28. Weber, M. G., Wagner, C. E., Best, R. J., Harmon, L. J. & Matthews, B. Evolution in a Community
Context: On Integrating Ecological Interactions and Macroevolution. *Trends Ecol. Evol.* **32**, 291–
304 (2017).

29. Kokko, H. *et al.* Can Evolution Supply What Ecology Demands? *Trends Ecol. Evol.* **32**, 187–197
(2017).

30. Turcotte, M. M., Reznick, D. N. & Daniel Hare, J. Experimental test of an eco-evolutionary dynamic
feedback loop between evolution and population density in the green peach aphid. *Am. Nat.* **181
Suppl 1**, S46–57 (2013).

31. Nuismer, S. *Introduction to Coevolutionary Theory*. (Macmillan Learning, 2017).

32. Stoltzfus, A. & McCandlish, D. M. Mutational Biases Influence Parallel Adaptation. *Mol. Biol. Evol.*
34, 2163–2172 (2017).

33. Payne, J. L. *et al.* Transition bias influences the evolution of antibiotic resistance in Mycobacterium
tuberculosis. *PLoS Biol.* **17**, e3000265 (2019).

34. Storz, J. F. *et al.* The role of mutation bias in adaptive molecular evolution: insights from convergent
changes in protein function. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **374**, 20180238 (2019).

35. Gomez, K., Bertram, J. & Masel, J. Mutation bias can shape adaptation in large asexual populations
experiencing clonal interference. *Proc. Biol. Sci.* **287**, 20201503 (2020).

36. Venkataram, S., Monasky, R., Sikaroodi, S. H., Kryazhimskiy, S. & Kacar, B. Evolutionary stalling
and a limit on the power of natural selection to improve a cellular module. *Proceedings of the
National Academy of Sciences* **117**, 18582–18590 (2020).

37. Hom, E. F. Y. & Murray, A. W. Niche engineering demonstrates a latent capacity for fungal-algal
mutualism. *Science* **345**, 94–98 (2014).

38. Wolfe, B. E. & Dutton, R. J. Fermented foods as experimentally tractable microbial ecosystems. *Cell

161, 49–55 (2015).

39. Chacón, J. M., Hammarlund, S. P., Martinson, J. N. V., Smith, L. B. & Harcombe, W. R. The
Ecology and Evolution of Model Microbial Mutualisms. *Annu. Rev. Ecol. Evol. Syst.* **52**, 363–384
(2021).

40. Blasche, S., Kim, Y., Oliveira, A. P. & Patil, K. R. Model microbial communities for ecosystems
biology. *Current Opinion in Systems Biology* **6**, 51–57 (2017).

41. Levy, S. F. *et al.* Quantitative evolutionary dynamics using high-resolution lineage tracking. *Nature*
519, 181–186 (2015).

42. Venkataram, S. *et al.* Development of a comprehensive genotype-to-fitness map of adaptation-
driving mutations in yeast. *Cell* **166**, 1585–1596.e22 (2016).

43. Jones, E. I., Bronstein, J. L. & Ferrière, R. The fundamental role of competition in the ecology and
evolution of mutualisms. *Ann. N. Y. Acad. Sci.* **1256**, 66–88 (2012).

44. Boyer, S., Hérissant, L. & Sherlock, G. Adaptation is influenced by the complexity of environmental
change during evolution in a dynamic environment. *PLoS Genet.* **17**, e1009314 (2021).

45. Blundell, J. R. *et al.* The dynamics of adaptive genetic diversity during the early stages of clonal
evolution. *Nature Ecology & Evolution* **3**, 293–301 (2019).

46. Good, B. H., Rouzine, I. M., Balick, D. J., Hallatschek, O. & Desai, M. M. Distribution of fixed
beneficial mutations and the rate of adaptation in asexual populations. *Proceedings of the National*
Academy of Sciences **109**, 4950–4955 (2012).

47. Good, B. H., Martis, S. & Hallatschek, O. Adaptation limits ecological diversification and promotes
ecological tinkering during the competition for substitutable resources. *Proceedings of the National*
Academy of Sciences **115**, E10407–E10416 (2018).

48. Schiffels, S., Szöllosi, G. J., Mustonen, V. & Lässig, M. Emergent neutrality in adaptive asexual
evolution. *Genetics* **189**, 1361–1375 (2011).

49. Nguyen Ba, A. N. *et al.* High-resolution lineage tracking reveals travelling wave of adaptation in
laboratory yeast. *Nature* **575**, 494–499 (2019).

50. Foster, K. R., Shaulsky, G., Strassmann, J. E., Queller, D. C. & Thompson, C. R. L. Pleiotropy as a
mechanism to stabilize cooperation. *Nature* **431**, 693–696 (2004).

51. Sachs, J. L., Mueller, U. G., Wilcox, T. P. & Bull, J. J. The evolution of cooperation. *Q. Rev. Biol.*
79, 135–160 (2004).

52. Harcombe, W. Novel cooperation experimentally evolved between species. *Evolution* **64**, 2166–2172
(2010).

53. Vasi, F., Travisano, M. & Lenski, R. E. Long-term experimental evolution in *Escherichia coli*. II.
Changes in life-history traits during adaptation to a seasonal environment. *Am. Nat.* **144**, 432–456
(1994).

54. MacArthur, R. H. & Wilson, E. O. *The theory of island biogeography*. (Princeton University Press,
2001).

55. Reznick, D., Bryant, M. J. & Bashey, F. *r*- and *K*-selection revisited: the role of population
regulation in life-history evolution. *Ecology* **83**, 1509–1520 (2002).

56. Mueller, L. D. & Ayala, F. J. Trade-off between r-selection and K-selection in Drosophila
populations. *Proc. Natl. Acad. Sci. U. S. A.* **78**, 1303–1305 (1981).

57. Novak, M., Pfeiffer, T., Lenski, R. E., Sauer, U. & Bonhoeffer, S. Experimental tests for an
evolutionary trade-off between growth rate and yield in *E. coli*. *Am. Nat.* **168**, 242–251 (2006).

58. Orivel, J. *et al.* Trade-offs in an ant-plant-fungus mutualism. *Proc. Biol. Sci.* **284**, (2017).

59.　Blount, Z. D., Lenski, R. E. & Losos, J. B. Contingency and determinism in evolution: Replaying
life's tape. *Science* **362**, (2018).
60.　Fukami, T. Historical contingency in community assembly: integrating niches, species pools, and
priority effects. *Annu. Rev. Ecol. Evol. Syst.* **46**, 1–23 (2015).
61.　Rainey, P. B. & Travisano, M. Adaptive radiation in a heterogeneous environment. *Nature* **394**, 69–
72 (1998).
62.　Meyer, J. R. *et al.* Repeatability and contingency in the evolution of a key innovation in phage
lambda. *Science* **335**, 428–432 (2012).
63.　Herron, M. D. & Doebeli, M. Parallel evolutionary dynamics of adaptive diversification in
Escherichia coli. *PLoS Biol.* **11**, e1001490 (2013).
64.　Hillesland, K. L. *et al.* Erosion of functional independence early in the evolution of a microbial
mutualism. *Proc. Natl. Acad. Sci. U. S. A.* **111**, 14822–14827 (2014).
65.　Meroz, N., Tovi, N., Sorokin, Y. & Friedman, J. Community composition of microbial microcosms
follows simple assembly rules at evolutionary timescales. *Nat. Commun.* **12**, 2891 (2021).
66.　MacLean, R. C. The tragedy of the commons in microbial populations: insights from theoretical,
comparative and experimental studies. *Heredity* **100**, 471–477 (2008).
67.　Dunn, B. *et al.* Recurrent rearrangement during adaptive evolution in an interspecific yeast hybrid
suggests a model for rapid introgression. *PLoS Genet.* **9**, e1003366 (2013).

Acknowledgements: We thank Gavin Sherlock and Katja Schwartz for providing the barcoded
yeast library, Stephen Mayfield and Frank Fields for laboratory equipment and help with algal
husbandry, Rachel Dutton and Manon Morin for help with sequencing, Scott Rifkin and Jessica
Bloom for help with microscopy, STARS students Jesse Yu and Sophia Rosemann for help with
experiments, Justin Meyer, Alena Martsul, Shohreh Sikaroodi for technical assistance, the
Kryazhimskiy, Meyer and Hwa labs, Damien Barrett, Josh Borin, Shermin de Silva, Susanne
Dunker, Nandita Garud, Stan Harpole, Canan Karakoç, Holly Moeller, Dmitri Petrov, and Peter
Zee for feedback on the manuscript. Sequencing was done in part at the UCSD IGM center
(University of California, San Diego, La Jolla, CA). We acknowledge the San Diego
Supercomputing Center for the use of the TSCC cluster for computing services. EFYH is funded
by National Science Foundation CAREER grant 1846376 and Deutsches Zentrum für Integrative
Biodiversitätsforschung (iDiv) grant DFG–FZT 118, 202548816. SK is funded by BWF Career
Award at the Scientific Interface grant 1010719.01, Alfred P. Sloan Foundation grant FG-2017-
9227 and the Hellman Foundation.

Author contributions: conceptualization (SV, EFYH, SK), methodology (SV, HYK, SK), data
acquisition (SV), analysis (SV, HYK, SK), initial manuscript (SV, SK), editing (SV, HYK,
EFYH, SK), supervision (EFYH, SK), funding (SK).

Competing interests: The authors declare no competing interests.

Materials and Correspondence: All raw sequencing data is available on the US National
Center for Biotechnology Information (NCBI) Sequence Read Archive (SRA) under BioProject
PRJNA735257. Other input data (e.g. growth data, variant calls, community yield etc) and
analysis scripts can be found in Data Files S1–S3 and on Dryad at
https://doi.org/10.6076/D14K5X. Strains and other biological materials are available by request
to SK.

Code availability: The latest version of the barcode counting software BarcodeCounter2 can be
found at https://github.com/sandeepvenkataram/BarcodeCounter2.git. All other analysis scripts
used for this study are available on Dryad at https://doi.org/10.6076/D14K5X.

Code and data availability for reviewers: As the data and code is not currently publicly
available on Dryad, reviewers can access this data through the following reviewer link:
https://datadryad.org/stash/share/g4RSTbYCCwCpQxidH0dhQDKf7SsVsijY8pjzcsy7Y88

678 Supplementary Materials

Supplementary Information Text

Figure S1. Lineage trajectories in yeast populations in the A-condition.

Figure S2. Lineage trajectories in yeast populations in the C-condition.

Figure S3. Estimation of selection coefficients of adapted lineages in the BLT data.

Figure S4. Evidence for pre-existing mutations in BLT experiments.

Figure S5. Lineage frequency trajectories in the competition assay.

Figure S6. Correlation between fitness estimates across replicates of the competition assay.

Figure S7. Mean and coefficient of variation of the fitness estimates in the competition assay.

Figure S8. Correlation between BLT and competitive fitness estimates.

Figure S9. An example illustrating the detection of large chromosomal duplications and
deletions in the genomic data.

Figure S10. Relationship between coverage and the proximity to chromosome ends.

Figure S11. Allele frequency spectrum for ancestral and adapted isolates.

Figure S12. Distribution of mutations per clone.

Figure S13. Distribution of adaptive mutations across driver loci.

Figure S14. The log-likelihood function for the estimation of the chrIII-3n beneficial mutation
rate.

Figures S15. Sampling probabilities for adaptive mutations at the most common driver loci.

Figures S16. Correlation between yield measurements across replicates.

Figures S17. Estimated yeast and alga yields among random A- and C-mutants.

Figure S18. Correlation between yeast mutant fitness in the original community and alga yield
in the mutant community

Figures S19. Representative microscopy image showing lack of physical associations between
yeast and alga cells.

Figures S20. Mutant growth curves in the A-condition.

Figures S21. Correlation between replicates for r and K measurements.

Figures S22. Relationship between growth parameters, yields and fitness.

Table S1. Statistics of mutational counts among sequenced mutants.

Table S2. Number of isolates carrying the most common adaptive mutations and the estimated
mutation rates.

Data S1. List of all sampled isolates, their fitness, phenotypes and yields.

Data S2. List of all identified mutations.

Data S3. Distribution of adaptive mutations across driver loci.

Supplementary Information for
"Mutualism-enhancing mutations dominate early adaptation in a microbial community"

Sandeep Venkataram, Huan-Yu Kuo, Erik F. Y. Hom, Sergey Kryazhimskiy*

January 23, 2022

Contents

---

*Correspondence: skryazhi@ucsd.edu

1 Data availability

All raw sequencing data is available on the US National Center for Biotechnology Information (NCBI) Sequence Read Archive (SRA) under BioProject PRJNA735257. Other input data (e.g. growth data, variant calls, community yield etc) and analysis scripts can be found on Dryad at

https://doi.org/10.6076/D14K5X.

The latest version of the barcode counting software BarcodeCounter2 can be found at

https://github.com/sandeepvenkataram/BarcodeCounter2.git.

2 Barcode lineage tracking (BLT) experiment and data analysis

2.1 Strains

We used the strain CC1690 of the alga *Chlamydomonas reinhardtii* as in Ref. [1], which can also be obtained from the Chlamydomonas Resource Center.

The barcoded library of the diploid yeast *Saccharomyces cerevisiae* strain GSY6699 is described in Ref. [2] and was kindly provided by Prof. Gavin Sherlock. This is a diploid, prototrophic strain derived from the BY genetic background, homozygous throughout the genome, except for locus YBR209W, where one copy of a DNA barcode was integrated, as described in Ref. [3]. Our starting library consists of about 5×10^5 clones, each of which carries a unique DNA barcode at this locus. In principle, the genomes of all clones should be identical everywhere else prior to our barcode lineage tracking (BLT) experiment. However, as discussed in detail below (see Sections 2.6.2 and 4), we found that our initial population already contains some pre-existing polymorphisms, which arose prior to our BLT experiment.

2.2 Growth conditions

Both yeast monocultures and yeast-alga communities were cultured in a defined basic medium ("CYM medium", see Ref. [1]) supplemented with 2% dextrose, 10mM KNO_2 and 0.5mM NH_4Cl, which we thereafter refer to as the "growth medium". All cultures were grown at room temperature (21°C) on a platform shaker with 70 foot-candles of constant light (three Feit Electric #73985 suspended approximately 24 inches above the platform shaker) shaking at 125 RPM, unless noted otherwise.

When yeast is grown alone in this medium (the A-condition), it reaches the final density of 2.49×10^6 cells per mL. When yeast and alga are co-cultured together (the C-condition), yeast and alga reach final densities (yields) of 2.72×10^6 cells per mL and 5.63×10^5 cells per mL, respectively. In all subsequent experiments, we grow our cultures in 10mL of the growth medium in 50mL flasks (the "standard conditions"), unless otherwise noted.

2.3   BLT experiment

2.3.1   Pre-cultures

Prior to the BLT experiment, yeast and alga were pre-cultured in 50mL of growth medium in 250mL delong baffled flasks (PYREX #C4446250) for two and 10 days respectively. Alga pre-cultures were started from colonies. To start yeast pre-cultures, the barcoded yeast library was thawed from frozen stock at room temperature, then 500µL were transferred into 50mL of the growth medium.

2.3.2   Initiation and propagation

We conducted five replicate BLT experiments for each of two treatments, yeast monoculture (the A-condition) and yeast + algae community (the C-condition). All BLT experiments were conducted in 10mL of the growth medium in 50mL Erlenmeyer flasks (FisherSci #FS2650050) capped with 50mL plastic beakers (VWR #414004-145) to maintain sterility. Each monoculture BLT experiment was initiated from 100µL of the yeast pre-culture. Each community BLT experiment was initiated from 100µL of the 1:1 (v/v) yeast and alga mixture. During the BLT experiment, cultures were grown on a platform shaker with constant light as described above for 5 days before being diluted 1:100 for the next growth cycle (100µL into 10mL fresh media). A total of 17 growth/dilution cycles were completed.

2.3.3   Culture preservation

Glycerol stocks were taken of the yeast pre-culture and yeast + algae inoculum mixture, as well as at the end of every odd growth cycle. Separate stocks were stored for DNA extraction and cell isolation purposes with two replicates each, for a total of 4 stocks per culture per time point. Cell isolation stocks were created by aliquoting 1.5mL of culture into 500µL of 80% glycerol, mixing by vortex and storing at $-80°$C. DNA stocks were created by removing the supernatant of the remaining 7mL of culture via centrifugation and resuspending in 2mL of 20% glycerol (80% glycerol diluted with 1x PBS), which was then stored as two separate 1mL stocks at $-80°$C.

2.4 Justification for ignoring adaptation in alga

Here we argue that ignoring adaptation in alga in our experiment is justified, based on our current knowledge of rates of adaptation in yeast and alga.

In the C-condition, that is, when yeast and alga are co-cultured together, yeast and alga reach final yields of 2.72×10^7 and 5.63×10^6 cells, respectively, with the bottleneck sizes being by a factor of 100 smaller. Since there is no evidence of alga or yeast death in this condition (Figure 1), we estimate that both species undergo on average $\log_2 100 = 6.64$ doublings per cycle. Numerous evolution experiments with yeast populations of comparable or smaller size across various environmental conditions have shown that adaptive mutations arise and reach high frequencies within 250 generations or less [4, 3, 5, 6, 7]. Although much less is known about the rates of adaptation in *Chlamydomonas reinhardtii*, one study reports a failure of the alga (strain CC2344) to adapt within 1000 generations of evolution at the bottleneck size of 10^5 cells, i.e., twice as large as in our experiment [8]. Another study reports 35% growth rate gain after 1880 generations of evolution in alga strain CC-503 cw92 mt+ [9], although the bottleneck size in that experiment was about 20 times larger than in ours and detectable gains appeared only after about 300 generations. These studies suggest that yeast has a much larger potential to gain adaptive mutations than alga, at least in laboratory conditions.

2.5 Barcode sequencing

2.5.1 DNA isolation

DNA stocks were thawed and DNA isolated using a "salting out" method, based on methods from Refs. [10, 11]. The thawed stocks were first centrifuged, and the supernatant removed. The pellet was resuspended in 300µL 3% SET buffer (3% SDS, 10mM EDTA, 30mM Tris) and incubated at 65°C for 15 minutes. The tube was cooled to room temperature by immersing in room temperature water, then 2.5µg of RNAse A was added. After vortexing, the mixture was incubated at 37°C for 1 hour, after which it was cooled on ice. 150µL of 3M Sodium Acetate was then added and mixed by inversion, after which it was cooled on ice for a further 5 minutes before centrifuging at maximum speed for 10 minutes on a tabletop centrifuge. The supernatant was transferred to a new tube and DNA was precipitated by the addition of 500µL isopropanol, which was mixed by inversion and then pelleted by centrifugation for 1 minute at maximum speed. The supernatant was removed and the pellet was washed with 200µL cold 70% ethanol without vortexing before being allowed to dry inverted for 30 minutes at room temperature before the DNA was resuspended in 50µL molecular biology grade water.

2.5.2 Sequencing library preparation

The barcode locus was amplified through a 2-step PCR protocol, slightly modified from the protocol described in Ref. [12]. The first amplification added inline indices for sample multiplexing, universal molecular identifiers for removing PCR duplicates during analysis and Illumina-compatible adapter sequences for a second round of amplification with standard Illumina Nextera XT primers. For the first reaction, 10µL of template was mixed with 25µL of OneTaq 2x Master Mix, 1µL of 25mM $MgCl_2$, 1µL each of the forward and reverse primers (at 10mM concentration) and 12µL of molecular biology grade water. Primer sequences are as described in Ref. [12]. This mixture was amplified using the following conditions:

1. 94°C for 10 min

2. 94°C for 3 min

3. 55°C for 1 min

4. 68°C for 1 min

5. Repeat steps 2–4 for a total of 8 cycles

6. 68°C for 1 min

7. Hold at 4°C

The amplified product was purified using Ampure XP magnetic beads following the protocol in Ref. [13] (with 50µL of beads used per sample). Then, 10µL of the purified product was used as the template for a second reaction along with Illumina Nextera XT primers (1µL of 10mM stock for each primer), 1µL of 25mM $MgCl_2$, 25µL of OneTaq 2x Master Mix and 12µL of water with the following reaction conditions:

1. 94°C for 5 min

2. 94°C for 30 sec

3. 62°C for 30 sec

4. 68°C for 30 sec

5. Repeat steps 2–4 for a total of 25 cycles

6. 68°C for 5 min

7. Hold at 4°C

The PCR products were purified using Ampure XP beads as before. 5µL of each sample was mixed to form a pool for Illumina sequencing, concentrated using Ampure beads as before with equal volume of beads as pooled sample, size selected via agarose gel extraction to isolate the correct amplicon before submitting for sequencing.

2.5.3 Sequencing

Populations 1 and 6 were initially sequenced on a MiSeq platform. All populations (including 1 and 6) were then also sequenced on a HiSeq platform. Data from both runs were combined for all downstream analysis of frequency trajectories. We obtained an average of 2.6 million paired-end reads per time point.

2.6 BLT data analysis

2.6.1 Barcode identification and counting

To identify and count DNA barcodes, we used a custom python pipeline BarcodeCounter2 available at https://github.com/sandeepvenkataram/BarcodeCounter2. The package first uses the BLASTn tool to identify sequences known to flank the barcode region within each read pair. If reads contain inline indices, samples can be demultiplexed. Universal molecular identifier (UMI) sequences can be extracted if present within the reads. If a sequence contains multiple barcode regions, these extracted regions are concatenated together. To account for sequencing errors, DNAClust [14] is then used to cluster the concatenated barcodes into clusters of nearly identical sequences which presumably originated from the same DNA molecule. The output of DNAClust is a FASTA database of all unique barcode sequences present in the library. Then, BWA [15] is used to map the barcode sequences from each sample onto the clustered FASTA database. Mapping of reads is necessary because the clustering process removes identifying information associating barcode sequences with samples from which they came. PCR duplicates are removed based on the UMI sequences, and the total number of unique reads corresponding to every barcode in the FASTA database is counted. The final output is a tab-delimited table of the read counts for every barcode in every sample. The software is designed to be user-friendly and highly customizable, with simple text files describing the input files, multiplexed samples and a sequence template describing the structure of the sequenced reads. The package is built using python3, and uses the popular BioPython package. The package has multithreading support, and can be run on both personal computers and supercomputing clusters.

When generating the database of all unique barcode sequences, we clustered sequences at 95% similarity, so that, given that the length of our barcode is 52bp, sequences with 3 or more basepair differences were merged into the same cluster. This set of clustered sequences was generated only once using all of the time points from all sequenced populations. Using this method, we identified a total of 557,087 unique barcodes in our data

with about 527,745 barcodes present at the initial time point, consistent with [3]. The number of detected barcodes gradually declines over time, as would be expected in the presence of demographic stochasticity and adaptive evolution (Figures S1B and S2B). Since only roughly 68,475 are detectable in any population by cycle 11 (i.e., about 13% of those present initially), we constrained our subsequent analysis to cycles 1 through 11.

2.6.2 Identification of adapted lineages and estimation of their fitness

The raw barcode frequency data for all our populations are shown in Figures S1, S2 (note that populations were sequenced only at odd cycles 1, 3, 5, 7, 9, 11, 13, 15 and 17). Our goal in analyzing these data was two fold. First, in order to physically isolate yeast clones that carry adaptive mutations (see Section 4), we needed to identify those yeast lineages that acquired an adaptive mutation (adapted lineages) among those that did not (neutral lineages). Second, in order to construct the distributions of fitness effects of beneficial mutations (bDFEs), we needed to estimate the competitive fitness of adapted lineages relative to the ancestoral yeast strain.

We were unable to successfully apply the BLT data analysis methods developed previously [3, 16] to our data primarily because of the high extinction rate of our barcodes. Thus, we developed a new iterative heuristic procedure described below that overcomes this problem. Direct fitness measurements of isolated clones (see Section 3) confirmed that the heuristic procedure works well.

Iterative heuristic procedure for identification of adapted lineages and estimating their selection coefficients.

1. **Initialization.** Neutral barcodes for iteration 1 are identified separately for every pair of consecutive time points. For a given time point pair, we define the set of neutral lineages at iteration 1 as those lineages whose frequency (a) does not exceed 10^{-4} at the earlier time point and (b) increases by less than 100-fold between cycle 1 and cycle 11.

2. **Estimation of mean fitness.** Given the set of neutral lineages at iteration $k-1$, we obtain their total frequency $x_{\text{neut}}^{(k-1)}(t)$ at each cycle t. The frequency of neutral lineages is governed by equation

$$x_{\text{neut}}(t+1) \approx x_{\text{neut}}(t)e^{-\bar{s}(t)},$$

where $\bar{s}(t)$ is the mean selection coefficient (per cycle) of the population at cycle t. Thus, we estimate the mean selection coefficient of the population at cycle t at iteration k as

$$\bar{s}^{(k)} = -\log \frac{x_{\text{neut}}^{(k-1)}(t+1)}{x_{\text{neut}}^{(k-1)}(t)}.$$

The mean fitness at cycle t is then defined as $1 + \bar{s}^{(k)}(t)$.

3. **Estimation of fitness of individual lineages.** The frequency $x_i(t)$ of lineage i with selection coefficient s_i (per cycle) relative to the ancestor is governed by the equation

$$x_i(t+1) = x_i(t)\, e^{s_i - \bar{s}(t)} = x_i(t)\, \frac{x_{\text{neut}}(t+1)}{x_{\text{neut}}(t)}\, e^{s_i}.$$

Thus, for every lineage i (including those that were called as neutral at the previous iteration) we estimate $s_i^{(k)}$ at iteration k at time point t as

$$s_i^{(k)}(t) = \log \frac{x_i(t+1)}{x_{\text{neut}}^{(k-1)}(t+1)} - \log \frac{x_i(t)}{x_{\text{neut}}^{(k-1)}(t)}. \tag{1}$$

These estimates are made for every pair of consecutive time points between cycles 1 and 11. If $x_i(t) = 0$ and $x_i(t+1) = 0$, the time point pair is excluded from the calculations. If only one of the two frequencies is 0, then this frequency is set to 0.5 / total read depth. To obtain the final estimate of the selection coefficient $s_i^{(k)}$ for lineage i at iteration k, we average all $s_i^{(k)}(t)$, weighting each $s_i^{(k)}(t)$ by the number of reads for lineage i at time t. We calculate the standard error $\sigma_i^{(k)}$ with the same weighting. The relative fitness of lineage i is then defined as $1 + s_i^{(k)}$.

4. **Calling adapted lineages.** Lineage i is called adapted at iteration k if $s_i^{(k)} > 2\sigma_i^{(k)}$.

5. **Updating the set of neutral lineages.** We calculate the coefficient of variation for each lineage i as $\text{CV} = \sigma_i^{(k)}/s_i^{(k)}$. We include any lineage i into the new set of neutral lineages at iteration k if (a) its maximum frequency does not exceed 10^{-4} between cycles 1 and 11, and (b) either the CV of the lineage is more the median CV across all lineages or if $s_i^{(k)} < 0$ (see Figure S3A).

6. **Termination.** The procedure is terminated when the set of neutral lineages at iteration k differs by less than 5% compared to the set at the iteration $k-1$.

Our procedure converges to a stable set of neutral lineages within 5 iterations. We report fitness values relative to the ancestral strain on a per-cycle basis, rather than the per-generation basis typically used in the literature because our cultures experience growth phases other than exponential growth, as discussed in Refs. [12, 17].

Validation. We performed two "sanity" checks to ensure that our analysis gives reasonable results. First, mean fitness is expected to monotonically increase over time in each population. We find that mean fitness does indeed increase monotonically during the first 11 cycles in all but one populations (Figure S3B). Second, we visually inspected how well individual lineage trajectories are fitted by equation (1) and found that these fits are generally very good (Figure S3C,D shows typical fits).

Pre-existing mutations. Some mutations could have arisen prior to the beginning of the experiment, specifically, prior to splitting the barcoded yeast library between five A and five C replicates. Such pre-existing mutations cause two issues with the downstream analysis and interpretation. First, they make it more difficult to identify causal mutations that increase fitness of adapted lineages. We discuss this problem and how we address it in Section 4. The second issue is that pre-existing beneficial mutations confound the inference of bDFEs and their comparison across treatments. Specifically, if multiple lineages (within the same population or in different replicate populations) carry the same pre-exisiting beneficial mutation and are identified as such, this mutation will be counted multiple times and will therefore inflate the size of a fitness class in the inferred bDFE.

To diagnose this problem, we note that there are two types of pre-existing mutations. Those that arose after the transformation that introduced barcodes into our strain (type I) and those that arose before this transformation (type II). All pre-existing mutations of type I are linked to different barcodes. Therefore, these mutations can be identified by comparing barcodes of lineages identified as adapted in different populations. We found that pairs of populations have on average 416 (15%) adapted lineages with a common barcode, whereas only 23 (0.8%) would be expected by chance (Figure S4). Barcodes are more often shared between A populations than other types of population pairs (22% vs 13%, t-test $P = 10^{-10}$). We suspect that this is because the majority of adapted mutants have higher relative fitness in the A-condition than the C-condition (Figure 2B in the main text), such that pre-existing mutations are more likely to reach high frequencies and be detected in the A-condition. We eliminate pre-existing beneficial mutations of type I from our bDFE analysis by constructing these distributions from lineages with unique barcodes.

The same pre-existing mutation of type II can be linked to multiple barcodes and is indistinguishable in the barcode data from multiple independently arising mutations. However, pre-existing mutations of type II can be identified in the genome sequencing data. As discussed in Section 4.5, we find no evidence of such mutations, suggesting that they must be rare in our populations.

2.6.3 Permutation tests for differences in bDFE across conditions

We randomly relabel adaptive lineages as being from either the A- or C-condition, and calculate the difference in bDFE median and IQR from this permuted data. This procedure was conducted 10,000 times, to generate a random distribution of median and IQR difference values. The observed median and IQR difference were larger than those of all permuted values.

2.7 Results

The median standard error in our estimates of fitness of adaptive mutants was 0.02, and median CV was 0.08 (Figure S3A). The standard deviation of the bDFEs is much larger, 0.22 for A populations and 0.33 for C populations, suggesting that there are multiple classes of adaptive mutations available to yeast in both conditions.

3 Competitive fitness assays

3.1 Clone isolation and barcode genotyping

We selected cycle 9 to isolate adapted mutants because the estimated fraction of adapted lineages in our evolving populations was large, but each adapted lineage was still at a low frequency (Figures S1 and S2). Specifically, 92% of a typical population in the A-condition consisted of adapted lineages, with the median frequency of an individual adapted lineage being 9×10^{-5} (~ 22 cells per lineage at the bottleneck), and 73% of a typical population in the C-condition consisted of adapted lineages, with the median frequency of an individual adapted lineage being 7×10^{-5} (~ 19 cells per lineage at the bottleneck).

Isolation of random clones.   To isolate adapted clones, frozen stocks of the monoculture and community populations from cycle 9 were thawed, plated onto standard 100mm Petri dishes with CYM + 1% agarose at a dilution of approximately 100 cells per dish, and incubated at 30°C for three days (algae do not grow at 30°C). 88 random colonies were isolated from each population, i.e., a total of 440 clones from the A- and C-condition each. Eight additional clones from each population were harvested at cycle 17 and are present in the pools described below, but they are not included in any of the analyses presented in this study. Each colony was transferred into a well of a 96-well plate (Corning 3370) with 200µL of CYM media and incubated for two days at 30°C. 10µLwas used for each of the "population", "row" and "column" pools described below. Then, 50µL of 80% glycerol was added to each well, and the plate was stored at −70°C.

Barcode genotyping.   The DNA barcodes of all isolated clones were identified by sequencing, using the Sudoku method [18, 19]. Specifically, 10µL of each clone was pooled into 10 "population" pools (one pool for each source population), eight "row" pools and 12 "column" pools. DNA barcodes in each pool were amplified and sequenced as described in Section 2.5. We expect that a given combination of row, column and population pools would have a single barcode in common, defining the isolate in the corresponding well of the appropriate plate. We determined all barcodes present in the intersection of each combinations of row, column population pool. If a single barcode is identified in the intersection, the corresponding well is assigned that barcode identity. If multiple barcodes or no barcode are identified, the associated clone is removed from further analysis.

Generation of A, C and N pools. We used the heuristic procedure described in Section 2.6.2 to classify clones with identified barcodes into three groups described below: non-adapted, adapted in the C-condition or adapted in the A-condition. Clones were pooled into three libraries defined by their membership in these three groups as follows. To construct each pool, we transferred 20µL of thawed frozen stock of each isolate into a 2mL 96-deep-well plate (Corning P-2ML-SQ-C) filled with 1.8mL of growth media supplemented to 10mM ammonia. After incubating these plates at 30°C for three days, we formed each pool by combining 200µL of individual saturated cultures. Three pools were stored in 20% glycerol at −70°C.

For the sake of efficiency, clone pooling was based on an earlier version of the heuristic procedure described in Section 2.6.2 used for the classification of lineages. The current version of the procedure (as described in Section 2.6.2) classifies lineages slightly differently. As a result, pools do not perfectly correspond to the classification of clones according to the current heuristic procedure, which is provided below. This minor discrepancy has no bearing on our results because the final classification of clones is based on fitness estimates from the competition assays (see Section 3.3). The *A pool* contains 214 clones that are classified as adapted in A populations. The *C pool* contains 223 clones that are classified as adapted in C populations. The *N pool* contains 144 clones, 84 of which are classified as neutral from the BLT analysis (i.e., not adapted in either A or C populations), 29 clones that are classified as adapted in the A-condition and 31 clones that are classified as adapted in the C-condition. Thus, we measured competitive fitness for a total of 581 clones.

3.2 Competition assay experiment

To conduct the competitive fitness assays, we pre-cultured each of the three pools (N, A and C; see above) separately in the growth media for two days. We also pre-cultured algae for 10 days, starting from colonies. We then combined A, C and N pools in the 1:1:18 ratio. We carried out three replicate competitions in the A-condition and three replicate competitions in the C-condition. To this end, we inoculated each of the six replicates with 100µL of the combined A/C/N pool. In addition, the three C-condition replicates were inoculated with 100µL of the algae preculture ($\sim 10^6$ cells / mL).

All replicates were propagated in conditions identical to the BLT experiment for a total of five growth cycles. Glycerol stocks were made at the end of each growth cycle after the dilution step by centrifuging the full culture volume, removing the spent media and resuspending the pellet in 2mL of 20% glycerol + PBS. Two 1mL aliquots of this glycerol suspension were stored at −70°C. One of these aliquots was harvested for DNA extraction and barcode sequencing using protocols described in Section 2.6.2.

3.3 Competition assay data analysis

Barcodes were identified and counted as described above (see Section 2.6.1). The resulting barcode count data were analyzed as described in Ref. [12] using software available at https://github.com/barcoding-bfa/fitness-assay-python. Briefly, the 84 non-adapted barcodes (as defined from the BLT analysis described in Section 2.6.2) from the N pool were used to estimate the mean fitness trajectories and the additive and multiplicative noise parameters for each pair of time points in each assay (see Ref. [12] for details). These estimates were used to estimate the fitness of every lineage the assay for each pair of neighboring time points along with the error in the estimate. The variance of an estimate for a given pair of time points was calculated as the inverse of the read depth at the earlier of the two timepoints + the estimated multiplicative noise parameter. Inverse variance weighting was then used to combine estimates across all time point pairs to generate a single fitness and error estimate for each lineage in each replicate. Replicate estimates were combined using further inverse variance weighting to generate the final fitness estimate for each isolate in the A- and C-conditions. For each mutant in each condition, we also calculated the 95% confidence interval around the fitness estimate based on the variability in fitness measurements between replicates (assuming that measurement errors are distributed normally).

3.4 Results

We obtained barcode frequency data at cycles 1, 2, 3, 4 and 5 (Figure S5). Fitness estimates of individual lineages were concordant between replicates (Figure S6), with relatively low estimation errors (Figure S7). Estimated fitness of sampled clones in both conditions along with errors and confidence intervals are provided in Data S1.

Determining valid clones for further analysis. Of the 581 clones that were pooled in the competition assays, we filtered out many clones from downstream analysis via a number of filters. First, clones that were sampled at cycle 17 were excluded from further analysis, as were clones that lacked a valid fitness measurement in either environment or had variance in fitness measurement in either environment ≥ 0.5. These filters removed 151 of 581 from consideration, leaving us with 430 clones for further analysis.

Calling adapted clones. We use the competition assay data to call adapted clones. A clone is called as adapted in a given environment if its estimated competitive fitness in that environment is more than two standard errors greater than 1 (2.3% FDR based on normal distribution). We identified 401 clones as adapted in the A-condition (see Data S1). 221 of them were sampled from the A-condition and we refer to them as the A-mutants. We identified 402 clones as adapted in the C-condition (see Data S1). 189 of them were sampled from the C-condition and we refer to them as the C-mutants. There

are 16 clones that are not adapted in either A or C condition, 13 clones that are adapted in the C-condition but not in the A-condition and 12 clones that are adapted in the A-condition but not in the C-condition. Of the 16 clones not adapted in either the A or C conditions, 12 come from lineages determined not to be adaptive in the BLT analysis, which we define as our neutral clones (Data S1). The genomes of 215 A-mutants, 181 C-mutants and eight neutral mutants were later sequenced, as described in Section 4.

Relationship between fitness estimated in competition assays and in BLT experiments. The correlation between fitness of adapted mutants (i.e., A- and C-mutants) estimated in the competition assays and those estimated from the BLT data is reasonably good (Figure S8, $R = 0.57$, $P = 8.77 \times 10^{-37}$), but there are also some systematic differences. Specifically, competition assays over-estimate BLT fitness in the A-condition and under-estimate it in the C-condition (Figure S8). We discuss possible reasons for these discrepancies below.

When we place mutants onto the bDFE in the non-home environment (as in Figure 2A in the main text) and when we estimate the probability of sampling a mutation in the non-home environment (see Section 5), we need to know the BLT fitness of mutants in their non-home environment. We have no direct measurements of BLT fitness of A- or C-mutants in their non-home environment, but we do have the non-home fitness estimates of all mutants in competition assays. However, directly substituting BLT fitness for competitive fitness would lead to biases due to the aforementioned discrepancies between the two estimates. To correct for these discrepancies, we linearly regress BLT fitness of A-mutants against their competitive fitness in the A-condition and we regress BLT fitness of C-mutants against their competitive fitness in the C-condition (see Figure S8). We then use these regressions to estimate the BLT fitness of C-mutants in the A-condition and A-mutants in the C-condition.

Difference in adaptive mutant fitness between conditions. Mutant's fitness were called as significantly different between A and C-conditions if the two 95% confidence intervals did not overlap. We discovered 178 such clones (108 A-mutants and 70 C-mutants), all of which were adaptive in both environments.

Possible reasons for the slight discrepancy between BLT and competitive fitness estimates. We can think of at least three possible reasons for this discrepancy. One possible reason is that we use a different set of reference lineages in the BLT experiments and competition assays. It is possible, for example, that some of the lineages that we use as reference in the competition assay are not in fact neutral in one or both of the conditions.

Another possible reason is that fitness is weakly frequency- dependent (this is expected in batch culture experiments [20]). Frequency dependence can manifest itself in a discrepancy between BLT and competition assays because lineages are present at much

higher frequencies in the competition assays than in the BLT experiments. Specifically, the median frequency of an adapted clone (as defined from the competition analysis) in the competition assay is 4×10^{-4} at the initial time point, whereas the median frequency of an adapted lineage is 3×10^{-6} at the initial time point (the frequency of the adapted mutant driving the lineage frequency must be even lower since many if not most adapted lineage also initially contain non-adapted individuals).

A third possible reason is that the media composition in the BLT experiments and in competition assays could be somewhat different at later stages of the growth cycle because population compositions are different and the media composition is determined by the collective metabolism of all variants present in the population.

Given the overall concordance of fitness estimates, we decided that dissecting these relatively minor effects was not particularly important within the scope of the present work.

4   Genome sequencing and analysis

We sequenced full genomes of 219 A-mutants, 187 C-mutants, 8 non-adapted evolved isolates and 24 ancestral isolates sampled from the inoculum population. Sequencing failed for four A-mutant and six C-mutants, leaving us with 215 A-mutants and 181 C-mutants with sequenced genomes, 8 non-adapted evolved isolates and 24 ancestral isolates (a total of 428 clones). For all remaining clones, we obtained high quality genome data ($> 4\text{x}$ coverage, mean coverage of 24x).

4.1   DNA extraction and library preparation and sequencing

DNA was extracted using a modified variant of the YeaStar yeast genomic DNA extraction kit Protocol I (Zymo research #D2002). In particular, a number of solutions were produced in-house, including YD Digestion buffer (1% SDS + 50mM Na_2PO_4), DNA Wash buffer (80% ethanol + 20mM NaCl) and Elution Buffer (10mM Tris-HCl). 0.2µL of 25mg/mL RNAse A (Zymo research #E1008-8) was used for each sample, as well as 1 Unit (0.2µL of 5U/µL stock solution) of Zymolyase (Zymo research #E1004).

Libraries were prepared using the method described in Ref. [13] and sequenced on the Illumina HiSeq4000 platform. Sequencing services were provided by Novogene Inc. and the UCSD Institute for Genomic Medicine.

4.2   Identification of new mutations

4.2.1   Small Variant calling

Reads were first trimmed using Trimmomatic with parameters

```
HEADCROP:10
ILLUMINACLIP:NexteraPE-PE.fa:2:30:10
LEADING:3
TRAILING:3
SLIDINGWINDOW:4:15
MINLEN:36
```

Reads were then mapped with bowtie2 (v. 2.3.4.3) using `-sensitive` parameters to the *Saccharomyces cerevisiae* reference genome (v. R64-2-1) with the addition of an extra "chromosome" defining the barcode locus. Reads were sorted, duplicates marked and short variants were called and filtered using GATK (v. 4.0.11.0) AddOrReplaceReadGroups, MarkDuplicates, HaplotypeCaller and VariantFiltration, respectively. Variant filtration used the filter expression

```
QD<10.0 || FS>20.0 || MQ<20.0 || AN>10 || AF<0.25 || QUAL<100.0 || DP<3
```

All other commands used default parameters. Variants were annotated with ENSEMBL Variant Effect Predictor using their command-line tool. As many variants had multiple possible annotations, coding sequence annotations ("missense_variant", "frameshift_variant", "stop_gained" and "stop_lost") were prioritized over synonymous annotations, which were prioritized over upstream noncoding annotations (within 2kb of a gene) and finally downstream noncoding annotations (again within 2kb of a gene). Variants further of 2kb of any gene or those within 2kb but with no annotations were removed as likely nonfunctional. Finally, variants with less than 3 reads of support for the derived allele were removed as putative false positives.

Additional filtering to remove erroneous and ancestral variants. The procedure described above identified 34,720 small variants across 428 sequenced isolates. We expect that many of these variants are sequencing and/or mapping errors that our procedure failed to remove, as well as fixed differences from the reference genome present in the ancestor of our experiment. To further filter out such spurious variants, we estimate the ancestral allele frequency spectrum from 24 sequenced ancestral clones and compare it with a typical allele frequency spectrum of 24 adapted mutants (averaged over 1000 random draws of 24 adapted mutants). As Figure S11 shows, the latter has an excess of variants that are present only in one clone, as expected for *de novo* mutations. The fact that there is no excess of mutations present in two or more adapted clones suggests that all or most mutations observed in two or more adapted clones are not adaptive. After removing 303 such variants, each of which was detected in 110 isolates on average, we are left with 1842 mutations across 428 sequenced strains (4.30 per clone), of which 33 strains carry no detectable derived small variants (Table S1 and Data S2).

4.2.2 Copy number variant calling

To identify large copy number variants (CNVs), we generate coverage plots for each sequenced clone by averaging read depth into 1kb windows with `bedtools genomecov`. An example plot is shown in Figure S9A. As coverage negatively correlates with the distance to telomeres (Figure S10), we re-calculate coverage after correcting for this variation (Figure S9B). We then manually identify CNVs from these corrected coverage plots by visualizing the coverage distribution at higher resolution (Figure S9C).

4.3 Identification of driver loci carrying small adaptive mutations

To differentiate adaptive "driver" mutations among residual ancestral and erroneous variants as well as nonadaptive "passenger" mutations, we rely on the idea of genetic parallelism, i.e., the fact that loci under selection gain mutations in independent lineages more often than expected by chance [21, 4]. We follow a similar procedure to the method described in Ref. [4]. For each gene, we define its *multiplicity* as the number of clones that carry a mutation in this gene. Since shorter genes require lower multiplicity to be called adaptive, we bin genes by their length into six 1kb bins plus one bin for genes with length $\geq$ 6kb. For each length bin $\ell = 1, 2, \ldots, 7$, we count the number of genes whose multiplicity is $m = 1, 2, \ldots$, denoting these counts by $k_{\ell m}$. We obtain the number of such genes $\langle k_{\ell m} \rangle$ expected in the absence of selection as follows.

We randomly and independently redistribute $N = 1718$ mutations (small mutations in adaptive clones after removing multiple mutations in the same gene in the same clone) across 7226 yeast genes 1000 times, with the probability for each mutation landing in a given gene being proportional to its length + 2kb. Then, the observed excess number of mutations with multiplicity m in length bin ℓ is $a_{\ell m} = \max \left\{ 0, k_{\ell m} - \langle k_{\ell m} \rangle \right\}$. These excess mutations are assumed to be adaptive. We then redistribute the remaining $\lfloor N - \sum_{\ell=1}^{7} \sum_m a_{\ell m} \rfloor$ potentially non-adaptive mutations as before, and identify additional excess mutations as adaptive. We repeat this procedure iteratively until the total number excess mutations is ≤ 1. We achieve convergence after 8 iterations and thereby obtain the final expected counts $\langle k_{\ell m} \rangle$.

Next, we would like to identify specific loci that carry adaptive mutations as those with multiplicities above some threshold. Specifically, we would like to determine multiplicity thresholds M_ℓ for each length bin $\ell = 1, \ldots, 7$, so that all loci in that bin with multiplicities $\geq M_\ell$ are called adaptive. To do so, for all ℓ we calculate the expected FDR at the given multiplicity threshold M_ℓ as

$$\beta(M_\ell) = \frac{\sum_{m \geq M_\ell} \langle k_{\ell m} \rangle}{\sum_{m' \geq M_\ell} k_{\ell m'}}.$$

We choose the multiplicity thresholds M_ℓ, so that $\beta(M_\ell) \leq \beta^*$ for all ℓ and for some desired β^*. We use $\beta^* = 10\%$.

We estimate the overall FDR as

$$\beta = \frac{\sum_{\ell=1}^{7} \sum_{m \geq M_\ell} \langle k_{\ell m} \rangle}{\sum_{\ell'=1}^{7} \sum_{m' \geq M_{\ell'}} k_{\ell' m'}}.$$

We conduct this analysis considering all 396 sequenced adaptive isolates together, as well as A- and C-mutants separately. Loci identified in any one of these three analyses are defined as putative adaptive loci. After identifying putative adaptive loci this way, we assume that all discovered mutations in these loci are adaptive (Data S3). We confirm that there is an excess of small adaptive mutations in adapted isolates when compared to ancestral isolates (Figure S12B).

4.4 Probability of genetic parallelism

We calculate the probability of genetic parallelism P_g for a set of mutants as follows. We consider every pair of mutants, and calculate the proportion that have a mutation in at least one common adaptive locus (including both small variants and CNVs). Adaptive isolates with no mutations at the identified adaptive loci are assumed to have an adaptive mutation at a locus that is not shared with any other isolate in the set.

To test whether A- and C-mutants have different probabilities of genetic parallelism, we reshuffle the evolution treatment label (i.e. Alone vs Community) across all A- and C-mutants and calculate P_g for both resulting groups. We obtain the absolute value of the difference in P_g between the two groups, $|\Delta P_g|$, in 1000 such permutations, and estimate the P-value as the fraction of permutations where $|\Delta P_g|$ exceeds the observed difference. We find that $P = 0.06$, suggesting that there is no significant difference in the probabilities of genetic parallelism between A- and C-mutants.

4.5 Results

Small mutations. The distributions of derived small variants per clone is shown in Figure S12A and their summary statistics are given in Table S1. These data show that the majority of small variants detected in the evolved clones are still likely ancestral or erroneous. We can estimate the expected number of adaptive small variants as follows. A typical adapted isolate carries 1.18 more mutations compared to an ancestral isolate (Figures S11, S12, Table S1). However, not all of these mutations may be adaptive. Indeed, given that small indels and single-nucleotide mutations occur at rate 3×10^{-3} per genome per generation [22], we expect 0.18 of such mutations to have occurred on the line of descent of any isolate sampled at cycle 9 (~ 60 generations). Based on this estimate, a typical adapted clone is expected to carry 1.18 - 0.18 = 1.00 adaptive mutation.

One potential problem with this estimate is that our yeast strain may have a somewhat different mutation rate than the strain used in Ref. [22]. We can obtain an alternative estimate for the number of adaptive mutations per clone by comparing adapted clones

with the neutral ones. We find that a typical neutral clone carries 1.96 ± 0.88 more mutations than a typical ancestral isolate[1] (Table S1), which is greater than 1.18 extra mutations carried by a typical adapted clone. Thus, it is possible that a typical adapted clone carries no beneficial small mutations.

The problem with the second estimate is that some of the clones identified as neutral may in fact carry adaptive mutations[2]. Therefore, we use both methods to obtain bounds on the number of adaptive mutations carried by a typical adapted clone and conclude that a typical adapted clone is expected to carry between 0 and 1 small adaptive mutations. Carrying out the same calculations for the A- and C-mutants separately, we estimate that a typical A-mutant carries between 0 and 1.34 small adaptive mutations (so that between 0 and 288 out of 1008 small mutations found in A-mutants are expected to be adaptive), and a typical C-mutant carries between 0 and 0.61 small adaptive mutations (so that between 0 and 110 out of 717 small mutations found in C-mutants are expected to be adaptive; Table S1).

Using genetic parallelism, we identify 185 mutations at 63 loci as adaptive (on average 0.47 mutations per clone), consistent with our expectation. 76 of these mutations at 39 loci are found in 60 A-mutants (on average 0.35 mutations per clone) and 109 mutations at 56 loci are found in 81 C-mutants (on average 0.60 mutations per clone; Table S1).

Large CNVs. We identified 176 CNVs across 167 strains (Data S2), of which 85 are aneuploidies. The distribution of CNVs per clone is shown in Figure S12C and summary statistics are given in Table S1. These data show that none of the ancestral or neutral isolates carry CNVs and that most adapted clones carry zero or one such mutation.

Since we found no CNV events in 32 sequenced ancestral and neutral isolates (Figure S12C, Table S1), we estimate the frequency of observing a non-adaptive CNV event as at most 1/32. Thus, we expect at most 12.4 such events among 396 sequenced adaptive clones. In fact, we found 176 CNV events, which suggests that all or almost all of them are adaptive (expected FDR ≤ 0.07; Data S3).

We found no statistical relationship between the number of CNV mutations and the number of identified small mutations per isolate ($R = -0.07$, $P = 0.18$).

Total number of identified adaptive mutations per clone. After combining small mutations and CNVs together, we find that a typical A-mutant carries 0.74 identified adaptive mutations with 100 A-mutants having no identified adaptive mutations (Figure S13 and Data S2). A typical C-mutant carries 1.11 identified adaptive mutations with 46 C-mutants having no identified adaptive mutations (Figure S13 and Data S2). Since all A- and C-mutants gained fitness in their home environment, each of them must

---

[1]This number is larger than 0.18 expected based on the yeast mutation rate, but the difference is not statistically significant ($P = 0.08$, t-test, expected $\mu = 0.18$).

[2]This is confirmed by the fact that we find two mutations at driver loci in the neutral clones (see Figure S12 and Table S1).

have at least one adaptive mutation. Therefore, the "unknown" sectors in Figure 2C in the main text refers to the numbers of A- or C-mutants without any identified adaptive mutations.

Pre-existing mutations. We use genetic information to test for the prevalence of preexisting mutations of type II, i.e., those that arose prior to the integration of the DNA barcodes, so that multiple lineages may carry an adaptive mutation identical by descent. If such mutations were prevalent, we would expect to observe an excess of adapted clones carrying identical genetic mutations, compared to ancestral isolates.

CNV events are not suitable for this analysis because they occur at very high rates. Indeed, many CNVs are aneuploidies, which occur at rate 9.7×10^{-4} per generation [22]. The only consistent segmental CNVs are ChrIV-1n and ChrIV-3n. ChrIV-1n mutations are localized to identical breakpoints across 64 mutants. However, these breakpoints are concordant with highly similar repetitive regions, YDRWTy2-2/YDRCTy1-2 and YDRWTy2-3/YDRCTy1-3. There are three different classes of ChrIV-3n amplifications across 10 mutants. At least some of these breakpoints also appear to be localized close to known repetitive regions, including YDRCTy2-1, YDRCTy1-1 and YDRWTy2-2. Thus, we suspect that these recombination-driven segmental events also occur at high rates and are not necessarily indicative of pre-existing variation.

As mentioned above, we find no excess of identical small mutations present in multiple adapted clones (Figure S11). In fact, an identical small mutation is found less frequently in multiple adapted clones than in multiple ancestral clones, suggesting that some of the pre-existing genetic variation may have been deleterious.

5 Simulations of evolutionary dynamics and the estimation of rates of adaptive mutations

We found that the sets of A- and C-mutants are genetically distinct (Figures 2C in the main text, S13 and Data S3), despite all of them being more fit than the ancestor in both environments. In this section, we show that this somewhat puzzling observation can be explained by the differences in the evolutionary dynamics in the concurrent mutation regime [23].

There are two key differences between the A- and C-conditions. First, the bDFEs are different (Figure 2A in the main text), resulting in different increases of population's mean fitness over time. Second, the fitness rank orders of adaptive mutations are also different (Figure 2A in the main text). These two facts imply that in the concurrent mutation regime the chances for a given mutation to escape drift while rare and reach a certain frequency and be sampled can be substantially different in the two conditions. Next, we develop a quantitative version of this argument.

As described in Section 4, we classify all discovered adaptive mutations into mutation classes by the type of CNV or the gene in or near which the mutation occurred (Data S3). Let k_{im}^A and k_{im}^C be the number of sequenced A- or C-mutants that carry a mutation from class m sampled from the replicate population $i = 1, 2, 3, 4, 5$ (see Table S2). We would like to know whether the differences between k_{im}^A and k_{im}^C can be explained by the observed differences in the bDFEs and by differences in the fitness benefits provided by mutations of class m in the A- and C-conditions.

The challenge is that the probability $\Pr \{k; s, U\}$ of observing k mutants of a certain type in a sample from a given population depends not only on the selection coefficients s of adaptive mutations of this type (which we have measured, as described in Section 3) but also on the unknown rate U at which such mutations arise. Therefore, we first find the mutation rates that fit our data and confirm that these rates are biologically plausible. We then ask how well the expected numbers of A- and C-mutants carrying specific mutations match the observed numbers k_{im}^A and k_{im}^C, given the estimated mutation rates.

5.1 Data

Exclusion of pre-existing mutations. A number of lineages are identified as adapted in multiple populations, indicative of pre-existing mutations of type I (see Section 2.6.2 for more details). Since such mutations do not carry information about the mutation rate, we exclude them from this analysis. That is, numbers k_{im}^A, k_{im}^C given in Table S2 and in Figure S15 are the numbers of sampled and sequenced clones with unique barcodes.

Selection coefficients. Fitness of individual mutants in the BLT experiments are estimated based on the competition assay data, as described above (see Section 3.4), and selection coefficients s_m^A and s_m^C are taken as averages over all mutants carrying a mutation from a given mutation class (Table S2).

Mean fitness trajectories. Our model described below takes into account the fact that the probability of a newly arisen beneficial mutation to survive genetic drift depends on how the mean fitness of the population changes over time. Thus, when simulating the evolutionary dynamics, we use the empirical mean selection coefficient trajectories $\bar{s}^A(t)$ and $\bar{s}^C(t)$ in the A- and C-conditions, respectively. $\bar{s}^A(t)$ and $\bar{s}^C(t)$ are computed by first averaging the values of the mean selection coefficient (estimated in as described in Section 2.6.2) over all five replicates and then fitting the logistic function to these data points (see Figure S3B).

5.2 Model

To keep the model as simple as possible, we assume that our population has the constant size $N = 2 \times 10^6$ in both A- and C-conditions, mutations from the mutation class m arise at (an unknown) rate U_m per division in both conditions. We assume that all mutation from class m confer the same (known) fitness benefits s_m^A and s_m^B relative to the ancestor in the A- and C-condition, respectively (Table S2. We also assume that all mutants are sampled at time $T = 60$ generations after the beginning of the experiments in both A- and C-conditions, which roughly corresponds to cycle 9, assuming $\log_2 100 = 6.64$ doublings per cycle (see Section 2.3).

To estimate the mutation rate U_m for mutation class m we define the log-likelihood function

$$L(U_m) = \frac{1}{5} \sum_{i=1}^{5} \log \Pr \left\{ k_{im}^A; s_m^A, U_m \right\} + \frac{1}{5} \sum_{i=1}^{5} \log \Pr \left\{ k_{im}^C; s_m^C, U_m \right\}. \tag{2}$$

The sampling probability $\Pr \{k; s, U\}$ depends on the number of lineages that independently arose in the population and that carry a mutation of the focal type, as well as on the frequencies of these lineages at the sampling time point T. It is difficult to obtain an analytical expression for this probability because it requires integrating over all these nuisance parameters. We therefore compute the sampling probability numerically, using population dynamic simulations described below.

5.2.1 Evolutionary dynamics simulations and the estimation of the sampling probability

As mentioned above, difference in the bDFE between the A- and C-conditions may contribute to the observed differences in the genetic composition of sampled mutants. The bDFE indirectly affects the survival probability of a newly arisen beneficial mutation by altering how the mean fitness of the population changes over time. Thus, we design our simulations so that they match the average empirical mean selection coefficient trajectories in the A- or C-condition (see Section 2.6.2 and Figure S3B). Matching these trajectories in simulations of the full Wright-Fisher where all mutation classes segregate simultaneously is difficult. Instead, we simulate the arrival and spread of mutations of each mutation class separately, while accounting for changes in mean fitness using our logistic fit of the observed mean fitness trajectories (Figure S3B). This approach also allows us to use branching process approximations described below in Section 5.2.2, which greatly speed up the calculations.

Since in this section we are concerned with mutants of one mutational class in one environment, we omit the subscript m and superscripts A/C. In other words, we assume that the mutations arrive at rate U per individual per generation and have selection coefficient s, and we sample 88 mutants from this population at time point $T = 60$.

In our simulations, we allow for new mutations to arise between 1 cycle prior to the beginning of the experiment up to cycle 9. Thus, we divide the time interval between -6.6 and $T = 60$ generations into 260 to 1300 segments of length $\Delta t = 0.01/s$ generations. For each time segment $(t_i, t_i + \Delta t)$, we draw the number of new mutants arising in the population in that segment from the Poisson distribution with rate $N U \Delta t$. Each of these mutants survives until the sampling time point T with probability $P_{\mathrm{surv}}(T; s, t_i)$, given by equation (3) below. For each mutant that survives, we draw its establishment time τ from the distribution given by equation (4) below. We then set the mutant's frequency at the sampling time point to $n(T; s_m, \tau)/N$, where n is given by equation (5) below. At the end of each simulation run, we obtain a list of frequencies of all independently arisen mutants carrying mutation of type m. We assume that each independently arisen mutants is linked to a distinct barcode. We then randomly sample 88 clones from the whole population and discard those that do not carry the focal adaptive mutation. If multiple clones are sampled from the same lineage, only one is retained. We further randomly sub-sample these clones with 97.5% success probability, simulating the small whole-genome sequencing failure rate, which results in the final number k of sampled clones that carry a mutation from the focal mutation class. We estimate the probability of sampling k clones, $\Pr\{k; s, U\}$, by running 10^4 simulations and recording the fractions of simulations where we observe k sampled clones.

5.2.2   Branching process approximations for mutant growth dynamics

Consider a mutant that arises at some point $t_0 < T$ on the ancestral background. The early population dynamics of such mutant lineage can be modeled as a branching process [23]. In particular, Desai and Fisher [23] used the branching process approximation to derive the probability that a mutant with selection coefficient s that arose at time zero in the ancestral population (i.e., whose mean selection coefficient is zero) has not gone extinct by time t and the related probability that the mutant "establishes" at time τ (see equations (11) and (17) in Ref. [23]). However, this model ignores the fact that the mean fitness of the population is changing over time while the focal mutant is still at low frequency. In our experiment, mean fitness changes very rapidly, as can be seen in Figure S3B, which can significantly alter mutant's survival probability and its establishment time. Assuming that the mean selection coefficient trajectory $\bar{s}(t)$ is known (see Section 5.1), we can model mutant dynamics analogous to Ref. [23] but with a generalized birth-death model with growth rate $1 + s - \bar{s}(t)$ and death rate 1. In this model, the probability that a mutant that arose at time t_0 survives until time t is given by

$$P_{\mathrm{surv}}(t; s, t_0) = \frac{\omega(t_0)}{1 + \omega(t_0) - e^{-\rho(t, t_0)}}, \tag{3}$$

where

$$\omega(t) = s - \bar{s}(t),$$

$$\rho(t, t_0) = \int_{t_0}^{t} \omega(t') \, dt',$$

provided that $\omega(t_0) > 0$. Conditional on surviving, the probability that this mutant establishes at time τ is given by

$$P_{\text{est}}(\tau; s, t_0) = \frac{\omega(t_0)}{1 + \omega(t_0)} \, \exp\left[-\rho(\tau, t_0) - \frac{e^{-\rho(\tau, t_0)}}{1 + \omega(t_0)}\right]. \tag{4}$$

Once the mutant establishes at time τ and provided that it is still beneficial, i.e., $\omega(\tau) > 0$, its subsequent dynamics are essentially deterministic, so that its size $n(t; \tau)$ at time t is approximately given by

$$n(t; s, \tau) = \frac{e^{\rho(t, \tau)}}{\omega(\tau)}, \tag{5}$$

which is analogous to equation (14) in Ref. [23] in the limit $t \gg 1/s$.

5.2.3 Estimation of mutation rates by maximum likelihood

For each mutation class m, we compute the likelihood function $L(U_m)$ using equation (2) for 40 discrete values of $u_m = \log_{10} U_m$ between -7 and -3 and interpolate between them using a polynomial of degree 4. The likelihood function for chrIII-3n mutations is shown as an example in Figure S14. We find the maximum of this function as a point where the polynomial approximation of L has a zero first derivative with respect to u_m. We estimate the standard error (SE) as the inverse of the square root of Fisher information [24].

5.3   Results

We estimated the rates of driver mutations at 9 loci that were mutated in at least five adapted clones, after excluding gene NUM1 where all mutations are likely pre-existing (see Figure S13, Table S2 and Data S3). Our estimates are generally consistent with those derived from a published mutation accumulation experiment by Zhu et al [22]. In particular, our estimate of the rate of small adaptive mutations in genes HEM1 and HEM2 is $\sim 3 \times 10^{-7}$ per generation, consistent with per basepair mutation rate of 1.67×10^{-10} estimated by Zhu et al [22]. Gene YIL169C is an exception, with an estimated rate of adaptive point mutations of 7.76×10^{-6} per generation. We estimate the rates of large CNV events to be $\sim 10^{-5}$ per generation, again consistent with the genome-wide rate of aneuploidies of 1.04×10^{-4} estimated by Zhu et al [22]. Our estimate of the segmental

duplications ChrIV-3n is much lower (3.24×10^{-7} per generation), also consistent with Ref. [22], although they do not provide a quantitative estimate of such events.

The probability that at least one sequenced mutant from all five replicate populations has a mutation with a given selection coefficient s and mutation rate U is shown in Figure S15A, for both A- and C-conditions. As expected, this function is different between conditions because the mean fitness dynamics are different (Figure S3B). Finally, we directly compare the observed numbers of mutations of each mutation class with the expected numbers in the A- and C-conditions and find a reasonable match (Figure S15B).

6 Phenotyping

For all phenotypic measurements, replicate measurements were conducted using distinct samples. In no case was a single sample measured repeatedly.

6.1 Measurement of ancestral yeast and alga growth (main text Figure 1)

Yeast and algae were thawed from frozen stocks ($-80°$C for yeast, LN2 storage for algae) and grown separately in standard growth conditions for one growth cycle. Communities were inoculated by mixing 100 µL of each species into 10 mL of our standard growth culture and further propagated for 2 growth cycles. On the third growth cycle, each culture (yeast alone, algae alone and the community, with two replicates each) was characterized daily by CFU counting. Each culture was plated onto both CYM media plates and TAP media (tris-acetate-phosphate) plates (1% agarose for each) with serial dilution. CYM plates were incubated at $30°$C while TAP plates were allowed to grow at room temperature under ambient lighting. Colonies were counted to estimate the density of both yeast and algae (as appropriate) in each culture at each timepoint.

6.2 Measurements of mutant yeast alone and in community (main text Figures 3-5)

We selected 31 C-mutants and 28 A-mutants for three types of phenotyping experiments: measurements of yields in the community, microscopy for detecting possible physical associations between yeast and algae, and growth-curve measurements of individual mutants in the A-condition. We selected these 59 mutants manually to cover a diversity of mutations and fitness values represented in the sampled adaptive mutants (see Data S3 for the number of mutants selected from each mutation class). Our reasoning for this selection rather sampling randomly is that mutants carrying a driver mutation at the same

locus would have similar phenotypic values[3]. By selecting clones for phenotyping randomly, we would not observe more rare phenotypes. To account for this over-dispersion in the selection of mutants we apply the mutation weighting procedure described below in Section 6.7.

6.3   Measurements of YYA and AYA

To estimate YYA, we inoculated all 60 yeast strains (including the ancestor) individually into the standard conditions (10 mL of media in 50 mL flasks) from frozen stocks and propagated them for two cycles (10 days). During the third cycle, we estimated yeast densities after 5 days of growth by plating and colony counting as described in Section 6.4. AYA estimate is obtained as described in Section 6.1. These measurements are shown in Figure S16A.

6.4   Measurements of YYC and AYC

We create mutants communities as follows. On day 0, we inoculate 50 mL growth media (in 250 mL non-baffled Erlenmeyer flask) with 1 mL of CC1690 *C. reinhardtii* stock stored in a liquid nitrogen freezer and incubate for 20 days. On day 20, 100 µL of this culture are transferred to the standard conditions (10 mL of fresh media in a 50 mL flask). Also on day 20, the 60 yeast strains (including the ancestor) are individually inoculated into the standard condition from frozen stocks. On day 25, we form the mutant communities by transferring 100 µL of each yeast culture and 200 µL of the algae culture into fresh media (200 µL of alga were used instead of 100 µL because the density of algae culture was approximately 50% of that at the initiation of the BLT experiment). These mutant communities are grown for one cycle in our standard conditions. On day 30, we transfer 100 µL of each mutant community into 10 mL fresh media, as in the BLT experiment. We estimate both yeast and alga density on day 35, as follows. We estimate alga density by measuring chlorophyll b fluorescence, which is linearly related to cell density [25]. To that end, we transfer 200µL of each culture into a well of a black-wall clear-bottom 96-well plate (Corning) and measure fluorescence in a plate reader (Molecular Devices Spectramax i3x, excitation at 435nm and observation at 670nm). AYC estimates are obtained by subtracting the background fluorescence value from the measured fluorescence values and multiplying the results by 1.267, which is the ratio of the CFU/mL to fluorescence in the ancestral community (see Section 6.1). To estimate YYC, we dilute the culture 1:1600,

---

[3]We have some support for this hypothesis. We tested whether mutants carrying the same mutation have more similar yeast and alga yields, r and K values than random mutants. We carried out four such tests for the 9 clones with chrIII-3n mutations, 5 clones with the chrIV-3n mutations, 3 clones with the chrIV-1n mutations, 3 clones with HEM2 mutations and 3 clones with HEM3 mutations. In all cases, clones carrying the same mutation had more similar phenotypes than random mutants, but in most cases the difference was not statistically significant, presumably due to lack of power.

plate 100 µL on Petri dishes with standard media + 1% agarose, grow at 30°C for 3 days and count colonies.

Yield estimates can be found in Data S1. Replicate measurements show reasonably good correlation (Figure S16). We find a positive correlation between Alga yield and yeast mutant fitness in both the A- and C-conditions (Figure S18).

6.5 Microscopy

To detect potential physical associations between algae and beneficial yeast mutants, we created mutant communities as described in Section 6.4. Communities were mounted on glass microscope slides (Fisher Scientific 12550143), sealed with Dow Corning high vacuum grease (Amazon B001UHMNW0) and imaged on a light microscope using a 20x objective with DIC. Figure S19 shows one representative community, the remaining 17 imaged mutants along with WT controls can be found in the Dryad data repository.

6.6 Mutant growth curve measurements

To estimate the growth parameters r and K for individual beneficial mutants, we carried out growth curve measurement experiments of individual yeast mutants and the ancestor in the A-condition. To this end, we inoculated all 60 yeast strains (including the ancestor) individually into the BLT condition (10mL of media in 50mL flasks) from frozen stocks and propagated them for two cycles (10 days). During the third cycle, we estimated yeast densities on days 10, 10.5, 11, 11.5, 12, 13, 14 and 15 by plating and colony counting as described in Section 6.4. The growth curves are shown in Figure S20 and the data are provided in Data S1.

We estimate r as the slope of the relationship between $\log\,(\mathrm{CFU/mL})$ and time (in hours) for the three measurements between 12 and 36 hours of growth. We estimate K as the maximum observed density (in CFU/mL). r and K estimates can be found in Data S1. Growth rate estimates are well correlated between replicates (Figure S21A). K measurements are more noisy (Figure S21B). Yield measurements had very poor replication from this experiment (Figure S20), so the experiment was repeated with two additional replicates specifically to estimate the yield of these yeast mutants when grown in isolation (Figure S21C). The relationship between these growth parameters, mutant fitness and species yields in mutant communities are shown in Figure 5 in the main text and Figure S22.

6.7 Statistical analysis

Mutation weighting. Even though not all A- or C-mutants were phenotyped, we would like to make certain statistical statements about the distribution of phenotypes among all sampled A- or C-mutants. We can do so by weighting each phenotyped mutant

according to its mutational class. We do so as follows. First, we associate each of the 59 phenotyped mutants with a single driver mutation. Mutants with multiple driver mutations are associated only with the driver mutation from the most common mutational class. Mutants with no identified driver mutations are associated with a unique unknown mutations. Then, for each of the phenotyped mutants, we weight its measured phenotypic value by the number of A- or C-mutants that share a mutation class with it, divided by the number of phenotyped A- or C-mutants that share a mutation class with it. Based on this reweighted distribution of phenotypes, we calculate various statistics such as kernel density estimates, correlation between phenotypes and probabilities of parallelism (see below).

Kernel density estimation. We use kernel density estimate (KDE) to determine how likely certain phenotypic trait values would occur among random A- and/or C-mutants. Specifically, we obtain the kernel density estimates $D_A(y, a)$ and $D_C(y, a)$ for the probabilities that a community formed by algae and a random A- or C-mutant, respectively, would produce a yeast yield y and alga yield a. To estimate D_A, we apply the `kde2d` function in R with bandwidth 1 along the x-axis and 4/3 along the y-axis to the mutation-weighted yield data for the A-mutants. We analogously obtain $D_C(y, a)$.

Accounting for measurement errors in statistical tests by permutation testing. As the measurement errors in our estimates of phenotypic values (r, K, yeast yield and algae yield) are quite large, we use a permutation and resampling procedure to determine the statistical significance in various tests involving these variables. In this procedure, we permute isolate labels and resample the phenotypic values associated with each isolate from the normal distribution with the estimated mean and with the standard deviation set to the estimated standard error of the mean. We carry out 1000 permutation and resampling instances in each test. For FDR analysis, we define a threshold p value (e.g. 0.05) and conduct the statistical test with both the resampled phenotypic values and the observed phenotypic values. We count the fraction of resampled values that are significant at our given threshold to estimate the false positive rate (averaged over the 1000 permutations), and then divide the number of observed positives by this estimate to compute the false discovery rate (FDR). Permutation tests of Pearson correlation significance are conducted as follows. For each permutation, isolate labels for each variable are permuted independently and phenotypic values are resampled as above. A randomized correlation coefficient is determined from each of the 1000 permutations, and significance is determined by the proportion of randomized R^2 values that exceed the observed R^2 value.

Phenotypic parallelism analysis. We quantify the degree of parallelism among a set of mutants with respect to a pair of quantitative traits X and Y by estimating the probability that, for two randomly selected mutants, trait X changes in the same direction

in both mutants and trait Y changes in the same direction in both mutants. Mathematically, if one randomly selected mutant has trait increments ΔX_i and ΔY_i relative to the ancestor and the other mutant has trait increments ΔX_j and ΔY_j, we estimate the probability that both $(\Delta X_i)(\Delta X_j) \geq 0$ and $(\Delta Y_i)(\Delta Y_j) \geq 0$. This measure of phenotypic parallelism emphasizes the direction of change rather than the magnitude, which, we believe, is more meaningful than some other quantitative measures, at least for the community and growth phenotypes that we consider in this study. When we compute this measure for the pair of yields of mutant communities, in which case X and Y are yeast and alga yields, we refer to it as the probability of ecological parallelism, P_e. When we compute this measure for the growth phenotypes, in which case X and Y are r and K, we refer to it as the probability of phenotypic parallelism, P_ph.

We obtain these probabilities of parallelism for the A- and C-mutants. We test the significance of the deviation of these probabilities from the expectation of 25% parallelism via a chi-square test. To determine the statistical significance of the difference between the probabilities of parallelism for the A- and C-mutants, we resample the phenotypic values of each of the 59 mutants from the normal distribution (see above) and permute mutant genotype and home-environment labels, so that the genotype and label are always associated with each other but dissociated from the phenotypic values. We then calculate the parallelism probabilities for these permuted and resampled data. We estimate the P-value by carrying out this permutation and resampling procedure 1000 times.

7 Possible function effects of mutations in *HEM1*, *HEM2* and *HEM3* genes

Three critical components of the heme biosynthesis pathway, HEM1, HEM2 and HEM3 are putative targets of adaptation in this study, and provide a substantially larger fitness benefit in the C-condition than in the A-condition (see Data S2 and Data S3). Most mutations found in these genes probably compromise the function of the encoded enzymes and lead to a decreased "siphoning" of succinyl-coenzyme A (sCoA) from the TCA cycle and the production of less heme and/or decoupling of aerobic-driven regulation from aspects of central metabolism. Although yeast is already capable of performing fermentation under ambient oxygen concentration, we speculate that these mutations enables yeast to ferment under even higher oxygen concentrations that results from alga photosynthesis.

7.1 HEM1

HEM1 encodes the mitochondrial 5-aminolevulinate synthase (ALAS) that catalyzes the first committed step of porphyrin through the condensation of glycine and sCoA using the cofactor pyridoxal 5'-phosphate (PLP):

$$\text{sCoA} + \text{glycine} + \text{PLPcofactor} \rightleftharpoons \text{5-aminolevulinate} + \text{CoA} + CO_2 + \text{PLPcofactor}$$

ALAS links heme/cytochrome production with the TCA cycle and aerobic respiration via sCoA, and thus plays a key role in cellular energetics [26]. ALAS functions as a homodimer (with subunits referred to as A and B below). A crystal structure of *S. cerevisiae* HEM1 is available (https://www.rcsb.org/3d-view/5TXR/1).

We found 5 adaptive mutations in the *HEM1* gene, four of which occurred in the C-mutants (Data S2 and Data S3). A nonsense mutation at residue 166 (out of 548) in HEM1/ALAS leads to a truncated coding sequence and presumably loss of function. All other mutations alter amino acids that are conserved in all sequenced *S. cerevisiae* strains, which we speculate may also significantly compromise normal HEM1/ALAS function, as described below.

His107Pro mutation does not seem to be involved in any substrate or co-factor binding directly, but a change from histidine to a proline, with significant backbone confirmational constraints may have serious consequences for the folding (and thus function) of the enzyme. Bracketing this position (107) nearby is Arg91, which plays a critical role in sCoA binding, and Asn121, which forms an alpha-carboxylate hydrogen bond with the glycine substrate [27]. Moreover, His107 (of subunit B) forms important structurally stabilizing side-chain interactions with Glu111 (subunit B) and Lys142 (subunit A) (see Figure S3 of [27]), which may very well be disrupted with a proline substitution (lacking a side chain).

Asn152Lys and Asn157Lys mutations occur in a region of the ALAS protein that becomes ordered upon PLP cofactor binding. N152 plays a direct role in coordinating sCoA [27], hydrogen bonding with the carboxylate (COO-) moiety of the sCoA succinyl group. Mutation from asparagine to a positively charged lysine may both disrupt the formation of this hydrogen bond and prevent a key side-chain interaction between Asn152 (subunit A) and Arg91 (subunit B) that stabilizes key structural elements needed for PLP cofactor binding (we expect the side chain of mutation Lys152 to repel that of Arg91 since both are positively charged). Asn157Lys is a few amino acids downstream from Ile153, which also plays a role in carboxylate/sCoA coordination like Asn121 [27]. Asn157 is adjacent to Ile158 (subunit A) that forms stabilizing main-chain and side-chain interactions upon PLP cofactor binding with Asn95 and Asn 97 (both on subunit B); mutation to Lys may disrupt these interactions, along with that of nearby stabilizing interactions between Arg98 (subunit B) and Ala147 (subunit A), due to positive charge repulsion between Arg98 and Asn157Lys (see Figure S3 of [27]).

Gly344Cys mutation occurs at a site that does not directly bind substrate or co-factor but is each bracketed by amino acids that do play key active site roles and could also have structural consequences as conformational flexibility is likely lost with the change away

from glycine. Gly344 is flanked (although several amino acids away) by K337 that forms a critical covalent pyridoxyl-lysine bond, and F365 which delineates the sCoA substrate-binding pocket [27]. Gly344 resides on the juncture between a beta-sheet and alpha-helix motif in the protein structure, with little space for a side chain. Mutation of this site to a cysteine may alter backbone flexibility and the folding of key structural elements. Gly344 is also in close proximity to Cys182 that forms stabilizing hydrogen bonds with amino acids Asn129, Thr275, and Gly 276; having another cysteine nearby may very likely interfere with that hydrogen bonding network.

7.2 HEM2

HEM2 encodes for the cytoplasmic/nuclear, homo-octameric aminolevulinate dehydratase (ALAD; porphobilinogen synthase), which catalyzes the second step in heme biosynthesis:

$$5\text{-aminolevulinate} \rightleftharpoons \text{porphobilinogen} + 2\,H_2O$$

A crystal structure of *S. cerevisiae* HEM2 is also available (https://www.rcsb.org/3d-view/1AW5/1).

We found 5 adaptive mutations in the *HEM2* gene, four of which also occurred in the C-mutants (Data S2 and Data S3). Two frame-shift mutations lead to premature stop codons ~ 100 out of 342 amino acids. The three remaining mutations Ile129Phe, Val132Phe, and Pro264Arg are predicted to have moderate impact.

Ile129   resides on a beta-strand with a side chain embedded in a relatively hydrophobic environment. It is unclear what the consequences of mutation Ile129Phe might be, as phenylalanine is a comparably bulky hydrophobic side chain to isoleucine that would at first glance seem to be a conservative substitution [28]. However, several aromatic amino acids are in the vicinity, including Trp30 and Tyr127; Ile129Phe may form aromatic ring-stacking interactions with these residues to disrupt fold structure.

Val132   is similarly on a beta-strand with a side chain embedded in another nearby hydrophobic pocket and interface with other non-polar amino acids on alpha-helix. The Val132Phe mutation could disrupt the tight packing at this interface as phenylalanine is substantially bulkier than valine. Moreover, Tyr168 is nearby which hydrogen bonds a key water molecule that is hydrogen-bonded to several other backbone atoms flanking Val132; the Val132Phe mutation may also disrupt this by interacting with Tyr168 through aromatic stacking interactions. Pro264 resides in a sharply kinked beta-turn between beta-strand and alpha-helix elements, which may be important for constraining and enabling key interactions of flanking residues with amino acids distributed across different structural elements: Val262/Ser265/Tyr287, Ser265/Glu292, and Lys263/Tyr287 (see https://www.rcsb.org/3d-view/1AW5/1).

Pro264Arg mutation is pronounced not just for the loss of backbone constraint provided by the imino acid proline [28], but it introduces a large positively charged side change that may form "inappropriate" interactions to disrupt monomer fold and subsequently oligomerization; these include interactions with nearby: negatively charged amino acids: Glu292 and Glu313 aromatic amino acids (through cation-pi interactions) Phe211 (normally interacting with Tyr268 via aromatic stacking), Tyr216 (normally interacting with the backbone of Phe211), Tyr268 (normally interacting with Glu313), and Tyr287 (normally interacting with Lys263 via cation-pi interactions).

7.3   HEM3

HEM3 encodes for the cytoplasmic/nuclear enzyme, porphobilinogen deaminase/hydroxymethylbilane synthase (HMBS), which catalyzes the third step in heme biosynthesis involving four molecules of porphobilinogen to make a linear hydroxymethybilane molecule that looks like a heme/tetrapyrole when "wrapped around":

$$4\,\text{porphobilinogen} + \text{H}_2\text{O} \rightleftharpoons \text{hydroxymethylbilane} + 4\,\text{NH}_3$$

We found 4 adaptive mutations in the *HEM3* gene, three of which occurred in the C-mutants (Data S2 and Data S3). However, since no structure exists for *S. cerevisiae* HEM3 protein and since the homology with the human ortholog for which the structure does exist (https://www.rcsb.org/3d-view/5M6R/1) is rather low ($\sim 40\%$ similarity), predicting the functional effects of these mutations is challenging.

8 Supplementary Figures

Figure S1. Lineage trajectories in yeast populations in the A-condition. First five panels show individual replicate populations A1 through A5. Barcode lineage frequencies are estimated at every odd cycle. Only the lineages identified as adapted are shown (black lines). Thick red lines show the total frequency of all non-adapted lineages. Last panel shows the number of nonextinct lineages over time for all populations.

Figure S2. Lineage trajectories in yeast populations in the C-condition. Notations as in Figure S1.

Figure S3. Estimation of fitness of adapted lineages in the BLT data. A. The mean lineage fitness (*x*-axis) is plotted against the coefficient of variation (CV) of the estimate (*y*-axis). Blue (red) points indicate lineages identified as adapted (non-adapted). **B.** Mean fitness of each population over time. Individual populations are shown in grey, the average mean fitness trajectory over all populations is shown in black, and a logistic fit to this average trajectory is shown in red. **C, D.** Examples of selection coefficient estimation for one lineage in one A population (C) and one C population (D).

Figure S4. Evidence for pre-existing mutations in the BLT experiments. For each population on the x-axis, we plot the fraction of lineages identified as adapted in that population which are also identified as adapted in every other population (y-axis, black points) as well as the expectation for this fraction (red points). The expected overlap between populations i and j is calculated by comparing the observed adaptive lineages in population i to a random set of lineages from population j that reach detectable frequency. All lineages are considered equally for sampling.

Figure S5. Lineage frequency trajectories in the competition assay. Reference lineages are shown in green, all other lineages are shown in orange.

Figure S6. Correlation between fitness estimates across replicates of the competition assay. Line shows the diagonal.

Figure S7. Mean and the coefficient of variation of the fitness estimates in the competition assay.

Figure S8. Correlation between BLT and competitive fitness estimates. Solid and dashed lines show linear regressions for the A-mutants (in the A-condition) and C-mutants (in the C-condition), respectively.

Figure S9. An example illustrating the detection of large chromosomal duplications and deletions in the genome sequence data. A. 1kb-window coverage plot for clone 386817 from population C1. **B.** Same data, after normalization using the relationship between coverage and chromosomal position shown in Figure S10. **C.** Focusing on chrIV to manually identify breakpoints for the chrIV-1n mutation (red lines).

Figure S10. Relationship between coverage and chromosome position. Mean coverage in 1kb windows for clone 386817 from population C1 is shown vs distance of locus from the end of a chromosome.

Figure S11. Allele frequency spectrum for ancestral and adapted isolates. Each bar shows the number of small mutations (point mutations and small indels) found in a given number of ancestral or adapted clones. The distribution for adapted clones is averaged over 1000 random draws of 24 clones.

Figure S12. Distribution of mutations per clone. A. Number of small mutations detected per sequenced isolate. Triangles on the x-axis indicate the means of each distribution (see also Table S1). **B.** Distribution of the number of identified adaptive small mutations per isolate. **C.** Distribution of the number of CNVs per isolate.

Figure S13. Distribution of adaptive mutations across the most common driver loci. Only driver loci with 5 or more mutations are shown (see Data S3 for the full distribution). Colors correspond to Figure 1 in the main text.

Figure S14. The log-likelihood function for the estimation of the chrIII-3n beneficial mutation rate.

Figure S15. Sampling probabilities for adaptive mutations at the most common driver loci. A. Shades of gray show the probability of sampling at least one clone at a locus with a given beneficial mutation rate and fitness in the A-condition. The most common loci are shown by points with the same colors as in Figure 1 in the main text. The beneficial mutation rate and the selection coefficient for each mutation class are given in Table S2. **B.** Same as A but for the C-condition. The mutation rate for each locus is assumed to be the same in both conditions, but the selection coefficients vary. **C.** The observed number of sequenced clones per replicate population with a mutation in each class (black points) and the numbers expected in our simulations (bars and whiskers) given the measured selection coefficients and the estimated mutation rates provided in Table S2. Bars show IQR (Q3 − Q1), whiskers show Q1 − 1.5 × IQR and Q3 + 1.5 × IQR.

Figure S16. Correlation between yield measurements across replicates. The solid line shows y=x. Normalization is calculated relative to the ancestral strain.

Figure S17. Estimated yeast and alga yields among random A- and C-mutants.
The heatmap shows the ratio of probabilities of observing a given pair of yields among random
C- and A-mutants. Data points are identical to Figure 3B in the main text. D_A and D_C were
estimated from these data as described in Section 6.7. Any points where either D_A or D_C fall
below 0.03 are colored gray. Normalization is calculated relative to the ancestral strain.

Figure S18. Correlation between yeast mutant fitness in the original community and alga yield in the mutant community. Normalization is calculated relative to the ancestral strain.

Figure S19. Representative microscopy image showing lack of physical associations between yeast and alga cells. Mutant culture formed by the C-mutant C2 109098 is shown. Some yeast and alga cells are indicated with arrows.

Figure S20. Mutant growth curves in the A-condition. Growth curves for the 59 adapted yeast mutants are shown, from which r and K values are estimated as described in Section 6.6. **A.** Replicate 1. **B.** Replicate 2.

Figure S21. Correlation between replicates for r and K measurements.

Figure S22. Relationship between growth parameters, yields and fitness. A. Δr vs fitness of mutants in the A-condition. **B.** ΔK vs fitness of mutants in the A-condition. **C.** Δr vs fitness in the C-condition. **D.** ΔK vs alga yield in mutant communities. **E.** Δr vs yeast yield in mutant communities. **F.** Δr vs alga yield in mutant communities. In all panels, normalization is calculated relative to the ancestral strain.

9   Supplementary Tables

Isolates		Small variants				CNVs	
				Adaptive			
Type	Number	Total	Per isolate	Expected	Identified	Total	Per isolate
Ancestral	24	76	3.17 ± 0.51	–	3	0	–
Neutral	8	41	5.13 ± 0.72	–	2	0	–
A-mutants	215	1008	4.69 ± 0.20	$(0, 288)$	76	84	0.39 ± 0.04
C-mutants	181	717	3.96 ± 0.22	$(0, 110)$	109	92	0.51 ± 0.04
Total	428	1842	–	–	190	176	–

Table S1. Statistics of mutational counts among sequenced isolates

Culture	Seq. isolates	Mutation class m/Locus								
		chrIV-1n	chrIII-3n	chrXIV-3n	chrIV-3n	chrIX-3n	chrXIII-3n	YIL169C	HEM1	HEM2
A1	13	0	2	0	0	1	0	0	0	0
A2	17	3	3	0	2	0	0	0	0	0
A3	12	3	3	0	1	0	0	0	0	0
A4	19	3	5	0	1	0	0	0	0	0
A5	18	5	1	0	0	1	0	1	0	0
C1	25	8	0	2	1	3	2	1	1	0
C2	16	1	1	6	0	2	0	0	0	1
C3	21	5	0	0	0	0	0	0	2	1
C4	11	5	0	0	0	0	0	0	1	1
C5	24	6	1	2	1	1	0	1	0	0
s_m^A		0.916	0.855	0.593	1.18	0.527	0.315	0.708	0.955	0.874
s_m^C		1.17	0.689	1.05	1.3	0.492	0.575	0.744	1.34	1.24
$\hat{u}_m$		-5.32	-4.77	-5.51	-6.49	-4.14	-4.58	-5.11	-6.65	-6.59

Table S2. Numbers of isolates carrying the most common adaptive mutations and the estimated mutation rates. Only clones with unique barcodes are shown. Column "Seq. isolates" shows the total number of sequenced isolates with unique barcodes sampled from each culture. Last row shows the estimated $u_m = \log_{10} U_m$.

References

[1] Hom EF, Murray AW. Niche engineering demonstrates a latent capacity for fungal-algal mutualism. Science. 2014;345(6192):94–98.

[2] Boyer S, Hérissant L, Sherlock G. Adaptation is influenced by the complexity of environmental change during evolution in a dynamic environment. PLoS Genet. 2021;17(1):e1009314.

[3] Levy SF, Blundell JR, Venkataram S, Petrov DA, Fisher DS, Sherlock G. Quantitative evolutionary dynamics using high-resolution lineage tracking. Nature. 2015;519(7542):181–186.

[4] Kryazhimskiy S, Rice DP, Jerison ER, Desai MM. Global epistasis makes adaptation predictable despite sequence-level stochasticity. Science. 2014;344(6191):1519–1522.

[5] Lang GI, Rice DP, Hickman MJ, Sodergren E, Weinstock GM, Botstein D, et al. Pervasive genetic hitchhiking and clonal interference in forty evolving yeast populations. Nature. 2013;500(7464):571–574.

[6] Kao KC, Sherlock G. Molecular characterization of clonal interference during adaptive evolution in asexual populations of *Saccharomyces cerevisiae*. Nature genetics. 2008;40(12):1499–1504.

[7] Blundell JR, Schwartz K, Francois D, Fisher DS, Sherlock G, Levy SF. The dynamics of adaptive genetic diversity during the early stages of clonal evolution. Nature ecology & evolution. 2019;3(2):293–301.

[8] Collins S, Bell G. Phenotypic consequences of 1,000 generations of selection at elevated CO_2 in a green alga. Nature. 2004;431(7008):566–569.

[9] Perrineau MM, Gross J, Zelzion E, Price DC, Levitan O, Boyd J, et al. Using natural selection to explore the adaptive potential of Chlamydomonas reinhardtii. PLoS One. 2014;9(3):e92533.

[10] Lee CK, Araki N, Sowersby DS, Lewis LK. Factors affecting chemical-based purification of DNA from *Saccharomyces cerevisiae*. Yeast. 2012;29(2):73–80.

[11] Newman SM, Boynton JE, Gillham NW, Randolph-Anderson BL, Johnson AM, Harris EH. Transformation of chloroplast ribosomal RNA genes in Chlamydomonas: molecular and genetic characterization of integration events. Genetics. 1990;126(4):875–888.

[12] Venkataram S, Dunn B, Li Y, Agarwala A, Chang J, Ebel ER, et al. Development of a comprehensive genotype-to-fitness map of adaptation-driving mutations in yeast. Cell. 2016;166(6):1585–1596.

[13] Baym M, Kryazhimskiy S, Lieberman TD, Chung H, Desai MM, Kishony R. Inexpensive multiplexed library preparation for megabase-sized genomes. PLoS One. 2015;10(5):e0128036.

[14] Ghodsi M, Liu B, Pop M. DNACLUST: accurate and efficient clustering of phylogenetic marker genes. BMC Bioinformatics. 2011;12:271.

[15] Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics. 2009;25(14):1754–1760.

[16] Nguyen Ba AN, Cvijović I, Echenique JIR, Lawrence KR, Rego-Costa A, Liu X, et al. High-resolution lineage tracking reveals travelling wave of adaptation in laboratory yeast. Nature. 2019;575(7783):494–499.

[17] Li Y, Venkataram S, Agarwala A, Dunn B, Petrov DA, Sherlock G, et al. Hidden Complexity of Yeast Adaptation under Simple Evolutionary Conditions. Curr Biol. 2018;28(4):515–525.e6.

[18] Barillot E, Lacroix B, Cohen D. Theoretical analysis of library screening using a N-dimensional pooling strategy. Nucleic Acids Res. 1991;19(22):6241–6247.

[19] Baym M, Shaket L, Anzai IA, Adesina O, Barstow B. Rapid construction of a whole-genome transposon insertion collection for Shewanella oneidensis by Knockout Sudoku. Nature communications. 2016;7(1):1–13.

[20] Manhart M, Adkar BV, Shakhnovich EI. Trade-offs between microbial growth phases lead to frequency-dependent and non-transitive selection. Proc Biol Sci. 2018;285(1872).

[21] Tenaillon O, Rodríguez-Verdugo A, Gaut RL, McDonald P, Bennett AF, Long AD, et al. The molecular diversity of adaptive convergence. Science. 2012;335(6067):457–461.

[22] Zhu YO, Siegal ML, Hall DW, Petrov DA. Precise estimates of mutation rate and spectrum in yeast. Proc Natl Acad Sci U S A. 2014;111(22):E2310–8.

[23] Desai MM, Fisher DS. Beneficial mutation-selection balance and the effect of linkage on positive selection. Genetics. 2007;176(3):1759–1798.

[24] Pawitan Y. In All Likelihood: Statistical Modelling and Inference Using Likelihood. Oxford science publications. OUP Oxford; 2001.

[25] Chen H, Zhou W, Chen W, Xie W, Jiang L, Liang Q, et al. Simplified, rapid, and inexpensive estimation of water primary productivity based on chlorophyll fluorescence parameter Fo. Journal of plant physiology. 2017;211:128–135.

[26] Hunter GA, Ferreira GC. Molecular enzymology of 5-aminolevulinate synthase, the gatekeeper of heme biosynthesis. Biochim Biophys Acta. 2011;1814(11):1467–1473.

[27] Brown BL, Kardon JR, Sauer RT, Baker TA. Structure of the Mitochondrial Aminolevulinic Acid Synthase, a Key Heme Biosynthetic Enzyme. Structure. 2018;26(4):580–589.e4.

[28] Betts MJ, Russell RB. Amino-acid properties and consequences of substitutions. In: Bioinformatics for Geneticists. Chichester, UK: John Wiley & Sons, Ltd; 2007. p. 311–342.