

Title page

The characteristics of the HIV-1 Env glycoprotein contribute to viral pathogenesis

Silvia Pérez-Yanes¹, Maria Pernas^{2,#}, Silvia Marfil^{3,#}, Romina Cabrera-Rodríguez¹, Raquel Ortiz³, Carla Roviroso³, Judith Estévez-Herrera¹, Isabel Olivares², Concepción Casado^{2,*}, Cecilio Lopez-Galindez^{2,*}, Julià Blanco^{3,4,*} and Agustin Valenzuela-Fernández^{1,*}

¹Laboratorio de Inmunología Celular y Viral, Unidad de Farmacología, Sección de Medicina, Facultad de Ciencias de la Salud de la Universidad de La Laguna (ULL), Campus de Ofra s/n, 38071 Tenerife, Spain.
(sperezya@ull.edu.es), (rcabrerr@ull.edu.es), (jesteveh@ull.edu.es), (avalenzu@ull.edu.es)

²Unidad de Virología Molecular, Laboratorio de Referencia e Investigación en Retrovirus. Centro Nacional de Microbiología. Instituto de Salud Carlos III. Majadahonda 28220 Madrid. Spain.
(ccasado@isciii.es), (mpernas@isciii.es), (iolivar@isciii.es), (ceciliolopezgalindez1462@gmail.com)

³Institut de Recerca de la Sida IrsiCaixa, Institut d'Investigació en Ciències de la Salut Germans Trias i Pujol (IGTP), 08916 Badalona, Spain.

⁴Universitat de Vic, Universitat Central de Catalunya, UVIC-UCC, 08500 Vic, Spain.
(smarfil@irsicaixa.es), (rortiz@irsicaixa.es), (croviroso@irsicaixa.es), (jblanco@irsicaixa.es)

#These authors contributed equally.

*Corresponding authors.

E-mail address for correspondece: avalenzu@ull.edu.es

1 **Abstract**

2 The understanding of HIV-1 pathogenesis and clinical progression is incomplete
3 because of the variable contribution of host, immune and viral factors. The
4 involvement of viral factors has been investigated in extreme clinical phenotypes
5 from rapid progressors to long-term non-progressors (LTNPs). Among HIV-1
6 proteins, the envelope glycoprotein complex (Env) has concentrated many
7 studies for its important role in the immune response and in the first steps of viral
8 replication. In this study, we analyzed the contribution of 41 Envs from 24 patients
9 with different clinical progression rates and viral loads (VLs), LTNP-Elite
10 Controllers (LTNP-ECs); Viremic LTNPs (vLTNPs), and non-controller's
11 individuals contemporary to LTNPs or recent, named Old and Modern
12 progressors. We analyzed the Env expression, the fusion and cell-to-cell transfer
13 capacities as well as viral infectivity. The sequence and phylogenetic analysis of
14 Envs were also performed. In every functional characteristic, the Envs from
15 subjects with viral control (LTNP-ECs and vLTNPs) showed significant lower
16 performance compared to those from the progressor individuals (Old and
17 Modern). Regarding sequence analysis, the variable loops of the gp120 subunit
18 of the Env (i.e., V2, V4 and mainly V5) of the progressor individuals showed
19 longer and more glycosylated sequences than controller subjects. Therefore,
20 HIV-1 Envs presenting poor viral functions and shorter sequences were
21 associated with viremic control and the non-progressor clinical phenotype,
22 whereas functional Envs were associated with the lack of virological control and
23 progressor clinical phenotypes. These correlations support the central role of Env
24 genotypic and phenotypic characteristics in the *in vivo* HIV-1 infection and
25 pathogenesis.

26 **Words: 250**

27 **IMPORTANCE**

28 The role of the virus in the pathogenesis of HIV-1 infection has not been
29 investigated in isolates from individuals with different progression rates. In this
30 work, we studied the properties of the envelope glycoprotein complex (Env) in
31 individuals with different progression rates to elucidate its role in pathogenesis.
32 We estimated the Env expression, the CD4 binding, the fusion and cell-to-cell
33 viral transfer capacities that affect the infectivity of the viral Envs in recombinant
34 viruses. The Envs from individuals which control viral replication and lack
35 clinical progression (LTNP-ECs and vLTNPs) showed lower functional capacities
36 than from subjects with clinical progression (Old and Modern). The functional
37 increase of the Envs characteristics was associated with an increase in viral
38 infectivity and in increased length of variable loops and the number of
39 glycosylation sites of the Env (gp120/SU). These results support the concept
40 that viral characteristics contribute to viral infection and pathogenesis.

41 **Words: 148**

42

43

44

45

46

47

48

49

50

51 **Introduction**

52 Pathogenesis of viral infections is the result of complex interactions between
53 host genetics, immune responses and viral factors. In human immunodeficiency
54 virus type 1 (HIV-1) infection and pathogenesis, the role of host (1-6), immune (6-
55 15) and viral factors (16-20) has been widely investigated. The interactions of
56 these factors have been primarily studied in extreme clinical phenotypes like
57 rapid progressors (RPs) (21, 22) or long-term non-progressors (LTNPs), LTNP-
58 Elite Controllers (LTNP-ECs), HIV controllers or Elite suppressors (ES) (17-19,
59 23, 24).

60 Due to these entangled interactions, the investigation of the role of viral proteins
61 and their specific properties in HIV-1 pathogenesis is challenging. Among the
62 viral proteins, the envelope glycoprotein complex (Env) has attracted numerous
63 studies because its essential role in the immune response and in the initial
64 events of the HIV-1 biological cycle (25-29), i.e the binding to the cellular
65 receptors (29-42). The binding efficiency of the viral Env to the CD4 receptor
66 determines further steps of the viral cycle: virus-cell signaling, fusion and cell-to-
67 cell virus transfer capabilities (18, 19, 43). HIV-1 Envs unable to stabilize
68 microtubules (i.e., increasing post-transductional acetylation of Lys⁴⁰ residue in
69 α -tubulin), to reorganize F-actin for the delineation of pseudopod-entry virus hot
70 zones present low CD4 binding, restricted fusion and low early infection (18, 19,
71 43-45).

72 There are few reports investigating the characteristics of viral Envs from HIV
73 individuals with different clinical characteristics. Lassen *et al.* studied the entry
74 efficiency of viral Envs from ES individuals relative to chronically infected
75 viremic and chronic progressors. Envs from ES showed decreased entry
76 efficacy and slower entry kinetics than those of chronic progressors (20). Our

77 group studied the CD4 binding, signaling capacity, fusogenicity of viral Envs
78 from viremic non-progressors (VNPs) that were similar to those of progressors
79 individuals (19). In previous reports, deficient viral Env glycoproteins, because
80 of poor CD4 binding, low transfer and signaling capacity (18) were identified in a
81 cluster of poor replicating viruses from a group of LTNP-ECs without clinical
82 progression for more than 20 years (17, 18). Thus, these works have established
83 that viral Env play an important role in the pathogenesis control in LTNPs (17-
84 20, 46, 47).

85 To further investigate the role of viral Env in HIV-1 infection and pathogenesis,
86 in this work, we expanded our previous studies to viral Envs from other sets of
87 LTNP-ECs and Viremic LTNPs (vLTNPs) in comparison with groups of chronic
88 progressors. Clonal full-length *env* genes derived from viruses of individuals in
89 these distinct clinical groups were analyzed for expression, CD4 dependent-
90 Env-mediated fusion, cell-to-cell viral transfer and infection efficiency. This
91 analysis permitted the establishment of a relationship between the initial events
92 of the viral replication cycle, mediated by the viral Env characteristics, with the
93 VL control and the clinical outcome and pathogenesis of the HIV-1 infection.

94

95

96 **Results**

97 **Analysis of the characteristics of viral envelopes of viruses from different** 98 **risk groups.**

99 For the investigation of the potential role of the HIV-1 Env in virological control
100 and pathogenesis, we studied the phenotypic characteristics of 41 Envs from 24
101 individuals without antiviral therapy and different VLs (**Table 1**). We analyzed 10
102 Envs from 6 LTNP-EC individuals with undetectable VL and infected in the late
103 80's and 90's; 10 viral clones from 6 Viremic LTNPs (vLTNPs) with VL <10,000
104 viral copies/mL and infected in the 90's. To ascertain that the characteristics of
105 the Envs from these LTNPs were not due to the sampling time, we compared
106 them with 10 Envs obtained from 6 HIV-1 individuals also infected in the same
107 period (90's), but with high VL >10⁵ viral copies/mL and chronic infection; these
108 Env were designated Old. Finally, we studied 11 viral clones from 6 chronic
109 individuals infected between 2013-2014 with VL >10⁴ viral copies/mL and named
110 Modern. The main characteristics of the participants are summarized in **Table**
111 **1**.

112 We first analyzed the potential differences in the expression between the Env
113 clones from the clinical groups, by measuring their cell-surface expression
114 levels in HEK-293T cells (**Figure 1A**, *shows study scheme*, and **Figure 2**).

115 Although we observed a progressive augmentation of Env expression in viral
116 clones derived from patients that do not control viremia (i.e., Old and Modern
117 patients) compared to LTNPs (EC and Viremic), this increase did not reach
118 statistical significance (**Figure 2**). Thus, the expression capability of the viral
119 Envs appears to not contribute to the differences in VL and pathogenesis
120 between groups.

121

122 **Analysis of cell-to-cell membrane fusion and viral transfer capacity of viral**
123 **envelopes.**

124 A key process for HIV Env-mediated infection is the interaction of the Env
125 complex with the CD4 receptor. When this interaction is functionally efficient,
126 viral transfer through synaptic contacts or fusion pore formation are triggered
127 during cell-to-cell or virus-to-cell contacts, respectively (18, 19, 43, 45, 48). We
128 examined the viral Env/CD4 interaction and the efficiency of subsequent
129 functions, measuring the membrane fusion capacity of the Envs (Figure 1B,
130 *shows study scheme*) in co-cultures between Env-expressing HEK-293T and
131 HIV-permissive target TZM-bl cells (Figure 3). To fully characterize our
132 experimental models, we used the Envs from reference HIV-1_{BaL} (CCR5-tropic)
133 and HIV-1_{NL4.3} (CXCR4-tropic) viruses (Figure 3 and 4). This fusion assay
134 yielded lower fusion values for Envs of viruses from LTNP-ECs and from
135 vLTNPs than for Old and Modern progressors, and attaining statistical
136 significance between LTNPs (EC and Viremic) and Modern Envs glycoproteins
137 (Figure 3B).

138 Next, we assayed the CD4-dependent cell-to-cell virus transfer capacity of the
139 viral envelopes. This experiment was performed co-culturing Env-expressing
140 HEK-293T cells with unstimulated primary CD4⁺ T lymphocytes as target cells
141 (Figure 1C, *shows study scheme, and Materials and methods*). In this assay,
142 we forced the formation of virological synapses between virus-effector HEK-
143 293T cells expressing the different Envs together with the structural HIV Gag
144 polyprotein, and fresh primary CD4⁺ T cells from healthy donors (Figure 1C,
145 *shows study scheme*). The Envs from the LTNPs (EC and Viremic) individuals
146 displayed a lower ability to transfer viral particles to primary CD4⁺ T
147 lymphocytes than Envs from Old individuals and significantly lower than from

148 Modern participants ($p < 0.0022$ between all groups) (**Figure 4**). These data
149 suggest that the Envs from LTNP-EC viruses had an impaired binding to the
150 cell-surface CD4 receptor and that this impairment was progressively overcome
151 in the Envs from individuals from the other groups with less control of viral
152 replication, and higher VL.
153 Thus, the phenotypic characterization of the Envs of viruses from subjects with
154 distinct progression rates confirmed that LTNP-ECs and vLTNPs presented
155 viruses with an impaired Env CD4-associated functions and a significant lower
156 fusogenic and transfer capacity, in comparison with viruses from the viremic
157 groups: These lower characteristics were also linked with the low VL detected in
158 these subjects (**Figures 3 and 4**). We also observed a functional improvement
159 in the viral Envs from the LTNP-EC and vLTNP individuals to those of chronic
160 Modern glycoproteins: These data support that the deficient Env fusion and
161 transfer capacities observed in the Envs of viruses from LTNP-EC and vLTNP
162 phenotypes have been enhanced in the viruses from individuals with
163 progressive infection, particularly in those of the Modern group.

164

165 **Infectivity of recombinant viruses with the analyzed envelopes.**

166 For the exploration of the potential consequences of these Env properties in
167 virus biology, we estimated the infectivity of recombinant viruses bearing the
168 Env from the different HIV+ phenotypic groups in TZM-bl cells (**Figure 5 and**
169 **Figure 1D**, *shows study scheme*). Viral Envs from the LTNP-EC group showed
170 the lowest infectivity values, whereas the Modern Envs produced the higher
171 titers. The viruses from vLTNPs displayed higher titers than LTNP-ECs but
172 lower than those from Old individuals. Recombinant viruses from individuals
173 with high VL and progressive infection (Old and Modern) have higher infectivity

174 rates than those with viral control (EC and Viremic). These results explain why
175 the viral properties analyzed (binding, fusion and transfer) have a significant
176 impact in viral infectivity with an important effect in the biology of HIV-1 and viral
177 pathogenesis.

178

179 **Correlation between viral characteristics of the envelopes.**

180 A significant correlation was observed between the HIV-1 Env-triggered cell-to-
181 cell transfer data, which is directly mediated by Env/CD4 binding, with Env-
182 mediated infectivity and fusogenicity (**Figure 6**). In all viral characteristics, the
183 Envs from subjects with virological control (EC and Viremic) showed the lower
184 values, whereas those from the non-controlling individuals (Old and Modern)
185 had the higher values. Therefore, HIV-1 Envs displaying poor viral functions,
186 because of the poor binding of the viral Env to the CD4, correlated with viremic
187 control and non-progressor clinical phenotypes. In contrast, functional Envs are
188 associated with the lack of viremic control and the progressor clinical
189 phenotypes. These statistical correlations support the role of viral properties in
190 the viral phenotype that contributes to HIV-1 infection, disease progression and
191 pathogenesis.

192

193 **Analysis of the viral envelope sequences.**

194 For the search of potential mechanisms involved in the changes of the
195 characteristics among the different Envs sets, we analyzed the Env amino-acid
196 (aa) sequences that could be associated with the distinct clinical phenotypes.
197 Initially, we performed a phylogenetic reconstruction from *env* aa sequences
198 together with other aa sequences obtained from HIV-1 Spanish individuals. All
199 aa sequences analyzed correspond to HIV-1 subtype B. This analysis did not

200 reveal phylogenetic relationships between the different groups analysed and no
201 clustering except for those aa sequences obtained from the same individual
202 (**Figure 7**). Envs from LTNP-ECs and one vLTNPs grouped in short branches,
203 as a consequence of the viral and evolutionary control, whereas long branch
204 length was observed in the sequences obtained from non-controller patients
205 (Old and Modern), because of the higher replication and viral evolution in these
206 individuals.

207 We then carried out a comprehensive study of the protein sequences focusing
208 in the variable loops and their associated potential N-linked glycosylation sites
209 (PNGs) in the gp120 subunit of the Env. In general, as previously reported,
210 there is a trend in the HIV-1 viral Env to gain length and glycosylation sites
211 along the epidemic ([49-51](#)). This increasing trend is also found in our work
212 where viruses from the LTNPs (EC, Viremic) and Old Envs isolated in the 90's
213 showed shorter lengths than those of the Modern group obtained in 2013-2014
214 (**Table 2**). The V3 loop was the most conserved and constant region in length
215 and glycosylation sites (**Table 2 and Figure 8**), while the other loops showed
216 length increases predominantly in the V2 and V5 loops that were reproduced in
217 the total length (**Table 2 and Figure 8**). The only statistical differences were
218 noticed between the total length in the LTNPs (EC and Viremic) versus Old and
219 Modern Envs in the V2 and V5 regions (**Figure 8**).

220 Regarding the PNGS in the sequences, many of the 24 relevant sites previously
221 described ([52-55](#)) were present in these set of viral glycoproteins. However,
222 major differences were observed in the aa extension of the loops with a
223 progressive acquisition of more PNGS in the Modern Envs (**Table 2**). Glycan at
224 N289 site was more present in LTNP-ECs, vLTNPs and Old viruses but is not
225 present in Modern ones. Position N362 which is N proximal to the CD4 binding

226 “DPE” motif (positions 368-370HXB2 sequence) was conserved in LTNP-EC,
227 Viremic and Old but was only present in two of the Modern Envs. It is interesting
228 to highlight that changes also occurred in the viral transmembrane gp41 protein
229 in glycan N816 that was dominant in LTNPs but not in chronic individuals (Old
230 and Modern).

231 It is interesting to mention that the trend in Env length increase follows the same
232 pattern that the functional growth of the Env shown in the distinct viral
233 characteristics (see **Figures 3 to 6**). We observed a good correlation between
234 the genetic distance to the subtype B ancestor sequence obtained from Los
235 Alamos National Laboratory HIV Database (LANL database,
236 <http://www.hiv.lanl.gov>) and the functionality of viral Env proteins analysed
237 (**Figure 9**). In general, the lower evolutionary sequences (less genetic distance
238 to subtype B ancestor) are those with lower functionality (LTNP-ECs) and the
239 higher evolutionary sequences are those with higher functionality (Moderns). In
240 summary, the viral Envs with the most efficient characteristics are found within
241 the Envs of the Modern group that also show the longer gp160 proteins, with
242 more glycosylated sites and higher distance to the subtype B ancestor.

243

244

245 **Discussion**

246 HIV-1 infected individuals display a wide spectrum of clinical progression rates.
247 The causes of this dispersion are multiple and associated with the operation of
248 numerous combinations of host genetic, immunological and viral factors. In this
249 work, we studied the potential contribution of viral Env glycoprotein
250 characteristics to the clinical outcome of HIV-1 infection in HIV+ individuals with
251 different clinical status.

252 The different groups of patients were defined by their clinical characteristics,
253 distinct VLs and isolation dates because several studies have described a clear
254 correlation between patients' VL and the likelihood of virus transmission,
255 disease progression and pathogenesis (56-63).

256 Although viral control in HIV-1 individuals has been linked to the host-immune
257 responses (10, 64), other researchers and our group, however, established, in
258 previous works, a direct connection between deficiencies in HIV-1 Env-
259 associated functions and long-term viremia control in LTNP-ECs (17, 18, 20).

260 The Envs from these LTNP-EC individuals were ineffective in the CD4 binding
261 and in the subsequent functions: viral signaling, fusion and cell entry. These
262 Env characteristics ensued in low replication and transmissibility of the virus
263 (18, 19, 43, 45). All these data strongly support the role of the viral Env in the
264 LTNP-EC phenotype and viral pathogenesis.

265 In the present work, we extended these observations to more Env from non-
266 progressor subjects, which are not associated with a cluster of infection, in
267 comparison to different sets of progressor chronic individuals. The Envs
268 characteristics from LTNP individuals (EC and Viremic) were compared with
269 those of individuals with progressive infection (Old and Modern). We

270 investigated the defects in the association of Envs with the CD4, membrane
271 fusion impairment and the cell-to-cell virus transfer and viral infection capacities.
272 Viral Envs from LTNPs showed the lower binding capacity to the CD4 receptor
273 and this initial inefficient Env/CD4 interaction led to a deficiency in membrane
274 fusion and virus cell-to-cell transfer capabilities. The properties of the Env from
275 LTNPs were not due to the ancestral origin of the LTNPs viruses isolated in the
276 late 80's and 90's, because the characteristics of the Old viruses which were
277 contemporary to the LTNPs did not showed these limited functional
278 characteristics. On the contrary, Envs from progressors (Old and Modern)
279 presented efficient CD4-mediated viral functionality that triggered an effective
280 membrane fusion and viral transfer. Thus, we disclosed that there is a clear
281 correlation between the level of viral fusion, the transfer capacity of the viral Env
282 and viral infectivity. The observed differences between the characteristics of the
283 Envs from these groups could not be associated with viral tropism, because all
284 the *env* nucleotide sequences from the studied viruses, showed an R5 tropism
285 (Web PSSM, <https://indra.mullins.microbiol.washington.edu/webpssm/>).
286 In summary, viral Envs from LTNPs exhibited non-functional characteristics
287 (**Figures 3-6**) in comparison with those from viruses of the progressive infection
288 groups, supporting the concept that the properties of the Envs were associated
289 with viral control and the clinical progression rate of the HIV-1 individuals.
290 In spite of the limited sampling, because of the difficult and laborious viral
291 characterization of the viral phenotypes, we observed statistically significant
292 differences between the characteristics of the Envs of viruses from LTNP-ECs
293 and the Moderns. Also, if we consider the Env characteristics from all clinical
294 groups, there is a consistent and recurrent tendency, although with no statistical
295 power in some cases, to gain functionality in the viral Envs from the LTNP

296 individuals (LTNP-ECs and vLTNPs), to those of the progressive groups (Old
297 and Modern).

298 Remarkably, the increase in Env functionality also correlated with longer and
299 more glycosylated proteins. The aa length and PNGs' profile of the Envs from
300 the individuals of the distinct clinical groups showed that the studied Envs tend
301 to increase length and glycosylation over the course of the epidemic as
302 previously described (see [\(49, 51\)](#)). We observed that Env changes
303 accumulated essentially in the V1, V2, V4 and V5 loops, as previously shown in
304 works relating the role of V1 and V4 loops in the CD4 binding and neutralization
305 [\(65-68\)](#) and viral cell-to-cell transfer capacity [\(50, 69, 70\)](#). Regarding specific
306 changes detected in our study, the loss of the N362 PNGs (position in the HXB2
307 isolate; group M, subtype B (HIV-1 M:B_HXB2R: NCBI:txid11706)) which was
308 prevalent in the EC, Viremic and Old but not in the Modern Envs groups could
309 be associated with the gain of functionality in the Envs. However, the opposite
310 effect with more efficient fusion and transfer capacity was found in Australian
311 viruses with the N362 glycosylation site [\(55\)](#). The potential role of the other
312 changes in PNGs detected in our study need to be further investigated. Besides
313 these important changes, it is clear that point mutations could have a significant
314 impact in the viral characteristics and HIV pathogenesis [\(71, 72\)](#). The variants
315 of concern (VOCs) of the pandemic severe acute respiratory syndrome
316 coronavirus (SARS-CoV-2) unfortunately are reminding us [\(73, 74\)](#). Thus, the
317 contribution of the individual mutations deserves further studies but it is now out
318 of the scope of the present work.

319 In contrast with the more significative changes detected in the V2 and V5 loops,
320 it is important to point the stability in length and glycosylation of the V3 loop.
321 This structure is key for viral tropism [\(75-79\)](#) and for the correct CD4 Env

322 binding as revealed with anti-V3 neutralizing antibodies that abrogate Env-CD4
323 interaction (80, 81).

324 In this study, we confirmed the inefficient functionality of the Envs from LTNP-
325 EC individuals previously described for a cluster of viruses (18, 20), but
326 extended to HIV+ individuals controlling viremia which are not clustered by the
327 same transmitted/founder (T/F) virus. Also, a gain of Envs functionality from
328 those of the LTNP individuals to the chronic not controlling individuals was
329 identified. This improvement was detected in every Env characteristic analyzed;
330 expression, fusion, virus transfer and infectivity. Interestingly, this functional
331 growth of viral Env was associated in this study with length and PNGs increases
332 in the variable loops. This increase was also reported in studies analyzing the
333 susceptibility, neutralization sensitivity, co-receptor binding, host range and viral
334 phenotype (49). This increase in the V1-V2 length and PNGs has also been
335 detected thorough chronic infections from early to late viral Env sampling like in
336 our work (49). Likewise in a group of individuals infected with closely related
337 viruses higher PNGs density has been observed in the V1-V5 region of the
338 gp120 during chronic infection compared to those observed during the early
339 acute infection phase (82). In viruses from the HIV-1 subtype B, it seems that
340 early after viral transmission to a new host a selection for viral variants with
341 shorter variable regions and a reduced degree of PNGs occurs (83). The growth
342 in functionality of the viral characteristics was also correlated with the genetic
343 distance of the sequences to the subtype B ancestor. Genetic variability in *env*
344 gene has been is associated with an increase in viral infectivity and replication
345 capacity (84-89). These changes could facilitate viral replication by increasing
346 viral fitness that favors the escape from the immune response and anti-retroviral
347 therapy (ART) failure (90-99).

348 The non-functional characteristics of the primary Envs of LTNP individuals (ECs
349 and Viremics) resulted in poor viral replication and very limited evolution that
350 could allow the efficient immune control of HIV-1 infection and pathogenesis. It
351 has been reported that in a LTNP-EC patient that followed discontinued ART,
352 the V1 domain of his HIV-1 strain that retained good infectivity and replicative
353 capacity included two additional N-glycosylation sites and was placed in the top
354 1% of lengths among the 6,112 Env sequences analyzed in the Los Alamos
355 National Laboratory online database (100).

356 Therefore, it is conceivable that the functional characterization of the inefficient
357 HIV-1 Envs could be significant in the development of a new generation of
358 immunogens. Indeed, attenuated HIV or simian immunodeficiency virus (SIV)
359 vaccines (LAHVs or LASVs) have been postulated as therapeutic vaccine
360 strategies (101-107). However, further antigenic and immunogenicity work is
361 needed to disclose the potential implications of these non-functional HIV Envs
362 in the vaccine/cure field.

363

364 In summary, in this work, we exposed that the characteristics of the viral Envs
365 from different groups of HIV-1 infected individuals could be associated with the
366 short or long-term VL control and the clinical progression rate of the infection.
367 The non-functional HIV-1 Envs could help in the development of new strategies
368 for functional cure and virus eradication. Our data support the hypothesis that
369 the functionality of viral Envs is a crucial characteristic for the control of viral
370 infection, replication and pathogenesis.

371

372

373 **Material and methods**

374 **Viral envelopes.**

375 Forty-one viral envelopes (Envs) were obtained from samples of different
376 origins: the HIV HGM BioBank integrated in the Spanish AIDS Research
377 Network (RIS-RETIC, ISCIII) (samples 1,2,3,6,7,8,13,14,15,16,17,18,19), the
378 Centro Sanitario Sandoval, Hospital Clínico San Carlos (samples
379 21,22,24,28,30,31,32,33,36,37,38,39,40,42,43,44,45,46,49,50,51,52), the
380 irsiCaixa Research Foundation (samples 9,10,11,12) and from Hospital Xeral
381 de Vigo (samples 26,27). Samples were obtained in three different phases of
382 the Spanish epidemic from 1993-94, 2004-2005 and 2013-2014. Samples were
383 processed following current procedures and frozen immediately after their
384 reception. All patients participating in the study gave their informed consent and
385 protocols were approved by institutional ethical committees. Identification
386 numbers and characteristics are found in [Table 1](#).

387

388 **Ethics Statement.**

389 Samples were obtained from participants who gave informed consent for
390 genetic analysis studies and they were registered as sample collection in the
391 Spanish National Registry of Biobanks for Biomedical Research with number
392 C.0004030. The consents were approved by the Ethical and Investigation
393 Committees of the “Centro Sanitario Sandoval” (Madrid) and the samples were
394 encoded and de-identified in these Centers. All clinical investigations were
395 conducted according to the principles expressed in the Declaration of Helsinki.
396 The studies were approved by the Comité de Ética de la Investigación y de
397 Bienestar Animal of the Instituto de Salud Carlos III with CEI PI 05_2010-v3 and
398 CEI PI 09-2013 numbers.

399 **Generation of *env* gene expression plasmids.**

400 The *env* genes were amplified at limiting dilution by nested PCR from proviral
401 DNA. The products were cloned into the pcDNA3.1D/V5-His's Topo expression
402 vector (Invitrogen) and NL4.3. The R5-tropic BaL.01-*env* (catalog number
403 11445) glycoprotein plasmid was from the NIH AIDS Research and Reference
404 Reagent Program. Ten viral Envs were derived from 6 LTNP-EC patients, 10
405 clones from 6 Viremic LTNPs, 10 clones from 6 "Old" individuals (contemporary
406 to LTNPs) and 11 clones from 10 recent "Modern" patients and NL4.3 and
407 BaL.01 reference clones expression plasmids were transformed in DH5 α cells,
408 and clones sequenced to check the correct insertion of the *env* gene.

409

410 **Env expression and fusion assays.**

411 The Env expression plasmids were used to transfect HEK-293T cells with X-
412 tremeGENE HP DNA Transfection Reagent (Sigma) in combination with either
413 a Tat expression plasmid pTat for Env expression and fusion assays, or with the
414 *env* defective HIV-1 backbone pSG3 plasmid for viral transfer assays ([18](#), [19](#),
415 [108](#)). As a negative control, HEK-293T cells were transfected only with pTat
416 and as a positive control we use the BaL and NL4.3 Envs. HEK-293T cells were
417 chosen as effector cells since they provide sensitive measures of fusion even
418 when using low fusogenic Env. 24 hours post-transfection, cells were collected,
419 and tested for Env surface expression and also fusion activity.

420 To test Env expression, 1×10^5 Env/Tat co-transfected HEK-293T cells were
421 incubated with 2G12 and IgGb12 monoclonal antibodies (mAbs; Polymun,
422 Viena, Austria) at 6 $\mu\text{g}/\text{mL}$ each for 45 minutes at RT. After washing the cells,
423 the PE-labeled goat anti-human IgG (Jackson ImmunoResearch Laboratories)
424 was added and incubated in the dark at room temperature for 15 minutes, as

425 similarly reported (18, 19). Cells were washed, fixed in formaldehyde 1%,
426 acquired in a Celesta flow cytometer (BD FACS Celesta) and analyzed using
427 the Flow-Jo software (Tree Star Inc.) The percentage of Env-positive cells and
428 the Mean Fluorescence Intensity (MFI) of these cells were used to evaluate Env
429 expression.

430 To test fusion activity, 1×10^4 Env/Tat-transfected or control Tat-transfected
431 HEK-293T cells were mixed (ratio 1:1) in 96-well plates with
432 $CD4^+CXCR4^+CCR5^+$ TZM-bl reporter cells for 6 hours at 37°C. Luciferase
433 activity was measured (Fluoroskan Accent, Labsystems) using Brite-Lite
434 (PerkinElmer) and normalized to BaL-Env-mediated fusion. NL4.3 and BaL-Env
435 expression plasmids were used as positive controls for Env staining and as
436 reference value for fusion activity (BaL = 100%), as similarly reported (19, 108)
437 (summarized in the scheme of Figure 1B).

438

439 **HIV-1 transfer/CD4 binding**

440 To test viral transfer activity, which exclusively depends on the binding of gp120
441 to the CD4 molecule, Env expression plasmids were co-transfected with the
442 Env-defective pSG3 plasmid in HEK-293T cells, as similarly reported (18, 19,
443 108). One day after transfection, 1×10^5 HEK-293T cells were mixed at a 1:1
444 ratio in 96-well plates with primary $CD4^+$ T lymphocytes freshly isolated from
445 healthy donors by negative selection ($CD4^+$ T-Cell Isolation Kit II, human,
446 Miltenyi Biotec). Viral transfer was assessed after 24 hours of incubation at
447 37°C in permeabilized (FIX & PERM Cell Permeabilization kit, Invitrogen Life
448 Technologies) and stained cells with the anti-HIV-1 p24 KC57 mAb (anti HIV
449 core antigen RD1 labelled, IZASA) for 20 minutes in the dark at RT. Then, the
450 cells were washed and fixed in formaldehyde 1%, and acquired in a Celesta

451 flow cytometer (BD FACS Celesta) and the content of p24 in gated CD4+ T
452 cells and gated HEK-293T cells was analyzed using the Flow-Jo software (Tree
453 Star Inc.). The percentage of p24+ HEK-293T cells was used as a control for
454 transfection efficiency and was similar among all experiments. Since co-
455 receptor binding or fusion activity are not required for viral transfer, the
456 frequency of p24+/CD4+ T cells was a direct measure of the amounts of HIV-1
457 virions bound to or taken up by target cells (summarized in the scheme of
458 **Figure 1C**).

459

460 **Infectivity assay**

461 Cloned viral Envs were used to generate pseudoviruses by co-transfection with
462 pSG3 plasmid of HEK-293T cells as indicated above and tested in TZM-bl cells
463 to determine the infectivity capacity. Serial Dilutions of the pseudoviruses
464 generated with the different Envs of the different groups of patients were made
465 in a 96-well plate. Then, 1×10^5 TZM-bl cells were added to the pseudoviruses
466 with DEAE dextran hydrochloride (Sigma) at 18 $\mu\text{g}/\text{mL}$. After 48 hours of
467 incubation at 37°C, luciferase activity was measured (Fluoroskan Accent,
468 Labsystems) using Brite-Lite (PerkinElmer). Uninfected TZM-bl cells were used
469 as a negative control. The TCID₅₀ (Median Tissue Culture Infectious Dose)
470 value was calculated with Montefiori template and normalized with the viral
471 concentrations (summarized in the scheme of **Figure 1D**).

472

473 **Phylogenetic Analysis.**

474 The evolutionary history was inferred by using the “maximum likelihood” (ML)
475 method and JTT matrix-based model (109). The tree with the highest log
476 likelihood (-49687,86) is shown. The percentage of trees in which the

477 associated taxa clustered together is shown next to the branches. Initial tree(s)
478 for the heuristic search was(were) obtained automatically by applying Neighbor-
479 Join and BioNJ algorithms to a matrix of pairwise distances estimated using the
480 JTT model, and then selecting the topology with superior log likelihood value. A
481 discrete Gamma distribution was used to model evolutionary rate differences
482 among sites (5 categories (+G, parameter = 0,6825)). The rate variation model
483 allowed for some sites to be evolutionarily invariable ([+I], 18,05% sites). The
484 tree is drawn to scale, with branch lengths measured in the number of
485 substitutions per site. This analysis involved 140 aa sequences. All positions
486 with less than 95% site coverage were eliminated (i.e., fewer than 5% alignment
487 gaps), and missing data and ambiguous bases were allowed at any position
488 (partial deletion option). There were a total of 829 positions in the final dataset.
489 Evolutionary analyses were conducted in MEGA X (110).

490 Nucleotide sequences have been deposited in GeneBank under the following
491 numbers: KC595156, KC595162, KC595225, KC595227, KC 595189,
492 MH605987, MH605986, KC595190, MH605988, MH605992, MH605991,
493 MH605970, MH605971, KC595223, KC595222, MH605972, MH605975,
494 MH605976, MH605978, MH605973, MH605979, MH605980, MH605981,
495 MH605982, MH605983, MH605984, MK394184, MK394185.

496

497 **Statistical analysis.**

498 Data and statistical analyses were performed using GraphPad Prism, version
499 6.07 (GraphPad Software). Significance when comparing groups was
500 determined with a nonparametric Kruskal-Wallis or by nonparametric Dunn's
501 test for multiple comparisons. A nonparametric Spearman test was used to
502 calculate correlations.

503 **Data Availability**

504 All “accession numbers” and “data” of this work are available.

505

506

507 **Acknowledgements**

508 We want to particularly acknowledge the patients in this study for their
509 participation and to the HIV BioBank integrated in the Spanish AIDS Research
510 Network and collaborating Centres ([http://hivhgmbiobank.com/donor-](http://hivhgmbiobank.com/donor-area/hospitals-and-centres-transferring-samples/?lang=en)
511 [area/hospitals-and-centres-transferring-samples/?lang=en](http://hivhgmbiobank.com/donor-area/hospitals-and-centres-transferring-samples/?lang=en)) for the generous
512 gifts of clinical samples used in this work. The HIV BioBank, integrated in the
513 Spanish AIDS Research Network, is partially funded by the RD16/0025/0019
514 project as part of the Plan Nacional R+D+I and cofinanced by ISCIII-
515 Subdirección General de Evaluación and el Fondo Europeo de Desarrollo
516 Regional (FEDER). The clinical follow-up of Drs. Carmen Rodriguez, Mar Vera
517 and Jorge Del Romero (Centro Sanitario Sandoval), Eulalia Grau (Hospital
518 Germans, Trias y Pujol; irsiCaixa, Badalona) is greatly appreciated.

519

520 **Funding**

521 This work is supported by Spanish AIDS network “Red Temática Cooperativa
522 de Investigación en SIDA” RD12/0017/0002, RD12/0017/0028,
523 RD12/0017/0034, RD16/0025/0011, RDCIII16/0002/0005 and RD16/0025/0041
524 as part of the Plan Nacional R+D+I and cofunded by Spanish “Instituto de Salud
525 Carlos III (ISCIII)-Subdirección General de Evaluación y el Fondo Europeo de
526 Desarrollo Regional (FEDER)”. J.B. is a researcher from “Fundació Institut de
527 Recerca en Ciències de la Salut Germans Trias i Pujol” supported by the Health
528 Department of the Catalan Government/Generalitat de Catalunya and ISCIII
529 grant numbers PI17/01318 and PI20/00093 (to JB). Work in CL-G’ and CC lab
530 was supported by grants SAF (2010-17226) and (2016-77894-R) from MINECO
531 (Spain) and FIS (PI 13/02269, ISCIII). A.V-F's Lab is supported by the
532 European Regional Development Fund (ERDF), RTI2018-093747-B-100

533 (“Ministerio de Ciencia e Innovación”, Spain), “Ministerio de Ciencia, Innovación
534 y Universidades” (Spain), ProID2020010093 (“Agencia Canaria de
535 Investigación, Innovación y Sociedad de la Información” and European Social
536 Fund), UNLL10-3E-783 (ERDF and “Fundación CajaCanarias”) and “SEGAI-
537 ULL”. S.P-Y is funded by “Fundación Doctor Manuel Morales” (La Palma,
538 Spain) and “Contrato Predoctoral Ministerio-ULL Formación de Doctores” (2019
539 Program) (“Ministerio de Ciencia, Innovación y Universidades”, Spain). R.C-R is
540 funded by RD16/0025/0011 and ProID2020010093 (“Agencia Canaria de
541 Investigación, Innovación y Sociedad de la Información” and European Social
542 Fund). J-E-H is funded by the Cabildo Tenerife “Agustin de Betancourt” 2017
543 Program.
544
545

546 **Figures legends**

547 **Figure 1. Outline of the experimental model used for the analysis of Env**
548 **expression, Env-mediated cell-to-cell fusion, viral transfer and viral**
549 **infectivity.** (A) Env expression: HEK-293T cells will be co-transfected with
550 primary of reference full-length viral *env* and a ptat Δenv HIV-1 expression
551 plasmid, allowing Env cell-surface expression in a viral production context. Cell-
552 surface Env expression will be then analyzed by flow cytometry using specific
553 anti-Env antibody. (B) Env-mediated fusion activity: after 24 hours, effector
554 HEK-293T cells producing HIV-1 particles bearing primary or reference Envs
555 will be co-cultured with TZM-bl cells to force synapsis formation and CD4-
556 mediated binding of budding particles to target cells. (C) Env-mediated viral
557 transfer: HEK-293T cells producing HIV-1 particles carrying primary or
558 reference Envs will be co-cultured with primary CD4+ T cells. Then, HIV-1
559 transfer will be analyzed by flow cytometry using specific anti-p24 antibody in
560 target CD4+ T cells. (D) Env-mediated viral infection: TZM-bl cells will be
561 infected with serial dilutions of viral particles obtained from transfected HEK-
562 293T and carrying the different primary or reference HIV-1 Envs. After 48 hours,
563 infectivity capacity will be analyzed by quantifying luciferase assay in infected
564 TZM-bl cells.

565

566 **Figure 2. Analysis of the expression of the different HIV-1-Env**
567 **glycoproteins from LTNP-EC, Viremic LTNP and control progressors**
568 **patients.**

569 Flow cytometry analysis of the cell-surface expression level of the assayed HIV-
570 1 Envs in HEK-293T cells from LTNP-EC (*gray bars*), vLTNP (*green bars*), Old
571 (*orange bars*) and Modern individuals (*red bars*) or reference HIV-1 viral strains

572 (ptat, No Ab2, NL4.3 and BaL, *black bars*). Env protein expression for each
573 patient (A) and Env protein expression in each group of patients comparing
574 mean values between each group (Kruskal-Wallis, Dunn's Multiple
575 Comparisons Test) (B); p value for comparison between all groups is shown,
576 *top left*. Values are mean \pm S.E.M. of three independent experiments.

577

578 **Figure 3. Analysis of membranes fusion-phenotypic features of HIV-1**

579 **Envs isolated from LTNP-EC, viremic LTNP and P individuals.** Analysis of
580 the ability to induce cell-to-cell fusion of HIV-1 Env proteins obtained from
581 LTNP-EC (*gray bars*), vLTNP (*green bars*), Old (*orange bars*) and Modern
582 individuals (*red bars*) or reference HIV-1 viral strains (ptat, NL4.3 and BaL,
583 *black bars*). (A) Env fusogenic activity for each patient in each group. (B)
584 Relative fusion activity of the full Env collection compared to the BaL control
585 established at 100% and grouped in the different groups of patients. Values are
586 mean \pm S.E.M. of three independent experiments. Statistical analysis was
587 performed using Kruskal-Wallis, Dunn's Multiple Comparisons Test; p value for
588 comparison between all groups is shown, *top left*.

589

590 **Figure 4. Analysis of HIV-1 Env-mediated cell-to-cell viral transfer.**

591 Analysis of the ability to induce cell-to-cell virus transfer of HIV-1 Env proteins obtained
592 from LTNP-EC (*gray bars*), vLTNP (*green bars*), Old Patients (*orange bars*), recent
593 patients (Moderns) (*red bars*) or reference HIV-1 viral strains (pSG3, CD4+ cells, NL4.3
594 and BaL, *black bars*). Analysis of HIV-1 Env-mediated cell-to-cell viral transfer for each
595 patient (A) and in each group where P values compare medians between groups using
596 a nonparametric Kruskal-Wallis Test (Kruskal-Wallis, Dunn's Multiple Comparisons

597 Test) (B); p value for comparison between all groups is shown, *top left*. Values
598 are mean \pm S.E.M. of two independent experiments.

599

600 **Figure 5. Viral infectivity of the viral Envs.**

601 Analysis of the infectivity (TCID₅₀ value normalized by viral p24 input) of the different of
602 HIV-1 Env proteins obtained from LTNP-EC (*gray bars*), vLTNP (*green bars*), Old
603 (*orange bars*) and Moderns (*red bars*) patients or reference HIV-1 viral strains (pSG3,
604 NL4.3 and BaL, *black bars*). Analysis of Env infectivity for each patient (A) and in each
605 group where P values compare medians between groups using a nonparametric
606 Kruskal-Wallis, Dunn's Multiple Comparisons Test (B); p value for comparison
607 between all groups is shown, *top left*. Values are mean \pm S.E.M. of three
608 independent experiments.

609

610 **Figure 6. Analysis of the correlation of the fusion, transfer and viral
611 infectivity Env characteristics between groups.**

612 (A) Correlation between Relative fusion and HIV Transfer of all Envs of the
613 different groups LTNP-EC (*gray circle*), vLTNP (*green circle*), Old patients
614 (*orange square*) and Modern patients (*red square*). The correlation was
615 calculated with a nonparametric Spearman test. (B) Correlation between
616 Relative fusion and Infectivity (TCID₅₀ value normalized by viral p24 input) of all
617 Envs of the different groups LTNP-EC (*gray circle*), vLTNP (*green circle*), Old
618 patients (*orange square*) and Modern patients (*red square*). The correlation was
619 calculated with a nonparametric Spearman test. (C) Correlation between
620 Infectivity and HIV Transfer of all Envs of the different groups LTNP-EC (*gray
621 circle*), vLTNP (*green circle*), Old patients (*orange square*) and recent patients
622 Moderns) (*red square*) is shown. The correlation was calculated with a

623 nonparametric Spearman test. Values are mean \pm S.E.M. of three independent
624 experiments; p value for comparison between all groups is shown, *top left*.

625

626 **Figure 7. Phylogenetic analysis of the vial Envs.**

627 The evolutionary history of the Env aa sequences was inferred as described in
628 Materials and Methods using the Maximum Likelihood method and JTT matrix-
629 based model (109). The tree with the highest log likelihood (-49687,86) is
630 shown. The percentage of trees in which the associated taxa clustered together
631 is shown next to the branches. Evolutionary analyses were conducted in MEGA
632 X (110).

633

634 **Figure 8. Analysis of the length and glycosylation sites in the loops of the**
635 **Envs from the different groups.**

636 Analysis of the length of each variable loops V1 (A), V2 (B), V3 (C), V4 (D), V5
637 (E) and all variable loops together (F). The results were grouped (LTNP-ECs:
638 *gray bar*, vLTNPs: *green bar*, Old patients: *orange bar*, and recent patients
639 (Moderns): *red bar*) and compared using a nonparametric Kruskal-Wallis,
640 Dunn's Multiple Comparisons Test; p value for comparison between all groups
641 is shown, *top left*. Values are mean \pm S.E.M. of three independent experiments.

642

643 **Figure 9. Correlation of the expression, fusion, transfer and viral**
644 **infectivity Env characteristics with the nucleotide genetic distance to**
645 **subtype B ancestor.**

646 Correlation between genetic distance to subtype B ancestor of all Envs of the
647 different groups and Env expression (A), Relative fusion (B), HIV Transfer (C)
648 and Infectivity (D). LTNP-ECs (*gray circle*), vLTNPs (*green circle*), Old patients

649 (*orange square*) and Modern patients (*red square*). The correlations were
650 calculated with a nonparametric Spearman test (p and r values are shown, *top*
651 *left*). Values of Env expression, Relative fusion, HIV transfer and Infectivity are
652 mean \pm S.E.M. of three independent experiments.

653

654

655 References

656

- 657 1. **Fellay J, Ge D, Shianna KV, Colombo S, Ledergerber B, Cirulli ET, Urban TJ, Zhang K,**
658 **Gumbs CE, Smith JP, Castagna A, Cozzi-Lepri A, De Luca A, Easterbrook P, Gunthard HF,**
659 **Mallal S, Mussini C, Dalmau J, Martinez-Picado J, Miro JM, Obel N, Wolinsky SM, Martinson**
660 **JJ, Detels R, Margolick JB, Jacobson LP, Descombes P, Antonarakis SE, Beckmann JS, O'Brien**
661 **SJ, Letvin NL, McMichael AJ, Haynes BF, Carrington M, Feng S, Telenti A, Goldstein DB,**
662 **Immunology NCFHAV.** 2009. Common genetic variation and the control of HIV-1 in humans.
663 *PLoS Genet* **5**:e1000791.
- 664 2. **Naranbhai V, Carrington M.** 2017. Host genetic variation and HIV disease: from
665 mapping to mechanism. *Immunogenetics* **69**:489-498.
- 666 3. **Migueles SA, Sabbaghian MS, Shupert WL, Bettinotti MP, Marincola FM, Martino L,**
667 **Hallahan CW, Selig SM, Schwartz D, Sullivan J, Connors M.** 2000. HLA B*5701 is highly
668 associated with restriction of virus replication in a subgroup of HIV-infected long term
669 nonprogressors. *Proc Natl Acad Sci U S A* **97**:2709-2714.
- 670 4. **Miura T, Brockman MA, Schneidewind A, Lobritz M, Pereyra F, Rathod A, Block BL,**
671 **Brumme ZL, Brumme CJ, Baker B, Rothchild AC, Li B, Trocha A, Cutrell E, Frahm N, Brander C,**
672 **Toth I, Arts EJ, Allen TM, Walker BD.** 2009. HLA-B57/B*5801 human immunodeficiency virus
673 type 1 elite controllers select for rare gag variants associated with reduced viral replication
674 capacity and strong cytotoxic T-lymphocyte [corrected] recognition. *J Virol* **83**:2743-2755.
- 675 5. **Migueles SA, Connors M.** 2015. Success and failure of the cellular immune response
676 against HIV-1. *Nat Immunol* **16**:563-570.
- 677 6. **Buckheit RW, 3rd, Allen TG, Alme A, Salgado M, O'Connell KA, Huculak S, Falade-**
678 **Nwulia O, Williams TM, Gallant JE, Siliciano RF, Blankson JN.** 2012. Host factors dictate
679 control of viral replication in two HIV-1 controller/chronic progressor transmission pairs. *Nat*
680 *Commun* **3**:716.
- 681 7. **Goulder P, Deeks SG.** 2018. HIV control: Is getting there the same as staying there?
682 *PLoS Pathog* **14**:e1007222.
- 683 8. **Cortes FH, de Paula HHS, Bello G, Ribeiro-Alves M, de Azevedo SSD, Caetano DG,**
684 **Teixeira SLM, Hoagland B, Grinsztejn B, Veloso VG, Guimaraes ML, Morgado MG.** 2018.
685 Plasmatic Levels of IL-18, IP-10, and Activated CD8(+) T Cells Are Potential Biomarkers to
686 Identify HIV-1 Elite Controllers With a True Functional Cure Profile. *Front Immunol* **9**:1576.
- 687 9. **Martin-Gayo E, Buzon MJ, Ouyang Z, Hickman T, Cronin J, Pimenova D, Walker BD,**
688 **Lichterfeld M, Yu XG.** 2015. Potent Cell-Intrinsic Immune Responses in Dendritic Cells Facilitate
689 HIV-1-Specific T Cell Immunity in HIV-1 Elite Controllers. *PLoS Pathog* **11**:e1004930.
- 690 10. **Deeks SG, Walker BD.** 2007. Human immunodeficiency virus controllers: mechanisms
691 of durable virus control in the absence of antiretroviral therapy. *Immunity* **27**:406-416.
- 692 11. **Pernas M, Casado C, Arcones C, Llano A, Sanchez-Merino V, Mothe B, Vicario JL, Grau**
693 **E, Ruiz L, Sanchez J, Telenti A, Yuste E, Brander C, Galindez CL.** 2012. Low-replicating viruses
694 and strong anti-viral immune response associated with prolonged disease control in a
695 superinfected HIV-1 LTNP elite controller. *PLoS One* **7**:e31928.

- 696 12. **Carrington M, Walker BD.** 2012. Immunogenetics of spontaneous control of HIV. *Annu*
697 *Rev Med* **63**:131-145.
- 698 13. **Migueles SA, Osborne CM, Royce C, Compton AA, Joshi RP, Weeks KA, Rood JE,**
699 **Berkley AM, Sacha JB, Cogliano-Shutta NA, Lloyd M, Roby G, Kwan R, McLaughlin M, Stallings**
700 **S, Rehm C, O'Shea MA, Mican J, Packard BZ, Komoriya A, Palmer S, Wiegand AP, Maldarelli F,**
701 **Coffin JM, Mellors JW, Hallahan CW, Follman DA, Connors M.** 2008. Lytic granule loading of
702 CD8+ T cells is required for HIV-infected cell elimination associated with immune control.
703 *Immunity* **29**:1009-1021.
- 704 14. **Goulder PJ, Phillips RE, Colbert RA, McAdam S, Ogg G, Nowak MA, Giangrande P,**
705 **Luzzi G, Morgan B, Edwards A, McMichael AJ, Rowland-Jones S.** 1997. Late escape from an
706 immunodominant cytotoxic T-lymphocyte response associated with progression to AIDS. *Nat*
707 *Med* **3**:212-217.
- 708 15. **Buckheit RW, 3rd, Salgado M, Siliciano RF, Blankson JN.** 2012. Inhibitory potential of
709 subpopulations of CD8+ T cells in HIV-1-infected elite suppressors. *J Virol* **86**:13679-13688.
- 710 16. **Blankson JN.** 2010. Effector mechanisms in HIV-1 infected elite controllers: highly
711 active immune responses? *Antiviral Res* **85**:295-302.
- 712 17. **Casado C, Pernas M, Sandonis V, Alvaro-Cifuentes T, Olivares I, Fuentes R, Martinez-**
713 **Prats L, Grau E, Ruiz L, Delgado R, Rodriguez C, del Romero J, Lopez-Galindez C.** 2013.
714 Identification of a cluster of HIV-1 controllers infected with low replicating viruses. *PLoS One*
715 **8**:e77663.
- 716 18. **Casado C, Marrero-Hernandez S, Marquez-Arce D, Pernas M, Marfil S, Borrás-**
717 **Granana F, Olivares I, Cabrera-Rodriguez R, Valera MS, de Armas-Rillo L, Lemey P, Blanco J,**
718 **Valenzuela-Fernandez A, Lopez-Galindez C.** 2018. Viral Characteristics Associated with the
719 Clinical Nonprogressor Phenotype Are Inherited by Viruses from a Cluster of HIV-1 Elite
720 Controllers. *mBio* **9**.
- 721 19. **Cabrera-Rodriguez R, Heilmann V, Marfil S, Pernas M, Marrero-Hernandez S, Cabrera**
722 **C, Urrea V, Casado C, Olivares I, Marquez-Arce D, Perez-Yanes S, Estevez-Herrera J, Clotet B,**
723 **Espert L, Lopez-Galindez C, Biard-Piechaczyk M, Valenzuela-Fernandez A, Blanco J.** 2019. HIV-
724 1 envelope glycoproteins isolated from Viremic Non-Progressor individuals are fully functional
725 and cytopathic. *Sci Rep* **9**:5544.
- 726 20. **Lassen KG, Lobritz MA, Bailey JR, Johnston S, Nguyen S, Lee B, Chou T, Siliciano RF,**
727 **Markowitz M, Arts EJ.** 2009. Elite suppressor-derived HIV-1 envelope glycoproteins exhibit
728 reduced entry efficiency and kinetics. *PLoS Pathog* **5**:e1000377.
- 729 21. **Dalmau J, Puertas MC, Azuara M, Marino A, Frahm N, Mothe B, Izquierdo-Useros N,**
730 **Buzon MJ, Paredes R, Matas L, Allen TM, Brander C, Rodrigo C, Clotet B, Martinez-Picado J.**
731 2009. Contribution of immunological and virological factors to extremely severe primary HIV
732 type 1 infection. *Clin Infect Dis* **48**:229-238.
- 733 22. **Jarrin I, Pantazis N, Dalmau J, Phillips AN, Olson A, Mussini C, Boufassa F, Costagliola**
734 **D, Porter K, Blanco J, Del Amo J, Martinez-Picado J.** 2015. Does rapid HIV disease progression
735 prior to combination antiretroviral therapy hinder optimal CD4+ T-cell recovery once HIV-1
736 suppression is achieved? *Aids* **29**:2323-2333.
- 737 23. **Casado C, Galvez C, Pernas M, Tarancon-Diez L, Rodriguez C, Sanchez-Merino V, Vera**
738 **M, Olivares I, De Pablo-Bernal R, Merino-Mansilla A, Del Romero J, Lorenzo-Redondo R, Ruiz-**

- 739 **Mateos E, Salgado M, Martinez-Picado J, Lopez-Galindez C.** 2020. Permanent control of HIV-1
740 pathogenesis in exceptional elite controllers: a model of spontaneous cure. *Sci Rep* **10**:1902.
- 741 24. **Lopez-Galindez C, Pernas M, Casado C, Olivares I, Lorenzo-Redondo R.** 2019. Elite
742 controllers and lessons learned for HIV-1 cure. *Curr Opin Virol* **38**:31-36.
- 743 25. **Checkley MA, Luttge BG, Freed EO.** 2011. HIV-1 envelope glycoprotein biosynthesis,
744 trafficking, and incorporation. *J Mol Biol* **410**:582-608.
- 745 26. **Berger EA.** 2015. Finding Fusin/CXCR4, the First "2nd Receptor" for HIV Entry. *Front*
746 *Immunol* **6**:283.
- 747 27. **Miedema F, Meyaard L, Koot M, Klein MR, Roos MT, Groenink M, Fouchier RA, Van't**
748 **Wout AB, Tersmette M, Schellekens PT, et al.** 1994. Changing virus-host interactions in the
749 course of HIV-1 infection. *Immunol Rev* **140**:35-72.
- 750 28. **Beitari S, Wang Y, Liu SL, Liang C.** 2019. HIV-1 Envelope Glycoprotein at the Interface
751 of Host Restriction and Virus Evasion. *Viruses* **11**.
- 752 29. **Engelman A, Cherepanov P.** 2012. The structural biology of HIV-1: mechanistic and
753 therapeutic insights. *Nat Rev Microbiol* **10**:279-290.
- 754 30. **Chan DC, Kim PS.** 1998. HIV entry and its inhibition. *Cell* **93**:681-684.
- 755 31. **Wyatt R, Sodroski J.** 1998. The HIV-1 envelope glycoproteins: fusogens, antigens, and
756 immunogens. *Science* **280**:1884-1888.
- 757 32. **Alkhatib G, Combadiere C, Broder CC, Feng Y, Kennedy PE, Murphy PM, Berger EA.**
758 1996. CC CKR5: a RANTES, MIP-1alpha, MIP-1beta receptor as a fusion cofactor for
759 macrophage-tropic HIV-1. *Science* **272**:1955-1958.
- 760 33. **Choe H, Farzan M, Sun Y, Sullivan N, Rollins B, Ponath PD, Wu L, Mackay CR, LaRosa**
761 **G, Newman W, Gerard N, Gerard C, Sodroski J.** 1996. The beta-chemokine receptors CCR3 and
762 CCR5 facilitate infection by primary HIV-1 isolates. *Cell* **85**:1135-1148.
- 763 34. **Oberlin E, Amara A, Bachelier F, Bessia C, Virelizier JL, Arenzana-Seisdedos F,**
764 **Schwartz O, Heard JM, Clark-Lewis I, Legler DF, Loetscher M, Baggiolini M, Moser B.** 1996.
765 The CXC chemokine SDF-1 is the ligand for LESTR/fusin and prevents infection by T-cell-line-
766 adapted HIV-1. *Nature* **382**:833-835.
- 767 35. **Feng Y, Broder CC, Kennedy PE, Berger EA.** 1996. HIV-1 entry cofactor: functional
768 cDNA cloning of a seven-transmembrane, G protein-coupled receptor. *Science* **272**:872-877.
- 769 36. **Deng H, Liu R, Ellmeier W, Choe S, Unutmaz D, Burkhart M, Di Marzio P, Marmon S,**
770 **Sutton RE, Hill CM, Davis CB, Peiper SC, Schall TJ, Littman DR, Landau NR.** 1996. Identification
771 of a major co-receptor for primary isolates of HIV-1. *Nature* **381**:661-666.
- 772 37. **Dalgleish AG, Beverley PC, Clapham PR, Crawford DH, Greaves MF, Weiss RA.** 1984.
773 The CD4 (T4) antigen is an essential component of the receptor for the AIDS retrovirus. *Nature*
774 **312**:763-767.
- 775 38. **Klatzmann D, Champagne E, Chamaret S, Gruest J, Guetard D, Hercend T, Gluckman**
776 **JC, Montagnier L.** 1984. T-lymphocyte T4 molecule behaves as the receptor for human
777 retrovirus LAV. *Nature* **312**:767-768.

- 778 39. **Blanco J, Valenzuela A, Herrera C, Lluís C, Hovanessian AG, Franco R.** 2000. The HIV-1
779 gp120 inhibits the binding of adenosine deaminase to CD26 by a mechanism modulated by
780 CD4 and CXCR4 expression. *FEBS Lett* **477**:123-128.
- 781 40. **Garcia-Perez J, Staropoli I, Azoulay S, Heinrich JT, Cascajero A, Colin P, Lortat-Jacob H,**
782 **Arenzana-Seisdedos F, Alcami J, Kellenberger E, Lagane B.** 2015. A single-residue change in
783 the HIV-1 V3 loop associated with maraviroc resistance impairs CCR5 binding affinity while
784 increasing replicative capacity. *Retrovirology* **12**:50.
- 785 41. **Colin P, Bénureau Y, Staropoli I, Wang Y, Gonzalez N, Alcami J, Hartley O, Brelot A,**
786 **Arenzana-Seisdedos F, Lagane B.** 2013. HIV-1 exploits CCR5 conformational heterogeneity to
787 escape inhibition by chemokines. *Proc Natl Acad Sci U S A* **110**:9475-9480.
- 788 42. **Herschhorn A, Gu C, Moraca F, Ma X, Farrell M, Smith AB, 3rd, Pancera M, Kwong PD,**
789 **Schön A, Freire E, Abrams C, Blanchard SC, Mothes W, Sodroski JG.** 2017. The β 20- β 21 of
790 gp120 is a regulatory switch for HIV-1 Env conformational transitions. *Nat Commun* **8**:1049.
- 791 43. **Valenzuela-Fernandez A, Alvarez S, Gordon-Alonso M, Barrero M, Ursa A, Cabrero JR,**
792 **Fernandez G, Naranjo-Suarez S, Yanez-Mo M, Serrador JM, Munoz-Fernandez MA, Sanchez-**
793 **Madrid F.** 2005. Histone deacetylase 6 regulates human immunodeficiency virus type 1
794 infection. *Mol Biol Cell* **16**:5445-5454.
- 795 44. **Garcia-Exposito L, Ziglio S, Barroso-Gonzalez J, de Armas-Rillo L, Valera MS, Zipeto D,**
796 **Machado JD, Valenzuela-Fernandez A.** 2013. Gelsolin activity controls efficient early HIV-1
797 infection. *Retrovirology* **10**:39.
- 798 45. **Barrero-Villar M, Cabrero JR, Gordon-Alonso M, Barroso-Gonzalez J, Alvarez-Losada**
799 **S, Munoz-Fernandez MA, Sanchez-Madrid F, Valenzuela-Fernandez A.** 2009. Moesin is
800 required for HIV-1-induced CD4-CXCR4 interaction, F-actin redistribution, membrane fusion
801 and viral infection in lymphocytes. *J Cell Sci* **122**:103-113.
- 802 46. **Kafando A, Martineau C, El-Far M, Fournier E, Doualla-Bell F, Serhir B, Kazienga A,**
803 **Sangare MN, Sylla M, Chamberland A, Charest H, Tremblay CL.** 2019. HIV-1 Envelope
804 Glycoprotein Amino Acids Signatures Associated with Clade B Transmitted/Founder and
805 Recent Viruses. *Viruses* **11**.
- 806 47. **Marconi V, Bonhoeffer S, Paredes R, Lu J, Hoh R, Martin JN, Deeks SG, Kuritzkes DR.**
807 2008. Viral dynamics and in vivo fitness of HIV-1 in the presence and absence of enfuvirtide. *J*
808 *Acquir Immune Defic Syndr* **48**:572-576.
- 809 48. **Blanco J, Bosch B, Fernandez-Figueras MT, Barretina J, Clotet B, Este JA.** 2004. High
810 level of coreceptor-independent HIV transfer induced by contacts between primary CD4 T
811 cells. *J Biol Chem* **279**:51305-51314.
- 812 49. **Curlin ME, Zioni R, Hawes SE, Liu Y, Deng W, Gottlieb GS, Zhu T, Mullins JI.** 2010. HIV-
813 1 envelope subregion length variation during disease progression. *PLoS Pathog* **6**:e1001228.
- 814 50. **Yuan T, Li J, Zhang MY.** 2013. HIV-1 envelope glycoprotein variable loops are
815 indispensable for envelope structural integrity and virus entry. *PLoS One* **8**:e69789.
- 816 51. **Sagar M, Wu X, Lee S, Overbaugh J.** 2006. Human immunodeficiency virus type 1 V1-
817 V2 envelope loop sequences expand and add glycosylation sites over the course of infection,
818 and these modifications affect antibody neutralization sensitivity. *J Virol* **80**:9586-9598.

- 819 52. **Gnanakaran S, Bhattacharya T, Daniels M, Keele BF, Hraber PT, Lapedes AS, Shen T,**
820 **Gaschen B, Krishnamoorthy M, Li H, Decker JM, Salazar-Gonzalez JF, Wang S, Jiang C, Gao F,**
821 **Swanstrom R, Anderson JA, Ping LH, Cohen MS, Markowitz M, Goepfert PA, Saag MS, Eron JJ,**
822 **Hicks CB, Blattner WA, Tomaras GD, Asmal M, Letvin NL, Gilbert PB, Decamp AC, Magaret**
823 **CA, Schief WR, Ban YE, Zhang M, Soderberg KA, Sodroski JG, Haynes BF, Shaw GM, Hahn BH,**
824 **Korber B.** 2011. Recurrent signature patterns in HIV-1 B clade envelope glycoproteins
825 associated with either early or chronic infections. *PLoS Pathog* **7**:e1002209.
- 826 53. **Wang S, Voronin Y, Zhao P, Ishihara M, Mehta N, Porterfield M, Chen Y, Bartley C, Hu**
827 **G, Han D, Wells L, Tiemeyer M, Lu S.** 2020. Glycan Profiles of gp120 Protein Vaccines from
828 Four Major HIV-1 Subtypes Produced from Different Host Cell Lines under Non-GMP or GMP
829 Conditions. *J Virol* **94**.
- 830 54. **Go EP, Irungu J, Zhang Y, Dalpathado DS, Liao HX, Sutherland LL, Alam SM, Haynes**
831 **BF, Desaire H.** 2008. Glycosylation site-specific analysis of HIV envelope proteins (JR-FL and
832 CON-S) reveals major differences in glycosylation site occupancy, glycoform profiles, and
833 antigenic epitopes' accessibility. *J Proteome Res* **7**:1660-1674.
- 834 55. **Sterjovski J, Churchill MJ, Ellett A, Gray LR, Roche MJ, Dunfee RL, Purcell DF, Saksena**
835 **N, Wang B, Sonza S, Wesselingh SL, Karlsson I, Fenyo EM, Gabuzda D, Cunningham AL, Gorry**
836 **PR.** 2007. Asn 362 in gp120 contributes to enhanced fusogenicity by CCR5-restricted HIV-1
837 envelope glycoprotein variants from patients with AIDS. *Retrovirology* **4**:89.
- 838 56. **Coffin JM.** 1995. HIV population dynamics in vivo: implications for genetic variation,
839 pathogenesis, and therapy. *Science* **267**:483-489.
- 840 57. **Fideli US, Allen SA, Musonda R, Trask S, Hahn BH, Weiss H, Mulenga J, Kasolo F,**
841 **Vermund SH, Aldrovandi GM.** 2001. Virologic and immunologic determinants of heterosexual
842 transmission of human immunodeficiency virus type 1 in Africa. *AIDS Res Hum Retroviruses*
843 **17**:901-910.
- 844 58. **Hecht FM, Hartogensis W, Bragg L, Bacchetti P, Atchison R, Grant R, Barbour J, Deeks**
845 **SG.** 2010. HIV RNA level in early infection is predicted by viral load in the transmission source.
846 *AIDS* **24**:941-945.
- 847 59. **Mellors JW, Kingsley LA, Rinaldo CR, Jr., Todd JA, Hoo BS, Kokka RP, Gupta P.** 1995.
848 Quantitation of HIV-1 RNA in plasma predicts outcome after seroconversion. *Ann Intern Med*
849 **122**:573-579.
- 850 60. **Mellors JW, Rinaldo CR, Jr., Gupta P, White RM, Todd JA, Kingsley LA.** 1996. Prognosis
851 in HIV-1 infection predicted by the quantity of virus in plasma. *Science* **272**:1167-1170.
- 852 61. **Pilcher CD, Joaki G, Hoffman IF, Martinson FE, Mapanje C, Stewart PW, Powers KA,**
853 **Galvin S, Chilongozi D, Gama S, Price MA, Fiscus SA, Cohen MS.** 2007. Amplified transmission
854 of HIV-1: comparison of HIV-1 concentrations in semen and blood during acute and chronic
855 infection. *AIDS* **21**:1723-1730.
- 856 62. **Quinn TC, Wawer MJ, Sewankambo N, Serwadda D, Li C, Wabwire-Mangen F,**
857 **Meehan MO, Lutalo T, Gray RH.** 2000. Viral load and heterosexual transmission of human
858 immunodeficiency virus type 1. Rakai Project Study Group. *N Engl J Med* **342**:921-929.
- 859 63. **Yue L, Prentice HA, Farmer P, Song W, He D, Lakhi S, Goepfert P, Gilmour J, Allen S,**
860 **Tang J, Kaslow RA, Hunter E.** 2013. Cumulative impact of host and viral factors on HIV-1 viral-
861 load control during early infection. *J Virol* **87**:708-715.

- 862 64. **Balasubramaniam M, Pandhare J, Dash C.** 2019. Immune Control of HIV. *J Life Sci*
863 (Westlake Village) **1**:4-37.
- 864 65. **van Gils MJ, Bunnik EM, Boeser-Nunnink BD, Burger JA, Terlouw-Klein M, Verwer N,**
865 **Schuitemaker H.** 2011. Longer V1V2 region with increased number of potential N-linked
866 glycosylation sites in the HIV-1 envelope glycoprotein protects against HIV-specific neutralizing
867 antibodies. *J Virol* **85**:6986-6995.
- 868 66. **Rong R, Bibollet-Ruche F, Mulenga J, Allen S, Blackwell JL, Derdeyn CA.** 2007. Role of
869 V1V2 and other human immunodeficiency virus type 1 envelope domains in resistance to
870 autologous neutralization during clade C infection. *J Virol* **81**:1350-1359.
- 871 67. **Moore PL, Gray ES, Choge IA, Ranchobe N, Mlisana K, Abdool Karim SS, Williamson C,**
872 **Morris L.** 2008. The c3-v4 region is a major target of autologous neutralizing antibodies in
873 human immunodeficiency virus type 1 subtype C infection. *J Virol* **82**:1860-1869.
- 874 68. **Castro E, Bélair M, Rizzardi GP, Bart PA, Pantaleo G, Graziosi C.** 2008. Independent
875 evolution of hypervariable regions of HIV-1 gp120: V4 as a swarm of N-Linked glycosylation
876 variants. *AIDS Res Hum Retroviruses* **24**:106-113.
- 877 69. **Davenport YW, West AP, Jr., Bjorkman PJ.** 2016. Structure of an HIV-2 gp120 in
878 Complex with CD4. *J Virol* **90**:2112-2118.
- 879 70. **Wang L, Izadmehr S, Kamau E, Kong XP, Chen BK.** 2019. Sequential trafficking of Env
880 and Gag to HIV-1 T cell virological synapses revealed by live imaging. *Retrovirology* **16**:2.
- 881 71. **Mishra N, Makhdoomi MA, Sharma S, Kumar S, Dobhal A, Kumar D, Chawla H, Singh**
882 **R, Kanga U, Das BK, Lodha R, Kabra SK, Luthra K.** 2019. Viral Characteristics Associated with
883 Maintenance of Elite Neutralizing Activity in Chronically HIV-1 Clade C-Infected Monozygotic
884 Pediatric Twins. *J Virol* **93**.
- 885 72. **Coffin J, Swanstrom R.** 2013. HIV pathogenesis: dynamics and genetics of viral
886 populations and infected cells. *Cold Spring Harb Perspect Med* **3**:a012526.
- 887 73. **Andreano E, Rappuoli R.** 2021. SARS-CoV-2 escaped natural immunity, raising
888 questions about vaccines and therapies. *Nat Med* **27**:759-761.
- 889 74. **Garcia-Beltran WF, Lam EC, St Denis K, Nitido AD, Garcia ZH, Hauser BM, Feldman J,**
890 **Pavlovic MN, Gregory DJ, Poznansky MC, Sigal A, Schmidt AG, lafrate AJ, Naranbhai V, Balazs**
891 **AB.** 2021. Multiple SARS-CoV-2 variants escape neutralization by vaccine-induced humoral
892 immunity. *Cell* **184**:2372-2383.e2379.
- 893 75. **Bieniasz PD, Fridell RA, Aramori I, Ferguson SS, Caron MG, Cullen BR.** 1997. HIV-1-
894 induced cell fusion is mediated by multiple regions within both the viral envelope and the CCR-
895 5 co-receptor. *Embo j* **16**:2599-2609.
- 896 76. **Isaka Y, Sato A, Miki S, Kawauchi S, Sakaida H, Hori T, Uchiyama T, Adachi A, Hayami**
897 **M, Fujiwara T, Yoshie O.** 1999. Small amino acid changes in the V3 loop of human
898 immunodeficiency virus type 2 determines the coreceptor usage for CXCR4 and CCR5. *Virology*
899 **264**:237-243.
- 900 77. **Cocchi F, DeVico AL, Garzino-Demo A, Cara A, Gallo RC, Lusso P.** 1996. The V3 domain
901 of the HIV-1 gp120 envelope glycoprotein is critical for chemokine-mediated blockade of
902 infection. *Nat Med* **2**:1244-1247.

- 903 78. **Speck RF, Wehrly K, Platt EJ, Atchison RE, Charo IF, Kabat D, Chesebro B, Goldsmith**
904 **MA.** 1997. Selective employment of chemokine receptors as human immunodeficiency virus
905 type 1 coreceptors determined by individual amino acids within the envelope V3 loop. *J Virol*
906 **71**:7136-7139.
- 907 79. **Wu L, Gerard NP, Wyatt R, Choe H, Parolin C, Ruffing N, Borsetti A, Cardoso AA,**
908 **Desjardin E, Newman W, Gerard C, Sodroski J.** 1996. CD4-induced interaction of primary HIV-1
909 gp120 glycoproteins with the chemokine receptor CCR-5. *Nature* **384**:179-183.
- 910 80. **Trkola A, Dragic T, Arthos J, Binley JM, Olson WC, Allaway GP, Cheng-Mayer C,**
911 **Robinson J, Maddon PJ, Moore JP.** 1996. CD4-dependent, antibody-sensitive interactions
912 between HIV-1 and its co-receptor CCR-5. *Nature* **384**:184-187.
- 913 81. **Valenzuela A, Blanco J, Krust B, Franco R, Hovanessian AG.** 1997. Neutralizing
914 antibodies against the V3 loop of human immunodeficiency virus type 1 gp120 block the CD4-
915 dependent and -independent binding of virus to cells. *J Virol* **71**:8289-8298.
- 916 82. **Pollakis G, Baan E, van Werkhoven MB, Berkhout B, Bakker M, Jurriaans S, Paxton**
917 **WA.** 2015. Association between gp120 envelope V1V2 and V4V5 variable loop profiles in a
918 defined HIV-1 transmission cluster. *Aids* **29**:1161-1171.
- 919 83. **Liu Y, Curlin ME, Diem K, Zhao H, Ghosh AK, Zhu H, Woodward AS, Maenza J, Stevens**
920 **CE, Stekler J, Collier AC, Genowati I, Deng W, Zioni R, Corey L, Zhu T, Mullins JI.** 2008. Env
921 length and N-linked glycosylation following transmission of human immunodeficiency virus
922 Type 1 subtype B viruses. *Virology* **374**:229-233.
- 923 84. **Dang LVP, Pham HV, Dinh TT, Nguyen TH, Vu QTH, Vu NTP, Le PTB, Nguyen LV, Le HT,**
924 **Vu PT, Olson L.** 2020. Characterization of envelope sequence of HIV virus in children infected
925 with HIV in Vietnam. *SAGE Open Med* **8**:2050312120937198.
- 926 85. **Fraser C, Lythgoe K, Leventhal GE, Shirreff G, Hollingsworth TD, Alizon S, Bonhoeffer**
927 **S.** 2014. Virulence and pathogenesis of HIV-1 infection: an evolutionary perspective. *Science*
928 **343**:1243727.
- 929 86. **Fischer W, Ganusov VV, Giorgi EE, Hraber PT, Keele BF, Leitner T, Han CS, Gleasner**
930 **CD, Green L, Lo CC, Nag A, Wallstrom TC, Wang S, McMichael AJ, Haynes BF, Hahn BH,**
931 **Perelson AS, Borrow P, Shaw GM, Bhattacharya T, Korber BT.** 2010. Transmission of single
932 HIV-1 genomes and dynamics of early immune escape revealed by ultra-deep sequencing. *PLoS*
933 *One* **5**:e12303.
- 934 87. **Quan Y, Liang C, Brenner BG, Wainberg MA.** 2009. Multidrug-resistant variants of HIV
935 type 1 (HIV-1) can exist in cells as defective quasispecies and be rescued by superinfection with
936 other defective HIV-1 variants. *J Infect Dis* **200**:1479-1483.
- 937 88. **Roche M, Jakobsen MR, Sterjovski J, Ellett A, Posta F, Lee B, Jubb B, Westby M, Lewin**
938 **SR, Ramsland PA, Churchill MJ, Gorry PR.** 2011. HIV-1 escape from the CCR5 antagonist
939 maraviroc associated with an altered and less-efficient mechanism of gp120-CCR5 engagement
940 that attenuates macrophage tropism. *J Virol* **85**:4330-4342.
- 941 89. **Keele BF, Giorgi EE, Salazar-Gonzalez JF, Decker JM, Pham KT, Salazar MG, Sun C,**
942 **Grayson T, Wang S, Li H, Wei X, Jiang C, Kirchherr JL, Gao F, Anderson JA, Ping LH, Swanstrom**
943 **R, Tomaras GD, Blattner WA, Goepfert PA, Kilby JM, Saag MS, Delwart EL, Busch MP, Cohen**
944 **MS, Montefiori DC, Haynes BF, Gaschen B, Athreya GS, Lee HY, Wood N, Seoighe C, Perelson**
945 **AS, Bhattacharya T, Korber BT, Hahn BH, Shaw GM.** 2008. Identification and characterization

- 946 of transmitted and early founder virus envelopes in primary HIV-1 infection. Proc Natl Acad Sci
947 U S A **105**:7552-7557.
- 948 90. **Hunt PW, Harrigan PR, Huang W, Bates M, Williamson DW, McCune JM, Price RW,**
949 **Spudich SS, Lampiris H, Hoh R, Leigler T, Martin JN, Deeks SG.** 2006. Prevalence of CXCR4
950 tropism among antiretroviral-treated HIV-1-infected patients with detectable viremia. J Infect
951 Dis **194**:926-930.
- 952 91. **Moore PL, Gray ES, Morris L.** 2009. Specificity of the autologous neutralizing antibody
953 response. Curr Opin HIV AIDS **4**:358-363.
- 954 92. **Kassaye S, Johnston E, McColgan B, Kantor R, Zijenah L, Katzenstein D.** 2009.
955 Envelope coreceptor tropism, drug resistance, and viral evolution among subtype C HIV-1-
956 infected individuals receiving nonsuppressive antiretroviral therapy. J Acquir Immune Defic
957 Syndr **50**:9-18.
- 958 93. **Shi B, Kitchen C, Weiser B, Mayers D, Foley B, Kemal K, Anastos K, Suchard M, Parker**
959 **M, Brunner C, Burger H.** 2010. Evolution and recombination of genes encoding HIV-1 drug
960 resistance and tropism during antiretroviral therapy. Virology **404**:5-20.
- 961 94. **Duenas-Decamp MJ, Peters P, Burton D, Clapham PR.** 2008. Natural resistance of
962 human immunodeficiency virus type 1 to the CD4bs antibody b12 conferred by a glycan and an
963 arginine residue close to the CD4 binding loop. J Virol **82**:5807-5814.
- 964 95. **Salazar-Gonzalez JF, Bailes E, Pham KT, Salazar MG, Guffey MB, Keele BF, Derdeyn**
965 **CA, Farmer P, Hunter E, Allen S, Manigart O, Mulenga J, Anderson JA, Swanstrom R, Haynes**
966 **BF, Athreya GS, Korber BT, Sharp PM, Shaw GM, Hahn BH.** 2008. Deciphering human
967 immunodeficiency virus type 1 transmission and early envelope diversification by single-
968 genome amplification and sequencing. J Virol **82**:3952-3970.
- 969 96. **Kitrinos KM, Hoffman NG, Nelson JA, Swanstrom R.** 2003. Turnover of env variable
970 region 1 and 2 genotypes in subjects with late-stage human immunodeficiency virus type 1
971 infection. J Virol **77**:6811-6822.
- 972 97. **Kitrinos KM, Nelson JA, Resch W, Swanstrom R.** 2005. Effect of a protease inhibitor-
973 induced genetic bottleneck on human immunodeficiency virus type 1 env gene populations. J
974 Virol **79**:10627-10637.
- 975 98. **Kitchen CM, Philpott S, Burger H, Weiser B, Anastos K, Suchard MA.** 2004. Evolution
976 of human immunodeficiency virus type 1 coreceptor usage during antiretroviral Therapy: a
977 Bayesian approach. J Virol **78**:11296-11302.
- 978 99. **Koot M, Keet IP, Vos AH, de Goede RE, Roos MT, Coutinho RA, Miedema F,**
979 **Schellekens PT, Tersmette M.** 1993. Prognostic value of HIV-1 syncytium-inducing phenotype
980 for rate of CD4+ cell depletion and progression to AIDS. Ann Intern Med **118**:681-688.
- 981 100. **Silver ZA, Dickinson GM, Seaman MS, Desrosiers RC.** 2019. A Highly Unusual V1
982 Region of Env in an Elite Controller of HIV Infection. J Virol **93**.
- 983 101. **Desrosiers RC.** 1998. Prospects for live attenuated HIV. Nat Med **4**:982.
- 984 102. **Almond N, Stott J.** 1999. Live attenuated SIV--a model of a vaccine for AIDS. Immunol
985 Lett **66**:167-170.

- 986 103. **Johnson RP.** 1999. Live attenuated AIDS vaccines: hazards and hopes. *Nat Med* **5**:154-
987 155.
- 988 104. **Murphey-Corb M.** 1997. Live-attenuated HIV vaccines: how safe is safe enough? *Nat*
989 *Med* **3**:17-18.
- 990 105. **Mills J, Desrosiers R, Rud E, Almond N.** 2000. Live attenuated HIV vaccines: a proposal
991 for further research and development. *AIDS Res Hum Retroviruses* **16**:1453-1461.
- 992 106. **Baba TW, Liska V, Khimani AH, Ray NB, Dailey PJ, Penninck D, Bronson R, Greene MF,**
993 **McClure HM, Martin LN, Ruprecht RM.** 1999. Live attenuated, multiply deleted simian
994 immunodeficiency virus causes AIDS in infant and adult macaques. *Nat Med* **5**:194-203.
- 995 107. **Blower SM, Koelle K, Kirschner DE, Mills J.** 2001. Live attenuated HIV vaccines:
996 predicting the tradeoff between efficacy and safety. *Proc Natl Acad Sci U S A* **98**:3618-3623.
- 997 108. **Curriu M, Fausther-Bovendo H, Pernas M, Massanella M, Carrillo J, Cabrera C, Lopez-**
998 **Galindez C, Clotet B, Debre P, Vieillard V, Blanco J.** 2012. Viremic HIV infected individuals with
999 high CD4 T cells and functional envelope proteins show anti-gp41 antibodies with unique
1000 specificity and function. *PLoS One* **7**:e30330.
- 1001 109. **Jones DT, Taylor WR, Thornton JM.** 1992. The rapid generation of mutation data
1002 matrices from protein sequences. *Comput Appl Biosci* **8**:275-282.
- 1003 110. **Kumar S, Stecher G, Li M, Knyaz C, Tamura K.** 2018. MEGA X: Molecular Evolutionary
1004 Genetics Analysis across Computing Platforms. *Mol Biol Evol* **35**:1547-1549.
1005
1006
1007

1008

Table 1. Epidemiological, clinical and host characteristics of the viral Envs.

Clinical Group	Sub-group	Env code ^a	Patient Identification code	Viral Load ^g (at sampling)	Diagnostic time	Sampling time	Viral dating ^b	HLA B		
LTNP	EC	1	2057906-3	< 50	1993	2004	1989	4901/5701		
		2	3227050	< 50	1988	2004	1991	0702/5201		
		3	3227058-3	< 50	1992	2004	1991	1402/1402		
		6	20044616-3	< 50	1998	2004	1999	1501/5703		
		7	10246788	< 50	1992	2005	1993	4402/5701		
		8		< 50	"	"	"	"		
		9	MDM ^c	507	1998	1996	1987	4402/3501		
		10	c	< 50	"	2011	1996	"		
		11	c	< 50	"	2005	"	"		
		12	c	< 50	"	2005	"	"		
		Viremic	13	4022834	3.710	1994	2004	ND	1401/4403	
			14	9684	2.557	1998	2005	1994	1302/4001	
	15		2988465	2.286	1993	2004	1999	1402/2705		
	16		38 17 5	418	1996	2014	1999	2705/5801		
	17				"	"	"	"		
	18				"	"	"	"		
	19				"	"	"	"		
	21		30	7.597	1989	1998	2000	1501/3501		
	22		64	11.926	1989	1999	1999	4402/4901		
	24			"	"	2002	"	"		
	Progressor		Old	26	V10 ^d	N.D.	1993	1994	1999	4002/4402
				27			"	"	"	
		28		V13	N.D.	1992	1994	1990	0702/1402	
		30		L10	89.000	---	1993	1993	1501/4901	
31				"	"	"	"	"		
32		L 11		42.000	1993	1993	2000	1801/5101		
33				"	"	"	"	"		
36		I14 ^d		130.000	1987	1994	2002	0702/3502		
37		d	"	"	"	"	"			
38		I18	170.000	1991	1994	1990	1402/4403			
Modern		39	ESI 17A	156.300	2014	2013	N.A. ^f	4201/4402		
		40		"	"	"	"	"		
		42	ESI 39A	137.700	2012	2014	N.A.	1517/3801		
		43		"	"	"	"	"		
		44	ESI 41A	129.700	2012	2014	N.A.	3503/5701		
		45		"	"	"	"	"		
	46	ESI 5A 2	49.107	2004	2007	N.A.	4102/4402			
	49	ESI 42 A	11.510	2011	2014	N.A.	1402/4403			
50		"	"	"	"	"				
51	ESI 42 B	41.090	2011	2014	N.A.	0702/1501				
52		"	"	"	"	"				

1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021

^aHIV-1 Env number used in this study and identification codes.

^bAccording to Bello et al. (2004). *J Gen Virol.* Feb;85(Pt 2):399-407. doi: 10.1099/vir.0.19365-0. PMID: 14769897.

^cDouble infected individual (Casado et al. (2007) *J Infect Dis.* 2007 Sep 15;196(6):895-9. doi: 10.1086/520885. Epub 2007 Aug 14. PMID: 17703421).

^dIndividuals with a short antiviral therapy (AZT (zidovudine) and DDI (didanosine) for V10 patient and AZT for I14 patient).

^eThe Modern Individuals have been infected within 3 years.

^fN.A.:Not applicable.

^gHIV RNA copies/mL

* Envs isolated from the same patient are indicated by brackets.

": same value than above.

1022

1023

1024

Table 2. Molecular characteristics of HIV-1 Envs: sequence length and N potential glycosylation sites (PNGs) in the variable loops (Vn) of the gp120 subunit.

Clinical Group	Subgroup	Env code	^a V1/G	V2/G	V3/G	V4/G	V5/G	^b ΣVn/G	^c Mean/G	^d Gp160	^e Mean		
LTNP	EC	1	28/4	43/2	37/2	28/4	12/1	148/13		848			
		2	33/5	41/2	37/2	31/4	12/1	154/14		853			
		3	33/5	41/2	37/2	31/3	12/2	154/14		853			
		6	28/3	41/2	37/2	34/4	12/1	152/12		852			
		7	32/5	47/2	37/2	30/4	11/2	157/15	151.1/14.4	859	851.8		
		8	32/5	47/2	37/2	30/4	11/2	157/15		859			
		9	24/4	43/2	36/2	28/4	12/1	143/14		843			
		10	27/4	42/2	37/2	29/5	14/2	149/16		851			
		11	27/5	42/3	37/2	29/5	13/2	148/17		850			
		12	27/4	42/3	37/2	32/5	12/1	150/14		850			
		Viremic		13	31/5	41/2	37/2	31/4	13/1	153/14		854	
				14	29/4	42/2	37/2	32/5	12/2	152/15		852	
15	34/5			41/2	37/2	36/5	12/1	160/15		860			
16	29/5			41/2	37/1	29/5	12/1	148/14		849			
17	29/5			41/2	37/2	29/5	12/1	148/16	150,3/14.1	849	851.5		
18	29/4			41/2	37/2	29/5	12/1	148/15		849			
19	29/4			41/2	37/2	29/5	12/1	148/14		849			
21	24/3			41/2	37/1	30/5	10/0	142/11		842			
22	28/4			41/2	37/2	32/5	12/1	150/14		850			
24	37/7			41/2	36/2	32/5	12/1	158/15		861			
Progressor	Old	26	31/4	41/3	37/2	39/7	14/2	160/18		862			
		27	31/5	48/3	37/2	28/5	14/2	158/16		858			
		28	25/5	41/2	36/2	33/5	12/2	145/15		848			
		30	33/4	41/2	37/2	27/4	11/1	150/14		852			
		31	33/5	41/2	37/2	36/5	13/2	158/16		860			
		32	28/5	44/2	36/2	30/5	15/2	151/15	153,8/15.2	853	855.8		
		33	31/4	44/2	36/2	30/5	15/2	156/14		856			
		36	28/4	46/1	37/2	34/5	14/2	157/15		859			
		37	28/4	46/2	37/2	34/5	14/2	157/16		859			
		38	30/4	41/3	37/1	31/4	13/2	150/13		851			
Modern ^e		39	31/4	41/2	37/2	29/4	12/2	149/14		849			
		40	31/4	41/2	37	29/4	17/2	154/14		849			
		42	29/4	48/2	37	36/6	17/2	167/13		878			
		43	29/4	48/3	37	30/4	15/2	159/15		872			
		44	28/4	47/3	37	31/4	15/2	158/15	158.1/14.7	859	862.0		
		45	28/4	47/2	37	31/4	15/2	158/14		859			

46	35/4	46/3	37	33/5	13/2	164/15	865
49	37/6	41/2	37	42/7	13/1	170/18	871
50	37/6	41/2	37	42/7	13/1	170/18	871
51	31/4	42/2	37	26/3	13/1	149/12	853
52	29/4	42/2	37	32/6	12/1	152/15	856

1025

1026

1027

1028

1029

1030

1031

1032

1033

1034

1035

1036

1037

1038

1039

1040

1041

1042

1043

1044

1045

1046

^aLength in amino acid (aa) and potential glycosylation sites (PNGs) of the Env-gp120 variable regions (Vn; from V1 to V5) expressed as Vn/G ratio.

^b $\Sigma Vn/G$ indicates the sum of the aa lengths of the Vn (n; from 1 to 5) and the potential G sites.

^cMean/G indicates the mean length and PNG value for each group of Envs.

^dGp160 shows the total length in aa of each Env including the gp41 subunit and the gp120 subunit.

^eMean gp160 length in aa for each group of Envs.

Table 1. Epidemiological, clinical and host characteristics of the viral Envs.

Clinical Group	Sub-group	Env code ^a	Patient Identification code	Viral Load ^g (at sampling)	Diagnostic time	Sampling time	Viral dating ^b	HLA B		
LTNP	EC	1	2057906-3	< 50	1993	2004	1989	4901/5701		
		2	3227050	< 50	1988	2004	1991	0702/5201		
		3	3227058-3	< 50	1992	2004	1991	1402/1402		
		6	20044616-3	< 50	1998	2004	1999	1501/5703		
		* 7	10246788	< 50	1992	2005	1993	4402/5701		
		8		< 50	"	"	"	"		
		9	MDM ^c	507	1998	1996	1987	4402/3501		
		10	^c	< 50	"	2011	1996	"		
		11	^c	< 50	"	2005	"	"		
		12	^c	< 50	"	2005	"	"		
		Viremic	13	4022834	3.710	1994	2004	ND	1401/4403	
			14	9684	2.557	1998	2005	1994	1302/4001	
	15		2988465	2.286	1993	2004	1999	1402/2705		
	16		38 17 5	418	1996	2014	1999	2705/5801		
	17				"	"	"	"		
	18				"	"	"	"		
	19				"	"	"	"		
	21		30	7.597	1989	1998	2000	1501/3501		
	22		64	11.926	1989	1999	1999	4402/4901		
	24				"	"	2002	"		
	Progressor		Old	26	V10 ^d	N.D.	1993	1994	1999	4002/4402
				27				"	"	
		28		V13	N.D.	1992	1994	1990	0702/1402	
		30		L10	89.000	----	1993	1993	1501/4901	
31				"	"	"	"	"		
32		L 11		42.000	1993	1993	2000	1801/5101		
33				"	"	"	"	"		
36		I14 ^d		130.000	1987	1994	2002	0702/3502		
37		^d	"	"	"	"	"			
38		I18	170.000	1991	1994	1990	1402/4403			
Modern		39	ESI 17A	156.300	2014	2013	N.A. ^f	4201/4402		
		40		"	"	"	"	"		
		42	ESI 39A	137.700	2012	2014	N.A.	1517/3801		
		43		"	"	"	"	"		
		44	ESI 41A	129.700	2012	2014	N.A.	3503/5701		
		45		"	"	"	"	"		
	46	ESI 5A 2	49.107	2004	2007	N.A.	4102/4402			
	49	ESI 42 A	11.510	2011	2014	N.A.	1402/4403			
50		"	"	"	"	"				
51	ESI 42 B	41.090	2011	2014	N.A.	0702/1501				
52		"	"	"	"	"				

^aHIV-1 Env number used in this study and identification codes.

^bAccording to Bello et al. (2004). *J Gen Virol.* Feb;85(Pt 2):399-407. doi: 10.1099/vir.0.19365-0. PMID: 14769897.

^cDouble infected individual (Casado et al. (2007) *J Infect Dis.* 2007 Sep 15;196(6):895-9. doi: 10.1086/520885. Epub 2007 Aug 14. PMID: 17703421).

^dIndividuals with a short antiviral therapy (AZT (zidovudine) and DDI (didanosine) for V10 patient and AZT for I14 patient).

^eThe Modern Individuals have been infected within 3 years.

^fN.A.:Not applicable.

^gHIV RNA copies/mL

^{*}Envs isolated from the same patient are indicated by brackets.

[“]: same value than above.

Table 2. Molecular characteristics of HIV-1 Envs: sequence length and N potential glycosylation sites (PNGs) in the variable loops (Vn) of the gp120 subunit.

Clinical Group	Subgroup	Env code	^a V1/G	V2/G	V3/G	V4/G	V5/G	^b ΣVn/G	^c Mean/G	^d Gp160	^e Mean		
LTNP	EC	1	28/4	43/2	37/2	28/4	12/1	148/13		848			
		2	33/5	41/2	37/2	31/4	12/1	154/14		853			
		3	33/5	41/2	37/2	31/3	12/2	154/14		853			
		6	28/3	41/2	37/2	34/4	12/1	152/12		852			
		7	32/5	47/2	37/2	30/4	11/2	157/15	151.1/14.4	859	851.8		
		8	32/5	47/2	37/2	30/4	11/2	157/15		859			
		9	24/4	43/2	36/2	28/4	12/1	143/14		843			
		10	27/4	42/2	37/2	29/5	14/2	149/16		851			
		11	27/5	42/3	37/2	29/5	13/2	148/17		850			
		12	27/4	42/3	37/2	32/5	12/1	150/14		850			
		Viremic	13	31/5	41/2	37/2	31/4	13/1	153/14		854		
			14	29/4	42/2	37/2	32/5	12/2	152/15		852		
	15		34/5	41/2	37/2	36/5	12/1	160/15		860			
	16		29/5	41/2	37/1	29/5	12/1	148/14		849			
	17		29/5	41/2	37/2	29/5	12/1	148/16	150,3/14.1	849	851.5		
	18		29/4	41/2	37/2	29/5	12/1	148/15		849			
	19		29/4	41/2	37/2	29/5	12/1	148/14		849			
	21		24/3	41/2	37/1	30/5	10/0	142/11		842			
	22		28/4	41/2	37/2	32/5	12/1	150/14		850			
	24		37/7	41/2	36/2	32/5	12/1	158/15		861			
	Progressor		Old	26	31/4	41/3	37/2	39/7	14/2	160/18		862	
				27	31/5	48/3	37/2	28/5	14/2	158/16		858	
		28		25/5	41/2	36/2	33/5	12/2	145/15		848		
		30		33/4	41/2	37/2	27/4	11/1	150/14		852		
31		33/5		41/2	37/2	36/5	13/2	158/16		860			
32		28/5		44/2	36/2	30/5	15/2	151/15	153,8/15.2	853	855.8		
33		31/4		44/2	36/2	30/5	15/2	156/14		856			
36		28/4		46/1	37/2	34/5	14/2	157/15		859			
37		28/4		46/2	37/2	34/5	14/2	157/16		859			
38		30/4		41/3	37/1	31/4	13/2	150/13		851			
Modern ^e		39	31/4	41/2	37/2	29/4	12/2	149/14		849			
		40	31/4	41/2	37	29/4	17/2	154/14		849			
		42	29/4	48/2	37	36/6	17/2	167/13		878			
		43	29/4	48/3	37	30/4	15/2	159/15		872			
		44	28/4	47/3	37	31/4	15/2	158/15	158.1/14.7	859	862.0		
		45	28/4	47/2	37	31/4	15/2	158/14		859			
		46	35/4	46/3	37	33/5	13/2	164/15		865			
		49	37/6	41/2	37	42/7	13/1	170/18		871			
		50	37/6	41/2	37	42/7	13/1	170/18		871			
		51	31/4	42/2	37	26/3	13/1	149/12		853			
52	29/4	42/2	37	32/6	12/1	152/15		856					

^a Length in amino acid (aa) and potential glycosylation sites (PNGs) of the Env-gp120 variable regions (Vn; from V1 to V5) expressed as Vn/G ratio.

^b $\Sigma Vn/G$ indicates the sum of the aa lengths of the Vn (n; from 1 to 5) and the potential G sites.

^c Mean/G indicates the mean length and PNG value for each group of Envs.

^d Gp160 shows the total length in aa of each Env including the gp41 subunit and the gp120 subunit.

^e Mean gp160 length in aa for each group of Envs.







A**C****B****D**

Figure 1
 Silvia Pérez-Yanes *et al.*

A**B**



A**B**

Figure 3
 Sílvia Pérez-Yanes *et al.*



A**B**

Figure 4
Silvia Pérez-Yanes *et al.*



A**B**

Figure 5
Silvia Pérez-Yanes *et al.*




A**B**

Figure 6
Silvia Pérez-Yanes *et al.*

A




Figure 7
Silvia Pérez-Yanes *et al.*









Figure 8
Silvia Pérez-Yanes *et al.*

