Long-range single-molecule mapping of chromatin modification in eukaryotes

Zhe Weng[2,*] ,Fengying Ruan[2,*] ,Weitian Chen[1,2,*], Zhe Xie[2,3], Yeming Xie [2], Chen Zhang[2], Zhichao Chen [1,2], Juan Wang[2], Yuxin Sun[2], Yitong Fang [2], Mei Guo [2], Yiqin Tong [2], Yaning Li [2], Chong Tang[2,4,5]

[1.] College of Life Sciences, University of Chinese Academy of Sciences, Beijing 100049, China
[2.] BGI Genomics, BGI-Shenzhen, Shenzhen 518083, China
[3.] Department of Biology, Cell Biology and Physiology, University of Copenhagen 13, 2100 Copenhagen，Denmark
[4.] Nantong University, Nantong, China, 226000
[5.] Nephrosis Precision Medicine Innovation Center, University of Beihua School of Medicine, Jilin 132011, China

[*] These authors contributed equally to this work.

Keywords: histone, H3K27me3, CTCF, methylation, m[6]A, methyltransferase

Running title: long-range profiling of H3K27me3 reveal the heterogenous histone modification

Correspondence:

Chong Tang

Director of technology, BGI Shenzhen, China

Phone: 8618025420976

Email: tangchong@bgi.com

Abstract

The epigenetic modifications of histones are essential marks related to the development

and disease pathogenesis, including human cancers. Mapping histone modification has

emerged as the widely used tool for studying epigenetic regulation. However, existing

approaches limited by fragmentation and short-read sequencing cannot provide

information about the long-range chromatin states and represent the average chromatin

status in samples. We leveraged the advantage of long read sequencing to develop a

method "BIND&MODIFY" for profiling the histone modification of individual DNA fiber.

Our approach is based on the recombinant fused protein A-EcoGII, which tethers the

methyltransferase EcoGII to the protein binding sites and locally labels the neighboring

DNA regions through artificial methylations. We demonstrate that the aggregated

BIND&MODIFY signal matches the bulk-level ChIP-seq and CUT&TAG, observe the

single-molecule heterogenous histone modification status, and quantify the correlation

between distal elements. This method could be an essential tool in the future third-

generation sequencing ages.

Introduction

The genome in each cell of a multi-cellular organism is identical and stays reasonably static, while the epigenome is very dynamic [1]. The epigenome varies in different cell types and plays significant roles in various biological processes, such as stem cell differentiation [2], nervous system [3], cell aging and disease [4] etc. However, the methods to study the epigenome are vague and may be less visible than the genome, which could be directly sequenced. The solutions proposed are based on the mechanisms to extract the epigenome signal by enzymatic or chemical means and isolate either the accessible or protected location.

Therefore, current interest is placed on collecting and comparing genome-wide chromatin accessibility and chromatin modification to locate epigenetic changes that accompany cell differentiation, environmental signaling, and disease development [5]. The next-generation sequencing using DNase I (DNase-seq) and MNase (MNase-seq) treatment provided the first demonstration that active chromatin coincides with nuclease hypersensitivity [6] [7]. The comparable method is ATAC-seq by using the transposon hyperactivity on the open chromatin [8, 9]. The most recent and advanced theory proposed to offer a single molecular long-read approach for assessing the chromatin accessibility by using third-generation sequencing-nanopore sequencings, such as SMACseq [10], NOME-seq [11], and nanoNOME [12]. The chromatin accessibility evaluation may have partly addressed the epigenetic problems, but the more specific question "which protein regulates chromatin accessibility" cannot be answered. The dynamic chromatin structure is tightly regulated by post-translational histone modifications and binding transcription factors [13]. To answer the question: "is a

transcription factor bound to a piece of DNA or locate the histone modification"

scientists use "ChIP-seq" to find out if a "protein of interest" is bound to a piece of

DNA [14]. The large-scale chromatin immunoprecipitation assay (ChIP-seq) based on

antibody precipitating down the DNA pieces crosslinked with targeted proteins and

direct ultra-high-throughput DNA sequencing [15]. ChIP-seq facilitated the finding of

protein-binding motifs and allowed us to identify noncanonical protein-binding motifs

further. In the literature, these have been extensively investigated to study the biological

functions of the histone acetylation and methylations, transcription factors, etc. [16-22].

Currently, the major challenge restricting the use of ChIP-seq is the need for high

input DNA amounts. The solutions in literature have pursued a variety of methods for

the low input DNA, for example, ChIPmentation [23] CUT&RUN [24] and CUT&TAG [25]. For

the CUT&RUN protocol, MNase is fused to protein A (pA-MNase) to guide the

chromatin cleavage to antibodies bound to the protein targets at their binding sites

across the genome [25]. Conceptually similar work has also been carried out by

CUT&TAG, in which transposon was used instead of the MNase. Many of the

techniques derived from the above assignments have improved to achieve the single-

cell DNA input [25-28]. However, the complex relationship between DNA methylation,

chromatin modification, and genome structure variation is often difficult to unravel with

a single omics tool. The most solutions to the problem are to apply the bisulfite

treatment to the immunoprecipitated DNA fragments in ChIP-seq, which guaranteed

both DNA methylation information and histone information [29-31]. The above literature

review describes the previous and current attempts to address the status of the

protein-bound on DNA.

Most ChIP-seq technologies used the short-read sequencing by NGS, and
downstream analysis was based on the peak calling algorithm with statistical analysis
of populated fragments [8, 9]. As a result, researchers are now faced with whether the
individual chromatin fibers have the same long-range physical organization as they
remove linkage between distal segments. Moreover, due to the short reads
sequencing, the methylation information and structure variation was limited to the
immunoprecipitated genome region. The limitations of these short-read sequencing
technologies lie in their lack of using non-cleavage labeling methods and long-read
sequencing to preserve all necessary epigenetic information in individual chromatin.
Our new approach BIND&MODIFY, by fusing methyltransferase to bind and modify
the targeted sites in long DNA molecule. In addition, we developed the single-
molecule long-read bound protein mapping sequencing. This single-molecule method
directly assays the protein binding regions, DNA methylation, and complex structure
within a single chromatin fiber at multikilobase scales. We used BIND&MODIFY to
study histone methylation and co-methylated histone status in cancer cells.
Moreover, we enumerate the detail regulation in transposon region, bivalent regulatory
states and observe distance coordinated changes in cancer cells. BIND&MODIFY also
allows for the footprinting of other specific transcription factors. We expect future
applications of BIND&MODIFY to enable new insights into the regulator dynamic
status on the genome in various experimental systems and sequencing platforms.

Results

The experimental overview of the BIND&MODIFY method

The rationale of the BIND&MODIFY method is indirectly labeling DNA regions bound

with protein of interest by an engineered recombinant fusion protein, protein A-M.EcoGII

(pA-M.EcoGII), whose methyltransferase activity can be locally controlled. The recently

characterized adenosine methyltransferase M.EcoGII could methylate a broad range of

genome DNA in a sequence non-specific manner[32] compared to DamID[33], which

methylated only rare GATC motif in the genome (Supplemental Figure 1). Our

BIND&MODIFY approach leverages the power of an engineered recombinant protein,

pA-M.EcoGII, which first tethered the methyltransferase close to the specific antibody-

bound protein of interest, then the adenines in DNA sequences adjacent to the protein

of interest were methylated at m^6A non-specific manner upon activation (Figure 1A). As

m^6A modifications were very few in the native background of mammalian chromosome

[34,35], the artificial m^6A modification, indicating the protein binding motif, could be directly

detected by the nanopore [36,37].

Details of recombinant protein pA-M.EcoGII construction design and purification steps

can be found in the methods section. Briefly, recombinant pA-M.EcoGII is designed by

cloning two immunoglobin binding domains of staphylococcal protein A fused N-

terminally with M.EcoGII (Figure 1A). The amino acid of the linker region between

protein A and M.EcoGII is DDDKEF. The recombinant pA-M.EcoGII was expressed in

the *E.coli* system and had a molecular weight of 61 kDa (Figure 1C). To access the

function of purified recombinant pA-M.EcoGII, the m^6A dependent restriction enzyme

DpnI digestion was used to test pA-M.EcoGII's methylation activity on lambda DNAs.

We found pA-M.EcoGII had similar m^6A methylation specificity compared to

commercially available M.EcoGII (NEB) (Figure 1D). To further evaluate the

effectiveness of purified recombinant pA-M.EcoGII, m^6A sensitive restriction enzymes

(EcoRV, PciI, PvuII) were used on pTXB1 plasmid DNAs treated by pA-M.EcoGII and

M.EcoGII. Both pA-M.EcoGII and commercially available M.EcoGII successfully

introduced the m^6A methylations into the plasmid, which inhibited the digestion activity

of m^6A sensitive restriction enzymes (Figure 1E). We also carried out an ELISA assay to

test our purified recombinant pA-M. EcoGII's function of IgG domain binding and results

showed that it had similar activity to commercially available protein A (Figure 1F). Taken

together, our purified recombinant pA-M.EcoGII has comparable enzymatic activity to

commercially available protein A and M.EcoGII.

Based on the recombinant pA-M.EcoGII protein, we developed the BIND&MODIFY

protocol (Figure 1G). Briefly, the cells are 1) lightly fixed and tethered to Concanavalin A

(ConA)-coated magnetic beads, 2) incubated with primary antibodies, 3) incubated with

pA-M.EcoGII with minimal washes, 4) addition of S-adenosylmethionine at 37 degrees

to initialize the methylation reaction for 30min then quenched by 0.1% SDS, 5)

extraction of DNA by phenol-chloroform and prepare the library for ONT nanopore

sequencing. After sequencing, the data was processed as genome alignment and m^6A

signal detection. The m^6A signal cut-off was determined based on background noise

(Supplemental Figure 2). The sensitivity and specificity of m^6A signal are around 0.92

and 0.99.

The validation of BIND&MODIFY on assigned location *in vitro*

Similar to previous studies, the occurrence frequency of adenosine on various

genomes is once in every 3bp [10], which was the theoretical resolution of M.EcoGII

treated areas. The resolution of BIND&MODIFY depends not only on the adenosine

frequency on the genome but also on the distance the tethered M.EcoGII could reach.

Here we present the test that evaluates the signal resolution of BIND&MODIFY. The

first step to developing the method is to synthesize the DNA with an assigned antibody

binding site. Then, the PCR was used to introduce the m^5C modification into the given

location on DNAs (Figure 2A). In the following steps, the pA-M.EcoGII recombinant

protein was tethered to the designed site with the help of the m^5C antibody. Then the

attached M.EcoGII methyltransferase transferred methyl group to neighboring

adenosine after activation. The N6-methyladenosine (m^6A) modified DNAs were

sequenced in Nanopore, and the data was analyzed in our house pipeline.

We calculated the m^6A sites, non-m^6A sites, and m^6A ratio within 50bp sliding windows

by using all the reads covered. In Figure 2B, the m^6A ratios were plotted with their

genomic location as an indicator of bound pA-M.EcoGII. The regions with high

methylation ratio (>0.53) correspond to the neighboring areas of the bound pA-

M.EcoGII, where the tethered M.EcoGII could methylate the genome sites at a close

distance (Figure 2B). The resolution of detecting pA-M.EcoGII binding site was around

100bp (Figure 2B), comparable to the conventional ChIP-seq peak size distribution of

100-300bp (Supplemental Figure 3).

Further investigation can be undertaken to explore the single molecular level

sensitivity and specificity of BIND&MODIFY. We calculated the methylation ratio within

50bp sliding windows on each detected DNA molecule. Then, we used the receiver

operating characteristic (ROC) curve analysis to determine the ability of the sectional

methylation ratio to classify or predict the pA-M.EcoGII binding location on the single

molecular level. When the methylation ratio cut-off is increased to 0.53, the sensitivity

decreases to 92 %, and the specificity increases to 79 % (Figure 2B). Therefore, the

BIND&MODIFY method has the potential ability to detect pA-M.EcoGII binding sites at

the single molecular level.

The secondary objective is to investigate the strand specificity of the pA-M.EcoGII

modifying. We plotted the methylation ratio, as the indicator of the pA-M.EcoGII

location, on both positive and negative strands (Figure 2C). We only found the pA-

M.EcoGII binding signal on the negative strand other than the positive strand. This

suggested that the tethered M.EcoGII might be spatial close to the negative strand of

DNA and preferentially modified the negative strand. These features make the

BIND&MODIFY method optimal for long-range histone modification studies on the

single molecular level and histone modification in a strand-specific manner.

BIND&MODIFY revealed the comparable strand-specific view of the epigenomic

regulator on genomic DNA

The ChIP-seq, CUT&RUN, and CUT&TAG detect protein–DNA binding events and

chemical modifications of histone proteins [38-40]. Several histone trimethylation states,

such as H3K4me3, are well-studied linked to active gene transcription, whereas the

relationship between H3K27me3 and transcriptional status appears less unambiguous

[41, 42]. Therefore, the H3K27me3 status in breast cell line MCF-7 is being explored by

ChIP-seq and BIND&MODIFY in our research. The average read length of Nanopore

is around 2kb (Supplemental Figure 4). The correlation of the experimental replicates

is 0.90 (Supplemental Figure 5).

To validate the H3K27me3 BIND&MODIFY signal, we first evaluate the consistency

between BIND&MODIFY and the conventional methods. Both the conventional

method (ChIP-seq) and the proposed BIND&MODIFY method illustrated the similar

H3K27me3 position signals in various genome scales (Figure 3A, Supplemental

Figure 6). Over 95% of high confidence position signals in ChIP-seq were also

observed BIND&MODIFY (Supplemental Figure 7). By analyzing the regional signal

strength, the BIND&MODIFY signal strength of m^6A counts was strongly correlated

with ChIP-seq peak signal intensity (Figure 3B). Taken together, the BIND&MODIFY

results exhibited a range of values comparable with conventional ChIP-seq.

Some studies observed an inverse relationship between H3K27me3 density in

transcription start sites and gene expression [41, 42]. To profile the enrichment of

H3K27me3 within genic regions, enrichment data were visualized using transcription

start site (TSS)-centered plots. Overall, the BIND&MODIFY results show a level of

agreement with ChIP-seq, that the H3K27me3 signal reaches the highest around

TSSs (Figure 3C). In the TSSs of low expression genes (bottom expression quantile),

H3K27me3 was prominently enriched around the TSS with a distinctive depletion

directly over the TSSs (Figure 3D). The bimodal pattern is relatively consistent in both

ChIP-seq and BIND&MODIFY results. Good agreement was also found when

comparing results from this work against published data [43] [44]. Further analysis was

done to see whether the histone trimethylation has strand specificity with the

BIND&MODIFY method. H3K27me3 was prominently enriched downstream of TSS in

the positive strand and upstream of TSS in the negative strand, contributing to form

the bimodal pattern around TSSs (Figure 3E). An example of the gene is illustrated in

Supplemental Figure 8. This is an exciting finding, and it could be hypothesized that

these two trimethylations spatially approach one of the DNA strands. In contrast, we

did not observe the significant strand-specific pattern around the TSSs of high

expression genes (Supplemental Figure 8).

As a critical regulator of genome organization, the CCCTC-binding factor (CTCF) has

been characterized as a DNA-binding protein with essential functions in maintaining

the topological structure of chromatin and inducing gene expression [45] (Supplemental

Figure 9A). In CUT&TAG data, we identified the strong CTCF binding sites 100bp

ahead of the TSSs of actively expressed genes (Supplemental Figure 9B). In contrast,

the CTCF binding signal around TSSs of inactive genes is distributed evenly

(Supplemental Figure 9C). Compared with CUT&TAG, BIND&MODIFY showed a

similar CTCF signal pattern around TSS in active and inactive genes. Further analysis

was done to see whether the antibody-bound CTCF domains spatially closed to either

one strand (Supplemental Figure 9D). We found that the CTCF signal on the negative

strand was sharper and more potent than on the positive strand and even more similar

to the CUT&TAG signal pattern, suggesting the antibody-bound domain is spatially

close to the negative strand. In the protein crystal structure analysis [46], the antibody-

bound domain is the C terminal of the CTCF and spatially closer to the negative

strand, further supporting this hypothesis (Supplemental Figure 10).

In conclusion, the BIND&MODIFY method has shown the comparable ability to

mapping the histone modification with conventional methods and highlights the

challenge of strand-specific view of epigenomic regulators on genome DNA

The long-range sequencing resolves the retrotransposon regions with high resolution

The retrotransposon composed of repeated sequence can be integrated elsewhere in

a genome and has various critical biological functions. The three major

retrotransposon orders are long terminal repeat (LTR) retrotransposons, long

interspersed elements (LINEs), and short interspersed elements (SINEs). Due to the

repeated sequence of retrotransposon, the short-read sequencing data showed the

40~60% multiple mapping rate in these complex regions (Supplemental Figure 11).

The multiple mapping in retrotransposons would cause signal noise or signal loss in

the ChIP-seq (Figure 4A). Compared with ChIP-seq, the BIND&MODIFY significantly

improves the unique mapping rate (Supplemental Figure 11), by taking advantage of

long-range sequencing, which generates read length that may span the entire length

of full transposon and avoid multiple mapping issues. With the advantages, the long-

range sequencing was also used to identify the novel transposon deletion and

insertions [47]. There were minimal studies about the H3K27me3 on retrotransposons [48].

We visualized the H3K27me3 status on LTR, SINE, and LINE using retrotransposon-

centered plots, including upstream/downstream 300bp of similar size retrotransposons

(Supplemental Figure 12). In the ChIP-seq, we observed the very high background

noise without any peak signal on LTR regions (Figure 4B). In contrast, the

BIND&MODIFY showed the H3K27me3 binding signal on the LTR region, with a 10X

better signal resolution (Figure 4B). In another type of retrotransposon SINE, the

ChIP-Seq loses the H3K27me3 signal in SINEs. However, the BIND&MODIFY

demonstrated the two clear peak signals in the SINEs (Figure 4C-D). Thus, the results

agree well with existing studies about H3K27me3 activity on SINEs [48].

The short interspersed nuclear element (SINE) B1 has insulator activity mediated by

the binding of specific transcription factors along with the insulator-associated protein

CTCF [49]. A genome-wide analysis of CTCF binding sites in the human and mouse

genomes discovered that many CTCF binding sites are derived from TE sequences [50].

Agreeing with previous reports, the BIND&MODIFY showed the clear CTCF signal in

the SINEs, which was lost in the ChIP-Seq (Supplemental Figure 13). In summary,

BIND&MODIFY takes advantage of the long-range sequencing to precisely map

histone modification and protein interaction in the challenging complex genome

regions.

The histone modification status on single-molecular resolution

The conventional CUT&TAG and ChIP-seq are based on statically calling the

peak of the enriched read in a specific region [51]. Recent single-molecule and

single-cell measurements of histone accessibility suggest that ATAC-seq on

cell populations represent an ensemble average of distinct nucleosome states

[52]. An essential attribute of the BIND&MODIFY measures the histone

modification in single-molecular resolution by taking advantage of the slight

variance (Supplemental Figure 14) and increased cumulative possibility in

segments (Supplemental Figure 15).

We then asked whether BIND&MODIFY could reveal the different H3K27me3

status by investigating the chr20:52223000-52225500 loci. The

chr20:52223000-52225500 loci modulated the lncRNA LOC105372672 and

Zinc finger protein 217 (ZNF217), shown to be a prognostic biomarker and

therapeutic target during breast cancer progression [53, 54]. The conventional

ChIP-Seq enriched the H3K27me3 bound motif by antibody-guided

amplification, which could be overrepresented (Figure 5A). The baseline signal

of this region is 6, which is thought to be the background noise in the

experiments (Figure 5A). In contrast, the BIND&MODIFY presents a striking

comprehensive picture of the H3K27me3 in this area. The super-resolution of

BIND&MODIFY uncovered remarkable three different epigenetic states (Figure

5B): repressed state with most histones methylated, which inhibits the gene

expression embodied in the genome region; the poised state with half histones

methylated, which transit from the inactive state to the active state; the active

state largely devoid of histone trimethylation. Some of the H3K27me3 only can

be observed in the subpopulation of the genome fibers, suggesting the highly

heterogenous histone methylation in cancer cells. The signal peaks (chr20:

52225000-52225500) in ChIP-seq sum up the H3K27me3 loci in two different states. The subpopulation of inactive fully methylated chromatin leads to the baseline signal in ChIP-seq, which was thought to be the background noise in the experiment. Our findings, three epigenetic states, align with the results reported by other publications in literature [55-58]. Many genome regions in cancer cells harbor a distinctive histone modification signature that combines the activating histone H3 Lys 4 trimethylation (H3K4me3) mark and the repressive H3K27me3 mark. The poised states with these bivalent domains, which are considered to poise expression of developmental genes, allowing timely activation (activate form) while maintaining repression (repressed state) in the absence of differentiation signals [55]. In contrast to the heterogeneous behaviors seen above, the gene desert areas without gene transcription activity have the only one repressed state with most histone trimethylated (Supplemental Figure 16). The previous bivalent model was drawn by finding the overlapped H3K4me3 and H3K27m3 average peaks in NGS but not observing the poised status. These findings reinforce the general belief that bivalent histone modification regulates gene expression.

However, the single molecular statuses in each gene are challenging to be quantified. Therefore, we could not get the big-picture view of global genes, summarizing a more comprehensive perspective with specific examples and details. Consequently, we used the mean methylation ratio of one molecule to present the methylation status of this molecule (Supplemental Figure 17), and the heterogeneity of the genes could be visualized by a series of mean

molecular methylations (Figure 5B, gradient line on the right panel). We

summarized the H3K27me3 heterogeneity of genes in Chr20 (Figure 5C). 95%

of genes display homogenous H3K27me3 regulation status with minimal

heterogeneity among molecules. Only one cluster with 31 genes (Figure 5C,

cluster 3) demonstrated the very high heterogeneity of H3K27me3 regulation

with active state, repressed state, and poised state. These genes are under

typical bivalent mode regulation of H3K27me3. By the gene ontology analysis,

we found that these genes were enriched in the breast cancer-related

pathways, for example, VEGF singling pathway[59], B cell receptor signaling

pathway[60], PD-L1 checkpoint pathway[61] ,etc. (Supplemental Figure 18). The

bivalent mode regulation in these pathways might enlighten the mechanism of

tumor heterogeneity, evolution, drug resistance, immune evasion, and the

cause of metastasis.

The BIND&MODIFY reveal the long-distance correlation of regulators

In the transition from the chromatin repressed state to the activated state, the

H3K27me3 changed in rhythmics. The region (chr20: 52223500-52224200)

firstly devoid of H3K27me3 and then other areas together erase the histone

trimethylation in the next step, suggesting the possibility of a synergic

regulatory switch at this location. To quantify distance (anti)correlation

between chromatin trimethylation states, we developed a modified correlation

coefficient (CC) metric for assessing the degree of trimethylation correlation

between genomic regions. Average CC profiles centered on positioned

nucleosomes reveal the detectable correlation between nucleosome positions

up to 100bp (Supplemental Figure 19). These observations are consistent with

nucleosomes imposing restrictions on one another, resulting in a short-range

correlation between nucleosome footprints that dephases over longer ranges

[10]. Furthermore, CC analysis of DNA confirmed the long-range positive

correlation between the promoter region and this upstream, downstream

element (Figure 6A, Supplemental Figure 20).

However, the CC index is only suitable to describe the distance correlation of

H3K27me3 for single genes. To further explore the biological meaning of

distance correlation, we developed the new DE index (Distance Effect index) to

quantify the distal effects of each genomic locus (Supplemental Figure 21).

The higher DE index means these genomic loci have a stronger distal

correlation, suggesting the regulation on these sites could impact a larger

area. Therefore, the genetic region could be presented as the serial of DE

index on the corresponding locus. We plotted the DE index of all the

transcription regions (2kb upstream of TSS) on chr20 (Figure 6B). There are

several DE patterns of H3K27me3 in the transcription region. Some

transcription initiation sites have a powerful DE index signal, which suggested

the rhythmic change in the more extensive area and the more substantial

regulator impact on the distal genomic region (Figure 6B). Consider that

H3K27me3 is associated with transcriptional repression [62], the strong DE index

signal of H3K27me3 was related to lower gene expression (Figure 6B). In the

gene ontology analysis, the dysregulation of these genes was associated with

cancer pathways, such as ubiquitin dysregulation, Apelin signaling,

proteasome, Oxytocin signaling, etc (Supplemental Figure 22).

We further plotted the distance correlation of transcription regions with CTCF

regulators. For the areas where the transcription initiation sites have strong DE

index signal (Figure 6C, Cluster 3), the CTCF on sites could synergically affect

the distal genome sites. Comparing with the gene expression profile, some

gene expression is strongly enhanced. By gene ontology analysis, these

enhanced distance regulations were related to the cancer-promoted pathways

such EGFR tyrosine kinase inhibitor resistance, GABAergic synapse, Hippo

signaling pathway, etc. (Supplemental Figure 23). In addition, the enhanced

distance regulation may enlighten the super-enhancer/suppressor effects,

which affect the gene region up to 8kb [1].

Overall, the BIND&MODIFY demonstrated the highly heterogenous status of

histone modification in the cancer cells and the long-range interaction of the

histone modification.

Discussion

ChIP-seq and CUT&TAG have been essential epigenetic study tools in literature for

many years, for example, histone modification, transcription factor binding, etc.

However, these methods suffer the limitations that the detected signals are represented

by the immunoprecipitation enriched DNA fragments in common without considering the

heterogeneity in DNA molecules. The short-read sequence also prevented the

possibility of studying the long-range interactions in multi-omics epigenome information.

We designed a new method, "BIND&MODIFY," by using the protein A-M.EcoGII

methyltransferase recombinant proteins to non-fragmented labeling the local DNA

regions. By this method, we simultaneously profiled the multi epigenome information on

a truly unbiased genome-wide scale, measured the underlying distribution of the histone

modification on single molecular resolution, and identified loci exhibiting significant

correlation.

The BIND&MODIFY generates the general trend of H3K27me3 loci similar to the widely

used ChIP-seq. Despite the significant changes in common, the relatively small different

signals were observed in terms of the variable peak signal strength. We postulate that

these are likely caused by the different fundamental principles between BIND&MODIFY

and conventional ChIP-seq. The BIND&MODIFY enables unbiased profiling of the DNA

molecule without immunoprecipitation enrichment and reflects the true nature of the

epigenome. Compared with that, the ChIP-seq amplifies the loci signals common in

most DNA molecules by immunoprecipitation enrichment and neglects the loci signal in

individual DNA molecules.

Moreover, the different measuring units in signal strength also led to the difference. One

drawback of the BIND&MODIFY is that it is unavoidable to sequence the non-signal

regions, requiring a higher sequencing depth. Fortunately, the nanopore throughput is

increasing rapidly, while selective enrichment methods are also becoming available.

Increase the read length are also helpful, especially for assaying the distance

correlation of distal regulatory elements.

We then compared our method with the alternative method CUT&TAG. The significant

difference is that the transposon in CUT&TAG is not reusable once inserted in the

DNAs. In contrast, the methyltransferase could methylate the multiple regional

adenosines, enhancing the signal strength. The CUT&TAG needed the secondary

antibody to tether more transposons to the location and increase the chance of DNA

insertion. In BIND&MODIFY, we found that the secondary antibody is not necessary.

Some may question this, and others suggest that the reusable methyltransferase may

increase the false-positive signals in non-target regions. *In vitro* evaluation, we found

the only scattered signal in the non-targeted area, but not clustered signals, which could

be obviously identified loci signal (Figure 2). *In vivo* evaluation, the genome DNA is not

movable by fixation, and the DNA methyltransferase can only activate locally (Figure 3).

Therefore, the false-positive signal of *in vitro* and *in vivo* can be safely neglected.

Base-calling is another area of future improvement. We used the Pacbio sequencing

data (specificity 0.99) to training our m^{6}A calling algorithm with nanopore sequencing.

Both sensitivity and specificity in nanopore m^{6}A detection are satisfactory. We are also

trying to call the native 5mC and artificially labeled m^{6}A simultaneously and analyze the

relationship between histone modification and DNA methylation. To our surprise, the

artificially labeled m^{6}A did not affect the 5mC detection efficiency significantly, and the

5mC correlation with bisulfite sequencing is 0.8 (Supplemental Figure 24). The

relationship between 5mC and histone modification in single molecules should be

determined in much deeper sequencing. Alternatively, we also could use the Pacbio,

which reads one nucleotide each time without the neighboring nucleotide signal

interference. Overall, the simultaneous detection of 5mC and regulator position in the

single-molecule is applicable with BIND&MODIFY.

The endogenous methylation in mammalian genomes also represents the potential

confounding factor in our analysis. By examining the data of non-treated original

genomic DNA, we found that the quantity of m^6A in original DNA is thousands of times

fewer than in the treated samples, suggesting the minimal effect of the endogenous

methylation. To determine the thresholds for the possibility of real m^6A and background

noise, the non-treated sample data was used to determine and filter out the background

noise signal distribution is below 0.53, which could significantly improve the m^6A calling

specificity and sensitivity.

However, there are also some species where m^6A occurs endogenously and strongly

correlated with the binding motif of the histone modification. Modifications such as 5mC,

4mC, cytidine deamination, or 5-glucomethyaltion are among the potential future

alternatives in such cases. Also, some artificial SAM could introduce the biotin to the

specific sites may strongly avoid the endogenously confounding signals. Finally, we

believe that the integration of BIND&MODIFY into a single-molecule multi-omics assay

represents a fruitful direction. We could simultaneously label the two or more proteins

located in the single DNA molecule with the different modification labels. Then the

protein interaction distance on the genome would be measurable in this method, which

could be the breakthrough in the protein-protein interaction in genome scales. In

principle, similar approaches may also apply to the individual RNA molecules to study

the RNA binding proteins. We expect this technology to prove the essential new class of

tools for the simultaneous study of multi-epigenomics.

Methods

Cell culture and antibodies

Human mammary gland carcinoma cell line MCF-7 were obtained from ATCC. MCF-7

were grown in DMEM (Gibco,11995065) supplemented with 10% FBS (Gibco,10099141),

0.01mg/ml insulin (HY-P1156, MedChemExpress), and 1% penicillin-streptomysin (Gibco,

15140122). Cell line was regularly checked for mycoplasma infection (Yeasen,

40612ES25). We used the following antibodies: H3K27me3 (Cell Signaling Technology,

9733), Guinea Pig anti-Rabbit IgG (Heavy & Light Chain) antibody, Anti-CTCF Antibody

(sigma,07-729-25UL), Protein A Antibody, pAb, Chicken (A00729-40, GenScript).

Recombinant protein preparation of protein A-M.EcoGII

pTXB1 vector (NEB, N6707S) was used as the protein expression backbone.

Downstream of the lac operator, a ribosome binding site and three FLAG epitope tags

were introduced, followed by two IgG-binding domains of staphylococcal protein A

encoding sequence, which was synthesized based on the previous work (Addgene,

124601). The M.EcoGII encoding sequence (Addgene, 122082) was also synthesized

based on its original discovery. The amino acid linker sequence between the C-terminus

of protein A and N-terminus of M.EcoGII is DDDKEF. The sequenced plasmid was

transformed into C3013 competent cells (NEB) following the manufacturer's protocol.

Each colony tested was inoculated into a 1 mL LB medium, and growth was continued at

37 °C for 2 h. That culture was used to start a 50 mL culture in 100 µg/mL carbenicillin

containing LB medium and incubated on a shaker until the cell density reached an A600

of 0.6, whereupon it was chilled on ice for 30 min. Fresh IPTG was added to 0.25 mM to

induce expression. Then the culture was incubated at 27 °C on a shaker for 16 h. The

culture was then collected by centrifugation at 10,000 rpm, 4 °C for 30 min, the

supernatant was discarded. The bacterial pellet was frozen in a dry ice-ethanol bath and

stored at −70 °C. The frozen pellet was resuspended in 20 mL chilled HEGX Buffer (20

mM HEPES-KOH at pH 7.2, 0.8 M NaCl, 1 mM EDTA, 10% glycerol, 0.2% Triton X-100)

including 1× Roche Complete EDTA-free protease inhibitor tablets and lysed using a high-

pressure cell disrupter (JNBIO, China). Cell debris was removed by centrifugation at

10000 rpm for 30 min at 4 °C, and the supernatant was loaded onto a column equipped

with chitin slurry resin (NEB, S6651S), then incubated the column on a rotator at 4 °C

overnight. The unbound soluble fraction was drained, and the columns were rinsed with

20 mL HEGX containing Roche Complete EDTA-free protease inhibitor tablets. The chitin

slurry was transferred to a 15 mL conical tube and resuspended in Elution buffer (10 mL

HEGX with 100 mM DTT). The tube was placed on the rotator at 4 °C for ~48 h. The

eluate was collected and concentrated using an Amicon Ultra-4 Centrifugal Filter Units

10 K (Millipore, UFC801096), and sterile glycerol was added to make a final 50% glycerol

stock of the purified protein. The fusion protein has stored the protein at -80°C.

Size characterization of recombinant protein pA-M.EcoGII

The size of purified pA-M.EcoGII recombinant protein was characterized by C Coomassie blue staining by resolving the protein in 7.5% SDS-PAGE gel.

ELISA of recombinant protein pA-M.EcoGII

To verify the binding efficiency of pA-M.EcoGII, we performed an ELISA assay in vitro. The purified recombinant protein or commercial M.EcoGII (NEB, M0603S) was diluted with coating buffer (0.05 M NaHCO$_3$ buffer, pH 9.2). The 96 well high binding plate (Greiner Bio-one, 655061) was coated with 100 µL pA-M.EcoGII or commercial M.EcoGII (NEB,M0603S) (1:120, 1:480 of stock 10mg/ml), negative control, commercial ProteinA (ThermoFisher, 21181) (1:120, 1:480 of stock 10mg/ml) in coating buffer per well for 4 h at room temperature. Further, 200 µL SuperBlock™ (TBS.) Blocking Buffer – Blotting (Invitrogen,37537) was added to each well and incubated for 2 h at room temperature. Then each well was washed 5X with 380 µL washing buffer (0.14 M NaCl; 0.01 M PO$_4$;0.05% Tween 20; pH 7.4). After that 95 µL Secondary Antibody (1:10000 in PBSB) was added to each well and incubated for 1 h at room temperature. The plate was washed 4X with 380 µL washing buffer (0.14 M NaCl; 0.01 M PO$_4$;0.05% Tween 20; pH 7.4). 90 µL TMB substrate solution was added and incubated for 15min at room temperature in the dark. Finally, 90 µL stop buffer (1.8N H$_2$SO$_4$) was added to stop the color development and read immediately at 450nm (yellow color) using FLUOstar® Omega Plate Reader by BMG LABTECH.

In vitro methtransferase activity of pA-M.EcoGII recombinant protein

To access the in vitro methylation efficiency of pA-M.EcoGII recombinant protein, m^6A

methylation-sensitive restriction enzyme DpnI was used to probe the adenine methylation

at GATC motif of 7 kb unmethylated dsDNA. The 7 kb dsDNA substrate was PCR

amplified from lambda DNA. For the methyltransferase reactions, each 50 µl reaction

volume was assembled on ice and contained 1µg 7kb unmethylated dsDNA, 1X Cutsmart

buffer, 640 µM SAM, and 4 µL of pA-M.EcoGII recombinant protein or commercial M.

EcoGII, then the mixture was incubated at 37 °C for 1 h. The methylated product was

purified using 0.6X Ampure XP (BECKMAN COULTER, A63882). 1 µl of DpnI (NEB) was

added to the reaction mixture to further incubate at 37 degrees for 10min. DpnI cutting

efficiency was examined by 1% agarose gel electrophoresis.

To assess the specificity of pA-M.EcoGII recombinant protein methylation and its

effectiveness at inhibiting restriction endonucleases, we carried out restriction analyses

using an unmethylated 7kb linear dsDNA template, which was PCR amplified from pTXB1.

One enzyme known to be insensitive to dA methylation (BamHI) and three enzymes

(EcoRV, PciI, PvuII) that cleave different base-pair sequences, the activities of which are

known to be blocked by adenine methylation. The commercial M.EcoGII and pA-

M.EcoGII recombinant protein methyltransferase reactions on the 7kb linear dsDNA were

carried out the same as described above. All restriction endonucleases used in this study

were purchased from NEB. For the restriction enzyme digest reaction, each 30 µL

reaction volume contained 500 ng methylated dsDNA, the appropriate digestion buffer,

time, and amount of enzyme following the manufacturer's protocol of NEB. The enzymes

were inactivated at 80 °C for 20 minutes. All sample was loaded to 1% agarose gel for

analysis.

In vitro pA-Tn5 transposome preparation

The pA-Tn5 was purchased from Vazyme (Vazyme, S603). To generate the pA-Tn5

adapter transposon, 7 µL of a 50 µM equimolar mixture of pre-annealed Tn5MEDS-A and

Tn5MEDS-B oligonucleotides, 40 µL of 7.5 µM pA-Tn5 fusion protein, and 28 µL coupling

buffer were mixed. The mixture was incubated for 1 h on a Thermocycler at room

temperature and then stored at −20 °C.

CUT&TAG for bench-top application

Gently resuspend and withdraw enough of the slurry such that there will be 10 µL for each

final sample. Place the tube on a magnet stand to clear (30 s to 2 min). Withdraw the

liquid, and remove it from the magnet stand. Add 1.5 mL Binding buffer (20 mM HEPES

pH 7.5, 10 mM KCl, 1 mM CaCl$_2$, and 1 mM MnCl$_2$), mix by inversion or gentle pipetting,

remove liquid from the cap and side with a quick pulse on a microcentrifuge. Resuspend

in a volume of Binding buffer equal to the volume of bead slurry (10 µL per final sample).

10 µL of activated beads were added per sample and incubated at room temperature for

15 min. Cells were harvested, counted and centrifuged for 3 min at 600×g at room

temperature. 500000 cells were washed 2X in 1.5 mL Wash Buffer (20mM HEPES pH

7.5, 150mM NaCl, 0.5mM Spermidine, 1× Protease inhibitor cocktail), after that the cells

were resuspended in 1.5 mL Wash Buffer by gentle pipetting in a 2mL tube. The unbound

supernatant was removed by placing the tube on the magnet stand to clear and pulling

off all of the liquid. The bead-bound cells were resuspended with 50 µL Dig-Wash Buffer

(20mM HEPES pH 7.5, 150mM NaCl, 0.5mM Spermidine, 1× Protease inhibitor cocktail,

0.05% Digitonin) containing 2mM EDTA and a 1:100 dilution of the appropriate primary

antibody. The primary antibody was incubated on a rotating platform overnight at 4 °C.

The primary antibody was removed by placing the tube on the magnet stand to clear and

pulling off all of the liquid. The secondary antibody was diluted 1:100 in 100 µL of Dig-

Wash buffer, and cells were incubated at room temperature for 1h. Cells were washed

using the magnet stand twice for 5 min in 1 mL Dig-Wash buffer to remove unbound

antibodies.0.04 µM of pA-tn5 was prepared in 150 µL Dig-Med Buffer (0.05% Digitonin,

20 mM HEPES, pH 7.5, 300 mM NaCl, 0.5 mM Spermidine,1× Protease inhibitor cocktail).

After removing the liquid on the magnet stand, 150 µL was added to the cells with gentle

vortexing, which was incubated with pA-Tn5 at room temperature for 1 h. Cells were

washed twice for 5 min in 1 mL Dig-Med Buffer to remove unbound pA-Tn5 protein. Next,

cells were resuspended in 300 µL Tagmentation buffer (10 mM $MgCl_2$ in Dig-Med Buffer)

and incubated at 37 °C for 1 h. To stop tagmentation, 15 µL EDTA,3 µL 10% SDS, 2.5 µL

20mg/ml proteinase K was added to 300 µL of the sample, which was incubated in a

water bath overnight at 55 °C. 320 µL PCI was added to the tube and mixed by full-speed

vortexing for 2 s. The upper phase was transferred to a phase-lock tube. 320 µL

Chloroform was added to the tube and inverted ~10x to mix. The tube was Centrifuged

for 3 min at room temperature at 16,000 x g. The aqueous layer was transferred to a fresh

1.5 mL tube containing 750 µL 100% ethanol and mixed by pipetting. The tube was

Chilled on ice and centrifuged for at least 10 min at 4 °C 16,000 x g. The liquid was

removed, and 1 mL 100% ethanol was added to the tube, then centrifuged 1 min at 4 °C

16,000 x g. The liquid was carefully poured off and air dry. When the tube is dry, 25 μL

10 mM Tris-HCl pH 8, 0.1 mM EDTA was added to the tube and vortex on full of dissolving

the genomics DNA

BIND&MODIFY for bench-top application

500,000 cells were used in each BIND&MODIFY assay. Cells were harvested, counted,

and centrifuged for 3 min at 600×g at room temperature. Cells were first lightly fixed by

adding formaldehyde (ThermoFisher, 28906) to a final concentration of 0.1% in 1.5ml

PBS, and incubated at room temperature for 15min. 2.5M Glycine was added to final

concentration of 0.125 M to quench the additional formaldehyde. Fixed cells were then

washed twice in 1.5 mL Wash Buffer (20mM HEPES pH 7.5, 150mM NaCl, 0.5mM

Spermidine, 1× Protease inhibitor cocktail) by gentle pipetting. 10 μL of activated beads

were added per sample and incubated at room temperature for 15 min. The unbound

supernatant was removed, bead-bound cells were resuspended in 100 μL Dig-Wash

Buffer (20 mM HEPES pH 7.5, 150 mM NaCl, 0.5 mM Spermidine, 1× Protease inhibitor

cocktail, 0.05% Digitonin) containing 2 mM EDTA and a 1:100 dilution of the appropriate

primary antibody. Primary antibody incubation was performed on a rotating platform

overnight at 4 °C. The primary antibody was removed by placing the tube on the magnet

stand to clear and pulling off all of the liquid. No secondary antibody was used for

BIND&MODIFY. Cells were washed twice using the magnet stand for 5 min in 1 mL Dig-

Wash buffer. 4 μL of pA-M.EcoGII was prepared in 150 μL Dig-Wash Buffer. After

removing the liquid on the magnet stand, 150 μL of pA-M.EcoGII containing Dig-Wash

buffer was added to the cells with gentle vortexing, which was incubated at room

temperature for 1 h. Cells were washed 2× for 5 min in 1 mL Dig-Wash Buffer and 1x for

5 min in 1mL Low salt buffer (20 mM HEPES pH 7.5, 0.5 mM Spermidine, 1× Protease

inhibitor cocktail, 0.05% Digitonin) to remove unbound pA-M.EcoGII protein. Next, cells

were resuspended in 300 μL Reaction buffer (20 mM HEPES pH 7.9, 0.5 mM Spermidine,

1× Protease inhibitor cocktail, 0.05% Digitonin, 10 mM 1M $Mgcl_2$, 300 mM sucrose) .The

reaction was activated by adding 5 μL SAM of 32 mM at 37°C in a thermmixer. Additional

5 μL SAM of 32 mM was added to the tube at 7.5 min and 15 min. The reaction was

stopped at 30 minutes by placing the tube on the magnet stand to clear and pulling off all

of the liquid. The bead-bound cells was resuspended with 300 μL Digestion Buffer(20 mM

HEPES pH 7.5, 300 mM NaCl, 0.5 mM Spermidine, 1× Protease inhibitor cocktail, 0.05%

Digitonin, 16.7 mM EDTA, 0.1% SDS, 0.167 mg/mL proteinase K) and incubated in water

bath overnight at 55 °C. The method for genomic DNA extraction was the same as

CUT&TAG for bench-top application.

CUT&TAG library preparation

To amplify libraries, 21 μL of CUT&TAG genomic DNA was mixed with 2 μL of a universal

i5 and a uniquely barcoded i7 primer, using a different barcode for each sample. A volume

of 25 μL KAPA HIFI ready mix (KAPA, KK2602) was added and mixed. The sample was

placed in a Thermocycler with a heated lid using the following cycling conditions: 72 °C

for 5 min; 98 °C for 30 s; 14 cycles of 98 °C for 10 s, 63 °C for 30 s and 72 °C for 15s;

final extension at 72 °C for 1 min and hold at 8 °C. Post-PCR clean-up was performed by

adding 1.3× volume of Ampure XP beads (Beckman Counter), and libraries were

incubated with beads for 5 min at RT, washed twice gently in 80% ethanol, and eluted in

30 µL 10 mM Tris pH 8.0. the library was analyzed using Agilent 2100 (2100 Bioanalyzer

Instrument, G2939BA). Then the library was sequenced in MGI2000 platform with

PE100+100+10 sequencing.

BIND&MODIFY library preparation

The BIND&MODIFY library was prepared following the manufacturer's protocol of SQK-

LSK109 (Nanopore, SQK-LSK109). The library was sequenced in the ONT PromethION

platform with R9.4.1 flow cell.

Basecalling and DNA methylation calling

Reads from the ONT data were performed using megalodon (V2.2.9), which used

Guppy basecaller to basecalling and Guppy model config

res_dna_r941_min_modbases-all-context_v001.cfg has been released into the Rerio

repository was used to identify DNA m⁶A methylation. megalodon_extras was used to

get per read modified_bases from the megalodon basecalls and mappings results. In

order to futher explore the accurate threshold of methylation probability, a control

sample with almost no m⁶A methylation was used as background noise, and Gaussian

mixture model was used to fit the methylation probability distribution generated by

megalodon.

Accessibility score

Hg19 genome and the gene elements were processed into 50bp bin sliding 5bp by

Bedtools (v2.27.1)[63]. The accessibility score over multi base-pair windows were

calculated as methylation ratio = m^6A bases in all covered reads under bin/ adenosine

bases in all covered reads under bin. And the accessibility score of each single

molecule in the bin was also calculated.

ChIP-seq data processing

Demultipexed fastq files were mapped to the hg19 genome using Bowtie2(2.4.1) [64]with

the following settings: bowtie2 --end-to-end --very-sensitive --no-mixed --no-discordant -

-phred33 -I 10 -X 700. peaks were called using MACS2 (v.2.1.0) [65] with the following

settings: -g 12000000-f BAMPE.

CUT&TAG data processing

Demultipexed fastq files were mapped to the hg19 genome using Bowtie2 with the

following settings: bowtie2 --end-to-end --very-sensitive --no-mixed --no-discordant --

phred33 -I 10 -X 700. Because of a constant amount of pATn5 is added to CUT&TAG

reactions and brings along a fixed amount of E. coli DNA, we used bowtie2 with the

parameters mentioned above to remove E. coli DNA and conduct normalization

according to the CUT&TAG tutorial [25]. CUT&TAG peaks were called using SEACR(1.3)

[66] with default parameters.

RNA-seq analysis

RNA-seq expected counts of the MCF-7 cell lines in all replicates were corrected to be

TPM, the mean TPM of all replicates was used as the expression level of each gene for

subsequent analysis.

Gene single molecular diversity

The accessibility of a single molecule in each gene was calculated, and the value in

each gene was sorted from small to large by the gene unit. Then the hierarchical

clustering was performed for the diversity of single molecule accessibility of each gene.

KOBAS3.0 [67] was used for KEGG and GO analysis for each cluster.

Assess replicate reproducibility

To study the reproducibility between replicates, the genome is split into 50 bp bins and

sliding 5bp, then a Pearson correlation of the log2-transformed values of m^6A

methylation ratio in each bin is calculated between replicate datasets.

SV calling

We used NGMLR (v0.2.7) [68] to compare the read of ONT to the human reference

genome of hg19 to get the BAM file for comparison. Then we used samtools (v1.2) to

sort the bam files.The sniffles (v1.0.12) [68] with the parameter --genotype -T 8 -S 8 were

then used to call the structural variation on the bam file created in the previous step.

Co-accessibility assessment

To evaluate co-accessibility patterns along the genome, we applied COA as follows.

Each chromosome in the genome was split into windows of size w. For each such

window (i, i + w), we identified another window (j,j+w) such that the span (i,j,w) was

covered by ≥N reads. For each single spanning molecule k, accessibility scores (A) in

each bin were then aggregated and binarized as described above. The local co-

accessibility matrix between two windows was calculated:

$$\text{COAi,j,w} = \sum_{i=1}^{n/w} \sum_{j=1}^{n/w} \left(0.5 - \frac{|\text{Ai,w} - \text{Aj,w}|}{\text{Ai,w} + \text{Aj,w}} \right) * \left(\frac{|\text{LAi} - \text{LAj}|}{n} + 1 \right)$$

where n is the length of selected region, L is the location of region.

Data availability

CUT&TAG for H3K27me3 and CTCF as well as Nanopore raw data are available at

China National GeneBank (CNGB) with project number of CNP0001299.

External sequencing datasets.

A number of previously published MCF-7 breast cancer datasets were used in this

study. ChIP-seq data for CTCF was downloaded from ENCODE with accession

ENCSR000DMR, ChIP-seq data for H3K27me3 was downloaded from ENCODE with

accession ENCSR761DLU. The RNA-seq data of MCF-7 was download from the Gene

Expression Omnibus (GEO) repository database with the accession number

GSE71862.

Other external datasets.

Hg19 genome, short interspersed nuclear elements (SINE) and long interspersed

nuclear elements (LINE) region were downloaded from NCBI. TES, TTS and other gene

elements were downloaded from the UCSC Table Browser. MCF-7 CTCF binding site

was downloaded from the CTCFBSDB_v2.0 [69].

Acknowledgment

This research was supported by the Science, Technology, and Innovation Commission

of Shenzhen Municipality (grant number JSGG20170824152728492). The supporter had

no role in designing the study, data collection, analysis, and interpretation, or in writing

the manuscript.

Author contributions

CT designed and supervised the experiments. ZW and FR perform the lab experiments;

WTC and CT perform the bioinformatics data analysis. All others joined the data analysis.

Competing interest

The authors declare no competing interests.

1.   What Is Epigenomics? | SpringerLink. (2021).
2.   Toh, T.B., Lim, J.J. & Chow, E.K. Epigenetics in cancer stem cells. *Mol Cancer* **16**, 29 (2017).
3.   Jiang, Y. et al. Epigenetics in the Nervous System. *The Journal of Neuroscience* **28**, 11753-11759 (2008).
4.   Oshima, M. & Iwama, A. Epigenetics of hematopoietic stem cell aging and disease. *Int J Hematol* **100**, 326-334 (2014).
5.   Tsompana, M. & Buck, M.J. Chromatin accessibility: a window into the genome. *Epigenetics & Chromatin* **7**, 33 (2014).
6.   Voong, L.N., Xi, L., Wang, J.-P. & Wang, X. Genome-wide Mapping of the Nucleosome Landscape by Micrococcal Nuclease and Chemical Mapping. *Trends in Genetics* **33**, 495-507 (2017).
7.   Ryu, Y.B. et al. Biflavonoids from Torreya nucifera displaying SARS-CoV 3CL(pro) inhibition. *Bioorg Med Chem* **18**, 7940-7947 (2010).
8.   Fejes, A.P. et al. FindPeaks 3.1: a tool for identifying areas of enrichment from massively parallel short-read sequencing technology. *Bioinformatics* **24**, 1729-1730 (2008).
9.   Wang, Z. et al. Combinatorial patterns of histone acetylations and methylations in the human genome. *Nat Genet* **40**, 897-903 (2008).
10.   Shipony, Z. et al. Long-range single-molecule mapping of chromatin accessibility in eukaryotes. *Nat Methods* **17**, 319-327 (2020).
11.   Lee, I. et al. Simultaneous profiling of chromatin accessibility and methylation on human cell lines with nanopore sequencing. *bioRxiv*, 504993 (2019).
12.   D., K. et al. The Architecture of SARS-CoV-2 Transcriptome. *Cell* **181** (2020).
13.   Shilatifard, A. Chromatin Modifications by Methylation and Ubiquitination: Implications in the Regulation of Gene Expression. *Annual Review of Biochemistry* **75**, 243-269 (2006).
14.   Mardis, E.R. ChIP-seq: welcome to the new frontier. *Nat Methods* **4**, 613-614 (2007).
15.   Johnson, D.S., Mortazavi, A., Myers, R.M. & Wold, B. Genome-Wide Mapping of in Vivo Protein-DNA Interactions. *Science* **316**, 1497 (2007).
16.   Mikkelsen, T.S. et al. Genome-wide maps of chromatin state in pluripotent and lineage-committed cells. *Nature* **448**, 553-560 (2007).
17.   Barski, A. et al. High-resolution profiling of histone methylations in the human genome. *Cell* **129**, 823-837 (2007).
18.   Schones, D.E. et al. Dynamic regulation of nucleosome positioning in the human genome. *Cell* **132**, 887-898 (2008).
19.   Park, P.J. ChIP-seq: advantages and challenges of a maturing technology. *Nat Rev Genet* **10**, 669-680 (2009).

20. Robertson, G. et al. Genome-wide profiles of STAT1 DNA association using chromatin immunoprecipitation and massively parallel sequencing. *Nat Methods* **4**, 651-657 (2007).

21. Kharchenko, P.V., Tolstorukov, M.Y. & Park, P.J. Design and analysis of ChIP-seq experiments for DNA-binding proteins. *Nat Biotechnol* **26**, 1351-1359 (2008).

22. Valouev, A. et al. Genome-wide analysis of transcription factor binding sites based on ChIP-Seq data. *Nat Methods* **5**, 829-834 (2008).

23. Schmidl, C., Rendeiro, A.F., Sheffield, N.C. & Bock, C. ChIPmentation: fast, robust, low-input ChIP-seq for histones and transcription factors. *Nat Methods* **12**, 963-965 (2015).

24. Skene, P.J. & Henikoff, S. An efficient targeted nuclease strategy for high-resolution mapping of DNA binding sites. *Elife* **6** (2017).

25. Kaya-Okur, H.S. et al. CUT&Tag for efficient epigenomic profiling of small samples and single cells. *Nat Commun* **10**, 1930 (2019).

26. Carter, B. et al. Mapping histone modifications in low cell number and single cells using antibody-guided chromatin tagmentation (ACT-seq). *Nat Commun* **10**, 3747 (2019).

27. Harada, A. et al. A chromatin integration labelling method enables epigenomic profiling with lower input. *Nat Cell Biol* **21**, 287-296 (2019).

28. Ai, S. et al. Profiling chromatin states using single-cell itChIP-seq. *Nat Cell Biol* **21**, 1164-1172 (2019).

29. Statham, A.L. et al. Bisulfite sequencing of chromatin immunoprecipitated DNA (BisChIP-seq) directly informs methylation status of histone-modified DNA. *Genome Res* **22**, 1120-1127 (2012).

30. Brinkman, A.B. et al. Sequential ChIP-bisulfite sequencing enables direct genome-scale investigation of chromatin and DNA methylation cross-talk. *Genome Res* **22**, 1128-1138 (2012).

31. Lhoumaud, P. et al. EpiMethylTag: simultaneous detection of ATAC-seq or ChIP-seq signals with DNA methylation. *Genome Biol* **20**, 248 (2019).

32. Murray, I.A. et al. The non-specific adenine DNA methyltransferase M.EcoGII. *Nucleic Acids Res* **46**, 840-848 (2018).

33. van Steensel, B. & Henikoff, S. Identification of in vivo DNA targets of chromatin proteins using tethered dam methyltransferase. *Nat Biotechnol* **18**, 424-428 (2000).

34. Douvlataniotis, K., Bensberg, M., Lentini, A., Gylemo, B. & Nestor, C.E. No evidence for DNA N (6)-methyladenine in mammals. *Sci Adv* **6**, eaay3335 (2020).

35. O'Brown, Z.K. et al. Sources of artifact in measurements of 6mA and 4mC abundance in eukaryotic genomic DNA. *BMC Genomics* **20**, 445 (2019).

36. Simpson, J.T. et al. Detecting DNA cytosine methylation using nanopore sequencing. *Nat Methods* **14**, 407-410 (2017).

37. Rand, A.C. et al. Mapping DNA methylation with high-throughput nanopore sequencing. *Nat Methods* **14**, 411-413 (2017).

38. Skene, P.J., Henikoff, J.G. & Henikoff, S. Targeted in situ genome-wide profiling with high efficiency for low cell numbers. *Nature Protocols* **13**, 1006-1019 (2018).

39. Kaya-Okur, H.S. et al. CUT&Tag for efficient epigenomic profiling of small samples and single cells. *Nature Communications* **10**, 1930 (2019).

40. Furey, T.S. ChIP–seq and beyond: new and improved methodologies to detect and characterize protein–DNA interactions. *Nature Reviews Genetics* **13**, 840-852 (2012).

41. Kouzarides, T. Chromatin modifications and their function. *Cell* **128**, 693-705 (2007).

42. Berger, S.L. The complex language of chromatin regulation during transcription. *Nature* **447**, 407-412 (2007).

43. Prickaerts, P. et al. Hypoxia increases genome-wide bivalent epigenetic marking by specific gain of H3K27me3. *Epigenetics & Chromatin* **9**, 46 (2016).

44. Young, M.D. et al. ChIP-seq analysis reveals distinct H3K27me3 profiles that correlate with transcriptional activity. *Nucleic Acids Research* **39**, 7415-7427 (2011).

45. Holwerda, S.J. & de Laat, W. CTCF: the protein, the binding partners, the binding sites and their chromatin loops. *Philos Trans R Soc Lond B Biol Sci* **368**, 20120369 (2013).

46. Yin, M. et al. Molecular mechanism of directional CTCF recognition of a diverse range of genomic sites. *Cell Res* **27**, 1365-1377 (2017).

47. Disdero, E. & Filée, J. LoRTE: Detecting transposon-induced genomic variants using low coverage PacBio long read sequences. *Mobile DNA* **8**, 5 (2017).

48. Pauler, F.M. et al. H3K27me3 forms BLOCs over silent genes and intergenic regions and specifies a histone banding pattern on a mouse autosomal chromosome. *Genome Res* **19**, 221-233 (2009).

49. Román, A.C. et al. Dioxin receptor and SLUG transcription factors regulate the insulator activity of B1 SINE retrotransposons via an RNA polymerase switch. *Genome Res* **21**, 422-432 (2011).

50. Kunarso, G. et al. Transposable elements have rewired the core regulatory network of human embryonic stem cells. *Nature Genetics* **42**, 631-634 (2010).

51. Buenrostro, J.D., Wu, B., Chang, H.Y. & Greenleaf, W.J. ATAC-seq: A Method for Assaying Chromatin Accessibility Genome-Wide. *Curr Protoc Mol Biol* **109**, 21.29.21-21.29.29 (2015).

52. Klemm, S.L., Shipony, Z. & Greenleaf, W.J. Chromatin accessibility and the regulatory epigenome. *Nat Rev Genet* **20**, 207-220 (2019).

53. Vendrell, J.A. et al. ZNF217 is a marker of poor prognosis in breast cancer that drives epithelial-mesenchymal transition and invasion. *Cancer Res* **72**, 3593-3606 (2012).

54. Krig, S.R. et al. ZNF217, a candidate breast cancer oncogene amplified at 20q13, regulates expression of the ErbB3 receptor tyrosine kinase in breast cancer cells. *Oncogene* **29**, 5500-5510 (2010).

55. Voigt, P., Tee, W.W. & Reinberg, D. A double take on bivalent promoters. *Genes Dev* **27**, 1318-1338 (2013).

56. Dunican, D.S. et al. Bivalent promoter hypermethylation in cancer is linked to the H327me3/H3K4me3 ratio in embryonic stem cells. *BMC Biol* **18**, 25 (2020).

57. Chapman-Rothe, N. et al. Chromatin H3K27me3/H3K4me3 histone marks define gene sets in high-grade serous ovarian cancer that distinguish malignant, tumour-sustaining and chemo-resistant ovarian tumour cells. *Oncogene* **32**, 4586-4592 (2013).

58.	Li, F. et al. Bivalent Histone Modifications and Development. *Curr Stem Cell Res Ther* **13**, 83-90 (2018).

59.	Adams, J. et al. Vascular endothelial growth factor (VEGF) in breast cancer: comparison of plasma, serum, and tissue VEGF and microvessel density and effects of tamoxifen. *Cancer Res* **60**, 2898-2905 (2000).

60.	Burger, J.A. & Wiestner, A. Targeting B cell receptor signalling in cancer: preclinical and clinical advances. *Nat Rev Cancer* **18**, 148-167 (2018).

61.	Schütz, F. et al. PD-1/PD-L1 Pathway in Breast Cancer. *Oncol Res Treat* **40**, 294-297 (2017).

62.	Hosogane, M., Funayama, R., Shirota, M. & Nakayama, K. Lack of Transcription Triggers H3K27me3 Accumulation in the Gene Body. *Cell Reports* **16**, 696-706 (2016).

63.	Hall, I.M.J.B. BEDTools: a flexible suite of utilities for comparing genomic features. **26**, 841 (2010).

64.	Langmead, B. & Salzberg, S.L.J.N.M. Fast gapped-read alignment with Bowtie 2. **9**, 357-359 (2012).

65.	Feng, J., Liu, T., Qin, B., Zhang, Y. & Liu, X.S.J.N.P. Identifying ChIP-seq enrichment using MACS. **7**, 1728 (2012).

66.	Meers, M.P., Tenenbaum, D., Henikoff, S.J.E. & Chromatin Peak calling by Sparse Enrichment Analysis for CUT&RUN chromatin profiling. (2019).

67.	Chen, X. et al. KOBAS 2.0: a web server for annotation and identification of enriched pathways and diseases. **39**, 316-322 (2011).

68.	Sedlazeck, F.J. et al. Accurate detection of complex structural variations using single-molecule sequencing. (2018).

69.	Lei, B., Mi, Z. & Cui, Y.J.N.A.R. CTCFBSDB: a CTCF-binding site database for characterization of vertebrate genomic insulators. **36**, D83-87 (2008).

A

pA-M.EcoGII

ConA Beads

Antibodies and pA-M.EcoGII fusion protein diffuse in

ConA Beads

pA-M.EcoGII methylates adjacent adenines in antibodies bond DNA

B

pTXB1-pA-M.EcoGII
8298 bp

ProteinA   Linker   M.EcoGII

C

MW kDa   pA-M.EcoGII

D

100bp ladder
M.EcoGII treated lambda DNA+DpnI
Untreated lambda DNA+DpnI
1kb ladder
pA-M.EcoGII treated lambda DNA+DpnI

E

BamHI Insensitive   EcoRV GATATC   PciI ACATGT   PvuII CAGCTG

Marker   M.EcoGII   pA-M.EcoGII   Untreated

F

Samples (OD450)	Dilution 1/120	Dilution 1/480
pA-M.EcoGII	2.238	1.939
M.EcoGII	0.121	0.122
Negative control	0.119	0.122
ProteinA	0.799	0.305

G

chromatin specific antibody

pA-M.EcoGII

SAM, 37°C

Stop buffer

Library preparation
Single molecule

non-m6A labeled adenines   m6A labeled adenines   non-m6A labeled adenines

Figure 1. The experiment concept and validation of the recombinant protein pA-M.EcoGII in BIND&MODIFY. (A)The experiment concept of the BIND&MODIFY. The cells were lightly fixed and permeabilized for the antibody and enzyme passage. The recombinant protein pA-M.EcoGII was located to desired sites under antibody guidance. Then the M.EcoGII activity was locally activated to modify the nearby regions to label the genomic DNA with targeted binding proteins. (B) The upper panel showed the plasmid map of the pA-M.EcoGII. The fused pA-M.EcoGII was cloned into pTXB1 plasmid and purified with compatible IMPACT protein purification system. The lower panel showed the expressed fusion protein structure: Protein A-linker-M.EcoGII-intein-CBD. (C) The Coomassie blue gel stain showed the purity of the purified protein A-M.EcoGII. (D) Methylation of linear lambda DNA by pA-M.EcoGII activates m6A-site dependent DpnI restriction endonuclease digestion. The PCR amplified unmethylated lambda DNA was treated with commercial M.EcoGII, no enzyme, and pA-M.EcoGII. The GATC m6A methylation dependent restriction endonuclease DpnI digestion showed the comparable methyltransferase activity of the commercial M.EcoGII and our recombinant proteins. DNA marker: 100bp ladder, 100-1510bp(left); 1kb ladder, 250-10,000bp(right). (E) The antibody affinity assay showed the recombinant pA-M.EcoGII had the affinity to the secondary antibody in two different dilutions (1/120,1/480, 10mg/ml). (F) Methylation of linear dsDNA by pA-M.EcoGII inhibits multiple site-specific methylation sensitive restriction endonucleases. The unmethylated DNA template was a 7kb linear dsDNA, which was PCR amplified from pTXB1 plasmid. The DNA template was treated with commercial M.EcoGII, pA-M.EcoGII and no enzyme. These treated DNA templates were each incubated with four restriction endonucleases (BamHI, EcoRV, PciI, PvuII).

The BamHI is the m6A methylation insensitive enzyme, and the EcoRV, PciI, PvuII are

the m6A methylation sensitive enzyme, with which the digestion could be blocked by

corresponding m6A site. Our pA-M.EcoGII recombinant protein showed digestion

inhibition on EcoRV, PciI, PvuII digested samples, better than commercial M.EcoGII, as

compared to untreated DNA template. DNA marker, 1kb ladder, 250-10,000bp. (G) The

experiment outlines of BIND&MODIFY. After light fixation and permeabilization, the cells

were tethered to Concanavalin A magnetic beads for the purification in the next steps.

Then the cells were incubated with antibody and pA-M.EcoGII with minimal washes.

The addition of S-adenosylmethionine to initialize the methylation reaction. The DNA

was extracted to prepare the library for ONT nanopore sequencing. After sequencing,

the data was processed as genome alignment and m6A base calling.

Figure 2. The experimental validation of the m6A calling in BIND&MODIFY *in vitro*.

(A) Schematic outline of *in vitro* validation experiment. A fragment of 700bp lambda

DNA was amplified by PCR, and 5mC was introduced at 5' end of forward strand only.

The 5mC labeled DNA was bound by 5mC antibody and was subsequently treated by

BIND&MODIFY method. (B) The methylation ratio of forward strand. When the

methylation cut-off was set at 0.53 (red line, sensitivity=0.92, specificity=0.79), high

methylation possibility region was observed in 3' end at resolution of about 100bp,

which was originally marked by 5mC at 3' end. (C) The methylation ratio of reverse

strand. No high methylation possibility region was observed above 0.53 cut-off.

Figure 3. The consistency of H3K27me3 pattern between ChIP-seq and

BIND&MODIFY in *vivo*. (A) The H3K27me3 signal, by BIND&MODIFY and ChIP-seq,

in genome scale view. (B) The scatter plot of BIND&MODIFY signal and ChIP-seq

signal in peak regions. The peak regions are identified H3K27me3 peaks in ChIP-seq.

The BIND&MODIFY signal means the m6A counts and the ChIP-seq signal means the

read counts. (C) The H3K27me3 pattern was plotted by the TSS centered plot for all the

genes. The plot covered the upstream/downstream 1000bp from TSSs. (D) The

H3K27me3 pattern was plotted by the TSS centered plot for the low expression genes.

The low expression genes are the genes in the bottom expression quantile. (E) The

H3K27me3 strand-specific view by the TSS centered plot for the low expression genes.

Figure 4. The BIND&MODIFY resolved H3K27me3 in the transposon areas with

higher resolution. (A-B) The LTRs with size 350~450bp were selected and centered.

The moving average H3K27me3 signals on the upstream/downstream 300bp of these

LTRs, including LTRs, were plotted with corresponding genomic sites. The y-axis (A)

indicated the normalized read counts of ChIP-seq. The y-axis (B) indicated the

normalized methylation ratio of m6A in BIND&MODIFY with nanopore sequencing.

Normalized peak signal resolution is defined by ratio of mean peak signal width

(maximum-minimum) and mean maximum peak signal, which is about 0.5. Normalized

methylation ratio resolution is defined by ratio of mean methylation ratio width

(maximum-minimum) and mean maximum methylation ratio, which is about 0.045. (C-

D) The SINEs with size 250~350bp were selected and centered. The moving average

H3K27me3 signal on the upstream/downstream 300bp of these SINEs, including

SINEs, were plotted with corresponding genomic sites. The y-axis (C) indicated the

normalized read counts of short-reads ChIP-seq. The y-axis (D) indicated the

normalized methylation ratio of m6A in BIND&MODIFY with nanopore sequencing.

A

B

chr20:52,183,610-52,210,378
(GRCh37/hg19 by Entrez Gene)

C

D

Figure 5. The BIND&MODIFY showed the heterogeneity of H3K27me3 regulation.

(A) The peak signal of H3K27me3 in ChIP-seq (Chr20:52,183,610-52,210,378). (B) The

single molecular resolution of the positive strand Chr20:52,183,610-52,210,378

visualize each molecular methylation statues of H3K27me3. The rows in heatmap

represented the different DNA molecules. The color indicated the methylation ratio,

which represented the H3K27me3 signal. The DNA molecules could be classified into

three states: repressed state, poised state, and active state based on their mean

methylation ratio. The right panel is the condensed illustration of individual DNA

molecules mean methylation ratio. Each block on the vertical line (molecule

heterogeneity line) represented the mean methylation of each DNA molecule and

representing heterogeneity in each DNA molecule. (C) H3K27me3 heterogeneity

pattern of genes in chr20. Mean H3K27me3 methylation ratio of individual DNA

molecule was plotted for all the genes on DNA region (Chr20:52,183,610-52,210,378).

Each pixel on each row corresponds to mean methylation ratio of each individual DNA

molecule, and each row corresponds to each gene. The methylation ratio was ranked

from low to high (left to right). (D) The right heatmap showed the magnified cluster 3 in

the left heatmap. Details of H3K27me3 heterogeneity calculation methods can be found

in supplementary figure 17.

Figure 6. The distance correlation of the epigenetic regulation. (A) The distance

correlation of the H3K27me3 in chr20: 52278000~52280500 (positive strand). The color

indicated the correlation coefficient (CC) metric among genome regions. The higher CC

metric means the stronger correlation in distance. (B) The global view of the H3K27me3

distance effect (DE) index for all the genetic transcription regions (upstream 2kb of

transcription start site) in chr20. The DE index is sum of the CC*distance, representing

this location impact on other sites. The red color means the stronger impact on distal

genomic sites. The left color bar indicated the clusters of these transcription regions.

The right color bar indicated the corresponding gene expression. (C) The global view of

the CTCF distance effect for all the genetic transcription in chr20. Details of correlation

coefficient (CC) metric and distance effect (DE) index calculation method can be found

in supplementary figure 21.