1   Lung adenocarcinoma cell-derived exosomal miR-328 promote osteoclastogenesis by

2   targeting Nrp2

Chengcheng Zhang[1, 2#], Jingru Qin[1#], Lu Yang[1#], Zhiyao Zhu[1], Jia Yang[1], Wan Su[1],

Haibin Deng[1*] & Zhongqi Wang[1*]

[1] Department of Medical Oncology, Longhua Hospital affiliated to Shanghai

University of Traditional Chinese Medicine, 725 Wanpingnan Road, Shanghai

200032, China.

[2] Cancer Institute, Longhua Hospital affiliated to Shanghai University of Traditional

Chinese Medicine, 725 Wanpingnan Road, Shanghai 200032, China.

Corresponding Author:

Haibin Deng, MD, Department of Medical Oncology, Longhua Hospital affiliated to

Shanghai University of Traditional Chinese Medicine, 725 Wanpingnan Road,

Shanghai 200032, China. (Tel: 008613661578200, E-mail: 13661578200@139.com)

Zhongqi Wang, MD, Department of Medical Oncology, Longhua Hospital affiliated

to Shanghai University of Traditional Chinese Medicine, 725 Wanpingnan Road,

Shanghai 200032, China. (Tel: 008618917763093, E-mail: aledx@sina.com)

[#]These authors contributed equally to this work, and both should be considered first

author.

Keywords: exosomes, miR-328, Nrp2, osteoclastogenesis, bone metastasis

Abstract

Bone metastasis of lung cancer and detailed mechanisms are still elusive, and the

roles of exosomes derived from lung adenocarcinoma cells in this process have

attracted much attention. In this study, we found that A549 cell-derived exosomes

(A549-Exos) promoted osteogenesis and bone resorption in vitro. Furthermore,

A549-Exos target bone in vivo and promoted bone resorption in vivo. Mechanistically,

A549-Exosomal miR-328 promoted bone resorption by targeting Nrp2 and

A549-Exos$^{\text{miR-328 Inhibitors}}$ inhibited bone resorption *in vivo*. Thus, A549-Exosomal

miR-328 promote osteoclastogenesis by targeting Nrp2 and A549-Exos$^{\text{miR-328 Inhibitors}}$

may serve as a potential nanomedicine for the treatment of bone metastasis.

Background

Bone metastasis is the most frequent site of lung cancer metastasis[1] and most lung cancer bone metastasis can be classified into osteolytic bone metastasis[2], which can cause pain, pathological, spinal instability, spinal cord compression and hypercalcemia and result in a decline in the quality of life[3,4]. However, lung cancer cells can not directly lead to osteolytic metastasis by proteolytic enzymes. Instead, lung cancer cells derived circulating miRNAs and exosomes played important roles by activating osteoclasts[5,6]. Thus, we focus on the impact of lung cancer derived exosomes on osteoclast differentiation.

Exosomes are small cup or sphere-shaped vesicles, and their diameters primarily ranged from 80 nm to 150 nm, that contains mRNAs, miRNAs[7], lipids[8] and proteins[9]. It has been reported that the role of exosomes in cellular-communication by the transfer of proteins[10], bioactive lipids and miRNAs[11]. Lung cancer derived exosomes have been attracted attention for its role in bone metastasis and prognosis[12]. Furthermore, exosomes may be potentially used as biomarkers, nanoparticle drugs and carriers of antitumor drugs. Recently, numerous investigations indicated that exosomal miRNAs played important roles in tumor growth, invasion and metastasis[13]. Therefore, finding the key exosomal miRNAs in the bone metastasis could be beneficial for treatment of bone metastasis and exosomal miRNAs may serve as a potential therapeutic target for bone metastasis[14].

In this study, we found that A549 cells derived exosomes are important extracellular vesicles in osteoclast differentiation and in bone metastasis. Furthermore, we found that A549-Exos promoted bone resorption in vitro and in vivo. We furthermore explored its mechanism and found that A549-Exosomal miR-328 promoted bone resorption by targeting Nrp2. Therefore, we generated A549-Exos loaded in miR-328 inhibitors and found that bone resorption was prevented by injecting A549-Exos$^{\text{miR-328 Inhibitors}}$. Therefore, we focused on the roles of A549 cells derived exosomes in bone resorption in this study.

Methods

Cell culture

A549 cell line was purchased from the American Type Culture Collection (ATCC)

and cultured in Dulbecco's modified Eagle's medium (DMEM) (Gibco, USA),

supplemented with 10% fetal bovine serum (FBS) (Gibco, USA). The Raw264.7 cell

line was cultured in DMEM (Gibco) supplemented with 10% FBS. We cultured cells

at 37 °C in a 5% CO_2 humidified incubator.

Exosome isolation and characterization

A549 cells were cultured in fresh complete medium containing EVs-free FBS

(Thermo, USA) for 48 h. Then, we collected the conditioned medium and centrifuged

them at 1500 g for 15 minutes at 4°C. After filtering through a 0.22 mm filter

(Millipore, Billerica, USA). The supernatant was ultracentrifuged twice at 110 000 g

for 1 hour at 4°C. Then, we resuspended the pellets in a right amount of PBS and

stored A549-Exos at –80°C for later use. We observed the morphology of A549-Exos

by transmission electron microscope (TEM). The size distribution of A549-Exos was

determined by using Nanosizer™ instrument (Malvern Instruments, Malvern, UK).

The expression of exosomal surface markers (CD9, TSG101) was analyzed by

western blot (WB).

Western blot (WB) analysis

Cells or A549-Exos were lysed in a RIPA lysis buffer supplemented with protease

inhibitor cocktail (CST). Then, we quantified total protein concentration by using the

BCA protein assay kit (Thermo, USA). Western blotting was performed as previously

described[15]. Primary antibodies were used in this study as follows: CD9 (CST, USA),

HSP70 (CST, USA), GAPDH (CST, USA) and OPG (CST, USA). All the secondary

antibodies (1:5000) were acquired from Abcam.

In vitro Osteoclastogenesis assays of RAW264.7 cells

We seeded RAW264.7 cells on 48-well plates ((1×10^4)) in D-MEM (Gibco, USA)

supplemented with 10% FBS (Gibco, USA). M-CSF (30ng/ml) and RANKL

(100ng/ml) were used to promote osteoclast maturation for 7 days. Then, we

performed TARP-staining according to general protocols. We considered TRAP

positive cells (more than 5 nucleus) considered as mature osteoclasts.

Luciferase reporter assay

We constructed the Nrp2 3'-UTR luciferase reporter construct by amplifying the

mouse Nrp2 mRNA 3'- UTR sequence (including the predicted miR-328 binding site)

by PCR. Then, the PCR products were purified and cloned into the XbaI site of the

pGL3-promoter vector (Promega) immediately downstream of the stop codon of

luciferase. The Nrp2 mutations were prepared by using a Quick- Change

Site-Directed Mutagenesis Kit (Stratagene) to get MUT-pGL3-Nrp2, which is

confirmed by sequencing. RAW264.7 cells were transfected with either WT or mutant

pGL3 construct and 40 ng of pRL-TKRenilla-luciferase plasmid and miR-328 mimics

or inhibitors for 48 h. Luciferase activities were measured as previously described[15].

Real-time PCR (RT-PCR)

Total RNA of RAW264.7 cells or A549-Exos were isolated by TRIzol reagent (Life

Technologies, USA) and synthesized complementary DNAs (cDNAs). Then,

Quantitative reverse transcriptase PCR (qRT-PCR) was performed as previously

described[15]. The primer sequences used for qRT-PCR were as follows: miR-328:

forward, 5'-CGG GCC TGG CCC TCT CTG CC-3'; reverse, 5'- CAG CCA CAA

AAG AGC ACA AT-3'; Ctsk: forward, 5'-CTC GGC GTT TAA TTT GGG AGA-3';

reverse, 5'-TCG AGA GGG AGG TAT TCT GAG T-3'; TRAF-6: forward, 5'-AAG

GTG GTG GCG TTA TAC TGC-3'; reverse, 5'-CTG GCA CAG CGG ATG TGA G-3'

and GAPDH: forward, 5'-AAT GGA TTT GGA CGC ATT GGT-3'; reverse, 5'-TTT

GCA CTG GTA CGT GTT GAT-3';

Animal studies

All animal experiments were conducted in accordance with the National Institute of

Health (NIH) Guide for the Care and Use of Laboratory Animals, with the approval of

Longhua hospital (No. LHERAW-19038). C57BL/6 mice (6-8 weeks old) were

obtained from the Animal Center of Longhua hospital. The mice were housed in

barrier housing conditions in the Animal Center of Longhua hospital. For delivery of

A549-Exos or A549-Exos[miR-328 Inhibitors], 100ng/ml in 1 ml PBS was injected into tail

vein once every 3 days for 4 weeks. The mice were sacrificed at 4 weeks after the

treatment.

Micro-CT analysis

femur samples dissected from mice were scanned and analyzed through micro-CT

(Quantum GX, PE, USA). We performed the micro-CT scans under the same

conditions: voltage 100 kV, current 80 μA, spatial resolution 12 μm, scanning 500

continuous sections. The data was collected and analyzed automatically through

computer software (Skyscan) to analyze the number of trabecular bones (Tb.N),

trabecular bone thickness (Tb.Th), trabecular bone space (Tb.Sp), bone volume

fraction (BV/TV) and three-dimensional reconstruction.

Exosomes labeling and tracking in vivo.

All animal experiments were conducted in accordance with the National Institute of

Health (NIH) Guide for the Care and Use of Laboratory Animals, with the approval of

Longhua hospital (No. LHERAW-19038). To visualize the target organ and the

metabolism process of exosomes, we labelled A549-Exos according to the

instructions of the manufacturer (Thermo, USA). First, we incubated 100ug of

exosomes with Vybrant DID (1:1000 in PBS) in the dark for 30 mins, and labelled

exosomes were washed in PBS, with centrifugation at $100,000 \times g$ for 60 min. Then,

the labelled exosomes were injected into the tail vein of mice (8 weeks old, n=3). The

mice were harvested at 3, 6, 12 hours after injection. The mean intensities for DID

signals were quantified from four random visual fields for each section.

HE & TRAP staining

For histological staining, we dissected femora from different groups and fixed them in

4% paraformaldehyde for 24 h, then decalcified them in ethylene diamine tetraacetic

acid (EDTA; pH=7.4) for 28 days, dehydrated them through ethanol and embedded in

paraffin. Then, we cut them into 10 μm thick longitudinally oriented sections and

performed hematoxylin and eosin (H&E) staining and TRAP staining as described

previously[15].

Statistical analysis

All data are shown as means ± standard deviation (SD) from at least three

independent experiments. Comparisons were performed using t test and one-way

ANOVA as appropriate. P < 0.05 was considered statistically significant.

Results

Characteristics of the isolated A549-Exos.

To investigate the role of lung adenocarcinoma cell in osteoclastogenesis, we first determined whether lung adenocarcinoma cell could affect the differentiation of osteoclasts, RAW264.7 cells were co-cultured with A549 cells in a transwell system. The differentiation of osteoclasts was enhanced when co-culturing with A549 cells (Figure 1A). Then, we performed TEM, Dynamic light scattering analysis and Western Blot analysis to charactize the exosomes (Exos) derived from A549. A549-Exos presented a cup or sphere-shaped morphology and their diameters primarily ranged from 80 nm to 150 nm (Figure 1B, 1C). Western Blot analysis showed that A549-Exos expressed exosomal marker proteins including CD9 and TSG101 (Figure 1D). Thus, these data suggested that Osteoclastic differentiation was enhanced when co-culturing with A549 cells and A549-Exos showed characteristic features of exosomes.

A549-Exos promoted osteoclast differentiation and bone resorption in vitro.

To investigate the role of A549-Exos in osteoclastogenesis in vitro. A549-Exos were labeled with lipophilic dye Dio and assessed whether A549-Exos could be transported to Raw264.7 cells. The fluorescence microscope revealed that A549-Exos could be successfully absorbed by Raw264.7 cells (Figure 2A). Then, we treated Raw264.7 cells with A549-Exos or an equal volume of Phosphate Buffer Saline (PBS). osteoclastic differentiation was upregulated as the concentration of exosomes derived from A549 cells increased (Figure 2B, 2C). Furthermore, we found that mRNA expression of Ctsk and Traf-6 were upregulated by A549-Exos compared to its' negative control at day 7 (Figure 2D, 2E). Taken together, these data suggested that A549-Exos promoted osteoclast differentiation in vitro.

A549-Exos target bone in vivo.

To determine the bio-distribution of A549-Exos in vivo. The tissue distribution of DID-labeled A549-Exos (A549-Exos-DID) was examined by fluorescence molecular tomography (FMT) imaging system. A549-Exos-DID was injected into the tail vein of 8-week-old male mice for 3 h, 6 h or 12 h. 3 hours after intravenous injection, the

major fluorescence signals were detected in the liver and lungs and the signals can be

detected in the limbs. After 6 h, the major fluorescence signals were still accumulated

in the liver and lung. Significantly, fluorescence signals in the limbs were enhanced,

but still weaker than those in the liver and lung. After 12 hours, fluorescence signals

still can be detected in the limbs, and fluorescence signals faded from liver and lungs.

The fluorescence signals can be rarely detected in the kidney (Figure 3A).

Furthermore, HE staining suggested that there was no obvious significant histologic

difference in heart, liver, lung, and kidney in different groups, suggesting that

A549-Exos are well-tolerated, with little to no acute, systemic toxicity (Figure S1).

Taken together, A549-Exos could be targeted for bone tissues in vivo without toxicity.

A549-Exos inhibited bone formation in vivo.

To investigate the role of A549-Exos in vivo, we injected A549-Exos or an equal

volume of PBS into tail vein of 8-week-old male mice third per week for 12 weeks.

Microcomputed tomography (micro-CT) showed significantly lower trabecular bone

volume, number, thickness, and higher trabecular separation in those mice injected

with A549-Exos compared to their controls (Figure 4A, 4B-4E). Furthermore, TRAP

staining showed that a greater number of osteoclasts on the trabecular bone surface in

A549-Exos groups than N.C., suggesting an agonist role of A549-Exos in the

osteoclast differentiation, which was consistent with our in vitro study (Figure 4F, 4H).

Our results indicated that A549-Exos promoted bone resorption in vivo.

A549-Exosomal miR-328 inhibited bone formation by targeting Nrp2.

Since exosomes may concentrate large numbers of miRNAs, which can regulate gene

expression by binding 3′-untranslated regions. MiR-328 has been reported to promote

osteoclast differentiation[16]. Subsequently, we examined the expression levels of

miR□328 in the exosomes, and qRT-PCR showed that A549 cells derived exosomes

had significantly higher expression of miR-328 in comparison with TC-1 cells derived

exosomes (Figure 5A).

To investigate osteoclastogenesis regulated by miR-328 in vitro. RAW264.7 cells

were transfected with miR-328 mimics or inhibitors. The miR-328 expression levels

were significantly upregulated by miR-328 mimic transfection and markedly inhibited

by miR-328 inhibitor transfection (Figure 5B). Furthermore, overexpression of

miR-328 suppress endogenous expression of OPG protein level (Figure 5C).

Consistent with the results of OPG protein expression, we found that miR-328 mimics

enhanced TRAP staining, whereas miR-328 inhibitors weakened TRAP staining

(Figure 5D). Overall, these data showed that osteoclast differentiation was promoted

by miR-328 in vitro.

To determine bone resorption regulated by miR-328 in vivo. we injected miR-328

inhibitors or their negative controls into tail vein of 8-week-old male mice third per

week for 8 weeks. Mice treated with miR-328 inhibitors showed increased trabecular

bone volume, number, and thickness, and decreased trabecular separation relative to

their negative groups. These data showed that bone resorption was regulated by

miR-328 in vivo.

To investigate target mRNAs of miR-328 in osteoclast differentiation process, we

have checked the miRNA-binding site in Targetscan and found that miR-328 binging

site on Nrp2 3′-UTR (Figure 5G). Previously, Nrp2 deficiency has been reported to

promote osteoclast differentiation and reduced osteoblast numbers[17]. Thus, luciferase

reporter constructs containing the WT or mutated predicted miRNA-binding sites of

Nrp2 were constructed. Then, RAW264.7 cells were transfected with

WT-Nrp2–3′-UTR or MUT-Nrp2–3′-UTR with miR-328 mimics or negative controls

and the luciferase activity of the Nrp2 reporter gene was inhibited by miR-328 mimics

and the Nrp2 reporter gene was promoted by miR-328 inhibitors, whereas

MUT-Nrp2–3′-UTR prevented this inhibition (Figure 5H). In conclusion, these data

showed that miR-328 directly targets Nrp2 and inhibited Nrp2 expression in

osteoclast cells.

A549-Exos[miR-328 Inhibitors] inhibited bone resorption *in vivo*.

To investigate the therapeutic potential of miR-328 on bone metastasis,

A549-Exos[miR-328 Inhibitors] was generated. A549 cells transfected with miR-328

inhibitors derived exosomes were labeled with lipophilic dye Dio and assessed

whether A549-Exos could carry miR-328 inhibitors and be transported to Raw264.7

cells. The fluorescence microscope revealed that A549-Exos[miR-328 Inhibitors] could be

successfully absorbed by Raw264.7 cells (Figure 6A). Then, we injected A549-Exos

or A549-Exos$^{miR-328\ Inhibitors}$ into tail vein of 8-week-old male mice third per week for

12 weeks. Mice treated with A549-Exos$^{miR-328\ Inhibitors}$ showed increased trabecular

bone volume, number, and thickness, and decreased trabecular separation relative to

A549-Exos group, which showed that A549-Exos$^{miR-328\ Inhibitors}$ could alleviate bone

loss caused by the bone metastasis (Figure 6B, 6C). In addition, TRAP staining

showed fewer number of osteoclasts on the trabecular bone surface in

A549-Exos$^{miR-328\ Inhibitors}$ groups than A549-Exos (Figure 6D, 6E), which revealed that

an inhibitory role of A549-Exos$^{miR-328\ Inhibitors}$ in the differentiation of osteoclasts. In

conclusion, these data suggested that intravenous injection of A549-Exos$^{miR-328\ Inhibitors}$

inhibited bone resorption in vivo and alleviate the osteopenia caused by bone

metastasis.

Discussion

In this study, we found that osteoclast differentiation was enhanced when co-culturing with A549 cells. A549-Exos promote osteoclast differentiation in vitro. Furthermore, A549-Exos could be targeted for bone tissues and promoted bone resorption in vivo. Moreover, we found that A549-Exosomal miR-328 promoted bone resorption by targeting Nrp2. In addition, A549-Exos[miR-328 Inhibitors] treatment in vivo ameliorated bone loss and prevented pathological fractures.

Previously, several cancer cells derived exosomes have been reported to be home to the bone microenvironment and cause metastatic lesions[18]. As murine prostate cancer line TRAMP-C1 inhibits the differentiation of the monocytic cell line RAW264.7 into osteoclasts[19], non-small cell lung cancer (NSCLC) cells secrete exosomes promoted osteoclast differentiation of murine RAW264.7 cells by activation of EGFR phosphorylation[20], exosomes released by multiple myeloma cells increase the viability and migration of osteoclast precursors, through the increasing of CXCR4 expression[21] and extracellular vesicles isolated from PC3 culture medium are internalized into osteoclast precursors and osteoblasts[22]. In this study, A549 cell-derived exosomal miR-328 is the regulator of bone resorption and A549-Exos[miR-328 Inhibitors] could ameliorate bone loss. Thus, A549 cells derived exosomes could load small molecule drugs and target bone tissues. Since a set of tumor cells derived exosomes could target bone tissues, we hypothesized that these exosomes can also be used as targeted drug carriers for the corresponding primary tumors. Our work might provoke interesting future works regarding the relationship between bone metastasis and tumor cells derived exosomes leading to new potential therapeutic targets.

A set of exosomal miRNAs has been reported to be correlated with bone formation or bone resorption[23]. For example, Exosomal miR-141-3p regulates osteoblast activity to promote the osteoblastic metastasis of prostate cancer[24], overexpression of hsa☐miR☐940 in tumor exosomes and its ability to ultimately lead to extensive osteoblastic lesions in the resulting tumor[25] and mi-192 cargo within exosome like vesicle transfer influences metastatic bone colonization[26]. These miRNAs may also

serve as therapeutic potential targets for the treatment of bone metastases from lung

cancers, which requires further investigation.

Conclusions

Lung adenocarcinoma cell-derived exosomes (A549-Exos) promoted osteogenesis

and bone resorption in vitro. Furthermore, A549-Exos target bone in vivo and

promoted bone resorption in vivo. Mechanistically, A549-Exosomal miR-328 promote

osteoclastogenesis by targeting Nrp2 and A549-Exos$^{\text{miR-328 Inhibitors}}$ may serve as a

potential nanomedicine for the treatment of bone metastasis.

Acknowledgements

Not applicable.

Author contributions:

Conceived and designed the review: HBD and ZQW. Collected the data: ZYZ and

CCZ. Analyzed the data: JY and WS. Supervised the study: JRQ and CCZ. Wrote the

paper: CCZ, JRQ and LY.

Fundings

This study was supported in part by National Natural Science Foundation of China

(No. 81904129), Innovation Program of Shanghai Municipal Education Commission

(No. 2017-01-07-00-10-E00064), Shanghai Hospital Development Center

(SHDC2020CR4050), Longhua Hospital Scholar Fund (NO.LYTD-33).

Availability of data and materials

Data sharing is not applicable to this article as no datasets were generated or analyzed

during the current study.

Ethics approval

All animal experiments were conducted in accordance with the National Institute of

Health (NIH) Guide for the Care and Use of Laboratory Animals, with the approval of

Longhua hospital (No.LHERAW-19038).

Consent for publication

All authors reached an agreement to publish the study in this journal.

Conflict of interests

The authors indicate no potential conflicts of interest.

References

1.   Fornetti, J., Welm, A.L. & Stewart, S.A. Understanding the Bone in Cancer Metastasis.

J Bone Miner Res **33**, 2099-2113 (2018).

2.   Zhang, L. & Gong, Z. Clinical Characteristics and Prognostic Factors in Bone

Metastases from Lung Cancer. *Med Sci Monit* **23**, 4087-4094 (2017).

3.   Altorki, N.K., *et al.* The lung microenvironment: an important regulator of tumour

growth and metastasis. *Nat Rev Cancer* **19**, 9-31 (2019).

4.   Wu, S., Pan, Y., Mao, Y., Chen, Y. & He, Y. Current progress and mechanisms of

bone metastasis in lung cancer: a narrative review. *Transl Lung Cancer Res* **10**,

439-451 (2021).

5.   Yin, J.J., Pollock, C.B. & Kelly, K. Mechanisms of cancer metastasis to the bone. *Cell*

Res **15**, 57-62 (2005).

6.   Xu, S., *et al.* Serum microRNA-139-5p is downregulated in lung cancer patients with

lytic bone metastasis. *Oncol Rep* **39**, 2376-2384 (2018).

7.   Hu, Y., *et al.* Exosomes from human umbilical cord blood accelerate cutaneous wound

healing through miR-21-3p-mediated promotion of angiogenesis and fibroblast

function. *Theranostics* **8**, 169-184 (2018).

8.   Record, M., Silvente-Poirot, S., Poirot, M. & Wakelam, M.J.O. Extracellular vesicles:

lipids as key components of their biogenesis and functions. *J Lipid Res* **59**, 1316-1324

(2018).

9.   Chen, C.Y., *et al.* Exosomal DMBT1 from human urine-derived stem cells facilitates

diabetic wound repair by promoting angiogenesis. *Theranostics* **8**, 1607-1623 (2018).

10.   Zeelenberg, I.S., *et al.* Targeting tumor antigens to secreted membrane vesicles in

vivo induces efficient antitumor immune responses. *Cancer Res* **68**, 1228-1235

(2008).

11.   Kumar, S. & Reddy, P.H. Are circulating microRNAs peripheral biomarkers for

Alzheimer's disease? *Biochim Biophys Acta* **1862**, 1617-1627 (2016).

12.   Li, F.X., *et al.* Role of tumor-derived exosomes in bone metastasis. *Oncol Lett* **18**,

3935-3945 (2019).

13.   Dai, J., *et al.* Exosomes: key players in cancer and potential therapeutic strategy.

Signal Transduct Target Ther **5**, 145 (2020).

14.   Zhang, J. & Wu, J. The Potential Roles of Exosomal miR-214 in Bone Metastasis of

Lung Adenocarcinoma. *Front Oncol* **10**, 611054 (2020).

15.   Wang, R., *et al.* miR-143 promotes angiogenesis and osteoblast differentiation by

targeting HDAC7. *Cell Death Dis* **11**, 179 (2020).

16.   Chen, J., *et al.* Identification of suitable reference gene and biomarkers of serum

miRNAs for osteoporosis. *Sci Rep* **6**, 36347 (2016).

17.   Verlinden, L., *et al.* Nrp2 deficiency leads to trabecular bone loss and is accompanied

by enhanced osteoclast and reduced osteoblast numbers. *Bone* **55**, 465-475 (2013).

18.   Rossi, M., Battafarano, G., D'Agostini, M. & Del Fattore, A. The Role of Extracellular

Vesicles in Bone Metastasis. *Int J Mol Sci* **19**(2018).

19.   Karlsson, T., Lundholm, M., Widmark, A. & Persson, E. Tumor Cell-Derived Exosomes

from the Prostate Cancer Cell Line TRAMP-C1 Impair Osteoclast Formation and

Differentiation. *PLoS One* **11**, e0166284 (2016).

20.   Taverna, S., *et al.* Amphiregulin contained in NSCLC-exosomes induces osteoclast

differentiation through the activation of EGFR pathway. *Sci Rep* **7**, 3170 (2017).

21.   Raimondi, L., *et al.* Involvement of multiple myeloma cell-derived exosomes in

osteoclast differentiation. *Oncotarget* **6**, 13772-13789 (2015).

22.   Inder, K.L., *et al.* Cavin-1/PTRF alters prostate cancer cell-derived extracellular

vesicle content and internalization to attenuate extracellular vesicle-mediated

osteoclastogenesis and osteoblast proliferation. *J Extracell Vesicles* **3**(2014).

23.   Luga, V., *et al.* Exosomes mediate stromal mobilization of autocrine Wnt-PCP

signaling in breast cancer cell migration. *Cell* **151**, 1542-1556 (2012).

24.   Ye, Y., *et al.* Exosomal miR-141-3p regulates osteoblast activity to promote the

osteoblastic metastasis of prostate cancer. *Oncotarget* **8**, 94834-94849 (2017).

25.   Hashimoto, K., *et al.* Cancer-secreted hsa-miR-940 induces an osteoblastic

phenotype in the bone metastatic microenvironment via targeting ARHGAP1 and

FAM134A. *Proc Natl Acad Sci U S A* **115**, 2204-2209 (2018).

26.   Valencia, K., *et al.* miRNA cargo within exosome-like vesicle transfer influences

metastatic bone colonization. *Mol Oncol* **8**, 689-703 (2014).

392
393
394

Figure legends

Figure 1 Characteristics of the isolated A549-Exos. (A) Representative images of
TRAP staining osteoclasts. Scale bar: 100 μm. (B) Representative TEM image of
A549-Exos (scale bar = 200 nm). (C) Representative NTA results showing the size
distribution of A549-Exos. (C) Detection of A549-Exos surface markers (CD9 and
HSP70) by Western Blot analysis.

Figure 2 A549-Exos promoted osteoclast differentiation and bone resorption in

vitro. (A) Fluorescence microscopy analysis of RAW264.7 cells treated with

DiO-labeled A549-Exos for 2h. (B) Representative images of TRAP staining of

mature osteoclasts treated with different concertrations of A549-Exos. Scale bar: 100

μm. (C) Quantitative analysis of positive TRAP staining osteoclasts in Figure 2B

(**P<0.01). (D) Relative levels of Ctsk (left) and TRAF-6 (right) mRNA expression

in Figure 2B (**P<0.01).

414
Figure 3. A549-Exos target bone in vivo without toxicity. (A) Representative FMT
images of the near-infrared fluorescence signals in organs isolated from mice
administered with PBS, A549-Exos for different periods of time (3 hours, 6 hours, and
12 hours).
419
420

Figure 4. A549-Exos inhibited bone formation in vivo. (A) Representative micro-CT images in femora from A549-Exos mices and negative control group (n = 8 per group). Scale bar: 100 µm. (B, C, D, E) Quantitative analysis of trabecular microarchitecture from A549-Exos group and negative control group. (F) Representative TRAP staining of femora from A549-Exos mice and negative control group. (G) Quantitative analysis of osteoclasts per bone surface in different groups. BV/TV, trabecular bone volume per tissue volume; Tb. Th, trabecular thickness; Tb. Sp, trabecular separation; Tb. N, trabecular number. (n=8 per group) (*P<0.05).

Figure 5. A549-Exosomal miR-328 promoted bone resorption by targeting Nrp2.
(A) miR-328 relative expressions in N.C. and LCC-Exos. (B) miR-328 levels in N.C.,
miR-328 mimics, I.N.C. and miR-328 inhibitors were measured by qRT-PCR. (C)
Western blot (WB) analysis of OPG and GAPDH in RAW264.7 cells transfected with
miR-328 mimics and miR-328 or their respective negative controls. (D)
Representative images of TRAP staining of mature osteoclasts treated with N.C.,
miR-328 mimics, I.N.C. and miR-328 inhibitors. (E) Representative micro-CT images
in femora from different groups. (F) Quantitative analysis of trabecular
microarchitecture from different groups. (G) Nrp2 might be the target of miR-328. (H)
Raw264.7 were transfected with a luciferase reporter carrying WT or MUT-Nrp2
3′-UTR in the miR-328 mimics (H left) and miR-328 (H right) or their respective
negative controls. *p < 0.05, **p < 0.01. Data are representative of at least three

443   independent experiments.

444
Figure 6. A549-Exos^{miR-328 Inhibitors} inhibited bone resorption *in vivo.* (A)
Fluorescence microscopy analysis of RAW264.7 cells treated with DiO-labeled
A549-Exos transfected with miR-328 inhibitors for 2h (B) Representative micro-CT
images in femora from A549-Exos group and A549-Exos^{miR-328 Inhibitors} group (n = 6
per group). (C) Quantitative analysis of trabecular microarchitecture from A549-Exos
group and A549-Exos^{miR-328 Inhibitors} group (n=8 per group). (D) Representative TRAP
staining of femora from different groups (n=8 per group). (E) Quantitative analysis of
osteoclasts per bone surface in different groups. (*P<0.05).
453
454

Figure S1 (A). Quantitative analysis of positive TRAP staining osteoclasts in Figure
1A. (B). The quantification analysis of the Western Blot of CD9 and HSP70. *P <
0.05, **P < 0.01.

Figure S2 (A). There was no histomorphometric change in the heart, liver, lung and
kidney.