Title

Multiple roles of heterogeneous nuclear ribonucleoprotein K during skeletal

muscle cell differentiation

Authors

Keisuke Hitachi[1], Yuri Kiyofuji[1], Masashi Nakatani[2], and Kunihiro Tsuchida[1,*]

Affiliations

[1]Division for Therapies against Intractable Diseases, Institute for Comprehensive

Medical Science (ICMS), Fujita Health University, Toyoake, Japan, [2]Faculty of

Rehabilitation and Care, Seijoh University, Tokai, Japan

*Correspondence: tsuchida@fujita-hu.ac.jp

Abstract

RNA-binding proteins (RBPs) regulate cell physiology via the formation of

ribonucleic-protein complexes with coding and non-coding RNAs. RBPs have multiple

functions in the same cells; however, the precise mechanism through which their

pleiotropic functions are determined remains unknown. In this study, we revealed the

multiple inhibitory functions of hnRNPK for myogenic differentiation. We first

identified hnRNPK as a lncRNA *Myoparr* binding protein. Gain- and loss-of-function

experiments showed that hnRNPK repressed the expression of *myogenin* at the

transcriptional level via binding to *Myoparr*. Moreover, hnRNPK repressed the

expression of a set of genes coding for aminoacyl-tRNA synthetases in a

Myoparr-independent manner. Mechanistically, hnRNPK regulated the eIF2α/Atf4

pathway, one branch of the intrinsic pathways of the endoplasmic reticulum sensors, in

differentiating myoblasts. Thus, our findings demonstrate that hnRNPK plays multiple

lncRNA-dependent and -independent roles in the inhibition of myogenic differentiation,

indicating that the analysis of lncRNA-binding proteins will be useful for elucidating

both the physiological functions of lncRNAs and the multiple functions of RBPs.

Keywords

transcriptional regulation, myogenic differentiation, RNA-binding protein, endoplasmic

reticulum stress

Introduction

Long non-coding RNAs (lncRNAs), which are >200 nucleotides (nt) in length and which do not encode more than 100 amino acids, are emerging as important regulators in diverse biological processes, including transcription, splicing, RNA stability, and translation (Statello et al., 2021). LncRNAs are pervasively transcribed from the noncoding genomic DNA; cis-regulatory regions, including promoter and enhancer, introns, 3' untranslated regions, and repetitive sequences (Chakraborty et al., 2014). LncRNAs are also expressed from the antisense direction of the coding genomic DNA (Hon et al., 2017). Thus, most of the genomic regions have the potential to express lncRNAs. Thus far, more than 260,000 lncRNA genes are registered in the LncBook, which is a curated knowledge-based database for human lncRNAs (Ma et al., 2019). Since lncRNAs exert their molecular functions by interacting with proteins (Hitachi and Tsuchida, 2020), their molecular functions differ widely, depending on the interacting partners.

As lncRNA-interacting factors, RNA-binding proteins (RBPs) are essential to determine the molecular function of each lncRNA (Statello et al., 2021). In the human genome, more than 1,500 genes encode RBPs (Gerstberger et al., 2014). RBPs consist of ribonucleoprotein complexes together with lncRNAs to regulate various biological aspects. For example, the association of Ddx5/Ddx17 with lncRNAs, such as *SRA*, *mrhl*, *MeXis*, or *Myoparr*, is required to activate the expression of downstream genes (Caretti et al., 2006; Hitachi et al., 2019; Kataruka et al., 2017; Sallam et al., 2018). RBPs, including NONO, SFPQ, FUS, and RBM14, associate with *Neat1*, which is a highly abundant lncRNA in mammals, to form a large membrane-less structure paraspeckle in the nucleus (Hirose et al., 2019). A ubiquitously expressed RBP, known as human

antigen R (HuR) associates with lncRNAs and regulates their stability; HuR increases the cytosolic *linc-MD1* levels in skeletal muscle cells (Legnini et al., 2014), whereas it promotes the decay of *lincRNA-p21* in HeLa cells (Yoon et al., 2012). Additionally, RBPs are also involved in RNA splicing, polyadenylation, RNA transport, and translation (Kelaini et al., 2021). The majority of RBPs are involved in multiple biological processes in concert with lncRNAs (Briata and Gherzi, 2020; Jonas et al., 2020; Nostrand et al., 2020), and mutations in genes coding for RBPs are associated with human genetic disorders (Gebauer et al., 2021).

Heterogeneous nuclear ribonucleoprotein K (hnRNPK), a member of the heterogeneous nuclear ribonucleoprotein family, has multiple roles, including chromatin remodeling, transcription, RNA splicing, and translation (Bomsztyk et al., 2004; Wang et al., 2020). hnRNPK acts together with *lincRNA-p21*, *EWSAT1*, and *lncRNA-OG*, to regulate the expression of downstream genes in mouse embryonic fibroblasts, Ewing sarcoma, and bone marrow-derived mesenchymal stem cells (Dimitrova et al., 2014; Howarth et al., 2014; Tang et al., 2018). However, the molecular function of hnRNPK in skeletal muscle cells has not been fully elucidated. LncRNA *Myoparr* is an essential regulator of skeletal muscle cell proliferation and differentiation (Hitachi et al., 2019). We previously identified hnRNPK as a candidate for *Myoparr*-associated protein in skeletal muscle cells (Hitachi et al., 2019), suggesting the unique functions of *Myoparr*-associated hnRNPK during myogenic differentiation. In the present study, we revealed the inhibitory role of hnRNPK as a *Myoparr*-associated protein in skeletal muscle cell differentiation. Moreover, by comparing the downstream genes regulated by *Myoparr* and hnRNPK, we also found a *Myoparr*-independent role of hnRNPK during myogenic differentiation. Our findings reveal that hnRNPK plays multiple

lncRNA-dependent and -independent roles in skeletal muscle cell differentiation and

will contribute to elucidating the complex roles of RBPs in cell differentiation.

Materials and Methods

Cell cultures, siRNA transfection, and ISRIB treatment

A mouse myoblast cell line, C2C12, was cultured in Dulbecco's modified

Eagle medium (DMEM) supplemented with 10% fetal bovine serum at 37°C under 5%

CO_2. Myogenic differentiation was induced by replacing the medium with the

differentiation medium, DMEM supplemented with 2% horse serum. C2C12 myoblasts

were transfected with 50 nM of Stealth RNAi (Thermo Fisher Scientific, Waltham, MA,

USA) using Lipofectamine 3000 (Thermo Fisher Scientific) according to the

manufacturer's protocol. The following siRNAs were used: Stealth RNAi siRNA

negative control (Negative Control, Med GC, Thermo Fisher Scientific), stealth RNAi

for *Myoparr*, and stealth RNAi siRNAs specific for *hnRNPK* (MSS205172 and

MSS205173, Thermo Fisher Scientific). The siRNA sequences are listed in

Supplemental Table 1. At 24 h after siRNA transfection, myogenic differentiation was

induced. At 24 h or 72 h after differentiation induction, cells were collected for the

analysis of RNAs and proteins. For ISRIB treatment, the differentiation medium was

added either with or without 1 µM ISRIB (Cayman Chemical Company, Ann Arbor, MI,

USA).

RNA isolation, reverse transcription reaction, and quantitative RT-PCR

Total RNA was extracted from C2C12 cells using ISOGEN II reagent

(Nippon Gene, Tokyo, Japan) according to the manufacturer's protocol. After DNase I

(Thermo Fisher Scientific) treatment, total RNA was used for reverse transcription

reaction using SuperScript III Reverse Transcriptase (Thermo Fisher Scientific) or

ProtoScript II Reverse Transcriptase (New England Biolabs (NEB), Beverly, MA, USA)

with random or oligo (dT) primers (Thermo Fisher Scientific). Quantitative real-time

PCR was conducted using SYBR Premix Ex Taq (Takara Bio Inc., Shiga, Japan) and a

Thermal Cycler Dice Real Time System TP800 (Takara Bio Inc.). The results were

normalized to the *Rpl26* expression. The primers used are listed in Supplementary Table

1.

Protein extraction and Western blotting

Cells were lysed in RIPA buffer (50 mM Tris-HCl (pH 8.0), 150 mM NaCl,

0.1% SDS, 1% Triton X-100, 0.5% sodium deoxycholate) containing protease inhibitors

(1 mM phenylmethylsulfonyl fluoride, 1 µg/ml aprotinin, 4 µg/ml leupeptin) and

phosphatase inhibitors (5 mM NaF, 5 mM β-glycerophosphate, 1 mM Na_3VO_4). The

protein concentration was measured using a Pierce BCA Protein Assay Kit (Thermo

Fischer Scientific). Equal amounts of protein were used for Western blotting. The

following primary antibodies were used: myogenin antibody (F5D) (sc-12732, Santa

Cruz Biotechnology, Dallas, TX, USA), MHC antibody (MF20, Developmental Studies

Hybridoma Bank (DSHB)), hnRNPK antibody (#4675, Cell Signaling Technology

(CST), Beverly, MA, USA), TIAR antibody (#8509, CST), and Atf4 antibody (693901,

BioLegend, San Diego, CA, USA). The following HRP-linked secondary antibodies

were used: anti-mouse IgG (#7076, CST) and anti-rabbit IgG (#7074, CST). Can Get

Signal Immunoreaction Enhancer Solution (Toyobo, Osaka, Japan) was used when

necessary. The signal was detected with ImmunoStar LD reagent (FUJIFILM Wako,

Osaka, Japan) using a cooled CCD camera system (Light-Capture, ATTO, Tokyo,

Japan).

Immunofluorescence assay

Immunofluorescence analyses of C2C12 cells were performed as previously

described (Hitachi et al., 2019). Briefly, 24 h or 72 h after the induction of

differentiation, cells were fixed with 4% PFA and permeabilized with 0.2% Triton

X-100. After blocking with 5% FBS, cells were stained with an anti-myogenin antibody

(F5D, DSHB) or with an anti-MHC antibody (MF20, DSHB). Nuclei were

counterstained with DAPI (Dojindo, Kumamoto, Japan). A DMI4000B microscope with

a DFC350FX CCD camera (Leica, Wetzlar, Germany) was used for visualization of

signals. Images were analyzed using the Image J software program (ver. 1.49).

Identification of *Myoparr*-binding proteins

Myoparr-binding proteins were collected with the RiboTrap Kit (Medical &

Biological Laboratories (MBL), Aichi, Japan) using BrU labeled RNAs, as described

previously (Hitachi et al., 2019). BrU-labeled *Myoparr* and *EGFP* RNA were prepared

using the Riboprobe System (Promega, Madison, WI, USA). The *Myoparr* or *EGFP*

RNA (50 pmol) were mixed with the nuclear extract from C2C12 myotubes for 2 h at

4°C. The RNA-protein complexes were collected by Protein G Plus Agarose (Thermo

Fisher Scientific) conjugated to an anti-BrdU antibody, and proteins were eluted by

adding BrdU. Purified proteins were detected by Western blotting using a specific

antibody, as described above.

RNA immunoprecipitation and the RNA pull-down assay

Immunoprecipitation of endogenous RNAs (*Myoparr* or *Xist*) was performed using the RIP-Assay Kit (MBL) using 2×10^7 C2C12 cells, as previously described (Hitachi et al., 2019). The following antibodies were used for RNA immunoprecipitation: normal rabbit IgG (#2729S, CST), anti-HNRNPK pAb (RN019P, MBL), or TIAR mAb (#8509, CST). After treatment with DNase I, immunoprecipitated RNAs was used for the reverse transcription reaction. The precipitation percentage (precipitated RNA vs. input RNA) was calculated by qRT-PCR using the primers listed in Supplementary Table 1.

An RNA pull-down assay was performed with a RiboTrap Kit. Various lengths of *Myoparr* were subcloned into a pGEM-Teasy vector (Promega). BrU-labeled *Myoparr* (10 pmol each) was bound to Protein G Plus Agarose conjugated to an anti-BrdU antibody and mixed with the *in vitro* transcribed/translated hnRNPK protein. After several washing steps, the binding of hnRNPK to *Myoparr* was analyzed by Western blotting using an hnRNPK antibody as described above.

Luciferase reporter assay

The upstream region of *myogenin* (-1650/+51) was PCR-amplified and cloned into the pGL4.20 vector (Promega). The *hnRNPK*-expressing plasmid was a kind gift from Dr. H. Okano (Yano et al., 2005). Proliferating C2C12 cells were transfected with the *myogenin* promoter and *hnRNPK*-expressing plasmid using Lipofectamine 2000 (Thermo Fisher Scientific). The total amount of DNA was kept constant by the addition of the pcDNA3 vector. Cells were collected at 24 h after the induction of differentiation and dissolved in Passive Lysis Buffer (Promega). The effect of hnRNPK on *myogenin*

promoter was measured using a Lumat LB 9507 luminometer (Berthold Technologies,

Bad Wildbad, Germany) with the Dual-Luciferase Reporter Assay System (Promega)

according to the manufacturer's protocol. The pGL4.74 vector (Promega) was used as

an internal control. The relative luciferase activity is shown as the firefly to Renilla

luciferase ratio.

To reconstitute the complex chromatin structure and epigenetic regulation *in*

vitro (Liu et al., 2001), the upstream regions of *myogenin* (-242/+51 and -1650/+51)

were subcloned into the episomal luciferase vector, pREP4-luc, and the -242-Luc and

-1650-Luc constructs were generated. To create the -1650Δccawmcc-Luc construct, the

upstream region (-971/-902) of *myogenin* was deleted from the -1650-Luc construct.

The region containing the putative hnRNPK-binding motif ccawmcc was identified by

RBPmap (Paz et al., 2014). The episomal luciferase vector, pREB7-Rluc, was used as

an internal control for the luciferase assay. The pREP4-luc and pREP7-Rluc vectors

were gifts from Dr. K. Zhao (Liu et al., 2001). Subconfluent C2C12 cells were

transfected with the indicated episomal luciferase vectors using Lipofectamine 2000.

After myogenic induction, cells were collected and used to measure the relative

luciferase activity.

Analysis of downstream genes regulated by *Myoparr* KD and *hnRNPK* KD

Downstream genes regulated by *Myoparr* KD and *hnRNPK* KD were identified as

described previously (Hitachi and Tsuchida, 2019) using our RNA-Seq raw data

(accession No. DRA005527). Briefly, statistical analysis of differentially expressed

genes by *Myoparr* KD and *hnRNPK* KD was performed using DESeq2 ver. 1.12.4

software (Love et al., 2014) with a Wald test (cut-offs: false discovery rate (adjusted

p-value, padj) < 0.05 and log 2 fold change > 0.75 or < -0.75). A pathway analysis of

significantly upregulated genes coding for cytosolic aminoacyl-tRNA synthetases was

performed based on the KEGG (Kanehisa et al., 2020). A Gene Ontology (GO) analysis

of differentially expressed genes was performed with DAVID ver. 6.8

(https://david.ncifcrf.gov/). An enrichment analysis was performed using Metascape

(Zhou et al., 2019).

Statistical analysis

Error bars represent standard deviation. Statistical analyses were performed

using unpaired two-tailed Student's *t*-tests. For comparisons of more than 2 groups, a

one-way ANOVA followed by Tukey's post hoc test was performed using Prism 9

(GraphPad Software, San Diego, CA, USA). Statistical significance is reported in the

Figures and Figure legends. *P* values of < 0.05 were considered statistically significant.

Results

Identification of hnRNPK as a *Myoparr*-binding protein in skeletal muscle cells

Our proteomics analysis identified both hnRNPK and TIAR as candidates for

Myoparr-associated proteins (Hitachi et al., 2019). To reveal whether hnRNPK and

TIAR are associated with the *Myoparr* function during myogenic differentiation, we

first examined the specific interaction of *Myoparr* with endogenous hnRNPK and TIAR.

In vitro synthesized *Myoparr* was labeled with 5-bromouridine (BrU) and mixed with

the nuclear extract from differentiating C2C12 myoblasts. After purification of *Myoparr*

by immunoprecipitation with a BrdU antibody, specific binding between *Myoparr* and

hnRNPK protein was confirmed by immunoblotting (Figure 1A). Since binding

between *EGFP* mRNA and hnRNPK was not observed, *EGFP* mRNA was used as a

negative control. Intracellular binding between endogenous *Myoparr* and hnRNPK was

shown by RNA immunoprecipitation using an hnRNPK-specific antibody without a

crosslinking (Figure 1B and C). This enrichment of *Myoparr* by hnRNPK was stronger

than that of *Xist*, which interacts with hnRNPK in embryonic stem cells (Chu et al.,

2015). Although TIAR protein was retrieved by synthesized *Myoparr*, intracellular

endogenous interaction between *Myoparr* and TIAR was not observed (Supplementary

Figure 1A-C). These results suggested that hnRNPK cooperatively works with *Myoparr*

during skeletal muscle differentiation.

hnRNPK represses the expression of *myogenin* in differentiating myoblasts

During myogenic differentiation, the expression of *Myoparr* gradually

increases and is required to activate the expression of *myogenin* (Hitachi et al., 2019).

Thus, we examined the changes in the expression of hnRNPK during C2C12 cell

differentiation. Although the expression of myogenin was highly increased after

myogenic induction, the expression of hnRNPK gradually decreased (Figure 2A),

suggesting the regulatory role of hnRNPK in the expression of *myogenin*. We next

knocked down *hnRNPK* using the small interfering RNAs (siRNAs) in differentiating

C2C12 cells. The knockdown (KD) of *hnRNPK* by two distinct siRNAs was associated

with a large increase in the expression of myogenin (Figure 2B). However, an

immunocytochemistry analysis of the myogenin expression after *hnRNPK* KD did not

show a sufficient increase in the ratio of myogenin-positive cells comparing to the

control (Figure 2C). These results indicated that *hnRNPK* KD is not enough to induce

the expression of myogenin in cells without the cell-intrinsic expression of myogenin.

We observed that the expression of *myogenin* was significantly increased by

hnRNPK KD (Figure 2D and E). Although the differences were not statistically

significant, the *Myoparr* expression levels also tended to be increased by *hnRNPK* KD

(Figure 2F). From these results, we surmise that hnRNPK negatively regulates the

expression of *myogenin* at the transcriptional level. The effect of hnRNPK on the

transcription of *myogenin* was examined using the *myogenin*-promoter-driven luciferase

assay. The overexpression of hnRNPK in differentiating C2C12 cells decreased the

myogenin promoter activity (Figure 2G), indicating that hnRNPK negatively regulates

the expression of *myogenin* via the *myogenin* promoter.

hnRNPK represses the expression of *myogenin* via binding to *Myoparr*

The hnRNPK-binding region of *Myoparr* was determined by RNA pull-down

experiments using the various forms of *Myoparr* (Figure 3A). Figure 3B showed that

hnRNPK bound to the full-length sense-strand of *Myoparr* (#3 in Figure 3B), but not to

the full-length antisense-strand of *Myoparr* (#4). Although deletion of the 5'-region (#5

and 6) or 3'-region (#2 and 7) of *Myoparr* did not affect binding to hnRNPK, the

deletion of the 3'-half of *Myoparr* (#1) completely diminished the binding to hnRNPK

(Figure 3B). These results indicated that an approximately 300-nt region (613-952 nt) of

Myoparr is indispensable for binding to hnRNPK. Searching the motif sequence of

RBPs from the 300-nt region revealed that there are 8 ccawmcc motifs, which are

recognized by hnRNPK (Figure 3C). The deletion of the motifs (660-729 nt) from the

full-length of *Myoparr* (#8) markedly weakened the binding to hnRNPK (Figure 3B).

Thus, the ccawmcc motifs on *Myoparr* were shown to be required for binding to

hnRNPK.

To clarify whether *Myoparr* is involved in regulating the expression of

myogenin by hnRNPK, we examined the effect of the ccawmcc motif on the *myogenin*

promoter activity. To imitate a chromatin structure and epigenetic regulation on the

plasmid DNA (Liu et al., 2001), the upstream region of *myogenin* (-1649 to +52)

including *Myoparr* was cloned into an episomal luciferase vector. In accordance with

our previous findings (Hitachi et al., 2019), the *myogenin* promoter showed high

activity in the presence of *Myoparr* (-1650-Luc) in comparison to the -242-Luc

construct, which only contains the *myogenin* promoter region in differentiating

myoblasts (Figure 3D). Intriguingly, the activity of the -1650-Luc construct was further

enhanced by the deletion of a region of approximately 70 bp (1650Δccawmcc-Luc),

which corresponds to the 660-729 nt region on *Myoparr* (Figure 3D). These results

indicate that hnRNPK represses the expression of *myogenin* through binding to the

ccawmcc motif on *Myoparr* during skeletal muscle differentiation.

hnRNPK inhibits skeletal muscle differentiation but is required for normal

myotube formation

The inhibitory role of hnRNPK in the expression of *myogenin* via the

ccawmcc motif on *Myoparr* suggested that hnRNPK and *Myoparr* have common

downstream genes. Our RNA-Seq analysis (Hitachi and Tsuchida, 2019) revealed that

hnRNPK KD significantly increased the expression of 226 genes and significantly

decreased the expression of 190 genes . We compared the downstream genes regulated

by *hnRNPK* KD and *Myoparr* KD, and the comparative heatmap analysis showed that

the gene regulated by *hnRNPK* KD showed the opposite direction to the *Myoparr* KD

(Figure 4A). Twenty percent of genes (84/416) altered by *hnRNPK* KD overlapped with

genes regulated by the *Myoparr* KD (Figure 4B and Supplementary Talbe 2). The

intersection of these genes was 12.3-fold greater than that expected by chance (p =

1.393037×10^{-45}). Although these 84 genes showed a low correlation coefficient

(R=0.0758323), we observed a negative correlation trend for one segment of them,

including *myogenin*; the expression of 50 genes belonging to this segment was

increased by *hnRNPK* KD and decreased by *Myoparr* KD (red frame in Figure 4C).

These genes were enriched in sarcomere organization, myofibril assembly, and muscle

contraction categories in GO terms (Figure 4D), indicating that hnRNPK inhibits

myogenic differentiation and maturation.

In accordance with the results of the RNA-Seq analysis, we observed a

significant increase in the expression of Myosin heavy chain (MHC), which is a later

marker of myogenic differentiation and maturation, with *hnRNPK* KD (Figure 4E),

indicating the inhibitory role of hnRNPK in myogenic differentiation and maturation.

Intriguingly, *hnRNPK* KD did not affect the percentage of differentiated cells; this was

shown by the fusion index (Figure 4F) as well as the results from immunocytochemistry

to detect myogenin (Figure 2C). Instead, we observed the appearance of locally

spherical myotubes following *hnRNPK* KD. These differed from the normal

tube-shaped myotubes (Figure 4F). Thus, these results indicated that in addition to its

inhibitory role in myogenic differentiation, hnRNPK is required for normal myotube

formation.

hnRNPK represses the expression of aminoacyl-tRNA synthetases via the

eIF2α/Atf4 pathway

The Venn diagram in Figure 4B indicates that 332 genes regulated by

hnRNPK KD were *Myoparr*-independent, suggesting the *Myoparr*-independent role of

hnRNPK in differentiating myoblasts. We performed an enrichment analysis of genes

regulated by *hnRNPK* KD and compared them with genes regulated by *Myoparr* KD.

As reported previously (Hitachi et al., 2019), genes related to cell cycle and cell division

were only enriched in genes regulated by *Myoparr* KD (Figure 5A). Skeletal

muscle-associated genes were regulated by both *Myoparr* KD and *hnRNPK* KD (Figure

5A). Intriguingly, genes coding for aminoacyl-tRNA synthetases were regulated

specifically by *hnRNPK* KD (red frame in Figure 5A). In mice, there are two-types of

aminoacyl-tRNA synthetases: cytosolic and mitochondrial aminoacyl-tRNA synthetase.

Our RNA-Seq analysis showed that *hnRNPK* KD significantly increased the expression

of 10 genes coding for cytosolic aminoacyl-tRNA synthetases, whereas it had little

effect on the expression of genes coding for mitochondrial aminoacyl-tRNA synthetases

(Supplementary Figure 2). To confirm the RNA-Seq results, we randomly picked up 5

genes coding for cytosolic aminoacyl-tRNA synthetases, and their expression changes

by *hnRNPK* KD were verified by qRT-PCR. *hnRNPK* KD using two distinct siRNAs

significantly increased the expression of *Aars*, *Gars*, *Iars*, *Nars*, and *Sars* (Figure 5B-F).

This regulation was not observed following *Myoparr* KD (Figure 5B-F), indicating that

hnRNPK regulates the expression of these genes in a *Myoparr*-independent manner.

The expression of genes coding for almost all cytosolic aminoacyl-tRNA

synthetases is activated by transcription factor Atf4 (Harding et al., 2003; Shan et al.,

2016). Thus, we examined whether *hnRNPK* KD altered the expression of Atf4 in

skeletal muscle cells. Although not statistically significant, the *Atf4* expression in

differentiating myoblasts tended to be increased by *hnRNPK* KD (Figure 5G). *Myoparr*

KD did not affect the expression of *Atf4* (Figure 5G). The effect of *hnRNPK* KD was

more pronounced in the expression of Atf4 protein. The amount of Atf4 protein was

highly increased by *hnRNPK* KD (Figure 5H). These results suggest that hnRNPK

regulates the expression of cytosolic aminoacyl-tRNA synthetases via the expression of

Atf4.

Under the condition of endoplasmic reticulum (ER) stress, *Atf4* mRNA is

translated more efficiently and contributes to the restoration of cell homeostasis via the

regulation of cytosolic aminoacyl-tRNA synthetases (Afroze and Kumar, 2017). Thus,

to reveal the molecular mechanism by which hnRNPK regulates the expression of Atf4,

we finally focused on ER stress. ISRIB is an inhibitor of eIF2α, which is a downstream

component of PERK signaling, one branch of the ER stress sensors. We investigated

whether ISRIB treatment could suppress the increase in the expression of Atf4 induced

by *hnRNPK* KD and found that ISRIB treatment completely rescued this

hnRNPK-KD-induced increase (Figure 6A). Furthermore, ISRIB treatment abrogated

the increased expression of *Aars*, *Gars*, *Iars*, *Nars*, and *Sars* by *hnRNPK* KD (Figure

6B-F and Supplementary Figure 3A-E). Thus, these results indicate that hnRNPK

regulates the expression of cytosolic aminoacyl-tRNA synthetases via the eIF2α/Atf4

pathway during myogenic differentiation.

Discussion

RBPs have multiple molecular functions, including RNA splicing,

transcription, translation, RNA stability, and the formation of the nuclear structure, to

regulate cell proliferation, differentiation, development, and diseases (Kelaini et al.,

2021). Although many RBPs have multiple functions in the cells (Briata and Gherzi,

2020; Jonas et al., 2020; Nostrand et al., 2020), it is still unclear how their pleiotropic

functions are determined. In this study, we revealed novel multiple functions of

hnRNPK, a member of the hnRNP family of RBPs, in skeletal muscle cells. By

focusing on a lncRNA *Myoparr*-associated protein, we found that hnRNPK repressed

the expression of *myogenin*, coding for one of the master regulators of muscle

differentiation, and inhibited myogenic differentiation through binding to the ccawmcc

motif on *Myoparr*. Moreover, our comparative analysis of the downstream genes of

hnRNPK and *Myoparr* showed that the function of hnRNPK was heterogenous. During

myogenic differentiation, hnRNPK repressed the expression of a set of genes coding for

cytosolic aminoacyl-tRNA synthetases via the eIF2α/Atf4 pathway. Taken together, our

study revealed multiple inhibitory roles of hnRNPK in skeletal muscle cells: one was

lncRNA-dependent and the other was lncRNA-independent (Figure 7). Recently, Xu et

al. reported that in hnRNPK, the deficiency of 36 amino acids diminished C2C12

differentiation (Xu et al., 2018). However, our results provided strong evidence to

support that hnRNPK has an inhibitory effect on muscle differentiation. In addition, we

observed the appearance of locally spherical myotubes following *hnRNPK* KD. The

morphology of these myotubes closely resembles myotubes with myofibril-assembly

defects (Wang et al., 2013), suggesting that hnRNPK may also be involved in the

regulation of the myofibril assembly in myotubes. Therefore, despite its

lncRNA-dependent and -independent roles in the inhibition of myogenic differentiation,

hnRNPK is apparently required for the formation of normal myotubes.

We observed that *hnRNPK* KD increased myogenin protein levels more

robustly than *myogenin* mRNA. The peak expression of myogenin protein is detected

1-2 days after that of *myogenin* mRNA in both *in vitro* and *in vivo* myogenesis (Angelis

et al., 1992; Figueroa et al., 2003), suggesting that a slight increase in *myogenin* mRNA

by *hnRNPK* KD in the early stages of myogenic differentiation eventually led to a

marked increase in myogenin protein. Therefore, the fine-tuning of the *myogenin*

expression by hnRNPK at the early stages of differentiation may have a significant

impact on the overall myogenic differentiation processes through the RNA-protein

network. Intriguingly, the expression of myogenin protein was increased by *hnRNPK*

KD, and the percentage of myogenin-positive cells was not changed, even if *hnRNPK*

was depleted. These results suggest that hnRNPK can specifically repress the

expression of *myogenin* in a subset of responding cells, rather than by simply turning off

the expression of *myogenin* in every myoblast. Our experiments showed that hnRNPK

repressed the expression of *myogenin* at the transcriptional level by binding to the

ccawmcc motif on *Myoparr*, suggesting that the existence of *Myoparr* would be

necessary for hnRNPK to inhibit the expression of *myogenin*. Since *myogenin* and

Myoparr share the same promoter region (Hitachi et al., 2019), *myogenin* and *Myoparr*

are likely expressed in the same cells. Thus, the expression of *myogenin* would not be

activated in the cells without the cell-intrinsic expression of *Myoparr*, even if *hnRNPK*

is depleted in every myoblast. Further studies are required to investigate the more

precise molecular mechanism by which hnRNPK regulates the expression of *myogenin*

via binding to *Myoparr*.

ER stress is induced by several perturbations disrupting cell homeostasis,

including protein misfolding, viral infection, and changes in intracellular calcium

concentration (Hetz et al., 2015). The cells recognize those stresses with three branches

of ER transmembrane sensors signaling, PERK, inositol-requiring protein 1 (IRE1), and

ATF6 (Afroze and Kumar, 2017). During myogenic differentiation, ATF6 signaling was

activated and led to apoptosis in myoblasts (Nakanishi et al., 2005). The increased

phosphorylation of eIF2α, a component of PERK signaling, was observed in myoblasts

at the early stage after the induction of differentiation (Alter and Bengal, 2011). In

addition, treatment with ER stress inducers enhanced apoptosis in myoblasts but led to

efficient myogenic differentiation in the remaining surviving cells (Nakanishi et al.,

2007). Recently, the deletion of PERK in satellite cells, which are adult muscle stem

cells, was shown to inhibit myogenic differentiation and led to impaired skeletal muscle

regeneration in adult mice (Xiong et al., 2017), indicating that ER stress promotes

myogenic differentiation. In this study, we showed that the *hnRNPK* KD in

differentiating myoblasts increased the expression of Atf4 and this effect was

diminished by treatment with ISRIB, a specific inhibitor of eIF2α. The expression levels

of a set of genes coding for cytosolic aminoacyl-tRNA synthetases, which are regulated

by Atf4, were also increased by the *hnRNPK* KD, and were completely rescued after

ISRIB treatment. The inhibitory effects of hnRNPK on Atf4 and cytosolic

aminoacyl-tRNA synthetases were independent of *Myoparr*. Therefore, our findings

suggest that hnRNPK fine-tunes the myogenic differentiation process by modulating ER

stress via eIF2α/Atf4 signaling in a lncRNA-independent manner. Since hnRNPK is

involved in the translational efficiency (Lynch et al., 2005; Yano et al., 2005), a

decrease in hnRNPK may induce the unfolded protein response via the translational

machinery in myoblasts and alter eIF2α/Atf4 signaling.

In conclusion, hnRNPK plays multiple lncRNA-dependent and -independent

roles in the inhibition of myogenic differentiation. Thus, the analysis of RBPs bound to

lncRNAs will be useful for elucidating both the physiological functions of lncRNAs and

the complex functions of RBPs in cell differentiation. Intriguingly, the *myogenin* gene is

responsible for the skeletal muscle atrophy caused by denervation (Moresi et al., 2010).

Atf4 is also involved in skeletal muscle atrophy caused by aging, fasting, and

immobilization (Ebert et al., 2012; Ebert et al., 2015). Induced ER stress, including

increased PERK signaling, was observed in skeletal muscle biopsy samples from

myotonic dystrophy 1 patients and in mdx mice, a model of Duchenne muscular

dystrophy (Hulmi et al., 2016; Ikezoe et al., 2007). Moreover, mutations in genes

coding for aminoacyl-tRNA synthetases are implicated in human neuromuscular

disorders (Benarroch et al., 2020), and autoantibodies against aminoacyl-tRNA

synthetases are found in autoimmune disease (Targoff et al., 1993). Collectively,

downstream genes of hnRNPK are strongly associated with neuromuscular and other

disorders in humans, suggesting that targeting hnRNPK to regulate the expression of

these genes and signaling may become a new therapeutic strategy for human diseases.

Acknowledgments

This work was supported in part by JSPS KAKENHI (19H03427 and 20K07315),

Intramural Research Grants (2-5) for Neurological and Psychiatric Disorders of NCNP,

and a Grant-in-Aid from the Mochida Memorial Foundation for Medical and

Pharmaceutical Research.

Author Contributions

K.H. and K.T. designed research; K.H. analyzed data; K.H., Y.K., and M.N. performed

research; K.H. and K.T. wrote the paper. All authors revised, edited, and read the

manuscript and approved the final manuscript.

Conflict of interest

The authors declare no competing interests in association with the present study.

Figure Legends

Figure 1. Intracellular interaction between *Myoparr* and hnRNPK in

differentiating C2C12 cells. (A) Synthesized *Myoparr* was mixed with C2C12 nuclear

extract. Following purification of *Myoparr* by immunoprecipitation, the interaction

between *Myoparr* and hnRNPK was confirmed by immunoblotting using an

hnRNPK-specific antibody. *EGFP* mRNA was used for the control. Note that the

smaller band would be a splicing isoform of hnRNPK (Makeyev and Liebhaber, 2002).

(B) RNA immunoprecipitation for *Myoparr* and *Xist* in differentiating C2C12 cells by

an hnRNPK antibody. The interaction between endogenous *Myoparr* and hnRNPK was

detected by qRT-PCR. Normal rabbit IgG was used for the control. The presence or

absence of reverse transcription reaction is shown by (RT+) or (RT-), respectively. n = 4,

mean ± SD. ***$p < 0.001$. (C) hnRNPK protein from C2C12 cells purified by

immunoprecipitation in (B) was confirmed by Western blotting.

Figure 2. hnRNPK represses the expression of myogenin at the transcriptional

level. (A-B) Western blots showing the hnRNPK and myogenin expression during

C2C12 differentiation (A) and in C2C12 cells 48 h after *hnRNPK* KD (B). The tubulin

expression served as an internal control. n = 3. (C) Immunocytochemistry for myogenin

48 h after *hnRNPK* KD in C2C12 cells. Nuclei were counterstained with DAPI. Bar,

100 μm. The percentage of the myogenin-positive cells is shown as the percentage of

the control. n = 4, mean ± SD. n.s., not significant. (D-E) Quantitative RT-PCR to detect

the expression of *hnRNPK* (D), *myogenin* (E), and *Myoparr* (F) after *hnRNPK*

knockdown. n = 3-4, mean ± SD. ***$p < 0.001$, *$p < 0.05$, n.s., not significant. (G)

Exogenous hnRNPK decreased the promoter activity of *myogenin* in differentiating

C2C12 cells. n = 3, mean ± SD. *$p < 0.05$.

Figure 3. hnRNPK represses the expression of *myogenin* via binding to *Myoparr*.

(A) A schematic diagram of various lengths of *Myoparr* used for RNA pull-down assays.

(B) *In vitro* transcribed/translated hnRNPK protein was pulled down by *Myoparr* and

then detected by Western blotting using an hnRNPK-specific antibody. Lane numbers

are consistent with (A). The RNA lanes indicate various forms of *Myoparr* used for

RNA pull-down. (C) The results of an RBPmap analysis searching the motif of

hnRNPK. Colored letters correspond to the hnRNPK motif. The red frame indicates the

ccawmcc motifs within the region deleted in Δccawmcc, #8 in (A). (D) The left panel

shows a schematic diagram of the constructs used for the luciferase assays. -242-Luc

contains the *myogenin* promoter region. -1650-Luc contains both the *myogenin*

promoter region and *Myoparr*. -1650Δccawmcc-Luc contains the *myogenin* promoter

region and *Myoparr* without the ccawmcc motifs, corresponding to #8 in A. The right

panel shows the relative luciferase activities of indicated constructs in differentiating

C2C12 cells. n = 4, mean ± SD. ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$.

Figure 4. hnRNPK inhibits myogenic differentiation but is required for normal

myotube formation. (A) A heatmap diagram showing the increased or decreased

expression of genes regulated by *Myoparr* KD and *hnRNPK* KD. (B) The intersection

of genes regulated by *Myoparr* KD and *hnRNPK* KD shows a significant (Fisher's exact

test) overlap, which is 12.3 times larger than would be expected by chance alone. (C)

Eighty-four genes with expression levels that were significantly altered by both

Myoparr KD and *hnRNPK* KD showed an un-correlated pattern (R = 0.0758323, log 2

ratio scale). The red frame indicates the 50 genes for which the expression was

decreased by *Myoparr* KD but increased by *hnRNPK* KD. (D) Enrichment GO

categories of the 50 genes, corresponding to the red frame in (C). (E) Western blot

showing the increased expression of MHC in C2C12 myotubes after *hnRNPK* KD. The

tubulin expression served as an internal control. n = 3. (F) Immunocytochemistry for

MHC 72 h after *hnRNPK* KD. Nuclei were counterstained with DAPI. Bar, 100 μm.

The fusion index is shown as the percent of the control. n = 3, mean ± SD. n.s., not

significant.

Figure 5. hnRNPK regulates the expression of genes coding for aminoacyl-tRNA

synthetases in a *Myoparr*-independent manner. (A) A Metascape analysis of genes

regulated by *Myoparr* KD and *hnRNPK* KD. The red frame indicates the GO terms that

were specifically enriched in *hnRNPK* KD. (B-G) qRT-PCR to detect the expression of

Aars (B), *Gars* (C), *Iars* (D), *Nars* (E), *Sars* (F), *Atf4* (G) after either *Myoparr* KD or

hnRNPK KD. n = 3, mean ± SD. $^{**}p < 0.01$, $^{*}p < 0.05$, n.s., not significant. (H) Western

blot showing the increased expression of Atf4 by *hnRNPK* KD. The tubulin expression

served as an internal control. n = 3.

Figure 6. hnRNPK regulates the expression of genes coding for aminoacyl-tRNA

synthetases via the eIF2α/Atf4 pathway. (A) Western blot showing the increased Atf4

expression after *hnRNPK* KD, and ISRIB treatment suppressed the increase. DMSO is

used as a control. n = 3. (B-F) qRT-PCR to detect the expression of *Aars* (B), *Gars* (C),

Iars (D), *Nars* (E), *Sars* (F) after *hnRNPK* KD and ISRIB treatment. n = 3, mean ± SD.

**$p < 0.01$, *$p < 0.05$, n.s., not significant.

Figure 7. Multiple inhibitory roles of hnRNPK during myogenic differentiation.

During myogenic differentiation, hnRNPK has two different downstream targets. One is

myogenin, which codes for a regulator of myogenic differentiation. hnRNPK represses

the *myogenin* expression by binding to the ccawmcc motifs on *Myoparr*. The other

target is aminoacyl-tRNA synthetases. hnRNPK regulates the expression of

aminoacyl-tRNA synthetases via the eIF2α/Atf4 pathway. In hnRNPK-depleted cells,

the hyperactivated expression of these genes may lead to the locally spherical formation

of myotubes.

Supplementary Table 1. Primer and siRNA sequences.

Supplementary Table 2. The value (log2 fold change) of the genes for which

expression levels were increased or decreased by *Myoparr* KD and *hnRNPK* KD.

Supplementary Figure 1. Detection of the interaction between *Myoparr* and TIAR

in differentiating C2C12 cells. (A) The interaction between synthesized *Myoparr* and

TIAR was detected by immunoblotting using a TIAR antibody. (B) qRT-PCR for the

detection of *Myoparr* and *Xist* following RNA-immunoprecipitation using a TIAR

antibody. The presence or absence of reverse transcription reaction is shown by (RT+)

or (RT-), respectively. Bars indicate the average of two independent experiments, and

circles and triangles represent the values of each experiment. (C) Purified TIAR protein

from C2C12 cells by immunoprecipitation using a TIAR antibody was confirmed by

Western blotting.

Supplementary Figure 2. RNA-Seq revealed the altered expression of a group of

genes coding for cytosolic aminoacyl-tRNA synthetases by *hnRNPK* KD. (A) The

KEGG pathway diagram of cytosolic aminoacyl-tRNA biosynthesis. The gene names

surrounded by the red frame indicate genes that are significantly upregulated by

hnRNPK KD. (B-C) The results of the RNA-Seq analysis showing the altered

expression of genes coding for cytosolic aminoacyl-tRNA synthetases (B) and

mitochondrial aminoacyl-tRNA synthetases (C) by *hnRNPK* KD. Red and blue indicate

genes that are significantly upregulated or downregulated, respectively, by *hnRNPK*

KD.

Supplementary Figure 3. hnRNPK regulates the expression of a group of genes

coding for aminoacyl-tRNA synthetases via the eIF2α/Atf4 pathway. (A-E) The

results of qRT-PCR for detecting the altered expression of *Aars* (A), *Gars* (B), *Iars* (C),

Nars (D), *Sars* (E) following *hnRNPK* KD (using a different siRNA from Figure 6) and

ISRIB treatment. n = 3, mean ± SD. *$p < 0.05$.

References

Afroze, D. and Kumar, A. (2017). ER stress in skeletal muscle remodeling and
myopathies. *Febs J* **286**, 379–398.
Alter, J. and Bengal, E. (2011). Stress-induced C/EBP Homology Protein (CHOP)
represses MyoD transcription to delay myoblast differentiation. *Plos One* **6**, e29498.
Angelis, M. G. C.-D., Lyons, G., Sonnino, C., Angelis, L. D., Vivarelli, E., Farmer,

K., Wright, W. E., Molinaro, M., Bouchè, M. and Buckingham, M. (1992).
MyoD, myogenin independent differentiation of primordial myoblasts in mouse
somites. *J Cell Biology* **116**, 1243–1255.

Benarroch, L., Bonne, G., Rivier, F. and Hamroun, D. (2020). The 2021 version of
the gene table of neuromuscular disorders (nuclear genome). *Neuromuscular Disord*
30, 1008–1048.

Bomsztyk, K., Denisenko, O. and Ostrowski, J. (2004). hnRNP K: One protein
multiple processes. *Bioessays* **26**, 629–638.

Briata, P. and Gherzi, R. (2020). Long non-coding RNA-ribonucleoprotein networks
in the post-transcriptional control of gene expression. *Non-coding Rna* **6**, 40.

Caretti, G., Schiltz, R. L., Dilworth, F. J., Padova, M. D., Zhao, P., Ogryzko, V.,
Fuller-Pace, F. V., Hoffman, E. P., Tapscott, S. J. and Sartorelli, V. (2006). The
RNA helicases p68/p72 and the noncoding RNA SRA are coregulators of MyoD
and skeletal muscle differentiation. *Dev Cell* **11**, 547–560.

Chakraborty, S., Deb, A., Maji, R. K., Saha, S. and Ghosh, Z. (2014). LncRBase: An
enriched resource for lncRNA information. *Plos One* **9**, e108010.

Chu, C., Zhang, Q. C., da Rocha, S. T., Flynn, R. A., Bharadwaj, M., Calabrese, J.
M., Magnuson, T., Heard, E. and Chang, H. Y. (2015). Systematic discovery of
Xist RNA binding proteins. *Cell* **161**, 404–416.

Dimitrova, N., Zamudio, J. R., Jong, R. M., Soukup, D., Resnick, R., Sarma, K.,
Ward, A. J., Raj, A., Lee, J. T., Sharp, P. A., et al. (2014). LincRNA-p21 activates
p21 in cis to promote polycomb target gene expression and to enforce the G1/S
checkpoint. *Mol Cell* **54**, 777–790.

Ebert, S. M., Dyle, M. C., Kunkel, S. D., Bullard, S. A., Bongers, K. S., Fox, D. K.,
Dierdorff, J. M., Foster, E. D. and Adams, C. M. (2012). Stress-induced skeletal
muscle Gadd45a expression reprograms myonuclei and causes muscle atrophy. *J*
Biol Chem **287**, 27290–27301.

Ebert, S. M., Dyle, M. C., Bullard, S. A., Dierdorff, J. M., Murry, D. J., Fox, D. K.,
Bongers, K. S., Lira, V. A., Meyerholz, D. K., Talley, J. J., et al. (2015).
Identification and small molecule inhibition of an activating transcription factor 4
(ATF4)-dependent pathway to age-related skeletal muscle weakness and atrophy. *J*
Biol Chem **290**, 25497–25511.

Figueroa, A., Cuadrado, A., Fan, J., Atasoy, U., Muscat, G. E., Muñoz-Canoves, P.,

623 **Gorospe, M. and Muñoz, A.** (2003). Role of HuR in skeletal myogenesis through
624  coordinate regulation of muscle differentiation genes. *Mol Cell Biol* **23**, 4991–5004.

625 **Gebauer, F., Schwarzl, T., Valcárcel, J. and Hentze, M. W.** (2021). RNA-binding
626  proteins in human genetic disease. *Nat Rev Genet* **22**, 185–198.

627 **Gerstberger, S., Hafner, M. and Tuschl, T.** (2014). A census of human RNA-binding
628  proteins. *Nat Rev Genet* **15**, 829–845.

629 **Harding, H. P., Zhang, Y., Zeng, H., Novoa, I., Lu, P. D., Calfon, M., Sadri, N., Yun,**
630  **C., Popko, B., Paules, R., et al.** (2003). An integrated stress response regulates
631  amino acid metabolism and resistance to oxidative stress. *Mol Cell* **11**, 619–633.

632 **Hetz, C., Chevet, E. and Oakes, S. A.** (2015). Proteostasis control by the unfolded
633  protein response. *Nat Cell Biol* **17**, 829–838.

634 **Hirose, T., Yamazaki, T. and Nakagawa, S.** (2019). Molecular anatomy of the
635  architectural NEAT1 noncoding RNA: The domains, interactors, and biogenesis
636  pathway required to build phase‐separated nuclear paraspeckles. *Wiley Interdiscip*
637  *Rev Rna* **10**, e1545.

638 **Hitachi, K. and Tsuchida, K.** (2019). Data describing the effects of depletion of
639  Myoparr, myogenin, Ddx17, and hnRNPK in differentiating C2C12 cells. *Data*
640  *Brief* **25**, 104172.

641 **Hitachi, K. and Tsuchida, K.** (2020). The Chemical Biology of Long Noncoding
642  RNAs. *Rna Technologies* 431–463.

643 **Hitachi, K., Nakatani, M., Takasaki, A., Ouchi, Y., Uezumi, A., Ageta, H., Inagaki,**
644  **H., Kurahashi, H. and Tsuchida, K.** (2019). Myogenin promoter-associated
645  lncRNA Myoparr is essential for myogenic differentiation. *Embo Rep* **20**, e47468.

646 **Hon, C.-C., Ramilowski, J. A., Harshbarger, J., Bertin, N., Rackham, O. J. L.,**
647  **Gough, J., Denisenko, E., Schmeier, S., Poulsen, T. M., Severin, J., et al.** (2017).
648  An atlas of human long non-coding RNAs with accurate 5′ ends. *Nature* **543**, 199
649  –204.

650 **Howarth, M. M., Simpson, D., Ngok, S. P., Nieves, B., Chen, R., Siprashvili, Z.,**
651  **Vaka, D., Breese, M. R., Crompton, B. D., Alexe, G., et al.** (2014). Long
652  noncoding RNA EWSAT1-mediated gene repression facilitates Ewing sarcoma
653  oncogenesis. *J Clin Invest* **124**, 5275–5290.

654 **Hulmi, J. J., Hentilä, J., DeRuisseau, K. C., Oliveira, B. M., Papaioannou, K. G.,**
655  **Autio, R., Kujala, U. M., Ritvos, O., Kainulainen, H., Korkmaz, A., et al.** (2016).

Effects of muscular dystrophy, exercise and blocking activin receptor IIB ligands on
the unfolded protein response and oxidative stress. *Free Radical Bio Med* **99**, 308–
322.

Ikezoe, K., Nakamori, M., Furuya, H., Arahata, H., Kanemoto, S., Kimura, T.,
Imaizumi, K., Takahashi, M. P., Sakoda, S., Fujii, N., et al. (2007). Endoplasmic
reticulum stress in myotonic dystrophy type 1 muscle. *Acta Neuropathol* **114**, 527–
535.

Jonas, K., Calin, G. A. and Pichler, M. (2020). RNA-binding proteins as important
regulators of long non-coding RNAs in cancer. *Int J Mol Sci* **21**, 2969.

Kanehisa, M., Furumichi, M., Sato, Y., Ishiguro-Watanabe, M. and Tanabe, M.
(2020). KEGG: integrating viruses and cellular organisms. *Nucleic Acids Res* **49**,
D545-D551.

Kataruka, S., Akhade, V. S., Kayyar, B. and Rao, M. R. S. (2017). Mrhl long
noncoding RNA mediates meiotic commitment of mouse spermatogonial cells by
regulating Sox8 expression. *Mol Cell Biol* **37**, e00632-16.

Kelaini, S., Chan, C., Cornelius, V. A. and Margariti, A. (2021). RNA-binding
proteins hold key roles in function, dysfunction, and disease. *Biology* **10**, 366.

Legnini, I., Morlando, M., Mangiavacchi, A., Fatica, A. and Bozzoni, I. (2014). A
feedforward regulatory loop between HuR and the long noncoding RNA linc-MD1
controls early phases of myogenesis. *Mol Cell* **53**, 506–514.

Liu, R., Liu, H., Chen, X., Kirby, M., Brown, P. O. and Zhao, K. (2001). Regulation
of CSF1 promoter by the SWI/SNF-like BAF complex. *Cell* **106**, 309–318.

Love, M. I., Huber, W. and Anders, S. (2014). Moderated estimation of fold change
and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550.

Lynch, M., Chen, L., Ravitz, M. J., Mehtani, S., Korenblat, K., Pazin, M. J. and
Schmidt, E. V. (2005). hnRNP K binds a core polypyrimidine element in the
eukaryotic translation initiation Factor 4E (eIF4E) promoter, and its regulation of
eIF4E contributes to neoplastic transformation. *Mol Cell Biol* **25**, 6436–6453.

Ma, L., Cao, J., Liu, L., Du, Q., Li, Z., Zou, D., Bajic, V. B. and Zhang, Z. (2019).
LncBook: a curated knowledgebase of human long non-coding RNAs. *Nucleic*
Acids Res **47**, D128-D134.

Makeyev, A. V. and Liebhaber, S. A. (2002). The poly(C)-binding proteins: A

multiplicity of functions and a search for mechanisms. *Rna* **8**, 265–278.

Moresi, V., Williams, A. H., Meadows, E., Flynn, J. M., Potthoff, M. J., McAnally,
J., Shelton, J. M., Backs, J., Klein, W. H., Richardson, J. A., et al. (2010).
Myogenin and class II HDACs control neurogenic muscle atrophy by inducing E3
ubiquitin ligases. *Cell* **143**, 35–45.

Nakanishi, K., Sudo, T. and Morishima, N. (2005). Endoplasmic reticulum stress
signaling transmitted by ATF6 mediates apoptosis during muscle development. *J*
Cell Biology **169**, 555–560.

Nakanishi, K., Dohmae, N. and Morishima, N. (2007). Endoplasmic reticulum stress
increases myofiber formation in vitro. *Faseb J* **21**, 2994–3003.

Nostrand, E. L. V., Freese, P., Pratt, G. A., Wang, X., Wei, X., Xiao, R., Blue, S. M.,
Chen, J.-Y., Cody, N. A. L., Dominguez, D., et al. (2020). A large-scale binding
and functional map of human RNA-binding proteins. *Nature* **583**, 711–719.

Paz, I., Kosti, I., Ares, M., Cline, M. and Mandel-Gutfreund, Y. (2014). RBPmap: a
web server for mapping binding sites of RNA-binding proteins. *Nucleic Acids Res*
42, W361–W367.

Sallam, T., Jones, M., Thomas, B. J., Wu, X., Gilliland, T., Qian, K., Eskin, A.,
Casero, D., Zhang, Z., Sandhu, J., et al. (2018). Transcriptional regulation of
macrophage cholesterol efflux and atherogenesis by a long noncoding RNA. *Nat*
Med **24**, 304–312.

Shan, J., Zhang, F., Sharkey, J., Tang, T. A., Örd, T. and Kilberg, M. S. (2016). The
C/ebp-Atf response element (CARE) location reveals two distinct Atf4-dependent,
elongation-mediated mechanisms for transcriptional induction of aminoacyl-tRNA
synthetase genes in response to amino acid limitation. *Nucleic Acids Res* **44**, 9719–
9732.

Statello, L., Guo, C.-J., Chen, L.-L. and Huarte, M. (2021). Gene regulation by long
non-coding RNAs and its biological functions. *Nat Rev Mol Cell Bio* **22**, 96–118.

Tang, S., Xie, Z., Wang, P., Li, J., Wang, S., Liu, W., Li, M., Wu, X., Su, H., Cen, S.,
et al. (2018). LncRNA-OG promotes the osteogenic differentiation of bone
marrow-derived mesenchymal stem cells under the regulation of hnRNPK. *Stem*
Cells **37**, 270–283.

Targoff, I. N., Trieu, E. P. and Miller, F. W. (1993). Reaction of anti-OJ
autoantibodies with components of the multi-enzyme complex of aminoacyl-tRNA

721  synthetases in addition to isoleucyl-tRNA synthetase. *J Clin Invest* **91**, 2556–2564.

722 **Wang, Z., Cui, J., Wong, W. M., Li, X., Xue, W., Lin, R., Wang, J., Wang, P.,**

723  **Tanner, J. A., Cheah, K. S. E., et al.** (2013). Kif5b controls the localization of

724  myofibril components for their assembly and linkage to the myotendinous junctions.

725  *Development* **140**, 617–626.

726 **Wang, Z., Qiu, H., He, J., Liu, L., Xue, W., Fox, A., Tickner, J. and Xu, J.** (2020).

727  The emerging roles of hnRNPK. *J Cell Physiol* **235**, 1995–2008.

728 **Xiong, G., Hindi, S. M., Mann, A. K., Gallot, Y. S., Bohnert, K. R., Cavener, D. R.,**

729  **Whittemore, S. R. and Kumar, A.** (2017). The PERK arm of the unfolded protein

730  response regulates satellite cell-mediated skeletal muscle regeneration. *Elife* **6**,

731  e22871.

732 **Xu, Y., Li, R., Zhang, K., Wu, W., Wang, S., Zhang, P. and Xu, H.** (2018). The

733  multifunctional RNA-binding protein hnRNPK is critical for the proliferation and

734  differentiation of myoblasts. 350–355.

735 **Yano, M., Okano, H. J. and Okano, H.** (2005). Involvement of Hu and heterogeneous

736  nuclear ribonucleoprotein K in neuronal differentiation through p21 mRNA

737  post-transcriptional regulation. *J Biol Chem* **280**, 12690–12699.

738 **Yoon, J.-H., Abdelmohsen, K., Srikantan, S., Yang, X., Martindale, J. L., De, S.,**

739  **Huarte, M., Zhan, M., Becker, K. G. and Gorospe, M.** (2012). LincRNA-p21

740  suppresses target mRNA translation. *Mol Cell* **47**, 648–655.

741 **Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A. H., Tanaseichuk, O.,**

742  **Benner, C. and Chanda, S. K.** (2019). Metascape provides a biologist-oriented

743  resource for the analysis of systems-level datasets. *Nat Commun* **10**, 1523.

Figure 1

Figure 2

Figure 3

A

1		1-612
2		1-952
3		1-1167 (sense)
4		1-1167 (antisense)
5		361-1167
6		596-1167
7		361-952
8		Δccawmcc (Δ660-729)

B

C

Protein: HNRNPK(Hs/Mm)

Position	Motif	Occurrence	Z-score	P-value
274	ccawmcc	gucagucccaugagacccuaaaga**ccuacca**cuaccacaucaggaccacuccagau	2.420	7.76e-03
295	ccawmcc	aagaccuaccacuaccacaucagga**ccacucc**agauuugggcgugugugugugugu	2.435	7.45e-03
674	ccawmcc	auuuucuucauuuacauuucaaaug**cuauccc**aaaaguucccauaccccccuuccc	2.884	1.96e-03
690	ccawmcc	uuucaaaugcuaucccaaaaguucc**cuauacc**cccccuccccccgccougcuccccu	2.986	1.41e-03
695	ccawmcc	aaugcuaucccaaaaguucccuaua**cccccc**ucccccgccougcucccccuaccca	2.957	1.55e-03
699	ccawmcc	cuauucccaaaaguucccuauaccc**cccucc**ccccgccougcuccccuacccacuca	3.246	5.85e-04
700	ccawmcc	uauucccaaaaguucccuauaccccc**ccucc**ccgccougcuccccuacccacucau	3.000	1.35e-03
706	ccawmcc	aaaaguucccuauacccccccucccc**ccgccc**ugcuccccuacccacucauuccccac	3.000	1.35e-03
718	ccawmcc	uacccccccuccccccgccougcucc**cccuaccc**cacucauuccccauucuuggcccug	2.957	1.55e-03
719	ccawmcc	acccccccuccccccgccougcucc**cuaccc**cacucauuccccauucuuggcccugg	3.319	4.52e-04

D

Relative luciferase activity

Figure 4

Figure 5

Figure 6

Figure 7

hnRNPK

Myoparr Atf4

myogenin aminoacyl-tRNA synthetases

hnRNPK

Myoparr Atf4

myogenin aminoacyl-tRNA synthetases

Normal myotubes

Abnormal myotubes

Supplementary Figure 1

Supplementary Figure 2

A

AMINOACYL-tRNA BIOSYNTHESIS

00970 2/4/16
(c) Kanehisa Laboratories

B Cytosolic aminoacyl-tRNA synthetases

Gene Name	Description	log2 Foldchange	padj
Aars (Aars1)	alanyl–tRNA synthetase	1.189688646	6.64E−11
Rars (Rars1)	arginyl–tRNA synthetase	0.717750186	0.000603064
Nars (Nars1)	asparaginyl–tRNA synthetase	1.037402961	1.73E−09
Dars (Dars1)	aspartyl–tRNA synthetase	0.298274648	0.325335115
Cars (Cars1)	cysteinyl–tRNA synthetase	0.872794782	2.21E−05
Qars (Qars1)	glutaminyl–tRNA synthetase	0.081217744	0.864024052
Eprs (Eprs1)	glutamyl–prolyl–tRNA synthetase	0.585228462	0.018216177
Gars (Gars1)	glycyl–tRNA synthetase	1.205275471	3.91E−13
Hars (Hars1)	histidyl–tRNA synthetase	0.413615028	0.200347566
Iars (Iars1)	isoleucine–tRNA synthetase	1.022779598	8.54E−07
Lars (Lars1)	leucyl–tRNA synthetase	0.709821097	0.000114578
Kars (Kars1)	lysyl–tRNA synthetase	0.349181856	0.168606343
Mars (Mars1)	methionine–tRNA synthetase 1	0.900197979	1.01E−05
Farsa	phenylalanyl–tRNA synthetase, alpha subunit	0.200736108	0.620163384
Farsb	phenylalanyl–tRNA synthetase, beta subunit	0.710725176	0.000803764
Sars (Sars1)	seryl–aminoacyl–tRNA synthetase	1.078993838	2.66E−06
Tars (Tars1)	threonyl–tRNA synthetase	0.905000211	8.27E−07
Wars (Wars1)	tryptophanyl–tRNA synthetase	0.878940288	5.01E−06
Yars (Yars1)	tyrosyl–tRNA synthetase	0.772263805	0.000703005
Vars (Vars1)	valyl–tRNA synthetase	0.593311919	0.003371758

C Mitochondrial aminoacyl-tRNA synthetases

Gene Name	Description	log2 Foldchange	padj
Aars2	alanyl–tRNA synthetase 2, mitochondrial	−0.781345711	0.012684624
Rars2	arginyl–tRNA synthetase 2, mitochondrial	0.020716521	0.976825609
Nars2	asparaginyl–tRNA synthetase 2 (mitochondrial)(putative)	0.356286565	0.378947013
Dars2	aspartyl–tRNA synthetase 2 (mitochondrial)	−0.118907946	0.852207722
Cars2	cysteinyl–tRNA synthetase 2 (mitochondrial)(putative)	−0.419664503	0.30297916
Ears2	glutamyl–tRNA synthetase 2, mitochondrial	0.432715891	0.343242224
Hars2	histidyl–tRNA synthetase 2	−0.125988909	0.813766049
Iars2	isoleucine–tRNA synthetase 2, mitochondrial	−0.111305242	0.796505735
Lars2	leucyl–tRNA synthetase 2, mitochondrial	−0.257452115	0.40655708
Mars2	methionine–tRNA synthetase 2 (mitochondrial)	−0.463741055	0.383638447
Fars2	phenylalanine–tRNA synthetase 2 (mitochondrial)	0.24527022	0.65503804
Pars2	prolyl–tRNA synthetase (mitochondrial)(putative)	0.203237774	0.784806846
Sars2	seryl–aminoacyl–tRNA synthetase 2	−0.782886579	0.039853007
Tars2	threonyl–tRNA synthetase 2, mitochondrial (putative)	0.25526333	0.552363803
Wars2	tryptophanyl tRNA synthetase 2 (mitochondrial)	0.172212967	0.789056335
Yars2	tyrosyl–tRNA synthetase 2 (mitochondrial)	0.078853278	0.90065489
Vars2	valyl–tRNA synthetase 2, mitochondrial	0.188587495	0.706779801

Supplementary Figure 3

Supplementary Table 1

Primers for real-time PCR analysis		
Target name	Forward	Reverse
myogenin	CAGTACATTGAGCGCCTACAG	GGACCGAACTCCAGTGCAT
Myoparr (1)	GTGCCCTATCGTCCATGGAG	CACTGACTTCACCTGACCCC
Myoparr (2)	GGGCAGGAAGGGAACAAGAA	CCCCAAATCTGGAGTGGTCC
Xist	CCCGCTGCTGAGTGTTTGAT	ATCCAGGCAATCCTTCTTCTTG
Atf4	AAGGAGGAAGACACTCCCTCT	CAGGTGGGTCATAAGGTTTGG
Aars	ATGGATGCCACTTTAACAGCA	TGGGTCGATTGTGTTTAGGAAGA
Gars	GGAGGCAGCACTTTATCCAAG	TCGGAAGCACTCTCCGTTCT
Iars	CCTCCCTTTGCTACTGGACTG	GGTGAGCGTATCTCGTAACGAT
Nars	GAGCTGTATGTATCTGACCGAGA	AAATGGTGGGAAATGGCTCTTT
Sars	CGGGTGGATAAAGGAGGGGA	TGCCCGAAATCTGCATCGTC
hnRNPK	CCGTACAGACTACAATGCCAG	GCCCTCTTCCAAGGTAGGGAT
Rpl26	GGTCTATGCCCATTCGGAAGG	TCGTTCGATGTAGATGACGTACT

siRNA		
Target name	Target Sequence	
Control siRNA	-	Stealth RNAi siRNA Negative Control, Med GC (Thermo Fisher Scientific)
Myoparr siRNA	GATGGACCCTGTCTGATGCTCTTAA	Thermo Fisher Scientific
hnRNPK siRNA-1	TCCCAAAGATTTGGCTGGATCTATT	MSS205172 (Thermo Fisher Scientific)
hnRNPK siRNA-2	GGAAGTGACTTTGATTGCGAGTTGA	MSS205173 (Thermo Fisher Scientific)