Myosin-driven Nucleation of Actin Filaments Drives Stereocilia Development Critical for

Hearing

Zane G. Moreland[#1,2,3], Fangfang Jiang[#1,2], Carlos Aguilar[#4], Melanie Barzik[#5], Rui Gong[7], Arik

Shams[5], Christian Faaborg-Andersen[5], Jesse C. Werth[5], Randall Harley[5], Daniel C. Sutton[5], Stacey

M. Cole[5], Andrew Parker[4], Susan Morse[4], Elizabeth Wilson[5], Yasuharu Takagi[6], James R. Sellers[6],

Steve D.M. Brown[4], Thomas B. Friedman[5], Gregory M. Alushin[7], Michael R. Bowl[4,8]* & Jonathan E.

Bird[1,2]*

[1]Department of Pharmacology and Therapeutics, and [2]Myology Institute, and [3]Graduate Program in

Biomedical Sciences, University of Florida, Gainesville, FL 32610, USA. [4]Mammalian Genetics Unit,

MRC Harwell Institute, Harwell Campus, OX11 0RD, UK. [5]Laboratory of Molecular Genetics, National

Institute on Deafness and Other Communication Disorders, National Institutes of Health, Bethesda,

MD 20814, USA. [6]Laboratory of Molecular Physiology, National Heart, Lung and Blood Institute,

National Institutes of Health, Bethesda, MD 20814, USA. [7]Laboratory of Structural Biophysics and

Mechanobiology, The Rockefeller University, New York, NY 10065, USA. [8]UCL Ear Institute,

University College London, 332 Gray's Inn Road, London, WC1X 8EE, UK

[#]Authors contributed equally

*Co-corresponding authors

Lead contact: j.bird@ufl.edu

Running title: A myosin motor regulates actin polymerization

Keywords: Actin, myosin, hearing, deafness, hair cell, stereocilia, molecular motor, MYO15, DFNB3

SUMMARY

The assembly and maintenance of actin-based mechanosensitive stereocilia in the cochlea is

critical for lifelong hearing. Myosin-15 (MYO15) is hypothesized to modulate stereocilia height by

trafficking actin regulatory proteins to their tip compartments, where actin polymerization must be

precisely controlled during development. We identified a mutation (p.D1647G) in the MYO15 motor-

domain that initially maintained trafficking, but caused progressive hearing loss by stunting stereocilia

growth, revealing an additional function for MYO15. Consistent with its maintenance of tip trafficking *in*

vivo, purified p.D1647G MYO15 modestly reduced actin-stimulated ATPase activity *in vitro*. Using

ensemble and single-filament fluorescence *in vitro* assays, we demonstrated that wild-type MYO15

directly accelerated actin filament polymerization by driving nucleation, whilst p.D1647G MYO15

blocked this activity. Collectively, our studies suggest direct actin nucleation by MYO15 at the

stereocilia tip is necessary for elongation *in vivo*, and that this is a primary mechanism disrupted in

DFNB3 hereditary human hearing loss.

INTRODUCTION

Cochlear hair cells are the primary transducers of sound in the mammalian inner ear and are fundamental for hearing. Each hair cell assembles approximately 100 individual stereocilia on its apical surface to form a mechano-sensitive hair bundle. Within each bundle, stereocilia are precisely graded into ranks of ascending height and this staircase architecture is critical for mechano-electric transduction (MET) (Barr-Gillespie, 2015). The regulation of stereocilia size is thus central for hearing and the disruption of hair bundle architecture is a common theme in hereditary deafness (Richardson et al., 2011). Stereocilia develop from microvilli by building a highly cross-linked para-crystalline core of actin filaments as an internal scaffold to confer shape and structural rigidity (Vélez-Ortega and Frolenkov, 2019). As stereocilia develop, the actin core thickens and elongates to reach its mature size, necessitating precise control of actin filament polymerization (Tilney et al., 1992; Kaltenbach et al., 1994; Krey et al., 2020). Actin filaments are uniformly polarized within stereocilia with rapidly growing barbed ends orientated towards the tip, the major site of actin polymerization and filament elongation, and also the site of MET (Flock and Cheung, 1977; Tilney et al., 1980; Beurg et al., 2009). Once stereocilia are fully assembled, actin polymerization continues primarily at the tip compartment, indicating an ongoing plasticity throughout adult life (Schneider et al., 2002; Zhang et al., 2012; Drummond et al., 2015; Narayanan et al., 2015). A number of proteins have been identified within the tip compartment that are essential for stereocilia growth, yet the molecular mechanisms governing actin polymerization are unknown. Identification of this mechanism is central to the acquisition and maintenance of hair cell mechano-sensitivity, and also for understanding hearing loss as the clinical manifestation of a cytoskeletal disease.

A key molecule in establishing stereocilia architecture is unconventional myosin 15 (MYO15), encoded by the gene *Myo15* in mice and *MYO15A* in humans. Mutations in *MYO15A* (MIM #602666) cause DFNB3 recessive hereditary hearing loss in humans (Friedman et al., 1995; Wang et al., 1998; Rehman et al., 2016). MYO15 is a member of the myosin superfamily of P-loop ATPases that generate contractile force on actin filaments to power cellular processes such as cytokinesis, endocytosis and vesicular trafficking (Hartman and Spudich, 2012; Houdusse and Sweeney, 2016). Two MYO15 isoforms are produced in auditory hair cells through alternative mRNA splicing (Liang et al., 1999; Anderson et al., 2000; Fang et al., 2015). A shorter isoform (MYO15-2, also known as MYO15-S) consists of the ATPase 'motor domain' and three light chain binding sites that associate with calmodulin-like proteins, in addition to myosin tail homology 4 (MyTH4), Src homology 3 (SH3), and protein 4.1, ezrin, radixin, moesin (FERM) domains (Fig. 1A). A larger isoform (MYO15-1, also known as MYO15-L) is identical to MYO15-2, except for the addition of a 133 kDa N-terminal domain

(Fig. 1A). Both MYO15 isoforms localize to the stereocilia tip compartment where actin polymerization

is concentrated (Schneider et al., 2002; Belyantseva et al., 2003; Drummond et al., 2015; Fang et al.,

2015; Narayanan et al., 2015). The prototypical *Myo15* mutant allele, the *shaker 2* (*Myo15*[sh2]),

prevents both isoforms from accumulating in stereocilia, blocking developmental elongation and

causing profound hearing loss from birth (Probst et al., 1998; Anderson et al., 2000; Belyantseva et

al., 2003; Fang et al., 2015). The use of an isoform-specific *Myo15* null allele revealed additional

functions, with MYO15-2 being necessary for stereocilia developmental elongation, whilst MYO15-1

independently maintains the postnatal size of shorter stereocilia (Fang et al., 2015) that are

structurally plastic in response to MET (Vélez-Ortega et al., 2017; Krey et al., 2020).

The MYO15-2 isoform associates with additional proteins essential for stereocilia elongation,

and by inference, actin polymerization. The 'elongation complex' consists of epidermal growth factor

receptor pathway substrate 8 (EPS8), whirlin (WHRN), G-protein signalling modulator 2 (GPSM2) and

G-protein subunit alpha$_{i3}$ (GNAI3). Similar to MYO15-2, these proteins are concentrated at the tips of

the tallest stereocilia (row 1) (Mburu et al., 2003; Belyantseva et al., 2005; Delprat et al., 2005; Manor

et al., 2011; Zampini et al., 2011; Tarchini et al., 2016; Mauriac et al., 2017; Tadenev et al., 2019).

Individual knock-out mouse mutants of *Eps8*, *Whrn*, *Gpsm2* or *Gnai3* recapitulate the *shaker 2*

phenotype, exhibit short stereocilia and are profoundly deaf (Mburu et al., 2003; Manor et al., 2011;

Zampini et al., 2011; Tarchini et al., 2016; Mauriac et al., 2017; Tadenev et al., 2019). There is strong

evidence that a key function of MYO15-2 is to traffic the elongation complex and concentrate it at the

stereocilia tips. First, elongation complex proteins are absent from the stereocilia of *Myo15 shaker 2*

hair cells, demonstrating their functional dependence upon MYO15-2 *in vivo* (Belyantseva et al., 2005;

Delprat et al., 2005; Manor et al., 2011; Mauriac et al., 2017; Tadenev et al., 2019). Second, EGFP-

tagged MYO15-2 actively traffics elongation complex proteins along filopodia in cell lines (Belyantseva

et al., 2005; Manor et al., 2011; Mauriac et al., 2017). Finally, enzymatic studies of the purified MYO15

ATPase domain reveal kinetic adaptations that enable long-range processive molecular trafficking

(Bird et al., 2014; Jiang et al., 2021). Together, these data support MYO15-2 delivering the elongation

complex to the stereocilia tips, where the elongation complex is hypothesized to regulate actin

polymerization. Despite its central role in promoting stereocilia growth, the specific molecular activity

of the elongation complex remains unknown.

Here, we describe a novel *Myo15* mutant mouse ('*jordan*') that exhibits progressive hearing

loss resulting from a missense substitution in the MYO15 motor domain. In striking contrast with

shaker 2 hair cells, MYO15 and the elongation complex proteins are still delivered to the stereocilia tip

compartment in *jordan* mutant hair cells, but stereocilia fail to elongate properly. These results

questioned the sufficiency of the elongation complex to drive stereocilia growth, and prompted us to

search for another role of MYO15 independent of delivering the elongation complex. We found that

purified MYO15 motor domain protein directly stimulated actin polymerization *in vitro*, and that the

jordan mutation blocked this activity, whilst only moderately affecting its ability to bind and move along

actin filaments. A companion structural study shows that the *jordan* mutation targets the actomyosin

binding interface and interferes with the ability of wild-type MYO15 to regulate structural plasticity

within the actin molecule itself (Gong et al., 2021). Combined, our results argue that MYO15 controls

stereocilia elongation by directly regulating F-actin conformation in stereocilia, and that this critical

activity is independent of the elongation complex. More broadly, our work suggests that in addition to

their classical roles generating force and motility, myosin motors have a physiological role regulating

actin polymerization *in vivo*.

RESULTS

A forward genetic screen identifies *jordan,* a *Myo15* allele causing progressive hearing loss

During a recent phenotype-driven ENU-mutagenesis screen (Potter et al., 2016), the MPC190

cohort (comprising 83 mice) was identified with 10 mice exhibiting severe hearing loss at 3-months of

age (Fig. 1B). A genome scan and single nucleotide polymorphism (SNP) mapping of third generation

(G3) deaf mice found linkage to a 16.7 Mb region on Chromosome 11 (Fig. S1A). Whole-genome

sequencing of a single deaf mouse identified a high confidence homozygous mutation within the

critical interval, consisting of an A-to-G transition at coding nucleotide 4940 of the *Myo15* gene

(ENSMUST00000071880). This variant was confirmed by Sanger sequencing (Fig. S1B) and leads to

the substitution of an evolutionarily conserved aspartate residue with a glycine (p.D1647G) in the

encoded MYO15 protein (Fig. 1C). We named this mutant allele *jordan* (*Myo15^{jd}*).

To confirm that the *Myo15^{jd}* substitution causes hearing loss, we performed a genetic

complementation test utilizing the *shaker-2* (*Myo15^{sh2}*) deafness allele *in trans* (Probst et al., 1998;

Stepanyan et al., 2006). We recorded ABRs from postnatal day 28 (P28) mice and found that

compound heterozygous *Myo15$^{jd/sh2}$* mice had elevated thresholds of >90 decibel sound pressure level

(dB SPL) at all frequencies (Fig. 1D). In contrast, *Myo15$^{jd/+}$* and *Myo15$^{sh2/+}$* littermates had normal

thresholds (< 40 dB SPL) (Fig. 1D). Failure of complementation in *Myo15$^{jd/sh2}$* mice confirms that the

p.D1647G mutation in *Myo15* is the cause of recessive deafness in the *jordan* pedigree.

All reported mutant *Myo15* mouse alleles cause profound deafness (MGI:1261811), measured
from P14 onwards (Probst et al., 1998; Anderson et al., 2000; Stepanyan et al., 2006; Fang et al.,
2015). ABR showed that several *Myo15^{jd/jd}* mice had residual hearing at 3 months (Fig. 1B, red
circles), suggesting a distinct mechanism of hearing loss. We investigated this using longitudinal ABR
measurements. At 4 weeks of age, *Myo15^{jd/jd}* mice had moderate hearing loss with broadband click
ABR threshold of 53 ± 11 dB SPL, compared to normal hearing *Myo15^{+/+}* (26 ± 6 dB SPL) and
Myo15^{+/jd} (24 ± 6 dB SPL) littermates (Fig. 1E). The hearing of *Myo15^{jd/jd}* mice progressively worsened,
with click ABR thresholds of 53 ± 11, 69 ± 9, 80 ± 9 and 81 ± 8 dB SPL at 4, 6, 9 and 12 weeks,
respectively (Fig. 1E,F, Fig. S1C-F). Click ABR thresholds for control *Myo15^{+/+}* and *Myo15^{jd/+}*
littermates ranged between 22 ± 4 and 26 ± 6 dB SPL; showing they did not develop hearing loss with
age (Fig. 1E,F, Fig. S1C-F). To investigate outer hair cell (OHC) function we measured distortion-
product otoacoustic emissions (DPOAEs), and found they were absent in *Myo15^{jd/jd}* mice at 12 weeks,
except for frequencies < 10 kHz, where they were significantly reduced compared to *Myo15^{+/+}* and
Myo15^{+/jd} littermates (Fig. S1G). The absence of DPOAEs shows that OHC function is impaired in
Myo15^{jd/jd} mice. As the only known mouse *Myo15* variant to cause progressive hearing loss, the *jordan*
allele is an important new model to explore the full spectrum of DFNB3 deafness, which presents
heterogeneously as either profound congenital, or progressive hearing loss in humans (Rehman et al.,
2016).
164
Stereocilia do not properly elongate in *jordan* mutant hair cells
To investigate the cellular pathology underlying hearing loss in *Myo15^{jd/jd}* mice, we used
scanning electron microscopy (SEM) to assess the structure of cochlear hair bundles. In wild-type
Myo15^{+/+} IHCs and OHCs at P8, stereocilia were graded into a staircase pattern with three distinct
rows (Fig. 2A,B). By comparison, bundle heights were reduced in *Myo15^{jd/jd}* littermates, whilst still
retaining the staircase architecture (Fig. 2C,D). Quantification of row 1 stereocilia heights showed that
Myo15^{jd/jd} IHCs and OHCs were significantly shorter than *Myo15^{+/+}* controls (Fig. 2G,H). Furthermore,
stereocilia at the lateral edge of *Myo15^{jd/jd}* OHCs, and to lesser extent IHCs, were shorter in height
such that the central stereocilia were tallest (Fig. 2C,D). *Myo15^{jd/jd}* hair cells consistently had 1 or 2
additional rows of stereocilia (Fig. 2C,D) when compared to the three well-defined rows in *Myo15^{+/+}*
hair cells (Fig. 2A,B). The additional row phenotype of *Myo15^{jd/jd}* hair cells was similar to *Myo15^{sh2/sh2}*
hair cells at P8 (Fig. 2E,F), however *Myo15^{jd/jd}* stereocilia lengths were significantly longer when
compared to *Myo15^{sh2/sh2}* hair cells (Fig. 2G,H). We conclude that stereocilia in *Myo15^{jd/jd}* hair cells
elongate more than *Myo15^{sh2/sh2}* hair cells, but fail to reach normal wild-type heights. The *Myo15*

shaker 2 and *jordan* allelic series shows that mutations in the motor domain led to altered stereocilia

heights and explain why *jordan* mice initially have less severe hearing loss than *shaker-2* mice.

To further investigate the progressive component of hearing loss, we next examined hair

bundle morphology at 12 weeks, when *jordan* mice were profoundly deaf (Fig. 1F). Compared with

Myo15$^{+/+}$, IHC bundles of *Myo15$^{jd/jd}$* mice had lost their staircase architecture (Fig. S2A) and were still

significantly shorter (Fig. S2B). Since the staircase was initially present at P8 in *Myo15$^{jd/jd}$* IHCs (Fig.

2D), these data suggested a postnatal resorption of stereocilia. Consistent with this hypothesis,

extremely short stereocilia at the hair bundle periphery were observed at 12 weeks in *Myo15$^{jd/jd}$* OHCs

(Fig. S2C,D). These data argue that the progressive hearing phenotype in the *jordan* mouse was due

to postnatal stereocilia bundle degradation. Overall, we conclude that the *jordan* allele causes hearing

loss by interfering with both initial elongation of stereocilia, and their postnatal maintenance. This was

consistent with the p.D1647G mutation targeting the motor domain of both MYO15 isoforms (Fang et

al., 2015).

MYO15 traffics the elongation complex in *jordan* hair cells

Hair bundle development requires MYO15 dependent trafficking of EPS8, WHRN, GNAI3 and

GPSM2 (the 'elongation complex'), and mutations that prevent MYO15 trafficking (*i.e. shaker-2*) cause

a short hair bundle phenotype (Belyantseva et al., 2005; Manor et al., 2011; Mauriac et al., 2017;

Tadenev et al., 2019). Our finding that stereocilia lengths in the *jordan* mouse were only marginally

longer than the *shaker-2*, led us to hypothesize that MYO15 trafficking was defective in *jordan* hair

cells. We used the previously validated pan-MYO15 antibody PB48 (Fig. 1A), that binds an epitope

common to all isoforms (Liang et al., 1999; Fang et al., 2015), to detect MYO15 in fixed cochleae from

Myo15$^{jd/jd}$ mutants and *Myo15$^{+/jd}$* littermates at P14. As expected, in *Myo15$^{+/jd}$* mice, PB48 labelling

was concentrated at the tips of all stereocilia rows in IHCs (Fig. 3A) (Belyantseva et al., 2003, 2005;

Rzadzinska et al., 2004). We confirmed that PB48 did not label the short stereocilia of *Myo15$^{sh2/sh2}$*

IHCs at P14 (Fig. 3A), consistent with MYO15 being absent from the bundle (Belyantseva et al.,

2003). In stark contrast with the *shaker-2*, we observed PB48 labelling at the tips of IHC stereocilia in

Myo15$^{jd/jd}$ mice at both P7 and P14 (Fig. 3A,B). These data support the *jordan* and *shaker-2* mutations

affecting stereocilia growth by profoundly different mechanisms: the *shaker-2* mutation blocks MYO15

trafficking, whilst the *jordan* mutation allows MYO15 to traffic but prevents stereocilia elongation.

We next investigated if the *jordan* mutation interfered with trafficking of the elongation complex

proteins. To test this, mutant *Myo15$^{jd/jd}$* and control *Myo15$^{+/jd}$* littermate cochleae at P7 were labelled

with validated antibodies to localize WHRN, EPS8, GPSM2 and GNAI3. All of the elongation complex

proteins were concentrated at the tips of the tallest stereocilia (row 1) in control *Myo15*^{+/jd} hair cells

(Fig. 3C), in agreement with previous work (Mburu et al., 2003; Manor et al., 2011; Zampini et al.,

2011; Tarchini et al., 2016; Mauriac et al., 2017). All four proteins in the elongation complex were also

targeted to the stereocilia tips in mutant *Myo15*^{jd/jd} hair cells (Fig. 3C), although the signal-to-noise of

GPSM2 and GNAI3 was lower, indicating a reduced overall accumulation (Fig. 3C). In older animals at

P14, EPS8 and WHRN were still correctly targeted in *Myo15*^{jd/jd} hair cells, however GNAI3 and

GPSM2 were no longer concentrated at stereocilia tips (Fig. S3). These data are consistent with

GPSM2-GNAI3 operating as a distinct complex with MYO15-EPS8 (Tadenev et al., 2019). We

conclude that the initial defective stereocilia elongation in *jordan* hair cells evident at P7 was not due

to gross disruption of elongation complex trafficking by MYO15.

Actin barbed end capping is unaffected in *jordan* mutant hair cells

226   The presence of the elongation complex at the stereocilia tips of *jordan* hair cells suggested

that the proteins known to stimulate growth were present, but insufficient to drive elongation. We

considered whether an overall inhibition of actin polymerization was preventing stereocilia elongation

in *jordan* hair cells. Capping proteins (CAPZ, TWF2) are present within stereocilia and regulate

filament polymerization by blocking barbed end elongation (Peng et al., 2009; Avenarius et al., 2017).

To detect free barbed ends, we monitored the incorporation of TMR-labelled actin monomers in

permeabilized hair cells at P6. In control *Myo15*^{+/jd} hair cells, TMR-actin was concentrated at row 2

stereocilia tips revealing uncapped barbed ends (Fig. 4A). TMR-actin was not consistently detected at

row 1 tips, arguing that barbed ends were inaccessible at that location (Fig. 4A). The *jordan* mutation

did not alter this distribution of free barbed ends and we observed a similar incorporation of TMR-actin

in *Myo15*^{jd/jd} hair cells compared to controls (Fig. 4A). In striking contrast, TMR-actin was incorporated

at the tips of all stereocilia rows in *Myo15*^{sh2/sh2} hair cells, indicating the abnormal presence of barbed

ends in row 1, and an overall loss of hair bundle row identity (Fig. 4A). In summary, *jordan* hair cells

retain row identity at P6, consistent with MYO15 correctly targeting the elongation complex to the

stereocilia tip compartment. These data argue that inappropriate actin filament capping is unlikely the

cause of short stereocilia in *jordan* hair cells. Instead, we hypothesized that a stimulatory factor

independent of the elongation complex was missing.

The *jordan* mutation affects the interaction of MYO15 with actin filaments

245   The *jordan* missense substitution is in the MYO15 motor domain helix-loop-helix (HLH) motif

that forms part of the direct binding interface with the actin filament (Gong et al., 2021). We

hypothesized that a defect in MYO15's interaction with actin might underlie the *jordan* phenotype. To

explore this, we examined MYO15-2 trafficking along filopodia; actin-based structures that protrude

from the periphery of heterologous cells (Belyantseva et al., 2003, 2005). We focused on MYO15-2 as

the isoform responsible for stereocilia growth (Fang et al., 2015). In transfected HeLa cells, EGFP-

tagged wild-type MYO15-2 accumulated at filopodia tips indicating robust anterograde myosin

movement along the filopodia shaft (Fig. 4B). Discrete puncta of MYO15-2 were observed along the

filopodia shaft, arising from retrograde actin filament treadmilling (Belyantseva et al., 2005; Kerber et

al., 2009; Bird et al., 2017). EGFP alone did not accumulate within filopodia, proving this distribution

required active myosin motility (Fig. 4B). In cells expressing the MYO15-2 *jordan* mutant, EGFP was

observed diffusely along filopodia shafts and was not concentrated at filopodia tips (Fig. 4B). This was

qualitatively similar to cells expressing the MYO15-2 *shaker-2* mutant (Fig. 4B), which was previously

shown to not traffic along filopodia (Belyantseva et al., 2005). The inability of the MYO15-2 *jordan*

mutant to traffic within filopodia and accumulate at filopodia tips was unexpected and contrary to

MYO15 protein accumulating at the stereocilia tips of *Myo15$^{jd/jd}$* hair cells (Fig. 3A).

Myosin motors are sensitive to actin filament topology (Nagy et al., 2008; Brawley and Rock,

2009; Reymann et al., 2012; Ropars et al., 2016) and we hypothesized that filopodia might not contain

the appropriate repertoire of actin-binding proteins (ABPs) to support motility. To test this hypothesis,

we used the porcine LLC-PK1-CL4 (CL4) epithelial cell line that generates microvilli and is a more

accurate model for stereocilia (Zheng et al., 2010). In CL4 cells transfected with wild-type EGFP-

MYO15-2, EGFP positive puncta localized at the tips of microvilli (Fig. 4C, orthogonal projections). In

striking contrast to HeLa cells, the MYO15-2 *jordan* mutant also concentrated into microvilli and was

indistinguishable from the wild-type in CL4 cells (Fig. 4C). No microvillar accumulation of the MYO15-

2 *shaker-2* mutant was observed (Fig. 4C). We conclude that whilst *jordan* and *shaker-2* mutants are

both immobile within filopodia, the *jordan* mutant can still actively concentrate in microvilli. These data

mimic our findings from hair cells *in vivo*, and support a change in actin binding as being central to the

stereocilia growth defect phenotype observed in *jordan* hair cells.

The ATPase activity of MYO15 is altered by the *jordan* mutation

To study the interaction between MYO15 and actin filaments in more detail, we characterized

the influence of the *jordan* mutation upon ATPase mechanochemistry (Bird et al., 2014; Jiang et al.,

2021). Motor domain proteins were expressed in *S. frugiperda* (*Sf*9) insect cells and purified by

chromatography (Fig. 5A,C). In contrast to the *shaker-2* variant, which aggregated within *Sf*9 cells,

both wild-type and *jordan* motor domains were soluble. Using size exclusion chromatography (SEC),

we found that FLAG-purified *shaker-2* motor domain eluted close to the void volume (Fig. 5B). In

contrast, the *jordan* motor domain eluted at a similar delay volume to the wild-type (Fig. 5B), which is

monomeric (Bird et al., 2014). We conclude that the *jordan* variant does not affect folding, whilst the

shaker-2 caused misfolding, consistent with the mutation being within the ATPase transducer core

(Gong et al., 2021).

To test for differences in enzymatic activity between wild-type and *jordan* motor domain

variants (Fig. 5A), we measured steady-state ATP hydrolysis using an enzyme-linked NADH assay.

The basal ATPase activity (k_{basal}) of the wild-type motor domain was 0.06 ± 0.01 s^{-1}, measured without

F-actin. The addition of F-actin caused a 97-fold increase in ATPase activity to an extrapolated

maximum catalytic rate (k_{cat}) of 5.8 ± 0.2 s^{-1} (Fig. 5D). Half-maximal activation of the ATPase activity

(K_{ATPase}) was reached at 29.1 ± 2.1 µM F-actin, as previously reported (Bird et al., 2014; Jiang et al.,

2021). Using identical assay conditions, the basal ATPase rate of the *jordan* variant was unchanged at

k_{basal} = 0.07 ± 0.01 s^{-1}. Strikingly, there was a reduced 7-fold maximal activation to k_{cat} = 0.87 ± 0.04 s^{-1}

(Fig. 5D). Furthermore, half-maximal ATPase activation was increased to K_{ATPase} = 114.3 ± 8.2 µM

actin, signifying a 4-fold reduction in the apparent actin affinity compared to wild-type.

The mechanical activity of wild-type and *jordan* motor domains was measured using an *in vitro*

gliding filament assay, where actin filaments are propelled across a microscope cover glass

functionalized with motor domain protein (Sellers et al., 1993). Due to the reduced apparent affinity of

the *jordan* motor domain for actin, we lowered the salt concentration to 10 mM KCl in these assays.

The wild-type motor domain robustly propelled actin filaments at 473 ± 67 nm·s^{-1} (Fig. 5E), consistent

with previous data (Bird et al., 2014). In contrast, the *jordan* mutant moved filaments at 216 ± 71 nm·s^{-1}

^{1}, a 2-fold reduction from the wild-type velocity (Fig. 5E). Overall, we conclude that whilst the *jordan*

mutation caused a significant ATPase defect and decreased the motor domain's apparent affinity for

actin, the motor domain was still mechanically active. Our data further establish a functional

correlation between motor domain activity and the severity of hearing loss in a *Myo15* allelic series.

The *jordan* motor domain retained partial activity consistent with this mutation causing intermediate

hearing loss, whilst the *shaker-2* motor domain was misfolded and associated with the most severe

phenotype.

The MYO15 motor domain directly stimulates actin polymerization

Our *in vitro* analysis of purified MYO15 motor activities were consistent with the *jordan* mutant

maintaining the ability to traffic the elongation complex and accumulate at stereocilia tips. As

stereocilia nevertheless fail to grow normally in *jordan* hair cells, we hypothesized that independent of

molecular trafficking, MYO15 had an additional function at the stereocilia tip critical for elongation.

Classic biochemical studies of muscle myosin have shown its motor domain (*e.g.* subfragment-1, S1)

can directly stimulate actin polymerization *in vitro*; however, the physiological relevance of this activity

in vivo, if any, is unknown (Yagi et al., 1965; Miller et al., 1988; Fievez and Carlier, 1993; Lheureux et

al., 1993). We hypothesized that the MYO15 motor domain could directly stimulate actin

polymerization to drive stereocilia elongation, and that this critical activity is disrupted by the *jordan*

deafness mutation.

We tested this hypothesis using pyrene-conjugated globular actin (G-actin) monomers that

increase in fluorescence as they polymerize into filamentous actin (F-actin) (Cooper et al., 1983). As a

control, 2 µM G-actin was polymerized with 1x KMEI (50 mM KCl, 1 mM $MgCl_2$, 1 mM EGTA, 10 mM

imidazole) while monitoring pyrene fluorescence. F-actin assembled with an initial lag phase

representing the kinetically unfavourable nucleation step (Fig. 6A, red line). When the reaction was

repeated with 1 µM wild-type motor domain + 2 µM G-actin + KMEI, actin polymerization initially

followed the same trajectory (Fig. 6A, blue line). However, following a delay of ~ 400 seconds, an

inflection was observed where the rate of actin polymerization reduced momentarily (Fig. 6A, blue

line), before rapidly increasing and reaching steady-state (Fig. 6A,B). We tested if MYO15-stimulated

actin polymerization was sensitive to the *jordan* mutation. Under identical experimental conditions

using 1 µM *jordan* MYO15 motor domain, the actin polymerization rate was initially indistinguishable

from the control reaction. However, following an inflection at ~ 600 seconds, the actin polymerization

rate decreased below that of the G-actin alone control (Fig. 6A). We conclude that the wild-type

MYO15 motor domain stimulated actin polymerization, whilst the deafness-causing *jordan* mutant

blocked this activity.

With either wild-type or *jordan* motor domains, we observed a delay before the actin

polymerization rate deviated from the control, with the delay longer for *jordan* (~ 600s) versus wild-

type (~ 400s). We hypothesized this difference arose from residual ATP (70 µM) being introduced

from the G-actin storage buffer; ATP binding would prevent the motor domain from occupying the

strongly actin bound (rigor) state. Furthermore, we expected this residual ATP to be hydrolysed faster

by the wild-type motor domain, compared with the *jordan* mutant (Fig. 5D). To test this, we repeated

the pyrene polymerization assay in the absence of free ATP. To ensure G-actin monomers were ATP-

bound, G-actin was desalted immediately prior to use to remove free ATP from solution. Using this

approach, we measured the polymerization of 2 µM G-actin + KMEI in the absence of free ATP (Fig.

6C) and observed kinetics comparable to before (Fig. 6A). Strikingly, pyrene fluorescence immediately

increased upon addition of 1 µM wild-type motor to the reaction, and rapidly saturated (Fig. 6C) with a

significantly reduced half-time of 124 ± 12 s, compared with 1091 ± 202 s for G-actin + KMEI alone

(Fig. 6D). While addition of 1 µM *jordan* motor domain to 2 µM G-actin + KMEI reduced the overall

extent of actin polymerization (Fig. 6C), the half-time to saturation was 1000 ± 65 s and not

significantly different to G-actin + KMEI alone (Fig. 6D). We conclude that the wild-type MYO15 motor

domain strongly accelerated actin filament polymerization in the absence of ATP, implicating the rigor

state as critical for this activity. Combined with experiments from our companion study showing that

ADP-bound MYO15 also stimulated polymerization (Gong et al., 2021), these data identify the strong

actin-binding states as the key MYO15 ATPase intermediates that stimulate actin polymerization.

The *jordan* deafness mutation inhibits *de novo* nucleation of actin filaments

To directly visualize how the MYO15 motor domain accelerates polymerization, we performed

actin polymerization assays using total internal reflection fluorescence microscopy (TIRFM) where the

elongation of individual filaments can be tracked (Fujiwara et al., 2002). A control time-lapse of 1 µM

G-actin + KMEI revealed a slow rate of filament nucleation, with short polymers attaching to surface

and elongating (Fig. 7A, top row). We repeated the experiment with 1 µM wild-type motor domain and

observed a large burst in filament density (Fig. 7A, middle row) that was statistically significant at 480

seconds when compared to the actin-alone control (Fig. 7B). In contrast, when polymerization

experiments were repeated with 1 µM *jordan* motor domain (Fig. 7A, bottom row), there was no

increase in actin filament density above the actin-alone control (Fig. 7C). Using kymographs to track

filament barbed ends, we found that the motor domain could influence actin filament elongation rates.

In control experiments, barbed ends elongated at 16.2 ± 2.6 nm·s^{-1} (Fig. 7C,D), consistent with

previous data using TMR-conjugated G-actin on Cys374 (Kuhn and Pollard, 2005). Elongation rates

were significantly reduced by the addition of either wild-type or *jordan* motor domain, to 13.3 ± 4.3

nm·s^{-1} and 13.4 ± 3.5 nm·s^{-1}, respectively (Fig. 7C,D). The decrease in elongation rate was dependent

upon timing of the filament burst (Fig. 7B). When data were binned as pre-burst (< 320 s), the addition

of wild-type or *jordan* motor domains did not significantly alter elongation rates compared to the

control (Fig. 7E). In contrast, the presence of either wild-type or *jordan* motor domain post-burst (> 720

s) both significantly, and equally, reduced elongation rates below the actin alone control (Fig. 7F).

Nucleation by wild-type MYO15 took an average of 480 seconds to be significantly increased

above the spontaneous nucleation rate of G-actin + KMEI alone (Fig. 7B). We considered whether the

delayed production of short filaments was caused by MYO15 severing actin filaments that nucleated

spontaneously. To test this hypothesis, we repeated our experiments in the absence of free ATP to

force the motor domain into rigor binding and accelerate polymerization. Free ATP was desalted from

1 µM G-actin monomers; this did not affect barbed-end elongated rates when polymerized by KMEI

(Fig. 7D,I). As expected, when 1 µM wild-type motor domain was included in the reaction, there was a

potent nucleation of short actin filaments observed within 90 s (Fig. 7G,H), confirming acceleration of

nucleation when the motor domain was forced into rigor. Under identical ATP free conditions, 1 µM

jordan motor domain did not nucleate additional filaments compared with the G-actin + KMEI control

(Fig. 7G,H). Similar to experiments where ATP was present, but presumably hydrolyzed (Fig. 7F, >

720 s), we saw that both wild-type and *jordan* motor domains significantly reduced barbed-end

elongation rates (Fig. 7I). We conclude that the MYO15 motor domain exerts multiple direct effects on

actin polymerization, including catalyzing *de novo* filament nucleation and reducing barbed end

elongation rates in a nucleotide-dependent fashion. Our data show that MYO15-induced actin

nucleation is a key process targeted by the *jordan* deafness mutation, and argue that this activity is

critical for stereocilia elongation and hearing.

DISCUSSION

Plasticity of the stereocilia actin core is central to hair bundle development, mechano-

sensitivity and hearing. Existing evidence that MYO15 moves along stereocilia actin filaments has led

to a cargo trafficking model, where the elongation complex is delivered to the stereocilia tips and

provides the actin regulation machinery necessary for elongation. Here, we uncover a previously

unknown activity of MYO15 that requires the cargo-trafficking model to be re-evaluated. *In vitro*, we

show that MYO15 can directly stimulate actin filament nucleation and that this activity is inhibited by

the *jordan* mutation, providing key evidence this activity is functionally linked to deafness pathology. *In*

vivo, the *Myo15 jordan* mutation stunted stereocilia growth, leading to an early-onset, progressive

hearing loss in mice. The effect of the *jordan* mutation to inhibit MYO15-induced actin nucleation *in*

vitro, and block stereocilia elongation *in vivo*, provides the functional link between these two

processes. We propose that in addition to trafficking key molecular cargos, MYO15 directly catalyses

actin filament nucleation at the stereocilia tip, and that both these processes combined are required

for normal stereocilia growth.

Using purified proteins and reconstituted assays, the MYO15 motor domain exerted multiple

effects upon actin filament polymerization *in vitro*. The motor domain: 1) nucleated actin filaments *de*

novo, and 2) slowed the elongation rate of existing actin filaments. No additional proteins were

required, demonstrating that these properties were intrinsic to the purified actomyosin system. The

jordan mutation completely blocked MYO15-induced actin filament nucleation, whilst filament
elongation rates were unaffected relative to wild-type MYO15. We conclude that MYO15's nucleation
activity is the key polymerization effect disrupted by the *jordan* deafness mutation. Similar to MYO15,
muscle myosin was shown in classic biochemical experiments to stimulate actin filament nucleation *in*
vitro (Yagi et al., 1965; Miller et al., 1988; Fievez and Carlier, 1993; Lheureux et al., 1993). The
MYO15 motor domain shares a common structural fold with muscle myosin (Gong et al., 2021), and
although the mechanism of muscle nucleation remains controversial (Lheureux et al., 1993), we
believe by homology to muscle that MYO15 nucleates actin through stabilizing inter-subunit contacts
that establish the protofilament (Fievez et al., 1997a, 1997b). In our companion study (Gong et al.,
2021), the MYO15 motor domain is further shown to influence structural plasticity within the DNaseI-
binding loop (D-loop), a sub-domain of actin that regulates the monomer to polymer transition
(Dominguez and Holmes, 2011). Combined, our data argue that nucleation is a shared function for at
least some members of the myosin superfamily, and that myosin motors can directly influence
cytoskeletal plasticity in addition to their canonical role generating force.
431
A cellular function for myosin-driven actin nucleation has, to the best of our knowledge, not
previously been reported in any tissue. Our study provides evidence that MYO15-driven nucleation
contributes to hair cell function *in vivo*, and furthermore is a cause of disease when disrupted. Actin
nucleation by MYO15 appears distinct from other nucleation factors, such as formins, Arp2/3, Spire
and Cobl (Machesky et al., 1994; Mullins et al., 1997, 1998; Sagot et al., 2002; Quinlan et al., 2005;
Ahuja et al., 2007). MYO15-driven nucleation was nucleotide-sensitive, and strongly promoted
nucleation under rigor (nucleotide-free) conditions, or when bound to ADP (Gong et al., 2021). Both
conditions populate the motor domain into strong actin binding states (Jiang et al., 2021). Conversely,
the presence of ATP in assays prevented nucleation and we hypothesize this is due to the motor
domain populating a weakly actin-bound state (Jiang et al., 2021). The nucleotide-sensing properties
of MYO15 may therefore couple local concentrations of ATP and ADP to actin polymerization. We
speculate this activity could form the basis of a tuning mechanism, where dynamic changes in
nucleotide concentrations control the height of developing stereocilia. The concentrations of ATP and
ADP in stereocilia are unknown, and whilst a creatine kinase ATP generation system has been
described in avian hair cells (Shin et al., 2007), its activity in mammals is unclear (Krey and Barr-
Gillespie, 2019). In addition to nucleotide state, MYO15-driven nucleation *in vitro* was observed at
high protein concentrations equimolar with actin, in agreement with data from muscle myosin (Miller et
al., 1988). The concentration of MYO15 in hair cells is unknown, however micromolar quantities may
be present at the stereocilia tips, especially as MYO15 actively traffics and self concentrates into this

subcellular compartment (Belyantseva et al., 2003). Other unconventional myosin motors critical for

hearing, including MYO1C, MYO3A, MYO6 and MYO7A (Avraham et al., 1995; Gibson et al., 1995;

Gillespie and Corey, 1997; Hasson et al., 1997; Holt et al., 2002; Salles et al., 2009; Ebrahim et al.,

2016), also concentrate in specific stereocilia sub-domains and it is exciting to consider whether they

also directly regulate actin polymerization as part of their cellular function.

In parallel with identifying MYO15 as an actin nucleator, we used the *jordan* mutant mouse to

probe sufficiency of the elongation complex to drive stereocilia elongation. Each protein in the

elongation complex is critical for growth, but existing models do not explain their molecular activty,

except that they must be delivered to the stereocilia tip via MYO15-driven trafficking. This model is

based on the *shaker-2* mouse, where stereocilia growth is inhibited and both MYO15 and the

elongation complex are absent from the hair bundle (Belyantseva et al., 2005; Delprat et al., 2005;

Manor et al., 2011; Zampini et al., 2011; Mauriac et al., 2017; Tadenev et al., 2019). Our data show

that MYO15-driven actin nucleation and elongation complex trafficking activities are simultaneously

lost in the *shaker-2*, leaving their relative contributions unknown. By contrast, MYO15-driven trafficking

of the elongation complex is initially (largely) preserved in the short stereocilia of the *jordan* mouse.

These data argue that the elongation complex proteins are necessary, but not sufficient for driving

stereocilia growth, and that MYO15-nucleation activity is critical *in vivo*. We cannot exclude the

possibility of reduced MYO15 trafficking in *jordan* hair cells; indeed, gliding filament velocities of the

mutant motor were ~ 50% of the wild-type *in vitro*. However, there is no evidence that the

concentration of MYO15 at the stereocilia tips determines stereocilia length. Neither the elimination of

MYO15-1 from shorter stereocilia rows (Fang et al., 2015), nor the over-expression of MYO15-2 alters

initial stereocilia lengths (Belyantseva et al., 2003, 2005). We also considered whether the *jordan*

mutation might interfere with MYO15 binding to another protein critical for stereocilia growth. Our

companion study (Gong et al., 2021) localizes the *jordan* missense substitution to the actomyosin

interface, and shows a direct structural effect within the actin monomer itself. Combined with our

study, these findings strongly suggest that MYO15 directly accelerates actin nucleation at the

stereocilia tip by structurally regulating the actin molecule.

Our study raises new questions concerning the stability and plasticity of the stereocilia

cytoskeleton, processes that must be tightly controlled for mechano-transduction and lifelong hearing

(Vélez-Ortega and Frolenkov, 2019). Stereocilia actin filaments are unidirectionally polarized with their

fast-growing barbed ends orientated towards the tip compartment, where new actin monomers are

incorporated (Flock and Cheung, 1977; Tilney et al., 1980; Schneider et al., 2002; Drummond et al.,

2015). Although actin capping proteins also contribute to stereocilia architecture (Peng et al., 2009;

Avenarius et al., 2017), we find that excessive barbed end capping is not the cause of short stereocilia

in *Myo15* mutant mice. Given the presence of barbed ends, why might actin nucleation be required to

extend pre-existing filaments at stereocilia tips? One possible explanation is that stereocilia elongation

occurs through direct end-to-end annealing of short actin polymers nucleated by MYO15, rather than

through addition of individual actin monomers at the barbed end. Pre-formed actin filaments can

anneal into longer filaments *in vitro* (Murphy et al., 1988; Kinosian et al., 1993; Sept et al., 1999), and

more recent *in vitro* studies have detected filament extension through incorporation of short polymers,

in addition to the dominant mode of monomer addition (Young et al., 2018). Interestingly, endocytosis

in yeast is proposed to use end-to-end actin filament annealing rather than monomer polymerization

(Okreglak and Drubin, 2010). Though speculative in hair cells, the extension of the stereocilia actin

core using short polymer annealing may allow for rapid changes in stereocilia length that are likely

needed to continually maintain the sensitivity of the MET machinery (Vélez-Ortega et al., 2017; Krey

et al., 2020).

Given the potent ability of MYO15 to nucleate actin filament polymerization *in vitro*, we infer

this activity must be tightly regulated in hair cells. There is extensive evidence for intramolecular

regulation throughout the myosin superfamily (Heissler and Sellers, 2016). For example, myosin 5

(MYO5A) exists in an autoinhibited conformer where the globular tail domain binds and inhibits motor

domain activity; binding of melanophilin (MLPH) to MYO5 releases this autoinhibition to activate the

motor domain (Li et al., 2005; Liu et al., 2006; Thirumurugan et al., 2006; Sato et al., 2007; Sckolnick

et al., 2013). Similar autoinhibitory regulation mechanisms have been described for other members of

the MyTH4 - FERM myosin family, *e.g.*, MYO7A and MYO10 (Umeki et al., 2009, 2011; Yang et al.,

2009; Sakai et al., 2011). We hypothesize that part of the elongation complex function may be to bind

MYO15 and shift the equilibrium between active and inhibited states. Intramolecular regulation may

also explain how different MYO15 isoforms independently exert their effects upon hair bundle

architecture (Fang et al., 2015). MYO15-1 and MYO15-2 possess an identical motor domain

competent to regulate actin polymerization, but differ by the addition of a 133 kDa N-terminal domain

(Fig. 1A). How the N-terminal domain affects the actin nucleation activity of the motor domain is an

important future question and will help uncover how the MYO15-1 isoform controls actin

polymerization in shorter stereocilia rows with active MET (Fang et al., 2015). In conclusion, we reveal

a new function for MYO15 and argue that defective nucleation of actin polymerization interferes with

stereocilia architecture and causes progressive hearing loss. We speculate that defective actin

nucleation by MYO15 is a central molecular pathology underlying DFNB3 hereditary hearing loss in

humans.

MATERIALS AND METHODS

Mice

Pedigree MPC190 was identified from a phenotype-driven mutagenesis screen undertaken at

the MRC Harwell Institute (Potter et al., 2016). Briefly, *N*-ethyl-*N*-nitrosourea (ENU) mutagenized

C57BL/6J males were mated with wild-type 'sighted C3H' (C3H.Pde6b$^+$) females. Resulting G_1 males

were crossed with C3H.Pde6b$^+$ females to produce G_2 females, which were screened for the *Cdh23^{ahl}*

allele. *Cdh23$^{+/+}$* G_2 females were backcrossed to their G_1 fathers to generate recessive G_3 pedigrees,

which entered a longitudinal phenotyping pipeline that included click box and ABR tests to assess

auditory function (Hardisty-Hughes et al., 2010). DNA from mice exhibiting hearing loss, and normal

hearing pedigree mates, was prepared from ear biopsies and used for linkage mapping utilizing the

Illumina GoldenGate Mouse Medium Density Linkage Panel (Gen-Probe Life Sciences Ltd, UK), which

identified a critical interval on chromosome 11. DNA was extracted from mouse MPC190/2.18a and

subject to whole genome sequencing employing the Illumina HiSeq platform (Oxford Genomics

Centre, Wellcome Trust Centre for Human Genetics). Subsequent alignment to the reference genome

(NCBIM38/mm10) identified a homozygous, non-synonymous coding lesion in the *Myo15* gene.

At the MRC Harwell Institute, mice were housed and maintained at the Mary Lyon Centre

under conditions outlined in the Home Office Code of Practice, with all animal procedures licenced by

the Home Office under the Animals (Scientific Procedures) Act 1986, UK and approved by the local

Ethical Review Committee (PBF9BD884 to MRB). At MRC Harwell, *jordan* mice were crossed to

C57BL/6N (*Cdh23$^{753A>G}$*) 'repaired' mice (Mianné et al., 2016) until congenic. Concurrently, *jordan*

mice were imported to the NIH and the University of Florida (UF) and maintained on a 'sightless C3H'

(C3H.Pde6b^{rd1}) background. Animal procedures were approved the Institutional Animal Care and Use

Committees (IACUC) at UF (#201910739 to JEB) and at the NIH (#1263-15 to TBF). Genomic DNA

from mouse tail biopsies was used as template in a PCR reaction with primers (5'-

CAGGAGGAGTACATCCGGG-3', 5'-AGACCACAGAAGTATCTGGGTCTT-3'). The resulting 161 bp

amplicon was analyzed by *MlsI* endonuclease digestion. Resulting restriction fragments lengths

unambiguously detected wild-type (161 bp) and mutant (116 bp + 45 bp) alleles.

Auditory phenotyping + behavioral testing

Auditory Brainstem Response (ABR) were recorded as previously described (Dunbar et al.,

2019). Briefly, mice were anaesthetised via intraperitoneal injection with ketamine hydrochloride (100

mg·kg^{-1}) and xylazine (10 mg·kg^{-1}). Anesthetized mice were placed inside a sound-attenuated

chamber (ETS-Lindgren) and recording electrodes (Grass Telefactor F-E2-12) placed sub dermally

over the vertex (active), right mastoid (reference), and left flank (ground). ABR responses were

collected using TDT system III hardware and BioSig software (Tucker Davis Technology). Stimuli were

presented free field from a ES1 transducer (TDT) calibrated at a distance of 1 cm from the right ear.

Stimuli consisted of 0.1 ms broadband clicks or 7 ms tone-bursts at 8 kHz, 16 kHz and 32 kHz. Each

stimulus was presented at a maximum 90 dB SPL, followed by decreasing steps of 5 - 10 dB SPL until

no replicable response peaks were observed. ABR operators were blind to genotype.

Distortion Product Oto-Acoustic Emissions (DPOAE) were recorded as a terminal procedure in

12 weeks old mice, as described (Dunbar et al., 2019). Mice were anaesthetised with a modified

ketamine/xylazine solution (see ABR), with the addition of acepromazine (2 mg·ml^{-1}, 8% v/v).

Anesthetized mice had a section of the pinna removed to access the external auditory meatus. Mice

were placed inside a sound-attenuated chamber (ETS Lindgren), and the DPOAE probe assembly

was inserted into the ear canal. In-ear calibration was performed before each test. An ER10B+ low-

noise probe microphone (Etymotic Research) was used to measure DPOAEs near the tympanic

membrane. Tone stimuli were presented via separate MF1 (Tucker Davis Technology) speakers, with

f1 and f2 at a ratio of f2/f1 = 1.2 (L1 = 65 dB SPL, L2 = 55 dB SPL). The f1 and f2 tones were

presented continuously in specific tone-bursts between 8 and 32 kHz.

Scanning electron microscopy

Inner ears were dissected and fixed overnight at 4ºC in 0.1 M phosphate buffer, 2.5%

glutaraldehyde (Sigma-Aldrich). Post-fixed ears were decalcified in 4.3% EDTA in phosphate buffer at

4ºC, before dissecting out the sensory epithelium. Samples were processed with alternating 1%

osmium tetroxide (Agar Scientific) in 0.1 M sodium cacodylate (Sigma-Aldrich) and 1%

thiocarbohydrazide (Sigma-Aldrich) treatments. Osmicated samples were dehydrated in graded

ethanols (25% to 100%) at 4ºC and stored in 100% acetone until critical point drying with liquid CO_2

(EM CPD300, Leica Microsystems Ltd.). Samples were mounted onto stubs using silver paint (Agar

Scientific), sputter-coated with platinum (Q150R S, Quorum Technologies) and visualised with a

scanning electron microscope (JSM-6010LV, JEOL).

For morphometric analyses, image pairs from the middle cochlear turn IHC and OHC bundles

were captured with a 5° tilt angle difference at a constant working distance of 20 µm. Stereocilia from

at least two bundles per animal were measured using ImageJ (https://imagej.nih.gov), with a minimum

of 14 stereocilia for each condition. Length measurements were taken from the rear aspect of the hair

bundle, so that the length of the tallest stereocilia (row 1) could be measured from the cuticular plate

insertion to the tip. Estimates for actual stereocilia length were calculated using a pseudo-eucentric

tilting approach (Bariani et al., 2005). A single measure x_1 (length of stereocilium) was taken from the

first image and measured again (x_2) on the corresponding 5°-tilted image pair. Perpendicular counter-

measures (y_1 and y_2), matched to (x_1 and x_2) were also recorded. Equation 1 was used to estimate

uncertainty (ζ) due to plane rotation. Using the uncertainty estimate ζ from Equation 1, every pair of

tilted measures (x_1 and x_2) was processed using Equation 2 to obtain a close approximation (ξ) of the

true length of each stereocilia measured.

$$\zeta = \frac{(\Delta y)cos\Delta\varphi + (2y_1(y_1 - \Delta y)/d)sin\Delta\varphi}{(1 + y_1(y_1 - \Delta y)/d^2 sin\Delta\varphi) + \left(\frac{\Delta y}{d}\right)\cos(2\Delta\varphi)} \qquad\qquad \xi = \frac{2d - 2\zeta cos\Delta\varphi}{\dfrac{d}{x_1} + \dfrac{d}{x_2}}$$

(Equation 1)   (Equation 2)

Where: ζ = uncertainty estimate, ξ = estimate of true size of structure of interest, $x_{1,2}$ = tilted

paired-measures of structure of interest, $y_{1,2}$ = perpendicular counter-measures to measures $x_{1,2}$, Δy =

arithmetic difference of counter-measures y_1 and y_2, $\Delta\varphi$ = tilting angle (5°), d = working distance (20

µm).

In addition, low-magnification (1500 - 2000X) fields of mid-turn sensory epithelium were

imaged and the OHC bundles counted and visually categorised as either: 'Intact', 'Abnormal /

Damaged' where the bundle had an abnormal shape or was missing stereocilia, or 'Missing Bundle'

where a cuticular plate was observed with no stereocilia. Cochleae from four different animals per

genotype were examined, with a total of 1344 wild type and 972 *jordan* OHC bundles.

Whole mount immunofluorescence and confocal microscopy

Mouse inner ears were fixed in 4% paraformaldehyde (EMS Diasum) in PBS for 2 hours at

room temperature, washed in PBS, and then micro-dissected to isolate the cochlear sensory

epithelium. Samples were permeabilized in 0.5% (v/v) Triton X-100 in PBS for 30 minutes at RT,

followed by blocking in 5% normal goat serum (Sigma-Aldrich), 2% bovine serum albumin (Fraction V,

Roche) in PBS for 1 hour at RT. Primary antibodies were diluted in blocking solution and incubated

with samples at 4°C overnight. Primary IgG antibodies used were: PB48 rabbit anti-MYO15 (Liang et

al., 1999), HL5136 rabbit anti-WHRN (Belyantseva et al., 2005), mouse anti-EPS8 (#610143, BD

Biosciences), rabbit anti-GPSM2 (Ezan et al., 2013) and rabbit anti-GNAI3 (#G4040, Sigma-Aldrich).

After washing in PBS, samples were labelled with Alexa Fluor 488 conjugated anti-IgG secondary

antibodies (Life Technologies) diluted in blocking solution for 1 hour at RT. Samples were co-labelled

with rhodamine phalloidin (Life Technologies) and mounted with high-precision #1.5 cover glass

(Zeiss) using Prolong Gold (Life Technologies). Images were captured using a 63x objective (1.4 NA,

plan apochromat, Zeiss) and a laser scanning confocal microscope (LSM780, Zeiss).

Actin barbed end incorporation assay

Cochleae from *shaker-2* or *jordan* mice were dissected at P6 in Leibowitz L-15 media (Life

Technologies) and adhered to the inside surface of a petri dish lid (Greiner CellStar 35 x 10 mm). The

stria vascularis was left intact, as described (Xiong et al., 2014). Cochleae were cultured for 6 hours in

DMEM / F-12 (Life Technologies) supplemented with 1% FBS (Atlanta Biologicals) and 1.5 $\mu g \cdot mL^{-1}$

ampicillin at 37°C, 5% CO_2. Cochlea were incubated for 2 minutes at RT in 0.5 μM TMR-actin

(Cytoskeleton) diluted in labelling buffer (concentrations in mM): HEPES (20), KCl (138), EGTA (3),

$MgCl_2$ (4), ATP (1), DTT (0.5), 1% (w/v) bovine serum albumin (BSA), 0.2 $mg \cdot mL^{-1}$ saponin, pH 7.5.

Cochleae were immediately fixed for 30 mins at RT in 4% paraformaldehyde (EMS Diasum) diluted in

PHEM buffer (concentrations in mM): PIPES (60), HEPES (25), EGTA (10), $MgCl_2$ (2), sucrose (120),

pH 7.0. Fixed cochleae were permeabilized in 0.5% (v/v) Triton X-100 in PBS for 30 minutes at RT,

labelled with Alexa 488 phalloidin (Life Technologies), and mounted using a high-precision #1.5 cover

glass (Zeiss) with Prolong Gold (Life Technologies). Confocal microscopy images were captured as

described above.

Expression of EGFP-MYO15-2 in mammalian cells

The pEGFP-C2-Myo15-2 and pEGFP-C2-Myo15-2(sh2) plasmids expressing an N-terminal

EGFP fusion with the mouse MYO15 isoform 2 coding sequence (NP_874357.2) were previously

reported (Belyantseva et al., 2003). The pEGFP-C2-Myo15-2(jd) plasmid was generated using site

directed mutagenesis (QuikChange II, Agilent) to introduce the jordan (c.4940A>G) non-synonymous

substitution. All expression plasmids were verified by Sanger sequencing and prepared as endotoxin-

free, transfection grade DNA (NucleoBond Xtra Maxi EF, TakaraBio). HeLa cells (#CCL2) were

obtained as an authenticated, low passage stock from the American Type Culture Collection (ATCC).

LLC-PK1-CL4 (CL4) cells were a kind gift from Dr. James Bartles at Northwestern University. HeLa

and CL4 cells were cultured in high-glucose DMEM (#11995, Life Technologies), supplemented with

1x GlutaMAX (Life Technologies) and 10% FBS (Atlanta Biologicals), and maintained at 37°C, 5%

CO_2. Transfection of HeLa cells or CL4 cells was performed using Lipofectamine 3000 (Life

Technologies) according to the manufacturer's protocol. pEGFP-C2 (Clontech) was used as an empty

vector control. Cells were plated onto fibronectin (10 µg·mL^{-1}) coated glass bottom culture dishes

(#1.5, MatTek Corp) and allowed to adhere, and in the case of CL4 cells, characteristic island

formation could be observed. Cells were fixed in 4% paraformaldehyde and 2% sucrose in PBS for 10

mins (EMS Diasum). Fixed cells were permeabilized/blocked in 0.2% (v/v) Triton X-100, 10% (v/v)

normal goat serum (NGS) in PBS for 1 hr at RT, followed by labelling with rhodamine phalloidin (Life

Technologies) and Hoechst 33342 (Life Technologies) in blocking buffer (2% NGS in PBS). Confocal

microscopy images were captured as described above.

Actin purification and labelling

Actin was extracted from rabbit skeletal acetone powder (Pel-Freeze, AZ) in chilled G-buffer

(concentrations in mM), Tris-HCl (2), ATP (0.2), $CaCl_2$ (0.1), NaN_3 (1), DTT (1), pH 8, using

established protocols (Spudich and Watt, 1971). Actin was additionally labelled on Cys 374 using

either N-(1-pyrene)-iodoacetamide (Life Technologies), or tetramethylrhodamine-5-maleimide

(Adipogen Life Sciences)(Criddle et al., 1985; Fujiwara et al., 2002). F-actin used for steady-state

ATPase and gliding filaments assays was purified through two rounds of polymerization and

depolymerization using ultracentrifugation. F-actin was dialyzed extensively against MOPS (4), $MgCl_2$

(1), EGTA (0.1), DTT (1), NaN_3 (1), pH 7.0, and its concentration measured at 290 nm (ε = 26,600 M$^-$

^{1}·cm^{-1}) prior to use. Unlabelled, pyrene- and TMR-labelled G-actin for polymerization studies were

further purified by size exclusion chromatography (Superdex 200, Cytiva) using isocratic elution in G-

buffer. Fractions were exclusively taken from the falling edge of the chromatogram, to ensure the

recovery of monomeric actin. Concentrations and dye-labelling efficiency was determined at 290 nm

(actin: ε = 26,600 M^{-1}·cm^{-1}), 344 nm (pyrene: ε = 22,000 M^{-1}·cm^{-1}) and 550 nm (rhodamine: ε = 96,900

M^{-1}·cm^{-1}). Correction factors were applied for pyrene actin, $A_{corr} = A_{290} - (0.127 * A_{344})$, and separately

for rhodamine actin, $A_{corr} = A_{290} - (0.208 * A_{550})$(Fujiwara et al., 2002). Biotinylated skeletal muscle

actin (#8109-01, HyperMol, Germany) was rehydrated, dialyzed against G-buffer and cleared by

ultracentrifugation for 60 minutes at 100k x g prior to use. For actin polymerization experiments

performed without free ATP in solution, G-actin stocks (with 0.2 mM ATP) were desalted (PD

SpinTrap G-25, Cytiva) into a modified G-buffer (no ATP): Tris-HCl (2), $CaCl_2$ (0.1), NaN_3 (1), DTT (1),

pH 8, immediately prior to use. Desalted G-actin (ATP) monomers were stored on ice and used within

3 hours.

Expression of MYO15 and deafness mutants in *Sf*9 cells using baculovirus

The baculoviral transfer vector pFastbac1 M15-2IQ-EGFP-FLAG, encoding the wild-type mouse MYO15 motor domain (NP_874357.2, aa. 1 - 743) as a C-terminal fusion with EGFP and FLAG moieties, was previously described (Bird et al., 2014). The expressed protein was 114 kDa. The *jordan* and *shaker-2* non-synonymous substitutions were separately introduced into pFastbac1 M15-2IQ-EGFP-FLAG by site-directed mutagenesis (QuikChange II, Agilent) and verified by Sanger sequencing. Plasmid DNA encoding wild-type, *jordan* and *shaker-2* was separately transformed into DH10Bac cells (Life Technologies) and recombinant bacmid DNA prepared following the manufacturer's protocol. First passage (P1) recombinant baculovirus was generated by transfecting *Sf*9 cells (Expression Systems) with bacmid DNA complexed using linear polyethylenimine (PEI MAX, 40,000 MW, Polysciences Inc., PA) at a 12:1 (PEI : DNA) molar ratio. *Sf*9 cells were maintained in suspension culture with ESF-921 medium (Expression Systems) in a shaking incubator at 27°C. P1 baculovirus was amplified in *Sf*9 cells using low multiplicity of infection (MOI = 0.1) to generate P2 baculovirus for expression. Dual-promoter baculovirus expressing mouse UNC45B + HSP90AA1 chaperones (Bird et al., 2014), and bovine smooth muscle essential (MYL6, also referred to as MLC17B / ELC) + chicken regulatory (MYL12B, also referred to as MLC20 / RLC) light chains, were previously described (Pato et al., 1996). All baculoviruses were tittered using an end-point dilution assay and the *Sf*9 Easy Titer cell line (Hopkins and Esposito, 2009). To express M15-2IQ, *Sf*9 cells were seeded at a density of 2×10^6 cells·mL^{-1} in ESF-921, and infected simultaneously with three baculoviruses at the following MOI: M15-2IQ (5), UNC45B / HSP90AA1 (5), ELC / RLC (5).The myosin chaperones UNC45B and HSP90AA1 we co-expressed to aid folding, in addition to essential (MYL6) and regulatory (MYL12B) muscle light chains to bind the LCBDs (Bird et al., 2014; Jiang et al., 2021). Identical expressions were performed for the *shaker-2* and *jordan* variants. Cells were harvested at 48 - 72 hours post-infection by centrifugation at 500 x *g* and flash frozen in liquid nitrogen.

Purification of the MYO15 motor domain

M15-2IQ (wild-type, *jordan*, *shaker-2*) motor domains were purified from frozen *Sf*9 cells following established protocols (Bird et al., 2014; Jiang et al., 2021). Cell pellets were lysed using a Dounce homogenizer in extraction buffer (concentrations in mM): MOPS (10), NaCl (500), EGTA (1), MgCl$_2$(10), ATP (2), PMSF (0.2), DTT (0.1), NaN$_3$ (1), leupeptin (2 µg·mL^{-1}), protease inhibitor cocktail (Halt EDTA-free; Thermo Scientific), pH 7.2. Cell lysates were cleared for 30 mins at 48k x *g* and the supernatant incubated with FLAG M2 affinity resin (Sigma-Aldrich) for 3 hours at 4°C. FLAG resin was

packed into a gravity flow column and washed with a high-salt buffer, MOPS (10), NaCl (500), EGTA

(1), MgCl$_2$(5), ATP (1), PMSF (0.1), DTT (0.1), NaN$_3$ (1), leupeptin (2 µg·mL^{-1}), pH 7.2, followed by a

low-salt buffer, MOPS (10), NaCl (60), EGTA (1), PMSF (0.1), DTT (0.1), NaN$_3$ (1), leupeptin (2

µg·mL^{-1}), pH 7.0. M15-2IQ protein was eluted using low-salt buffer supplemented with 0.2 mg·mL^{-1} 3x

FLAG peptide (American Peptide, CA). For assessment of hydrodynamic radius, purified motor

domains (wild-type, *jordan*, *shaker-2*) were concentrated by centrifugation (10'000 MWCO; Amicon,

EMD-Millipore) and directly analyzed using size exclusion chromatography (see below), alongside

protein size standards (Thyroglobulin (1), Ferritin (2), Aldolase (3), Conalbumin (4), Ovalbumin (5),

Carbonic anhydrase (6), Ribonuclease A (7); HMW + LMW gel filtration calibration kit (Cytiva). For

preparative scale protein production, FLAG-eluted motor domains (wild-type + *jordan*) were bound to a

strong anion exchanger (5/50 MonoQ GL; Cytiva) using a Purifier 10 chromatography system (GE

Healthcare). The column was washed with MOPS (10), NaCl (100), EGTA (1), PMSF (0.1), DTT (1),

pH 7.0, and a 160 CV gradient elution performed to 1M NaCl (100% B). Fractions eluting at ~31

mS·cm^{-1} were concentrated (10,000 MWCO) and further purified using size exclusion

chromatography (Superdex 200, Cytiva) with isocratic elution in MOPS (10), KCl (100), EGTA (0.1),

NaN$_3$ (1), PMSF (0.1), DTT (1), leupeptin (1 µg·mL^{-1}), pH 7.0. The M15-2IQ : ELC : RLC complex

(1:1:1) eluted as a single peak and was concentrated (10,000 MWCO) before determining complex

concentration at 280 nm (ε = 88,020 M^{-1}·cm^{-1}).

ATPase assays

739   Steady-state actin activated ATPase assays were measured using a NADH-coupled assay, as

previously reported (Bird et al., 2014). Briefly, wild-type M15-2IQ (30 nM) or *jordan* (150 nM) protein

was assayed in the following reaction buffer (concentration in mM): MOPS (10), KCl (50), MgCl$_2$ (5),

EGTA (0.1), MgATP (2), 40 U·mL^{-1} lactate dehydrogenase (Sigma-Aldrich), 200 U·mL^{-1} pyruvate

kinase (Sigma-Aldrich), phosphoenolpyruvate (1) (Sigma-Aldrich), NADH (0.2), pH 7.0 at 20° ± 0.1 ℃.

Due to its lower activity, a higher concentration (150 nM) of M15-2IQ *jordan* protein was used. The

concentration of F-actin in the reaction was titrated from 0 to 100 µM. The absorbance of NADH (ε =

6,220 M^{-1}·cm^{-1}) at 340 nm was measured using a dual-beam spectrophotometer (UV-1800, Shimadzu)

and the ATP hydrolysis rate calculated from the change in absorbance over time. Hydrolysis rates

were corrected for basal M15-2IQ ATPase activity in the absence of F-actin, and also for the ATPase

activity of F-actin. ATP hydrolysis rates were fit to the Michaelis-Menten equation to estimate k_{cat} and

k_{ATPase} using Prism (GraphPad).

Gliding filament motility assay

Motility chambers were assembled by coating a clean cover glass (#1.5) with 0.1%

nitrocellulose in amyl acetate (Ladd Research Industries), and attaching it to a microscope slide with

two strips of double-sided adhesive tape (Scotch, 3M) to form a channel approximately 3 mm wide.

The chamber was incubated for 5 mins in 0.1 mg·mL^{-1} anti-GFP (clone GFP-20, Sigma-Aldrich)

diluted with motility buffer (MB) (concentrations in mM): MOPS (20), KCl (10), MgCl$_2$ (5), EGTA (0.1),

pH 7.0. The surface was blocked using 1 mg·mL^{-1} BSA (Sigma-Aldrich) diluted in MB and incubated

for 1 min. After washing the chamber with MB, 1 µM M15-2IQ (wild-type or *jordan*) diluted in MB was

incubated for 1 min to functionalize the anti-GFP coated surface. The chamber was washed with 1

mg·mL^{-1} BSA in MB, followed by MB alone. Finally, the chamber was incubated for 2 mins with TMR-

phalloidin stabilized actin filaments (5 nM) diluted in MB, and subsequently washed in MB. Filaments

were visualized using an inverted epifluorescence microscope (Olympus IX-51) and motility recorded

in MB supplemented with 50 mM DTT, 2 mM ATP, 3 mM glucose, 100 µg·mL glucose oxidase, and 20

µg·mL catalase at 30 ± 0.5 °C. Actin filament velocities were analyzed as described (Sellers et al.,

1993).

Polymerization assay

Actin polymerization was measured using G-actin labelled on Cys 374 with N-(1-pyrene)-

iodoacetamide (see above), and a cuvette-based fluorometer (PTI Quantamaster 400, HORIBA

Scientific) used to excite pyrene at 365 nm and capture fluorescence emission at 407 nm. Gel filtered

G-actin (10% pyrene labelled, with free ATP, or desalted) was converted to the physiological Mg^{2+}

bound form by addition of 50 µM MgCl$_2$ and 0.2 mM EGTA for exactly two minutes at room

temperature. The polymerization reaction was initiated by mixing G-actin (3 x stock) with KMEI buffer

(1.5x stock) in a 1:2 ratio, respectively. Myosin (wild-type or *jordan*) was included in the 1.5x KMEI

buffer as needed. Final concentrations were 2 µM G-actin and 1 µM myosin, in assay buffer

(concentrations in mM): KCl (50), MgCl$_2$ (1), EGTA (1), imidazole (10), pH 7.0 at 25° ± 0.1 ºC. Data

were corrected for dead-time and fluorescence recorded until the transient reached plateau, or for a

maximum of 3 hours. Fluorescence derivatives with respect to time were processed in Prism

(GraphPad). Half times were calculated as described (Hansen et al., 2013).

TIRF single-filament polymerization assay

High-tolerance cover slips (24 × 50 mm, #1.5, Marienfeld Superior, Germany) were cleaned by

sequential sonication (10 mins each) in 2% Hellmanex III (Hellma, Germany), 1M KOH, 100% ethanol,

and finally Milli-Q water. Cover slips were dried under a nitrogen stream and processed for 10 mins

under argon plasma (ZEPTO, Diener Electronic, Germany). A mixture of mPEG-silane (2 mg·mL^{-1},

Laysan Bio, AL) and biotin–PEG–silane (2 µg·mL^{-1}, Laysan Bio) was prepared in 96% ethanol, 0.1%

(v/v) HCl. Cover slips were coated with 100 µL of the mPEG mixture and baked at 70°C for 1 hour.

Cover slips were rinsed twice in 96% ethanol, sonicated for 10 mins, followed by two rinses in Milli-Q,

sonicated for 10 mins in Milli-Q, and finally dried under a nitrogen stream. Flow chambers were

assembled using double-sided sticky tape to create a 3 mm wide channel on a glass slide.

Functionalized coverslips were placed over the tape and firmly pushed down to seal the flow chamber.

Immediately prior to use, flow cells were washed with buffer T50 (concentrations in mM): Tris·HCl

(10), KCl (50), DTT (1), pH 8.0. Flow cells were sequentially washed with 0.1 mg·mL^{-1} (Neutravidin,

Thermo Scientific) in T50 for 1 min, followed by a wash with 1 mg·mL^{-1} bovine serum albumin (A0281,

Sigma Aldrich) in T50 for 30 s, and a final wash of T50. Experiments were performed in the following

reaction buffer (final concentration in mM): KCl (50), MgCl$_2$ (1), EGTA (1), imidazole (10), ATP (0.025),

DTT (10), glucose (15), 0.5% methylcellulose, 20 µg·mL catalase, 100 µg·mL glucose oxidase, pH 7.0.

G-actin (1 µM total) was included in the reaction with TMR-actin (20%) and biotin-actin (10%) doping.

Purified myosin motor domain (1 µM) was optionally added, as needed. The reaction was loaded into

the flow cell and immediately mounted onto an inverted microscope (Nikon Ti-E2) equipped with an oil

immersion objective (CFI Apochromat TIRF 100x, 1.49 N.A., Nikon) for objective-style total internal

reflection fluorescence (TIRF) microscopy (H-TIRF, Nikon). The flow cell was illuminated using a 561

nm laser line, and emission filtered using a bandpass filter (ET630/75m, Chroma). Time-lapse images

were captured on an EM-CCD camera (iXon Ultra 888, Andor) controlled by NIS-Elements (AR

version 5.2, Nikon). Samples were imaged at 21 ± 1°C. The assay dead-time was typically 45 - 60 s,

and was included in data analyses.

Images were pre-processed in FIJI (https://fiji.sc)(Schindelin et al., 2012) by performing

background subtraction and image registration (descriptor-based series registration, 2d/3d +t). Actin

filament densities were quantified using the Analyze Particle command (size > 3 pixel^2, circularity: 0.0-

1.0) to count discrete particles within a 50 x 50 µm region of interest (ROI) that was randomly selected

from the field of view. A minimum of 3 experiments, from two independent protein preparations, were

analyzed for each condition. Filament elongation rates were calculated from time-lapse TIRF imaging

sequences using kymographs generated in Elements Software (version 5.2, Nikon). At least 40

filaments from two independent experiments were analyzed for each condition.

Statistical analyses

All statistical calculations and non-linear regressions were performed in Prism (GraphPad).

Regression fit parameters are quoted as mean ± SEM, and data points as mean ± SD, unless

otherwise stated. Statistical significance: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$.

AUTHOR CONTRIBUTIONS

824 Conceptualization: JEB, MRB

825 Investigation: ZGM, FJ, CA, MB, CFA, AS, JW, RH, DCS, SMC, AP, SM, EW,

826   YT, MRB, JEB

827 Formal analysis: ZGM, FJ, CA, MB, RG, AS, CFA, JW, RH, DCS, SMC, AP, SM,

828   EW, YT, JRS, SDMB, GMA TBF, MRB, JEB

829 Writing - original draft: ZGM, FJ, CA, MB, MRB, JEB

830 Writing – review and editing: All authors.

831 Project administration: JEB

832 Funding acquisition: JRS, SDMB, TBF, MRB, JEB

ACKNOWLEDGEMENTS

We thank Mireille Montcouquiol for the gift of GPSM2 antisera, James Bartles for the gift of

LLC-PK1-CL4 cells, Tracy Fitzgerald at the NIDCD Mouse Auditory Testing Core for expert technical

assistance, W. Clay Smith for use of fluorometry equipment, Peter Höök, H. Lee Sweeney and Tom

Pollard for insightful discussions, and Peter Barr-Gillespie for critical feedback on the manuscript. This

research was supported (in part) by the Intramural Research Program of the NIH (NIDCD DC000039

to TBF, NHLBI HL006049 to JRS), the Medical Research Council (MC_U142684175 to SDMB,

MC_UP_1503/2 to MRB) and National Institute on Deafness and Other Communication Disorders of

the National Institutes of Health under Award Number R01DC018827 (to JEB). The content is solely

the responsibility of the authors and does not necessarily represent the official views of the National

Institutes of Health. The funding bodies had no role in study design, data collection or analysis,

decision to publish, or preparation of the manuscript.

ACKNOWLEDGEMENTS

848 The authors declare that no competing interests exist.

REFERENCES

Ahuja R, Pinyol R, Reichenbach N, Custer L, Klingensmith J, Kessels MM, Qualmann B (2007)
Cordon-bleu is an actin nucleation factor and controls neuronal morphology. Cell 131:337–350.

Anderson DW, Probst FJ, Belyantseva IA, Fridell RA, Beyer L, Martin DM, Wu D, Kachar B, Friedman
TB, Raphael Y, Camper SA (2000) The motor and tail regions of myosin XV are critical for normal
structure and function of auditory and vestibular hair cells. Hum Mol Genet 9:1729–1738.

Avenarius MR, Krey JF, Dumont RA, Morgan CP, Benson CB, Vijayakumar S, Cunningham CL,
Scheffer DI, Corey DP, Müller U, Jones SM, Barr-Gillespie PG (2017) Heterodimeric capping protein
is required for stereocilia length and width regulation. J Cell Biol 216:3861–3881.

Avraham KB, Hasson T, Steel KP, Kingsley DM, Russell LB, Mooseker MS, Copeland NG, Jenkins
NA (1995) The mouse Snell's waltzer deafness gene encodes an unconventional myosin required for
structural integrity of inner ear hair cells. Nat Genet 11:369–375.

Bariani P, De Chiffre L, Hansen HN, Horsewell A (2005) Investigation on the traceability of three
dimensional scanning electron microscope measurements based on the stereo-pair technique. Precis
Eng 29:219–228.

Barr-Gillespie P-G (2015) Assembly of hair bundles, an amazing problem for cell biology. Mol Biol Cell
26:2727–2732.

Belyantseva IA, Boger ET, Friedman TB (2003) Myosin XVa localizes to the tips of inner ear sensory
cell stereocilia and is essential for staircase formation of the hair bundle. Proc Natl Acad Sci U S A
100:13958–13963.

Belyantseva IA, Boger ET, Naz S, Frolenkov GI, Sellers JR, Ahmed ZM, Griffith AJ, Friedman TB
(2005) Myosin-XVa is required for tip localization of whirlin and differential elongation of hair-cell
stereocilia. Nat Cell Biol 7:148–156.

Beurg M, Fettiplace R, Nam J-H, Ricci AJ (2009) Localization of inner hair cell mechanotransducer
channels using high-speed calcium imaging. Nat Neurosci 12:553–558.

Bird JE, Barzik M, Drummond MC, Sutton DC, Goodman SM, Morozko EL, Cole SM, Boukhvalova
AK, Skidmore J, Syam D, Wilson EA, Fitzgerald T, Rehman AU, Martin DM, Boger ET, Belyantseva
IA, Friedman TB (2017) Harnessing molecular motors for nanoscale pulldown in live cells. Mol Biol
Cell 28:463–475.

Bird JE, Takagi Y, Billington N, Strub M-P, Sellers JR, Friedman TB (2014) Chaperone-enhanced
purification of unconventional myosin 15, a molecular motor specialized for stereocilia protein
trafficking. Proc Natl Acad Sci U S A 111:12390–12395.

Brawley CM, Rock RS (2009) Unconventional myosin traffic in cells reveals a selective actin
cytoskeleton. Proc Natl Acad Sci U S A 106:9685–9690.

Cooper JA, Walker SB, Pollard TD (1983) Pyrene actin: documentation of the validity of a sensitive
assay for actin polymerization. J Muscle Res Cell Motil 4:253–262.

Criddle AH, Geeves MA, Jeffries T (1985) The use of actin labelled with N-(1-pyrenyl)iodoacetamide
to study the interaction of actin with myosin subfragments and troponin/tropomyosin. Biochem J
232:343–349.

Delprat B, Michel V, Goodyear R, Yamasaki Y, Michalski N, El-Amraoui A, Perfettini I, Legrain P,
Richardson G, Hardelin J-P, Petit C (2005) Myosin XVa and whirlin, two deafness gene products

required for hair bundle growth, are located at the stereocilia tips and interact directly. Hum Mol Genet
14:401–410.

Dominguez R, Holmes KC (2011) Actin structure and function. Annu Rev Biophys 40:169–186.

Drummond MC, Barzik M, Bird JE, Zhang D-S, Lechene CP, Corey DP, Cunningham LL, Friedman
TB (2015) Live-cell imaging of actin dynamics reveals mechanisms of stereocilia length regulation in
the inner ear. Nat Commun 6:6873.

Dunbar LA et al. (2019) Clarin-2 is essential for hearing by maintaining stereocilia integrity and
function. EMBO Mol Med 11:e10288.

Ebrahim S, Avenarius MR, Grati M, Krey JF, Windsor AM, Sousa AD, Ballesteros A, Cui R, Millis BA,
Salles FT, Baird MA, Davidson MW, Jones SM, Choi D, Dong L, Raval MH, Yengo CM, Barr-Gillespie
PG, Kachar B (2016) Stereocilia-staircase spacing is influenced by myosin III motors and their cargos
espin-1 and espin-like. Nat Commun 7:10833.

Ezan J, Lasvaux L, Gezer A, Novakovic A, May-Simera H, Belotti E, Lhoumeau A-C, Birnbaumer L,
Beer-Hammer S, Borg J-P, Le Bivic A, Nürnberg B, Sans N, Montcouquiol M (2013) Primary cilium
migration depends on G-protein signalling control of subapical cytoskeleton. Nat Cell Biol 15:1107–
1115.

Fang Q, Indzhykulian AA, Mustapha M, Riordan GP, Dolan DF, Friedman TB, Belyantseva IA,
Frolenkov GI, Camper SA, Bird JE (2015) The 133-kDa N-terminal domain enables myosin 15 to
maintain mechanotransducing stereocilia and is essential for hearing. eLife 4.

Fievez S, Carlier MF (1993) Conformational changes in subdomain-2 of G-actin upon polymerization
into F-actin and upon binding myosin subfragment-1. FEBS Lett 316:186–190.

Fievez S, Carlier MF, Pantaloni D (1997a) Mechanism of myosin subfragment-1-induced assembly of
CaG-actin and MgG-actin into F-actin-S1-decorated filaments. Biochemistry 36:11843–11850.

Fievez S, Pantaloni D, Carlier MF (1997b) Kinetics of myosin subfragment-1-induced condensation of
G-actin into oligomers, precursors in the assembly of F-actin-S1. Role of the tightly bound metal ion
and ATP hydrolysis. Biochemistry 36:11837–11842.

Flock A, Cheung HC (1977) Actin filaments in sensory hairs of inner ear receptor cells. J Cell Biol
75:339–343.

Friedman TB, Liang Y, Weber JL, Hinnant JT, Barber TD, Winata S, Arhya IN, Asher JH (1995) A
gene for congenital, recessive deafness DFNB3 maps to the pericentromeric region of chromosome
17. Nat Genet 9:86–91.

Fujiwara I, Takahashi S, Tadakuma H, Funatsu T, Ishiwata S (2002) Microscopic analysis of
polymerization dynamics with individual actin filaments. Nat Cell Biol 4:666–673.

Gibson F, Walsh J, Mburu P, Varela A, Brown KA, Antonio M, Beisel KW, Steel KP, Brown SD (1995)
A type VII myosin encoded by the mouse deafness gene shaker-1. Nature 374:62–64.

Gillespie PG, Corey DP (1997) Myosin and adaptation by hair cells. Neuron 19:955–958.

Gong R, Jiang F, Moreland ZG, Reynolds M, Espinosa de los Reyes S, Gurel PS, Shams A, Bowl MR,
Bird JE, Alushin GM (2021) Structural basis for tunable control of actin dynamics by myosin-15 in
mechanosensory stereocilia. bioRxiv 2021.07.09.451843.

Hansen SD, Zuchero JB, Mullins RD (2013) Cytoplasmic actin: purification and single molecule
assembly assays. Methods Mol Biol Clifton NJ 1046:145–170.

Hardisty-Hughes RE, Parker A, Brown SDM (2010) A hearing and vestibular phenotyping pipeline to
identify mouse mutants with hearing impairment. Nat Protoc 5:177–190.

Hartman MA, Spudich JA (2012) The myosin superfamily at a glance. J Cell Sci 125:1627–1632.

Hasson T, Gillespie PG, Garcia JA, MacDonald RB, Zhao Y, Yee AG, Mooseker MS, Corey DP (1997)
Unconventional myosins in inner-ear sensory epithelia. J Cell Biol 137:1287–1307.

Heissler SM, Sellers JR (2016) Various Themes of Myosin Regulation. J Mol Biol 428:1927–1946.

Holt JR, Gillespie SKH, Provance DW, Shah K, Shokat KM, Corey DP, Mercer JA, Gillespie PG
(2002) A chemical-genetic strategy implicates myosin-1c in adaptation by hair cells. Cell 108:371–381.

Hopkins R, Esposito D (2009) A rapid method for titrating baculovirus stocks using the Sf-9 Easy Titer
cell line. BioTechniques 47:785–788.

Houdusse A, Sweeney HL (2016) How Myosin Generates Force on Actin Filaments. Trends Biochem
Sci 41:989–997.

Jiang F, Takagi Y, Shams A, Heissler SM, Friedman TB, Sellers JR, Bird JE (2021) The ATPase
mechanism of myosin 15, the molecular motor mutated in DFNB3 human deafness. J Biol Chem
296:100243.

Kaltenbach JA, Falzarano PR, Simpson TH (1994) Postnatal development of the hamster cochlea. II.
Growth and differentiation of stereocilia bundles. J Comp Neurol 350:187–198.

Kerber ML, Jacobs DT, Campagnola L, Dunn BD, Yin T, Sousa AD, Quintero OA, Cheney RE (2009)
A novel form of motility in filopodia revealed by imaging myosin-X at the single-molecule level. Curr
Biol CB 19:967–973.

Kinosian HJ, Selden LA, Estes JE, Gershman LC (1993) Actin filament annealing in the presence of
ATP and phalloidin. Biochemistry 32:12353–12357.

Krey JF, Barr-Gillespie PG (2019) Molecular Composition of Vestibular Hair Bundles. Cold Spring
Harb Perspect Med 9:a033209.

Krey JF, Chatterjee P, Dumont RA, O'Sullivan M, Choi D, Bird JE, Barr-Gillespie PG (2020)
Mechanotransduction-Dependent Control of Stereocilia Dimensions and Row Identity in Inner Hair
Cells. Curr Biol CB 30:442-454.e7.

Kuhn JR, Pollard TD (2005) Real-time measurements of actin filament polymerization by total internal
reflection fluorescence microscopy. Biophys J 88:1387–1402.

Lheureux K, Forné T, Chaussepied P (1993) Interaction and polymerization of the G-actin-myosin
head complex: effect of DNase I. Biochemistry 32:10005–10014.

Li X-D, Ikebe R, Ikebe M (2005) Activation of myosin Va function by melanophilin, a specific docking
partner of myosin Va. J Biol Chem 280:17815–17822.

Liang Y, Wang A, Belyantseva IA, Anderson DW, Probst FJ, Barber TD, Miller W, Touchman JW, Jin
L, Sullivan SL, Sellers JR, Camper SA, Lloyd RV, Kachar B, Friedman TB, Fridell RA (1999)
Characterization of the human and mouse unconventional myosin XV genes responsible for hereditary
deafness DFNB3 and shaker 2. Genomics 61:243–258.

Liu J, Taylor DW, Krementsova EB, Trybus KM, Taylor KA (2006) Three-dimensional structure of the
myosin V inhibited state by cryoelectron tomography. Nature 442:208–211.

Machesky LM, Atkinson SJ, Ampe C, Vandekerckhove J, Pollard TD (1994) Purification of a cortical
complex containing two unconventional actins from Acanthamoeba by affinity chromatography on
profilin-agarose. J Cell Biol 127:107–115.

Manor U, Disanza A, Grati M, Andrade L, Lin H, Di Fiore PP, Scita G, Kachar B (2011) Regulation of
stereocilia length by myosin XVa and whirlin depends on the actin-regulatory protein Eps8. Curr Biol
CB 21:167–172.

Mauriac SA, Hien YE, Bird JE, Carvalho SD-S, Peyroutou R, Lee SC, Moreau MM, Blanc J-M, Geyser
A, Medina C, Thoumine O, Beer-Hammer S, Friedman TB, Rüttiger L, Forge A, Nürnberg B, Sans N,
Montcouquiol M (2017) Defective Gpsm2/Gαi3 signalling disrupts stereocilia development and growth
cone actin dynamics in Chudley-McCullough syndrome. Nat Commun 8:14907.

Mburu P et al. (2003) Defects in whirlin, a PDZ domain molecule involved in stereocilia elongation,
cause deafness in the whirler mouse and families with DFNB31. Nat Genet 34:421–428.

Mianné J, Chessum L, Kumar S, Aguilar C, Codner G, Hutchison M, Parker A, Mallon A-M, Wells S,
Simon MM, Teboul L, Brown SDM, Bowl MR (2016) Correction of the auditory phenotype in C57BL/6N
mice via CRISPR/Cas9-mediated homology directed repair. Genome Med 8:16.

Miller L, Phillips M, Reisler E (1988) Polymerization of G-actin by myosin subfragment 1. J Biol Chem
263:1996–2002.

Mullins RD, Heuser JA, Pollard TD (1998) The interaction of Arp2/3 complex with actin: nucleation,
high affinity pointed end capping, and formation of branching networks of filaments. Proc Natl Acad
Sci U S A 95:6181–6186.

Mullins RD, Stafford WF, Pollard TD (1997) Structure, subunit topology, and actin-binding activity of
the Arp2/3 complex from Acanthamoeba. J Cell Biol 136:331–343.

Murphy DB, Gray RO, Grasser WA, Pollard TD (1988) Direct demonstration of actin filament
annealing in vitro. J Cell Biol 106:1947–1954.

Nagy S, Ricca BL, Norstrom MF, Courson DS, Brawley CM, Smithback PA, Rock RS (2008) A myosin
motor that selects bundled actin for motility. Proc Natl Acad Sci U S A 105:9616–9620.

Narayanan P, Chatterton P, Ikeda A, Ikeda S, Corey DP, Ervasti JM, Perrin BJ (2015) Length
regulation of mechanosensitive stereocilia depends on very slow actin dynamics and filament-severing
proteins. Nat Commun 6:6855.

Okreglak V, Drubin DG (2010) Loss of Aip1 reveals a role in maintaining the actin monomer pool and an in vivo oligomer assembly pathway. J Cell Biol 188:769–777.

Pato MD, Sellers JR, Preston YA, Harvey EV, Adelstein RS (1996) Baculovirus expression of chicken nonmuscle heavy meromyosin II-B. Characterization of alternatively spliced isoforms. J Biol Chem 271:2689–2695.

Peng AW, Belyantseva IA, Hsu PD, Friedman TB, Heller S (2009) Twinfilin 2 regulates actin filament lengths in cochlear stereocilia. J Neurosci Off J Soc Neurosci 29:15083–15088.

Potter PK et al. (2016) Novel gene function revealed by mouse mutagenesis screens for models of age-related disease. Nat Commun 7:12444.

Probst FJ, Fridell RA, Raphael Y, Saunders TL, Wang A, Liang Y, Morell RJ, Touchman JW, Lyons RH, Noben-Trauth K, Friedman TB, Camper SA (1998) Correction of deafness in shaker-2 mice by an unconventional myosin in a BAC transgene. Science 280:1444–1447.

Quinlan ME, Heuser JE, Kerkhoff E, Mullins RD (2005) Drosophila Spire is an actin nucleation factor. Nature 433:382–388.

Rehman AU, Bird JE, Faridi R, Shahzad M, Shah S, Lee K, Khan SN, Imtiaz A, Ahmed ZM, Riazuddin S, Santos-Cortez RLP, Ahmad W, Leal SM, Riazuddin S, Friedman TB (2016) Mutational Spectrum of MYO15A and the Molecular Mechanisms of DFNB3 Human Deafness. Hum Mutat 37:991–1003.

Reymann A-C, Boujemaa-Paterski R, Martiel J-L, Guérin C, Cao W, Chin HF, De La Cruz EM, Théry M, Blanchoin L (2012) Actin network architecture can determine myosin motor activity. Science 336:1310–1314.

Richardson GP, de Monvel JB, Petit C (2011) How the genetics of deafness illuminates auditory physiology. Annu Rev Physiol 73:311–334.

Ropars V, Yang Z, Isabet T, Blanc F, Zhou K, Lin T, Liu X, Hissier P, Samazan F, Amigues B, Yang ED, Park H, Pylypenko O, Cecchini M, Sindelar CV, Sweeney HL, Houdusse A (2016) The myosin X motor is optimized for movement on actin bundles. Nat Commun 7:12456.

Rzadzinska AK, Schneider ME, Davies C, Riordan GP, Kachar B (2004) An actin molecular treadmill and myosins maintain stereocilia functional architecture and self-renewal. J Cell Biol 164:887–897.

Sagot I, Rodal AA, Moseley J, Goode BL, Pellman D (2002) An actin nucleation mechanism mediated by Bni1 and profilin. Nat Cell Biol 4:626–631.

Sakai T, Umeki N, Ikebe R, Ikebe M (2011) Cargo binding activates myosin VIIA motor function in cells. Proc Natl Acad Sci U S A 108:7028–7033.

Salles FT, Merritt RC, Manor U, Dougherty GW, Sousa AD, Moore JE, Yengo CM, Dosé AC, Kachar B (2009) Myosin IIIa boosts elongation of stereocilia by transporting espin 1 to the plus ends of actin filaments. Nat Cell Biol 11:443–450.

Sato O, Li X-D, Ikebe M (2007) Myosin Va becomes a low duty ratio motor in the inhibited form. J Biol Chem 282:13228–13239.

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C,
Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A
(2012) Fiji: an open-source platform for biological-image analysis. Nat Methods 9:676–682.

Schneider ME, Belyantseva IA, Azevedo RB, Kachar B (2002) Rapid renewal of auditory hair bundles.
Nature 418:837–838.

Sckolnick M, Krementsova EB, Warshaw DM, Trybus KM (2013) More than just a cargo adapter,
melanophilin prolongs and slows processive runs of myosin Va. J Biol Chem 288:29313–29322.

Sellers JR, Cuda G, Wang F, Homsher E (1993) Myosin-specific adaptations of the motility assay.
Methods Cell Biol 39:23–49.

Sept D, Xu J, Pollard TD, McCammon JA (1999) Annealing accounts for the length of actin filaments
formed by spontaneous polymerization. Biophys J 77:2911–2919.

Shin J-B, Streijger F, Beynon A, Peters T, Gadzalla L, McMillen D, Bystrom C, Van der Zee CEEM,
Wallimann T, Gillespie PG (2007) Hair Bundles Are Specialized for ATP Delivery via Creatine Kinase.
Neuron 53:371–386.

Spudich JA, Watt S (1971) The regulation of rabbit skeletal muscle contraction. I. Biochemical studies
of the interaction of the tropomyosin-troponin complex with actin and the proteolytic fragments of
myosin. J Biol Chem 246:4866–4871.

Stepanyan R, Belyantseva IA, Griffith AJ, Friedman TB, Frolenkov GI (2006) Auditory
mechanotransduction in the absence of functional myosin-XVa. J Physiol 576:801–808.

Tadenev ALD, Akturk A, Devanney N, Mathur PD, Clark AM, Yang J, Tarchini B (2019) GPSM2-GNAI
Specifies the Tallest Stereocilia and Defines Hair Bundle Row Identity. Curr Biol CB 29:921-934.e4.

Tarchini B, Tadenev ALD, Devanney N, Cayouette M (2016) A link between planar polarity and
staircase-like bundle architecture in hair cells. Dev Camb Engl 143:3926–3932.

Thirumurugan K, Sakamoto T, Hammer JA, Sellers JR, Knight PJ (2006) The cargo-binding domain
regulates structure and activity of myosin 5. Nature 442:212–215.

Tilney LG, Cotanche DA, Tilney MS (1992) Actin filaments, stereocilia and hair cells of the bird
cochlea. VI. How the number and arrangement of stereocilia are determined. Dev Camb Engl
116:213–226.

Tilney LG, Derosier DJ, Mulroy MJ (1980) The organization of actin filaments in the stereocilia of
cochlear hair cells. J Cell Biol 86:244–259.

Umeki N, Jung HS, Sakai T, Sato O, Ikebe R, Ikebe M (2011) Phospholipid-dependent regulation of
the motor activity of myosin X. Nat Struct Mol Biol 18:783–788.

Umeki N, Jung HS, Watanabe S, Sakai T, Li X, Ikebe R, Craig R, Ikebe M (2009) The tail binds to the
head-neck domain, inhibiting ATPase activity of myosin VIIA. Proc Natl Acad Sci U S A 106:8483–
8488.

Vélez-Ortega AC, Freeman MJ, Indzhykulian AA, Grossheim JM, Frolenkov GI (2017)
Mechanotransduction current is essential for stability of the transducing stereocilia in mammalian
auditory hair cells. eLife 6.

Vélez-Ortega AC, Frolenkov GI (2019) Building and repairing the stereocilia cytoskeleton in
mammalian auditory hair cells. Hear Res 376:47–57.

Wang A, Liang Y, Fridell RA, Probst FJ, Wilcox ER, Touchman JW, Morton CC, Morell RJ, Noben-
Trauth K, Camper SA, Friedman TB (1998) Association of unconventional myosin MYO15 mutations
with human nonsyndromic deafness DFNB3. Science 280:1447–1451.

Xiong W, Wagner T, Yan L, Grillet N, Müller U (2014) Using injectoporation to deliver genes to
mechanosensory hair cells. Nat Protoc 9:2438–2449.

Yagi K, Mase R, Sakakibara I, Asai H (1965) FUNCTION OF HEAVY MEROMYOSIN IN THE
ACCELERATION OF ACTIN POLYMERIZATION. J Biol Chem 240:2448–2454.

Yang Y, Baboolal TG, Siththanandan V, Chen M, Walker ML, Knight PJ, Peckham M, Sellers JR
(2009) A FERM domain autoregulates Drosophila myosin 7a activity. Proc Natl Acad Sci U S A
106:4189–4194.

Young G et al. (2018) Quantitative mass imaging of single biological macromolecules. Science
360:423–427.

Zampini V, Rüttiger L, Johnson SL, Franz C, Furness DN, Waldhaus J, Xiong H, Hackney CM, Holley
MC, Offenhauser N, Di Fiore PP, Knipper M, Masetto S, Marcotti W (2011) Eps8 regulates hair bundle
length and functional maturation of mammalian auditory hair cells. PLoS Biol 9:e1001048.

Zhang D-S, Piazza V, Perrin BJ, Rzadzinska AK, Poczatek JC, Wang M, Prosser HM, Ervasti JM,
Corey DP, Lechene CP (2012) Multi-isotope imaging mass spectrometry reveals slow protein turnover
in hair-cell stereocilia. Nature 481:520–524.

Zheng L, Zheng J, Whitlon DS, García-Añoveros J, Bartles JR (2010) Targeting of the hair cell
proteins cadherin 23, harmonin, myosin XVa, espin, and prestin in an epithelial cell model. J Neurosci
Off J Soc Neurosci 30:7187–7201.

FIGURE LEGENDS

FIGURE 1

A) Schematic showing the protein domains of the long (MYO15-1) and short (MYO15-2) isoforms

encoded by the *Myo15* gene. The *jordan* and *shaker-2* deafness mutations are shown. **B)** ABR

phenotyping of the *jordan* pedigree at 3 months identified 10 mice with statistically elevated hearing

thresholds (red circles) for click and at 8, 16 and 32 kHz stimuli, compared to their normal hearing

pedigree mates (n = 73, black circles). Statistical outliers were detected using robust regression and

outlier removal (red circles, ROUT, Q = 1%). Thresholds of affected mice that did not respond to the

highest intensity stimulus (90 dB SPL) are recorded as 95 dB SPL. Data are mean ± SD. **C)**

Evolutionary conservation of the aspartate (D) residue of MYO15 altered to glycine (G) in *jordan* mice

that causes hearing loss. Residue positions refer to mouse MYO15-1 (NP_034992.2). **D)** ABR

recordings of *Myo15*$^{jd/sh2}$ compound heterozygotes at P28 shows profound hearing loss, with

thresholds elevated compared with normal hearing *Myo15*$^{+/jd}$ *or Myo15*$^{+/sh2}$ littermates. **E-F)**

Longitudinal auditory phenotyping of *jordan* mice at 1- (E) and 3- (F) months of age. ABR recordings

show that *Myo15*$^{jd/jd}$ mice (red, n=10) exhibit a progressive, moderate-to-severe hearing loss affecting

all frequencies, whereas age-matched *Myo15*$^{+/+}$(black, n=10) and *Myo15*$^{+/jd}$ (grey, n=15) littermate

controls have normal thresholds (15 - 45 dB SPL). *Myo15*$^{+/+}$ *vs Myo15*$^{jd/jd}$ comparison, ANOVA with

Tukey's multiple comparisons test.

FIGURE 2

A-F) Representative SEM images of stereocilia bundles from *Myo15*$^{+/+}$, *Myo15*$^{jd/jd}$, and *Myo15*$^{sh2/sh2}$

OHCs (A,C,E) and IHCs (B,D,F) at P8. In *Myo15*$^{+/+}$ mice, both IHC and OHC bundles display the

characteristic staircase architecture with 3 stereocilia ranks of increasing height. IHC and OHC

bundles from either *Myo15*$^{jd/jd}$ and *Myo15*$^{sh2/sh2}$ mice are shorter in height than the wild-type bundles.

Myo15$^{jd/jd}$ stereocilia taper in height towards the periphery of the bundle (white arrows). **G)** Projected

heights of tallest (row 1) OHC stereocilia at P8 are 1.2 ± 0.1 µm (*Myo15*$^{+/+}$, 58 stereocilia from 4 mice),

0.8 ± 0.1 µm (*Myo15*$^{jd/jd}$, 60 stereocilia from 4 mice), and 0.4 ± 0.1 µm (*Myo15*$^{sh2/sh2}$, 50 stereocilia from

2 mice). **H)** Projected heights of tallest (row 1) IHC stereocilia at P8 are 2.2 ± 0.2 µm (*Myo15*$^{+/+}$, 47

stereocilia from 4 mice), 1.3 ± 0.1 µm (*Myo15*$^{jd/jd}$, 47 stereocilia from 4 mice), and 0.9 ± 0.1 µm

(*Myo15*$^{sh2/sh2}$, 30 stereocilia from 2 mice). Brown-Forsythe and Welch ANOVA with Dunnett's T3

multiple comparisons test. Images are from mid-cochlear turn. Scale bars, 1 µm.

FIGURE 3

A,B) Immunofluorescence (IF) confocal images showing anti-MYO15 (green, PB48) in control *Myo15$^{+/jd}$*,

Myo15$^{sh2/sh2}$ and *Myo15$^{jd/jd}$* IHCs fixed at P14 (A), or P7 (B). Phalloidin was used to label F-actin

(magenta). **C)** IF confocal images of elongation complex proteins (green) in control *Myo15$^{+/jd}$* and

Myo15$^{jd/jd}$ IHCs fixed at P7, overlaid with phalloidin labelled F-actin (magenta). Images are

representative of data from at least two independent animals per genotype / antibody, and are shown

as maximum intensity projections reconstructed from z-stacks. Scale bars, 10 µm.

FIGURE 4

A) Actin barbed-end assay in detergent-permeabilized hair cells from mouse cochlear explant cultures

at P6. TMR-labelled G-actin (green) was added prior to fixation to identify uncapped barbed ends.

Phalloidin labelling of F-actin (magenta) is overlaid. In both *Myo15$^{jd/jd}$* and littermate *Myo15$^{+/jd}$* controls,

barbed-ends were detected at row 2 stereocilia tips, and at the tips of all stereocilia rows in *Myo15$^{sh2/sh2}$*

hair cells. **B)** HeLa cells were transfected with EGFP-tagged *Myo15-2* expression constructs or EGFP

alone (green) as indicated, fixed and probed with phalloidin (magenta) and Hoechst (blue). EGFP-M15

(wt) trafficked to filopodia tips (red arrowheads), while EGFP-M15 (jd) and EGFP-M15 (sh2) did not.

Boxed regions are magnified (inverted grayscale). **C)** CL4 cells were transfected with EGFP-tagged

Myo15-2 and prepared as above. Both EGFP-M15 (wt) and EGFP-M15 (jd) concentrated at microvillar

tips, while EGFP-M15 (sh2) did not. Orthogonal projections are shown (inverted grayscale). Images are

representative from at least three independent experiments. Scale bars, 2 µm (A); 20 µm (B+C).

FIGURE 5

A) Cartoon of MYO15 motor domains expressed in *Sf*9 cells, consisting of the ATPase and two light

chain binding domains (LCBD). **B)** Size exclusion chromatography analysis of FLAG/IEX purified

M15(wt) and M15(jd) proteins. M15(sh2) elutes close to the void volume (arrow) and was aggregated.

Protein calibration standards (dotted lines), FLAG peptide (asterisk). **C)** SDS-PAGE analysis of SEC

purified motor domain proteins. The motor domain (arrow) co-purifies with RLC and ELC light chains

for all variants. M15(sh2) was misfolded and extracted from *Sf*9 cells at low yield. **D)** Steady-state

ATPase activation of M15(wt) and M15(jd) motor domains measured using a NADH-coupled assay at

20 ± 0.1°C. Reactions were performed with [F-actin] as shown. Rectangular hyperbola fits are shown

for M15(wt) (blue, k_{cat} = 5.8 ± 0.2 s^{-1}, k_{ATPase} = 29.1 ± 2.1 µM) and for M15(jd) (green, k_{cat} = 0.87 ± 0.04

s^{-1}, k_{ATPase} = 114.3 ± 8.2 µM, n = 4). **E)** Frequency histogram of F-actin velocities in a gliding filament

assay at 30 ± 0.1°C. Gaussian fits (dotted line) are overlaid for M15(wt) (473 ± 67 nm·s^{-1}, n = 5449

filaments, mean ± SD) and M15(jd) (216 ± 71 nm·s^{-1}, n = 2844 filaments). Data are from 2

independent experiments.

FIGURE 6

A) Fluorescent time-course of 2 μM G-actin (10% pyrene) measured in a fluorimeter (red trace).

Polymerization was induced by addition of KMEI (1x) at t = 0 s. Introduction of 1 μM M15(wt) (blue) at t

= 0 s increases in actin polymerization following an inflexion point (arrow), whilst 1 μM M15(jd) (green)

does not (arrow). Reactions contains 70 μM ATP carried over from the actin storage G-buffer. **B)** First

derivative of pyrene fluorescence with respect to time for G-actin + M15(wt)(blue) and G-actin +

M15(jd)(green). **C)** Fluorescent time course of 2 μM G-actin (10% pyrene) induced to polymerize by

KMEI buffer in the absence of free ATP (red trace). G-actin monomers were desalted immediately

prior to use, ensuring they were ATP bound. Addition of 1 μM M15(wt) (blue) triggers an almost

immediate increase in actin polymerization. **D)** Quantification of time to reach half maximal

fluorescence (n = 4).

FIGURE 7

A) TIRFM visualization of actin filaments polymerizing on PEG-biotin-neutravidin functionalized cover

glass. Polymerization of 1 μM G-actin (20% TMR + 10% biotin labelled) was induced by 1 x KMEI (50

mM KCl, 1 mM MgCl$_2$, 1 mM EGTA, 10 mM imidazole, pH 7.0) at t = 0 s, in the presence of 25 μM ATP.

Representative time-lapses shown for: 1 μM G-actin (top), 1 μM G-actin + 1 μM M15(wt) (middle), and

1 μM G-actin + 1 μM mutant M15(jd) (bottom). **B)** Quantification of actin filament density shows delayed

nucleation activity of MYO15 in the presence of ATP. **C)** Kymographs of actin filament elongation. **D**)

Barbed end elongation rates for G-actin alone (red), G-actin + M15(wt) (blue), G-actin + M15(jd) (green).

E + F) Elongation rate data (from D) re-binned as before (E), or after (F) nucleation. The G-actin + KMEI

control data set (from D) is reproduced identically as a comparator in E and F. **G)** Time-lapse of actin

filaments polymerization induced by KMEI at t = 0 s, with no ATP in solution. G-actin (ATP) monomers

were prepared by desalting into ATP-free G-buffer. **H)** Actin filament density shows nucleation activity

of MYO15 is accelerated in the absence of ATP. **I)** Barbed end filament rates in the absence of free

ATP. Reaction deadtimes were typically 50 s and included in quantification. TIRFM images are shown

as inverted grayscale. Scale bars are 10 μm (A,G). Data are from 4 experiments, using 2 independent

protein preparations.

FIGURE S1

A) Whole genome SNP mapping of genomic DNA from ten MPC190 mice exhibiting hearing loss (red, HL) and three hearing (H) littermate controls. The genotype of each mouse is either homozygous for C57BL/6J (black box), C3H (white box) or heterozygous (grey box) for each marker. The analysis defined a 16.5Mb critical interval on Chromosome 11 between markers rs29410974 and rs26906764 (Chr11:50420012-67162951, GRCm38). **B)** Sanger sequencing confirmed the only WGS-identified, coding lesion (arrow) within the critical interval, in exon 17 (ENSMUSE00000244718) of the *Myo15* gene (ENSMUSG00000042678). A nucleotide transition (c.4940A>G) at codon 1647 alters the wild-type aspartate (D) codon to a mutant glycine (G). Example electropherograms are shown for *Myo15*$^{+/+}$, *Myo15*$^{+/jd}$ and *Myo15*$^{jd/jd}$. **C-F)** Minimum ABR detection thresholds for a longitudinal cohort of *Myo15*$^{jd/jd}$ mice (n = 10), *Myo15*$^{+/+}$ (n=10) and *Myo15*$^{+/jd}$ (n=15) littermates at 4, 6, 9 and 12 weeks reveal progressive hearing loss. Data points represent individual mice, with the mean value indicated. **** p < 0.0001; two-way ANOVA with Tukey's multiple comparisons test. **G)** Distortion Product Otoacoustic Emission (DPOAE) responses are greatly reduced at 12 weeks in *Myo15*$^{jd/jd}$ mutants (red, n = 10) at all frequencies tested compared with *Myo15*$^{+/+}$ (black, n = 9) and *Myo15*$^{+/jd}$ (gray, n = 14) littermates. Data are mean ± SD.

FIGURE S2

A) SEM analyses of IHC bundles from *Myo15*$^{jd/jd}$ and *Myo15*$^{+/+}$ littermates at 12 weeks. **B)** Quantification of the tallest (row 1) stereocilia at 12 weeks in IHCs from *Myo15*$^{+/+}$ mice (45 stereocilia from 3 animals) and *Myo15*$^{jd/jd}$ mice (45 stereocilia from 3 animals). **** p < 0.0001, unpaired, two-tailed *t*-test. **C)** SEM analyses of *Myo15*$^{jd/jd}$ OHC bundles at 12 weeks. OHC bundles were qualitatively categorized as "intact", "abnormal/damaged" or "missing". Representative examples are shown. In the "abnormal/damaged" group, stereocilia were typically resorbed from the periphery of the hair bundle. **D)** Quantification of OHC bundle categories at 12 weeks in *Myo15*$^{+/+}$ (1344 OHCs from 4 animals) and *Myo15*$^{jd/jd}$ littermates (972 OHCs from 4 animals). All images and quantification are from the mid-cochlear turn. Scale bars, 1 μm.

FIGURE S3

Immunofluorescence (IF) confocal images of anti-EPS8, anti-WHRN, anti-GNAI3 labelling or anti-GPSM2 (green) in control *Myo15*$^{+/jd}$ and *Myo15*$^{jd/jd}$ IHCs fixed at P14. Phalloidin was used to label F-

236 actin (magenta). Images are representative of data from two independent animals per genotype and

237 antibody combination. Scale bars, 10 µm.

FIGURE 1

FIGURE 2

FIGURE 3

FIGURE 4

FIGURE 5

FIGURE 6

FIGURE 7

FIGURE S1

FIGURE S2

A

Myo15 (+/+)   *Myo15 (jd/jd)*

IHCs

Myo15 (jd/jd)

IHCs

C

Myo15 (+/+)   *Myo15 (jd/jd)*

OHCs   *Intact*   *Intact*

Myo15 (jd/jd)

OHCs   *Abnormal/Damaged*   *Abnormal/Damaged*

B

12 weeks IHCs

****

Height of IHC stereocilia row 1 (μm)

Myo15(+/+)   *Myo15(jd/jd)*

D

12 weeks OHCs

% of OHC bundles

Intact Bundle   Abnormal / Damaged   Missing Bundle

Myo15(+/+)   *Myo15 (jd/jd)*

FIGURE S3