Thalamic bursting and the role of timing and synchrony in thalamocortical signaling in the awake mouse

Peter Y Borden[1*], Nathaniel C Wright[1*], Arthur E Morrissette[2],

Dieter Jaeger[2], Bilal Haider[1], Garrett B Stanley[1]

[1] Wallace H Coulter Department of Biomedical Engineering, Georgia Institute of Technology and

Emory University, Atlanta, GA, USA

[2] Emory University, Department of Biology, Atlanta, Georgia, USA

*Denotes co-first authorship

Abbreviated Title: Thalamic burst timing and synchronization in awake mouse
13
14   With 45 text pages, 9 Figures, 5 Supplemental Figures and corresponding text
15

16   Correspondence: Garrett B Stanley
17   Coulter Department of Biomedical Engineering
18   Georgia Institute of Technology & Emory University
19   313 Ferst Drive
20   Atlanta GA 30332-0535
21   USA

22   Phone: 404-385-5037
23   Fax: 404-385-5044
24   Email: garrett.stanley@bme.gatech.edu

25
26   **Conflict of Interest**: The authors declare no competing financial interests.
27

Acknowledgements: This work was supported by NIH National Institute of Neurological
Disorders and Stroke Grant R01-NS048285 (GBS), R01-NS104928 (GBS), U01-NS094302 (GBS
and DJ), and NIH National Institute of Neurological Disorders and Stroke Pre-doctoral NRSA
NS098691 (PYB). The authors would like to thank Katie Borden, Ilya Kolb, Will Stoy, and Precision
Biosystems Lab at Georgia Tech, for additional support and advice on analysis. Additionally, we
thank all members of the Stanley Laboratory for the Control of Neural Systems at Georgia Tech
for feedback and experimental design which made this work possible

Summary
The thalamus controls transmission of sensory signals from periphery to cortex, ultimately
shaping perception. Despite this significant role, dynamic thalamic gating and the
consequences for downstream cortical sensory representations have not been well
studied in the awake brain. We optogenetically modulated the ventro-posterior medial
thalamus in the vibrissa pathway of the awake mouse, and measured spiking activity in
the thalamus, and activity in primary somatosensory cortex (S1) using extracellular
electrophysiology and genetically encoded voltage imaging. Thalamic hyperpolarization
significantly enhanced thalamic sensory-evoked bursting, yet surprisingly the S1 cortical
response was not amplified, but instead timing precision was significantly increased,
spatial activation more focused, and there was an increased synchronization of cortical
inhibitory neurons. A thalamocortical network model implicates the modulation of precise
timing of feedforward thalamic population spiking, presenting a highly sensitive, timing-
based gating of sensory signaling to cortex.
49

INTRODUCTION

We explore the world through our sensory periphery, where sensors transduce the signals
that ultimately give us perception of the world. The mammalian sensory thalamus gates
information from the periphery to primary sensory cortices, controlling what signals do
and do not make their way to cortex, thus playing a very critical role in sensing. Far from
a static relay, the thalamus is under continuous influence by modulatory inputs from
brainstem and feedback mechanisms from cortex (Sherman and Guillery, 2002;
Sherman, 2005), most extensively explored in the context of sleep and epileptic seizures
(Huguenard and McCormick, 2007; Fogerson and Huguenard, 2016).   The large majority
of detailed thalamic studies focusing on its potential role in sensory signaling have been
either in ex-vivo/slice preparations or in intact/in-vivo preparations under anesthesia,
where prominent thalamic properties such as synchronization of convergent
thalamocortical projections and tonic/burst gating have been shown to strongly boost
signaling under these conditions (Murray Sherman, 2001; Sherman, 2001; Swadlow and
Gusev, 2001; Swadlow, 2002; Lesica and Stanley, 2004; Lesica et al., 2006; Wang et al.,
2010; Stanley et al., 2012; Whitmire et al., 2016). Recent studies exploring the role of
thalamus in the awake, unanesthetized brain have confirmed the critical role of thalamus
in grossly modulating sensory cortex (Halassa et al., 2011; Poulet et al., 2012; Lewis et
al., 2015; Reinhold et al., 2015; Yu et al., 2016; Wright et al., 2021). However, whether
increases in thalamic excitability act to boost cortical sensory-evoked responses in the
awake brain as predicted from ex-vivo and anesthetized studies is unknown.
71
Serving as the primary input to sensory cortex, properties of thalamic activity strongly
shape the spontaneous, baseline activity of cortex, as well as the corresponding cortical
response to ascending sensory inputs. It has been shown that even small changes in
baseline membrane potential have appreciable effects on spontaneous firing of thalamic
neurons (Béhuret et al., 2015), setting the overall tone of synaptic drive to cortex. The
use of pharmacology to directly modulate thalamus (Godwin et al., 1996; Hirata and
Castro-Alamancos, 2010; Poulet et al., 2012) or opto/microstimulation and pharmacology
to indirectly affect thalamus through cortical (Olsen et al., 2012; Mease et al., 2014;
Crandall et al., 2015; Li and Ebner, 2016; Kirchgessner et al., 2020) and subcortical
(Aguilar and Castro-Alamancos, 2005) inputs has further causally revealed the extreme
sensitivity of cortex to overall thalamic drive. One prominent characteristic of neurons in
the thalamus is the presence of T-type calcium channels that are normally inactivated,
but become de-inactivated through prolonged hyperpolarization (Jahnsen and Llinás,
1984a, 1984b; Sherman and Koch, 1986; Suzuki and Rogawski, 1989). Subsequent
depolarizing inputs lead to calcium-mediated "bursts" of action potentials characterized
by transient, high-frequency spiking, which is distinct from "tonic" spiking mediated
through Na+/K+ channel dynamics. Most actively investigated in the context of sleep
states and rhythmic discharge, the role of this mechanism in sensory signaling remains

unclear, despite decades of speculation. At the synaptic level, thalamocortical high
frequency bursting events have been shown to have a significant impact on downstream
cortical activation (Swadlow and Gusev, 2001), where thalamic bursts are associated with
an amplified post-synaptic response in recipient cortical neurons, thought to arise
primarily from properties of the thalamocortical synapse. Coupled with sensitivity of
cortex to the timing of thalamic inputs via the "window of opportunity" established by the
disynaptic feedforward cortical inhibition (Pinto et al., 2000, 2003; Wehr and Zador, 2003;
Wilent and Contreras, 2004, 2005; Gabernet et al., 2005; Isaacson and Scanziani, 2011),
the gating of thalamic signaling through the aggregate effects of all these properties
across the thalamic population is hypothesized to serve a critical role in processing
sensory information (Crick, 1984; Pinto et al., 2000; Sherman and Guillery, 2002; Wilent
and Contreras, 2004; Sherman, 2005). However, this has not been investigated
extensively in the intact brain due to the lack of methodological approaches to precisely
measure and manipulate thalamic properties. To date, no studies have utilized temporally
precise, repeatable and reversible modulation to precisely measure the effects of thalamic
burst/tonic gating in awake cortex at the level of population signals and single neuron
firing.

Here, we directly determined how thalamic gating properties control sensory-evoked
thalamic and cortical responses in the vibrissa pathway of the awake, head-fixed mouse.
Instead of driving or silencing neural activity, optogenetic manipulation was used to
modulate thalamus while recording extracellular thalamic and cortical activity and
acquiring widefield cortical voltage imaging, using the voltage indicator ArcLight (Jin et
al., 2012; Borden et al., 2017). We found that baseline thalamic firing rate was
surprisingly invariant to optogenetic thalamic hyperpolarization through a transition from
tonic to burst firing activity, and the baseline activity in S1 cortex was correspondingly
invariant to thalamic hyperpolarization, following a brief transient increase in firing activity.
In response to sensory stimuli, the thalamic hyperpolarization significantly enhanced the
sensory-evoked thalamic bursting; however, the magnitude of the response in S1 cortex
was not amplified, but instead slightly attenuated relative to baseline, despite the burst-
dominated thalamic input. Notably, the sensory evoked response was also significantly
more brief and spatially focused, and accompanied by an increase in the synchronization
of the putative cortical Fast-spiking (FS) inhibitory neurons. A thalamocortical network
model replicated these findings, and implicated changes in the precise timing and
synchronization across the thalamic population as a likely mechanism underlying the
experimental observations. Taken together, the results here point to timing rather than
response magnitude as a fundamental feature of the thalamocortical circuit, presenting a
dynamic, timing-based gating of sensory signaling to cortex.

RESULTS

All experiments were conducted in the vibrissa pathway of the mouse (Figure 1A). To
directly test the predictions of previous in-vitro and anesthetized work, we conducted the
first experiment in the isoflurane-anesthetized mouse (illustrated in Figure 1B). In the
lightly anesthetized mouse, we utilized extracellular electrodes to record whisker-evoked
spiking activity in VPm thalamus, in the presence (LED, ~17mW/mm^2, 590nm) and
absence (Control) of a hyperpolarizing optogenetic modulation of excitatory thalamic relay
neurons expressing halorhodopsin (eNphR3.0, see Supplemental Figure S1), through a
small fiber optic cable attached to the electrode (Figure 1B). Note that the inhibitory opsin
was engaged with a constant illumination at a range of relatively low light levels, to induce
sustained hyperpolarization, as opposed to complete inactivation. Note also that the total
amount of light delivered here was in a range that has been previously shown to not
induce significant heating of the surrounding tissue (see Methods), and control
experiments indicate that there is no effect of the light alone in the absence of opsin
expression (Supplemental Figure S4). Although the in-vivo optogenetic implementation
precludes precise knowledge of the degree of hyperpolarization of the thalamic neurons
due to variations in opsin expression, position of optic fiber relative to cells, etc., a
separate set of in-vitro, brain slice experiments showed that VPm neurons were
hyperpolarized by ~15-25 mV for the light levels utilized (see Supplemental Figure S2).
Given that in the in-vitro experiments, light was presented more directly to the VPm
neurons expressing halorhodopsin, and that in the in-vivo experiments the presentation
of light did not completely silence the neurons, it is likely that the in-vitro experiments were
an upper-bound for the in-vivo case, and the actual hyperpolarization induced in-vivo was
less than that of the in-vitro experiments.

Neurons in VPm thalamus responded to a single, computer-controlled punctate whisker
stimulation (1200 deg/sec, sawtooth waveform) with a brief, transient sequence of action
potentials (Figure 1D, see Methods for single unit and multi-unit classification).
Optogenetic thalamic hyperpolarization significantly increased the sensory-evoked
thalamic bursting, as shown in Figure 1D (all spikes in black, putative T-type calcium
"burst" spiking in red, classified according to the criterion in Figure 1C). Shown in the
insets are the low baseline firing rates in the anesthetized condition. Overall, there was
an increase in the evoked response from the Control to the LED condition (Figure 1E left,
black, p=0.0046, paired Wilcoxon signed-rank, n=29 units), in large part driven by the
increase in sensory evoked thalamic bursting (Figure 1E left, red, p=4.5e-4, paired
Wilcoxon signed-rank, n=29 units). In a subset of experiments, we recorded the
downstream cortical multiunit activity (Figure 1E right), and found a corresponding
increase during the period of hyperpolarization (n=102 trials, 2 recordings, 1 mouse).
Additionally, we simultaneously recorded cortical S1 activation at a meso-scopic scale
with optical voltage imaging in primary somatosensory (barrel) cortex using a Genetically

Engineered Voltage Indicator (GEVI, ArcLight, Borden et al., 2017) during optogenetic
hyperpolarization of thalamus (see Methods). This approach enables the characterization
of the overall spatiotemporal changes in S1 due to the thalamic manipulation
(Supplemental Figure S4). The imaging was focused at a cortical depth consistent with
layer 2/3 (see Methods).   The fluorescence imaging of cortical S1 confirmed a
corresponding amplification of the sensory evoked response during thalamic
hyperpolarization (raw %dF/F$_o$), as shown in an example session for the fluorescence
images in Figure 1F and further quantified across the larger dataset in the left panel of
Figure 1G (normalized relative peak evoked response, ~60% increase, 1G inset raw
relative peak evoked %dF/F$_o$, p=0.017, paired Wilcoxon signed-rank, n=13 recordings
across 9 mice). There was an amplification in the peak response, followed by a relatively
fast transient decrease in fluorescence, followed by a gradual return to baseline over
approximately ~200ms. The results here using optogenetic manipulation of thalamus and
wide-field voltage imaging of cortex are consistent with previous observations of
spontaneous thalamic bursting and the impact on downstream synaptic targets in cortical
layer 4 (Swadlow and Gusev, 2001). Importantly, the amplification in the peak response
observed here under these conditions reflects what would have been expected from the
observations in thalamus, if the cortical sensory representation were strongly by the
feedforward thalamic bursting.
188
Thalamic Hyperpolarization and Ongoing Thalamic and Cortical Activity in the
Awake Brain
In a next set of experiments, we sought to characterize the influence of thalamic
properties on gating of signaling to cortex in the awake mouse. Before turning to the
sensory-evoked responses, we first probed the effect thalamic hyperpolarization has on
baseline, ongoing activity that is significantly higher in the awake state as compared to
under anesthesia. The baseline firing rate in the awake state is significantly higher than
in the anesthetized condition, and we thus expect to see some effect of the optogenetic
hyperpolarization on baseline firing. An example single-unit VPm recording (experimental
setup in Figure 2A) is shown in the raster plot in Figure 2B. VPm neurons in awake mice
exhibited moderate ongoing firing activity (4.9 Hz for this example). Upon presentation of
a ramping optical input to thalamus (LED, 590nm, ~17mW/mm^2), the spiking activity
underwent a qualitative change to a more sparse pattern of multiple action potentials,
typical across all recorded cells. Bursts were present in the baseline (Control, no light)
condition, but substantially more prevalent in the presence of the hyperpolarizing
optogenetic input. This effect is summarized across all clearly identified single-units in
Figure 2C (n=51 single VPm units), showing the average instantaneous firing rate for all
spiking in black, and for the putative burst spiking in red (bands represent +/- 1 standard
error of the mean, SEM). This qualitatively shows that following an initial decrease in
firing rate at the onset of the light ramp, the VPm firing activity briefly rose above pre-

stimulus baseline firing rate, before returning to a level near baseline. This was
accompanied by an increase in the burst firing rate, suggesting that the return to pre-
stimulus baseline firing rate was due to a compensatory transition from tonic to burst
firing. Quantitatively, following the initial transient, the single-unit firing rate was relatively
invariant from the Control (0 mW/mm^2) to the hyperpolarized condition, with a slight
increase for only the strongest of LED conditions (Figure 2D left, single-unit, Control vs
LED p=0.2, Control vs Double LED p=0.02, unpaired two-sample t-test, n=54 SU
recordings for Control, and n=51 for each of the LED conditions; ~17mW/mm^2 - LED or
~35mW/mm^2 - Double LED). Note that the firing rate in the LED conditions was calculated
from the 250-750ms window, after the initial transient decrease in firing rate. There was
a corresponding significant increase in the firing rate associated with bursting (red,
Control vs LED p=3e-13, Control vs Double LED p= 1e-15, unpaired t-test), indicating a
compensatory shift from tonic to burst firing with the thalamic hyperpolarization resulted
in a significant increase in the burst ratio metric from the Control to hyperpolarized
condition (Fig. 2D, right, Control vs LED p=9e-13, Control vs Double LED p=1.8e-11,
Wilcoxon rank-sum test). Further examination showed that increasing hyperpolarization
also increased the number of spikes per burst (illustrated in left of Figure 2E), which
partially offsets the loss of tonic spiking (right of Figure 2E, Control vs LED p=2.3e-5,
Control vs Double LED p=1.7e-6, Wilcoxon rank-sum test, n=54 for the Control and n=51
for both LED conditions).
229
Although it is not surprising that the thalamic burst mechanism was engaged by the
optogenetic hyperpolarization, it is surprising that the net baseline firing rate was not
decreased, but was instead relatively invariant to the thalamic hyperpolarization. This
effect was not as apparent in the anesthetized condition when the baseline thalamic firing
rate is near zero, but emerges in the awake state where baseline thalamic firing rate is
significantly higher. To investigate possible causes, we constructed an integrate-and-fire
or burst (IFB) model of thalamic firing that has been utilized in previous studies (Smith et
al., 2000; Lesica and Stanley, 2004; Lesica et al., 2006), where the resting membrane
potential was set by the level of activation of the hyperpolarizing opsin (see Methods).
When the model neuron was set to have a spontaneous, baseline firing rate consistent
with our observations (~5 Hz), the effect of the hyperpolarizing opsin activation indeed
replicated this phenomenon (Figure 2F). An initial decrease in firing rate at the onset of
the light ramp was followed by a return to pre-light baseline level of firing (compare with
Figure 2C). Moreover, as in the experimental observation, the return to the baseline firing
rate in the hyperpolarized condition is dominated by the increase in bursting activity in the
IFB model (Figure 2F, red). The exact combination of the baseline activity and intensity
of the hyperpolarizing input strongly influenced the net resultant effect (Supplemental
Figure S3). Prolonged periods of halorhodopsin activation have been shown to potentially
alter the reversal potential of chloride (Raimondo et al., 2012). However, due to the short

timescales of LED stimulation used in this work (1-1.5s), and the large transition to burst
firing, the effect shown here is more likely driven by low voltage activation of T-type
calcium channels, further reinforced by the IFB model.
252
To uncover the downstream effects of the above observations in VPm, we performed the
same optogenetic manipulations during multi-electrode recording of single-unit activity in
S1 in the awake mouse (Figure 3A). Note that for this analysis, while these S1 recordings
were combined across cortical layer and cell type, subsequent analyses demonstrate
similar trends across cell types and for targeted layer 4 recordings (see Figure 7 and
Supplemental Figure 5). Shown in the top panel of Figure 3B is a raster plot from one
example session (10 trials each from 15 simultaneously recorded units), in response to
the optogenetic hyperpolarization of thalamus with the LED (again 590nm, 17mW/mm^2).
This example demonstrates a clear transient increase in spiking with the thalamic
hyperpolarization, typical of the cortical recordings. The bottom panel of Figure 3B shows
the aggregate single-unit firing rate, with a clear transient increase in firing rate at the light
onset (bands represent +/- 1 SEM), with a gradual return to the baseline, pre-
hyperpolarization level of firing rate at steady-state. Figure 3C summarizes this trend,
showing the mean single-unit firing rate across recorded cortical single-units in the
Control condition (0mW/mm^2) versus activation of halorhodopsin in thalamus (LED -
17mW/mm^2 or Double LED - 35mW/mm^2), for the initial increase in firing rate during a
transient period (300-350 ms, open symbols) and the subsequent return to baseline firing
rate at steady state (700-750 ms, closed symbols). There was a significant transient
increase in the cortical firing rate across both led conditions (transient, Control vs LED
p=5.4e-14, Control vs Double LED p= 1.4e-15, Wilcoxon signed-rank test, n=118 units).
After the transient increase, we observed a subsequent return to near baseline, either
with no net effect or a marginal increase in the rate (steady-state, Control vs LED p=0.2,
Control vs Double LED p= 0.013, Wilcoxon signed-rank test). Interestingly, following
thalamic hyperpolarization, the cortical firing rate appeared to begin its initial transient
increase while the thalamic activity remained relatively low (Figure 2), which could reflect
an increased sensitivity of the thalamocortical synapse following the initial period of
quiescence (Swadlow and Gusev, 2001; Swadlow et al., 2005).
280
Taken together, the results here suggest that in response to the optogenetically induced
thalamic hyperpolarization, the thalamic VPm neurons in the awake mouse exhibit a
transient decrease in baseline firing rate, followed by a return to approximately pre-
hyperpolarization level that is at least partially explained by a trade-off between tonic and
burst firing, while the downstream activity in cortical S1 exhibits a transient increase in
baseline firing rate, followed by an eventual return to approximately the original baseline
rates.
288

Thalamic Hyperpolarization and Sensory Evoked Thalamic and Cortical Activity
We next recorded the sensory evoked single-unit activity in VPm thalamus in the
presence and absence of thalamic hyperpolarization in the awake mouse (Fig. 4A).
Neurons in VPm thalamus responded to a single, computer-controlled punctate whisker
stimulation (1200 deg/sec, sawtooth waveform) with a brief, transient increase in single-
unit activity, which was significantly reshaped by thalamic hyperpolarization (Fig. 4B;
PSTH of all spikes (black) across single-units, 2 ms bins, LED approximately 17mW/mm^2,
n=54 VPm single units for control, n=51 for LED). Also shown are the PSTHs comprised
of putative burst spikes only (red), present in the Control condition but appearing to be
significantly increased in the LED condition. To further quantify this effect, we tested a
range of LED intensities to determine the impact of various levels of thalamic
hyperpolarization on the evoked response (Figure 4C; LED approximately 17mW/mm^2,
and Double LED approximately 35 mW/mm^2). Overall, there was a significant increase in
sensory-evoked VPm bursting with thalamic hyperpolarization (Figure 4C, left, burst
spikes (red), Control vs LED p=0.003, Control vs Double LED p=1.6e-4, unpaired
Wilcoxon rank sum, n=54 VPm single-units for Control and n=51 for both LED conditions).
However, for the same set of recordings there was a surprisingly invariant overall
sensory-evoked response magnitude across VPm single units (Figure 4C, left, all spikes
(black), Control vs LED p=0.9, Control vs Double LED p=0.4, unpaired two sample t-test).
There was a corresponding significant increase in single-unit burst ratio (Figure 4C, right,
Control vs LED p=1.3e-5, Control vs Double LED p=2.4e-6, Wilcoxon rank-sum, n=54
VPm single-units for Control and n=51 VPm single-units for both LED conditions). Taken
together, single-unit VPm analyses reveal a boosting of the sensory evoked thalamic
bursting with thalamic hyperpolarization, yet a surprisingly invariant overall mean evoked
rate.
314
Note that upon more careful inspection of the VPm PSTHs, thalamic hyperpolarization
results in a qualitative decrease in the sensory-evoked response in the first 10 ms, and a
subsequent increase in the sensory-evoked response at later times. This could indicate
changes in synchronous thalamic firing over the course of the sensory response, which
is a critical factor in determining the downstream cortical response. To explicitly quantify
this, we analyzed a large subset of the VPm recordings that were obtained from multi-
electrode arrays. The multi-electrode arrays yielded a portion of simultaneously recorded
VPm single-units, thus enabling us to analyze pair-wise properties. Figure 4D illustrates
the formation of the spike cross-correlogram (CCG) from the spiking activity of a
simultaneously recorded pair of neurons (see Methods). We applied this analysis across
pairs of simultaneously recorded VPm pairs, restricting the spiking activity to three non-
overlapping windows following the whisker stimulus (0-10ms, 10-20ms, and 20-30ms).
The resulting grand mean CCGs for the Control and LED condition are shown in Figure
4E. Qualitatively we observe a decrease in the central peak of the CCG for the earliest

post-stimulus time window (0-10ms, left) from the Control to the LED condition, and very
little change for the other two time windows (10-20ms middle, 20-30ms right).   We
quantified this observation by computing the area under the CCG in the central window [-
5ms, +5ms], representing the amount of synchronous spiking across the pairs. Most
notably, the LED induced a decrease in the synchronous firing for the earliest post-
stimulus time window (0 – 10 ms), when synchronous spike rates were generally highest
(Control vs. LED, $p < 1e-3$, n=187 pairs from 46 VPm units, comparison of confidence
intervals from re-sampling with replacement, see Methods).   Synchronous spike rates
also changed in the two later response windows, when synchronous rates were generally
lower, with a slight increase for the (10-20 ms) window and a slight decrease for the (20-
30 ms) window (10 – 20 ms: $p < 0.01$; 20 – 30 ms: $p < 0.01$, n=187 pairs from 46 VPm
units).   Thus, taken together, the VPm data reveal a significant boosting of sensory-
evoked burst firing with thalamic hyperpolarization, accompanied by a relatively invariant
overall sensory-evoked response magnitude, and a surprising decrease in the synchrony
of the prominent early thalamic response.
344
To examine the downstream consequences of the observed changes in the sensory-
evoked response in thalamus, we recorded cortical S1 activation with wide-field optical
voltage imaging during optogenetic hyperpolarization of thalamus in the awake mouse
(illustrated in Figure 5A, see Methods). Note further that fluorescent imaging approaches
like this are differential in nature (responses are based on relative not absolute changes
in fluorescence), and are thus not well suited to capture overall absolute levels of ongoing
background activity, but instead are targeting evoked responses relative to background.
Based on previous findings (Swadlow and Gusev, 2001), the results in the anesthetized
mouse, and the significant boosting of sensory-evoked bursting in thalamus with thalamic
hyperpolarization in the awake mouse, we expected to see a corresponding amplification
of the cortical sensory-evoked response. Shown in Figure 5A is a trial averaged sequence
of the sensory-evoked fluorescence images. Following the deflection at time 0, a localized
change in fluorescence began to emerge 10-15 ms later. Surprisingly, thalamic
hyperpolarization did not result in an increased cortical response, but instead induced an
apparent modest decrease in sensory evoked S1 activity. As quantified in Figure 5B, the
peak fluorescence showed an approximately 20-40% decrease with this level of thalamic
hyperpolarization (Control vs LED p=0.020, Control vs Double LED p=0.0039, paired
Wilcoxon sign rank test, 2 mice over 9 sessions for LED conditions). Note that the right
panel of Figure 5B is the same data as in the left panel, with each dataset normalized to
the Control condition, to show a percent decrease in the peak fluorescence.
365
Finally, in a separate set of experiments, we hyperpolarized thalamus while
simultaneously recording the S1 Cortical SU responses using a multichannel electrode
(illustrated in Figure 5C). The corresponding PSTHs for responses to a punctate whisker

stimulus (1200 deg/sec, sawtooth waveform) are shown in Figure 5D (Control
(0mW/mm^2) vs. LED (17mW/mm^2) conditions). The average absolute evoked cortical
response remained approximately invariant during LED conditions across all cortical
recorded units (Figure 5E, left, 5 to 100ms post-stimulus, n=118). However, the evoked
cortical spiking relative to baseline decreased with increasing LED intensities (Control vs
LED p=0.20, Control vs Double LED p=0.034, paired Wilcoxon signrank, n=118 units,
across 3 mice), consistent with the GEVI recordings (Figure 5B), which are relative
measures by construction. Note that the recorded cortical activity here was aggregated
across cell type and cortical layer, but subsequent analyses will address cell type and
layer (see Figure 6 and Supplemental Figure S5). Importantly, similar trends were
observed through analysis restricted to putative layer 4 cortical neurons (Supplemental
Figure S5A and B).

To summarize, given the boosting of thalamic bursting induced by thalamic
hyperpolarization that serves as the primary feedforward inputs to cortical S1, it was
surprising to observe in the awake mouse not an increase, but instead a sensory evoked
cortical response that was invariant in absolute amplitude and decreased relative to
background activity.

Thalamic Bursting and the Spatiotemporal Shaping of the Sensory Evoked Cortical
Response

The GEVI imaging of cortex enables further investigation of the spatiotemporal
characteristics of the observed phenomenon. Figure 6A shows the imaging frame
associated with the peak sensory-evoked fluorescence, for the Control (top) and thalamic
hyperpolarization (LED, bottom) conditions. What is apparent qualitatively from the
images is that there was an overall attenuation of the cortical evoked response with
thalamic hyperpolarization, as previously shown in Figure 4. Less immediately obvious
is the effect of the thalamic hyperpolarization on the spatial extent of the sensory evoked
cortical activation, as an overall loss in amplitude would trivially produce an apparent loss
in spatial activation, as illustrated conceptually in the top panel of Figure 6B – often
referred to as the "iceberg" effect. The bottom panel of Figure 6B shows conceptually the
alternative - that the reduction in spatial area of activation may not just be due to the
reduction in amplitude of the evoked response, but instead can reflect a spatial
sharpening of the sensory evoked response with increasing thalamic hyperpolarization
after accounting for the decrease in peak amplitude of fluorescence. For the GEVI
recordings here, the latter is indeed the case - this is summarized as a reduction in the
evoked normalized area of activation in the left panel of Figure 6D (Control vs LED
p=0.025, Control vs Double LED p=0.046, paired t-test, 2 mice over 9 recording sessions
for LED conditions).

409
In addition to the spatial characteristics of the sensory evoked response revealed by the
voltage imaging, the voltage imaging is also useful in revealing relatively fast temporal
dynamics of the aggregate cortical activity. Figure 6C shows the integrated fluorescence
signal within a barrel column in the presence (LED condition) or absence (Control) of light
input to activate halorhodopsin in thalamus, giving a picture of the temporal dynamics of
the sensory-evoked clear response to the punctate whisker stimulus (2 mice over 9
recording sessions). We again found a decrease of the sensory evoked response (note
that the fluorescence in Fig. 5C is normalized) despite a significant boosting of the
sensory-evoked thalamic bursting (Fig. 4). Also apparent is the presence of a more rapid,
post-peak decay in the fluorescence with thalamic hyperpolarization, and a prolonged
period of sub-baseline fluorescence before returning to baseline after several hundred
milliseconds. To quantify this property of the sensory evoked response, we calculated the
temporal width of the evoked response as the time between the fluorescence peak and
the first return to baseline level (defined as peak-to-baseline, Figure 6C), as shown in the
middle panel of Figure 6D, revealing a clear narrowing/shortening of the temporal
response as a function of thalamic hyperpolarization from ~150ms in the Control
condition, to ~67ms in the thalamic hyperpolarized condition (Control vs LED p=0.0059,
Control vs Double LED p=0.0019, unpaired t-test, LED conditions n=9, Control n=8
recording sessions, note one Control recording did not return to a pre-stimulus baseline).
To quantify the prolonged post-stimulus undershoot of activity, we calculated the overall
fluorescence in a window from 120 to 400ms post-stimulus (defined as mean recovery in
Figure 6C, Control vs LED p=0.016, Control vs Double LED p=0.020, paired t-test, n=9
recording sessions), as shown in the right panel of Figure 6D. The increasingly negative
relative fluorescence in this window revealed a clear increase in the undershoot with
increasing thalamic hyperpolarization. Taken together, these S1 voltage imaging results
show that in addition to the attenuation of the sensory-evoked response with thalamic
hyperpolarization, there is a corresponding sharpening of the spatial activation of S1 and
a temporal narrowing/shortening in the form of a more transient evoked response with an
inhibitory undershoot.
439
Thalamic Bursting and Cortical Spike Timing
Taken together, the results from Figure 6 revealed that the thalamic hyperpolarization
strongly shapes both the spatial and temporal aspects of the cortical sensory evoked
response in the awake brain. To further probe this phenomenon, we parsed our cortical
single-units into putative excitatory neurons (regular spiking units, RSUs) and putative
inhibitory neurons (fast spiking units, FSUs). Specifically, single-unit activity was
identified (see Methods) and parsed based on RSU and FSU classification and combined
across cortical layers, the result of which is shown in Figure 7A. Note that while this
analysis involved combining recordings across the electrode array, we conducted a

parallel analysis restricted to a smaller population of identified cortical layer 4 RSUs and
FSUs (see below, and Supplemental Figure S5). The top panel of Figure 7 shows the
waveform for typical putative RSU (red) and FSU (blue), where the solid line represents
the mean waveform, and the band represents the standard deviation around the mean.
The bottom panel shows the distribution of the time from the trough to the peak for the
recorded waveforms. A threshold was set at 0.4 ms based on prior literature ((Guo et al.,
2017), see Methods), for classifying the individual units into RSUs or FSUs.
456
Apparent in the aggregate PSTH is the expected higher baseline/ongoing activity and
sensory evoked responses in the FSUs compared to the RSUs, as shown in Figure 7B
(~17 HZ for FSU, n=32; ~3 Hz for RSU, n=86), along with several other observations
quantified in Figure 7C. As shown in the left panels of Figure 7C, in response to the
thalamic hyperpolarization (LED ~17 mW/mm^2, double LED ~35 mW/mm^2), there was a
relatively modest increase in baseline (pre-stimulus, -50 – 0 ms) firing rate for RSUs
(Control vs LED p=5.3e-4, Control vs Double LED p=0.012, Wilcoxon signed-rank test,
n=86), while FSUs showed no significant change (Control vs LED p=0.15, Control vs
Double LED p=0.30, Wilcoxon signed-rank test, n=32). The response to the punctate
sensory stimulus was a relatively short latency, transient increase in firing activity in both
cell types, appearing to be relatively unchanged in overall magnitude with thalamic
hyperpolarization, further quantified in the middle panels of Figures 6C (early response,
5-50ms post stimulus). We observed a slight increase in the RSU evoked firing rate (less
than ~10% increase, Control vs LED p=0.067, Control vs Double LED p=0.0082,
Wilcoxon signed-rank test, n=86); however, FSUs displayed no statistical difference
(Control vs LED p=0.30, Control vs Double LED p=0.45, Wilcoxon signed-rank test,
n=32).
474
Although the overall magnitude of the early phase of the sensory evoked response was
relatively unchanged by the thalamic hyperpolarization for both the FSUs and RSUs,
there was a qualitative change in shape of the PSTH, with an increased latency to peak
in the early phase, and change in the later phase of the evoked response. Specifically,
thalamic hyperpolarization induced a reduction in the later phase of the cortical response,
and a dip below the baseline firing rate in both the FSUs and RSUs, as shown in the
PSTHs in Figure 7B. This was reflected in a decrease in the late phase (60-100ms) of the
evoked response from the Control to the LED condition as shown in the right panels of
Figure 7C (RSU - Control vs LED p=0.017, Control vs Double LED p=0.014, n=86, FSU
Control vs LED p=2.2e-4, Control vs Double LED p=4.5e-4, n=32, Wilcoxon signed-rank
test).
486
The preceding analysis was conducted by aggregating cortical units across probe
recording sites, combining neurons across cortical layers. However, as cortical layer 4 is

the primary recipient of synaptic input from VPm projections, we repeated the above
analyses for putative cortical layer 4 neurons based on analysis of the multi-unit activity,
local field potential, and corresponding current-source density (see Methods). In the
cortical layer 4 analysis, we observe very similar results to those shown for the larger
aggregate dataset, and the trends we observed were qualitatively the same as shown in
Figure 7 (see Supplemental Figure S5).
495
The changes in timing and shape of the PSTHs of the cortical neurons suggest the
possibility that thalamic hyperpolarization could affect synchronization within the cortical
network. We analyzed synchrony across simultaneously recorded cortical single-units.
Synchrony in firing across a pair of Neurons 1 and 2 is evaluated by examining the spike
times of Neuron 2 relative to a particular spike of Neuron 1, across all spikes of Neuron
1, as illustrated in Figure 8A (see Methods). This forms the spike cross-correlogram
(CCG), from which the synchrony is calculated as the integrated area within a +/- 7.5 ms
window (see Methods, (Wang et al., 2010; Whitmire et al., 2016)), as illustrated in the
bottom of Figure 8A. Due to the sensitivity of the synchrony metric to firing rate, we only
examined pairs with a robust measurement (more than 50 synchronous events) to control
for measurement accuracy (experimental results were invariant with different thresholds,
see Methods), which restricted the analysis to FSUs in our dataset. Figure 8B shows
rasters from an example pair of FSUs in the Control and double LED conditions (top),
along with the spikes identified to be synchronous by this criterion (bottom). Figure 8C
shows the aggregate cross-correlograms across 99 FSU pairs for the sensory evoked
response, calculated from the spiking in the 100 ms window following the delivery of the
sensory stimulus. There is an increase in the concentration of mass around 0 lag with
thalamic hyperpolarization, which is indicative of an increased synchrony. This is
summarized in Figure 8D, showing a significant increase across FSU units in the
measured synchrony (Control vs LED, p=1.5e-8, Control vs Double LED, p=3.5e-13,
Wilcoxon signed-rank test, n=99 FSU pairs). For comparison, the synchrony was also
computed for the ongoing, spontaneous activity, revealing the synchronizing effect of the
transient sensory input. These results were qualitatively similar for various synchrony
window sizes (not shown).
520
521 **Modeling of the thalamic burst driven cortical E-I circuit**
522
We next sought to understand the mechanistic basis of two key experimental results: 1)
the nearly invariant absolute S1 sensory response in the thalamic hyperpolarized (LED)
condition in the awake mouse, despite the increase in sensory-evoked burst spikes in
VPm; and 2) the increase in sensory-evoked cortical synchrony in FS neurons in the
thalamic hyperpolarized (LED) condition. To explore the potential role of various thalamic
and cortical mechanisms, we constructed a simple model of the thalamocortical network,

as described previously (Wright et al., 2021). This network mimics the numerical
expansion of neurons at the thalamocortical junction, and incorporates several known
properties of thalamocortical and intracortical connectivity (see Methods and
Supplemental Information on Cortical E-I Modeling). We previously tuned model
parameters to produce qualitatively realistic velocity tuning curves (Wright et al., 2021),
using those parameters here. The network was composed of a single model VPm
"barreloid" (40 independent spike trains) projecting to a single cortical barrel "column",
modeled as an interconnected network of 800 excitatory and 100 inhibitory single-
compartment leaky integrate-and-fire neurons with clustered connectivity (Litwin-Kumar
and Doiron, 2012; Bujan et al., 2015; Wright et al., 2017b, 2017a). To create the model
input, ongoing and sensory-evoked VPm tonic and burst spikes were generated based
on the empirical VPm PSTHs (Figure 9A, top two rows, see Methods). Generating
thalamic input spikes in this manner qualitatively reproduced the high rate of synchronous
spikes in the short-latency response window. Across conditions the rate of synchronous
spikes in the model VPm input matched those of our experimental observations,
confirmed using the approach described above for the empirical VPm data, with a slight
but significant decrease in the synchronous spiking in the short-latency response window
from Control to LED (Fig. 9B), consistent with what we observed experimentally. Because
we sought to investigate the effects of such changes in synchronous thalamic spiking and
single-neuron bursting on downstream network activity, we did not impose activity-
dependent depression at the thalamocortical synapse. Given the very short intervals
between burst spikes, the relatively slow recovery of a given thalamocortical synapse
following the first spike in a burst would in principle diminish the efficacy with which
subsequent spikes in the burst drive postsynaptic targets. Yet we show below that this
consideration is not required to match experimental observations. For a more detailed
description of the model, see Supplemental Information on Cortical E-I Modeling.

This model network succeeded in qualitatively reproducing the two results identified
above. First, despite the substantial increase in burst spikes in the thalamic input driving
the model cortical network (constructed to match empirical rates – Figure 9A, top), the
network cortical response was relatively invariant with thalamic hyperpolarization, with
only a very slight increase for the LED condition (Figure 9A, bottom two rows, and Figure
9C). Note that unlike the experimental data, synchrony measures from simulated cortical
activity were not limited by lack of firing events, enabling synchrony measurements across
both excitatory and inhibitory sub-populations. We found that the evoked pairwise
synchrony of S1 neurons was higher in the LED condition for both excitatory and inhibitory
model neurons (Fig. 9D).   In short, the mechanisms incorporated in this simple model
were sufficient to predict the two key experimental results.

Having reproduced these results, we next sought to test the hypothesis that the nearly
invariant mean cortical response despite the boosting of thalamic bursting represented
the net effect of two opposing mechanisms: enhanced thalamic bursting in the LED
condition supporting more robust thalamocortical synaptic drive, counterbalanced by a
reduction in the rate of spikes that are synchronous across thalamic neurons. To this
end, we manipulated the rate of synchronous thalamic input spikes, while maintaining the
mean rate of tonic and burst spikes, analogous to our previous work (Wang et al., 2010;
Wright et al., 2021). Specifically, an artificial condition was simulated in which tonic spike
times were probabilistically shifted away from the early PSTH peak, such that the peak
firing rate of this manipulated condition approximately matched that of the LED condition,
referring to this manipulation as "randomized" (Fig. 9E, top). The grand VPm input CCG
was again generated, confirming that this manipulation reduced the rate of synchronous
spikes in the input spike trains (Fig. 9F). Because the rate of tonic spikes in the 30 ms
post-stimulus window was not altered and used the burst spike distribution from the
Control condition, this enabled the effects of such changes in synchronous spiking alone
to be inferred, without a possibly counterbalancing increase in bursting. In contrast to the
LED condition, shifting even small percentages of tonic VPm spikes (i.e., reducing the
rate of synchronous VPm spikes) resulted in a significant decrease in cortical network
response rates (Fig. 9G) and synchrony (Fig. 9H), with both measures decreasing
monotonically with increasing percentage of shifted VPm spikes. These simulations thus
support the notion that while thalamic bursting per se likely provides highly efficacious
synaptic drive to cortex, the loss of short-latency, synchronous tonic firing in the LED
condition counterbalances this effect, resulting in a nearly invariant S1 response. Further,
the trends observed in cortical synchrony (Fig. 9H) suggest thalamic bursts play an
important role in synchronizing cortical neurons.
593
Taken together, the modeling results support the notion that cortical sensory
representations are highly sensitive to thalamic firing. The degree of i) thalamic bursting
in individual neurons and ii) synchronous spiking across the thalamic population together
determine the efficacy with which VPm spiking drives S1.   This predicts the
counterintuitive empirical result that the enhanced VPm bursting induced by
hyperpolarization does not result in a substantially boosted S1 response.
600
601

DISCUSSION

The sensory thalamus controls the flow of signaling from the periphery to cortex, ultimately gating what we do and do not perceive about the outside world. Despite its critical role in sensing, how this circuit controls signaling remains poorly understood. Here, through a range of experimental approaches in the awake, head-fixed mouse, we show that optogenetic thalamic hyperpolarization significantly enhances sensory-evoked bursting, yet the baseline thalamic firing rate and sensory evoked magnitude are both relatively invariant. Sensory cortex subsequently exhibits a surprisingly invariant absolute evoked response despite the potent thalamic burst input, instead demonstrating increased timing precision, a focusing of spatial activation, and increased synchrony of spiking. Thalamocortical network modeling further supports the assertion that the bursting-induced changes in thalamic spike timing and thalamic population synchrony are sufficient to explain the increase in cortical synchronization and the invariant cortical response amplitude, respectively. The findings here present a highly sensitive, timing-based gating of sensory signaling to cortex.

Thalamocortical Gating of Sensory Signals

One surprising observation was that the absolute cortical sensory evoked response amplitude was invariant to thalamic hyperpolarization in awake mice, despite a significant increase in thalamic bursting. At first glance, this appears to contradict previous work clearly demonstrating the increased efficacy with which bursts in individual thalamic neurons impact monosynaptically-connected cortical neurons at the single synapse level (Swadlow and Gusev, 2001). Importantly, however, the efficacy with which a population of thalamic neurons drives cortical spiking depends on relative spike timing both within a given thalamic neuron (i.e., tonic vs. burst) and across convergent thalamic neurons (i.e., degree of synchronization). We found that while thalamic hyperpolarization promoted single-neuron bursting, there was a surprising decrease in synchronous thalamic firing in the critical early sensory-evoked response. The nearly invariant cortical response makes sense in light of this observation; cortex is extremely sensitive to thalamic spike timing over this 10ms period established by the disynaptic inhibition mediated "window of opportunity" (Pinto et al., 2000; Wilent and Contreras, 2004, 2005; Gabernet et al., 2005). The findings here thus coexist with the potency of thalamic bursts in driving post-synaptic cortical responses at the single synapse level.

While the absolute cortical response amplitude was nearly invariant to thalamic hyperpolarization, the timing precision and synchrony of the cortical response were enhanced by our manipulation. Such changes in cortical representations suggest a more potent input for subsequent downstream signaling, and perhaps enhanced stimulus detectability - despite the invariance of S1 firing rate – but corresponding recordings of

the S1 recipient regions and a behavioral assay would be needed to test this hypothesis
directly. We also note that the observed changes in FS synchrony likely play a role in
shaping the overall cortical response we observe, as well as shaping the cortical
sensitivity to timing of the thalamic input through direct interactions with the excitatory
cortical sub-population, but a full exploration of the excitatory/inhibitory interactions within
cortex was beyond the scope of what was investigated here. Further, the nature of the
short ISIs within thalamic bursts almost certainly means that depression at the
thalamocortical synapse plays a role in determining the potency of the sensory-evoked
cortical response, as has been previously shown. However, the modeling results here
suggest that single-neuron bursting and across-neuron synchronous thalamic firing, and
how they interact with the timing sensitivity of cortex established by the disynaptic
feedforward inhibition, are the key role players in the observed dynamics, although more
extensive experiments involving causal manipulations of these mechanisms would need
to be conducted to more conclusively establish this explanation. Finally, though not
explicitly tested in these experiments, the results here would suggest that important
coding aspects of this particular pathway, notably velocity and direction tuning, would be
strongly shaped through changes in the timing properties of the thalamic population, as
we have previously shown in the context of rapid sensory adaptation (Wang et al., 2010).
660
In several of the analyses here, we combined recorded cortical units across layers. While
the optical GEVI imaging approach is focused on S1 layer 2/3, it also likely captures
activity above and below layer 2/3, as well as any cross-laminar processes, and further
obscures any differences across excitatory and inhibitory sub-populations due to the
indiscriminate viral targeting.   Neurons in VPm thalamus send axonal projections
differentially across laminae in S1 (Sermet et al., 2019), likely resulting in variations in the
effects of thalamic hyperpolarization on S1 neurons in different cortical layers. To account
for the possibility that the pooled dataset was not representative of thalamorecipient
neurons, we repeated a subset of our analyses for putative S1 layer 4 neurons, and in
each case observed the same trends as in the aggregate data (Supplemental Figure S5).
Note that the identification of L4 neurons was conducted conservatively, and thus some
of the non-L4 neurons in the aggregate population are likely also L4. Importantly, the
putative L4 neurons had a larger sensory evoked response than the non-L4 neurons, as
is apparent in comparing Supplemental Figure S5 to Figures 5 and 7, thus likely
dominating the aggregate analysis. Further, note that while the measurements of
synchrony in S1 here were limited to the putative FS neurons due to the relatively low
firing rates of the RS neurons, the modeling results suggest that the RS neurons would
also exhibit an increase in synchrony with thalamic hyperpolarization, but this must be
directly measured in future studies.   Regardless, the qualitative consistency of our
cortical results across cell types, laminar location, and recording modalities suggests that

our observations provide a representative – though by no means comprehensive – view
of the net effects of thalamic spike timing on cortical sensory responses.
683
Invariance in Baseline Thalamic Firing Activity
In this study, we utilized optogenetic hyperpolarization to bias thalamic sensory
responses toward bursting, without significantly changing other dynamics that might
indirectly impact sensory responses. One particularly surprising finding here was that
despite optogenetically induced hyperpolarization of thalamus, the overall baseline firing
rate of thalamic neurons was not suppressed, but instead unchanged. Following a
transient decrease in firing rate, the recorded VPm neurons returned to their original
baseline firing rate. This finding is, however, consistent with other reported observations.
In the visual pathway for example, optogenetic excitation of the thalamic reticular nucleus
(TRN) transiently silences the lateral geniculate nucleus, followed by a return to the
original firing rate at steady-state (Reinhold et al., 2015). In the somatosensory pathway,
strong photoinhibition attenuated VPm thalamic firing rate, but failed to quench activity
altogether (Halassa et al., 2011; Poulet et al., 2012; Lewis et al., 2015; Reinhold et al.,
2015; Yu et al., 2016). The most parsimonious explanation given the observations here
is that moderate amounts of hyperpolarization serve to engage the dynamics of the T-
type calcium channels that are inactive at normal baseline conditions, effectively
compensating for the loss in tonic spiking due to the hyperpolarization. Although the
optogenetic approach here does not enable direct observation of the magnitude of the
hyperpolarizing input, separate intracellular in-vitro slice experiments where we repeated
the protocol while patching on to VPm neurons revealed relatively modest amounts of
hyperpolarization that were well-sustained during constant light illumination. This is
important, as recent studies have suggested that prolonged activation of specific opsins
can have unintended consequences, notably here the possibility of changes in the
reversal potential for chloride (Raimondo et al., 2012). This effect could theoretically
result in changes in the degree of hyperpolarization, although halorhodopsin as a pump
is less directly affected by immediate (local) changes in reversal potential as compared
to channel-based optogenetics. Further, with halorhodopsin there is the potential for
photoinactivation and decreased photocurrents (Zhang et al., 2019), which could also
change the degree of hyperpolarization over time. Through the combination of the
intracellular control experiments and replication of the primary result through the integrate
and fire or burst (IFB) model, the likelihood that these possible effects played a primary
role here is low, especially over the relatively short timescales considered here.
716
Importantly, beyond the engagement of the T-type calcium channel burst mechanism
upon initial hyperpolarization, it would seem that additional hyperpolarization would
further push the neuron away from threshold, making it more difficult to burst, which would
predict a corresponding decrease in firing rate. However, we found that increased

hyperpolarization increased the number of spikes per burst, which served to offset the
decrease in the number of bursts with increasing hyperpolarization. The result is a
surprisingly resilient mechanism in response to this perturbation. It should be noted that
other elements of the circuit likely play a role in the observation here – for example, the
initial decrease in VPm firing rate would decrease excitation of TRN, subsequently
decreasing inhibition of VPm, which would work synergistically with the intrinsic properties
of the burst mechanism in this compensatory action.

The Potential Role of Thalamic Bursts in Sensory Signaling
Almost four decades ago, Crick proposed a provocative hypothesis – that the thalamic-
reticular complex serves as a dynamic gate for attentional control of sensory signaling to
cortex (Crick, 1984). Further refinement of this idea suggested that the switching between
tonic and burst firing modes of thalamic neurons that is facilitated by the unique dynamics
of the T-type calcium channels that are prevalent in the thalamus could establish a context
dependent signaling (Sherman and Guillery, 2002; Sherman, 2005). In this framework,
burst spiking would promote the detection of salient sensory features, while tonic spiking
would promote the transmission of details about the nature of the sensory stimulus.
Furthermore, the thalamic burst would also potentially provide a "wake-up call" to cortex
(Sherman, 2001), garnering attentional resources that ultimately would serve to switch
the thalamic mode of firing to tonic through depolarizing corticothalamic feedback. As
attractive a framework as this is, it is also a daunting theory to test experimentally due to
the complexity of the circuit, the required specificity of recording and manipulation, and
ultimately the need to cast in the context of changing states of arousal during trained
behaviors. Previous anesthetized studies have made substantial progress toward this
goal. For example, sensory-evoked thalamic bursting under anesthesia has been shown
to be well-driven by the appearance of salient sensory features (Lesica and Stanley,
2004; Alitto et al., 2005; Denning and Reinagel, 2005), which promotes the detection of
change in the sensory input from the perspective of an ideal observer of thalamic spiking
(Lesica and Stanley, 2004), and this sensitivity is strongly shaped by thalamic state
(Lesica et al., 2006). Further studies showing the sensitivity of sensory cortex to
spontaneous (non-sensory) thalamic bursting during wakefulness (Swadlow and Gusev,
2001) seem to set the stage for at least part of the overarching coding scheme in which
the cortical response would be amplified by the thalamic bursts in "detect" mode.

In opposition to this view, some have noted that because ongoing thalamic spontaneous
bursting events are particularly prominent during slow wave sleep and under anesthesia,
they likely play no role in sensory signaling during wakefulness. However, this has been
largely disproven, with low rates of bursts occurring both spontaneously and during
naturalistic stimuli in awake somatosensory (Swadlow and Gusev, 2001; Stoelzel et al.,
2009; Wright et al., 2021), visual (Niell and Stryker, 2010), and auditory systems

(Massaux et al., 2004). Further, we have shown previously in the awake, head-fixed rat
(Whitmire et al. 2016) and mouse (Wright et al. 2021) that rapid sensory adaptation has
a particularly strong effect on sensory-evoked burst firing, suggesting that the adaptive
changes in bursting may play an important role in perceptual adaptation during
wakefulness. Thus, it is critical to precisely probe the effects of sensory-evoked thalamic
bursting on cortical sensory representations during wakefulness, and specifically to test
the decades-old "wake-up call" hypothesis.
768
To this end, an optogenetic manipulation approach was adopted to gain systematic
control over thalamic burst/tonic firing modes, and the functional effects were precisely
quantified at the level of thalamus and primary sensory cortex.   During normal
physiological conditions, the thalamus receives a range of complex, excitatory and
inhibitory inputs that interact with the intrinsic cell properties to collectively set the baseline
membrane potential and firing rate of these neurons as a function of behavioral state.
Cortex also exhibits profound changes in ongoing and sensory-evoked firing across
states of wakefulness, and there is strong evidence that thalamic activity itself is a strong
driver of cortical state (Poulet et al., 2012). But because thalamus and cortex are densely
interconnected through feedforward and feedback projections, and are both subject to a
range of other modulatory inputs, it is generally difficult to infer directions of causality.
The approach here enabled the disentangling of these interactions by probing the
downstream effects of sensory-evoked thalamic bursting and synchronous firing per se,
independent of the many other naturally-occurring fluctuations. We do not suggest that
the thalamic hyperpolarization mimics a particular behavioral state, but rather that it
provides an opportunity to bias thalamic sensory responses toward burst firing mode, so
that we might probe the consequences for cortex. Such targeted manipulations are
critical to understanding state-dependent sensory signaling in this complex circuit more
generally.
788
In the experiments presented here, in the absence of optogenetic manipulation, bursting
represents approximately 15% of the total sensory evoked response, as compared to
greater than 30% during imposed thalamic hyperpolarization. Thus, even in the absence
of optogenetic manipulation, the bursting is not insignificant, and determining the effects
of thalamic bursts on cortex is relevant even under normal conditions. It is also important
to note that awake, head-fixed mice likely explore a much narrower range of behavioral
(e.g., arousal) states than occur in naturalistic settings. It is well-known that states of low-
arousal are directly correlated with thalamic burst firing modes, thus predictive of more
sensory-evoked burst firing than we observed here. While the experimental preparation
used here provides the requisite stability to perform precise stimulation and recording, a
complementary scenario for exploring the role of bursting across states of wakefulness
would involve freely moving rodents across a wider range of behavioral states. These

kinds of studies, in concert with those providing even more precise control of thalamic
firing modes (e.g., via closed-loop feedback control of neural activity (Bolus et al., 2018,
2021)) need to be employed to more strongly, effectively, and comprehensively explore
the coupling across brain regions and behavioral states.
805

Although the results here in the awake brain demonstrate that the absolute S1 sensory
evoked response was invariant to thalamic hyperpolarization and even diminished relative
to background cortical activity despite potent sensory-evoked thalamic bursting, what
emerges is increased timing precision, increased focus of spatial activation, and a
corresponding synchronization of the cortical sensory-evoked response that collectively
could promote detectability. Given the likely timing sensitivity of down-stream brain
structures in the sensorimotor arc, the synchronization of cortical activity may in fact be a
more critical element of cortical signaling than overall magnitude, supported by behavioral
work demonstrating the importance of cortical synchrony over firing rate (Jadhav et al.,
2009). Taken together, the results here point to timing rather than response magnitude
as a fundamental currency of the thalamocortical circuit, presenting a dynamic, timing-
based gating of sensory signaling to cortex that has strong implications for detectability
and discriminability in complex sensory environments.

819
820

References

Aguilar JR, Castro-Alamancos M a (2005) Spatiotemporal gating of sensory inputs in thalamus during quiescent and activated states. J Neurosci 25:10990–11002.

Alitto HJ, Weyand TG, Usrey WM (2005) Distinct properties of stimulus-evoked bursts in the lateral geniculate nucleus. J Neurosci 25:514–523.

Barthó P, Slézia A, Mátyás F, Faradzs-Zade L, Ulbert I, Harris KD, Acsády L (2014) Ongoing network state controls the length of sleep spindles via inhibitory activity. Neuron 82:1367–1379.

Béhuret S, Deleuze C, Bal T (2015) Corticothalamic Synaptic Noise as a Mechanism for Selective Attention in Thalamic Neurons. Front Neural Circuits 9:1–21.

Bolus MF, Willats AA, Rozell CJ, Stanley GB (2021) State-space optimal feedback control of optogenetically driven neural activity. J Neural Eng 18:036006.

Bolus MF, Willats AA, Whitmire CJ, Rozell CJ, Stanley GB (2018) Design strategies for dynamic closed-loop optogenetic neurocontrol in vivo. J Neural Eng 15:026011.

Borden PY, Ortiz AD, Waiblinger C, Sederberg AJ, Morrissette AE, Forest CR, Jaeger D, Stanley GB (2017) Genetically expressed voltage sensor ArcLight for imaging large scale cortical activity in the anesthetized and awake mouse. Neurophotonics 4:031212.

Brecht M, Sakmann B (2002) Whisker maps of neuronal subclasses of the rat ventral posterior medial thalamus, identified by whole-cell voltage recording and morphological reconstruction. J Physiol:495–515.

Bruno RM, Simons DJ (2002) Feedforward mechanisms of excitatory and inhibitory cortical receptive fields. J Neurosci 22:10966–10975.

Bujan AF, Aertsen A, Kumar A (2015) Role of Input Correlations in Shaping the Variability and Noise Correlations of Evoked Activity in the Neocortex. J Neurosci 35:8611–8625.

Crandall SRR, Cruikshank SJJ, Connors BWW, Crandall SRR, Cruikshank SJJ, Connors BWW (2015) A Corticothalamic Switch: Controlling the Thalamus with Dynamic Synapses. Neuron 86:1–15.

Crick F (1984) Function of the thalamic reticular complex: The searchlight hypothesis. Proc Natl Acad Sci U S A 81:4586–4590.

Cruikshank SJ, Lewis TJ, Connors BW (2007) Synaptic basis for intense thalamocortical activation of feedforward inhibitory cells in neocortex. Nat Neurosci 10:462–468.

Denning KS, Reinagel P (2005) Visual control of burst priming in the anesthetized lateral geniculate nucleus. J Neurosci 25:3531–3538.

Fogerson PM, Huguenard JR (2016) Tapping the Brakes: Cellular and Synaptic Mechanisms that Regulate Thalamic Oscillations. Neuron 92:687–704.

Gabernet L, Jadhav SP, Feldman DE, Carandini M, Scanziani M (2005) Somatosensory Integration Controlled by Dynamic Thalamocortical Feed-Forward Inhibition. Neuron 48:315–327.

Gentet LJ, Avermann M, Matyas F, Staiger JF, Petersen CCH (2010) Membrane potential dynamics of GABAergic neurons in the barrel cortex of behaving mice. Neuron 65:422–435.

Godwin DW, Vaughan JW, Sherman SM (1996) Metabotropic glutamate receptors switch visual response mode of lateral geniculate nucleus cells from burst to tonic. J Neurophysiol 76:1800–1816.

Guo W, Clause AR, Barth-Maron A, Polley DB (2017) A Corticothalamic Circuit for Dynamic
Switching between Feature Detection and Discrimination. Neuron 95:180-194.e5.
Halassa MM, Siegle JH, Ritt JT, Ting JT, Feng G, Moore CI (2011) Selective optical drive of
thalamic reticular nucleus generates thalamic bursts and cortical spindles. Nat Neurosci
14:1118–1120.
Hirata A, Castro-Alamancos MA (2010) Neocortex Network Activation and Deactivation States
Controlled by the Thalamus. J Neurophysiol 103:1147–1157.
Huguenard JR, McCormick D a. (2007) Thalamic synchrony and dynamic regulation of global
forebrain oscillations. Trends Neurosci 30:350–356.
Isaacson JS, Scanziani M (2011) How Inhibition Shapes Cortical Activity. Neuron 72:231–243.
Jadhav SP, Wolfe J, Feldman DE (2009) Sparse temporal coding of elementary tactile features
during active whisker sensation. Nat Neurosci 12:792–800.
Jahnsen H, Llinás R (1984a) Voltage-dependent burst-to-tonic switching of thalamic cell activity:
an in vitro study. Arch Ital Biol 122:73–82.
Jahnsen H, Llinás R (1984b) Electrophysiological properties of guinea-pig thalamic neurones: an
in vitro study. J Physiol 349:205–226.
Jin L, Han Z, Platisa J, Wooltorton JRA, Cohen LB, Pieribone VA (2012) Single Action Potentials
and Subthreshold Electrical Events Imaged in Neurons with a Fluorescent Protein Voltage
Probe. Neuron 75:779–785.
Khatri V, Hartings J a, Simons DJ (2004) Adaptation in thalamic barreloid and cortical barrel
neurons to periodic whisker deflections varying in frequency and velocity. J Neurophysiol
92:3244–3254.
Kimura F, Itami C, Ikezoe K, Tamura H, Fujita I, Yanagawa Y, Obata K, Ohshima M (2010) Fast
activation of feedforward inhibitory neurons from thalamic input and its relevance to the
regulation of spike sequences in the barrel cortex. J Physiol 588:2769–2787.
Kirchgessner MA, Franklin AD, Callaway EM (2020) Context-dependent and dynamic functional
influence of corticothalamic pathways to first- and higher-order visual thalamus. Proc Natl
Acad Sci 117:13066–13077.
Lesica N a, Stanley GB (2004) Encoding of natural scene movies by tonic and burst spikes in the
lateral geniculate nucleus. J Neurosci 24:10731–10740.
Lesica N a, Weng C, Jin J, Yeh C-I, Alonso J-M, Stanley GB (2006) Dynamic encoding of natural
luminance sequences by LGN bursts. PLoS Biol 4:e209.
Lewis LD, Voigts J, Flores FJ, Ian Schmitt L, Wilson MA, Halassa MM, Brown EN (2015) Thalamic
reticular nucleus induces fast and local modulation of arousal state. Elife 4:1–23.
Li L, Ebner FF (2016) Cortex dynamically modulates responses of thalamic relay neurons
through prolonged circuit-level disinhibition in rat thalamus in vivo. J Neurophysiol
116:2368–2382.
Litwin-Kumar A, Doiron B (2012) Slow dynamics and high variability in balanced cortical
networks with clustered connections. Nat Neurosci 15:1498–1505.
Lu SM, Guido W, Sherman SM (1992) Effects of membrane voltage on receptive field properties
of lateral geniculate neurons in the cat: Contributions of the low-threshold Ca2+
conductance. J Neurophysiol 68:2185–2198.
Lustig BR, Friedman RM, Winberry JE, Ebner FF, Roe AW (2013) Voltage-sensitive dye imaging
reveals shifting spatiotemporal spread of whisker-induced activity in rat barrel cortex. J

Neurophysiol 109:2382–2392.
Ma Y, Shaik MA, Kim SH, Kozberg MG, Thibodeaux DN, Zhao HT, Yu H, Hillman EMC (2016)
Wide-field optical mapping of neural activity and brain haemodynamics: considerations
and novel approaches. Philos Trans R Soc B Biol Sci 371:20150360.
Massaux A, Dutrieux G, Cotillon-Williams N, Manunta Y, Edeline J-M (2004) Auditory Thalamus
Bursts in Anesthetized and Non-Anesthetized States: Contribution to Functional
Properties. J Neurophysiol 91:2117–2134.
McCormick DA, Connors BW, Lighthall JW, Prince DA (1985) Comparative electrophysiology of
pyramidal and sparsely spiny stellate neurons of the neocortex. J Neurophysiol 54:782–
806.
Mease RA, Krieger P, Groh A (2014) Cortical control of adaptation and sensory relay mode in
the thalamus. Proc Natl Acad Sci 111:6798–6803.
Millard DC, Whitmire CJ, Gollnick CA, Rozell CJ, Stanley GB (2015) Electrical and Optical
Activation of Mesoscale Neural Circuits with Implications for Coding. J Neurosci 35:15702–
15715.
Murray Sherman S (2001) Tonic and burst firing: Dual modes of thalamocortical relay. Trends
Neurosci 24:122–126.
Niell CM, Stryker MP (2010) Modulation of Visual Responses by Behavioral State in Mouse
Visual Cortex. Neuron 65:472–479.
Olsen SR, Bortone DS, Adesnik H, Scanziani M (2012) Gain control by layer six in cortical circuits
of vision. Nature 483:47–52.
Owen SF, Liu MH, Kreitzer AC (2019) Thermal constraints on in vivo optogenetic manipulations.
Nat Neurosci 22:1061–1065.
Pinto DJ, Brumberg JC, Simons DJ (2000) Circuit Dynamics and Coding Strategies in Rodent
Somatosensory Cortex. J Neurophysiol 83:1158–1166.
Pinto DJ, Hartings JA, Brumberg JC, Simons DJ (2003) Cortical damping: analysis of
thalamocortical response transformations in rodent barrel cortex. Cereb Cortex 13:33–44.
Poulet JF a, Fernandez LMJ, Crochet S, Petersen CCH (2012) Thalamic control of cortical states.
Nat Neurosci 15:370–372.
Raimondo J V., Kay L, Ellender TJ, Akerman CJ (2012) Optogenetic silencing strategies differ in
their effects on inhibitory synaptic transmission. Nat Neurosci 15:1102–1104.
Reinhold K, Lien AD, Scanziani M (2015) Distinct recurrent versus afferent dynamics in cortical
visual processing. Nat Neurosci 18:1789–1797.
Sederberg AJ, Pala A, Zheng HJ V., He BJ, Stanley GB (2019) State-aware detection of sensory
stimuli in the cortex of the awake mouse Macke JH, ed. PLOS Comput Biol 15:e1006716.
Sermet BS, Truschow P, Feyerabend M, Mayrhofer JM, Oram TB, Yizhar O, Staiger JF, Petersen
CCH (2019) Pathway-, layer- and cell-type-specific thalamic input to mouse barrel cortex.
Elife 8:1–28.
Sherman SM (2001) A wake-up call from the thalamus. Nat Neurosci 4:344–346.
Sherman SM (2005) Thalamic relays and cortical functioning. In: Progress in Brain Research, pp
107–126.
Sherman SM, Guillery RW (2002) The role of the thalamus in the flow of information to the
cortex Adams P, Guillery RW, Sherman SM, Sillito AM, eds. Philos Trans R Soc London Ser B
Biol Sci 357:1695–1708.

Sherman SM, Koch C (1986) The control of retinogeniculate transmission in the mammalian
lateral geniculate nucleus. Exp Brain Res:1–20.
Smith GD, Cox CL, Sherman SM, Rinzel J (2000) Fourier Analysis of Sinusoidally Driven
Thalamocortical Relay Neurons and a Minimal Integrate-and-Fire-or-Burst Model. J
Neurophysiol 83:588–610.
Speed A, Del Rosario J, Burgess CP, Haider B (2019) Cortical State Fluctuations across Layers of
V1 during Visual Spatial Perception. Cell Rep 26:2868-2874.e3.
Stanley GB, Jin J, Wang Y, Desbordes G, Wang Q, Black MJ, Alonso J-MJ-M (2012) Visual
orientation and directional selectivity through thalamic synchrony. J Neurosci 32:9073–
9088.
Stoelzel CR, Bereshpolova Y, Swadlow H a (2009) Stability of Thalamocortical Synaptic
Transmission across Awake Brain States. J Neurosci 29:6851–6859.
Stujenske JM, Spellman T, Gordon JA (2015) Modeling the Spatiotemporal Dynamics of Light
and Heat Propagation for In Vivo Optogenetics. Cell Rep 12:525–534.
Suzuki S, Rogawski M a (1989) T-type calcium channels mediate the transition between tonic
and phasic firing in thalamic neurons. Proc Natl Acad Sci U S A 86:7228–7232.
Swadlow H a, Gusev AG (2001) The impact of "bursting" thalamic impulses at a neocortical
synapse. Nat Neurosci 4:402–408.
Swadlow HA. (2002) Thalamocortical control of feed–forward inhibition in awake
somatosensory 'barrel' cortex Adams P, Guillery RW, Sherman SM, Sillito AM, eds. Philos
Trans R Soc London Ser B Biol Sci 357:1717–1727.
Swadlow HA, Bezdudnaya T, Gusev AG (2005) Spike timing and synaptic dynamics at the awake
thalamocortical synapse. In: Progress in Brain Research, pp 91–105.
Taub AH, Katz Y, Lampl I (2013) Cortical Balance of Excitation and Inhibition Is Regulated by the
Rate of Synaptic Activity. J Neurosci 33:14359–14368.
Urbain N, Salin PA, Libourel PA, Comte JC, Gentet LJ, Petersen CCHH (2015) Whisking-Related
Changes in Neuronal Firing and Membrane Potential Dynamics in the Somatosensory
Thalamus of Awake Mice. Cell Rep 13:647–656.
Wang Q, Webber RM, Stanley GB (2010) Thalamic synchrony and the adaptive gating of
information flow to cortex. Nat Neurosci 13:1534–1541.
Wehr MS, Zador AM (2003) Balanced inhibition underlies tuning and sharpens spike timing in
auditory cortex. Nature 426:442–446.
Whitmire CJ, Waiblinger C, Schwarz C, Stanley GB (2016) Information Coding through Adaptive
Gating of Synchronized Thalamic Bursting. Cell Rep 14:795–807.
Wilent WB, Contreras D (2004) Synaptic responses to whisker deflections in rat barrel cortex as
a function of cortical layer and stimulus intensity. J Neurosci 24:3985–3998.
Wilent WB, Contreras D (2005) Dynamics of excitation and inhibition underlying stimulus
selectivity in rat somatosensory cortex. Nat Neurosci 8:1364–1370.
Wright NC, Borden PY, Liew YJ, Bolus MF, Stoy WM, Forest CR, Stanley GB (2021) Rapid Cortical
Adaptation and the Role of Thalamic Synchrony during Wakefulness. J Neurosci 41:5421–
5439.
Wright NC, Hoseini MS, Wessel R (2017a) Adaptation modulates correlated subthreshold
response variability in visual cortex. J Neurophysiol 118:1257–1269.
Wright NC, Hoseini MS, Yasar TB, Wessel R (2017b) Coupling of synaptic inputs to local cortical

activity differs among neurons and adapts after stimulus onset. J Neurophysiol 118:3345–
3359.
Yang H, Kwon SE, Severson KS, O'Connor DH (2016) Origins of choice-related activity in mouse
somatosensory cortex. Nat Neurosci 19:127–134.
Yu J, Gutnisky DA, Hires SA, Svoboda K (2016) Layer 4 fast-spiking interneurons filter
thalamocortical signals during active somatosensation. Nat Neurosci 19:1–14.
Yu J, Hu H, Agmon A, Svoboda K (2019) Recruitment of GABAergic Interneurons in the Barrel
Cortex during Active Tactile Behavior. Neuron 104:412-427.e4.
Zhang C, Yang S, Flossmann T, Gao S, Witte OW, Nagel G, Holthoff K, Kirmse K (2019) Optimized
photo-stimulation of halorhodopsin for long-term neuronal inhibition. BMC Biol 17:95.
1013
1014

FIGURE CAPTIONS

Figure 1. Thalamic hyperpolarization amplifies thalamic bursting and sensory-evoked cortical response in the anesthetized mouse. A. Pathway of the mouse vibrissa system from the facial vibrissae in the periphery, to brainstem, to thalamus, to S1. **B.** Experimental setup. Mice were injected with the viral vector eNphR3.0 (AAV5-CamIIKianse-eNphR3.0-mCherry) targeting the ventral posterior-medial (VPm) region of the thalamus and the viral vector ArcLight (AAV1-hysn1-ArcLightD- SV40) in the vibrissa region of the primary somatosensory cortex (S1). For optogenetic hyperpolarization, thalamic units were presented with constant illumination (LED, 590 nm, approx. 17 mW/mm^2) from a 200µm optic fiber, with LED illumination starting at 0.5s preceding stimulus (t= -0.5s) and ending 0.5s after stimulus (t=0.5), while recording simultaneously with a single tungsten electrode. Note that unlike for the awake recordings, light presentation was not ramped. For cortical GEVI imaging, the entire cortical area was illuminated through the thinned skull at 465 nm with an LED and imaged with a CCD imaging setup (see Methods). **C.** Thalamic bursts were identified as two or more spikes with inter-spike interval (ISIs) less than 4ms, preceded by silence for 100ms or more. Thalamic spikes that were part of an identified burst were classified as putative burst spikes (red), and all else were classified as tonic spikes (black). **D.** Peri-stimulus time histogram (PSTH) for thalamic spiking in response to a single, punctate whisker stimulus (t=0) for the Control (no light, top panel) and thalamic hyperpolarized (LED, bottom panel) conditions, 29 units. Instantaneous firing rates (bin size 2 ms) shown for all spikes (black) and putative burst spikes (red). **E. Left** Mean sensory-evoked thalamic firing rate (n=29 thalamic units) over the 0 – 30 ms time window increased from the Control (no light) to the LED conditions, for all spikes (black, p= 0.0046, paired Wilcoxon signed-rank) and burst spikes (red, p= 4.5e-4, paired Wilcoxon signed-rank). Error bars represent mean +/- SEM. **Right**. Mean sensory-evoked Cortical multiunit (MU) firing rate from 102 trials (across 2 recordings, 1 mouse, 5ms bin size). **F.** Example session of GEVI imaging following the delivery of a punctate whisker stimulus at time t=0, for the Control (no light, top row) and LED (bottom row) conditions. Images are averaged across 51 trials. Black bar represents 1mm. **G. Left** - Mean normalized peak GEVI sensory-evoked response was larger for the LED as compared to the Control condition (n= 9 mice, 13 recording sessions). For this plot, animals and sessions were each normalized to their control levels, and the LED condition is reported relative to the control. **Left Inset** - raw relative peak evoked %dF/Fo, p=0.017, paired Wilcoxon signed-rank, n=13 recordings across 9 mice. **Right** - Time series of the normalized sensory-evoked GEVI signal, generated from the integrated fluorescence within the 0.2 x 0.2mm ROI in the Control (gray) and LED (orange) conditions. Error bars represent mean +/- SEM.

Figure 2. Thalamic baseline firing rate is invariant to optogenetic hyperpolarization via a tonic/burst switch. A. Experimental setup. Mice were injected with the viral vector eNphR3.0 (AAV5-CamIIKianse-eNphR3.0-mCherry) targeting the ventral posterior-medial (VPm) region of the thalamus. On each trial, thalamic units were presented with light for 1.5 s (590 nm, LED approx. 17mW/mm^2 ramp, Double LED approx. 35mW/mm^2 ramp – see Methods) from a 200µm optic fiber and recorded simultaneously with a single tungsten electrode. **B.** Example extracellular single trial rasters depicting effects of thalamic hyperpolarization on spiking in VPm thalamus. Black indicates tonic spikes; red indicates burst spikes. Recorded spike waveform

shown at top. **C.** Aggregate single-unit (SU, n=51 units) PSTH for all spikes (black) and burst spikes
(red). Apparent is an initial decrease in overall firing rate following presentation of light, followed
by a return to the pre-hyperpolarization level. Bands represent are mean +/- SEM. **D. Left -** Mean
single-unit thalamic firing rate over the 250-750ms time window from the Control (no light) to
the LED condition, for all spikes (black, Control vs LED p=0.2, Control vs Double LED p=0.02,
unpaired two-sample t-test, n=54 for Control, and n=51 for both LED conditions) and burst spikes
(red, Control vs LED p=3e-13, Control vs Double LED p=1e-15, unpaired t-test). **Right -** Increase in
burst ratio from the Control to LED condition, where burst ratio is defined as the number of burst
spikes divided by the total number of spikes (Control vs LED p=9e-13, Control vs Double LED
p=1.8e-11, unpaired Wilcoxon rank sum test). **E. Left -** Example bursts for the Control and LED
conditions. **Right -** Increase in the mean number of spikes per burst from the Control to the LED
condition (Control vs LED p=2.3e-5, Control vs Double LED p=1.7e-6, unpaired Wilcoxon rank sum
test, n=54 for Control and n=51 for each of the LED conditions). Error bars represent mean +/-
SEM.   **F.** Simulated spontaneous activity from Integrate-and-fire-or-burst (IFB) model with
hyperpolarizing input at time t=0, showing all spikes (black) and burst spikes (red). See Methods
and Supplemental Figure S3.

1075

Figure 3. Cortical S1 baseline firing rate is invariant to optogenetic hyperpolarization of
thalamus after transient increase. A. Experimental setup. Mice were injected with the viral
vector eNphR3.0 (AAV5-CamIIKianse-eNphR3.0-mCherry) targeting the ventral posterior-medial
(VPm) region of the thalamus. Thalamic units were continuously presented with light (590 nm,
LED approx. 17mW/mm^2 ramp, Double LED approx. 35mW/mm^2 ramp) from a 200μm optic fiber,
while cortical single-units were recorded simultaneously with a laminar multi-electrode within
an identified cortical column (barrel) – see Methods. **B. Top -** Example extracellular rasters of
cortical activity, depicting effects of thalamic hyperpolarization on cortical spiking, across trials
and simultaneously recorded units (10 trials each from 15 units). **Bottom -** Aggregate PSTH across
all recorded trials and cortical single-units (n=118). Bands represent mean +/- SEM. Highlighted
are the transient and steady-state portions of the cortical response to the light. **C.** Mean cortical
firing rate in the Control and LED conditions, showing an increase in the transient (open symbol,
300-350ms, Control vs LED p=5.4e-14, Control vs Double LED p=1.4e-15, Wilcoxon signed-rank
test) and a return to near steady-state (closed symbol, 700-750ms, Control vs LED p=0.2, Control
vs Double LED p=0.013, Wilcoxon signed-rank test) portions of the response. Error bars represent
mean +/- SEM.

1092

Figure 4. Thalamic hyperpolarization enhances the thalamic sensory-evoked bursting response
in the awake mouse. A. Experimental setup. Mice were injected with the viral vector eNphR3.0
(AAV5-CamIIKianse-eNphR3.0-mCherry) targeting the ventral posterior-medial (VPm) region of
the thalamus and the viral vector ArcLight (AAV1-hysn1- ArcLightD- SV40) in the vibrissa region
of the primary somatosensory cortex (S1). For optogenetic hyperpolarization, on each trial
thalamic units were presented with light for 1.5 s (590 nm, LED 29approx. 17mW/mm^2 ramp,
Double LED 29approx. 35mW/mm^2 ramp – see Methods) from a 200μm optic fiber, with LED
illumination beginning at 0.75s preceding stimulus delivery, and continuing for 0.75s after
stimulus delivery, while recording simultaneously with a single tungsten electrode. For cortical
GEVI imaging, the entire cortical area was illuminated through the thinned skull at 465 nm with

an LED and imaged with a CCD imaging setup (see Methods). **B.** Grand single unit PSTH for
thalamic spiking in response to a single, punctate whisker stimulus (t=0) for the Control (no light,
top panel, n=54 units) and LED (bottom panel, n=51 units) conditions. Instantaneous firing rates
(bin size 2 ms) shown for all spikes (black) and putative burst spikes (red). **C. Left** – Mean sensory-
evoked thalamic response over 0-30ms time window for single-unit VPm recordings, for all spikes
(black, Control vs LED p=0.9, Control vs Double LED p=0.4, unpaired two-sample t-test, n=54 for
Control, and n=51 for both LED conditions), and for burst spikes (red, Control vs LED p=0.003,
Control vs Double LED p=1.6e-4, unpaired Wilcoxon rank sum, n=54 for Control, n=51 for both
LED conditions). Error bars represent mean +/- SEM. **Right** – Mean single-unit sensory-evoked
thalamic burst ratio over the 0–30 ms time window (Control vs LED p=1.3e-5, Control vs Double
LED p=2.4e-6, unpaired Wilcoxon rank sum, n=54 for Control, and n=51 for both LED conditions).
D. Construction of spike cross-correlogram (CCG) from pair-wise VPm spiking. **E.** Grand mean
CCGs of the sensory-evoked response for the Control and LED conditions, when restricting spiking
data to the [0-10ms], [10-20ms], and [20-30ms] post-stimulus time windows (n=187
simultaneously recorded pairs from 46 VPm single units, across 6 recording sessions).   **F.** Mean
shuffle-corrected synchronous spike count for the [0-10ms], [10-20ms], and [20-30ms] post-
stimulus time windows. All reported values are mean synch AP count +/- 99.95% confidence
interval, resampling relative spike times with replacement ([0-10 ms], Control to LED, p < 0.001;
[10-20 ms], p < 0.01; [20-30 ms], p < 0.01).

1122
Figure 5. Cortical sensory-evoked response magnitude is invariant to thalamic
hyperpolarization in the awake mouse. **A.** Example session of GEVI imaging following the
delivery of a punctate whisker stimulus at time t=0, for the Control (no light, top row) and LED
(bottom row) conditions. Images are averaged across 51 trials. **B. Left** – Peak GEVI sensory-
evoked response (between 0-110ms post stimulus) slightly decreased from the Control to the
LED condition (Control vs LED p=0.020, Control vs Double LED p=0.0039, paired Wilcoxon
signrank, n=9 recordings from 2 mice). **Right –** Mean normalized peak GEVI sensory-evoked
response for the Control and LED conditions (n=9 recordings). Before combining, animals and
sessions were each normalized to their Control levels, and the LED condition is reported relative
to the Control. **C.** In separate experiments, we hyperpolarized the thalamus while simultaneously
recording the S1 Cortical response using a multichannel electrode. **D.** Cortical Grand PSTH evoked
sensory response (t=0) across all recorded single units (n=118 units, 5ms bins) for Control (black)
and LED (orange) conditions. **E. Left** Average absolute evoked cortical response remained
approximately invariant during LED on conditions across all cortical recorded units (n=118).
Evoked sensory response period defined as between 5-100ms post stimulus. **Right.** Relative
evoked cortical response decreased with increasing LED intensities (Control vs LED p=0.20,
Control vs Double LED p=0.034, paired Wilcoxon signrank, n=118 units). Relative evoked response
defined as the absolute response minus the preceding baseline activity -50-0ms pre-stimulus.
Error bars represent mean +/- SEM.

1142
Figure 6. Thalamic hyperpolarization results in a spatially sharpened and temporally narrowed
cortical sensory evoked response. A. GEVI imaging frames at peak response for the control (top)
and LED (bottom) conditions for an example session. **B.** Cartoon illustration of the coupling
between amplitude and spatial area in cortical imaging. **Top -** A reduction in evoked fluorescence

amplitude can be qualitatively perceived as a reduction in spatial area of activation, due to the
"iceberg" effect. **Bottom -** However, the spatial area of activation above a fixed threshold after
normalization to the peak reveals the true effects on the area. **C.** Normalized amplitude time
series of fluorescence within ROI for Control (black) and thalamic hyperpolarized (orange, LED),
n= 9 recordings from 2 mice) conditions. Bands represent mean +/- SEM. **D.** Summary analyses
(Error bars represent mean +/- SEM). Note, for this figure all conditions are normalized to their
respective peak responses, as portrayed in B.   **Left** - Mean spatial activation area of the
normalized sensory-evoked response decreased from the Control to the LED condition (Control
vs LED p=0.025, Control vs Double LED p=0.046, paired t-test, n= 9 recordings). **Middle** - The peak-
to-baseline (defined in Figure 6C), calculated as the mean time from the peak of the fluorescence
to the first return to pre-stimulus baseline (ms), decreased from the Control to the LED condition
(Control vs LED p=0.0059, Control vs Double LED p=0.0019, unpaired t-test, n= 8 Control
recordings, 9 LED recordings, note one of the Control recordings did not return to baseline). **Right**
– The recovery (defined in Figure 6C), calculated as the relative mean fluorescence during the
time duration following the peak stimulus-evoked response (120-400ms), decreased from the
Control to the LED condition (Control vs LED p=0.016, Control vs Double LED p=0.020, paired t-
test, n= 9 recordings). LED approx. 17mW/mm^2 ramp, Double LED approx. 35mW/mm^2 ramp –
see Methods. Error bars represent mean +/- SEM.

1165

Figure 7. Thalamic hyperpolarization affects timing but not magnitude of sensory-evoked
responses in cortex. A. Cortical single-units were classified as Regular Spiking Units (RSUs, red)
or Fast Spiking Units (FSUs, blue) based on the time interval from the peak-to-trough (see
Methods). Example RSU and FSU waveforms are shown at the top, where bands represent +/- 1
SD. The distribution of time intervals from peak-to-trough for the spike waveforms is shown at
the bottom. **B.** PSTHs of the aggregate putative RSUs (left, n=86) and FSUs (right, n=32) in
response to a punctate whisker deflection at time t=0 (bin size 2ms) for the Control (black) and
double LED (orange) conditions. **C.** Summary analyses (Error bars represent mean +/- SEM) for
RSUs (top row) and FSUs (bottom row). **Left** - Mean baseline (-50 to 0ms) RSU (Control vs LED
p=5.3e-4, Control vs Double LED p=0.012, Wilcoxon signed-rank test, n=86) and FSU (Control vs
LED p=0.15, Control vs Double LED p=0.30, Wilcoxon signed-rank test, n=32) firing rates for the
LED relative to the Control condition. **Middle** - RSU (Control vs LED p=0.067, Control vs Double
LED p=0.0082, Wilcoxon signed-rank test, n=86) and FSU (Control vs LED, p=0.30, Control vs
Double LED p=0.45, Wilcoxon signed-rank test, n=32) early sensory response (5-50ms) for the
LED compared to the Control conditions. **Right** – Late sensory response (60-100ms) for the LED
compared to the Control condition for the RSU (Control vs LED p=0.017, Control vs Double LED
p=0.014, n=86, Wilcoxon signed-rank test) and FSU (Control vs LED p=2.2e-4, Control vs Double
LED p=4.5e-4, n=32, Wilcoxon signed-rank test) populations. LED approx. 17mW/mm^2 ramp,
Double LED approx. 35mW/mm^2 ramp – see Methods. Error bars represent mean +/- SEM.

1185

Figure 8. Thalamic hyperpolarization enhances cortical synchrony. A. The spike cross-
correlogram (CCG) was calculated as a histogram of spike times of cell 2 relative to a target spike
of cell 1, repeated for all spikes of cell 1. The pair-wise synchrony was calculated from the CCG as
the area of the CCG between +/- 7.5ms, normalized by the total area of the CCG. **B. Top** Example
Raster of two simultaneously collected FSU units, Cell 1 (black), Cell 2 (purple) for Control and

LED conditions. **Bottom**. Synchronous spiking events only for the same neural pair (red, firing
within 7.5ms of each other). **C.** Aggregate cross-correlograms for FSUs (n=99 pairs, see Methods)
for the control (black) and LED (orange) conditions. Note that for the LED condition, the light level
was 35 mW/mm^2 to better emphasize the change in synchrony with thalamic hyperpolarization.
Cross-correlograms have been smoothed via a moving average filter, 2.5ms window. **D.** Mean
levels of synchrony FSU pairs, for the spontaneous/baseline cortical firing as compared to the
sensory-evoked response for the Control and LED conditions. Relative to the Control condition,
thalamic hyperpolarization (LED, 35mW/mm^2) resulted in an increase in synchrony for the FSU
(Control vs LED, p=1.5e-8, Control vs Double LED, p=3.5e-13, Wilcoxon signed-rank test, n=99
pairs) populations. LED approx. 17mW/mm^2 ramp, Double LED approx. 35mW/mm^2 ramp – see
Methods. Error bars represent mean +/- SEM.
1202
Figure 9. A thalamocortical network model reproduces key experiment results
implicating thalamic spike timing. A. Thalamocortical model schematic (see Methods). **Top**
inset - A clustered network of excitatory (red) and inhibitory (blue) neurons received inputs from
a model VPm barreloid, as well as random excitatory inputs. **Top** - Grand mean PSTHs for VPm
tonic and burst spikes. **Bottom** – Grand mean PSTHs for network excitatory and inhibitory
neurons, for the Control (black) and LED (orange) conditions. **B. Left** – Grand spike cross-
correlograms for VPm spike train inputs to the model cortical network, for three post-stimulus
windows. **Right** – Mean synchronous spike counts for VPm spike train inputs to the model
cortical network. Error bars indicate 95% confidence intervals from re-sampling relative spike
times with replacement. **C.** Grand mean (+/- SEM) rates for all model network excitatory (red)
and inhibitory (blue) neurons for the early (0 – 30 ms post-stim) response window. Excitatory
control vs LED: 4.6% increase, p = 6.85 x 10^{-9}, Wilcoxon signed-rank test. Inhibitory control vs.
LED: 3.6% increase, p = 2.15 x 10^{-13}, Wilcoxon signed-rank test. **D.** Top – Grand cross-
correlograms for 200 randomly-selected, valid pairs of network inhibitory neurons (see
Methods). **Inset** – Same, but for window used to calculate synchrony (+/- 7.5 ms). **Bottom** -
Grand mean (+/- SEM) pairwise synchrony for 200 valid inhibitory pairs (left) and 147 valid
excitatory pairs (right). Inh-inh control vs. LED: 22.4% increase, p = 1.44 x 10^{-34}, Wilcoxon signed-
rank test. Exc-exc control vs. LED: 8.4% increase, p = 1.74 x 10^{-7}, Wilcoxon signed-rank test. **E.**
Same as in (A), but including "randomized" condition (gray PSTHs), in which P = 15% of drawn
tonic spike times occurring near the early PSTH peak were shifted to a later, random time. **F.**
Same as in (B, left), but including "randomized" condition (gray CCG). **G.** Same as in (C), but for
various choices of P (percent randomized VPm tonic spike times). **H.** Same as in (D, bottom), but
for various choices of P. * Indicates 0.01 ≤ p < 0.05, ** indicates 0.001 ≤ p < 0.01, *** indicates
p < 0.001, Wilcoxon signed-rank test.
1227
1228
1229
1230

STAR METHODS

Resource Availability

Lead Contact: Further information and requests for data or code should be directed to
and will be fulfilled by the lead contact, Garrett Stanley
(garrett.stanley@bme.gatech.edu).

Materials Availability: This study did not generate new unique reagents.

Data and Code Availability:

All data reported in this paper will be shared by the lead contact upon request.

All original code will be deposited in a public repository and will be publicly available as
of the date of publication. DOIs will be listed in the key resources table.

Any additional information required to reanalyze the data reported in this paper is
available from the lead contact upon request.

Experimental Model and Subject Details

All procedures were approved by the Institutional Animal Care and Use Committee at the
Georgia Institute of Technology and were in agreement with guidelines established by the
National Institutes of Health. Experiments were performed on adult (6 – 15 weeks) male
and female Mice. Mice strains included C57BL/6J (Jackson Laboratories) and the
transgenic NR133-Cre (Gerfen et al., 2013) on a C57BL/6J background mice.

Method Details

AAV Delivery: At least 5 weeks prior to experimentation, 6 week old female C57BL/6J
(Jackson Laboratories) mice were injected with different viral constructs either in the
ventral posteromedial (VPm) thalamic region with AAV-5-CamKinaseII-eNph3.0 (UNC
Viral Vector core) for optogenetic modulation, in the primary somatosensory (S1) cortex
with AAV-1-hsyn1-ArcLight (UPenn Viral Vector Core) for optical voltage imaging, or both.
For surgical procedures, mice were anesthetized using isoflurane (3-5%). After the
mouse was fully anesthetized, small craniotomies were placed over the regions of interest
and were aligned using stereotaxic measurements (For VPm, 1.8mm lateral from midline
by 1.8mm caudal from bregma). For cortical expression, either single or multiple injection
sites were used surrounding the barrel cortex (centered at 1.5mm caudal from bregma
and 3mm lateral from midline). The virus was loaded into a modified Hamilton syringe
(701-N) with a ~35 micron borosilicate glass pipette tip or a Hamilton Neuros Syringe.
The syringe was initially lowered to the corresponding depth below the surface (for VPm:
3mm and For S1: 0.5mm) and the tissue was allowed to rest for 1 minute before injection.

Both sites received injections of 0.5-1µl of viral construct at a flow rate of 0.1µl/minute.
After injection, the pipette remained in place for an additional 5 minutes before slowly
being removed from the brain. The bore holes were filled with either bone wax or left to
close naturally. Throughout injection, mice were kept warm using a water heating system
to maintain body temperature. See Supplemental Figure S1 for histological validation of
expression of halorhodopsin in thalamus and ArcLight in S1.
1270

Awake Animal Preparation: At least four weeks after ArcLight and eNphR3.0 viral
injection, mice were anesthetized under isoflurane and were implanted with a head-plate.
Over the course of 5-14 days preceding the first imaging experiment, mice were routinely
handled to gain familiarity with the imaging system and immobilization device. During this
acclimatization period, mice were head fixed for increasingly longer periods of time, from
15 minutes to 1.5 hours. During stimulation of the whisker, mice were prevented from
interacting with the whisker stimulator by obstructing the path from the paws to the
whisker. Mice were rewarded with sweetened milk (Nestle, Ltd.) throughout imaging. After
at least 5 days of handling and acclimating, mice became tolerant to immobilization in the
headplate restraint system. During passive stimulation of the whiskers, the mice often
actively moved their whiskers. Therefore, the whisker stimulator was placed relatively
close (5mm) to the face to prevent the whisker from slipping out of the manipulator;
however, the amplitude of the deflection was adjusted to maintain a consistent angular
velocity (1200 deg/s).
1285

Whisker Stimulation: Whisker stimulation was similar to that utilized previously (Borden
et al., 2017). Briefly, individual vibrissae of the mice were deflected by a high fidelity (1
KHz) galvanometer system (Cambridge Technologies). A whisker stimulus was applied
by positioning the custom designed galvanometer 5-10mm from the face and delivering
an exponential sawtooth (rise and fall time = 5ms). The waveform stimulus velocity was
taken by averaging the time to peak velocity of the stimulus. The velocity was adjusted
based on distance from the face.
1293

Thalamic and Cortical Electrophysiology: For thalamic electrophysiology, a small
craniotomy was made over the primary whisker sensitive thalamic ventral-posterior
medial (VPm) region of the mouse, around the injection site, using stereotactic
coordinates (see above). VPm was then mapped under anesthesia using either a
2MOhm tungsten electrode (FHC) or 32-channel silicon probe (NeuroNexus
Technologies). The mapping electrode was slowly lowered below the cortical surface
during manual stimulation of the whiskers, while spiking activity (threshold crossings of
high-pass filtered voltages) was monitored using the data acquisition system. We
stopped descending when we recorded whisker-driven spiking activity at a depth
consistent with VPm (typically 2900 – 3600 µm below the surface). We then used the
galvanometer to present precise single-whisker stimulation, and confirmed that multi-unit
activity was consistent with electrophysiological features of VPm. Specifically, the

electrode was determined to be located in VPm if the peri-stimulus time histogram (PSTH)
contained a peak response 3ms - 10ms after a 1200 degree/s punctate single whisker
stimulus and did not have a latency shift by more than 20ms after 1s of a 10Hz adapting
stimulus (Wang et al., 2010). The principal whisker was determined by the largest 30ms
PSTH response of multiple neighboring whiskers.   We then noted the depth and
stereotactic coordinates of this recording site and slowly retracted the mapping electrode,
sealed the craniotomy, and returning the mouse to its home cage.   During awake
recording sessions, we targeted the same location for recording, briefly repeating the
above steps to confirm electrode location. We recorded thalamic spiking using either a
2MOhm tungsten electrode (FHC) with 200 µm attached optical fiber, or a 32-channel
silicon probe array with 100 µm attached optical fiber (A1x32-Poly3-5mm-25s-177-
OA32LP, NeuroNexus Technologies).   For cortical recordings, initial mapping was
conducted using cortical ArcLight voltage imaging or intrinsic imaging (see below). Once
the target cortical column (barrel) was identified and confirmed, a 32-channel linear silicon
probe (A1x32-5mm-25-177, NeuroNexus Technologies) or single tungsten electrode
(FHC) was inserted. For both thalamic and cortical electrophysiology, after the conclusion
of the study either a small 7uA 10s lesion, or a fluorescent dye was placed near the
recording location and confirmed using post-mortem histological validation. Neuronal
signals were band-pass filtered (500Hz–5KHz), digitized at either 24.414 or 30
KHz/channel and collected using either a 96-channel (Blackrock Microsystems, Salt Lake
City, UT, USA), or a 64-channel (Tucker-Davis Technologies, Alachua, FL, USA) data-
acquisition system.

1328
1329

<u>Awake Cortical Fluorescent ArcLight Imaging</u>: ArcLight transfected mice were imaged
through the thinned or removed skull using a two camera system: a Scimedia Imaging
system to measure cortical ArcLight spatial activity, and a custom camera to measure
hemodynamic activity for subtraction. The cortex was imaged using a 184 x123 pixel CCD
Camera, MiCam2 HR Camera (Scimedia, Ltd) to capture ArcLight, and a Basler Ace
(acA1920-155um) 480 x 180 pixel (4x4 binned) CMOS Camera to capture auto-
fluorescence, at 200 Hz with a tandem lens microscope. The entire cortical area was
illuminated at 465 nm with a 400 mW/cm2 LED system (Scimedia, Ltd.) to excite the
ArcLight fluorophore and background auto-fluorescence. The excitation light was
projected onto the cortical surface using the first dichroic mirror (bandpass: 475/625nm,
Semrock, Inc.). Collected light was passed through a second dichroic mirror (Longpass
cutoff: 495 nm, Semrock, Inc.) for collection of the ArcLight and auto-fluorescence signal.
The auto-fluorescence signal was filtered with a bandpass filter between the wavelengths
of 465/75 nm (Semrock, Inc).   The ArcLight signal was filtered with a bandpass emission
filter between wavelengths of 520⁄35 nm (Semrock, Inc.). The imaging system was
focused approximately 300µm below the surface of the brain to target cortical layer 2/3,

although the imaging likely captures fluorescence from the cell bodies as well as neuropil
above and below layer 2/3.

1348

<u>Anesthetized Cortical Fluorescent ArcLight & Intrinsic Imaging:</u> ArcLight transfected mice
were imaged through the thinned or removed skull using a Scimedia Imaging system to
measure cortical spatiotemporal activity (leveraging a single camera setup). The cortex
was imaged using a 184x123 pixel CCD Camera, MiCam2 HR Camera (Scimedia, Ltd)
at 200 Hz, and a tandem lens macroscope. The entire cortical area was illuminated at
465 nm with a 400 mW/cm^2 LED system (Scimedia, Ltd.) to excite the ArcLight
fluorophore. The excitation light was further filtered (cutoff: 472-430 nm bandpass filter,
Semrock, Inc.) and projected onto the cortical surface using a dichroic mirror (cutoff: 495
nm, Semrock, Inc.). Collected light was filtered with a bandpass emission filter between
wavelengths of 520-535 nm (Semrock, Inc.). The imaging system was focused
approximately 300µm below the surface of the brain to target cortical layer 2/3. For
intrinsic imaging of the hemodynamic response, the cortical surface was illuminated by a
625nm red LED (ThorLabs), and imaged with the same camera system as above, at a
temporal resolution of 10Hz. During intrinsic imaging, no emission filters were used. In
order to evoke a cortical intrinsic response, the whisker was repetitively stimulated at
10Hz for 6 seconds.
1365

<u>Functional Fluorescent Mapping of Barrel Cortex:</u> The mouse's whisker system was first
mapped by imaging the rapid ArcLight response to a high velocity (1200 Deg/s) sensory
stimulus separately applied to three different whiskers. The resulting whisker response
averaged over 20 trials was determined to be associated with a principal whisker, and
barrel, if the evoked response was spatially limited to roughly a 0.2 mm x 0.2 mm area
25-30ms after stimulation. Additionally, the response was determined to be originating
from the barrel field if the center of mass of activation moved consistently with the
histologically defined barrel field and was within the standard stereotaxic location of S1
(~3mm lateral, 0.5-1.5mm from bregma). After mapping, a single whisker was deflected
in a way as to emulate a high velocity slip-stick event (1200 deg/s), either with or without
thalamic optogenetic hyperpolarization.
1377

<u>Simultaneous Imaging and Thalamic Optogenetic Manipulation:</u> After mapping both the
thalamic and cortical regions, an optrode (2M Ohm tungsten electrode mounted to a 200
µm optic fiber) was positioned to the stereotaxic locations of the pre-mapped thalamic
region and lowered to the corresponding depth. Once a single thalamic unit was identified
using the above constraints, the unit was determined to be sensitive to optical stimulation
by briefly (1-2s) hyperpolarizing the cells using ~17mW/mm^2 (LED condition) or
~35mW/mm^2 (double LED condition) (unless otherwise noted) at 590nm from an LED
light source (Thorlabs, M590-F1). Each cell was determined to be a thalamic optically
sensitive unit if the lighted caused a transient decrease in firing rate or if the cessation of
the 590nm light caused a rebound burst (Brecht and Sakmann, 2002). After identifying

an optically sensitive thalamic unit, the whisker stimulus was presented under various
light conditions. Light stimulation was presented 500-750ms preceding and following
whisker deflection. There was at least a four second interval between stimulus deliveries
to allow for recovery of halorhodopsin (eNphR3.0). Each session imaged 200ms-1s of
frames preceding whisker stimulation to measure spontaneous activity. Prior to use, light
power was measured from the tip of the ground optical fiber before each experiment to
maintain approximate light intensities delivered to each cell. During light delivery, the
downstream cortical response was recorded using either electrodes for cortical
electrophysiology or voltage imaging as described above. The optogenetic and viral
expression of each experiment was verified through confocal and brightfield imaging of
fixed slices.

1399

The LED light intensity used for optogenetic stimulation is within the published range of
light stimulation ($35mW/mm^2$ is estimated from our maximum power measure of ~1.1mW
through a 200μm fiber) (Stujenske et al., 2015; Owen et al., 2019). Further, studies that
have directly quantified the effects of optical stimulation on local tissue heating and neural
activation (in the absence of opsin expression) have found no significant difference in the
firing rate change for 1mW light intensity (Stujenske et al., 2015) or minor firing rate
changes for 3mW light intensity, but no behavioral effects (Owen et al., 2019). See
Supplemental Figure S4 for controls that demonstrate a lack of light effects and lack of
confounding interactions between optogenetic activation of VPm thalamus and GEVI
imaging in cortical S1.

1410
1411

Anesthetized Electrophysiology: A subset of experiments was conducted with mice under
light anesthesia, as a control. These mice were initially anesthetized using isoflurane (3-
5%) and then placed on a heated platform (FHC, Inc.) in a stereotaxic nose cone to
maintain anesthesia. A large incision was placed over the animal's skull, and
the connective tissue and muscle surrounding the skull was removed using a
fine scalpel blade. A modified headplate was attached using dental acrylic (Metabond)
and secured to the skull. For cortical imaging, the skull was thinned with a dental drill until
transparent, or removed entirely and covered with saline or ringers solution. After surgery,
the isoflurane levels were dropped to ~<1% for imaging and electrophysiology, the
procedures for which were identical to those for the awake animal. The animal's vitals
(heart rate and respiratory rate) were constantly measured for tracking anesthesia depth.

1423

Histology: Histological procedures were similar to those utilized previously (Borden et al.,
2017), to validate ArcLight in S1 and/or opsin expression in VPm thalamus. Histological
samples were prepared by perfusing the animal transcardially with phosphate buffered
saline (PBS) followed by 4% paraformaldehyde. Brains were post-fixed overnight in 4%
paraformaldehyde then transferred to PBS before sectioning. Thick sections were cut

using a vibratome (100 μm, Leica, VTS 1000) and either directly mounted or saved for
staining.   See Supplemental Figure S1 for histological validation of expression of
halorhodopsin in VPm and ArcLight expression in cortex.

1432

Integrate & Fire or Burst (IFB) Modeling: To further explore the surprising finding of
relatively invariant baseline firing rates during hyperpolarization of VPm thalamus, we
utilized a biophysically inspired model of thalamic burst/tonic firing.   Specifically, we
suggest that this finding was a consequence of the burst mechanism, and not the result
of possible confounds related to previously reported changes in reversal potential of
chloride during prolonged periods of halorhodopsin activation (Raimondo et al., 2012).
The Integrate and Fire or Burst (IFB) model was derived from previously published
models of thalamic function from the LGN (Smith et al., 2000; Lesica and Stanley, 2004;
Lesica et al., 2006).   In order to simulate the experimental parameters and account for
changes in thalamic activity, some additional terms and parameters were added and
adjusted. Additionally, we generated ongoing activity using two methods, either injected
current noise or synaptic events, with both showing the same results. The results shown
here use the synaptic event model where IPSCs and EPSCs are modulated as fixed
inputs. The model itself was written and analyzed using custom scripts in Matlab 2016a.

1447

1448   The model is based on modifications of the standard integrate and fire model representing
1449   the effects of integrated synaptic currents on membrane voltage:

1450

1451
$$C \frac{dV}{dt} = I_{EPSC} - I_{IPSC} - I_L - I_T - I_{Halo}$$

1452
$$I_L = g_L(V - V_L), \ I_T = g_T m_\infty h(V - V_T)$$

1453
$$\frac{dh}{dt} = \begin{cases} \dfrac{-h}{\tau_h^-}, (V > V_h) \\ \dfrac{(1-h)}{\tau_h^+}, (V < V_h) \end{cases}$$

1454
$$\frac{dI_{EPSC}}{dt} = \frac{I_{EPSC}}{\tau_s}$$

1455
$$\frac{dI_{IPSC}}{dt} = \frac{I_{IPSC}}{\tau_s}$$

1456

The following parameters were used to simulate thalamic activity: $C = 2\text{uF}/cm^2$, $g_L =$
$0.035 \text{ mS}/cm^2$, $g_T = 0.07 \text{ mS}/cm^2$, $V_L = -65\text{mV}$, $V_{reset} = -45\text{mV}$, $V_h = -68\text{mV}$, $V_T =$
120mV, $\tau_h^+ = 0.1\text{s}$, $\tau_h^- = 0.02\text{s}$, $\tau_s = 1\text{e} - 2\text{uF}/cm^2$, $Threshold = -35\text{mV}$, $I_{Halo} = 0 -$
$1uA/cm^2$.   EPSPs occurred at a rate $\text{EPSP}_{rate} = 1 - 25\text{Hz}$, where each excitatory post-
synaptic event had a peak of $3\text{nA}/cm^2$, and decayed according to the above first-order
differential equation for I_{EPSC}. Similarly, IPSPs occurred at a rate of $\text{IPSP}_{rate} = 0 - 5\text{Hz}$,

where each inhibitory post-synaptic event had a peak of $1uA/cm^2$, and decayed
according to the above first-order differential equation for I_{IPSC}. To simulate the different
levels of thalamic activity, we varied the rates of EPSP inputs on the thalamic model
(based on published ranges of thalamic activity (Urbain et al., 2015)). IPSPs were
simulated at a much lower rate (20% of EPSP rate) to add additional variability to baseline
activity. The model outputs represent the average response of 100 simulated thalamic
neurons in response to various levels of thalamic hyperpolarization and baseline activity.
The model was updated at 1ms steps. The absolute refractory period was set to 1ms.
1471
<u>Cortical E-I Modeling:</u> We constructed a simple model of the thalamocortical network
using custom scripts written in Python 3.6.10, as described previously (Wright et al., 2021)
We modeled a single cortical barrel as a clustered network of excitatory and inhibitory
single-compartment leaky integrate-and-fire (LIF) neurons, subject to excitatory thalamic
and non-thalamic synaptic inputs. For each condition, we simulated 50 trials (200 ms per
trial), with a time-step of 0.05 ms. A detailed description of the model can be found in
Cortical E-I Modeling Supplemental, Section 1. All code is freely available upon request.
1479
We created VPm input spiking to match both the experimentally observed sensory-
evoked firing and the pair-wise synchrony of thalamic firing, across conditions.
Specifically, we modeled a single VPm barreloid as forty independent "neurons", or spike
trains. Spike times were drawn from the empirical VPm PSTHs, re-scaled for each neuron
to give a broad distribution of firing rates. Bursts were modeled as triplets of spikes with
1.25 ms inter-spike interval.
1486
1487   Non-zero thalamocortical (TC) synaptic weights were broadly distributed, and we
1488   implemented differential TC connectivity by imposing higher TC convergence (Bruno and
1489   Simons, 2002; Cruikshank et al., 2007) and shorter synaptic latencies (Cruikshank et al.,
1490   2007; Kimura et al., 2010) for inhibitory neurons, and requiring that VPm neurons with the
1491   highest mean rates synapsed only onto inhibitory neurons (Bruno and Simons, 2002).
1492
We modeled a single cortical column as a network of 800 excitatory and 100 inhibitory
LIF neurons, with relatively strong inhibitory-to-excitatory synapses (Gabernet et al.,
2005). We imposed spatial clustering via "small-world" network connectivity (Litwin-
Kumar and Doiron, 2012; Bujan et al., 2015; Wright et al., 2017b, 2017a), with 10% re-
wiring probability. Inhibitory LIF neurons had shorter membrane time constants (Gentet
et al., 2010) and refractory periods than excitatory neurons, which – together with the TC
connection properties described above – supported higher firing rates in inhibitory
neurons, as observed here (Fig. 7) and in previous work (Bruno and Simons, 2002; Khatri
et al., 2004; Gentet et al., 2010; Taub et al., 2013). Excitatory neurons were subject to

an inhibitory spike-rate adaptation conductance, which helped to stabilize network
activity.

1504

We employed alternate models to parse the roles played by changes in thalamic bursting
per se, and by changes in synchronous thalamic spiking. In these alternate models, we
manually manipulated tonic spike times to effect changes in the rate of synchronous tonic
input spikes, while holding mean tonic and burst rates fixed. Specifically, for each trial
and spike train, we probabilistically relocated tonic spikes that occurred within +/- 5 ms of
the early PSTH peak to a later (random) time within approximately the first 30 ms post-
stimulus.

1512

For each model, we calculated the grand mean +/- SEM firing rates for all neurons, for an
"early" (0 – 30 ms post-stimulus) and "late" (60 – 100 ms post-stimulus) response window,
where stimulus onset time represents the time of galvo deflection onset, or $t = 0$ in the
empirical VPm PSTH. For cortical network synchrony analysis, we randomly selected
200 excitatory and 200 inhibitory neuron pairs. We then calculated synchrony for each
"valid" pair, i.e., each of these pairs with at least one relative spike time in the 30 ms post-
stimulus window across all trials. For VPm spike train synchrony analysis, we randomly
selected 50 VPm neuron pairs, and repeated the procedure described above for the
experimental VPm data. Note that we used this calculation to confirm that drawing
thalamic spike times from the empirical VPm PSTHs, and assuming independence across
these input spike trains, was sufficient to reproduce the empirical trends in synchronous
thalamic spiking.

Quantification and Statistical Analysis

Thalamic Electrophysiology Data Analysis - Mean Response, Burst Ratio, Synchronous
Spike Counts: We report several different basic measurements of spiking activity from
our thalamic units including evoked response and evoked bursting response. Thalamic
firing activity was reported as PSTHs with units of firing rate in Hz, calculated as the
number of spikes within a bin, divided by the size of the bin (see individual plots for bin
size). We determined thalamic evoked response as the initial response (0-30ms) to
sensory stimuli, reported as the number of spikes per stimulus averaged over many trials
(9-102 trials). The corresponding evoked bursting response was determined as number
of bursting spikes per trial in that same post stimulus period. Bursting spikes were defined
as 2 or more spikes that fire at most 4ms apart preceded by 100ms of silence. The 100ms
pre-stimulus activity is based on reported values for T-type calcium bursts (Lu et al., 1992;
Swadlow and Gusev, 2001; Whitmire et al., 2016). All data analysis of the recorded
extracellular thalamic units was accomplished using custom Matlab and Python scripts.

1539

We calculated the mean synchronous spike counts using all pairs of sensory-responsive
thalamic neurons recorded simultaneously using silicon probes, as described previously
(Wright et al., 2021).   Briefly, for each pair of simultaneously-recorded, sensory-
responsive single-units, we considered the spike times in a brief (10 ms) post-stimulus
window, and calculated the relative spike times (which ranged from +20 ms to -20 ms).
We repeated this for all pairs, populating a grand set of relative spike times, separately
for each light level, generating the spike cross-correlogram (CCG). We then summed the
CCG between +5 ms and -5 ms, and divided by the number of contributing pairs. We
repeated these steps using an equivalent number of random relative spike times, and
subtracted the synchronous spike counts for these shuffled relative spike times from the
true synchronous count. This yielded the "mean synchronous spike count (shuffle-
corrected)", or the average number of synchronous spike counts per pair, beyond what
would be predicted from pairs of neurons with randomized relative spike times. We
calculated 95% confidence intervals by re-sampling the relative spike times with
replacement. We performed this analysis for three separate windows: 0 – 10 ms, 10 –
20 ms, and 20 – 30 ms post-stimulus.
1556
Thalamic and Cortical Electrophysiology Data Analysis: Offline spike sorting was
accomplished using either the Plexon Offline Spike Sorter v4 (Plexon, Inc, for tungsten
electrode thalamic recordings), or KiloSort2, followed by curation in Phy (for silicon probe
recordings in thalamus and cortex). We then required each unit to satisfy various criteria
in order to be considered a well-isolated single-unit. We first calculated the signal-to-
noise ratio (SNR), or the amplitude of the mean waveform (trough-to-peak) divided by the
standard deviation.   Second, we calculated the inter-spike-interval (ISI) violation
percentage, or the percentage of all spikes within the 0 – 1 ms inter-spike interval. We
required the SNR to be greater than or equal to 2 (3), and the ISI violation percentage to
be less than or equal to 1 (1.5) for cortex (thalamus). Cortical single-units were classified
as putative Regular Spiking Units (RSUs) or Fast Spiking Units (FSUs) based on the
characteristics of the spike waveform (McCormick et al., 1985; Niell and Stryker, 2010;
Guo et al., 2017; Speed et al., 2019; Yu et al., 2019). Specifically, units with a spike-width
exceeding 0.4 ms (defined as trough-to-peak) were classified as RSUs, and below this
as FSUs. For thalamic single-units, we required the spike-width to exceed 0.3 ms, as
narrower waveforms could reflect activity at TRN synaptic terminals (Barthó et al., 2014).
For multi-unit data, we measured threshold crossings from the continuously recorded
thalamic or cortical activity. Thalamic multiunit activity was captured using a threshold
criterion of 5 standard deviations over the entire recording (Yang et al., 2016). For cortical
multiunit recordings, a manual threshold was set based on each experiment. Additional
data analysis utilized custom scripts using Matlab (Mathworks, Inc).
1578
Cortical units were initially analyzed together, across recording sites, as the dataset was
not sufficient to confidently assess any cross-laminar effects. However, we did repeat

some analyses using only putative layer 4 (L4) units, identified using methods similar to
those described previously (Sederberg et al., 2019), as L4 was particularly important to
this study as the primary thalamorecipient cortical layer. Briefly, for each experiment, we
first calculated the across-trial average local field potential (LFP, or 2 – 200 Hz bandpass-
filtered voltage traces) and multi-unit (MU, threshold crossings of 150 Hz highpass-filtered
voltage traces) responses to punctate whisker deflections. We then calculated the current
source density (CSD) profile of the LFP responses. Finally, we used a combination of the
amplitudes and latencies of LFP deflections, the locations and latencies of CSD sinks,
the amplitudes and latencies of MU firing, and the approximate probe depth to estimate
the center of L4. Specifically, L4 is expected to be 400 – 600 µm below the cortical
surface, and exhibit short-latency, large-amplitude LFP and MU responses, and a short-
latency CSD sink. We required at least two of the three electrophysiological signals (LFP,
CSD, MU) to provide qualitative agreement on the L4 center, and for this center to be
between 400 and 600 µm below the estimated cortical surface, which excluded one of
the six experiments from this analysis. Finally, we considered L4 channels to be those
between 100 µm above and 100 µm below the central channel. A cortical single-unit was
deemed a putative L4 unit if the channel on which its mean waveform was largest was
included in the list of putative L4 channels.

Cortical firing activity was reported as PSTHs with units of firing rate in Hz, calculated as
the number of spikes within a bin, divided by the size of the bin (see individual plots for
bin size). All cortical spikes within a 1ms ISI were removed from analysis (and this was
<1% of total). Synchrony across cortical single-units was computed from spike cross-
correlograms across recorded pairs. Specifically, synchrony was defined as proportion of
spikes from the full (+/- 100ms) cross-correlogram that were in a central +/- 7.5ms window
(Wang et al., 2010; Whitmire et al., 2016).   To determine the number of needed
synchronous events to accurately measure synchrony, we simulated two neurons with a
~5% change in synchrony (assuming a normal distribution). We found that approximately
50 events were required to accurately separate the two distributions.

<u>Voltage Imaging Data Processing:</u> Raw images were loaded and converted from the
SciMedia ". gsd" format using custom scripts and down-sampled by a factor of two. Each
dataset was first normalized to a $\%\Delta F/F_o$ measurement by subtracting and dividing each
trial by the temporal average of the frames 0 to 200ms preceding light delivery (F_o). In
two instances, where 200ms of preceding LED onset activity was not captured, the F_o
was taken as an average 200ms period across no-stimulus trials. Hemodynamic noise
was removed using a PCA background subtraction method. As described in detail
(Borden et al., 2017), *in vivo* ArcLight imaging overlaps with the hemoglobin absorption
spectrum, and therefore contains hemodynamic noise that must be removed for analysis.
Imaging the wildtype mouse cortical surface using the same blue excitation and ArcLight

filter set revealed similar patterns of oscillatory activity, likely through auto-fluorescence
and effects of hemodynamic absorption and blood flow (Ma et al., 2016). The background
PCA subtraction utilizes the auto-fluorescence signal from non-ArcLight transfected
regions to predict the hemodynamic signal across the recorded space. Specifically, the
method uses principal component analysis of non-expressing low background auto-
fluorescence regions (determined from the maximum fluorescence from a non-injected
animal) to find the ongoing hemodynamic components on a single trial basis. Additionally,
the background fluorescence regions were selected at least 1mm away from the recorded
whisker evoked response (Borden et al., 2017). Ideally, these criteria would create a
spatially defined region with little or no ArcLight fluorescence to isolate the hemodynamic
signal from the signal of interest. Each frame is first spatially averaged by either a 200
μm x 200 μm circular averaging (pillbox) filter or a media filter to reduce noise. On a single
trial, the corresponding top five principal components of the low background regions
(which contains approximately 85% of the variance explained) are projected on a pixel by
pixel basis across the entire recording using a lasso regression method with
regularization. The lasso regression utilizes a cross-validated approach to determine the
minimum number of components to develop the model of hemodynamic noise. In order
to prevent the removal of any stimulus evoked activity, each pixel was fit on pre-stimulus
activity (either before light onset for experiments involving optogenetics, or immediately
preceding stimulus delivery). The final predicted hemodynamic signal for each pixel was
subtracted across the entire recording on a pixel by pixel basis. Due to the complex
waveform of the hemodynamic response, a simple notch filter is not effective at
separating the signal from the noise (Borden et al., 2017). We found that the background
PCA subtraction method greatly reduced hemodynamic signal across the entire frame,
compared to the off-ROI method (Borden et al., 2017). In some instances, brief onset and
offset light artifacts of the 590nm light was visible in the recorded ArcLight cortical signal.
To account for any optogenetic transient light artifacts, we only considered the relative
changes in fluorescence during steady state light levels. Both raw and processed images
showed qualitatively similar results.
1650
<u>Awake Voltage Imaging Data Analysis – Dual Camera:</u> In the awake animal, we utilized
a dual camera imaging system to capture a background fluorescence signal for
hemodynamic subtraction. Two different cameras were used to capture the ArcLight and
auto-fluorescence signals, and therefore, pixels could not be directly registered for
subtraction for pixel by pixel correction. Instead, we utilized the same Background PCA
subtraction method to find and develop models of the hemodynamic response based on
the global PCA signal derived from the background image. For the dual camera data,
each component was fit over the entire recording for subtraction of the hemodynamic
noise. Both raw and processed images showed qualitatively similar results. Unless

otherwise noted, each dataset was processed with the Background PCA or Dual Camera
subtraction method as stated above.
1662
Imaging Data Analysis – Peak amplitude, Normalized Peak, and Temporal Properties:
We measured the effect of the optogenetic stimulation on the peak amplitude of the
evoked mean ArcLight fluorescence in the determined cortical barrel. The cortical barrel
region of interest (ROI) for each stimulated barrel and each data set was selected as the
~200 μm x 200 μm region with the largest response 30ms after stimulus delivery. This
determined ROI was used for all subsequent analyses of the temporal response. To better
isolate the evoked amplitude, the frame preceding stimulus delivery (t=-5ms) was
subtracted from the resulting recorded signal. For each recording, the peak amplitude
was defined as the $\Delta F/F_o$ at the time of the maximum average response between 0 and
110ms for the strongest stimuli (1200 Deg/s) presented under control and various
optogenetic conditions. In order to measure the temporal properties of the evoked
response, we concentrated on the timeseries data from the determined cortical barrel
ROI. For normalized fluorescence (Norm $\Delta F/F_o$), each session's peak response was
divided by the average peak response to the strongest stimulus (1200 Deg/s) under the
control condition. The normalization allows for a better comparison across animals which
may have different levels of ArcLight expression. Peak time was defined as the time of
the maximum fluorescence between 0 and 110 ms post stimulus, and the time of return
to baseline as the time when the fluorescence crossed the pre-stimulus baseline value
following the peak. Peak-to-baseline was then calculated as the time between the
fluorescence peak and the return of the fluorescence to baseline. Recovery was defined
as the average fluorescence in the 120-400ms window following the stimulus.
1684
Imaging Data Analysis – Area Measurements: In addition to measuring the peak
response, we also measured the effect of different thalamic states on the evoked area of
sensory cortical activity. We measured the activated area as the number of pixels
exceeding a threshold using the average response at the peak frame (0-110ms)
preceding stimulus delivery. Similar to other studies (Lustig et al., 2013; Millard et al.,
2015), we measured the spatial activation using the 70% threshold. To compare the area
independent of amplitude changes, we normalized the peak frame by dividing by the peak
fluorescence in each condition (Control and LED). In order to isolate the evoked activity
from ongoing activity, we subtracted the frame preceding stimulus delivery (t=-5ms).
Different thresholds had no effect on the observed trends.
1695
Statistical Analysis: All tests were conducted using the MATLAB Statistics Toolbox
(Mathworks, Inc.). For all measurements, we determined if the specific data sets were
normally distributed using the Lilliefors test for normality. If the data were normal, we used
the appropriate (paired or unpaired) t-test for statistical difference. If the population was
determined to have non-normal distributions, we conducted non-parametric Wilcoxon
signed-rank tests to determine statistical significance. All sample sizes are reported in the

figure captions and Results text, along with an indication of particular test and the
corresponding statistical significance level (* - p<0.05, ** - p<0.01, *** - p<0.001 in
figures).
1705
1706

Figure 1

A
Facial Vibrissae
S1 Cortex
VPm Thalamus
Brainstem

B
CCD camera
S1
whisker sensory stimulus
VPm
tungsten
LED
isoflurane anesthesia

C
$t_{silence}$ t_{isi}
tonic   burst

burst spikes
$t_{isi} < 4$ ms
$t_{silence} > 100$ ms

D anesth. VPm grand PSTH

all spikes
burst spikes
Control
LED

Rate (Hz) Rate (Hz)
sensory stim onset
time (ms)

E *anesth. VPM evoked response*

Firing Rate (Hz)
**
***
Control   LED

anesth. cortical MU recordings
LED
Control
sensory stim onset
time (ms)

F anesth. CTX evoked GEVI response

Control
LED
5ms 10ms 15ms 20ms 25ms
sensory stim onset
-%dF/Fo

G anesth. CTX evoked GEVI response

Pk. Norm. %dF/ Fo
Pk.%dF/Fo
*
Control   LED

LED
Control
Pk. Norm. %dF/ Fo
sensory stim onset
time (ms)

Figure 2

Figure 3

A silicon probe S1 LED VPm

B S1 example session

Cell 15 Cell 10 Cell 5 Cell 1

cell and trial

LED

S1 single unit activity

transient

steady-state

mean rate (Hz)
18 16 14 12 10 8 6

time (s)
-0.6 -0.4 -0.2 0 0.2 0.4 0.6

C

○ transient
● steady-state

mean rate (Hz)
20 15 10 5

***

*

n.s.

Control   LED   ⚡LED

Figure 4

Figure 5

A awake mean S1 response (example session)

Control

LED

sensory stim onset

5ms 10ms 15ms 20ms 25ms

1 mm

-0.2 0 0.2 0.4 0.6
-%dF/Fo

B

rel. Evoked Peak S1

Peak -%dF/Fo

**

*

Control LED LED

rel. Evoked Norm. Peak S1

norm. Peak -%dF/Fo

Control LED LED

C

A32 Probe

S1

VPm

LED

whisker sensory stimulus

D awake S1 CTX grand PSTH

Control
LED

mean rate (Hz)

sensory stim onset

time (ms)

E

abs. Evoked (5-100ms)

mean rate (Hz)

n.s.

Control LED LED

rel. Evoked (Response-Baseline)

mean rate (Hz)

*

n.s.

Control LED LED

Figure 6

A — mean GEVI response (example session)

1mm

LED

-%dF/Fo
-0.4 -0.3 -0.2 -0.1 0 0.1

B

Control   LED

max amp

space

normalize

normed amp
1.0
0.7

space

C — normalized amplitude
mean GEVI ROI time-series (example session)

peak-to-baseline

mean recovery
(avg. 120 - 400 ms)

norm. %dF/Fo
1

Control
LED

stim onset

-0.1   0.1   0.2   0.3   0.4   0.5

D

evoked norm. area >70%
Area (mm²)
0.25
0.15
0.05
*
Control LED ⚡LED

peak-to-baseline (ms)
Time (ms)
200
150
100
50
**
Control LED ⚡LED

mean recovery
Norm. Response
0
-1
-2
*
Control LED ⚡LED

Figure 7

Figure 8

A

single trial
target spike

cell 1

cell 2

time rel to target spike (ms)
-100   0   100

repeat for
all spikes
and trials

pairwise cross-
correlogram (CCG)

count →

Lag (ms)
-100   0   100

$$synchrony = \frac{sum(CCG[-7.5ms:7.5ms])}{sum(CCG)}$$

B Example FSU Pair
Spiking Activity

Trial

Control

LED

stim onset   Time(ms)

Synchronous Spikes
only

Trial

Control

LED

C

Avg. Population
Pairwise CCG

Control

LED

Time(ms)
-25   0   25

D

***

Synchrony

Spont.   Control   LED   LED

Figure 9