A tail of plumage colouration: disentangling geographic, seasonal, and dietary effects on

plumage colour in a migratory songbird

Sean M. Mahoney[1*], Matthew W. Reudink[1], Andrea Contina[2], Kelly A. Roberts[3], Veronica T.

Schabert[3], Emily G. Gunther[3], Kristen M. Covino[4]

[1]M.W. Reudink and S.M. Mahoney, Department of Biological Sciences, Thompson Rivers
University, Kamloops, BC, Canada. — [2]A. Contina, Department of Integrative Biology,
University of Colorado Denver, Denver, CO, USA. — [3] K.A Roberts, V.T. Schabert, and E.G.
Gunther, Department of Animal Behavior, Ecology, and Conservation, Canisius College,
Buffalo, NY, USA — [4]K.M. Covino (Kristen.covino@lmu.edu) Biology Department, Loyola
Marymount University, Los Angeles, CA,USA.

*Corresponding author: smahoney@tru.ca
Short Title: American redstart colour variation

Abstract

Plumage ornamentation in birds serves critical inter- and intra-sexual signaling functions. While

carotenoid-based plumage colouration is often viewed as a classic condition-dependent sexually

selected trait, plumage colouration can be influenced by a wide array of both intrinsic and

extrinsic factors. Understanding the mechanisms underlying variation in colouration is especially

important for species where the signaling function of ornamental traits is complex or the

literature conflicting. Here, we examined variation in the yellow/orange tail feathers of American

redstarts (*Setophaga ruticilla*) passing through two migratory stopover sites in eastern North

America during both spring and fall migration to assess the role of geographic variation and

seasonality in influencing feather colouration. In addition, we investigated whether diet during

moult (inferred via stable isotope analysis of feather δ^{15}N and δ^{13}C) influenced plumage

colouration. Our findings indicate that geographic variation, season, and diet all influence

individual differences in American redstart colouration. However, the extent to which these

factors influence colour expression is largely dependent on the colour metric under study, likely

because different colour metrics reflect different attributes of the feather (e.g., structural

components vs. pigment deposition). The effects of diet (δ^{15}N) and season were pronounced for

brightness, suggesting a strong effect of diet and feather wear/degradation on feather structure.

Though hue, a metric that should strongly reflect pigment deposition, was also associated with a

reduction in ornamentation from spring to fall, that effect was dependent on age, with only adults

experiencing a reduction in ornamentation. Taken together, our results highlight the numerous

sources of variation of plumage coloration and underscores the difficulty of unraveling complex

visual signaling systems, such as those in American redstarts.

Keywords: carotenoid, plumage, tetrahedral colourspace, stable isotope, nitrogen, carbon

Introduction

The colourful plumage exhibited by birds serves important roles for both inter-and intrasexual communication (reviewed in Hill 2006; Santos et al. 2011). The literature linking variation in plumage colouration to individual quality and reproduction is multitudinous (Griffith and Pryke 2006; Hill 2006), yet the patterns are often complex and may vary across seasons and populations (Marini et al. 2015; Reudink et al. 2015). As such, much research has been dedicated to understanding the mechanisms underlying variation in the expression of ornamental plumage colouration (Siefferman and Hill 2005; Griggio et al. 2009; Newton and Dawson 2011; Doutrelant et al. 2012; Reudink et al. 2015).

Plumage colouration in birds can be produced by various mechanisms, including the deposition of pigments (e.g., carotenoids, melanins, and porphyrines) and the arrangement of melanosomes within the feather microstructure producing structural (purple, blue, green, and iridescent) colours. Carotenoid colouration (yellow, orange, red plumage) has received by far the most attention, in large part because the expression of carotenoid-based colouration is directly dependent on birds acquiring carotenoids in the diet (the pigments cannot be synthesized *de novo*) and subsequently modifying dietary pigments through a costly physiological process (Lozano et al. 1994; Hill and Montgomerie 1994; Weaver et al. 2018). Thus, carotenoid-based colouration is often viewed as a classic honest indicator of individual condition and quality (Hill 2011; Hill and Johnson 2012; Weaver et al. 2018; but see Simons et al. 2015).

Geographic variation in sexual signals has been observed across many taxa (Hill 2006; Norrris et al. 2007; Badyaev et al. 2012). These differences have been attributed to genetic drift, differing natural or sexual selection pressures (Hill 2006; Wilkins et al. 2013), arbitrary mate choice (Prum 2010), or differences in environmental conditions (e.g., dietary carotenoid

availability; Hill 1993). Regardless of the mechanism, understanding variation within and among

populations is critical for understanding the function and evolution of ornamentation.

Diet during feather growth has been directly linked to ornamentation and feather quality

in both experimental (Hill 2000; Peters et al. 2008; Koch et al. 2017) and observational studies

(Slagsvold and Lifjeld 1985; Arriero and Fargallo 2006; Eeva et al. 2009; but see Ferns and

Hinsley 2008). For observational studies of free-living birds, one of the challenges is

determining dietary conditions during moult. However, stable isotopes of carbon (δ^{13}C) and

nitrogen (δ^{15}N) vary predictably in the environment and are incorporated into growing tissues

(Fry 2006; Bowen 2010). Because feathers are inert once grown, they can provide insight into

diet during tissue growth. δ^{13}C has been used extensively in dietary studies because carbon

isotope ratios vary based on plant water stress and photosynthetic system (Lajtha and Marshall

1994). Because these isotope elements are transferred up the food chain and eventually

incorporated into animal tissues, they can provide important insight into habitat and diet during

tissue growth (Hobson 1999). Nitrogen is commonly employed in food web studies, as ^{15}N is

preferentially incorporated into animal tissues as one moves up the food chain (Post 2002;

Poupin et al. 2011). Thus, analysis of δ^{15}N tissue values can provide insight into the trophic level

of animals within the environment. However, δ^{15}N values can also be influenced by

environmental conditions, as they tend to be positively associated with temperature and

negatively associated with rainfall (Craine et al. 2009).

A recent study on Bullock's orioles (*Icterus bullockii*) demonstrated that feather δ^{15}N

values were directly associated with feather carotenoid content and hue (Sparrow et al. 2017). In

the case of Bullock's orioles, birds with lower feather δ^{15}N values expressed more orange-shifted

hue. Because Bullock's orioles rely on both insects and fruit during moult, the authors suggested

that individuals with a relatively higher amount of fruit in their diet (or those moulting in habitats

that resulted in lower $\delta^{15}N$ values) may have been in better condition and better able to

metabolically convert dietary carotenoids to red or orange keto-carotenoids (McGraw 2006;

Sparrow et al. 2017; Weaver et al. 2018).

Though feathers are metabolically inert once grown, feather colours are not necessarily

static across the life of the feather. Feather wear can modify feather colouration; for example,

European starlings (*Sturnus vulgaris*) lose their distinctive winter spots through wear and

abrasion on the feather tips leaving only the melanin-based feather structure behind for the

breeding season. For most species, these changes are less dramatic but can still have important

fitness consequences. Feather abrasion, ultraviolet (UV) radiation, and feather degrading bacteria

can all lead to changes in ornamentation after initial feather growth (Örnberg et al. 2002;

Shawkey and Hill 2004). For example, carotenoid-based red colouration in House finches

(*Haemorhous mexicanus*) becomes increasingly more saturated and hue becomes more yellow-

shifted in the spring following moult (McGraw and Hill 2004). Melanin-based plumage in Fox

sparrows (*Passerella iliaca*) also changes colour (Weckstein et al. 2002), possibly due to UV

exposure or accumulation of dirt and oils on the feathers (Montgomerie 2006). Additionally,

colour in many species changes with age — sometimes dramatically in through delayed plumage

maturation (Hawkins et a. 2012) and sometimes more subtly as individuals age (Marini et al.

2015; Ward et al. 2021); therefore, it is important for studies to assess the environmental,

biological, ontological, and temporal influences on feather colour to help unravel the

mechanisms s regulating this phenotypic trait.

For the American redstart (*Setophaga ruticilla*), the brilliant orange and black plumage

ornamentation expressed by males in definitive plumage has been extensively studied (e.g.,

Reudink et al. 2009a,b; Marini et al. 2015; Reudink et al. 2015), yet the role of plumage for

inter- and intra-sexual signaling remains frustratingly complicated and unresolved (see Marini et

al. 2015; Table 1). As such, recent research on this species has focused on understanding the

mechanisms underlying variation in plumage colouration (Marini et al. 2015; Reudink et al.

2015). In an 11-year study, Reudink et al. (2015) found that rainfall and temperature during

moult was directly linked variation in feather hue; however, individuals themselves also appear

to change throughout life, with males expressing most orange-shifted hue expressed during their

first breeding season in definitive plumage (Marini et al. 2015). At a broader scale, American

redstarts also exhibit marked geographic variation in plumage ornamentation across the range

(Norris et al. 2007). Adding to the complexity of this system, American redstarts exhibit delayed

plumage maturation, with young (hatch-year [HY] and second-year [SY]) males exhibiting

female-like grey and yellow plumage throughout their first year of life, while adult (after hatch

year [AHY] and after second year [ASY]) birds display jet black and orange plumage. As such,

male redstarts of different age classes appear to utilize different metabolic pathways prior to

depositing ingested dietary carotenoids. Adding to the complexity, these potentially important

signals are grown at different times of year and under different conditions — in the nest in early-

to-mid summer for young birds and after breeding in late summer for adult birds. Thus,

American redstarts in different age- and sex-classes are likely differentially affected by both

intrinsic and extrinsic factors.

Our goal was to examine the relative importance of the several factors that may underlie

variation in the expression of ornamental colouration in the American redstart. We investigated

plumage colouration of American redstarts passing through two different migration stopover

sites in eastern North America (Fig. 1, Braddock Bay Bird Observatory (BBBO) and Appledore

Island Migration Station (AIMS)) during both spring and fall migration and examined the stable

isotope ratios of $\delta^{13}C$ and $\delta^{15}N$ in the feathers of each individual. As such, we were able to

concurrently determine the relative effects of geographic variation (site), temporal variation

(season), biological (sex), ontological (age), environment and diet ($\delta^{13}C$ and $\delta^{15}N$), along with

age and sex, in the expression of plumage ornamentation. We predicted there would be

differences in plumage colouration between sites and that birds captured in fall would exhibit a

greater degree of ornamentation (increased red chroma, orange-shifted hue, higher brightness)

compared to those captured in spring. In addition, we predicted that birds with more negative

$\delta^{13}C$ feather isotope values (indicating habitats with lower water stress) and more positive $\delta^{15}N$

feather isotope values (indicating feeding at a higher trophic level) would exhibit increased

ornamentation.

Methods

Field Methods

We collected feather samples from American redstarts, a sexually dimorphic Nearctic-

Neotropical migrant that breeds in deciduous forests in the United States and Canada (Sherry et

al. 2020). Our collection sites were two locations of similar latitude but different proximity to the

Atlantic coast: Braddock Bay Bird Observatory (43° 9'N, 77° 36'W) in New York State and

Appledore Island Migration Station (42°58′N, 70°36′ W) off the coast of Maine. Braddock Bay

Bird Observatory (BBBO) is an inland site just south of Lake Ontario and the Appledore Island

Migration Station (AIMS) is located seven miles offshore from the Maine-New Hampshire

border (Fig. 1).

Field researchers captured birds at AIMS using mist nets (weather permitting) checked at

least every 20 minutes from sunrise to sunset from 11 May to 7 June during spring migration and

14 August to 15 September during fall migration 2016 and 2017. During 2016 and 2017 at

BBBO field researchers captured birds with mist nets checked every half hour starting at sunrise

and for 6 hours thereafter from 14 April through 8 June (spring) and 12 August through 2

November (fall). A United States Geological Service leg band was placed on each bird and sex

and age were determined according to Pyle (1997) based on plumage, molt, and/or degree of

skull pneumatization. In fall, age was determined to be hatch-year (hereafter young) or after-

hatch-year (hereafter adult). In spring, age was determined to be second-year (young) or after-

second-year (adult). We sampled the left and right fifth rectrix feathers (second from the outside

on each side of the tail) provided the feathers were not replaced outside of regular molt, in which

case we did not sample feathers and that individual was excluded from this study. In total, we

collected feathers from 134 individuals; 59 from BBBO and 75 from AIMS (Table S1).

Sample preparation and stable isotope analysis

We conducted analysis of stable isotope ratios of carbon ($^{13}C/^{12}C$) and nitrogen ($^{15}N/^{14}N$)

on keratin tissues obtained from feathers plucked during mist-netting efforts at AIMS and

BBBO. We cleaned each feather to remove oils and debris with immersion in a 2:1 chloroform-

methanol solution followed by air-drying for 24-h under a fume hood. We then washed each

feather with a 1:30 solution of detergent (Fisher Scientific Versa-Clean #04-0342) and deionized

water before rinsing three times with deionized water (Paritte and Kelly 2009; Chew et al. 2019).

We cut a 350 µg fragment and packed in tin capsules placed in a Carlo Erba NC2500 elemental

analyzer (CE Instruments, Milano, Italy), interfaced with a Thermo Delta V+ isotope ratio mass

spectrometer (Thermo-Finnigan, Bremen, Germany) operated by the Central Appalachians

Stable Isotope Facility (CASIF) at the Appalachian Laboratory (Frostburg, Maryland, USA). We

matched each sample with powdered keratin (porcine) from Spectrum Chemicals, a quality

control (QC) in-house standard. We report feather carbon (δ^{13}C) and nitrogen (δ^{15}N) as mean $\pm$

SD in delta notation of parts per thousand (‰) from the standards ($\delta D_{sample} = [(R_{sample}/R_{standard}) -$

$1] \times 1,000$) compared to the Vienna PeeDee Belemnite (V-PDB) and AIR scales, respectively.

The standard deviation of the reference material was ± 0.12‰ (δ^{13}C) and ± 0.11‰ (δ^{15}N).

Colour Variation

Prior to cutting and packaging feathers (above) but after the above-described cleaning

steps, feather colouration was determined by measuring the reflectance within the avian visual

range (300-700nm) with an Ocean Optics Jaz Spectrometer (Reudink et al. 2015). The yellow-

orange vane sections of each rectrix were randomly scanned ten times each. Upon each use, the

spectrometer was calibrated by scanning an Ocean Optics WS-1 white standard and a Colorline

Ebony #142 dark standard. Additionally, the spectrometer was also calibrated between each

feather using the Ocean Optics WS-1 white standard.

We tested how American redstart colour variation was explained by site, season, sex, age,

and isotopic signatures at sampling using separate mixed models in the *lme4* package in R. We

used brightness, red chroma, red hue, and "colourspace" in our models. Colourspace was derived

from a Principal Components Analysis (PCA) of an individual redstart colour in tetrahedral

space (see Tetrahedral Models below). Either hue, chroma, brightness, or colourspace PC1 was

included as the response variable, site, season, sex, age, δ^{13}C and δ^{15}N were main effects, and the

interaction between age and season because redstarts exhibit delayed colour maturation

(Hawkins et al. 2012). Year was included as a random effect. We then undertook model

reduction for all mixed models based on the change in Akaike information criterion corrected for

small sample sizes (AICc, Burnham and Anderson 2003) between the full model and each

reduced model. We chose our final model based on the lowest AIC (Burnham and Anderson

2003). ΔAIC values within 4 were considered competitive and we selected the best fitting model

using ANOVAs.

Tetrahedral Models

We assessed visual discrimination of statistical differences using tetrahedral colour

models, which correct for avian visual systems (Vorobyev et al. 1998), in the R package *pavo2*

(Maia et al. 2019). Although the visual system of American redstarts is unknown, most

passerines are ultraviolet sensitive (Ödeen and Håstad 2003), so our tetrahedral models use this

visual system. We used a receptor-noise discrimination model to calculate the photon catch of

each cone type used in avian colour vision. We used a value of 0.1 for colour (chromatic) and

luminance (achromatic) discrimination and the photoreceptor ratios to 1:2:2:4 (ultraviolet,

shortwave, mediumwave, longwave) in our models (Maier and Bowmaker 1993), which were

selected based on a recent review of noise receptor estimates (Olsson et al. 2018). Finally, we

summarized the colour variables derived from the tetrahedral model using a PCA to generate a

single value representing the "colourspace" for each individual (see above *Colour variation*

Methods).

Results

Colour Variation

Our AICc model selection identified the reduced models for brightness, chroma, and hue and the full model for colourspace PC1 as the top models (Table 2 and Table S2). American redstart brightness was explained by site, season, and δ^{15}N (Table 2). In both the spring and fall, American redstart feathers were brighter at the AIMS site (Fig. 1, site: $F_{1,134}=6.45$, P=0.012; Season: $F_{1,134}=12.82$, P<0.0001). Brightness was positively related to δ^{15}N (Fig. 1. $F_{1,134}=11.78$, P=0.001). Red chroma did not vary among any of our predictor variables (Table 2).

Hue was explained by season, sex, age, and the interaction between season and age (Table 2 and Table S2). Hue was more red-shifted in males (Fig. 3, $F_{1,134}=12.39$, P=0.001) and in adults (Fig. 3, $F_{1,134}=9.43$, P=0.003) and following moult in the fall (Fig. 3, $F_{1,134}=26.09$, P<0.0001). There was an interaction between season and age indicating that although hue in young birds did not vary between seasons, adult hue was more red-shifted in the fall (Fig. 3, $F_{1,134}=7.96$, P=0.006).

From our PCA of tetrahedral colourspace, PC1 (81% of variation) was associated with hue length Φ, which describes latitudinal hue shifts in tetrahedral space, that is, shifts towards medium or ultraviolet wavelengths (Table S3; Stoddard and Prum 2008). PC2 (11% of variation) was associated with hue length θ, which describes longitudinal shifts in tetrahedral space and indicates shifts towards longer or shorter wavelengths (Table S3; Stoddard and Prum 2008). In our PCA, individuals with higher PC1 and 2 scores exhibited more yellow-shifted hues and individuals with lower scores had more red-shifted hues (Table S3). Colourspace PC1 was explained by site, season, age, sex (Table 2 and Table S2). Adults generally had lower

colourspace PC1 scores, indicating red-shifted plumage colour and young birds had higher

colourspace PC1 scores indicating more yellow-shifted colour plumage. Young and adult

redstarts occupied different colourspaces during the spring and fall (Fig. 4, Age: $F_{1,132.1}=27.07$,

P<0.0001; Season: $F_{1,133.3}=90.91$, P<0.0001). Colourspace also varied by site and sex (Fig. 4.

Site: $F_{1,133.7}=11.20$, P=0.001; Sex: $F_{1,132}=10.79$, P=0.001), where males had higher colourspace

PC1 scores than females, and individuals at the AIMS site had lower PC1 scores on average.

Discussion

Carotenoid-based colouration serves important inter- and intra-sexual signaling functions;

however, for some species like the American redstart, the relationship between fitness-related

components and colour metrics can be perplexing and seemingly contradictory (Reudink et al.

2015). Part of the complexity likely stems from our incomplete understanding of the mechanisms

underlying variation in American redstart colouration. Here, we demonstrate that geographic

variation, season, and diet all underly intra-individual differences in American redstart

colouration. However, the extent to which these factors influence colour expression is largely

dependent on the colour metric under study, likely because different colour metrics reflect

different aspects of feather quality (e.g., structural components vs. pigment deposition).

Geographic variation in plumage ornamentation has been widely documented across

numerous species (Hill 2006; Badyaev et al. 2012) including the American redstart (Norris et al.

2007). Norris et al. (2007) found that individuals moulting at higher latitudes expressed plumage

with higher red chroma compared to those at lower latitudes. In this case, differences in

colouration were only detected between our two study sites (separated by 580km), and only for

brightness and colourspace PC1, not hue or chroma. All birds in our study were captured during

migration and, as such, we lack information in the specific moult origins for each individual.

Norris et al (2007) suggested that geographic variation in American redstart colouration was

likely driven by differences in dietary carotenoid availability or condition-related effects (e.g.,

parasite load) mediating carotenoid metabolism. Hue and red chroma should be most closely

linked to carotenoid content (Saks et al. 2003; Montgomerie 2006) and while brightness is

negatively associated with feather carotenoid content, it is strongly mediated by feather structure

(Shawkey and Hill 2005). Thus, while the site-specific differences we observed may be due to

differences in carotenoid availability at the moult location, these differences may also reflect

longitudinal geographic variation in condition, as one site was inland while the other was more

coastal, but both were of similar latitude.

As expected, American redstarts expressed differences in brightness, hue, and

colourspace PC1 between seasons, with higher brightness, more red-shifted hue, and a

colourspace PC1 associated with more red/orange-shifted colouration during the fall, following

molt. However, variation in hue was dependent on age; while adults were more red-shifted in fall

than spring, we detected no differences in first-year birds, potentially indicating that the orange

colouration expressed by adult males may be more susceptible to the effects of UV degradation,

feather-degrading bacteria, or feather wear between seasons (Örnborg et al. 2002; Delhey et al.

2006; Shawkey et al. 2007; Shawkey et al. 2009). Alternatively, if birds passing through

migratory stopover sites in spring and fall differ in their moult location, differences between

spring and fall plumage could simply reflect geographic variation in moult location.

Only brightness was associated with $\delta^{15}N$; birds with higher feather $\delta^{15}N$ values had

feathers with higher brightness. Because $\delta^{15}N$ is directly linked to diet (Pearson et al. 2003;

Gómez et al. 2018), these results suggest that American redstarts eating arthropods enriched with

^{15}N (e.g., *Heteromurus nitidus*, *Drosophilia melanogastor;* Oelbermann and Scheu 2002) likely

are not acquiring additional dietary carotenoids; however, these dietary sources may be

advantageous during feather growth, allowing the birds to create feathers with a high-quality

microstructure. Whether or not feather brightness *per se* is a signal used by American redstarts is

unclear; however, Reudink et al. (2009b) demonstrated that males that occupied higher-quality

winter-territories expressed feathers with higher brightness and Reudink et al. (2009a) found that

polygynous males had higher feather brightness than monogamous males. Because polygyny

requires the ability to hold multiple territories or a single large territory, Reudink et al. (2009a)

suggested that feather brightness may act as a social signal during both the breeding and

wintering season, directly reflecting a male's competitive ability. Thus, diet during moult may

potentially impact intrasexual signaling.

In a study of Bullock's orioles, Sparrow et al. (2017) found that feather δ^{15}N values were

associated with feather colouration and feather carotenoid content. The authors found that while

δ^{15}N values were positively associated with overall carotenoid concentration, birds with more

orange-shifted hue had feathers with lower δ^{15}N values. The authors suggested that low δ^{15}N

values could indicate a diet with a higher fruit:insect ratio and birds with more orange-shifted

hue, though they had lower overall carotenoid concentrations, may have had a higher abundance

of specific orange or red carotenoids. It is important to note that American redstarts and

Bullock's oriole diets differ considerably; orioles have a more varied diet of both fruits and

insects (Flood et al. 2016), while redstarts are primarily insectivores (Robinson and Holmes

1982). Thus, the way in which δ^{15}N is associated with the expression of carotenoid-based

colouration is likely highly dependent on the dietary ecology of the species. It is also important

to consider that our observed differences may be due to accumulation of dirt in feathers or due to

degradation caused by UV exposure (Örnborg et al. 2002). The decrease in brightness and hue

between the fall and spring is consistent with previous studies (Prum et al. 1999). Interestingly,

in our study red chroma did not significantly vary, as was previously found in carotenoid-red

colouration as a result of abrasion or soiling (McGraw and Hill 2004). However, without

estimates of dirt on feathers, microscopy of feather nanostructure, or repeated measures, we can

only speculate on this effect and encourage future work to consider adventitious colouration.

Literature Cited

Arriero, E., and Fargallo, J. A. 2006. Habitat structure is associated with the expression of
carotenoid-based coloration in nestling blue tits *Parus caeruleus*. – Naturwissenschaften
93: 173–180.

Badyaev, A. V., Belloni, V., and Hill, G. E. 2012. House Finch (*Haemorhous mexicanus*. – In
Rodewald, P. G. (eds), The Birds of North America. Cornell Lab of Ornithology. Retrieved
from: https://birdsna.org/ Species-Account/bna/species/houfin. http://dx.doi.org/10.2173/
bna.46.

Bowen, G.J., 2010. Isoscapes: spatial pattern in isotopic biogeochemistry. Annu. Rev. Earth
Planet. Sci. – 38: 161-187.

Burnham, K. P., and Anderson, D. R. 2002. Model selection and multimodel inference: A
practical information-theoretic approach. New York (USA)

Chew, B., Kelly, J., and Contina, A. 2019. Stable isotopes in avian research: a step by step
protocol to feather sample preparation for stable isotope analysis of carbon (δ^{13}C), nitrogen
($\delta1^{5}$N), and hydrogen (δ^{2}H). Version 1.1. protocolsio.
dx.doi.org/10.17504/protocols.io.z2uf8ew.

Craine, J. M., Elmore, A. J., Aidar, M. P. M., Bustamante, M., Dawson, T. E., Hobbie, E. A., and
Wright, I. J. 2009. Global patterns of foliar nitrogen isotopes and their relationships with
climate, mycorrhizal fungi, foliar nutrient concentrations, and nitrogen availability. – New
Phytologist 183: 980–992.

Delhey, K., Peters, A., Johnsen, A., and Kempenaers, B. 2006. Seasonal changes in blue tit
crown color: do they signal individual quality? – Behav. Ecol. 17: 790-798.

Doutrelant, C., Gregoire, A., Midamegbe, A., Lambrechts, M. M., and Perret, P. 2012. Female
plumage colouration is sensitive to the cost of reproduction. An experiment in blue tits. – J.
Anim Ecol. 81: 87–96.

Eeva, T., Sillanpaa, S., and Salminen, J. P. 2009. The effects of diet quality and quantity on
plumage colour and growth of great tit *Parus major* nestlings: A food manipulation
experiment along a pollution gradient. – J. Avian Biol. 40: 491–499.

Ferns, P. N., and Hinsley, S. A. 2008. Carotenoid plumage hue and chroma signal different
aspects of individual and habitat quality in tits. – IBIS 150: 152–159.

Flood, N. J., Schlueter, C. L., Reudink, M. W., Pyle, P. Patten, M. A., Rising, J. D., and
Williams, P. L. 2016. Bullock's Oriole (*Icterus bullockii*), version 3.0. In The Birds of
North America (P. G. Rodewald, editor). Cornell Lab of Ornithology, Ithaca, New York,
USA.

Fry, B., 2006. Stable isotope ecology. Vol. 521. New York: Springer 318 p.
363
364   Gómez, C., Larsen, T., Popp, B., Hobson, K. A. and Cadena, C. D., 2018. Assessing seasonal
365   changes in animal diets with stable-isotope analysis of amino acids: a migratory boreal
366   songbird switches diet over its annual cycle. – Oecologia, 187: 1-13.
367
368   Griffith, S. C., and Pryke, S. R., 2006. Benefits to females of assessing color displays. – In: Hill,
369   G. E., McGraw, K. J. (eds) Bird coloration, vol 2. Function and evolution. Harvard
370   University Press, Cambridge, 233-279.
371
372   Griggio, M., Serra, L., Licheri, D., Campomori, C., and Pilastro, A. 2009. Moult speed affects
373   structural feather ornaments in the blue tit. – J. Evol. Biol. 22:782–792.
374
375   Hawkins, G. L., Hill, G. E., and Mercadante, A. 2012. Delayed plumage maturation and delayed
376   reproductive investment in birds. – Biol. Rev. 87: 257–274
377
378   Hill, G. E. 1993. Geographic variation in the carotenoid plumage pigmentation of male house
379   finches (*Carpodacus mexicanus*). – Biol. J. Linn. Soc. 49: 63-86.
380
381   Hill, G. E. 2000. Energetic constraints on expression of carotenoid-based plumage coloration. –
382   – J. Avian Biol. 31: 559-566.
383
384   Hill, G. E. 2006. Female mate choice for ornamental coloration. In: Hill, G. E., McGraw, K. J.
385   (eds) Bird coloration, Vol. 2. Function and evolution. Harvard University Press,
386   Cambridge, pp. 137–200.
387
388   Hill, G. E., and Johnson, J. D. 2012 The vitamin A-redox hypothesis: a biochemical basis for
389   honest signaling via carotenoid pigmentation. – Am. Nat. 180: E127–E150.
390
391   Hobson, K. A., 1999. Stable-carbon and nitrogen isotope ratios of songbird feathers grown in
392   two terrestrial biomes: implications for evaluating trophic relationships and breeding
393   origins. – The Condor 101: 799-805.
394
395   Koch, R. E., Josefson, C. C., and Hill, G. E. 2017. Mitochondrial function, ornamentation, and
396   immunocompetence. – Biol. Rev. 92: 1459– 1474.
397
398   Lajtha, K., and Marshall, J. D. 1994. Sources of variation in isotopic composition in plants. – In:
399   Lajtha, K., and Marshall, J. D. (eds), Stable isotopes in ecological and environmental
400   sciences. Oxford University Press. pp. 1–21.
401
402   Lozano, G. A. 1994. Carotenoids, parasites, and sexual selection. – Oikos 70: 309–311.
403
404   Marini, K. L., McKellar, A. E., Ratcliffe, L. M., Marra, P. P., and Reudink, M. W. 2015. Age-
405   related change in carotenoid-based plumage of the American Redstart (*Setophaga
406   ruticilla*). – J. Ornithol. 156: 783-793.
407

McGraw, K. J., and Hill, G. E. 2004. Plumage color as a dynamic trait: carotenoid pigmentation
of male house finches (*Carpodacus mexicanus*) fades during the breeding season. – Can. J.
Zool. 82: 734-738.

Maier, E. J., and Bowmaker, J. K. 1993. Colour vision in the passeriform bird, *Leiothrix lutea*:
correlation of visual pigment absorbance and oil droplet transmission with spectral
sensitivity. – J. Comp. Phys. A 172: 295–301.

McGraw, K. J. 2006. The mechanics of carotenoid coloration in birds. – In: Hill, G.E. and &
McGraw, K. J. (eds), Bird coloration, Vol. 1: mechanisms and measurements. Harvard
University Press. 177–242.

Montgomerie, R. 2006. Cosmetic and adventitious colors. – In: Hill, G.E. and & McGraw, K. J.
(eds), Bird coloration, Vol. 1: mechanisms and measurements. Harvard University Press.
399-427.

Newton, I., and Dawson, A. 2011. Seasonal changes in moult, body mass and reproductive
condition in siskin *Carduelis spinus* exposed to daylength regimes simulating different
latitudes. – J. Avian Biol. 42: 22–28.

Ödeen, A., and Håstad, O. 2003. Complex distribution of avian color vision systems revealed by
sequencing the SWS1 opsin from total DNA. – Molec. Biol. and Evol. 20: 855–861.

Oelbermann, K., and Scheu, S. 2002. Stable isotope enrichment (δ^{15}N and δ^{13}C) in a generalist
predator (*Pardosa lugubris*, Araneae: Lycosidae): effects of prey quality. – Oecologia,
130: 337-344.

Olsson, P., Lind, O. and Kelber, A. 2018. Chromatic and achro- matic vision: parameter choice
and limitations for reliable model predictions. – Behav. Ecol. 29: 273–282.

Örnborg, J., Andersson, S., Griffith, S. C. and Sheldon, B. C. 2002. Seasonal changes in a
ultraviolet structural colour signal in blue tits, *Parus caeruleus*. – Biol. J. Linnean Soc. 76:
237-245.

Norris, D. R., Marra, P. P., Kyser, T. K., Ratcliffe, L. M. and Montgomerie, R. 2007. Continent-
wide variation in feather colour of a migratory song- bird in relation to body condition and
moulting locality. – Biol. Lett. 3: 16–19.

Paritte, J. M. and Kelly, J. F. 2009. Effect of cleaning regime on stable-isotope ratios of feathers
in Japanese Quail (*Coturnix japonica*). – The Auk 126:165–174.

Pearson, S. F., Levey, D. J., Greenberg, C. H. and Del Rio, C. M. 2003. Effects of elemental
composition on the incorporation of dietary nitrogen and carbon isotopic signatures in an
omnivorous songbird. – Oecologia 135: 516-523.

Peters, A., Delhey, K., Andersson, S., Van Noordwijk, H., and Förschler, M. I. 2008.
Condition☐dependence of multiple carotenoid☐based plumage traits: an experimental
study. – Functional Ecology 22: 831-839.

Post, D. M. 2002. Using stable isotopes to estimate trophic position: Models, methods, and
assumptions. – Ecology 83: 703–718.

Poupin, N., Bos, C., Mariotti, F., Huneau, J.-F., Tomé, D. and Fouillet, H. 2011. The nature of
the dietary protein impacts the tissue-to-diet ^{15}N discrimination factors in laboratory rats. –
PLoS ONE 6: e28046.

Prum, R. O. 2010. The Lande–Kirkpatrick mechanism is the null model of evolution by
intersexual selection: implications for meaning, honesty, and design in intersexual signals.
– Evolution 64: 3085–3100

Prum, R. O., Torres, R, Kovach, C., Williamson, S. and Goodman, S. M. 1999. Coherent light
scattering by nanostructured collagen arrays in the caruncles of the Malagasy asities
(Eurylaimidae: Aves). – J. Experimental Biol. 202: 3507-3522.

Pyle, P. 1997. Identification guide to North American birds: a compendium of information on
identifying, ageing, and sexing" near-passerines" and passerines in the hand. Slate Creek
Press, 1997.

Reudink, M. W., Marra, P. P., Boag, P. T. and Ratcliffe, L. M. 2009a. Plumage colouration
predicts paternity and polygyny in the American redstart. – Anim. Behav. 77: 495–501.

Reudink, M. W., Studds, C. E., Kyser, T. K., Marra, P. P. and Ratcliffe, L. M. 2009b. Plumage
brightness predicts non-breeding season territory quality in a long-distance migratory
songbird, the American redstart *Setophaga ruticilla*. – J. Avian Biol. 40: 34–41.

Reudink, M. W., Van Wilgenburg, S. L., Steele, L., Pillar, A. G., Marra, P. P. and McKellar, A.
E. 2015. Patterns of migratory connectivity in Vaux's swifts at a northern migratory roost:
A multi-isotope approach. – Condor 117: 670–682.

Robinson, S. K., and Holmes, R. T. 1982. Foraging behavior of forest birds: the relationships
among search tactics, diet, and habitat structure. – Ecology 63: 1918-1931.

Saks, L., McGraw, K. and Hõrak, P. 2003. How feather colour reflects its carotenoid content. –
Func. Ecol. 17: 555–561.

Santos, E. S. A., Scheck, D. and Nakagawa, S. 2011. Dominance and plumage traits: meta-
analysis and metaregression analysis. – Anim. Behav. 82: 3-19

Shawkey, M. D. and Hill, G. E. 2004. Feathers at a fine scale. – The Auk 121: 652-655.

Shawkey, M. D. and Hill, G. E. 2005. Carotenoids need structural colours to shine. – Biol. Lett.

1: 121-124.

Shawkey, M. D., Pillai, S. R., Hill, G. E., Siefferman, L. M. and Roberts, S. R. 2007. Bacteria as
an agent for change in structural plumage color: correlational and experimental evidence. –
Am. Nat. 169: S113-S121.

Shawkey, M. D., Pillai, S. R. and Hill, G. E. 2009. Do feather-degrading bacteria affect sexually
selected plumage color? – Naturwissenschaften 96: 123-128.

Sherry, T. W., Holmes, R. T., Pyle, P. and Patten, M. A. 2020. American Redstart (*Setophaga*
ruticilla), version 1.0. In Birds of the World (P. G. Rodewald, Editor). Cornell Lab of
Ornithology, Ithaca, NY, USA.

Siefferman L. and Hill, G. E. 2005. Male Eastern Bluebirds trade future ornamentation for
current reproductive investment. – Biol. Lett. 1: 208–211

Simons, M. J. P., Groothuis, T. G. G. and Verhulst, S. 2015. An appraisal of how the vitamin A-
redox hypothesis can maintain honesty of carotenoid-dependent signals. – Ecol. Evol. 5:
224-228.

Slagsvold, T. and Lifjeld, J. T. 1985. Variation in plumage colour of the great tit *Parus major* in
relation to habitat, season and food. – J. Zool. 206: 321–328.

Sparrow, K. L., Donkor, K. K., Flood, N. J., Marra, P. P., Pillar, A. G. and Reudink, M. W. 2017.
Conditions on the Mexican moulting grounds influence feather colour and carotenoids in
Bullock's orioles (*Icterus bullockii*). – Ecol. Evol. 7: 2643-2651

Stoddard, M. C., and Prum, R. O. 2008. Evolution of avian plumage color in a tetrahedral color
space: a phylogenetic analysis of new world buntings. – The American Naturalist, 171:
755-776.

Vorobyev, M., Osorio, D., Bennett, A. T., Marshall, N. J. and Cuthill, I. C. 1998. Tetrachromacy,
oil droplets and bird plumage colours. – J. Comp. Phys. A 183: 621–633.

Ward, G. M., Mahoney, S. M., Joly, S. and Reudink, M. W. 2021. Effects of age and weather
during moult on mountain bluebird *Sialia currucoides* structural colouration. – J. Avian
Biol. 52: https://doi.org/10.1111/jav.02616

Weaver, R. J., Santos, E. S., Tucker, A. M., Wilson, A. E. and Hill, G. E., 2018. Carotenoid
metabolism strengthens the link between feather coloration and individual quality. –
Nature Comm. 9: 1-9.

Weckstein, J. D., Kroodsma, D. E., Faucett, R. C., Poole, A. F. and Gill, F. B. 2002. Fox
Sparrow (*Passerella iliaca*), version 2.0. The Birds of North America (PG Rodewald, ed.).
Cornell Lab of Ornithology, Ithaca, New York, USA. https://doi. org/10.2173/bna, 715.

Wilkins, M. R., Seddon, N., and Safran, R. J. 2013. Evolutionary divergence in acoustic signals:
causes and consequences. – Trends Ecol. Evol. 28: 156-166.
547

Table 1. Linear mixed effects model selection results using a sample-size corrected Akaike's

Information Criterion (AICc) method. Models were built to predict the effects of American

redstart colouration (brightness, red and green chroma, hue, tetrahedral colourspace PC1) based

on site, season, age, $\delta^{15}N$, and $\delta^{13}C$. Final models used in analyses are indicated in bold. ΔAIC:

Change between full and reduced model; AICcWt: AICc weights, indicating the probability a

model is the most parsimonious model; LL: Log-likelihood.

Variable	Model	K	AICc	ΔAICc	AICcWt	LL
Brightness	**Reduced**	**6**	**-473.7**	**0.0**	**1.0**	**243.2**
	Full	11	-466.7	7.0	0.0	245.4
Red Chroma	**Reduced**	**4**	**-537.5**	**0.0**	**1.0**	**272.9**
	Full	11	-529.0	8.5	0.0	276.6
Hue	**Reduced**	**7**	**79.3**	**0.0**	**0.9**	**-32.2**
	Full	11	84.4	5.1	0.1	-30.1
Colourspace	**Full**	**11**	**-74.2**	**0.0**	**0.6**	**49.2**
PC1	Reduced	6	-73.1	1.1	0.4	42.9

Table 2. Model results demonstrating the effect of AICc selected fixed effects of site, season, sex, age, $\delta^{15}N$, and $\delta^{13}C$ on American redstart colouration (brightness, red and green chroma, hue, and tetrahedral colourspace PC1), and variances of the random effects of year. Significant effects are in bold.

Variable	Fixed effect	Model df	Residual df	*F*	*P*	Random effect σ^2
Brightness	**Site**	**1**	**134.0**	**6.45**	**0.012**	
	Season	**1**	**134.0**	**12.82**	**<0.0001**	0
	$\delta^{15}N$	**1**	**134.0**	**11.78**	**0.001**	
Red Chroma	Site	1	134.0	3.16	0.078	0
Hue	**Season**	**1**	**134.0**	**26.09**	**<0.0001**	
	Age	**1**	**134.0**	**9.43**	**0.003**	0
	Sex	**1**	**134.0**	**12.39**	**0.001**	
	Season x Age	**1**	**134.0**	**7.96**	**0.006**	
Colourspace PC1	**Site**	**1**	**133.7**	**11.20**	**0.001**	
	Season	**1**	**133.3**	**90.91**	**<0.0001**	
	Age	**1**	**132.1**	**27.07**	**<0.0001**	
	Sex	**1**	**132.0**	**10.79**	**0.001**	0.05
	$\delta^{13}C$	1	132.5	1.79	0.183	
	$\delta^{15}N$	1	131.9	0.47	0.494	
	Season x Age	1	131.9	0.42	0.520	
	Sex x Age	1	131.9	2.98	0.087	

Fig. 1. Map showing study sites in New York, US (Braddock Bay Bird Observatory) and off the coast of Maine, US (Appledore Island Migration Station). Inset photos show adult female (top) and adult male (bottom) American redstarts (photo credit: Kristen M. Covino).

Fig. 2. Left panels: American redstart feather brightness in two sampling sites (AIMS and BBBO) in Fall and Spring. Right panel: Brightness and $\delta^{15}N$ sampled from American redstart feathers. Females are indicated by red dots and males are indicated by blue dots.

Fig. 3. Female and male American redstart feather hue in adult (left panels) and young (right panels) birds during fall (top) and spring (bottom).during fall (top) and spring (bottom).

Fig. 4. American redstart colourspace PC1 derived from a principal components analysis of tetrahedral colourspace. Left panels: (Top) Female young and adult colourspace in two sites (AIMS and BBBO). (Bottom) Male young and adult colourspace in two sites (AIMS and BBBO). Right panels: young and adult colourspace in fall and spring. Females are indicated by red dots and males are indicated by blue dots.

Table S1. Sample sizes, means and SDs for brightness, red chroma, hue, colourspace (PC1 from a PCA of tetrahedral colour variables), $\delta13C$, $\delta15N$ values, for each site, season, sex, and age. Note there were no samples for adult females at the AIMs site during fall migration.

Season	Sex	Age	n	Brightness	SD Brightness	Chroma	SD Chroma	Hue	SD Hue	Colourspace	SD Colourspace	$\delta13C$	SD $\delta13C$	$\delta15N$
Fall	F	Young	16	0.28	0.06	0.3	0.04	0.74	0.42	-0.17	0.15	-23.33	0.61	6.88
	M	Adult	1	0.25	-	0.28	-	-0.01	-	-0.55	-	-23.46	-	6.32
		Young	13	0.28	0.03	0.3	0.04	0.59	0.46	-0.22	0.19	-22.94	0.61	6.72
Spring	F	Adult	5	0.26	0.02	0.31	0.05	0.82	0.34	0.09	0.04	-23.28	0.53	5.59
		Young	18	0.27	0.06	0.29	0.03	0.76	0.29	0.17	0.18	-22.85	0.71	7.15
	M	Adult	6	0.24	0.04	0.31	0.05	0.71	0.41	-0.07	0.2	-23.75	0.42	7.42
		Young	16	0.24	0.04	0.28	0.02	0.69	0.28	0.16	0.23	-22.99	1.03	7.35
Fall	F	Adult	7	0.28	0.06	0.27	0.02	0.33	0.28	-0.26	0.26	-23.86	1.35	7.23
		Young	10	0.25	0.02	0.28	0.03	0.63	0.25	-0.04	0.16	-23.36	0.65	6.62
	M	Adult	2	0.26	0.02	0.3	0.02	0.1	0.11	-0.38	0.17	-23.23	0.33	7.02
		Young	9	0.25	0.04	0.27	0.01	0.4	0.11	-0.17	0.12	-23.4	0.54	6.23
Spring	F	Adult	5	0.23	0.03	0.31	0.04	0.97	0.3	0.16	0.23	-23.43	0.72	7.36
		Young	10	0.25	0.02	0.3	0.03	0.94	0.32	0.26	0.22	-23.11	0.66	7.92
	M	Adult	7	0.22	0.03	0.28	0.01	0.51	0.17	-0.04	0.16	-23.34	0.74	5.54
		Young	9	0.24	0.04	0.28	0.01	0.68	0.15	0.18	0.11	-24.2	0.71	7.16

Table S2. Parameter estimates from linear mixed effects model selection results using a sample-size corrected Akaike's Information Criterion (AICc) method. Models were built to predict the effects of American redstart colouration (brightness, red and green chroma, hue, tetrahedral colourspace PC1) based on site, season, age, $\delta^{15}N$, and $\delta^{13}C$.

Variable	Fixed effect	Estimate	SE	df	t	P
Brightness	Intercept	0.22	0.02	134	12.11	<0.0001
	Site	-0.02	0.01	134	-2.54	0.012
	Season	-0.02	0.01	134	-3.58	<0.0001
	$\delta^{15}N$	0.01	0.00	134	3.43	0.001
Red Chroma	Intercept	0.29	0.00	134	80.61	<0.0001
	Site	-0.01	0.01	134	-1.78	0.078
Hue	Intercept	0.70	0.05	134	13.75	<0.0001
	Season	0.15	0.06	134	2.48	0.015
	Age	-0.39	0.11	134	-3.63	<0.0001
	Sex	-0.19	0.05	134	-3.52	0.001
	Season x Age	0.37	0.13	134	2.82	0.006
Colourspace PC1	Intercept	0.48	0.44	133.7	1.10	0.275
	Site	0.11	0.03	133.7	3.35	0.001
	Season	0.33	0.03	133.4	9.64	<0.0001
	Age	-0.16	0.06	132.0	-2.48	0.015
	Sex	-0.05	0.03	131.9	-1.60	0.111
	$\delta^{13}C$	0.03	0.02	132.5	1.34	0.183
	$\delta^{15}N$	-0.01	0.01	131.9	-0.69	0.494
	Season x Age	0.05	0.07	131.9	0.64	0.520
	Age x Sex	-0.12	0.07	131.9	-1.73	0.087

Table S3. Factor loadings for the first two principal components derived from tetrahedral colour models of female and male American Redstart adults and young.

Tetrahedral colourspace variable	PC1	PC2
hue Φ	-0.952	0.172
r.achieved	-0.114	-0.018
u	-0.067	0.017
u.r	-0.067	0.017
z	-0.067	0.017
Luminance	-0.060	-0.031
x	-0.024	-0.050
r.vector	0.001	-0.029
l	0.001	-0.072
l.r	0.001	-0.072
y	0.002	0.055
m.r	0.024	0.046
m	0.024	0.046
s.r	0.041	0.009
s	0.041	0.009
r.max	0.160	-0.100
hue θ	0.181	0.968
Variation	81%	11%
Eigenvalue	0.06	0.008