A tail of plumage colouration: disentangling geographic, seasonal, and dietary effects on

plumage colour in a migratory songbird

Sean M. Mahoney[1*], Matthew W. Reudink[1], Andrea Contina[2], Kelly A. Roberts[3], Veronica T.

Schabert[3], Emily G. Gunther[3], Kristen M. Covino[4]

[1]M.W. Reudink and S.M. Mahoney, Department of Biological Sciences, Thompson Rivers
University, Kamloops, BC, Canada. — [2]A. Contina, Department of Integrative Biology,
University of Colorado Denver, Denver, CO, USA. — [3] K.A Roberts, V.T. Schabert, and E.G.
Gunther, Department of Animal Behavior, Ecology, and Conservation, Canisius College,
Buffalo, NY, USA — [4]K.M. Covino (Kristen.covino@lmu.edu) Biology Department, Loyola
Marymount University, Los Angeles, CA,USA.

*Corresponding author: smahoney@tru.ca
Short Title: American redstart colour variation

Abstract

Plumage ornamentation in birds serves critical inter- and intra-sexual signaling functions. While

carotenoid-based plumage colouration is often viewed as a classic condition-dependent sexually

selected trait, plumage colouration can be influenced by a wide array of both intrinsic and

extrinsic factors. Understanding the mechanisms underlying variation in colouration is especially

important for species where the signaling function of ornamental traits is complex or the

literature conflicting. Here, we examined variation in the yellow/orange tail feathers of American

redstarts (*Setophaga ruticilla*) passing through two migratory stopover sites in eastern North

America during both spring and fall migration to assess the role of geographic variation and

seasonality in influencing feather colouration. In addition, we investigated whether diet during

moult (inferred via stable isotope analysis of feather $\delta^{15}N$ and $\delta^{13}C$) influenced plumage

colouration. Our findings indicate that geographic variation, season, and diet all influence

individual differences in American redstart colouration, represented by both traditional and

tetrahedral colour variables. The extent to which these factors influence colour expression

however is largely dependent on the colour metric under study, likely because different colour

metrics reflect different attributes of the feather (e.g., structural components vs. pigment

deposition). The effects of diet ($\delta^{15}N$) and season were pronounced for brightness, suggesting a

strong effect of diet and feather wear/degradation on feather structure. Though hue, a metric that

should strongly reflect pigment deposition, also changed from spring to fall, that effect was

dependent on age, with only adults experiencing a reduction in ornamentation. Taken together,

our results highlight the numerous sources of variation behind plumage coloration and

underscores the difficulty of unraveling complex visual signaling systems, such as those in

American redstarts.

Keywords: carotenoid, plumage, tetrahedral colourspace, stable isotope, nitrogen, carbon

Introduction

The colourful plumage exhibited by birds serves important roles for both inter-and intrasexual communication (reviewed in Hill 2006; Santos et al. 2011). The literature linking variation in plumage colouration to individual quality and reproduction is multitudinous (Lope and Møller 1993; Griffith and Pryke 2006; Hill 2006; Henschen et al. 2017; Ferree 2019; LaBarbera et al. 2019; Enbody et al. 2018), yet the patterns are often complex and may vary across seasons and populations (Marini et al. 2015; Reudink et al. 2015). As such, much research has been dedicated to understanding the mechanisms underlying variation in the expression of ornamental plumage colouration (Siefferman and Hill 2005; Griggio et al. 2009; Newton and Dawson 2011; Doutrelant et al. 2012; Reudink et al. 2015).

Plumage colouration in birds can be produced by various mechanisms, including the deposition of pigments (e.g., carotenoids, melanins, and porphyrines) and the arrangement of melanosomes within the feather microstructure producing structural (purple, blue, green, and iridescent) colours. Carotenoid colouration (yellow, orange, red plumage) has received by far the most attention, in large part because the expression of carotenoid-based colouration is directly dependent on birds acquiring carotenoids in the diet (the pigments cannot be synthesized *de novo*) and subsequently modifying dietary pigments through a costly physiological process (Lozano et al. 1994; Hill and Montgomerie 1994; Weaver et al. 2018). Thus, carotenoid-based colouration is often viewed as a classic honest indicator of individual condition and quality (Hill 2011; Hill and Johnson 2012; Weaver et al. 2018; but see Simons et al. 2015).

Diet during feather growth has been directly linked to ornamentation and feather quality

in both experimental (Hill 2000; Peters et al. 2008; Koch et al. 2017) and observational studies

(Slagsvold and Lifjeld 1985; Arriero and Fargallo 2006; Eeva et al. 2009; but see Ferns and

Hinsley 2008). For observational studies of free-living birds, one of the challenges is

determining dietary conditions during moult (Besozzi et al. 2021). However, stable isotopes of

carbon (δ^{13}C) and nitrogen (δ^{15}N) vary predictably in the environment and are incorporated into

growing tissues (Fry 2006; Bowen 2010). Because feathers are inert once grown, they can

provide insight into diet during tissue growth. δ^{13}C has been used extensively in dietary studies

because carbon isotope ratios vary based on plant water stress and photosynthetic system (Lajtha

and Marshall 1994). Further, since these isotope elements are transferred up the food chain and

eventually incorporated into animal tissues, they can provide important insight into habitat and

diet during tissue growth (Hobson 1999). Nitrogen is commonly employed in food web studies,

as ^{15}N is preferentially incorporated into animal tissues as one moves up the food chain (Post

2002; Poupin et al. 2011). Thus, analysis of δ^{15}N tissue values can provide insight into the

trophic level of animals within the environment. However, δ^{15}N values can also be influenced by

environmental conditions, as they tend to be positively associated with temperature and

negatively associated with rainfall (Craine et al. 2009).

A recent study on Bullock's orioles (*Icterus bullockii*) demonstrated that feather δ^{15}N

values were directly associated with feather carotenoid content and hue (Sparrow et al. 2017). In

the case of Bullock's orioles, birds with lower feather δ^{15}N values expressed more orange-shifted

hue. Because Bullock's orioles rely on both insects and fruit during moult, the authors suggested

that individuals with a relatively higher amount of fruit in their diet (or those moulting in habitats

that resulted in lower δ^{15}N values) may have been in better condition and better able to

metabolically convert dietary carotenoids to red or orange keto-carotenoids (McGraw 2006;

Sparrow et al. 2017; Weaver et al. 2018).

Though feathers are metabolically inert once grown, feather colours are not necessarily

static across the life of the feather. Feather wear can modify feather colouration; for example,

European starlings (*Sturnus vulgaris*) lose their distinctive winter spots through wear and

abrasion on the feather tips leaving only the melanin-based feather structure behind for the

breeding season. For most species, these changes are less dramatic but can still have important

fitness consequences. Feather abrasion, ultraviolet (UV) radiation, and feather degrading bacteria

can all lead to changes in ornamentation after initial feather growth (Örnberg et al. 2002;

Shawkey and Hill 2004). For example, carotenoid-based red colouration in House finches

(*Haemorhous mexicanus*) becomes increasingly more saturated and hue becomes more yellow-

shifted in the spring following moult (McGraw and Hill 2004). Melanin-based plumage in Fox

sparrows (*Passerella iliaca*) also changes colour over time (Weckstein et al. 2002), possibly due

to UV exposure or accumulation of dirt and oils on the feathers (Montgomerie 2006).

Additionally, colour in many species changes with age — sometimes dramatically through

delayed plumage maturation (Hawkins et a. 2012) and sometimes more subtly as individuals age

(Marini et al. 2015; Ward et al. 2021); therefore, it is important for studies to assess the

environmental, biological, ontological, and temporal influences on feather colour to help unravel

the mechanisms regulating this phenotypic trait.

For the American redstart (*Setophaga ruticilla*), the brilliant orange and black plumage

ornamentation expressed by males in definitive plumage has been extensively studied (e.g.,

Reudink et al. 2009a,b; Marini et al. 2015; Reudink et al. 2015), yet the role of plumage for

inter- and intra-sexual signaling remains frustratingly complicated and unresolved (see Marini et

al. 2015; Table 1). As such, recent research on this species has focused on understanding the

mechanisms underlying variation in plumage colouration (Marini et al. 2015; Reudink et al.

2015). In an 11-year study, Reudink et al. (2015) found that rainfall and temperature during

moult was directly linked variation in feather hue; however, individuals themselves also appear

to change throughout life, with males expressing the most orange-shifted hue during their first

breeding season in definitive plumage (Marini et al. 2015). At a broader scale, American

redstarts also exhibit marked geographic variation in plumage ornamentation across the range

(Norris et al. 2007). Adding to the complexity of this system, American redstarts exhibit delayed

plumage maturation, with young (hatch-year [HY] and second-year [SY]) males exhibiting

female-like grey and yellow plumage throughout their first year of life, while adult (after hatch

year [AHY] and after second year [ASY]) birds display jet black and orange plumage. As such,

male redstarts of different age classes appear to utilize different metabolic pathways prior to

depositing ingested dietary carotenoids. Adding to the complexity, these potentially important

signals are grown at different times of year and under different conditions — in the nest in early-

to-mid summer for young birds and after breeding in late summer for adult birds. Thus,

American redstarts in different age- and sex-classes are likely differentially affected by both

intrinsic and extrinsic factors.

Our goal was to examine the relative importance of the several factors that may underlie

variation in the expression of ornamental colouration in the American redstart. We investigated

plumage colouration of American redstarts passing through two different migration stopover

sites in eastern North America (Fig. 1, Braddock Bay Bird Observatory (BBBO) and Appledore

Island Migration Station (AIMS)) during both spring and fall migration and examined the stable

isotope ratios of δ^{13}C and δ^{15}N in the feathers of each individual. As such, we were able to

concurrently determine the relative effects of geographic (site), temporal (season), biological

(sex), ontological (age), environmental and dietary variation (δ^{13}C and δ^{15}N), along with age and

sex, in the expression of plumage ornamentation. We predicted there would be differences in

plumage colouration between sites and that birds captured in fall would exhibit a greater degree

of ornamentation (increased red chroma, orange-shifted hue, higher brightness) compared to

those captured in spring. In addition, we predicted that birds with more negative δ^{13}C feather

isotope values (indicating habitats with lower water stress) and more positive δ^{15}N feather

isotope values (indicating feeding at a higher trophic level) would exhibit increased

ornamentation.

Methods

Field Methods

We collected feather samples from American redstarts, a sexually dimorphic Nearctic-

Neotropical migrant that breeds in deciduous forests in the United States and Canada (Sherry et

al. 2020). Our collection sites were two locations of similar latitude but different proximity to the

Atlantic coast: Braddock Bay Bird Observatory (43° 9'N, 77° 36'W) in New York State and

Appledore Island Migration Station (42°58′N, 70°36′ W) off the coast of Maine. Braddock Bay

Bird Observatory (BBBO) is an inland site just south of Lake Ontario and the Appledore Island

Migration Station (AIMS) is located seven miles offshore from the Maine-New Hampshire

border (Fig. 1).

Field researchers captured birds at AIMS using mist nets (weather permitting) checked at

least every 20 minutes from sunrise to sunset from 11 May to 7 June during spring migration and

14 August to 15 September during fall migration 2016 and 2017. During 2016 and 2017 at

BBBO field researchers captured birds with mist nets checked every half hour starting at sunrise

and for 6 hours thereafter from 14 April through 8 June (spring) and 12 August through 2

November (fall). A United States Geological Service leg band was placed on each bird and sex

and age were determined according to Pyle (1997) based on plumage, molt, and/or degree of

skull pneumatization. In fall, age was determined to be hatch-year (hereafter young) or after-

hatch-year (hereafter adult). In spring, age was determined to be second-year (young) or after-

second-year (adult). We sampled the left and right fifth rectrix feathers (second from the outside

on each side of the tail), provided the feathers were not replaced outside of regular molt, in

which case we did not sample feathers and that individual was excluded from this study. In total,

we collected feathers from 134 individual American redstarts; 59 from BBBO and 75 from

AIMS (Table S1).

Sample preparation and stable isotope analysis

We conducted analysis of stable isotope ratios of carbon ($^{13}C/^{12}C$) and nitrogen ($^{15}N/^{14}N$)

on keratin tissues obtained from feathers plucked during mist-netting efforts at AIMS and

BBBO. We cleaned each feather to remove oils and debris with immersion in a 2:1 chloroform-

methanol solution followed by air-drying for 24-h under a fume hood. We then washed each

feather with a 1:30 solution of detergent (Fisher Scientific Versa-Clean #04-0342) and deionized

water before rinsing three times with deionized water (Paritte and Kelly 2009; Chew et al. 2019).

We cut a 350 µg (± 10 µg) fragment and packed in tin capsules placed in a Carlo Erba NC2500

elemental analyzer (CE Instruments, Milano, Italy), interfaced with a Thermo Delta V+ isotope

ratio mass spectrometer (Thermo-Finnigan, Bremen, Germany) operated by the Central

Appalachians Stable Isotope Facility (CASIF) at the Appalachian Laboratory (Frostburg,

Maryland, USA). We matched each sample with powdered keratin (porcine) from Spectrum

Chemicals, a quality control (QC) in-house standard. We report feather carbon (δ^{13}C) and

nitrogen (δ^{15}N) as mean ± SD in delta notation of parts per thousand (‰) from the standards

($\delta D_{sample} = [(R_{sample}/R_{standard}) - 1] \times 1{,}000$) compared to the Vienna PeeDee Belemnite (V-PDB)

and AIR scales, respectively. The standard deviation of the reference material was ±0.12‰

(δ^{13}C) and ±0.11‰ (δ^{15}N).

Colour Variation

Prior to cutting and packaging feathers (above) but after the above-described cleaning

steps, feather colouration was determined by measuring the reflectance within the avian visual

range (300-700nm) with an Ocean Optics Jaz Spectrometer (Reudink et al. 2015). The yellow-

orange vane sections of each rectrix were randomly scanned ten times each. Upon each use, the

spectrometer was calibrated by scanning an Ocean Optics WS-1 white standard and a Colorline

Ebony #142 dark standard. Additionally, the spectrometer was also calibrated between each

feather using the Ocean Optics WS-1 white standard.

We quantified plumage colouration using traditional colourimetric values (e.g.

Montgomerie 2006) and using tetrahedral models which correct for the avian visual system (e.g.

Stoddard and Prum 2008; Fig. 2). Because nearly all studies on American redstarts use

traditional colourimetric variables, we chose to include them in our study to allow comparison

with the previous literature. To generate traditional color metrics of brightness, red chroma, and

hue from Montogmerie (2006), we utilized the R-based program RCLR v.28. Reflectance spectra

were imported into RCLR v.28 where we first performed smoothing using a LOESS function to

eliminate noise and local peaks in the curves. Next, we calculated color variables of brightness,

red chroma, and hue from the smoothed curves following Montgmerie (2006). Brightness (mean

$R_{300-700}$) was calculated as the mean amount of light reflected from 300–700 nm. Red chroma

($R_{605-700}$/ $R_{300-700}$) was measured as the proportion of light reflected in the orange-red segment of

the spectrum relative to the overall light reflected. Hue (arctan ($[(R_{415-510}-R_{320-415})/R_{320-700}]$/

$[(R_{605-700}-R_{400-500})/R_{320-700}]$)) was measured via segment classification to account for the

proportion of light reflected in different regions of the spectrum (Saks et al. 2003). High hue

values reflect dominance of shorter wavelengths (yellow-shifted), while low hue represent a

dominance of longer wavelengths (red-shifted). Next, we averaged the colour variables from the

ten reflectance spectra generated from each individual to obtain a single value of brightness, red

chroma, and hue for each individual. These techniques follow those used in previous work by

our group on American redstarts (Reudink et al. 2009a; Osmond et al. 2013; Tonra et al. 2014).

We also quantified colour using tetrahedral models which correct for avian visual

systems (Vorobyev et al. 1998), in the R package *pavo2* (Maia et al. 2019). Although the visual

system of American redstarts is unknown, most passerines are ultraviolet sensitive (Ödeen and

Håstad 2003), so our tetrahedral models use this visual system. We used a receptor-noise

discrimination model to calculate the photon catch of each cone type used in avian colour vision.

We used a value of 0.1 for colour (chromatic) and luminance (achromatic) discrimination and the

photoreceptor ratios to 1:2:2:4 (ultraviolet, shortwave, mediumwave, longwave) in our models

(Maier and Bowmaker 1993), which were selected based on a recent review of noise receptor

estimates (Olsson et al. 2018). The models calculate hue θ, hue Φ, r.achieved, and luminance.

Hue θ represents the angle of the color vector in the longitudinal plane while hue Φ represents

the angle in the latitudinal plane and together hue θ and Φ describe the hue or the direction of

colour vector in tetrahedral space (Stoddard and Prum 2008). r. achieved represents the chroma

or saturation, or how different the colour is from the achromatic center of the tetrahedron

(Stoddard and Prum 2008). Finally, luminance is analogous to brightness and represents the

overall amount of light reflected from the feather sample and calculated independent of colour.

Although this may seem redundant to include luminance, because all other previous redstart

papers include traditional brightness, we thought it important to include both brightness and

luminance in our study.

Statistical Analyses

We tested how American redstart colour variation was explained by site, season, sex, age,

and isotopic signatures at sampling using separate mixed models in the *lme4* package in R. In our

models, we used traditional colourimetric variables (brightness, red chroma, and red hue) and

colour modelled in tetrahedral space (hue θ, hue Φ, r.achieved, and luminance). In separate

models, each colour variable was included as the response variable. Site, season, sex, age, $\delta^{13}C$

and $\delta^{15}N$ were main effects, and the interaction between age and season was included because

redstarts exhibit delayed colour maturation (Hawkins et al. 2012). Year was included as a

random effect. We then undertook model reduction for all mixed models based on the change in

Akaike information criterion corrected for small sample sizes (AICc, Burnham and Anderson

2003) between the full model and each reduced model. We chose our final model based on the

lowest AIC (Burnham and Anderson 2003). ΔAIC values within 4 were considered competitive

and we selected the best fitting model using ANOVAs.

Results

Colour Variation

Our AICc model selection identified the reduced models for all response variables as the top models (Table 2 and Table S2). American redstart brightness was explained by site, season, and δ^{15}N (Table 2). In both the spring and fall, American redstart feathers were brighter at the AIMS site (Fig. 3, site: $F_{1,134}=6.45$, P=0.012; Season: $F_{1,134}=12.82$, P<0.0001). Brightness was positively related to δ^{15}N (Fig. 3. $F_{1,134}=11.78$, P=0.001). Red chroma did not vary among any of our predictor variables (Table 2).

Hue was explained by season, sex, age, and the interaction between season x age (Table 2 and Table S2). Hue was more red-shifted in males (Fig. 4, $F_{1,134}=12.39$, P=0.001) and in adults (Fig. 4, $F_{1,134}=9.43$, P=0.003) and following moult in the fall (Fig. 4, $F_{1,134}=26.09$, P<0.0001). There was an interaction between season and age indicating that although hue in young birds did not vary between seasons, adult hue was more red-shifted in the fall (Fig. 4, $F_{1,134}=7.96$, P=0.006).

Our analyses found redstart feather reflectance modelled in tetrahedral colourspace varied by several of our predictor variables (Table 2 and Table S2). Hue θ (Fig. 5) was predicted by site ($F_{1,133.9}=5.93$, P=0.02) and the interactions between season x age ($F_{3,133.9}=30.11$, P<0.0001) and sex x age ($F_{2,133.9}=49.50$, P<0.0001). Hue Φ (Fig. 5) was predicted by site ($F_{1,133.8}=9.86$, P=0.002), season ($F_{1,133.6}=129.14$, P<0.0001), age ($F_{1,132.0}=24.12$, P<0.0001), and sex ($F_{1,132.0}=5.89$, P=0.02). Season ($F_{1,134.0}=33.69$, P<0.0001), age ($F_{1,131.9}=6.89$, P=0.01), and δ^{15}N

260   (Fig. 6, $F_{1,131.9}$=6.86, P=0.01) were significant predictors of r.achieved. Finally, luminance was

261   predicted by δ^{15}N (Fig. 6, $F_{1,134.0}$=5.59, P=0.02).

Discussion

Carotenoid-based colouration serves important inter- and intra-sexual signaling functions;

however, for some species like the American redstart, the relationship between fitness-related

components and colour metrics can be perplexing and seemingly contradictory (Reudink et al.

2015). Part of the complexity likely stems from our incomplete understanding of the mechanisms

underlying variation in American redstart colouration. Here, we demonstrate that geographic

variation, season, and diet all underly intra-individual differences in American redstart

colouration. However, the extent to which these factors influence colour expression is largely

dependent on the colour metric under study, likely because different colour metrics reflect

different aspects of feather quality (e.g., structural components vs. pigment deposition).

Geographic variation in plumage ornamentation has been widely documented across

numerous species (Hill 2006; Badyaev et al. 2012) including the American redstart (Norris et al.

2007). Norris et al. (2007) found that individuals moulting at higher latitudes expressed plumage

with higher red chroma compared to those at lower latitudes. In our study, we found differences

in colouration between our two study sites (separated by 580km at similar latitudes) in three

variables: brightness, hue θ, and hue Φ. All birds in our study were captured during migration

and, as such, we lack information in the specific moult origins for each individual. Norris et al.

(2007) suggested that geographic variation in American redstart colouration was likely driven by

differences in dietary carotenoid availability or condition-related effects (e.g., parasite load)

mediating carotenoid metabolism. Hue and chroma variables should be most closely linked to

carotenoid content (Saks et al. 2003; Montgomerie 2006) and while brightness is negatively

associated with feather carotenoid content, it is strongly mediated by feather structure (Shawkey

and Hill 2005). Thus, while the site-specific differences we observed may be due to differences

in carotenoid availability at the moult location, these differences may also reflect longitudinal

geographic variation in condition, as one site was inland while the other was more coastal, but

both were of similar latitude.

As expected, American redstarts expressed differences in brightness, hue θ, hue Φ, and

r.achieved with higher brightness, and more red-shifted hue during the fall, following molt.

However, variation in hue was dependent on age; while adults were more red-shifted in fall than

spring, we detected no differences in first-year birds, potentially indicating that the orange

colouration expressed by adult males may be more susceptible to the effects of UV degradation,

feather-degrading bacteria, or feather wear between seasons (Örnborg et al. 2002; Delhey et al.

2006; Shawkey et al. 2007; Shawkey et al. 2009). Alternatively, if birds passing through

migratory stopover sites in spring and fall differ in their moult location, differences between

spring and fall plumage could simply reflect geographic variation in moult location.

Our brightness metrics (brightness and luminance) were associated with $\delta^{15}N$; birds with

higher feather $\delta^{15}N$ values had feathers with higher brightness. Because $\delta^{15}N$ is directly linked to

diet (Pearson et al. 2003; Gómez et al. 2018), these results suggest that American redstarts eating

arthropods enriched with ^{15}N (e.g., *Heteromurus nitidus*, *Drosophilia melanogastor;*

Oelbermann and Scheu 2002) likely are not acquiring additional dietary carotenoids; however,

these dietary sources may be advantageous during feather growth, allowing the birds to create

feathers with a high-quality microstructure. Whether or not feather brightness and luminance *per*

se is a signal used by American redstarts is unclear; however, Reudink et al. (2009b)

demonstrated that males that occupied higher-quality winter-territories expressed feathers with

higher brightness and Reudink et al. (2009a) found that polygynous males had higher feather

brightness than monogamous males. Because polygyny requires the ability to hold multiple

territories or a single large territory, Reudink et al. (2009a) suggested that feather brightness may

act as a social signal during both the breeding and wintering season, directly reflecting a male's

competitive ability. Thus, diet during moult may potentially impact intrasexual signaling.

In a study of Bullock's orioles, Sparrow et al. (2017) found that feather $\delta^{15}N$ values were

associated with feather colouration and feather carotenoid content. The authors found that while

$\delta^{15}N$ values were positively associated with overall carotenoid concentration, birds with more

orange-shifted hue had feathers with lower $\delta^{15}N$ values. The authors suggested that low $\delta^{15}N$

values could indicate a diet with a higher fruit:insect ratio and birds with more orange-shifted

hue, though they had lower overall carotenoid concentrations, may have had a higher abundance

of specific orange or red carotenoids. It is important to note that American redstarts and

Bullock's oriole diets differ considerably; orioles have a more varied diet of both fruits and

insects (Flood et al. 2016), while redstarts are primarily insectivores (Robinson and Holmes

1982). Thus, the way in which $\delta^{15}N$ is associated with the expression of carotenoid-based

colouration is likely highly dependent on the dietary ecology of the species. It is also important

to consider that our observed differences may be due to accumulation of dirt in feathers or due to

degradation caused by UV exposure (Örnborg et al. 2002). The decrease in brightness and hue

between the fall and spring is consistent with previous studies (Prum et al. 1999). Interestingly,

while our tetrahedral metric for chroma (r.achieved) did vary in our study, our traditional

colourimetric variable (red chroma) did not significantly vary, as was previously found in

carotenoid-red colouration as a result of abrasion or soiling (McGraw and Hill 2004). However,

without estimates of dirt on feathers, microscopy of feather nanostructure, or repeated measures,

we can only speculate on this effect and encourage future work to consider adventitious

colouration.

Literature Cited

Arriero, E., and Fargallo, J. A. 2006. Habitat structure is associated with the expression of
carotenoid-based coloration in nestling blue tits *Parus caeruleus*. – Naturwissenschaften
93: 173–180.

Badyaev, A. V., Belloni, V., and Hill, G. E. 2012. House Finch (*Haemorhous mexicanus*. – In
Rodewald, P. G. (eds), The Birds of North America. Cornell Lab of Ornithology. Retrieved
from: https://birdsna.org/ Species-Account/bna/species/houfin. http://dx.doi.org/10.2173/
bna.46.

Besozzi, E., Chew, B., Allen, D.C. and Contina, A., 2021. Stable isotope analysis of an aberrant
Painted Bunting (Passerina ciris) feather suggests post-molt movements -- The Wilson
Journal of Ornithology. DOI: 10.1676/20-00140

Bowen, G.J., 2010. Isoscapes: spatial pattern in isotopic biogeochemistry. Annu. Rev. Earth
Planet. Sci. – 38: 161-187.

Burnham, K. P., and Anderson, D. R. 2002. Model selection and multimodel inference: A
practical information-theoretic approach. New York (USA).

Chew, B., Kelly, J., and Contina, A. 2019. Stable isotopes in avian research: a step by step
protocol to feather sample preparation for stable isotope analysis of carbon (δ^{13}C), nitrogen
(δ^{15}N), and hydrogen (δ^2H). Version 1.1. protocolsio.
dx.doi.org/10.17504/protocols.io.z2uf8ew.

Craine, J. M., Elmore, A. J., Aidar, M. P. M., Bustamante, M., Dawson, T. E., Hobbie, E. A., and
Wright, I. J. 2009. Global patterns of foliar nitrogen isotopes and their relationships with
climate, mycorrhizal fungi, foliar nutrient concentrations, and nitrogen availability. – New
Phytologist 183: 980–992.

Delhey, K., Peters, A., Johnsen, A., and Kempenaers, B. 2006. Seasonal changes in blue tit
crown color: do they signal individual quality? – Behav. Ecol. 17: 790-798.

Doutrelant, C., Gregoire, A., Midamegbe, A., Lambrechts, M. M., and Perret, P. 2012. Female
plumage colouration is sensitive to the cost of reproduction. An experiment in blue tits. – J.
Anim Ecol. 81: 87–96.

Eeva, T., Sillanpaa, S., and Salminen, J. P. 2009. The effects of diet quality and quantity on
plumage colour and growth of great tit *Parus major* nestlings: A food manipulation
experiment along a pollution gradient. – J. Avian Biol. 40: 491–499.

Enbody, E. D., Boersma, J., Schwabl, H., and Karubian, J. 2018. Female ornamentation is
associated with elevated aggression and testosterone in a tropical songbird. Behav. Ecol. -
- 29: 1056-1066.

Ferree, E. D. 2019. An experimental test of the effect of male attractiveness on maternal investment in Dark-eyed Juncos. Auk: Ornithol. Adv. -- 136: 1-12.

Ferns, P. N., and Hinsley, S. A. 2008. Carotenoid plumage hue and chroma signal different aspects of individual and habitat quality in tits. – IBIS 150: 152–159.

Flood, N. J., Schlueter, C. L., Reudink, M. W., Pyle, P. Patten, M. A., Rising, J. D., and Williams, P. L. 2016. Bullock's Oriole (*Icterus bullockii*), version 3.0. In The Birds of North America (P. G. Rodewald, editor). Cornell Lab of Ornithology, Ithaca, New York, USA.

Fry, B., 2006. Stable isotope ecology. Vol. 521. New York: Springer 318 p.

Gómez, C., Larsen, T., Popp, B., Hobson, K. A. and Cadena, C. D., 2018. Assessing seasonal changes in animal diets with stable-isotope analysis of amino acids: a migratory boreal songbird switches diet over its annual cycle. – Oecologia, 187: 1-13.

Griffith, S. C., and Pryke, S. R., 2006. Benefits to females of assessing color displays. – In: Hill, G. E., McGraw, K. J. (eds) Bird coloration, vol 2. Function and evolution. Harvard University Press, Cambridge, 233-279.

Griggio, M., Serra, L., Licheri, D., Campomori, C., and Pilastro, A. 2009. Moult speed affects structural feather ornaments in the blue tit. – J. Evol. Biol. 22:782–792.

Hawkins, G. L., Hill, G. E., and Mercadante, A. 2012. Delayed plumage maturation and delayed reproductive investment in birds. – Biol. Rev. 87: 257–274

Henschen, A. E., Whittingham, L. A., and Dunn, P. O. 2017. The relationship between blood parasites and ornamentation depends on the level of analysis in the common yellowthroat. J. Avian Biol. -- 48: 1263-1272.

Hill, G. E. 1993. Geographic variation in the carotenoid plumage pigmentation of male house finches (*Carpodacus mexicanus*). – Biol. J. Linn. Soc. 49: 63-86.

Hill, G. E. 2000. Energetic constraints on expression of carotenoid-based plumage coloration. – – J. Avian Biol. 31: 559-566.

Hill, G. E. 2006. Female mate choice for ornamental coloration. In: Hill, G. E., McGraw, K. J. (eds) Bird coloration, Vol. 2. Function and evolution. Harvard University Press, Cambridge, pp. 137–200.

Hill, G. E., and Johnson, J. D. 2012 The vitamin A-redox hypothesis: a biochemical basis for honest signaling via carotenoid pigmentation. – Am. Nat. 180: E127–E150.

419 Hobson, K. A., 1999. Stable-carbon and nitrogen isotope ratios of songbird feathers grown in
420   two terrestrial biomes: implications for evaluating trophic relationships and breeding
421   origins. – The Condor 101: 799-805.
422
423 Koch, R. E., Josefson, C. C., and Hill, G. E. 2017. Mitochondrial function, ornamentation, and
424   immunocompetence. – Biol. Rev. 92: 1459– 1474.
425
426 Lajtha, K., and Marshall, J. D. 1994. Sources of variation in isotopic composition in plants. – In:
427   Lajtha, K., and Marshall, J. D. (eds), Stable isotopes in ecological and environmental
428   sciences. Oxford University Press. pp. 1–21.
429

430 LaBarbera, K., Hayes, K. R., Langhans, K. E., and Lacey, E. A. 2019. Elevation affects extra-
431   pair paternity but not a sexually selected plumage trait in dark-eyed juncos. Behav. Ecol.
432   Sociobiol. -- 73: 82.
433

434 Lope, F. and Møller, A. P. 1993. Female reproductive effort depends on the degree of
435   ornamentation of their mates. – Evolution. 4: 1152-1160.
436

437 Lozano, G. A. 1994. Carotenoids, parasites, and sexual selection. – Oikos 70: 309–311.
438
439 Marini, K. L., McKellar, A. E., Ratcliffe, L. M., Marra, P. P., and Reudink, M. W. 2015. Age-
440   related change in carotenoid-based plumage of the American Redstart (*Setophaga*
441   *ruticilla*). – J. Ornithol. 156: 783-793.
442
443 McGraw, K. J., and Hill, G. E. 2004. Plumage color as a dynamic trait: carotenoid pigmentation
444   of male house finches (*Carpodacus mexicanus*) fades during the breeding season. – Can. J.
445   Zool. 82: 734-738.
446
447 Maier, E. J., and Bowmaker, J. K. 1993. Colour vision in the passeriform bird, *Leiothrix lutea*:
448   correlation of visual pigment absorbance and oil droplet transmission with spectral
449   sensitivity. – J. Comp. Phys. A 172: 295–301.
450
451 McGraw, K. J. 2006. The mechanics of carotenoid coloration in birds. – In: Hill, G.E. and &
452   McGraw, K. J. (eds), Bird coloration, Vol. 1: mechanisms and measurements. Harvard
453   University Press. 177–242.
454
455 Montgomerie, R. 2006. Cosmetic and adventitious colors. – In: Hill, G.E. and & McGraw, K. J.
456   (eds), Bird coloration, Vol. 1: mechanisms and measurements. Harvard University Press.
457   399-427.
458
459 Newton, I., and Dawson, A. 2011. Seasonal changes in moult, body mass and reproductive
460   condition in siskin *Carduelis spinus* exposed to daylength regimes simulating different
461   latitudes. – J. Avian Biol. 42: 22–28.
462

Ödeen, A., and Håstad, O. 2003. Complex distribution of avian color vision systems revealed by sequencing the SWS1 opsin from total DNA. – Molec. Biol. and Evol. 20: 855–861.

Oelbermann, K., and Scheu, S. 2002. Stable isotope enrichment (δ^{15}N and δ^{13}C) in a generalist predator (*Pardosa lugubris*, Araneae: Lycosidae): effects of prey quality. – Oecologia, 130: 337-344.

Olsson, P., Lind, O. and Kelber, A. 2018. Chromatic and achro- matic vision: parameter choice and limitations for reliable model predictions. – Behav. Ecol. 29: 273–282.

Örnborg, J., Andersson, S., Griffith, S. C. and Sheldon, B. C. 2002. Seasonal changes in a ultraviolet structural colour signal in blue tits, *Parus caeruleus*. – Biol. J. Linnean Soc. 76: 237-245.

Norris, D. R., Marra, P. P., Kyser, T. K., Ratcliffe, L. M. and Montgomerie, R. 2007. Continent-wide variation in feather colour of a migratory song- bird in relation to body condition and moulting locality. – Biol. Lett. 3: 16–19.

Paritte, J. M. and Kelly, J. F. 2009. Effect of cleaning regime on stable-isotope ratios of feathers in Japanese Quail (*Coturnix japonica*). – The Auk 126:165–174.

Pearson, S. F., Levey, D. J., Greenberg, C. H. and Del Rio, C. M. 2003. Effects of elemental composition on the incorporation of dietary nitrogen and carbon isotopic signatures in an omnivorous songbird. – Oecologia 135: 516-523.

Peters, A., Delhey, K., Andersson, S., Van Noordwijk, H., and Förschler, M. I. 2008. Condition□dependence of multiple carotenoid□based plumage traits: an experimental study. – Functional Ecology 22: 831-839.

Post, D. M. 2002. Using stable isotopes to estimate trophic position: Models, methods, and assumptions. – Ecology 83: 703–718.

Poupin, N., Bos, C., Mariotti, F., Huneau, J.-F., Tomé, D. and Fouillet, H. 2011. The nature of the dietary protein impacts the tissue-to-diet ^{15}N discrimination factors in laboratory rats. – PLoS ONE 6: e28046.

Prum, R. O. 2010. The Lande–Kirkpatrick mechanism is the null model of evolution by intersexual selection: implications for meaning, honesty, and design in intersexual signals. – Evolution 64: 3085–3100

Prum, R. O., Torres, R, Kovach, C., Williamson, S. and Goodman, S. M. 1999. Coherent light scattering by nanostructured collagen arrays in the caruncles of the Malagasy asities (Eurylaimidae: Aves). – J. Experimental Biol. 202: 3507-3522.

Pyle, P. 1997. Identification guide to North American birds: a compendium of information on identifying, ageing, and sexing" near-passerines" and passerines in the hand. Slate Creek

Press, 1997.

Reudink, M. W., Marra, P. P., Boag, P. T. and Ratcliffe, L. M. 2009a. Plumage colouration
predicts paternity and polygyny in the American redstart. – Anim. Behav. 77: 495–501.

Reudink, M. W., Studds, C. E., Kyser, T. K., Marra, P. P. and Ratcliffe, L. M. 2009b. Plumage
brightness predicts non-breeding season territory quality in a long-distance migratory
songbird, the American redstart *Setophaga ruticilla*. – J. Avian Biol. 40: 34–41.

Reudink, M. W., Van Wilgenburg, S. L., Steele, L., Pillar, A. G., Marra, P. P. and McKellar, A.
E. 2015. Patterns of migratory connectivity in Vaux's swifts at a northern migratory roost:
A multi-isotope approach. – Condor 117: 670–682.

Robinson, S. K., and Holmes, R. T. 1982. Foraging behavior of forest birds: the relationships
among search tactics, diet, and habitat structure. – Ecology 63: 1918-1931.

Saks, L., McGraw, K. and Hõrak, P. 2003. How feather colour reflects its carotenoid content. –
Func. Ecol. 17: 555–561.

Santos, E. S. A., Scheck, D. and Nakagawa, S. 2011. Dominance and plumage traits: meta-
analysis and metaregression analysis. – Anim. Behav. 82: 3-19

Shawkey, M. D. and Hill, G. E. 2004. Feathers at a fine scale. – The Auk 121: 652-655.

Shawkey, M. D. and Hill, G. E. 2005. Carotenoids need structural colours to shine. – Biol. Lett.
1: 121-124.

Shawkey, M. D., Pillai, S. R., Hill, G. E., Siefferman, L. M. and Roberts, S. R. 2007. Bacteria as
an agent for change in structural plumage color: correlational and experimental evidence. –
Am. Nat. 169: S113-S121.

Shawkey, M. D., Pillai, S. R. and Hill, G. E. 2009. Do feather-degrading bacteria affect sexually
selected plumage color? – Naturwissenschaften 96: 123-128.

Sherry, T. W., Holmes, R. T., Pyle, P. and Patten, M. A. 2020. American Redstart (*Setophaga*
ruticilla), version 1.0. In Birds of the World (P. G. Rodewald, Editor). Cornell Lab of
Ornithology, Ithaca, NY, USA.

Siefferman L. and Hill, G. E. 2005. Male Eastern Bluebirds trade future ornamentation for
current reproductive investment. – Biol. Lett. 1: 208–211

Simons, M. J. P., Groothuis, T. G. G. and Verhulst, S. 2015. An appraisal of how the vitamin A-
redox hypothesis can maintain honesty of carotenoid-dependent signals. – Ecol. Evol. 5:
224-228.

Slagsvold, T. and Lifjeld, J. T. 1985. Variation in plumage colour of the great tit *Parus major* in
relation to habitat, season and food. – J. Zool. 206: 321–328.
556
557 Sparrow, K. L., Donkor, K. K., Flood, N. J., Marra, P. P., Pillar, A. G. and Reudink, M. W. 2017.
558   Conditions on the Mexican moulting grounds influence feather colour and carotenoids in
559   Bullock's orioles (*Icterus bullockii*). – Ecol. Evol. 7: 2643-2651
560
561 Stoddard, M. C., and Prum, R. O. 2008. Evolution of avian plumage color in a tetrahedral color
562   space: a phylogenetic analysis of new world buntings. – The American Naturalist, 171:
563   755-776.
564
565 Vorobyev, M., Osorio, D., Bennett, A. T., Marshall, N. J. and Cuthill, I. C. 1998. Tetrachromacy,
566   oil droplets and bird plumage colours. – J. Comp. Phys. A 183: 621–633.
567
568 Ward, G. M., Mahoney, S. M., Joly, S. and Reudink, M. W. 2021. Effects of age and weather
569   during moult on mountain bluebird *Sialia currucoides* structural colouration. – J. Avian
570   Biol. 52: https://doi.org/10.1111/jav.02616
571
572 Weaver, R. J., Santos, E. S., Tucker, A. M., Wilson, A. E. and Hill, G. E., 2018. Carotenoid
573   metabolism strengthens the link between feather coloration and individual quality. –
574   Nature Comm. 9: 1-9.
575
576 Weckstein, J. D., Kroodsma, D. E., Faucett, R. C., Poole, A. F. and Gill, F. B. 2002. Fox
577   Sparrow (*Passerella iliaca*), version 2.0. The Birds of North America (PG Rodewald, ed.).
578   Cornell Lab of Ornithology, Ithaca, New York, USA. https://doi. org/10.2173/bna, 715.
579
580 Wilkins, M. R., Seddon, N., and Safran, R. J. 2013. Evolutionary divergence in acoustic signals:
581   causes and consequences. – Trends Ecol. Evol. 28: 156-166.
582

Table 1. Linear mixed effects model selection results using a sample-size corrected Akaike's

Information Criterion (AICc) method. Models were built to predict the effects of American

redstart colouration (traditional colourimetric values (from Montgomerie 2006): brightness, red

chroma, red hue and tetrahedral colourspace values (from Stoddard and Prum 2008): hue θ, hue

Φ, r.achieved, and luminance) based on sex, site, season, age, δ^{15}N, and δ^{13}C. Final models used

in analyses are indicated in bold. ΔAIC: Change between full and reduced model; AICcWt: AICc

weights, indicating the probability a model is the most parsimonious model; LL: Log-likelihood.

Variable	Model	K	AICc	ΔAICc	AICcWt	LL
Brightness	**Reduced**	**6**	**-473.7**	**0**	**0.8**	**243.2**
	Full	9	-470.8	3	0.2	245.1
Red Chroma	**Reduced**	**4**	**-537.5**	**0**	**1**	**272.9**
	Full	11	-529	8.5	0	276.6
Red Hue	**Reduced**	**7**	**79.3**	**0**	**0.9**	**-32.2**
	Full	11	84.4	5.1	0.1	-30.1
Hue θ	**Reduced**	**9**	**-327.1**	**0**	**0.9**	**173.3**
	Full	11	-322.5	4.7	0.1	173.3
Hue Φ	**Reduced**	**7**	**-93.9**	**0**	**0.9**	**54.4**
	Full	11	-88.9	5.0	0.1	56.5
r.achieved	**Reduced**	**6**	**-424.2**	**0**	**0.8**	**218.4**
	Full	11	-421.6	2.6	0.2	222.9
Luminance	**Reduced**	**4**	**-444.5**	**0.0**	**0.9**	**226.4**
	Full	11	-440.3	4.2	0.1	232.2

Table 2. Model results demonstrating the effect of AICc selected fixed effects of site, season, sex, age, δ^{15}N, and δ^{13}C on American redstart colouration (brightness, red chroma, red hue, and tetrahedral colourspace hue θ, Φ, r.achieved, and luminance), and variances of the random effects of year. Significant effects are in bold.

Variable	Fixed effect	Model *df*	Residual *df*	*F*	*P*	Random effect σ^2
Brightness	**Site**	**1**	**134.0**	**6.45**	**0.012**	
	Season	**1**	**134.0**	**12.82**	**<0.0001**	0
	δ15N	**1**	**134.0**	**11.78**	**0.001**	
Red Chroma	Site	1	134.0	3.16	0.078	0
Red Hue	**Season**	**1**	**134.0**	**26.09**	**<0.0001**	
	Age	**1**	**134.0**	**9.43**	**0.003**	0
	Sex	**1**	**134.0**	**12.39**	**0.001**	
	Season x Age	**3**	**134.0**	**7.96**	**0.006**	
Hue θ	**Site**	**1**	**133.9**	**5.93**	**0.02**	
	Season x Age	**3**	**133.9**	**30.11**	**<0.0001**	0
	Sex x Age	**2**	**133.9**	**49.50**	**<0.0001**	
Hue Φ	**Site**	**1**	**133.8**	**9.86**	**0.002**	
	Season	**1**	**133.6**	**129.14**	**<0.0001**	0.03
	Age	**1**	**132.0**	**24.12**	**<0.0001**	
	Sex	**1**	**132.0**	**5.89**	**0.02**	
r.achieved	**Season**	**1**	**134.0**	**33.69**	**<0.0001**	
	Age	**1**	**131.9**	**6.89**	**0.01**	0.002
	δ15N	**1**	**131.9**	**6.86**	**0.01**	
Luminance	**δ15N**	**1**	**134.0**	**5.59**	**0.02**	0.002

Fig. 1. Map showing study sites in New York, US (Braddock Bay Bird Observatory) and off the coast of Maine, US (Appledore Island Migration Station). Inset photos show adult female (top) and adult male (bottom) American redstarts (photo credit: Kristen M. Covino).

Fig. 2. American redstart plumage colour varied by age and sex. Left panels: Reflectance plots showing the percentage (±SE) of light reflected off the feather surface. Right panel: Tetrahedral plot depicting age and sex plumage colour variation after correcting for the avian visual system. Grey dot represents the achromatic center of tetrahedral space.

Fig. 3. Left panels: American redstart feather traditional brightness (from Montgomerie 2006) in two sampling sites (AIMS and BBBO) in Fall and Spring. Right panel: Brightness and $\delta^{15}N$ sampled from American redstart feathers. Females are indicated by red dots and males are indicated by blue dots.

Fig. 4. Female (F) and male (M) American redstart feather red hue (from Montgomerie 2006) in adult (left panels) and young (right panels) birds during fall (top) and spring (bottom) during fall (top) and spring (bottom).

Fig. 5. Female (red) and male (blue) American redstart tailfeather colour variation as modelled in tetrahedral colourspace. Hue θ represents the angle of the color vector in the longitudinal plane, while hue Φ represents the angle in the latitudinal plane. Together hue θ and hue Φ describe the hue or the direction of colour vector in tetrahedral space (Stoddard and Prum 2008). Top panel: hue θ varied by season, age, and sex. Bottom panel: Hue Φ similarly varied by season, age, and sex.

Fig. 6. American redstart feather reflectance as modelled in tetrahedral colourspace (Stoddard and Prum 2008). Females are shown in red and males in blue. Top panel: r.achieved, representing the total length of the colour vector from the achromatic center of the tetrahedron, decreased with increasing δ^{15}N. Bottom panel: Luminance, representing the total amount of light reflected from the feather sample increased with δ^{15}N.