Modulation in Alpha Band Activity Reflects Syntax Composition:

An MEG Study of Minimal Syntactic Binding

Sophie M. Hardy[1,2]*, Ole Jensen[1], Linda Wheeldon[3],

Ali Mazaheri[1,4], Katrien Segaert[1,4]*

[1]Centre for Human Brain Health, University of Birmingham, B15 2TT, UK; [2]Department of

Psychology, University of Warwick, CV4 7AL, UK; [3]Department of Foreign Languages and

Translations, University of Agder, 4630, Norway, [4]School of Psychology, University of

Birmingham, B15 2TT, UK

Corresponding authors: sophie.hardy@warwick.ac.uk; k.segaert@bham.ac.uk

Abstract

Successful sentence comprehension requires the binding, or composition, of multiple words into larger structures to establish meaning. Using magnetoencephalography, we investigated the neural mechanisms involved in binding at the syntax level, in a task where contributions from semantics were minimized. Participants were auditorily presented with minimal sentences that required binding (pronoun and pseudo-verb with the corresponding morphological inflection; *"she grushes"*) and pseudo-verb wordlists that did not require binding (*"cugged grushes"*). Relative to no binding, we found that syntactic binding was associated with a modulation in alpha band (8-12 Hz) activity in left-lateralized language regions. First, we observed a significantly smaller increase in alpha power around the presentation of the target word (*"grushes"*) that required binding (-0.05s to 0.1s), which we suggest reflects an expectation of binding to occur. Second, during binding of the target word (0.15s to 0.25s), we observed significantly decreased alpha phase-locking between the left inferior frontal gyrus and the left middle/inferior temporal cortex, which we suggest reflects alpha-driven cortical disinhibition serving to strengthen communication within the syntax composition neural network. Together, our findings highlight the critical role of rapid spatial-temporal alpha band activity in controlling the allocation, transfer and coordination of the brain's resources during syntax composition.

Keywords: alpha oscillations; left-lateralized language network; inter-regional connectivity; sentence comprehension

Modulation in Alpha Band Activity Reflects Syntax Composition:

An MEG Study of Minimal Syntactic Binding

The expressive power of human language is largely derived from our ability to combine multiple words into larger syntactic structures with more complex meaning. Such binding, or compositional, processes occur during the comprehension of even the most basic two-word phrases (e.g., *"she walks"*). Characterizing the neural processes involved in composition – also referred to as *Unification* (Hagoort, 2003) or *Merge* (Chomsky, 1995) – has been a central topic of research for many years (Hagoort, 2019; Matchin & Hickok, 2020; Pylkkänen, 2019). In the present study we provide novel insight into the neural mechanisms involved in syntax composition using a task in which syntactic binding is dissociable from semantic composition. We use the term *syntactic binding* to specifically refer to the neural processes involved in the combining of individual words into larger structures. We employed a minimal phrase paradigm involving pseudo-words (i.e., follow the rules of a language, but have no given meaning).

Sentential compositional processes predominantly occur within a left-lateralized network of brain regions, including the inferior frontal gyrus (IFG) and angular gyrus (Friederici et al., 2000; Humphries et al., 2006; Matchin et al., 2017; Pallier et al., 2011). Within this network, modulations in alpha and beta frequencies are thought to be crucial for higher-order linguistic functions (Bastiaansen et al., 2010; Meyer, 2018; Prystauka & Lewis, 2019). However, the precise neural mechanisms of syntax composition, relating to frequency modulations, remain elusive. This is because while some studies have found compositional processing to be associated with increased alpha and beta power (Meyer et al., 2013; Segaert et al., 2018), others have found sentence unification to be associated with a power *de*crease (Gastaldon et al., 2020; Lam et al., 2016; Wang et al., 2012). Moreover, it is unclear how functional connectivity (i.e., phase-locked) between the neural oscillations in different brain

regions may contribute towards successful syntax composition. Current evidence suggests that functional connectivity between regions implicated in compositional processes, such as the left IFG, anterior temporal lobe (ATL) and posterior superior temporal gyrus, is beneficial for sentence comprehension (Lopopolo et al., 2021; Schoffelen et al., 2017; Vassileiou et al., 2018). However, the studies discussed thus far have typically used complex sentence structures – this means that other cognitive processes, such as working memory, are also involved in comprehending the sentence stimuli (Pylkkänen, 2019), thereby making it difficult to functionally isolate the neural processes involved in sentence composition alone. In this study, therefore, we aim to characterize the neural mechanisms of syntax composition, both in terms of power modulations and phase-locked connectivity, at its most basic two-word level.

A short two-word sentence (e.g., *"red boat"*) is a traceable linguistic unit which can be used to decompose the brain networks implicated in the processing of lengthier sentences (Pylkkänen, 2019). Research involving minimal sentences has identified a network of left-lateralized brain areas that underlie composition, most notably including the ATL (Bemis & Pylkkänen, 2011, 2013; Blanco-Elorrieta et al., 2018; Fyshe et al., 2019; Pylkkänen et al., 2014; Schell et al., 2017; Westerlund et al., 2015; Zaccarella et al., 2017; Zhang & Pylkkänen, 2015, 2018; Ziegler & Pylkkänen, 2016; but cf. Kochari et al., 2021). Importantly, in order to more precisely identify the neural processes involved in syntactic binding that are dissociable from semantics, one further approach is to use pseudo-words within a minimal phrase paradigm. Using fMRI, Zaccarella and Friederici (2015) found increased hemodynamic responses in the anterior part of the left pars opercularis (part of the IFG) during comprehension of determiner-noun phrases involving pseudo-nouns (*"this flirk"*) compared to wordlists involving one pseudo-noun (*"apple flirk"*). Using EEG, Segaert et al. (2018) found increased alpha and beta power (centralized over frontal-central

electrodes) during comprehension of minimal phrases involving pseudo-verbs (*"she grushes"*) compared to wordlists of two pseudo-verbs (*"cugged grushes"*), which they interpreted as reflecting syntactic binding (see also Poulisse et al., 2020). We aim to build on these findings and use magnetoencephalography (MEG) to precisely characterize and localize the rapid temporal features and functional connectivity within a spatially distributed network of brain regions that support syntax composition independent of semantics.

Materials and Methods

The methods, hypotheses and planned analyses of this study were pre-registered on the Open Science Framework prior to data collection (https://osf.io/ntszu).

Participants

We recruited 25 healthy participants: all were right-handed and native monolingual British-English speakers. One participant was excluded due to excessive movement artefacts during the MEG test session (> 50% trials removed), meaning that a sample of 24 participants was used in the time-frequency analyses (13 female / 11 male, $M = 24.2$yrs, $SD = 4.1$yrs). Anatomical T1 brain scans were acquired for 21 of the participants (two participants did not attend the MRI session, and another did not complete the MRI session due to unexpected discomfort). Further technical issues with MRI-MEG co-registration with two participants meant that a sample of 19 participants was used for the source localization and connectivity analyses (10 female / 9 male, $M = 24.1$yrs, $SD = 4.2$yrs). The study was approved by the University of Birmingham Ethical Review Committee. All participants provided written informed consent and were compensated monetarily.

Experimental Design and Stimuli

We employed a simple design of two experimental conditions: the sentence condition consisting of a minimal two-word phrase (pronoun plus pseudo-verb) for which syntactic binding occurred (e.g., *"she grushes"*); and the wordlist condition consisting of two pseudo-verbs for which no syntactic binding occurred (e.g., *"cugged grushes"*). Syntactic binding occurred in the sentence condition (but not the wordlist condition) because the correct morphological inflection (i.e., -es) cued binding with the corresponding pronoun, in a way that is similar to how an intransitive verb phrase may be interpreted (e.g., *"she grushes"* is akin to *"she walks"*).

Behavioural evidence from previous use of this paradigm has shown that participants judge pseudo-verb sentences with the correct morphological inflection to be valid sentences (e.g., *"she grushes"*), but judge sentences with the incorrect inflection (e.g., *"she grush", "I grushes"*) and pseudo-verb wordlists (e.g., *"cugged grushes"*) to be invalid sentences (Poulisse et al., 2019). This is evidence that listeners do not attempt to bind the two pseudo-verbs together when in a wordlist as, if they did, they would judge it to be a valid sentence (e.g., if they had interpreted *"cugged grushes"* as a deverbal adjective and noun pairing). We are therefore confident that participants in our study were engaging in syntactic binding when a minimal sentence was presented, but not when a wordlist was presented.

To construct the experimental items, we used a set of 20 pseudo-verbs created by Ullman et al. (1997).[1] All pseudo-verbs were monosyllabic and could be inflected according to the grammatical rules of regular English verbs. We combined each pseudo-verb with three different morphological affixes (no affix; +s; +ed) to create 60 possible pseudo-verb-affix combinations. In English, only certain pronouns may be combined with certain affixes (e.g*., "she grushes"* is acceptable, but *"I grushes"* is not). Using a list of six pronouns (I, you, he,

---

[1] Pseudo-verbs (root form, un-inflected): brop, crog, cug, dotch, grush, plag, plam, pob, prap, prass, satch, scash, scur, slub, spuff, stoff, trab, traff, tunch, vask.

she, they, we), we created 120 sentence items by pairing each pseudo-verb-affix with two different pronouns that were syntactically appropriate for the corresponding affix, such that syntactic binding may plausibly occur (e.g., *"I dotch", "she grushes", "they cugged"*). To create the wordlist items, we paired together two different pseudo-verb-affix stimuli for which no syntactic binding occurred (e.g., *"cugged grushes", "dotch traffed"*). Each pseudo-verb-affix stimulus occurred twice as the first word in a pair and twice as the second word in a pair, creating a total of 120 wordlist items. We ensured that the two words within each wordlist pair always consisted of a different pseudo-verb and a different affix. All auditory stimuli were spoken by a native English male speaker and normalised to 1db volume.

We also created 120 filler items. Sixty of the fillers consisted of reversed speech and were included as a detection task for the participants. We reversed the speech of each of the 60 pseudo-verb-affix combinations and then paired each with either a non-reversed pseudo-verb-affix or pronoun (half as the first word of the pair and half as the second word). A further 60 filler items were used to increase the variability of stimuli presented to participants (Roland et al., 2007). Thirty such items consisted of two pronouns (e.g., *"she I"*); this contrasted the experimental sentence items in which a pronoun was followed by a pseudo-verb (e.g., *"she grushes"*). The other 30 items consisted of a pseudo-verb-affix stimulus followed by one of five possible adverbs (early, promptly, quickly, rarely, safely; e.g., *"cugged quickly"*); this contrasted the experimental wordlist items in which a pseudo-verb was followed by another pseudo-verb (e.g., *"cugged grushes"*). Overall, this set of fillers created a task and variability for the participants – the filler items did not act as a control to the experimental items and were not analysed.

Experimental Procedure

The participants' task was to detect the reversed speech (which only occurred on filler trials). On each trial, participants were auditorily presented with a two-word phrase (**Figure 1**). Participants were instructed to press a button if part of the speech was reversed (half of the participants used their left index finger, and half used their right index finger), but to do nothing if the speech was not reversed. There was no difference in response decision processes (i.e., no button press) between the critical experimental conditions of interest (sentence vs. wordlist). Each participant completed 360 trials (consisting of 240 experimental trials and 120 filler trials) in a unique randomised order, divided into six blocks of 60 trials each. Before beginning the task, participants completed 23 practice trials that were similar to the experimental and filler items used in the main task.

Figure 1. Stimuli presentation timings per trial and the related epoch window. Stimuli presentation and trigger signals were controlled using E-Prime (Schneider et al., 2002). Visual stimuli were presented using a PROPixx projector, and auditory stimuli were presented using the *Elekta* audio system and MEG-compatible ear plugs. Participants' motor responses were recorded using a NAtA button pad.

As expected, participants were highly accurate at detecting the reverse speech on the filler trials ($M = 94.6\%$, $SD = 2.3\%$, $Range = 82\text{-}98\%$), indicating that they were closely

listening to the filler and experimental stimuli throughout (since they did not know when the reversed speech would be presented).

Data Acquisition

During the task, ongoing MEG data were recorded using the TRIUX^{TM} system from *Elekta* (Elekta AB, Stockholm, Sweden). This system has 102 magnetometers and 204 planer gradiometers. These are placed at 306 locations, each having one magnetometer and a set of two orthogonal gradiometers. The data were collected using a sampling rate of 1000 Hz and was stored for offline analyses. Prior to sampling, a lowpass filter of ~250 Hz was applied. Scalp surface data were acquired using a Polhemus 3D digitiser to facilitate later co-registration with anatomical brain scans. Additional electrooculography (EOG) and electrocardiogram (ECG) data were collected using methods compatible with the TRIUX^{TM} system.

Anatomical high-resolution T1 brain images were acquired for participants at a later session using a MAGNETOM Prisma 3T MRI system from *Siemens* (Siemens Healthcare, Erlangen, Germany). These images were used for the reconstruction of individual head shapes to create forward models for the source localization analyses.

MEG Pre-processing

The offline processing and analyses of the data were performed using functions from the Fieldtrip software package (Oostenveld et al., 2011) and custom scripts in the MATLAB environment. First, we applied a 0.1 Hz high-pass filter to the MEG data to remove slow frequency drift in the data. The data were segmented into epochs aligned to the onset of the auditory presentation of the second word from -2.7s to 1.9s (see **Figure 1**). We visually inspected the waveforms of each experimental trial and removed trials that contained

excessive signal artefacts (e.g., large sensor jumps or gross motor movement by the participant); on average, we removed 16% of experimental trials (38.3/240) per participant (*Range* = 4-47%). We also removed any persistently poor channels that contained excessive noise or flatlined (*Mean* channels removed per participant = 2.46; *Range* = 0-9). We then used a spline interpolation weighted neighbourhood estimate to interpolate across the removed channels per participant. Ocular and cardiac artifacts were removed from the data using an independent component analysis (ICA) (*Mean* artifacts removed per participant = 1.75; *Range* = 1-3). We identified these components from their stereotypical topography and time course, as well as by comparisons with the recorded ECG and EOG time courses.

We removed all filler trials as we were specifically interested in the difference in neural responses between the experimental sentence and wordlist conditions (i.e., our contrast of interest). We further removed trials for which the participant incorrectly responded with a button press (i.e., indicated that the speech was reversed when it was not) and trials during which the participant made an accidental button press before the response screen. Once all data pre-processing was complete, there was an average of 101 sentence trials and 100 wordlist trials per participant that were usable for the analyses (out of a maximal 120 per experimental condition).

Statistical Analyses

Time-frequency

For the frequency range 1-30 Hz (1 Hz steps), we obtain time-frequency representations (TFRs) of power for each trial using sliding Hanning tapers with an adaptive time window of three cycles for each frequency ($\Delta T = 3/f$). This approach has also been used in a number of previous studies (e.g., Segaert et al., 2018; van Diepen & Mazaheri, 2017; Whitmarsh et al., 2011). For each participant, the data for the planar gradiometer pairs was

added to create a 102-channel combined planar map in sensor space, and we baseline-corrected the data using the oscillatory activity in a 0.5s period of the fixation cross presentation (specifically, -1.9s to -1.4s in the epoch window). We calculated the TFRs separately per experimental condition for each participant and then averaged across all participants.

We assessed the statistical differences in time-frequency power between the sentence and wordlist conditions across participants using a cluster-level randomization test (incorporated in the Fieldtrip software), which circumvents the type-1 error rate in a situation involving multiple comparisons (i.e., multiple channels and time-frequency points; Maris & Oostenveld, 2007). This approach first clusters the data in sensor space depending on whether the contrast between the two conditions exceeds a dependent samples t-test threshold of $p <$ .05 (two-tailed). We used the following pre-defined frequency bands: alpha (8-12 Hz); low beta (15-20 Hz) and high beta (25-30 Hz). We considered a cluster to consist of at least two significant adjacent combined planar gradiometers. A Monte Carlo p-value of a cluster was then obtained by calculating the number of times the t-statistics in the shuffled distribution is higher than the original t-statistic obtained when contrasting conditions. We performed the analyses within the time window of interest, centred around the presentation of the second word (-0.5s to 1s of the epoch).

Source Localization

A realistically shaped description of each participant's brain was constructed using individual head models obtained using the Polhemus 3D digitiser and the acquired MRI anatomical brain scan (where available). Source estimation of the time-locked MEG data was performed using a frequency-domain beam-forming approach (dynamic imaging of coherent sources [DICS]; Gross et al., 2001), which uses adaptive spatial filters to localize power in

the entire brain. The brain volume of each individual participant was discretized to a grid with a 0.8cm resolution and the lead field was calculated for each grid point. A common filter was calculated for the sentence and wordlist condition and then applied for the data separately for the individual conditions (Mazaheri et al., 2014; Whitmarsh et al., 2011). The source estimates of the individual participants' functional data along with the individual anatomical MRI images were warped into a standard Montreal Neurological Institute (MNI) standard brain (Quebec, Canada; http://www.bic.mni.mcgill.ca/brainweb) before averaging and statistics.

We performed cluster-based randomization tests to identify the grid-points in which there was a significant difference in power between the two experimental conditions, guided by the findings of the time-frequency analyses (i.e., time-windows). We used these clusters to identify specific regions of interest (ROIs) of the condition difference. We derived anatomical labels of these regions from Brodmann's map and from the Automated Anatomical Labelling Atlas (Tzourio-Mazoyer et al., 2002).

Inter-regional Connectivity

We performed inter-regional connectivity analyses on four distinct ROIs identified as showing a significant condition difference in the source localization analyses. TFRs of the complex Fourier spectra for each trial at each ROI was obtained by using a sliding Hanning tapers with an adaptive time window of three cycles for each frequency ($\Delta T = 3/f$). We calculated the inter-regional phase-locked values of the oscillatory activity (1-30 Hz) between the four ROIs (creating six inter-regional connections) for the time period of interest (-0.5s to 1s of the epoch) separately for the sentence condition and wordlist condition (following Bastos & Schoffelen, 2016; Lachaux et al., 1999). We used the following formula to calculate the phase locking value (PLV):

$$PLV = \frac{1}{N} \left| \sum_{n=1}^{N} e^{i(\emptyset_{(n,t)} - \psi_{(n,t)})} \right|$$

Applied to our study, the PLV reflects the consistency of the phase difference of oscillatory activity across trials between two ROIs. Here: N is the number of trials; $\emptyset_{(n,t)}$ is the phase (obtained from the complex spectra) at time t in trial n in one ROI; and $\psi_{(n,t)}$ is the phase at time t in trial n in the other ROI. A PLV of 0 indicates no phase locking between the activities of the two ROIs, whereas a PLV of 1 is indicative of perfect phase locking.

We then performed cluster-level randomization tests (Maris & Oostenveld, 2007) on the PLVs in order to identify the inter-regional connections in which there was a significant difference in phase-locked values between the two experimental conditions (for a similar statistical approach see, Schmidt et al., 2014). The use of a cluster-level randomization test, in which a Monte Carlo p-value is obtained, enables the control of multiple comparisons of different ROIs and time-frequency points. To correct for the multiple analyses performed across the six different inter-regional connections, we applied a Bonferroni correction (α / n comparisons) to our critical p value of interest (.05/6 = .0083). We used the same pre-defined frequency bands as for the oscillatory time-frequency analyses: alpha (8-12 Hz); low beta (15-20 Hz) and high beta (25-30 Hz).

Results

We compared participants' MEG activity during the comprehension of minimal sentences that required binding (a pronoun combined with a pseudo-verb with the corresponding morphological inflection; *"she grushes"*) to wordlists that did not require binding (two pseudo-verbs; *"cugged grushes"*). Our findings reveal two key mechanisms of syntactic binding.

Less alpha power in the left-lateralized brain network when syntactic binding occurs

The grand-average of the time-frequency representations (TFR) of power averaged across all sensors aligned to the onset of the second word are summarized in **Figure 2A** for the sentence condition in which syntactic binding occurred, and the wordlist condition in which no binding occurred. In both conditions, there are power increases in alpha and low beta surrounding the presentation of the second word (at 0s; *"grushes"*). Approximately 0.5s after the second word presentation, there is a slight decrease in alpha and low beta power in both conditions.

The statistical results of the cluster-based permutation tests (which controlled for multiple comparisons of different channels and time-frequency points) revealed that there was a significant condition difference in the alpha frequency range (8-12 Hz) surrounding the presentation of the second word where alpha power was lower in the sentence condition, compared to the wordlist condition ($p = .021$, see **Figure 2B-C**). This difference was maximal around the onset of the second word (-0.05s to 0.1) over a cluster of sensors predominantly in the left-frontal region (**Figure 2B**). Importantly, this oscillatory effect of alpha was distinct from the evoked fields as we did not observe any significant condition differences in the event-related fields during this time period ($p > .05$). We found no significant difference in oscillatory activity within the other analysed frequency bands: low beta (15-20 Hz) and high beta (25-30 Hz).

Source analyses co-registered on the participants' anatomical MRI brain scans indicated that the significant condition difference in alpha power (8-12 Hz, -0.05s to 0.1s) was localized to a network of left-lateralized brain regions which are typically associated with language function (**Figure 3A**). Within this network, we identified four brain areas (see **Figure 3B**) in which significant differences were found between the sentence condition and wordlist condition: the left inferior frontal gyrus (BA44; peak coordinates [-42 7 17]); the left

angular gyrus (BA39; peak coordinates [-58 -48 29]); the left middle/inferior temporal cortex (BA21; peak coordinates [-60 -30 -18]); and the left anterior frontal gyrus (BA46; peak coordinates [-40 43 7]).

Figure 2. (A) Time-frequency representations of power averaged across all sensors, expressed as an absolute change from the baseline period (i.e., -1.9s to -1.4s before the onset of the second word) for the sentence condition [left panel] in which syntactic binding occurred (e.g., *"she grushes"*), and the wordlist condition [right panel] in which no binding occurred (e.g., *"cugged grushes"*). Time relates to the main time period of interest aligned to the onset of the second word (at 0s). The rectangle highlights the time period where we maximally observed the significant difference in alpha power (8-12 Hz) between the two conditions (-0.05s to 0.1s; *p* = .021). **(B)** The scalp topography of the condition contrast

(sentence minus wordlist) of the averaged alpha power activity in the time window (-0.05s to 0.1s) where we maximally observed the significant difference in alpha power between the two conditions. The black dots illustrate where this effect was largest at the scalp level. For visualization purposes, the condition contrast of the non-baseline corrected data is shown. **(C)** The time course of the alpha power envelope for the sensors showing a significant difference in power between the sentence and wordlist conditions. The shaded grey area corresponds to the time window in which the difference between conditions is maximal centred around the presentation of the second word (-0.05s to 0.1s).

Figure 3. Source localization estimates of the condition difference in alpha power (8-12 Hz) of the sentence condition (e.g., *"she grushes"*) minus the wordlist condition (e.g., *"cugged grushes"*) surrounding the presentation of the target word (-0.15s to 0.15s) as shown for a surface **(A)** and sliced **(B)** view of the brain. The displays are masked for significant clusters only ($p < .05$). The condition difference was maximal over the left-frontal areas of the brain, with significant differences observed in the left inferior frontal gyrus, the left angular gyrus, the left middle/inferior temporal cortex, and the left anterior frontal gyrus.

During syntactic binding, there is decreased alpha phase-locking between the left inferior frontal gyrus and the left middle/inferior temporal cortex

We calculated the inter-regional phase-locked values between the peak coordinates of the four brain areas identified as displaying significant condition differences in the source localization analyses (**Figure 4A**). The statistical results of the cluster-based permutation tests (which controlled for multiple comparisons of different ROIs and time-frequency points) revealed a significant condition difference ($p = .005$) in inter-regional phase-locking in alpha activity (8-12 Hz) between the left inferior frontal gyrus (BA44) and the left middle/inferior temporal cortex (BA21) following the presentation of the second word that required binding (around 0.15s to 0.25s). During this time period, there was significantly less alpha phase-locking between these two brain regions in the sentence condition, compared to the wordlist condition (**Figure 4B**). While we are aware that amplitude may correlate with PLV (van Diepen & Mazaheri, 2018), we consider that it is unlikely that our observed condition difference in the PLVs is driven by amplitude differences alone because it occurred at a time interval when there was *not* the greatest amplitude difference between the two conditions, and because we only observed the effect between two specific ROIs, not the whole analysed left-lateralized brain network.

We did not find any other significant condition differences in inter-regional phase-locking in the other connections between our ROIs. Qualitatively though, across both conditions, inter-regional phase-locking does appear to be strongest between the left angular gyrus and the left middle/inferior temporal cortex, as well as within the alpha frequency range between the left anterior frontal gyrus and the left inferior frontal gyrus (**Figure 4C**).

Figure 4. (A) Inter-regional phase-locking differences of oscillatory activity (2-30 Hz) were calculated between four brain areas in the left hemisphere for the time period of interest (-0.5s to 1s; second word presented at 0s). **(B)** A significant condition difference in inter-regional phase-locking was observed between the left inferior frontal gyrus (BA44) and the

left middle/inferior temporal cortex (BA21) in the alpha frequency range (8-12 Hz). Time-frequency estimates of the phase-locking value are shown for the sentence condition (e.g., *"she grushes"*) [left upper panel], the wordlist condition (e.g., *"cugged grushes"*) [left lower panel] and the condition contrast (sentence minus wordlist) [right panel]. There was significantly greater phase-locking of alpha activity between BA44 and BA21 in the wordlist condition compared to the sentence condition when subjected to a cluster-randomisation analysis, maximally observed around 0.15s to 0.25s following the presentation of the target word ($p = .005$), as highlighted by the rectangle. **(C)** The inter-regional phase-locking values between the other brain regions of interest. Here, no other significant differences were found between the sentence and wordlist conditions.

Discussion

The findings of our MEG study suggest that minimal syntax composition is associated with distinct oscillatory changes in alpha band activity and the engagement of the left-lateralized language network of the brain. In the reported experiment we compared minimal pseudo-verb sentences for which syntactic binding may plausibly occur (e.g., *"she grushes"*) and wordlists for which no binding occurred (e.g., *"cugged grushes"*). We found that surrounding (-0.05s to 0.1s) the presentation of the target word (*"grushes"*), alpha power was significantly less in the sentence, compared to the wordlist, condition. The sources of this condition difference were localized to left-lateralized brain regions, including the inferior frontal gyrus (IFG), angular gyrus, middle/inferior temporal cortex, and anterior frontal gyrus. Moreover, following the presentation of the target word (0.15s to 0.25s), we observed decreased alpha phase-locking between the LIFG and the left middle/inferior temporal cortex in the sentence, compared to the wordlist, condition, indicating that syntactic binding is associated with a decoupling of alpha activity between these regions.

Modulation in alpha power in a network of left-lateralized language regions reflects an expectation of binding to occur

Compared to baseline, we observed an increase in alpha power in both conditions during the comprehension of the second word, but critically this power increase was *less* when syntactic binding was required, compared to when no binding was required. This modulation in alpha power is consistent with existing evidence of alpha oscillatory activity in compositional processing (Gastaldon et al., 2020; Lam et al., 2016; Segaert et al., 2018) and the view that neural oscillations subserve the processing of syntactic information within language-relevant cortices (Meyer, 2018; Prystauka & Lewis, 2019). Our findings further demonstrate that alpha operates in auditory language processing in addition to its role in visual processing in occipital areas (Mazaheri et al., 2014; Zumer et al., 2014), supporting a varied functionality of alpha oscillations in multiple sensory systems in different cortical regions (Foxe & Snyder, 2011).

We suggest that, in our experiment, the observed lesser alpha power in the binding context around the presentation of the target word reflects an expectation of binding to occur. When comprehending linguistic input, we build expectations in order to predict upcoming words (Chang et al., 2006; Kuperberg & Jaeger, 2016); indeed, probability estimates are considered to be inherent within the neural linguistic system (Kutas et al., 2011). Thus, if the first word was a pronoun, as opposed to a pseudo-verb, the participant may reasonably expect that binding was likely to be required given their existing knowledge (based on language use in everyday life) about the properties and syntactic function of pronouns. This interpretation is consistent with studies that have found greater alpha suppression (i.e., decreased alpha power compared to baseline) when participants comprehend a highly predictive, compared to a less predictive, sentence (Piai et al., 2014; Rommers et al., 2017; Wang et al., 2017) and the proposed role of alpha power decreases in controlling the allocation of the brain's resources

(Jensen & Mazaheri, 2010; Klimesch, 2012). In particular, decreased alpha power limits cortical excitability, thereby allowing for continual processing regardless of alpha phase (medium-to-high attentional state); whereas, when alpha power is higher, cortical processing is more discontinuous as it depends on the phase of the alpha rhythm (rhythmic attentional state) (Alavash et al., 2021; Jensen et al., 2014; van Diepen et al., 2019). Our finding of less alpha power around the presentation of the target word that required binding (compared to a no binding context) may therefore reflect the initiation of anticipatory binding processes (for which a medium-to-high attentional state is required), along with the increased engagement of brain regions involved in syntactic binding.

The observed condition difference in alpha power was localized to a left-lateralized network of brain regions, consistent with established neurobiological models of linguistic processing (Friederici, 2011; Hagoort, 2003; Tyler & Marslen-Wilson, 2008). The implication of the LIFG (BA44) is expected given the region's proposed role in managing the combination of words into a coherent syntactic structure (Hagoort, 2005; Segaert et al., 2012; Snijders et al., 2008; Uddén et al., 2019) and in computing dependency structures (Lopopolo et al., 2021). Indeed, the LIFG has been identified in previous studies of minimal syntactic binding (Zaccarella et al., 2017; Zaccarella & Friederici, 2015) and top-down predictive processing (Matchin et al., 2017; Strijkers et al., 2019). The other three left-lateralized regions we identified – angular gyrus (BA39), anterior frontal gyrus (BA46), and middle/inferior temporal cortex (BA21) – have also been found to contribute towards syntactic processing (Giraud et al., 2004; Humphries et al., 2006; Matchin et al., 2017; Menenti et al., 2011; Segaert et al., 2012). Our findings therefore suggest that successful composition of the syntactic properties between words in a sentence is driven by the engagement of a distributed network of left-lateralized regions (including the frontal gyri, temporal cortex and angular gyrus), and critically their coordination (as we discuss in more

detail in the next section). Moreover, given that we did not observe any effect of compositionality in the anterior temporal lobe (ATL) in our task in which contributions from semantics were minimized, our findings further add to the current evidence that the functional role of the ATL relates primarily to semantic, not syntactic, composition (Del Prato & Pylkkänen, 2014; Kim & Pylkkänen, 2019; Pylkkänen, 2019; Wilson et al., 2014).

However, our findings of alpha power decrease is somewhat at odds with Segaert et al. (2018) who, using a similar paradigm but with EEG, found syntactic binding to be associated with alpha power increases. One explanation for this difference may reflect MEG vs. EEG differences in spatial coherence and sensitivity to deeper brain tissues that can lead to differences in detectable power (Bénar et al., 2019; Lopes da Silva, 2013). In particular, MEG and EEG differ in their sensitivity to the radical and tangential components of the dipolar sources in the brain, which can lead to the two producing diverging estimates for the same cognitive process (Dehghani et al., 2010; Edgar et al., 2003; Fonteneau et al., 2015; Ross et al., 2020). An alternative explanation for the study differences may relate to morpho-syntactic differences between English (used in this study) and Dutch (used in Segaert et al., 2018). Compared to English, Dutch is more morphologically complex in that there are a greater number of possible verb inflections (to signal its binding to a preceding pronoun) and there are more rules on inflection usage (Booij, 2019). This could mean that Dutch listeners focus more on morphological congruency, whereas English speakers focus more on overall sentential composition, leading to group differences in detectable alpha power. Further work is therefore needed to uncover how differences in the complexity of the inflection system of a language may drive differences in the oscillatory signature of syntax composition even at its most basic two-word level. Nevertheless, the findings of our study and Segaert et al. (2018) should not be considered as directly opposed, but instead reflective of the different components of the wider neural processes involved in syntax composition (the intricacies of

which may differ between languages or which may be differently detected by MEG vs. EEG),

and the more general functionality of alpha oscillations in syntax processing (Meyer, 2018;

Prystauka & Lewis, 2019).

Alpha band decoupling in the language network reflects strengthened network communication required for successful syntactic binding

Inter-regional connectivity analyses revealed that syntactic binding was associated

with a decoupling of alpha between the LIFG and the left middle/inferior temporal cortex: we

found *less* alpha phase-locking in the sentence, compared to the wordlist, condition following

the presentation of the target word (0.15s to 0.25s). The time window of the effect, after the

auditory processing of the target word (which typically occurs within the first 0.1s;

Zouridakis et al., 1998), suggests that it reflects the underlying syntax composition

mechanisms taking place as opposed to an expectation of binding to occur. Our finding is

consistent with evidence that alpha band decoupling, or desynchronisation, between relevant

brain regions is beneficial for language comprehension and can predict successful sentence

encoding (Alavash et al., 2021; Becker et al., 2013; Magazzini et al., 2016; Vassileiou et al.,

2018). This functional connectivity reflects the dynamic interaction among distributed brain

regions that are subserved by deep white matter pathways and which communicate through

frequency-specific networks (Sarubbo et al., 2020; Schoffelen et al., 2017; White et al.,

2018). This communication may operate through top-down mechanisms of cortical

disinhibition, such that alpha band desynchronisation serves to functionally disinhibit the

cortex to enable information to be transferred from and to specific areas for processing

(Jensen & Mazaheri, 2010; Klimesch, 2012; Sadaghiani & Kleinschmidt, 2016).

We suggest that similar mechanisms of frequency-specific communication and alpha-

driven disinhibition were operating in our task when participants comprehended sentences

that needed binding (*"she grushes"*). In order to bind the two words together into a minimal

syntactic structure, increased information, such as the labels of syntactic components

(Murphy, 2015), needed to be transferred between the LIFG and the middle/inferior temporal

cortex. Thus, the less alpha band synchronisation between these two regions, the greater the

cortical disinhibition; this, in turn, strengthens the communication within the cortical and

oscillatory network involved in syntax composition, thereby enhancing participants'

processing. Our findings are therefore indicative of a top-down downregulation of alpha

oscillations that operate as part of a broader attentional control network throughout the

brain's sensory channels, which during speech comprehension act to promote the spread of

information between language relevant cortices (Alavash et al., 2021; Sadaghiani &

Kleinschmidt, 2016). This interpretation further fits with the wider understanding of the role

of alpha oscillations in controlling the access, transfer and storage of information within the

brain (Bonnefond et al., 2017; Klimesch, 2012).

Summary

In sum, when comparing minimal pseudo-verb sentences that required binding (*"she

grushes"*) to no binding wordlists (*"cugged grushes"*), we first found evidence of less alpha

power in left-lateralized brain regions which we suggest reflects an expectation of binding to

occur. Second, we found that during syntactic binding there was decreased alpha phase-

locking between the LIFG and the left middle/inferior temporal cortex; we suggest that this

results from increased information transfer and the strengthening of the neural network

involved in syntax composition. The observed oscillatory modulations (including power

decreases and decoupling) occurred in brain regions in the left hemisphere known to be

relevant for language processing. These regions are not uniquely selective for syntactic

processing; rather, we suggest that each region plays a contributing role in syntax

composition mechanisms, and that coordination between these regions (particularly the LIFG and left middle/inferior temporal cortex) is critical for successful syntactic binding. Together, our findings contribute to the wider understanding of the rapid spatial-temporal dynamics of syntactic processing in the brain that occur independently of semantic processing.

Acknowledgments

This research was supported by an Economic and Social Research Council (ESRC) studentship awarded to Sophie M. Hardy from the University of Birmingham Doctoral Training Centre (grant number: ES/J50001X/1). We thank Jonathon Winter and Nina Salman for their help with data collection. The authors report no potential conflict of interest.

References

Alavash, M., Tune, S., & Obleser, J. (2021). Dynamic large-scale connectivity of intrinsic cortical oscillations supports adaptive listening in challenging conditions. *PLoS Biology*, in press. https://doi.org/10.1101/2021.02.22.432315

Bastiaansen, M., Magyari, L., & Hagoort, P. (2010). Syntactic unification operations are reflected in oscillatory dynamics during on-line sentence comprehension. *Journal of Cognitive Neuroscience*, *22*(7), 1333–1347. https://doi.org/10.1162/jocn.2009.21283

Bastos, A. M., & Schoffelen, J.-M. (2016). A tutorial review of functional connectivity analysis methods and their interpretational pitfalls. *Frontiers in Systems Neuroscience*, *9*. https://doi.org/10.3389/fnsys.2015.00175

Becker, R., Pefkou, M., Michel, C. M., & Hervais-Adelman, A. G. (2013). Left temporal alpha-band activity reflects single word intelligibility. *Frontiers in Systems Neuroscience*, *7*. https://doi.org/10.3389/fnsys.2013.00121

Bemis, D. K., & Pylkkänen, L. (2011). Simple composition: A magnetoencephalography investigation into the comprehension of minimal linguistic phrases. *Journal of Neuroscience*, *31*(8), 2801–2814. https://doi.org/10.1523/jneurosic.5003-10.2011

Bemis, D. K., & Pylkkänen, L. (2013). Basic linguistic composition recruits the left anterior temporal lobe and left angular gyrus during both listening and reading. *Cerebral Cortex*, *23*(8), 1859–1873. https://doi.org/10.1093/cercor/bhs170

Bénar, C. G., Grova, C., Jirsa, V. K., & Lina, J. M. (2019). Differences in MEG and EEG power-law scaling explained by a coupling between spatial coherence and frequency: A simulation study. *Journal of Computational Neuroscience*, *47*(1), 31–41. https://doi.org/10.1007/s10827-019-00721-9

Blanco-Elorrieta, E., Ferreira, V. S., Del Prato, P., & Pylkkänen, L. (2018). The priming of basic combinatory responses in MEG. *Cognition*, *170*, 49–63. https://doi.org/10.1016/j.cognition.2017.09.010

Bonnefond, M., Kastner, S., & Jensen, O. (2017). Communication between brain areas based on nested oscillations. *ENeuro*, *4*(2). https://doi.org/10.1523/ENEURO.0153-16.2017

Booij, G. (2019). *The morphology of Dutch*. Oxford University Press.

Chang, F., Dell, G. S., & Bock, K. (2006). Becoming syntactic. *Psychological Review*, *113*(2), 234–272. https://doi.org/10.1037/0033-295X.113.2.234

Chomsky, N. (1995). *The minimalist program*. The MIT Press. https://doi.org/10.1017/S0022226797006889

Dehghani, N., Cash, S. S., Chen, C. C., Hagler, D. J., Huang, M., Dale, A. M., & Halgren, E. (2010). Divergent cortical generators of MEG and EEG during human sleep spindles suggested by distributed source modeling. *PLoS ONE*, *5*(7), e11454. https://doi.org/10.1371/journal.pone.0011454

Del Prato, P., & Pylkkänen, L. (2014). MEG evidence for conceptual combination but not numeral quantification in the left anterior temporal lobe during language production. *Frontiers in Psychology*, *5*. https://doi.org/10.3389/fpsyg.2014.00524

Edgar, J. C., Huang, M. X., Weisend, M. P., Sherwood, A., Miller, G. A., Adler, L. E., & Cañive, J. M. (2003). Interpreting abnormality: An EEG and MEG study of P50 and the auditory paired-stimulus paradigm. *Biological Psychology*, *65*(1), 1–20. https://doi.org/10.1016/S0301-0511(03)00094-2

Fonteneau, E., Bozic, M., & Marslen-Wilson, W. D. (2015). Brain network connectivity during language comprehension: Interacting linguistic and perceptual subsystems. *Cerebral Cortex*, *25*(10), 3962–3976. https://doi.org/10.1093/cercor/bhu283

Foxe, J. J., & Snyder, A. C. (2011). The role of alpha-band brain oscillations as a sensory suppression mechanism during selective attention. *Frontiers in Psychology*, *2*. https://doi.org/10.3389/fpsyg.2011.00154

Friederici, A. D. (2011). The brain basis of language processing: From structure to function. *Physiological Reviews*, *91*(4), 1357–1392. https://doi.org/10.1152/physrev.00006.2011

Friederici, A. D., Meyer, M., & Von Cramon, D. Y. (2000). Auditory language comprehension: An event-related fMRI study on the processing of syntactic and lexical information. *Brain and Language*, *74*(2), 289–300. https://doi.org/10.1006/brln.2000.2313

Fyshe, A., Sudre, G., Wehbe, L., Rafidi, N., & Mitchell, T. M. (2019). The lexical semantics of adjective-noun phrases in the human brain. *Human Brain Mapping*, *40*(15), 4457–4469. https://doi.org/10.1002/hbm.24714

Gastaldon, S., Arcara, G., Navarrete, E., & Peressotti, F. (2020). Commonalities in alpha and beta neural desynchronizations during prediction in language comprehension and production. *Cortex*, *133*, 328–345. https://doi.org/10.1016/j.cortex.2020.09.026

Giraud, A. L., Kell, C., Thierfelder, C., Sterzer, P., Russ, M. O., Preibisch, C., & Kleinschmidt, A. (2004). Contributions of sensory input, auditory search and verbal comprehension to cortical activity during speech processing. *Cerebral Cortex*, *14*(3), 247–255. https://doi.org/10.1093/cercor/bhg124

Gross, J., Kujala, J., Hamalainen, M., Timmermann, L., Schnitzler, A., & Salmelin, R. (2001). Dynamic imaging of coherent sources: Studying neural interactions in the human brain. *Proceedings of the National Academy of Sciences*, *98*(2), 694–699. https://doi.org/10.1073/pnas.98.2.694

Hagoort, P. (2003). How the brain solves the binding problem for language: A neurocomputational model of syntactic processing. *NeuroImage*, *20*, S18–S29. https://doi.org/10.1016/j.neuroimage.2003.1796

Hagoort, P. (2005). On Broca, brain, and binding: A new framework. *Trends in Cognitive Sciences*, *9*(9), 416–423. https://doi.org/10.1016/j.tics.2005.07.004

Hagoort, P. (2019). The neurobiology of language beyond single-word processing. *Science*, *366*(6461), 55–58. https://doi.org/10.1126/science.aax0289

Humphries, C., Binder, J. R., Medler, D. A., & Liebenthal, E. (2006). Syntactic and semantic modulation of neural activity during auditory sentence comprehension. *Journal of Cognitive Neuroscience*, *18*(4), 665–679.

Jensen, O., Gips, B., Bergmann, T. O., & Bonnefond, M. (2014). Temporal coding organized by coupled alpha and gamma oscillations prioritize visual processing. *Trends in Neurosciences*, *37*(7), 357–369. https://doi.org/10.1016/j.tins.2014.04.001

Jensen, O., & Mazaheri, A. (2010). Shaping functional architecture by oscillatory alpha activity: Gating by inhibition. *Frontiers in Human Neuroscience*, *4*, 186. https://doi.org/10.3389/fnhum.2010.00186

Kim, S., & Pylkkänen, L. (2019). Composition of event concepts: Evidence for distinct roles for the left and right anterior temporal lobes. *Brain and Language*, *188*, 18–27. https://doi.org/10.1016/j.bandl.2018.11.003

Klimesch, W. (2012). Alpha-band oscillations, attention, and controlled access to stored information. *Trends in Cognitive Sciences*, *16*(12), 606–617. https://doi.org/10.1016/j.tics.2012.10.007

Kochari, A. R., Lewis, A. G., Schoffelen, J.-M., & Schriefers, H. (2021). Semantic and syntactic composition of minimal adjective-noun phrases in Dutch: An MEG study. *Neuropsychologia*, *155*, 107754. https://doi.org/10.1016/j.neuropsychologia.2021.107754

Kuperberg, G. R., & Jaeger, T. F. (2016). What do we mean by prediction in language comprehension? *Language, Cognition and Neuroscience*, *31*(1), 32–59. https://doi.org/10.1080/23273798.2015.1102299

Kutas, M., DeLong, K. A., & Smith, N. J. (2011). A look around at what lies ahead: Prediction and predictability in language processing. In M. Bar (Ed.), *Predictions in the brain: Using our past to generate a future* (pp. 190–207). Oxford University Press.

Lachaux, J.-P., Rodriguez, E., Martinerie, J., & Varela, F. J. (1999). Measuring phase synchrony in brain signals. *Human Brain Mapping*, *8*(4), 194–208. https://doi.org/10.1002/(SICI)1097-0193(1999)8:4<194::AID-HBM4>3.0.CO;2-C

Lam, N. H. L., Schoffelen, J. M., Uddén, J., Hultén, A., & Hagoort, P. (2016). Neural activity during sentence processing as reflected in theta, alpha, beta, and gamma oscillations. *NeuroImage*, *142*, 43–54. https://doi.org/10.1016/j.neuroimage.2016.03.007

Lopes da Silva, F. (2013). EEG and MEG: Relevance to neuroscience. *Neuron*, *80*(5), 1112–1128. https://doi.org/10.1016/j.neuron.2013.10.017

Lopopolo, A., van den Bosch, A., Petersson, K.-M., & Willems, R. M. (2021). Distinguishing syntactic operations in the brain: Dependency and phrase-structure parsing. *Neurobiology of Language*, *2*(1), 152–175. https://doi.org/10.1162/nol_a_00029

Magazzini, L., Ruhnau, P., & Weisz, N. (2016). Alpha suppression and connectivity modulations in left temporal and parietal cortices index partial awareness of words. *NeuroImage*, *133*, 279–287. https://doi.org/10.1016/j.neuroimage.2016.03.025

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data. *Journal of Neuroscience Methods*, *164*(1), 177–190. https://doi.org/10.1016/j.jneumeth.2007.03.024

Matchin, W., Hammerly, C., & Lau, E. (2017). The role of the IFG and pSTS in syntactic prediction: Evidence from a parametric study of hierarchical structure in fMRI. *Cortex*, *88*, 106–123. https://doi.org/10.1016/j.cortex.2016.12.010

Matchin, W., & Hickok, G. (2020). The cortical organization of syntax. *Cerebral Cortex*, *30*(3), 1481–1498. https://doi.org/10.1093/cercor/bhz180

Mazaheri, A., van Schouwenburg, M. R., Dimitrijevic, A., Denys, D., Cools, R., & Jensen, O. (2014). Region-specific modulations in oscillatory alpha activity serve to facilitate processing in the visual and auditory modalities. *NeuroImage*, *87*, 356–362. https://doi.org/10.1016/j.neuroimage.2013.10.052

Menenti, L., Gierhan, S. M. E., Segaert, K., & Hagoort, P. (2011). Shared language: Overlap and segregation of the neuronal infrastructure for speaking and listening revealed by functional MRI. *Psychological Science*, *22*(9), 1173–1182. https://doi.org/10.1177/0956797611418347

Meyer, L. (2018). The neural oscillations of speech processing and language comprehension: State of the art and emerging mechanisms. *European Journal of Neuroscience*, *48*(7), 2609–2621. https://doi.org/10.1111/ejn.13748

Meyer, L., Obleser, J., & Friederici, A. D. (2013). Left parietal alpha enhancement during working memory-intensive sentence processing. *Cortex*, *49*(3), 711–721. https://doi.org/10.1016/j.cortex.2012.03.006

Murphy, E. (2015). The brain dynamics of linguistic computation. *Frontiers in Psychology*, *6*, 1515. https://doi.org/10.3389/fpsyg.2015.01515

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J. M. (2011). FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational Intelligence and Neuroscience*, 156869. https://doi.org/10.1155/2011/156869

Pallier, C., Devauchelle, A. D., & Dehaene, S. (2011). Cortical representation of the constituent structure of sentences. *Proceedings of the National Academy of Sciences*, *108*(6), 2522–2527. https://doi.org/10.1073/pnas.1018711108

Piai, V., Roelofs, A., & Maris, E. (2014). Oscillatory brain responses in spoken word production reflect lexical frequency and sentential constraint. *Neuropsychologia*, *53*(1), 146–156. https://doi.org/10.1016/j.neuropsychologia.2013.11.014

Poulisse, C., Wheeldon, L., Limachya, R., Mazaheri, A., & Segaert, K. (2020). The oscillatory mechanisms associated with syntactic binding in healthy ageing. *Neuropsychologia*, *146*, 107523. https://doi.org/10.1016/j.neuropsychologia.2020.107523

Poulisse, C., Wheeldon, L., & Segaert, K. (2019). Evidence against preserved syntactic comprehension in healthy aging. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *45*(12), 2290–2308. http://dx.doi.org/10.1037/xlm0000707

Prystauka, Y., & Lewis, A. G. (2019). The power of neural oscillations to inform sentence comprehension: A linguistic perspective. *Language and Linguistics Compass*, *13*(9), e12347. https://doi.org/10.1111/lnc3.12347

Pylkkänen, L. (2019). The neural basis of combinatory syntax and semantics. *Science*, *366*(6461), 62–66. https://doi.org/10.1126/science.aax0050

Pylkkänen, L., Bemis, D. K., & Blanco-Elorrieta, E. (2014). Building phrases in language production: An MEG study of simple composition. *Cognition*, *133*(2), 371–384. https://doi.org/10.1016/j.cognition.2014.07.001

Roland, D., Dick, F., & Elman, J. L. (2007). Frequency of basic English grammatical structures: A corpus analysis. *Journal of Memory and Language*, *57*(3), 348–379. https://doi.org/10.1016/j.jml.2007.03.002

Rommers, J., Dickson, D. S., Norton, J. J. S., Wlotko, E. W., & Federmeier, K. D. (2017). Alpha and theta band dynamics related to sentential constraint and word expectancy. *Language, Cognition and Neuroscience*, *32*(5), 576–589. https://doi.org/10.1080/23273798.2016.1183799

Ross, B., Tremblay, K. L., & Alain, C. (2020). Simultaneous EEG and MEG recordings reveal vocal pitch elicited cortical gamma oscillations in young and older adults. *NeuroImage*, *204*, 116253. https://doi.org/10.1016/j.neuroimage.2019.116253

Sadaghiani, S., & Kleinschmidt, A. (2016). Brain networks and α-Oscillations: Structural and functional foundations of cognitive control. *Trends in Cognitive Sciences*, *20*(11), 805–817. https://doi.org/10.1016/j.tics.2016.09.004

Sarubbo, S., Tate, M., De Benedictis, A., Merler, S., Moritz-Gasser, S., Herbet, G., & Duffau, H. (2020). Mapping critical cortical hubs and white matter pathways by direct electrical stimulation: An original functional atlas of the human brain. *NeuroImage*, *205*, 116237. https://doi.org/10.1016/j.neuroimage.2019.116237

Schell, M., Zaccarella, E., & Friederici, A. D. (2017). Differential cortical contribution of syntax and semantics: An fMRI study on two-word phrasal processing. *Cortex*, *96*, 105–120. https://doi.org/10.1016/j.cortex.2017.09.002

Schmidt, B. T., Ghuman, A. S., & Huppert, T. J. (2014). Whole brain functional connectivity using phase locking measures of resting state magnetoencephalography. *Frontiers in Neuroscience*, *8*. https://doi.org/10.3389/fnins.2014.00141

Schneider, W., Eschman, A., & Zuccolotto, A. (2002). *E-prime user's guide*. Psychology Software Tools Inc.

Schoffelen, J.-M., Hultén, A., Lam, N., Marquand, A. F., Uddén, J., & Hagoort, P. (2017). Frequency-specific directed interactions in the human brain network for language. *Proceedings of the National Academy of Sciences*, *114*(30), 8083–8088. https://doi.org/10.1073/pnas.1703155114

Segaert, K., Mazaheri, A., & Hagoort, P. (2018). Binding language: Structuring sentences through precisely timed oscillatory mechanisms. *European Journal of Neuroscience*, *48*(7), 2651–2662. https://doi.org/10.1111/ejn.13816

Segaert, K., Menenti, L., Weber, K., Petersson, K. M., & Hagoort, P. (2012). Shared syntax in language production and language comprehension—An fMRI study. *Cerebral Cortex*, *22*(22), 1662–1670. https://doi.org/10.1093/cercor/bhr249

Snijders, T. M., Vosse, T., Kempen, G., Van Berkum, J. J. A., Petersson, K. M., & Hagoort, P. (2008). Retrieval and unification of syntactic structure in sentence comprehension: An fMRI study using word-category ambiguity. *Cerebral Cortex*, *19*(7), 1493–1503. https://doi.org/10.1093/cercor/bhn187

Strijkers, K., Chanoine, V., Munding, D., Dubarry, A.-S., Trébuchon, A., Badier, J.-M., & Alario, F.-X. (2019). Grammatical class modulates the (left) inferior frontal gyrus within 100 milliseconds when syntactic context is predictive. *Scientific Reports*, *9*(1), 4830. https://doi.org/10.1038/s41598-019-41376-x

Tyler, L. K., & Marslen-Wilson, W. (2008). Fronto-temporal brain systems supporting spoken language comprehension. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *363*(1493), 1037–1054. https://doi.org/10.1098/rstb.2007.2158

Tzourio-Mazoyer, N., Landeau, B., Papathanassiou, D., Crivello, F., Etard, O., Delcroix, N., Mazoyer, B., & Joliot, M. (2002). Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *NeuroImage*, *15*(1), 273–289. https://doi.org/10.1006/nimg.2001.0978

Uddén, J., Hultén, A., Schoffelen, J.-M., Lam, N., Harbusch, K., Bosch, A. van den, Kempen, G., Petersson, K. M., & Hagoort, P. (2019). Supramodal sentence processing in the human brain: FMRI evidence for the influence of syntactic complexity in more than 200 participants. *BioRxiv*, 576769. https://doi.org/10.1101/576769

Ullman, M. T., Corkin, S., Coppola, M., Hickok, G., Growdon, J. H., Koroshetz, W. J., & Pinker, S. (1997). A neural dissociation within language: Evidence that the mental dictionary is part of declarative memory, and that grammatical rules are processed by the procedural system. *Journal of Cognitive Neuroscience*, *9*(2), 266–276. https://doi.org/10.1162/jocn.1997.9.2.266

van Diepen, R. M., Foxe, J. J., & Mazaheri, A. (2019). The functional role of alpha-band activity in attentional processing: The current zeitgeist and future outlook. *Current Opinion in Psychology*, *29*, 229–238. https://doi.org/10.1016/j.copsyc.2019.03.015

van Diepen, R. M., & Mazaheri, A. (2017). Cross-sensory modulation of alpha oscillatory activity: Suppression, idling, and default resource allocation. *European Journal of Neuroscience*, *45*(11), 1431–1438. https://doi.org/10.1111/ejn.13570

van Diepen, R. M., & Mazaheri, A. (2018). The caveats of observing inter-trial phase-coherence in cognitive neuroscience. *Scientific Reports*, *8*(1), 2990. https://doi.org/10.1038/s41598-018-20423-z

Vassileiou, B., Meyer, L., Beese, C., & Friederici, A. D. (2018). Alignment of alpha-band desynchronization with syntactic structure predicts successful sentence comprehension. *NeuroImage*, *175*, 286–296. https://doi.org/10.1016/j.neuroimage.2018.04.008

Wang, L., Hagoort, P., & Jensen, O. (2017). Language prediction is reflected by coupling between frontal gamma and posterior alpha oscillations. *Journal of Cognitive Neuroscience*, *30*(3), 432–447. https://doi.org/10.1162/jocn_a_01190

Wang, L., Jensen, O., Van den Brink, D., Weder, N., Schoffelen, J. M., Magyari, L., Hagoort, P., & Bastiaansen, M. (2012). Beta oscillations relate to the N400m during language comprehension. *Human Brain Mapping*, *33*(12), 2898–2912. https://doi.org/10.1002/hbm.21410

Westerlund, M., Kastner, I., Al Kaabi, M., & Pylkkänen, L. (2015). The LATL as locus of composition: MEG evidence from English and Arabic. *Brain and Language*, *141*, 124–134. https://doi.org/10.1016/j.bandl.2014.12.003

White, E. J., Nayman, C., Dunkley, B. T., Keller, A. E., Valiante, T. A., & Pang, E. W. (2018). Addressing the language binding problem with dynamic functional connectivity during meaningful spoken language comprehension. *Frontiers in Psychology*, *9*. https://doi.org/10.3389/fpsyg.2018.01960

Whitmarsh, S., Nieuwenhuis, I. L. C., Barendregt, H. P., & Jensen, O. (2011). Sensorimotor alpha activity is modulated in response to the observation of pain in others. *Frontiers in Human Neuroscience*, *5*, 91. https://doi.org/10.3389/fnhum.2011.00091

Wilson, S. M., DeMarco, A. T., Henry, M. L., Gesierich, B., Babiak, M., Mandelli, M. L., Miller, B. L., & Gorno-Tempini, M. L. (2014). What role does the anterior temporal lobe play in sentence-level processing? Neural correlates of syntactic processing in semantic variant primary progressive aphasia. *Journal of Cognitive Neuroscience*, *26*(5), 970–985. https://doi.org/10.1162/jocn_a_00550

Zaccarella, E., & Friederici, A. D. (2015). Merge in the human brain: A sub-region based functional investigation in the left pars opercularis. *Frontiers in Psychology*, *6*, 1818. https://doi.org/10.3389/fpsyg.2015.01818

Zaccarella, E., Meyer, L., Makuuchi, M., & Friederici, A. D. (2017). Building by syntax: The neural basis of minimal linguistic structures. *Cerebral Cortex*, *27*(1), 411–421. https://doi.org/10.1093/cercor/bhv234

Zhang, L., & Pylkkänen, L. (2015). The interplay of composition and concept specificity in the left anterior temporal lobe: An MEG study. *NeuroImage*, *111*, 228–240. https://doi.org/10.1016/j.neuroimage.2015.02.028

Zhang, L., & Pylkkänen, L. (2018). Composing lexical versus functional adjectives: Evidence for uniformity in the left temporal lobe. *Psychonomic Bulletin & Review*, *25*(6), 2309–2322. https://doi.org/10.3758/s13423-018-1469-y

Ziegler, J., & Pylkkänen, L. (2016). Scalar adjectives and the temporal unfolding of semantic composition: An MEG investigation. *Neuropsychologia*, *89*, 161–171. https://doi.org/10.1016/j.neuropsychologia.2016.06.010

Zouridakis, G., Simos, P. G., & Papanicolaou, A. C. (1998). Multiple bilaterally asymmetric cortical sources account for the auditory N1m component. *Brain Topography*, *10*(3), 183–189. https://doi.org/10.1023/A:1022246825461

Zumer, J. M., Scheeringa, R., Schoffelen, J.-M., Norris, D. G., & Jensen, O. (2014). Occipital alpha activity during stimulus processing gates the information flow to object-selective cortex. *PLoS Biology*, *12*(10), e1001965. https://doi.org/10.1371/journal.pbio.1001965