1   **Neurons in the Nucleus Tractus Solitarius are Selective to the Orientation of Gastric**

2   **Electrical Stimulation**

4   Jiayue Cao[1], Xiaokai Wang[1], Terry L. Powley[3], Zhongming Liu[*1,2]

5   [1]Department of Biomedical Engineering, University of Michigan Ann Arbor

6   [2]Department of Electrical Engineering and Computer Science, University of Michigan Ann Arbor

7   [3]Department of Psychological Sciences, Purdue University West Lafayette

* Correspondence

Zhongming Liu, Ph.D.

Associate Professor

Room 2119, Carl A Gerstacker Building

2200 Bonisteel Blvd.

Ann Arbor, MI 48109

Phone: (734)764-2073

Email: zmliu@umich.edu

Abstract

Gastric electrical stimulation (GES) is a bioelectric intervention for gastroparesis, obesity, and other functional gastrointestinal disorders. In a potential mechanism of action, GES activates the nerve endings of vagal afferent neurons and induces the vago-vagal reflex through the nucleus tractus solitarius (NTS) in the brainstem. However, it is unclear where and how to stimulate in order to optimize the vagal afferent responses. To address this question with electrophysiology in rats, we applied mild electrical currents to two serosal targets on the distal forestomach with dense distributions of vagal intramuscular arrays that innervated the circular and longitudinal smooth muscle layers. During stimulation, we recorded single and multi-unit responses in NTS and evaluated how the recorded responses depended on the stimulus orientation and amplitude. We found that NTS responses were highly selective to the stimulus orientation for a range of stimulus amplitudes. The strongest responses were observed when the applied current flowed in the same direction as the intramuscular arrays in parallel with the underlying smooth muscle fibers. Our results suggest that gastric neurons in NTS may encode the orientation-specific activity of gastric smooth muscles relayed by vagal afferent neurons. This finding suggests that the orientation of GES is critical to effective engagement of vagal afferents and should be considered in light of the structural phenotypes of vagal terminals in the stomach.

Significance Statement

The brain monitors and regulates the stomach in part through vagal neural pathways. Although there has

been increasing progress describing the structural phenotypes of vagal projections to the stomach, very

little is known about the specific gastric information transmitted to and encoded by the brain. In this

study, we report that gastric neurons in the nucleus tractus solitarius encode the orientation-specific

activity of gastric smooth muscles. This neural code is relayed through vagal afferent neurons with their

nerve terminals along circular or longitudinal muscle fibers. This finding provides a critical foundation

for understanding gastric interoception. It brings up a new consideration that the orientation of gastric

electrical stimulation is critical to engaging vagal afferent pathways for modulating gastric functions.

Introduction

The vagus is the primary neural pathway that connects the brain and the stomach (Browning & Travagli, 2011; Levinthal & Strick, 2020; Powley et al., 2019). It enables rapid stomach-brain interactions (Andermann & Lowell, 2017; Carabotti et al., 2015; Furness and Stebbing, 2018; Holtmann & Talley, 2014; Mayer, 2011; Williams et al., 2016) and reflexive control of the gastrointestinal tract (Babic & Browning, 2014; Travagli & Anselmi, 2016). Neural signaling via the vagus is bi-directional. The afferents ascend sensory signals to the nucleus tractus solitarius (NTS) (Rogers et al., 1995; Powley & Phillips, 2002). The efferents descend motor commands from the dorsal motor nucleus of the vagus (DMV) (Travagli et al., 2006). Despite progress in structural characterization of vagal projections to the stomach (Brookes et al., 2013; Powley et al., 2019), very little is known about the specific gastric information transmitted to and encoded by the brain. In particular, it is unclear how NTS, as the first nucleus for visceral sensation, represents and integrates gastric information relayed through vagal afferents.

Both vagal afferents and efferents innervate the stomach with extensive and specialized terminals in the stomach wall (Brookes et al., 2013; Furness et al., 2020; Powley & Phillips, 2002). The afferent terminals exhibit distinctive morphologies and chemical codes sensitized to various gastric signals (Berthoud et al., 1997; Berthoud & Powley, 1992; Powley et al., 2019). In particular, the intramuscular array (IMA) runs in parallel with muscle fibers and serves as mechanoreceptors to report on the stretch (or length) of gastric smooth muscles (Berthoud & Powley, 1992; Wang & Powley, 2000; Phillips & Powley, 2000; Powley et al., 2011). It has two subtypes, namely longitudinal and circular IMAs, which target longitudinal and circular muscles, respectively. Both subtypes show regional distributions overlapping in the antrum but distinctive in the forestomach (or fundus) (Powley et al., 2016; Tan et al., 2021).

Stimulation targeted to vagal afferent terminals is one mechanism of gastric electrical stimulation (GES) (Cheng et al., 2021). Electrical current pulses with a low amplitude, high frequency, and short duration

are inadequate to activate the muscle or pace the stomach, but can potentially relieve nausea and vomiting

and treat motor dysfunctions for drug-resistant gastroparesis (Abell et al., 2003; Lal et al., 2015; Levinthal

& Bielefeldt, 2017; McCallum et al., 2010; O'Grady et al., 2009). Despite its approved use in clinics, the

efficacy of GES varies across patients and its parameterization remains empirical and suboptimal. A

potential strategy to optimize GES is to engage the central nervous system via vagal afferent signaling

(Gottfried-Blackmore et al., 2020; Mayer et al., 2006; McCallum et al., 2010; Payne et al., 2019; Qin et

al., 2005; Tang et al., 2006). However, it is unclear where and how to apply GES in order to maximize the

vagal afferent responses.

In this study, we targeted GES to longitudinal and/or circular IMAs and recorded neuronal responses from

NTS. Since IMAs were orientation specific (Powley et al., 2016), we hypothesized that the GES-evoked

vagal afferent responses were dependent on the stimulus orientation. To test this hypothesis in rats, we

chose two serosal targets in the distal forestomach with high IMA densities and applied short current

stimuli with different orientations and amplitudes. We recorded the neuronal spiking responses in NTS

and evaluated the responses across various stimulus settings. By testing this hypothesis, we aimed to

characterize the functional specialization of gastric neurons in NTS and to offer additional clues as to how

to apply GES in order to modulate the stomach-brain neuroaxis.

Materials and Methods

Subjects

This study used seven Sprague–Dawley rats (male, weight: 250–400 g; Envigo RMS, Indianapolis, IN)

according to a protocol approved by the Institutional Animal Care and Use Committee and the Laboratory

Animal Program at Purdue University. All animals were housed in a strictly controlled environment

(temperature: 21 ± 1 °C; 12 h light-dark cycle with light on at 6:00 a.m. and off at 6:00 p.m.). Every

animal received acute surgical implantation of patch electrodes for gastric electrical stimulation and depth

electrodes for neuronal recordings in the brainstem.

Gastric electrical stimulation

Every animal received abdominal surgery to be implanted with patch electrodes on the serosal surface of

the distal forestomach. The animal was initially anesthetized with 5% isoflurane mixed with oxygen (flow

rate: 1 L/min). The dose of isoflurane was then reduced to 2% to keep a surgical plane. Then, the animal

was placed in a supine position on the surgical bench. Following a toe-pinch test, an incision was made

starting from the xiphoid and moving 4 cm caudally. Skin and muscle layers were retracted and separated

to expose the ventral forestomach.

Patch electrodes (Microprobes, Gaithersburg, MD, USA) were sutured at two target locations, namely

Target 1 and 2, on the serosal surface of the distal forestomach (Figure 1A). Target 1 was on the greater

curvature and about 4 mm away from the limiting ridge. Target 2 was on the ventral stomach, 2 mm away

from the limiting ridge, and halfway in between the lesser curvature and the greater curvature. According

to our previously mapped distributions of longitudinal and circular IMAs (Powley et al., 2016; Tan et al.,

2021), Target 1 had a high density of longitudinal IMAs but a low density of circular IMAs, whereas

Target 2 had a high density of circular IMAs but a low density of longitudinal IMAs (Figure 1B). Target

1 and Target 2 were chosen as complementary targets selective to the "hotspots" of longitudinal and

circular IMA, respectively.

Each patch electrode consisted of either one or two bipolar pairs of contacts. Each contact was made of a

Pt/Ir foil (12 mm^2) on a thin perylene substrate (77 mm^2) and was positioned 2 mm away from the center

of the substrate. Each bipolar pair was oriented for current delivery along either the longitudinal or

circular direction. We attached two pairs of electrodes on Target 1 for four animals, two pairs of

electrodes on Target 2 for four animals, and one pair of electrodes on Target 1 for three animals (Table 1).

We fixed the electrodes by securing the four corners of the substrate on the serosal layer. After

implantation of the electrodes, the muscle layer and the skin were closed with sutures, while the leads

were kept outside of the abdomen and were connected to a current stimulator (model 2200, A-M Systems,

Sequim, USA).

For each target location, electrical current was delivered in four plausible orientations: longitudinal

pointing towards the proximal (0°) or distal stomach (180°) with one pair of electrodes, or circular

pointing toward the lesser (90°) or greater (270°) curvature with the other pair of electrodes (Figure 2B).

Current stimuli were delivered as a train of rectangular pulses (pulse width: 0.3 ms, pulse amplitude: 0.6,

0.8, or 1 mA with alternating polarity, and inter-stimulus-interval: 100 ms for two animals and 250 ms for

the other five animals). A pulse train lasted 10 s followed by a stimulus-free rest period of >20 s.

Different pulse trains were configured to stimulate in either the longitudinal or circular direction in an

order randomized and counterbalanced across animals.

Neurophysiological recording

To record neuronal spiking activity, we implanted a 32-channel silicon probe (Neuronexus, Ann Arbor,

USA) in NTS for each animal. The surgery was performed by placing the animal in the prone position on

a stereotaxic frame (Stoelting Co., Wool Dale, IL, USA). To expose the skull, a midline skin incision was

made from between the eyes to the neck. Two stainless steel screws were drilled into the skull above the

olfactory bulb and were used as the ground and reference. A 5x5 mm^2 cranial window was created at the

bottom of the occipital bone to expose the brainstem until the obex was identifiable.

The silicon probe was inserted at about 0-0.8mm left and 0-0.2mm rostral to the obex (Figure 1C). The

depth of insertion was determined by using a searching process to identify gastric neurons in NTS.

Specifically, gastric stimulation was delivered (pulse width = 0.3 ms, current = 1.0 mA, pulse frequency =

1 Hz) while the probe was being inserted into NTS. The insertion was stopped once the stimulus-evoked

spiking activity was observable in one or multiple channels. The searching process was repeated multiple

times to identify different groups of neurons, while each time the probe was pulled out and re-inserted to

different depths (from 0.5 to 1.3 mm). At each depth of insertion, neural data was recorded from all 32

channels by using a broadband recording system (Tucker Davis Technologies, Alachua, USA) with a

sampling rate of 24 kHz.

Single and multi-unit firing rate

We applied a high-pass filter (>300 Hz) to the raw signals to extract the neuronal spiking activity. First,

unsorted spikes were identified if they surpassed a threshold (between 50 and 100 **μ**V) defined for each

animal. The unsorted spikes were stored by channel as the multi-unit activity (MUA). Second, spike

sorting was applied separately to each channel to extract the single-unit activity (SUA). The spike sorting

extracted from each spike two features based on the peak-trough differences: the difference between the

first peak and the trough (Feature 1) and the difference between the second peak and the trough (Feature

2). As illustrated in Figure 1F, the spike features were used to group every spike into distinct clusters

where each cluster corresponded to a neuron. The clustering was based on the k-means algorithm (kmeans

in Matlab), where the distance between spikes was their Euclidean distance in the feature space. The

number of clusters was determined based on the distributions of spike representations in the feature space.

The sorted spikes across all channels were stored as SUA.

Based on the SUA or MUA, we evaluated the neuronal responses to GES in the level of either single or

multiple units, respectively. For each neuron, the SUA response was quantified as the number of spikes

within a 100-ms post-stimulus period in each trial of GES parameterized by its location (Target 1 & 2),

direction (0^o, 90^o, 180^o, 270^o), and amplitude (0.6, 0.8, 1 mA). The single-trial SUA response was

averaged across repeated stimuli with the same parameters and then across animals. Similarly, the time-

averaged MUA response was quantified for each channel.

For a given location and amplitude of GES, the time-averaged MUA response was compared across four

stimulus orientations (0^o, 90^o, 180^o, 270^o). The orientation that resulted in the highest MUA response was

considered as the preferred orientation for each channel under evaluation. Then, the MUA response was

averaged across channels and animals, to obtain the group averaged NTS responses as a function of the

current amplitude (0.6, 0.8, 1 mA).

In addition to the time-averaged SUA or MUA response, we further evaluated the response as a function

of the post-stimulus latency. The time-resolved response was based on the post-stimulus time histogram

(PSTH). Specifically, the number of spikes was calculated within a 5 ms sliding window stepping by one

time point (1/24 kHz) until reaching 100 ms after the stimulus. The firing rate (i.e., the number of spikes

per 5ms) was averaged across repeated trials of the same stimulus setting. Neurons (for SUA) or channels

(for MUA) that showed their maximal PSTH less than 0.05 spikes per 5 ms were discarded and excluded

from subsequent analyses. Further, the PSTH was averaged within four coarsely defined post-stimulus

periods: 0-13 ms, 13-29 ms, 29-73 ms, and 73-100 ms.

Statistical analysis and orientation selectivity

We used a paired t-test to evaluate the statistical significance of any response difference given GES

applied in any two different orientations. The t-test was separately performed for the mean SUA or MUA

response averaged over the 100-ms post-stimulus period. Bonferroni correction was used to correct for

multiple comparisons (i.e., 6 pairs of different orientations under comparison). The significance level α

was set as 0.05. Similarly, we also used a paired t-test to evaluate the difference between GES applied to

two different locations (i.e., Target 1 and 2).

We used an orientation selectivity index (OSI) to measure how a neuron (SUA) or neuronal ensemble

(MUA) was selective to GES applied in the preferred orientation vs. other orientations. Specifically, let

N_{PO} be the firing rate in response to GES in the preferred orientation (e.g., 0^o), and let N_i (i=1 to 3) be the

firing rate in response to GES in the non-preferred orientations (e.g., 90^o, 180^o, 270^o). The OSI was the

average difference in the firing rate between the preferred orientation and every non-preferred orientation,

as expressed by Equation 1. The OSI was in a range from 0 to 1, while negative values were set to 0. A

higher value of OSI indicated that the neuron or channel under evaluation was more selective to the GES

in the preferred orientation.

$$OSI = \frac{1}{3}\sum_{i=1}^{3}(N_{po} - N_i)$$ (1)

We used a permutation test to evaluate the statistical significance of the OSI. Specifically, the SUA or

MUA responses were randomly shuffled into different groups defined by the stimulus orientation. The

shuffled dataset disrupted any real response-stimulus association. After repeating the process 100,000

times and at each time evaluating the OSI with the shuffled dataset, we generated a null distribution of

OSI, against which the actual OSI was compared for calculation of the *p-value*.

Results

In seven rats, we recorded single and multi-unit activity from gastric neurons in NTS while applying GES

to two complementary regions of interest in the distal forestomach (Figure 1A): Target 1 with a high

density of longitudinal IMA (Figure 1B left) and Target 2 with a high density of circular IMA (Figure 1B

right), according to recent studies (Powley et al., 2016; Tan et al., 2021). We evaluated the GES-evoked

neuronal responses as a function of the stimulus orientation, location, and amplitude. In particular, our

emphasis was on the orientation selectivity of NTS. This was motivated by the fact that the IMA, as an

array of vagal afferent receptors (Berthoud and Powley, 1992), typically shows a parent arbor branching

into elongated and parallel receptors directly innervating muscle fibers (Wang and Powley, 2000). Figure

1B middle shows a digitized example of IMA. Figure 2A and 2B further show the topographic

distributions of longitudinal and circular IMA samples (Powley et al., 2016), respectively.

Orientation dependence of NTS population responses to GES

When GES was delivered as a current pulse with a short width (0.3 ms) and a low amplitude (<1 mA), the stimuli were too mild to directly pace the stomach or induce muscle contractions (Du et al., 2009; Li & Chen, 2010; Cheng et al., 2021; Forster et al., 2001; McCallum et al., 1998). However, such GES could evoke population spikes (Figure 1D & E) observable with a 32-channel silicon probe inserted into NTS (Figure 1C). Figure 1D shows a typical MUA recording in NTS during a train of alternating currents delivered to Target 1 in two opposing directions (i.e., positive and negative pulses). In this example, many more spikes were observed following the negative pulses than following the positive pulses (Figure 1D). The spikes occurred shortly after each negative pulse (Figure 1E). By spike sorting (Figure 1F), two neurons were identified (Figure 1G) and their responses were observed mostly in the first 20 ms following the pulse (Figure 1H). This observation led us to hypothesize that gastric neurons in NTS were selective to the orientation of GES.

We extended the above observation by performing quantitative comparisons across four orientations and two locations, which covered the non-overlapping "hotspots" (Figure 1B) of longitudinal and circular IMAs shown in Figure 2A and 2D, respectively. For Target 1, when a 1mA current stimulus was applied in 0^o (i.e., flowing in the longitudinal direction towards the proximal stomach), the stimulus induced multiple population spikes with spike timing highly consistent across repeated trials (Figure 2B). For an example MUA recorded from one channel, one spike occurred around 10 ms; a few spikes occurred from 40 to 60 ms. When the current stimuli were applied in other directions, fewer spikes were evoked and were less consistent across trials (Figure 2B). Beyond this example, the population response in NTS, in terms of the number of spikes per 100 ms averaged across channels and animals, was 4.09 ± 0.50 (mean $\pm$ sem) for GES at 0^o, which was significantly higher than those for other orientations (2.52 ± 0.45 for 270^o; 2.26 ± 0.37 for 180^o; 1.88 ± 0.34 for 90^o), as shown in Figure 2C. In summary, 1-mA GES at Target 1 evoked reliable population spikes in NTS when the stimulus was delivered in 0^o (longitudinal towards the proximal stomach).

Similarly, for Target 2, we observed much stronger and more reliable population responses when 1-mA

GES was applied in 270^{o} (circular towards the greater curvature) relative to other orientations. Given the

preferred orientation, Figure 2E shows an example of the population spikes observed from one channel.

The spike timing was highly consistent across repeated trials. The population spikes were fewer and much

less consistent given GES in non-preferred orientations. The mean firing rate was 3.83 ± 0.39 per 100 ms

(mean $\pm$ sem) for GES at 270^{o}, which was significantly higher than those for other orientations (1.56 ± 0.17

for 0^{o}; 1.99 ± 0.25 for 180^{o}; 1.56 ± 0.19 for 90^{o}) (Figure 2F).

Effects of the current amplitude

We asked whether the orientation selectivity of NTS neurons observed with 1-mA GES was generalizable

to GES with lower amplitudes. To address this question, we compared the neuronal population responses

in NTS given GES with three different current amplitudes (0.6, 0.8, and 1 mA). For both Target 1 (Figure

3A) and Target 2 (Figure 3D), the time-averaged population responses in NTS were always greater when

GES was applied in its preferred orientation (0^{o} for Target 1, 270^{o} for Target 2), despite the difference in

the current amplitude used in this study (Figure 3B & E). In its preferred orientation, a stimulus with a

higher amplitude tended to induce stronger population responses in NTS. Stimuli in other non-preferred

orientations did not show significant amplitude-dependent effects on the resulting population responses.

We further calculated the orientation selectivity index (OSI) as the average difference in the neuronal

response to GES between the preferred and non-preferred orientations described above. For both Target 1

and 2, increasing the current amplitude led to more pronounced orientation selectivity (i.e., increasingly

higher OSI) (Figure 3C). For Target 1, the OSI at 0.6 mA was significantly lower than the OSI at 0.8 mA

($p<0.005$) or 1 mA ($p<0.001$) based on a permutation test with Bonferroni correction. The observations

and statistics were similar for Target 2 (Figure 3F). See Table 2 and Table 3 for statistical results in

further details. In summary, orientation selectivity was generalizable across different current amplitudes

(0.6, 0.8, and 1 mA).

Orientation selectivity of single-unit responses

We further tested whether the orientation selectivity was applicable to the level of single neurons in NTS.

We applied spike sorting to identify individual neurons activated by GES and analyzed their firing rates

as a function of the stimulus orientation and location. In total, we identified 106 neurons activated by

GES at Target 1 and 119 neurons by GES at Target 2.

Most neurons activated by GES manifested high selectivity for one stimulus orientation. In Figure 4B, an

example neuron showed reliable responses only when GES at Target 1 was in 0^o, whereas virtually no

responses were observed for other orientations. In total, 67 out of 106 neurons all shared the same

preferred orientation (i.e., 0^o) and showed significantly higher firing rates given GES in 0^o than in other

orientations (Figure 4C). Neurons with a preference for other orientations (i.e., 270^o, 180^o, 90^o) were not

only fewer but also less orientation selective (i.e., significantly lower OSI) (Figure 4D & E). Similarly for

GES at Target 2, 86 out of 119 neurons were highly selective to the same preferred orientation of 270^o

(Figure 4F through J). In summary, gastric neurons in NTS were selective to 0^o (longitudinal towards the

proximal stomach) for GES at Target 1 and were selective to 270^o (circular towards the greater curvature)

for GES at Target 2.

Orientation selectivity at different post-stimulus latencies

The example in Figure 1E shows that GES evoked stronger and more reliable neuronal responses at some

post-stimulus latencies than others. We asked whether the orientation selectivity was dependent on the

response latency. To answer this question, we calculated the PSTH to resolve SUA in time and evaluated

the time dependence of orientation selectivity (Figure 5). When Target 1 was stimulated in 0^o, the time-

resolved firing rate (PSTH) of the activated gastric neurons was on average higher around 10 and 43 ms

than around 20 ms or after 73 ms (Figure 5B). The neuronal responses to GES in other orientations did

not demonstrate similarly strong time dependence (Figure 5B).

According to the PSTH, we further divided the post-stimulus period into four periods (0-13, 13-29, 29-73,

73-100 ms) and evaluated the orientation selectivity of individual neurons separately for each period. Out

of 106 neurons activated by GES at Target 1, 77 neurons were activated from 0 to 13 ms, 65 neurons from

13 to 29 ms, 79 from 29 to 73 ms, and 48 neurons from 72 to 100 ms. For two periods (0-13 ms and 29-73

ms), not only were more neurons activated, but most of the activated neurons were also selective to the

same orientation (Figure 5C). The firing rate of each activated neuron manifested a stronger preference

for GES in 0^o during the periods of 0-13 ms and 29-73 ms, but much less so for other periods (Figure 5D).

These results suggest that gastric neurons in NTS were selective to the orientation of GES at Target 1

with respect to their responses at two specific delays (about 8 ms and 43 ms) relative to the stimulus

onset. For GES at Target 2, the firing rate averaged across all activated neurons was higher when the

stimulus was in 270^o at nearly all post-stimulus latencies (Figure 5F). The peak responses were noticeable

in two periods (0-13 ms and 29-73 ms). In particular, the period from 29 to 73 ms showed that 77 out of

119 neurons were selective to the same orientation (270^o), whereas slightly lower but comparable

orientation selectivity was also noticeable for other periods. These results suggest that neurons activated

by GES at Target 2 showed consistent selectivity to 270^o for all post-stimulus periods, showing a lesser

degree of time dependence compared to the responses to GES at Target 1. In summary, when Target 2 is

stimulated, more NTS neurons prefer the orientation of 270^o within 29-73ms after the stimulus.

Compare neural responses after the stimulus on different targets

We further asked whether stimulating different gastric locations could activate the same neurons within

NTS, and if yes, whether their responses were distinctive and dependent on the stimulus location. To

answer these questions, we implanted electrodes to both Target 1 and Target 2 in four rats (Figure 6A).

When applying 1-mA current stimuli in the preferred orientation (0^o for Target 1 or 270^o for Target 2), we

identified the single units that responded to either or both of the locations.

In total, 100 neurons were identified: 88 with Target 1, 63 with Target 2, and 51 with both. On average,

the single-unit firing rate per 100 ms was 1.72 ± 0.15 (mean $\pm$ standard error) for Target 1 and 0.91 ± 0.10

for Target 2. For the 51 neurons responsive to both targets, their average firing rate was 2.12 ± 0.21 for

Target 1 and 0.92 ± 0.10 for Target 2. The time-averaged neuronal responses were significantly higher for

Target 1 than for Target 2 (paired t-test: t=7.35, p=1.7e-9, dof=50). Only seven neurons showed higher

responses to Target 2, whereas the majority showed higher responses to Target 1 (Figure 6C).

We further resolved the single-unit firing rate in time and averaged the response across the 51 neurons

that were responsive to both targets. Figure 6B shows the firing of a single neuron across repeated stimuli.

In this example, the neuron fired at different times in response to the stimulus at Target 1 vs. Target 2.

Activation with Target 1 was mostly between 30 and 60 ms, whereas activation with Target 2 was later,

especially around 80 ms (Figure 6B). We averaged the firing rate across neurons, and found that the

average neuronal response was higher for Target 1 for the first 60 ms, whereas it was higher for Target 2

between 65 and 100 ms (Figure 6D).

These results suggest that gastric electrical stimulation selectively applied to longitudinal or circular

IMAs at different locations may activate a common group of neurons in NTS but at different times. These

neurons are more responsive to stimulation applied to the longitudinal IMAs on the greater curvature at

early times (before 60 ms) but are more responsive to the circular IMAs at later times (after 65 ms).

Discussions

In summary, this paper presents neurophysiological evidence that gastric neurons in NTS are highly

selective to the orientation of gastric electrical stimulation applied to the hotspots of longitudinal and

circular IMAs in the distal forestomach. To activate NTS, it is preferable to stimulate the longitudinal

IMAs by delivering current pulses to the greater curvature in the longitudinal direction towards the

proximal stomach, or to stimulate the circular IMAs by delivering current pulses in between the lesser and

greater curvatures in the circular direction towards the greater curvature. This orientation preference is

largely generalizable across different stimulus amplitudes (from 0.6 to 1 mA) and dependent on the post-

stimulus latency. Further, stimulating longitudinal and circular IMA may activate a shared subset of

neurons in NTS, although the activated neurons are relatively more responsive to stimulation of

longitudinal IMAs. These findings suggest that gastric neurons in NTS, at least in part, encode

orientation-specific muscle activity relayed through vagal afferent neurons that innervate gastric muscle

layers through topologically distinctive and orientation selective intramuscular arrays. The scientific and

clinical implications of our findings are discussed as below.

The brain sees the gut as a sensory organ (Furness et al., 2013). The primary function of the stomach lies

in its motor events effectuated by smooth muscle cells in the gastric wall. The number of gastric smooth

muscle cells is orders of magnitude greater than the number of sensory neurons that relay muscle activity

to the brain (Powley et al., 2019). The mismatch prevents sensory neurons from uniformly innervating the

stomach in a one-to-one manner and necessitates "sparse coding" for efficient gastric representations in

NTS. Our results provide functional evidence that NTS encodes and integrates orientation specific muscle

activity from the stomach.

Because the stomach moves primarily along two orientations (longitudinal and circular), it makes

intuitive sense for NTS to mainly represent longitudinal and circular movements while neglecting other

orientations that are less functionally significant. Such orientation selectivity is enabled by highly

specialized morphologies of the afferent nerve terminals (i.e., IMAs) that selectively target longitudinal

and circular muscles (Wang and Powley, 2000). As they resemble elongated and parallel arrays, IMAs are

uniquely suited to serve as the mechanoreceptors for orientation selective neural coding (Powley and

Phillips, 2002).

In addition, the non-uniform distributions of circular and longitudinal IMAs (Powley et al., 2016) may

further support sparse coding. Muscle activity is coordinated across regions on the stomach (Lu et al.,

2017). Muscle cells are actively coupled to and paced by interstitial cells of Cajal (ICC), and intrinsically

modulated by enteric neurons (Furness et al., 2020). Therefore, spatially uniform sampling of muscle

activity would otherwise be redundant, rather than sparse. The fact that the IMAs show regional

"hotspots" allows the brain to sample only a few regions while likely being able to infer about other

regions by reasonable extrapolation given the stomach's intrinsic organization.

In general, electrical current stimulation is not cell type specific and thus possible to activate other cells,

axons, or receptors other than IMAs. Although we did not fully rule out such possibilities, the activation

of IMAs is the primary source that drives the orientation selective neural responses at NTS. The current

pulses used in this study were too brief and weak to activate smooth muscles. Prior studies suggest that a

current pulse should be at least 350 ms and 2 to 4 mA in order to directly cause muscle activation (Du et

al., 2009; Hocking et al., 1992; Li & Chen, 2010; McCallum et al., 1998; Tomita, 1966), whereas we used

a current pulse of 0.3 ms and 0.6 to 1 mA. In addition, the longitudinal and circular smooth muscles are

present in both Target 1 and 2, unlike the differentiated distributions of longitudinal and circular IMAs. If

the stimuli were to directly activate muscles, they would have activated both longitudinal and circular

muscles and thus could not explain the differential orientation selectivity for GES applied to the two

targets. Arguably, the current pulses might have directly activated the axons of vagal afferents. However,

the axonal activation is unlikely to explain the observed NTS responses. The axons of vagal afferents run

towards the esophagus and ascend to the brain. Direct activation of those axons would prefer the current

flowing towards the lesser curvature. This was opposite to our observation that neurons in NTS preferred

the current at Target 2 to flow towards the greater curvature. If the stimulation were applied primarily to

the axons, the effects would be more likely to block the ascending activation. Lastly, the current stimuli

could also have activated other afferent receptors, e.g., intra-ganglionic laminar endings (IGLE). The

morphology of IGLE is like a plate of lamelliform terminal puncta in contact with the ganglion between

longitudinal and circular muscle layers (Powley et al., 2019). Such a structure is unlikely to transmit

orientation specific signals. Also note that the preferred orientation varies between the two target

locations, in agreement with the different regional distributions of IMAs (Powley et al., 2016), but not

the distribution of IGLE (Tan et al., 2021).

Results from this study also suggest that neurons in NTS integrate sensory information from different

gastric regions. A single neuron could be activated with stimulation applied to different targets and in

different orientations. Such a neuron thus has a broad receptive field and a multitude of orientation tuning.

Speculatively, the integrative function may arise from second-order neurons that receive converging

projections of vagal afferent neurons, as well as higher-order neurons in NTS (Powley, 2000; Whitehead,

1988). The dynamic interplay among different subnuclei may likely account for single-unit responses at

different latencies (Figure 6). Our speculation agrees with the current understanding that individual

subnuclei in NTS do not form a fine-grained topographic map of the stomach, but progressively integrates

sensory input (Powley, 2000; Rogers et al., 1995; Travagli & Anselmi, 2016).

The orientation selectivity of neurons in NTS is likely to have a profound implication to where and how

GES should be applied. To the best of our knowledge, prior studies on characterizing the effects of GES

with various parameters have never addressed the effects of the stimulus orientation. In clinic practice,

GES is applied to the corpus or antrum (Abell et al., 2003; Hasler, 2009). This choice was perhaps

historical, dating back to the initial attempt of using direct currents to pace gastric contractions (Kelly &

La Force, 1972). While gastric pacing has not been widely adopted (Hasler, 2009; Yin, & Chen, 2008),

the primary goal of GES has focused on modulating neural circuits that control the stomach. However,

clinical applications with GES largely or entirely ignore the morphological, topological or chemical

phenotypes of different kinds of nerve terminals.

Results from this study suggest that the orientation is a key parameter for GES, likely more important

than other parameters, e.g., pulse width, amplitude, or frequency. For GES to activate NTS, two effective

locations are in the distal forestomach: one on the greater curvature and the other in between the lesser

and greater curvatures, as informed by the regional distributions of IMAs. For the former, the optimal

orientation is the longitudinal direction towards the distal stomach. For the latter, the optimal orientation

is the circular direction towards the greater curvature. GES applied with such optimal locations and

orientations may trigger a cascade of integrative processing within and beyond NTS, and likely give rise

to therapeutic benefits. This expected outcome remains speculative and awaits confirmation in future

preclinical or clinical studies.

Acknowledgment

The authors thank Dr. Robert Phillips for his guidance on the electrode implementation for gastric

electrical stimulation, thank Drs. Zhenjun Tan, Kun-Han Lu for initial discussions in this project, and

thank Drs. Mark Sayles and Xueguo Zhang for their initial assistance in the electrophysiological

recording. This study is supported by NIH grants (OD023847, OD030538), the University of Michigan,

and Purdue University.

Table 1 The number and orientation of the implanted electrodes in each animal. The number of electrodes

is specified with respect to the location (Target 1 vs. 2) and the direction (longitudinal vs. circular).

Animal ID	Target 1		Target 2	
	Longitudinal	Circular	Longitudinal	Circular
1	2	2	0	0
2	2	2	0	0
3	2	2	0	0
4	2	2	2	2
5	2	0	2	2
6	2	0	2	2
7	2	0	2	2

455 **Table 2** Statistical difference in the mean population firing rate (i.e., the number of spikes per 100 ms)

456 given GES applied to Target 1 with a preferred orientation (0^o) vs. other orientations (270^o, 180^o, 90^o).

457 Each row corresponds to a different current amplitude. The statistics below are based on paired t-tests

458 with Bonferroni correction for multiple comparisons.

current	Firing rate at 0^o	0^o vs 270^o	0^o vs 180^o	0^o vs 90^o
0.6mA	2.91±0.42 /100ms	t(91)=3.48; p=0.0023	t(91)=0.54; p=0.59	t(91)=3.56; p=0.0018
0.8mA	3.51±0.45 /100ms	t(92)=4.74; p=2.3e-5	t(92)=7.26; p=3.7e-10	t(92)=6.32; p=2.8e-8
1mA	4.09±0.50 /100ms	t(92)=3.82; p=7.2e-4	t(92)=6.93; p=1.7e-9	t(92)=7.63; p=6.2e-11

462 **Table 3** Statistical difference in the mean population firing rate (i.e., the number of spikes per 100 ms)

463 given GES applied to Target 2 with a preferred orientation (270^o) vs. other orientations (0^o, 180^o, 90^o).

464 Each row corresponds to a different current amplitude. The statistics below are based on paired t-tests

465 with Bonferroni correction for multiple comparisons.

current	Firing rate at 270^o	270^o vs 0^o	270^o vs 180^o	270^o vs 90^o
0.6mA	2.60±0.34 / 100ms	t(71)=4.62; p=5.0e-5	t(71)=4.60; p=5.4e-5	t(71)=4.57; p=6.0e-5
0.8mA	3.72±0.40 / 100ms	t(71)=7.19; p=1.6e-9	t(71)=6.98; p=3.9e-9	t(71)=7.45; p=8.3e-10
1mA	3.83±0.39 / 100ms	t(93)=7.49; p=1.2e-10	t(93)=6.39; p=1.9e-8	t(93)=7.86; p=2e-11

    Figure Caption

    Figure 1 Experiment protocol and analysis pipeline. A shows the experimental setup and the location

    of the implanted electrodes for gastric electrical stimulation. **B** shows the distributions and morphology of

    IMAs. The distributions of longitudinal and circular IMAs are shown in the left and right respectively

    (Tan et al., 2021). The middle shows an example of a parent axon branching out to parallel arrays of

    longitudinal IMAs. **C** illustrates the position of the silicon probe inserted into the NTS relative to the area

    postrema (AP) and Obex in the axial and coronal view, respectively. **D** shows an example time series of

    MUA (bottom) during a train of alternating current pulses (top) applied to Target 1 in the longitudinal

    direction. Positive pulses deliver the current toward the distal stomach. Negative pulses deliver the current

    toward the proximal stomach. The population spikes are marked with red stars. **E** shows the zoom-in of

    the same time series within 100 ms following a single negative pulse. The stimulus onset is at time 0. **F**

    shows the feature representations of individual spikes. Each dot is a spike described in terms of two spike

    features, which are illustrated in the red plot in the bottom-right inset. Two clusters of spikes are

    highlighted in yellow and blue. Each cluster corresponds to a single neuron. **G** shows the individual

    spikes (in grey) that belong to each cluster and their average within the cluster (in red). The boundary of

    the top or bottom plot is color-matched to the blue or yellow cluster, respectively. **H** shows the post-

    stimulus time histogram of the firing rate (the average number of spikes per 5 ms) for each of the two

    neurons sorted in **F** and **G**.

    Figure 2 GES-evoked population spikes in NTS given different stimulus orientations. A and **D** show

    the locations of GES in comparison with the digitized samples of longitudinal and circular IMA based on

    prior work (Powley et al., 2016; Tan et al., 2021). **B** and **E** show the pairing of the anode (red) and the

    cathode (black) for delivering a current pulse in one of four different directions (0^{o}, 270^{o}, 180^{o}, and 90^{o}).

    In **B** (or **E**), the raster displays the timing of the population spikes observed from one channel in response

    to a single current pulse across repeated trials. Each black dot marks the time of a spike. Each row

indicates a single trial. Time 0 is the stimulus onset. Data in **B** and **E** are from different channels. **C** (or **F**)

plots the average number of spikes during the (100 ms) post-stimulus period given each orientation, after

averaging across trials. The error bar indicates the standard error of the mean. ** means $p<0.001$ and ***

means $p<0.0001$ for paired t-test with Bonferroni correction.

Figure 3 Population responses and orientation selectivity in NTS given GES with different current

amplitudes. **A** (or **D**) shows the electrodes implanted to Target 1 (or 2). **B** (or **E**) shows the mean firing

rate in response to GES at Target 1 (or 2) in each of the four orientations with the current amplitude at

0.6, 0.8, or 1 mA. The error bar indicates the standard error of the mean. **C** (or **F**) shows the orientation

selectivity index (OSI) given different current amplitudes for Target 1 (or 2). Grey dots are individual

samples. ** means p<0.005 and *** means p<0.001 (permutation test with Bonferroni correction).

Figure 4 Orientation selectivity for single neurons in NTS. For 1-mA GES at Target 1 (**A**) and Target

2 (**F**), **B** and **G** show the firing of a neuron (right) in response to a current pulse delivered in four different

orientations (left). The raster displays the timing of SUA across repeated trials. Each dot marks the time

of a spike. Each row is one stimulus trial. Time 0 is the stimulus onset. Data shown in **B** and **G** are from

different neurons. **C** and **H** plot the average firing rate in terms of the number of spikes per 100 ms given

GES in different orientations. The error bars represent the standard errors of the mean. **D** and **I** show the

box plots of the orientation selectivity index (OSI) for neurons selective to each of the four orientations.

Grey dots represent individual neurons. **E** and **J** show the orientation tuning for neurons selective to each

of the four orientations. The radii of the concentric circles indicate different firing rates: 0.5, 1, and 1.5

spikes per 100 ms. The black lines connect the firing rates in response to GES with different orientations.

Results shown in **A** through **E** correspond to GES at Target 1. Those in **F** through **J** correspond to GES at

Target 2.

Figure 5 Time dependence of orientation selectivity. For 1-mA GES at Target 1 (**A**) and Target 2 (**E**),

B and **F** plot the single-unit post-stimulus time histogram (PSTH) averaged across neurons with

stimulation applied in 0^o, 270^o, 180^o, and 90^o. **C** and **G** show the number of neurons that preferred each of

the four orientations in terms of their responses at four different post-stimulus periods. **D** and **H** show the

number of spikes (coded as the radius) observed during each of the four periods (color-coded as the

circles) given GES in each of the four orientations (coded as the angle). Regardless of the stimulus

orientation, the total number of neurons activated during each period is denoted as n.

Figure 6 Location dependence of single-unit responses. A shows 1-mA GES applied in the preferred

orientations for both of the two target locations. In **B**, the left shows the raster displays of the same

neuron's firing pattern in response to repeated trials of GES at Target 1 (top) and Target 2 (bottom); the

right shows the corresponding PSTH. **C** plots the relative changes in the time-averaged single-unit

responses given GES at Target 1 vs. Target 2. Each line corresponds to a single neuron. For each neuron,

the time-average firing rate averaged across trials is always normalized to 1, regardless of whether it is

greater when stimulation is at Target 1 or Target 2. The corresponding y-axis is on the left. The grey bar

plot shows the averaged firing rate, with the y-axis on the right. **D** shows the PSTH averaged across

neurons given GES at either Target 1 (black) or Target 2 (gray). The black (or gray) dots highlight the

time points where the response to GES at Target 1 (or Target 2) is significantly higher than the response

to GES at Target 2 (or Target 1) based on paired t-tests separately evaluated for each time point with $\square$ =

0.05.

Reference

Abell, T., McCallum, R., Hocking, M., Koch, K., Abrahamsson, H., LeBlanc, I., ... & Starkebaum, W.

(2003). Gastric electrical stimulation for medically refractory gastroparesis. *Gastroenterology*, *125*(2),

421-428.

Andermann, M. L., & Lowell, B. B. (2017). Toward a wiring diagram understanding of appetite control.

Neuron, *95*(4), 757-778.

Babic, T., & Browning, K. N. (2014). The role of vagal neurocircuits in the regulation of nausea and

vomiting. *European journal of pharmacology*, *722*, 38-47.

Berthoud, H. R., & Powley, T. L. (1992). Vagal afferent innervation of the rat fundic stomach:

morphological characterization of the gastric tension receptor. *Journal of Comparative Neurology*,

319(2), 261-276.

Berthoud, H. R., Patterson, L. M., Neumann, F., & Neuhuber, W. L. (1997). Distribution and structure of

vagal afferent intraganglionic laminar endings (IGLEs) in the rat gastrointestinal tract. *Anatomy and*

embryology, *195*(2), 183-191.

Brookes, S. J., Spencer, N. J., Costa, M., & Zagorodnyuk, V. P. (2013). Extrinsic primary afferent

signalling in the gut. *Nature reviews Gastroenterology & hepatology*, *10*(5), 286-296.

Browning, K. N., & Travagli, R. A. (2011). Central nervous system control of gastrointestinal motility

and secretion and modulation of gastrointestinal functions. *Comprehensive physiology*, *4*(4), 1339-1368.

Carabotti, M., Scirocco, A., Maselli, M. A., & Severi, C. (2015). The gut-brain axis: interactions between

enteric microbiota, central and enteric nervous systems. *Annals of gastroenterology: quarterly publication*

of the Hellenic Society of Gastroenterology, *28*(2), 203.

Carmagnola, S., Cantù, P., & Penagini, R. (2005). Mechanoreceptors of the proximal stomach and

perception of gastric distension. *American Journal of Gastroenterology*, *100*(8), 1704-1710.

Cheng, L.K., Nagahawatte, N.D., Avci, R., Du, P., Liu, Z., Paskaranandavadivel, N. (2021). Strategies to

refine gastric stimulation and pacing protocols: experimental and modeling approaches. *Frontiers in*

Neuroscience, 15:645472.

Davis, C. J., Harding, R. K., Leslie, R. A., & Andrews, P. L. R. (1986). The organisation of vomiting as a

protective reflex. In *Nausea and vomiting: Mechanisms and treatment* (pp. 65-75). Springer, Berlin,

Heidelberg.

De Ponti, F., Azpiroz, F., & Malagelada, J. R. (1987). Reflex gastric relaxation in response to distention

of the duodenum. *American Journal of Physiology-Gastrointestinal and Liver Physiology*, *252*(5), G595-

G601.

Du, P., Li, S., O'Grady, G., Cheng, L. K., Pullan, A. J., & Chen, J. D. (2009). Effects of electrical

stimulation on isolated rodent gastric smooth muscle cells evaluated via a joint computational simulation

and experimental approach. *American Journal of Physiology-Gastrointestinal and Liver Physiology*,

297(4), G672-G680.

Ehrlein, H. J., & Schemann, M. (2005). Gastrointestinal motility. *Technische Universität München,*

Munich.

Furness, J. B., Rivera, L. R., Cho, H-J, Bravo, D. M., & Callaghan, B (2013) The gut as a sensory organ.

Nature Reviews Gastroenterology and Hepatology. 10: 729-740.

Furness, J. B., & Stebbing, M. J. (2018). The first brain: Species comparisons and evolutionary

implications for the enteric and central nervous systems. *Neurogastroenterology & motility*, *30*(2),

e13234.

Furness, J. B., Di Natale, M., Hunne, B., Oparija-Rogenmozere, L., Ward, S. M., Sasse, K. C., Powley, T.

L. Stebbing, M. J., Jaffey D., & Fothergill, L. J. (2020). The identification of neuronal control pathways

supplying effector tissues in the stomach. *Cell and Tissue Research*, 1-13.

Forster, J., Sarosiek, I., Delcore, R., Lin, Z., Raju, G. S., & McCallum, R. W. (2001). Gastric pacing is a

new surgical treatment for gastroparesis. *The American journal of surgery*, *182*(6), 676-681.

Gottfried-Blackmore, A., Adler, E. P., Fernandez-Becker, N., Clarke, J., Habtezion, A., & Nguyen, L.

(2020). Open-label pilot study: non-invasive vagal nerve stimulation improves symptoms and gastric

emptying in patients with idiopathic gastroparesis. *Neurogastroenterology & Motility*, *32*(4), e13769.

Goyal, R. K., Guo, Y., & Mashimo, H. (2019). Advances in the physiology of gastric emptying.

Neurogastroenterology & Motility, *31*(4), e13546.

Hocking, M. P., Vogel, S. B., and Sninsky, C. A. (1992). Human gastric myoelectric activity and gastric

emptying following gastric surgery and with pacing. *Gastroenterology* 103, 1811–1816.

Holtmann, G., & Talley, N. J. (2014). The stomach–brain axis. *Best Practice & Research Clinical*

Gastroenterology, *28*(6), 967-979.

Lal, N., Livemore, S., Dunne, D., & Khan, I. (2015). Gastric electrical stimulation with the enterra

system: a systematic review. *Gastroenterology research and practice*, *2015*.

Levinthal, D. J., & Bielefeldt, K. (2017). Systematic review and meta-analysis: Gastric electrical

stimulation for gastroparesis. *Autonomic Neuroscience*, *202*, 45-55.

Levinthal, D. J., & Strick, P. L. (2020). Multiple areas of the cerebral cortex influence the stomach.

Proceedings of the National Academy of Sciences, *117*(23), 13078-13083.

Li, S., & Chen, J. D. (2010). Cellular effects of gastric electrical stimulation on antral smooth muscle cells

in rats. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, *298*(6),

R1580-R1587.

Lu, K-H, Cao, J., Oleson, S. T., Powley, T. L., Liu, Z. (2017) Contrast enhanced magnetic resonance

imaging of gastric emptying and motility in rats. *IEEE Transactions on Biomedical Engineering*, 64(11),

2546-2554.

Mayer, E. A. (2011). Gut feelings: the emerging biology of gut–brain communication. *Nature Reviews*

Neuroscience, *12*(8), 453-466.

Mayer, E. A., Tillisch, K., & Bradesi, S. (2006). modulation of the brain–gut axis as a therapeutic

approach in gastrointestinal disease. *Alimentary pharmacology & therapeutics*, *24*(6), 919-933.

McCallum, R. W., Chen, J. D. Z., Lin, Z., Schirmer, B. D., Williams, R. D., & Ross, R. A. (1998). Gastric

pacing improves emptying and symptoms in patients with gastroparesis. *Gastroenterology*, *114*(3), 456-

461.

McCallum, R. W., Dusing, R. W., Sarosiek, I., Cocjin, J., Forster, J., & Lin, Z. (2010). Mechanisms of

symptomatic improvement after gastric electrical stimulation in gastroparetic patients.

Neurogastroenterology & Motility, *22*(2), 161-e51.

McCallum, R. W., Snape, W., Brody, F., Wo, J., Parkman, H. P., & Nowak, T. (2010). Gastric electrical

stimulation with Enterra therapy improves symptoms from diabetic gastroparesis in a prospective study.

Clinical Gastroenterology and Hepatology, *8*(11), 947-954.

O'Grady, G., Egbuji, J. U., Du, P., Cheng, L. K., Pullan, A. J., & Windsor, J. A. (2009). High-frequency

gastric electrical stimulation for the treatment of gastroparesis: a meta-analysis. *World journal of surgery*,

33(8), 1693-1701.

Payne, S. C., Furness, J. B., & Stebbing, M. J. (2019). Bioelectric neuromodulation for gastrointestinal

disorders: effectiveness and mechanisms. *Nature reviews Gastroenterology & hepatology*, *16*(2), 89-105.

Powley, T. L. (2000). Vagal circuitry mediating cephalic-phase responses to food. *Appetite*, *34*(2), 184-

188.

Powley, T. L., & Phillips, R. J. (2002). I. Morphology and topography of vagal afferents innervating the

GI tract. *American Journal of Physiology-Gastrointestinal and Liver Physiology*, *283*(6), G1217-G1225.

Powley, T. L., Hudson, C. N., McAdams, J. L., Baronowsky, E. A., & Phillips, R. J. (2016). Vagal

intramuscular arrays: the specialized mechanoreceptor arbors that innervate the smooth muscle layers of

the stomach examined in the rat. *Journal of Comparative Neurology*, *524*(4), 713-737.

Powley, T.L., Jaffey, D.M., McAdams, T.A., Baronoswky, E.A., Black D., Chesney, L., Evans, C.,

Phillips, R.J. (2019). Vagal innervation of the stomach reassessed: brain-gut connectome uses smart

terminals. *Annals of The New York Academy of Sciences*, 1454(1): 14-30.

Powley, T. L., & Phillips, R. J. (2002). I. Morphology and topography of vagal afferents innervating the

GI tract. *American Journal of Physiology-Gastrointestinal and Liver Physiology*, *283*(6), G1217-G1225.

Powley, T. L., Spaulding, R. A., & Haglof, S. A. (2011). Vagal afferent innervation of the proximal

gastrointestinal tract mucosa: chemoreceptor and mechanoreceptor architecture. *Journal of Comparative*

Neurology, *519*(4), 644-660.

Phillips, R. J., & Powley, T. L. (2000). Tension and stretch receptors in gastrointestinal smooth muscle:

re-evaluating vagal mechanoreceptor electrophysiology. *Brain Research Reviews*, *34*(1-2), 1-26.

Qin, C., Sun, Y., Chen, J. D. Z., & Foreman, R. D. (2005). Gastric electrical stimulation modulates

neuronal activity in nucleus tractus solitarii in rats. *Autonomic Neuroscience*, *119*(1), 1-8.

Rogers, R. C., McTIGUE, D. M., & Hermann, G. E. (1995). Vagovagal reflex control of digestion:

afferent modulation by neural and endoneurocrine factors. *American Journal of Physiology-*

Gastrointestinal and Liver Physiology, *268*(1), G1-G10.

Tan, Z.T., Ward, M.P., Phillips, R.J., Zhang, X., Jaffey, D.M., Chesney, L., Rajwa, B., Baronowsky,

E.A., McAdam, J., Powley, T.L. (2021) Stomach region stimulated determines effects on duodenal

motility in rats. *American Journal of Physiology, Regulatory, Integrative and Comparative Physiology*,

320(3): R331-R341.

Tang, M., Zhang, J., & Chen, J. D. Z. (2006). Central mechanisms of gastric electrical stimulation

involving neurons in the paraventricular nucleus of the hypothalamus in rats. *Obesity surgery*, *16*(3), 344-

352.

Tomita, T. (1966). Electrical responses of smooth muscle to external stimulation in hypertonic solution.

The Journal of physiology, *183*(2), 450-468.

Travagli, R. A., & Anselmi, L. (2016). Vagal neurocircuitry and its influence on gastric motility. *Nature*

Reviews Gastroenterology & Hepatology, *13*(7), 389.

Travagli, R. A., Hermann, G. E., Browning, K. N., & Rogers, R. C. (2006). Brainstem circuits regulating

gastric function. *Annu. Rev. Physiol.*, *68*, 279-305.

U.S. Food and Drug Administration, Center for Devices and Radiological Health. Gastric Electrical

Stimulation (GES) System H990014 approval letter, March 31, 2000. Retrieved March 29, 2021, from

https://www.accessdata.fda.gov/cdrh_docs/pdf/H990014A.pdf.

Van Oudenhove, L., Demyttenaere, K., Tack, J., & Aziz, Q. (2004). Central nervous system involvement

in functional gastrointestinal disorders. *Best Practice & Research Clinical Gastroenterology*, *18*(4), 663-

680.

Wang, F. B., & Powley, T. L. (2000). Topographic inventories of vagal afferents in gastrointestinal

muscle. *Journal of Comparative Neurology*, *421*(3), 302-324.

Whitehead, M. C. (1988). Neuronal architecture of the nucleus of the solitary tract in the hamster. *Journal*

of Comparative Neurology, *276*(4), 547-572.

Williams, E. K., Chang, R. B., Strochlic, D. E., Umans, B. D., Lowell, B. B., & Liberles, S. D. (2016).

Sensory neurons that detect stretch and nutrients in the digestive system. *Cell*, *166*(1), 209-221.

A

Antrum Forestomach

Corpus

Target 2

Target 1 Electrical stimulator

B

Longitudinal IMAs   Circular IMAs

high   high

1 mm

low   low

C

Axial left   Recording electrode

0.5-1.3 mm

AP   AP

NTS   200um

Obex

Recoding site

0-0.8 mm   Coronal
Left

D

Stimuli

Voltage (uV)

400
200
0
-200
-400

0   0.1   0.2   0.3   0.4   0.5   0.6   0.7   0.8   0.9   1
Time (s)

E

Stimulus

Voltage (uV)

400
200
0
-200
-400

0   10   20   30   40   50   60   70   80   90   100
Time (ms)

F

Spikes in feature domain

Feature 2

0   200   400   600   800
Feature 1

Feature 1   Feature 2

G

Voltage (uV)

400
200
0
-200
-400

0.2   0.4   0.6

Voltage (uV)

400
200
0
-200
-400

0.2   0.4   0.6
Time (ms)

H

PSTH

Number of spikes /5ms

0.8
0.4
0

0   20   40   60   80   100

0.8
0.4
0

0   20   40   60   80   100
Time (ms)

Spike train of MUA

B

0°

270°

180°

90°

Repetitions

Time (ms)

C

**p<0.001; ***p<0.0001
(Bonferroni corrected)

Number of spikes /100 ms

E

0°

270°

180°

90°

Repetitions

Time (ms)

F

***p<0.0001
(Bonferroni corrected)

Number of spikes /100 ms

Target 1

Target 2